Grant# DE-FG05-87ER13739

DE93 000676

James N. Siedow, Department of Botany, Duke University
Molecular Studies of Functional Aspects of Plant Mitochondria

The initial (two-year) funding period for this grant runs from August, 1990 through July, 1992. Manuscripts derived from this period are listed at the end of this report. The goal of this research, as outlined in the original proposal to DOE, is to characterize the mechanism by which a protein encoded by the mitochondrial genome of *cms*-T maize (URF13) interacts with a family of compounds produced by certain fungi (T-toxins) to permeabilize biological membranes. The research carried out in the funding period covered by this report as well as that which will be carried out in the upcoming budget period will continue to follow lines of investigation outlined in the original DOE proposal.

Research Progress, 8/90-present:

Results obtained with previous DOE support had indicated that cloning and expressing the gene encoding URF13 into *E. coli* confers T-toxin sensitivity on the resulting bacterial cells (1). Further, site-directed mutagenesis of URF13 expressed in *E. coli* led to 1) the identification of a specific residue, Asp-39, that appears to be essential for maintaining the T-toxin sensitivity of URF13 and 2) the demonstration that truncation of the normal 115 amino acid-containing URF13 molecule to a species containing less than 83 amino acids resulted in the loss of T-toxin sensitivity (2,3).

Because URF13 is an integral membrane protein that is capable of generating a hydrophilic pore across biological membranes following interaction with T-toxin, it was important to understand the relationship between the structural organization of URF13 in the membrane and its function (i.e., its ability to permeabilize the membrane). Theoretical considerations of the structural features of URF13 based upon its derived amino acid sequence (specifically, hydrophobicity and hydrophobic moment profiles) in conjunction with the experimental results obtained from the directed mutagenesis studies (2) led to the development of a model of the structure of the URF13 monomer within the membrane (4). In this model, URF13 is postulated to contain three membrane-spanning alpha-helical domains (Fig. 1), with helix I consisting of a hydrophobic membrane-spanning region and helices II and III forming a pair of amphipathic alpha-helices. Pore formation following interaction between URF13 and T-toxin requires the alignment of 6 to 8 amphipathic helices which suggests that the functional form of URF13 within the membrane is an oligomeric structure composed of at least three to four URF13 monomers.

Fig. 1. Proposed model of URF13 disposition within the membrane, indicating the location of the three putative membrane-spanning helical domains. Asp(D)-39 (site of DCCD reaction) is shaded.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

J.N. Siedow, Cont:
DE-FG05-87ER13739

A major focus of the research carried out over the funding period covered in this report involved attempts to test the three-helical/oligomeric model of URF13. To that end, an URF13 fusion protein was constructed that contained 14 amino acids, linked to the amino terminus of URF13, derived from the phage T7 sl0 coat protein that contained a 10-amino acid epitope to a commercially available antibody. Given that an antibody to a 15-amino acid region in the carboxy-terminus of URF13 had already been generated, the fusion protein possessed antigenic tags at either end of the URF13 molecule. Protease digestion studies were then used to determine the accessibility of the amino and carboxyl termini of this URF13 fusion protein in E. coli membrane preparations having either right-side-out or inside-out orientations (5). Consistent with the three-helical model, the amino and carboxyl ends of URF13 were found to be localized on opposite sides of the membrane. Somewhat surprisingly, these studies also indicated that URF13 has two topological orientations in the E. coli plasma membrane, with 60-75% of the carboxy termini oriented toward the cytoplasmic surface and the remaining 25-40% localized on the outer, periplasmic surface of the membrane. This dual orientation was not unique to the fusion protein; the carboxy terminus of standard URF13 expressed in E. coli showed a similar mixed orientation. Interestingly, similar protease studies using right-side-out and inside-out membrane preparations from isolated cms-T maize mitochondria indicated that URF13 has only one orientation in mitochondria, with the carboxy terminus located exclusively on the matrix side of the inner mitochondrial membrane (which, electrochemically, corresponds to the preferred orientation in E. coli).

The possibility that URF13 might exist in the membrane in an oligomeric state was first suggested from the results of studies looking at the binding of radiolabeled (³H) T-toxin to E. coli cells expressing URF13 (6). The binding of T-toxin to URF13 was found to be specific and to display a marked cooperativity, having a Hill coefficient that varied between 1.3 and 1.7 from one preparation to the next. To test more directly for the presence of URF13 oligomers, studies were carried out using a lysine-specific, hydrophobic bifunctional cross-linking reagent (5). These studies directly confirmed that a portion of the URF13 molecules in both E. coli and cms-T mitochondria exist in the membrane in an oligomeric state. With the lysine-specific cross-linking reagent, dimers and trimers appear predominantly, but more recent studies (See below) indicate that there is considerable tetramer formation and even some pentamers and hexamers. Addition of T-toxin did not change the oligomeric pattern seen following cross-linking, indicating that the changes in URF13 structure associated with pore formation are more subtle than the simple association of URF13 monomers (or even dimers) into higher-order structures. When cross-linking studies were carried out in conjunction with proteolysis experiments on the URF13 fusion protein, the results indicated that in any given multimeric complex, all the participating monomers have the same orientation within the membrane (5).

In addition to establishing the cooperative (and specific) nature of the binding of T-toxin to URF13, the binding studies with radiolabeled T-toxin demonstrated that cells expressing the directed mutations of URF13 lacking

J.N. Siedow, Cont:
DE-FG05-87ER13739

toxin sensitivity (the truncated 82-amino acid species and a substitution of Val for Asp at residue 39) also showed reduced levels of T-toxin binding compared to cells expressing unmodified URF13 (6). Current research along this line involves studies being carried out to better establish the reason for the lowered T-toxin binding in these (and other, See below) mutants of URF13. A summary of the current understanding of the structure of URF13 and its interaction with T-toxin, as well as the molecular basis of disease susceptibility in cms-T maize was recently put together by the PI (7).

Current Unpublished Research:

As a followup to the cross-linking and toxin binding studies described above, we began to look at how the structure of URF13 was affected by treatment with the carboxylate-specific reagent, dicyclohexylcarbodiimide (DCCD). Previous studies had shown that addition of DCCD to isolated cms-T mitochondria led to a loss of sensitivity to added T-toxin [Holden, M.J. & H. Sze (1989) Plant Physiol. 91: 1296-1302], and we had obtained analogous results following DCCD treatment of *E. coli* cells expressing URF13 (2). Lysine-specific cross-linking of DCCD treated cells initially suggested that such treatment led to an enhanced formation of higher-order (trimer and tetramer) oligomers, but subsequent control experiments established that DCCD alone was capable of cross-linking URF13, and did so more effectively than the lysine-specific cross-linking reagent. Since the earlier mutagenesis studies had established that Asp-39 was the residue that reacts with DCCD (2), these results suggest that Asp-39 is directly involved in monomer:monomer interactions (probably by binding to a residue on an adjacent monomer via H-bonding or, less likely, an ionic interaction) in the URF13 oligomeric complex. As expected, toxin-insensitive mutations that substitute for the aspartate residue at position 39 (e.g., val but not glu) lead to a loss of cross-linking in the presence of added DCCD. Less expected, but of marked interest, was the observation that Val-39 also shows a loss of cross-linking with the lysine-specific reagent (i.e., this mutation has led to a loss of ability of URF13 to oligomerize within the membrane). Interestingly, Glu-39, which is also toxin-insensitive, continues to show the presence of higher-order oligomers with either DCCD or the lysine-specific cross-linker. In aggregate, these results clearly implicate residue 39 as having a role in oligomer formation in URF13. It will remain for future studies to establish the amino acid to which Asp-39 becomes linked following reaction with DCCD. Among other things, it should help to further define the structure model of URF13 in the membrane since the nature of the bond formed between a DCCD-activated carboxylate residue and a displacing nucleophile (i.e., amides and esters being the two most likely), requires that there be a reasonably close proximity between Asp-39 and its reaction partner.

Of further interest in this line of investigation is the effect of DCCD on T-toxin binding to URF13. Two changes take place in the observed binding pattern: 1) there is a complete loss of cooperativity, the binding becomes strictly hyperbolic and 2) the level of toxin binding to URF13 decreases by a

J.N. Siedow, Cont:
DE-FG05-87ER13739

factor of almost exactly two. Interestingly, the intrinsic K_d for the binding of T-toxin to URF13 is not affected by DCCD treatment. Taken together, these results appear to indicate that DCCD cross-linking locks the oligomeric URF13 complexes into a "closed" conformation that is not able to undergo a change to the "open," pore-forming conformation following binding of T-toxin. Less clear is whether this failure to undergo the conformational change to the open state leads to a loss of half the toxin binding sites or whether the failure to bind the second complement of T-toxin leads to the loss of the conformational change that causes pore formation. Current studies are attempting to address this issue. The studies outlined above looking at the effects of DCCD on URF13 are presently being written up and should be submitted for publication by the end of May (C.I. Kaspí and J.N. Siedow, manuscript in preparation).

Future Research:

In conjunction with the DCCD studies outlined above, we are presently carrying out a series of experiments looking at DCCD cross-linking in an attempt to establish a relationship between the observed cross-linking pattern and the loss of URF13 sensitivity to T-toxin. Such experiments include studies varying both the concentration of DCCD used to cross-link URF13 as well as following the time course of cross-linking in the presence of a constant concentration of DCCD. Developmental studies of the relationship between the appearance of toxin sensitivity and higher-order oligomers following the induction of URF13 in *E. coli* are also being carried out. While correlative in nature, preliminary results of these studies point to trimers and tetramers as being the molecular species associated with the appearance of T-toxin sensitivity (i.e., monomers and dimers are apparently not effective in pore formation). This work will continue in the upcoming funding period.

Additional studies related to this project that are presently not far enough along to warrant detailed discussion in this document but which will be the focus of much of our efforts in the coming year include the following: 1) Attempts to purify URF13 for use in direct physical studies of its structure (e.g., CD spectroscopy and, ultimately, NMR spectroscopy) by standard membrane protein isolation/purification techniques have not been successful, primarily due to the very low levels of URF13 protein expressed in *E. coli* (or *cms*-T mitochondria for that matter). In an attempt to overcome this problem, we are taking advantage of the use of amino-terminal specific fusions to URF13 that will, hopefully, allow us to either undertake affinity purification of the URF13 (using a polyhistidine fusion peptide coupled to purification on a Ni-containing affinity column) or change the level and nature of expression of URF13 to the point that inclusion bodies will result (using an amino-terminal fusion of glutathione-S-transferase). To date, the protein with the polyhistidine amino terminus has been constructed, and it does produce toxin-sensitive *E. coli* cells. 2) Characterization of the numerous toxin-insensitive mutants of URF13 generated by G. Ward in the Levings' laboratory through the use of saturation mutagenesis of URF13 will be carried out during

J.N. Siedow, Cont:
DE-FG05-87ER13739

this period. We are presently screening these toxin-insensitive mutants for their ability to bind labeled T-toxin. Interestingly, most of the toxin-insensitive mutations obtained to date do not bind toxin. We will carry out detailed, quantitative analyses of the T-toxin binding properties of those mutants that do bind toxin, specifically to characterize the level of toxin binding and to ascertain whether cooperativity of T-toxin binding is maintained or not in these mutations.

Publications (Resulting from DOE sponsored research)

Publications from funding prior to 8/90:

1. Dewey, R.E., J.N. Siedow, D.H. Timothy and C.S. Levings, III (1988) A 13-kilodalton maize mitochondrial protein in *E. coli* confers sensitivity to *Bipolaris maydis* toxin. *Science* 239: 293-295.
2. Braun, C.J., J.N. Siedow, M.E. Williams and C.S. Levings, III (1989) Mutations in the maize mitochondrial T-urf13 eliminate sensitivity to a fungal pathogen. *Proc. Natl. Acad. Sci. USA* 86: 4435-4439.
3. Braun, C.J., J.N. Siedow and C.S. Levings, III (1989) The T-urf13 gene is responsible for toxin sensitivity in maize and *E. coli*. In: "The Molecular Basis of Plant Development" (R.B. Goldberg, Ed.) pp. 79-85. Alan R. Liss, New York.

Publications from 8/90 to present (copies included with this report):

4. Korth, K.L., F. Struck, C.I. Kaspi, J.N. Siedow and C.S. Levings, III (1991) Topological orientation of the membrane protein URF13. In "Plant Molecular Biology," (R.G. Herrmann and B.A. Larkins, Eds.) pp. 375-381. Plenum Press, London.
5. Korth, K.L., C.I. Kaspi, J.N. Siedow and C.S. Levings, III (1991) URF13, a maize mitochondrial pore-forming protein, is oligomeric and has a mixed orientation in *E. coli* plasma membranes. *Proc. Natl. Acad. Sci. USA* 88: 10865-10869.
6. Braun, C.J., J.N. Siedow and C.S. Levings, III (1990) Fungal toxins bind to the URF13 protein in maize mitochondria and *E. coli*. *The Plant Cell* 2: 153-161.
7. Levings, C.S. III and J.N. Siedow (1992) Molecular basis of disease susceptibility in the Texas cytoplasm of maize. *Plant Mol. Biol.* - In Press.

PROGRESS REPORT

Grant# DE-FG05-87ER13739 - Covering the period 8/1/90 to present

The experiments outlined in this proposal are part of a collaborative effort between my laboratory and that of C. S. Levings III (Department of Genetics, North Carolina State University) to understand the origin of the unique susceptibility of cms-T maize to infection by the fungi Bipolaris maydis race T and Phyllosticta maydis. One attractive feature of this collaboration is that it combines my expertise in the study of membrane bioenergetics and membrane proteins with Levings' considerable experience in molecular genetics.

Introduction:

Following the appearance of the Southern corn leaf blight in 1969-70, a number of laboratories looked at the relationship between the cms-T genotype in maize and its sensitivity to a set of pathotoxins (collectively referred to as T-toxins) produced by the fungus responsible for the leaf blight. This research established that T-toxins specifically affect cms-T maize mitochondria, and the nature of the effects of the toxins on cms-T mitochondria (inhibition of malate oxidation, uncoupling of oxidative phosphorylation and the leakage of ions and molecules such as NAD⁺) led to the concept that the interaction of T-toxins with cms-T mitochondria results in a permeabilization of the inner mitochondrial membrane (Reviewed in Levings and Siedow, 1992).

The research carried out in conjunction with this grant has focused on studies of a mitochondrially-encoded 13 kDa protein (URF13) that is correlated with both male sterility and T-toxin sensitivity in cms-T maize. The goal of this research has been to characterize the mechanism by which URF13 interacts with T-toxins to permeabilize biological membranes. Results obtained during the funding period prior to the current one demonstrated that expressing the gene encoding URF13 in E. coli confers T-toxin sensitivity on the resulting bacterial cells (Dewey et al., 1988). Proof that T-toxin and URF13 were permeabilizing the bacterial cells, analogous to the effect seen in cms-T mitochondria, was provided by the observation that accumulated ⁸⁶Rb was immediately lost when E. coli cells expressing URF13 were exposed to T-toxin (Braun et al., 1989a). Subsequently, site-directed mutagenesis of URF13 expressed in E. coli led to 1) the identification of a specific residue, Asp-39, that appears to be essential for maintaining the T-toxin sensitivity of URF13 and 2) the demonstration that truncation of the normal 115-amino acid URF13 molecule to a species having fewer than 83 amino acids resulted in the loss of T-toxin sensitivity (Braun et al., 1989a/b).

Research Progress, 8/90 to present:

During the funding period associated with the present grant, studies have primarily involved attempts to better characterize the structure of URF13. Because URF13 is an integral membrane protein, capable of generating a hydrophilic pore across biological membranes following interaction with T-toxin, it was important to understand the relationship between the structural organization of URF13 in the membrane and its function (i.e., its ability to permeabilize the membrane). Theoretical considerations of the structural

features of URF13 based upon its derived amino acid sequence (e.g., hydrophobicity and hydrophobic moment profiles) in conjunction with experimental results obtained from the directed mutagenesis studies (Braun et al., 1989a) led to the development of a model of the structure of monomeric URF13 within the membrane (Korth et al., 1991; Levings and Siedow, 1992). In this model (Fig. 1; NOTE - citations refer to figures appearing in Appendix B), URF13 was postulated to contain three membrane-spanning alpha-helical domains, with helix I consisting of a hydrophobic membrane-spanning region (Fig. 2) and helices II and III forming two amphipathic membrane-spanning alpha-helices (Fig. 3). Given their amphipathic nature, helices II and III are capable of lining the hydrophilic pore formed by URF13. Because molecules having molecular weights up to 650 can cross the membrane following the interaction between URF13 and T-toxin, the resulting pore requires the association of six to eight amphipathic helices. This suggests that at least some of the functional URF13 in the membrane must be in an oligomeric structure composed of up to three to four URF13 monomers (Levings and Siedow, 1992).

The focus of the research carried out over the two years associated with the current funding period has involved attempts to verify the three-helical structure of URF13 and to demonstrate the presence of URF13 oligomers in the membrane. Attempts to support the three-helix model required localization of the carboxy and amino termini of URF13 within the membrane. To that end, an URF13 fusion protein was constructed that contained 14 amino acids linked to the amino terminus of URF13, derived from the phage T7 s10 coat protein. This sequence contained an 11-amino acid epitope that reacted with a commercially available antibody against the s10 protein. Given that an antibody to a 15-amino acid region in the carboxy terminus of URF13 already existed, the resulting s10:URF13 fusion protein provided antigenic tags at both ends of the molecule. Protease digestion studies were then used to determine the accessibility of the amino and carboxy termini of the fusion protein in E. coli membrane preparations having either right-side-out or inside-out orientations. The amino and carboxy ends of URF13 were found to be localized on opposite sides of the membrane (Korth et al., 1991), indicating that URF13 exists in the E. coli membrane with an odd number of membrane-spanning regions. Because the resulting proteolytic products were not compatible with a single membrane-spanning region, these results were taken as being supportive of the three-helical model. Somewhat surprisingly, these studies also indicated that URF13 has two topological orientations in the E. coli plasma membrane, with 60-75% of the carboxy termini oriented toward the cytoplasmic surface and the remaining 25-40% localized on the outer, periplasmic surface of the membrane. This dual orientation was not unique to the fusion protein; the carboxy terminus of standard URF13 expressed in E. coli showed a similar mixed orientation. However, protease studies using right-side-out and inside-out membrane preparations from cms-T mitochondria indicated that in its native setting, URF13 has only one orientation, with the carboxy terminus located exclusively on the matrix side of the inner mitochondrial membrane (which, electrochemically, corresponds to the preferred orientation in E. coli).

The possibility that URF13 might exist in the membrane in an oligomeric state was first suggested from the results of studies looking at the binding of radiolabeled (³H) T-toxin (specifically, reduced Pm toxin from P. maydis) to E. coli cells expressing URF13 (Braun et al., 1990). The binding of Pm toxin to URF13 was found to be specific and saturable, with saturation appearing at 250-500 pmole Pm toxin/mg E. coli protein and half-maximal binding falling in

the range of 50-90 nM. In addition, Pm toxin binding displayed a marked cooperativity, having a Hill coefficient that varied between 1.3 and 1.7 from one preparation to the next. While a Hill coefficient greater than 1.0 could have several explanations, one distinct possibility was the presence of binding cooperativity within an URF13 oligomer having multiple toxin binding sites. To test directly for the presence of URF13 oligomers, studies were carried out using lysine-specific, bifunctional cross-linking reagents. The best results were obtained with a 16.1 Å hydrophobic cross-linker, EGS (Korth *et al.*, 1991). These studies demonstrated that a subset of the URF13 molecules in both *E. coli* and *cms*-T mitochondria exist in the membrane in an oligomeric state, with dimers and some trimers appearing. More recent studies indicate that there is considerable trimer and some tetramer formation (See below). Addition of T-toxin did not change the oligomeric pattern seen upon cross-linking, indicating that T-toxin does not effect the association of URF13 monomers (or dimers) into higher-order structures, but that it operates through more subtle changes within previously existing oligomers. Finally, even though URF13 is present in a mixed orientation in the *E. coli* membrane, cross-linking studies carried out in conjunction with proteolysis experiments on the sl0:URF13 fusion protein, indicated that in any given multimeric complex, all the participating monomers have the same orientation within the membrane (Korth *et al.*, 1991).

In addition to establishing the cooperative (and specific) nature of the interaction between T-toxin and URF13, the Pm toxin binding studies demonstrated that two of the three known classes of toxin-resistant mutations of URF13, the truncated 82-amino acid species and substitutions at Asp-39, showed reduced levels of toxin binding compared to cells expressing standard URF13 (Braun *et al.*, 1990). However, the third toxin-insensitive mutation (a ten-residue deletion of amino acids 2-11 at the N-terminus) maintained levels of toxin binding comparable to those of standard URF13. Recently, we have examined more closely the nature of Pm toxin binding appearing in toxin-resistant URF13 mutants (Fig. 5). For mutants modified at residue 39, not only did the level of binding decrease (<50%) relative to standard URF13, but there was also a complete loss of binding cooperativity (Kaspi and Siedow, 1992). In addition, there was a two- to three-fold increase in the intrinsic dissociation constant (K_d) for T-toxin binding to the mutants. As part of the work outlined in this proposal, the Pm toxin binding properties of other toxin-resistant mutants will be studied to attempt to localize a specific toxin binding domain within URF13.

As a follow-up to the cross-linking and toxin binding studies described above, we investigated how the structure of URF13 was affected by treatment with the carboxylate-specific reagent, dicyclohexylcarbodiimide (DCCD). Previous studies had shown that addition of DCCD to isolated *cms*-T mitochondria led to a loss of sensitivity to added T-toxin [Holden, M.J. & H. Sze (1989) *Plant Physiol.* 91: 1296-1302.], and we had obtained analogous results following DCCD treatment of *E. coli* cells expressing URF13 (Braun *et al.*, 1989a). During the current funding period, we have found that DCCD is capable of cross-linking URF13 (Fig. 4) and does so more effectively than EGS, leading to the appearance of more higher-order (trimer and tetramer) species (Kaspi and Siedow, 1992). Our earlier mutagenesis studies had established that Asp-39 was the residue that reacts with DCCD (Braun *et al.*, 1989a). The nature of the reaction between a DCCD-activated carboxylate residue (i.e., Asp-39) and the displacing epsilon amino group requires that they be closely juxtaposed. Therefore, the appearance of DCCD cross-linking suggests that Asp-39 is either directly

involved in monomer:monomer interaction (i.e., by interacting with an epsilon amino group on an adjacent monomer) or, at the least, is proximal to a lysine residue on an adjacent URF13 monomer in the oligomeric complex. As expected, toxin-insensitive mutations that substitute for Asp-39 (e.g., D39V) lead to a loss of cross-linking by DCCD. However, the mutant D39E, which is also toxin-resistant but still has a carboxylate residue at position 39, did show cross-linking with DCCD. One of the goals of the present proposal will be to determine the residue that Asp-39 reacts with during DCCD-mediated cross-linking. Among other things, this will help to define the specific helix:helix interactions involved in the URF13 oligomeric structure.

Of additional interest in this line of investigation was the effect of DCCD on Pm toxin binding to URF13 (Kaspi and Siedow, 1992). Two changes appeared in the observed binding: 1) there was a complete loss of cooperativity and 2) the level of toxin binding to URF13 decreased by a factor of almost exactly two (analogous to the results seen with the toxin-insensitive substitution mutations at Asp-39). Interestingly, and unlike the Asp-39 mutants, the intrinsic K_d for toxin binding was not affected by DCCD treatment. Taken together, these results indicate that DCCD cross-linking locks the oligomeric URF13 complexes into a "closed" conformation that is not able to undergo a change to the "open," pore-forming conformation following binding of T-toxin. Less clear at present is whether this failure to undergo the conformational change to the open state leads to a loss of half the toxin binding sites or whether the failure to bind the second complement of T-toxin leads to the loss of the conformational change that causes pore formation.

Finally, we are attempting to use the DCCD reaction to establish a relationship between the observed DCCD cross-linking pattern and the loss of URF13 sensitivity to T-toxin. Such experiments include studies varying the concentration of DCCD used to cross-link URF13 as well as following the time course of cross-linking in the presence of a constant concentration of DCCD. We are also taking advantage of the lag phase associated with the expression URF13 in *E. coli* in studies looking at the relationship between the time course of appearance of higher-order oligomers as ascertained by cross-linking and the onset of T-toxin sensitivity. While correlative in nature, the initial results of these studies point to cross-linked trimers and tetramers as being the molecular species associated with the loss of T-toxin sensitivity following DCCD treatment and further work is ongoing.

Summary:

The research carried out during the current funding period has focused on the structure of URF13, and the results support the validity of the three-helix model of URF13 (Fig. 1) and provide direct evidence for the oligomeric nature of at least some of the URF13 molecules in the membrane. In addition, the toxin binding studies have provided insight into the dynamic nature of the T-toxin:URF13 interaction and the extent to which Asp-39 is crucial to the interaction that leads to membrane pore formation. Additional knowledge of the structure of URF13 is needed if the nature of the interaction between URF13 and T-toxin to produce a hydrophilic pore within the membrane is to ultimately be understood. The goal of the research outlined in the present proposal is to extend these studies to attempt to better define the structure of URF13 and to further ascertain amino acid residues that are involved in both oligomer formation and the interaction with T-toxin.

Publications Resulting from DOE Sponsored Research

Publications since submission of the previous DOE proposal (copies included with this proposal):

Braun, C.J., J.N. Siedow and C.S. Levings III (1990) Fungal toxins bind to the URF13 protein in maize mitochondria and E. coli. The Plant Cell 2: 153-161.

Korth, K.L., F. Struck, C.I. Kaspi, J.N. Siedow and C.S. Levings III (1991) Topological orientation of the membrane protein URF13. In "Plant Molecular Biology" (Herrmann, R.G. and Larkins, B.A., eds.) pp. 375-381. Plenum Press, London.

Korth, K.L., C.I. Kaspi, J.N. Siedow and C.S. Levings III (1991) URF13, a maize mitochondrial pore-forming protein, is oligomeric and has a mixed orientation in E. coli plasma membranes. Proc. Natl. Acad. Sci. USA 88: 10865-10869.

Levings, C.S. III and J.N. Siedow (1992) Molecular basis of disease susceptibility in the Texas cytoplasm of maize. Plant Mol. Biol. 18:
- IN PRESS.

Kaspi, C.I. and J.N. Siedow (1992) Cross-linking of URF13 by dicyclohexylcarbodiimide and its effect on T-toxin-mediated membrane pore formation. J. Biol. Chem. - SUBMITTED.

Relevant publications from funding prior to 8/90:

Dewey, R.E., J.N. Siedow, D.H. Timothy and C.S. Levings III (1988) A 13-kilodalton maize mitochondrial protein in E. coli confers sensitivity to Bipolaris maydis toxin. Science 239: 293-295.

Braun, C.J., J.N. Siedow, M.E. Williams and C.S. Levings III (1989a) Mutations in the maize mitochondrial T-urf13 eliminate sensitivity to a fungal pathogen. Proc. Natl. Acad. Sci. USA 86: 4435-4439.

Braun, C.J., J.N. Siedow and C.S. Levings III (1989b) The T-urf13 gene is responsible for toxin sensitivity in maize and E. coli. In: "The Molecular Basis of Plant Development" (Goldberg, R.B., ed.) pp. 79-85. Alan R. Liss, NY.

SCIENTIFIC BACKGROUND

URF13, cms-T and T-toxin Sensitivity:

Cytoplasmic male sterility is a maternally inherited trait that leads to the formation of non-viable pollen during microsporogenesis (1). Three cms types (T, C and S) can be distinguished in maize based upon their interaction with different nuclear genes that are able to restore fertility (1,2). Maize carrying cms-T cytoplasm, in widespread use in the U.S. in the late 1960s for the production of hybrid seed, is uniquely susceptible to infection by the fungus Bipolaris maydis, race T (formerly Helminthosporium maydis), leading to a disease commonly known as Southern corn leaf blight (3-5). A second fungus, Phyllosticta maydis, produces a similar disease, yellow leaf blight, on cms-T maize. These fungi each produce a set of structurally related pathotoxins, referred to collectively as T-toxins (6), that specifically affect cms-T maize mitochondria, but not mitochondria isolated from normal (fertile) maize or any of the other maize cms types (7-9). The numerous reported effects of T-toxins on cms-T mitochondria (7-11) have led to the view that these compounds produce their toxic effect by permeabilizing the inner mitochondrial membrane (3,10). Methomyl (S-methyl-N-[(methylcarbamoyl)oxy]thioacetimidate), an insecticide that bears no obvious structural resemblance to T-toxin, mimics all the observed effects of T-toxin on cms-T mitochondria but at 10^4 -fold higher concentrations (12). Leaver et al. (13,14) initially demonstrated that a 13 kDa polypeptide was specifically synthesized by isolated cms-T mitochondria. Subsequently, Dewey et al. (15) characterized a unique mitochondrial gene associated with this 13 kDa protein (T-urf13) that encodes a 115-amino acid polypeptide, URF13 (Fig. 2). Using an antibody specific for a 15-amino acid sequence in the carboxy terminus, URF13 was shown to be expressed in cms-T mitochondria and associated with the mitochondrial membrane (16).

Sensitivity to T-toxin and cytoplasmic male sterility in cms-T maize both appear to be correlated with the presence of URF13. Fertile cms-T maize plants can be derived from spontaneous revertants in cms-T cell culture lines (17,18). True breeding revertant plants regenerated from such cell lines are both male fertile and disease (or T-toxin)-resistant (18). Genetic analyses of such revertants indicate that they are associated with either the loss (19,20) or modification (21) of the T-urf13 gene. Definitive proof of the relationship between T-toxin sensitivity and the presence of URF13 was provided by Dewey et al. (22) who expressed URF13 in E. coli. The resulting bacteria were sensitive to T-toxin (or methomyl) as evidenced by the inhibition of cellular respiration (22), the massive and immediate loss of accumulated ^{86}Rb from the cells (23,24), and subsequent cell death. Additional support for the relationship between URF13 expression and T-toxin sensitivity has been provided by the expression of URF13 in baker's yeast (25,26). Huang et al. (25) constructed a chimeric gene consisting of the complete URF13 coding region fused at its amino terminus to a mitochondrial transit peptide. Growth of yeast cells expressing this chimeric gene was inhibited by T-toxin or methomyl. Interestingly, yeast cells transformed with full-length T-urf13 lacking any mitochondrial transit peptide were insensitive to T-toxin, although immunoblotting indicated that URF13 was being expressed in these cells (25).

The results outlined above suggest that URF13 is capable of permeabilizing biological membranes when it interacts with T-toxin but do not delineate

whether a specific association between URF13 and some other mitochondrial protein is needed to develop this response. Associations between URF13 and both the F_1F_0 -ATP synthase complex (27) and cytochrome c oxidase (28) have been reported based upon co-immunoprecipitation studies, leaving open the possibility that URF13 interacts with homologous complexes (29,30) in the bacterial plasma membrane. However, fractionation of the individual mitochondrial electron transfer complexes of cms-T mitochondria (31) showed some URF13 associated with every mitochondrial complex (K.L. Korth, personal communication). Whether purified URF13 incorporated into a lipid bilayer can bring about a T-toxin-induced permeabilization of the membrane in the absence of any additional protein remains to be demonstrated.

A reasonable explanation for the mechanism of action of T-toxin is that it binds to URF13 and brings about a change in the structure of URF13 such that a hydrophilic pore results. We investigated the binding of tritiated, reduced T-toxin from P. maydis (Pm toxin) to E. coli expressing URF13 (as outlined in the PROGRESS REPORT) and found that URF13 specifically bound Pm toxin and did so in a cooperative manner (32). We also observed a specific binding of Pm toxin to cms-T mitochondria, but the level of binding was over an order of magnitude lower (per mg protein) than the binding seen with E. coli. This very low level of binding could explain an earlier failure to detect specific T-toxin binding in cms-T mitochondria (33). These results established that T-toxin interacts directly with URF13. Further, toxin binding provides a more quantitative measure of the nature of the interaction between URF13 and T-toxin than that associated with simple toxin sensitivity assays. Continued use of Pm toxin binding will be made in the studies outlined in this proposal.

URF13 Structural Features:

The ability to express URF13 in E. coli provided the opportunity to carry out directed mutagenesis studies to characterize specific functional domains of URF13. Initially, three categories of toxin-resistant mutants were identified (Fig. 2; 22,24): 1) URF13 species containing fewer than 83 amino acids (24); 2) mutations at Asp-39, changing this residue to any other amino acid resulted in a toxin-resistant protein (24); and 3) a deletion mutation involving a ten-amino acid region near the URF13 amino terminus ($\Delta 2-11$) (23).

The current paradigm for hydrophilic pore formation by proteins in biological membranes postulates a series of membrane-spanning, amphipathic alpha-helices associated in a cylindrical fashion perpendicular to the plane of the membrane (34-37). The amphipathic helices are localized in the membrane such that their polar faces are oriented inward, lining the hydrophilic channel (36,37). These considerations were utilized to develop a model for the structure of URF13 (3,38) that envisioned a three-membrane-spanning, alpha-helical arrangement (Fig. 1). A hydrophobicity plot of URF13 (Fig. 2) indicates a hydrophobic membrane-spanning region between residues 13 and 32 (Fig. 2). Based upon hydrophobic moment calculations (Fig. 3) and helical wheel plots, two membrane-spanning amphipathic helices can be generated between residues 35 and 83. Based on this model, the failure of residues beyond Leu-83 to contribute to T-toxin sensitivity is attributed to their localization in the aqueous phase outside of the three membrane-spanning regions. Asp-39, which is essential for T-toxin sensitivity (24), is unusual in that it is a polar amino acid located on the hydrophobic face of helix II. However, this localization

is consistent with the ability of Asp-39 to form a stable adduct after reaction of URF13 with dicyclohexylcarbodiimide (DCCD) (24, See also below). As outlined in the PROGRESS REPORT, studies of the topological orientation of URF13 in the *E. coli* plasma membrane indicate that the amino and carboxy termini are localized on opposite sides of the membrane (39), consistent with the three-membrane-spanning helix model (Fig. 1).

Because NAD^+ ($M_r = 663$) can pass through the pores generated by URF13 (7), a pore diameter of 0.8-1.5 nm is indicated (40). Even with a pore diameter of 0.8 nm, six amphipathic helices would be needed to form the channel lining (40,41). If the three-helix model is correct, pore formation would require the association of some URF13 monomers into at least a trimeric structure to accommodate the largest observed pores. The possibility that URF13 could exist as an oligomer was first suggested by the appearance of cooperativity in the binding of Pm toxin to URF13 in *E. coli* (32) and by the appearance of apparent URF13 dimeric species on SDS gels (42). Our covalent cross-linking studies directly confirmed that a fraction of the URF13 in the membrane exists in an oligomeric state (39). More recently, DCCD has been shown to effectively cross-link URF13 (Fig. 4; 43). The observation that DCCD cross-links URF13 is interesting because this reagent has also been found to protect *cms*-T mitochondria (44,45) and *E. coli* cells expressing URF13 (24) against the effects of added T-toxin. In addition, [^{14}C]DCCD covalently labels URF13, and directed mutagenesis studies established that the site of DCCD labeling was Asp-39 (24).

In order to understand how DCCD acts in URF13, it is necessary to point out that DCCD reacts with carboxylate side chains on proteins to form an activated, DCCD-carboxylate intermediate that can either: 1) be hydrolyzed to an unmodified carboxylate with the loss of N,N'-dicyclohexylurea, 2) undergo rearrangement to form a covalently bonded DCCD adduct, or 3) undergo reaction with a suitably positioned nucleophile, displacing the DCCD as a dialkylurea and forming a covalent linkage between the displacing nucleophile and the carboxylate residue (46-48). The second reaction is slow relative to hydrolysis (#1), so rearrangement to form a stable DCCD adduct primarily occurs with carboxylate residues localized in hydrophobic environments (46,49,50). With respect to the third reaction, epsilon amino groups (e.g., lysines) represent the kinetically most active displacing nucleophile in proteins (47,48), giving rise to an amide bond with the DCCD-reactive carboxylate residue. Reconciling the observations that [^{14}C]DCCD can both covalently label and cross-link URF13 came with the realization that the labeled DCCD adduct only appeared in monomeric URF13 species on SDS gels, while the cross-linked, multimeric URF13 species were never labeled with DCCD (e.g., DCCD was displaced as a result of the intermolecular cross-linking reaction). This suggested that, within URF13 multimers, the intermolecular cross-linking reaction is kinetically more favored and only in monomeric URF13 is the lifetime of the activated DCCD/Asp-39 intermediate sufficiently long to permit rearrangement to form a covalent DCCD adduct. The DCCD cross-linking reaction therefore presents the opportunity to determine the residue that is in proximity to Asp-39 on an adjacent monomer as outlined in PROPOSED RESEARCH.

The response of URF13 to T-toxin requires that URF13 1) has the capacity to bind T-toxin and 2) is able to undergo the structural changes needed to generate a pore. Loss of either function will lead to a toxin-resistant species. With that in mind, we have investigated the Pm toxin binding

properties of both toxin-resistant URF13 mutants and DCCD-treated URF13 (43). Mutants at Asp-39 (D39V, D39E) (Fig. 5) and DCCD-treated URF13 both show a marked reduction ($\leq 50\%$) in the level of T-toxin binding relative to standard URF13, and there is a complete loss of binding cooperativity (43). As more toxin-resistant mutants of URF13 are generated (See PROPOSED RESEARCH), binding studies will be carried out on them to attempt to delineate specific residues and/or local domains that are associated with toxin binding versus those more directly involved in pore formation (See PROPOSED RESEARCH).

While this literature review has focused on URF13, it should be noted that a large number of proteins are known that are able to bring about the formation of hydrophilic channels or pores within biological membranes (37). These include diphtheria (51,52), tetanus (52,53) and botulinum toxins (54), exotoxin A (55) and several colicins (56,57,58), among others. Recently, the three-dimensional structures of the water-soluble forms of several of these channel-forming toxins, as determined by x-ray crystallography (59,60,61) and NMR spectroscopy (62), have appeared. In not focusing on this large literature as part of this SCIENTIFIC BACKGROUND, I do not mean to underestimate its relevance to our studies of URF13. Space limitations simply didn't allow its inclusion. However, I do keep track of the results coming out of this research and, where applicable, integrate them into our thinking about URF13.

SIGNIFICANCE

Since the appearance of the x-ray structure of the photosynthetic reaction center of R. viridus, attention has become focused on the marked advances being made in the study of membrane protein structure and structure-function relationships. Our research on URF13 was initiated to characterize the nature of T-toxin sensitivity in cms-T mitochondria. However, the project has evolved into a broader attempt to understand the structural features of URF13 that contribute to membrane pore formation and how T-toxin interacts with URF13 to elicit this response. In the work outlined under PROPOSED RESEARCH, I intend to draw upon the prior knowledge gained in studying URF13 as well as other channel-forming protein toxins to help advance our understanding of both URF13-T-toxin and URF13-membrane interactions during pore formation. In addition, this research takes advantage of some of the unique aspects of URF13 to help enhance our general understanding of pore-forming proteins. These advantages of URF13 include its relatively simple structure (practically, 83 amino acids cannot accommodate more than three membrane-spanning regions), its unique response to T-toxin and its ease of manipulation in E. coli. Channel-forming proteins play many roles in biology. While the specifics associated with the structures of different channels vary, there are, undoubtedly, some principles of both protein structure and membrane-protein interaction that apply generally to a number of functionally dissimilar protein channels. Learning more about the specifics of the URF13-T-toxin interaction and subsequent pore formation will allow us to gain further insight into some of these principles.

PROPOSED RESEARCH

URF13 Structural Studies:

Our working model of the structure of URF13 (Fig. 1) postulates the presence of three membrane-spanning helices, two of which are amphipathic and participate directly in pore formation. Given the ability of URF13 to form oligomeric structures in the membrane (39,43), there must be specific helix-helix interactions between adjacent URF13 monomers. In the absence of purified URF13, attempts to directly determine the structure of URF13, using NMR spectroscopy for example, are not possible. Therefore, we will continue to apply biochemical/molecular approaches to further develop the structural model of URF13, focusing on the nature of the helix-helix interactions involved in oligomer formation. More specifically, the potential number of interactions between the three membrane-spanning helices of URF13 within a given oligomer is limited, particularly if the two amphipathic helices, II and III, line the hydrophilic pore. The cross-linking studies outlined below should delimit if the monomer:monomer associations involve helix II:II and III:III or helix II:III interactions and the extent to which the hydrophobic helix I associates with either of the amphipathic helices on adjacent monomers. Identification of the specific residues (and their associated helices) involved in URF13 cross-linking reactions will constrain the possible orientations that the helices can take and ultimately lead to a better picture of the URF13 structure.

The ability of DCCD to cross-link URF13 (Fig. 4; 43) provides a useful starting point for investigating intermolecular helix-helix interactions, since one of the two reaction partners (Asp-39) is already established (24) and the other is most likely a lysine (46-48), which must be sufficiently close to Asp-39 to displace DCCD and form a stable amide linkage (46). Of the five lysine residues in URF13, three are found within the functional, pore-forming domain (i.e., residues 1-83 (24)) and, of these, two (Lys-32 and -37) are postulated to be in the vicinity of Asp-39 (Fig. 1). Lys-32 is located in the interhelical region between helices I and II while Lys-37 is located near the amino-terminal end of helix II. To determine which of these residues is involved in DCCD cross-linking, site-directed mutagenesis of each lysine, as well as the two in combination, will be carried out on URF13 in the pET5.13T construct using the procedure of Kunkel (63), as applied previously to URF13 (24,39). In addition to testing each construct for cross-linking with DCCD (43), the level of expression of URF13 (24), T-toxin sensitivity (22) and the binding of labeled Pm toxin (32) will all be measured.

If cross-linking is not seen with DCCD treatment it will be important to ascertain whether the resulting URF13 is still able to form oligomers. The lysine-specific cross-linker, EGS, could be used for this purpose, but the fact that we are modifying potential EGS reactive sites (See below) could confound this control. Therefore a second cross-linker that does not react with either lysine or carboxylate residues will be used to circumvent this potential problem. One possibility is *p*-azidophenyl glyoxal (APG), a bifunctional reagent that reacts with arginine residues through the phenyl glyoxal moiety and then as a non-specific photoaffinity reagent in the subsequent cross-linking reaction (64). URF13 contains five arginine residues located within both postulated helices II and III that should make URF13 amenable to APG cross-linking. Should APG prove unsuitable as a cross-linker of URF13, other reagents are commercially available (65), and these would be tested.

Because of the specificity of EGS toward lysine residues, the mutants developed above to look at the lysine involved in the DCCD cross-linking reaction can also be utilized in studies designed to ascertain which lysine residues are involved in EGS cross-linking. Among the three membrane-associated lysines, the two at positions #32 and 37 appear to be best positioned to be involved in EGS cross-linking (Fig. 1). However EGS (16.1 Å) is sufficiently long to allow it to extend out of the membrane and interact with two lysine residues (#s 102 and 104) that are located within the hydrophilic tail region, in the middle of the carboxy-terminal epitope. Reaction of either of these lysines with EGS could lead to a loss of antigenicity, which may explain why EGS does not appear to give rise to as much of the higher-order oligomers as DCCD does (unpublished observations). This potential problem will be surmounted by using pET5.13T which has a second epitope on its amino terminus that does not contain any lysines (39). This will allow verification of EGS cross-linking using immunoblotting even if the carboxy-terminal epitope has been modified. Shorter-armed, lysine-specific cross-linkers DSP (12 Å) and DST (6.4 Å) (65) will also be tested in these studies to further delimit intermolecular lysine-lysine distances.

The goals of this research are to establish: 1) what lysine residue(s), on adjacent monomers, are in proximity to the Asp-39 in helix II and 2) determine which lysine residues are involved in cross-linking by EGS. The information that comes out of these studies should allow some indication of what specific helix-helix associations exist in URF13 oligomers and allow us to begin to place constraints on the possible arrangements of the helices within the oligomeric structure. The results of these studies will also allow us to refine the URF13 structural model (Fig. 1) as new information appears.

Intermolecular interactions between helices I and I as well as helices I and II will also be tested. The only cysteine residue (Cys-27) in URF13 is located at the carboxy-terminal end of helix I (Fig. 1). The presence of helix I/I interactions on adjacent URF13 monomers can be tested for by seeing whether iodine treatment [which causes disulfide bond formation between proximally oriented cysteine residues (66)] of *E. coli* cells expressing URF13 results in the appearance of any cross-linked dimers on SDS gels. To look for helix I/II interactions, heterobifunctional cross-linking reagents that have specificities for cysteine and lysine residues on their respective ends will be used (65). These include succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC, 11.6 Å) and succinimidyl 4-(p-maleimidophenyl)butyrate (SMPB, 14.5 Å), both of which have proved useful in cross-linking other proteins (67,68). The postulated proximity of Cys-27 to lysines 32 and 37 (Fig. 1) makes the latter two residues candidates for intermolecular cross-linking with Cys-27 using these reagents, if an appropriately oriented interaction between helices I and II exists. If cross-linking is observed with either of these reagents, the involvement of Lys-32 and/or -37 can be tested directly using the site-directed mutants at these two residues described above for the DCCD and EGS cross-linking studies. With respect to all of these cross-linking studies, it is worth emphasizing that the three-dimensional geometric relationship between any two adjacent helices could effectively preclude the appearance of cross-linking even though the helices do interact. Negative results will therefore not be taken as evidence for the absence of a specific helix-helix interaction.

Recently, Pakula and Simon (66) reported the development of a strategy for the analysis of membrane protein structure in which directed mutagenesis was

used to randomly create individual cysteine substitutions within postulated trans-membrane helical regions of the Tar receptor of *E. coli*. Iodine treatment of the resulting cells and analysis of the formation of cross-linking between Tar receptor dimers allowed the determination of which cysteine residues were sufficiently close to form intermolecular disulfide cross-links (66). The results provided insights into the arrangement of the two trans-membrane helices within the Tar receptor dimeric structure. While no small undertaking, this procedure is applicable to URF13 and would allow us, among other things to address the question of the extent to which helix III is involved in interhelical interactions in conjunction with oligomer formation. None of the studies outlined above allow us to determine anything with respect to the interactions between helix III and helices I and/or II. While this study will directly address the question of URF13 structure, all cysteine substitution mutants will be tested for T-toxin sensitivity and their ability to oligomerize using the standard cross-linking reagents (EGS, DCCD). This study will be a major undertaking and won't be initiated until the structural studies outlined above are in process. However, it offers the opportunity of allowing us to deduce the complete structural arrangement of the helices within the URF13 oligomeric complexes using a systematic approach.

T-toxin Binding Studies:

The binding of [³H]Pm toxin to URF13 has proved informative in characterizing the interaction between URF13 and T-toxin, providing the first suggestion that pore formation may involve an oligomeric species (32). More recently, quantitative analysis of the binding of URF13 to toxin-resistant mutations at Asp-39 have helped to establish a correlation between the cooperativity of T-toxin binding and membrane pore formation (Fig. 5; 43). Currently, G. Ward and M. Williams in the Levings' laboratory have been carrying out a saturation mutagenesis study on URF13 that involves a selection for methomyl-resistant phenotypes. They have now saturated much of the URF13 primary sequence and have found toxin-resistant mutations involving 25 different URF13 residues, located within all three postulated membrane-spanning regions (Ward et al., unpublished observation and personal communication). They are presently undertaking a qualitative screen of these mutants to distinguish those that bind labeled T-toxin from those that do not. This analysis is ongoing, but preliminary indications are that most of the toxin-resistant mutants are deficient in binding T-toxin. As part of the collaborative effort between our two laboratories, my group will carry out quantitative binding studies on these mutants (See Letter of Collaboration, Appendix A). Also of interest is the extent to which any of the toxin-resistant mutants show variations in their ability to undergo oligomerization, as evidenced by cross-linking upon treatment with either EGS or DCCD. It is likely that no single amino acid substitution would be sufficient to eliminate oligomer formation if several pairs of interactions are involved (69,70), but screening for oligomer formation is quite straightforward. As noted previously, lysine-specific mutants from the mutagenesis studies outlined above will also be looked at for their T-toxin binding properties.

While the near term goal will involve the mapping of URF13 residues that directly affect T-toxin binding versus those more indirect in their effects on the T-toxin/URF13 interaction, the quantitative toxin binding studies should provide additional information. For example, T-toxin binding studies of the mutations at position 39 have indicated changes in cooperativity, the level

of toxin binding and the intrinsic K_d (Fig. 5; 43). As characterization of the toxin-resistant mutants becomes more thorough, it is expected that related binding properties associated with residues in either specific stretches of the URF13 primary sequence or localized regions of the URF13 tertiary structure (Fig. 1) will begin to emerge.

We would also like to address the question of which URF13 species actually bind T-toxin. For example, it is presently not known whether monomeric URF13 is capable of binding T₃-toxin. We will attempt to develop a membrane binding system to investigate [³H]Pm toxin interactions with electrophoretically separated URF13 oligomers. Detergent-solubilized URF13 oligomers will either be separated by native gel electrophoresis (71), or first cross-linked with DCCD and separated on SDS/PAGE, followed by transfer to an appropriate membrane [nitrocellulose (72) or polyvinylidene difluoride (73)]. The resulting membranes will then be probed with labeled Pm toxin to determine if binding is associated with specific oligomeric species. Binding will be determined by fluorography and/or excision of the region of interest and scintillation counting. It's clear that these experiments contain numerous potential pitfalls: e.g., native gel electrophoresis may not resolve the oligomeric species (74) or renaturation of URF13 after SDS/PAGE may not be sufficient to allow specific toxin binding (75). Further, with DCCD-treated protein, the effects of DCCD cross-linking on the toxin binding properties of URF13 cannot be neglected (43). However, these experiments do not represent the primary thrust of our binding studies; the goal is simply to attempt to ascertain, qualitatively, which species of URF13 are capable of binding Pm toxin.

Purification of URF13:

To ultimately gain a complete understanding of the structure of URF13 and the mechanism of pore formation will require our having a purified protein. Preliminary solubilization and purification procedures have been applied to URF13. Based upon these studies, we know that URF13 can be solubilized using 1% lauryl maltoside, and standard protein purification procedures have been initiated using this material. The Pm toxin binding studies indicate that 250-500 pmoles of toxin bind per mg *E. coli* protein (32). Assuming one binding site per URF13 monomer, this corresponds to only about 3-6 ug URF13/mg *E. coli* protein. If we scale up the growth of *E. coli*, a 10 l culture would give rise to the equivalent of 5-10 mg of URF13. This is consistent with the generally low levels of expression of eukaryotic membrane proteins in *E. coli* for reasons that are presently not well understood (76). I should point out that during the course of our studies on URF13, no fewer than four different promoter systems have been used (22,24,39 & Ward, *et al.*, unpublished), all giving comparable levels of URF13 expression. Some success with the over expression of eukaryotic membrane proteins has been attained using the insect-baculovirus system (77,78), but attempts to express URF13 using this system have indicated that the production of URF13 is toxic to the insect cell cultures, even in the absence of T-toxin (K. L. Korth and C. S. Levings, personal communication). Needless to say, numerous laboratories are presently working on procedures to obtain higher levels of expression of eukaryotic membrane proteins in bacteria, and we will take advantage of any breakthroughs that might appear.

The low level of URF13 expression in the starting material (0.5% of total bacterial cell protein) makes it difficult to apply standard purification procedures, given the inevitable losses that appear during many purification

steps. To overcome this problem, we need to develop an affinity procedure that will allow us to carry out a single, high-yield and specific purification step for URF13. This will put us in a position to more effectively apply subsequent standard procedures and still have a reasonable amount of product at the end of the purification. While not presented under PROGRESS REPORT, during the course of the current funding period, we have attempted to use the monoclonal antibody specific for the carboxy-terminal region of URF13 (38,39) to develop an immunoaffinity purification of detergent-solubilized URF13. URF13 was successfully adsorbed to this column, but it could not be efficiently eluted from the column using a variety of standard elution conditions. We will continue attempts to refine the procedure for eluting URF13 from the immunoaffinity column, but also initiate the development of other affinity purification procedures.

Several approaches will be taken to develop other affinity purification procedures for URF13. We will take advantage of the fact that several commercial systems are presently available for constructing chimeric proteins in which the protein of interest is fused to a second peptide that is amenable to affinity chromatography. Two such systems will be tried initially. One is the QIAexpress system (79,80) in which a string of six histidine residues plus a protease cleavage site will be linked, in frame, to the amino terminus of URF13. The resulting fusion protein can be purified by specific adsorption to a Ni-chelate affinity column (Ni-NTA agarose) and subsequently eluted under mildly acidic conditions (pH 4.5-5.0) or in the presence of 250 mM imidazole at pH 7.0 (79,81). Removal of the amino-terminal extension is accomplished by treating the purified protein with the protease, Factor Xa (79). We will begin with the amino terminal fusion because of our previous finding that a 14-residue amino-terminal extension had no effect on the T-toxin sensitivity of URF13 in our topology studies (39). If fusing the (His)₆ sequence to the amino terminus is not effective, we will also try making a carboxy-terminal fusion. As a second option, several kits are available for developing chimeric constructs in which an entire protein is fused to the amino terminus of URF13. We will initially use the Pharmacia system (82,83) for fusing the glutathione S-transferase (GST) gene to the amino terminus of URF13 with the idea of purifying the subsequent product using affinity chromatography of the GST on a glutathione-Sepharose column. Elution from the column requires 5 mM glutathione and the GST can be removed by proteolysis using Factor Xa, analogous to the (His)₆ fusion. While we will start out with these two systems, as others come onto the market, we will try any that look particularly suitable for use with membrane proteins. In addition to our maintaining a close collaboration with the Levings laboratory, which is a decided plus in developing these constructs, Dr. David Rhoads will be joining my laboratory as an NSF Plant Biology Fellow on the URF13 project, and he comes with a strong background in molecular biology as a result of his Ph.D. research in Lee McIntosh's laboratory at Michigan State (84).

An alternative to the above approaches to developing an affinity procedure for purifying URF13 will also be considered, particularly if the molecular approaches prove unsuccessful. Although methomyl has a low affinity for URF13 (8,10), Gauvrit and coworkers (85) have shown that methomyl analogues with long-chain (C-13) alkyl moieties on the carbamate methyl group (instead of a methyl) have a biological activity (on a concentration basis) similar to that of T-toxin. If such a long-chain analogue were synthesized with an amino function at the end of the alkyl tail, it should be suitable for coupling to an affinity matrix such as Activated CH-Sepharose 4B without interfering with the

functional end of the molecule (85). Before getting into the organic chemistry involved here, I will see if Gauvrit is interested in a collaborative effort by which he synthesizes the compound and we use it to generate an affinity matrix for the purification of URF13. If he is not interested, other organic chemists will be approached. While molecular methods outlined above will be given first priority, if they don't show satisfactory progress, synthesis of a methomyl affinity column will be undertaken.

Concurrent with attempts to purify URF13, we will develop a liposome reconstitution assay to test for the functional integrity of the solubilized URF13. The reconstitution of colicins and the diphtheria-type toxins into unilamellar vesicles (liposomes) has been a very useful tool in the study of channel formation by these proteins (51,57). Further, it has not been demonstrated directly that URF13 acts independently of other proteins, and this can only be tested by reconstituting purified URF13 into an artificial membrane system. We will develop a reconstitution system using liposomes because we have experience using such vesicles, and because they have proved so generally useful in the past for studying isolated membrane proteins (86,87). Initially, a chloride efflux procedure utilized by Davidson *et al.* (88) with colicin E1 will be tried instead of using membrane potential-sensitive dyes. Numerous variables can affect the functional incorporation of protein into artificial membranes (87,88), and these will need to be systematically investigated. Further, if URF13 pore formation requires certain oligomers that are normally present in very low amounts, a functional reconstitution may only emerge upon reaching a certain level of purification.

Once URF13 has been purified, several lines of experimentation can be pursued. For example, circular dichroism (CD) spectroscopy has proved useful for measuring the relative proportions of alpha-helix and beta-sheet present in proteins (89,90). Our model for URF13 (Fig. 1) predicts a 60% alpha-helical content. CD spectroscopy will be carried out on purified URF13 to ascertain the amount of alpha-helix present (90) in both solubilized and membrane-bound URF13 as well as look at the effects of the addition of T-toxin and DCCD on its secondary structure. Care will be taken to minimize the effects of light scattering and absorption flattening of URF13 in unilamellar vesicles (91) and the uv absorption of T-toxin (6) when making these measurements.

Finally, it should be recalled that the ultimate goal of this research is to understand the three-dimensional structure of URF13 and how the latter interacts with T-toxin to form a membrane pore. Given the unlikelihood of ever getting URF13 crystals, NMR spectroscopy seems the best candidate for directly measuring the URF13 structure. Smith and Peersen recently reviewed the status of solid-state NMR spectroscopy in the study of membrane protein structure (92) and pointed out both the utility and drawbacks of using orientation-dependent measurements and dipolar coupling approaches to probe membrane protein secondary structure, helix-helix interactions and the behavior of specific (¹³C-labeled) residues (e.g., following T-toxin binding). We clearly cannot do this kind of work without the collaboration of a solid-state NMR spectroscopist. However, once NMR-scale amounts (10-20 mg) of pure URF13 are available, we will give a high priority to establishing a collaboration with such a laboratory. The unique and interesting biological features associated with the URF13-T-toxin interaction should make this a very attractive system for use with solid-state NMR, which is how I would propose to sell it.

BIBLIOGRAPHY

1. Hanson, M. R. and Conde, M. F. (1985) *Intl. Rev. Cytol.* 94: 213-267.
2. Laughnan, J. R. and Gabay-Laughnan, S. (1983) *Annu. Rev. Genet.* 17: 27-48.
3. Levings, C. S., III and Siedow, J. N. (1992) *Plant Mol. Biol.* - IN PRESS.
4. Levings, C. S., III (1991) *Science* 250: 942-947.
5. Pring, D. R. and Lonsdale, D. M. (1989) *Annu. Rev. Phytopathol.* 27: 483-502.
6. Danko, S. J., Kono, Y., Daly, J. M., Suzuki, Y., Takeuchi, S. and McCrery, D. A. (1984) *Biochemistry* 23: 759-766.
7. Matthews, D. E., Gregory, P. and Gracen, V. E. (1979) *Plant Physiol.* 63: 1149-1153.
8. Berville, A., Gahzi, A., Charbonner, M. and Bonavent, J. F. (1984) *Plant Physiol.* 76: 508-517.
9. Holden, M. J. and Sze, H. (1984) *Plant Physiol.* 75: 235-237.
10. Klein, R. R. and Koeppe, D. E. (1985) *Plant Physiol.* 77: 912-916.
11. Holden, M. J. and Sze, H. (1987) *Plant Physiol.* 84: 670-676.
12. Humaydan, H. S. and Scott, E. W. (1977) *Hortic. Sci.* 12: 312-313.
13. Forde, B. G., Oliver, R. J. C. and Leaver, C. J. (1978) *Proc. Natl. Acad. Sci. USA* 75: 3841-3845.
14. Forde, B. G. and Leaver, C. J. (1980) *Proc. Natl. Acad. Sci. USA* 77: 418-422.
15. Dewey, R. E., Levings, C. S., III and Timothy, D. H. (1986) *Cell* 44: 439-449.
16. Dewey, R. E., Timothy, D. H. and Levings, C. S., III (1987) *Proc. Natl. Acad. Sci. USA* 84: 5374-5378.
17. Brettell, R. I. S., Thomas, E. and Ingram, D. S. (1980) *Theor. Appl. Genet.* 58: 55-58.
18. Umbeck, P. F. and Gegenbach B. G. (1983) *Crop Sci.* 23: 584-588.
19. Fauron, C. M-R., Havlik, M. and Brettell, R. I. S. (1990) *Genetics* 124: 423-428.
20. Rottmann, W. H., Brears, T., Hodge, T. P. and Lonsdale, D. M. (1987) *EMBO J.* 6: 1541-1546.

21. Wise, R. P., Pring, D. R. and Gengenbach, B. G. (1987) *Proc. Natl. Acad. Sci. USA* 84: 2858-2862.
22. Dewey, R. E., Siedow, J. N., Timothy, D. H. and Levings, C. S., III (1988) *Science* 239: 293-295.
23. Braun, C. J., Siedow, J. N. and Levings, C. S., III (1989) in "The Molecular of Plant Development: UCLA Symposium on Molecular Biology." (Goldberg, R. B., ed.) pp. 79-85. Alan R. Liss, New York.
24. Braun, C. J., Siedow, J. N., Williams, M. E. and Levings, C. S., III (1989) *Proc. Natl. Acad. Sci. USA* 86: 4435-4439.
25. Huang, J., Lee, S-H., Lin, C., Medici, R., Hack, E. and Myers, A. M. (1990) *EMBO J.* 9: 339-347.
26. Glab, N., Wise, R. P., Pring, D. R., Jacq, C. and Slonimski, P. (1990) *Mol. Gen. Genet.* 223: 24-32.
27. Wise, R. P., Fliss, A. E., Pring, D. R. and Gengenbach, B. G. (1987) *Plant Mol. Biol.* 9: 121-126.
28. Leaver, C. J., Isaac, P. G., Small, I. D., Bailey-Serres, J., Liddell, A. D. and Hawkesford, M. P. (1988) in "Mitochondrial Biogenesis." (Leaver, C. J. and Lonsdale, D. M., eds.) pp. 81-92. Royal Society Press, London.
29. Pederson, P. L. and Carafoli, E. (1987) *Trends Biochem. Sci.* 12: 146-150.
30. Cherupf, V., Lemieux, L., Au, D. C.-T. and Gennis, R. B. (1990) *J. Biol. Chem.* 265: 11185-11192.
31. Peiffer, W. E., Ingle, R. T. and Ferguson-Miller, S. (1990) *Biochemistry* 29: 8696-8701.
32. Braun, C. J., Siedow, J. N. and Levings, C. S., III (1990) *The Plant Cell* 2: 153-161.
33. Frantzen, K. A., Daly, J. M. and Knoche, H. W. (1987) *Plant Physiol.* 83: 863-868.
34. Fox, R. O. and Richards, F. M. (1982) *Nature (London)* 300: 325-330.
35. Hall, J. E., Vodyanoy, I., Balasubramian, T. M. and Marshall, G. R. (1983) *Biophys. J.* 45: 233-247.
36. Guy, H. R. (1984) *Biophys. J.* 49: 249-259.
37. Ojcius, D. M. and Young, J. D-E. (1991) *Trends Biochem. Sci.* 16: 225-229.
38. Korth, K. L., Struck, F., Kaspf, C. I., Siedow, J. N. and Levings, C. S., III (1991) in "Plant Molecular Biology." (Herrmann, R. G. and Larkins, B. A., eds.) pp. 375-381. Plenum Press, London.

39. Korth, K. L., Kaspi, C. I., Siedow, J. N. and Levings, C. S., III (1991) Proc. Natl. Acad. Sci. USA 88: 10865-10869.
40. Davidson, V. L., Brunden, K. R., Cramer, W. A. and Cohen, F. S. (1984) J. Membr. Biol. 79: 105-118.
41. Spach, G., Heitz, F. and Trudelli, Y. (1983) in "Physical Chemistry of Transmembrane Ion Motions." (Spach, G., ed.) pp. 375-383. Elsevier, Amsterdam.
42. Hack, E., Lin, C., Yang, H and Horner, H. T. (1991) Plant Physiol. 95: 861-870.
43. Kaspi, C. I. and Siedow, J. N. (1992) J. Biol. Chem. - SUBMITTED
44. Bouthyette, P.-Y., Spitsberg and Gregory, P. (1985) J. Exptl. Bot. 36: 511-528.
45. Holden, M. J. and Sze, H. (1989) Plant Physiol. 91: 1296-1302.
46. Nalecz, M. J., Casey, R. P. and Azzf, A. (1986) Methods Enzymol. 125: 86-108.
47. Albertson, N. F. (1962) Organic Reactions 12: 205-213.
48. Carraway, K. L. and Koshland, D. E. (1972) Methods Enzymol. 25: 616-623.
49. Futai, M., Noumi, T. and Maeda, M. (1987) Adv. Biophys. 23: 1-37.
50. Pick, U. and Racker, E. (1979) Biochemistry 18: 108-113.
51. Stenmark, H. Arfansen, S., Afansiev, B. V. and Olsnes, S. (1992) J. Biol. Chem. 267: 8957-8962.
52. O'Keefe, D. O. and Collier, R. J. (1989) Proc. Natl. Acad. Sci. USA 86: 343-346.
53. Menestrina, G., Forti, S. and Gambale, F. (1989) Biophys. J. 55: 393-405.
54. Donovan, J. J. and Middlebrock, J. L. (1986) Biochemistry 25: 2872-2876.
55. Farahbaakhsh, Z. T. and Wisniewski, B. J. (1989) Biochemistry 28: 580-585.
56. Lakey, J. H., Massotte, D., Heitz, F., Dasseux, J.-L., Faucon, J.-F., Parker, M. W and Pattus, F. (1991) Eur. J. Biochem. 196: 599-607.
57. Cramer, W. A., Cohen, F. S., Merrill, A. R. and Song, H. Y. (1990) Mol. Microbiol. 4: 519-526.
58. Merrill, A. R., Cohen, F. S. and Cramer, W. A. (1990) Biochemistry 29: 5829-5836.
59. Parker, M. W., Postma, J. P. M., Pattus, F., Tucker, A. D. and Tsernoglou, D. (1992) J. Mol. Biol. 224: 639-657.

60. Choe, S., Bennett, M. J., Fujii, G., Curmi, P. M. G., Kantardjieff, K. A., Collier, R. J. and Eisenberg, D. (1992) *Nature* 357: 216-222.
61. Allured, V. S., Collier, R. J., Carroll, S. F. and McKay, D. B. (1986) *Proc. Natl. Acad. Sci. USA* 83: 1320-1324.
62. Wormald, M. R., Merrill, A. R., Cramer, W. A. and Williams, R. J. P. (1990) *Eur. J. Biochem.* 191: 155-161.
63. Kunkel, T. (1985) *Proc. Natl. Acad. Sci. USA* 82: 488-492.
64. Ngo, T. T., Yam, C. F., Lenhoff, H. M. and Ivy, J. (1981) *J. Biol. Chem.* 256: 11313-11818.
65. Pierce Chemical Co. (1992) "ImmunoTechnology Catalog and Handbook" Section E, pp. 8-39.
66. Pakula, A. A. and Simon, M. I. (1992) *Proc. Natl. Acad. Sci. USA* 89: 4144-4148.
67. Mahan, D. E., Morrison, L., Watson, L. and Haugeland, L. S. (1987) *Anal. Biochem.* 162: 163-176.
68. Gitman, A. G., Kahane, I. and Loyter, A. (1985) *Biochemistry* 24: 2762-2768.
69. Popot, J.-L. and de Vitry, C. (1990) *Annu. Rev. Biophys. Biophys. Chem.* 19: 369-403.
70. Popot, J.-L. and Engelman, D. M. (1990) *Biochemistry* 29: 4031-4037.
71. Dewald, B., Dulaney, J. T. and Touster, O. (1974) *Methods Enzymol.* 32: 82-91.
72. Towbin, H., Staehlin, T. and Gordon, J. (1979) *Proc. Natl. Acad. Sci. USA* 76: 4350-4354.
73. Pilatte, Y., Hammer, C., Frank, M. and Fries, L. (1989) *J. Immunol. Methods* 120: 37-43.
74. Ludwig, B., Downer, N. W. and Capaldi, R. A. (1979) *Biochemistry* 18: 1401-1407.
75. Gershonfi, J. M. and Palade, G. E. (1983) *Anal. Biochem.* 131: 1-15.
76. Ferreira, G. C. and Pederson, P. L. (1992) *J. Biol. Chem.* 267: 5460-5466.
77. Page, M. J., Hall, A., Rhodes, S., Skinner, R. H., Murphy, V., Sydenham, M. and Lowe, P. N. (1989) *J. Biol. Chem.* 264: 19147-19154.
78. Vissavajjhala, P. and Ross, A. H. (1990) *J. Biol. Chem.* 265: 4746-4752.
79. QIAGEN Inc. (1991) "The QIAexpressionist"

80. Stuber, D., Matile, H. and Garotta, G. (1990) in "Immunological Methods." Vol. IV. (Lefkovits, I. and Pernis, B., eds.) pp. 121-152. Academic Press, New York.
81. Hochuli, E., Bannwarth, W., Dobeli, H., Gentz, R. and Stuber, D. (1988) Bfo/Technology 6: 1321-1325.
82. Smith, D. B. and Johnson, K. S. (1989) Gene 67: 31-40.
83. Johnson, K. S. (1989) Nature 338: 585-587.
84. Rhoads, D. M. and McIntosh, L. (1991) Proc. Natl. Acad. Sci. USA 88: 2122-2126.
85. Aranda, G., Durlin, P. and Gauthier, C. (1987) Phytochemistry 26: 1909-1913.
86. Hirata, H. (1986) in "Techniques for the Analysis of Membrane Proteins." (Ragan, C. I. and Cherry, R. J., eds.) pp. 77-96. Chapman Hall, London.
87. Silvius, J. R. (1992) Annu. Rev. Biophys. Biomol. Struct. 21: 323-348.
88. Davidson, V. L., Cramer, W. A., Bishop, L. J. and Brunden, K. R. (1984) J. Biol. Chem. 259: 594-600.
89. Johnson, W. C. (1988) Annu. Rev. Biophys. Biophys. Chem. 17: 145-166.
90. Provencher, S. W. and Glockner, J. (1981) Biochemistry 20: 33-37.
91. Mao, D. and Wallace, B. A. (1984) Biochemistry 23: 2667-2673.
92. Smith, S. O. and Peersen, O. B. (1992) Annu. Rev. Biophys. Biomol. Struct. 21: 25-47.

CURRICULUM VITAE

Name: James N. Siedow

Rank: Professor of Botany

Degrees: B.A. (with high honors) - University of Texas, Austin; 1969
Ph.D. - Indiana University, Bloomington; 1972

Employment: Fellow, Microbiology Training Grant, with A. San Pietro,
Department of Plant Sciences, Indiana University; 1969-72.
NIH Postdoctoral Fellow, with G. Palmer, Biophysics Research
Division, University of Michigan; 1973.
NIH Postdoctoral Fellow and Research Associate, with G. Palmer,
Department of Biochemistry, Rice University; 1974-76.
Assistant Professor, Department of Botany, Duke University,
1976-81.
Associate Professor, Department of Botany, Duke University,
1981-87.
Professor, Department of Botany, Duke University, 1987-present.

Additional: Visiting Professor, Department of Genetics, North Carolina State
University; 1987.
Program Director, Cellular Biochemistry Program, National Science
Foundation; 1988/89.

Awards: Trinity College of Arts & Sciences Award for Excellence in
Teaching; 1985.
NIH Postdoctoral Fellowship; 1973-74.
Woodrow Wilson Fellow; 1969
Phi Beta Kappa, University of Texas; 1969.

Publications (within the past five years):

Mack, A. J., T. K. Peterman and J. N. Siedow (1987) Lipoxygenase in higher
plants: biochemical properties and physiological role. *in* "Isozymes: Current
Topics in Biological and Medical Research, Vol-13" pp. 127-154. Alan R.
Liss, Inc., New York.

Sugawara, F., G. Strobel, R. N. Strange, J. N. Siedow, G. D. van Duyne and J.
Clardy (1987) Phytotoxins from the pathogenic fungi Drechslera maydis and
Drechslera sorghicola. Proc. Natl. Acad. Sci. USA 84: 3081-3085.

Berthold, D. A., D. J. Fluke and J. N. Siedow (1987) A determination of the
aroid alternative oxidase by radiation inactivation analysis. *in* "Plant
Mitochondria: Structural, Functional and Physiological Aspects" (Moore, A. L.
and R. B. Beechey, eds.) pp. 113-116. Plenum Press, New York.

Siedow, J. N. and M. E. Musgrave (1987) The significance of cyanide-resistant
respiration to carbohydrate metabolism in higher plants. *in* "Plant
Mitochondria: Structural, Functional and Physiological Aspects" (Moore, A. L.
and R. B. Beechey, eds.) pp. 351-359. Plenum Press, New York.

- Siedow, J. N., D. A. Berthold, D. J. Fluke, M. E. Musgrave and S. J. Stegink (1987) The cyanide-resistant oxidase in higher plant mitochondria. in "Advances in Membrane Biochemistry and Bioenergetics" (Kim, C. H., H. Tedeschi, J. J. Diwan and J. C. Salerno, eds.) pp. 227-233. Plenum Press, New York.
- Musgrave, M. E., C. O. Miller and J. N. Siedow (1987) Do some plant responses to cytokinins involve the cyanide-resistant respiratory pathway? *Planta* 172: 330-335.
- Kennedy, R. A., T. C. Fox and J. N. Siedow (1987) Activities of isolated mitochondria and mitochondrial enzymes from aerobically and anaerobically germinated barnyard grass (Echinochloa) seedlings. *Plant Physiol.* 85: 474-480.
- Koepppe, D. E. and J. N. Siedow (1987) Mitochondrial electron transport: Chemiosmosis. in "Models in Plant Physiology, Biochemistry and Technology, Vol-1" (Newman, D. and K. G. Wilson, eds.) pp. 105-108. CRC Press, Boca Raton
- Musgrave, M. E. and J. N. Siedow (1987) The cyanide-resistant pathway. in "Models in Plant Physiology, Biochemistry and Technology, Vol-1" (Newman, D. and K. G. Wilson, eds.) pp. 109-112. CRC Press, Boca Raton
- Dewey, R. E., J. N. Siedow, D. H. Timothy and C. S. Levings III (1988) A 13-kilodalton maize mitochondrial protein in E. coli confers sensitivity to Bipolaris maydis toxin. *Science* 239: 293-295.
- Berthold, D. A., D. J. Fluke and J. N. Siedow (1988) Determination of molecular mass of the aroid alternative oxidase by radiation-inactivation analysis. *Biochem. J.* 252: 73-77.
- Braun, C. J., J. N. Siedow and C. S. Levings III (1989) The T-urf13 gene is responsible for toxin sensitivity in maize and E. coli. in "The Molecular Basis of Plant Development" (Goldberg, R. B., ed.) pp. 79-85. Alan R. Liss, Inc., New York.
- Braun, C. J., J. N. Siedow, M. E. Williams and C. S. Levings III (1989) Mutations in the maize mitochondrial T-urf13 gene eliminate sensitivity to a fungal pathotoxin. *Proc. Natl. Acad. Sci. USA* 86: 4435-4439.
- Siedow, J. N. (1990) Regulation of the cyanide-resistant respiratory pathway. in "Perspectives in Biochemical and Genetic Regulation of Photosynthesis" (Zelitch, I., ed.) pp. 355-366. Wiley-Liss, New York.
- Braun, C. J., J. N. Siedow and C. S. Levings III (1990) Fungal toxins bind to the URF13 protein in maize mitochondria and Escherichia coli. *The Plant Cell* 2: 153-161.
- Crumbliss, A. L., K. L. McLachlan, J. N. Siedow and S. P. Walton (1990) Electrochemistry of heat-extracted methanogenic bacterial cofactor F₄₃₀. *Inorg. Chim. Acta* 170: 161-163.
- Kalt, W., C. B. Osmond and J. N. Siedow (1990) Malate metabolism in the dark after ¹³C₂ fixation in the Crassulacean plant Kalanchoe tubiflora. *Plant Physiol.* 94: 826-832.

- Kramer, P. J., J. N. Siedow and J. S. MacFall (1990) Nuclear magnetic resonance research on plants. in "Measurement Techniques in Plant Science" (Hashimoto, Y., P. J. Kramer, H. Nonami and B. R. Strain, eds.) pp. 403-431. Academic Press, San Diego.
- Siedow, J. N. (1991) Plant lipoxygenase: structure and function. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42: 145-188.
- Korth, K. L., C. I. Kaspi, J. N. Siedow and C. S. Levings III (1991) Topological orientation of the membrane protein URF13. in "Plant Molecular Biology" (Herrmann, R. G. and B. A. Larkins, eds.) pp. 375-381. Plenum Press, London.
- Moore, A. L. and J. N. Siedow (1991) The regulation and nature of the cyanide-resistant alternative oxidase of plant mitochondria. *Biochim. Biophys. Acta* 1059: 121-140.
- Korth, K. L., C. I. Kaspi, J. N. Siedow and C. S. Levings III (1991) URF13, a maize mitochondrial pore-forming protein, is oligomeric and has a mixed orientation in Escherichia coli plasma membranes. *Proc. Natl. Acad. Sci. USA* 88: 10865-10869.
- Moore, A. L. and J. N. Siedow (1992) The nature and regulation of the alternative oxidase of plant mitochondria. *Biochem. Soc. Trans.* 20: 361-363.
- Levings, C. S., III and J. N. Siedow (1992) Molecular basis of disease susceptibility in the Texas cytoplasm of maize. *Plant Mol. Biol.* 18: - IN PRESS.
- Siedow, J. N., J. Whelan, A. Kearns, J. T. Wiskich and D. A. Day (1992) Topography of the alternative oxidase in soybean mitochondria. in "Molecular, Biochemical and Physiological Aspects of Plant Respiration" (Lambers, H. and L. H. W. van der Plas, eds.) - IN PRESS.
- Siedow, J. N. and A. L. Moore (1992) Regulation of electron transfer through the alternative respiratory pathway. in "Molecular, Biochemical and Physiological Aspects of Plant Respiration" (Lambers, H. and L. H. W. van der Plas, eds.) - IN PRESS.
- Kaspi, C. I. and J. N. Siedow (1992) Cross-linking of URF13 by dicyclohexylcarbodiimide and its effect on T-toxin-mediated pore formation. *J. Biol. Chem.* - SUBMITTED.
- Siedow, J. N. and A. L. Moore (1992) A kinetic model for the regulation of electron transfer through the cyanide-resistant pathway in plant mitochondria. *Biochim. Biophys. Acta* - SUBMITTED.
- Berthold, L. A. and J. N. Siedow (1992) Purification of the cyanide-resistant alternative oxidase of skunk cabbage (Symplocarpus foetidus) mitochondria. *Plant Physiol.* - SUBMITTED.
- Robinson, S. A., D. Yakir, M. Ribas-Carbo, L. Giles, C. B. Osmond, J. N. Siedow and J. A. Berry (1992) Measurements of the engagement of cyanide-resistant respiration in the CAM plant Kalanchoe diadema using on-line oxygen isotope discrimination. *Plant Physiol.* - SUBMITTED.

FACILITIES AND EQUIPMENT

Facilities:

My laboratory at Duke University consists of three rooms, two of which are adjoining, equipped for biochemical studies, with a total area of 930 ft². All rooms are on the basement floor of the Biological Sciences Building. One of the laboratories is designated for use of radioisotopes and contains a high-exhaust fume hood. Additional facilities include a walk-in cold room for biochemical manipulations at 5°C, a controlled temperature Tyler box for growing etiolated maize, if needed, and a dishwasher room.

Equipment:

Major items of equipment housed in my laboratory include: a Shimadzu UV-265 uv-vis spectrophotometer; a Shimadzu RF5000U spectrofluorophotometer; a Sorvall RC-5B preparative centrifuge; a Beckman model L-265B ultracentrifuge; two microcentrifuges; three oxygen monitoring assemblies (electrode/reaction cell/amplifier/recorder) to measure bacterial and mitochondrial respiratory activities; a Waters Model M6000A dual-pump HPLC with Shimadzu SPD-6A uv detector and Hewlett-Packard 3390A integrator; a Gilson microfractionator and an ISCO Golden Retriever for fractionating small and large chromatographic columns, respectively; a New Brunswick Microferm ten-liter fermentor; and three refrigerated circulating water baths. Additional equipment in the laboratory includes a pH meter, micro- and macro-analytical balances and a lot of electrophoretic equipment (four complete Hoefer SE-400 vertical slab gel units, one Hoefer SE-200 mini-gel unit, a Baserunner nucleic acid sequencing unit, four power supplies and a Biorad Transblot apparatus).

I also have access to a complete Pharmacia FPLC system (including Mono Q, Mono S and Mono P columns), an LKB Rackbeta liquid scintillation counter, a Beckman TL-100 tabletop ultracentrifuge, a Perkin-Elmer Cetus DNA Thermal Cycler for carrying out PCR reactions, and a Konica automated x-ray film developer, all of which are housed within two shared instrumentation rooms on the same floor as my laboratory. In addition, the Department of Botany houses the University's primary oligonucleotide synthesis facility which offers two-day turnaround service at a cost of \$2.50 per base.

An additional item of equipment required of the proposed research is a CD spectrophotometer, which exists in the Department of Biochemistry as part of its Protein Structure Facility. They have a Jobin-Yvon Dichrography Mark V instrument interfaced with an Apple IIe computer and it is available to the general Duke community for a nominal user fee (\$10/sample).

APPENDIX A

Letter of Collaboration

North Carolina State University

Department of Genetics
College of Agriculture and Life Sciences

Box 7614
Raleigh, NC 27695-7614
(919) 515-~~8302~~ 7115
FAX: (919) 515-~~8355~~ 7115

June 24, 1992

Dr. Jim Siedow
Department of Botany
Duke University
Durham, NC 27706

Dear Jim:

I am pleased to continue our collaboration in the study of the URF13 protein. In my view, our joint efforts have made significant progress in understanding how URF13 interacts with toxins to permeabilize membranes, and I look forward to our future collaborations.

Gerty C. Ward and Mark E. Williams of my lab have nearly completed the saturation mutagenesis study of T-*urf13*. We have identified many single amino acid mutations that are no longer able to confer toxin sensitivity to *E. coli*. To date, we have over 40 mutations that are toxin insensitive. This mutant collection should be valuable for your studies, and I am pleased to provide this collection for your future studies of URF13.

Yours truly,

A handwritten signature in cursive script, reading "C. S. Levings, III".

Charles S. Levings, III
Professor of Genetics

CSL:sq

APPENDIX B

Figures 1-5

Fig. 1 Proposed model of URF13 disposition within the membrane, indicating the location of the three putative membrane-spanning helical domains. Asp-39 (site of the DCCD reaction) is indicated in bold outline. Taken from ref. 3.

Fig. 2 A. Hydropathy profile of URF13. Predicted hydrophobic regions are indicated by the shaded areas above the single-letter amino acid sequence. The arrows show the location of regions where mutations cause a loss of T-toxin sensitivity. B. Schematic representation of the origin of the DNA in the *T-urf13* open reading frame. Taken from ref. 24.

Fig. 3 Hydropathic moment profile of URF13. The hydropathic moment was calculated at 100° according to Eisenberg [Eisenberg, D. (1984) Ann. Rev. Biochem. 53: 595-623.] using the consensus scale of Eisenberg and a sliding window of 19 amino acids. Numbers on the abscissa refer to the URF13 primary sequence. Taken from ref. 3.

Fig. 4: Immunoblots of DCCD treated intact *E. coli* cells and isolated spheroplasts and isolated *cms-T* mitochondria. Cells and spheroplasts were treated with 1.0 mM DCCD while mitochondria were treated with 50 μM DCCD. All incubations were at 1.0 mg protein/ml for 10 min at room temperature with gentle agitation. Samples were separated on SDS/PAGE and the resulting blots probed with the URF13 specific antibody, MAb-C (39) and visualized with a chemiluminescent reagent.

Fig. 5: Scatchard plot of [^3H]Pm toxin binding to *E. coli* cells expressing standard URF13 and mutants at Asp-39. (●) - standard URF13; (○) - D39E; (▼) - D39V. Aliquots of *E. coli* cells (0.5 mg protein/ml) were incubated for 30 min with varying concentrations of tritiated Pm toxin and the cells were centrifuged through silicon oil to separate bound and unbound toxin as described previously (32). Specifically bound toxin is expressed as pmoles Pm toxin bound per mg *E. coli* protein. Calculated binding parameters are as follows: URF13 - $K_d = 90$ nM, $P_t = 475$ pmol/mg, Hill coefficient = 1.47; D39E - $K_d = 145$ nM, $P_t = 198$ pmol/mg; D39V - $K_d = 226$ nM, $P_t = 130$ pmol/mg.

reprint removed.
als

END

**DATE
FILMED**

12 / 14 / 92