DOE/ER/13855--T1

Final Report

DOE/ER/13855--T1

DE93 000908

Stable Isotope Studies

Contracts: AT(11-1)-3127, E(11-1)-3127, EY-76-S-02-3127
DE-AC02-80ER10612, DE-FG02-88ER13855

between

Division of Research and Office of Basic Energy Sciences
of
U. S. Atomic Energy Commission
U. S. Energy Research and Development Administration
U.S. Department of Energy

and

Research Foundation of the City University of New York
for and in conjunction with
Brooklyn College of the City University of New York

and with

Research Foundation of the State University of New York
for and in conjunction with
State University of New York at Stony Brook

Takanobu Ishida
Principal Investigator

March 1, 1972 - February 29, 1992

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Table of Contents

	page
Acknowledgement	iii
Summary	v
Personnel	ix
Curriculum Vitae	xiii
Publication	xv
I. Introduction	1
II. Correlation of isotope effects with molecular structures and molecular forces	5
II-A. Development of theoretical tools	6
II-A-1. Direct orthogonal polynomial approximations	6
II-A-2. Orthogonal expansion coupled with diagonal approximation	8
II-B. Correlation studies	11
II-B-1. End-atom isotope effects	11
Deviations from the first rule of the mean	14
II-B-2. Central-atom isotope effects	15
Isotope effects at the center of highly symmetric molecules	16
III. Correlation of zero-point energy with molecular structures and molecular forces	19
III-A. Zero-point energy as an explicit function of molecular structures and forces	19
III-A-1. Development of mathematical tools	19
III-A-2. Zero-point energy of n-alkanes	23
III-A-3. Hydrogen isotope effects in ZPE and linear frequency sum rules	24
III-A-4. Additivity for zero-point energy	26
III-A-5. Other applications	28
III-B. Isotope effect on the zero-point energy shift upon condensation	34
IV. Vapor pressure isotope effects	37
IV-A. Vapor pressure isotope effects in liquid fluoromethanes	37
IV-B. Vapor pressure isotope effects in liquid and solid ammonia	45
IV-C. Medium cluster model of liquid for vapor pressure isotope effects	50
V. Fractionation of stable isotopes	53
V-A. Fractionation of nitrogen-15	53
V-A-1. Equilibrium properties of NO/N ₂ O ₃ system	54
V-A-2. Nitrogen-15 fractionation based on the NO/N ₂ O ₃ exchange system	55

	page
V-B. Hydrophobic catalysts for hydrogen isotope exchange between dihydrogen and water	57
V-B-1. Hydrophobicity of polycrystalline platinum surfaces	58
V-B-2. Chemical poisoning and regeneration of platinum surfaces	59
V-B-3. Electrochemical modification of platinum surfaces by poly(p-phenylene) .	60
V-B-4. Electrochemical modification of platinum surfaces by other organic films .	67
V-B-5. Capacitance measurement for determination of effective surface area of edge plane of pyrolytic graphite	67
V-C. Transient behavior of square cascade of isotope separation stages	73
VI. References	77

Acknowledgement

This research has been a collaborative effort with a number of senior colleagues, post-doctoral associates, graduate students, and undergraduate students.

Major funding for the research has been provided by the U. S. Atomic Energy Commission, Energy Research and Development Agency and Department of Energy, whose continuous support for this program over the twenty-year period from March, 1972 to February, 1992 has made it possible. The City University of New York supported the program over the period, March, 1972 through August, 1979, with grants for personnel and purchase of instruments through its Faculty Research Award Program and for computing time through its Computing Centers. The Max Planck Society of Germany financed Principal Investigator's leave in 1986. The Science and Technology Agency of Japan supported a postdoctoral associate. The Tokyo Club provided a support for a postdoctoral fellow for two years. Brookhaven National Laboratory allowed an access to an isotope ratio mass spectrometer and other major equipment in early seventies.

I express my sincere appreciation to all who have made this research possible.

**Takanobu Ishida
Professor of Chemistry
State University of New York
Stony Brook, New York**

[blank page]

Summary

The research has been conducted by the Principal Investigator, Professor Takanobu Ishida, in collaboration with other colleagues and students in the general area of isotope chemistry and isotope separation during the period, March 1, 1972 ~ February 29, 1992. The research has been carried out principally at Brooklyn College of the City University of New York, March, 1972 ~ August, 1979, and at the State University of New York at Stony Brook, September, 1979 ~ February, 1992. The Principal Investigator spent the period, January, 1986 ~ July, 1986, in Mainz, Germany, as a Max Planck Institute Visiting Scholar.

The research has been in four general areas: (1) correlation of isotope effects with molecular forces and molecular structures, (2) correlation of zero-point energy and its isotope effects with molecular structure and molecular forces, (3) vapor pressure isotope effects, and (4) fractionation of stable isotopes.

(1) Correlation of isotope effects with molecular structure and molecular forces:

- ♦ Mathematical correlation method for the isotopic reduced partition function ratio with molecular structures and molecular forces were developed on the basis of the orthogonal functional theorem, and prototype molecules, CO_2 , HCHO , CH_4 , C_2H_4 , C_2H_6 , and C_6H_6 were studied.
- ♦ For end-atom isotope effects and, in particular, the hydrogen isotope effects:
 - The isotopic reduced partition function ratio can be expressed as a sum of contributions of bond-stretching, angle-bending and other intra-molecular motions; the contributions are additive.
 - The contributions by stretching and bending motions largely determine the isotope effects.
 - The contributions of bond-stretching coordinates are proportional to the stretching force constant, while the contributions of bending motions are not as proportional.
 - A simple rule for computing the isotope effects was developed.
- ♦ For central-atom isotope effects:
 - Bond-stretching forces contribute over seventy percent of total magnitudes of isotope effects.
 - The valence coordinate contributions to the isotopic reduced partition function ratio is not as additive as for the end-atom isotope effects, mainly due to the higher-order kinetic energy interactions.
- ♦ For the isotope substitution at the center of highly symmetric molecules, it is possible to obtain an explicit and straightforward correlation of the isotope effects with the molecular parameters.

- ♦ A systematics of D-matrix was developed to predict and identify the sources of deviations from given isotopic rules of the mean.

(2) Correlation of zero-point energy with molecular structures and molecular forces:

- ♦ A mathematical tool for the correlation of zero-point energy (ZPE) with the molecular parameters was developed on the basis of the orthogonal functional theorem.
- ♦ Linear frequency sum rule was established, which holds approximately but with high precision for the sets of isotopic molecules for which a second-order sum rule holds. The rules for hydrogen isotope substitutions hold with precisions of few wavenumbers.
- ♦ Zero-point energy and hydrogen isotope shifts in zero-point energy can be expressed as a sum of contributions of bond-stretching, angle-bending and other molecular motions: The contributions are additive. Tables of transferrable frequency parameters for the additivity were constructed for the homologous series, n-alkanes and n-alkyl chlorides. Sums of the parameters give the zero-point energy within 0.7% of exact values.
- ♦ Within each main group of periodic table, the contribution of H-X bond-stretching to hydrogen isotope shifts in zero-point energy for the molecule HX is a linear function of electronegativity of X, where X is a molecular moiety.
- ♦ The effect of London dispersion forces on the isotope effect of zero-point energy shifts due to condensation is expressible as a product of an isotope-independent factor, which only depends on properties of the condensed phase, and an isotope-dependent factor, which only depends on properties of the gas. Rules of thumb for predicting what molecular features would lead to a larger or smaller isotope-shifts in the condensation-shift were developed.

(3) Vapor Pressure Isotope Effects:

- ♦ Vapor pressure isotope effects (VPiE) for D/H- and $^{13}\text{C}/^{12}\text{C}$ -substitutions in liquid methyl fluoride, methylene difluoride, and fluoroform were measured, and the results were compared with the VPiEs in liquid methane. All observed results are explainable in terms of molecular symmetry, relative magnitudes of intermolecular forces in the liquids, and temperature-dependent force constants on the basis of simple cell model.
 - Large normal D/H-VPiEs observed in the fluoromethanes are due to the blue-shifts in the C-H stretching frequencies upon condensation.
 - Negative curvatures in the plots of $T \ln(f_c/f_g)$ vs. $1/T$ for the D/H-VPiE in the fluoromethanes are due to the blue shifts, the magnitudes of which increase with decreasing temperature.

- An anomalously large, normal D/H-VPIE in methylene difluoride is also attributable to an extensive interaction between the blue-shifting C-H stretching modes and the strong external forces.
 - The $^{13}\text{C}/^{12}\text{C}$ -VPIEs are inverse, mainly due to interactions of the internal modes with hindered rotation.
- ♦ Vapor pressure isotope effects for D/H- and $^{15}\text{N}/^{14}\text{N}$ -substitutions in liquid and solid ammonia were measured at temperatures between 163K and 243K.
 - The triple points for $^{14}\text{NH}_3$, $^{14}\text{ND}_3$, and $^{15}\text{NH}_3$ were determined.
 - The linear dependence of $T \ln(f_e/f_p)$ on $1/T$ is due to the lattice modes, especially, to the hindered rotation.
 - The difference in the vapor pressures of NH_3 and ND_3 is approximately constant between the triple point temperatures of NH_3 and ND_3 .
 - The close-packed structure of solid ammonia persists more or less not only through the phase change but also over the liquid temperature range that we studied.
 - The external force in liquid ammonia is much more directional on the hydrogen-side of ammonia molecule than on the lone-pair side of the molecule.
- ♦ Medium cluster model of liquid was developed. In the model, a cluster consists of a central molecule and a number (m) of indistinguishable molecules that surround the central molecule, and the intermolecular interactions among these molecules are explicitly taken into account.
 - The model contains a built-in mechanism by which the observed VPIEs can be represented better than the cell model without significantly increasing the number of force constant parameters.
 - As m is increased beyond approximately nine, qualitative features of computational results of the medium cluster model approaches those of the cell model, and the results due to the artifacts of limited cluster sizes become insignificant relative to the experimental precisions attainable by the presently available techniques. A close representation of an observed VPIE data by the cell model should, in principle, be possible through careful attention to details of interaction force constant parameters within the framework of the cell model.

(4) Fractionation of stable isotopes:

- ♦ Fractionation of ^{15}N was studied using a nitrogen isotope exchange between a liquid, which is representable as a mixture of liquid N_2O_3 and liquid N_2O_4 , and a gas phase in equilibrium with the liquid. The system was studied at temperatures, $-9^\circ\text{C} \sim 20^\circ\text{C}$, and under pressures, $1 \sim 4$ atm.

- The effective unit stage separation factor, α_{eff} , increases with increasing pressure, e.g., at 15°C, from 1.006 at 1 atm to 1.030 at 2.7 atm, and then decreases as the increasing pressure starts to reduce the average oxidation state of nitrogen in the liquid.
- The height equivalent of theoretical plate (HETP) is a linear function of the gas flow rate. The overall exchange rate is diffusion-controlled.
- The HETP of this NO/N₂O₃ system is smaller than that of the Nitrox process, the process based on the exchange between nitric oxide and nitric acid, by an order of magnitude. Thus, the NO/N₂O₃-based plant will have a significantly smaller plant than the Nitrox plant in spite of the smaller α_{eff} of the NO/N₂O₃ system.
- The product-end refluxing of NO/N₂O₃-based plant will require one-third the amount of sulfur dioxide needed for the product-end refluxing in the Nitrox plant.
- ◆ Development of electrode-based hydrophobic catalyst for the hydrogen-isotope exchange between dihydrogen and water was carried out.
 - Poly(p-phenylene) film, prepared by electropolymerization of p-terphenyl in acetonitrile and tetraethylammonium tetrafluoroborate is mechanically durable, electrochemically stable, and as hydrophobic as poly(tetrafluoroethylene). A hydrogen-isotope exchange column can be packed with a conductive packing material, and catalyst metal (e.g., platinum) and poly(p-phenylene) are electro-codeposited on it; an *in situ* control of hydrophobicity is possible.
 - Chemically poisoned platinum surface can be depoisoned easily by subjecting it to an electrode potential excursion, thus making an *in situ* regeneration of the catalyst bed possible.
 - Effective surface area of the edge plane of pyrolytic graphite can be determined from measurements of its electric capacitance using a triangular potential wave of 1 Hz to 10 Hz.
- ◆ The first-order solution of a partial differential equation representing a material balance in a square cascade of close-separation stages derived by Karl Cohen for the total reflux condition was extended to a second-order solution, thus lowering the time limit of applicability of Cohen's solution.
 - The transient behavior is not symmetric with respect to change of signs in ϵ ($\equiv \alpha - 1$).
 - The second-order term is especially important in the domain of inverse isotope effect in which ϵ is negative.
 - An isotope separation plant based on an inverse isotope effect phenomenon should have a shorter start-up time if all other plant parameters are same.
 - It will also be easier to design a product-end refluxer having a small holdup, if the desired isotope enriches in a gas phase.

Personnel

Present Affiliation

Principal Investigator:

Takanobu Ishida

**Professor
State University of New York at Stony Brook
Stony Brook, New York**

Scientific Collaborators:

Jacob Bigeleisen

**Distinguished Professor Emeritus
State University of New York at Stony Brook**

Francis T. Bonner

**Professor
State University of New York at Stony Brook**

Karl Heinzinger

**Max-Planck Institut für Chemie
Mainz, Germany**

Hidetake Kakihana

**Professor Emeritus
Sophia University, Tokyo, Japan**

Myung W. Lee

**Savannah River Laboratory
E. I. duPont de Nemours
Aiken, South Carolina**

Takao Oi

**Associate Professor
Department of Chemistry
Sophia University, Tokyo, Japan**

William Spindel

**Senior Consultant
National Academy of Sciences
Washington, D.C.**

Postdoctoral Research Associates:

Motonari Adachi

**Assistant Professor
Institute of Atomic Energy
Kyoto University, Japan**

Madhav Ghatge

**Chief, Process & Project Engineering
Clean Coal Project Division
Morgantown Energy Technology Center
U.S. Department of Energy**

Koichi Iinuma	Assistant Professor Department of Nuclear Engineering Tohoku University, Sendai, Japan
Masaaki Musashi	Postdoctoral research associate Department of Chemistry State University of New York at Stony Brook
Masato Murata	Advanced Isotope Separation and Application Section Power Reactor and Nuclear Fuel Development Corp. Tokai, Japan
Anthony Popowicz	Assistant Director of Computing Services Rockefeller University
Fukuji Yamashita	Associate Professor Department of Chemical Engineering Kanagawa Institute of Technology

Graduate Students:

Lester Borodinsky, Ph.D.	Staff Scientist Keller and Heckman Washington, D.C.
Min Chu, Ph.D.	Postdoctoral Fellow Department of Chemistry University of Illinois, Urbana-Champaign
Lisette Cuevas	Graduate student Department of Chemistry State University of New York
Susan Hillenbrand, M.S.	Lecturer Department of Chemistry Long Island University, Southampton, New York
Pat Jayanta, M.S.	Molecular Biosystems San Diego, California
Arundhati Kanungo, Ph.D.	Mrs. Michael Prencipe
Tseng Ven King, Ph.D.	Patent Examiner Superconductivity Group U.S. Patent Office
Zvi Kornblum, Ph.D.	Professor and Chair Department of Chemistry Cooper Union, New York
Leonard Krebs, M.S.	

Peter Lew, Ph.D.	Exploitation Technology U. S. Central Intelligence Agency
Steven Minick, M.S.	Environmental Measurement Laboratory U.S. Department of Energy, New York
James Pollin, Ph.D.	Engineering Manager Dallas Gas & Electric, Inc.
Michael Prencipe, Ph.D.	Section Head, Abrasives Research Section Colgate Palmolive, New Jersey
Jan Shulman, Ph.D.	Rohm and Haas Co. Delaware Valley
Henry Wieck, Ph.D.	Division Leader Chemistry Division P. A. Consulting Group, New Jersey

Undergraduate Students:

Mohammad Bahavar	Ph.D. Candidate Department of Chemical Engineering Clarkson University
Paul E. Bigeleisen, M.D.	Associate Professor Department of Anesthesiology University of Minnesota
Steven Brandt, Ph.D.	Director of Pharmaceutical Chemistry Sepractor, Inc. Marlborough, Massachusetts
Isaac Cohen, Ph.D.	Chief, Development Division Estee Lauder Inc. Melville, New York
Jeffrey Feiner, M.D.	South Orange County Cardiology Group Laguna Hills, California
David Finston, Ph.D.	Associate Professor Department of Mathematics New Mexico State University
Edward Gabalski, M.D.	
Mark Gilbert, M.D.	
Catherine Harrison, Ph.D.	
Richard Hom, M.D./Ph.D.	

William Limm, Ph.D.

Food and Drug Administration
Washington, D.C.

Darrell Mayfield, M.S.

Geologist
Axem Resources
Denver, Colorado

Riaz Rouhani, Ph.D.

Postdoctoral Research Associate
Department of Biochemistry
New York Hospital-Cornell Medical Center

Ellen Sekreta, Ph.D.

Postdoctoral Researcher Associate
Department of Chemistry
University of California, Irvine

Claudio Spadacenta, B.S.

David Trauber, M.D.

Private practice
Brooklyn, New York

Technical Personnel:

Armand Gazes

Associate Director of Computing Services
Rockefeller University

Curriculum Vita

Takanobu Ishida

B.S. (Chemistry) Kyoto University, 1953; M.S. (Chemistry) Kyoto University, 1955; M.S. (Nuclear Engineering), New York University, 1958; Ph.D. (Nuclear Engineering) Massachusetts Institute of Technology, 1964.

Research Associate, Chemistry Department, Brookhaven National Laboratory, 1964 - '66.

Research Associate, Chemistry Department, Belfer Graduate School of Science, Yeshiva University, 1966 - '68.

Brooklyn College, City Univeristy of New York, Assistant Professor to Professor of Chemistry, 1968 - '79.

University of Rochester, Visiting Associate Professor to Visiting Professor of Chemistry, 1973 - '79.

Professor of Chemistry, State University of New York at Stony Brook, 1979 - .

Visiting Scholar, Max Planck Institut für Chemie, Mainz, 1986.

Research Collaborator, Brookhaven National Laboratory 1972 - .

Member, American Chemical Society

Member, American Physical Society

[blank page]

Publication

Papers

1. Correlation of isotope effects with molecular forces. II. Triatomic molecules", Bigeleisen, J.; Ishida, T.; Spindel, W., *J. Chem. Phys.*, **1971**, *55*, 5021.
2. "Isotope chemistry and molecular structure. A simplified theory of end atom isotope effects", Bigeleisen, J.; Ishida, T., *J. Am. Chem. Soc.*, **1973**, *95*, 6155.
3. "Vibrational reduced partition function of the Morse oscillator", Ishida, T., *J. Chem. Phys.*, **1974**, *61*, 3009.
4. "Isotope chemistry and molecular structure. Total deuterium isotope effects", Bigeleisen, J.; Ishida, T., *J. Chem. Phys.*, **1975**, *62*, 80.
5. "Isotope chemistry and molecular structure. Carbon and oxygen isotope chemistry", Bigeleisen, J.; Horn, R. C.; Ishida, T., *J. Chem. Phys.*, **1976**, *64*, 3303.
6. "Isotope chemistry and molecular structure. Deviations from the first rule of the mean", Ishida, T.; Bigeleisen, J., *J. Chem. Phys.*, **1976**, *64*, 4775.
7. Medium cluster model for vapor pressure isotope effect", Pollin, J. S.; Ishida, T., *J. Chem. Phys.*, **1977**, *66*, 4433.
8. "The influence of molecular orientation on the vapor pressure isotope effect in liquid methanes", Pollin, J. S.; Ishida, T., *J. Chem. Phys.*, **1977**, *66*, 4442.
9. "Theory of inverse isotope effect", Ishida, T.; Kornblum, Z. C.; Pollin, J. S., Zentralinstitut für Isotopen- und Strahlenforschung Mitteilungen, Leipzig, Nr. 5, "Arbeitstagung, Isotope in der Natur, Gera, 22-26 September, 1975", **1976**, Akademie der Wissenschaften der DDR, Leipzig, pp. 40-111.
10. "Kinetics of a Square Cascade of Close-Separation Stages under Total Reflux", Wieck, H. J.; Ishida, T., *Separation Science*, **1977**, *12* (No. 6), 587.
11. "Carbon-13 vapor pressure isotope effect in CHF_3 ", Borodinsky, L.; Wieck, H. J.; Mayfield, D.; Ishida, T., *J. Chem. Phys.*, **1978**, *68*, 3279.
12. "Approximation of isotopic reduced partition function ratios by the WINIMAX polynomial perturbation method (WIMPER)", Bigeleisen, J.; Lee, M. W.; Ishida, T.; Bigeleisen, P. E., *J. Chem. Phys.*, **1978**, *68*, 3501.
13. "Isotope effect on zero-point energy shift upon condensation. I. Formulation and application to ethylene, methane, and fluoromethanes", Kornblum, Z. C.; Ishida, T., *J. Chem. Phys.*, **1978**, *69*, 1814.

14. "Correlation of the isotope chemistry of hydrogen, carbon, and oxygen with molecular forces by the WIMPER(2) method", Bigeleisen, J.; Ishida, T.; Lee, M.W., *J. Chem. Phys.*, **1981**, *74*, 1799.
15. "Studies in the vibrational spectroscopy of sodium trioxodinitrate", Bonner, F. T.; Akhtar, M. J.; King, T. V.; Chen, L. H.; Ishida, T., *J. Phys. Chem.*, **1981**, *85*, 4051.
16. "Ab initio molecular orbital study of dimer CHF₃", Popowicz, A.; Ishida, T., *Chem. Phys. Lett.*, **1981**, *83*, 520.
17. "Vapor pressure isotope effects in liquid fluoroform", Popowicz, A.; Oi, T.; Shulman, J.; Ishida, T., *J. Chem. Phys.*, **1982**, *76*, 3732.
18. "Synthesis and purification of methyl-¹³C fluoride", Cohen, I.; Shulman, J.; Popowicz, A.; Ishida, T. *J. Labelled Compd. Radiopharm.*, **1982**, *19*, 631.
19. "Correlation of zero-point energy with molecular structure and molecular forces. I. Development of approximation", Oi, T.; Ishida, T., *J. Phys. Chem.*, **1983**, *87*, 1067.
20. "Ab initio molecular orbital calculation on clusters of methyl fluoride", Oi, T.; Sekreta, E.; Ishida, T., *J. Phys. Chem.*, **1983**, *87*, 2323.
21. "Vapor pressure isotope effects in liquid methyl fluoride", Oi, T.; Shulman, J.; Popowicz, A.; Ishida, T., *J. Phys. Chem.*, **1983**, *87*, 3153.
22. "Correlation of zero-point energy with molecular structure and molecular forces. II. Approximation for n-paraffins", Oi, T.; Ishida, T., *J. Phys. Chem.*, **1984**, *88*, 1507.
23. "Correlation of zero-point energy with molecular structure and molecular forces. III. Approximation for H/D isotope shifts and linear frequency sum rule", Oi, T.; Ishida, T., *J. Phys. Chem.*, **1984**, *88*, 2057.
24. "Nitrogen-15 fractionation by countercurrent exchange between liquid N₂O₃-N₂O₄ mixture and their vapor phases under pressured conditions", Prencipe, M.; Spindel, W.; Ishida, T., *Sep. Sci. Tech.*, **1985**, *20*, 489.
25. "Additivity of vibrational zero-point energy", Oi, T.; Popowicz, A.; Ishida, T., *J. Phys. Chem.*, **1986**, *90*, 3090.
26. "Valence coordinate contributions to zero-point energy shifts due to hydrogen isotope substitution", Oi, T.; Ishida, T., *Bull. Chem. Soc. Jpn.*, **1986**, *59*, 373.
27. "Orthogonal approximation and classification of molecules for zero-point energy shifts due to H/D isotope substitutions", Oi, T.; Ishida, T., *Bull. Chem. Soc. Jpn.*, **1986**, *59*, 3669.
28. "Vapor pressure isotope effects in liquid methylene difluoride", Kanungo, A.; Oi, T.; Popowicz, A.; Ishida, T., *J. Phys. Chem.*, **1987**, *91*, 4198.
29. "Zero-point energy shifts in hydrides of the elements upon H/D isotope substitutions and their periodicity", Oi, T.; Aikawa, M.; Kidetake, H.; Ishida, T., *Z. Naturforsch.* **1988**, *43a*, 511.
30. "Molecular dynamics study of a lithium ion in ammonia", Hannongbua, S. V.; Ishida, T.; Spohr, E.; Heinzinger, K., *Z. Naturforsch.*, **1988**, *43a*, 572.

31. "Vapor pressure isotope effects in liquid and solid ammonia", King, T. V. ; Oi, T.; Popowicz, A.; Heinzinger, K.; Ishida, T., *Z. Naturforsch.*, **1989**, *44a*, 359.
32. "Isotope Separation", Spindel, W. ; Ishida, T. in *"Encyclopedia of Physics"*, Lerner, R. G.; Trigg, G. L., Eds., **1991**, VCH Publishers, New York.
33. "Isotope Separation", Spindel, W.; Ishida, T., *J. Chem. Educ.*, **1991**, *68*, 312.
34. "Molecular forces, isotope effects, and isotope separation", Ishida, T. in *"Isotope Separation and Chemical Exchange Uranium Enrichment"*, *Bull. Res. Lab. Nucl. React.*, Special Issue No. 1, **1992**, Fujii, Y.; Ishida, T.; Takeuchi, K., Eds., Tokyo Institute of Technology, Tokyo.
35. "Correlation of vibrational zero-point energy shifts of hydrides upon H/D isotope substitutions with electronegativity", Oi, T.; Ishida, T. [Submitted to *J. Chem. Soc. Jpn.*]

Editor/Co-Editor

1. *"Isotope Chemistry: Festschrift für Jacob Bigeleisen"*, *Z. Naturforsch.* **1989**, *44a*, Ishida, T.; Saunders, M.; Van Hook, W. A.; Wolfsberg, M., Eds.
2. *"Isotope Separation and Chemical Exchange Uranium Enrichment"*, *Bull. Res. Lab. Nucl. React.*, Special Issue No. 1, **1992**, Fujii, Y.; Ishida, T.; Takeuchi, K., Eds., Tokyo Institute of Technology.

Manuscripts in Preparation

201. "Influence of surface oxides on contact angle of water on polycrystalline platinum", Krebs, L. C.; Ishida, T.
202. "Electrochemical modification of polycrystalline platinum surface by poly(p-phenylene)", Krebs, L. C.; Ishida, T.
203. "Effective surface area and capacitance of edge plane of pyrolytic graphite in acetonitrile solutions", Minick, S. K.; Ishida, T.

I. Introduction

General objective of this project has been investigation of isotope effect phenomena and development of fractionation methods for stable isotopes. This part of present report consists of the following four sections:

Section II. Theoretical correlation of isotope effects with molecular forces and molecular structure.

Section III. Theoretical correlation of zero-point energy and its isotope effects with molecular forces and molecular structure.

Section IV. Experimental and theoretical studies of vapor pressure isotope effects.

Section V. Experimental and theoretical studies of fractionation of stable isotopes.

Section V-B deals with studies of hydrophobic catalysts for the hydrogen isotope exchange between dihydrogen and water. Results of these studies will be reported in some detail since they have not yet been published.

Following convention will be used for the reference numbers: The numbers below 100 refer to the publications that have resulted from the Department of Energy support, the numbers in 100's are other publications, and those in 200's refer to our studies performed under this support but the results of which have not been published. List of the first category is found under Publications on pages xv - xvii, one for the second category under References on pages 77-79, and one for the third category under Manuscripts in Preparation on page xvii.

Isotope effects look their simplest when viewed from the perspective point of the *reduced partition function ratio* (RPFR).⁽¹⁰¹⁾ The RPFR is the ratio of the reduced partition functions (Q_r and Q'_r) for a pair of isotopic molecules

$$\frac{s}{s'} f = \frac{Q_r}{Q'_r} = \frac{\left(\frac{Q}{Q'}\right)_{qm}}{\left(\frac{Q}{Q'}\right)_c} \quad (1)$$

In Eq. (1), Q and Q' are the partition functions of the heavier and lighter isotopic molecules in the isotopic pair under consideration, respectively; the subscripts qm , cl , and r refer to the quantum mechanical, classical, and reduced partition functions, respectively; s and s' are the symmetry numbers of the heavier and lighter isotopic molecules.

The RPFR is a property of a particular substitution at a given atomic position in a given substance. Value of RPFR directly depends on the molecular environment surrounding the isotopically substituted

atom. It is also indirectly influenced by other environmental factors such as the physical states, temperature, pressure, density, and intermolecular forces.

Both the equilibrium and kinetic isotope effects on all physicochemical properties are expressible as a ratio of, usually, two RPFRs. The great utility of *isotope effect* (IE) studies lie in the fact that, to the extent the Born Oppenheimer approximation holds, many structural and counting factors of partition functions cancel for the RPFR between the pair of isotopic molecules [cf. Eq. (1)]. Additional factors cancel for the ratio of two RPFRs. Two examples of such ratio of RPFRs will be given below. These examples represent a large majority of the isotope effects that have been studied in this project.

The equilibrium constant for an isotope exchange equilibrium,

is given by

$$K = \frac{[Q'(AX')/Q(AX)]_{eq}}{[Q'(BX')/Q(BX)]_{eq}} = \frac{[(s/s')f]_{BX}}{[(s/s')f]_{AX}} \cdot \frac{\left(\frac{A'}{A}\right)_{BX}}{\left(\frac{A'}{A}\right)_{AX}}. \quad (3)$$

Generally, A and B in Eq. (2) are polyatomic moieties of respective molecules. The isotope fractionation factor, α , due to Reaction (2) is expressible as

$$\alpha = \frac{[(s/s')f]_{BX}}{[(s/s')f]_{AX}}. \quad (4)$$

In both Eqs. (3) and (4) the mass- and moment of inertia factors contained in the classical partition functions cancel out for the balanced chemical equation such as Eq. (2).⁽¹⁰¹⁾ For an isotope exchange equilibrium in the gas phase, the only remaining factors of f_{BX} and f_{AX} are those due to the internal vibrations of the isotopic molecular pairs, AX/AX' and BX/BX'. For an equilibrium in a condensed phase, f_{BX} and f_{AX} also contain contributions from intermolecular interactions. The high-temperature limit of K is the classical equilibrium constant, K_{cl} ,

$$K_{cl} = \frac{(s'/s)_{BX}}{(s'/s)_{AX}}, \quad (5)$$

so that

$$\alpha = \frac{K}{K_{cl}}. \quad (6)$$

Both K and α are functions of temperature. However, the ratio of α and K is always a constant, which only depends on the symmetry numbers.

The second example is the isotopic vapor pressure ratio, which is expressible as

$$\ln \frac{P'}{P} = \left[\ln \frac{f_c}{f_s} \right] \cdot [\text{correction factors}], \quad (7)$$

where P' and P are the vapor pressures of the lighter and heavier isotopic substances, f_c/f_g is the ratio of the RPFR of the condensed phase to that of the vapor phase, and the "correction factors" correct for relatively small corrections such as the gas imperfection and liquid molar volume effects.

As a result of the cancellations of the otherwise dominant factors when the ratio of the ratios is evaluated, only those intra- and intermolecular interactions that are affected by the isotopic substitution manifest themselves in the isotope effects. Isotopic substitutions at strategic locations in a molecular substance is thus an extremely sensitive probe for elucidating for the molecular structures and molecular forces.

[blank page]

II. Correlation of Isotope Effects with Molecular Forces and Structure

The isotopic *reduced partition function ratio* (RPFR) of an ideal gas is

$$\ln (s/s')f = \sum_{i=1}^f \ln \frac{u_i}{u'_i} \frac{e^{-u_i/2} / (1 - e^{-u_i})}{e^{-u'_i/2} / (1 - e^{-u'_i})} , \quad (8)$$

where

$$u_i = \frac{h c \nu_i}{k T} , \quad (9)$$

and $f = 3N-6$ or $3N-5$. A self-consistent set of vibrational frequencies ν_i' (for the lighter isotopic molecule) and ν_i (for the heavier molecule) are obtained from vibrational eigenvalues λ_i' and λ_i ;

$$\lambda_i' = 4 \pi^2 \nu_i'^2 \quad \text{and} \quad \lambda_i = 4 \pi^2 \nu_i^2 \quad (10)$$

as solutions of

$$| H' - \lambda I | = 0 \quad \text{and} \quad | H - \lambda I | = 0 \quad (11)$$

where I is a unit matrix,

$$H' = \{ h'_{ij} \} \quad \text{and} \quad H = \{ h_{ij} \} , \quad (12)$$

and

$$H' = G' F \quad \text{and} \quad H = G F . \quad (13)$$

In Eq. (13), G is Wilson's G -matrix, and F is the corresponding matrix of harmonic force constants:

$$G' = \{ g'_{ij} \} , \quad G = \{ g_{ij} \} , \quad \text{and} \quad F = \{ f_{ij} \} . \quad (14)$$

Except for the molecules which possess a very high symmetry, Eq. (11) requires a numerical solution involving an algorithm which consists of a simultaneous diagonalization of two square, positive definite matrices, F and G . Such a procedure represents a black-box, as far as our physical understanding of isotope effects in terms of the molecular forces and structures is concerned.. We were in need of an analytic expression of $\ln (s/s')f$ in explicit terms of the elements of F and G matrices for such elucidations when this project was started.

II-A. Development of Theoretical Tools

The wanted tool is an approximation of an acceptable precision for the function, $\ln (s/s')f$, of Eq. (8), directly expressed in terms of f_{ij} 's, g_{ij} 's and the temperature. The first attempt toward this goal was Bigeleisen's Taylor series expansion⁽¹⁰³⁾:

$$\ln \frac{s}{s'} f = \sum_{m=1}^{\infty} \frac{(-1)^{m+1} B_{2m-1}}{2m (2m)!} \sum_{i=1}^f \delta u_i^2 \quad [All u \text{ and } u' < 2\pi] \quad , \quad (15)$$

where B_k are Bernoulli numbers ($B_1 = 1/6$, $B_3 = 1/30$, $B_5 = 1/42$, etc.), and

$$\delta u_i^{2m} = u_i'^{2m} - u_i^{2m} \quad . \quad (16)$$

This infinite series converges only when the largest of all u_i 's is smaller than 2π . This means that, for instance, at 300K all vibrational frequencies must be lower than 1310 cm^{-1} . The series is thus useful only at considerably low temperatures, e.g., $T < 200\text{K}$.

We have developed, in collaboration with the Bigeleisen group, several new approximations which eliminated the severe limitation on frequencies and improved the precision of the approximation in several steps.^(2, 7, 12, 14, 105, 106) They are all based on the principles of orthogonal polynomial approximation. In the following, these methods will be summarized in two groups, that is, the methods based on a direct expansion of $\ln (s/s')f$, and a second group of methods which utilize the diagonal matrices, $G_o = \{g_{ij} \delta_{ij}\}$ and $F_o = \{f_{ij} \delta_{ij}\}$ [δ_{ij} = Kronecker delta], as a zero-th order approximation.

II-A-1. Direct Orthogonal Polynomial Approximations

In an orthogonal expansion, a function $f(x)$ is expressed as a series of terms of mutually orthogonal functions. An orthogonal polynomial oscillates within a region of orthogonality. The error of a truncated series approximation of $f(x)$ thus oscillates in the domain in which the orthogonality is defined, rather than the error gradually increasing with the variable, x , moving away from the center of expansion.

The RPFR of Eq. (8) can be written as

$$\ln \frac{s}{s'} f = \sum_i [\ln b(u_i') - \ln b(u_i)] \quad , \quad (17)$$

where

$$\ln b(u) = -\ln u + \frac{u}{2} + \ln(1 - e^{-u}) \quad (18a)$$

$$= \sum_{k=1}^{\infty} \ln \left[1 + \left(\frac{u}{2\pi k} \right)^2 \right] \quad (18b)$$

An application of Lanczos' τ -method⁽¹⁰⁴⁾ on the function, $\ln(1+x)$, leads to the following orthogonal polynomial approximation^(105, 106) for the RPFR:

$$\ln \frac{s}{s'} f = \sum_{m=1}^n W_m(n, u'_{\max}, L) A_m \sum_{i=1}^f \delta u_i^{2m}, \quad (19)$$

where n is the order of the expansion, f the degree of freedom, L an integer parameter for the expansion used to isolate several lowest-order terms, and

$$A_m = \frac{(-1)^{2m+1} B_{2m-1}}{2^n (2m)!}, \quad (20)$$

and

$$W_m(n, u'_{\max}, L) = T(n, m, R_{L+1}) + \frac{1}{z(m)} \sum_{k=1}^L \frac{T(n, m, R_k) - T(n, m, R_{L+1})}{k^{2m}} \quad (21)$$

in which

$$T(n, m, R) = \frac{\sum_{p=m}^n \frac{(-1)^p C_n^p}{p^R}}{\sum_{p=0}^n \frac{(-1)^p C_n^p}{p^R}}, \quad (22)$$

and

$$R_k = \frac{1}{k^2} \left(\frac{u'_{\max}}{2\pi} \right)^2 \quad [k = L+1, L+2, \dots] \quad (23)$$

In Eqs. (21) and (22), $z(m)$ is the Riemann zeta function, and C_n^p is the coefficient of the orthogonal polynomial $P_n(x)$ used for the approximation,

$$P_n(x) = \sum_{m=0}^n C_n^m x^m \quad (24)$$

The usefulness of the orthogonal expansion, Eq. (19), lies in the fact that it converges for any frequency at any temperature and that, as in the Bigeleisen series of Eq. (15), the term $\sum_i \delta u_i^{2m}$ is expressible in terms of the trace of the m -th power of the product of F and G -matrices:

$$\begin{aligned} \sum_{i=1}^f \delta u_i^{2m} &= \left(\frac{\hbar c}{kT} \right)^{2m} \sum_{i=1}^f \delta \lambda_i^m \\ &= \left(\frac{\hbar c}{kT} \right)^{2m} \{ \text{Tr} [(FG')^m] - \text{Tr} [(FG)^m] \} \\ &= \left(\frac{\hbar c}{kT} \right)^{2m} [\delta \text{Tr} (H^m)] \end{aligned} \quad (25)$$

Through Eq. (25), $\ln(s/s')f$ and thus all isotope effects can be directly correlated to the molecular forces via the F -matrix and to the molecular geometry via the G -matrix. For instance, for $m = 1$ and 2 , we have

$$\sum_i \delta \lambda_i = \sum_i \delta(g_{ii}) f_{ii} + 2 \sum \sum_{i < j} \delta(g_{ij}) f_{ij} \quad , \quad (26)$$

and

$$\begin{aligned} \sum_i \delta \lambda_i^2 = & \sum_i \delta(g_{ii}^2) f_{ii}^2 + \sum \sum_{i \neq j} \delta(g_{ii} g_{jj}) f_{ij}^2 \\ & + 4 \sum \sum_{i \neq j} \delta(g_{ii} g_{jj}) f_{ii} f_{jj} \\ & + 2 \sum \sum_{i \neq j} \delta(g_{ij}^2) (f_{ii} f_{jj} + f_{ij}^2) \\ & + 2 \sum \sum \sum_{i \neq j \neq k} \delta(g_{ii} g_{jk}) f_{ij} f_{ik} \\ & + 2 \sum \sum \sum_{i \neq j \neq k} \delta(g_{ij} g_{ik}) (f_{ii} f_{jk} + f_{ij} f_{ik}) \\ & + \sum \sum \sum \sum_{i \neq j \neq k \neq l} \delta(g_{ij} g_{kl}) (f_{ii} f_{jk} + f_{ik} f_{jl}) \quad . \end{aligned} \quad (27)$$

The approximation for $\ln b(u)$ that corresponds to Eq.(19) for $\ln(s/s')$ is

$$\ln b(u) = \sum_{m=1}^n W_m(n, u'_{\max}, L) A_m u^{2m} \quad . \quad (28)$$

When the shifted Chebyshev polynomials of the first kind is used, examples of the errors that Eq. (28) yields are, with $L=0$:

6.6% (n=2),	0.12% (n=3),	and 0.0033% (n=4)	at u = 6;
7.1% (n=2),	2.0% (n=3),	and 1.6% (n=4)	at u = 16;
19.6% (n=2),	7.2% (n=3),	and 1.7% (n=4)	at u = 18.

A general improvement in the approximation was achieved by constructing an optional set of Jacobi polynomials for this mathematical problem.⁽¹⁰⁶⁾ The Jacobi polynomials, $P_n^{(\gamma, \delta)}(x)$, are a class of Gaussian hypergeometric functions characterized by the order, n, of the polynomial and two parametric constants, γ and δ . The optimal set of Jacobi polynomials and Eq. (28) yielded a three- to ten-fold improvements in the precision. For instance, with $L = 5$;

0.62% (n=2),	0.083% (n=3),	and 0.011% (n=4)	at u = 6;
2.8 % (n=2),	2.8 % (n=3),	and 0.66% (n=4)	at u = 12;
5.7 % (n=2),	2.1 % (n=3),	and 0.82 % (n=4)	at u = 18.

II-A-2. Orthogonal Expansion Methods with Diagonal Approximation

The precision of approximation of the orthogonal expansion methods was further improved without losing the explicitness of Eq. (19) and the contact with the physical significance^(12, 14) by recognizing a fact that the diagonal part of F-matrix,

$$F_0 = \{ f_{ij} \delta_{ij} \} \quad , \quad (29)$$

and the diagonal part of G-matrix,

$$G_0 = \{ g_{ij} \delta_{ij} \} \quad , \quad (30)$$

can be used as the zero-th order approximation for the actual matrices, $F = \{ f_{ij} \}$ and $G = \{ g_{ij} \}$, respectively. This fact was used by Wolfsberg in his perturbation approximation theory⁽¹¹⁾ for RPFR. Each element of the zero-th order approximation represents the contribution of each valence coordinate regarded to be an independent harmonic oscillator, and the higher order terms are contributions of inter-coordinate interactions. The RPFR for the hypothetical polyatomic molecule corresponding to the $\{F_0, G_0\}$ set is

$$\begin{aligned} \ln \frac{s}{s'} f &= \sum_{m=1}^{\infty} W_m A_m \left(\frac{hc}{kT} \right)^{2m} \sum_{i=1}^f \lambda_{i,0}^m \\ &= \sum_{m=1}^{\infty} W_m A_m \text{Tr}(\mathbf{H}_0^m) \end{aligned} \quad (31)$$

where

$$\text{Tr}(\mathbf{H}_0^m) = \sum_{i=1}^f f_{ii}^m g_{ii}^m \quad (32)$$

Since the magnitudes of the diagonal elements, f_{ii} and g_{ii} , are much greater than those of the off-diagonal elements, $\ln(s/s')f_0$ is a good zero-th order approximation for the real $\ln(s/s')f$. Then, the orthogonal expansion methods described in the preceding Section could be used to approximate the remainder. Thus,

$$\begin{aligned} \ln \frac{s}{s'} f &= \ln \frac{s}{s'} f_0 + \left(\ln \frac{s}{s'} f - \ln \frac{s}{s'} f_0 \right) \\ &= \ln \frac{s}{s'} f_0 + \sum_m W_m A_m \left(\frac{hc}{kT} \right)^{2m} \delta\Delta_0 \text{Tr}(\mathbf{H}^m) \end{aligned} \quad (33)$$

where

$$\begin{aligned} \delta\Delta_0 \text{Tr}(\mathbf{H}^m) &= \delta [\text{Tr}(\mathbf{H}^m) - \text{Tr}(\mathbf{H}_0^m)] \\ &= [\text{Tr}(\mathbf{H}'^m) - \text{Tr}(\mathbf{H}_0'^m)] - [\text{Tr}(\mathbf{H}^m) - \text{Tr}(\mathbf{H}_0^m)] \end{aligned} \quad (34a, b)$$

in which the prime refers to the lighter of the two isotopic molecule, as usual. All terms contained in the last expression of Eq. (34b) are generally much smaller than the terms $f_{ii} g_{ii}$ of $\ln(s/s')f_0$. For $m=1$ and $m=2$, Eq. (34) become

$$\delta\Delta_0 \text{Tr}(\mathbf{H}) = \sum_{i \neq j} \delta(g_{ij}) f_{ij} \quad (35)$$

and

$$\begin{aligned} \delta\Delta_0 \text{Tr}(\mathbf{H}^2) &= \sum_{i \neq j} [4\delta(g_{ii} g_{ij}) f_{ii} f_{ij} + \delta(g_{ij}^2)(f_{ij}^2 + f_{ii} f_{jj}) + \delta(g_{ii} g_{jj}) f_{ij}^2] \\ &+ \sum_{i \neq j \neq k} [2\delta(g_{ii} g_{jk}) f_{ij} f_{ik} + 2\delta(g_{ij} g_{ik})(f_{ii} f_{jk} + f_{ij} f_{ik})] \\ &+ \sum_{i \neq j \neq k \neq l} \delta(g_{ik} g_{jl}) f_{ii} f_{jj} \end{aligned} \quad (36)$$

In a variation of the diagonal-element approximation, Eq. (33), the zero-th order term $\ln (s/s')f_0$, is replaced by $\ln (s/s')f_d$, which is evaluated by using the diagonal terms of H_d , a matrix obtained by setting all off-diagonal term of the product of G and F matrices to zero:

$$H_d \equiv \{ h_{ij} \delta_{ij} \} \quad , \quad (37)$$

where

$$h_{ij} = \sum_k f_{ik} g_{kj} \quad , \quad (38)$$

so that the i-th eigenvalue of $|H_d - \lambda I| = 0$ is

$$\lambda_{i,d} = h_{ii} = \sum_k f_{ik} g_{ki} \quad . \quad (39)$$

Thus,

$$\ln \frac{s}{s'} f = \ln \frac{s}{s'} f_d + \sum_m W_m A_m \left(\frac{hc}{kT} \right)^{2m} \delta \Delta_d \text{Tr}(H^m) \quad , \quad (40)$$

where

$$\begin{aligned} \delta \Delta_d \text{Tr}(H^m) &\equiv \delta [\text{Tr}(H^m) - \text{Tr}(H_d^m)] \\ &= [\text{Tr}(H'^m) - \text{Tr}(H_d'^m)] - [\text{Tr}(H^m) - \text{Tr}(H_d^m)] \quad . \end{aligned} \quad (41)$$

An advantage of the $\ln (s/s')f_d$ approximation over the $\ln (s/s')f_0$ approximation is that the zero-th order approximation in the former takes into account some of the effects of the first-order interactions terms. For $m=1$ and $m=2$, Eq. (41) becomes

$$\delta \Delta_d \text{Tr}(H) = 0 \quad , \quad (42)$$

$$\begin{aligned} \Delta_d \text{Tr}(H^2) &= \sum_{i \neq j} [2\delta(g_{ii} g_{ij}) f_{ii} f_{ij} + \delta(g_{ij}^2) f_{ii} f_{jj} + \delta(g_{ii} g_{jj}) g_{ij}^2] \\ &+ \sum_{i \neq j \neq k} [2\delta(g_{ii} g_{jk}) f_{ij} f_{ik} + \delta(g_{ij} g_{ik}) (2f_{ii} f_{jk} + f_{ij} f_{ik})] \\ &+ \sum_{i \neq j \neq k \neq l} \delta(g_{ik} g_{jl}) f_{il} f_{jk} \end{aligned} \quad (43)$$

Although one is able to set a general mathematical error bound for the approximations for $\ln b(u)$, the same is not true for the approximations of $\ln (s/s')f$. This is because $\ln (s/s')f$ depends on particular distributions of the vibrational frequencies of molecules and their isotope shifts. In spite of these limitations, the precisions of the approximations are sufficiently high to have allowed us to find useful correlations of isotope effects with various structural features of real molecules. These results will be summarized in next section.

II-B. Correlation Studies

In terms of the orthogonal polynomial approximation, effects of molecular geometry on the isotope effect phenomena manifest themselves through the various elements of Wilson's G-matrix.⁽¹⁰⁸⁾ Hydrogen atoms in hydrogenous molecules are terminal atoms except in hydrogen-bonding structures. An isotope substitution at the positions of these hydrogen atoms and other end-atom substitutions lead to a type of isotope effects which are simple to analyze, mainly due to relatively simple differences in the G-matrices of these isotopic molecules.^(1-4, 6, 12, 14, 105-107) The substitutions at positions other than an end position, on the other hand, cause far-reaching effects down several chemical bonds and thus generally require a more careful attention.^(5, 12, 14, 34, 105, 106)

II-B-1. End-Atom Isotope Effects

In Figure 1, atoms 1, 4 and 5 are examples of end-atom, and atoms 2 and 3 are examples of non-end atom. Elements of G-matrix are generally given by⁽¹⁰⁸⁾

$$g_{ij} = \sum_{\text{All atoms}} \mu_{\alpha} \vec{s}_{i\alpha} \cdot \vec{s}_{j\alpha} \quad , \quad (44)$$

where μ_{α} is the reciprocal mass of atom α , and $\vec{s}_{i\alpha}$ is Wilson's s-vector for the coordinate i and atom α .

Referring to Figure 2, we have

$$\vec{s}_{r1,1} = -\vec{e}_{12} \quad , \quad \vec{s}_{r1,2} = \vec{e}_{12} \quad , \quad \vec{s}_{r1,3} = \vec{0} \quad , \quad (45)$$

and

$$\vec{s}_{r2,1} = \vec{0} \quad , \quad \vec{s}_{r2,2} = \vec{e}_{32} \quad , \quad \vec{s}_{r2,3} = -\vec{e}_{32} \quad . \quad (46)$$

Figure 1

Figure 2

where $\vec{e}_{\alpha\beta}$ is a unit vector directed from atom α to atom β . For the bending coordinate ϕ , the vectors $\mathbf{s}_{\phi,1}$ and $\mathbf{s}_{\phi,3}$ are perpendicular to the unit vectors $\mathbf{e}_{12}$ and $\mathbf{e}_{31}$, respectively, but the vector $\mathbf{s}_{\phi,2}$ is not perpendicular to either bond, 1-2 or 3-2, unless the equilibrium bond angle is 90° .

Thus, the diagonal elements are;

$$g_{11} = \mu_1 + \mu_2, \quad g_{22} = \mu_2 + \mu_3, \quad g_{\phi\phi} = \mu_1 \rho_1 + \mu_2 \rho_2 + \mu_3 \rho_3 \quad (47)$$

where

$$\rho_1 = (r_0/r_{12})^2, \quad \rho_3 = (r_0/r_{23})^2, \quad \rho_2 = r_0^2 (1/r_{12}^2 + 1/r_{32}^2 - 2 \cos \phi_0 / r_{12}r_{23}) \quad (48)$$

and the off-diagonal elements are;

$$g_{11,\phi} = (\mathbf{s}_{11,2} \cdot \mathbf{s}_{\phi,2}) \mu_2 = -\mu_2 (r_0 / r_{12}) \sin \phi_0 \quad (49a)$$

$$g_{22,\phi} = (\mathbf{s}_{22,2} \cdot \mathbf{s}_{\phi,2}) \mu_2 = -\mu_2 (r_0 / r_{23}) \sin \phi_0 \quad (49b)$$

$$g_{11,2} = (\mathbf{s}_{11,2} \cdot \mathbf{s}_{22,2}) \mu_2 = -\mu_2 \cos \phi_0 \quad (49c)$$

We note that the kinetic energy interaction between a bond-stretching motion and an angle-bending motion depends only on the mass of the central atom such as atom 2 in Figure 2 and not on the terminal atoms such as atoms 1 and 3 in Figure 2. This is due to the fact that the vector $\mathbf{s}_{\phi,1}$ is perpendicular to the vector $\mathbf{e}_{12}$. The isotopic difference $\delta g_{11,\phi}$ for an isotope substitution at Position 1 vanishes. In general, $\delta g_{ij} = 0$ between a stretching coordinate and a bending coordinate for an end-atom substitution. Furthermore, all off-diagonal elements g_{ij} between a bond-stretching motion and a torsion of an adjacent bond vanish due to the similar perpendicularity of the corresponding $\mathbf{s}$ -vectors.

We studied end-atom isotope effects in $\text{CO}_2^{(1)}$ and, H_2O , CH_2O , CH_4 , C_2H_4 , C_2H_6 and C_6H_6 ^(4, 12, 14) in the light of the orthogonal polynomial approximations of RPFR. For an end-atom substitution the only intercoordinate kinetic energy couplings (g_{ij}) that contribute to b_1 (s/s')f are the adjacent bend-bend interactions such as those among the three HCH angle bends sharing a C-H bond in methane. All other g_{ij} 's ($i \neq j$) vanish for the end-atom isotope effect. Furthermore, the contributions of interactions among the bending coordinates to isotope effect are small compared to the diagonal terms such as $f_{ii} \delta g_{ii}$ and $f_{ii}^2 \delta g_{ii}$ [cf. Eq. (26), (27), (35), (36), (42), (43)], since $g_{ij} f_{ij} \ll g_{ii} f_{ii}$, $g_{ij}^2 \ll f_{ii} f_{jj}$, and $f_{ij}^2 \ll f_{ii} f_{jj}$. Consequently, the diagonal element approximations [cf. Eq. (34) and (41)] which use only the diagonal terms of F- and G-matrices are generally within 1% of the exact values of b_1 (s/s')f at 300K for this type of isotopic substitution.

Hydrogen isotope effects are attributable to individual internal valence coordinates ⁽⁴⁾, and the isotope effects attributed to individual valence-coordinate force constants are nearly additive at room temperature. The point is illustrated in Table 1. The bond-stretching force constants account for at least 65% of the total H/D isotope effect at 300K and become even more dominant at higher temperature. The relative importance of stretching and bending forces in an end-atom substitution is due to their dominance in the first two term (the terms of $m=1$ and 2) in the orthogonal expansion.

Table 1. Contributions of Various Valence Coordinates to H/D-Isotope Effects in Hydrocarbons at 300K ^(a)

Isotopic par	ln (s/s')f @ 300K	Σ "Contri- bution" (%) ^(a)	C-H stretch f _{1j}	UC ^(c)	H-C-H bend f _{1j}	UC	H-C-C bend f _{1j}	UC	C-CH ₃ pwag ^(d) f _{1j}	UC	H-C-CC f _{1j}	UC	pwag f _{1j}	UC	H ₂ C=CH ₂ , C' f _{1j}	UC	C-CH ₃ , OPLA f _{1j}	UC	H-C-CC OPLA f _{1j}	UC	Adj HCHxHCH f _{1j}	UC	wag x CC-Tors f _{1j}	UC
Methane ^(a)																								
(d ₁ /d ₀) [JM]	2.4562	101.5	5.495	0.318	0.475	0.522																		
(d ₁ /d ₀) [HS]	2.3119	96.5	4.918	0.325	0.370	0.487															-0.078	-0.193		
Ethylene (d ₁ /d ₀)																								
(d ₁ /d ₀) [JM]	2.3206	106.2	5.026	0.333	0.598	0.594			1.133	0.178					0.626	0.159	0.802	0.165						
Ethylene (cis-d ₂ /d ₀)	4.6474	106.2	5.026	0.329	0.598	0.595			1.133	0.178					0.626	0.319	0.802	0.165						
Ethylene (gem-d ₂ /d ₀)	4.6492	104.5	5.926	0.331	0.598	1.182 [*]			1.133	0.332 [*]					0.626	0.305 [*]	0.802	0.325 [*]						
Ethylene (trans-d ₂ /d ₀)	4.6488	105.6	5.026	0.331	0.598	0.594			1.133	0.178					0.626	0.327	0.802	0.165						
Ethane (d ₁ /d ₀)																								
(d ₁ /d ₀)	2.3534	98.1	4.703	0.331	0.506	0.572	0.506	0.572																
Benzene ^(a)																								
(d ₁ /d ₀) [PI]	2.1441	102.8	5.063	0.322									0.870	0.658							0.378	0.748		
(d ₁ /d ₀) [nPI]	0.2656	111.3																						-0.087 -0.150

(a) For the sources of the molecular geometry and F-matrix, see Ref. (4). All f_{1j}'s are in units of mdyn / Å. Each bending coordinate is weighted by an appropriate equilibrium bond length.

(b) Sum of exact total contributions of all f_{1j}'s, expressed in percentage of exact ln (s/s')f at 300K.

(c) Pwag = In-plane wagging, CT = Crawford's torsion, OPLA = Out of plane wagging, Tors = Wilson's torsion.

(d) UC = Unit Contribution to ln (s/s')f by the f_{1j} at 300K, defined as the contribution of the f_{1j} divided by f_{1j} and by the number of times that the corresponding non-vanishing δg_{1j} occurs in the m = 1 term.

(e) [JM] = F-matrix by Jones and McDowell. [HS] = F-matrix by Hershorn and Shiner. [PI] = In-plane vibrations only. [nPI] = Out-of-plane vibrations only.

* In ethylene (gem-d₂), these coordinates contain double D-for-H substitutions.

In Table 1, it is noted that the unit contribution of C-H stretching force constant per unit value of (mdyn.Å⁻¹) is relatively constant among all molecules, while those for the H-C-H bending coordinate are more scattered. This is a consequence of relatively small effects of the second-order (m=2) terms involving the C-H coordinate compared to the first-order term. The only significant contribution of C-H stretch is (1/24) W_i(hc/kT)f_i δg_i, where f_i is the diagonal force constant for the C-H stretching and δg_i = 1/μ_H - 1/μ_D = 0.496 amu.

Generally as far as the H/D isotope effect are concerned, the hydrogenous bond stretching as a valence coordinate is a very good approximation of a normal coordinate. This is because, hydrogen atoms being the lightest atoms, do most of the vibrational motions against the other atoms, and the motions do not kinetically interact with other valence coordinates except between the adjacent bends that involve the isotopically substituted hydrogen.

Analysis of end-atom isotope effects other than the hydrogen isotope effect is similarly simple and the additivity holds well, although the bond-stretching valence coordinates involving other end-atoms are not as good an approximation of a normal coordinate as the stretching coordinate involving a hydrogen atom.

These observations lead to a simple method for calculating the end-atom isotope effects.⁽²⁾ It is possible to compute the RPFR at room temperature and above with a reasonable accuracy simply from the atomic masses and the stretching and bending force constants. The method is based on the first-order approximation to the method of Eq. (19):

$$\ln \frac{s}{s'} f \cong \frac{1}{24} \left(\frac{\hbar c}{kT} \right)^2 \sum_{\text{stretches \& bends}} (W'_i g'_{ii} - W_i g_{ii}) f_{ii} , \quad (50)$$

where W_i' and W_i are the first-order (m=1) modulating coefficients for the lighter and heavier isotopic molecules and found from one of the tables reported in References 105 and 106. Here it is sufficient to use λ_i' ≡ (μ_{end}' + μ) f_{ii} and λ_i ≡ (μ_{end} + μ) f_{ii} as a basis for the look-up values of the range of expansion in the W-tables. Here, the assumption, W_i' ≡ W_i, is adequate.

Deviations from the First Rule of The Mean

One of the most critical tests of various isotope effect data on a given molecule for the data's internal consistency is the tests on the sum rules for the isotopic vibrational frequencies and those on the rules of the mean for the isotopic thermodynamic data.

The first order sum rule⁽¹⁰⁸⁾ for the frequencies states that, for certain sets of isotopic molecules, a certain algebraic sum of the molecular sums of the squares of harmonic frequencies over the set vanishes. The set of isotopic molecules must satisfy the superposition rule, which requires that the set of molecules can be geometrically superimposed with appropriate signs in such a way that the atoms vanish at all positions. The sum rule is a direct and exact consequence of the assumptions of a harmonic potential, its

invariance with respect to isotopic substitution, and the equivalence of isotopically substituted positions. The higher order sum rules^(103, 109) are also based on the same assumptions.

There is a statistical mechanical analogue of each of the sum rules.^(103, 105, 109, 110) The most common ones are the rule of the mean related to the first order sum rule and the temperature-independent pre-exponential factor related to the Teller-Redlich product (3n-6) rule.⁽¹⁰³⁾ Thus, because of the first order sum rule, the equilibrium constant at room temperature for an isotopic disproportionation reaction such as

is very close to the high temperature classical limit, $s(d_0) s(d_2) / s(d_1)^2 = 4$. The usually small deviations from the rule of the mean value can be interpreted⁽⁶⁾ in the light of the $m = 2$ and higher order terms of orthogonal expansion of $\ln (s/s')$. It is possible to develop a systematics of the deviations from any n -th order rule of the mean and express the deviation in terms of the reciprocal masses of the substituted atoms and the harmonic force constants.

To facilitate a simple identification of the valence coordinates that would make non-vanishing contributions to the m -th order deviation from an n -th order rule of the mean and of those which do not contribute to the deviation, a new matrix, D-matrix, was introduced.⁽⁶⁾ The elements of a D-matrix are zero or unity, depending on whether an interaction between two valence coordinates makes a zero or finite contribution to the deviation. A systematic method for constructing D-matrix was developed. In general, presence of off-diagonal elements in F-matrix among the primarily bending motions, such as bond-angle bending, wagging and bond-torsion, tends to increase the deviation even in the cases of D/H substitutions. In the cases of multiple isotope substitutions in a molecule, proximity of the substituted atomic positions promotes significant couplings of higher order and leads to deviations from the rule of the mean.

II-B-2. Central Atom Isotope Effects

When an isotope substitution is made at a non-end atom position such as positions 2 and 3 in Figure 1, number of non-vanishing cross-interaction terms in the $m=2$ and higher terms increases exponentially. The $m=2$ term in the orthogonal expansion of $\ln (s/s')$ contains the isotopic difference in the trace of $(\text{FG})^2$;

$$\delta \text{Tr}(\text{H}^2) = \text{Tr}[(\text{FG}')^2] - \text{Tr}[(\text{FG})^2] \quad (51)$$

which further consists of the following seven types of terms: (1) $f_{ii}^2 \delta g_{ii}^2$, (2) $f_{ij}^2 \delta(g_{ii} g_{jj})$, (3) $f_{ii} f_{ij} \delta(g_{ii} g_{ij})$, (4) $(f_{ii} f_{ij} + f_{ij}^2) \delta(g_{ij}^2)$, (5) $f_{ij} f_{ik} \delta(g_{ii} g_{jk})$, (6) $(f_{ij} f_{ik} + f_{ii} f_{jk}) \delta(g_{ij} g_{ik})$, and (7) $f_{ij} f_{ik} \delta(g_{ij} g_{ik})$. In the end-atom isotope effects, all of these vanish except for those which only involve the diagonal elements

such as g_{ii} and g_{jj} and g_{ij} 's in which (i, j) are adjacent angle-bending coordinates. This is so, even when all off-diagonal f_{ij} 's were non-vanishing. In reality, the f_{ij} elements tend to be negligible between i and j that are few bonds away from each other.

In contrast, in the central atom isotope effects the following g_{ij} 's are in general non-vanishing and isotope-dependent:

- (i, j) = two stretches sharing the central atom,
- (i, j) = a stretch and a bend sharing a bond involving the central atom,
- (i, j) = a stretch and a bend sharing the central atom but not a bond,
- (i, j) = two bends sharing the central atom,

and (i, j) = two bends sharing a bond involving the central atom.

This has a consequence of extending the effects of an isotope substitution at a non-terminal atomic position down several bonds instead of merely to one or two bonds. The question of down how many chemical bonds an isotope substitution at an atomic position carries its effects depends on an interplay between the off-diagonal g_{ij} and interaction force constants f_{ij} . We examined these effects using prototype molecules, CO_2 , H_2O , CH_4 , C_2H_4 , C_2H_6 and C_6H_6 .^(5, 12, 15) The stretching forces dominate the isotope chemistry of carbon and oxygen, accounting for 70% to 90% of the RPFR among them. In general, valence-bond contributions for the central atom isotope effects are far less additive than those of the end-atom isotope effect. Contributions of individual force constants to the isotope effect do not obey the rules of the means as well as those for the end-atom isotope effects. The method of D-matrix is especially useful for finding the molecular features that cause the deviations from the rules of the means.

Isotope Effects at the Center of Highly Symmetric Molecules

Although the correlation of isotope effects with various features of a molecule is generally not a straightforward task, the analysis of effects of an isotopic substitution at an atomic position located on all symmetry elements of a highly symmetric molecule leads to a clear-cut correlation due, in large part, to an extensive factorization of the vibrational secular equation.⁽³⁴⁾ The structure of block-factoring of the secular equation is isotope-independent for these substitutions. This leads to a simple formula for the isotope shift in every vibrational frequency, in which all terms are expressed as an explicit function of the elements of F- and G-matrices.

In the highly symmetric molecules such as those in point groups T_d , O_h , and D_{3h} , many chemical bonds around the central atom are at 90° from each other, and many planes contained in the molecule are perpendicular with each other. This leads to many vanishing g_{ij} 's. A central atom substitution such as one at X in XY_4 , XY_6 , and XY_2Z_2 , does not change the molecular symmetry. Thus,, for the X'/X -isotope effect in XZ_2Y_2 , the irreducible representation,

$$\Gamma = 2 A_1' + 2 A_2' + 3 E_1' + 2 E_2' + E_1'' + E_2'' \quad , \quad (52)$$

is unchanged by the X'/X -substitution. Furthermore, only the two-dimensional block of A_2' , the three-dimensional block of E_1' and the two-dimensional block of E_2' are isotope-dependent, and every element of G -matrix in the A_2' , E_1' and E_2' blocks contains a term of μ_X , the reciprocal mass of atom X . Thus, each element of the corresponding H -matrix is a uniformly linear function of μ_X :

$$H_{ij} = a_{ij} \mu_X + b_{ij} \quad (53)$$

Since all algebraic equations of the orders up to three are analytically solvable, we have a means for obtaining the expressions for all vibrational eigenvalues of both isotopic molecules and their isotope shifts. For example, take a 2-dimensional block. The vibrational secular equation is of the form,

$$\lambda^2 - \alpha \lambda + \beta = 0 \quad , \quad (54)$$

where α and β are

$$\alpha = G_{11}F_{11} + G_{22}F_{22} + 2 G_{12}F_{12} \quad , \quad (55a)$$

$$\text{and} \quad \beta = (G_{11}F_{11} - G_{12}^2)(F_{11}F_{22} - F_{12}^2) \quad . \quad (55b)$$

They are in the forms,

$$\alpha = b_1 \mu_X + c_1 \quad , \quad (56a)$$

$$\text{and} \quad \beta = a_0 \mu_X^2 + b_0 \mu_X + c_0 \quad , \quad (56b)$$

in which the coefficient a_0 is a function of the force constants only, while b_1 and c_1 are functions of force constants and also linear functions of μ_Y and μ_Z .

Differentiating Eq. (54) and solving for $\delta\lambda$, one obtains

$$\delta\lambda = \frac{\lambda\delta\alpha - \delta\beta}{2\lambda - \alpha} \quad (57)$$

where $\delta\lambda = \lambda' - \lambda$, $\delta\alpha = \alpha' - \alpha$, and $\delta\beta = \beta' - \beta$. Thus,

$$\delta\alpha = b_1 \delta\mu_X \quad (58a)$$

$$\delta\beta \equiv (2a_0\mu_X + b_0) \delta\mu_X \quad (58b)$$

and

$$\left(\frac{\hbar c}{kT}\right)^2 \delta\lambda = \delta u^2 \equiv 2u\delta u \quad (59)$$

Equations (58b) and (59) hold well when X is a heavy atom. We see that $\delta\mu_X$ factors out of the numerator on the right-hand side of Eq. (57). The λ 's in Eq. (57) can be expressed in terms of the same geometric and force constant parameters as those used for the expressions of α and β . We thus have a means for explicitly correlating λ and $\delta\lambda$, and consequently u and δu , with the parameters such as molecular geometry, isotopic masses and the force constants.

Similarly, for a 3-dimensional blocks of vibrational secular equation, λ and $\delta\lambda$ can be expressed in terms of the molecular parameters. ⁽³⁴⁾

Then, the method of using the symmetry coordinates for the highly symmetric molecules can be combined with Bigeleisen's $G(u)$ - $S(u)$ approximation^(101, 103) for $\ln (s/s')f$;

$$\ln \left(\frac{s}{s'} \right) f = \sum_i G(u_i) \delta u_i \left[1 + \frac{S(u_i)}{2 G(u_i)} \frac{\delta u_i}{u_i} + \dots \right] \quad (60)$$

where

$$G(u) = \frac{1}{2} - \frac{1}{u} + \frac{1}{e^u - 1} \quad (61)$$

and

$$S(u) = \frac{1}{u} - \frac{u e^u}{(e^u - 1)^2} \quad (62)$$

If u_i and δu_i were calculable, even the first term of the $G(u)\delta u$ -approximation would generally give an order of magnitude better precision than any other correlation method. However, the precision required for a satisfactory evaluation of δu is far more stringent than the requirement for u . The present method for the explicit relationships for λ_i and $\delta \lambda_i$ presented in this section provides a reliable means for isotope effect correlation for the substitutions at the highly symmetric center of molecules having a high symmetry.

III. Correlation of Zero-Point Energy with Molecular Forces and Structures

We studied various theoretical aspects of vibrational *zero-point energy* (ZPE), its isotope shift, $\delta(\text{ZPE})$, and phase-shifts of the isotope shift, $\Delta\delta(\text{ZPE})$.^(13, 19, 22, 23, 25-27, 29, 36)

The ZPE is defined as

$$\text{ZPE} = \frac{1}{2} h \sum_i^f \nu_i = \frac{\pi}{2} \sum_i^f \lambda_i^{1/2} \quad , \quad (63)$$

where f is vibrational degree of freedom, and λ_i is the i -th vibrational eigenvalue obtained by solving

$$| \mathbf{H} - \lambda \mathbf{I} | = | \mathbf{G} \mathbf{F} - \lambda \mathbf{I} | = 0 \quad . \quad (64)$$

Since it is difficult to accurately approximate the function $f(x) = x$ in terms of powers of $x^{1/2}$, there had not been an accurate approximation for the ZPE in terms of the elements of $\mathbf{G}$ and $\mathbf{F}$ matrices. We developed such an approximation method based on the orthogonal polynomial series.

III-A. The ZPE as Explicit Function of Molecular Structure and Molecular Forces

In this part, we will first present the development of the mathematical tool and then give a summary of results of various applications to various isotope effect and non-isotope effect problems.

III-A-1. Development of Mathematical Tools

We based our development⁽¹⁹⁾ of mathematical theory for this purpose on Lanczos' orthogonal polynomial expansion⁽¹⁰⁴⁾ of function $x^{1/2}$ as follows:

$$x^{1/2} = \sum_{m=0}^n \alpha_m x^m \quad [0 \leq x \leq 1] \quad (65)$$

where

$$\alpha_m = \frac{C_n^m}{2m-1} \left[\sum_{j=0}^n \frac{C_n^j}{2j-1} \right]^{-1} \quad , \quad (66)$$

in which C_n^m is the coefficient of the m-th order term in a Jacobi polynomial of order n defined in the range, $0 \leq x \leq 1$.

Let the normal frequency spectrum of an N-atomic molecule be represented by the following ascending sequence of vibrational eigenvalues;

$$\lambda_1, \lambda_2, \dots, \lambda_i, \dots, \lambda_f (= \lambda_{\max}) \quad (67)$$

in which degenerate eigenvalues are counted separately. Corresponding to the sequence (67), we create a normalized sequence, that is, normalized to a certain value λ_0 , which would be determined later:

$$x_1, x_2, \dots, x_i, \dots, x_f (= x_{\max}) \quad (68)$$

$$\text{where } x_i = \lambda_i / \lambda_0 \quad (69)$$

$$\text{and } x_{\max} = x_f = \lambda_{\max} / \lambda_0 \quad (70)$$

The parameter λ_0 is completely arbitrary at this point. In order to take advantage of Eq. (65), however, we define the range of expansion, ξ , such that

$$x_{\max} \leq \xi \quad (71)$$

Then, the ZPE can be approximated by

$$ZPE = \frac{\hbar}{2} \sum_i \lambda_i^{1/2} \equiv \frac{\hbar}{2} \lambda_0 \sum_{m=0}^n \frac{\alpha_m}{\lambda_0^m} \sum_i \lambda_i^m \quad (72)$$

where α_m is

$$\alpha_m = \frac{1}{2m-1} \frac{C_n^m}{\xi^m} \left[\sum_{j=0}^n \frac{1}{2j-1} \frac{C_n^j}{\xi^j} \right]^{-1} \quad (73)$$

Further, we have

$$\sum_i \lambda_i^m = \text{Tr}(\mathbf{H}^m) = \text{Tr}[(\mathbf{F}\mathbf{G})^m] \quad (74)$$

For n=2, Eq. (72) becomes

$$ZPE = \frac{\hbar}{2} \sum_{i=1}^f [b_0(\lambda_0, \xi) + b_1(\lambda_0, \xi) \lambda_i + b_2(\lambda_0, \xi) \lambda_i^2] \quad (75)$$

where

$$b_0 = a_0 \lambda_0^{1/2} \xi^2 / D \quad (76)$$

$$b_1 = a_1 \lambda_0^{-1/2} \xi / D \quad (77)$$

$$b_2 = a_2 \lambda_0^{-3/2} / D \quad (78)$$

in which

$$D = a_0 \xi^2 + a_1 \xi + a_2 \quad (79)$$

The best values of the parameters λ_0 and ξ would be those that minimize the error defined by

$$S = \sum_i w(\lambda_i) \Delta_i^2 \quad (80)$$

where $w(\lambda_i)$ is a weighting function, which we took to be proportional to λ_i^k ,

$$w(\lambda_i) = A \lambda_i^k \quad (81)$$

and

$$\Delta_i = b_0(\lambda_0, \xi) + b_1(\lambda_0, \xi)\lambda + b_2(\lambda_0, \xi)\lambda^2 - \lambda_i^{1/2} \quad (82)$$

For a given k , an optimum set of (λ_0, ξ) is obtained by simultaneously solving

$$\partial S / \partial \lambda_0 = 0 \quad (83a)$$

$$\text{and} \quad \partial S / \partial \lambda_0 = 0 \quad (83b)$$

Three cases were explored: Case A, for which a constraint $\lambda_0 = \lambda_{\max}$ was set; case B, another special case in which the range of expansion ξ was equated to $x_{\max}$; and case C in which ξ and $\eta (= \lambda_{\max} / \lambda_0)$ were treated as independent of each other.

The optimization of S yields the following sets of solutions. Case A: three positive roots for ξ for every k value tested, ranging from $k = 0$ to 10^6 . Each of three roots, $\xi(a1)$, $\xi(a2)$, and $\xi(a3)$, leads to a distinct approximation. Case B: three k -dependent positive roots for ξ (and the corresponding values of λ_0), which are designated as $\xi(b1)$, $\xi(b2)$ and $\xi(b3)$, and two k -independent roots, $\xi(b4)$ and $\xi(b5)$. The three roots, $\xi(b1)$, $\xi(b2)$, and $\xi(b3)$, lead to an identical approximation, while $\xi(b4)$ and $\xi(b5)$ give two distinct approximations. Case C: two positive roots, $[\xi(c1)$ and $\eta(c2)]$ and $[\xi(c2)$, $\eta(c2)]$. The two roots produce an identical approximation.

Some representative values of the solutions of Eqs. (83a, b) for each of the three cases are shown in Tables 2, 3, and 4.

Table 2. Best Values of ξ for Case A, $\lambda_0 = \lambda_{\max}$ ⁽¹⁹⁾

k	$\xi(a1)$	$\xi(a2)$	$\xi(a3)$
0.0	0.849 12	1.834 37	2.147 21
0.5	0.876 48	1.665 32	3.179 50
1.0	0.898 83	1.706 55	3.752 41
1.5	0.918 03	1.759 90	4.160 45
2.0	0.934 77	1.810 41	4.470 11
2.5	0.949 49	1.855 61	4.714 30
3.0	0.962 53	1.895 49	4.912 21
3.5	0.974 15	1.930 65	5.076 03
4.0	0.984 57	1.961 72	5.213 94
4.5	0.993 96	1.989 31	5.331 67
5.0	1.002 45	2.013 94	5.433 38
10.0	1.057 29	2.164 22	5.997 85
15.0	1.085 31	2.235 18	6.237 23
20.0	1.102 31	2.276 35	6.369 54
∞^a	1.171 57	2.430 50	6.828 43

Table 3. Best Values of ξ for Case B, $\xi = \lambda_{\max} / \lambda_0$ ⁽¹⁹⁾

k	$\xi(b1)$	$\xi(b2)$	$\xi(b3)$
0.0	0.867 17	1.746 08	19.836 36
0.5	0.897 93		
1.0	0.926 22	1.566 35	20.751 94
1.5	0.950 22		
2.0	0.970 20	1.463 61	21.227 86
2.5	0.986 72		
3.0	1.000 30	1.404 27	21.472 68
3.5	1.011 40		
4.0	1.020 39	1.368 67	21.604 78
4.5	1.027 63		
5.0	1.033 38	1.347 15	21.678 12
6.0	1.041 40		
7.0	1.046 02		
8.0	1.048 36		
9.0	1.049 21		

^a The value of k higher than 10^6 , beyond which the roots are within 10^{-5} of their asymptotic limits.

Table 4. Best Values of ξ and η for Case C. ⁽¹⁹⁾

k	$\xi(c1)^a$	$\eta(c1)^a$	$\xi(c2)$	$\eta(c2)$
0.172663	2.438 38	1.443 47		
0.5	3.258 79	1.319 60		
1.0	3.856 21	1.246 17	11.566 89	3.737 96
1.5	4.264 91	1.202 28		
2.0	4.570 27	1.172 15	9.982 92	2.560 34
2.5	4.808 92	1.149 98		
3.0	5.001 20	1.132 93	9.202 11	2.084 57
3.5	5.159 66	1.119 39		
4.0	5.292 64	1.108 36	8.732 96	1.828 81
5.0	5.405 88	1.099 20		
7.0	5.788 69	1.069 74		
10.0	6.042 37	1.051 42		
15.0	6.269 62	1.035 75		
20.0	6.394 96	1.027 40		
∞	6.828 43	1.000 00

Figure 3. Relative errors in the ZPE approximations for the model system using various roots of Eqs. (83a,b) ⁽¹⁹⁾

For the model system which has its frequency distribution according to Eq. (82), the relative error, σ , of an approximation can be written as

$$\frac{\sigma_{ZPE}}{ZPE} = \frac{2k+3}{2k+2} \left(\frac{S}{\lambda_{\max}} \right)^{1/2} \quad (84)$$

This function is plotted in Figure 3.

III-A-2. Zero-Point Energy Approximation for n-Alkanes

For real molecules, actual precision of approximation depends on the particular distribution of frequencies. The best second-order approximation based on the Chebyshev polynomial $T_n^*(x)$ for n-pa-

Table 5. Best Approximations for n-Paraffins ⁽²²⁾

approx		complete spectra	out-of-plane spectra
A	parameters	a1 ($k = 0$)	a3 ($k = 0$)
	precision	-3.1% (C_1) to -0.3% (C_{10})	-1.3% (C_3) to -0.5% (C_{14})
B	parameters	b1 ($k = 0$)	b1 ($k = 0$)
	precision	-2.5% (C_1) to -0.03% (C_7)	-3.1% (C_3) to -2.7% (C_{14})
C	parameters	c1 ($k = 0.173$)	c1 ($k = 0.173$)
	precision	+0.1% (C_1) to +5.8% (C_{10})	+0.6% (C_3) to 1.4% (C_{14})

Figure 4. Absolute error and distribution of λ_i as functions of x for Case B at $k = 0$. ⁽²²⁾

ffins up to $C_{14}H_{30}$ are summarized in Table 5. Among all cases we studied⁽²²⁾ the best results for n-paraffins were obtained by equating the range of expansion to the largest normalized eigenvalue (Case-B) and setting $k = 0$; the absolute error is 2.5% for C_1 and lower than 1% for the molecules larger than propane. These results which are entirely different from the one expected from Figure 3 are due to the difference in the frequency distribution. Figure 4 illustrates this for Case B at $k = 0$. The unit interval of the histogram is $\Delta x = 0.01$, and the heights are normalized to unity for the most populous interval.

III-A-3. Hydrogen Isotope Effects in ZPE and Linear Frequency Sum Rules

For an isotope shift in ZPE, let us choose to use the same set of approximation parameters for the ZPE's of both isotopic molecules. Then, Eq. (75) becomes

$$\delta(\text{ZPE}) \equiv \frac{\hbar}{2} [b_1 \delta(\sum \lambda_i) + b_2 \delta(\sum \lambda_i^2)] \quad (85)$$

where $\delta(\sum \lambda_i) = \text{Tr}(H') - \text{Tr}(H)$, (86a)

$$\delta(\sum \lambda_i^2) = \text{Tr}(H'^2) - \text{Tr}(H^2) \quad , \quad (86b)$$

and $H' = G' F$ and $H = G F$. (87a,b)

The best approximation parameters for all single and multiple D-for-H substitutions in H_2O , HCHO , CH_4 , C_2H_4 , C_2H_6 , and C_6H_6 were found among the Cases of C. ⁽²³⁾ In Figure 5, the ordinate is

$$\frac{\Delta}{\lambda_{\max}^{1/2}} = \frac{\frac{1}{2} \hbar v_{\text{approx}} - \frac{1}{2} \hbar v}{\frac{1}{2} \hbar v_{\max}} \quad (88)$$

The actual errors for $\delta(\text{ZPE})$ of real molecules are strongly dependent on the distribution of eigenvalue shifts.

Generally, the approximations for $\delta(\text{ZPE})$ of all molecules can be analyzed by means of classification of the valence coordinates into four categories;

- n_b = number of H-X bonds,
- n_a = number of independent H-X-Y angles,
- n_t = number of torsional and other out-of-plane coordinates involving hydrogen,
- n_r = number of coordinates which do not directly involve motions of hydrogens.

Figure 5. Normalized error $\Delta/\lambda_{\max}^{1/2}$ as a function of normalized eigenvalue $\lambda/\lambda_{\max}$ for approximations of Type C. Numbers on the curves are k values. ⁽²³⁾

Thus, at $k = 0.173$ for Case C, the least-squares fit of the best-k data on the prototype molecules we studied yields⁽²³⁾

$$k_{\text{best}} = 0.019 n_s - 0.082 (n_s + n_t) + 0.994 \quad (89)$$

Linear frequency sum rule was derived on the basis of these orthogonal expansion.⁽²³⁾ The rule approximately holds for any set of isotopic molecules that satisfies a second-order frequency sum rule. The first-order⁽¹¹²⁾ and second-order⁽¹⁰⁹⁾ frequency sum rules exactly holds under the Born-Oppenheimer approximation and, respectively, correlate the sums of the squares and the fourth powers of vibrational frequencies of isotopic molecules. From Eq. (85), we have

$$\delta \Sigma v_i \equiv \frac{1}{2\pi} [b_1 \delta \Sigma \lambda_i + b_2 \delta \Sigma \lambda_i^2] \quad (90)$$

The second-order sum rule may be generally written as

$$\Sigma_j \alpha_j (\delta \Sigma \lambda_i^2)_j = 0 \quad (91)$$

where α_j is a sum-rule coefficient for the j -th isotopic pair, and the sum over j is taken over all pairs involved in the sum rule. When a second-order rule holds, then a first-order rule also holds among the same set of isotopic molecules and with the same coefficient. Thus, if Eq. (91) holds, then we also have

$$\Sigma_j \alpha_j (\delta \Sigma \lambda_i)_j = 0 \quad (92)$$

If we define a sum rule difference, Δ_s , by

$$\Delta_s = \Sigma_j \alpha_j (\delta \Sigma \lambda_i)_j \quad (93)$$

Eq. (90) then leads to

$$\Delta_{s, \text{approx}} = \frac{1}{2\pi} [b_1 \Sigma_j \alpha_j (\delta \Sigma \lambda_i) + b_2 \Sigma_j \alpha_j (\delta \Sigma \lambda_i^2)] \quad (94)$$

Then, $\Delta_{s, \text{approx}}$ vanishes if Eq. (91) and, consequently, Eq. (92) hold. If $\Delta_{s, \text{approx}}$ is a good approximation of Δ_s , the latter should vanish approximately. Thus

$$\Sigma_j \alpha_j (\delta \Sigma v_i)_j \equiv 0 \quad (95)$$

We called this relation an approximate linear frequency sum rule.

The error made in equating $\Sigma \alpha_j (\delta \Sigma v_i)_j$ to zero, $\varepsilon(\Delta_s)$, is

$$\varepsilon(\Delta_s) = \Delta_s - \Delta_{s, \text{approx}} = \Sigma_j \alpha_j \varepsilon[(\delta \Sigma v_i)_j] \quad (96)$$

where $\varepsilon[(\delta \Sigma v_i)_j]$ is the error in $(\delta \Sigma v_i)_j$. Since the relative error for $\delta \Sigma v_i$ varies approximately linearly with the extent of deuteration, the absolute error may be written as

$$\varepsilon[(\delta \Sigma v_i)_j] \equiv (\beta j + \gamma) (\delta \Sigma v_i)_{j, \text{approx}} \quad (97)$$

where j may be taken as j in (d_{j+1}/d_j) or in (d_j/d_0) . Any sum rule expression can be rewritten by using only the pairs of the form (d_{j+1}/d_j) or only the pairs of the form (d_j/d_0) . Therefore, no generality is lost by the approximate linear rule, Eq. (95), by attaching the special meaning to j as introduced by Eq. (97). Substitution of Eq. (97) into Eq. (96) and the use of Eqs. (90) - (94) yield

$$\varepsilon(\Delta_s) \equiv \beta \Sigma_j \alpha_j (\delta \Sigma v_i)_j \quad (98)$$

$(\delta \Sigma \nu_i)_j$'s for monosubstitutions (d_{j+1}/d_j) of equivalent atoms are alike, $\sum j \alpha_j$ is zero for any set satisfying the first-order sum rule, and the factor β is on the order of 10^{-2} . Therefore, $\varepsilon(\Delta_s)$ is likely to be a generally small quantity. Some examples follow:

Methane: $3(d_2/d_1) - (d_3/d_0)$

$$\Delta_s = 3 \times 1313.97 - 3940.77 = 1.14$$

$$\Delta_{s, \text{approx}} = 3 \times 1326.27 - 3978.80 = 0.01 \quad (\text{Case C1; } k = 0.6)$$

Ethylene: $2(d_3/d_1) - (d_4/d_0)$

$$\Delta_s = 2 \times 2602.27 - 5202.97 = -1.57$$

$$\Delta_{s, \text{approx}} = 2 \times 2628.05 - 5252.10 = 0.00 \quad (\text{Case C1; } k = 0.173)$$

Ethane: $(d_1/d_0) + (d_6/d_5) - (\text{trans-}d_2/d_1) - (d_3/1,1,2,2-d_4)$

$$\Delta_s = 1319.20 + 1349.61 - 1321.58 - 1348.20 = -0.97$$

$$\Delta_{s, \text{approx}} = 1319.19 + 1322.43 - 1319.26 - 1322.27 = -0.01 (\text{Case C1; } k = 0.3)$$

The "law of the mean" for the isotopic ZPEs⁽¹¹³⁾ incorrectly predicts that the sums of isotopic frequencies vanish (unconditionally) for all combinations of the isotopic pairs for which the number of deuteriums cancel. The rules presented here are applicable only for the sets on which a second-order rule holds and yield a precision which is below 0.1%.

III-A-4. Additivity for Zero-Point Energy

The concept of additivity for the vibrational ZPE is an old one.⁽¹¹⁴⁻¹²²⁾ However, the earlier attempts were empirical in nature and without much theoretical foundation. Most of them used a least-squares procedure on the experimentally observed fundamentals and thus lack internal consistency and a common basis for comparison between different authors.

The present orthogonal approximation for the ZPE has provided the needed theoretical foundation.⁽²⁵⁾ We showed that the earlier rules with varying levels of sophistication can be derived as special cases of the more general rule. From Eq. (75), we obtain

$$\begin{aligned} \frac{2}{\hbar} (ZPE) \equiv & \sum_i C_i^0 + b_1 \sum_{i < j} C_{ij}^1 + b_2 \sum_{i < j} \sum_{m=1}^4 C_{ij}^2(m) \\ & + b_2 \sum_{i \neq j} \sum_{(j < k)} \sum_{m=1}^3 C_{ijk}^2(m) + b_2 \sum_{i < j} \sum_{k < l} \sum_{i < k} \sum_{j < l} \sum_{m=1}^2 C_{ijkl}^2(m) \quad , \end{aligned} \quad (99)$$

where

$$C_i^0 = (g_{ii} f_{ii})^{1/2} \quad , \quad (100)$$

$$C_{ij}^1 = 2 g_{ij} f_{ij} \quad , \quad (101)$$

$$C_{ij}^2(1) = 2 g_i g_j f_{ij}^2 \quad , \quad (102)$$

$$C_{ij}^2(2) = 4 g_{ij} f_{ij} (g_i f_{ii} + g_j f_{jj}) \quad , \quad (103)$$

$$C_{ij}^2(3) = 2 g_{ij}^2 f_{ij}^2, \quad (104)$$

$$C_{ij}^2(4) = 2 g_{ij}^2 f_{ii} f_{jj}, \quad (105)$$

$$C_{ijk}^2(1) = 4 g_{ij} g_{jk} f_{ij} f_{ik}, \quad (106)$$

$$C_{ijk}^2(2) = 4 g_{ij} g_{ik} f_{ii} f_{jk}, \quad (107)$$

$$C_{ijk}^2(3) = 4 g_{ij} g_{jk} f_{ij} f_{ik}, \quad (108)$$

$$C_{ijk}^2(4) = 4 g_{ij} g_{ik} f_{ii} f_{jk}, \quad (109)$$

$$C_{ijk}^2(5) = 4 g_{ij} g_{jk} f_{ii} f_{jj}, \quad (110)$$

Each of these C-terms has a physical meaning and can be logically prorated and assigned to the valence coordinates involved in them. For instance, one half of the term, $C_{ij}^2 = 2 g_{ij} f_{ij}$ can be assigned to each of the coordinates i and j . A generalization of such proration for all first-order and second-order C-terms contained in Eq. (99) leads to an expression,

$$\text{ZPE} \equiv \frac{\hbar}{2} \left[\sum_p N_p \Gamma_p + \sum_{(pq)} N_{(pq)} \Gamma_{(pq)} \right], \quad (111)$$

where N_p is the number of valence coordinates of type p (e.g., C-H stretch, H-C-H bend in CH_2 group) contained in the molecule, and Γ_p is the diagonal (i.e., non-interactive) contribution of one of the equivalent coordinates of type p . The quantities $N_{(pq)}$ and $\Gamma_{(pq)}$ are similar to the diagonal terms except that the one with the subscript (pq) refer to an interaction of a particular type between a valence coordinate of type p and another of type q . There can be more than one type of interaction between a given set of coordinate types, p and q . For instance, a C-H stretch and an H-C-H bend may share a terminal H-atom in one type of interaction, or share the C-atom in another interaction type, or share the C-H bond in still another. Table 6 is an example of the additivity parameters, Γ_p and $\Gamma_{(pq)}$, for n -alkanes.

Table 6. Contributions per Coordinate and Major Corrections per Intercoordinate Interactions to the Frequency Sum of $\text{C}_n\text{H}_{2n+2}$.⁽²⁵⁾

types of coordinates and interactions			unit contributions and corrections, ^a cm^{-1}			
SS-1 ^a notation	description ^b of p and (pq)	N_p and $N_{(pq)}$ ^c	zeroth order	first order	second order	tot $\Gamma_p + \Gamma_{(pq)}$
r	CH str (CH_3)	$6, (n \geq 2)$	2931.2	0	-35.2	2896.1
d	CH str (CH_2)	$2(n-2), (n \geq 3)$	2881.6	0	-40.7	2840.9
R	CC str ($\text{CH}_3\text{-CH}_3$)	$1, (n = 2)$	1118.9	0	-29.2	1089.7
$R(I)$	CC str ($\text{CH}_3\text{-CH}_2\text{-}$)	$2, (n \geq 3)$	1106.9	0	-28.1	1078.8
$R(II)$	CC str ($\text{-CH}_2\text{-CH}_2\text{-}$)	$n-3, (n \geq 4)$	1119.4	0	-34.0	1085.4
α	HCH bend (CH_3)	$6, (n \geq 2)$	1301.4	0	-79.2	1222.3
δ	HCH bend (CH_2)	$n-2, (n \geq 3)$	1313.5	0	-77.8	1235.7
β	HCC bend (CH_3)	$6, (n \geq 2)$	1042.0	0	-58.2	983.7
γ	HCC bend (CH_2)	$4(n-2), (n \geq 3)$	1065.4	0	-60.9	1004.5
ω	CCC bend	$n-2, (n \geq 3)$	500.9	0	-31.0	469.9
τ	C-C torsion	$3(n-1), (n \geq 2)$	165.9	0	-3.9	162.0
$R\beta, R\gamma$	(CC) $\times$ (HCC); CC common	$2(2n-1), (n \geq 2)$	0	-176.8	14.8	-162.0
$R\omega$	(CC) $\times$ (CCC); CC common	$2(n-2), (n \geq 3)$	0	-168.8	11.6	-157.2
$\gamma\gamma$	(HCC) $\times$ (HCC); CH common (CH_2)	$4(n-2), (n \geq 3)$	0	-94.8	5.8	-89.0
$\omega\omega(I)$	(CCC) $\times$ (CCC); CC common	$n-3, (n \geq 4)$	0	82.2	-5.2	77.0
$\beta\omega$	(HCC) $\times$ (CCC); CC common, H and C in trans	$2, (n \geq 3)$	0	66.8	-5.8	61.0
$\beta\omega(I)^*$	(HCC) $\times$ (CCC); CC common, H and C in gauche	$4(n-2), (n \geq 3)$	0	27.0	-2.4	24.6
$\beta\omega(I)^\dagger$	(HCC) $\times$ (CCH); CC common, H and H in trans	$2(n-1) + m, (n \geq 2)^*$	0	103.2	-16.8	86.4

^a Schachtschneider and Snyder's notation. ^b AX str means an A-X stretching coordinate. $(p) \times (q)$ denotes an interaction between a coordinate of type p and a coordinate of type q , and specification for a particular kind of interaction is given after a semicolon. ^c N_p = number of coordinates of kind p . $N_{(pq)}$ = number in intercoordinate interaction of kind (pq) . The constraints are given in the parentheses. ^d Γ_p is the contribution to the frequency sum per coordinate of type p , expressed in cm^{-1} . $\Gamma_{(pq)}$ is the correction to the frequency sum per interaction, expressed in cm^{-1} . ^e Also, $\gamma\omega(g)$. ^f Also, $\beta\gamma(I)$ and $\gamma\gamma(I)$. ^g $m = 1$ for $n = 2$; $m = 0$ for $n \geq 3$.

As Table 7 illustrates, these parameters yield the results which are within 0.7% of exact values of ZPE.

Table 7. Approximations for the Frequency Sums of n-Alkanes ⁽²⁵⁾

n	Exact cm ⁻¹	Additivity	
		Value cm ⁻¹	% error
2	31,487	31,475	- 0.038
3	43,677	43,395	- 0.65
4	55,809	55,429	- 0.68
5	67,917	67,435	- 0.71
6	80,011	79,442	- 0.71
7	92,101	91,448	- 0.71
8	104,187	103,454	- 0.70
9	116,271	115,469	- 0.69
10	128,354	127,467	- 0.69

Several simpler additivity rules of various precisions were also derived starting with Eqs. (99) and (111). The table of additivity parameters were also constructed for alkyl chlorides. ⁽²⁵⁾

III-A-5. Other Applications

The expressions for ZPE and $\delta(\text{ZPE})$ presented here formed a basis for various correlation studies. The following is an account of these studies.

Valence Coordinates and Hydrogeon Isotope Effects in ZPE ⁽²⁶⁾

From Eq. (85), we have for the isotopic difference in the frequency sums,

$$\delta \Sigma \omega_i \equiv \beta_1' \delta \Sigma \lambda_i + \beta_2' \delta \Sigma \lambda_i^2 \equiv \beta_1' S_1 + \beta_2' S_2, \quad (112)$$

where β_1' and β_2' are related to b_1 and b_2 in the obvious manner, and S_1 and S_2 represent the sums of the first-order and second-order contributions, respectively.

Using the theory for the ZPE-additivity reported above, the isotope shift in the frequency sum can also be expressed as a sum of contributions of the valence coordinates. Thus,

$$S_1 = \Sigma_p T_1(p) + \Sigma_{(pq)} T_1(pq) \quad (113)$$

$$\text{and} \quad S_2 = \Sigma_p T_2(pq) + \Sigma_{(pq)} T_2(pq) \quad (114)$$

in which $T_1(pq)$ is the sum of all terms of the form $2 f_{ij} \delta g_{ij}$ where i is a coordinate of type p , and j a coordinate of type q , while $T_2(pq)$ is the sum of all second-order interaction terms, $T_2^{mn}(pq)$, each containing different types of geometric factor as in Eq. (27). Although we can conceive of various additivity formula for $\delta \Sigma \omega_i$, the best one that gave the physical insight without becoming too cumbersome and without losing the precision is

$$\delta \Sigma \omega_i \equiv S_1 + \Sigma_p T_2^1(p) = S_1 + \Sigma_p n_p f_p^2 \delta g_p^2, \quad (115)$$

where n_p , f_p , and g_p are the number of type- p coordinates and the diagonal elements of F- and G-matrices, respectively. The precision is 0.7% or better. This is illustrated by Table 8 ⁽²⁶⁾ for d_1/d_0 -substitution in methane.

Table 8. Major Contributions to Isotope Shift in the Frequency Sum for CH_3D / CH_4 ^(a)

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
Coordinate type (p) or Interaction type (pq)	f_p or f_{pq} /mdyn Å ⁻¹	n_p or n_{pq}	$T_1(p)^{b,c}$ or $T_1(pq)$	$T_2(p)^{b,c}$ or $T_2(pq)$	$T_3(p)^{b,c}$ or $T_3(pq)$	$\frac{(4)^d}{(2) \times (3)}$	$T_4(p)^{b,c}$	$TF(p)$ or $TF(pq)$, mÅ ⁻¹ ^e	(4)+(8) ^b
C-H stretch	4.918	1	1137.4 (92.4)	-202.1 (-16.4)	935.3 (76.0)	231.3	-202.4 (-16.4)		935.1 (75.9)
HCH bend ^f	0.370	3	256.7 (20.8)	-11.8 (-1.0)	244.9 (19.9)	231.3	-8.1 (-0.7)	[3]=-3.4; [4]=-1.5 (-0.3) (-0.1)	248.5 (20.2)
(HCH)X(HCH); -0.0779 adjacent ^g		3	54.1 (4.4)	-2.9 (-0.2)	51.2 (4.2)	-231.3	0.0 (0.0)	[2]=-1.5; [3]=-3.2; [5]=2.2 (-0.1) (-0.3) (0.2)	54.1 (4.4)
Total (approx) ^d			1448.2 (117.6)	-216.8 (-17.6)	1231.4 (100.0)		-210.5 (-17.1)		1237.7 (100.5)
Total (exact) ^h			1448.2 (117.6)	-216.4 (-17.6)	1231.8 (100.0)		-210.5 (-17.1)		1237.7 (100.5)

^a F-matrix taken from Hartshorn and Shiner. ⁽¹²³⁾ The approximation of Case C1 is used; $k = 0.77$, $\xi = 3.614329$, $\eta = 1.274406$, $\omega = 3022.31$ cm⁻¹ and $\delta\Sigma\omega(\text{exact}) = 1241.45$ cm⁻¹. ^b All T terms are in cm⁻¹. The numbers in parentheses below the wavenumber values are percent of $\delta\Sigma\omega(\text{exact})$. ^c In units of Å mdyne⁻¹ cm⁻¹. ^d The sum of the contributions tabulated. ^e The sum of all contributions, both tabulated and non-tabulated. Contributions whose absolute magnitudes are less than 0.1% of $\delta\Sigma\omega(\text{exact})$ are not tabulated. ^f Number in the brackets is the type of interaction, m. ^g All bending coordinates are weighted by the equilibrium C-H bond length. The adjacent bends are two bends that share a C-H bond.

Classification of Molecules According to $\delta(ZPE)$ for H/D-Isotope Effect ⁽²⁷⁾

Although all H-X stretching coordinates are nearly independent of other vibrations in the molecule, $\delta(ZPE)$ is still affected by the molecular geometry near the site of the H/D substitution. A classification presented in Table 9 facilitated a finer tuning for a higher precision and a better correlation of the isotope effect. The basis of Table 9 is a study of a few dozens of hydrogenous molecules. In the Table, Y generally represents a molecular moiety rather than an atom.

Table 9. Classification of Molecules for H/D Isotope Shifts in Zero-Point energies^{a)}

Group	Description	$\bar{\lambda}_0$ ^{b)}	Representative molecule				
			molecule	pair	λ_0	% error at $\bar{\lambda}_0$	Ref.
Ia	HX-Y [Linear]	0.277	HCN	d_0/d_1	0.259	-0.72	4
Ib1	HX-Y-Z [Linear]: (Z=H)	0.264	C ₂ H ₂	d_0/d_1	0.266	0.07	5
Ib2	(Z≠H)	0.197	HCCF	d_0/d_1	0.206	0.37	5
Ic	HX-Y [Bent]	0.427	HOCl	d_0/d_1	0.442	0.40	6
Id	HX...Y [Planar]	0.278	C ₂ H ₄	d_0/d_1	0.269	-0.35	7
Ie	HX...Y [Nonplanar] ^{d)}	0.322 (0.278)	CHF ₃	d_0/d_1	0.409	2.78	8
			CHCl ₃	d_0/d_1	0.299	-0.90 (0.86) ^{g)}	5
			CHBr ₃	d_0/d_1	0.257	-2.56 (-0.84) ^{g)}	5
IIa	H ₂ X	1.052	H ₂ O	d_0/d_1	1.005	-0.28	9
IIb	H ₂ X-Y [Planar]	0.718	H ₂ CO	d_0/d_1	0.742	0.31	10
IIc	H ₂ XY ₂ [Nonplanar]	0.441	CH ₂ F ₂	d_0/d_2	0.498	1.38	11
IIId1	H ₂ X-YZ ₂ [Planar]: (Z=H)	0.398	CH ₂ CH ₂	CH ₂ /CD ₂	0.401	0.08	12
IIId2	(Z≠H)	0.272	CH ₂ CF ₂	d_0/d_2	0.276	0.16	5
IIe	H ₂ X...Y [Nonplanar] ^{e)}	0.400	CH ₂ CH ₂ Cl	d_0/d_2	0.390	-0.30	5
			CH ₂ CH ₂ CH ₃	CH ₂ /CD ₂	0.410	0.26	5
IIIa	H ₂ X	0.723	NH ₃	d_0/d_1	0.677	-0.64	13
IIIb	H ₂ X-Y [Nonplanar] ^{d)}	0.462 (0.437) ^{e)}	CH ₃ F	d_0/d_2	0.537	1.69	14
			CH ₃ Cl	d_0/d_2	0.463	0.46 (0.65) ^{g)}	5
			CH ₃ Br	d_0/d_2	0.432	-0.76 (-0.13) ^{g)}	5
			CH ₃ I	d_0/d_2	0.415	-1.21 (-0.58) ^{g)}	5
IIIc	H ₂ X-Y-Z...W [X-Y-Z Linear]	0.528	CH ₃ CCH	d_0/d_2	0.574	0.91	5
IIId	H ₂ X-Y...Z [Around Y nonlinear] ^{e)}	0.419	CH ₂ CH ₂ Cl	CH ₂ /CD ₂	0.436	0.46	5
			CH ₂ CH ₂ CH ₃	CH ₂ /CD ₂	0.442	0.63	5
IV	H ₄ X	0.686	CH ₄	d_0/d_1	0.733	0.64	15
				d_0/d_2	0.646	-0.59	

^{a)} Case-B approximation. ^{b)} The group average of the optimum k . ^{c)} The first-named isotopic species is the reference molecules from which the $\bar{\lambda}_{\text{ref}}$ was obtained. ^{d)} With no possibility of torsion involving the H atom being substituted. ^{e)} Alternate k_0 value obtained by excluding fluoro-compounds. ^{f)} The numbers in the parentheses are the percent errors obtained by using the alternate k_0 listed in the parentheses in the third column. ^{g)} With possibility of torsions involving the H atom being substituted.

$\delta(\text{ZPE})$ in Hydrides and Electronegativity^(29, 35)

As expected from the additivity for $\delta(\text{ZPE})$, a linear relationship was found between the square root of A-H stretching force constants in AH_n ($n=1\sim 4$) molecules and $\delta(\text{ZPE})$ for H/D-substitution.⁽²⁹⁾ Figure 6 demonstrates the fact that the contributions of f_{ii} ($i = \text{stretch}$) to $\delta(\text{ZPE})$ through all interaction terms such as f_{ii} , f_{ij} , $\delta(g_{ii}g_{jj})$ and $f_{ii}f_{jj}$, $\delta(g_{ij}^2)$ are negligible. A similar plot, Figure 7, for the H-A-H bending contributions is more scattered, reflecting some significant contributions by these interaction terms.

Figure 6. Correlation between stretching force constant and its contribution to $\delta(\text{ZPE})$.

Figure 7. Correlation between bending force constant and its contribution to $\delta(\text{ZPE})$.

Since the first introduction of the concept of electronegativity by Pauling⁽¹²⁸⁾ in 1930's, various attempts have been made for improved definitions of electronegativity. Among them are; ones by Wilmshurst (χ_w)^(129, 134), by Inomata and Masuda (χ_i)⁽¹³⁰⁾, and by Boyd and Edgecome (χ_B)⁽¹³¹⁾, and the unshielded core potential (V_B) as a measure of electronegativity by Luo and Benson.⁽¹³²⁾ Gordy's scale is based on numbers of electrons in the valence shells and the single bond covalent radii of the atoms.⁽¹³³⁾ Wilmshurst extended Gordy's formula to include the electronegativity calculations for various molecular

moieties. Boyd and Edgecombe obtained their atomic and group electronegativity values on the basis of the topological properties of the electron density distribution in molecules.

It can be shown⁽³⁵⁾ that the contribution of H-X stretching motions to $\delta(\text{ZPE})$ increases linearly with the electronegativity of X, where X is either an atom or a group of atoms. Wilmshurst's χ_w is

$$\chi_w = g (1 + m_H / m_X)^{-1/2} \nu_{\alpha} - h \quad , \quad (116)$$

where m_α is the mass of atom α , and g and h are constant property of X. No isotope effects were considered by these authors. However, since all electronegativities must be isotope-independent in accordance with Born-Oppenheimer principle, the parameter g must be a function of isotopic mass in such a way to cancel out the isotope dependence of the factors, $(1 + m_H / m_X)^{-1/2}$ and ν_{α} . If one assumes that the bond, H-X, in a polyatomic environment are independent harmonic oscillator (which assumption we know is a good one, as we have discussed throughout Sections II-A and II-B of the present report), the parameter g for the H/D isotope effects must be in a form,

$$g = g_0 \sqrt{m_H} \quad \text{or} \quad = g_0 \sqrt{m_D} \quad , \quad (117a,b)$$

while h is independent of isotope substitutions. With this modification of χ_w , the stretching frequency shift, $\delta\nu_{\alpha}$, must have a linear dependence on the electronegativity. Thus, for the D/H-isotope effect, we have

$$\delta(\text{ZPE})_{\alpha} = r \chi + s \quad . \quad (118)$$

Table 10 summarizes our comparison, in which

$$\delta(\text{ZPE}) \equiv \delta(\text{ZPE})_{\alpha} + \delta(\text{ZPE})_{\text{bend}} \quad (119)$$

Figure 8 illustrates a linearity for each main group, IVA, VA, VIA, and VIIA, of the periodic table. The average deviation from the linear line is better than several wavenumbers.

Table 10. Components of $\delta(\text{ZPE})$ and electronegativity

Molecule	$\delta(\text{ZPE})$	$\delta(\text{ZPE})_{\text{str}}$	$\delta(\text{ZPE})_{\text{bond}}$	CORR	Atom or group	χ_{W}	χ_{I}	χ_{S}	V_{B}
NF	568.9	568.9			F	3.94	3.99	4.00	9.915
NCI	422.9	422.9			Cl	3.00	3.00	3.05	7.04
NBr	381.8	381.8			Br	2.68	2.68	2.75	6.13
NI	334.6	334.6			I	2.36	2.36		5.25
H ₂ O	632.5	545.9	87.5	-0.9	-OH	3.86	3.72	3.64	8.11
H ₂ S	463.3	393.8	69.5	0.0	-SH	2.61	2.66	2.63	5.77
H ₂ Se	407.6	349.5	58.2	-11.1	-SeH	2.37	2.36	2.47	5.13
NH ₃	659.0	500.0	158.7	2.3	-NH ₂	3.40	3.02	3.10	6.67
PH ₃	485.0	355.7	133.0	-3.7	-PH ₂	2.20	2.19	2.17	4.55
AsH ₃	450.4	338.0	156.7	-44.3	-AsH ₂	2.03	2.05	2.26	4.20
SbH ₃	396.4	304.6	116.3	-24.5	-SbH ₂	1.76	1.86		3.55
CH ₄	645.4	440.4	203.7	1.3	-CH ₃	2.63	2.47	2.56	5.19
SiH ₄	474.3	343.0	131.5	-0.2	-SiH ₃	1.76	1.91	1.91	3.41
GeH ₄	464.3	337.8	127.2	-0.7	-GeH ₃	1.69	1.88	2.05	3.24
SnH ₄	405.0	308.0	97.8	-0.8	-SnH ₃	1.51	1.71		2.83

* All the $\delta(\text{ZPE})$, $\delta(\text{ZPE})_{\text{str}}$, $\delta(\text{ZPE})_{\text{bond}}$ and CORR values are given in units of cm^{-1} .

Figure 8. Plot of $\delta(\text{ZPE})_{\text{str}}$ vs. Wilmsbury's electronegativity

III-B. Isotope Effects in Zero-Point Energy Shifts upon Condensation

One of the most important factors that determine the isotopic difference in the vapor pressures is the isotope shift in the ZPE upon condensation of gaseous molecules, $\delta\Delta(\text{ZPE})$;

$$\delta\Delta(\text{ZPE}) = \frac{1}{2} hc \sum_i [\delta\omega_i(\text{g}) - \delta\omega_i(\text{c})] \quad (120a)$$

$$\text{and} \quad \Delta\delta(\text{ZPE}) = \frac{1}{2} hc \sum_i [\Delta\omega_i' - \Delta\omega_i] \quad (120b)$$

$$\text{where} \quad \delta\omega_i = \omega_i' - \omega_i \quad (121a)$$

$$\text{and} \quad \Delta\omega_i = \omega_{i,\text{gas}} - \omega_{i,\text{condensed}} \quad (121b)$$

$$\text{so that} \quad \delta\Delta(\text{ZPE}) = \Delta\delta(\text{ZPE}) \quad (122)$$

Ordinarily, $\delta\Delta(\text{ZPE})$ as defined here is positive. See Figure 9. For the condensed phase systems in which the only intermolecular force is London dispersion forces, the sum of the condensation shifts in the vibrational eigenvalues, $\Delta\lambda_{iA}$, of a representative molecule A is

$$\sum_i \Delta\lambda_{iA} = -2 \frac{\alpha_B}{S_{AB}^6} \sum_i \left(\frac{\partial \vec{P}}{\partial Q_{iA}} \right)^2 \quad (123)$$

where α_B is the electronic polarizability of a neighbor molecule B in its ground state, S_{AB} is the equilibrium average intermolecular separation in the condensed phase, and $(\partial \vec{P} / \partial Q_{iA})$ is the first derivative of the dipole moment vector $\vec{P}$ with respect to the i -th normal coordinate Q_i of the representative molecule.⁽¹²⁴⁾

Figure 9. Isotope effect on ZPE shift upon condensation.⁽¹³⁾

A combination of Eqs. (120) and (123) with a $\delta(\text{ZPE})$ approximation such as Eqs. (111), (112) and (115) provides a direct means of evaluating $\delta\Delta(\text{ZPE})$ using properties of the gaseous molecule. At the time of our study of this problem, the only available relationship between $\Sigma\omega_i$ and $\Sigma\lambda_i$ was that of Bigeleisen and Goldstein.⁽¹²⁵⁾ The first term of this infinite series is

$$\sum_i \delta\omega_i \equiv \frac{1}{8\pi^2 c^2} \frac{1}{\omega^2} \sum_i \delta\lambda_i \quad (124)$$

so that the vapor pressure isotope effect parameter, $B = \delta\Delta(\text{ZPE})$, becomes

$$B \equiv \frac{1}{8\pi^2} \frac{h}{c k} \left(\frac{C}{\omega_0} \right) \left(\frac{\alpha}{S^6} \right) \sum_i \left[\left(\frac{\partial \vec{P}}{\partial Q_i} \right)^2 - \left(\frac{\partial \vec{P}}{\partial Q_i} \right)^2 \right]_{\text{gas}} \quad (125)$$

where C is a condensed-phase structural constant which accounts for an overall influence of all molecules in the first and outer coordination spheres and is well known for various crystal lattice structures.

The magnitude of $(\partial P/\partial Q)^2$ can be evaluated by transforming the polar tensor,

$$\vec{P}_{x(\alpha)} = \begin{pmatrix} \partial P_x / \partial X_\alpha & \partial P_x / \partial Y_\alpha & \partial P_x / \partial Z_\alpha \\ \partial P_y / \partial X_\alpha & \partial P_y / \partial Y_\alpha & \partial P_y / \partial Z_\alpha \\ \partial P_z / \partial X_\alpha & \partial P_z / \partial Y_\alpha & \partial P_z / \partial Z_\alpha \end{pmatrix}, \quad (126)$$

into the center-of-mass/ principal axis coordinate system. The partial derivatives in Eq. (126) are the cartesian components of the derivatives for atom α . Then, it can be shown⁽¹³⁾ that

$$\left(\frac{\partial \vec{P}}{\partial Q_i} \right)^2 = \sum_{\alpha}^N \mu_{\alpha} \xi_{\alpha}^2 - \Omega - 2D, \quad (127)$$

where μ_{α} is the reciprocal mass of atom α , and ξ_{α} is the effective atomic charge on atom α as defined by Person and Newton,⁽¹²⁷⁾

$$\xi_{\alpha}^2 = \text{Tr} [\vec{P}_{x(\alpha)} \vec{P}_{x(\alpha)}'] \quad (128)$$

where $\vec{P}_{x(\alpha)}'$ is the transpose matrix. The term Ω and D are correction terms which, respectively, represent the effects of molecular rotations and external-internal kinetic energy interactions. The terms of $\vec{P}_{x(\alpha)}$ can be evaluated from a quantum mechanical algorithm.

By means of Eqs. (124)-(138), $\delta\Delta(\text{ZPE})$ can be expressed as a product of an isotope-independent condensed-phase factor and an isotopic difference in the gas-phase properties.⁽¹³⁾ The theory was tested⁽¹³⁾ by calculating the effective atomic charges for carbon and hydrogen in ethylene using the CNDO/2 molecular orbital program. The theory was applied to the D/H- and $^{13}\text{C}/^{12}\text{C}$ - isotope effects in methane and fluoromethanes, and the results were compared with experimental values of the total IR absorption intensities and $\delta\Delta(\text{ZPE})$. This led to a set of rules of thumb for predicting a large value of $\delta\Delta(\text{ZPE})$.⁽¹³⁾

The following properties and features of a molecule generally increase $\delta\Delta(\text{ZPE})$.

- Highly polar bonds, because they increase ξ_{α}^2 . The effect is more marked for a central atom surrounded by polar bonds than for a terminal atom bonded to such a central atom. Thus, for a $^{13}\text{C} / ^{12}\text{C}$ substitution, the B factor for CF_4 should be greater than that for CH_4 .
- Larger difference in the reciprocal masses of the substituted isotopic atoms.
- Molecular properties that make $\delta\Omega$ smaller, which include smaller electric dipole moments of the molecule, larger principal moments of inertia, and central isotope substitution rather than peripheral isotope substitution.

- d) Absence of hydrogen atoms, which, for instance, makes the Bigeleisen-Goldstein characteristic frequency ω_0 smaller.
- e) More closely packed liquid structure, or smaller liquid molal volume, which tends to increase the effective value of the Lennard-Jones and Ingham crystal constant and to decrease the nearest-neighbor distances of liquid molecules.

As presented in Section IV below, the results of our studies of VPIE of fluoromethanes qualitatively agree with these general rules.

IV. Vapor Pressure Isotope Effects

The *vapor pressure isotope effect* (VPIE) is a useful tool for elucidating molecular forces in condensed phases. We studied the VPIE in the following specific areas:

- a) VPIE of fluoromethanes, $\text{CH}_{4-n}\text{F}_n$ ($n = 1 \sim 3$): References (13, 16, 17, 18, 20, 21, 28)
- b) VPIE of ammonia: References (31, 30)
- c) Theories related to VPIE: References (7, 8, 9, 13)

IV-A. Vapor Pressure Isotope Effects in Fluoromethanes

Vapor pressure isotope effects were measured using the precision cryostat⁽¹⁷⁰⁾, which provides a computer-control for stability and uniformity ($< \pm 0.001\text{K}$) of cryogenic temperatures ($3 \sim 300\text{K}$) and for ease of operation. Table 11 lists the isotopic molecules and the temperature ranges that we studied..

Table 11. Fluoromethanes Studied

<u>Molecules</u>	<u>Isotopes</u>	<u>Temperature range</u>
Methyl fluorides (liq)	$^{12}\text{CH}_3\text{F}$, $^{12}\text{CD}_3\text{F}$, $^{13}\text{CH}_3\text{F}$	132.48 ~ 213.12 K
Methylene difluorides (liq)	$^{12}\text{CH}_2\text{F}_2$, $^{12}\text{CD}_2\text{F}_2$, $^{13}\text{CH}_2\text{F}_2$	149.36 ~ 244.82 K
Fluoroforms (liquid)	$^{12}\text{CHF}_3$, $^{12}\text{CDF}_3$, $^{13}\text{CHF}_3$	125.95 ~ 212.16 K

For each isotopic pair, the data on the vapor pressure P' of $^{12}\text{CH}_{4-n}\text{F}_n$ taken as the reference, and δP , the isotopic difference defined as $P' - P$, were converted to the ratio of the RPFR of the condensed phase, f_c , to that of the gas phase, f_g , by a conversion;

$$\ln \frac{f_c}{f_g} = \left[1 + P' \left(B_0 - \frac{V_c}{RT} \right) \right] \ln \frac{P'}{P} - \ln f_{\text{ref}}^g, \quad (129)$$

where B_0 and V_c , are, respectively, the second virial coefficient and the molal volume of the condensed phase, and f_{rot}^s is non-classical rotational correction.⁽¹³⁸⁾ Figures 10 ~ 15 are plots of $T \ln(f_c/f_p)$ against T^{-1} . The differences between $\ln(f_c/f_p)$ and $\ln(P'/P)$ are mostly within 1 to 2% of each other.

Figure 10. $^{13}\text{C}/^{12}\text{C}$ VPIE in methyl fluoride⁽²¹⁾: The solid line was computed from F-matrices of Tables 12 and 13.

Figure 11. D/H VPIE in methyl fluoride⁽²¹⁾: The solid line computed from F-matrices of Tables 12 & 13.

Figure 12. $^{13}\text{C}/^{12}\text{C}$ VPIE in methylene difluoride⁽²²⁾: The solid line computed from F-matrices of Tables 14 and 15.

Figure 13. D/H VPIE in methylene difluoride⁽²²⁾: The solid line was computed from F-matrices of Tables 14 and 15.

Figure 16 summarizes the plots of Figures 10 ~ 15 and compare them with the similar plots made from the data of VPIE's of methane.^(136, 137) The following general observations are made.

- 1) All hydrogen isotope effects in the fluoromethanes are normal ($P' > P$), while the D/H effects in methane are inverse ($P' < P$).
- 2) All carbon isotope effects in the fluoromethanes are inverse, while the carbon isotope effects in methane is normal.

Figure 14. $^{13}\text{C}/^{12}\text{C}$ VPiE in fluoroform.⁽⁷⁾ The solid line was computed from F-matrix of Tables 16 & 17.

Figure 15. D/H VPiE in fluoroform.⁽⁷⁾ The solid line was computed from F-matrix of Tables 16 and 17.

Figure 16. Comparison of VPiEs in liquid fluoromethanes and liquid methane⁽²⁸⁾. Solid lines are $^{13}\text{C}/^{12}\text{C}$ effects; dashed lines are the effects of perdeuterio substitutions.

- 3) The carbon isotope effects in CH_2F_2 are roughly of the same order of magnitude as the average of the carbon isotope effects in CH_3F and CHF_3 .
- 4) The D/H effects in CH_2F_2 is unusually large compared to the D/H effects in the other fluoromethanes:

$$\frac{1}{2} [T \ln (P'/P)_{\text{CH}_2\text{F}_2/\text{CD}_2\text{F}_2}] \gg \frac{2}{3} [T \ln (P'/P)_{\text{CH}_3\text{F}/\text{CD}_3\text{F}}] \quad (130)$$

and

$$\frac{1}{2} [T \ln (P'/P)_{\text{CH}_2\text{F}_2/\text{CD}_3\text{F}_2}] \gg \frac{2}{1} [T \ln (P'/P)_{\text{CHF}_3/\text{CDF}_3}] \quad (131)$$

- 5) The curvature of every D/H-VPiE plot for the fluoromethanes is negative, while that of every $^{13}\text{C}/^{12}\text{C}$ -VPiE in the fluoromethanes is positive.

If each condensed phase molecule could be regarded as a point mass, then ΔH_{vap} for the lighter isotopic molecule would be smaller than ΔH_{vap} of the heavier molecule, and one would have expected $P' > P$. This is the normal VPIE. The external motion tends to lead to a normal VPIE. The inverse VPIE ($P' < P$) is due to the internal molecular structure and its vibrational motions that interact with the intermolecular forces in the condensed phase.

If all molecular forces were independent of temperature, then we would have the following as the simplest correlation between the RPFRs and the molecular parameters:

$$\ln(f_c/f_g) = \sum_i^{\text{external}} \delta \ln b(u_i) - \sum_i^{\text{internal}} \Delta \delta \ln b(u_i) \equiv \frac{A}{T^2} - \frac{B}{T} \quad (132)$$

where $b(u)$ is defined in Eq. (18a), $\delta \ln b(u) = \ln b(u') - \ln b(u)$, $\Delta \delta \ln b(u) = \delta \Delta \ln b(u)$ as in Eqs. (120) ~ (122), and

$$A = \frac{1}{24} \left(\frac{\hbar c}{kT} \right)^2 \sum_i^{\text{External}} (v_i'^2 - v_i^2)_{\text{condensed phase}} \quad (133)$$

$$B = \frac{1}{2} \left(\frac{\hbar c}{kT} \right) \sum_i^{\text{Internal}} [\delta v_{i, \text{gas}} - \delta v_{i, \text{condensed}}] \quad (134)$$

The term of A/T^2 is the contribution of the external oscillatory motions of the whole molecule, which could be taken either as in the liquid simple cell model or as a central molecule in the liquid cluster model. The liquid cluster model will be presented in Section IV-C of this report.

As the temperature decreases, going from left toward right in Figures 10 ~ 16, the liquids contract, the external forces increase, the magnitudes of the translational and rotational frequencies generally increase, and thus the factor A of Eqs. (132) and (133) would increase. The curvature of a plot of $T \ln(f_c/f_g)$ against $1/T$ would then be positive.

The negative curvatures of the D/H-plots of Figures 11, 13, and 15 (Observation 5) thus imply dominant influence of the temperature on the B-term in Eq. (132): The B-factor should algebraically increase with the decreasing temperature in order for the curvature to be negative.

To quantitatively understand these phenomena, we conducted a series of normal coordinate analyses for the gaseous and liquid fluoromethanes.^(17, 21, 28) The best sets of F-matrices thus obtained are summarized in Tables 12 ~ 17. The solid lines drawn through experimental points in Figures 10 ~ 15 were calculated using these F-matrices. The T-dependence of the interaction force constants between the external and internal motions and the diagonal external elements for some of the external coordinates are necessary in order to reproduce the observed VPIE for both hydrogen and carbon substitutions, in particular, for good reproduction of;

- a) the steep slopes and the negative curvature of the D/H-VPIE plots and the very small slope of the carbon isotope plots, and
- b) the large normal VPIEs for the D/H substitutions and the small inverse VPIEs of the $^{13}\text{C}/^{12}\text{C}$ substitutions.

Table 12. F-matrix of Gaseous Methyl Fluoride ^(a)

diagonals			off-diagonals	
description	notation	value, mdyn Å ⁻¹	notation	value, mdyn Å ⁻¹
C-H stretch	f_d	4.686	f_{dd}	-0.036
C-F stretch	f_D	5.662	f_{dD}	0.602
HCH bend	f_α	0.408	$f'_{d\alpha}$	-0.125
HCF bend	f_β	0.617	$f_{d\alpha}$	-0.225
			$f_{D\alpha}$	-0.258
			$f_{\alpha\alpha}$	-0.054
			$f'_{d\beta}$	0.118
			$f_{d\beta}$	0.015
			$f_{D\beta}$	0.258
			$f'_{\alpha\beta}$	-0.143
			$f_{\alpha\beta}$	-0.119
			$f_{\beta\beta}$	-0.144

* All bending coordinates are weighted by the equilibrium C-H bond length. An F-matrix element with one subscript is a diagonal element. When two off-diagonal elements are listed to a similar interaction, e.g., $f'_{d\alpha}$ and $f_{d\alpha}$, the one without a prime refers to the interaction between two coordinates which share a common C-H bond, while the one with a prime refers to the interaction between two coordinates which do not share a common C-H bond.

Table 13. F-matrix of Liquid Methyl Fluoride ^(a)

element	value, ^a mdyn Å ⁻¹ or mdyn Å
f_{T_x}, f_{T_y}	0.120
f_{T_z}	0.300
f_{R_x}, f_{R_y}	$8.1527 \times 10^{-3} - (6.8040 \times 10^{-4})\Delta + (3.6677 \times 10^{-6})\Delta^2$
f_{R_z}	$f_{R_x}/8$
f_{DT_x}	$9.4169 \times 10^{-3} + (1.0788 \times 10^{-3})\Delta + (1.6879 \times 10^{-5})\Delta^2$
f_{DT_z}	$1.2045 \times 10^{-1} + (7.1425 \times 10^{-4})\Delta + (1.4527 \times 10^{-6})\Delta^2$
$f_{\alpha T_z}$	$-f_{DT_z}$

* All elements are in units of mdyn Å⁻¹ except f_R 's, which are in units of mdyn Å. $\Delta \equiv T - 153$.

Table 14. F-matrix of Gaseous Methylene Difluoride ^(a)

diagonals			off-diagonals	
description	notation	value, mdyn Å ⁻¹	notation	value, mdyn Å ⁻¹
C-H stretch	f_d	4.829	f_{dd}	C-H, C-H 0.130
C-F stretch	f_D	6.020	f_{dD}	C-H, C-F 0.106
HCH bend	f_α	0.462	$f_{d\alpha}$	C-H, HCH 0.105
FCF bend	f_β	1.000	$f_{d\beta}$	C-H, FCF -0.136
HCF bend	f_γ	0.635	$f_{d\gamma}$	C-H, HCF -0.063
			$f'_{d\alpha}$	C-H, HCF 0.073
			f_{DD}	C-F, C-F 0.996
			$f_{D\alpha}$	C-F, HCH -0.305
			$f_{D\beta}$	C-F, FCF 0.346
			$f_{D\gamma}$	C-F, HCH 0.249
			$f'_{D\alpha}$	C-F, HCF -0.253
			$f_{\alpha\beta}$	HCH, FCF -0.286
			$f_{\alpha\gamma}$	HCH, HCF -0.053
			$f_{\beta\gamma}$	FCF, HCF -0.160
			$f_{\alpha\alpha}$	HCF, HCF -0.048
			$f_{\alpha\beta}$	HCF, HCF -0.161
			$f_{\alpha\gamma}$	HCF, HCF -0.222

* All bending coordinates are weighted by the equilibrium C-H bond length. An F matrix element with one subscript is a diagonal element. When two off-diagonal elements are listed to a similar interaction, e.g., $f_{d\alpha}$ and $f'_{d\alpha}$, the one without a prime refers to the interaction between two coordinates which share a common C-H bond, while the one with a prime refers to the interaction between two coordinates which do not share a common bond. * For $f_{\alpha\alpha}$ and $f_{\alpha\beta}$ they share a C-H bond and a C-F bond, respectively.

Table 15. F-matrix of Liquid Methylene Difluoride ^(a)

diagonals		off-diagonals	
notation	value, ^a mdyn Å ⁻¹	notation	value, ^a mdyn Å ⁻¹
f_d	4.882	f_{DD}	0.131
f_D	5.959	f_{dD}	0.106
f_α	0.462	$f_{d\alpha}$	0.105
f_β	0.983	$f_{d\beta}$	-0.136
f_γ	0.630	$f_{d\gamma}$	-0.063
		$f'_{d\alpha}$	0.073
		f_{DD}	0.869
f_{T_x}	0.300	$f_{D\alpha}$	-0.314
f_{T_y}	0.200	$f_{D\beta}$	0.398
f_{T_z}	0.300	$f_{D\gamma}$	0.239
f_{R_x}	0.200	$f'_{D\alpha}$	-0.263
f_{R_y}	0.070	$f_{\alpha\beta}$	-0.283
f_{R_z}	0.300	$f_{\alpha\gamma}$	-0.053
		$f_{\beta\gamma}$	-0.157
		$f_{\alpha\alpha}$	-0.055
		$f_{\alpha\beta}$	-0.156
		$f_{\alpha\gamma}$	-0.217

$$f_{\alpha 11 R_x} = f_{\alpha 12 R_x} = -f_{\alpha 13 R_x} = -f_{\alpha 22 R_x}, \quad 3.059 \times 10^{-2} - (1.529 \times 10^{-4})\Delta + (1.130 \times 10^{-6})\Delta^2$$

$$f_{\alpha 11 R_y} = f_{\alpha 22 R_y} = -f_{\alpha 13 R_y} = -f_{\alpha 21 R_y}, \quad -4.920 \times 10^{-2} + (1.120 \times 10^{-3})\Delta$$

* All elements are in mdyn Å⁻¹, except f_R and $f_{\alpha R}$ which are in mdyn Å and mdyn, respectively. $\Delta = T - 190$.

Table 16. F-matrix of Gaseous Fluoroform⁽¹⁷⁾

Diagonals			Off-diagonals		
Description	Notation	Value (mdyn/Å)	Notation	Value (mdyn/Å)	
C-H stretch	f_D	5.0000	f_{Dd}	1.1720	
C-F stretch	f_d	6.0179	f_{dd}	0.7499	
HCF bend	f_α	0.4913	$f_{D\alpha}$	-0.1674	
FCF bend	f_β	0.7646	$f_{D\beta}$	0.1674	
			$f_{d\alpha}$	0.1943	
			$f_{d\alpha'}$	-0.2657	
			$f_{d\beta}$	0.2560	
			$f_{d\beta'}$	-0.1808	
			$f_{\alpha\alpha}$	0.0393	
			$f_{\alpha\beta}$	-0.0973	
			$f_{\alpha\beta'}$	-0.1517	
			$f_{\beta\beta'}$	-0.2667	

^aThe internal coordinate system consists of one CH bond stretch D , three CF bond stretches d , three HCF angle bends α , and three FCF angle bends β . (See Fig. 4.) All bending coordinates are weighted by the equilibrium C-F bond length of 1.332 Å. An F matrix element with one subscript is a diagonal element, and one with two subscripts is an off-diagonal element. When two off-diagonal elements are listed for a similar interaction, e.g., $f_{d\alpha}$ and $f_{d\alpha'}$, the one without a prime refers to the interaction between two coordinates which share a common C-F bond, while the one with a prime refers to the interaction between two coordinates which do not share a common C-F bond.

Table 17. F-matrix of Liquid Fluoroform⁽¹⁷⁾

Diagonals			Off-diagonals		
Description	Notation	Value (mdyn/Å) ^b	Notation	Value (mdyn/Å) ^b	
C-H stretch	f_D	5.0726	f_{Dd}	1.0400	
C-F stretch	f_d	5.9097	f_{dd}	0.5650	
HCF bend	f_α	0.4872	$f_{D\alpha}$	-0.1674	
FCF bend	f_β	0.7650	$f_{D\beta}$	0.1674	
			$f_{d\alpha}$	0.1943	
			$f_{d\alpha'}$	-0.2657	
			$f_{d\beta}$	0.0256	
			$f_{d\beta'}$	-0.1808	
			$f_{\alpha\alpha}$	0.0393	
			$f_{\alpha\beta}$	-0.0973	
			$f_{\alpha\beta'}$	-0.1517	
			$f_{\beta\beta'}$	-0.2667	
	f_{T_x}	0.20			
	f_{T_y}	0.20			
	f_{T_z}	0.40	f_{DT_z}	$-0.73 + 5.0 \times 10^{-3} \times (T - 165 \text{ K})$	
	f_{R_x}	0.20	$f_{\alpha T_x}$	0.13	
	f_{R_y}	0.20	$f_{\alpha T_y}$	0.13	
	f_{R_z}	0.05	$f_{\beta T_z}$	-0.13	

^aFor definitions of the internal coordinates and F-matrix element notations, see the footnote of Table IV.

^bAll F-matrix elements are in the units of mdyn/Å except f_{R_i} 's, which are in the units of mdyn Å.

From the first-order orthogonal approximation (cf. Section II of present report), we expect the effects of non-isotopic substituents on isotope effects to be additive to the extent that the force constants are transferrable among different chemical species. Then, $T \ln (P'/P)$ per D-for-H substitution would be approximately similar. This is roughly the case between the D/H-VPiE's of methyl fluoride and fluoroform, but it is not so with methylene difluoride (Observation 4). Table 18 provides a quantitative explanation. The methane data were taken from References (136 - 138). From Table 18,⁽²⁸⁾ the following points are noted.

- The magnitudes of the shift in $\delta(\text{ZPE})$ upon condensation, $\Sigma \delta \Delta v_i$, for the hydrogen isotope effects is significantly greater than that for the carbon isotope substitutions, as expected.
- The shifts, $\Sigma \delta \Delta v_i$ (D/H) for the fluoromethanes are negative, while that in methane is positive. The negative shifts, which is an anomaly rather than a norm, are due mainly to the large blue shifts in the C-H stretching frequencies upon condensation, i.e., v_1 in CHF_3 , v_1 and v_6 in CH_2F_2 , and v_1 and v_4 in CH_3F . The blue-shifts in fluoroform and methyl fluoride have been qualitatively explained on a basis of hydrogen bonding in molecular orbital studies.^(16, 20) This explains why the D/H-effects in the fluoromethanes are normal and large while the D/H-effects in methane is inverse (Observation 1).

Table 18. Comparison of Frequency Shifts in Fluoromethanes ^(a)

fluoroform ^b at 165 K	¹² CHF ₃		¹³ CDF ₃		¹³ CHF ₃	
	ν_i' (gas)	$\Delta\nu_i'$	$\Delta\nu_i$	$\delta\Delta\nu_i$	$\Delta\nu_i$	$\delta\Delta\nu_i$
$\nu_1(A_1)$	3036.2	-26.3	-18.3	-8.0	-26.2	-0.1
$\nu_2(A_1)$	1138.9	20.4	22.6	-2.2	18.8	1.6
$\nu_3(A_1)$	701.4	4.6	3.0	1.6	4.3	0.3
$\nu_4(E)$	1381.0	5.4	-6.1	11.5	5.8	-0.4
$\nu_5(E)$	1151.8	-7.9	1.7	-9.6	-8.1	0.2
$\nu_6(E)$	507.7	-0.2	0.0	-0.2	-0.2	0.0
totals		-6.7	-1.5	-5.2	-8.1	1.4

methylene difluoride ^c at 190 K	¹² CH ₂ F ₂		¹³ CD ₂ F ₂		¹³ CH ₂ F ₂	
	ν_i' (gas)	$\Delta\nu_i'$	$\Delta\nu_i$	$\delta\Delta\nu_i$	$\Delta\nu_i$	$\delta\Delta\nu_i$
$\nu_1(A_1)$	2947.0	-16.0	-10.7	-5.3	-16.1	0.1
$\nu_2(A_1)$	1507.1	-0.1	15.3	-15.4	-1.3	1.2
$\nu_3(A_1)$	1107.3	28.4	11.5	16.9	27.8	0.6
$\nu_4(A_1)$	532.3	1.1	1.3	-0.2	1.2	-0.1
$\nu_5(A_2)$	1260.6	-2.2	1.1	-3.3	-2.6	0.4
$\nu_6(B_1)$	3014.2	-15.8	-10.2	-5.6	-15.9	0.1
$\nu_7(B_1)$	1183.3	13.0	15.7	-2.7	12.3	0.7
$\nu_8(B_2)$	1436.9	-0.4	-5.5	5.1	-0.2	-0.2
$\nu_9(B_2)$	1081.2	-7.9	-2.8	-5.1	-7.8	-0.1
totals		0.1	15.7	-15.6	-2.6	2.7

methyl ^d fluoride at 153 K	¹² CH ₃ F		¹³ CD ₃ F		¹³ CH ₃ F	
	ν_i' (gas)	$\Delta\nu_i'$	$\Delta\nu_i$	$\delta\Delta\nu_i$	$\Delta\nu_i$	$\delta\Delta\nu_i$
$\nu_1(A_1)$	2909.8	-5.3	3.5	-8.8	-6.0	0.7
$\nu_2(A_1)$	1462.6	5.6	18.8	-13.2	6.1	-0.5
$\nu_3(A_1)$	1056.8	61.7	28.9	32.8	61.5	0.2
$\nu_4(E)$	3006.7	-23.6	-12.8	-10.8	-24.0	0.4
$\nu_5(E)$	1479.3	-1.0	-3.1	2.1	-0.7	-0.3
$\nu_6(E)$	1183.0	1.3	0.9	0.4	1.3	0.0
totals		15.4	21.2	-5.8	14.8	0.6

methane ^e	¹² CH ₄		¹³ CD ₄		¹³ CH ₄	
	ν_i' (gas)	$\Delta\nu_i'$	$\Delta\nu_i$	$\delta\Delta\nu_i$	$\Delta\nu_i$	$\delta\Delta\nu_i$
$\nu_1(A_1)$	3143.7	11.5	8.2	3.3	11.5	0.0
$\nu_2(E)$	1574.2	3.9	2.8	1.1	3.9	0.0
$\nu_3(F_2)$	3154.1	13.8	11.5	2.3	13.6	0.2
$\nu_4(F_2)$	1357.4	4.9	3.0	1.9	4.9	0.0
totals		75.4	57.3	18.1	74.8	0.6

^a All values are calculated and in the unit of cm⁻¹. $\Delta\nu_i = \nu_i(\text{gas}) - \nu_i(\text{liq})$. $\delta\Delta\nu_i = \Delta\nu_i' - \Delta\nu_i = \delta\nu_i(\text{gas}) - \delta\nu_i(\text{liq})$ where $\delta\nu_i = \nu_i' - \nu_i$.

The differences in the D/H- and ¹³C/¹²C-effects between fluoromethanes and methane are due to the differences in the molecular symmetry and the relative magnitudes of intermolecular forces. The carbon atom in methane does not move at all in all but one of the F₂ mode. This results in the insignificant contribution to the zero-point energy term. The dominant term for the carbon-VPIE in methane is the one due to the translational lattice modes. In contrast, the peripheral D/H substitutions in methane lead to the large, positive $\delta\Delta(\text{ZPE})$, which is responsible for the inverse D/H-VPIE in methane.

Fluoromethanes being of lower molecular symmetry, the carbon atom actively participates in all normal modes, and many interactions between the internal vibrations and the lattice modes are

symmetry-allowed. In addition, the intermolecular forces are greater in the liquid fluoromethanes than in methane, as Table 19 implies. This and the blue shifts are responsible for the large, normal D/H-VPIE's

Table 19. Some Properties of Liquid Fluoromethanes ⁽²⁰⁾

	normal bp, °C	ΔH vap at nbp, kJ/mol	dipole moment (gas), D	polarizability (gas), Å ³
CH ₄	-161.5	8.4	0	2.60
CH ₃ F	-78.4	17.2	1.85	2.67
CH ₂ F ₂	-51.6	19.2	1.97	2.74
CHF ₃	-82.0	16.7	1.65	2.81
CF ₄	-128.0	12.1	0	2.89

in the fluoromethane. For the carbon VPIE, the inverse effect are due to the red-shifts and the fact that the isotopic substitutions are made near the centers of mass and near the principal axes.

An examination of the Figures and Tables of the present section reveals that the rules of thumb discussed in Section IIIB in terms of the theory of $\Delta\delta(\text{ZPE})$ are in agreement with the experimental results.

The unusually large blue-shift in the D/H-effect of methylene difluoride (Table 18) is responsible for the disparity in the VPIE per D/H substitution (Observation 4). The large shift is primarily due to two reasons. First, CH₂F₂ being of the lowest molecular symmetry, the molecules interact more extensively in the liquid. Secondly, since CH₂F₂ has two hydrogen atoms and two hydrogen acceptor atoms, the highest possible average number of hydrogen bonds per CH₂F₂ molecule is two. In contrast, the highest possible average number of hydrogen bonds per CHF₃ or CH₃F molecule is one. In CHF₃ and CH₃F there are inbalance between the donor and acceptor atoms. These observations and Table 19 are also consistent with the fact that the thermal expansion coefficients of liquid methylene difluoride is the smallest of the three fluoromethanes: at 173K, for instance, $\beta = 2.2$ for CH₃F, 1.8 for CH₂F₂, and 2.4 for CHF₃.

All observed D/H- and ¹³C /¹²C-VPIE's can be explained in terms of molecular symmetry and intermolecular interactions in the liquid.

IV-B. Vapor Pressure Isotope Effects in Liquid and Solid Ammonia

We studied hydrogen- and nitrogen-VPIEs in liquid and solid ammonia.^(30,31) Both VPIEs are normal at temperatures between 163K and 243K. There have been earlier investigations of the VPIEs of ammonia.⁽¹³⁵⁾ Figures 17 and 18 compare our results⁽³¹⁾ with the earlier data. The solid lines drawn through our data (the dots) are the least-squared fit lines in the form,

$$T \ln \frac{p'}{p} = \frac{A}{T} - B \quad . \quad (135)$$

Figure 17. Observed D/H vapor pressure isotope effects in ammonia. Dots; present work, Circles (139); Crosses (140).

Figure 18. Observed ¹⁵N/¹⁴N vapor pressure isotope effects in ammonia. Dots; present work. Squares (141).

The parameters A and B are tabulated in Table 20 and may be physically interpreted by in Eqs. (133) and (134). The vapor pressures of the individual isotopic ammonia can be represented by

$$\text{and} \quad \ln P(\text{torr}) = a - \frac{b}{T} - cT \quad [\text{liquid}] \quad (136)$$

$$\ln P(\text{torr}) = a - \frac{b}{T} \quad [\text{solid}] \quad (137)$$

where T is in K. The least-squares fit parameters are summarized in Table 21.

Table 20. Least-squares fits of isotopic vapor pressure ratios to Eq. (135).⁽¹¹⁾

Isotopic Pair	Temperature range (K)	A	B	σ
liq d_0 /liq d_3	$T_u < T$	15041.2 ± 42.9	33.1387 ± 0.1976	0.1025
liq d_0 /sol d_3	$T_u' < T < T_u$	161084 ± 5414	768.112 ± 27.426	0.778
sol d_0 /sol d_3	$T < T_u'$	16768.7 ± 152.5	29.9744 ± 0.8487	0.4715
liq 14/liq 15	$T_u < T$	364.550 ± 3.105	0.83480 ± 0.01438	0.00795
sol 14/sol 15	$T < T_u'$	1477.21 ± 20.90	6.40848 ± 0.11658	0.06242

T_u' and T_u are the triple points (cf: Table 22) of the first- and second-named isotopic species, respectively. σ is the root mean squares deviation.

Species	Phase	σ (log P)	a	b	c	T(K)	Number of pts
$^{14}\text{NH}_3$	Liq	0.00091	9.94326	1471.633	0.0038492	196–243	94
		[0.000167	10.07133	1485.600	0.0041449	199–242	11]
		(0.00123	9.95028	1473.17	0.0038603	196–250	)
	Sol	0.00251	10.01883	1633.785		163–195	109
		[0.000669	10.00528	1630.993		177–195	7]
		(0.00326	9.98379	1627.22		173–195	)
$^{14}\text{ND}_3$	Liq	0.00028	10.30672	1541.893	0.0043762	199–243	67
	Sol	0.00289	10.23684	1700.623		163–198	135
$^{15}\text{NH}_3$	Liq	0.00060	9.94265	1472.232	0.0038514	196–243	69
	Sol	0.00314	10.04193	1638.775		163–195	102

Table 21. Vapor pressure equations for individual isotopic ammonias:⁽¹¹⁾
Eq. (136) for the solid,
Eq. (137) for the liquid.

The triple points obtained by this work are compared with previous results in Table 22. For comparison,

the triple point for NH_3 obtained by Overstreet and Giaque⁽¹⁴³⁾ is 195.41K (45.58 torr) after a 0.05K temperature scale correction. Also, Armstrong's triple point temperatures were based on the ice point of 273.16K.

Table 22. Triple points of isotopic ammonia⁽¹¹⁾

Isotopic species	T_u (K)		P_u (torr)	
	Armstrong	This work	Armstrong	This work
$^{14}\text{NH}_3$	195.45	195.41	45.58	45.49
$^{14}\text{ND}_3$	199	198.96	48.22	48.35
$^{15}\text{NH}_3$	—	195.58	—	48.83

It is notable that the difference in the vapor pressures between NH_3 and ND_3 does not change much at the temperatures between the triple points of NH_3 and ND_3 , while the magnitudes of the vapor pressures change considerably (P' from about 45.5 torr to about 60.4 torr) in the same range. See Figure 19.⁽³¹⁾ The difference in the pressures in this range can be represented by

$$\delta P \text{ (torr)} = P' - P = -(0.0251 \pm 0.0323)T(\text{K}) + (16.388 \pm 6.381) \quad (138)$$

From Clausius-Clapeyron equation, this means $P'\Delta H'_{\text{vap}} \equiv P\Delta H_{\text{vap}}$ and $P'/P \equiv \Delta S_{\text{sub}}/\Delta S'_{\text{vap}} > 1$. From the data, we have $\Delta H(\text{sublimation})/\Delta H(\text{vaporization}) = 1.286 \pm 0.012$, while the average of P'/P is 1.275. This phenomenon is still being investigated.

The experimental data of P'/P was reduced to f_i/f_e using the same method that has been described in this report in relation to the VPIE of fluoromethanes. A normal coordinate analysis of the condensed phase ammonia molecules resulted in the F-matrices which are summarized in Table 23 for the solid and

Table 24 for the liquid. For the gas phase, we used Špirko's anharmonic function,⁽¹⁴⁴⁾ which well reproduces all known frequencies of the gaseous isotopic ammonia including the doublets resulting from the umbrella-type inversion. The other F-matrices reproduce all known internal and lattice frequencies of solid ammonia (NH_3 and ND_3). There are no experimental lattice frequency data for

$^{15}\text{NH}_3$. Relative contributions of the

internal and external motions to the VPIE vary with the temperature. Table 25 illustrates the magnitudes of these contributions at typical temperatures.

We derived the following conclusions.

- 1) Unlike the VPIE in liquid fluoromethanes, the slope of the plots of Figures 17 and 18 are attributable primarily to the translational and rotational motions of the ammonia molecules. The contributions from the internal vibrational motions are very significant but not as temperature-dependent as those in the liquid fluoromethanes.
- 2) The rotational motions of ammonia molecules in the liquid are strongly hindered. The degree of the hindrance against rotation in the liquid is about the same as in the solid.
- 3) The rotation of ammonia in the liquid is strongly hindered mainly because of the external forces acting against the motions of hydrogen atom effecting the molecular rotation and not as much because of the forces acting against nitrogen motions.
- 4) There are significant differences between the liquid and the solid in the forces that resist movements of the nitrogen atom in the directions perpendicular to the molecular figure axis.
- 5) There are strong interaction between the internal and external modes, especially in the solid.

These conclusions are in qualitative agreement with the following experimental and theoretical results in the fields other than isotope effects.

- a) The x-ray studies on the liquid over a wide range of temperatures (199 ~ 277K) shows that the close-packed structure (12 nearest neighbors) of the solid persists, more or less, not only through the phase change but also over a range of liquid temperature.

Figure 19. $\Delta P = P(\text{NH}_3) - P(\text{ND}_3)$ vs. T near triple points.

Table 23. F-matrix of solid ammonia in valence coordinates*

Description	Notation	Value (mdyn Å ⁻¹ or mdyn Å)
N'-H stretch	f_D	5.835
N-H stretch	f_A	5.834
HN'H bend	f_2	0.569
HN'H bend	f_3	0.569
Str'-Str' interact	f_{DD}	-0.041
Str'-Str'	f_{AA}	-0.041
Bend'-Bend'	f_{aa}	-0.0040
Bend'-Bend'	f_{ff}	-0.0041
Str'-Bend'	f_{Da}	0.046
Str'-Bend'	f_{Df}	-0.0426
Str'-Bend'	f_{Af}	0.046
Str'-Bend'	f_{Df}	-0.0426
Translation	$f_{T_i}^c$	$0.158 - (1.8061 \times 10^{-3}) \Delta$ $+ (8.7505 \times 10^{-6}) \Delta^2$
	$f_{T_{xx}}^c$	$0.1154 - (7.4132 \times 10^{-4}) \Delta$
	$f_{T_{yy}}^c$	$0.3360 - (3.4349 \times 10^{-3}) \Delta$ $+ (1.5771 \times 10^{-6}) \Delta^2$
	$f_{T_{zz}}^c$	$0.1012 - (5.2011 \times 10^{-4}) \Delta$
Rotation	$f_{R_i}^c$	$0.1558 - (5.1828 \times 10^{-4}) \Delta$
	$f_{R_{xx}}^c$	8.5272×10^{-2} $- (2.5960 \times 10^{-4}) \Delta$
	$f_{R_{yy}}^c$	1.0989×10^{-1} $- (3.3455 \times 10^{-4}) \Delta$
	$f_{R_{zz}}^c$	$0.2054 - (6.6728 \times 10^{-4}) \Delta$
Ext-Ext	$f_{T_i-T_j}^c$	9.9600×10^{-2} $- (4.4630 \times 10^{-4}) \Delta$ $+ (7.4074 \times 10^{-6}) \Delta^2$
	$f_{R_i-R_j}^c$	6.5534×10^{-2} $- (2.5868 \times 10^{-4}) \Delta$
Int-Ext	$f_{a-T_i}^c$	-1.000×10^{-3} $- (4.1183 \times 10^{-3}) \Delta$ $- (8.4982 \times 10^{-5}) \Delta^2$
	$f_{f-T_i}^c$	1.000×10^{-3} $+ (4.0429 \times 10^{-3}) \Delta$ $- (2.5714 \times 10^{-4}) \Delta^2$
	$f_{D-T_i}^c$	$(5.8181 \times 10^{-3}) \Delta$ $+ (2.6252 \times 10^{-4}) \Delta^2$
	$f_{A-T_i}^c$	$(-1.0315 \times 10^{-3}) \Delta$ $- (6.8580 \times 10^{-4}) \Delta^2$

* All elements are in units of mdyn Å⁻¹, except f_D 's and the interactions among f_A 's, which are in units of mdyn Å. $\Delta = T(K) - 180$. The superscripts, c and f, denote the corner and face molecules, respectively. The primed f (e.g., f_D) refers to an interaction between an NH stretch and an HNH bend in which the nitrogen atom is the only atom shared by the two coordinates, while the unprimed f (e.g., f_D) refers to the one between the two coordinates in which an N-H bond is shared. For the external modes, the molecular symmetry axis is the z-axis. The notation such as f_{T_x} means that the element applies to the translation in the molecular xy-plane, while $f_{T_x-T_y}$ refers to an interaction between the coordinates T_x and T_y .

Table 24. F-matrix of liquid ammonia in valence coordinates*

Description	Notation	Value (mdyn Å ⁻¹ or mdyn Å)
N-H stretch	f_D	6.015
HNH bend	f_2	0.529
	f_3	-0.003
	f_{Da}	-0.049
	f_{Df}	-0.0416
	f_{Af}	0.044
Translation	f_{T_i}	$f_{T_{xx}} = 0.1421$ $- 3.4177 \times 10^{-4} \Delta$
Rotation	f_{R_i}	$f_{R_{xx}} = 0.1040 - 8.6319 \times 10^{-5} \Delta$ $+ 1.8428 \times 10^{-6} \Delta^2$
Internal-External Interaction	f_{DT_i}	+0.0268
	f_{AT_i}	+0.0042

All elements are in units of mdyn Å⁻¹ except f_A 's which are in units of mdyn Å. $\Delta = T - 217$.

Table 25. Comparison of various contributions to $T \ln (f_c/f_g)$ in isotopic ammonias. All values in K.

	$T \ln (f_c/f_g)$ at 195.41 K		$T \ln (f_c/f_g)$ at 198.98 K	
	d_0/d_3	14/15	d_0/d_3	14/15
Translations	3.0	6.07	2.6	1.68
Rotations	84.4	1.48	76.7	0.19
Internal vibrations	-31.9	-6.41	-37.1	-0.88
Non-classical rotations	1.5	0.0	1.5	0.0
Total	57.0	1.14	43.8	0.99

- b) A molecular dynamics study⁽³⁰⁾ based on a flexible, four-site model of liquid ammonia confirmed the coordination number of 12, with a slightly more pronounced first neighbor structure at lower temperatures. Other molecular dynamics and Monte Carlo studies also reported the coordination numbers ranging from 12 to 13.
- c) According to our molecular dynamics study,⁽³⁰⁾ there is little directional force on the lone-pair side of the nitrogen atom, while the hydrogens behave as if they are tied down by the nitrogen atoms of the neighbor molecules. There are approximately three such nearest neighbor molecules on the hydrogen side..

Thus, the directional preference of the external force on the hydrogen side of an ammonia molecule in the liquid seems to be responsible for the significant rotational contribution in the D/H-VPIE, while the practical absence of directionality of the force on the lone-pair side causes the small rotational contributions in the nitrogen VPIE.

IV-C. Medium Cluster Model for Vapor Pressure Isotope Effects

All experimental data on the VPIEs have been explained unexpectedly well by the theories based on the simple cell model of liquid. The model assumes that each liquid molecule is entrapped in an isotropic potential cage, and the surrounding molecules form a structureless environment around the molecule that represents the liquid phase. We developed a new model called *medium cluster model*^(7,8) for liquid, which explicitly takes the intermolecular interactions into account and investigated the reasons why the cell model, which is an obvious oversimplification of liquid structure, apparently works well in explaining the VPIE phenomena.

According to the medium cluster model, a liquid phase is regarded as consisting of a cluster of somewhat regularly arranged molecules. A central molecule is surrounded by a variable number of molecules in the first coordination shell. The method of normal coordinate analysis is used to determine the modes of vibration of the condensed phase cluster taken as a unit, and the RPFR is calculated by using an appropriate number of normal frequencies of the cluster that are attributable to the motions of the central molecule. Clusters of 3, 7, 9 and 13 methane molecules in three basic relative orientations of the constituent molecules were examined, and their carbon and hydrogen isotope effects were studied.

Consider a liquid phase consisting of N molecules to be an assembly of independent clusters with m molecules each. If the clusters are indistinguishable, the condensed phase partition function can be written as

$$Q_c = \frac{q_{cluster}^{N/m}}{(N/m)!} \quad , \quad (139)$$

where $q_{cluster}$ is the complete molecular partition function for the cluster regarded as a macromolecule. Each cluster consists of one molecule at the central position in the aggregate and $m - 1$ indistinguishable shell molecules in the first coordination sphere. Using $q_{central}$ and q_{shell} for the molecular partition functions of each type, the cluster partition function may be expressed as

$$q_{cluster} = \frac{q_{shell}^{m-1}}{(m-1)!} q_{central} \quad . \quad (140)$$

If, in addition, one assumes that the shell and central species are in equilibrium with each other, one has

$$\frac{q_{central}}{q_{shell}} = \frac{1}{m-1} \quad . \quad (141)$$

Substitution of Eqs. (140) and (141) into Eq. (139) and use of the Stirling approximation yield

$$Q_c = \left\{ \frac{e^{1/m} (m-1)^{(m-1)/m} q_{central}}{[(m-1)!] \left(\frac{N}{m}\right)!} \right\}^N \quad . \quad (142)$$

Then,

$$\ln \frac{s}{s'} f = \frac{1}{N} \ln \left(\frac{Q_c}{Q_c'} \right)_{q_m} - \ln \left(\frac{q_c}{q_c'} \right)_c \quad . \quad (143)$$

In accordance with the Born-Oppenheimer approximation, the cluster size m is isotope-independent and , thus,

$$\ln \frac{f_c}{f_g} = \ln \frac{f_c}{f_g} f_{\text{central}} - \ln \frac{f_c}{f_g} f_{\text{gas}} , \quad (144)$$

which appears to be identical in its form with the cell-model expression. The counting factors such as those appearing on the right hand side of Eq. (142) have cancelled out between the isotopic molecules. However, the evaluation of f_{central} in Eq. (144) explicitly takes into account the intermolecular interactions with its closest neighbors.

Since the shell molecules in a cluster are being used to simulate the entire bulk phase, it would be unrealistic to assign the same force constants to the central and shell molecules. The shell force constants must reflect the existence of additional hierarchical coordination spheres and the distribution of energy among central and shell molecules. We have correlated the potential energy surfaces of the central and shell molecules with each other by apportioning the shift in the total electronic kinetic energy upon condensation between the two molecular species of the model through an application of the localized virial theorem.^(146, 147) Thus, the F-matrices of the gas (F_g), central (F_c), and shell (F_s) molecules are related by

$$F_s = F_g + \frac{1}{m-1} (F_c - F_g) \quad . \quad [m > 1] \quad (145)$$

The model was applied to the VPIE of ^{13}C , ^{14}C , D, and T-substituted methanes using $m = 3, 7, 9$, and 13. The clusters, $m=7, 9$, and 13, respectively correspond to the simple cubic, body-centered cubic, and face-centered cubic lattice structures. Three equilibrium orientation of methane molecules were considered. They are illustrated in Figure 20. The results obtained can be summarized as follows.

- 1) Medium cluster model generally gives better agreement than the cell model with experimental values of the first quantum correction (i.e., A term of Eqs. (132) and (133)) and the ZPE shift term (i.g., B term of Eqs. (132) and (134)) without significantly increasing the number of force constant parameters beyond those needed for the simple cell model.
- 2) For each equilibrium orientation of Figure 20, both A- and B-terms exhibit a cyclic behavior as a function of conformational angles, as expected. For the spherical-top isotopic methanes, the cyclic variations are mainly and clearly due to interactions of molecular rotation with bond-angle bending and bond-stretching motions.
- 3) The effects of changing molecular orientations diminish with increasing cluster size (m), until the variations become negligibly small compared to experimental uncertainties. This is illustrated in Figure 21 for d_e / d_0 -effects in the gear geometry and in Figure 22 for $^{13}\text{C}/^{12}\text{C}$ -effects in the rocket geometry. In Figures 21 and 22, the intermolecular direction is the y-axis, and δ is rotational angle of the central molecule around the y-axis.

- 4) Thus, the formalism of simple cell model can be regarded to implicitly account for intermolecular interactions beyond the nearest neighbor molecules.

Figure 20. Relative orientation of adjacent methane molecules, A is the rocket geometry, B the antiparallel geometry, and C the gear geometry.⁽⁷⁾

Figure 21. VPIE of $\text{CD}_4 / \text{CH}_4$ for various cluster sizes in the gear geometry as a function of central molecular rotation.⁽⁷⁾

Figure 22. VPIE of $^{13}\text{CH}_4 / ^{12}\text{CH}_4$ for various cluster sizes in the rocket geometry as a function of central molecular rotation.⁽⁷⁾

V. Fractionation of Stable Isotopes

Studies of isotope fractionation will be presented in three sections:

- 1) Experimental and theoretical studies of ^{15}N fractionation.^(24, 148)
- 2) Basic studies for development of hydrophobic catalysts for the exchange reaction of hydrogen isotopes between dihydrogen and water.^(162, 168, 169, 201, 202, 203)
- 3) Theoretical studies of transient behaviors of a square cascade for isotope fractionation.⁽¹⁰⁾

V-A. Fractionation of Nitrogen Isotopes

The cryogenic distillation of nitric oxide and the Nitrox process have been the main processes for producing highly enriched nitrogen-15. The latter utilizes the exchange of nitrogen isotopes between nitric oxide gas and nitric acid, the main exchange reaction being

for which the unit stage separation factor α (cf: Eq. (4) on page 2) is 1.055 at 25°C and 1 atm pressure when 10M nitric acid is used. Typical values of α for the NO distillation range from 1.03 at 110.5K to 1.007 at 172.8K.⁽¹⁴⁹⁾

In the Nitrox process, the product-end refluxing is most commonly provided by a chemical reduction using SO_2 , the main reaction being

To produce ^{15}N of a given enrichment, the rate of product withdrawal is limited by the minimum reflux ratio, R_{min} , i.e., the ratio of the refluxed stream flow rate (i.e., nitric oxide here) to that of the product withdrawal rate. For instance, R_{min} for the Nitrox process yielding 99.9% ^{15}N from the natural abundance of 0.37% in one square-cascade is about 5,400. This means that, to produce one mole of ^{15}N , $5,400 \times (3/2) = 8,100$ moles of SO_2 will be needed. The cost of SO_2 is by far the largest single non-personnel expense for the operation of a Nitrox plant. If this cost could be reduced, the chemical exchange method for ^{15}N would become more attractive.

To explore this possibility, we studied ^(24, 148) a chemical exchange process based on an isotope exchange between a liquid phase, which is primarily a mixture of liquid N_2O_3 , and a gas phase, which is in phase and chemical equilibria with the liquid phase. Under the experimental conditions that we studied, the liquid is primarily N_2O_3 and the gas is primarily NO, so that this system will be referred to as the *NO/ N_2O_3 exchange system*.

The amount of SO_2 that is stoichiometrically required for the NO / N_2O_3 system is one-third of that needed for the nitrox system. Furthermore, as shown in this report, it turned out⁽²⁴⁾ that the HEPT (the height equivalent of theoretical plate) for the NO/ N_2O_3 packed-column system is almost an order of magnitude smaller than the HEPT for the NO/ HNO_3 operating under similar conditions.

V-A-1. Equilibrium Properties of NO/ N_2O_3 System

First, the phase and chemical equilibrium data in literature on the liquid and gaseous system involving NO, NO_2 , N_2O_3 , and N_2O_4 were systematized to obtain a self-consistent data basis.⁽¹⁴⁸⁾ The only assumption made was that the liquid phase only consists of N_2O_3 and N_2O_4 . This limits the present model to the liquid whose average oxidation number ranges from 3 and 4. The average atom fraction of +4 nitrogen in the liquid and gas phase are denoted x and y , respectively, the mole fractions of N_2O_3 and N_2O_4 in the liquid are x_3 and x_4 , respectively, and the mole fractions of NO, NO_2 , N_2O_3 , and N_2O_4 in the gas phase are y_1 , y_2 , y_3 , and y_4 , respectively. The calculational model successfully reproduced the following experimental data⁽¹⁴⁸⁾:

a) Total pressure, P , over the liquid mixture as a function of x and T ,

b) equilibrium constant of

as a function of temperature and pressure,

c) equilibrium constant of

as a function of temperature and pressure, and

d) ratio of partial pressures, p_3 (for N_2O_3) to p_4 (for N_2O_4) to the liquid molefractions, x_3 and x_4 , as a function of temperature.

From the model, it was possible to evaluate all partial pressures and y for a given x and T . From this data basis the effective stage separation factor, α_{eff} , was computed as a weighted average of the relevant isotope exchange reaction;

$$\alpha_{eff} = \frac{[\sum_{i=1}^2 x_i (s/s') f_i]_{liquid}}{[\sum_{i=1}^4 y_i (s/s') f_i]_{gas}}, \quad (150)$$

where the sum is taken over all chemical species present in each phase. The effective separation factor as a function of the liquid composition, the total pressure, and temperature are summarized in Figures 23 and 24.

Figure 23. Effective separation factor as a function of liquid composition, x , at various temperatures.⁽¹⁴⁸⁾

Figure 24. Effective separation factor as a function of total pressure at various temperatures.⁽¹⁴⁸⁾

The plots go over a maximum at every temperature. The effective separation factor is increased by the addition of nitric oxide to the system, as long as the added NO decreases the denominator of Eq. (150) for the vapor phase more than the numerator for the liquid phase. As the composition of the vapor phase approaches that of pure NO, a point is reached where further increase in the relative amount of +2 nitrogen in the system affects the gas phase composition less than the composition of the liquid phase; the numerator of Eq. (150) then decreases more rapidly than the denominator, and α_{eff} thus decreases. These tendencies are in good agreement with experiments.⁽¹⁵⁰⁾

V-A-2. Nitrogen-15 Fractionation Based on the NO/N₂O₃ Exchange System

The NO/N₂O₃ exchange system for ¹⁵N separation was investigated⁽²⁴⁾ at temperatures between -9°C and 20°C and under various pressures up to 4 atm, and α_{eff} , HETP and the liquid and gas phase compositions were determined.

Some typical results are presented in Table 26 and Figure 25.⁽²⁴⁾ A comparison with the Nitrox

Table 26. Comparison of Effects of Operating Parameters on the Size of NO/N₂O₃ Single Exchange Column System:
Basis = 1 g of 99% ¹⁵N / day ^(a)

	7A	10A	15	16	12B
Operating conditions:					
T (°C)	-9	+15.5	+14.5	+15.0	+20.0
P _i (atm)	1.00	1.00	3.40	4.08	2.00
λ/A (mL/cm ² ·min)	0.35	0.42	0.37	0.48	0.42
L/A (mmol N/cm ² ·min)	12.7	13.5	13.1	17.4	13.8
Parameters obtained:					
α _{eff}	1.36	1.006	1.032	1.030	1.020
HETP	1.02	0.75	1.11	1.07	1.02
Minimum required parameters:					
n _{min}	289	1706	324	345	515
(L/P) _{min} × 10 ⁻³	7.81	45.5	8.75	9.31	13.8
L _{min} (mmol N/min)	361	2105	405	431	640
H _{min} (cm)	295	1280	360	369	525
A _{min} (cm ²)	28.4	157	30.9	24.8	46.0
V _{min} (cm ³)	8.4 × 10 ³	2.0 × 10 ⁵	1.1 × 10 ⁴	9.2 × 10 ³	2.4 × 10 ⁴

^aλ = liquid flow rate (mL/min); A = inside cross-sectional area of the exchange column (cm²); L = interstage flow rate (mmol N/min); n_{min} = minimum number of stages; (L/P)_{min} = minimum reflux ratio; L_{min} = minimum interstage flow (mmol N/min); H_{min} = minimum height of exchange column = (n_{min})(HETP); A_{min} = minimum inside cross-sectional area of exchange column = L_{min}/(λ/A); V_{min} = minimum inside volume = (H_{min})(A_{min}).

Figure 25. α_{eff} as a function of total pressure at 15°C.
The solid lines are computed from various F-matrices.

Table 27. Comparison of the NO/NO System and Nitrox System ^(a)

	NO/HNO ₃ ^b			NO/N ₂ O ₃ ^c		
Operating parameters:						
T (°C)	25			14.5		
P _i (atm)	1			3.5		
λ/A (mL/cm ² ·min)	1.6	3.0	5.0	0	0.75 ^d	1.1 ^d
L/A (mmol N/cm ² ·min)	17	32	53	26		39
α _{eff}	1.055			1.032		
HETP	2.8	4.0	5.8	0.85		0.60
Minimum required parameters:						
n _{min}	191			324		
(L/P) _{min} × 10 ⁻³	4.90			8.75		
L _{min} (mmol N/min)	228			405		
H _{min} (cm)	535	764	1100	275		194
A _{min} (cm ²)	13	7.1	4.3	15.6		10.4
V _{min} × 10 ⁻³ (cm ³)	7.2	5.4	4.8	4.3		2.0

^a For the symbols, see the footnote of Table 26.

^b From Reference (15).

^c Based on Run 15.

system is given in Table 27.⁽²⁴⁾ For instance, at 15°C, we note the following.

- At 15°C, α_{eff} increases from 1.006 at 1 atm to 1.030 at 2.7 atm and levels off thereafter.
- The HETP correlates well with the linear flow rate of the gas stream, showing a dominant diffusion-control of the overall exchange rate.
- At 15°C, the HETP of the NO/N₂O₃ system under 3~4 atm can be smaller by a factor of 7 to 10 than that of the Nitrox system running under similar conditions, rendering the NO/N₂O₃ system a possibility of a significantly smaller plant in spite of its smaller separation factor.

The smaller volume of the NO/N₂O₃ system reduces not only the capital construction cost but also the holdup of the enriched isotope in the plant, thus lowering the plant operating cost.

V-B. Hydrophobic Catalysts for Hydrogen Isotope Exchange between Dihydrogen and Water

The exchange reactions of isotopes of hydrogen, H, D, and T, between gaseous dihydrogen and water such as,

are of importance in the nuclear energy technology. The reaction requires a catalyst containing Group VIII metals such as platinum. However, the nucleophilic water molecules tend to preempt the catalyst surfaces, excluding the hydrogen molecules from the active sites and resulting in an extensive retardation of the isotope exchange reactions on the surface.

One solution of the problem is a hydrophobic catalyst such as those developed by Chalk River. However, all heterogeneous catalyst could potentially suffer a loss of catalytic activity due to deterioration of crystallographic surface structure and chemical poisoning.

The goal of the present study is development of a catalyst of new type, which would provide an active control over the surface hydrophobicity and also afford *in situ* regeneration of the catalytic activity. The concept is as follows. A bed of an electrically conducting porous support material serves as (a) a packing material for a countercurrent, isotope exchange column and (b) a working electrode. The surface of the substrate electrode is chemically modified by electrodeposition of a conducting organic film and a catalyst metal. Basis of this design is two recent developments in electrochemistry.

- 1) The wettability of electroactive modifiers appears to be adjustable through electrochemical control of the average redox state of the modifier molecule.^(151, 152) They tend to be hydrophilic in the oxidized states and hydrophobic in the reduced states.
- 2) Fine aggregates of metals can be electrochemically imbedded in a polymer film formed on a substrate electrode.^(153 - 156) This was found in an effort to prepare conductive polymers.

These facts suggest that, if an electrode that is modified by an electrochemically active film is used as a support for a catalyst metal such as platinum, one may be able to control its hydrophobic environment for the catalyst surfaces *in situ*. This is neither electrolysis nor conventional electrocatalysis in that no significant electric current flows through the catalyst bed. Another advantage is the possibility of *in situ* regeneration of the catalytic activity by periodic potential excursions. It is well known that the catalytic activities of platinized platinum electrodes can be restored by repeated cycling of the potential between cathodic and anodic voltages.⁽¹⁵⁷⁾ The regeneration is at least partially due to crystallographic restructuring of the catalytic sites^(158, 159) and removal of chemical poisons from the surface.^(158, 160)

An account of our investigations for this goal will be presented in five sections; first on a highly controlled measurement of the hydrophobicity of polycrystalline platinum surfaces, second on chemical poisoning and regeneration of platinum surfaces, third on electrochemical modifications of platinum surfaces by films of poly (p-phenylene), fourth on modifications of platinum surface by other organic films, and fifth on capacitance measurement for the effective surface area of the edge plane of pyrolytic graphite.

Since none of these results have been published, we will report on these results in more detail than usual. The manuscripts in preparation for publication will bear reference numbers 2xx and be tabulated under "Manuscripts in Preparation" on page xvii.

V-B-1. Hydrophobicity of Polycrystalline Platinum Surfaces

In order to evaluate the hydrophobicity of an electrochemically modified platinum surface, it is first necessary to know the hydrophobicity of a clean platinum surface to serve as a reference point. The literature cites widely scattered values of contact angle formed between water and "clean" platinum surface, ranging from 0° (e.g, complete wetting) to an upper limit of about 61°. In the literature, the values differing from 0° had been assumed to be due mainly to presence of organic contaminants. Several investigators developed elaborate procedures to reduce and eliminate even the grease or oil originating in the atmosphere which come in contact during preparation of the solid surface. Many others developed several methods for preparation of a "reproducible and clean" platinum surfaces. The methods include mechanical abrasion, heat treatments, electropolishing, and uses of 30% hydrogen peroxide, aqua regia, hot and concentrated nitric acid, and hot dichromate-sulfuric acid. But, the discrepancies had still persisted.

As a result of our investigation reported below, we now believe that the literature values ranging from 55° to 62° represent clean, oxide-free platinum surfaces, while the surfaces that give the very small contact angles are, if not completely, very well covered with the oxides and hydroxyl species of platinum.

Preparation of Ultra-Pure Water

The most widely accepted ultra-high purity water for electrochemical works is prepared by Conway's pyrocatalytic distillation of water.⁽¹⁶¹⁾ The method involves a slow distillation through a rhodium-based catalyst bed heated to 750 °C in an all-quartz apparatus. We prepared the water by the Conway method and by a new, less time-consuming, distillation procedure involving acidic permanganate solution.^(162,201)

Figure 26 is a cyclic voltammogram of polycrystalline platinum surface in 0.5 M sulfuric acid prepared from the APD (acid permanganate distillation) water and 18 MΩ·cm sulfuric acid purchased from

AESAR. The platinum is 99.9985% Puratronic from Johnson Matthey, preconditioned by a newly developed procedure [162, 201]. The voltammogram shows all prominent anodic and cathodic peaks of a well-formulated surface of platinum. The slight shoulder on Trace "b" in the oxygen region near 1.0V corresponds to 1×10^{-4} gram-equivalents per liter of impurity. A similar voltammogram obtained in sulfuric acid made from the PCD (pyrocatalytically distilled) water contains three times as much impurity in this region. The APD water passed our purity test involving potentiostating at $-0.28 \text{ V}_{\text{RCE}}$, which is more stringent than Conway's test.

Figure 26. Cyclic voltammogram of polycrystalline platinum in $0.5 \text{ M H}_2\text{SO}_4$ prepared from APE water and AESAR sulfuric acid: Sweep rate = 100 mV/s . (a) freshly prepared and conditioned Pt, (b) after potentiostating at -0.28 V with nitrogen-stirring for 2.5 hrs, (c) after two potential cycles at 100 mV/s between -0.225 V and 1.25 V .

The APE water is much less tedious to prepare than the PCD water, and the apparatus for APD water is much less expensive than that required for the PCD process. We used the PCD water for all of our present investigation.

Hydrophobicity of Platinum

Figure 27 correlates the contact angle and the EOC (electrode open circuit) potential with the potential at which the platinum electrode was potentiostated (for 15 minutes) prior to the measurements. A $1/4"$ rod of 99.99% Pt was polished in a specially made mount using a series of garnet and diamond powers in APD water and dried in an oil-free, dry nitrogen stream before potentiostating. The Figure shows that both the wettability and oxidation state of Pt surface remain relatively constant until an applied potential

Figure 27. Contact angle and EOC vs. applied potential.

of about $0.9 V_{\text{SCD}}$ is reached. It is above this potential that the surface "oxides" are known to begin to accumulate. As the degree of surface oxidation increases, as indicated by the rise in the electrode open circuit potential, the wettability also increases. After being potentiostated at +1.7V or above, the surface shows a completely wetting.

V-B-2. Chemical Poisoning and Regeneration of Platinum Surfaces

Reactivation of a heterogeneous catalyst in a process equipment usually involves a shutdown of the equipment and reformation of the catalyst bed. This is an especially time-consuming and costly procedure for an isotope separation equipment because of the long startup period (cf. Section V-C in this report). The cost of shutdown and regeneration of the catalyst bed could be avoided, if the catalytic activity could be regenerated through manipulations of the electrochemical potential associated with the catalyst bed. As far as we are aware, there had been no systematic study of removal of chemical poisons from platinum surfaces.

We carried out such a study ^(162, 201) on the polycrystalline platinum electrodes in sulfuric acid prepared using the AESAR sulfuric acid and the APD water. A platinum working electrode was poisoned in a controlled manner. For each of the catalyst poisons, Ag(I), Cu(II), Cd(II), Pb(II), Cr(III), Mn(II), Fe(II), Co(II), Ni(II), Al(III), and bisulfite, which could be present in the industrial process water, the electrode was first potentiostated at $-0.25V_{\text{SCE}}$ for 15 minutes in a N_2 -stirred, $1 \times 10^{-6} M$ solution. Then, a series of triangular potential waves between $-0.24 V_{\text{SCE}}$ and $+1.25 V_{\text{SCE}}$ was imposed, while the electrode was still in the same poison-bearing solution. All solutions were sulfate solution except for Pb(II), which was a perchlorate solution. The cleaning scans were interspersed with an analytical CV scan. An example of the results is shown in Figure 28.

Figure 28. Recovery of polycrystalline Pt from Ag-poisoning

- 1st CV immediately after poisoning
- - - - 2nd CV after 2 cleaning cycles following the 1st CV
- 3rd CV after 5 more cleaning cycles following the 2nd CV
- 4th CV after 9 more cleaning cycles following the 3rd CV

In general, the cathodic deposits can be removed and the catalytic activity regenerated by several linear potential scans between these potential even when the solution is quiescent. The reactivation is achieved in one cleaning scan, if the solution is well-stirred.

V-B-3. Electrochemical Modification of Pt Surface by Poly(p-Phenylene) Film

The chemical modifier suitable for the present purpose has to satisfy the following requirements:

- 1) **Electrochemical polymerization:** The ability for the monomers to polymerize electrochemically allows for a close control over the polymerization process and a uniformity of the modifier film.
- 2) **Wide potential range:** The potential range for the electrochemical redox reactions of the modifier film should be sufficiently wide to accommodate various potential excursions, e.g., - 0.24 to +1.25 V_{SCE} in aqueous solutions.
- 3) **Stability:** The film should possess chemical, electrochemical, and mechanical stabilities in various aqueous solution.
- 4) **High hydrophobicity:** Contact angles between the surface and water or dilute aqueous solutions of more than 90° are preferred. In comparison, poly(tetrafluoroethylene) films give a contact angle of about 112°. In addition, a strong dependence of the hydrophobicity on applied potential would make its use more flexible.
- 5) **High electrical conductivity:** Under ideal conditions, a high conductivity of the reduced form of the modifier would not be needed for a steady state operation of the isotope exchange column. In order to implement an *in situ* catalyst regeneration, however, it would be necessary that the film has some degree of electrical conductivity in its reduced form.

We found one such chemical modifier in poly (p-phenylene).^(162, 202) Preparation and characterization of the film on platinum surface will be summarized below.

Number of Electrons Involved per Monomer in the Electrochemical Polymerization

The electropolymerization of p-terphenyl has been found to proceed via an intermediate dihydro-oligomer dication radicals,

which lose two protons to form a neutral oligomer.

The oxidation of p-terphenyl in this mechanism involves two electrons per monomer. However, the electropolymerization requires more than two electrons per monomer to accommodate incorporation (i.e., intercalation) of supporting electrolyte anions (i.e., counter-ions). The overall number of electrons per monomers, n , was found by Delahay's method⁽¹⁶³⁾ of plotting $\log [(\tau^{1/2} - t^{1/2}) / t^{1/2}]$ vs. potential, using a chronopotentiographic data. Here, τ is the transition time, and the slope of a straight line of the plot is equal to $-nF/2.303RT$. This method can be applied to every plateau of the chronopotentiogram, and every plateau has a corresponding peak in a cyclic voltammogram. The method was applied to a chrono-

Figure 29. Cyclic voltammogram of poly(p-phenylene) Sweep rate = 1.00 V/s. The polymer film was grown on a Pt disk in a solution of 2 mM p-terphenyl with 0.1 M TEATFB over a period of 2 seconds at a potential of +1.40 V_{Ag/Ag+}.

potentiographic plateau corresponding to the intercalation shoulder near the anodic switching potential of the cyclic voltammogram such as the one shown in Figure 29. This yields $n_{\text{interval}} = 0.2 \pm 0.04$ when the counter ion is BF_4^- , making the total n about 2.2 in tetraethylammonium tetrafluoroborate (TEATFB) as supporting electrolyte.

Nature of the CV-Peaks

For the cathodic peak at 1.1V in Figure 29, a plot of the peak current, i_p , as a function of the square root of the CV sweep rate, $v^{1/2}$, yielded a parabolic plot representable by

$$i_p (\mu\text{A}) = 103.8 v - 93.6 v^{1/2} \quad , \quad (152)$$

where v is in seconds. From this, we conclude that the cathodic peak around 1.1V has two equally contributing components; the one which is proportional to v is due to reduction of a surface-immobilized species, which we believe is the polymer, and the other which is proportional to $v^{1/2}$ due to deintercalation of a mobile species, which we think is tetrafluoroborate ion.

The peak current for the +0.4V reduction wave is proportional to the sweep rate, indicating that this wave is due to reduction of an immobilized species, which is contrary to a general believe that this is due to reduction of triphenylenic moieties present in the film.⁽¹⁶⁴⁾ We think this peak represents reduction of the polymer, that is, the reduction of a cationic form of the polymer to a neutral form.

Using Anson's method⁽¹⁶⁵⁾ for determination of n for a surface-bound species, we obtained $n = 1.1 \pm 0.3$ for the reduction wave at +0.4V. We also found that the charge consumed by this reduction wave is almost independent of the duration of polymerization time, while that for the wave at 1.1V increases with the polymerization time. Thus, the peak at 0.4V only depends on the surface area of platinum. From this, we obtained the surface coverage to be 2.5×10^{-14} monomer units per cm^2 of surface area.

We conclude that

- the reduction peak at 1.1V is half due to the reduction of the polymer and half due to the deintercalation of BF_4^- ,
- the reduction peak at 0.4V is solely due to the reduction of the polymer, and
- for the peak at 0.4V, $n = 1.1 \pm 0.3$, and the charge consumed depends only on the surface area, which fact can be used to determine the surface coverage and the film thickness.

Effects of Supporting Electrolyte

The three supporting electrolytes shown in Table 28 have sufficiently high solubility in acetonitrile and high stability against oxidation within our potential range of interest, i.e., 0.0V to +1.4V_{Ag/Ag⁺}.

Table 28. Properties of Acceptable Supporting Electrolytes

Supporting Electrolyte	Reduction potential ^(a) V (Ag/Ag ⁺)	Oxidation ^(b) potential V (Ag/Ag ⁺)	Solubility	Estimated ^(c) size (Å ³)
TEATFB ^(d)	-2.0	>+2.0	v.s.	52
TEAP ^(e)	-2.0	>+1.8	v.s.	62
TEA-TRF ^(f)	-2.0	>+1.6	v.s.	57

^(a) Potential measured using 1/4" diameter platinum disk electrode.

^(b) Lowest potential attempted.

^(c) Estimated using covalent bond radii and without solvation.

^(d) Tetraethylammonium tetrafluoroborate

^(e) Tetraethylammonium perchlorate

^(f) Tetraethylammonium triflate: $(\text{Et}_4\text{N})^+(\text{CF}_3\text{SO}_3)^-$

Figure 30. Cyclic voltammogram of poly(p-phenylene) film formed in different supporting electrolytes: Sweep rate = 1.0V/s. The CVs were taken in solutions containing the same supporting electrolyte as the polymerization solutions. The films were grown over a period of 10 s.

The polymerization was performed in each electrolyte. The polymerization current in TEATFB is significantly lower than those in the other two. In Figure 30 we compare the CVs of the polymer films taken in solutions containing the same supporting electrolyte as one in which they were grown.

The following conclusions were obtained.

- a) The differences in the shape of the oxidation/reduction couples, their positions, and their intensities are due to differences of stabilities of the films in the counter-ions and to differences in the effects of anion size on intercalation/deintercalation processes.
- b) Rousseau *et al.* ⁽¹⁶⁶⁾ observed that porphyrin π dications are stable in the presence of ClO_4^- but react with BF_4^- . They suggest that tetrafluoroborate ion can react with electrochemically generated cations and radical cations. This could be the reason for our observation of the lower polymerization current. An instability of tetrafluoroborate ion could lead to side reactions between the radical cations and anion during the polymerization and result in fewer intercalation sites. On the other hand, the higher intercalation peaks (cf: Figure 30) in the TEATFB solution than in the others is due to the smaller size of BF_4^- .
- c) The polymerization in the perchlorate solution produced a conductive film, which is contrary to other studies. ⁽¹⁶⁷⁾

Stability of Poly (p-Phenylene) Films

Another criterion for a desirable chemical modifier is a mechanical, chemical and electrochemical stability. All poly (p-phenylene) films that we prepared passed the following stability tests.

- a) Wrighton's scotch tape peel test: The color of the film, which ranges from bright blue-green for thin films to bright golden to brown for thicker films, do not change by the test. The CVs do not change by the test.
- b) Potential cycling in 0.05 M sulfuric acid between - 0.24 V_{SCE} and +1.25 V_{SCE} at a rate of 1V/s for 15 hours: The CVs taken before and after the tests in an aqueous solution and in an acetonitrile solution are presented in Figures 31 and 32, respectively.

Figure 31 shows the great enhancement of the features of catalytically active sites of platinum by the potential-cycling, indicating the beneficial restructuring of the exposed platinum surfaces that coexist with the poly(p-phenylene) film.

Figure 31. Cyclic voltammogram of platinum wire electrode coated with poly (p-phenylene) in 0.05 M sulfuric acid: Sweep rate = 1.0V/s, potential range = - 0.24 to +1.25 V_{SCE}. (a) --- before cycling, (b) — after 15 hr of cycling. The polymer film was grown in acetonitrile solution of 2mM p-terphenyl and 0.1 M TEATFB at +1.30 V (Ag/Ag⁺) for 20 seconds.

Figure 32. Cyclic voltammogram of platinum wire electrode coated with poly (p-phenylene) in 0.1 M TEATFB: Sweep rate = 1.0 V/s, potential range = 0.00 to +1.25 V (a) --- before the cycling in sulfuric acid, (b) — after the treatment (cf: footnote for Figure 31). The polymer film was prepared in the same way as in Figure 31.

Figure 32 shows an exceptionally good retention of poly (p-phenylene) characteristics after the very extended potential-excursion period of 15 hours. This is far superior than any other chemical modifiers that have been reported in literature for any purpose.

Hydrophobicity of Poly (p-Phenylene) Films

The contact angles between APD water and poly (p-phenylene) film on platinum measured in a water vapor-saturated nitrogen atmosphere are summarized in Table 29. The choice of supporting electrolyte has a definite effect on the wettability of the film. The films produced in TEATFB become increasingly more hydrophobic as the average thickness increases and reaches a value of 112° at 1 μm, which is the same contact angle as poly(tetrafluoroethylene). In contrast, the film grown in TEAP is hydrophilic.

In this connection, we also note that, while the deintercalation peak near 1.1 V in the CV of the film grown in TEATFB (cf: Figure 29) increases with the increasing film thickness, the similar cathodic peak in the CV of the film grown in the perchlorate environment (not shown) is almost independent of the film

Table 29. Contact Angles of Poly(p-phenylene) on Pt with water

Polymerization time (second)	Tetrafluoroborate		Perchlorate	
	Film thickness (μm)	Contact angle (degrees)	Film thickness (μm)	Contact angle (degrees)
0	--	57	--	57
2	0.13	80	0.11	77
4	0.21	89	0.18	54
6	0.28	100	0.24	52
8	0.36	102	0.28	50
10	0.41	107	0.33	48
20	0.65	112	--	--
30	0.91	114	--	--
40	1.33	117	--	--
50	1.68	115	--	--

* Film thickness was calculated using the experimentally determined coverage of 2.5×10^{14} units/cm² and $n = 2.2$ for the polymerization reaction in the TEATFB solution.

thickness. This is because, perchlorate ions being larger in size than tetrafluoroborate ions, they are trapped in the polymer cage as the polymer grows and thus do not participate in the charge transport as actively as BF_4^- .

Conclusion

We conclude that, among those we have studied, the best chemical modifier for our stated purpose is poly (p-phenylene) film grown in acetonitrile solution containing tetraethylammonium tetrafluoroborate as supporting electrolyte. It is durable and withstands an excessive potential cycling in neutral to acid aqueous media. It possesses a hydrophobicity that is at least as high as "Teflon." Although the polymer itself does not seem to conduct electric charges through the conjugated double-bond chain, the film is sufficiently porous to allow the transport of BF_4^- ions to the extent that the potential cycling yields the beneficial restructuring of the platinum surfaces.

V-B-4. Electrochemical Modification of Pt Surfaces by Other Organic Films

We have also studied the following other possible surface modifiers: ⁽¹⁶⁸⁾

- (a) poly (2,6-dimethylphenol)
- (b) poly (2, 3, 4, 5, 6-pentafluorostyrene)
- (c) tetrafluoro- 1,4-benzoquinone
- (d) poly (2, 6-difluorophenol)

The most interesting of these is (a). Polymorization on a platinum disk electrode was performed by linearly scanning the electrode potential from - 0.5V to 1.0V(vs. SCE/MeOH) in methanol solution containing 0.1 M 2,6-dimethylphenol and 0.2 M NaOH. The film of the PPO (polyphenylene oxide) looks blue on the platinum surface. The polymerization current goes over two maxima as the imposed potential was increased linearly as a function of time, indicating a preadsorption mechanism involving lower oligomers. The stability of the PPO film against a potential-cycling in 0.05 M H₂SO₄ as determined by cyclic voltammographic and FTIR measurements is satisfactory, although the stability of poly (p-phenylene) films is far superior.

Poly-(2,3,4,5,6-pentafluorostyrene), polyPFS, can be electrochemically prepared by potentiostating a platinum electrode in a 5 mM PFS + 0.2 M TEATFB in acetonitrile, but the polymer does not stay on the platinum surface well. The quinones and hydroquinones do not electrochemically polymerize. Poly-(2,6-difluorophenol), poly-DFP, adheres to the platinum surface well and is fairly stable against potential excursions.

V-B-5. Capacitance Measurement for Determination of Effective Surface Area of the Edge Plane of Pyrolytic Graphite

Forms of pyrolytic graphite such as reticulated vitreous carbon (RVC) are potential candidate for an acceptable substrate material for the catalyst bed envisioned in the present study. A catalyst, such as platinum, and a chemical modifier, such as poly(p-phenylene) would be codeposited on the substrate, which could also act as a column packing in an isotope exchange column. For further studies we would have to have a means of determination of the effective surface area of such a substrate. In order to provide a basis for such determination, we have investigated ^(169, 203) the electric capacitance measurement methods for the area of the edge plane of the pyrolytic graphite.

Several techniques exist for the measurement of effective area of solid surfaces. Determination of effective area of the edge plane of pyrolytic graphite is especially challenging, because the microfissure openings usually exist on the surface.

The method for the surface area of an electrode based on the electrical capacitance measurement has an important advantage over other common methods such as those using the surface adsorption of

gaseous molecules in that the capacitance measurement in a solution is non-destructive and tends to yield results which are more representative of the environment in which the electrode will be eventually used. The concept of capacitance measurement is not new, however, a satisfactory procedure has not been established. Here is a report on a method which we believe is promising.

Some explanations of the basics are in order. Suppose first that a conductor electrode is dipped in a solution which contains nothing but an electrochemically inactive electrolyte in a sufficiently high concentration so that the resistance across the solution in the cell is negligible and the capacitance charging is instantaneous. Further suppose that the electrode surface is perfectly planar. Then, the capacitance is

$$\frac{1}{C} = \frac{1}{C_H} + \frac{1}{C_d} \quad , \quad (153)$$

where C_H and C_d are the capacitances of the Helmholtz plane and the diffusion double layer, respectively. Suppose further that a triangular voltage wave is imposed on the electrode; the voltage starts at a cathodic value, V_i , say, $V_i < 0$ relative to a reference electrode, linearly increases through $V = 0$ until it reaches an anodic switching potential $V_f = -V_i$, and then linearly decreases through $V = 0$ until it gets back to V_i . For this idealized electrochemical cell, the output current $i(t)$ looks like Figure 33a, and the capacitance could be calculated from $C = i_c / v$, where i_c is indicated in Figure 33a and v is the voltage scan rate in volts/sec. If a faradaic current, i_f , is present, but if the input voltage excursion is not so excessive as to assure that the faradaic current is nowhere near the limiting current, then $i(t)$ would look like Figure 33b, and the capacitance can be obtained as $C = (i_1 + i_2) / (2v)$, where $i_1 (= i_c + i_f)$ and $i_2 (= i_c - i_f)$ are indicated in Figure 33b. In the Cases (a) and (b) a graphical interpretation would directly yield the capacitance. If, in Case (a), the resistive load through the electrolyte solution is finite, the charging current would follow the exponential form,

$$i(t) = vC(1 - e^{-t/RC}) \quad , \quad (154)$$

where v is again the slope of the voltage scan.

On the other hand, if the electrode surface has sufficiently complicated structure such as the fissures on the real graphite's edge plane, the charging current should be regarded as consisting of two or more parallel capacitors, each of which may be charged at a different rate. It is reminded that only parallel capacitors, and not the capacitors connected in series, can be charged at different rates. Let us call the electrode surface capacitances that charge at independent rates *distributed capacitance*. Figures 33c and 33d respectively depict such a distributed capacitance without and with a faradaic current.

The charging current plot that we obtained for the edge planes of pyrolytic graphite in acetonitrile solutions of tetraethylammonium tetrafluoroborate (TEATFB) is of Type (d). Previous works on the cases of distributed capacitance have either addressed themselves with complex modelling⁽¹⁷⁰⁾ or used

Figure 33. Charging/Discharging Curves: (a) Current response of an ideal capacitor without a faradaic current. (b) Current response in presence of a faradaic current. (c) Current response in presence of a distributed capacitance and in absence of a faradaic current. (d) Current response in presence of a distributed capacitance and a faradaic current.

extremely artificial graphical methods.⁽¹⁷¹⁾ For instance, Figure 34 illustrates one of the graphical methods by Soffer⁽¹⁷¹⁾, in which the charging/discharging currents are plotted against the applied voltage. Using Soffer's terminology, the "fast charging" ends at Point a where the curve is supposed to suddenly start to deviate from a "vertical line." The vertical line is supposed to represent the fast charging. The region between Points a and b is due to "slow charging." The difference in the current between Points a and c is taken to be equal to twice i_f , and that between Points b and c is taken to be equal to twice $(i_c + i_f)$,

Figure 34. Soffer's graphical method: The vertical axis is current, and the horizontal axis is potential.

where i_f and i_s are the currents due to the fast and slow charging, respectively. The method suffers from the difficulty and subjectiveness of determination of Point a in Figure 34.

We have fitted the experimental charging curve of the Type d (Figure 33d) to a functional form,

$$i(t) = \nu C_1(1 - e^{-t/R_1 C_1}) + \nu C_2(1 - e^{-t/R_2 C_2}) + a, \quad (155)$$

where, again, ν is the voltage sweep rate, and a is a constant representing the background current. The results are summarized below.

- a) All charging/discharging current data on the edge plane of our pyrolytic graphite can be fitted satisfactorily using only two exponential terms. An example is shown in Figure 35. The saw-tooth trace is the raw experimental data, the saw teeth being due to those superimposed on the input potential ramp produced by a software. The data was deconvoluted to the form of Eq. (155). A single set of parameters fit both the charging and discharging currents equally well, indicating a reversible nature of the transport processes.
- b) Inclusion of a faradaic current on the right-hand side of Eq. (155), i.e., a term proportional to the sweep rate, ν , and time, does not improve the goodness of the fit. This is as expected under our experimental conditions.
- c) Figure 36 summarizes the fast and slow charging capacitances of many specimens of vacuum-heat-treated pyrolytic graphite as a function of constant potential superimposed on the analytic potential ramp. The geometric area is 0.0792 cm^2 , and all capacitance measurements were conducted in 0.1 M TEATFB in acetonitrile. Each point in Figure 26 represents a measurement, in each of which an input triangular wave with an amplitude of 5 mV and sweep rate of 1 Hz was applied. The average values of the capacitance correspond to $280 \pm 30 \text{ } \mu\text{f/cm}^2$ for the fast-charging component and $90 \pm 30 \text{ } \mu\text{f/cm}^2$ for the slow-charging component. If the thicknesses of the electric double layers of both components are alike, this result would mean that about one quarter of the surface areas of the edge-plane of our specimens of pyrolytic graphite are due to the intra-structure.
- d) The vacuum heat treatment of graphite tends to decrease the capacitance, which appear to be due to a decrease in the surface oxygen functionalities.

- e) We have also measured the capacitance as a function of potential scan rate. While the fast capacitance is practically independent of the scan rates in the range between 1 Hz to 10 Hz, the slow capacitance roughly dropped two-folds as the scan rate was increased from 1 Hz to 5 Hz and ten-folds as ν was increased from 1 Hz to 10 Hz. This observation is consistent with a notion that the slow process is due to a charging of an internal structure such as the microfissures underneath the edge plane surface.

Figure 35. Two-component fit of charging/discharging currents of edge plane of pyrolytic graphite in 0.1 M TEATFB in acetonitrile: S = slow component, F = fast component

Figure 36. Capacitances of edge plane of pyrolytic graphite as a function of applied potential: Geometric area = 0.0792 cm², supporting electrolyte = 0.1 M TEATFB in acetonitrile. (a) fast-charging component, (b) slow-charging component.

Conclusion

- a) The capacitance measurement is reproducible to within 10%.
- b) Since any capacitors connected in series must be charged all at the same rate, the two exponential terms means that there are at least two, significant, parallel charging pathways on the edge- plane surface of the pyrolytic graphite. There are significant internal surface area detectable by the capacitance measurement technique that we have developed.

V-C. Transient Behavior of Square Cascade of Isotope Separation Stages

The transient behavior of a square cascade is important for an efficient operation of isotope separation plants, especially because the stage separation factors (α) of isotope separation processes are close to unity, *i.e.*, $|\epsilon| = |\alpha - 1| \ll 1$. The behavior was theoretically analyzed by Cohen.⁽¹⁷²⁾ The theory is useful in a number of ways, for instance, in analyzing the behavior of a plant during a cold start-up period, and in determination of α from laboratory scale experiments. We refined Cohen's theory⁽¹⁰⁾ and pointed out⁽⁹⁾ one of its possible consequences regarding the processes based on inverse isotope effects.

Consider the material balance of the isotope that one wants to enrich between the product stage and the s -th stage in a square cascade of separation stages. The consideration leads to the following partial differential equation:⁽¹⁷⁰⁾

$$\lambda \frac{\partial N}{\partial t} = \frac{\partial^2 N}{\partial s^2} - \epsilon \frac{\partial}{\partial s} [\Psi N + N(1 - N)] , \quad (156)$$

where $N = N(s, t)$ is the average atom fraction of the target isotope in the input streams for the s -th stage at time t after the start-up time of the cascade. The stage numbers are counted starting at the feed point ($s = 0$) toward the product end ($s = n$). Also, $\lambda = 2h$, where h is the holdup per stage per unit flow, or the average process time per stage, and $\Psi = 2P / (\epsilon L)$, where P is the product withdrawal rate, and L the total interstage flow rate at stage s . In a square cascade, L is independent of s . Under the total reflux, L is also equal to twice the reflux rate.

The solution of Eq. (156) by Cohen is in the form of Eq. (157) for the case of $N \ll 1$ and the following initial and boundary conditions; $N(s, t=0) = N_0$ for all s -values, $N(0, t) = N_0$ at all t , $P = 0$ at all t , and the holdup in the product refluxer = H :

$$\frac{N(n, t) - N_0}{N_{\infty} - N_0} = 1 - Ae^{-B\tau} , \quad (157)$$

where $N_{\infty} = e^{\epsilon}$ which is the steady state atom fraction at the product end, $\tau = t / \lambda n^2$ is a dimensionless time, and both A and B depend on two dimensionless parameters; $A = A(\epsilon n, K/\lambda n)$ and $B = B(\epsilon n, K/\lambda n)$, in which $K = 2H/L$ and $K/\lambda n = H/Lhn$. Cohen mapped A and B for $K/\lambda n = 0.1$ (0.1) 0.5 and for $\epsilon n = 0.0$ (0.1) 1.2 and stated that this solution is applicable to $t > \lambda n^2/10$.

We extended the time limit to well below $\lambda n^2/10$ by deriving a higher-order solution of Eq. (156) in the form,

$$\frac{N(n, t) - N_0}{N_{\infty} - N_0} = 1 - A_1 e^{-B_1\tau} - A_2 e^{-B_2\tau} , \quad (158)$$

and tabulated A_1 , B_1 , A_2 and B_2 for $K/\lambda n = 0.0$ (0.05) 0.5 and $\epsilon n = -1.2$ (0.2) 1.2. The relative importance of the second-order term is demonstrated by Figure 37.⁽⁹⁾ The Figure shows that the second-order term is significant even at the time, t , which is within the Cohen's limit of $\tau > 0.1$.

Figure 37. Relative importance of the first and second terms of Eq. (158).

In particular, the second-order term is very significant in all cases of inverse isotope effect. The isotope of light elements that we are usually interested in enriching, e.g., D, ^{13}C , ^{15}N , ^{34}S , is heavier than the naturally most abundant isotopes. Thus, for instance, in a distillation process, the desired isotope usually enriches in the liquid phase, which corresponds to a positive ϵn and to a normal vapor pressure isotope effect. If the VPIE is inverse, then ϵn is negative, and the second term of the solution, Eq. (37), is not at all negligible.

More generally, the transient behavior of a separation cascade is not symmetric with respect to change of signs in ϵ . The importance of this fact is illustrated by Figure 38⁽⁹⁾, in which the time it takes for the plant to attain 95% of a steady state separation, $\tau_{0.95}$, is plotted as a function of ϵn and $k/\lambda n$. The transient time is a lot shorter in the domain of inverse isotope effect than in the domain of normal isotope effect.

Among other implications of this finding, a plant based on an inverse isotope effect will take a lot shorter time to reach a steady state and thus allow a product withdrawal sooner than a plant based on a normal isotope effect having the same absolute magnitude of separation factor and the same number of separative stages.⁽¹⁰⁾ Besides, it is easier to construct a product-end refluxer with low refluxer-holdup, if the desired isotope enriches in the gas phase.

Figure 38. The 95% equilibrium time

[blank page]

VI. References

101. Bigeleisen, J.; Mayer, M. G., *J. Phys. Chem.* **1947**, *15*, 261.
102. Bigeleisen, J., *J. Phys. Chem.* **1963**, *38*, 769.
103. Bigeleisen, J., "Proceedings of the Amsterdam Symposium on Isotope Separation," p. 121, 1991, North-Holland Publishing Company, Amsterdam.
104. Lanczos, C., "Applied Analysis," 1956, Prentice Hall, Englewood Cliffs, New Jersey.
105. Bigeleisen, J.; Ishida, T. *J. Chem. Phys.* **1968**, *48*, 1311.
106. Ishida, T.; Spindel, W.; Bigeleisen, J., *Adv. Chem. Ser.*, **1969**, *89*, 192.
107. Bigeleisen, J.; Ishida, T.; Spindel, W., *Proc. Natl. Acad. Sci. USA* **1970**, *67*, 113.
108. Wilson, E. B.; Decius, J. C.; Cross, P. C., "Molecular Vibrations," 1955, McGraw-Hill, New York.
109. Bigeleisen, J., *J. Chem. Phys.* **1958**, *28*, 694.
110. Bigeleisen, J., *J. Chem. Phys.* **1955**, *23*, 2264.
111. Singh, G.; Wolfsberg, M., *J. Chem. Phys.* **1976**, *62*, 4165; Skaron, S. A.; Wolfsberg, M., *J. Am. Chem. Soc.* **1977**, *99*, 5253.
112. Decius, J. C.; Wilson, E. B., Jr., *J. Chem. Phys.* **1951**, *19*, 1409.
113. Bernstein, H. J.; Pullin, A. D. E., *J. Chem. Phys.* **1953**, *21*, 2188.
114. Pitzer, K. S., *J. Chem. Phys.* **1937**, *5*, 473, *ibid.*, 1940, *8*, 711, *Chem. Rev.* **1940**, *27*, 39.
115. Cottrell, T. L. *J. Chem. Soc.* **1948**, 1448.
116. Pitzer, K. S.; Catalano, E. *J. Am. Chem. Soc.* **1956**, *78*, 4844.
117. Bernstein, H. J. *J. Chem. Phys.* **1952**, *20*, 263, 1328.
118. Evans, J. C.; Bernstein, H. J. *Can. J. Chem.* **1955**, *33*, 1171.
119. Bernstein, H. J. *Can. J. Chem.* **1956**, *34*, 617.
120. Bernstein, H. J. *J. Chem. Phys.* **1956**, *24*, 911.
121. Bernstein, H. J. *J. Phys. Chem.* **1965**, *69*, 1550.
122. Fujimoto, T.; Shingu, H. *J. Chem. Soc. Jpn.* **1962**, *83*, 19.
123. Hartshorn, S. R.; Shiner, V. J., Jr., *J. Am. Chem. Soc.* **1972**, *94*, 9002.
124. Wolfsberg, M., *J. Chim. Phys.* **1963**, *60*, 15.
125. Bigeleisen, J.; Goldstein, P., *Z. Naturforsch.*, **1963**, *A18*, 205.
126. Bigeleisen, J.; Weston, R. E., Jr.; Wolfsberg, M., *Z. Naturforsch.* **1963**, *A18*, 210.
127. Person, W. B.; Newton, J. H., *J. Chem. Phys.* **1974**, *61*, 1040.

128. Pauling, L., *"The Nature of the Chemical Bond,"* 3rd ed., 1960, Cornell University, NY.
129. Wilmshurst, J. K.; *J. Chem. Phys.* **1958**, *28*, 733.
130. Inomata, N.; Masuda, S., *Chem. Lett.*, **1982**, 1003.
131. Boyd, R. J.; Edgecombe, K. E., *J. Am. Chem. Soc.* **1988**, *110*, 4182.
132. Luo, Y. R.; Benson, S. W., *J. Phys. Chem.* **1989**, *93*, 3306.
133. Gordy, W., *Phys. Rev.* **1946**, *61*, 604.
134. Wilmshurst, J. K., *J. Chem. Phys.* **1957**, *27*, 1129.
135. Jancso, G.; Van Hook, W. A., *Chem. Rev.* **1974**, *74*, 691.
136. Armstrong, G. T.; Brickwedde, F. G.; Scott, R. B., *J. Res. Natl. Bur. Stand.* **1955**, *55*, 39.
137. Johns, T. F., *"Proceedings of the International Symposium on Isotope Separation,"* **1958**, North-Holland Publishing Co., Amsterdam.
138. Bigeleissen, J.; Cragg, C. B.; Jeevanandam, M., *J. Chem. Phys.* **1967**, *47*, 4335.
139. Kirshenbaum, I.; Urey, H. C., *J. Chem. Phys.* **1942**, *10*, 706.
140. Wolff, H.; Höpfner, A., *Ber. Bunsenges. Phys. Chem.* **1969**, *73*, 480.
141. Thode, H. G., *J. Am. Chem. Soc.* **1940**, *62*, 581.
142. Armstrong, G. T., *Natl. Bur. Standards Report No. 2626*, June, **1953**.
143. Overstreet, R.; Giaque, W. F., *J. Am. Chem. Soc.* **1937**, *59*, 254.
144. Spirko, V., *J. Mol. Spectrosc.* **1983**, *101*, 30.
145. Stern, M. J.; Van Hook, W. A.; Wolfsberg, M., *J. Chem. Phys.* **1963**, *39*, 3179.
146. Bader, R. F. W.; Beddall, P. M., *J. Chem. Phys.* **1972**, *56*, 3320.
147. Bader, R. F. W.; Beddall, P. M.; Peslak, J., Jr., *J. Chem. Phys.* **1973**, *58*, 557; Bader, R. F. W.; Henneker, W. H.; Cade, P. E., *ibid.* **1967**, *46*, 3341.
148. Ishida, T.; Spindel, W., *J. Chem. Eng. Data*, **1970**, *15* (No. 1), 107.
149. Eshelman, D. M.; Torre, F. J.; Bigeleisen, J., *J. Chem. Phys.* **1974**, *60*, 420.
150. Monse, E. U.; Kauder, L. N.; Spindel, W., *J. Chem. Phys.* **1964**, *41*, 3898; Monse, E. U.; Kauder, L. N.; Spindel, W., *Z. Naturforsch.* **1963**, *18a*, 235; Monse, E. U.; Spindel, W.; Kauder, L. N.; Taylor, T. I. *J. Chem. Phys.* **1960**, *32*, 1557.
151. Willman, K. W.; Murray, R. W., *Anal. Chem.* **1983**, *55*, 1139.
152. Hernandez R.; Diaz, A. F.; Waltman, R.; Bargon, J. *J. Phys. Chem.* **1984**, *88*, 3333.
153. Dao, W. H.; Kuwana, T., *J. Am. Chem. Soc.* **1984**, *106*, 473.
154. Weisshaar, D. E.; Kuwana, T., *J. Electroanal. Chem.* **1984**, *163*, 395.
155. Bookbinder, D. C.; Bruce, J. A.; Dominey, R. N.; Lewis, N. S.; Wrighton, M. S. *Proc. Natl. Acad. Sci. USA* **1980**, *77*, 6280.
156. Tourillon, G.; Garnier, F., *J. Phys. Chem.* **1984**, *88*, 5281.
157. Hammett, L. P., *J. Am. Chem. Soc.* **1924**, *46*, 7.

158. Kinoshita, K. in *Modern Aspects of Electrochemistry*; 1982, Bockris, J. O'M., Ed., Vol. 14, Chapter 8, Plenum, New York.
159. Chialvo, A. C.; Triaca, W. E.; Arvia, A. J. *J. Electroanal. Chem.* **1983**, *146*, 93.
160. Bilmes, S. A.; De Taconi, N. R.; Arvia, A. J. *J. Electroanal. Chem.* **1984**, *164*, 129.
161. Conway, B. E.; Angerstein-Kozlowska, H.; Sharp, W. B. A.; Criddle, E. E., *Anal. Chem.* **1973**, *45*, 1331.
162. Krebs, L. C.; Ishida, T.; "Characterization of Electrochemically Modified Polycrystalline Platinum Surfaces," 1991, U.S. Department of Energy Report, DOE/ER/13855-6.
163. Delahay, P.; Mattax, C. C., *J. Am. Chem. Soc.* **1954**, *26*, 874.
164. Schiavon, G.; Zecchin, S.; Zotti, G.; Cattarin, S., *J. Electroanal. Chem.* **1986**, *213*, 53.
165. Brown, A. P.; Koval, C.; Anson, F. C., *J. Electroanal. Chem.* **1976**, *72*, 379.
166. Rousseau, K.; Farrington, G. C.; Dolphin, D., *J. Org. Chem.* **1972**, *37* (24), 3968.
167. Aeiyaeh, S.; Dubois, J. E., Lacaze, P. C., *J. Chem. Soc., Chem. Commun.* **1986**, 1608; Aeiyaeh, S.; Lacaze, P. C., *J. Poly. Sci., Part A: Poly. Chem.* **1989**, *27*, 515.
168. Jayanta, P. S.; Ishida, T., "Preparation of Polymer-Modified Electrodes: A Literature and Experimental Study," 1991, U. S. Department of Energy Report, DOE/ER/13855-5.
169. Minick, S. K.; Ishida, T., "Capacitance of Edge Plane of Pyrolytic Graphite in Acetonitrile Solutions," 1991, U. S. Department of Energy Report, DOE/ER/13855-4.
170. Bigeleisen, J.; Brooks, F. P.; Ishida, T.; Ribnikar, S. V., *Rev. Sci. Instr.* **1968**, *39*, 356.
171. Gagnon, E. G., *J. Electrochem. Soc.* **1974**, *121* (11), 1444; Gagnon, E. G., *ibid.*, **1975**, *122* (4), 525; Gagnon, E. G. *ibid.*, **1974**, *121*, 512; Austin, L. G.; Gagnon, E. G., *ibid.* **1973**, *120* (2), 251.
172. Tobias, H.; Soffer, A., *J. Electroanal. Chem.* **1983**, *148*, 221; Oren, Y.; Soffer, A., *ibid.* **1985**, *186*, 63.
173. Cohen, K., "The Theory of Isotope Separation as Applied to the Large-Scale Production of U-235," 1951, McGraw-Hill, New York.

END

**DATE
FILMED**

12 / 18 / 92