DOE/ER/13214--7

DE92 018399

SUMMARY PROGRESS REPORT**U.S. DEPARTMENT OF ENERGY GRANT No. DE-FG02-84ER13214****EFFECTS OF FREEZING AND COLD ACCLIMATION
ON THE PLASMA MEMBRANE OF ISOLATED PROTOPLASTS****May 16, 1987 - June 1, 1991**

**Peter L. Steponkus
Department of Soil, Crop and Atmospheric Sciences
609 Bradfield Hall
Cornell University
Ithaca, New York 14853
607-255-1707**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER**DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED**

Summary Statement: The focus of this project is on lesions in the plasma membrane of protoplasts isolated from non-acclimated rye leaves (NA protoplasts) that occur during freezing to temperatures below -5°C and result in alterations in the semipermeable characteristics (leakiness) of the plasma membrane and preclude osmotic expansion following thawing of the suspending medium. This form of injury, referred to as "loss of osmotic responsiveness", is associated with the formation of large, aparticle domains in the plasma membrane, aparticle lamellae subtending the plasma membrane, and lamellar-to-hexagonal_{II} phase transitions in the plasma membrane and subtending lamellae. Lamellar-to-hexagonal_{II} phase transitions do not occur during freeze-induced dehydration of protoplasts isolated from cold acclimated leaves (ACC protoplasts). The goals of this project are (i) to provide a mechanistic understanding of the mechanism by which freeze-induced dehydration effects the formation of aparticle domains and lamellar-to-hexagonal_{II} phase transitions and (ii) to determine the mechanisms by which cold acclimation and cryoprotectants preclude or diminish these ultrastructural changes. Our working hypothesis, which has evolved from both experimental and theoretical studies, is as follows: The formation of aparticle domains and lamellar-to-hexagonal_{II} phase transitions in the plasma membrane and subtending lamellae are manifestations of hydration-dependent bilayer-bilayer interactions. At temperatures of -10°C , the chemical potential of ice is of sufficient magnitude (-12 MPa) to remove water from the hydrophilic surfaces of membranes resulting in the close apposition of the plasma membrane with various endomembranes (e.g., the chloroplast envelope and endoplasmic reticulum). The formation of large aparticle domains is a consequence of the exclusion of membrane proteins from regions of close approach because of their large steric/hydration forces. Further dehydration results in the demixing of lipid components that differ in their hydration characteristics and results in the formation of domains that are enriched in H_{II} -forming lipids, which then undergo the lamellar-to- H_{II} phase transition. Lamellar-to-hexagonal_{II} phase transitions are precluded in ACC protoplasts because of alterations in membrane lipid composition: The differential propensity for dehydration-induced H_{II} formation is also observed in liposomes prepared from the plasma membrane lipids of non-acclimated and cold acclimated rye leaves. And, enrichment of the plasma membrane of NA protoplasts with dilinoleoylphosphatidylcholine precludes H_{II} formation in the plasma membrane. The decreased propensity for H_{II} formation following cold acclimation or artificial enrichment with di-unsaturated species of phosphatidylcholine is attributed to an increase in the hydration characteristics of the plasma membrane, which results in a greater spatial separation between the plasma membrane and endomembranes during freeze-induced dehydration. During cold acclimation there is an increase in the proportion of highly hydrated lipid species (phospholipids) and a decrease in the proportion of poorly hydrated species (cerebrosides) in the plasma membrane.

A. Phenomenology of Freezing Injury: Loss of Osmotic Responsiveness and Hexagonal_{II} Formation

Survival of protoplasts isolated from non-acclimated leaves (NA protoplasts) decreases from $>90\%$ to $<10\%$ over the range of -2°C to -10°C , (LT_{50} of -5°C). Over the range of -2°C to -5°C , injury is a consequence of expansion-induced lysis (EIL); at temperatures below -5°C , loss of osmotic responsiveness (LOR) is the predominant form of injury - its incidence increasing from 12% to 87% over the range of -5°C to -10°C (Uemura & Steponkus, 1989b). After one week of cold acclimation, survival increases to $>65\%$ at -10°C , with approximately 35% survival at -30°C (LT_{50} of -13°C). After 2, 3, and 4 weeks of cold acclimation there are progressive increases in survival at all temperatures over the range of -10 to -30°C , with the LT_{50} increasing to -26°C . The large increase in survival at -10°C after one week of cold acclimation is associated with a decrease in the propensity for dehydration-induced lamellar-to-hexagonal_{II} phase transitions (Sugawara & Steponkus, 1990). Freezing over the range of -7.5 to -12.5°C resulted in the formation of aparticle domains in the plasma membrane, the formation of aparticle lamellae subtending the plasma membrane, and lamellar-to-hexagonal_{II} phase transitions in the plasma membrane and subtending lamellae. The aparticle lamellae were usually observed as one or more stacked layers subtending the plasma membrane. The lamellae appeared to be formed as a result of the close approach of other membranes (chloroplast outer envelope, vesicles) to the plasma membrane. H_{II} was also observed in the outer envelope of the chloroplasts and the ER. After 1 week of cold acclimation, neither large, aparticle domains nor hexagonal_{II} phase are observed in the plasma membrane or endomembranes of protoplasts frozen to either -10°C (65% survival) or -25°C (15% survival).

B. Alterations in the Plasma Membrane Lipid Composition and Lamellar-to-Hexagonal_{II} Phase Transitions

A molecular species analysis of the plasma membrane lipids of rye leaves has been reported by Lynch & Steponkus (1987). In the plasma membrane of non-acclimated leaves, $>50\text{ mol}\%$ of the total lipid consists of free sterols ($33\text{ mol}\%$), sterylglucosides ($15\text{ mol}\%$), and acylated sterylglucosides ($4\text{ mol}\%$); after cold acclimation, the total proportion is similar - but the proportion of free sterols increases ($44\text{ mol}\%$) and the sterylglucosides ($6\text{ mol}\%$) and acylated sterylglucosides ($1\text{ mol}\%$) decrease. Phospholipids, predominantly phosphatidylcholine ($15\text{ mol}\%$) and phosphatidylethanolamine ($11\text{ mol}\%$), with lesser amounts of phosphatidylglycerol, phosphatidylserine, and

phosphatidylinositol, comprise 32 mol% of the total plasma membrane lipid in non-acclimated leaves. After cold acclimation, the phospholipid content increases to 42 mol%. Although the relative proportions of the individual phospholipids did not change appreciably, di-unsaturated species of both phosphatidylcholine and phosphatidylethanolamine accounted for the net increase. Glucocerebrosides, which were comprised of a long-chain hydroxy fatty acid (primarily C₂₂ and C₂₄) and a long-chain base (predominantly either hydroxysphinganine or sphingadienine) and glucose, decreased from 16 to 7 mol% after cold acclimation. Within these classes, more than 100 different molecular species have been identified - however, none are unique to either non-acclimated or cold acclimated leaves.

Because of the complexity of the lipid composition of the plasma membrane and the fact that the proportion of virtually every molecular species changed during cold acclimation, the role of these lipid changes cannot be determined from a correlative analysis. Therefore, more direct approaches were taken to establish that the alterations in the lipid composition are responsible for the decreased propensity for dehydration-induced lamellar-to-hexagonal_{II} phase transitions after cold acclimation. Dehydration of liposomes (MLVs) comprised of lipids isolated from the plasma membrane fractions of non-acclimated leaves results in the formation of H_{II} phase; under similar conditions the H_{II} phase was not observed in liposomes prepared from the plasma membrane lipids of cold acclimated leaves (Cudd & Steponkus, 1988). Although these studies established that the differential propensity for dehydration-induced H_{II} formation was a consequence of alterations in the lipid composition of the plasma membrane, the question remained as to which of the many changes were responsible for the differential behavior.

In subsequent studies, a protoplast-liposome fusion technique (Arvinte & Steponkus, 1988) was used to selectively enrich the plasma membrane of NA protoplasts with individual lipids (Uemura & Steponkus, in preparation). Dehydration-induced H_{II} formation in the plasma membrane of NA protoplasts is precluded by artificial enrichment of the plasma membrane with dilinoleoylphosphatidylcholine (Sugawara & Steponkus, in preparation). Following fusion and freezing of the protoplasts to -10°C, the frequency and size of the aparticle domains in the plasma membrane were greatly reduced and no H_{II} was observed in the plasma membrane. However, aparticle lamellae subtending the plasma membrane were still apparent, and H_{II} was observed in membranes other than the plasma membrane (e.g., the chloroplast outer envelope and the ER). This appearance of H_{II} in the endomembranes was not unexpected since the fusion procedure only results in alteration of the plasma membrane lipid composition. Nevertheless, the results suggest that the decreased propensity for H_{II} formation following the first week of cold acclimation is associated with increases in the unsaturated species of phosphatidylcholine.

C. Hydration Characteristics of Plasma Membrane Lipids

As dehydration is central to both cellular injury during freezing and the formation of the H_{II} phase, we have undertaken an examination of the hydration characteristics of the lipid components common to the rye plasma membrane. It is our working hypothesis that (i) the formation of aparticle domains and lamellar-to-hexagonal_{II} phase transitions in NA protoplasts are the result of the close approach of bilayers during freeze-induced dehydration and (ii) that the decreased propensity for H_{II} formation in ACC protoplasts is, in part, a consequence of alterations in the hydration characteristics of the plasma membrane (see Steponkus & Lynch, 1989; Steponkus *et al.*, 1990). To test this hypothesis, we determined the hydration characteristics of plasma membrane lipids isolated from non-acclimated and cold acclimated rye leaves. In preliminary studies, we determined that plasma membrane lipids isolated from cold acclimated rye leaves were more hydrated than those isolated from non-acclimated leaves (Lynch & Steponkus, 1989a). Desorption isotherms (water content vs. water potential) were constructed following equilibration of the total plasma membrane lipid extracts at known vapor pressures. Over the range of -3 to -300 MPa, the plasma membrane lipids from cold-acclimated leaves were significantly more hydrated than those isolated from non-acclimated leaves.

To establish the compositional differences in lipid composition that are responsible for the differential hydration characteristics of NA and ACC plasma membrane lipids, we determined the hydration characteristics of lipid species representative of the various lipid classes and mixtures of these components (Lynch & Steponkus, 1989a, b; Steponkus *et al.*, 1990). The major classes of the rye plasma membrane lipids are sterols, sterylglucosides, cerebrosides, and phospholipids, including phosphatidylcholine (PC) and phosphatidylethanolamine (PE). We have constructed desorption isotherms to examine the hydration characteristics of these lipids dispersed in water. Sterols, sterylglucosides, and cerebrosides are weakly hydrated, binding 6, 5, and 1 wt % water at -10.9 MPa, or 1.5, 1.7, and 1 moles of water/mole of lipid, respectively. In contrast, phosphatidylcholine species are highly hydrated. For example,

dioleoylphosphatidylcholine (DOPC) bilayers retain 18.5 wt % water at -12.8 MPa, a value corresponding to 10.2 moles water/mole lipid. In contrast, phosphatidylethanolamine species are poorly hydrated; with 1-palmitoyl-2-oleoylphosphatidylcholine (POPE) retaining 3.6 wt % water (1.5 moles of water/mole lipid at -12.8 MPa). The hydration characteristics of phospholipids are primarily determined by the headgroup composition, and are not significantly affected by the fatty acid component for lipids in the same phase (Lynch and Steponkus, unpubl.; Webb and Steponkus, unpubl.). As a first approximation, the differences in the hydration characteristics of the plasma membrane lipids of non-acclimated and cold-acclimated leaves reflect differences in the proportions of the poorly hydrated and highly hydrated species. For example, during cold acclimation, the levels of cerebrosides decrease from 16 mol% to 6 mol% and phospholipids increase from 32 mol% to 42 mol% (Lynch and Steponkus, 1987). However, there are some exceptions in that the proportion of free sterols (a poorly hydrated species) also increases following cold acclimation, but this is offset by decreases in the sterylglucosides and acylated sterylglucosides, which are also poorly hydrated species. Thus, the net change in the hydration characteristics of the total lipid fraction appears to be a consequence of the decrease in cerebrosides and the increase in phospholipids, especially phosphatidylcholine.

Subsequent studies (Webb & Steponkus, 1990) have been directed to the hydration characteristics of lipid mixtures characteristic of the plasma membrane. Desorption of aqueous dispersions of DOPE, DOPC, and their 3:1, 1:1, and 1:3 (mol:mol) mixtures above saturated salt solutions at 20°C resulted in a linear decrease in the water content of the lipids with the ln of the vapor pressure. DOPC is significantly more hydrated than DOPE at equivalent vapor pressures; at -13 MPa, DOPC had a water content of 18.5 wt% compared to 6.7 wt% for DOPE. The vapor pressure at which the lipid bilayers reached zero water content, P_0 , was -380 MPa for DOPC compared to only -65 MPa for DOPE. The water content of DOPE:DOPC mixtures in 1:3, 1:1, and 3:1 molar proportions increases non-linearly with increasing proportion of DOPC. At -13 MPa, the water content of dispersions containing 0, 25, 50, 75, and 100 mol% DOPC was 6.7, 12.5, 13.8, 15.5, and 18.5 wt% respectively.

We have extended these studies to more complex ternary and quaternary lipid mixtures whose compositions more closely resemble those of the acclimated and non-acclimated rye plasma membranes (Webb & Steponkus, 1991b). Specifically, using the DOPE:DOPC 1:1 mixture as a starting point we have examined the effect of added plant sterols and rye cerebrosides on bilayer hydration. The addition of plant sterols, primarily β -sitosterol and campesterol, to the DOPE:DOPC 1:1 dispersion at either 33 or 50 mol% significantly reduced bilayer hydration from 15.0 wt% to 13.1 wt% and 11.1 wt% water, respectively, at -10 MPa. Our results are consistent with the low hydration of sterols described above and that of cholesterol as measured by others. Addition of small amounts of rye plasma membrane cerebrosides (CMH) to the 1:1:2 DOPE:DOPC:ST dispersion generates a mixture whose composition models that of the acclimated rye plasma membrane: DOPE:DOPC:ST:CMH 22:22:50:5. This dispersion showed a slight reduction in water content compared to the 1:1:2 mixture, having 10.7 wt% water at -10 MPa. A further increase in CMH levels to 20 mol%, and concomitant decrease in the total phospholipid levels to 30 mol% (DOPE:DOPC:ST:CMH 15:15:50:20), yielded a dispersion whose composition mimics that of the non-acclimated rye plasma membrane. This mixture showed a significant reduction in water content compared to that of the acclimated plasma membrane model, having 6.8 wt% water at -10 MPa. Reduced hydration of these dispersion containing cerebrosides is consistent with the low hydration of cerebrosides. These data suggest that the hydration of plasma membrane lipid extracts of the composition previously reported (Lynch and Steponkus, 1987) should be significantly different. It is expected that the lipids from the plasma membrane of acclimated rye plants should have significantly higher water contents than those from non-acclimated rye plants when dehydrated at equivalent vapor pressures.

D. Lyotropic Phase Behavior of Plasma Membrane Lipids

A major goal of this project is to provide a mechanistic understanding of dehydration-induced lamellar-to-hexagonal_{II} phase transitions in NA protoplasts and to determine the factors that preclude H_{II} formation in ACC protoplasts. Given that the differential propensity for H_{II} formation is, in large part, a consequence of alterations in the lipid composition of the plasma membrane, a major effort has been directed to determination of the lyotropic phase behavior of the plasma membrane lipids (individual lipid classes/species, simple mixtures, and complex mixtures).

Individual Lipids. The lyotropic phase behavior of the major species of phosphatidylcholine has been determined (Lynch & Steponkus, 1989c). This was of special interest because it has been suggested by others that in mixtures containing PC and PE, lyotropic L _{α} - L _{β} phase transitions in PC during dehydration could result in the enrichment of unsaturated PE to levels sufficient to facilitate the lamellar-to-hexagonal II (L _{α} -to-H_{II}) phase transition. This suggestion is based on the fact that dehydration increases the L _{α} -to-L _{β} phase transition temperature (T_m) of PC

species. Although the T_m of the major unsaturated species of PC in the plasma membrane increased by 30 to 40°C over the range of water contents between 20 and 5 wt%, examination of the desorption isotherms for these species revealed that at -12 MPa (the vapor pressure of ice at -10°C, which is the temperature at which H_{II} phase occurs in NA protoplasts), the T_m s for 16:0/18:2 PC, 18:2/18:2 PC, and 18:3/18:3 PC were -16°C, -38°C, and -47°C, respectively. Therefore, it is unlikely that a lyotropic L_α -to- L_β phase separation in the PC component of the NA plasma membrane contributes to the formation of the H_{II} phase at -10°C.

Calorimetric and x-ray diffraction studies of the phase behavior of fully hydrated and dry rye glucocerebrosides have been completed (Lynch *et al.*, 1991). The phase behavior of the rye cerebrosides are significantly different from bovine brain cerebrosides. On initial heating, aqueous dispersions of rye cerebrosides show a high enthalpy transition at 56°C, which X-ray diffraction shows is the transition from a highly ordered lamellar crystalline phase (L_β) to a liquid crystalline phase (L_α), with a limiting repeat spacing of 5.5 nm. Subsequent cooling, below the T_m value the cerebrosides exhibit a series of irreversible transitions and gel phase metastability, depending on the thermal history. In the dry state, rye cerebrosides undergo an initial endothermic transition at 130°C to an H_{II} phase.

Binary Lipid Mixtures. Examination of a variety of simple binary lipid mixtures, of compositions relevant to the rye plasma membrane, have been performed by DSC, desorption, freeze-fracture electron microscopy (FFEM), and X-ray diffraction. Mixtures of rye cerebrosides (CMH) and POPC in excess water demonstrate complex thermal behavior (Lynch & Steponkus, 1987c). Mixtures containing >30 mol% CMH exhibit nonideal mixing and gel phase immiscibility at $\geq 2^\circ\text{C}$, resulting in the co-existence of CMH-rich and POPC-rich domains. Similarly, CMH and DOPE appear to be immiscible in the gel phase and exhibit limited miscibility in the L_α phase. Nevertheless, the addition of CMH increased the T_{bh} of the PE regardless if the cerebroside was in the L_α or L_β phase. The influence of sterol glucosides on the calorimetric-determined phase behavior with various molecular species of PC has been investigated (Lynch & Steponkus, 1988). The miscibility of the sterol glucosides with the various PC molecular species decreased in the order: POPC>DPPC>DOPC. Addition of SG to PC resulted in a decrease in both the T_m and the transition enthalpy with increasing mol% SG in the mixture. This behavior was different from that of cholesterol:PC dispersions and suggests that the glucosylation of the β -hydroxyl group of the sterol alters the interactions of the sterol with phospholipids.

Analysis of the effect of dehydration on the phase behavior of binary DOPE:DOPC dispersions has been performed by FFEM, DSC, and x-ray diffraction (Webb & Steponkus, 1990, 1991a). For pure PC systems, the temperature of the main gel-liquid crystalline phase transition (T_m), is known to increase with decreasing lipid hydration. For DOPC, the measured T_m was -18°C at ≥ 17 wt% water and increased to 1°C at 1 wt% water. For pure DOPE, the T_m value of -6°C was observed at ≥ 12 wt% water and this value increased to -2°C at 3 wt% water. For mixtures of DOPE:DOPC the lyotropic increases of T_m occurred when the water content was below 13-15 wt%. For all of these mixtures, the DSC data indicated that the temperature of the main gel-liquid crystalline phase transition was $\leq 5^\circ\text{C}$. Therefore, all lamellar phases observed in samples desorbed at 20°C represent the liquid crystalline lamellar phase (L_α). This conclusion was experimentally confirmed by wide-angle X-ray diffraction of a DOPE:DOPC 1:1 dispersion dehydrated to water contents between 6 wt% and 40 wt%. For all hydration levels, a diffuse Bragg reflection centered at a spacing of 0.46 nm, characteristic of disordered acyl chains in the L_α phase, was observed and the sharp Bragg reflection at 0.42 nm, characteristic of acyl chains in the gel phase, did not occur. FFEM studies of desorbed DOPE:DOPC dispersions showed that the lyotropic L_α - H_{II} phase transition in these lipid mixtures was dependent on both the lipid composition of the dispersion and on the water content of the system. Construction of a phase diagram from these data indicates that the tendency of a mixed lipid system to undergo a lyotropic transition to the H_{II} phase is dependent on both lipid composition of the dispersion and on the water content of the system. The latter variable is a function of both lipid composition (headgroup hydration) and vapor pressure. The H_{II} phase was favored by higher proportions of DOPE and/or lower water contents.

We have performed experiments designed to determine if DOPE and DOPC might undergo fluid-fluid separation as a result of their different hydration characteristics when under dehydration stress (Webb & Steponkus, 1991a). Such separation has been predicted to occur during vesicle-vesicle close approach (Rand, 1981) and in dehydrated lamellae (Bryant and Wolfe, 1989). Aqueous dispersions of DOPE:DOPC 1:1, at 85 wt% water, were loaded into X-ray capillary tubes and dehydrated to water contents between 6 wt% and 40 wt%. At 40 wt% and 25 wt% water, single fluid lamellar repeat spacings of 6.20 nm and 5.01 nm were observed. At water contents of 18 wt% to 6 wt%, the diffractions resolve into two separate fluid lamellar spacings. At 18 wt% water two different lamellar repeat spacing were observed at 4.45 nm and 4.12 nm. At 11 wt% water, two lamellar spacing of 4.44 nm and 3.55 nm occurred and

at 6 wt% water, the observed spacings were 4.29 nm and 4.00 nm. This hydration-dependent appearance of two separate fluid lamellar repeat spacings is not observed in either pure DOPC or pure egg PE at 25°C over the same water content range. These data show that the lyotropic L_α -to- H_{II} transition in this PE:PC system occurred without an L_α -to- L_β phase separation of the PC species as a mechanism of enriching PE species to levels sufficient to form the H_{II} phase. Rather these studies suggest that lipids of very different hydration characteristics may separate in the fluid phase during dehydration stress and that this demixing may facilitate the transition to the H_{II} phase (Bryant and Wolfe, 1989). Experiments are in progress to determine if such separations occur in more complex lipid mixtures and if they may contribute to H_{II} phase formation in the freeze-dehydrated plasma membrane lipids.

4. Phase Behavior of Complex Lipid Mixtures. The phases adopted by phospholipid systems are markedly affected by the presence of other non-phospholipid lipid components (*i.e.* sterols) present in membranes. Therefore, analysis of the phase behavior of the mixtures described above has been performed by DSC and freeze-fracture electron microscopy. Desorption of DOPE:DOPC 1:1 results in a lyotropic increase in T_m of 10°C over the range of water contents from 85 to 3 wt% H_2O . Addition of plant sterols to levels of 33 mol% (DOPE:DOPC:ST, 1:1:1) or 50 mol% (DOPE:DOPC:ST, 1:1:2) reduced the increase in the T_m to 4° and 2°C, respectively. Thus, the presence of plant sterols at levels similar to those occurring in the rye plasma membrane (33 mol% in non-acclimated, 44 mol% in acclimated plasma membranes; Lynch and Steponkus, 1987) precludes the lyotropic increase in T_m observed in pure phospholipid systems. Because cholesterol is thought to act as a "lateral spacer" between phospholipids, it is likely that the increased phospholipid acyl packing order and decreased molecular area, which occur during dehydration and promote the L_β phase, are prevented by the presence of sterols. It should also be noted that the transition enthalpy is also reduced at decreased water contents for all of these mixtures. As a result of the lowered transition enthalpies, an L_α - L_β phase transition would be more easily affected by other factors such as solute and pH changes. For mixtures containing cerebroside, the observations described above also occurred. In DOPE:DOPC:ST:CMH 22:22:50:5, the mixture modelled after the acclimated plasma membrane composition, the T_m showed no lyotropic increase with decreasing water contents: observed T_m values were -14°C to -15°C between 2 wt% water and 85 wt% water. No other transitions, indicative of phase separation of the high-melting point cerebroside (Lynch *et al.*, 1991), were observed up to 80°C. The transition enthalpy for this mixture also decreased with decreasing hydration. Similar results were obtained for the mixture that represented the lipid composition of non-acclimated plasma membranes, DOPE:DOPC:ST:CMH, 15:15:50:20. Transition temperatures for this mixture were stable at -18°C at all hydrations between 4 wt% and 85 wt% water. No other phase separations were observed between -50°C and 80°C. Therefore, the lyotropic increase of T_m observed for pure phospholipid systems is precluded in complex lipid mixtures containing sterols.

Determination of the tendency of these dispersions to adopt the H_{II} phase upon dehydration was performed by FFEM. It was expected that addition of plant sterols would promote the lyotropic L_α -to- H_{II} transition as a result of increased lateral packing pressures in the acyl region of the bilayers coupled with a negligible increase in packing pressures in the headgroup domain as well as the effect of cholesterol on decreasing the overall bilayer hydration. A lowering of the thermotropic T_{bh} value by the addition of cholesterol to fully hydrated DOPE:DOPC bilayers has been observed by others. As outlined in the previous section, the DOPE:DOPC 1:1 mixture was observed to be exclusively in the L_α phase at water contents ≥ 8.8 wt%. Both the L_α and H_{II} phases co-existed at 5.7 wt% and 5.0 wt% water and only the H_{II} phase was present at water contents ≤ 2.6 wt%. With the addition of sterols to 50 mol% (DOPE:DOPC:ST, 1:1:2), desorption at -13 MPa (10.3 wt% water) yielded a dispersion exclusively in the lamellar phase in the FFEM studies, which was determined to be the L_α phase based on DSC studies. Further desorption at -76 MPa (4.5 wt% water) resulted in the co-existence of both lamellar (L_α) and H_{II} phases. With the addition of rye cerebroside to 5 mol% in this mixture, the acclimated rye plasma membrane model mixture DOPE:DOPC:ST:CMH 22:22:50:5 desorbed at -13 MPa (9.8 wt% water) was present only as the lamellar (L_α) phase lipid. Further desorption of this lipid mixture at -39 MPa (5.9 wt% water) induced the formation of co-existing lamellar and H_{II} phases. In lipid dispersions of a composition representative of the non-acclimated rye plasma membrane and showing elevated cerebroside levels (DOPE:DOPC:ST:CMH, 15:15:50:20), desorption at -3 MPa (9.6 wt% water) yielded a dispersion present only in the lamellar (L_α) phase. Desorption of this lipid mixture at -13 MPa (6.3 wt% water) gave a dispersion predominantly in the lamellar phase, but also with significant amounts of lipid in the H_{II} phase. Further dehydration at -39 MPa (3.9 wt% water) gave a dispersion with high levels of both lamellar and H_{II} phase.

From these preliminary results it appears that the tendency of the complex lipid mixtures to enter the H_{II} as a function of water content is very similar. That is, for all of these mixtures at approximately 9 to 10 wt% water (9.6 wt% for

1:1, 10.3 wt% for 1:1:2, 9.8 wt% for 22:22:50:5, and 9.6 wt% for 15:15:50:5) only the lamellar phase was observed. However, when all of these mixtures approach water contents near 6 wt% (5.7 wt% for 1:1, 4.5 wt% for 1:1:2, 5.9 wt% for 22:22:50:5, and 6.3 wt% for 15:15:50:20) the first observations of the H_{II} phase were made. The difference between these samples in their tendency towards the lyotropic H_{II} phase is the vapor pressure required in order to achieve the 6 wt% water at which the H_{II} phase first occurs. For 1:1 DOPE:DOPC and 1:1:2 DOPE:DOPC:ST this pressure was -76 MPa, for the DOPE:DOPC:ST:CMH 22:22:50:5 this water content was attained at -39 MPa, and for the DOPE:DOPC:ST:CMH 15:15:50:20, it was attained at -13 MPa. We interpret these results to suggest that sterols and cerebroside do not contribute to H_{II} phase formation through effects on lateral packing pressures as the H_{II} phase was not observed to occur at higher water contents in the presence of these components. However, both sterols and cerebroside appear to promote the lyotropic L_{α} -to- H_{II} transition by reducing bilayer hydration such that the vapor pressure required for H_{II} phase formation is less extreme. Therefore, the lower hydration of NA plasma membrane lipids should facilitate the L_{α} -to- H_{II} transition by requiring relatively minor dehydrations to reach the critical water contents. Conversely, the higher hydration of ACC plasma membrane lipids would act to prevent or preclude H_{II} phase formation by requiring more severe dehydration to reach the water content for H_{II} formation.

Plasma Membrane Vesicles. In previous studies, it was demonstrated that the differential propensity for H_{II} formation in MLVs formed from the total plasma membrane lipid extract (Cudd & Steponkus, 1988). Studies are now in progress to determine the temperature dependence of this behavior. Dispersions of both the acclimated and non-acclimated rye plasma membrane lipids have been desorbed and examined by freeze-fracture electron microscopy for their tendency to adopt the H_{II} phase during dehydration. Non-acclimated rye plasma membrane lipid dispersions desorbed at -3 MPa and 20°C were present only in the lamellar phase. When desorbed at -13 MPa the non-acclimated lipids phase separated into both lamellar and H_{II} phases. At more extreme dehydrations (-39 MPa) the non-acclimated plasma membrane lipids were predominantly in the H_{II} phase. Dehydration of the acclimated rye plasma membrane lipids gave results very similar to those obtained for the non-acclimated plasma membrane lipids. At -3 MPa only the lamellar phase was observed, with no evidence for non-lamellar structures nor regions of bilayer-bilayer interactions. As was observed for the non-acclimated plasma membrane lipids, the acclimated lipids dehydrated at -13 MPa were in a mixture of both lamellar and H_{II} phases. It is likely that, for the lipids from both non-acclimated and cold-acclimated tissues, the desorption temperature of 20°C is above the T_{bh} value for the lipids. Under these conditions, it is not surprising that we observe no difference between the different plasma membrane lipids extracts in their propensity to undergo the L_{α} -to- H_{II} phase transition; studies at lower temperatures are in progress.

D. Determination of Freezable/Unfreezable Water Associated With Lipids

Recently, Crowe *et al.* (1990) suggested that freezing and dehydration are fundamentally different stress vectors. This suggestion was, in part, based on the assumption that the water associated with the polar headgroups of phospholipids is unfreezable (*i.e.*, 0.25 g H_2O /g phosphatidylcholine) as determined from differential scanning calorimetric (DSC) studies by Chapman *et al.* (1967). In contrast to this interpretation, we submit that a portion of the water associated with the headgroup of phosphatidylcholine is freezable, but may remain unfrozen depending on the thermal history of the sample. To experimentally verify this contention, DSC studies were conducted with multilamellar liposomes that were dehydrated over a saturated solution of NaH_2PO_4 at 20°C, which yielded a water content of 0.22 g H_2O /g of DPPC. To initiate ice formation, the samples first were cooled to -80°C at 100°C/min. With both dehydrated samples and samples that contained excess water, ice nucleation occurred at approximately -20°C and -45°C, with the first and the second nucleation events attributed to heterogeneous and homogeneous ice nucleation, respectively. Subsequently, the samples were warmed at a rate of 100°C/min to either -30 or -40°C for an isothermal storage period of 0 to 60 min. The samples were then cooled at 100°C/min to -80°C prior to the calorimetric measurements during warming from -80 to 30°C at 3°C/min. To provide an accurate estimate of the amount of water that was frozen, corrections were made for the decrease in the heat of fusion of ice at subzero temperatures and for the shift in the base line that occurs during ice melting because of the difference in the specific heats of ice and water. With this protocol, only a portion of freezable water actually crystallized during initial cooling to -80°C at 100°C/min because of the amount of water that crystallized increased during the first 10 and 20 min of storage at -30°C and -40°C, respectively. In samples that were maintained at -30°C for 30 min, 0.11 ± 0.02 and 0.15 ± 0.01 g of water per 1 g of lipid remained unfrozen for the dehydrated suspension and the suspension with excess of water, respectively. The somewhat higher value of unfrozen water obtained for suspensions in excess water is, in part, attributable to a greater total amount of unfrozen water remaining between ice crystals. The dependence of the amount of unfrozen water on subzero temperature was calculated from the partial area of the endotherm as a function of the temperature. The values obtained for the dehydrated samples were in agreement with the desorption isotherm for DPPC equilibrated

over saturated salt solutions; the values obtained for samples in excess water were somewhat higher. These studies demonstrate that a portion of the water associated with the polar headgroups of phosphatidylcholine is freezable; whether or not it freezes is dependent on the thermal history of the sample.

F. Influence of Cryoprotectants on Lamellar-to-Hexagonal_H Phase Transitions

Because both conventional cryopreservation procedures and vitrification procedures result in extreme cell dehydration, we have extended our studies of dehydration-induced H_H formation to include considerations that are relevant to cryopreservation of plant germplasm. Although vitrification procedures offer considerable promise for the cryopreservation of a diverse range of biological materials, a major impediment is the apparent toxicity of the vitrification solutions. As a result, there is much interest in formulating vitrification solutions that are less toxic. However, most studies have focused on characterization of the physical properties of the vitrification solutions with relatively little attention given to the biological aspects of toxicity. It is commonly assumed that toxicity is a consequence of "biochemical" rather than "osmotic" effects, with osmotic effects often inferred to be restricted to volumetric changes that occur during osmotic excursions. However, a major cause of toxicity of vitrification solutions is attributable to the large osmotic stress that is imposed during the dehydration step (Langis & Steponkus, 1991). For example, with NA protoplasts, dehydration in sorbitol solutions of 5.34 osmolal or greater results in the formation of aparticle domains and lamellar-to-hexagonal_H phase transitions in the plasma membrane and the subtending lamellae. With more concentrated solutions of sorbitol (13 osmolal) these changes occur after very short periods of time: aparticle domains were apparent in 50% of the plasma membrane fracture faces within one minute and the hexagonal_H phase appears after two minutes (Steponkus & Fujikawa, 1991). When dehydration is effected in a vitrification solution composed of 24 wt% ethylene glycol + 30 wt% sorbitol + 6 wt% bovine serum albumin (18 osmolal), aparticle domains were apparent in 65% of the plasma membrane fracture faces after just 30 seconds, but hexagonal_H phase was observed in only 10% of the fracture faces after three minutes. These differences are attributable to the presence of ethylene glycol in the vitrification solution because loading the protoplasts with ethylene glycol prior to dehydration in a 13 osmolal sorbitol solution also ameliorates these changes in membrane ultrastructure: the frequency and size of the aparticle domains is diminished and hexagonal_H phase is not observed. When loaded with ethylene glycol and dehydrated in the vitrification solution, aparticle domains were observed in only 10% of the protoplasts after a 3-minute incubation period. Prolonged exposure (5 minutes or longer) to the vitrification solution resulted in the blistering, vesiculation, and dissolution of the plasma membrane. These changes were not observed in protoplasts dehydrated in 13 osmolal sorbitol solutions for up to 30 minutes. In conventional cryopreservation procedures (freeze-induced dehydration), equilibration in DMSO prior to freezing decreases the temperature at which aparticle domains and H_H phase are first observed (Pihakaski & Steponkus, 1987).

G. Osmotic Adjustment in Rye

During cold acclimation of Puma rye, the intracellular osmolality increases from 0.53 to 1.03, which corresponds to an decrease in the solute potential from -12 MPa to -23 MPa. The increase in intracellular osmolality is progressive during the four-week acclimation period and parallels the progressive increase in freezing tolerance from -5°C to -25°C. The increased concentration of intracellular solutes may contribute to the increased freezing tolerance in two ways: i) a lower intracellular solute potential will decrease the extent of cellular dehydration at subfreezing temperatures and ii) various solutes (e.g., soluble sugars, glycine betaine, and proline) minimize membrane destabilization during dehydration. Therefore, studies of osmotic adjustment during cold acclimation were initiated to complement our analysis of alterations in the lipid composition of the plasma membrane (Koster *et al.*, 1989). The total soluble sugar content more than doubled by the second week of cold acclimation (the amount increases from 11 to 26 mg/g FW). Most of this increase was attributed to the accumulation of sucrose and raffinose, which increased from 2.4 and 0 mg/g FW to 11 and 5 mg/g FW, respectively. The monosaccharides (predominantly glucose) remained at a fairly constant level (approx. 10 mg/g FW) throughout the course of acclimation. The level of glycine betaine increased from 2.45 to 11 µmol/g FW following cold acclimation. Proline accumulation began during the third week of cold acclimation. Unlike the soluble sugars, proline accumulated throughout the cold acclimation period and did not plateau. It would appear that the amounts of proline measured in Puma rye (from 27 to 580 µg/g FW) contribute only slightly to the overall increase in intracellular osmolality.

H. Cryo-SEM Observations of *in situ* Ice Formation in Rye Leaves

Although it is universally acknowledged that ice formation occurs in the extracellular spaces of plant tissues, there are remarkably few published studies documenting the location and morphology of the ice crystals and the morphology

of the cells in the frozen state. In collaboration with N.R. Read at the University of Edinburgh, preliminary studies using low-temperature scanning electron microscopy (cryo-SEM) were conducted to determine the location and morphology of ice crystals formed in rye leaves frozen *in situ* and their effects on cell morphology (Allen *et al.*, 1990). In unfrozen leaves of winter rye, the mesophyll cells form a complex, porous network of interconnected, thin-walled cells. In leaves frozen at -2.5°C , extracellular ice does not encapsulate the cells. Rather, large, discrete and often angular extracellular ice crystals are formed within the intercellular air spaces, and only a small proportion of the surface of the individual cells are in direct contact with the ice crystals. Nevertheless, the cells are substantially dehydrated. Because only a small proportion of the surface area of the cells is in contact with the ice crystals, cell dehydration occurs by vapor phase equilibration with the ice crystals. The diffusion of water from the surface of the cells to the surface of the remote ice crystals is a function of the gradient in the concentration of the water phase at each interface. Although diffusion in the gas phase is more rapid than in the liquid phase, the distance between the cell surface and the surface of the ice crystals is considerably greater than when the cells are encapsulated in ice. As a result, the diffusion of water from the cell surface to the ice crystals can be two to three orders of magnitude slower than that which would occur if the cells were in a partially frozen aqueous medium at the same temperature. This large difference in water diffusion would, in large part, account for the large differences in the critical cooling rate necessary to preclude intracellular ice formation in isolated protoplasts suspended in an aqueous medium ($3^{\circ}\text{C}/\text{min}$) in comparison to that which is required to prevent intracellular ice formation in intact leaves ($\sim 3^{\circ}\text{C}/\text{hr}$). Thus, the porosity of the leaf tissue and the characteristic pattern of ice crystal formation will greatly affect the rate of cell dehydration and hence the critical cooling rate required to preclude intracellular ice formation.

I. Dehydration-Induced Alterations in the Plasma Membrane of Protoplasts of Cold-Acclimated Rye Leaves

The primary focus of this project has been on freeze-induced lamellar-to- H_{II} phase transitions in the plasma membrane, which is the principal cause of injury NA protoplasts frozen to temperatures below -5°C . This form of injury is precluded after the first week of cold acclimation. Recently, we initiated studies to characterize the manifestations and cause(s) of injury in protoplasts isolated from rye leaves that are in intermediate stages of acclimation and in protoplasts isolated from fully acclimated leaves to determine what factors limit the maximum freezing tolerance of winter rye. These studies have resulted in identification of the fracture-jump lesion as the principal freeze-induced ultrastructural modification in the plasma membrane of ACC protoplasts and established that its occurrence closely coincides with injury of protoplasts isolated from rye leaves at both intermediate and maximum levels of freezing tolerance (Fujikawa and Steponkus, 1990). The results of these studies are presented in the Background section of the renewal proposal as they serve as the foundation for our future studies.

PUBLICATIONS:

- Allan, W.T.G., N.D. Read, C.E. Jeffree and P.L. Steponkus. 1990. Cryo-scanning electron microscopy of ice formation in rye leaves. *Cryobiology* 27:664. (Abstract)
- Bronshteyn, V.L. and P.L. Steponkus. 1991. Unfrozen versus unfreezable water in phospholipid vesicles. *Cryobiology* 28:in press. (Abstract; manuscript in preparation).
- Bryant, G. and J. Wolfe. 1989. Can hydration forces induce lateral phase separations in lamellar phases? *Eur. J. Biophys.* 16:369-372.
- Cudd, A. and P.L. Steponkus. 1987. Osmotic dehydration-induced lamellar to hexagonal $_{\text{II}}$ phase transitions in liposomes of rye plasma membrane lipids. *Cryobiology* 24:556-557. (Abstract)
- Cudd, A. and P.L. Steponkus. 1988. Lamellar-to-hexagonal $_{\text{II}}$ phase transitions in liposomes of rye plasma membrane lipids after osmotic dehydration. *Biochim. Biophys. Acta* 941:278-286.
- Dowgert, M.F., J. Wolfe and P.L. Steponkus. 1987. The mechanics of injury to isolated protoplasts following osmotic contraction and expansion. *Plant Physiol.* 83:1001-1007.
- Fujikawa, S. and P.L. Steponkus. 1990. Freeze-induced alterations in the ultrastructure of the plasma membrane of rye protoplasts isolated from cold-acclimated leaves. *Cryobiology* 27:665-666. (Abstract) (Manuscript in preparation for submission to *Planta*).

Hatton, B. and P.L. Steponkus. 1987. Computerized cryomicroscopic video image analysis. *Cryobiology* 24:555. (Abstract)

Koster, K.L., P.L. Steponkus and D.V. Lynch. 1989. Solute accumulation during cold acclimation of rye. *Plant Physiol.* 89:S-26. (Abstract) (Manuscript in preparation)

Langis, R. and P.L. Steponkus. 1991. Vitrification of isolated rye protoplasts: protection against dehydration injury by ethylene glycol. *Cryo-Letters* 12:107-112.

Lynch, D.V. and P.L. Steponkus. 1987. Plasma membrane lipid alterations associated with cold acclimation of winter rye seedlings (*Secale cereale* L. cv Puma). *Plant Physiol.* 83:761-767.

Lynch, D.V. and P.L. Steponkus. 1987. Thermotropic phase behavior of glucocerebroside from rye leaves. *Cryobiology* 24:555-556. (Abstract)

Lynch, D.V. and P.L. Steponkus. 1987. Plasma membrane lipid alterations following cold acclimation: relevance to freezing tolerance. In: *Metabolism, Function and Structure of Plant Lipids*, edited by P.K. Stumpf. pp. 213-215. Plenum Publ. Co., New York.

Lynch, D.V. and P.L. Steponkus. 1988. Thermotropic phase behavior of steryl glucoside-phospholipid mixtures. *Plant Physiol.* 86:S-53. (Abstract)

Lynch, D.V. and P.L. Steponkus. 1988. Lyotropic phase behavior of rye plasma membrane phosphatidylcholine species. *Cryobiology* 25:521-522. (Abstract)

Lynch, D.V. and P.L. Steponkus. 1989. Lyotropic phase behavior of unsaturated phosphatidylcholine species: relevance to the mechanism of plasma membrane destabilization and freezing injury. *Biochim. Biophys. Acta* 984:267-272.

Lynch, D.V. and P.L. Steponkus. 1989. The behavior of rye plasma membrane lipids at low hydrations. *Cryobiology* 26:556. (Abstract)

Lynch, D.V., M. Caffrey, J.L. Hogan and P.L. Steponkus. 1991. Calorimetric and X-ray diffraction studies of rye glucocerebroside mesomorphism. *Biophys.J.* (in press).

Pihakaski, K. and P.L. Steponkus. 1987. Freeze-induced phase transitions in the plasma membrane of isolated protoplasts. *Physiol. Plant.* 69:666-674.

Steponkus, P.L. 1990. Cold acclimation and freezing injury from a perspective of the plasma membrane. In: *Environmental Injury to Plants*, edited by F. Katterman. pp. 1-16. Academic Press, New York.

Steponkus, P.L. 1991. Behavior of the plasma membrane during osmotic excursions. In: *Endocytosis, Exocytosis, and Vesicle Traffic in Plants*, edited by C. Hawes, J. Coleman and D. Evans. pp. (in press). SEB Symp. Series, Cambridge.

Steponkus, P.L. and S. Fujikawa. 1991. Membrane destabilization during vitrification procedures. *Cryobiology* 28:(in press) (Abstract).

Steponkus, P.L. and D.V. Lynch. 1989. Freeze/thaw-induced destabilization of the plasma membrane and the effects of cold acclimation. *J. Bioenerg. Biomembr.* 21:21-41.

Steponkus, P.L. and D.V. Lynch. 1989. The behavior of large unilamellar vesicles of rye plasma membrane lipids during freeze/thaw-induced osmotic excursions. *Cryo-Letters* 10:43-50.

Steponkus, P.L., D.V. Lynch and M. Uemura. 1990. The influence of cold acclimation on the lipid composition and cryobehaviour of the plasma membrane of isolated rye protoplasts. *Philos. Trans. R. Soc. Lond. B* 326:571-583.

- Steponkus, P.L., D.V. Lynch, M. Uemura, R.A. Balsamo and T. Arvinte. 1988. Plant cryobiology: cellular and molecular aspects of freezing injury and cold acclimation. In: *Low Temperature Biotechnology: Emerging Applications and Engineering Contributions*. A.S.M.E. Publication BED-Vol. 10; HTD-Vol. 98, edited by J.J. McGrath and K.R. Diller. pp. 47-56. American Society of Mechanical Engineers, New York.
- Steponkus, P.L. and M. Uemura. 1989. Behavior of the plasma membrane during osmotic excursions: the effect of alterations in the plasma membrane lipid composition. In: *Plant Water Relations and Growth Under Stress*, edited by M. Tazawa, M. Katsumi, Y. Masuda and H. Okamoto. pp. 75-82. MYU KK, Tokyo.
- Steponkus, P.L. and M.S. Webb. 1991. Freeze-induced dehydration and membrane destabilization in plants. In: *Water and Life: Comparative Analysis of Water Relationships at the Organismic, Cellular and Molecular Level*, edited by G. Somero and B. Osmond. pp. (in press). Springer-Verlag, Berlin.
- Sugawara, Y. and P.L. Steponkus. 1990. Effect of cold acclimation and modification of the plasma membrane lipid composition on lamellar-to-hexagonal_{II} phase transitions in rye protoplasts. *Cryobiology* 27:667. (Manuscript in preparation for submission to *Plant Physiology*).
- Uemura, M., R.A. Balsamo and P.L. Steponkus. 1988. Alteration of the freezing tolerance of rye protoplasts by modification of the plasma membrane lipid composition. *Plant Physiol.* 86:S-53. (Abstract)
- Uemura, M., E.B. Cahoon, D.V. Lynch and P.L. Steponkus. 1989. Temporal aspects of alterations in the cryobehavior and lipid composition of the plasma membrane during cold acclimation of winter rye. *Cryobiology* 26:585-586.
- Uemura, M. and P.L. Steponkus. 1987. A comparison of freezing vs. hypertonic stress on the ATPase activity and protein composition of the plasma membrane of winter rye seedlings. *Cryobiology* 24:557-558. (Abstract)
- Uemura, M. and P.L. Steponkus. 1989. Parallel effects of freezing and osmotic stress on the ATPase activity and protein composition of the plasma membrane of winter rye seedlings. *Plant Physiol.* 91:961-969.
- Uemura, M. and P.L. Steponkus. 1989. Effect of cold acclimation on the incidence of two forms of freezing injury in protoplasts isolated from rye leaves. *Plant Physiol.* 91:1131-1137.
- Uemura, M. and P.L. Steponkus. 1989. Alteration of the freezing tolerance of rye protoplasts by selective lipid enrichment of the plasma membrane. *Cryobiology* 26:585. (Abstract) (Manuscript in preparation)
- Webb, M.S. and P.L. Steponkus. 1990. Hydration characteristics of phospholipid mixtures. *Cryobiology* 27:665.
- Webb, M.S. and P.L. Steponkus. 1990. Dehydration-induced hexagonal_{II} phase formation in phospholipid bilayers. *Cryobiology* 27:666-667.
- Webb, M.S. and P.L. Steponkus. 1991a. Dehydration-induced lamellar-to-hexagonal_{II} phase transitions in DOPE:DOPE mixtures. (Final draft completed - manuscript to be submitted to *Biochim. Biophys. Acta*)
- Webb, M.S. and P.L. Steponkus. 1991b. Hydration and phase properties of mixtures of DOPE, DOPE, plant cerebrosides and plant sterols. (Manuscript in preparation for *Biochim. Biophys. Acta*)

Protoplasmic (1) and exoplasmic (2) fracture faces of the plasma membrane of a protoplast isolated from 4-week cold-acclimated leaves of rye and frozen to -25°C before cryo-fixation in liquid propane. Asterisks mark regions where the fracture-jump lesion has revealed aparticulate membranes in close apposition to the plasma membrane. Both micrographs at 40,000x.

END

**DATE
FILMED**

9 / 15 / 92