WSRC-RP--91-1213

DE93 000638

**BIOLOGICAL MONITORING OF
UPPER THREE RUNS CREEK,
SAVANNAH RIVER SITE
AIKEN COUNTY, SOUTH CAROLINA**

MARCH 1990 - JULY 1991

CHEM-NUCLEAR LABORATORY SERVICES, INC.

**REPORT TO: WESTINGHOUSE SAVANNAH RIVER COMPANY
AIKEN COUNTY, SOUTH CAROLINA**

REPORT DATE: NOVEMBER 1991

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

	<u>PAGE</u>
1.0 <u>INTRODUCTION</u>	1
3.0 <u>METHODS</u>	4
3.1 <u>Macroinvertebrate Assessment</u>	4
3.1.1 Sampling	4
3.1.2 Sample Processing	5
3.1.3 Taxonomic Identification	5
3.1.4 Biomass	6
3.1.5 Data Analysis	6
3.2 <u>Water Chemistry Analysis</u>	8
3.2.1 Field Measurements	8
3.2.2 Sampling	9
4.0 <u>RESULTS</u>	14
4.1 <u>Biological Monitoring</u>	14
4.1.1 Richness Parameters	14
4.1.2 Abundance Parameters	15
4.1.3 Pollution Tolerance Parameters	18
4.1.4 Functional Group Analysis	19
4.1.5 Total Biomass	20
4.1.6 Evenness Parameters	20
4.2 <u>Water Chemistry</u>	21
4.2.1 General Parameters	21
4.2.1.1 Total Suspended Solids	21
4.2.1.2 Specific Conductance	21
4.2.1.3 Hardness	22
4.2.1.4 Alkalinity	22
4.2.1.5 Residual Chlorine	22
4.2.2 Metals	22
4.2.2.1 Chromium	22
4.2.2.2 Copper	23
4.2.2.3 Lead	23
4.2.2.4 Manganese	23

4.2.2.5	Mercury	23
4.2.2.6	Sodium	23
4.2.2.7	Uranium	24
4.2.2.8	Zinc	24
4.2.3	Nutrients	24
4.2.3.1	Nitrites	24
4.2.3.2	Nitrates	24
4.2.3.3	Ammonia Nitrogen	25
4.2.3.4	Total Phosphate	25
4.2.3.5	Ortho-Phosphate	25
4.2.4	Organics	26
4.2.4.1	Tributyl Phosphate	26
4.2.4.2	Kerosene	26
4.2.5	Field Parameters	26
4.2.5.1	pH	26
4.2.5.2	Dissolved Oxygen	26
4.2.5.3	Temperature	26
4.3	<u>Chronic Toxicity Testing</u>	27
5.0	<u>SUMMARY AND CONCLUSIONS</u>	80
6.0	<u>REFERENCES</u>	83

LIST OF TABLES

		<u>Page</u>
Table 1	Sampling Dates. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	10
Table 2	Taxonomic References. Qualitative and Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	11
Table 3	Water Chemistry Sample Containers and Sample Volumes. Upper Three Runs Creek Biomonitoring. Aiken County, SC. March 1990-July 1991.	12
Table 4	Methods for Water Chemistry Analysis Upper Three Runs Creek Biomonitoring. Aiken County, SC. March 1990-July 1991.	13
Table 5	Species List. Upper Three Runs Creek Biomonitoring. March 1990-July 1991.	38
Table 6	Summary of Biomonitoring Parameter Results. Quantitative and Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	48
Table 7a	Species Richness. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	49
Table 7b	Species Richness. Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	50
Table 8	Average # of Species per Replicate. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	51
Table 9a	Total # of Organisms per Site. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	52
Table 9b	Total # of Organisms per Site. Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	53

LIST OF TABLES (Con't)

		<u>Page</u>
Table 10a	Mean Density of Organisms per m ² . Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	54
Table 10b	Density by Major Taxonomic Group. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	55
Table 11a	Percent Composition by Major Taxonomic Group. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	58
Table 11b	Percent Composition by Major Taxonomic Group. Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	61
Table 12a	Dominant Species, Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	63
Table 12b	Dominant Species, Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	64
Table 13a	EPT Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	65
Table 13b	EPT Index. Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	66
Table 14a	Hisenhoff Biotic Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	67
Table 14b	Hisenhoff Biotic Index. Qualitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	68
Table 15a	Functional Group Analysis - % by Pop. #. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	69
Table 15b	Functional Group Analysis - % by Biomass. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	71
Table 16	Biomass. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	74

LIST OF TABLES (Con't)

		<u>Page</u>
Table 17a	Shannon Weaver Diversity Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	75
Table 17b	Equitability Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC, March 1990 - July 1991	76
Table 18	Summary of Water Chemistry Data. Upper Three Runs Creek Biomonitoring, Aiken County, SC. March 1990 - July 1991	77
Table 19	Summary of Quarterly Chronic <u>Ceriodaphnia</u> Toxicity Testing. F/H Area ETF Effluent. March 1990 to July 1991.	79

LIST OF FIGURES

		<u>Page</u>
Figure 1	Map of Sampling Sites. Upper Three Runs Creek Blomonitoring. Aiken County, SC. March 1990 - July 1991.	3
Figure 2	Species Richness. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	28
Figure 3	Average # of Species Per Replicate. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	29
Figure 4	Density. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	30
Figure 5	Percent Composition By Major Taxonomic Group By # of Organisms. All Dates Combined. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	31
Figure 6	EPT Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	32
Figure 7	Hilsenhoff Biotic Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	33
Figure 8	Functional Group Analysis - % by Pop. #. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	34
Figure 9	Functional Group Analysis - % By Biomass. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	35
Figure 10	Shannon Weaver Diversity Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	36
Figure 11	Equitability Index. Quantitative Macroinvertebrate Sampling Upper Three Runs Creek, Aiken County, SC.	37

LIST OF APPENDICES

- A. Species Lists by Replicate, Site and Date. Qualitative Macroinvertebrate Sampling. Upper Three Runs Creek, Aiken County, SC. March 1990 - March 1991.**
- B. Species Lists by Replicate, Site and Date. Quantitative Macroinvertebrate Sampling. Upper Three Runs Creek, Aiken County, SC. March 1990 - July 1991.**
- C. Water Chemistry Quarterly Data. Upper Three Runs Creek and F/H Area ETF Effluent. Aiken County, SC. March 1990 - July 1991.**
- D. Statistical Analysis Results. Upper Three Runs Creek Biomonitoring, Aiken County, SC.**
- E. Hilsenhoff Biotic Index Tolerance Values and Functional Feeding Groups**
- F. Taxonomic Identification Modifications.**

1.0 INTRODUCTION

In anticipation of the fall 1988 start up of effluent discharges into Upper Three Runs Creek by the F/H Area Effluent Treatment Facility of the Savannah River Site, Aiken, SC, a two and one half year biological study was initiated in June 1987. Upper Three Runs Creek is an intensively studied fourth order stream known for its high species richness. Designed to assess the potential impact of F/H area effluent on the creek, the study included qualitative and quantitative macroinvertebrate stream surveys at five sites (see map), chronic toxicity testing of the effluent, water chemistry and bioaccumulation analysis.

In a March 1990 study of the potential impact of F/H Area effluent on the macroinvertebrate communities of Upper Three Runs Creek was extended, with reductions in the number of sites to be sampled and in the frequency of water chemistry sampling. This report presents the results of macroinvertebrate stream surveys at three sites, chronic toxicity testing of the effluent and water chemistry analysis of the three stream sites and the effluent from March 1990 to July 1991.

2.0 SAMPLING LOCATIONS

Three locations were sampled during the study period. Included were one upstream location (Site 1) and two downstream locations (Sites 2 and 5). The upstream location was situated on Upper Three Runs Creek adjacent to the Aquatic Ecology Laboratory, 50m upstream from the confluence of Tim's Branch and 60m upstream from Road C and the F/H Area effluent (H-016) discharge point. Site 2 was located on Upper Three Runs Creek 20 m downstream from the discharge and Road C. Site 5 was sited on Upper Three Runs Creek approximately 3 km downstream from the discharge.

Minor habitat differences were noted among the sites, which primarily affected qualitative sampling. Snag habitats were more abundant and depth was greater at Site 5 than at other sites. Site 2 was characterized by a relatively open canopy, shallow depth, and extensive sand banks along the eastern shoreline. Site 1 exhibited a paucity of snag habitat, hard, steeply sloped banks, and was the most shaded by riparian vegetation.

A map of sampling sites is provided in Figure 1.

3.0 METHODS

3.1 Macroinvertebrate Assessment

3.1.1 Sampling

Five replicate multiplate samplers were set in the stream at each site on a quarterly schedule for quantitative sampling as shown in Table 1.

Each sampler was comprised of 14 circular plates with a diameter of 7.5 cm and thickness of 0.3 cm (area: 0.12 M²). The plates were made of masonite board and were separated by masonite board spacers. A metal bolt through the plates and spacers held each assembly together.

Samplers were suspended in the water by rope from floats; each sampler being approximately 25 cm below the water surface. Proceeding from left to right when facing up stream replicates A through E were placed at equal spacing across the stream. Similar flow regimes were selected in determining exact site location.

Samplers were retrieved by swiftly lifting each sampler from the water, placing it in a labelled zip-lock type plastic bag and cutting the support rope. Approximately 200 ml of stream water was placed in each bag as well as a site and replicate label. Bags with samplers were placed in a cooler with ice to chill the samples until processing in the laboratory.

Qualitative sampling was conducted on a semi-annual basis, on the following dates: March 23, 1990, October 30, 1990 and March 28, 1991. Three samples of 500 cm³ were collected at each site including a) snag/vascular plant habitat, b) sand habitat c) detritus/mud/leaf pack habitat. Samples were collected with the use of an aquatic dip net (used for sweeping, bottom sampling, and as a kick net), a U.S. Standard #30 brass sieve, a white collecting pan and hand collecting of various substrates such as moss, bark, and stick/log scrapings. Approximately 15 minutes of sampling effort by a team of two

biologists was expended at each site. Each sample was placed in a 500 ml plastic labelled container, with 80% ethanol for a fixative.

3.1.2 Sample Processing

Multiplate Samplers/Quantitative:

Each sampler and the contents of the sampler bag were placed in a white collecting pan. Samplers were disassembled and each plate and spacer was scraped/brushed to remove all organisms. Subsequently the contents of each collecting pan were rinsed in a U.S. Standard #30 sieve (to remove sediment) and placed back in the pan. All organisms in the pan were then placed in labelled vials of 80% ethanol preservative, using an illuminated magnifier to facilitate sorting.

Qualitative Sampling:

Sorting qualitative samples involved processing each sample container in approximately five 100 cm³ portions until each container was completed. Each portion was placed in a U.S. Standard #30 sieve, rinsed with tap water and transferred to a white collecting pan. Macroinvertebrates were removed from the pan with the aid of an illuminated magnifier and placed in labelled vials of 80% ethanol preservative.

3.1.3 Taxonomic Identification

Each vial of sorted organisms was emptied into a glass petri dish (3" diameter) with 80% ethanol and examined under a WILD M3 stereomicroscope at 16x magnification. Organisms were initially separated by order for most taxa and family for true flies (Diptera). After species identification, organisms collected by multiplate were returned to the labelled vials for biomass determination, except for voucher specimens. One to three organisms of each species collected in the survey were placed in the reference collection, as voucher specimens. Some midges, black flies and annelids were mounted on slides with CMC-10, and examined with a WILD M20 compound microscope (400X Power). A list of references used for identification is given in Table 2.

3.1.4 Biomass

After macroinvertebrate species identification, organisms were generally separated by functional group for each site and placed in labelled vials. Due to the assignment of many taxa to more than one functional group, biomass determination categories included taxa such as orthoclads, tanypodines, chironomini, and Pteronarcys. Values obtained were divided among the appropriate functional groups. All five replicates per site were combined. Each sample for each group at each site was dried at 105°C, weighed and subsequently placed in a weighed crucible at 550°C for ashing (30 minutes in thermolyne furnace). This was followed by 30 minutes in the dessicator for equilibration to room temperature. Subsequently the crucibles were re-weighed, the initial dry weight subtracted and the ash-free dry weights calculated. Weight determinations were made with a Mettler AE200 electronic balance reading to four places (0.0001g).

3.1.5 Data Analysis

A number of methods to evaluate the data obtained in qualitative and quantitative macroinvertebrate assessments have been developed. Several approaches to analyzing the data are used in this report. These approaches concern the richness, density, evenness, pollution tolerance, % composition by major taxonomic group, biomass, functional group analysis and dominant species.

Community richness is the total number of species collected at a site. Pollution by an effluent discharge generally reduces the number of species (Weber, 1973). A mean density of organisms is calculated per M². When depressed, the mean density can reflect the effects of toxic pollutants (Weber, 1973). When elevated, it can suggest nutrient enrichment (Weber, 1973).

Community evenness is based upon the concept that a stream unaffected by pollution will support many species of macroinvertebrates, each represented by a few organisms, whereas a stream impacted by organic pollution will have only a few tolerant species with large numbers of organisms (Weber, 1973). Organisms intolerant to organic pollution are

killed or drift downstream to a more tolerable habitat. With toxic wastes, a similar phenomenon occurs, except that the reduction in the numbers of intolerant species occurs with a decrease in the species population (Weber, 1973). Two indices involving evenness are used; diversity and equitability. Both are computed using the Shannon-Weaver functions and the Margalef measure, as described by Weber 1973). Species diversity is an index taking into account the number of species and the distribution of individual organisms within a sampled segment of waterway. In severely polluted water, diversity usually is less than one (Weber, 1973). In unpolluted water, it is higher and may attain four or more. The higher the diversity the less stressful the aquatic environment to the organisms contained therein (Weber, 1973). Equitability is an index that compares the aquatic community under consideration with an idealized community. Such a community is found in unpolluted habitats where many species are present, few are relatively abundant in individuals, and several are represented within the community by only a few individuals. Equitability has been found to be a more sensitive test of environmental stress than diversity (Weber, 1973). Levels above 0.5 are indicative of good water quality (Weber, 1973). These indices are generally only applicable to quantitative samples of more than 100 organisms.

Community pollution tolerance evaluates the macroinvertebrate community based upon the concept that certain taxa of aquatic organisms are considered to be more pollution intolerant than others. Intolerant taxa include most species of mayflies, caddisflies and stoneflies, as well as, certain species of beetles (e.g., some elmids), midges (e.g., Potthastia) and Megaloptera (Hilsenhoff, 1987). Other taxa are more tolerant of pollution, including libellulid dragonfly nymphs, true bugs and fly larvae (including many species of midges) (Hilsenhoff, 1987). Two ways of measuring pollution tolerance are provided here; the Hilsenhoff Biotic Index and EPT index. A biotic index gives an indication of the average pollution tolerance of organisms present at a stream site. Each species is given a tolerance rating between 0 and 10, with 0 indicating a very intolerant organism and 10 an organism with high tolerance of pollution. The biotic index used here was produced by Hilsenhoff (1987) and designed for northern streams. It has been modified slightly for several midge species to account for known tolerances in this region. The EPT index is

simply the total number of species at a site which are in the pollution intolerant orders Ephemeroptera (mayflies), Plecoptera (stoneflies) or Trichoptera (caddisflies).

Functional Group Analysis involves classifying each species by feeding mode, determining the percentage of the total number of organisms in each feeding mode, and the biomass of the organisms in each feeding mode. Species are assigned to a functional group based upon information in Merritt & Cummins (1984).

Functional groups include shredders (herbivores and detritivores), scrapers, predators, piercer-herbivores and collectors (filterers and gatherers). In as much as piercer-herbivores never comprised $\geq 1\%$ of the total number of biomass of organisms at any site/date combination, this category is ignored in discussion of functional groups. A shift in the proportions of the functional groups between sites may reflect the presence of a pollution source or changes in the trophic structure of the ecosystem. It must however, be carefully analyzed as a difference in habitat can also produce a shift.

Results of each biological monitoring parameter are discussed separately, then summarized. As might be expected for these types of data, the data sets proved not to be normally distributed despite a square root transformation of the data $\sqrt{y + 1/2}$. As a result, the data were analyzed by non-parametric methods for paired data, using a significance level of $p=0.05$. Comparisons of data means among sites were accomplished using Friedman's Test (Gilbert, 1987). For the percent composition by major taxonomic group and functional group analysis, each group was separately subjected to Friedman's Test. Results of statistical analyses are provided in Appendix D.

3.2 Water Chemistry Analysis

3.2.1 Field Measurements

At each site measurements were taken of dissolved oxygen (ppm), pH, conductivity, and temperature (°C). Dissolved oxygen was measured with a YSI Model 54ARC D.O. meter with a YSI Model A77003 probe. The meter was recalibrated with the elevation

scale/calibration knob and temperature at each site. A Fisher Model 951 pH/temperature meter with a probe was used to measure pH and temperature. The meter was calibrated, using pH 4 and pH 7 standard buffer solutions, at each site. Temperature readings were made to 0.1°C. Conductivity was analyzed using a YSI Salinity-Conductivity-Temperature meter (Model 33). All readings were made 2-6" below the water surface and recorded in a bound field record book.

3.2.2 Sampling

Samples for chemical analysis were taken at each site as grab samples just below the water surface. Containers and sample volumes are listed in Table 3. After collection, samples were placed in coolers, chilled with ice and returned to the laboratory.

Chemical analyses were conducted according to the procedures in "Methods for Chemical Analysis of Water and Wastes" (U.S. EPA, 1979) and are referenced, along with detection limits and preservatives in Table 4.

TABLE 1
 Sampling Dates
 Quantitative Macroinvertebrate Sampling
 Upper Three Runs Creek
 Aiken County, SC
 March 1990 - July 1991

Quarter	# Reps/Site	# Sites	Sample Retrieval Date
Spring 1990	5	5	April 20, 1990
Summer 1990	5	3	July 26, 1990
Fall 1990	5	3	October 23, 1990
Winter 90/91	5	3	January 15, 1991
Spring 1991	5	3	April 17, 1991
Summer 1991	5	3	July 25, 1991

Samplers were placed in the creek 28-30 days prior to retrieval.

TABLE 2
Taxonomic References Used for Identification of Macroinvertebrates
Quantitative and Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
April 1990 - July 1991

Taxon	Reference No. ¹
Annelida	17
Mollusca	27, 38
Crustacea	7, 27, 15, 45
Hydracarina	27
Collembola	20
Ephemeroptera	2, 4, 20, 18, 21, 24
Odonata	4, 20, 25
Plecoptera	4, 20, 37
Hemiptera	4, 20
Megaloptera/Neuroptera	4
Lepidoptera	4
Trichoptera	4, 20, 30, 36, 44
Coleoptera	1, 4, 20
Diptera	4, 8, 10, 16, 19, 20, 34
Chironomidae	3, 4, 5, 6, 11, 13, 14, 26, 28
	29, 31, 32, 33, 35, 42, 43

¹ See References - Section 6.0.

TABLE 3
 Water Chemistry Sample Containers and Sample Volumes
 Upper Three Runs Creek Biomonitoring
 Aiken County, SC
 April 1990 - July 1991

Sampler Container	Volume	Parameters	Preservative
Glass, amber	250 ml	Res Cl	none
Plastic	1/2 gallon	cond, TSS, Hard, Alk, NO2	none
Plastic	250 ml	O-P04	none
Plastic	500 ml	Metals	HN03
Plastic	500 ml	Dis. Metals	none
Plastic	1/2 gallon	NO3, NH3, T-P04	HN03
Glass, amber	1 liter	Kerosene, TBP	none

TABLE 4
Methods for Water Chemistry Analysis
Upper Three Runs Creek Biomonitoring
Aiken, SC
April 1990 - July 1991

<u>Parameter</u>	<u>EPA Method #</u>	<u>Detection Limit</u>	<u>Preservative</u>
pH	150.1	0.1 Units	On-Site Analysis
Dissolved Oxygen	360.1	0.1 mg/l	On-Site Analysis
Temperature	170.1	1°C	On-Site Analysis
Specific			
Conductance	120.1	10 umhos/cm	None
Total Hardness	130.2	1 mg/l	None
Alkalinity	310.1	1 mg/l	None
Total Suspended			
Solids	160.2	1 mg/l	None
Total Residual			
Chlorine	330.1	0.1 mg/l	None
Nitrate (As N)	352.1	0.1 mg/l	H ₂ SO ₄ to pH <2.0
Nitrate (As N)	354.1	0.02 mg/l	None
Ammonia (As N)	350.3	0.01 mg/l	H ₂ SO ₄ to pH <2.0
Total Phosphorus			
(As P)	365.2	0.01 mg/l	H ₂ SO ₄ to pH <2.0
Ortho Phosphate			
(As P)	365.2	0.01 mg/l	Filter on Site
Chromium	218.2	0.005 mg/l	HNO ₃ to pH <2.0
Copper	220.2	0.005 mg/l	HNO ₃ to pH <2.0
Lead (Furnace)	234.2	0.005 mg/l	HNO ₃ to pH <2.0
Manganese	234.2	0.005 mg/l	HNO ₃ to pH <2.0
Mercury (water)	245.1	0.005 mg/l	HNO ₃ to pH <2.0
Sodium	273.1	0.002 mg/l	HNO ₃ to pH <2.0
Uranium	711 B	0.001 mg/l	HNO ₃ to pH <2.0
Zinc	289.1	0.005 mg/l	HNO ₃ to pH <2.0
* Kerosene	SPE/GC	0.1 mg/l	Qt. Amber Glass with Teflon Lined Lid
* Tributyl Phosphate	SPE/GC	0.05 mg/l	Qt. Amber Glass with Teflon Lid

* Solid Phase Extraction GC Analysis Method as Provided by DuPont SRP.

4.0 RESULTS

4.1 Biological Monitoring

A list of species collected from March 1990 to July 1991 is provided in Table 5. Quantitative sampling yielded a total of 104 species, ranging from 69 species at Site 5 to 77 species at Site 2. Qualitative sampling resulted in a total of 163 species, varying from 104 species at Site 1 to 110 species at Site 2. A summary of the biological monitoring parameter results for each site is found in Table 6.

4.1.1 Richness Parameters

Species Richness

Species richness for quantitative sampling methods ranged varied a minimum of 21 at Site 5 (October 1990) to a maximum of 40 at Site 2 in April 1991 (Table 7a; Figure 2). Mean species richness was similar at all three sites, ranging from 29.8 at Site 1 to 33.2 at Site 2. A Friedman's non-parametric test indicated no statistically significant differences among sites ($p = 0.142$) during the study period (April 1990 to July 1991). Highest species richness values were obtained in April 1991. Combined data from all quarters showed 70 species at Site 1 (Table 7a), 77 species at Site 2 and 69 species at Site 5.

Qualitative sampling yielded higher species richness values than quantitative sampling. The minimum species richness value was 50 (Table 7b), at Site 5 in October 1990. The maximum species richness value was 70, at Site 2 in March 1990. Highest species richness was encountered during spring sampling periods. An exception occurred in March 1991 at Site 2, in which relatively low species richness was noted. The total number of species from all sampling periods combined ranged from 104 species at Site 1 to 110 species at Site 2.

Average Number of Species per Replicate

The mean number of macroinvertebrate species colonizing the multiplate samplers after one month of exposure ranged from 12.8 at site 1 to 15.1 at Site 2 (Table 8). The minimum mean number of species per replicate was 8 species, at Site 5 in October 1990. In April 1991, at Site 2, a maximum mean of 19 species was found. A Friedmans nonparametric test of the data indicated no statistically significant differences among sites ($p = 0.072$). Seasonal variations were noted, with the greatest numbers of species per replicate in the winter and spring and lowest numbers of species in the fall. No temporal trends in mean species richness per replicate were found.

4.1.2 Abundance Parameters

Total Number of Organisms per Site

A total of 3684 organisms were collected on the multiplate samplers at the three sites between March 1990 and July 1991 (Table 9a). Populations colonizing the multiplate samplers were lowest in the summer and fall of 1990 and highest in spring 1990. The mean number of organisms per site for all sampling periods combined ranged from 178.3 at Site 1 to 230.8 at Site 2.

Mean numbers of organisms collected at each site and data combination by qualitative methods were slightly greater than obtained by quantitative methods (Table 9b). As with quantitative methods the greatest mean numbers of organisms were collected at Site 2 (306.3) and the least at Site 1 (201.0). Numbers of organisms collected in March 1991 (643) were substantially lower than in March 1990 (1196). The fewest organisms were collected in the fall.

Density

Macroinvertebrate density associated with the multiplate samples ranged from a minimum of 130 organisms per square meter at Site 5 in October 1990 to 610 organisms per square

meter at Site 2 in January 1991 (Table 10a; Figure 4). Mean density for all sampling periods combined was greatest at Site 2 (369.3 organisms/m²) and lowest at Site 1 (285.3 organisms/m²). No significant differences in density among sites were found (Friedman's Test = $p = 0.072$). Density values were higher during 1991 than in 1990, with the exception of Site 5 during spring sampling. The most substantial increases from 1990 to 1991 occurred in the winter, with a 1990 mean density of 236.7 organisms/m², and a 1991 mean density of 508 organisms/m², and in the summer, with a 1990 mean density of 183 organisms/m² and a 1991 mean density of 389.7 organisms/m². Density in the spring and fall did not vary significantly between 1990 and 1991.

Density/Percent Composition By Major Taxonomic Group

The most abundant organisms colonizing multiplate samplers in Upper Three Runs Creek were midges, with an average density of 127.2/m² (Table 10b) comprising 38.8% of the community. (Table 11a; Figure 5). Highest mean densities of midges were found at Site 2 (146.9/m²; 39.8%), although the differences were not statistically significantly greater than at other sites. These elevations in mean midge densities and percentages at Site 2 were a function of more substantial populations of Chironominae midges (Polypedilum, Rheotanytarsus, Tanytarsus) than at Sites 1 and 5. Midge densities peaked in the winter quarter and fell to a minimum in the summer quarter of both 1990 and 1991.

Caddisflies were second in abundance of the major taxonomic groups collected in quantitative sampling (Table 10b), with a mean density of 58.8/m², 17.9% of the community (Table 11a). Hydropsychid caddisflies were dominant among the Trichoptera. Both downstream sites (Sites 2 and 5) showed lower mean densities of caddisflies (54.9 - 56.0/m²) than the control site 1 (65.3/m²). The difference was not statistically significant (Friedman's Test: $p = 0.956$) and is attributable to a single sampling period (July 1991), in which particularly large numbers of Hydropsyche were collected upstream (Appendix B). Caddisflies were most abundant in the winter quarter and least abundant during the summer.

Stoneflies and beetles were, on average, equally abundant on the multiplate samplers during March 1990 to July 1991. Both exhibited densities of approximately $46/\text{m}^2$, 14-15% of the community (Tables 10b; 11a). Mean densities of stoneflies were lower at Sites 1 and 2 (40 to $43.7/\text{m}^2$) than at Site 5 ($54.7/\text{m}^2$). Differences among sites were not statistically significant (Friedman's Test: $p = 0.74$). The elevated stonefly densities at Site 5 occurred only during the spring, when stonefly populations on the multiplate samplers reached a peak. Beetles (primarily adult Elmidae) exhibited marked variability among seasons years and sites. During the winter beetles were largely absent from the multiplate samplers. Conversely, beetle densities were highest during the summer. Densities of beetles were substantially increased during the spring and summer 1991 sampling periods as compared to the spring and summer of 1990. Site 2 supported statistically significant elevated mean densities of beetles in relation to Sites 1 and 5. (Friedman's Test: $p = 0.029$).

Mayflies were substantially less abundant than midges, caddisflies, stoneflies or beetles, with a mean density of $19.9/\text{m}^2$, 6.26% of the community (Table 10b, 11a). Mean densities were highest in the winter and lowest in the fall. During most sampling periods the density of mayflies at Site 2 (mean $26.1/\text{m}^2$) was greater than at Site 1 (mean $14.7/\text{m}^2$). This difference was not statistically significant (Friedman's Test: $p = 0.252$). The inverse relationship of mayfly density and (elmid) beetle density, coupled with the similar trophic role in the community (both are scrapers and collector-gatherers) suggests the two groups fill the same niche at offset times of year. Due to the larger size of mayflies, the system cannot support as great a density of these nymphs as the density of beetles.

Density of non-Chironomid dipterans (mainly Simulium) varied from a mean of $13.9/\text{m}^2$ in July 1990 to $37.3/\text{m}^2$ in April 1991. Mean densities were highest in the spring. No statistically significant differences among sites were noted.

Qualitative sampling indicated a macroinvertebrate community structure similar to that reflected by quantitative sampling. Midges were again the dominant component, with a higher proportion of the community (45.8% Table 11-b) than indicated by quantitative sampling (38.8%). Other important taxa were caddisflies (14.4%), stoneflies (12.6%) and

mayflies (8.5%). Beetles comprised a much smaller proportion of the sample (3.3%) than demonstrated by quantitative sampling methods (14.4%).

Dominant Species

Due to the strong flow regime of Upper Three Runs Creek, rheophilic organisms were dominant among the macroinvertebrates colonizing multiplate samplers. The most commonly collected organism at all sites was Hydropsyche elissoma, a net-spinning caddisfly (Table 12a). Other dominant species noted at all three sites were the midges Polypedilum convictum and Rheocricotopus robacki. The blackfly larva Simulium jonesi was a dominant species at Sites 1 and 2 and the midge Corynoneura sp. was a dominant species at Sites 1 and 3. A riffle beetle, Stenelmis crenata, was the second most commonly collected species at Site 2. As noted in earlier phases of Upper Three Runs Creek biomonitoring, most dominant organisms were collector-filterers. No distinct shift in community dominance were noted among sites.

Qualitative sampling yielded many of the same dominant species of macroinvertebrates as quantitative sampling. Although the filtering caddisfly, Hydropsyche elissoma, was the most commonly collected species at Site 1, midges such as Tvetenia discoloripes gp. and Polypedilum convictum tended to predominate overall (Table 12b). The predatory stonefly, Perlesta placida, was one of the dominant species at Site 1 and Site 2. The order of species, in terms of abundance, was different at each site. Nevertheless, no clear shifts in community structure are noted.

4.1.3 Pollution Tolerance Parameters

EPT Index

The number of species of Ephemeroptera, Plecoptera and Trichoptera collected by quantitative sampling varied from a minimum of 7 at Sites 1 and 2 in July 1991 to a maximum of 18 at Site 5 in April 1990 (Table 13a; Figure 6). The highest mean EPT Index was found at Site 2 (EPT = 13.2). No statistically significant differences in EPT richness

were noted among sites (Friedman's Test). As with species richness, the greater number of species were collected in the spring quarter, with a minimum in the autumn quarter. However, the July 1991 EPT values were uncharacteristically low for the summer quarter, especially at Sites 1 and 2.

Qualitative sampling produced EPT values ranging from 10 at Site 5 in October 1990 to 28 at Site 2 in March 1990 (Table 13b). The highest mean EPT value was at Site 2 (EPT = 19.7). March 1991 EPT values were abnormally low for the spring quarter, particularly at Site 2.

Hilsenhoff Biotic Index

The biotic index for quantitative samples ranged from a low of 3.67 at Site 5 in October 1990 (Table 14a; Figure 7) to a high of 5.95 in January 1991 at Site 1. The overall mean biotic index was 4.54, indicative of good to very good water quality. No statistically significant differences among sites were found (Friedman's Test). Biotic index values increased (deteriorated) in 1991, as compared to 1990. Highest (poorest) biotic index values were in the winter quarter, in conjunction with greater midge abundance.

Greater midge abundance also resulted in higher biotic index values for qualitative sampling (Table 14b) than for quantitative sampling. Mean biotic index values ranged from 4.84 at Site 1 to 5.35 at Site 2. Higher biotic index values at Site 2, as compared to Site 1, were noted during all three sampling events.

4.1.4 Functional Group Analysis

Collectors of fine particulate organic matter (FPOM) numerically dominated the macroinvertebrate community of Upper Three Runs Creek (Table 15a; Figure 8), comprising 62% of the total number of organisms collected. This total was evenly split between collector-gatherers (30.7%) and collector-filterers (31.3%). Predators comprised the next largest segment of the community (18.9%), followed by scrapers (12.3%) and shredders (7.2%). The proportions of the functional groups at each of the three sites were

similar. The only statistically significant difference encountered was the mean percentage of predators at Sites 2 and 5. A significantly smaller proportion of predators was collected at Site 2 (Friedman's Test; $p = 0.052$). Seasonally, the proportions of collector-filterers and collector-gatherers were inversely related, with highest collector-gatherer and lowest collector-filterer percentages in the winter quarter. The proportion of predators peaked in the spring quarter of 1990 and 1991. In July 1991 the proportion of predators fell to a level of 11.6%. This value was considerable less than in July 1990. Scrapers were a minimum proportion of the macroinvertebrate community in the winter quarter and a maximum in the summer quarter. Shredders consistently comprised a major portion of the community.

Functional Group analysis using biomass showed predators to be dominant, generally representing 50-80% of the total biomass. July 1991 data were an exception, however, with predators comprising 0.4% to 32% of the total biomass (Table 15b).

4.1.5 Total Biomass

Comparing overall means, total biomass ranged from a mean of 0.3701 g/m² (AFDW) at Site 1 to 0.5405 g/m² at Site 5 (Table 16). Differences among sites were not statistically significant ($p=0.43$; Appendix D). Highest biomass values were obtained in the spring quarter of each year, as many species reach the penultimate instar. Lowest biomass values occurred during the summer quarter, when many early instar nymphs and larvae were collected. Biomass values in July 1991 were particularly low at Sites 1 and 2.

4.1.6 Evenness Parameters

Shannon-Weaver Diversity Index

Shannon-Weaver diversity values were indicative of diverse macroinvertebrate communities at all sites, ranging from a mean of 4.08 at Site 1 to 4.24 at Site 2. No statistically significant differences among sites were measured. Mean diversity values dropped lower (3.53) in July 1991 than in any other quarter of the current phase of the study.

Equitability Index

Equitability varied from a value of 0.59 at Site 5 in January 1991 to 1.65 at Site 2 in July 1991. Mean equitability varied only slightly among sites. No significant differences among sites were noted. (Friedman's Test). Equitability was lowest during the winter quarter.

4.2 Water Chemistry

4.2.1 General Parameters

4.2.1.1 Total Suspended Solids

Total suspended solids ranged in value from <5.0 to 32.0 mg/L for the entire period sampled (See Appendix C1). The highest concentrations of total suspended solids for all sampling dates were measured in July 1991 at all three sites, with Site 5 having the highest concentration (32.0 mg/L) among the three locations. Lowest concentrations for suspended solids were found during January 1991, and values among sites were comparable. A comparison of mean concentrations among sampling locations showed that Site 5 had the highest average amount of suspended solids throughout the entire sampling period, while Site 1 had the lowest. Differences in average total suspended solids among sites were not statistically significant ($F=5.58$; $p=0.072$; Appendix D).

4.2.1.2 Specific Conductance

Specific conductance values were found to be low at all sites sampled. Values ranged from 20.0 to 58.0 micromhos/cm for the entire period sampled (See Appendix C1). For all sites combined, the highest and lowest values across sites were measured during July 1990. Site 2 had the highest average specific conductance, but this was only slightly more than that of the other two sites and was not found to be statistically significant ($F=4.08$; $p=0.184$).

4.2.1.3 Hardness

Hardness values were determined to be in the soft classification (0-75 mg/L CaCO_3), with concentrations ranging from 5.0 to 33.0 mg/L CaCO_3 (See Appendix C1). Lowest hardness concentrations were typically found during October 1990 at all three sites, with the highest values occurring during July 1990 at all three sites. The highest average hardness was measured at Site 1 with a value of 11.7 mg/L CaCO_3 , but this was not considerably higher than the other two sites, and no statistical differences among sites were found ($F=3.58$; $p=0.252$).

4.2.1.4 Alkalinity

Alkalinity values for all three sites combined ranged from <1.0 to 5.0 mg/L and were well below the U.S. EPA Aquatic Life Criterion of 20.0 mg/L as CaCO_3 (U.S. EPA 1986). Sampling results indicated that for all three sites, the highest average alkalinity was measured during April 1990, while the lowest was found during July 1990. Average alkalinity at Site 1 was lowest relative to the other two groups, but average concentrations among groups were not found to be statistically significant ($F=0.75$; $p=0.956$).

4.2.1.5 Residual Chlorine

Residual chlorine concentrations ranged from Not Detected (ND) to <0.05 mg/L, with Not Detected found both in July 1990 and July 1991 (see Appendix C1). All other sites and time periods evaluated were found to have concentrations of <0.05. Mean values among sites were identical; therefore, no statistical analysis was performed.

4.2.2 Metals

4.2.2.1 Chromium

Both dissolved and total Chromium was less than detectable limits at all sites and collection dates (Table 18; Appendix C2).

4.2.2.2 Copper

Copper was detected at levels of 0.019 mg/L at Site 1 in April 1990. This concentration of copper was close to the detection limit. No copper was detected at any other site and date combination.

4.2.2.3 Lead

No lead was detected at any of the three sampling sites or in the effluent.

4.2.2.4 Manganese

Manganese was detected at levels ranging as high as 0.147 mg/L (total) at Site 5 in July 1990. No statistically significant differences in concentration of manganese among sites were noted (Appendix D). Manganese levels in the effluent were generally much lower than in samples from stream sites.

4.2.2.5 Mercury

At detection limits ranging from 2 ug/L to 30 ug/L, no mercury was detected from any stream site or effluent sample.

4.2.2.6 Sodium

Mean Sodium (Total) values ranged from 1.324 mg/L at Site 1 to 1.601 mg/L at Site 5. Differences among the sites were not statistically significant (Appendix D). Levels of Total Sodium were substantially higher in the F/H Area effluent, at a mean of 17.57 mg/L.

4.2.2.7 Uranium

Uranium was not detected in most samples collected from the effluent and the three stream sites. Low levels of uranium was detected at Site 5 in July 1990 (4.8 ug/L). Differences among sites were not statistically significant (Appendix D).

4.2.2.8 Zinc

Zinc levels were generally at or below detection limits. Relatively high zinc levels (0.1 mg/L) were detected, however, at Site 1 in July 1991.

4.2.3 Nutrients

4.2.3.1 Nitrites

Nitrite concentrations did not vary considerably among sites. Concentrations ranged from <.01 to <0.10 for all sites during the entire sampling period (See Appendix C3). The highest concentration (<0.10 mg/L) was measured during January and March 1991 at all three sites. Mean estimated concentrations of nitrites were similar for all sites and were not found to be statistically significant ($F=1.0$; $p=0.74$).

4.2.3.2 Nitrates

Nitrate concentrations were low at all three sites and ranged from 0.10 to 0.38 mg/L (See Appendix C3). The lowest concentration of nitrates (0.10 mg/L) was measured during several sampling periods. The highest concentration was measured at Site 2 during October 1990 which corresponded to a high nitrate concentration (26 mg/l) in the F/H ETF effluent. Mean concentrations among sites were comparable and were not found to be statistically significant when tested ($F=4.08$; $p=0.184$).

4.2.3.3 Ammonia Nitrogen

Concentrations of ammonia nitrogen ranged from <0.10 to 0.38 mg/L (See Appendix C3). The lowest concentration was measured during April 1990 at both Sites 2 and 5. The highest concentration was measured at Site 5 during July 1991. For the most part, concentrations among sites were comparable through all sampling periods and average concentrations were found to be similar. No statistical differences among sites were determined when tested ($F=0.58$; $p=0.956$).

4.2.3.4 Total Phosphate

Total Phosphate (T-PO₄) concentrations were found to be high and ranged from 0.02 to 0.310 mg/L (20 to 310 ug/L). Lowest concentrations were measured during April 1990 at all three sites, while the highest concentration was measured in July 1991 at Site 5. Among sites, total phosphate was comparable during all sampling periods. Average concentrations of total phosphates were lowest at Site 1. No statistical differences among sites were found when tested ($F=1.58$; $p=0.57$). Although no criteria have been developed, the U.S. EPA has recommended that total phosphates as phosphorous should not exceed 100 ug/L so as to prevent eutrophication.

4.2.3.5 Ortho-Phosphate

Concentrations of Ortho-Phosphate during the period sampled ranged from <0.01 to 0.05 mg/L (See Appendix C3). Among groups and time periods, concentrations were found be comparable. Average concentrations were similar and no statistical differences among sites were found ($F=1.58$; $p=0.57$).

4.2.4 Organics

4.2.4.1 Tributyl Phosphate

No tributyl phosphate was detected during any sampling period at any site (See Appendix C4). Mean concentrations among sites were identical; therefore, no statistical analysis was performed.

4.2.4.2 Kerosene

No kerosene was detected in any sampling period at any site (See Appendix C4). No variation in measured concentrations among sites during any sampling period was found. Average concentrations of Kerosene were comparable for all three sites, therefore, no statistical analysis was conducted.

4.2.5 Field Parameters

4.2.5.1 pH

pH in Upper Three Runs Creek ranged from a mean of 5.48 at Site 2 to 5.74 at Site 5. F/H Area Effluent pH was more alkaline than the stream, with a mean pH of 7.31.

4.2.5.2 Dissolved Oxygen

Dissolved oxygen remained at acceptable levels (>5.0 mg/L) throughout the study period at all sites.

4.2.5.3 Temperature

Little temperature variation was noted among sites. Effluent temperatures ranged from 17.8 to 30.4°C and were substantially warmer than stream temperatures during the winter and spring quarters.

4.3 Chronic Toxicity Testing

Six quarterly Ceriodaphnia dubia Survival and Reproduction Tests were conducted on F/H Area ETF Effluent during the study period (Table 19). Results of all tests showed the effluent to be acutely toxic at 100% concentration. With the exception of March 1990 the effluent was neither acutely nor chronically toxic at 30% concentration. As a result, 7 Day LC₅₀ values ranged from 39.5-54.8% effluent concentration and Chronic NOEC values were 3 to 30% effluent concentration.

Figure 2
Species Richness
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 3
Average # Of Species Per Replicate
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 4
Density (#/m²)
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 5
 Percent Composition By Major Taxonomic Group
 By # of Organisms - All Dates Combined
 Quantitative Macroinvertebrate Sampling
 Upper Three Runs Creek
 Alken County, SC

March 1990 - July 1991

Figure 6
EPT Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 7
Hilsenhoff Biotic Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC

March 1990 - July 1991

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 8
Functional Group Analysis - % By Pop. #
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC

March 1990 - July 1991

CG = Collector Gathered
CF = Collector Filterers
P = Predators
SC = Scrapers
SH = Shredder Herbivores
SD = Shredder Detritivores

Site 1

Site 2

Site 5

Figure 9
Functional Group Analysis - % By Biomass
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

March 1990 - July 1991

CG = Collector Gatherers
CF = Collector Filterers
P = Predators
SC = Scrapers
SH = Shredder Herbivores
SD = Shredder Detritivores

Site 1

Site 2

Site 5

Figure 10
Shannon - Weaver Diversity Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC

March 1990 - July 1991

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Figure 11
Equitability Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Alken County, SC

Winter Quarter

Spring Quarter

Summer Quarter

Autumn Quarter

Table 5

Species List – Upper Three Runs Creek Biomonitoring
April 1990 – July 1991

	Quantitative Sampling			Qualitative Sampling					
Taxon	Site	1	2	5	Site	1	2	5	
EPHEMEROPTERA (mayflies)									
Baetis ephippiatus		X	X	X		X	X		
Baetis frondalis							X		
Baetis intercalaris			X						
Baetis propinquus						X			
(Baetis sp.)				X					
Centroptilum sp.							X		
Dannella simplex			X			X	X	X	
Dolania americana						X	X	X	
Ephemerella argo		X	X	X		X	X	X	
Ephemerella catawba group						X		X	
Ephemerella dorothea		X	X	X		X	X	X	
Ephemerella invaria/rotunda		X	X	X		X	X	X	
Eurylophella temporalis							X	X	
(Eurylophella sp.)							X		
Heptagenia flavescens		X	X	X					
Hexagenia limbata						X			
Neophemera youngi			X	X		X	X	X	
Paraleptophlebia sp.			X			X	X	X	
Pseudocloeon bimaculatus			X	X				X	
Pseudocloeon dubium			X	X			X		
Pseudocloeon punctiventris		X	X			X	X	X	
Stenonema modestum/smithae		X	X	X		X	X	X	
Tricorythodes sp.		X	X						
Number of Mayfly Species by Site		8	13	9	Number of Mayfly Species by Method		13	14	12
				14	18				

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

	Quantitative Sampling				Qualitative Sampling			
Taxon	Site	1	2	5	Site	1	2	5

ODONATA (dragonflies,damselflies)
Boyeria vinosa
Calopteryx spp.
Gomphus (G.) lividus
Neurocordulia sp.
Perithemis sp.
Progomphus sp.
Tetragoneuria costalis/cynosura

X		X

X	X	X
X	X	X
	X	X
X		
X		
	X	
		X

Number of Odonate Species by Site
Number of Odonate Species by Method

1	0	1
1		

4	4	4
7		

PLECOPTERA (stoneflies)
Acroneuria abnormis
Acroneuria mela
Amphinemura delosa
Haploperla brevis
Helopicus subvarians
Isoperla bilineata
Isoperla dicala
Isoperla holochlora/lata
(Isoperla sp.)
Leuctra sp.
Paragnetina kansensis
Perlesta placida
Pteronarcys dorsata
Taeniopteryx sp.

X	X	X
X		X
X	X	X
X	X	
X	X	X
		X
	X	X
X	X	X
X	X	X
X	X	X
	X	

X	X	X
		X
	X	
		X
	X	
X	X	X
X	X	
X		
X	X	X
X	X	X
X	X	X
X	X	X

Number of Stonefly Species by Site
Number of Stonefly Species by Method

8	8	8
11		

7	9	8
11		

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Quantitative Sampling					Qualitative Sampling				
Taxon	Site	1	2	5	Site	1	2	5	
HETEROPTERA (true bugs)									
Gerris remigis								X	
Metrobates hesperius								X	
Ochterus sp.							X		
Rhagovelia obesa						X	X	X	
<i>Number of True Bug Species by Site</i>		0	0	0	<i>Number of True Bug Species by Method</i>		1	2	3
		0					4		
MEGALOPTERA (hellgrammites)									
Chauliodes rastricornis							X		
Corydalis cornutus		X	X	X		X			
Nigronia serricornis			X	X				X	
<i>Number of Megaloptera Species by Site</i>		1	2	2	<i>Number of Megaloptera Species by Method</i>		1	1	1
		2					3		
LEPIDOPTERA (moths)									
Parapoynx obscuralis		X	X				X		
<i>Number of Lepidoptera Species by Site</i>		1	1	0	<i>Number of Lepidoptera Species by Method</i>		0	1	0
		1					1		

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Quantitative Sampling					Qualitative Sampling				
Taxon	Site	1	2	5	Site	1	2	5	
<hr/>									
TRICHOPTERA (caddisflies)									
Agarodes libalis							X		
Anisocentropus pyraloides								X	
Brachycentrus numerosus		X	X	X		X	X		
Ceraclea sp.							X		
Cheumatopsyche spp.		X	X	X		X	X	X	
Chimarra sp.		X		X			X		
Diplectrona modesta			X						
Hydropsyche elissoma		X	X	X		X	X	X	
(Hydropsyche sp.)							X	X	
Hydroptila sp.		X	X	X			X		
Lepidostoma sp.							X		
Lype diversa		X	X	X		X			
Macrostemum carolina						X			
Mayatrichia ayama		X		X					
Micrasema rusticum		X	X						
Micrasema wataga		X	X			X		X	
Neureclipsis sp.							X	X	
Nyctiophylax sp.		X							
Oecetis sp.						X	X		
Phylocentropus sp.		X					X	X	
Polycentropus sp.		X				X	X	X	
Ptilostomis sp.							X		
Pycnopsyche sp.						X		X	
Triaenodes spp.						X	X	X	
<hr/>									
Number of Caddisfly Species by Site		12	8	7	13		19		
Number of Caddisfly Species by Method									

**Species List - Upper Three Runs Creek Biomonitoring
March 1990 - July 1991**

42

Table 5

Species List - Upper Three Runs Creek Biomonitoring
March 1990 - July 1991

Taxon	Quantitative Sampling			Qualitative Sampling					
	Site	1	2	5	Site	1	2	5	
<hr/>									
DIPTERA (true flies)									
Non-Chironomidae									
Antocha sp.						X			
Atherix lantha		X	X	X		X	X	X	
Bezzia sp. 2						X		X	
Chelifera sp.		X							
Clinocera sp.				X					
Culex sp.							X	X	
Dolichopodidae						X			
(Empididae)			X	X			X	X	
Erioptera/Ormosia sp.						X	X	X	
Hemerodromia sp. 1		X	X	X			X	X	
Hemerodromia sp. 2							X		
Hexatoma sp.						X			
Palpomyia sp. 4								X	
Probezzia sp.						X	X	X	
Simulium jonesi		X	X	X		X	X	X	
Simulium snowi								X	
Simulium tuberosum		X	X	X			X		
Stilobezzia sp.							X		
Telmatoscopus sp.		X							
Tipula sp. A								X	
Tipula sp. 1							X	X	
Tipula (Yamato.) sp. 7			X					X	
Tipula (Yamato.) nr strepens						X			
Wiedemannia sp.						X		X	
<hr/>									
Number of Fly Species by Site									
Number of Fly Species by Method									
		6	5	5			10	10	13
				9					20

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Quantitative Sampling			Qualitative Sampling				
	Site	1	2	5	Site	1	2	5
Chironomidae – Tanypodinae								
<i>Ablabesmyia janta/parajanta</i>						X		
<i>Ablabesmyia mallochi</i>						X	X	X
<i>Clinotanypus pinguis</i>						X	X	
<i>Conchapelopia</i> sp.		X	X	X		X	X	X
<i>Meropelopia</i> sp.						X	X	X
<i>Nilotanypus fimbriatus</i>		X		X			X	
<i>Paramerina</i> sp.		X				X	X	X
<i>Procladius</i> sp.								X
Chironomidae – Diamesinae								
<i>Potthastia longimana</i>						X		X
Chironomidae – Orthocladiinae								
<i>Brillia flavifrons</i>			X	X		X	X	X
<i>Corynoneura</i> nr. <i>taris</i>		X	X	X				
<i>Corynoneura</i> sp. 3		X	X	X		X		
<i>Corynoneura</i> sp. 4			X					X
<i>Cricotopus bicinctus</i>		X	X	X		X		X
<i>Cricotopus vierriensis</i>			X	X				
(<i>Cricotopus</i> sp.)		X						
<i>Eukiefferiella</i> sp.				X				
<i>Lopescladius</i> sp.								X
<i>Nanocladius</i> sp.		X	X	X			X	
<i>Orthocladius annectens</i>						X		
<i>Orthocladius carlatus</i>						X	X	X
<i>Orthocladius lignicola</i>		X	X	X			X	X
(<i>Orthocladius</i> sp.)		X	X	X				X
<i>Parakiefferiella</i> sp. A		X	X			X	X	
<i>Parakiefferiella</i> sp. 4						X		
<i>Parametriocnemus lundbecki</i>		X	X	X		X	X	X
<i>Rheocricotopus robacki</i>		X	X	X		X	X	X
<i>Rheocricotopus tuberculatus</i>		X						

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Site	Quantitative Sampling			Site	Qualitative Sampling		
		1	2	5		1	2	5
Rheosmittia sp.			X			X		
Synorthocladius semivirens		X	X			X	X	X
Thienemanniella fusca group		X	X	X		X	X	X
Thienemanniella xena group		X	X	X		X	X	X
Tvetenia discoloripes group		X	X	X		X	X	X
Tvetenia paucunca		X	X	X		X	X	X
Unniella multivirga						X	X	
Chironomidae – Chironomini								
Cryptochironomus blarina							X	X
Cryptochironomus fulvus group			X	X		X	X	X
Kloosia dorsenna							X	X
Lauterborniella agrayloides								X
Microtendipes pedellus						X	X	X
Microtendipes nr. rydalensis							X	
Pagastiella ostansa			X			X	X	X
Paracladopelma loganae							X	
Paracladopelma undine						X	X	X
Paralauterborniella nigrohalteralis							X	X
Polypedilum convictum		X	X	X		X	X	X
Polypedilum fallax		X	X	X				X
Polypedilum halterale						X	X	X
Polypedilum illinoense		X	X	X				
Polypedilum scalaenum			X			X	X	X
Pseudochironomus sp.			X					
Stenochironomus hiliaris		X		X		X		X
Tribelos jucundum						X	X	X
Xylotopus par								

Table 5

Species List – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

	Quantitative Sampling			Qualitative Sampling				
Taxon	Site	1	2	5	Site	1	2	5
Chironomidae – Tanytarsini								
Cladotanytarsus sp.		X				X	X	X
Rheotanytarsus distinctissimus group		X	X	X		X	X	X
Rheotanytarsus exiguus group		X	X	X		X	X	X
Tanytarsus glabrescens group		X	X	X		X	X	X
Tanytarsus sp. I Cantrell						X	X	X
Tanytarsus sp. IX Rutter						X		
Tanytarsus sp. XII Rutter							X	
Tanytarsus sp. XIII Rutter				X			X	X
Tanytarsus sp. XIV Rutter			X					
Tanytarsus sp. XVI Rutter			X			X	X	X
(Tanytarsus sp.)		X		X		X	X	
Number of Midge Species by Site		26	31	26	39		40	40
Number of Midge Species by Method					39	54		
CRUSTACEA								
Asellus sp.						X		
Cambarinae						X	X	X
Palaemonetes paludosus						X	X	X
Number of Crustacea Species by Site		0	0	0	3		2	2
Number of Crustacea Species by Method					0	3		

Table 5

Species List - Upper Three Runs Creek Biomonitoring
March 1990 - July 1991

Taxon	Quantitative Sampling			Qualitative Sampling
	Site	1	2	5
ANNELIDA (worms)				
(Naididae)				
Pristina/Stephensoniana sp.				
Spirosperma sp.				
(Tubificidae)				
<i>Number of Worm Species by Site</i>				
<i>Number of Worm Species by Method</i>				
MOLLUSCA (snails, clams)				
Campeloma sp.				
Elliptio sp.				
Ferrissia sp.				
Gillia altilis				
<i>Number of Worm Species by Site</i>				
<i>Number of Worm Species by Method</i>				
ACARI (water mites)				
Hydracarina				
<i>Number of Mite Species by Site</i>				
<i>Number of Mite Species by Method</i>				
Total # of Macroinvertebrate Species				
<i>By Site</i>				
<i>By Method</i>				

Taxa in parentheses were too immature to allow more definitive identification or were damaged.

Table 6

Summary of Biomonitoring Parameter Results
Quantitative and Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Quantitative
Sampling

Parameter	
Species Richness	
Average # of Species per Sampler	
Total # of Organisms per Site	
Density (per m ²)	
EPT Index	
Biotic Index	
Biomass (g/m ²)	
Diversity Index	
Equitability Index	

Site

Quantitative
Sampling

1	2	5
29.8	33.2	31.5
12.8	15.1	14.2
178.3	230.8	204.8
285.3	369.3	327.7
11.8	13.2	12.7
4.62	4.65	4.43
0.3701	0.4151	0.5405
4.08	4.24	4.16
0.95	0.98	0.96

Site

Quantitative
Sampling

1	2	5
56.7	58.3	58.7
N/A	N/A	N/A
201.0	306.3	237.7
N/A	N/A	N/A
18.0	19.7	16.3
4.84	5.35	5.04
N/A	N/A	N/A
N/A	N/A	N/A
N/A	N/A	N/A

Table 7a

Species Richness
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
June–July 1987	36.0	45.0	56.0	50.0	47.0	74.0	46.8
September–October 1987	39.0	43.0	34.0	31.0	37.0	67.0	36.8
January–February 1988	35.0	51.0	45.0	52.0	45.0	80.0	45.6
March–April 1988	57.0	57.0	56.0	48.0	53.0	89.0	54.2
June–July 1988	37.0	38.0	36.0	46.0	43.0	74.0	40.0
September–October 1988	29.0	34.0	36.0	32.0	27.0	61.0	31.6
Overall Mean	38.8	44.7	43.8	43.2	42.0	74.2	42.5

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
December–January 1989	39.0	37.0	38.0	37.0	38.0	65.0	37.8
March–April 1989	38.0	54.0	44.0	40.0	45.0	73.0	44.2
June–July 1989	34.0	31.0	29.0	24.0	22.0	55.0	28.0
September–October 1989	30.0	29.0	26.0	30.0	27.0	58.0	28.4
December–January 1990	35.0	30.0	31.0	27.0	32.0	50.0	31.0
Overall Mean	35.2	36.2	33.6	31.6	32.8	60.2	33.9

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March–April 1990	28.0	35.0	35.0	27.0	36.0	63.0	32.2
June–July 1990	32.0	33.0			31.0	50.0	32.0
September–October 1990	26.0	28.0			21.0	37.0	25.0
December–January 1991	32.0	35.0			37.0	49.0	34.7
March–April 1991	39.0	40.0			37.0	55.0	38.7
June–July 1991	22.0	28.0			27.0	37.0	25.7
Mean	29.8	33.2			31.5	48.5	31.4
Standard Deviation	5.9	4.6			6.5	10.2	5.3
Total	70.0	77.0			69.0	104.0	72.0

Table 7b

Species Richness
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
November 1987	40.0	71.0	66.0	65.0	67.0	139.0	61.8
March 1988	79.0	76.0	59.0	96.0	63.0	168.0	74.6
October 1988	45.0	55.0	41.0	45.0	34.0	106.0	44.0
Overall Mean	54.7	67.3	55.3	68.7	54.7	137.7	60.1

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March 1989	78.0	81.0	80.0	83.0	73.0	146.0	79.0
October 1989	30.0	27.0	32.0	39.0	37.0	71.0	33.0
Overall Mean	54.0	54.0	56.0	61.0	55.0	108.5	56.0

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2
March 1990	58.0	70.0
October 1990	52.0	54.0
March 1991	60.0	51.0
Mean	56.7	58.3
Standard Deviation	4.2	10.2
Total	104.0	110.0

Site 5	Total	Mean
62.0	108.0	63.3
50.0	90.0	52.0
64.0	98.0	58.3
58.7	98.7	57.9
7.6	9.0	5.7
107.0	163.0	107.0

Table 8

Average # of Species per Replicate
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5		Mean
June–July 1987	17.0	21.8	29.2	26.0	25.4		23.9
September–October 1987	20.8	22.0	16.6	15.6	16.8		18.4
January–February 1988	16.6	26.4	22.4	22.4	23.8		22.3
March–April 1988	28.0	29.4	26.8	29.8	27.6		28.3
June–July 1988	16.8	16.8	16.6	18.6	19.2		17.6
September–October 1988	10.8	12.8	12.6	15.8	10.8		12.6
Overall Mean	18.3	21.5	20.7	21.4	20.6		20.5

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5		Mean
December–January 1989	20.4	19.6	18.4	17.8	19.2		19.1
March–April 1989	21.0	24.0	22.2	18.6	22.0		21.6
June–July 1989	12.6	12.0	11.2	10.6	8.6		11.0
September–October 1989	9.0	10.6	11.4	14.4	9.6		11.0
December–January 1990	16.8	14.6	14.6	11.6	14.4		14.4
Overall Mean	16.0	16.2	15.6	14.6	14.8		15.4

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5		Mean
March–April 1990	9.6	16.0	19.4	15.4	16.6		15.4
June–July 1990	12.4	12.6			12.2		12.4
September–October 1990	10.4	12.0			8.0		10.1
December–January 1991	15.6	17.6			17.5		16.9
March–April 1991	16.0	19.0			17.4		17.5
June–July 1991	12.8	13.2			13.6		13.2
Mean	12.8	15.1			14.2		14.3
Standard Deviation	2.6	2.9			3.7		2.8

Table 9a

Total # of Organisms per Site
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
June–July 1987	241	300	527	423	413	1904	380.8
September–October 1987	733	699	260	230	346	2268	453.6
January–February 1988	267	599	461	539	274	2140	428.0
March–April 1988	555	501	542	632	670	2900	580.0
June–July 1988	207	174	242	252	275	1150	230.0
September–October 1988	152	141	101	197	122	713	142.6
Overall Mean	359.2	402.3	355.5	378.8	350.0	1845.8	369.2

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
December–January 1989	343	442	469	271	303	1828	365.6
March–April 1989	328	335	352	319	299	1633	326.6
June–July 1989	104	126	130	138	93	591	118.2
September–October 1989	71	122	120	175	80	568	113.6
December–January 1990	160	144	142	122	142	710	142.0
Overall Mean	201.2	233.8	242.6	205.0	183.4	1066.0	213.2

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March–April 1990	125	194	273	287	278	1157.0	231.4
June–July 1990	122	113			108	343.0	114.3
September–October 1990	121	140			81	342.0	114.0
December–January 1991	280	381			291	952.0	317.3
March–April 1991	185	277			257	719.0	239.7
June–July 1991	237	280			214	731.0	243.7
Mean	178.3	230.8			204.8	707.3	210.1
Standard Deviation	68.0	100.6			89.8	325.3	80.4
Total	1070	1385			1229	4244	

Table 9b

Total Number of Organisms per Site
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
November 1987	123.0	498.0	258.0	458.0	215.0	1552.0	310.4
March 1988	325.0	489.0	285.0	778.0	245.0	2122.0	424.4
October 1988	147.0	233.0	212.0	210.0	83.0	885.0	177.0
Overall Mean	198.3	406.7	251.7	482.0	181.0	1519.7	303.9

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March 1989	447.0	465.0	435.0	336.0	261.0	1944.0	388.8
October 1989	94.0	146.0	169.0	118.0	102.0	629.0	125.8
Overall Mean	270.5	305.5	302.0	227.0	181.5	1286.5	257.3

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2
March 1990	299.0	594.0
October 1990	122.0	136.0
March 1991	182.0	189.0
Mean	201.0	306.3
Standard Deviation	90.0	250.5
Total	603.0	919.0

Site 5	Total	Mean
303.0	1196.0	399.0
138.0	396.0	132.0
272.0	643.0	214.3
237.7	745.0	248.4
87.7	409.6	136.7
713.0	2235.0	

Table 10a

Mean Density of Organisms per Square Meter
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
June-July 1987	385.6	480.0	843.2	676.8	660.8	3046.4	609.3
September-October 1987	1172.8	1118.4	416.0	368.0	553.6	3628.8	725.8
January-February 1988	427.2	958.4	737.6	862.4	438.4	3424.0	684.8
March-April 1988	888.0	801.6	867.2	1011.2	1072.0	4640.0	928.0
June-July 1988	331.2	278.4	387.2	403.2	440.0	1840.0	368.0
September-October 1988	243.2	225.6	161.6	315.2	211.2	1156.8	231.4
Overall Mean	574.7	643.7	568.8	606.1	562.7	2956.0	591.2

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
December-January 1989	548.8	707.2	750.4	433.6	484.8	2924.8	585.0
March-April 1989	524.8	536.0	563.2	510.0	478.4	2612.4	522.5
June-July 1989	233.3	210.0	216.7	230.0	155.0	1045.0	209.0
September-October 1989	118.3	203.3	200.0	291.7	133.3	946.6	189.3
December-January 1990	266.7	240.0	236.7	203.3	236.7	1183.4	236.7
Overall Mean	338.4	379.3	393.4	333.7	297.6	1742.4	348.5

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March-April 1990	200	310	437	459	445	955.2	318.3
June-July 1990	195	181			173	548.8	183.0
September-October 1990	194	224			130	547.2	182.7
December-January 1991	448	610			466	1523.2	508.0
March-April 1991	296	443			411	1150.4	383.3
June-July 1991	379	448			342	1169.6	389.7
Mean	285.3	369.3			327.7	982.4	327.5
Standard Deviation	108.8	160.9			143.6	383.1	127.7
Total	1712.0	2216.0			1967.0	5894.4	

Mean and total columns for 1990-1991 do not include Sites 3 & 4.

Table 10b

Density by Major Taxonomic Group (Number/m²)

Quantitative Macroinvertebrate Sampling

Upper Three Runs Creek

Aiken County, SC

March 1990 – July 1991

Ephemeroptera (Mayflies)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	1.6	24.0	(56.0)	(22.4)	33.6	19.7	59.2
June–July 1990	20.8	17.6			17.6	18.7	56.0
September–October 1990	1.6	22.4			4.8	9.6	28.8
December–January 1991	35.2	38.4			12.8	28.8	86.4
March–April 1991	14.4	27.2			22.4	21.3	64.0
June–July 1991	14.4	27.2			22.4	21.3	64.0
Mean (Mar 90 – Jul 91)	14.7	26.1			18.9	19.9	59.7
Standard Deviation	12.7	7.0			9.8		18.5

Plecoptera (Stoneflies)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	59.2	59.2	(100.8)	(73.6)	110.4	76.3	228.8
June–July 1990	43.2	16.0			38.4	32.5	97.6
September–October 1990	25.6	30.4			33.6	29.9	89.6
December–January 1991	36.8	48.0			32.0	38.9	116.8
March–April 1991	56.0	44.8			72.0	57.6	172.8
June–July 1991	41.6	41.6			41.6	41.6	124.8
Mean (Mar 90 – Jul 91)	43.7	40.0			54.7	46.1	138.4
Standard Deviation	12.4	15.0			31.0		53.0

Trichoptera (Caddisflies)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	64.0	44.8	(108.8)	(105.6)	81.6	63.5	190.4
June–July 1990	59.2	70.4			40.0	56.5	169.6
September–October 1990	60.8	52.8			30.4	48.0	144.0
December–January 1991	16.0	41.6			16.0	24.5	73.6
March–April 1991	35.2	33.6			51.2	40.0	120.0
June–July 1991	156.8	86.4			116.8	120.0	360.0
Mean (Mar 90 – Jul 91)	65.3	54.9			56.0	58.8	176.3
Standard Deviation	48.5	19.8			37.1		98.7

Table 10b

Density by Major Taxonomic Group (Number/m²)

Quantitative Macroinvertebrate Sampling

Upper Three Runs Creek

Aiken County, SC

March 1990 – July 1991

Coleoptera (Beetles)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	9.6	41.6	(20.8)	(30.4)	14.4	21.9	65.6
June–July 1990	30.4	32.0			43.2	35.2	105.6
September–October 1990	8.0	12.8			9.6	10.1	30.4
December–January 1991	0.0	0.0			1.6	0.5	1.6
March–April 1991	49.6	107.2			80.0	78.9	236.8
June–July 1991	94.4	217.6			96.0	136.0	408.0
Mean (Mar 90 – Jul 91)	32.0	68.5			40.8	47.1	141.3
Standard Deviation	35.5	82.0			39.5		154.3

Diptera (Non–Chironomidae)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	20.8	38.4	(48.0)	(132.6)	32.0	30.4	91.2
June–July 1990	17.6	11.2			12.8	13.9	41.6
September–October 1990	36.8	24.0			14.4	25.1	75.2
December–January 1991	19.2	8.0			30.4	19.2	57.6
March–April 1991	14.4	49.6			48.0	37.3	112.0
June–July 1991	22.4	40.0			20.8	27.7	83.2
Mean (Mar 90 – Jul 91)	21.9	28.5			26.4	25.6	76.8
Standard Deviation	7.8	16.8			13.2		24.9

Chironomidae (Midges)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean	Total
March–April 1990	41.6	100.8	(48.0)	(132.6)	172.8	105.1	315.2
June–July 1990	19.2	14.4			20.8	18.1	54.4
September–October 1990	60.8	81.6			33.6	58.7	176.0
December–January 1991	340.8	473.6			369.6	394.7	1184.0
March–April 1991	123.2	179.2			137.6	146.7	440.0
June–July 1991	48.0	32.0			40.0	40.0	120.0
Mean (Mar 90 – Jul 91)	105.6	146.9			129.1	127.2	381.6
Standard Deviation	120.4	170.2			133.1		417.0

Table 10b

Density by Major Taxonomic Group (Number/m²)

Quantitative Macroinvertebrate Sampling

Upper Three Runs Creek

Aiken County, SC

March 1990 – July 1991

Chironomidae – Tanypodinae	Site 1	Site 2
March–April 1990	1.6	1.6
June–July 1990	0.0	0.0
September–October 1990	1.6	1.6
December–January 1991	1.6	4.8
March–April 1991	4.8	1.6
June–July 1991	0.0	0.0
Mean (Mar 90 – Jul 91)	1.6	1.6
Standard Deviation	1.8	1.8

Site 5	Mean	Total
0.0	1.1	3.2
0.0	0.0	0.0
0.0	1.1	3.2
1.6	2.7	8.0
1.6	2.7	8.0
3.2	1.1	3.2
1.1	1.4	4.3
1.3	1.0	3.1

Chironomidae – Orthocladiinae	Site 1	Site 2
March–April 1990	19.2	64.0
June–July 1990	1.6	6.4
September–October 1990	36.8	54.4
December–January 1991	241.6	232.0
March–April 1991	84.8	124.8
June–July 1991	36.8	24.0
Mean (Mar 90 – Jul 91)	70.1	84.3
Standard Deviation	88.5	83.0

Site 5	Mean	Total
97.6	60.3	180.8
9.6	5.9	17.6
17.6	36.3	108.8
267.2	246.9	740.8
100.8	103.5	310.4
28.8	29.9	89.6
86.9	80.4	241.3
96.9	88.0	264.0

Chironomidae – Chironominae	Site 1	Site 2
March–April 1990	20.8	35.2
June–July 1990	17.6	8.0
September–October 1990	22.4	25.6
December–January 1991	97.6	236.8
March–April 1991	33.6	52.8
June–July 1991	11.2	8.0
Mean (Mar 90 – Jul 91)	33.9	61.1
Standard Deviation	32.1	87.8

Site 5	Mean	Total
75.2	43.7	131.2
11.2	12.3	36.8
16.0	21.3	64.0
100.8	145.1	435.2
35.2	40.5	121.6
11.2	10.1	30.4
41.6	45.5	136.5
37.9	50.8	152.3

Mean and total density values for March–April 1990 exclude Site 3 and Site 4.

Table 11a

Percent Composition by Major Taxonomic Group
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Ephemeroptera (Mayflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	7.5%	7.6%	11.5%	9.1%	6.7%	9.6%
Phase 2 (Dec 88 – Feb 90)	11.6%	8.5%	6.3%	6.2%	9.4%	8.5%
March–April 1990	0.8%	7.7%	12.8%	4.9%	7.6%	7.4%
June–July 1990	10.7%	9.7%			10.2%	10.2%
September–October 1990	0.8%	10.0%			3.7%	5.3%
December–January 1991	7.9%	6.3%			2.8%	5.7%
March–April 1991	4.9%	6.1%			5.1%	5.4%
June–July 1991	3.8%	6.1%			6.5%	5.5%
Combined (Mar 1990 – Jul 1991)	5.1%	7.1%			5.8%	6.2%
Standard Deviation	3.9%	1.8%			2.7%	1.9%

Plecoptera (Stoneflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	7.0%	10.0%	10.3%	9.2%	8.5%	9.3%
Phase 2 (Dec 88 – Feb 90)	12.1%	13.8%	13.4%	15.0%	14.6%	13.8%
March–April 1990	29.6%	19.1%	23.1%	16.0%	24.8%	21.8%
June–July 1990	22.1%	8.9%			22.2%	17.8%
September–October 1990	13.2%	13.6%			25.9%	16.4%
December–January 1991	8.2%	7.9%			6.9%	7.7%
March–April 1991	18.9%	10.1%			17.5%	15.0%
June–July 1991	11.0%	9.3%			12.2%	10.7%
Combined (Mar 1990 – Jul 1991)	15.3%	10.8%			16.7%	14.2%
Standard Deviation	8.0%	4.2%			7.6%	5.1%

Trichoptera (Caddisflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	7.1%	6.5%	8.0%	9.2%	10.6%	8.2%
Phase 2 (Dec 88 – Feb 90)	9.7%	13.4%	11.9%	14.4%	11.7%	12.4%
March–April 1990	32.0%	14.4%	24.9%	23.0%	18.4%	21.9%
June–July 1990	30.3%	38.9%			23.2%	30.9%
September–October 1990	31.4%	23.6%			23.5%	26.3%
December–January 1991	3.6%	6.8%			3.4%	4.8%
March–April 1991	11.9%	7.6%			12.5%	10.4%
June–July 1991	41.4%	19.3%			34.1%	30.8%
Combined (Mar 1990 – Jul 1991)	22.9%	14.9%			17.1%	17.9%
Standard Deviation	14.3%	12.0%			10.5%	10.9%

Table 11a

Percent Composition by Major Taxonomic Group
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Coleoptera (Beetles)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	2.6%	4.7%	8.3%	4.0%	4.9%	5.2%
Phase 2 (Dec 88 – Feb 90)	2.6%	8.3%	6.8%	8.7%	6.0%	7.1%
March–April 1990	4.8%	13.4%	4.8%	6.6%	3.2%	6.3%
June–July 1990	15.6%	17.7%			25.0%	19.2%
September–October 1990	4.1%	5.7%			7.4%	5.6%
December–January 1991	0.0%	0.0%			0.3%	0.1%
March–April 1991	16.8%	24.2%			19.5%	20.6%
June–July 1991	24.9%	48.6%			28.0%	34.9%
Combined (Mar 1990 – Jul 1991)	11.2%	18.6%			12.4%	14.4%
Standard Deviation	9.5%	17.1%			11.8%	12.9%
Diptera (Non-Chironomidae)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	6.9%	3.4%	2.3%	4.9%	5.1%	4.3%
Phase 2 (Dec 88 – Feb 90)	4.4%	3.8%	4.1%	15.5%	11.0%	7.6%
March–April 1990	10.4%	12.4%	11.0%	28.9%	7.2%	14.7%
June–July 1990	9.0%	6.2%			7.4%	7.6%
September–October 1990	19.0%	10.7%			11.1%	13.7%
December–January 1991	4.3%	1.3%			6.5%	3.8%
March–April 1991	4.9%	11.2%			11.7%	9.7%
June–July 1991	5.9%	8.9%			6.1%	7.1%
Combined (Mar 1990 – Jul 1991)	7.7%	7.7%			8.1%	7.8%
Standard Deviation	5.5%	4.1%			2.4%	4.2%
Chironomidae (Midges)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
Phase 1 (Pre-Operational)	68.8%	67.4%	59.2%	63.1%	64.0%	63.1%
Phase 2 (Dec 88 – Feb 90)	59.1%	51.6%	57.0%	40.0%	46.8%	50.1%
March–April 1990	20.8%	32.5%	23.4%	20.6%	38.9%	27.7%
June–July 1990	9.8%	8.0%			12.0%	9.9%
September–October 1990	31.4%	36.4%			25.9%	32.2%
December–January 1991	76.1%	77.7%			79.4%	77.7%
March–April 1991	42.2%	40.4%			33.5%	38.4%
June–July 1991	12.7%	7.1%			12.6%	10.5%
Combined (Mar 1990 – Jul 1991)	37.0%	39.8%			39.4%	38.8%
Standard Deviation	24.6%	25.9%			24.8%	24.9%

Table 11a

Percent Composition by Major Taxonomic Group
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Chironomidae – Tanypodinae	Site 1	Site 2
March–April 1990	0.8%	0.5%
June–July 1990	0.0%	0.0%
September–October 1990	0.8%	0.7%
December–January 1991	0.4%	0.8%
March–April 1991	1.6%	0.4%
June–July 1991	0.0%	0.0%
Combined (Mar 1990 – Jul 1991)	0.6%	0.4%
Standard Deviation	0.6%	0.3%

Site 5
0.0%
0.0%
0.0%
0.0%
0.3%
0.4%
0.2%
0.2%

All Sites
0.2%
0.0%
0.6%
0.5%
0.7%
0.3%
0.4%
0.3%

Chironomidae – Orthocladiinae	Site 1	Site 2
March–April 1990	9.7%	20.6%
June–July 1990	0.8%	3.6%
September–October 1990	19.0%	24.4%
December–January 1991	54.0%	38.0%
March–April 1991	29.1%	28.3%
June–July 1991	9.7%	5.3%
Combined (Mar 1990 – Jul 1991)	24.6%	22.8%
Standard Deviation	19.1%	13.4%

Site 5
22.0%
5.5%
13.5%
57.4%
24.5%
8.4%
26.5%
18.9%

All Sites
9.8%
3.2%
19.9%
48.6%
27.0%
7.7%
24.6%
16.8%

Chironomidae – Chironominae	Site 1	Site 2
March–April 1990	10.3%	11.4%
June–July 1990	9.0%	4.4%
September–October 1990	11.6%	11.3%
December–January 1991	21.7%	38.9%
March–April 1991	11.5%	11.7%
June–July 1991	3.0%	1.8%
Combined (Mar 1990 – Jul 1991)	11.8%	16.6%
Standard Deviation	6.1%	13.2%

Site 5
16.9%
6.5%
12.4%
21.7%
8.6%
3.3%
12.7%
6.9%

All Sites
7.1%
6.7%
11.7%
28.6%
10.6%
2.6%
13.8%
9.1%

Table 11b

Percent Composition by Major Taxonomic Group
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Ephemeroptera (Mayflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	17.9%	12.5%	24.4%	4.1%	18.1%	13.2%
March 1988	15.2%	13.4%	4.6%	7.7%	15.4%	10.6%
October 1988	4.8%	2.2%	4.3%	5.2%	3.6%	4.0%
March 1989	10.7%	9.9%	7.8%	7.7%	10.3%	9.3%
October 1989	14.9%	4.8%	5.9%	7.6%	9.8%	8.0%
March 1990	7.7%	8.6%			3.3%	7.0%
October 1990	7.4%	22.8%			13.8%	14.9%
March 1991	12.1%	2.1%			7.4%	7.2%
Combined (1990–1991)	9.0%	9.4%			6.9%	8.5%

Plecoptera (Stoneflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	15.4%	7.2%	6.6%	5.5%	17.7%	8.7%
March 1988	10.9%	16.0%	20.6%	16.6%	17.0%	16.1%
October 1988	0.7%	3.9%	7.1%	10.5%	10.8%	6.3%
March 1989	16.3%	17.2%	16.8%	22.3%	14.9%	17.5%
October 1989	3.2%	2.7%	8.3%	8.5%	5.9%	5.9%
March 1990	17.7%	12.0%			6.6%	12.0%
October 1990	16.4%	2.9%			13.8%	10.9%
March 1991	20.3%	6.9%			16.2%	14.6%
Combined (1990–1991)	18.2%	9.6%			11.6%	12.6%

Trichoptera (Caddisflies)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	28.5%	20.1%	11.6%	7.4%	5.6%	13.5%
March 1988	6.5%	24.3%	18.1%	24.9%	6.6%	19.0%
October 1988	27.9%	26.2%	21.2%	24.8%	12.1%	23.6%
March 1989	6.7%	15.9%	7.8%	14.9%	9.6%	11.0%
October 1989	29.8%	25.3%	11.8%	22.0%	12.8%	19.7%
March 1990	22.4%	22.4%			8.6%	18.9%
October 1990	10.7%	8.8%			2.2%	7.1%
March 1991	14.3%	7.4%			10.3%	10.6%
Combined (1990–1991)	17.6%	17.3%			8.0%	14.4%

Table 11b

Percent Composition by Major Taxonomic Group
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Coleoptera (Beetles)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	1.6%	1.2%	1.5%	1.1%	1.9%	1.4%
March 1988	5.3%	1.2%	3.5%	2.4%	2.3%	2.7%
October 1988	3.4%	4.7%	3.8%	3.3%	2.4%	3.7%
March 1989	1.2%	3.4%	2.5%	5.7%	3.1%	3.2%
October 1989	3.2%	3.4%	1.8%	2.5%	2.9%	2.7%
March 1990	1.3%	1.5%			3.0%	1.8%
October 1990	11.5%	8.1%			10.1%	9.8%
March 1991	1.7%	1.1%			2.9%	2.0%
Combined (1990–1991)	3.5%	2.4%			4.3%	3.3%

Diptera (Non-Chironomidae)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	15.4%	4.5%	7.0%	11.4%	4.7%	7.8%
March 1988	5.6%	4.3%	5.3%	5.7%	4.2%	5.1%
October 1988	15.0%	8.6%	17.0%	8.1%	22.9%	12.9%
March 1989	5.4%	4.5%	12.9%	8.0%	13.4%	8.4%
October 1989	8.5%	4.1%	8.3%	22.9%	9.8%	10.3%
March 1990	7.4%	5.7%			8.3%	6.8%
October 1990	9.8%	5.2%			13.0%	10.9%
March 1991	4.4%	7.9%			7.0%	6.5%
Combined (1990–1991)	7.0%	6.1%			8.7%	7.2%

Chironomidae (Midges)	Site 1	Site 2	Site 3	Site 4	Site 5	All Sites
November 1987	17.1%	51.2%	45.7%	69.0%	40.0%	51.5%
March 1988	54.0%	39.3%	45.4%	40.2%	47.9%	43.6%
October 1988	39.5%	50.6%	43.4%	45.2%	25.3%	43.4%
March 1989	57.5%	44.5%	51.0%	37.2%	46.0%	47.9%
October 1989	27.7%	43.2%	32.0%	25.4%	42.2%	34.3%
March 1990	39.1%	43.6%			66.7%	48.3%
October 1990	34.4%	22.8%			24.6%	24.7%
March 1991	42.3%	66.7%			50.0%	52.7%
Combined (1990–1991)	39.1%	45.3%			52.2%	45.8%

Table 12a

Dominant Species
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken Co., SC
March 1990 – July 1991

SITE 1

Rank	Species	#	%
1	Hydropsyche elissoma	183	17.1
2	Corynoneura sp.	62	5.8
3	Simulium jonesi	59	5.5
4	Polypedium convictum	57	5.3
5	Rheocricotopus robacki	51	4.8

SITE 2

Species	#	%
Hydropsyche elissoma	152	10.9
Stenelmis crenata	110	7.9
Polypedium convictum	93	6.7
Rheocricotopus robacki	88	6.4
Simulium jonesi	81	5.8

SITE 5

Species	#	%
Hydropsyche elissoma	165	13.4
Polypedium convictum	95	7.7
Tvetenia discoloripes gp.	86	7.0
Rheocricotopus robacki	78	6.3
Corynoneura sp.	56	4.6

The number shown represents all sampling periods combined.

Only species comprising approximately 5% or more of the total number of organisms collected at a site are included.

Table 12b

Dominant Species
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken Co., SC
March 1990 – July 1991

SITE 1

Rank	Species	#	%
1	Hydropsyche elissoma	63	10.4
2	Tvetenia discoloripes gp.	46	7.6
3	Perlesta placida	41	6.8
4	Polypedilum convictum	39	6.5
5	Rheocricotopus robacki	37	6.1

SITE 2

Species	#	%
Tvetenia discoloripes gp.	90	9.8
Polypedilum convictum	83	9.0
Hydropsyche elissoma	80	8.7
Perlesta placida	49	5.3
Cheumatopsyche spp.	47	5.1

SITE 5

Species	#	%
Polypedilum convictum	66	9.3
Rheotanytarsus distinctissimus	51	7.2
Tvetenia discoloripes gp.	48	6.7
Rheocricotopus robacki	40	5.6
Hydropsyche elissoma	32	4.4

The number shown represents all sampling periods combined.

Only species comprising approximately 5% or more of the total number of organisms collected at a site are included.

Table 13a

EPT Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
June-July 1987	11	14	17	18	14	14.8
September-October 1987	8	11	11	8	7	9.0
January-February 1988	12	19	19	18	17	17.0
March-April 1988	17	23	24	18	18	20.0
June-July 1988	14	15	13	13	15	14.0
September-October 1988	8	10	9	8	5	8.0
Overall Mean	11.7	15.3	15.5	13.8	12.7	13.8

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
December-January 1989	16	18	18	15	20	17.4
March-April 1989	16	19	14	13	16	15.6
June-July 1989	17	11	13	14	10	13.0
September-October 1989	12	11	10	16	10	11.8
December-January 1990	15	17	16	14	16	15.6
Overall Mean	15.2	15.2	14.2	14.4	14.4	14.7

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Total	Mean
March-April 1990	12	13	17	13	18	25	14.6
June-July 1990	15	15			14	22	14.7
September-October 1990	10	12			7	14	9.7
December-January 1991	13	16			13	18	14.0
March-April 1991	14	16			14	20	14.7
June-July 1991	7	7			10	11	8.0
Mean	11.8	13.2			12.7	18.3	12.6
Standard Deviation	3.8	4.0			3.9	4.7	3.9
Total	28	29			24	38	27.0

Table 13b

EPT Index
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
November 1987	16	20	18	21	17	18.4
March 1988	27	28	23	36	21	27.0
October 1988	10	14	12	11	10	11.4
Overall Mean	17.7	20.7	17.7	22.7	16.0	18.9

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
March 1989	24	24	26	26	18	23.6
October 1989	14	8	9	14	13	11.6
Overall Mean	19.0	16.0	17.5	20.0	15.5	17.6

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2
March 1990	23	28
October 1990	13	20
March 1991	18	11
Mean	18.0	19.7
Standard Deviation	5.0	8.5
Total	30.0	37.0

Site 5	Total	Mean
20	37	23.7
10	24	14.3
19	26	16.0
16.3	29.0	18.0
5.5	7.0	5.0
29.0	48.0	32.0

Table 14a

Biotic Index

Quantitative Macroinvertebrate Sampling

Upper Three Runs Creek

Aiken County, SC

March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
June–July 1987	4.56	4.73	5.31	5.00	4.81	4.88
September–October 1987	5.52	5.91	5.35	5.54	5.29	5.52
January–February 1988	5.18	5.30	5.12	5.33	4.90	5.17
March–April 1988	5.17	4.88	4.84	4.92	5.20	5.00
June–July 1988	4.50	4.64	4.38	4.60	4.49	4.52
September–October 1988	5.63	4.66	4.18	5.14	3.80	4.68
Overall Mean	5.09	5.02	4.86	5.09	4.75	4.96

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
December–January 1989	5.10	5.21	5.12	4.59	5.04	5.01
March–April 1989	4.69	4.89	4.83	4.38	4.09	4.58
June–July 1989	4.43	4.53	4.79	4.13	4.40	4.46
September–October 1989	4.73	3.91	4.48	3.93	4.10	4.23
December–January 1990	4.24	3.94	4.27	3.75	3.54	3.95
Overall Mean	4.64	4.50	4.70	4.16	4.23	4.44

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
March–April 1990	4.20	3.95	4.01	3.87	4.18	4.03
June–July 1990	3.80	4.40			3.97	4.05
September–October 1990	4.20	4.25			3.67	4.09
December–January 1991	5.95	5.63			5.62	5.62
March–April 1991	4.89	5.00			4.74	4.88
June–July 1991	4.66	4.64			4.41	4.57
Mean	4.62	4.65			4.43	4.54
Standard Deviation	0.76	0.60			0.69	0.63

Table 14b

Biotic Index
Qualitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
November 1987	4.49	5.10	4.45	4.88	4.54	4.69
March 1988	5.01	4.88	4.75	4.74	4.73	4.82
October 1988	6.13	5.33	4.44	4.92	4.92	5.15
Overall Mean	5.21	5.10	4.55	4.85	4.73	4.89

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
March 1989	4.71	4.94	4.86	4.84	5.59	4.99
October 1989	5.50	5.47	5.44	4.20	5.21	5.16
Overall Mean	5.11	5.21	5.15	4.52	5.40	5.08

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 5	Mean
March 1990	4.92	5.13	5.28	5.12
October 1990	4.75	5.49	4.96	5.18
March 1991	4.84	5.43	4.87	5.05
Mean	4.84	5.35	5.04	5.12
Standard Deviation	0.09	0.19	0.22	0.07

Table 15a

Functional Group Analysis-% by Pop. #
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 - July 1991

Collector-Gatherers	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	48.7%	50.1%	46.8%	44.1%	47.0%	47.5%
Phase 2 (Dec 88 - Feb 90)	45.2%	37.1%	38.8%	29.7%	35.7%	37.6%
March-April 1990	16.0%	32.0%	24.0%	19.0%	30.0%	25.0%
June-July 1990	19.0%	21.0%			25.0%	22.0%
September-October 1990	18.0%	29.0%			21.0%	23.0%
December-January 1991	54.0%	36.0%			52.0%	54.0%
March-April 1991	33.0%	35.0%			29.0%	33.0%
June-July 1991	24.0%	32.0%			24.0%	27.1%
Mean (Mar 1990 - Jul 1991)	27.3%	30.8%			30.2%	30.7%
Standard Deviation	14.4%	5.4%			11.2%	12.1%

Collector-Filterers	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	22.3%	14.6%	13.5%	19.5%	18.4%	17.5%
Phase 2 (Dec 88 - Feb 90)	19.7%	22.2%	27.3%	32.5%	24.5%	24.2%
March-April 1990	44.0%	28.0%	34.0%	53.0%	25.0%	37.0%
June-July 1990	30.0%	38.0%			28.0%	32.0%
September-October 1990	45.0%	33.0%			35.0%	38.0%
December-January 1991	18.0%	33.0%			16.0%	18.0%
March-April 1991	25.0%	24.0%			27.0%	25.0%
June-July 1991	48.0%	28.0%			40.0%	38.0%
Mean (Mar 1990 - Jul 1991)	35.0%	30.7%			28.5%	31.3%
Standard Deviation	12.4%	5.0%			8.3%	8.2%

Predators	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	15.1%	18.7%	16.9%	17.6%	17.0%	17.0%
Phase 2 (Dec 88 - Feb 90)	17.1%	20.7%	19.9%	18.7%	20.8%	19.4%
March-April 1990	31.0%	22.0%	26.0%	18.0%	26.0%	24.0%
June-July 1990	27.0%	17.0%			21.0%	22.0%
September-October 1990	21.0%	19.0%			27.0%	22.0%
December-January 1991	15.0%	17.0%			18.0%	15.0%
March-April 1991	23.0%	14.0%			22.0%	19.0%
June-July 1991	10.0%	9.0%			14.0%	11.6%
Mean (Mar 1990 - Jul 1991)	21.2%	16.3%			21.3%	18.9%
Standard Deviation	7.7%	4.5%			4.9%	4.8%

::

Table 15a

Functional Group Analysis by Site
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Scrapers	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	5.5%	7.1%	11.6%	8.4%	7.5%	8.1%
Phase 2 (Dec 88 – Feb 90)	10.6%	10.8%	7.8%	10.0%	9.5%	9.7%
March–April 1990	4.0%	11.0%	10.0%	7.0%	8.0%	8.0%
June–July 1990	17.0%	17.0%			18.0%	18.0%
September–October 1990	8.0%	10.0%			10.0%	9.0%
December–January 1991	5.0%	4.0%			3.0%	5.0%
March–April 1991	12.0%	15.0%			12.0%	13.0%
June–July 1991	15.0%	28.0%			19.0%	21.0%
Mean (Mar 1990 – Jul 1991)	10.2%	14.2%			11.7%	12.3%
Standard Deviation	5.3%	8.1%			6.1%	6.2%
Shredder–Herbivores	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	7.8%	8.8%	9.9%	9.2%	8.4%	8.9%
Phase 2 (Dec 88 – Feb 90)	3.9%	6.2%	6.3%	5.1%	6.1%	5.5%
March–April 1990	4.0%	5.0%	4.0%	3.0%	7.0%	5.0%
June–July 1990	5.0%	6.0%			4.0%	5.0%
September–October 1990	7.0%	8.0%			5.0%	7.0%
December–January 1991	7.0%	9.0%			11.0%	7.0%
March–April 1991	3.0%	5.0%			4.0%	4.0%
June–July 1991	2.0%	2.0%			2.0%	2.0%
Mean (Mar 1990 – Jul 1991)	4.7%	5.8%			5.5%	5.0%
Standard Deviation	2.1%	2.5%			3.1%	1.9%
Shredder–Detritivores	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
Phase 1 (Pre-Operational)	0.6%	0.8%	1.2%	1.2%	1.7%	0.9%
Phase 2 (Dec 88 – Feb 90)	3.4%	2.9%	4.9%	4.0%	3.3%	3.6%
March–April 1990	1.0%	2.0%	1.0%	1.0%	4.0%	2.0%
June–July 1990	2.0%	1.0%			4.0%	2.0%
September–October 1990	1.0%	1.0%			2.0%	1.0%
December–January 1991	1.0%	1.0%			0.0%	1.0%
March–April 1991	4.0%	7.0%			6.0%	6.0%
June–July 1991	1.0%	1.0%			1.0%	1.0%
Mean (Mar 1990 – Jul 1991)	1.7%	2.2%			2.8%	2.2%
Standard Deviation	1.2%	2.4%			2.2%	1.9%

Table 15B

Functional Group Analysis – Biomass Data
 Quantitative Macroinvertebrate Sampling – Site 1
 Upper Three Runs Creek Biomonitoring
 Savannah River Site
 Aiken County, SC
 March 1990 – July 1991

Biomass in Grams

Date	CG		Functional Group *			
			P	SC	SH	SD
April 1990	0.0035	0.0410	0.2052	0.0044	0.0019	0.0019
July 90	0.0069	0.0068	0.1270	0.0167	0.0107	0.0101
October 90	0.0033	0.0031	0.0607	0.0094	0.0066	0.0066
January 91	0.0202	0.0095	0.1723	0.0744	0.0511	0.0508
April 91	0.0182	0.0253	0.3627	0.0155	0.0005	0.0000
July 91	0.0012	0.0000	0.0020	0.0016	0.0007	0.0007

% of Total Biomass

Date	CG	CF	P	SC	SH	SD
April 1990	1.4%	15.9%	79.6%	1.7%	0.7%	0.7%
July 90	3.9%	3.8%	71.2%	9.4%	6.0%	5.7%
October 90	3.7%	3.4%	67.7%	10.5%	7.4%	7.3%
January 91	5.3%	2.5%	45.5%	19.7%	13.5%	13.4%
April 91	4.3%	6.0%	85.9%	3.7%	0.1%	0.0%
July 91	19.4%	0.0%	32.3%	25.8%	11.3%	11.3%

* CG – collector gatherer; CF – collector filter; P – predator; SC – scraper
 SH – shredder herbivore; SD – shredder detritivore

Table 15B

Functional Group Analysis – Biomass Data
 Quantitative Macroinvertebrate Sampling – Site 2
 Upper Three Runs Creek Biomonitoring
 Savannah River Site
 Aiken County, SC
 March 1990 – July 1991

Biomass in Grams

Date	CG	CF	Functional Group *			
			P	SC	SH	SD
April 90	0.0162	0.0254	0.4765	0.0145	0.0012	0.0007
July 90	0.0084	0.0072	0.0970	0.0087	0.0007	0.0007
October 90	0.0047	0.0097	0.1130	0.0161	0.0067	0.0064
January 91	0.0162	0.0125	0.1428	0.0319	0.0132	0.0139
April 91	0.0338	0.0272	0.3163	0.0312	0.0007	0.0000
July 91	0.0036	0.0052	0.0086	0.0078	0.0079	0.0078

% of Total Biomass

Date	CG	CF	P	SC	SH	SD
April 1990	3.0%	4.8%	89.2%	2.7%	0.2%	0.1%
July 90	6.8%	5.8%	79.1%	7.1%	0.6%	0.6%
October 90	3.0%	6.2%	72.2%	10.3%	4.3%	4.0%
January 91	7.0%	5.4%	62.0%	13.8%	5.7%	6.0%
April 91	8.3%	6.6%	77.3%	7.6%	0.2%	0.0%
July 91	8.8%	12.7%	21.0%	19.1%	19.3%	19.1%

* CG – collector gatherer; CF – collector filter; P – predator; SC – scraper
 SH – shredder herbivore; SD – shredder detritivore

Table 15B

Functional Group Analysis - Biomass Data
Quantitative Macroinvertebrate Sampling - Site 5
Upper Three Runs Creek Biomonitoring
Savannah River Site
Aiken County, SC
March 1990 - July 1991

Biomass in Grams

Date	CG	CF	Functional Group *			
			P	SC	SH	SD
April 1990	0.0179	0.0406	0.3064	0.0147	0.0027	0.0021
July 90	0.0126	0.0158	0.0673	0.0262	0.0151	0.0152
October 90	0.0029	0.0044	0.1253	0.0171	0.0123	0.0122
January 91 **	NA	NA	NA	NA	NA	NA
April 91	0.0316	0.0293	0.0005	0.0529	0.0001	0.0002
July 91	0.0316	0.0293	0.0005	0.0529	0.0001	0.0002

% of Total Biomass

Date	CG	CF	P	SC	SH	SD
April 1990	4.7%	10.6%	79.7%	3.8%	0.7%	0.5%
July 90	8.3%	10.4%	44.2%	17.2%	9.9%	10.0%
October 90	1.7%	2.5%	71.9%	9.8%	7.1%	7.0%
January 91	NA	NA	NA	NA	NA	NA
April 91	4.1%	7.5%	84.8%	3.6%	0.1%	<0.1
July 91	27.4%	25.6%	0.4%	46.3%	0.1%	0.2%

* CG - collector gatherer; CF - collector filter; P - predator; SC - scraper
SH - shredder herbivore; SD - shredder detritivore

** Biomass not divided into functional groups.

Table 16

Total Biomass (AFDW in g/m²)
 Quantitative Macroinvertebrate Sampling
 Upper Three Runs Creek
 Aiken County, SC
 March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5		Mean
June–July 1987	0.1120	0.2712	0.2152	0.1123	0.1347		0.1691
September–October 1987	0.2406	0.5138	0.2424	0.1675	0.2938		0.2916
January–February 1988	0.3282	0.4144	0.3914	0.4867	0.8299		0.4901
March–April 1988	1.1550	0.7107	0.9030	0.5171	0.4277		0.7427
June–July 1988	0.4226	0.2616	0.6826	0.2050	0.5064		0.4156
September–October 1988	0.0434	0.2224	0.7998	0.2694	0.2414		0.3153
Overall Mean	0.3836	0.3990	0.5391	0.2930	0.4057		0.4041

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5		Mean
December–January 1989	0.6976	0.6003	0.5058	0.3898	0.4514		0.5290
March–April 1989	0.4162	0.5755	0.3346	0.2507	0.3933		0.3941
June–July 1989	0.2682	0.5118	0.1637	0.2440	0.1388		0.2653
September–October 1989	0.3091	0.3053	0.4175	0.2771	0.2112		0.3040
December–January 1990	0.9242	1.0700	0.6025	0.3865	0.6528		0.7272
Overall Mean	0.5231	0.6126	0.4048	0.3096	0.3695		0.4439

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2
March–April 1990	0.4298	0.8908
June–July 1990	0.2970	0.2045
September–October 1990	0.1495	0.2610
December–January 1991	0.6305	0.3842
March–April 1991	0.7037	0.6820
June–July 1991	0.0103	0.0682
Mean	0.3701	0.4151
Standard Deviation	0.2706	0.3121

Site 5		Mean
0.6407		0.3923
0.2537		32.0000
0.2903		0.2336
0.6123		0.5423
1.2553		0.8803
0.1905		0.0897
0.5405		5.6897
0.3984		12.8922

Table 17a

Shannon Weaver Diversity Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
June-July 1987	4.33	4.72	4.70	4.93	4.74	4.68
September-October 1987	4.11	4.13	4.29	4.19	4.23	4.19
January-February 1988	4.17	4.49	4.28	4.17	4.32	4.29
March-April 1988	4.83	4.92	4.80	4.72	4.70	4.79
June-July 1988	4.57	4.68	4.44	4.67	4.50	4.57
September-October 1988	3.95	4.27	4.71	4.27	4.01	4.24
Overall Mean	4.33	4.54	4.54	4.49	4.42	4.46

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
December-January 1989	4.41	4.29	4.14	4.42	4.31	4.31
March-April 1989	4.44	5.01	4.50	4.32	4.60	4.57
June-July 1989	4.56	4.18	3.72	3.87	3.83	4.03
September-October 1989	4.28	3.92	3.97	3.99	4.13	4.06
December-January 1990	4.40	4.32	4.34	4.22	4.54	4.36
Overall Mean	4.42	4.34	4.13	4.16	4.28	4.27

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
March-April 1990	3.86	4.40	4.30	3.58	4.32	4.41
June-July 1990	4.46	4.32			4.36	4.70
September-October 1990	3.92	4.20			3.91	4.25
December-January 1991	4.00	3.95			3.92	4.12
March-April 1991	4.89	5.00			4.74	4.88
June-July 1991	3.34	3.56			3.69	3.53
Mean	4.08	4.24			4.16	4.32
Standard Deviation	0.53	0.48			0.39	0.48

Table 17b

Equitability Index
Quantitative Macroinvertebrate Sampling
Upper Three Runs Creek
Aiken County, SC
March 1990 – July 1991

Phase 1 (Pre-Operational)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
June–July 1987	0.83	0.87	0.70	0.92	0.85	0.81
September–October 1987	0.64	0.60	0.85	0.87	0.76	0.73
January–February 1988	0.74	0.65	0.64	0.50	0.64	0.64
March–April 1988	0.74	0.79	0.73	0.81	0.74	0.76
June–July 1988	0.95	1.00	0.89	0.83	0.77	0.88
September–October 1988	0.79	0.97	1.08	0.88	0.89	0.89
Overall Mean	0.83	0.76	0.78	0.76	0.76	0.78

Phase 2 (Dec 1988 – Feb 1990)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
December–January 1989	0.79	0.78	0.68	0.86	0.76	0.77
March–April 1989	0.84	0.89	0.75	0.73	0.80	0.79
June–July 1989	1.03	0.87	0.66	0.88	0.68	0.86
September–October 1989	0.93	0.76	0.88	0.77	0.96	0.85
December–January 1990	0.89	0.97	0.97	1.00	1.06	0.97
Overall Mean	0.91	0.83	0.77	0.82	0.88	0.83

Phase 3 (Mar 1990 – Jul 1991)	Site 1	Site 2	Site 3	Site 4	Site 5	Mean
March–April 1990	0.75	0.89	0.83	0.63	0.81	0.96
June–July 1990	1.03	0.88			0.97	1.22
September–October 1990	0.85	0.96			1.05	1.12
December–January 1991	0.72	0.66			0.59	0.75
March–April 1991	0.89	0.83			0.93	0.88
June–July 1991	1.47	1.65			1.42	1.51
Mean	0.95	0.98			0.96	1.07
Standard Deviation	0.28	0.34			0.28	0.27

Table 18

Summary of Water Chemistry Data
 Quarterly Water Chemistry Sampling
 Upper Three Runs Creek ; F/H Area ETF Effluent
 Aiken County, SC
 March 1990 – July 1991

N = 6

Mean Value

Parameter

General Parameters	Units	Site 1	Site 2	Site 5	Effluent
Total Suspended Solids	mg/L	9.8	10.5	17.3	4.3
Specific Conductance	umhos/cm	27.0	28.8	28.5	109.7
Hardness	mg/L	11.7	8.9	11.3	1.3
Alkalinity	mg/L	2.50	2.75	2.82	9.63
Residual Chlorine	mg/L	< 0.05	< 0.05	< 0.05	< 0.05

Metals (Total)	Units	Site 1	Site 2	Site 5	Effluent
Chromium	mg/L	< 0.011	< 0.010	< 0.010	< 0.010
Copper	mg/L	< 0.013	< 0.011	< 0.015	< 0.011
Lead	mg/L	< 0.028	< 0.028	< 0.028	< 0.028
Manganese	mg/L	0.046	0.033	0.050	0.011
Mercury	ug/L	0.7	0.7	0.7	0.7
Sodium	mg/L	1.324	1.517	1.601	17.567
Uranium	ug/L	< 0.8	< 1.3	< 1.6	< 50.0
Zinc	mg/L	0.167	0.024	0.076	0.050

Metals (Dissolved)	Units	Site 1	Site 2	Site 5	Effluent
Chromium	mg/L	< 0.011	< 0.011	< 0.010	< 0.010
Copper	mg/L	< 0.013	< 0.011	< 0.011	< 0.011
Lead	mg/L	< 0.028	< 0.028	< 0.028	< 0.028
Manganese	mg/L	0.032	0.031	0.033	0.007
Mercury	ug/L	0.7	0.7	0.7	0.7
Sodium	mg/L	1.610	1.825	1.932	16.960
Uranium	mg/L	< 0.8	< 0.9	< 1.1	< 0.8
Zinc	mg/L	0.035	0.025	0.025	0.055

Nutrients	Units	Site 1	Site 2	Site 5	Effluent
Nitrite Nitrogen	mg/L	< 0.05	< 0.05	< 0.05	< 0.13
Nitrate Nitrogen	mg/L	0.13	0.17	0.19	7.62
Ammonia Nitrogen	mg/L	0.29	0.26	0.28	0.24
Total Phosphate	mg/L	< 0.09	< 0.11	< 0.16	< 0.07
Ortho Phosphate	mg/L	< 0.04	< 0.04	< 0.05	< 0.04

Table 18

Summary of Water Chemistry Data
Quarterly Water Chemistry Sampling
Upper Three Runs Creek ; F/H Area ETF Effluent
Aiken County, SC
March 1990 – July 1991

N = 6

Organics	Units	Site 1	Site 2	Site 5	Effluent
Tri-Butyl Phosphate	ug/L	< 28.0	< 28.0	< 28.0	< 28.0
Kerosene	ug/L	< 4.63	< 4.63	< 4.63	< 4.63

Field Parameters	Units	Site 1	Site 2	Site 5	Effluent
pH	units	5.63	5.48	5.74	7.31
Dissolved Oxygen	mg/L	9.2	9.4	9.3	8.0
Temperature	C	16.1	16.0	16.0	22.8

Table 19

Ceriodaphnia Toxicity Testing Results
Upper Three Runs Creek;F/H Area ETF Effluent
Savannah River Site
Aiken, South Carolina
April 1990 – June 1991

	<i>Acute Toxicity</i> <i>LC50</i>	<i>Chronic Toxicity</i> <i>NOEC</i>
<i>Date</i>		
March 1990	39.54	3.0
June 1990	42.41	30.0
September 1990	46.70	30.0
December 1990	44.38	30.0
March 1991	54.77	30.0
June 1991	54.77	30.0
Mean	47.10	25.50
Standard Deviation	N/A	N/A

Note:

Acute Toxicity calculated by Trimmed Spearman Karber Analysis.

Chronic Toxicity calculated by Dunnett's t Test with Bonferroni's Adjustment.

5.0 SUMMARY AND CONCLUSIONS

Biological and chemical monitoring of Upper Three Runs Creek to assess the impact of F/H-Area-ETF effluent was maintained on a quarterly basis during the period from March 1990 to July 1991.

Biological Monitoring

Biological monitoring included quarterly definitive toxicity testing of the effluent, quarterly quantitative macroinvertebrate stream assessments and semi-annual qualitative macroinvertebrate stream assessments. Macroinvertebrate stream assessments were conducted at three sites, including an upstream control site (Site 1) and two downstream sites (Site 2 and Site 5).

Macroinvertebrate stream assessments provided a species list of aquatic taxa as well as abundance data for each site. All three sampling sites proved to support species rich communities of macroinvertebrates. Both quantitative and qualitative sampling indicated the greatest species richness values at downstream Site 2, although differences among sites were not statistically significant.

Macroinvertebrate density averaged 248.4 organisms per square meter in Upper Three Runs Creek over the study period. Similarly to the species richness pattern, macroinvertebrate density was greatest at downstream Site 2, although differences among sites were not significant. Midges were the dominant major taxonomic group in both quantitative samples (32.7%) and qualitative samples (45.4%), followed by Trichoptera (quantitative 20.9%; qualitative 14.4%), Plecoptera (quantitative 14.9%; qualitative = 12.6%), and beetles (quantitative 14.4%; qualitative 3.3%). Percentages of major taxonomic groups did not vary significantly among sites except for Coleoptera, which were significantly more abundant at Site 2 than at other sites. Rheophilic organisms were the most abundant in both quantitative and qualitative samples, including the net-spinning caddisfly Hydropsyche elissoma, the blackfly larva Simulium jonesi and the midges: Rheocricotopus robacki, Polypedilum convictum, Corynoneura sp. and Tvtenia discoloripes. Predatory stoneflies (Perlesta placida) were among the most commonly collected organisms in qualitative sampling. No distinct shifts in community structure among sites was noted.

A high proportion of pollution intolerant organisms was found at all sites. A mean EPT value of 12.6 was obtained for quantitative sampling and 18.0 for qualitative sampling. Hilsenhoff biotic index values (quantitative data) ranged from a mean of 4.43 at Site 5 to 4.65 at Site 2, indicating good to very good water quality at all sites. No significant differences in EPT values or biotic index values were found among sites.

Total biomass (g/m^2) varied seasonally, peaking in the spring quarter and reaching a minimum in the summer quarter. No statistically significant differences among sites were found. Biomass values were particularly low in July 1991.

Functional group analysis of the density data showed the macroinvertebrate community of Upper Three Runs Creek was dominated by collectors of fine particulate organic matter (62%), evenly split into collector-filterers and collector-gatherers. Predators and scrapers were also abundant, numerically comprising 18.9% and 12.3% of the community respectively. Shredders were a minor component of the macroinvertebrate community. The proportions of functional groups did not vary significantly among sites with the exception of predators, which comprised a significantly smaller percentage of the community at Site 2. Predators were the dominant functional group, in terms of biomass, at all sites.

Biological parameters exhibited considerable seasonal variation during the study period. This was most evident for species richness and diversity, which reached maximum levels in the spring, and in the relative proportion of major taxonomic groups. Caddisflies, mayflies and midges comprised a larger percentage of the community in the winter than in other seasons. Stoneflies reached a peak in relative abundance in the spring and beetles in the summer. There were also temporal trends which were not seasonally related. Inter-year differences were observed in density values, which were substantially higher in all 1991 sampling periods than in 1990, and in biotic index values, which were higher (poorer) in 1991. A number of biological parameter values were aberrant in July 1991. Species richness, caddisfly density, and riffle beetle density (and hence scrapers) all were higher in July 1991 than the previous five sampling periods, especially in light of seasonal fluctuations. In contrast, the diversity of the community and density of predators and shredder-herbivores was uncharacteristically low in July 1991. Although a shift in community structure occurred, no clear evidence of water quality degradation was found.

Water Chemistry

Data from all three sites confirmed the acidic, low hardness characteristics of Upper Three Runs Creek. Temperature values followed normal seasonal cycles and dissolved oxygen remained at acceptable levels throughout the year. Evidence of elevated nutrient levels was found, particularly Nitrate-Nitrogen and Total Phosphate. No differences in nutrient levels among sites, however, were noted. Detectable levels of metals were generally not encountered, with the exception of manganese and sodium. The only metal with elevated concentrations in the effluent was sodium.

Chronic Toxicity Testing

F/H area effluent was found to be toxic to the test organism, Ceriodaphnia dubia, at 100% effluent concentration. At 30% effluent concentration no acute or chronic toxicity was found, with the exception of the March 1990 test. Based upon these results, the effluent would not be expected to be toxic to stream fauna at either the mean or 7Q10 discharge of the receiving stream, which result in effluent dilutions of 1:370 and 1:225, respectively.

Conclusion

Analysis of the quarterly biomonitoring data indicates no evidence of impact upon the macroinvertebrate communities of Upper Three Runs Creek by the F/H Area ETF effluent.

6.0 REFERENCES

1. Barr, Cheryl B. and Joan B. Chapin. 1988. The Aquatic Dryopoidea of Louisiana (Coleoptera: Psephenidae, Dryopidae, Elmidae). *Tulane Studies in Zoology and Botany*. 26 (2): 89-164.
2. Berner L. and M. Pescador, 1988. *The Mayflies of Florida*. University Presses of Florida, Gainesville, Florida, 415 pp.
3. Bode, R.W. 1983. Larvae of North American Eukiefferiella and Tvetenia (Diptera:Chironomidae). *NYS Mus. Bull.* 452:1 - 40.
4. Brigham, A.R., W.U. Brigham, and A. Gnilka (eds.) 1982. *The Aquatic Insects and Oligochaetes of North and South Carolina*. Midwest Aquatic Enterprises, Mahomet, IL. 837 pp.
5. Caldwell, B.A. 1986. Description of the Immature Stages and Adult Female of Unniella multivirga Saether (Diptera:Chironomidae) with Comments on Phylogeny. *Aquatic Insects*.
6. Epler, J.H. 1987. Biosystematics of the Genus Dicrotendipes Kieffer, 1913, (Diptera:Chironomidae:Chironominae) of the World. *Mem. Amer. Entom. Soc.* 36:1 - 214.
7. Fitzpatrick, J.F. 1983. *How to Know the Freshwater Crustacea*. Wm. C. Brown Co. Publishers, Dubuque, Iowa. 227 pp.
8. Gelhaus, John. 1986. Larvae of the Crane Fly Genus Tipula. *Univ. Kansas. Science Bull.*
9. Gilbert, R.O. 1987. *Statistical Methods for Environmental Pollution Monitoring*. Van Nostrand Reinhold Company, New York. 320 pp.
10. Glukhova, V.M. 1979. Larval Midges of the Subfamilies Palpomyiinae and Ceratopogoninae of the Fauna of the USSR (Translation). Nauka Publishers, St. Petersburg. 250 pp.
11. Grodhaus, G. 1987. Endochironomus Kieffer, Tribelos Townes, Synendotendipes n. gen., and Endotribelos, n. gen. (Diptera:Chironomidae) of the Nearctic region. *J. Kansas Ent. Soc.* 60:167 - 247.
12. Hilsenhoff, William L., 1987. An Improved Biotic Index of Organic Stream Pollution. *Great Lakes Entomologist*. 20(1):31 - 29.

13. Hirvenoja, M. 1973. Revision der Gattung *Cricotopus* van der Wulp und ihrer Verwandten (Diptera, Chironomidae). *Ann. Zool. Fenn.* 10:1 - 363.
14. Hudson, P.L., D.R. Lenat, B.A. Caldwell and D. Smith. 1990. Chironomidae of the Southeastern United States: A Checklist of Species and Notes on Biology, Distribution and Habitat. Fish and Wildlife Research Series - U.S. Dept. of Interior. 7: 1-46.
15. Holsinger, J.R. 1972. The Freshwater Amphipod Crustaceans (Gammaridae) of North America. EPA Identification Manual #5.
16. Johannsen, O.A. 1937 - 1939. Aquatic Diptera I-IV. Cornell University. Agric. Exp. Stn.
17. Klemm, D.J. 1985. A Guide to the Freshwater Annelida (Polychaeta, Naidid and Tubificid Oligochaeta, and Hirudinea) of North America. Kendall/Hunt Publishing Co., Dubuque Iowa. 198 pp.
18. Lewis, P.A. 1974. Taxonomy and Ecology of *Stenonema* Mayflies (Heptageniidae: Ephemeroptera). U.S. EPA, Environmental Monitoring Ser. Rept. EPA-670/4-74-006. 81 pp.
19. McAlpine et al. 1981. Manual of Nearctic Diptera, Vol. 1. Res. Branch, Agric. Can. Monogr. 27. 674 pp.
20. Merritt, R.W. and K.W. Cummins (eds.). 1984. An Introduction to Aquatic Insects of North America. Kendall/Hunt Publishing Co., Dubuque, Iowa. 722 pp.
21. Morihara, D.K. and W.P. McCafferty. 1979. The *Baetis* Larvae of North America (Ephemeropter:Baetidae). *Trans. Am. Ent. Soc.* 105:139 - 221.
22. Morse, J.C., J.W. Chapin, D.D. Herlong and R.S. Harvey. 1980. Aquatic Insects of Upper Three Runs Creek, Savannah River Plant, South Carolina. Part I: Orders Other than Diptera. *J. Georgia Entomol. Soc.* 15(1):73 - 101.
23. Morse et al. 1985. Aquatic Insects of Upper Three Runs Creek, Savannah River Plant, South Carolina. Part II: Diptera. *J. Georgia Entomol. Soc.*
24. Needham, J.G., J.R. Traver, and Y.C. HSU. 1935. The Biology of Mayflies with a Systematic Account of North American Species. Comstock, Ithaca. 759 pp.
25. Needham, J.G. and M.J. Westfall, Jr. 1955. A Manual of the Dragonflies on North America (Anisoptera) Including the Greater Antilles and the Provinces of the Mexican Border. Univ. Calif. Press, Berkley. 615 pp.
26. Oliver, D.R. and Roussel, M.E. 1983. Redescription of *Brillia* Kieffer (Diptera:Chironomidae) with Descriptions of Nearctic Species. *Can. Ent.* 115:257 - 279.

27. Pennak, R.W. 1978. Freshwater Invertebrates of the United States (2nd Ed.). J. Wiley and Sons, NY. 803 pp.
28. Roback, S.S. 1981. The Immature Chironomids of the Eastern United States. V. Pentaneurini - Thienemannimyia group. Proc. Acad. Nat. Sci. Philad. 133:73 - 128.
29. Roback, S.S. 1985. The Immature Chironomids of the Eastern United States. VI. Pentaneurini - Genus Ablabesmyia. Proc. Acad. Nat. Sci. Philad. 137:153 - 213.
30. Ross, H.H. 1944. The Caddisflies, or Trichoptera, of Illinois. Bull. Ill. Nat. Hist. Surv. 23:1 - 326.
31. Saether, O.A. 1977. Taxonomic Studies on Chironomidae: Nanocladius, Pseudochironomus and the Harnischia complex. Bull. Fish. Res. Bd. Can. 196:1 - 143.
32. Saether, O.A. and J.E. Sublette 1983. A Review of the Genera Doithrix n. gen., Georthocladius Strenzke, Parachaetocladius Wulker, and Pseudorthocladius Goethebuer. (Diptera:Chironomidae, Orthocladinae). Ent. Scand. Suppl. 20:1 - 100.
33. Saether, O.A. 1985. A Review of Rheocricotopus Thienemann and Harnisch, 1932, with the Description of Three New Species (Diptera, Chironomidae). Spixiana Suppl. 11:59 - 108.
34. Snoddy, E.L. and R. Noblet. 1976. Identification of the Immature Black Flies (Dipter:Simuliidae) of the Southeastern U.S. with Some Aspects of the Adult Role in Transmission of Leucocytozoon smithi to Turkeys. S.C. Agric. Exp. Sta., Clemson, S.C. Techn. Bull. 1057. 58pp.
35. Soptonis, A.R. 1977. A Review of the Nearctic Species of Orthocladius (Orthocladius) Van Der Wulp (Diptera:Chironomidae). Mem. Ent. Soc. Can. 102:1 - 187.
36. Scheffer, P.W. and G.B. Wiggins. 1986. A Systematic Study of the Nearctic Larvae of the Hydropsyche Morosa Group (Trichoptera:Hydropsychidae). Alger Press, Toronto, Canada. 94 pp.
37. Stewart, K. W. and B.P. Stark. 1988. Nymphs of the North American Stonefly Genera (Plecoptera). Entomol. Society of America.
38. Thompson, F.G. 1984. The Freshwater Snails of Florida. A Manual for Identification. University Presses of Florida, Gainesville, Florida. 94 pp.
39. Usinger, R.L., Ed. 1956. Aquatic Insects of California. University of California Press, Los Angeles, California. 508 pp.

40. U.S. EPA. 1979. Methods for Chemical Analysis of Water and Wastes. EPA-600/4-79-020.
41. Weber, C.I. (Ed.). 1973. Biological Field and Laboratory Methods for Measuring the Quality of Surface Waters and Effluents. Cincinnati: Nat. Envir. Res. Center. U.S. Environmental Protection Agency. 176 pp.
42. Wiederholm, T. (Ed.) 1983. Chironomidae of the Holarctic Region. Keys and diagnoses. Part 1 - Larvae. Ent. Scand. Suppl. 19:1 - 457.
43. Wiederholm, T. (Ed.) 1986. Chironomidae of the Holarctic Region. Keys and diagnoses. Part 2 - Pupae. Ent. Scand. Suppl. 28:1 - 482.
44. Wiggins, G.B. 1977. Larvae of the North American Caddisfly Genera. Univ. Toronto Press, Toronto. 401 pp.
45. Williams, W.D. 1972. Freshwater Isopods (Asellidae) of North America. U.S. EPA Identification Manual #7.

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 23, 1990
 CLIENT: S.R.S.
 STREAM: Upper Three Runs Creek
 LOCATION: Aiken Co., SC

ALL SITES

Site 1	Site 2	Site 5	TOTAL
--------	--------	--------	-------

EPHEMEROPTERA (mayflies)

Baetis ephippius	0	2	0	2
Dannella simplex	1	1	1	3
Dolania americana	0	1	2	3
Ephemerella argo	1	2	1	4
Ephemerella invaria/rotunda	8	24	1	33
Eurylophella temporalis	0	2	2	4
Hexagenia limbata	6	0	0	6
Neophemera youngi	3	1	2	6
Paraleptophlebia sp.	1	0	0	1
Pseudocloeon dubium	0	2	0	2
Pseudocloeon punctiventris	0	4	1	5
Stenonema modestum/smithae	3	12	0	15

ODONATA (dragonflies)

Boyeria vinosa	1	1	0	2
Calopteryx sp.	0	1	0	1
Gomphus (Gomphus) lividus	0	3	2	5
Perithemis sp.	1	0	0	1

PLECOPTERA (stoneflies)

Acroneuria abnormis	1	0	0	1
Haploperla brevis	0	0	1	1
Isoperla bilineata	0	2	0	2
Isoperla holochlora/lata	2	10	0	12
Isoperla dicala	5	1	10	16
Leuctra sp.	1	2	1	4
Paragnetina kansensis	2	2	3	7
Perlesta placida	31	43	4	78
Pteronarcys dorsata	11	11	1	23

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 23, 1990
 CLIENT: S.R.S.
 STREAM: Upper Three Runs Creek
 LOCATION: Aiken Co., SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
HEMIPTERA (true bugs)				
Rhagovelia obesa	0	0	1	1
MEGALOPTERA				
Corydalus cornutus	2	0	0	2
Nigronia serricornis	0	0	1	1
TRICHOPTERA (caddisflies)				
Agarodes libalis	0	1	0	1
Cernotina sp.	0	1	0	1
Cheumatopsyche sp.	9	44	2	55
Chimarra sp.	0	1	0	1
Hydropsyche elissoma	44	69	16	129
Hydropsyche sp.	0	1	1	2
Hydroptila sp.	0	2	0	2
Lype diversa	1	0	0	1
Macrostemum carolina	1	0	0	1
Micrasema wataga	3	0	1	4
Neureclipsis sp.	0	1	0	1
Oecetis sp.	1	1	0	2
Phylocentropus sp.	0	6	2	8
Polycentropus sp.	4	0	1	5
Pycnopsyche sp.	1	0	0	1
Triaenodes sp.	3	6	3	12
LEPIDOPTERA (moths)				
Parapoynx obscuralis	0	1	0	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 23, 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

ALL SITES

Site 1	Site 2	Site 5	TOTAL
--------	--------	--------	-------

COLEOPTERA (beetles)

Ancyronyx variegatus	0	1	0	1
Dineutus discolor	0	5	0	5
Macronychus glabratus	3	1	0	4
Optioservus sp.	1	2	3	6
Sperchopsis tessellatus	0	0	1	1
Stenelmis (larvae)	0	0	5	5

DIPTERA - misc. (true flies)

Antocha sp.	1	0	0	1
Atherix lantha	1	1	2	4
Bezzia sp. 2	3	0	2	5
Empididae	0	1	4	5
Hemerodromia sp.1	0	0	2	2
Hexatoma sp.	1	0	0	1
Probezzia sp. 1	2	4	2	8
Palpomyia sp. 4	0	0	3	3
Simulium jonesi	14	27	9	50
Simulium tuberosum	0	1	0	1
Tipula sp.1	0	0	1	1

DIPTERA- Chironomidae (midges)

Ablabesmyia janta	1	0	0	1
Ablabesmyia mallochi	0	1	2	3
Brillia flavifrons	4	0	0	4
Clinotanypus pinguis	6	1	0	7
Conchapelopia sp.	1	9	8	18
Cricotopus bicinctus	1	0	0	1
Cryptochironomus fulvus gp.	2	9	4	15
Lopescladius sp.	0	0	1	1
Microtendipes pedellus	1	3	0	4
Nanocladius sp.	0	1	0	1

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: March 23, 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
<i>Nilotanypus fimbriatus</i>	0	1	0	1
<i>Orthocladius carlatus</i>	0	0	3	3
<i>Orthocladius lignicola</i>	0	1	1	2
<i>Orthocladius</i> sp.	0	0	6	6
<i>Pagastiella ostansa</i>	0	1	0	1
<i>Paracladopelma undine</i>	1	2	1	4
<i>Parakiefferiella</i> sp. A	1	0	0	1
<i>Parakiefferiella</i> sp. 4	1	0	0	1
<i>Paralauterborniella nigrohalteralis</i>	0	0	2	2
<i>Parametriocnemus lundbecki</i>	1	7	10	18
<i>Paramerina</i> sp.	0	0	2	2
<i>Polypedilum convictum</i>	19	60	22	101
<i>Polypedilum halterale</i>	1	0	3	4
<i>Polypedilum scalaenum</i>	2	6	6	14
<i>Pothastia longimana</i>	0	0	3	3
<i>Procladius</i> sp.	0	0	2	2
<i>Rheocricotopus robacki</i>	21	25	8	54
<i>Rheotanytarsus distinctissimus</i> gp.	14	41	50	105
<i>Rheotanytarsus exiguus</i> gp.	1	2	0	3
<i>Stenochironomus hilaris</i>	1	0	2	3
<i>Synorthocladius semivirens</i>	0	2	0	2
<i>Tanytarsus glabrescens</i> gp.	0	1	2	3
<i>Tanytarsus</i> sp. XIII Rutter	0	2	1	3
<i>Thienemanniella fusca</i> gp.	0	2	0	2
<i>Thienemanniella xena</i> gp.	0	1	0	1
<i>Tribelos jucundum</i>	1	1	24	26
<i>Tvetenia discoloripes</i> gp.	37	79	38	154
<i>Tvetenia paucunca</i>	0	1	1	2

CRUSTACEA

<i>Cambarinae</i>	0	0	1	1
<i>Palaemonetes paludosus</i>	3	13	0	16

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

ALL SITES

Site 1	Site 2	Site 5	TOTAL
--------	--------	--------	-------

ANNELIDA (worms)

Spirosperma sp.	0	1	1	2
Tubificidae	1	1	1	3

MOLLUSCA (snails, clams)

Campeloma sp.	5	10	0	15
Gillia altilis	0	6	4	10

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 1

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Dannella simplex	0	1	0	1
Ephemerella argo	0	1	0	1
Ephemerella invaria/rotunda	1	7	0	8
Hexagenia limbata	0	0	6	6
Neophemera youngi	0	3	0	3
Paraleptophlebia sp.	0	1	0	1
Stenonema modestum/smithae	1	1	1	3
ODONATA (dragonflies)				
Boyeria vinosa	1	0	0	1
Perithemis sp.	0	1	0	1
PLECOPTERA (stoneflies)				
Acroneuria abnormis	0	1	0	1
Isoperla holochlora/lata	0	2	0	2
Isoperla dicala	0	4	1	5
Leuctra sp.	0	1	0	1
Paragnetina kansensis	0	2	0	2
Perlesta placida	0	31	0	31
Pteronarcys dorsata	0	11	0	11
MEGALOPTERA				
Corydalus cornutus	0	2	0	2

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 1

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
TRICHOPTERA (caddisflies)				
Cheumatopsyche sp.	0	7	2	9
Hydropsyche elissoma	0	41	3	44
Lype diversa	0	0	1	1
Macrostemum carolina	0	1	0	1
Micrasema wataga	0	3	0	3
Oecetis sp.	0	1	0	1
Phylocentropus sp.	0	0	4	4
Pycnopsyche sp.	0	0	1	1
Triaenodes sp.	0	1	2	3
COLEOPTERA (beetles)				
Macronychus glabratus	0	3	0	3
Optioservus sp.	0	1	0	1
DIPTERA - misc. (true flies)				
Antocha sp.	0	1	0	1
Atherix lantha	0	1	0	1
Bezzia sp. 2	1	0	2	3
Hexatoma sp.	0	0	1	1
Probezzia sp.	2	0	0	2
Simulium jonesi	0	14	0	14

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

SITE 1

	Sand	Snag	Mud	TOTAL
<hr/>				
DIPTERA- Chironomidae (midges)				
<i>Ablabesmyia janta</i>	0	0	1	1
<i>Brillia flavifrons</i>	0	4	0	4
<i>Clinotanytus pinguis</i>	0	0	6	6
<i>Conchapelopia</i> sp.	0	0	1	1
<i>Cricotopus bicinctus</i>	0	0	1	1
<i>Cryptochironomus fulvus</i> gp.	2	0	0	2
<i>Microtendipes pedellus</i>	0	0	1	1
<i>Paracladopelma undine</i>	0	0	1	1
<i>Parakiefferiella</i> sp. A	0	0	1	1
<i>Parakiefferiella</i> sp.4	0	0	1	1
<i>Parametrioctenemus lundbecki</i>	0	0	1	1
<i>Polypedilum convictum</i>	0	17	2	19
<i>Polypedilum halterale</i>	0	0	1	1
<i>Polypedilum scalaenum</i>	1	0	1	2
<i>Rheocricotopus robacki</i>	0	20	1	21
<i>Rheotanytarsus distinctissimus</i> gp.	0	13	1	14
<i>Rheotanytarsus exiguus</i> gp.	0	0	1	1
<i>Stenochironomus hilaris</i>	0	1	0	1
<i>Tribelos jucundum</i>	0	0	1	1
<i>Tvetenia discoloripes</i> gp.	0	34	3	37
CRUSTACEA				
<i>Palaemonetes paludosus</i>	0	0	3	3
ANNELIDA (worms)				
Tubificidae	0	0	1	1
MOLLUSCA (snails, clams)				
<i>Campeloma</i> sp.	0	0	5	5

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Baetis ephippiatus	0	2	0	2
Dannella simplex	0	0	1	1
Dolania americana	1	0	0	1
Ephemerella argo	0	2	0	2
Ephemerella invaria/rotunda	0	24	0	24
Eurylophella temporalis	0	2	0	2
Neophemera youngi	0	1	0	1
Pseudocloeon dubium	0	2	0	2
Pseudocloeon punctiventris	0	4	0	4
Stenonema modestum/smithae	0	12	0	12
 ODONATA (dragonflies)				
Boyeria vinosa	0	1	0	1
Calopteryx sp.	0	0	1	1
Gomphus (Gomphus) lividus	0	0	3	3
 PLECOPTERA (stoneflies)				
Isoperla bilineata	0	2	0	2
Isoperla holochlora/lata	0	10	0	10
Isoperla dicala	0	1	0	1
Leuctra sp.	0	2	0	2
Paragnetina kansensis	0	2	0	2
Perlesta placida	0	43	0	43
Pteronarcys dorsata	0	11	0	11

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 2

DATE: March 23, 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
TRICHOPTERA (caddisflies)				
Agarodes libalis	0	1	0	1
Cernotina sp.	0	0	1	1
Cheumatopsyche sp.	0	43	1	44
Chimarra sp.	0	1	0	1
Hydropsyche elissoma	0	69	0	69
Hydropsyche sp. A	0	1	0	1
Hydroptila sp.	0	2	0	2
Neureclipsis sp.	0	1	0	1
Oecetis sp.	0	1	0	1
Phylocentropus sp.	1	0	5	6
Triaenodes sp.	0	4	2	6
LEPIDOPTERA (moths)				
Parapoynx obscuralis	0	1	0	1
COLEOPTERA (beetles)				
Ancyronyx variegatus	0	1	0	1
Dineutus discolor	0	5	0	5
Macronychus glabratus	0	1	0	1
Optioservus sp.	0	2	0	2
DIPTERA - misc. (true flies)				
Atherix lantha	0	1	0	1
Empididae	0	1	0	1
Probezzia sp.	0	0	4	4
Simulium jonesi	1	26	0	27
Simulium tuberosum	0	1	0	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
DIPTERA- Chironomidae (midges)				
<i>Ablabesmyia mallochi</i>	0	1	0	1
<i>Clinotanytus pinguis</i>	0	0	1	1
<i>Conchapelopia</i> sp.	0	6	3	9
<i>Cryptochironomus fulvus</i> gp.	3	3	3	9
<i>Microtendipes pedellus</i>	0	0	3	3
<i>Nanocladius</i> sp.	0	0	1	1
<i>Nilotanytus fimbriatus</i>	0	1	0	1
<i>Orthocladius lignicola</i>	0	1	0	1
<i>Pagastiella ostansa</i>	0	0	1	1
<i>Paracladopelma undine</i>	0	0	2	2
<i>Parametrioctenus lundbecki</i>	0	5	2	7
<i>Polypedilum convictum</i>	1	59	0	60
<i>Polypedilum scalaenum</i>	3	1	2	6
<i>Rheocricotopus robacki</i>	0	25	0	25
<i>Rheotanytarsus distinctissimus</i> gp.	0	41	0	41
<i>Rheotanytarsus exiguus</i> gp.	0	1	1	2
<i>Synorthocladius semivirens</i>	0	2	0	2
<i>Tanytarsus glabrescens</i> gp.	0	1	0	1
<i>Tanytarsus</i> sp. XIII Rutter	1	1	0	2
<i>Thienemanniella fusca</i> gp.	0	2	0	2
<i>Thienemanniella xena</i> gp.	0	1	0	1
<i>Tribelos jucundum</i>	0	1	0	1
<i>Tvetenia discoloripes</i> gp.	1	78	0	79
<i>Tvetenia paucunca</i>	0	1	0	1
 CRUSTACEA				
<i>Palaemonetes paludosus</i>	0	7	6	13
 ANNELIDA (worms)				
<i>Spirosperma</i> sp.	0	0	1	1
Tubificidae	1	0	0	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: March 23, 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
MOLLUSCA (snails, clams)				
Campeloma sp.	0	0	10	10
Gillia altilis	0	0	6	6

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: March 23, 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Dannella simplex	1	0	0	1
Dolania americana	2	0	0	2
Ephemerella argo	1	0	0	1
Ephemerella invaria/rotunda	0	1	0	1
Eurylophella temporalis	2	0	0	2
Neophemera youngi	2	0	0	2
Pseudocloeon punctiventris	0	1	0	1
ODONATA (dragonflies)				
Gomphus (Gomphus) lividus	1	1	0	2
PLECOPTERA (stoneflies)				
Haploperla brevis	0	0	1	1
Isoperla dicala	0	10	0	10
Leuctra sp.	0	1	0	1
Paragnetina kansensis	0	3	0	3
Perlesta placida	1	3	0	4
Pteronarcys dorsata	0	1	0	1
HEMIPTERA (true bugs)				
Rhagovelia obesa	0	1	0	1
MEGALOPTERA				
Nigronia serricornis	0	0	1	1

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 5

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
<hr/>				
TRICHOPTERA (caddisflies)				
Cheumatopsyche sp.	0	2	0	2
Hydropsyche elissoma	0	16	0	16
Hydropsyche sp. A	0	1	0	1
Micrasema wataga	1	0	0	1
Phylocentropus sp.	0	0	2	2
Polycentropus sp.	1	0	0	1
Triaenodes sp.	3	0	0	3
COLEOPTERA (beetles)				
Optioservus sp.	0	3	0	3
Sperchopsis tessellatus	0	0	1	1
Stenelmis (larvae)	0	4	1	5
DIPTERA - misc. (true flies)				
Atherix lantha	0	2	0	2
Bezzia sp. 2	0	1	1	2
Empididae	0	4	0	4
Hemerodromia sp. 1	0	2	0	2
Probezzia sp.	1	0	1	2
Palpomyia sp. 4	3	0	0	3
Simulium jonesi	0	9	0	9
Tipula sp.1	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
DIPTERA- Chironomidae (midges)				
<i>Ablabesmyia mallochi</i>	0	0	2	2
<i>Conchapelopia</i> sp.	4	2	2	8
<i>Cryptochironomus fulvus</i> gp.	2	1	1	4
<i>Lopescladius</i> sp.	1	0	0	1
<i>Orthocladius carlatus</i>	0	3	0	3
<i>Orthocladius lignicola</i>	0	1	0	1
<i>Orthocladius</i> sp.	1	5	0	6
<i>Paracladopelma undine</i>	0	0	1	1
<i>Paralauterborniella nigrohalteralis</i>	0	0	2	2
<i>Parametriochnemus lundbecki</i>	4	5	1	10
<i>Polypedilum convictum</i>	10	9	3	22
<i>Polypedilum halterale</i>	0	1	2	3
<i>Polypedilum scalaenum</i>	0	0	6	6
<i>Paramerina</i> sp.	0	2	0	2
<i>Potthastia longimana</i>	0	3	0	3
<i>Procladius</i> sp.	0	2	0	2
<i>Rheocricotopus robacki</i>	3	5	0	8
<i>Rheotanytarsus distinctissimus</i> gp.	23	27	0	50
<i>Stenochironomus hilaris</i>	0	2	0	2
<i>Tanytarsus glabrescens</i> gp.	0	1	1	2
<i>Tanytarsus</i> sp. XIII Rutter	0	1	0	1
<i>Tribelos jucundum</i>	0	0	24	24
<i>Tvetenia discoloripes</i> gp.	0	37	1	38
<i>Tvetenia paucunca</i>	0	1	0	1
CRUSTACEA				
<i>Cambarinae</i>	0	0	1	1
ANNELIDA (worms)				
<i>Spirosperma</i> sp.	0	0	1	1
<i>Tubificidae</i>	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: March 23, 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Sand	Snag	Mud	TOTAL
MOLLUSCA (snails, clams)				
Gillia altilis	0	0	4	4

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Baetis ephippiatus	1	3	0	4
Baetis frondalis	0	1	0	1
Centroptilum sp.	0	1	0	1
Dolania americana	1	3	0	4
Eurylophella sp.	0	1	0	1
Neophemera youngi	1	4	7	12
Paraleptophlebia sp.	3	11	2	16
Stenonema modestum/smithae	3	7	10	20
ODONATA (dragonflies)				
Boyeria vinosa	0	1	3	4
Calopteryx sp.	0	2	1	3
Gomphus (Gomphus) lividus	0	0	1	1
Neurocordulia sp.	1	0	0	1
PLECOPTERA (stoneflies)				
Acroneuria abnormis	6	2	5	13
Isoperla sp.	1	0	0	1
Leuctra sp.	1	1	1	3
Paragnetina kansensis	9	0	6	15
Pteronarcys dorsata	3	1	7	11
HEMIPTERA (true bugs)				
Metrobates hesperius	0	0	1	1
Gerris remigis	0	0	1	1
Ochterus sp.	0	1	0	1
Rhagovelia obesa	0	5	1	6

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
MEGALOPTERA				
Corydalis cornutus	1	0	0	1
Chauliodes rastricornis	0	1	0	1
Nigronia serricornis	0	0	1	1
TRICHOPTERA (caddisflies)				
Agarodes libalis	0	1	0	1
Anisocentropus pyraloides	0	0	1	1
Brachycentrus numerosus	3	1	0	4
Cheumatopsyche sp.	0	1	0	1
Hydropsyche elissoma	5	3	1	9
Lepidostoma sp.	0	1	0	1
Oecetis sp.	0	2	0	2
Ptilostomis sp.	0	1	0	1
Polycentropus sp.	0	1	0	1
Pycnopsyche sp.	0	0	1	1
Triaenodes sp.	5	1	0	6
LEPIDOPTERA (moths)				
Parapoynx obscuralis	0	2	0	2

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM ASSESSMENT.

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

ALL SITES

Site 1	Site 2	Site 5
--------	--------	--------

TOTAL

COLEOPTERA (beetles)

Ancyronyx variegatus	1	0	0	1
Anchytarsus bicolor	1	0	0	1
Copelatus glyphicus	1	0	1	2
Cyphon sp.	0	2	0	2
Dineutus discolor	4	0	2	6
Gyrinus sp. (larvae)	1	0	3	4
Hydaticus sp.	0	0	1	1
Hydroporus sp.	2	3	0	5
Macronychus glabratus	0	2	1	3
Optioservus sp.	0	2	2	4
Oulimnius latisculus	1	0	0	1
Peltodytes sp.	0	1	0	1
Sperchopsis tessellatus	1	1	2	4
Stenelmis sp. (larvae)	2	0	2	4

DIPTERA - misc. (true flies)

Atherix lantha	5	1	3	9
Culex sp.	0	1	1	2
Erioptera /Ormosia sp.	1	1	3	5
Hemerodromia sp. 1	0	1	0	1
Probezzia sp.	1	0	0	1
Simulium jonesi	4	1	3	8
Simulium snowi	0	0	3	3
Simulium tuberosum	0	1	0	1
Tipula (Yamato.) nr strepens	1	0	0	1
Tipula sp.1	0	1	4	5
Tipula (Yamato.) sp.7	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

ALL SITES

Site 1	Site 2	Site 5
--------	--------	--------

TOTAL

DIPTERA- Chironomidae (midges)

Ablabesmyia mallochi	1	2	1	4
Conchapelopia sp.	2	3	1	6
Corynoneura sp. 3	3	0	0	3
Corynoneura sp. 4	0	0	2	2
Cryptochironomus fulvus gp.	1	2	0	3
Microtendipes pedellus	1	0	0	1
Paralauterborniella nigrohalteralis	0	3	0	3
Paramerina sp.	2	8	0	10
Parametriocnemus lundbecki	1	0	1	2
Polypedilum convictum	5	9	13	27
Rheocricotopus robacki	10	0	4	14
Rheosmittia sp.	1	0	0	1
Rheotanytarsus distinctissimus gp.	2	0	1	3
Rheotanytarsus exiguus gp.	5	0	2	7
Stenochironomus hilaris	2	0	0	2
Tanytarsus sp.	1	1	0	2
Tanytarsus sp. XVI Rutter	0	0	1	1
Thienemanniella fusca gp.	3	0	2	5
Thienemanniella xena gp.	0	0	3	3
Tribelos jucundum	0	0	2	2
Tvetenia discoloripes gp.	1	1	1	3
Unniella multivirga	1	2	0	3

CRUSTACEA

Caecidotea sp.	1	0	0	1
Cambarinae	1	2	2	5
Palaemonetes paludosus	2	14	6	22

ANNELIDA (worms)

Pristina sp.	0	0	1	1
Tubificidae	4	4	8	16

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

ALL SITES

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

Site 1	Site 2	Site 5
--------	--------	--------

TOTAL

MOLLUSCA (snails, clams)

Cameloma sp.	1	6	0	7
Gillia altilis	1	1	5	7
Elliptio sp.	0	1	0	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Baetis ephippiatus	0	1	0	1
Dolania americana	0	1	0	1
Neophemera youngi	1	0	0	1
Paraleptophlebia sp.	0	0	3	3
Stenonema modestum/smithae	0	3	0	3
ODONATA (dragonflies)				
Neurocordulia sp.	0	1	0	1
PLECOPTERA (stoneflies)				
Acroneuria abnormis	0	6	0	6
Isoperla sp.	0	1	0	1
Leuctra sp.	0	1	0	1
Paragnetina kansensis	0	9	0	9
Pteronarcys dorsata	1	2	0	3
MEGALOPTERA				
Corydalus cornutus	0	1	0	1
TRICHOPTERA (caddisflies)				
Brachycentrus numerosus	0	1	2	3
Hydropsyche elissoma	1	4	0	5
Triaenodes sp.	0	0	5	5

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 1

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
COLEOPTERA (beetles)				
Anchytarsus bicolor	0	1	0	1
Ancyronyx variegatus	1	0	0	1
Copelatus glyphicus	0	1	0	1
Dineutus discolor	0	4	0	4
Gyrinus sp.	0	0	1	1
Hydroporus sp.	0	1	1	2
Oulimnius laticulus	1	0	0	1
Sperchopsis tessellatus	0	1	0	1
Stenelmis sp. (larvae)	0	2	0	2
DIPTERA - misc. (true flies)				
Atherix lantha	0	5	0	5
Erioptera /Ormosia sp.	0	0	1	1
Probezzia sp.	0	0	1	1
Simulium jonesi	1	3	0	4
Tipula (Yamato.) nr strepens	0	0	1	1
DIPTERA- Chironomidae (midges)				
Ablabesmyia mallochi	0	1	0	1
Conchapelopia sp.	0	2	0	2
Corynoneura sp. 3	0	2	1	3
Cryptochironomus fulvus gp.	0	0	1	1
Microtendipes pedellus	0	0	1	1
Parametriocnemus lundbecki	0	1	0	1
Paramerina sp.	0	2	0	2
Polypedilum convictum	0	5	0	5
Rheocricotopus robacki	0	10	0	10
Rheosmittia sp.	1	0	0	1
Rheotanytarsus distinctissimus gp.	0	2	0	2
Rheotanytarsus exiguus gp.	0	0	5	5
Stenochironomus hilaris	0	2	0	2
Tanytarsus sp.	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
Thienemanniella fusca gp.	0	3	0	3
Tvetenia discoloripes gp.	0	1	0	1
Unniella multivirga	0	0	1	1
CRUSTACEA				
Asellus sp.	0	1	0	1
Cambarinae	0	0	1	1
Palaemonetes paludosus	0	1	1	2
ANNELIDA (worms)				
Tubificidae	0	4	0	4
MOLLUSCA (snails, clams)				
Gillia altilis	0	0	1	1
Campeloma sp.	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Baetis ephippiatus	0	0	3	3
Baetis frondalis	0	0	1	1
Centroptilum sp.	0	0	1	1
Dolania americana	0	0	3	3
Eurylophella sp.	0	1	0	1
Neophemera youngi	0	3	1	4
Paraleptophlebia sp.	1	7	3	11
Stenonema modestum/smithae	0	5	2	7
 ODONATA (dragonflies)				
Boyeria vinosa	0	1	0	1
Calopteryx sp.	0	1	1	2
 PLECOPTERA (stoneflies)				
Acroneuria abnormis	0	2	0	2
Leuctra sp.	0	0	1	1
Pteronarcys dorsata	0	1	0	1
 HEMIPTERA (true bugs)				
Ochterus sp.	0	1	0	1
Rhagovelia obesa	0	0	5	5
 MEGALOPTERA				
Chaulioides rastricornis	0	0	1	1

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 2

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
TRICHOPTERA (caddisflies)				
Agarodes libalis	0	0	1	1
Brachycentrus numerosus	0	1	0	1
Cheumatopsyche sp.	0	1	0	1
Hydropsyche elissoma	0	3	0	3
Lepidostoma sp.	0	1	0	1
Oecetis sp.	0	2	0	2
Polycentropus sp.	0	1	0	1
Ptilostomis sp.	0	0	1	1
Triaenodes sp.	0	0	1	1
LEPIDOPTERA (moths)				
Parapoynx obscuralis	0	0	2	2
COLEOPTERA (beetles)				
Cyphon sp.	1	0	1	2
Hydroporus sp.	0	0	3	3
Macronychus glabratus	0	2	0	2
Optioservus sp.	0	2	0	2
Sperchopsis tessellatus	0	1	0	1
Peltodytes sp.	0	1	0	1
DIPTERA – misc. (true flies)				
Atherix lantha	0	1	0	1
Culex sp.	0	0	1	1
Erioptera /Ormosia sp.	1	0	0	1
Hemerodromia sp. 1	0	1	0	1
Simulium jonesi	1	0	0	1
Simulium tuberosum	0	1	0	1
Tipula sp. 1	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
DIPTERA- Chironomidae (midges)				
<i>Ablabesmyia mallochi</i>	0	1	1	2
<i>Conchapelopia</i> sp.	0	1	2	3
<i>Cryptochironomus fulvus</i> gp.	2	0	0	2
<i>Paralauterborniella nigrohalteralis</i>	1	0	2	3
<i>Paramerina</i> sp.	1	5	2	8
<i>Polypedilum convictum</i>	0	7	2	9
<i>Tanytarsus</i> sp.	1	0	0	1
<i>Tvetenia discoloripes</i> gp.	0	1	0	1
<i>Unniella multivirga</i>	0	1	1	2
CRUSTACEA				
<i>Cambarinae</i>	0	2	0	2
<i>Palaemonetes paludosus</i>	0	7	7	14
ANNELIDA (worms)				
<i>Tubificidae</i>	2	1	1	4
MOLLUSCA (snails, clams)				
<i>Campeloma</i> sp.	1	4	1	6
<i>Gillia atilis</i>	0	0	1	1
<i>Elliptio</i> sp.	0	0	1	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
EPHEMEROPTERA (mayflies)				
Neophemera youngi	0	7	0	7
Paraleptophlebia guttata	1	1	0	2
Stenonema modestum/smithae	1	5	4	10
ODONATA (dragonflies)				
Boyeria vinosa	0	2	1	3
Calopteryx sp.	0	1	0	1
Gomphus (Gomphus) lividus	1	0	0	1
PLECOPTERA (stoneflies)				
Acroneuria abnormis	0	5	0	5
Leuctra sp.	0	0	1	1
Pteronarcys dorsata	0	6	1	7
Paragnetina kansensis	0	5	1	6
HEMIPTERA (true bugs)				
Gerriss remigis	1	0	0	1
Metrobates hesperius	1	0	0	1
Rhagovelia obesa	0	1	0	1
MEGALOPTERA				
Nigronia serricornis	0	1	0	1
TRICHOPTERA (caddisflies)				
Anisocentropus pyraloides	0	1	0	1
Hydropsyche elissoma	0	1	0	1
Pycnopsyche sp.	0	0	1	1

APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 5

DATE: OCTOBER 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
COLEOPTERA (beetles)				
Copelatus glyphicus	0	1	0	1
Dineutus discolor	0	0	2	2
Gyrinus sp. (larvae)	0	2	1	3
Hydaticus sp.	0	1	0	1
Macronychus glabratus	0	0	1	1
Optioservus sp.	0	2	0	2
Sperchopsis tessellatus	0	2	0	2
Stenelmis sp. (larvae)	0	0	2	2
DIPTERA - misc. (true flies)				
Atherix lantha	0	2	1	3
Culex sp.	0	0	1	1
Erioptera /Ormosia sp.	0	0	3	3
Simulium jonesi	0	1	2	3
Simulium snowi	0	1	2	3
Tipula sp.1	0	3	1	4
Tipula (Yamato.) sp.7	0	1	0	1
DIPTERA- Chironomidae (midges)				
Ablabesmyia mallochi	0	0	1	1
Conchapelopia sp.	0	0	1	1
Corynoneura sp. 4	0	1	1	2
Parametriocnemus lundbecki	0	0	1	1
Polypedilum convictum	0	1	12	13
Rheocricotopus robacki	0	0	4	4
Rheotanytarsus distinctissimus gp.	0	0	1	0
Rheotanytarsus exiguus gp.	0	2	0	2
Tanytarsus sp. XVI Rutter	0	0	1	1
Thienemanniella fusca gp.	0	0	2	2
Thienemanniella xena gp.	0	1	2	3
Tribelos jucundum	0	0	2	2
Tvetenia discoloripes gp.	0	1	0	1

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	sand	snag	mud	TOTAL
<hr/>				
CRUSTACEA				
Cambarinae	1	1	0	2
Palaemonetes paludosus	0	6	0	6
ANNELIDA (worms)				
Pristina sp.	1	0	0	1
Tubificidae	0	5	3	8
MOLLUSCA (snails, clams)				
Gillia altilis	1	0	4	5

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 1991

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Baetis propinquus	2	0	0	2
Dannella simplex	1	0	0	1
Dolania americana	0	0	2	2
Ephemerella argo	1	1	2	4
Ephemerella dorothea	3	3	2	8
Ephemerella catawba gp.	4	0	3	7
Hexagenia limbata	3	0	0	3
Neophemera youngi	1	0	2	3
Pseudocloeon bimaculatus	0	0	1	1
Pseudocloeon punctiventris	1	0	1	2
Stenonema modestum/smithae	6	0	7	13
ODONATA (dragonflies)				
Boyeria vinosa	0	0	1	1
Calopteryx sp.	1	0	0	1
Progomphus sp.	0	1	0	1
Tetragoneuria costalis/cynosura	0	0	1	1
PLECOPTERA (stoneflies)				
Acroneuria abnormis	3	0	1	4
Acroneuria mela	0	0	1	1
Amphinemura delosa	0	1	0	1
Isoperla lata/holochlora	0	2	0	2
Isoperla dicala	16	1	11	28
Leuctra sp.	0	0	4	4
Paragnetina kansensis	6	3	6	15
Perlesta placida	10	6	21	37
Pteronarcys dorsata	2	0	0	2

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Site 1	Site 2	Site 5	TOTAL
HEMIPTERA (true bugs)				
Rhagovelia obesa	1	0	0	1
MEGALOPTERA				
Nigronia serricornis	0	0	1	1
TRICHOPTERA (caddisflies)				
Cheumatopsyche sp.	10	2	10	22
Hydropsyche elissoma	14	8	15	37
Micrasema wataga	1	0	1	2
Neureclipsis sp.	0	0	1	1
Phylocentropus sp.	0	3	1	4
Triaenodes sp.	1	1	0	2
COLEOPTERA (beetles)				
Ancyronyx variegatus	0	1	0	1
Dineutus assimilis	0	0	1	1
Dineutus discolor	0	0	2	2
Gonielmis dietrichi	1	0	0	1
Hydroporus sp.	0	0	1	1
Macronychus glabratus	1	1	2	4
Stenelmis sp. (larvae)	1	0	2	3

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

Site 1	Site 2	Site 5
--------	--------	--------

TOTAL

DIPTERA – misc. (true flies)

Antocha sp.	1	0	0	1
Atherix lantha	1	0	1	2
Bezzia sp.	0	0	1	1
Dolichopodidae	1	0	0	1
Erioptera /Ormosia sp.	0	1	0	1
Hemerodromia sp. 1	3	1	7	11
Hemerodromia sp. 2	0	1	0	1
Probezzia sp.	1	8	0	9
Stilobezzia sp.	0	1	0	1
Simulium jonesi	0	3	7	10
Tipula sp. A	0	0	1	1
Tipula (Yamatotipula) sp.7	0	0	1	1
Wiedemannia sp.	1	0	1	2

DIPTERA– Chironomidae (midges)

Ablabesmyia mallochi	2	0	2	4
Brillia flavifrons	0	3	2	5
Cladotanytarsus sp. A	2	1	2	5
Cricotopus bicinctus	6	0	5	11
Cryptochironomus fulvus gp.	1	7	3	11
Cryptochironomus blarina	0	1	1	2
Kloosia sp.	0	2	1	3
Lauterborniella agrayloides	0	0	1	1
Meropelopia americana	3	3	7	13
Microtendipes nr. rydalensis	0	1	0	1
Orthocladius carlatus	4	2	5	11
Orthocladius annectens	1	0	0	1
Pagastiella ostansa	0	3	0	3
Paramerina sp.	0	0	4	4
Paracladopelma loganae	0	1	0	1
Paracladopelma undine	1	2	1	4
Parakiefferiella sp. A	1	1	0	2
Paralauterborniella nigrohalteralis	0	9	2	11

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 1991

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Site 1	Site 2	Site 5	TOTAL
<i>Parametriocnemus lundbecki</i>	1	2	10	13
<i>Polypedilum convictum</i>	15	14	31	60
<i>Polypedilum fallax</i>	0	0	1	1
<i>Polypedilum halterale</i>	0	12	3	15
<i>Polypedilum scalaenum</i>	0	8	2	10
<i>Potthastia longimana</i>	1	0	0	1
<i>Rheocricotopus robacki</i>	6	4	28	38
<i>Rheotanytarsus distinctissimus</i> gp.	11	9	3	23
<i>Rheotanytarsus exiguus</i> gp.	1	17	3	21
<i>Synorthocladus semivirens</i>	1	0	1	2
<i>Tanytarsus glabrescens</i> gp.	1	1	0	2
<i>Tanytarsus</i> sp. I Cantrell	1	2	0	3
<i>Tanytarsus</i> sp. IX Rutter	1	0	0	1
<i>Tanytarsus</i> sp. XII Rutter	0	2	0	2
<i>Tanytarsus</i> sp. XVI Rutter	1	3	2	6
<i>Thienemanniella fusca</i> gp.	2	0	2	4
<i>Thienemanniella xena</i> gp.	1	0	0	1
<i>Tribelos jucundum</i>	1	5	3	9
<i>Tvetenia discoloripes</i> gp.	8	10	9	27
<i>Tvetenia paucunca</i>	2	0	2	4
<i>Unniella multivirga</i>	1	1	0	2
<i>Xylotopus par</i>	1	0	0	1
CRUSTACEA				
<i>Palaemonetes paludosus</i>	3	4	8	15
ANNELIDA (worms)				
Tubificidae	0	2	2	4

**APPENDIX A : LIST OF SPECIES FOR QUALITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: March 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Site 1	Site 2	Site 5	TOTAL
MOLLUSCA (snails, clams)				
Gillia altilis	1	1	0	2
Campeloma sp.	2	7	4	13
ACARI				
Hydracarina	1	0	0	1

March 1991 qualitative macroinvertebrate samples were not separated by habitat.

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE
IN-STREAM ASSESSMENT.**

DATE: April 20, 1990
 CLIENT: S.R.S.
 STREA Upper Three Runs Creek
 LOCATION: Aiken Co. SC

ALL SITES

Site 1	Site 2	Site 3	Site 4	Site 5	TOTAL
--------	--------	--------	--------	--------	-------

EPHEMEROPTERA (mayflies)

Baetis intercalaris	0	0	1	0	0	1
Ephemerella argo	0	3	5	2	2	12
Ephemerella dorothea	1	3	9	1	2	16
Ephemerella invaria/rotunda	0	6	13	9	13	41
Heptagenia flavescens	0	1	0	2	1	4
Neophemera youngi	0	0	0	0	2	2
Pseudocloeon punctiventris	0	1	1	0	0	2
Stenonema modestum/smithae	0	1	6	0	1	8

PLECOPTERA (stoneflies)

Acroneuria abnormis	4	11	12	6	5	38
Acroneuria mela	1	0	1	2	4	8
Helopicus subvarians	1	0	0	0	0	1
Isoperla bilineata	11	0	0	0	0	11
Isoperla lata/holochlora	0	0	0	0	1	1
Isoperla dicala	3	6	8	8	11	36
Leuctra sp.	0	0	0	0	1	1
Paragnetina kansensis	8	10	9	6	5	38
Perlesta placida	5	4	25	18	24	76
Pteronarcys dorsata	4	6	8	6	18	42

MEGALOPTERA

Corydalus cornutus	1	1	0	0	0	2
--------------------	---	---	---	---	---	---

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE
IN-STREAM ASSESSMENT.**

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

Site 1	Site 2	Site 3	Site 4	Site 5	TOTAL
--------	--------	--------	--------	--------	-------

TRICHOPTERA (caddisflies)

Brachycentrus numerosus	0	0	5	2	6	13
Cheumatopsyche sp.	5	3	8	3	7	26
Chimarra sp.	0	0	1	0	0	1
Hydropsyche elissoma	34	25	53	61	37	210
Hydroptila sp.	0	0	0	0	1	1
Lype diversa	1	0	0	0	0	1
Micrasema wataga	0	0	1	0	0	1

LEPIDOPTERA (moths)

Parapoynx obscuralis	1	0	0	0	0	1
----------------------	---	---	---	---	---	---

COLEOPTERA (beetles)

Anchytarsus bicolor	0	0	0	0	1	1
Ancyronyx variegatus	0	1	0	0	0	1
Dineutus (larvae)	0	1	0	0	1	2
Gonielmis dietrichi	0	0	1	0	0	1
Macronychus glabratus	3	13	2	2	2	22
Optioservus sp.	1	0	0	2	0	3
Stenelmis crenata	0	4	3	6	0	13
Stenelmis convexula	2	4	0	6	0	12
Stenelmis (larvae)	0	3	7	3	5	18

DIPTERA - misc. (true flies)

Atherix lantha	0	0	1	0	0	1
Clinocera sp.	0	0	0	0	2	2
Chelifera sp.	0	0	0	1	0	1
Hemerodromia	0	0	6	0	2	8
Simulium jonesi	13	21	23	80	16	153
Simulium tuberosum	0	1	0	2	0	3
Tipula (Yamato.) sp.7	0	2	0	0	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE
IN-STREAM ASSESSMENT.**

ALL SITES

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

Site 1	Site 2	Site 3	Site 4	Site 5	TOTAL
--------	--------	--------	--------	--------	-------

DIPTERA- Chironomidae (midges)

<i>Brillia flavifrons</i>	0	4	1	0	7	12
<i>Cladotanytarsus</i> sp.	1	0	0	0	0	1
<i>Conchapelopia</i> sp.	1	1	2	2	0	6
<i>Corynoneura</i> sp. 3	0	0	0	0	2	2
<i>Cryptochironomus fulvus</i> gp.	0	1	0	0	0	1
<i>Orthocladius</i> sp.	0	1	0	0	0	1
<i>Pagastiella ostanta</i>	0	1	0	0	0	1
<i>Parakiefferiella</i> sp. A	1	0	0	4	0	5
<i>Parametriocnemus lundbecki</i>	1	0	2	0	5	8
<i>Polypedilum convictum</i>	8	14	22	22	37	103
<i>Polypedilum fallax</i>	0	0	0	1	1	2
<i>Polypedilum illinoense</i>	0	0	1	0	0	1
<i>Rheocricotopus robacki</i>	1	6	2	0	8	17
<i>Rheotanytarsus distinctissimus</i> gp.	2	3	4	6	5	20
<i>Rheotanytarsus exiguus</i> gp.	1	0	0	0	0	1
<i>Tanytarsus glabrescens</i> gp.	0	2	3	0	4	9
<i>Tanytarsus</i> sp. XVI Rutter	0	1	0	0	0	1
<i>Thienemanniella fusca</i> gp.	0	1	1	0	0	2
<i>Thienemanniella xena</i> gp.	0	4	2	3	6	15
<i>Tvetenia discoloripes</i> gp.	9	24	19	21	30	103
<i>Tvetenia paucunca</i>	1	0	5	0	3	9

SITE 1 **DATE:** April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

SITE 1

DATE: April 20, 1990

CLIENT: S.R.S.

STREA Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
EPHEMEROPTERA (mayflies)						
<i>Ephemerella dorothea</i>	0	0	1	0	0	1
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	0	3	0	0	1	4
<i>Acroneuria mela</i>	1	0	0	0	0	1
<i>Helopicus subvarians</i>	0	0	0	1	0	1
<i>Isoperla bilineata</i>	0	0	1	10	0	11
<i>Isoperla dicala</i>	0	0	0	0	3	3
<i>Paragnetina kansensis</i>	0	3	1	3	1	8
<i>Perlesta placida</i>	0	3	2	0	0	5
<i>Pteronarcys dorsata</i>	0	0	4	0	0	4
MEGALOPTERA						
<i>Corydalus cornutus</i>	0	0	1	0	0	1
TRICHOPTERA (caddisflies)						
<i>Cheumatopsyche</i> sp.	0	1	0	1	3	5
<i>Hydropsyche elissoma</i>	0	6	21	1	6	34
<i>Lype diversa</i>	1	0	0	0	0	1
LEPIDOPTERA (moths)						
<i>Parapoynx obscuralis</i>	0	1	0	0	0	1
COLEOPTERA (beetles)						
<i>Macronychus glabratus</i>	1	1	0	0	1	3
<i>Optioservus</i> sp.	0	0	1	0	0	1
<i>Stenelmis convexula</i>	0	2	0	0	0	2
DIPTERA – misc. (true flies)						

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
Simulium jonesi	5	0	7	1	0	13
DIPTERA- Chironomidae (midges)						
Cladotanytarsus sp.	0	1	0	0	0	1
Conchapelopia sp.	0	1	0	0	0	1
Parakiefferiella sp. A	1	0	0	0	0	1
Parametriocnemus lundbecki	1	0	0	0	0	1
Polypedilum convictum	0	1	5	2	0	8
Rheocricotopus robacki	1	0	0	0	0	1
Rheotanytarsus distinctissimus gp.	0	0	1	0	1	2
Rheotanytarsus exiguus gp.	0	0	1	0	0	1
Tvetenia discoloripes gp.	0	2	6	0	1	9
Tvetenia paucunca	0	0	1	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Ephemerella argo</i>	0	1	0	2	0	3
<i>Ephemerella dorothea</i>	0	0	1	2	0	3
<i>Ephemerella invaria/rotunda</i>	0	0	2	2	2	6
<i>Heptagenia flavescens</i>	0	0	1	0	0	1
<i>Pseudocloeon punctiventris</i>	0	0	0	1	0	1
<i>Stenonema modestum/smithae</i>	0	0	0	1	0	1
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	3	3	3	2	0	11
<i>Isoperla dicala</i>	0	1	3	2	0	6
<i>Paragnetina kansensis</i>	4	3	1	1	1	10
<i>Perlesta placida</i>	0	0	2	2	0	4
<i>Pteronarcys dorsata</i>	0	0	1	5	0	6
MEGALOPTERA						
<i>Corydalus cornutus</i>	0	0	0	1	0	1
TRICHOPTERA (caddisflies)						
<i>Cheumatopsyche</i> sp.	0	1	2	0	0	3
<i>Hydropsyche elissoma</i>	1	3	7	11	3	25
COLEOPTERA (beetles)						
<i>Ancyronyx variegatus</i>	1	0	0	0	0	1
<i>Dineutus</i> (larvae)	0	0	0	0	1	1
<i>Macronychus glabratus</i>	1	2	8	2	0	13
<i>Stenelmis crenata</i>	0	0	3	1	0	4
<i>Stenelmis convexula</i>	0	0	4	0	0	4
<i>Stenelmis</i> (larvae)	1	0	2	0	0	3

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA – misc. (true flies)						
Simulium jonesi	0	5	3	7	6	21
Simulium tuberosum	0	0	1	0	0	1
Tipula (Yamato.) sp.7	0	1	1	0	0	2
DIPTERA– Chironomidae (midges)						
Brillia flavifrons	1	0	1	2	0	4
Conchapelopia sp.	1	0	0	0	0	1
Cryptochironomus fulvus gp.	0	0	0	1	0	1
Orthocladius sp.	0	0	1	0	0	1
Pagastiella ostansa	1	0	0	0	0	1
Polypedilum convictum	3	3	4	3	1	14
Rheocricotopus robacki	2	0	4	0	0	6
Rheotanytarsus distinctissimus gp.	0	1	1	1	0	3
Tanytarsus glabrescens gp.	2	0	0	0	0	2
Tanytarsus sp. XVI Rutter	1	0	0	0	0	1
Thienemanniella fusca gp.	0	0	1	0	0	1
Thienemanniella xena gp.	0	1	1	2	0	4
Tvetenia discoloripes gp.	1	2	14	6	1	24

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 3

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
EPHEMEROPTERA (mayflies)						
<i>Baetis intercalaris</i>	0	0	0	1	0	1
<i>Ephemerella argo</i>	1	1	1	1	1	5
<i>Ephemerella dorothea</i>	4	0	0	3	2	9
<i>Ephemerella invaria/rotunda</i>	4	1	2	4	2	13
<i>Pseudocloeon punctiventris</i>	0	0	0	0	1	1
<i>Stenonema modestum/smithae</i>	1	0	2	1	2	6
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	2	1	1	5	3	12
<i>Acroneuria mela</i>	0	0	0	0	1	1
<i>Isoperla dicala</i>	0	1	5	0	2	8
<i>Paragnetina kansensis</i>	3	5	0	1	0	9
<i>Perlenta placida</i>	7	1	4	10	3	25
<i>Pteronarcys dorsata</i>	2	1	1	3	1	8
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	1	0	1	2	1	5
<i>Cheumatopsyche</i> sp.	2	0	3	3	0	8
<i>Chimarra</i> sp.	0	0	0	1	0	1
<i>Hydropsyche elissoma</i>	20	3	13	8	9	53
<i>Micrasema wataga</i>	0	0	0	1	0	1
COLEOPTERA (beetles)						
<i>Gonielmis dietrichi</i>	0	0	0	0	1	1
<i>Macronychus glabratus</i>	0	1	0	0	1	2
<i>Stenelmis crenata</i>	0	1	1	0	1	3
<i>Stenelmis</i> (larva)	2	0	2	1	2	7

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 3

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA - misc. (true flies)						
Atherix lantha	1	0	0	0	0	1
Hemerodromia sp. 1	0	0	0	6	0	6
Simulium jonesi	1	0	2	6	14	23
 DIPTERA- Chironomidae (midges)						
Brillia flavifrons	1	0	0	0	0	1
Conchapelopia sp.	0	0	1	0	1	2
Parametriocnemus lundbecki	1	0	1	0	0	2
Polypedilum convictum	6	0	5	9	2	22
Polypedilum illinoense	0	0	0	0	1	1
Rheocricotopus robacki	1	0	1	0	0	2
Rheotanytarsus distinctissimus gp.	2	1	0	1	0	4
Tanytarsus glabrescens gp.	1	0	2	0	0	3
Thienemanniella fusca gp.	0	0	0	0	1	1
Thienemanniella xena gp.	2	0	0	0	0	2
Tvetenia discoloripes gp.	5	0	10	1	3	19
Tvetenia paucunca	2	0	0	1	2	5

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 4

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Ephemerella argo</i>	0	1	1	0	0	2
<i>Ephemerella dorothea</i>	0	1	0	0	0	1
<i>Ephemerella invaria/rotunda</i>	0	2	7	0	0	9
<i>Heptagenia flavescens</i>	0	0	1	1	0	2
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	2	2	1	0	1	6
<i>Acroneuria mela</i>	0	1	0	1	0	2
<i>Isoperla dicala</i>	2	1	3	0	2	8
<i>Paragnetina kansensis</i>	2	1	2	1	0	6
<i>Perlesta placida</i>	5	4	6	2	1	18
<i>Pteronarcys dorsata</i>	1	0	1	1	3	6
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	0	1	0	0	1	2
<i>Cheumatopsyche</i> sp.	0	2	0	0	1	3
<i>Hydropsyche elissoma</i>	10	17	27	1	6	61
COLEOPTERA (beetles)						
<i>Macronychus glabratus</i>	0	0	1	0	1	2
<i>Optioservus</i> sp.	1	0	1	0	0	2
<i>Stenelmis crenata</i>	0	0	3	1	2	6
<i>Stenelmis convexula</i>	1	0	2	3	0	6
<i>Stenelmis</i> (larvae)	0	0	0	2	1	3
DIPTERA - misc. (true flies)						
<i>Chelifera</i> sp.	1	0	0	0	0	1
<i>Simulium jonesi</i>	1	12	64	1	2	80
<i>Simulium tuberosum</i>	0	0	2	0	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 4

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	0	0	0	0	2	2
Parakiefferiella sp. A	0	0	2	0	2	4
Polypedilum convictum	5	2	8	3	4	22
Polypedilum fallax	0	0	0	0	1	1
Rheotanytarsus distinctissimus gp.	0	1	1	0	4	6
Thienemanniella xena gp.	0	0	1	0	2	3
Tvetenia discoloripes gp.	1	4	10	3	3	21

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Ephemerella argo</i>	2	0	0	0	0	2
<i>Ephemerella dorothea</i>	0	2	0	0	0	2
<i>Ephemerella invaria/rotunda</i>	10	0	3	0	0	13
<i>Heptagenia flavescens</i>	0	0	0	0	1	1
<i>Neophemera youngi</i>	2	0	0	0	0	2
<i>Stenonema modestum/smithae</i>	0	0	0	1	0	1
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	1	2	0	2	0	5
<i>Acroneuria mela</i>	0	1	2	1	0	4
<i>Isoperla holochlora/lata</i>	1	0	0	0	0	1
<i>Isoperla dicala</i>	6	2	2	1	0	11
<i>Leuctra</i> sp.	1	0	0	0	0	1
<i>Paragnetina kansensis</i>	2	1	1	1	0	5
<i>Perlesta placida</i>	10	6	5	3	0	24
<i>Pteronarcys dorsata</i>	6	9	3	0	0	18
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	4	1	1	0	0	6
<i>Cheumatopsyche</i> sp.	5	0	1	1	0	7
<i>Hydropsyche elissoma</i>	11	9	11	6	0	37
<i>Hydroptila</i> sp.	0	1	0	0	0	1
COLEOPTERA (beetles)						
<i>Anchytarsus bicolor</i>	0	0	1	0	0	1
<i>Dineutus</i> sp. (larvae)	1	0	0	0	0	1
<i>Macronychus glabratus</i>	1	0	1	0	0	2
<i>Stenelmis</i> (larvae)	2	2	1	0	0	5

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: April 20, 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA – misc. (true flies)						
<i>Clinocera</i> sp.	0	2	0	0	0	2
<i>Hemerodromia</i> sp. 1	0	2	0	0	0	2
<i>Simulium jonesi</i>	11	1	2	2	0	16
 DIPTERA– Chironomidae (midges)						
<i>Brillia flavifrons</i>	4	2	0	0	1	7
<i>Corynoneura</i> sp. 3	0	2	0	0	0	2
<i>Parametriocnemus lundbecki</i>	0	4	0	0	1	5
<i>Polypedilum convictum</i>	20	10	2	3	2	37
<i>Polypedilum fallax</i>	0	0	0	0	1	1
<i>Rheocricotopus robacki</i>	3	4	1	0	0	8
<i>Rheotanytarsus distinctissimus</i> gp.	0	3	2	0	0	5
<i>Tanytarsus glabrescens</i> gp.	1	1	2	0	0	4
<i>Thienemanniella xena</i> gp.	0	5	1	0	0	6
<i>Tvetenia discoloripes</i> gp.	6	9	5	9	1	30
<i>Tvetenia paucunca</i>	1	2	0	0	0	3

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM ASSESSMENT.

DATE: July 1990
 CLIENT: S.R.S.
 STREA Upper Three Runs Creek
 LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Baetis sp.	0	0	1	1
Dolania americana	0	0	1	1
Heptagenia flavescens	6	3	7	16
Neophemera youngi	0	0	1	1
Pseudocloeon dubium	0	0	1	1
Stenonema modestum/smithae	6	6	0	12
Tricorythodes sp.	1	2	0	3
PLECOPTERA (stoneflies)				
Acroneuria abnormis	6	5	8	19
Leuctra sp.	0	1	1	2
Paragnetina kansensis	16	1	4	21
Perlesta placida	0	1	3	4
Pteronarcys dorsata	5	2	8	15
MEGALOPTERA				
Corydalus cornutus	2	6	0	8
Nigronia serricornis	0	2	0	2
TRICHOPTERA (caddisflies)				
Brachycentrus numerosus	3	1	1	5
Cheumatopsyche sp.	14	10	9	33
Chimarra sp.	2	0	1	3
Diplectrona modesta	0	2	0	2
Hydropsyche elissoma	10	26	14	50
Hydroptila sp.	2	1	0	3
Lype diversa	1	2	0	3
Micrasema wataga	3	2	0	5
Nyctiophylax sp.	1	0	0	1
Phylocentropus sp.	1	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: July 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
LEPIDOPTERA (moths)				
Parapoynx obscuralis	0	3	0	3
COLEOPTERA (beetles)				
Ancyronyx variegatus	1	0	1	2
Dineutus (larvae)	1	0	0	1
Macronychus glabratus	4	7	3	14
Stenelmis crenata	6	6	8	20
Stenelmis convexula	5	6	13	24
Stenelmis (larvae)	2	1	2	5
DIPTERA - misc. (true flies)				
Atherix lantha	2	1	1	4
Hemerodromia sp. 1	3	2	2	7
Simulium jonesi	4	4	4	12
Simulium tuberosum	1	0	1	2
Telmatoscopus sp.	1	0	0	1
DIPTERA- Chironomidae (midges)				
Cricotopus vierriensis	0	1	1	2
Cryptochironomus fulvus gp.	0	0	1	1
Nanocladius sp.	0	1	0	1
Parametriocnemus lundbecki	0	0	2	2
Polypedilum convictum	7	2	2	11
Polypedilum fallax	0	0	1	1
Polypedilum scalaenum	0	1	0	1
Pseuchironomus sp.	0	1	0	1
Rheocricotopus robacki	1	1	1	3
Rheotanytarsus distinctissimus gp.	1	1	1	3
Stenochironomus hilaris	2	0	2	4

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: July 1990

CLIENT: S.R.S.

STREA Upper Three Runs Creek

LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
Tanytarsus glabrescens gp.	1	0	0	1
Tvetenia discoloripes gp.	0	1	2	3
ANNELIDA (worms)				
Tubificidae	0	1	0	1
MOLLUSCA (snails, clams)				
Ferrissia sp.	1	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: July 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Heptagenia flavescens	0	0	4	0	2	6
Stenonema modestum/smithae	5	1	0	0	0	6
Tricorythodes sp.	0	1	0	0	0	1
PLECOPTERA (stoneflies)						
Acroneuria abnormis	1	2	0	1	2	6
Paragnetina kansensis	0	0	3	9	4	16
Pteronarcys dorsata	0	2	2	1	0	5
MEGALOPTERA						
Corydalus cornutus	0	2	0	0	0	2
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	0	2	0	1	0	3
Cheumatopsyche sp.	5	1	4	0	4	14
Chimarra sp.	1	1	0	0	0	2
Hydropsyche elissoma	2	2	4	0	2	10
Hydroptila sp.	1	1	0	0	0	2
Lype diversa	0	1	0	0	0	1
Micrasema wataga	0	3	0	0	0	3
Nyctiophylax sp.	1	0	0	0	0	1
Phylocentropus sp.	1	0	0	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: July 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
COLEOPTERA (beetles)						
Ancyronyx variegatus	0	1	0	0	0	1
Dineutus (larvae)	1	0	0	0	0	1
Macronychus glabratus	0	2	1	0	1	4
Stenelmis crenata	0	2	2	2	0	6
Stenelmis convexula	0	1	4	0	0	5
Stenelmis (larvae)	0	1	0	0	1	2
DIPTERA – misc. (true flies)						
Atherix lantha	0	1	0	0	1	2
Hemerodromia sp. 1	1	1	1	0	0	3
Simulium jonesi	1	2	0	0	1	4
Simulium tuberosum	0	0	1	0	0	1
Telmatoscopus sp.	0	1	0	0	0	1
DIPTERA– Chironomidae (midges)						
Polypedilum convictum	0	6	0	0	1	7
Rheocricotopus robacki	0	1	0	0	0	1
Rheotanytarsus distinctissimus gp.	0	1	0	0	0	1
Stenochironomus hiliaris	0	2	0	0	0	2
Tanytarsus glabrescens gp.	0	1	0	0	0	1
MOLLUSCA (snails, clams)						
Ferrissia sp.	1	0	0	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: July 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Heptagenia flavescens	0	1	0	1	1	3
Stenonema modestum/smithae	0	2	0	1	3	6
Tricorythodes sp.	0	0	1	0	1	2
PLECOPTERA (stoneflies)						
Acroneuria abnormis	2	0	2	1	0	5
Leuctra sp.	0	1	0	0	0	1
Paragnetina kansensis	0	0	1	0	0	1
Perlesta placida	0	0	1	0	0	1
Pteronarcys dorsata	0	0	2	0	0	2
MEGALOPTERA						
Corydalus cornutus	1	3	2	0	0	6
Nigronia serricornis	0	1	1	0	0	2
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	0	0	0	0	1	1
Cheumatopsyche sp.	0	0	0	2	8	10
Diplectrona modesta	0	1	1	0	0	2
Hydropsyche elissoma	2	4	1	5	14	26
Hydroptila sp.	0	0	0	1	0	1
Lype diversa	1	0	0	0	1	2
Micrasema wataga	0	0	2	0	0	2
LEPIDOPTERA (moths)						
Parapoynx obscuralis	0	0	2	0	1	3

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: July 1990

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
COLEOPTERA (beetles)						
Macronychus glabratus	0	3	0	3	1	7
Stenelmis crenata	3	0	1	1	1	6
Stenelmis markeli	0	2	1	3	0	6
Stenelmis (larvae)	0	0	1	0	0	1
DIPTERA – misc. (true flies)						
Atherix lantha	0	0	0	1	0	1
Hemerodromia sp.1	0	0	1	0	1	2
Simulium jonesi	0	1	1	1	1	4
DIPTERA– Chironomidae (midges)						
Cricotopus vierriensis	1	0	0	0	0	1
Nanocladius sp.	0	1	0	0	0	1
Polypedilum convictum	0	1	1	0	0	2
Polypedilum scalaenum	0	1	0	0	0	1
Pseudochironomus sp.	0	0	0	1	0	1
Rheocricotopus robacki	0	0	1	0	0	1
Rheotanytarsus distinctissimus gp.	1	0	0	0	0	1
Tvetenia discoloripes gp.	0	0	1	0	0	1
ANNELIDA (worms)						
Tubificidae	0	1	0	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: July 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Baetis sp.	0	0	0	0	1	1
Dolania americana	0	0	0	1	0	1
Heptagenia flavescens	1	3	3	0	0	7
Neophemera youngi	0	0	0	0	1	1
Pseudocloeon dubium	0	0	1	0	0	1
PLECOPTERA (stoneflies)						
Acroneuria abnormis	3	1	0	2	2	8
Leuctra sp.	0	0	0	0	1	1
Paragnetina kansensis	1	1	1	0	1	4
Perlesta placida	0	0	0	2	1	3
Pteronarcys dorsata	0	0	3	4	1	8
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	0	0	1	0	0	1
Cheumatopsyche sp.	0	0	1	4	4	9
Chimarra sp.	0	0	0	1	0	1
Hydropsyche elissoma	1	2	0	9	2	14
COLEOPTERA (beetles)						
Ancyronyx variegatus	0	0	0	1	0	1
Macronychus glabratus	1	1	0	1	0	3
Stenelmis crenata	2	1	0	2	3	8
Stenelmis convexula	3	0	5	3	2	13
Stenelmis (larvae)	1	0	0	1	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: July 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA - misc. (true flies)						
Atherix lantha	0	1	0	0	0	1
Hemerodromia sp. 1	0	1	0	0	1	2
Simulium jonesi	0	1	1	1	1	4
Simulium tuberosum	0	0	0	1	0	1
DIPTERA- Chironomidae (midges)						
Cricotopus vierriensis	0	0	0	0	1	1
Cryptochironomus fulvus gp.	0	0	0	0	1	1
Parametriocnemus lundbecki	0	0	0	2	0	2
Polypedilum convictum	0	0	0	0	2	2
Polypedilum fallax	1	0	0	0	0	1
Rheocricotopus robacki	0	0	0	1	0	1
Rheotanytarsus distinctissimus gp.	0	0	0	1	0	1
Stenochironomus hilaris	0	0	0	2	0	2
Tvetenia discoloripes gp.	0	1	0	1	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Heptagenia flavescens	0	1	0	1
Neophemera youngi	0	7	1	8
Paraleptophlebia sp.	0	1	0	1
Stenonema modestum/smithae	1	5	2	8
PLECOPTERA (stoneflies)				
Acroneuria abnormis	3	6	7	16
Paragnetina kansensis	10	7	9	26
Perlesta placida	0	1	0	1
Pteronarcys dorsata	3	5	5	13
MEGALOPTERA				
Corydalus cornutus	0	0	1	1
TRICHOPTERA (caddisflies)				
Brachycentrus numerosus	3	10	5	18
Cheumatopsyche sp.	2	2	0	4
Hydropsyche elissoma	26	20	14	60
Lype diversa	4	1	0	5
Micrasema rusticum	1	0	0	1
Polycentropus sp.	2	0	0	2
COLEOPTERA (beetles)				
Macronychus glabratus	1	3	1	5
Stenelmis sp. 3	0	1	0	1
Stenelmis crenata	3	2	5	10
Stenelmis convexula	1	0	0	1
Stenelmis sp. (larvae)	0	2	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
DIPTERA - misc. (true flies)				
Atherix lantha	4	2	0	6
Simulium jonesi	19	13	9	41
DIPTERA- Chironomidae (midges)				
Conchapelopia sp.	0	1	0	1
Corynoneura nr. taris	3	4	0	7
Corynoneura sp. 3	1	0	2	3
Paramerina sp.	1	0	0	1
Parakiefferiella sp. A	1	0	0	1
Polypedilum convictum	8	8	5	21
Polypedilum fallax	0	1	1	2
Polypedilum illinoense	0	1	0	1
Rheocricotopus robacki	11	20	3	34
Rheotanytarsus distinctissimus gp.	3	5	2	10
Rheotanytarsus exiguus gp.	2	1	0	3
Tanytarsus sp.	1	0	2	3
Thienemanniella fusca gp.	5	7	4	16
Thienemanniella xena gp.	2	1	1	4
Tvetenia discoloripes gp.	0	2	1	3
ACARI				
Hydracarina	0	0	1	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
EPHEMEROPTERA (mayflies)						
Stenonema modestum/smithae	0	0	1	0	0	1
PLECOPTERA (stoneflies)						
Acroneuria abnormis	0	3	0	0	0	3
Paragnetina kansensis	2	2	1	2	3	10
Pteronarcys dorsata	2	0	0	1	0	3
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	1	1	1	0	0	3
Cheumatopsyche sp.	1	1	0	0	0	2
Hydropsyche elissoma	15	2	9	0	0	26
Lype diversa	0	1	1	0	2	4
Micrasema rusticum	0	0	1	0	0	1
Polycentropus sp.	1	0	1	0	0	2
COLEOPTERA (beetles)						
Macronychus glabratus	0	0	1	0	0	1
Stenelmis crenata	0	0	1	1	1	3
Stenelmis convexula	0	0	1	0	0	1
DIPTERA - misc. (true flies)						
Atherix lantha	0	0	0	0	4	4
Simulium jonesi	4	14	0	1	0	19

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Corynoneura nr. taris	3	0	0	0	0	3
Corynoneura sp. 3	0	0	0	1	0	1
Paramerina sp.	1	0	0	0	0	1
Parakiefferiella sp. A	0	0	0	0	1	1
Polypedilum convictum	4	0	2	2	0	8
Rheocricotopus robacki	6	0	4	1	0	11
Rheotanytarsus distinctissimus gp.	1	0	1	1	0	3
Rheotanytarsus exiguus gp.	0	0	2	0	0	2
Tanytarsus sp.	0	0	0	0	1	1
Thienemanniella fusca gp.	3	0	0	1	1	5
Thienemanniella xena gp.	0	0	1	0	1	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

**DATE: OCTOBER 1990
CLIENT: WSRC
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC**

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Heptagenia flavescens	0	1	0	0	0	1
Neophemera youngi	3	0	2	0	2	7
Paraleptophlebia sp.	0	1	0	0	0	1
Stenonema modestum/smithae	0	5	0	0	0	5
PLECOPTERA (stoneflies)						
Acroneuria abnormis	0	2	1	1	2	6
Paragnetina kansensis	4	0	3	0	0	7
Perlesta placida	0	0	0	1	0	1
Pteronarcys dorsata	0	1	1	1	2	5
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	8	0	1	1	0	10
Cheumatopsyche sp.	1	1	0	0	0	2
Hydropsyche elissoma	3	12	0	2	3	20
Lype diversa	0	0	1	0	0	1
COLEOPTERA (beetles)						
Macronychus glabratus	0	0	1	1	1	3
Stenelmis crenata	0	0	1	1	0	2
Stenelmis sp. 3	1	0	0	0	0	1
Stenelmis sp. (larvae)	0	2	0	0	0	2
DIPTERA - misc. (true flies)						
Antocha sp.	0	0	0	0	2	2
Simulium jonesi	0	9	0	4	0	13

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

**DATE: OCTOBER 1990
CLIENT: WSRC
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC**

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	0	1	0	0	0	1
Corynoneura nr. taris	0	1	0	3	0	4
Polypedilum convictum	1	4	0	2	1	8
Polypedilum fallax	0	0	1	0	0	1
Polypedilum illinoense	0	1	0	0	0	1
Rheocricotopus robacki	3	6	1	6	4	20
Rheotanytarsus distinctissimus gp.	0	5	0	0	0	5
Rheotanytarsus exiguus gp.	0	1	0	0	0	1
Thienemanniella fusca gp.	1	2	0	2	2	7
Thienemanniella xena gp.	0	1	0	0	0	1
Tvetenia discoloripes gp.	0	0	1	1	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

**DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC**

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Neophemera youngi</i>	0	0	1	0	0	1
<i>Stenonema modestum/smithae</i>	0	1	1	0	0	2
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	1	2	2	2	0	7
<i>Paragnetina kansensis</i>	2	3	2	2	0	9
<i>Pteronarcys dorsata</i>	2	2	1	0	0	5
MEGALOPTERA						
<i>Corydalus cornutus</i>	0	0	1	0	0	1
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	2	0	0	3	0	5
<i>Hydropsyche elissoma</i>	0	7	3	4	0	14
COLEOPTERA (beetles)						
<i>Macronychus glabratus</i>	0	1	0	0	0	1
<i>Stenelmis crenata</i>	0	4	1	0	0	5
DIPTERA – misc. (true flies)						
<i>Simulium jonesi</i>	0	1	2	6	0	9

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 5

DATE: OCTOBER 1990
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Corynoneura sp. 3	0	0	0	2	0	2
Polypedilum convictum	0	2	0	2	1	5
Polypedilum fallax	0	1	0	0	0	1
Rheocricotopus robacki	0	2	0	1	0	3
Rheotanytarsus distinctissimus gp.	0	2	0	0	0	2
Tanytarsus sp.	0	0	0	2	0	2
Thienemanniella fusca gp.	0	3	0	0	1	4
Thienemanniella xena gp.	0	0	0	1	0	1
Tvetenia discoloripes gp.	0	1	0	0	0	1
ACARI						
Hydracarina	0	0	0	1	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

All Sites

DATE: JANUARY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co., 1991

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Ephemerella invaria/rotunda	7	3	3	13
Pseudocloeon dubium	0	1	1	2
Stenonema modestum/smithae	15	20	4	39
PLECOPTERA (stoneflies)				
Acroneuria abnormis	4	4	2	10
Acroneuria mela	0	0	1	1
Helopicus subvarians	2	2	1	5
Isoperla bilineata	1	1	0	2
Isoperla dicala	5	5	4	14
Leuctra sp.	0	1	0	1
Paragnetina kansensis	1	2	1	4
Perlesta placida	6	13	8	27
Pteronarcys dorsata	4	1	3	8
Taeniopteryx sp.	0	1	0	1
TRICHOPTERA (caddisflies)				
Brachycentrus numerosus	3	6	2	11
Cheumatopsyche sp.	1	2	0	3
Hydropsyche elissoma	5	17	7	29
Lype diversa	1	0	1	2
Micrasema rusticum	0	1	0	1
COLEOPTERA (beetles)				
Macronychus glabratus	0	0	1	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: JANUARY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co., 1991

All Sites

Site 1	Site 2	Site 5	TOTAL
--------	--------	--------	-------

DIPTERA - misc. (true flies)

Atherix lantha	1	1	1	3
Chelifera sp.	1	0	0	1
Empididae	0	1	0	1
Hemerodromia sp. 1	0	0	2	2
Simulium jonesi	1	1	2	4
Simulium tuberosum	9	2	14	25

DIPTERA- Chironomidae (midges)

Conchapelopia sp.	1	3	1	5
Corynoneura nr. taris	27	19	11	57
Corynoneura sp. 3	58	29	51	138
Cricotopus bicinctus	1	0	1	2
Cricotopus vierriensis	0	5	0	5
Nanocladius sp.	0	1	1	2
Orthocladius lignicola	1	0	0	1
Orthocladius sp.	0	1	3	4
Parakiefferiella sp. A	0	1	0	1
Parametriochnemus lundbecki	1	0	0	1
Polypedilum convictum	26	48	38	112
Polypedilum fallax	1	0	1	2
Polypedilum illinoense	0	0	1	1
Rheocricotopus robacki	26	50	55	131
Rheosmittia sp.	0	1	0	1
Rheotanytarsus distinctissimus gp.	22	42	12	76
Rheotanytarsus exiguus gp.	12	58	10	80
Tanytarsus sp. XIII Rutter	0	0	1	1
Thienemanniella fusca gp.	5	7	1	13
Thienemanniella xena gp.	18	26	21	65
Tvetenia discoloripes gp.	13	5	20	38
Tvetenia paucunca	1	0	3	4

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: JANUARY 1991

CLIENT: S.R.S.

STREA Upper Three Runs Creek

LOCATION: Aiken Co., 1991

All Sites

	Site 1	Site 2	Site 5	TOTAL
ANNELIDA (worms)				
Naididae	0	0	1	1
MOLLUSCA (snails, clams)				
Ferrissia sp.	0	0	1	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: JANUARY 1991

CLIENT: S.R.S.

STREA Upper Three Runs Creek

LOCATION: Aiken Co., S.C.

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
EPHEMEROPTERA (mayflies)						
<i>Ephemera invaria/rotunda</i>	0	0	3	1	3	7
<i>Stenonema modestum/smithae</i>	0	6	1	1	8	16
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	2	0	1	1	0	4
<i>Helopicus subvarians</i>	0	2	0	0	0	2
<i>Isoperla bilineata</i>	0	0	0	0	1	1
<i>Isoperla dicala</i>	0	1	0	2	2	6
<i>Paragnetina kansensis</i>	1	0	0	0	0	1
<i>Perlesta placida</i>	1	1	0	0	4	
<i>Pteronarcys dorsata</i>	2	0	1	1	0	4
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	0	2	1	0	0	3
<i>Cheumatopsyche sp.</i>	0	0	0	0	1	1
<i>Hydropsyche elissoma</i>	0	0	0	1	4	6
<i>Lype diversa</i>	0	0	0	0	1	1
DIPTERA – misc. (true flies)						
<i>Atherix lantha</i>	0	0	0	0	1	1
<i>Chelifera sp.</i>	0	0	0	0	1	1
<i>Simulium jonesi</i>	0	0	0	1	0	1
<i>Simulium tuberosum</i>	0	0	1	6	3	9

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: JANUARY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co., S.C.

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	0	0	0	0	1	1
Corynoneura nr. taris	7	0	14	3	3	27
Corynoneura sp. 3		1	36	14	2	68
Cricotopus bicinctus	0	0	1	0	0	1
Orthocladius lignicola	0	0	0	1	0	1
Parametriocnemus lundbecki	1	0	0	0	0	1
Polypedilum convictum	8	4	1	8	6	2
Polypedilum fallax	0	0	0	0	1	1
Rheocricotopus robacki		4	3	6	8	2
Rheotanytarsus distinctissimus gp.	0	4	2	6	11	22
Rheotanytarsus exiguus gp.	0	2	1	1	8	12
Thienemanniella fusca gp.	0	0	6	0	0	6
Thienemanniella xena gp.	2	3	6	4	4	18
Tvetenia discoloripes gp.	3	2	4	2	2	13
Tvetenia paucunca	0	0	1	0	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: JANUARY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
EPHEMEROPTERA (mayflies)						
<i>Ephemerella invaria/rotunda</i>	1	0	1	0	1	3
<i>Pseudocloeon dubium</i>	0	0	0	0	1	1
<i>Stenonema modestum/smithae</i>	6	9	3	1	1	20
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	1	2	1	0	0	4
<i>Helopicus subvarians</i>	0	1	0	1	0	2
<i>Isoperla bilineata</i>	0	0	1	0	0	1
<i>Isoperla dicala</i>	1	2	1	0	1	5
<i>Leuctra sp.</i>	1	0	0	0	0	1
<i>Paragnetina kansensis</i>	1	0	0	1	0	2
<i>Perlesta placida</i>	7	5	0	1	0	13
<i>Pteronarcys dorsata</i>	0	1	0	0	0	1
<i>Taeniopteryx sp.</i>	0	0	0	0	1	1
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	2	3	1	0	0	6
<i>Cheumatopsyche sp.</i>	2	0	0	0	0	2
<i>Hydropsyche elissoma</i>	13	3	0	1	0	17
<i>Micrasema rusticum</i>	1	0	0	0	0	1
DIPTERA - misc. (true flies)						
<i>Atherix lantha</i>	0	1	0	0	0	1
<i>Empididae</i>	0	1	0	0	0	1
<i>Simulium jonesi</i>	0	0	0	0	1	1
<i>Simulium tuberosum</i>	1	0	0	0	1	2
DIPTERA- Chironomidae (midges)						
<i>Conchapelopia sp.</i>	1	1	1	0	0	3
<i>Corynoneura nr. taris</i>	6	2	2	8	1	19

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: JANUARY 1991

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<i>Corynoneura</i> sp. 3	0	12	2	7	8	29
<i>Cricotopus vierriensis</i>	0	0	0	0	5	5
<i>Nanocladius</i> sp.	0	1	0	0	0	1
<i>Orthocladius</i> sp.	0	1	0	0	0	1
<i>Parakiefferiella</i> sp. A	0	0	0	1	0	1
<i>Polypedilum convictum</i>	12	7	14	5	10	48
<i>Rheocricotopus robacki</i>	20	10	5	4	11	50
<i>Rheosmittia</i> sp.	0	1	0	0	0	1
<i>Rheotanytarsus distinctissimus</i> gp.	17	6	9	0	10	42
<i>Rheotanytarsus exiguus</i> gp.	20	3	13	3	19	58
<i>Thienemanniella fusca</i> gp.	2	2	1	2	0	7
<i>Thienemanniella xena</i> gp.	4	4	5	5	8	26
<i>Tvetenia discoloripes</i> gp.	2	0	2	1	0	5

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: JANUARY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Ephemerella invaria/rotunda</i>	2	0	0	1	0	3
<i>Pseudocloeon dubium</i>	0	0	0	1	0	1
<i>Stenonema modestum/smithae</i>	2	0	0	1	1	4
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	1	0	0	0	1	2
<i>Acroneuria mela</i>	0	0	0	0	1	1
<i>Helopicus subvarians</i>	0	0	0	1	0	1
<i>Isoperla dicala</i>	1	0	0	1	2	4
<i>Paragnetina kansensis</i>	0	0	1	0	0	1
<i>Pteronarcys dorsata</i>	2	0	0	1	0	3
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	0	0	0	2	0	2
<i>Hydropsyche elissoma</i>	2	0	1	3	1	7
<i>Lype diversa</i>	0	0	0	0	1	1
COLEOPTERA (beetles)						
<i>Macronychus glabratus</i>	0	0	0	0	1	1
DIPTERA - misc. (true flies)						
<i>Atherix lantha</i>	1	0	0	0	0	1
<i>Hemerodromia sp. 1</i>	2	0	0	0	0	2
<i>Simulium jonesi</i>	0	0	1	0	1	2
<i>Simulium tuberosum</i>	1	0	7	6	0	14

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: JANUARY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co., SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	1	0	0	0	0	1
Corynoneura nr. taris	5	0	1	5	0	11
Corynoneura sp. 3	21	0	3	25	2	51
Cricotopus bicinctus	0	0	0	1	0	1
Nanocladius sp.	1	0	0	0	0	1
Orthocladius sp.	0	0	0	3	0	3
Polypedilum convictum	8	0	4	8	18	38
Polypedilum fallax	0	0	0	0	1	1
Polypedilum illinoense	0	0	0	0	1	1
Rheocricotopus robacki	27	0	4	13	11	55
Rheotanytarsus distinctissimus gp.	0	0	2	9	1	12
Rheotanytarsus nr. exiguus	0	0	0	8	2	10
Tanytarsus sp. XIII Rutter	0	0	0	0	1	1
Thienemanniella fusca gp.	1	0	0	0	0	1
Thienemanniella xena gp.	6	0	0	11	4	21
Tvetenia discoloripes gp.	4	0	2	9	5	20
Tvetenia paucunca	0	0	1	2	0	3
ANNELIDA (worms)						
Naididae	0	0	0	0	1	1
MOLLUSCA (snails, clams)						
Ferrissia sp.	0	0	0	1	0	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: April 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Baetis ephippiatus	1	1	0	2
Baetis intercalaris	0	1	0	1
Dannella simplex	0	1	0	1
Ephemerella argo	1	0	1	2
Ephemerella dorothea	1	0	3	4
Ephemerella catawba	0	2	4	6
Ephemerella invaria/rotunda	0	2	3	5
Heptagenia flavescens	2	2	0	4
Pseudocloeon dubium	0	1	0	1
Pseudocloeon bimaculatus	0	2	1	3
Pseudocloeon punctiventris	1	1	0	2
Stenonema modestum/smithae	3	4	1	8
ODONATA (dragonflies)				
Perithemis sp.	1	0	1	2
PLECOPTERA (stoneflies)				
Acroneuria abnormis	7	5	9	21
Acroneuria mela	2	0	1	3
Isoperla dicala	5	1	13	19
Paragnetina kansensis	5	7	12	24
Perlesta placida	16	15	9	40
Pteronarcys dorsata	0	0	1	1
MEGALOPTERA				
Corydalus cornutus	0	1	0	1
TRICHOPTERA (caddisflies)				
Cheumatopsyche sp.	2	5	3	10
Chimarra sp.	2	0	0	2
Hydropsyche elissoma	18	16	29	63

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: April 1991

CLIENT: S.R.S.

STREAM: Upper Three Runs Creek

LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
COLEOPTERA (beetles)				
Gonielmis dietrichi	0	2	0	2
Macronychus glabratus	6	11	13	30
Stenelmis crenata	20	35	23	78
Stenelmis antennalis	5	18	14	37
Stenelmis sp. (larvae)	0	1	0	1
DIPTERA – misc. (true flies)				
Empididae	0	0	2	2
Simulium jonesi	8	16	12	36
Simulium tuberosum	1	15	16	32
DIPTERA- Chironomidae (midges)				
Brillia flavifrons	0	0	2	2
Conchapelopia sp.	2	1	1	4
Corynoneura nr. taris	7	0	2	9
Corynoneura sp. 3	3	2	1	6
Corynoneura sp. 4	0	1	0	1
Cricotopus bicinctus	0	6	11	17
Cricotopus sp.	1	0	0	1
Cricotopus vierriensis	0	1	0	1
Eukeifferiella sp.	0	0	1	1
Nilotanytus fimbriatus	1	0	0	1
Orthocladius lignicola	7	19	14	40
Orthocladius sp.	3	0	1	4
Parametriocnemus lundbecki	2	5	0	7
Polypedilum convictum	5	19	11	35
Polypedilum fallax	1	0	0	1
Rheocricotopus robacki	9	10	4	23
Rheocricotopus tuberculatus	1	0	0	1
Rheotanytarsus distinctissimus gp.	12	11	9	32
Rheotanytarsus exiguus gp.	2	2	1	5
Synorthocladius semivirens	3	1	0	4

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

DATE: April 1991

CLIENT: S.R.S.

STRE. M: Upper Three Runs Creek

LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
Tanytarsus glabrescens gp.	1	1	0	2
Tanytarsus sp.	0	0	1	1
Thienemanniella fusca gp.	2	1	1	4
Thienemanniella xena gp.	7	2	2	11
Tvetenia discoloripes gp.	9	28	24	61
Tvetenia paucunca	0	2	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1	DATE:	April 1991
	CLIENT:	S.R.S.
	STREA	Upper Three Runs Creek
	LOCATION:	Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Baetis ephippiatus	1	0	0	0	0	1
Ephemerella argo	0	0	1	0	0	1
Ephemerella dorothea	0	1	0	0	0	1
Heptagenia flavescens	1	0	1	0	0	2
Pseudocloeon punctiventris	0	1	0	0	0	1
Stenonema modestum/smithae	0	0	0	2	1	3
ODONATA (dragonflies)						
Perithemis sp.	0	0	1	0	0	1
PLECOPTERA (stoneflies)						
Acroneuria abnormis	3	1	0	2	1	7
Acroneuria mela	0	0	0	2	0	2
Isoperla dicala	1	1	0	3	0	5
Paragnetina kansensis	2	0	3	0	0	5
Perlesta placida	2	4	3	0	7	16
TRICHOPTERA (caddisflies)						
Cheumatopsyche sp.	0	0	0	2	0	2
Chimarra sp.	0	2	0	0	0	2
Hydropsyche elissoma	0	5	6	7	0	18
COLEOPTERA (beetles)						
Macronychus glabratus	0	6	0	0	0	6
Stenelmis crenata	3	2	10	1	4	20
Stenelmis antennalis	1	0	4	0	0	5

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 1

DATE: April 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA - misc. (true flies)						
Simulium jonesi	0	2	2	3	1	8
Simulium tuberosum	0	0	1	0	0	1
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	0	0	1	0	1	2
Corynoneura nr. taris	0	0	1	6	0	7
Corynoneura sp. 3	0	0	0	2	1	3
Cricotopus sp.	0	0	0	0	1	1
Nilotanytus fimbriatus	0	0	0	0	1	1
Orthocladius lignicola	0	1	0	6	0	7
Orthocladius sp.	0	0	2	1	0	3
Parametrioctenus lundbecki	2	0	0	0	0	2
Polypedilum convictum	2	2	0	0	1	5
Polypedilum fallax	0	0	0	0	1	1
Rheocricotopus robacki	1	1	3	2	2	9
Rheocricotopus tuberculatus	0	0	0	1	0	1
Rheotanytarsus distinctissimus gp.	2	1	0	7	2	12
Rheotanytarsus exiguus gp.	0	0	0	0	2	2
Synorthocladius semivirens	0	0	0	3	0	3
Tanytarsus glabrescens gp.	0	0	0	0	1	1
Thienemanniella fusca gp.	0	0	0	2	0	2
Thienemanniella xena gp.	0	2	1	3	1	7
Tvetenia discoloripes gp.	1	2	1	1	4	9

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: April 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Baetis ephippiatus</i>	1	0	0	0	0	1
<i>Baetis intercalaris</i>	1	0	0	0	0	1
<i>Dannella simplex</i>	0	1	0	0	0	1
<i>Ephemerella catawba</i> gp.	0	2	0	0	0	2
<i>Ephemerella invaria/rotunda</i>	2	0	0	0	0	2
<i>Heptagenia flavescens</i>	0	2	0	0	0	2
<i>Pseudocloeon dubium</i>	0	1	0	0	0	1
<i>Pseudocloeon punctiventris</i>	0	1	0	0	0	1
<i>Pseudocloeon bimaculatus</i>	1	0	1	0	0	2
<i>Stenonema modestum/smithae</i>	0	2	0	2	0	4
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	0	0	1	3	1	5
<i>Isoperla dicala</i>	0	0	0	1	0	1
<i>Paragnetina kansensis</i>	2	2	1	0	2	7
<i>Perlesta placida</i>	6	0	5	1	3	15
MEGALOPTERA						
<i>Corydalus cornutus</i>	1	0	0	0	0	1
TRICHOPTERA (caddisflies)						
<i>Cheumatopsyche</i> sp.	1	0	1	1	2	5
<i>Hydropsyche elissoma</i>	2	5	8	1	0	16
COLEOPTERA (beetles)						
<i>Gonielmis dietrichi</i>	0	2	0	0	0	2
<i>Macronychus glabratus</i>	5	0	3	2	1	11
<i>Stenelmis crenata</i>	3	11	6	11	4	35
<i>Stenelmis antennalis</i>	3	5	2	6	2	18
<i>Stenelmis</i> (larvae)	0	0	0	0	1	1

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 2

DATE: April 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA – misc. (true flies)						
<i>Simulium jonesi</i>	0	3	9	3	1	16
<i>Simulium tuberosum</i>	10	2	3	0	0	15
DIPTERA– Chironomidae (midges)						
<i>Conchapelopia</i> sp.	0	0	0	1	0	1
<i>Corynoneura</i> sp. 3	0	0	1	0	1	2
<i>Corynoneura</i> sp. 4	1	0	0	0	0	1
<i>Cricotopus bicinctus</i>	0	6	0	0	0	6
<i>Cricotopus vierriensis</i>	0	0	0	0	1	1
<i>Orthocladius lignicola</i>	2	4	2	2	9	ERR
<i>Parametrioctenus lundbecki</i>	0	0	1	2	2	5
<i>Polypedilum convictum</i>	0	4	5	7	3	19
<i>Rheocricotopus robacki</i>	4	1	2	0	3	10
<i>Rheotanytarsus distinctissimus</i> gp.	2	1	2	6	0	11
<i>Rheotanytarsus exiguus</i> gp.	1	0	0	0	1	2
<i>Synorthocladius semivirens</i>	1	0	0	0	0	1
<i>Tanytarsus glabrescens</i> gp.	1	0	0	0	0	1
<i>Thienemanniella fusca</i> gp.	0	1	0	0	0	1
<i>Thienemanniella xena</i> gp.	0	0	0	1	1	2
<i>Tvetenia discoloripes</i> gp.	7	7	7	4	3	28
<i>Tvetenia paucunca</i>	1	0	1	0	0	2

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: April 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Ephemerella argo</i>	0	0	1	0	0	1
<i>Ephemerella dorothea</i>	1	2	0	0	0	3
<i>Ephemerella catawba</i> gp.	0	1	2	1	0	4
<i>Ephemerella invaria/rotunda</i>	0	1	0	1	1	3
<i>Pseudocloeon bimaculatus</i>	0	0	0	1	0	1
<i>Stenonema modestum/smithae</i>	0	0	0	0	1	1
ODONATA (dragonflies)						
<i>Perithemis</i> sp.	0	0	0	0	1	1
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	4	2	2	1	0	9
<i>Acroneuria mela</i>	1	0	0	0	0	1
<i>Isoperla dicala</i>	2	6	4	1	0	13
<i>Paragnetina kansensis</i>	2	1	3	3	3	12
<i>Perlesta placida</i>	1	2	4	0	2	9
<i>Pteronarcys dorsata</i>	0	0	0	0	1	1
TRICHOPTERA (caddisflies)						
<i>Cheumatopsyche</i> sp.	0	2	0	0	1	3
<i>Hydropsyche elissoma</i>	0	9	5	6	9	29
COLEOPTERA (beetles)						
<i>Macronychus glabratus</i>	1	3	0	2	7	13
<i>Stenelmis crenatus</i>	0	7	4	3	9	23
<i>Stenelmis antennalis</i>	0	2	1	2	9	14

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: April 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA – misc. (true flies)						
Empididae	1	0	1	0	0	2
Simulium jonesi	0	3	3	3	3	12
Simulium tuberosum	0	1	10	4	1	16
DIPTERA– Chironomidae (midges)						
Brillia flavifrons	1	0	1	0	0	2
Conchapelopia sp.	0	1	0	0	0	1
Corynoneura nr. taris	0	0	0	2	0	2
Corynoneura sp. 3	0	1	0	0	0	1
Cricotopus bicinctus	0	1	0	10	0	11
Eukiefferiella sp.	0	0	0	0	1	1
Orthocladius lignicola	0	0	3	9	2	14
Orthocladius sp.	0	0	1	0	0	1
Polypedilum convictum	6	4	1	0	0	11
Rheocricotopus robacki	0	0	3	0	1	4
Rheotanytarsus distinctissimus gp.	2	0	2	3	2	9
Rheotanytarsus exiguus gp.	0	0	1	0	0	1
Tanytarsus sp.	0	0	0	1	0	1
Thienemanniella fusca gp.	0	0	1	0	0	1
Thienemanniella xena gp.	0	0	2	0	0	2
Tvetenia discoloripes gp.	0	9	6	6	3	24

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: JULY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

	Site 1	Site 2	Site 5	TOTAL
EPHEMEROPTERA (mayflies)				
Heptagenia flavescens	8	10	3	21
Stenonema modestum/smithae	1	7	11	19
PLECOPTERA (stoneflies)				
Acroneuria abnormis	15	15	16	46
Paragnetina kansensis	5	4	5	14
Pteronarcys dorsata	6	7	5	18
MEGALOPTERA				
Corydalis cornutus	1	1	0	2
Nigronia serricornis	0	1	1	2
TRICHOPTERA (caddisflies)				
Brachycentrus numerosus	0	2	3	5
Cheumatopsyche sp.	7	4	2	13
Chimarra sp.	0	0	2	2
Hydropsyche elissoma	90	48	64	202
Lype diversa	0	0	1	1
Mayatrichia ayama	1	0	1	2
COLEOPTERA (beetles)				
Dineutus (larvae)	0	1	0	1
Gonielmis dietrichi	0	2	2	4
Macronychus glabratus	24	34	20	78
Stenelmis antennalis	0	1	1	2
Stenelmis crenata	18	63	14	95
Stenelmis convexula	14	34	22	70
Stenelmis (larvae)	3	1	1	5

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

DATE: JULY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

ALL SITES

Site 1	Site 2	Site 5
--------	--------	--------

TOTAL

DIPTERA - misc. (true flies)

Hemerodromia sp. 1	0	1	0
Simulium jonesi	14	23	13
Simulium tuberosum	0	1	0

1
50
1

DIPTERA- Chironomidae (midges)

Cricotopus bicinctus	0	1	0
Cricotopus vierriensis	0	5	0
Conchapelopia sp.	0	0	1
Nanocladius sp.	1	0	0
Nilotanytus fimbriatus	0	0	1
Parametriocnemus lundbecki	5	1	2
Polypedilum convictum	3	2	2
Polypedilum fallax	0	1	0
Polypedilum illinoense	1	0	0
Rheocricotopus robacki	3	1	7
Rheotanytarsus distinctissimus gp.	2	1	3
Stenochironomus hiliaris	0	0	1
Tanytarsus glabrescens gp.	1	1	1
Thienemanniella fusca gp.	2	0	0
Tveteria discoloripes gp.	12	7	9

1
5
1
1
1
8
7
1
1
11
6
1
3
2
28

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: JULY 1991

CLIENT: S.R.S.

STREA Upper Three Runs Creek

LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Heptagenia flavescens	0	2	3	1	2	8
Stenonema modestum/smithae	0	0	1	0	0	1
PLECOPTERA (stoneflies)						
Acroneuria abnormis	4	4	5	2	0	15
Paragnetina kansensis	0	2	0	1	2	5
Pteronarcys dorsata	1	0	2	1	2	6
MEGALOPTERA						
Corydalus cornutus	0	1	0	0	0	1
TRICHOPTERA (caddisflies)						
Cheumatopsyche sp.	0	3	1	1	2	7
Hydropsyche elissoma	8	17	20	28	17	90
Mayatrichia ayama	0	1	0	0	0	1
COLEOPTERA (beetles)						
Macronychus glabratus	5	5	5	2	7	24
Stenelmis crenata	1	8	8	1	0	18
Stenelmis convexula	0	7	5	1	1	14
Stenelmis (larvae)	0	1	1	0	1	3
DIPTERA – misc. (true flies)						
Simulium jonesi	0	6	5	3	0	14

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 1

DATE: JULY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Nanocladius sp.	0	0	0	1	0	1
Parametriocnemus lundbecki	2	1	1	1	0	5
Polypedilum convictum	0	2	0	1	0	3
Polypedilum illinoense	1	0	0	0	0	1
Rheocricotopus robacki	1	0	0	2	0	3
Rheotanytarsus distinctissimus gp.	0	0	0	2	0	2
Tanytarsus glabrescens gp.	0	0	0	1	0	1
Thienemanniella fusca gp.	1	1	0	0	0	2
Tvetenia discoloripes gp.	0	2	7	1	2	12

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 2

DATE: JULY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
Heptagenia flavescens	1	3	2	3	1	10
Stenonema modestum/smithae	0	1	0	0	6	7
PLECOPTERA (stoneflies)						
Acroneuria abnormis	5	3	3	4	0	15
Paragnetina kansensis	1	1	0	2	0	4
Pteronarcys dorsata	0	1	3	0	3	7
MEGALOPTERA						
Corydalus cornutus	0	0	0	0	1	1
Nigronia serricornis	0	0	0	1	0	1
TRICHOPTERA (caddisflies)						
Brachycentrus numerosus	0	0	0	0	2	2
Cheumatopsyche sp.	0	0	3	0	1	4
Hydropsyche elissoma	5	7	6	10	20	48
COLEOPTERA (beetles)						
Dineutus (larva)	0	0	0	1	0	1
Gonielmis dietrichi	0	0	0	1	1	2
Macronychus glabratus	8	12	4	3	7	34
Stenelmis antennalis	0	0	0	0	1	1
Stenelmis crenata	22	7	14	10	10	63
Stenelmis convexula	10	7	8	3	6	34
Stenelmis (larvae)	1	0	0	0	0	1
DIPTERA – misc. (true flies)						
Hemerodromia sp. 1	0	0	0	1	0	1
Simulium jonesi	1	9	7	4	2	23
Simulium tuberosum	0	0	1	0	0	1

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 2

DATE: JULY 1991
CLIENT: S.R.S.
STREAM: Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Cricotopus bicinctus	0	0	0	1	0	1
Cricotopus vierriensis	0	0	0	4	1	5
Parametrioctenus lundbecki	1	0	0	0	0	1
Polypedilum convictum	0	2	0	0	0	2
Polypedilum fallax	0	0	0	0	1	1
Rheocricotopus robacki	0	0	0	0	1	1
Rheotanytarsus distinctissimus gp.	0	0	0	1	0	1
Tanytarsus glabrescens gp.	0	0	1	0	0	1
Tvetenia discoloripes gp.	1	0	0	4	2	7

**APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.**

SITE 5

DATE: JULY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
EPHEMEROPTERA (mayflies)						
<i>Heptagenia flavescens</i>	2	1	0	0	0	3
<i>Stenonema modestum/smithae</i>	3	5	1	2	0	11
PLECOPTERA (stoneflies)						
<i>Acroneuria abnormis</i>	4	2	5	3	2	16
<i>Paragnetina kansensis</i>	1	1	0	3	0	5
<i>Pteronarcys dorsata</i>	1	2	1	1	0	5
MEGALOPTERA						
<i>Nigronia serricornis</i>	0	0	1	0	0	1
TRICHOPTERA (caddisflies)						
<i>Brachycentrus numerosus</i>	0	1	2	0	0	3
<i>Cheumatopsyche</i> sp.	2	0	0	0	0	2
<i>Chimarra</i> sp.	0	1	1	0	0	2
<i>Hydropsyche elissoma</i>	7	19	17	21	0	64
<i>Lype diversa</i>	1	0	0	0	0	1
<i>Mayatrachia ayama</i>	0	0	0	1	0	1
COLEOPTERA (beetles)						
<i>Gonielmis dietrichi</i>	0	0	0	2	0	2
<i>Macronychus glabratus</i>	4	9	4	1	2	20
<i>Stenelmis antennalis</i>	0	0	1	0	0	1
<i>Stenelmis crenata</i>	3	2	7	2	0	14
<i>Stenelmis convexula</i>	3	8	4	6	1	22
<i>Stenelmis</i> (larvae)	0	1	0	0	0	1
DIPTERA - misc. (true flies)						
<i>Simulium jonesi</i>	5	2	3	3	0	13

APPENDIX B : LIST OF SPECIES FOR QUANTITATIVE MACROINVERTEBRATE IN-STREAM
ASSESSMENT.

SITE 5

DATE: JULY 1991
CLIENT: S.R.S.
STREA Upper Three Runs Creek
LOCATION: Aiken Co. SC

	Rep A	Rep B	Rep C	Rep D	Rep E	TOTAL
<hr/>						
DIPTERA- Chironomidae (midges)						
Conchapelopia sp.	1	0	0	0	0	1
Nilotanypus fimbriatus	0	1	0	0	0	1
Parametriocnemus lundbecki	1	1	0	0	0	2
Polypedilum convictum	0	1	1	0	0	2
Rheocricotopus robacki	2	3	0	1	1	7
Rheotanytarsus distinctissimus gp.	1	0	2	0	0	3
Stenochironomus hilaris	0	0	0	0	1	1
Tanytarsus glabrescens gp.	0	1	0	0	0	1
Tvetenia discoloripes gp.	0	4	4	0	1	9

Appendix C1

General Parameters (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
Total Suspended Solids (TSS)				
April 1990	7.0	7.0	14.0	< 4.0
July 1990	15.0	17.0	27.0	2.0
October 1990	< 5.0	< 5.0	< 12.0	< 5.0
January 1991	< 5.0	7.0	< 5.0	< 5.0
March 1991	< 5.0	< 5.0	14.0	< 5.0
July 1991	22.0	22.0	32.0	< 5.0
Mean	< 9.8	< 10.5	< 17.3	< 4.3
Standard Deviation	N/A	N/A	N/A	N/A

	Site 1	Site 2	Site 5	Effluent
Specific Conductance (umhos/cm)				
April 1990	21.0	21.0	22.0	110.0
July 1990	54.0	58.0	55.0	67.0
October 1990	23.0	27.0	27.0	280.0
January 1991	22.0	22.0	22.0	89.0
March 1991	22.0	22.0	23.0	36.0
July 1991	20.0	23.0	22.0	76.0
Mean	27.0	28.8	28.5	109.7
Standard Deviation	13.3	14.4	13.1	87.0

	Site 1	Site 2	Site 5	Effluent
Hardness				
April 1990	7.0	7.0	7.0	1.0
July 1990	33.0	20.0	30.0	< 1.0
October 1990	6.0	5.0	6.0	< 1.0
January 1991	7.5	7.0	6.0	< 1.0
March 1991	6.9	6.9	11.0	2.5
July 1991	10.0	7.5	8.0	< 1.0
Mean	11.7	8.9	11.3	1.3
Standard Deviation	10.5	5.5	9.3	0.6

Appendix C1

General Parameters (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
<i>Alkalinity</i>				
April 1990	5.0	5.0	5.0	19.0
July 1990	< 1.0	< 1.0	< 1.0	< 1.0
October 1990	4.0	4.0	4.0	8.5
January 1991	2.0	1.9	2.0	11.0
March 1991	1.9	2.0	2.4	9.8
July 1991	< 1.1	< 2.6	< 2.5	8.5
Mean	< 2.50	< 2.75	< 2.82	9.63
Standard Deviation	N/A	N/A	N/A	N/A

	Site 1	Site 2	Site 5	Effluent
<i>Residual Chlorine</i>				
April 1990	< 0.05	< 0.05	< 0.05	< 0.05
July 1990	ND	ND	ND	ND
October 1990	< 0.05	< 0.05	< 0.05	< 0.05
January 1991	< 0.05	< 0.05	< 0.05	< 0.05
March 1991	< 0.05	< 0.05	< 0.05	< 0.05
July 1991	ND	ND	ND	ND
Mean	< 0.05	< 0.05	< 0.05	< 0.05
Standard Deviation	N/A	N/A	N/A	N/A

Metals (mg/L)
Quarterly Water Chemistry Sampling
Upper Three Runs Creek ; F/H Area ETF Effluent
Aiken County, SC
March 1990 – July 1991

[illegible]

Appendix C2

Metals (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County; SC

March 1990 – July 1991

<i>Manganese (Mn)</i>	<i>Site 1</i>		<i>Site 2</i>		<i>Site 5</i>		<i>Effluent</i>	
	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>
April 1990	0.005	0.019	0.005	0.019	0.004	0.026	< 0.015	< 0.015
July 1990	0.124	0.122	0.120	0.124	0.133	0.147	< 0.005	< 0.005
October 1990	0.021	0.023	0.020	0.024	0.020	0.025	< 0.001	< 0.001
January 1991	0.018	0.024	0.020	0.024	0.019	0.026	< 0.005	< 0.005
March 1991	0.012	0.017	0.012	0.023	0.011	0.028	< 0.005	< 0.633
July 1991	< 0.010	0.070	< 0.010	0.050	< 0.010	0.050	< 0.010	< 0.010
Mean	< 0.032	< 0.046	< 0.031	< 0.044	< 0.033	< 0.050	< 0.007	< 0.112
Standard Deviation	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

<i>Mercury (Hg)</i>	<i>Site 1</i>		<i>Site 2</i>		<i>Site 5</i>		<i>Effluent</i>	
	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>
April 1990	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005
July 1990	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002
October 1990	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002
January 1991	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002
March 1991	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002	< 0.0002
July 1991	< 0.0030	< 0.0030	< 0.0030	< 0.0030	< 0.0030	< 0.0030	< 0.0030	< 0.0030
Mean	< 0.0007	< 0.0007	< 0.0007	< 0.0007	< 0.0007	< 0.0007	< 0.0007	< 0.0007
Standard Deviation	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

<i>Sodium (Na)</i>	<i>Site 1</i>		<i>Site 2</i>		<i>Site 5</i>		<i>Effluent</i>	
	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>	<i>Dissolved</i>	<i>Total</i>
April 1990	1.360	1.450	1.540	1.260	2.150	1.590	No data	No data
July 1990	1.770	* 0.122	1.650	* 0.120	1.700	* 0.124	15.400	15.400
October 1990	1.350	1.390	1.910	1.940	1.710	1.730	45.300	50.900
January 1991	1.610	* 1.440	2.230	* 2.070	1.780	* 1.740	19.200	19.700
March 1991	1.400	* 1.350	1.440	1.450	1.680	2.060	6.960	* 4.300
July 1991	2.170	2.190	2.180	2.260	2.570	* 2.360	14.900	15.100
Mean	1.610	* 1.324	1.825	* 1.517	1.932	* 1.601	16.960	17.567
Standard Deviation	0.320	0.668	0.334	0.782	0.359	0.775	15.508	17.950

Appendix C2

Metals (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1		Site 2		Site 5		Effluent	
Uranium (U)	Dissolved	Total	Dissolved	Total	Dissolved	Total	Dissolved	Total
April 1990	< 0.0010	< 0.0010	0.0020	0.0030	< 0.0010	< 0.0010	< 0.0010	< 0.0010
July 1990	< 0.0010	< 0.0010	< 0.0010	< 0.0010	0.0033	0.0048	< 0.0010	0.0011
October 1990	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005	< 0.0005
January 1991	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010
March 1991	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010	< 0.0010
July 1991	No data	< 0.0010	No data	< 0.0010	No data	< 0.0010	No data	< 0.0010
Mean	< 0.0008	< 0.0009	< 0.0009	< 0.0013	< 0.0011	< 0.0016	< 0.0008	* 0.0500
Standard Deviation	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

	Site 1		Site 2		Site 5		Effluent	
Zinc (Zn)	Dissolved	Total	Dissolved	Total	Dissolved	Total	Dissolved	Total
April 1990	0.033	< 0.030	< 0.030	< 0.030	< 0.030	< 0.030	< 0.030	< 0.030
July 1990	< 0.035	0.026	< 0.035	0.031	< 0.040	0.038	0.010	0.012
October 1990	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005	< 0.005
January 1991	< 0.010	< 0.010	< 0.010	< 0.010	< 0.010	< 0.010	< 0.010	< 0.010
March 1991	0.014	0.012	0.018	0.018	0.014	0.321	0.015	0.030
July 1991	0.110	0.920	0.050	< 0.050	< 0.050	< 0.050	0.260	< 0.070
Mean	< 0.035	< 0.167	< 0.025	< 0.024	< 0.025	< 0.076	< 0.055	* 0.050
Standard Deviation	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

* Dissolved metal value exceeds total metal value; data suspect.

Appendix C3

Nutrients (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
Nitrites (NO₂)				
April 1990	< 0.01	< 0.01	< 0.01	< 0.01
July 1990	< 0.05	< 0.05	< 0.05	< 0.12
October 1990	< 0.01	< 0.01	< 0.01	0.08
January 1991	< 0.10	< 0.10	< 0.10	0.20
March 1991	< 0.10	< 0.10	< 0.10	< 0.10
July 1991	< 0.01	< 0.01	< 0.01	0.30
Mean	< 0.05	< 0.05	< 0.05	< 0.13
Standard Deviation	N/A	N/A	N/A	N/A

	Site 1	Site 2	Site 5	Effluent
Nitrates (NO₃)				
April 1990	0.10	0.10	0.20	4.00
July 1990	0.22	0.17	0.21	2.18
October 1990	0.11	0.38	0.27	26.00
January 1991	0.10	0.10	0.20	6.40
March 1991	0.10	0.10	0.11	1.34
July 1991	0.13	0.14	0.15	5.80
Mean	0.13	0.17	0.19	7.62
Standard Deviation	0.05	0.11	0.05	9.22

	Site 1	Site 2	Site 5	Effluent
Ammonia Nitrogen (NH₃)				
April 1990	0.18	0.10	< 0.10	< 0.10
July 1990	< 0.30	< 0.30	< 0.30	< 0.30
October 1990	< 0.30	< 0.30	< 0.30	< 0.30
January 1991	< 0.30	< 0.30	< 0.30	< 0.30
March 1991	< 0.30	< 0.30	< 0.30	< 0.30
July 1991	0.37	0.26	0.38	0.12
Mean	0.29	0.26	0.28	0.24
Standard Deviation	N/A	N/A	N/A	N/A

Appendix C3

Nutrients (mg/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
Total Phosphate (T-PO₄)				
April 1990	0.03	0.02	0.03	0.02
July 1990	0.08	0.08	0.29	0.02
October 1990	< 0.10	< 0.10	< 0.10	< 0.10
January 1991	< 0.10	< 0.10	< 0.10	< 0.10
March 1991	< 0.10	< 0.10	< 0.10	< 0.10
July 1991	0.13	0.23	0.31	0.06
Mean	< 0.09	< 0.11	< 0.16	0.07
Standard Deviation	N/A	N/A	N/A	N/A

	Site 1	Site 2	Site 5	Effluent
Ortho-Phosphate (O-PO₄)				
April 1990	< 0.01	< 0.02	< 0.02	< 0.02
July 1990	0.03	0.02	0.05	< 0.01
October 1990	0.02	0.02	0.02	0.01
January 1991	< 0.05	< 0.05	< 0.05	< 0.05
March 1991	< 0.05	< 0.05	< 0.05	< 0.05
July 1991	0.02	0.02	0.02	0.01
Mean	< 0.04	< 0.04	< 0.05	< 0.04
Standard Deviation	N/A	N/A	N/A	N/A

Appendix C4

Organics (ug/L)

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
<i>Tributyl Phosphate (TBP)</i>				
April 1990	< 50.00	< 50.00	< 50.00	< 50.00
July 1990	< 50.00	< 50.00	< 50.00	< 50.00
October 1990	< 10.00	< 10.00	< 10.00	< 10.00
January 1991	< 10.00	< 10.00	< 10.00	< 10.00
March 1991	< 10.00	< 10.00	< 10.00	< 10.00
July 1991	< 10.00	< 10.00	< 10.00	< 10.00
Mean	< 28.00	< 28.00	< 28.00	< 28.00
Standard Deviation	N/A	N/A	N/A	N/A

	Site 1	Site 2	Site 5	Effluent
<i>Kerosene</i>				
April 1990	< 0.10	< 0.10	< 0.10	< 0.10
July 1990	< 0.05	< 0.05	< 0.05	< 0.05
October 1990	< 1.00	< 1.00	< 1.00	< 1.00
January 1991	< 1.00	< 1.00	< 1.00	< 1.00
March 1991	< 1.00	< 1.00	< 1.00	< 1.00
July 1991	< 20.00	< 20.00	< 20.00	< 20.00
Mean	< 4.63	< 4.63	< 4.63	< 4.63
Standard Deviation	N/A	N/A	N/A	N/A

Appendix C5

Field Parameters

Quarterly Water Chemistry Sampling

Upper Three Runs Creek ; F/H Area ETF Effluent

Aiken County, SC

March 1990 – July 1991

	Site 1	Site 2	Site 5	Effluent
pH (units)				
April 1990	5.38	5.15	5.15	No data
July 1990	4.35	4.04	4.32	6.77
October 1990	5.87	5.82	5.72	No data
January 1991	6.04	6.44	6.69	7.57
March 1991	6.89	5.98	6.78	7.46
July 1991	5.24	5.42	5.80	7.45
Mean	5.63	5.48	5.74	7.31
Standard Deviation	0.86	0.83	0.93	0.37

	Site 1	Site 2	Site 5	Effluent
Dissolved Oxygen (mg/L)				
April 1990	10.1	10.2	9.1	9.2
July 1990	8.5	8.5	8.3	7.1
October 1990	10.1	10.0	10.4	No data
January 1991	8.9	9.2	9.1	9.6
March 1991	10.1	11.8	12.0	8.2
July 1991	7.2	6.6	6.8	5.8
Mean	9.2	9.4	9.3	8.0
Standard Deviation	1.2	1.8	1.8	1.6

	Site 1	Site 2	Site 5	Effluent
Temperature (C)				
April 1990	15.3	15.3	15.1	19.3
July 1990	23.7	23.8	23.8	30.4
October 1990	12.9	12.8	12.8	No data
January 1991	9.4	9.4	9.4	17.8
March 1991	13.1	12.9	13.1	18.5
July 1991	22.0	21.5	22.0	28.0
Mean	16.1	16.0	16.0	22.8
Standard Deviation	5.6	5.6	5.7	5.9

APPENDIX D. Statistical Analysis Results. Friedman's Test (Non-Parametric) Upper Three
Runs Creek Biomonitoring, Aiken County, SC. March 1990-July 1991.

Parameter	Probability (p)	F Value
<u>Biological - Quantitative</u>		
Species Richness	0.142	4.33
Average # of Species/Replicate	0.072	5.33
Density	0.072	5.33
% Ephemeroptera	0.57	1.33
% Plecoptera	0.184	4.00
% Trichoptera	0.43	2.33
% Coleoptera	0.142	5.08
% Diptera (non-midges)	0.956	0.330
% Chironomidae	0.956	0.330
EPT Index	0.430	2.33
Biotic Index	0.184	4.00
% Collector - Gatherers (By Abundance)	0.43	2.58
% Collector - Filterers (By Abundance)	0.43	2.33
% Predators (By Abundance)	0.052*	6.33
% Scrapers (By Abundance)	0.252	3.0
% Shredder - Herbivores (By Abundance)	0.252	3.0
% Shredder - Detritivores (By Abundance)	0.57	1.33
Diversity Index	0.57	1.33
Equitability Index	1.00	0.0
Ephemeroptera Density	0.252	3.58
Plecoptera Density	0.74	1.08
Trichloptera Density	0.956	0.58
Coleoptera Density	0.29*	7.58
Diptera Density	0.956	0.33
Chironomidae Density	0.74	1.0

* Statistically significant at p - 0.05.

**APPENDIX D Statistical Analysis Results. Friedman's Test (Non-Parametric) Upper Three
Runs Creek Biomonitoring, Aiken County, SC. March 1990-July 1991.**

Parameter	Probability (p)	F Value
<u>Chemical Analysis</u>		
Total Suspended Solids	0.072	5.58
Specific Conductance	0.184	4.08
Hardness	0.252	3.58
Alkalinity	0.956	0.75
Residual Chlorine	N/A	N/A
Nitrites	0.74	1.00
Nitrates	0.184	4.08
Ammonia	0.956	0.58
Total Phosphate	0.570	1.58
Ortho Phosphate	0.570	1.58
Tri-Butyl Phosphate	N/A	N/A
Kerosene	N/A	N/A
Manganese (Total)	0.142	5.25
Sodium (Total)	0.072	5.33

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
EPHEMEROPTERA (mayflies)	
Baetis ephippiatus	5
Baetis frondalis	5
Baetis intercalaris	6
Baetis propinquus	5
(Baetis sp.)	5
Centroptilum sp.	2
Dannella simplex	2
Dolania americana	2
Ephemerella argo	1
Ephemerella catawba group	1
Ephemerella dorothea	1
Ephemerella invaria/rotunda	1
Eurylophella temporalis	5
(Eurylophella sp.)	5
Heptagenia flavescens	4
Hexagenia limbata	6
Neoephemera youngi	5
Paraleptophlebia sp.	1
Pseudocloeon bimaculatus	4
Pseudocloeon dubium	4
Pseudocloeon punctiventris	5
Stenonema modestum/smithae	5
Tricorythodes sp.	4

ODONATA (dragonflies,damselflies)	
Boyeria vinosa	2
Calopteryx spp.	5
Gomphus (G.) lividus	5
Neurocordulia sp.	5
Perithemis sp.	8
Progomphus sp.	1
Tetragoneuria costalis/cynosura	5

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
-------	--------------

PLECOPTERA (stoneflies)
<i>Acroneuria abnormis</i>
<i>Acroneuria mela</i>
<i>Amphinemura delosa</i>
<i>Haploperla brevis</i>
<i>Helopicus subvarians</i>
<i>Isoperla bilineata</i>
<i>Isoperla dicala</i>
<i>Isoperla holochlora/lata</i>
(<i>Isoperla</i> sp.)
<i>Leuctra</i> sp.
<i>Paragnetina kansensis</i>
<i>Perlesta placida</i>
<i>Pteronarcys dorsata</i>
<i>Taeniopteryx</i> sp.

0
0
3
1
1
4
2
2
2
0
1
5
0
2

HETEROPTERA (true bugs)
<i>Gerris remigis</i>
<i>Metrobates hesperius</i>
<i>Ochterus</i> sp.
<i>Rhagovelia obesa</i>

8
8
8
8

MEGALOPTERA (hellgrammites)
<i>Chauliodes rastricornis</i>
<i>Corydalis cornutus</i>
<i>Nigronia serricornis</i>

4
6
0

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
TRICHOPTERA (caddisflies)	
Agarodes libalis	3
Anisocentropus pyraloides	3
Brachycentrus numerosus	1
Ceraclea sp.	3
Cheumatopsyche spp.	5
Chimarra sp.	4
Diplectrona modesta	0
Hydropsyche elissoma	6
(Hydropsyche sp.)	6
Hydroptila sp.	6
Lepidostoma sp.	1
Lype diversa	2
Macrostemum carolina	3
Mayatrichia ayama	4
Micrasema rusticum	2
Micrasema wataga	2
Neureclipsis sp.	7
Nyctiophylax sp.	5
Oecetis sp.	8
Phylocentropus sp.	5
Polycentropus sp.	6
Ptilostomis sp.	5
Pycnopsyche sp.	4
Triaenodes spp.	6

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
COLEOPTERA (beetles)	
Anchytarsus bicolor	4
Ancyronyx variegatus	6
Copelatus glyphicus	8
Cyphon sp.	8
Dineutus assimilis	8
Dineutus discolor	8
(Dineutus – larvae)	8
Goniemis dietrichi	4
Gyrinus sp.	8
Hydroporus sp.	8
Hydaticus sp.	8
Macronychus glabratus	4
Oulimnius latisculus	4
Optioservus sp.	4
Peltodytes sp.	8
Sperchopsis tessellatus	8
Stenelmis antennalis	5
Stenelmis convexula	5
Stenelmis crenata	5
Stenelmis sp. 3	5
(Stenelmis – larvae)	5

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
DIPTERA (true flies)	
Non-Chironomidae	
Antocha sp.	3
Atherix lantha	2
Bezzia sp. 2	6
Chelifera sp.	6
Clinocera sp.	6
Culex sp.	10
Dolichopodidae	4
(Empididae)	6
Erioptera/Ormosia sp.	7
Hemerodromia sp. 1	6
Hemerodromia sp. 2	6
Hexatoma sp.	2
Palpomyia sp. 4	6
Probezzia sp.	6
Simulium jonesi	2
Simulium snowi	2
Simulium tuberosum	4
Stilobezzia sp.	6
Telmatoscopus sp.	6
Tipula sp. A	5
Tipula sp. 1	4
Tipula (Yamato.) sp. 7	4
Tipula (Yamato.) nr strepens	4
Wiedemannia sp.	6
Chironomidae – Tanypodinae	
Ablabesmyia janta/parajanta	5
Ablabesmyia mallochi	8
Clinotanypus pinguis	8
Conchapelopia sp.	6
Meropelopia sp.	8
Nilotanypus fimbriatus	6
Paramerina sp.	8
Procladius sp.	9

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
Chironomidae – Diamesinae	
Potthastia longimana	2
Chironomidae – Orthoclaadiinae	
Brillia flavifrons	5
Corynoneura nr. taris	7
Corynoneura sp. 3	7
Corynoneura sp. 4	7
Cricotopus bicinctus	8
Cricotopus vierriensis	6
(Cricotopus sp.)	7
Eukiefferiella sp.	8
Lopescladius sp.	5
Nanocladius sp.	3
Orthocladius annectens	6
Orthocladius carlatus	6
Orthocladius lignicola	6
(Orthocladius sp.)	6
Parakiefferiella sp. A	6
Parakiefferiella sp. 4	6
Parametriocnemus lundbecki	5
Rheocricotopus robacki	6
Rheocricotopus tuberculatus	6
Rheosmittia sp.	6
Synorthocladius semivirens	6
Thienemanniella fusca group	6
Thienemanniella xena group	6
Tvetenia discoloripes group	5
Tvetenia paucunca	5
Unniella multivirga	5

Appendix E

Biotic Index - Upper Three Runs Creek Biomonitoring
March 1990 - July 1991

Taxon	Biotic Index
-------	--------------

Chironomidae - Chironomini	
Cryptochironomus blarina	8
Cryptochironomus fulvus group	8
Kloosia dorsenna	6
Lauterborniella agrayloides	6
Microtendipes pedellus	6
Microtendipes nr. rydalensis	6
Pagastiella ostanta	6
Paracladopelma loganae	7
Paracladopelma undine	7
Paralauterborniella nigrohalteralis	8
Polypedilum convictum	6
Polypedilum fallax	5
Polypedilum halterale	8
Polypedilum illinoense	8
Polypedilum scalaenum	8
Pseudochironomus sp.	5
Stenochironomus hilaris	5
Tribelos jucundum	5
Xylotopus par	2

Chironomidae - Tanytarsini	
Cladotanytarsus sp.	7
Rheotanytarsus distinctissimus group	6
Rheotanytarsus exiguus group	6
Tanytarsus glabrescens group	6
Tanytarsus sp. I Cantrell	6
Tanytarsus sp. IX Rutter	6
Tanytarsus sp. XII Rutter	6
Tanytarsus sp. XIII Rutter	6
Tanytarsus sp. XIV Rutter	6
Tanytarsus sp. XVI Rutter	6
(Tanytarsus sp.)	6

Appendix E

Biotic Index – Upper Three Runs Creek Biomonitoring
March 1990 – July 1991

Taxon	Biotic Index
CRUSTACEA	
Asellus sp.	
Cambarinae	
Palaemonetes paludosus	
ANNELIDA (worms)	
(Naididae)	
Pristina/Stephensoniana sp.	
Spirosperma sp.	
(Tubificidae)	
MOLLUSCA (snails, clams)	
Campeloma sp.	
Elliptio sp.	
Ferrissia sp.	
Gillia altilis	
ACARI (water mites)	
Hydracarina	

Taxa in parentheses were too immature to allow more definitive identification or were damaged.

APPENDIX F

Taxonomic Identification Modifications

The following species have been more definitively identified due to the availability of additional specimens and/or taxonomic references during the study period.

<u>Previously Identified As</u>	<u>Currently Identified As</u>
Ephemerella invaria/rotunda	E. catawba group
Isoperla nr. nana	I. holochlora/lata
Stenelmis markeli	Stenelmis crenata
Stenelmis sinuata	S. convexula
Orthocladius curtiseta	O. carlatus
Symposiocladius lignicola	Orthocladius lignicola
Caecidotea sp.	Asellus sp.

END

**DATE
FILMED**

12 / 8 / 92