DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DOE/ER/60715--3

DE93 000896

SPATIAL SCALING: ITS ANALYSIS AND EFFECTS ON ANIMAL MOVEMENTS IN SEMIARID LANDSCAPE MOSAICS

Final Report

for Period 1 September 1988 - 31 May 1992

John A. Wiens

Department of Biology
Colorado State University
Fort Collins, CO 80523

September 1992

OCT 07 1992

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

Prepared for

THE U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-88ER60715

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Abstract. The research conducted under this agreement focused in general on the effects of environmental heterogeneity on movements of animals and materials in semiarid grassland landscapes, on the form of scale-dependency of ecological patterns and processes, and on approaches to extrapolating among spatial scales. The findings are summarized in a series of published and unpublished papers that are included as the main body of this report. We demonstrated the value of "experimental model systems" employing observations and experiments conducted in small-scale microlandscapes to test concepts relating to flows of individuals and materials through complex, heterogeneous mosaics. We used fractal analysis extensively in this research, and showed how fractal measures can produce insights and lead to questions that do not emerge from more traditional scale-dependent measures. We developed new concepts and theory to deal with scale-dependency in ecological systems and with integrating individual movement patterns into considerations of population and ecosystem dynamics.

Landscape ecology, which deals with spatially explicit mosaics, has not yet developed much formal theory. In contrast, there is a large body of theory dealing with patchiness in ecological systems, although little of it deals with patches located in a more complex spatial mosaic. Moreover, the expression of ecological patterns and their relation to underlying processes may change as a function of spatial or temporal scale. Our research has focused on understanding the effects of spatial heterogeneity on ecological systems and how these effects may change with scale. In particular, we considered (a) how the explicit locational relationships of patch elements in spatial mosaics influence the movements of individual organisms; (b) how these individual-level mechanisms translate into features of higher ecological levels; (c) the form of scale-dependency of spatial patterns in mosaics; and (d) how this scale-dependency affects the linkage between individual mechanisms and the characteristics of ecological systems. Throughout this work, we have emphasized a close interplay between the development of ecological theory and the testing and exploration of this theory in empirical and experimental studies. The substance of our work is reported in a series of publications and submitted manuscripts, which are appended as part of this report. We report our major findings by summarizing each of these papers, in sequence:

Wiens, J.A. 1989. Spatial scaling in ecology. *Functional Ecology* 3: 385-397.

I explore the scale-dependency of ecological phenomena. Both the variance of system attributes and the predictability of system behavior are related to spatial and temporal scale. The selection of an appropriate scale of investigation depends on the organisms and systems studied and the objectives of the study. Scale-dependent dynamics are not continuous, but vary discontinuously among "domains" of scale. Within a domain, pattern-process linkages are tight and system dynamics can be extrapolated among scales; between domains, system behavior becomes qualitatively different, as different processes act to influence patterns. Extrapolation between domains is difficult. I end by calling upon ecologists to investigate scaling as a subject of study in its own right.

Wiens, J.A., and B.T. Milne. 1989. Scaling of 'landscapes' in landscape ecology, or, landscape ecology from a beetle's perspective. *Landscape Ecology* 3: 87-96.

We advocate a "microlandscape" approach to studying the effects of landscape structure on ecological phenomena. We examine how the fractal structure of microlandscapes affects the movements of tenebrionid beetles. Beetles moved nonrandomly in fractal mosaics, avoiding areas in which the landscape structure fell within a certain range of fractal dimensions. These findings highlight scales and patterns to which beetles may be particularly responsive. The approach leads to different sorts of questions than are normally asked by ecologists.

Wiens, J.A. 1990. On the use of 'grain' and 'grain size' in ecology. *Functional Ecology* 4: 720.

I comment on the terminology of "grain" in ecology. The conventional use of the term, indicating a form of habitat response (or habitat structure) is dependent on the scale of investigation. It is better to define "grain" operationally, in terms of the finest scale of resolution of an investigation. This lessens the dependence on scale and may facilitate comparisons among investigations that are standardized according to scale.

Wiens, J.A. 1990. Habitat fragmentation and wildlife populations: the importance of autecology, time, and landscape structure. *Transactions of the 19th IUGB Congress, Trondheim 1989*: 381-391.

Habitat fragmentation, an area of major concern in conservation biology, involves more than simply a loss of habitat area and increasing isolation of habitat fragments. Ecological differences among species, past history and time lags in response to changes, and the landscape context of habitat fragments will all affect population and community responses to fragmentation. Because of these complex effects, long-term, experimental studies of fragmentation are needed.

Kotliar, N.B., and J.A. Wiens. 1990. Multiple scales of patchiness and patch structure: a hierarchical framework for the study of heterogeneity. *Oikos* 59: 253-260.

Habitat heterogeneity exists at multiple scales. We develop a hierarchical conceptualization of patchiness that emphasizes how patch contrast and patch aggregation vary among scales. The scale(s) at which patchiness is expressed is a function of both landscape structure and of the perception of organisms.

Wiens, J.A. 1990. Ecology 2000: an essay on future directions in ecology. *Revista Chilena de Historia Natural* 63: 309-315.

I consider future directions that may be particularly important in ecology. These are (1) determining how individual-based mechanisms underlie higher-level patterns; (2) analyzing the scale-dependency of ecological phenomena; (3) understanding how heterogeneity effects are expressed in complex landscape mosaics; and (4) avoiding drawing a distinction between basic and applied research.

- Wiens, J.A. 1992. Ecological flows across landscape boundaries: a conceptual overview. Pp. 217-235 *in* *Landscape Boundaries: Consequences for Biotic Diversity and Ecological Flows* (A.J. Hansen and F. diCastri, eds.). Springer-Verlag, New York.

I develop concepts for exploring what happens at boundaries between elements in landscape mosaics. I focus on the role of landscape structure in producing inequities in passive diffusion patterns, and on the active behavioral responses of animals to boundary conditions. These considerations lead to the development of a series of preliminary functions describing relations among the components of flows between landscape elements. Patch attributes, landscape characteristics, and features of organisms all contribute to these relations.

- Milne, B.T., M.G. Turner, J.A. Wiens, and A.R. Johnson. 1992. Interactions between the fractal geometry of landscapes and allometric herbivory. *Theoretical Population Biology* 41: 337-353.

The scale at which herbivorous mammals forage is determined in part by home range area, population density, metabolic rate, movement speed, transport costs, and ingestion rate. These features vary allometrically with body mass. We consider the interactions between allometric herbivory and landscape structure, expressed as the fractal dimensions of landscape patterns. In simulations, the transition from aggregated foraging to increased dispersion was modified by landscape pattern, and it was modified further by the scale at which the foragers operated.

- Johnson, A.R., J.A. Wiens, B.T. Milne, and T.O. Crist. 1992. Animal movements and population dynamics in heterogeneous landscapes. *Landscape Ecology* 7: 63-75.

The movements of organisms determine how they may respond to environmental heterogeneity. These responses, in turn, influence population dynamics, community structure, and ecosystem processes. We review theoretical approaches based on diffusion processes and animal-movement models. We emphasize (1) scaling relationships derived from percolation theory and fractal geometry; (2) extensions of traditional patch-based metapopulation models; and (3) individual-based, spatially explicit models governed by local rules. We also emphasize the usefulness of "microlandscape" investigations.

Johnson, A.R., B.T. Milne, and J.A. Wiens. in press. Diffusion in fractal landscapes: simulations and experimental studies of tenebrionid beetle movements. *Ecology*.

We consider the generation of anomalous diffusion movements in heterogeneous (fractal) landscapes, using both field studies and simulation models. In grassland mosaics, diffusion patterns changed at particular scale thresholds that may define domains.

Wiens, J.A., N.C. Stenseth, B. Van Horne, and R.A. Ims. in press. Ecological mechanisms and landscape ecology. *Oikos*.

We focus on how individual-level mechanisms operating in a heterogeneous mosaic may produce ecological patterns that are spatially dependent. We develop theory that relates population density and distribution among patches to (1) within-patch movements of individuals; (2) emigration from patches as a function of population density, patch configuration, patch context, and within-patch movement; and (3) loss of individuals as they disperse through landscape elements. This approach is then related to empirical studies employing "experimental model systems," small-scale analogs of larger systems that are amenable to experimental manipulation.

Wiens, J.A. in press. What is landscape ecology, really? *Landscape Ecology*.

On the basis of a review of papers published in *Landscape Ecology* volumes 1-5, I conclude that, as it has been practiced, landscape ecology is mostly conducted at broad spatial scales. Although the discipline involves a fair amount of conceptual work, these concepts are rarely formalized into mathematical theory, and are hardly ever exposed to experimental approaches. In general, landscape ecology is more a qualitative than a quantitative science. I argue that these characteristics must change if landscape ecology is to become a rigorous basis for resource management or ecological investigations.

Crist, T.O., D.S. Guertin, J.A. Wiens, and B.T. Milne. in press. Animal movement in heterogeneous landscapes: an experiment with *Eleodes* beetles in shortgrass prairie. *Functional Ecology*.

We examine how beetles respond to vegetational heterogeneity at several scales. Variations in vegetation structure have strong influences on beetle movement patterns, and these effects differ among beetle species. The fractal dimension of pathways, however, does not differ either among species or with variations in microlandscape heterogeneity. Simulation exercises show that both net displacement and fractal dimension differ from expectations based on a correlated random walk. These findings suggest that, although beetles of different size respond to environmental

structure on different scales, the complexity of their movement patterns is insensitive to scale and environmental structure, within the range of scales we studied.

Crist, T.O., J.A. Wiens, and B.T. Milne. submitted. Patch scaling and seed harvesting by ants. *Ecology*.

We examine seed harvesting by *Pogonomyrmex* ants over a hierarchical series of scales. Harvest rates differ among colonies according to macrovegetation features, and within colonies according to differences in fine-scale vegetational cover. There are additional effects of seed species, distance of seed sources from colonies, and patch types on local harvesting rates. Ant foraging pathways differ according to vegetation structure at both macro and micro scales. Predator activities may also modify movement pathways and activity times, and thence seed-harvesting rates.

Wiens, J.A., T.O. Crist, and B.T. Milne. submitted. On quantifying insect movements. *Environmental Entomology*.

We consider how insect movement pathways may be quantified, noting at the outset that investigations need to be scaled appropriately with regard to the organisms studied and the questions asked. We advocate the use of surveying instruments to record pathway locations, of renormalization procedures to remove spatial interdependence among observations, and of fractal measures to express pathway complexity.

Wiens, J.A., T.O. Crist, K.A. With, and B.T. Milne. submitted. Fractal patterns of insect movement in semiarid grasslands. *Nature*.

We suggest that the fractal dimension (D) and the intercept of the fractal regression (K) can be used to express both scale-dependent and scale-independent features of movement pathways. The fractal dimension of movement pathways of ants in semiarid grasslands is greater than that of beetles and grasshoppers occupying the same microlandscape mosaics. Pathways of grasshoppers differ in tortuosity among species and life stages, but those of congeneric beetle species do not. Over a range of scales from cm to m, pathway complexity is unrelated to variations in mosaic heterogeneity. The intercept of the fractal regression (K), which indexes the length of a pathway, varies widely among the taxa. Differences in D and K among taxa may relate to differences in the resources used by the insect and in the scales on which they view resource distribution.

reprints & preprints
removed.
ds

END

**DATE
FILMED**

12 / 8 / 92