DOE/ER/60846--3

DE92 018800

ANNUAL TECHNICAL PROGRESS REPORT

REPORT ON RESEARCH FOR:

*Colloids in Groundwater: Their Mobilization, Subsurface
Transport, and Sorption Affinity for Toxic Chemicals*
[DE-FG02-89 ER60846]

SUBMITTED BY:

Philip M. Gschwend
Associate Professor
Water Resources and Environmental Engineering
Department of Civil and Environmental Engineering
Massachusetts Institute of Technology
Cambridge, MA 02139

SUBMITTED TO:

Martin A. Langsam
U.S. Department of Energy
Chicago Field Office
9800 South Cass Ave
Argonne, IL 60439

SUBMITTED: July 10, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Introduction

We have been pursuing a series of studies to elucidate the roles of colloidal phases in subsurface environments. Most notably, we have discovered the critical role of secondary cementitious phases like goethite (Ryan and Gschwend, 1992). As part of this effort, we developed a new procedure for measuring "surface iron oxides" which is a great improvement over earlier methodologies (Ryan and Gschwend, 1991). Currently, we are developing a unifying model with which we can predict the mobilization of colloids to groundwater flowing through such porous media. Also, we have been able to show that groundwater samples contain colloidal phases in sufficient quantities and of suitable properties to enhance the mobile load of toxic compounds like polycyclic aromatic hydrocarbons (Backhus and Gschwend, 1990). Finally, we have synthesized our varied field investigations from sites in Massachusetts, New Jersey, Delaware, Nevada, Connecticut, and New York to yield a protocol for other researchers interested in evaluating colloids in groundwater (Backhus *et al.*, submitted). These accomplishments are discussed in more detail below. Through these efforts, we have become increasingly convinced of the central role played by colloidal phases in numerous subsurface phenomena controlling contaminant fates.

Research Accomplishments

Our originally stated objectives (proposal of July 1988) included: [1] development of a quantitative source function for colloid mobilization to groundwater, and [2] assessment of the most important colloidal phases to which nonpolar compounds sorb.

The first major accomplishment was to identify the central role played by secondary

cementitious phases like iron oxides in binding colloids to immobile subsurface solids (Ryan and Gschwend, 1992). This recognition of particular, intimate solid phase associations produces a very powerful basis from which to build an understanding of both how colloids are removed from groundwater flow as well as how they may be returned. In a related, originally unproposed effort, we developed a new method for measuring the abundance of secondary iron oxides in aquifer solids which utilizes Ti(III) to selectively extract this Fe fraction (Ryan and Gschwend, 1991). Now we can assess a critical property of subsurface solids which is needed for calculating the release of colloids. For example, although an anoxic sediment sample that we investigated had more total iron in it than an oxic sample from the same geological formation, this anoxic sample was less capable of retaining colloidal kaolinite. By applying our new Ti(III) extraction procedure, it was easy to show that the oxic sample, which contained only 70% as much total iron, actually had almost 2,000% as much secondary cementitious iron in the form critical to limiting colloid transport.

Based on our conceptualization of how colloids are intimately bound to immobile subsurface solids, we are in a position to postulate the system responses to changes in groundwater solution chemistry. For example, if solution pH were increased above the pH_{zpc} of goethite, all solids would be negatively charged and colloids should be mobilized; this is what we see (Figure 1)! If a reductant is applied to convert Fe(III) to Fe(II) and thereby dissolve goethite, one should see an enhanced dispersion of clay colloids upon subsequent flushes with suitable aqueous solutions (e.g., low ionic strength); again, this is precisely what we see! If an anionic surfactant solution is applied at levels known to reverse iron oxide surface charges, then colloids should be released into the flow. Again, that is precisely what we see. To synthesize

Figure 1. Enhanced rate of clay-sized particle release from New Jersey Pine Barrens aquifer solids as a function of solution pH (ionic strength held constant at 0.01 M). Also shown is the continuously more negative surface charge of the released colloids at higher pHs.

this variety of experimental data, we have been working to calculate the detachment energy which limits colloid release as a function of the solids involved and the solution ionic strength and pH (Ryan, 1992; Ph.D. Thesis). Correlating this result with observed colloid release rates appears very promising (Figure 2). Notably, the same concepts that dictate E_{detach} in our model also apply to the attachment process. Thus, our progress on quantitating colloid release from real-world subsurface solids may well also permit us to predict the surface chemical factor limiting heterocoagulation.

In addition to elucidating colloid transport phenomena, we have also sought to use the real-world colloids that we find to evaluate the tendency of organic compounds to sorb to those solids. Two major accomplishments were achieved. First, we have been able to add trace quantities of specially chosen polycyclic aromatic hydrocarbons to groundwater samples and ascertain whether colloidal sorption is significant (Backhus and Gschwend, 1990). This procedure only minimally changes the groundwater solution and colloid chemistry (e.g., no colloid preconcentration is used which might alter the colloid's properties). In some groundwater samples, colloid-bound forms of chemicals like benzo(a)pyrene would account for about half of this substance's mobile load. The second major accomplishment involved our demonstration that aromatic compound sorption to inorganic colloids can be important too (Backhus, 1990; Ph.D. Thesis). Sorption isotherms were linear over very wide ranges; sorption intensity was clearly a function of the sorbate's aqueous activity coefficient; and indications are that sorption may not be a function of solid phase composition (only surface area). These intriguing results have caused us to critically review literature in this area, and there appears to be support for our hypothesis that sorption of nonpolar compound to inorganic surfaces in groundwater situations

Figure 2. Correlation between clay-sized particle release rates and estimated detachment energies for a wide variety of flushing solutions which differ in ionic strength (I), pH, ascorbic acid content (H_2Asc), and combinations of these.

could be important (Backhus *et al.*, in preparation).

Our final major accomplishment has been to aggregate all our groundwater sampling experiences so as to recommend a set of procedures to other researchers interested in colloidal phases or colloid-bound chemicals in the subsurface (Backhus *et al.*, submitted). This document reflects field studies at six widely different sites, and it demonstrates the ubiquity of colloids in groundwater where subsurface geochemistry has recently changed.

Bibliography

- Backhus, D.A., and P.M. Gschwend. 1990. Fluorescent polycyclic aromatic hydrocarbons as probes for studying the impact of colloids on pollutant transport in groundwater. *Environ. Sci. Technol.*, 24:1214-1223.
- Backhus, D.A. 1990. *Colloids in Groundwater: Laboratory and Field Studies of Their Influence on Hydrophobic Organic Contaminants*. Ph.D. Dissertation. 198 pp.
- Ryan, J.N., and P.M. Gschwend. 1991. Extraction of iron oxides from sediments using reductive dissolution by titanium (III). *Clays and Clay Minerals*, 5: 509-518.
- Ryan, J.N., P.M. Gschwend. 1992. Effect of iron diagenesis on the transport of colloidal clay in an unconfined sand aquifer. *Geochim. Cosmochim. Acta*, 56 (in press).
- Ryan, J.N. 1992. *Clay Colloid Mobilization in an Iron Oxide-Coated Sand*. Ph.D. Dissertation.
- Backhus, D.A., J.N. Ryan, D.M. Groher, J.K. MacFarlane, and P.M. Gschwend. 1992. Sampling colloids and colloid-associated contaminants in groundwater. *Ground Water* (submitted).
- Ryan, J.N., and P.M. Gschwend. Effect of ionic strength, pH, Fe(III) oxide reduction, and surfactant adsorption on clay colloid release from an iron oxide-coated sand (in preparation).
- Backhus, D.A., P.M. Gschwend, and S.J. Eisenreich. Sorption of hydrophobic organic chemicals to inorganic surfaces: A review. (in preparation).

END

**DATE
FILMED
9/22/92**