The Regge Spectroscopy of Charmed Mesons[†]

J. Finkelstein
Department of Physics
Columbia University
New York, New York 10027

Stephen S. Pinsky
Department of Physics
Ohio State University
Columbus, Ohio 43210

Abstract

The implication of assigning the newly discovered particles to Regge trajectories is discussed. The Ademollo-Veneziano-Weinberg relation is used to determine the trajectory parameters and exchange degeneracy is used to predict a tensor particle at 2264 MeV and an axial vector particle at 2140 MeV.

[†] This work is supported in part by the U.S. Energy Research and Development Administration

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

The discovery of new heavy mesons produced in e^+e^- annihilation has recently been announced.^{1,2} In this note we will explore the assignment of those mesons to Regge trajectories and the consequent predictions of other mesons lying higher on those same trajectories. Our analysis will be independent of any detailed quark model of the new mesons, but will turn out to be reasonably consistent with the quark model calculation of ref. (3).

We begin by assuming that the particle detected in the $K\pi$, $K\pi\pi$ and $K\pi\pi\pi$ modes at 1870 MeV.^{1,2} (we take the average mass of the charged and neutral states) which is denoted by D is a pseudoscalar. The peak in the recoil spectrum to the D observed at 2010 MeV. we will assume represents a vector particle denoted by D^* . These two assignments together with the relationship between the vector and pseudoscalar trajectories proposed by Ademollo Veneziano and Weinberg (AVW)⁴ enable us to determine the parameters of the Regge trajectories.

The version of the rule proposed by AVW that is relevant to us can be stated as

$$\alpha_V(m_p^2) = \frac{1}{2} \quad (1)$$

where α_V is a vector trajectory and m_p is the mass of the appropriate pseudoscalar particle. This rule is empirically extremely successful, as can be seen in the two examples $\alpha_\rho(m_\pi^2) = \frac{1}{2}$ and $\alpha_K^*(m_K^2) = \frac{1}{2}$. In our case the rule becomes

$$\alpha_{D^*}(m_D^2) = \frac{1}{2} \quad (2)$$

Using $m_D = 1870$ MeV., the vector assignment of the D^* ; $\alpha_D^* (m_D^{2*}) = 1$ and a linear trajectory we determine the slope of D^* trajectory to be $.92 \text{ GeV.}^{-2}$. We note that this value is quite consistent with the value of the slope of most well established Regge trajectories. The ρ trajectories for example in a recent charge exchange experiment ⁵ was found to have a slope of $.93 \text{ GeV.}^{-2}$ though different determinations vary somewhat. ⁶ This is considerably different than the value of $.33 \text{ GeV}^{-2}$ that one obtains by placing the $\psi(3100)$ and the $J^P = 2^+ \chi(3550)$ on the same linear trajectory. If we are willing to believe a linear extrapolation of the D^* trajectory to vanishing invariant mass we obtain an intercept of -2.72 thus the trajectory take the form

$$\alpha_D^*(s) = -2.72 + .92 s \quad (3)$$

The next highest particle on the D^* trajectory is, if we assume weak exchange degeneracy, a tensor particle which we denote by D^{**} . Using the trajectory of the D^* given in eqn (3) we find $m_{D^{**}} = 2264$ MeV. This value is slightly below the value of 2330 MeV. predicted in ref. (3). By comparison with decays of known tensor mesons we would estimate the partial width of $D^{**} \rightarrow D \pi$ at about 8 MeV. and $D^{**} \rightarrow D^* \pi$ at about 42 KeV. The full width would therefore be in the 10 - 20 MeV. range since there is relatively little phase space for multi-pion final states. These estimates are sensitive to the assumed mass of the D^{**} ; for example $m_{D^{**}} = 2.33 \text{ GeV.}$ implies a partial width of $D^{**} \rightarrow D \pi$ of 17 MeV.

Even though exchange-degenerate Regge analysis is less reliable for unnatural than for natural spin parity trajectories, it is amusing to consider that if the D trajectory were parallel to the D^* trajectory given in equation (3), it would pass through the value one at a mass of 2141 MeV. and give rise to an axial vector meson D_A^* at that mass. A peak at just about this mass is in fact seen in the missing mass spectrum reported in ref. (1); however, this peak has an alternative explanation³ as a reflection from $D_V^* \bar{D}_V^*$ production. While perhaps both effects are present, the $D_A^* \bar{D}$ has the advantage over the other mode of being an s-wave decay.

It can be observed that two-body thresholds seem to be closely associated with particles or at least with prominent structure in the e^+e^- annihilation cross section. Compare the $D \bar{D}^*$ threshold at 3.74 GeV. with the ψ' at a mass 3.68 GeV; the $D \bar{D}_V^*$ threshold at 3.88 GeV. with the possible structure at 3.95 GeV., and the $D_V^* \bar{D}_V^*$ threshold at 4.02 GeV. with the structure at 4.03 GeV. There is also the analogous case of the $K \bar{K}$ threshold at .99 GeV. to be compared with the ϕ at a mass of 1.019 GeV. With our speculation on the masses of the D^{**} and D_A^* we note that the $D^{**} \bar{D}$ and $D^* \bar{D}_A^*$ thresholds are at 4.13 GeV. and 4.15 GeV. respectively close to the structure in e^+e^- cross section at 4.1 GeV., and the $D^{**} \bar{D}_A^*$ threshold is at 4.4 GeV. in agreement with the structure at 4.4 GeV.

There also appears to be an interesting parallel between the A_1 and the D_A^* since both are closely associated with kinematic enhancements. The A_1 is suspected to be associated with a deck effect in the $p\pi$ channel. Similarly we estimate the D_A^* to have about the same mass as a kinematic reflection of $D^* \pi$.³ It may in fact be that these are dual descriptions.

The authors have benefited from conversation with G. Goldhaber, and would like to acknowledge the hospitality of the Aspen Center for Physics where this work was done.

References

1. G. Goldhaber, F.M. Pierre et.al., Phys. Rev. Lett. 37, 255 (1976).
2. I. Peruzzi et.al., SLAC - PUB 1776.
3. A. De Rujula, H. Georgi and S.L. Glashow, Harvard preprints, "Is Charm Found" and "Charm Spectroscopy via Electron-positron Annihilation".
4. M. Ademollo, G. Veneziano and S. Weinberg, Phys. Rev. Lett. 22, 83 (1969).
5. A.V. Barnes et.al., Phys. Rev. Lett. 37, 76 (1976).
6. V. Barger, XVII International Conference on High Energy Physics, London 1974 (P I-210).