DOE/PC/79938--T12

DE92 040285

Tenth Quarterly Report

January, February, March, 1990

Studies of Coupled Chemical and Catalytic Coal Conversion Methods

Contract 22-87PC79938
Department of Chemistry
The University of Chicago
Chicago, Illinois 60637

Leon M. Stock, Principal Investigator
Kuntal Chatterjee and Michael Ettinger
Research Staff

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

RECEIVED
USDOE/PET
50 APR 16 PM 1:03
ACQUISITION & ASSOCIATION

MASTER

Abstract

This report concerns our research on base-catalyzed coal solubilization and a new approach for hydrogen addition.

The work on base-catalyzed, chemical solubilization is continuing with Mr. Kuntal Chatterjee.

This report is focused on the hydrogenation research of Mr. Michael Ettinger. Specifically it deals with the use of arene chromium carbonyl complexes as reagents for the addition of dideuterium to coal molecules. In one phase of the work, he has established that the aromatic hydrocarbons in a representative coal liquid can be converted in very good yield to arene chromium carbonyl compounds. In a second phase of the work directly related to our objective of improved methods for catalytic hydrogenation, he has established that the aromatic constituents of the same coal liquid add dideuterium in the presence of added naphthalene chromium carbonyl.

The study is continuing.

Introduction.

When an arene is coordinated to $\text{Cr}(\text{CO})_3$, profound changes in its reactivity occur^{1,2}. Ring and benzylic hydrogen atoms exhibit enhanced acidity, and typical arene reactions such as electrophilic aromatic substitutions are quenched. In contrast, the ring carbon atoms are activated toward attack by nucleophiles¹. These changes in reactivity are manifestations of the forceful electron withdrawing nature of the $\text{Cr}(\text{CO})_3$ component, and of the ability of this group to stabilize charged intermediates. In addition, arenechromium tricarbonyl complexes have been proven to be good catalysts in the hydrogenation of dienes³. While most simple complexes such as toluenechromium tricarbonyl require the use of high temperatures and dihydrogen pressures, arenechromium tricarbonyl complexes with naphthalene, anthracene, and phenanthrene show remarkable activity under extremely mild conditions. An investigation is under way in which the chemistry of coal arene groups to which $\text{Cr}(\text{CO})_3$ groups have been attached is being studied. Coal arenechromium tricarbonyl complexes may prove to be good internal hydrogenation catalysts for the hydrogenation of surrounding molecules. Another aspect of the chemistry of coal arenechromium tricarbonyl complexes is the susceptibility of the complexed aromatic ring to nucleophilic substitution which would provide a means of alkylating the complexed aromatic rings to either enhance the reactivity of the rings or to provide bulky alkyl groups to disrupt the π - π ring interactions that play such an important role in holding coal molecules together. As a result, via hydrogenation of aromatic rings or via the chemical modification of aromatic rings, a very low severity hydrogen addition reaction may be realized.

Results and Discussion.

In order to expedite the initial stages of investigation, preliminary experiments are being carried out using a coal liquid distilled at approximately 850°C obtained from run 250D at the Wilsonville pilot plant using Illinois #6 coal. In contrast to solid coals and heavy distillates, this material is soluble in all common organic solvents. When aromatic carbon atoms are complexed to the $\text{Cr}(\text{CO})_3$ group, their ^{13}C chemical shift moves upfield in the range of 80-100ppm. Any complexation of the coal arene groups will be evident using this coal liquid because the baseline is very clean in this region. The ^{13}C NMR spectrum of the coal liquid is shown in Figure 1.

The first stage of research centered on the conversion of the aromatic hydrocarbons in a coal liquid into arenechromium carbonyl compounds. Reactions to form complexes with the coal liquid arenes using the method of Mahaffy and Pauson⁴ in which the arene is refluxed in dioxane with chromium hexacarbonyl required long reaction times and large excesses of chromium hexacarbonyl to achieve acceptable levels of complexation. To alleviate these problems, the use of an alternative reagent, triaminechromiumtricarbonyl, was investigated. This reagent is easy to synthesize⁵, equation 1, and has been used to obtain a number of organochromium complexes⁶. Reactions of the coal liquid with triamine-

chromiumtricarbonyl proved to be very successful. High yields of complexation with the coal liquid arene molecules were obtained to the point where the product was orange colored and would crystallize. The ^{13}C NMR spectrum of one such product is shown in Figure 1. The carbonyl resonance can be seen at 234 ppm as well as the shift in the aromatic carbons to 80-100 ppm.

The second stage of the investigation of the chemistry of the coal liquid arene chromiumtricarbonyl complexes will study the hydrogenation capacity of the complexes. Studies have been carried out using simple arenechromium tricarbonyl complexes^{1,7} as catalysts in the hydrogenation of methyl 2,4-hexadienoate in which the severity of reaction conditions varied from 150°C at 700 psi dihydrogen pressure for toluenechromium tricarbonyl to 27°C at 57 psi dihydrogen pressure for naphthalenechromium tricarbonyl. The proposed explanation for the dramatic increase in activity using polycyclic arenechromium tricarbonyl complexes is that there exist two Cr-C bonds with the bridgehead carbon atoms out of the six Cr-C bonds that are longer than the others with bond orders of less than one⁸. Thus, the longer bonds should therefore be more easily displaced by the incoming diene ligand in the rate determining step. This advantage is not realized in the case of the monocyclic arenechromium tricarbonyl complexes such as benzenechromium tricarbonyl where all of the Cr-C bonds are equal length with bond orders close to unity. The effect of solvent was also examined. Coordinating solvents such as tetrahydrofuran or acetone increased the rate of the hydrogenation reaction presumably by facilitating the initial ligand substitution step.

Hydrogenation reactions were carried out under mild conditions in a glass micro hydrogenation apparatus at pressures lower than 200 psi and temperatures less than 50°C using dideuterium gas. The use of dideuterium gas allows the addition of any deuterium to be detected by means of deuterium NMR. The first reactions were carried out with methyl 2,4-hexadienoate with dideuterium catalyzed by naphthalenechromium tricarbonyl in acetone as described in the literature in order to insure that the procedure and

apparatus were satisfactory. The first reaction shown in equation 2 yielded

the desired product methyl hex-3-enoate-2,5-d₂. A similar reaction was carried out in the presence of small amount of coal liquid to test whether or not the coal liquid would poison the catalyst. This reaction, shown in equation 3, proved to be successful yielding a 63% conversion of methyl

2,4-hexadienoate to methyl hex-3-enoate-2,5-d₂. The ¹H and ²H NMR spectra of the products are shown in Figure 2. It is evident that the presence of the coal liquid does nothing significant to inhibit the catalyst.

The next set of experiments were carried out using the coal liquid arenechromium tricarbonyl complexed material. Direct hydrogenation of the material proved to be unsuccessful. In part, this could be due to the fact that all of the possible active sites for hydrogenation may of been rendered unreactive by complexation with Cr(CO)₃. Experiments were carried out on a mixture of the coal liquid arenechromium tricarbonyl complex and unreacted

coal liquid, to provide a substrate, as shown in equation 4. A comparison of

the ^2H NMR spectra can be made between the unreacted coal liquid and the coal liquid arenechromium tricarbonyl reaction showed that no significant deuteration had taken place, Figure 3. Two possible reasons for this are that the coal liquid arenechromium tricarbonyl complex is not an active catalyst at the employed conditions, or that there are no suitable sites for hydrogenation in the unreacted coal liquid.

In order to test the catalytic activity of the coal liquid arenechromium tricarbonyl complex, experiments were carried out using methyl 2,4-hexadienoate as the substrate and the coal liquid arenechromium tricarbonyl complex as the catalyst as shown in equation 5. The ^2H NMR of the reaction product is shown in Figure 4.

Although the activity of the coal liquid arenechromium tricarbonyl complex as a hydrogenation catalyst appears to be small for methyl 2,4-hexadienoate and unreacted coal liquids under the mild conditions employed, it may well prove to be viable under moderately more severe conditions. In addition, the nature of the aromatic rings in this particular coal liquid may actually lead to very inactive coal liquid arenechromium tricarbonyl complexes, as many monocyclic complexes are. A slightly heavier distillate may provide a higher degree of polycyclic aromatics

yielding more active complexes while still affording us the conveniences provided by the use of a coal liquid.

In order to test the latter hypothesis that significant hydrogenation substrates do not exist in the coal liquid, a preliminary reaction was carried out using naphthalenechromium tricarbonyl, an active catalyst, and unreacted coal liquid as the substrate, equation 6. A comparison of the ^2H NMR spectra

of the unreacted coal liquid before and after the reaction, Figure 5, shows that some dideuterium addition has taken place. First, the relative aromatic deuterium content has decreased indicating that aliphatic deuterium content has increased. Second, the two major resonances of the unreacted coal liquid at approximately 1.54 ppm and 1.16 ppm are no longer the dominant resonances. The major resonances in the reacted coal liquid have shifted to approximately 2.27 ppm and 1.45 ppm. In addition, new peaks have appeared at 3.57 ppm and 2.71 ppm. Thus, at a preliminary level, it can be concluded that there are some hydrogenation sites available in the coal liquid.

Research in this area is continuing to test the capacity of coal liquid arenechromium tricarbonyl complexes to catalyze dihydrogen addition. At this point, a move towards more severe conditions and towards the use of another coal substrate appears warranted.

Another aspect of the chemistry of coal liquid arenechromium tricarbonyl complexes is the addition of carbon nucleophiles to the complexed aromatic rings. The electrophilic nature of the complexed arenes makes possible the coupling of carbon nucleophiles with the aromatic rings^{9,10}. Addition of the nucleophile to the aromatic ring produces stable η^5 -cyclohexadienyl

chromium complexes which can be oxidized to induce the loss of the chromium yielding an overall nucleophilic substitution for hydrogen. Alternatively, the intermediate can be protonated and the resulting cyclohexa-1,3-diene can be detached from the chromium yielding an overall nucleophilic addition with the reduction of one the aromatic double bonds.

In this future stage of the investigation, the chemical modification of aromatic rings in coal will be studied to see the effects on the break up of intermolecular forces that bind coal molecules together. Solubilization of highly aromatic coal has been attained by the alkylation of aromatic rings^{11,12}. In one set of experiments, Stock and his coworkers found that the highly aromatic, highly insoluble Pocahontas coal could be 90% solubilized in pyridine. This was achieved by alkylation, using sodium amide, of the aromatic ring with n-octyl groups to the extent of 2.9 octyl groups per 100 carbon atoms. The alkylation of aromatic rings in coal arenechromium tricarbonyl complexes with t-butyl groups, one of the most successful nucleophiles, along with the reduction of one of the aromatic double bonds should prove to be an extremely effective and efficient means of coal solubilization.

Experimental.

The coal liquid was obtained through Amoco Research Center from the Wilsonville Pilot Plant, Run 250D. The ^{13}C NMR spectra were recorded on a General Electric GN 300 operating at 300MHz. The ^2H NMR spectra were recorded on a Varian XL 1000 operating at 400MHz. The ^1H NMR spectra were recorded on the University of Chicago 500MHz spectrometer. Tetrahydrofuran and dioxane were freshly distilled from sodium benzophenone ketyl immediately before use. All solvents were deoxygenated by means of standard freeze-pump-thaw methods. KOH, acetone, $\text{Cr}(\text{CO})_6$, were used as received.

Triaminechromium tricarbonyl

Chromium hexacarbonyl (17.6g, 0.08mol), KOH (32.0g), ethanol (80ml), n-butanol (80ml) and water (25ml) were placed in a flame dried 500mL three neck RB flask fitted with a stirrer and a reflux condenser. The contents of the flask were warmed gradually to 150°C. Chromium hexacarbonyl, which sublimed initially, was washed out later by the solvent and no more sublimation took place. The reaction solution gradually turned red. After 4h of refluxing, the mixture was cooled, 150mL of concentrated aqueous ammonia solution was added and the mixture was slowly stirred for two hours. A yellow precipitate was filtered through a glass filter, washed with aqueous ammonia and ether.

Coal liquid arenechromium tricarbonyl

$(\text{NH}_3)_3\text{Cr}(\text{CO})_3$ (4.68g, 0.025mol), coal liquid (5g), and dioxane (25mLs) were loaded into a 100 mL flask provided with a magnetic stirrer, and a reflux condenser. The solution was refluxed on an oil bath for 48 hrs. The reaction mixture was filtered through Celite in a glass filter and the flask and Celite were washed with ether. The product was purified over a column of silica gel with a 1:1:4 CHCl_3 : CH_2Cl_2 : Hexane.

Napthalenechromium tricarbonyl

$C_{10}H_8Cr(CO)_3$ was prepared by a slightly modified procedure¹⁰ to that described by Peuson⁴. A mixture of naphthalen (5.3g, 0.042mol) and $Cr(CO)_6$ (4.4g, 0.020mol) were refluxed in dibutyl ether (100 mL), hexane (10mL), and THF (1mL) for 70 hrs under nitrogen. The heating bath was maintained at 160oC for 15 hrs and at 145oC for the remaining time. Solvent and excess naphthalene were removed under vaccum and the product purified over a column of silica gel with a 1:1:4 $CHCl_3$: CH_2Cl_2 : Hexane.

Hydrogenation reactions

Hydrogenation reactions were carried out on a 10mL micro hydrogenation apparatus. Typically, the reactants and dried, deoxygenated acetone were loaded into the reactor and it was evacuated and flushed with nitrogen three times, deuterium three times and then brought up to the final pressure at the desired temperature. At the end of the reaction time, the mixture was filtered through Celite in a glass filter and concentrated under vaccum.

References.

1. Semmelheck, M. F. J. Organom. Chem. Library 1976, 1, 371.
2. Jaouen, G. in 'Transition Metal Organometallics in Organic Synthesis', Vol. 2, H. Alper (ed.), Academic Press 1978.
3. Cais, M., Fraenkel, D., Weidenbaum, K. Coord. Chem. Rev. 1975, 16, 27.
4. Mahaffy, C. A., Pauson, P. L. Inorg. Synth. 1979, 19, 154.
5. Razuvaev, G. A., Artemov, A. N., Aladjin, A. A., Sirotkin, N. I.
J. Organometal. Chem., 1976, 111, 131.
6. Moser, G. A., Rausch, M. D. Synth. React. Inorg. Met. -Org. Chem., 1955,
280, 255.
7. Cais, M., Rejoan, A. Inorg. Chim. Acta, 1970, 4, 509.
8. Muir, K. W., Ferguson, G., Sim, G. A. Inorg. Chem. 1966, 5, 1851.
9. Semmelhack, M. F., Clark, G. R., Harrison, J. J., Thebtarononth, Y., Wulff, W.,
Yamashita, A. Tetrahedron, 1981, 37, 3957.
10. Kundig, E. P., Desobry, V., Simmons, D. P., Wenger, E. J. Am. Chem. Soc.,
1989, 111, 1804.
11. Gawlak, M., Carson, B., Ignasiak, T., Strausz, O. P., Preprints of papers,
International Conference on Coal Science, Maastricht, The Netherlands,
1987, 57.
12. Stock, L. M., Miyake, M. Energy Fuels, 1988, 2, 815.

Figure 1. (a) ^{13}C NMR Spectrum Of Wilsonville Run 250D Coal Liquids in CHCl_3 .
 (b) ^{13}C NMR Spectrum Of Coal Liquid Arene- $\text{Cr}(\text{CO})_3$ in CHCl_3

Figure 2. (a) ^1H NMR Spectrum of Methyl hexa-2,4-dienoate and Methyl hex-3-enoate-2,5- d_2 in CHCl_3 .

(b) ^2H NMR Spectrum of Methyl hex-3-enoate-2,5- d_2 in CH_2Cl_2 .

Figure 3 (a) ^2H NMR of Unreacted Coal Liquid.
(b) ^2H NMR of Coal Liq Arene- $\text{Cr}(\text{CO})_3$ Reaction Product.

Figure 4. ^2H NMR Spectrum of Equation 5 Product in CH_2Cl_2 showing unreacted Methyl hexa-2,4-dienoate and Coal Liq- $\text{Cr}(\text{CO})_3$.

Figure 5.(a) ^2H NMR Spectrum of Unreacted Coal Liquid in CH_2Cl_2
(b) ^2H NMR Spectrum of Equation 6 Product in CH_2Cl_2 .

END

DATE
FILMED

10/27/92