NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Resonances decaying to a photon and a $J(\psi)$ particle*

John Babcock and Jonathan L. Rosner

School of Physics and Astronomy
University of Minnesota
Minneapolis, Minnesota 55455

MASTER

Abstract

If $L = 1$ orbital excitations $^3P_{2,1,0}$ of the $J(\psi)$ particle exist (assuming it to be a $^3S_1$ quark-antiquark state), the decays $^3P_{2,1} \rightarrow J(\psi) + \gamma$, $J(\psi) \rightarrow (\text{lepton})^+ + (\text{lepton})^-$ are predicted to yield $W(^3P_{2,1}, \theta) \sim 1 + (0.4 \text{ to } 0.6) \cos^2 \theta$ and $W(^3P_{1,0}, \theta) \sim \sin^2 \theta$. Here $W(\theta)$ is the distribution in angle θ between the photon and either lepton in the $J(\psi)$ rest frame. The decay $^3P_0 \rightarrow J(\psi) + \gamma$ (which must have $W(^3P_0, \theta) \sim 1 + \cos^2 \theta$) is predicted to be suppressed.

Among the varied interpretations of the $J(\psi)$ particle of mass around 3.1 GeV, perhaps the most straightforward and widely publicized is its assignment as a $^3S_1$ state of a new quark c and the corresponding antiquark.² If this is true, the $c\bar{c}$ system must have orbital excitations, the lowest of which ($^3P_{0,1,2}$) should have appreciable transitions to $^3S_1 + \gamma$.³

In this letter predictions are made for the polarization of the $J(\psi)$ in the decays $^3P_{2,1} \rightarrow J(\psi) + \gamma$ and for the relative magnitudes of the transitions $^3P_{2,1,0} \rightarrow J(\psi) + \gamma$. These predictions are intended to determine whether the $J(\psi)$ has a single set of orbital excitations (as in the $c\bar{c}$ model), none (as in models where its vector nature is somehow "fundamental,") or more than one (as in more complicated color⁴ or six-quark⁵ models.)

The predictions assume that (i) $J(\psi)$ is the lowest $^3S_1$ state of $c\bar{c}$, i.e., a relative of the f , ω , and ϕ , (ii) the $^3P_{2,1,0}$ states may be classified in a badly broken symmetry along with the A_2 , A_1 , and δ , (iii) single-quark selection rules describe the decays, as is known to hold for lower-mass particles,⁶⁻¹⁰ (iv) reduced matrix elements for $(A_2, A_1, B) \rightarrow \gamma$ can be related to those for $(A_2, A_1) \rightarrow \rho\pi$ and $B \rightarrow \omega\pi$ by vector dominance.⁹⁻¹¹

The polarization of the $J(\psi)$ can be measured by its subsequent decay to lepton pairs. If θ is the angle between either lepton and the photon in the $J(\psi)$ rest frame as shown in Fig. 1, one then expects the following angular distributions $W(\theta)$:

(a) For $^3P_2 \rightarrow J(\psi) + \gamma$, $W(\theta) \sim 1 + (0.4 \text{ to } 0.6 \cos^2 \theta)$. This should be a prominent decay.

*Work supported in part by the Energy and Research Development Administration under Contract No. AT(11-1)1764.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

(b) For ${}^3P_1 \rightarrow J(\psi) + \gamma$, $W(\theta) \sim \sin^2 \theta$, corresponding to $J(\psi)$

polarized longitudinally with respect to the photon direction.

(c) For ${}^3P_0 \rightarrow J(\psi) + \gamma$, it is clear that $J(\psi)$ must be transversely polarized with respect to the photon direction. We predict a suppression of this matrix element.

These properties are sufficiently distinctive that we feel it important to draw attention to them as early as possible. The cascade process $\psi' (3.7) \rightarrow {}^3P_{2,1,0} + \gamma \rightarrow J(\psi) + 2\gamma$ does not seem to be present,¹² suggesting that the ${}^3P_{2,1,0}$ states may be above $\psi' (3.7)$ or may not be composed of the same quarks. Instead, we suggest that in all processes in which the $J(\psi)$ has been observed (through its decay into lepton pairs $\ell^+\ell^-$) one look directly for resonances in the $\ell^+\ell^-\gamma$ system. We now prove that under the assumptions (i) - (iv), such resonances will have the properties (a) - (c).

The decays of $(q\bar{q}, L=1)$ mesons to $(q\bar{q}, L=0)$ mesons and photons have been discussed previously for the "light" quarks u, d, s .^{7,8} The processes which will concern us directly are the heavy-quark analogues of $A_2 (J^{PC} = 2^{++}) \rightarrow \gamma \rho$, $A_1 (1^{++}) \rightarrow \gamma \rho$, and $\delta (0^{++}) \rightarrow \gamma \rho$. In the limit of $U(4)$ symmetry (which we shall adopt) the matrix elements for the heavy-quark transitions will be expressible in terms of the same reduced matrix elements that affect the above A_2, A_1 , and δ transitions as well as the decays $A_2 \rightarrow \gamma \pi$, $A_1 \rightarrow \gamma \pi$, and $B \rightarrow \gamma \pi$. These relations are summarized in Table I. The matrix elements of the dipole operator $D_+^{e.m.}$ enter into partial widths as⁸

$$\Gamma(A, \lambda \rightarrow \gamma_{\lambda=1} + \{J(\psi) \text{ or } \pi\}) = \frac{e^2}{\pi} \frac{b_Y^3}{2J_A+1} \sum_{\lambda} | \langle A, \lambda | D_+^{e.m.} | J(\psi) \text{ or } \pi, \lambda=1 \rangle |^2. \quad (1)$$

Here the labels $\lambda, \lambda=1$ correspond to spin projections along the direction of photon emission in the rest frame of the parent particle A ; E_γ is the photon energy in this frame. The reduced matrix elements in Table I, $D(W_Y, W_Y^2)$, correspond to

$$D(0,0) \longleftrightarrow (8,1) + (1,8), \quad \Delta L_Z = 1 \quad (2)$$

$$D(1,1) \longleftrightarrow (3, \bar{3}), \quad \Delta L_Z = 0 \quad (3)$$

$$D(1,0) \longleftrightarrow (8,1) - (1,8), \quad \Delta L_Z = 1 \quad (4)$$

in the notation of Refs. 7 and 8. Also shown are the (normalized) distributions in θ arising from each helicity.

Two simple relations based on vector dominance convert Table I to a powerful spectroscopic tool.

1. Consistency relation. In the last three processes of Table I, let us take the photon off shell to the nearest vector meson pole and demand that the results for $A_2^+ \rightarrow \rho^0 \pi^+$, $A_1^+ \rightarrow \rho^0 \pi^+$, and $B \rightarrow \omega \pi$ be consistent with those obtained by calculating the matrix element of Q_5 .^{6,13} These results are expressible in terms of two reduced matrix elements rather than three.¹⁴ Hence there must exist one relation among the $D(W_Y, W_Y^2)$.¹⁰ This relation is^{9,11}

$$D(1,0) = \frac{3\sqrt{2}}{4} D(0,0), \quad (5)$$

whose consequences are also shown in Table I.^{15,16} Eq. (5) has two interesting consequences:

(1a) The decay ${}^3P_1 \rightarrow \gamma + J(\psi)$ leads to a pure $\sin^2 \theta$ distribution in lepton pairs.

(1b) The decay ${}^3P_2 \rightarrow \gamma + J(\psi)$ is of the form

$$W({}^3P_2, \theta) \sim (1 + \cos^2 \theta) + r \sin^2 \theta, \quad (6)$$

where

$$0 \leq r \leq \frac{7}{4} \quad (7)$$

In particular a pure $\sin^2 \theta$ distribution is excluded for the ${}^3P_2$ decay.¹⁷

2. Proportionality relation. The reduced matrix elements $D(0,0)$ and $D(1,0)$ refer to transitions with $\Delta L_z = \pm 1$ while $D(1,1)$ refers to transitions with $\Delta L_z = 0$. Again assuming vector dominance, the ratio of the $\Delta L_z = 0$ to $\Delta L_z = \pm 1$ reduced matrix elements may be related to that involved (say) in $B \rightarrow \omega \pi$. Empirically the $\Delta L_z = 0$ pionic transitions from $L = 1$ to $L = 0$ multiplets all are very small compared to $\Delta L_z = \pm 1$ transitions.¹⁸ Thus we are led to conjecture that

$$D(1,1) \simeq 0 \quad (9)$$

(A similar relation indeed is roughly compatible with data on baryon photoproduction.⁹) This, taken together with Eq. (5), has two further consequences:

(2a) The decay ${}^3P_0 \rightarrow \gamma + J(\psi)$ is forbidden.

(2b) The angular distribution in the decay ${}^3P_2 \rightarrow \gamma + J(\psi) \rightarrow \gamma + \ell^+ + \ell^-$

is

$$W({}^3P_2, \theta) \sim 1 + \frac{3}{5} \cos^2 \theta \quad (10)$$

This is close to $1 + \cos^2 \theta$ (the characteristic distribution for ${}^3P_0$ decay) but probably can be distinguished from the ${}^3P_0$ transition by its being considerably stronger.

Finally, using vector dominance and fits¹⁹ to the processes $A_2 \rightarrow \rho \pi$, $B \rightarrow \omega \pi$ etc., one can obtain quantitative estimates of partial widths. These are sensitive to a number of assumptions we have not made previously, such as $Q(c) = 2/3$, the particular kinematic factor in Eq. (1) (which is replaced by $p_\gamma^3/m_{\text{quark}}^2$ times a Gaussian cutoff factor in most quark models), and the masses chosen for the ${}^3P_J$ states.

The predictions for partial widths are summarized in Table II for various choices of mass of the decaying $c\bar{c}$, $L = 1$ state. The fact has been used^{19,20} that $\Gamma\{B \rightarrow \omega (\lambda_\omega = 0) + \pi\}$ does not vanish precisely, and that $D(1,1)$ is not exactly zero. Note the wide range of variation of predicted widths for different states: the ${}^3P_2 \rightarrow {}^3S_1 + \gamma$ width is roughly 40 times the ${}^3P_0 \rightarrow {}^3S_1 + \gamma$ width. (The latter is rather sensitive to the exact choice of parameters.) For comparison, we have listed two earlier predictions³ in which the differences among various $L = 1$ states are not taken into account.

The fact that $D(1,1)$ is nonzero in Table II leads to a slight alteration of the prediction (10):

$$W({}^3P_2, \theta) \Big|_{D(1,1) \neq 0} \sim 1 + 0.4 \cos^2 \theta \quad (11)$$

since $D(1,1)$ and $D(0,0)$ interfere constructively in the $\lambda = 1$ amplitude.

The relative enhancement of $^3P_2$ decays may be even more prominent than shown in Table II as a result of the production process. If the ^3P_J states are produced via electromagnetic cascades from a $^3S_1$ parent, one might expect similar ratios of reduced matrix elements, leading to $^3P_2$ predominance. Similarly, applying vector dominance once again to relate $^3P_J \rightarrow J(\psi) + \gamma$ to $^3P_J \rightarrow \gamma + \gamma$, one finds that the dominant $L = 1$ state produced in two-photon processes will be the $^3P_2$ with $\lambda = 2$. The most naive $U(4)$ estimate based on Table I gives $\Gamma(^3P_2 \rightarrow \gamma + \gamma) = O(1 \text{ MeV})$. However, the $\gamma - J(\psi)$ coupling seems to be about $1/\sqrt{3}$ of that predicted by $U(4)$.^{1,2} Hence one might expect suppression of single-photon rates by about 3 and two-photon rates by about 9 from those quoted above.

To summarize: if the $J(\psi)$ is a heavy quark-antiquark composite, one expects distinctive variations in partial widths and angular distributions of the lowest resonances decaying to $J(\psi)$ and a photon. The dominant state should have spin 2, and should decay to $J(\psi) + \gamma$ primarily (but not entirely) in a state of helicity 2.

We thank Professors Murray Gell-Mann and Frederik Zachariasen for extending the hospitality of the California Institute of Technology during the inception of this work.

References

1. J. J. Aubert, et.al., Phys. Rev. Letters 33, 1404 (1974); J.-E. Augustin, et.al., Ibid., 1406 (1974); C. Bacci, et.al., Ibid., 1408, 1649(E) (1974); W. Braunschweig, et.al., Phys. Letters   , (1975).

2. S. L. Glashow, J. Iliopoulos, and L. Maiani, Phys. Rev. D2, 1295 (1970); C. E. Carlson and P. G. O. Freund, Phys. Letters 39B, 349 (1972); G. Altarelli and L. Maiani, Rome preprint, 1974 (unpublished); M. K. Gaillard, B. W. Lee, and J. L. Rosner, Rev. Mod. Phys. 47, (1975); Thomas Appelquist and H. David Politzer, Phys. Rev. Letters 34, 43 (1975); A. De Rujula and S. L. Glashow, Ibid., 46 (1975).
3. These transitions have been discussed by many authors, among them C. G. Callan, et.al., Phys. Rev. Letters 34, 52 (1975); Thomas Appelquist, et.al., Ibid., 365 (1975); and E. Eichten, et.al., Ibid., 369 (1975). To our knowledge no such sources of monochromatic photons have as yet been identified.
4. O. W. Greenberg, Phys. Rev. Letters 13, 598 (1964); M. Y. Han and Y. Nambu, Phys. Rev. 139, B1006 (1965); M. Gell-Mann, Acta Physica Austriaca Suppl. IX, 733 (1972); I. Bars and R. D. Peccei, Phys. Rev. Letters 34, 985 (1975).
5. R. Michael Barnett, Phys. Rev. Letters 34, 365 (1975); H. Harari, Stanford Linear Accelerator Center report SLAC-PUB-1568, March, 1975 (unpublished).
6. Frederick J. Gilman, Moshe Kugler and Sydney Meshkov, Phys. Rev. D9, 715 (1974).
7. A. J. G. Hey and J. Weyers, Phys. Letters 48B, 69 (1974).
8. Frederick J. Gilman and Inga Karliner, Phys. Rev. D10, 2194 (1974).
9. John Babcock and Jonathan L. Rosner, California Institute of Technology report CALT-68-485, March, 1975, submitted to Annals of Physics.
10. J. L. Rosner, rapporteur's talk at XVII International Conference on High Energy Physics, London, England, July 1-10, 1974, proceedings edited by J. R. Smith, Rutherford High Energy Laboratory, Chilton, England, 1974, p. II-171.

11. Wesley P. Petersen, University of Minnesota report, March, 1975, submitted to Phys. Rev.
12. Robert Hofstadter, invited talk at Session GD, Washington Meeting of the American Physical Society, April 30, 1975.
13. A. J. G. Hey, J. Rosner, and J. Weyers, Nucl. Phys. B61, 205 (1973).
14. Richard H. Capps, Phys. Rev. 165, 1899 (1968); L. Micu, Nucl. Phys. B10, 521 (1969); E. W. Colglazier and J. L. Rosner, Nucl. Phys. B27, 349 (1971).
15. For decays to an $L = 0$ multiplet and a pion, there is one reduced matrix element with $\Delta L_z = \pm 1$ and one with $\Delta L_z = 0$. Photonic decays, in order to be consistent with this picture and with vector dominance, thus must be described by only one $\Delta L_z = \pm 1$ reduced matrix element rather than by two. This is achieved precisely via Eq. (5).
16. Another way to obtain Eq. (5) very simply is to consider the decay $D(J^{PC} = 1^{++}) \rightarrow \rho + \gamma$. If vector dominance applies, this process is dominated by $D \rightarrow \rho \rho$. As shown by C. N. Yang, Phys. Rev. 77, 242 (1950), a spin-1 particle cannot decay to two identical transversely polarized spin-1 particles. The $\lambda = 0$ decay amplitude for $D \rightarrow \rho \gamma$ (and hence also for $^3P_1 \rightarrow J(\psi) + \gamma$) must then vanish, leading again to Eq. (5).
17. In an experiment of this type, in which one integrates over production variables, one cannot separate $\lambda = 0$ and $\lambda = 2$ from one another in $^3P_2$ decay.
18. See, e.g., Colglazier and Rosner, Ref. 14, and Gilman, et.al., Ref. 6.
19. We use the specific numbers quoted by J. Rosner, Physics Reports 11C, 189 (1974).
20. S. U. Chung, et.al., Phys. Letters 47B, 526 (1973).

Table I. Matrix elements of $D_+^{c.m.}$ for decays

$(A, \lambda) \rightarrow \gamma + (B, \lambda-1)$. Multiply by overall factor of $Q(c)/4$ for $c\bar{c}$ decays and by overall factor of $1/24$ for A_2 , A_1 , and B decays.

State A, B	λ	$W(\theta)$	$D(0,0)$	$D(1,1)$	$D(1,0)$
$c\bar{c}(^3P_2);$ $c\bar{c}(^3S_1)$	2	$\frac{3}{8}(1+\cos^2\theta)$	$\begin{cases} \sqrt{6} \\ 2\sqrt{6}^a \end{cases}$	0	$4\sqrt{3}/3$
	1	$\frac{3}{4}\sin^2\theta$	$\sqrt{3}$	$-\sqrt{3}$	0
	0	$\frac{3}{8}(1+\cos^2\theta)$	$\begin{cases} 1 \\ 0^a \end{cases}$	-2	$-2\sqrt{2}/3$
$c\bar{c}(^3P_1);$ $c\bar{c}(^3S_1)$	1	$\frac{3}{4}\sin^2\theta$	$\sqrt{3}$	$\sqrt{3}$	0
	0	$\frac{3}{8}(1+\cos^2\theta)$	$\begin{cases} \sqrt{3} \\ 0^a \end{cases}$	0	$-2\sqrt{6}/3$
$c\bar{c}(^3P_0);$ $c\bar{c}(^3S_1)$	0	$\frac{3}{8}(1+\cos^2\theta)$	$\begin{cases} \sqrt{2} \\ 0^a \end{cases}$	$\sqrt{2}$	$-4/3$
$c\bar{c}(^1P_1);$ $c\bar{c}(^1S_0)$	1	-	$\sqrt{6}$	0	0
$A_2^+; \pi^+$	1	-	$\begin{cases} 0 \\ 3\sqrt{3}^a \end{cases}$	$3\sqrt{3}$	$2\sqrt{6}$
$A_1^+; \pi^+$	1	-	$\begin{cases} 0 \\ 3\sqrt{3}^a \end{cases}$	$-3\sqrt{3}$	$2\sqrt{6}$
B; π	1	-	$\sqrt{6}$	0	0

^{a)} Second line, where shown, indicates coefficients subject to the constraint $D(1,0) = (3\sqrt{2}/4) D(0,0)$ [Eq. (5) of text].

Table II. Partial widths based on Eqs. (1) and (5) and Table I for $^3P_J \rightarrow \gamma + J(\psi)$ and $^1P_1 \rightarrow \gamma + ^1S_0$. Reduced matrix elements estimated from decays such as $A_2 \rightarrow \rho\pi$, $B \rightarrow \omega\pi$ using vector dominance and $O(4)$ symmetry. Use of experimental $\gamma - J(\psi)$ coupling would suppress predicted numbers (not in brackets) by a factor of 3.

$c\bar{c}$ state ↓	$m(L=1)$ GeV →				
	3.4	3.5 (a)	3.5 (b)	3.7	4.0
$^3P_2$	1.2 MeV	[3 MeV]	[240 keV]	8.6 MeV	26 MeV
$^3P_1$	100 keV	[3 MeV]	[240 keV]	730 keV	2.3 MeV
$^3P_0$	34 keV	[3 MeV]	[240 keV]	250 keV	770 keV
$^1P_1$	400 keV	[3 MeV]	[240 keV]	2.9 MeV	9.0 MeV

(a) Appelquist, et.al., Ref. 3

(b) Eichten, et.al., Ref. 3

Figure Caption

Figure 1: Definition of angle θ in decays $^3P_J \rightarrow J(\psi) + \gamma \rightarrow \ell^- + \ell^+ + \gamma$.