DOE/ER/60245--T1

DE93 001748

ANNUAL PROGRESS REPORT
for
July 15, 1984 through January 15, 1986

RESEARCH ON THE SEASONAL SNOW OF THE ARCTIC SLOPE

from

Professor Carl S. Benson
Geophysical Institute
University of Alaska
Fairbanks, Alaska 99775

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

OVERVIEW

Introduction

This project deals with the seasonal snow on Alaska's Arctic Slope. It is concentrated on snow of the R₄D project area. However, an important aspect of this study is to relate the snow cover of this area with the rest of the Arctic Slope. The goals include determination of the amount of precipitation which comes as snow, the wind transport of this snow and its depositional pattern as influenced by drifting, the physical properties of the snow, the physical processes which operate in it, the proportions of it which go into evaporation, infiltration and runoff, and the biological role of the snow cover.

Seasonal Snow in Alaska

Alaska is famous for its glaciers which cover an area of nearly 10^5 km². However, glaciers cover only 5% of Alaska's total area (1.5×10^6 km²) all of which is subjected to seasonal snow. The physical properties of this snow, its thickness, the length of time it remains on the surface, and the physical processes which occur within it, above it, and below it, are of fundamental importance to the physical and biological environment. Alaskan snow varies significantly within its four climatic zones, but can be described according to three extreme types (Benson, 1982).

1. Tundra Snow - primarily above timberline and on the entire arctic slope.
2. Taiga Snow - primarily in the interior, between the Brooks Range and the Alaska Range

3. Maritime Snow - primarily in the coastal mountains of southeastern and southcentral Alaska.

Tundra Snow of the Arctic Slope

Alaska's seasonal snow lasts longest on the Arctic Slope where it is present for three quarters of the year. It differs from the hydrologically important snow of the Western United States in that its temperatures are lower, steeper temperature gradients occur in it, and there is less of it per unit area, but it lasts longer and enters more directly into the ecology (including most human activity) as snow itself rather than serving primarily as a cold storage water reservoir. This snow plays an essential role in physical and biological processes on the Arctic Slope.

In addition to protecting the tundra from damage by vehicles, snow also provides vital protection for microtine rodents during most of their lives. Furthermore, since the Arctic Slope is underlain by permafrost, its hydrology is significantly different from non-permafrost regions where snow dominates the hydrology (such as in parts of the western U.S.).

The importance of snow on the Arctic Slope as a dominant feature of the environment is being increasingly recognized. The U.S. Soil Conservation Service (SCS) has the prime responsibility for snow surveys in Alaska. It began with a network of ten snow courses in 1961 and now operates over 150 of them. The first measurements in the Arctic began with a Wyoming gage installed by the Geophysical Institute of the University of Alaska in 1975, today 10 or 12 of these gages operate in the Arctic. In spite of this, the SCS currently receives more requests for information about snow on the Arctic Slope than for any other region in Alaska.

From a scientific point of view, there is much to learn about snow on the Arctic Slope. Among the primary questions are:

- (1) Quantity - Most precipitation on the Arctic Slope comes in the form of snow and its quantity has been underestimated in the Weather Bureau records by a factor of about three. Overall means, in the Weather Bureau records, from 1961 to 1979 are 11.3 cm for Barrow and 14.4 cm for Barter Island; our corrected values are 25 cm for Barrow and 32 cm for Barter Island (Benson, 1982; Clagett and Benson, 1983).
- (2) Wind action - Snow on the Arctic Slope is moved by the wind to a significant but largely unknown extent. The directions of movement, the quantity of mass moved, the distance of movement and the formation of snow drifts are all subjects which need to be studied. As seen by the distribution of snow drifts, two wind directions, one from east-northeast, and the other nearly in opposition from the west, transport virtually all of the wind-blown snow on the Arctic Slope. In the west, at Atkasook, measurements indicate that the flux of wind blown snow from the eastern winds is about twice that from the western winds, and that the values vary significantly from year to year (Benson, 1982). In the southern part of the Arctic Slope, katabatic winds from the Brooks Range blow from the south; this is most pronounced in large river valleys such as the Killik River valley.
- (3) Distribution of snow - Significant precipitation gradients exist. From west to east, the snowfall at Barrow is about 80% of that at Barter Island. From north to south, the snowfall at Barter Island is about 60% of the 50 cm measured on the McCall Glacier 100 km south in the Romanzof Mountains. There are regions with minimum snow cover which lie on the slope between the foothills and the Arctic coast (Holmgren, Benson and

Weller, 1975). The boundaries of these regions, and the reasons for their existence need to be investigated. They are of special interest because of their use as prime calving ground by caribou (Kuropat and Bryant, 1980).

- (4) Physical Properties - The tundra snow on the Arctic Slope is dry and its wind-blown sastrugi surface strongly resembles the year-round surfaces of the Greenland and Antarctic Ice Sheets, or the winter snow surface of the adjacent Arctic Ocean. In many places the structure of the tundra snow resembles the top annual stratigraphic unit of the dry-snow facies of an ice sheet. It consists of a hard, wind-packed layer overlying a low density depth hoar layer.

Perhaps the most important feature of the snow pack is its control of the radiation balance by its simple presence or absence. The virtually unbroken tundra snow surface has an albedo (surface reflectivity) of about 80%. When the snow melts, in late May or early June, it disappears rapidly and the albedo drops to about 15% (Holmgren et al., 1975). This five-fold decrease in albedo occurs at the same time when the amount of incident radiation is rapidly increasing, and results in a dramatic change in the amount of absorbed radiation at the surface (Weller et al., 1972; Benson et al., 1975). Similar observations were made on the Canadian tundra by McFadden and Ragotski (1967) who estimated a sixfold increase in absorbed radiation at the surface when the snow melted away.

- (5) Physical processes (including gas migration in the snow, evaporation, infiltration and runoff) - The snow, the overlying air and the underlying soil and vegetation form a three-layer system in which heat and mass transfer processes are active. These processes lead to hard, wind-packed

layers and soft, low-density depth hoar layers (Benson, 1969). The latter are of special biological significance because animals which live under the snow (like lemmings) depend on it when they make tunnels. Animals which feed by digging through the snow (like Caribou) depend on it to yield food more easily than would hard-packed snow.

An especially interesting phenomenon is the behavior of trace gases in the snow. In particular, CO_2 is more concentrated within the snow pack (with highest values at the bottom) than in the overlying ambient air. Not only are the values higher, but they show short-term increases by a factor of two or three in the springtime (Kelley, Weaver and Smith, 1968; Coyne and Kelley, 1974). The reasons underlying these phenomena are not understood.

A significant, but unknown, part of the snow deposited on the Arctic Slope is lost directly to the atmosphere by sublimation and evaporation. Much of this loss occurs while the blowing snow is airborne (Schmidt, 1972; Tabler, 1976), the rest is lost from the surface, especially during spring after vegetation perforates the snow surface (Benson et al., 1975; Dingman et al., 1980).

Because the underlying ground is frozen, infiltration is severely limited and runoff in the streams, which are not fed by glaciers, is characterized by a large, single pulse during spring breakup (Dingman et al., 1980).

The above phenomena are involved in questions relating to the abiotic environment of natural biological systems and to man-made disturbances of the tundra. Roads and buildings interact with winds to cause variations in the pattern of snow drifts, this in turn modifies temperature and moisture

conditions at the base of the snow pack. In particular drifts formed along roads produce ponding, or wet areas parallel to the roads and the physical processes associated with this require attention. Impurities on the snow, such as road dust, will affect melting and evaporative rates. Unfortunately, we do not understand the undisturbed melt and evaporative losses. In order to deal with these problems it is necessary to know how much snow falls, and how it is moved by wind. In general, this is a difficult problem which requires continued development of theoretical models of snow transport and deposition (Schmidt, 1972; Tabler and Schmidt, 1973; Tabler, 1980), in addition to more field research.

Original Objectives

The original objectives of this research included a regional study of snow on the entire Arctic Slope. During the first year the scope was restricted to the R₄D area. The need for some comparison with other regions became clear as the work progressed and was discussed during the April R₄D meetings in San Diego. Some broadening of the scope will be presented below.

Mapping

The need for accurate topographic maps of the R₄D research area was apparent in this project as well as for all other R₄D projects. A mapping project was added to this project because of the investigator's experience with preparation of aerial photogrammetric maps. The ground control was to be surveyed and pre marked so that photos could be taken and mapping initiated in the fall of 1984.

ACCOMPLISHMENTS DURING 1984 - 85

Mapping

Ground control for aerial photogrammetry was established by precision surveying done by Glen Liston and helpers during July and August of 1984. The primary instrument used was a Zeiss "Elta 2" Total Station (theodolite - Electronic Distance measuring system). A total of 50 points were surveyed, 13 of which were pre-marked for aerial photogrammetry. The 50 points are listed here in seven subsets, including the azimuth point Δ "Lake" which is off the map.

1. The USGS Bench Mark: Δ Gal. This point has accurately determined latitude, longitude and altitude and served to locate the map and to define an azimuth. The azimuth was determined by sighting from Δ Gal to the USGS bench mark: Δ Lake.

The absolute coordinates of Δ Gal are: $68^{\circ}37'13.045''N$; $149^{\circ}16'53.509''W$,
Altitude, 968.04m.

The coordinates of Δ Lake are: $68^{\circ}29'22.396''N$; $149^{\circ}19'39.209''W$,
Altitude, 1502.05m.

The azimuth between Δ Gal and Δ Lake is a bearing of $187^{\circ}21'45.91''$
(from True North).

In our survey we have used Δ Gal as an origin point and assigned it coordinates:

$x = 10,000.000$ $y = 10,000.000$ $z = 968.045$ (all values in meters).

These values were chosen to avoid negative x and y values. The grid of coordinates shows on the map.

2. Six pre-marked photo-control points for the large maps of the entire area, scale 1:6000 with 5m contour interval.
3. Six pre-marked photo-control points for the small maps of the intensive area; scale 1:1000 with 1m contour interval.

The pre-marked points were marked by white panels of fabric in the form of a cross with arms 6 m long. The arms show on the photos and can be used to identify 12 m lines for scale. The white panels are each 4 m long and there is 4 m gap between them with the point in the center. The arms are oriented True N-S and E-W so they provide azimuth on the photos as well as scale.

4. Eleven surveyed points in the drainage valley lying between the Toolik and Kuparuk rivers. These were not pre-marked for aerial photography but are marked in the field by wooden stakes bearing the designation "T-K". They are spaced at intervals of 500 m in the drainage valley. Each of them is fastened to a steel rebar which goes more than 1 m into the ground; the rebar pieces have marked survey caps on them.
5. Twelve surveyed points in the Toolik drainage valley. These were not pre-marked for aerial photography but are marked in the field by wooden stakes bearing the designation "T". Each is fastened to marked steel rebar as with the T-K series.
6. Three NGS-Alaska Bench marks along the haul road.
7. Twelve extra points required to complete the survey.

The coordinates of these survey positions are listed in Table 1.

TABLE 1. Coordinates of survey points established in the field for the R₄D photogrammetric maps.

	X	Y	Z
1) <u>USGS Bench Mark Control</u>			
GAL	10,000.000	10,000.000	968.045
2) <u>Large Map Control Points</u>			
1	6620.440	14,303.380	832.261
2	10,342.548	15,236.551	828.282
3	10,764.286	11,344.228	864.040
4	10,601.854	7067.927	972.881
5	7055.405	6913.080	794.571
6	6615.277	10,948.864	787.614
3) <u>Intensive Map Control Points</u>			
S-1	8232.099	9932.757	899.385
S-2	9167.113	9968.324	936.082
S-3	9415.168	9157.225	938.842
S-4	9453.845	8500.804	946.183
S-5	8651.619	8208.378	903.245
S-6	8280.463	9298.622	893.319
4) <u>Toolik - Kuparuk Drainage Stakes</u>			
T-K Culvert	7591.206	12,510.083	830.202
T-K 1	7783.596	12,048.756	835.711
T-K 2	7976.125	11,587.096	840.799
T-K 3	8168.567	11,125.657	847.503
T-K 4	8360.932	10,664.432	856.453
T-K 5	8553.322	10,203.096	865.909
T-K 6	8614.109	9706.934	871.755
T-K 7	8674.912	9210.688	882.689
T-K 8	8735.707	8714.490	891.926
T-K 9	8893.039	8240.000	899.226
T-K 10	9050.412	7765.385	904.921

5) Toolik River Drainage Stakes

T-Culvert	8554.495	13,063.064	839.996
T-1	8690.366	12,581.379	846.518
T-2	8826.201	12,100.825	854.002
T-3	8962.054	11,619.704	861.590
T-4	9097.980	11,138.302	870.229
T-5	9233.719	10,657.577	881.087
T-6	9369.520	10,176.635	895.396
T-7	9505.393	9695.568	910.317
T-8	9641.251	9217.574	925.170
T-9	9777.005	8733.785	939.989
T-10	9912.840	8252.708	947.123
T-11	10,048.686	7771.590	955.215

6) NGS - Alyeska Bench Marks

H153	7876.283	12,598.361	843.822
J153	9236.657	13,485.151	852.722
K153	10,606.090	14,374.994	834.478

7) Extra Points Required to Complete Survey

TM	9081.070	9965.045	936.170
MS117 Sub Pt	7674.713	11,302.565	865.131
MS117 Gate	7220.838	12,413.415	860.868
Pt 1 Sub Pt	6869.021	13,658.660	840.630
E.T. Hill	8979.280	13,321.220	853.012
J. Hill	9323.033	13,544.071	853.012
K. Flats	10,459.595	14,272.483	838.621
S-4 A	9596.121	8375.936	953.982
S-4 B	10,268.485	7996.386	968.852
T-K5 A	8599.166	10,093.194	868.852
T-K5 C	8590.403	9900.429	871.987
T-8A	9727.422	8909.389	935.013

Vertical aerial photographs were taken after all the pre-marked survey points were established. Both black and white and color Infra Red photography was done. Because of persistent bad weather in August and September 1984 it was difficult to complete the surveying and the aerial photography. The black and white photos were flown twice before a usable set, free of snow cover, was obtained. The color I.R. photos were not as good as they could be because the time of maximum chlorophyll had passed. Therefore, color I.R. photos were taken again in the summer of 1985 with excellent results.

We worked closely with the photogrammetrists at North Pacific Aerial Surveys (NPAS) in Anchorage to construct stereomodels for the photogrammetry. All survey points were verified and internally-consistent stereo models permitted us to make several products.

1. Orthophoto negatives were prepared and orthophoto maps were made at the scales of 1:6000 and 1:1000.
2. Topographic contour maps were made of the overall area at scale of 1:6000 with 5 m contour interval. These contour maps were printed on top of the orthophoto maps. Two map sheets at this scale cover the overall area.
3. Topographic contour maps were made of the intensive study area at scale of 1:1000 with 1 m contour interval. These contour maps were printed on the 1:1000 scale orthophoto maps. Four map sheets at this scale cover the intensive area.

Field Research on Seasonal Snow: (1) Summary of Field Work

12-14 Jan: Snow depths were measured along selected probe lines. Snow pits were dug, yielding information on snow stratigraphy, density, and temperature. Wind slabs were identified and measured in the area and the

temperature at the snow-soil interface was measured at several places.

At Prudhoe Bay the ARCO Snow Fence drift was mapped, and stratigraphy and snow densities were determined within the drift. Snow depths in the general Prudhoe Bay area were measured while we were enroute to the R₄D area.

1-2 Mar: Glen Liston and Larry Hinzman worked together on data loggers for joint use in the projects of Kane and Benson. Wind erosion blocks were installed.

At Prudhoe Bay the ARCO Snow Fence Drift was mapped again and snow depths measured in the region.

24-30 Mar: 3 Km, E-W probe line was established, with measurements taken at 5 m intervals.

An Anemometer was installed and the volume of the large drift trap on North cliffs was measured.

A Wind-drift direction survey was conducted by measuring drift orientations.

Four, 500 m prob lines were measured at various locations within the research site.

Depth measurements and Ramm Hardness profiles were made at several other identifiable locations within site.

Snow pits were dug in the same general locations as during 12-14 Jan.

Photos were taken from selected sites.

6 Apr:

Flight from Fairbanks to Prudhoe Bay. Aerial photos were taken of prevailing snow drift patterns and drift directions.

These photographs showed the winter baseline snow depth-distribution within the R₄D site.

19-21 Apr:

Installation of a Meteorological Station was done by Larry Hinzman and Glen Liston. The station included pyranometer, albedometer, anemometer, wind vane, net radiometer, hygrometer, and thermometer.

Photos were taken from selected sites, of which six were primary sites used for panorama photos.

Snow survey work was continued.

9 May - 2 June:

The primary spring field work was done during this time. This included Anemometer maintainance, chart charge, time checks.

Periodic Wyoming Precip Gage time and data checks

10 May, Pyranometer installed

10 May - 1 June, Daily weather observations were made including daily estimation of percent snowcover across intensive and complete site.

11 May, Movie camera installed at T-K 9, looking down valley.

11,15,18,22 May, Snowpits were dug (see 12-14 Jan) to measure snow properties including the progression of the 0°C isotherm through the snow as melting progressed.

11 May - 1 June, Ground photographic survey network was established, covering entire site, with photos taken at 2-5 day intervals.

12-13 May, Depth and water equivalent survey was done across the entire T-K drainage basin. This was done cooperatively by Glen Liston and Larry Hinzman.

Samples were taken at 10 m intervals.

13-30 May, Daily photographs taken of 3 small, 10 m x 10 m plots, observing micro snow-melt characteristics.

15,19,24 May, Mapping of equivalent backbody temperature distribution across bare patches and in adjacent snow cover using Barnes Precision Radiation Thermometer Model PRT-5, while concurrently taking micro meteorological measurements.

16 May, Prudhoe Bay: ARCO Snow Fence Drift mapped.

17 May, OM-2, 35 mm, automatic camera was installed looking up T-K valley. 1 frame per hour. Film changed daily, 16 rolls were used between 17 May and 1 June.

18 May, Rebar was placed at T-K and T drainage stakes for late summer-early fall installation, and Ramm

Hardness profiles were measured in the snowpack.

19,22,24 May, Albedo measurements were made on various vegetative coverings and snow cover.

20 May, Second movie camera established looking up T-K valley.

21 May, Aerial photos taken of R₁₄D site, Mid-Melt.

23,28 May, Topographic Mapping Project ground control flags were removed.

23-31 May, Periodic Stream Gage readings were taken, and stream velocity profiles were done, cooperatively by Glen Liston and Larry Hinzman.

24-30 May, Water samples were taken cooperatively for analyses by Kaye Everett.

25 May, A new stream gage was installed, cooperatively by Glen Liston and Larry Hinzman, it was moved on 26 May and permanently established on 28 May.

28 May, Aerial photos taken of R₁₄D sites. End of Melt Period.

29,30 May, snow depth measurements were made on persistent drifts.

1 June, Pyranometer removed, OM-2 camera removed, Movie camera removed.

6-8 Sept, 6 foot long 5/8" diam. rebar was installed at T-K and T drainage stakes, marking caps were placed on rebar and 2" x 2" wooden poles were attached to them. Depth of the active layer measured. Soil samples were collected to obtain moisture content of active layer.

Field Research on Seasonal Snow: (2) Preliminary Results

The distribution of snow across the entire R₄D area was measured by combining several techniques. Snow depth was probed along selected traverses and pit studies were made to study extremes of snow types. Photographs were taken with automatic cameras set up to watch the progression of snow melt; the cameras included 35 mm black and white cameras and 8 mm color movie cameras. Aerial photographs were taken over the site on 6 April, 21 May and 28 May. From these photographs, taken from the air and from fixed points on the ground, and measurements of snow properties, maps were made of the snow cover. These maps showed the progression of snow melt and from them we are constructing a map of the water equivalent of the snow pack at its maximum. Figure 1 is an example of the snow distribution on 28 May.

The snow drifting is from south winds in the R₄D area. The normal East-West winds of the Arctic Slope begin about 20 km north of the site. The boundary between winds from the south and winds from the East and West is not well-known on the Arctic Slope. However, it is clear that the south winds are associated with the Brooks Range and do not extend far from it.

The sensitivity of snow drifting to topography is pronounced. Snow was thickest on lee slopes and thinnest on windward slopes. The windward slope leading up to the cliffs, north of the intensive study site, was blown clear of snow while the north-facing lee slopes had the largest drifts in the R₄D area. Some of the variability observed in Figure 1 would not have been easily related to topography if our detailed topographic maps were not available. This is especially evident immediately to the north of the intensive study site.

23 MAY 1965

G264-43a

100% snowcover

Vegetation showing,
> 50% snowcover

snow patches,
< 50% snowcover

SCALE: 1:17,000

N

170m 340m

Figure 1

Snow was highly variable in quantity and structure throughout the site. A well-developed depth-hoar layer was present nearly everywhere. Two very hard wind slabs were present in some places. The snow covered areas in Figure 1 correspond roughly to the wind slab locations. These wind slabs had integrated Ram hardness profiles exceeding 8000 joules. In some places without wind slabs the integrated hardness profiles were less than 300. (In wind-free parts of interior Alaska the integrated Ram profiles are less than 10 joules.) Density values reached maximums of 0.54 g cm^{-3} in some wind slabs and minimum values of less than 0.20 g cm^{-3} in depth hoar layers.

The variability in water equivalent on the ground is maximized because regions with deeper snow also had the higher density values. The water equivalent of the winter snow is being assessed to provide a total for the R₄D area. The average on the surface is about 15 cm water equivalent. In places with thick wind slabs the water content exceeded 20 cm water equivalent.

The amount of snow lost by evaporation is significant and appears in excess of 50%. Analyses are still underway to refine this value for evaporation. Micrometeorological measurements were made during the time before, during, and after the snow melt to contribute to this understanding. These measurements were part of the combined projects by Kane and Benson. Our measurements with the Barnes Radiation thermometer showed surface temperatures varying from 0°C over snow patches to more than 30°C on adjacent snow-free areas. Thus, the ability of the air to hold moisture varies by a factor of 5 over horizontal distances of only a few meters.

The depth-hoar layer referred to above was exceptionally well developed in many places. Vertical channels were apparent to a degree greater than expected. During the 1984-85 winter this channeling was better developed in the R₄D area than in interior Alaska. However, the depth hoar layer was

thicker in the interior. The heat flux from soil to the overlying snow ends on the Arctic Slope when the active layer is frozen. In interior Alaska, where permafrost is absent or deep, the heat flux from soil to snow continues all winter. The total amount of heat flow from soil to snow during the winter is one of the parameters we are currently determining.

~~PROPOSED RESEARCH FOR SIX-MONTH EXTENSION~~

~~Research on the Winter snow and its disappearance will be done in cooperation with the project on the active layer by Dr. Kane. Observations of the formation and evolution of the snow pack will be made throughout the winter. The distribution of wind slabs, thicker snow packs and depth hoar will be compared with their distributions measured during the 1984-85 winter.~~

~~Cooperative work on gas flow through the snow will be done with Dr. Kelley's project. This research is important for two reasons. First, it will contribute to better understanding of water vapor flux and depth hoar growth in the snow. Secondly, it will be directed at the problem of CO₂ variations in the snow. The problem of gas flow in the snow will be addressed by measuring CO₂, CO, and CH₄ under the snow and in the air above. In addition, SF₆ will be injected into the snow and detected as it diffuses away from its point source. This will supplement our measurements on the flux of water vapor through the snow pack. These measurements may shed light on the drying of vegetation under snowpack and possibly some on the behavior of small mammals living under the snow. These measurements will be done in cooperation with Dr. Kelley and are being budgeted through his proposal.~~

~~Based on our research last winter, we will use photogrammetry in combination with snow surveys and pit studies to determine the water~~

~~equivalent and structure of the snow pack. A snow probe was useful and~~

END

**DATE
FILMED**

12 / 1 / 92