Twelfth Quarterly Report

DOE/PC/79938--T14

July, August, September 1990

DE92 040287

Studies of Coupled Chemical and Catalytic Coal Conversion Methods

Contract 22-87PC79938
Department of Chemistry
The University of Chicago
Chicago, Illinois 60637

Leon M. Stock, Principal Investigator
Kuntal Chatterjee, Michael Ettinger, and Carlos Cheng
Research Staff

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

db
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ANIONIC FRAGMENTATION REACTIONS IN COAL. REACTIONS OF MODEL COMPOUNDS
IN STRONG BASES.

During the previous quarter, a new approach for the depolymerization of the coal macromolecule was tried.¹ This was aimed towards carbon-carbon bond cleavage in presence of strong bases. Such bond cleavage reactions are well known with the alkali metals.² Electron transfer reactions take place from the metals to the aromatic nuclei resulting in the formation of anion-radicals (or dianions) which subsequently undergo carbon-carbon bond cleavage.² In our work instead of using the alkali metals, we have used strong bases to cleave the carbon-carbon bonds in hydrocarbons, and have suggested that hydrocarbon elimination reactions occur.

Such anionic fragmentation reactions involving strong bases are not very well established. However, we have obtained circumstantial evidence that such bond cleavage reactions do take place in some coal systems. For example, when the high rank Lower Kittanning coal, PSOC 1197 was treated with Lochmann's base (equimolar mixture of n-butyllithium and potassium tert-butoxide) in refluxing heptane and quenched with ammonium chloride and methanol, the pyridine solubility of the product increased from 5% (raw coal) to 39%.³ A similar increase in solubility due to base treatment was also observed in a separate study for another high rank coal, Pocahontas No. 3 from the Argonne National Laboratory Premium Sample Program. The solubility of the coal increased from 5% to 32% after the base treatment. These results, together with the absence of literature studies on base promoted fragmentation reactions prompted us to perform reactions on some pure compounds to assess this concept in more detail.

This report will involve some results regarding base promoted fragmentation reactions in hydrocarbons. The base used for all the

reaction sequences was Lochmann's base and, as with the coal, the anionic hydrocarbon was quenched with dilute aqueous acid. For some cases, to trap the reactive anions, the basic material was also quenched with a deuterium oxide-methyl alcohol-d mixture or an alkyl halide to obtain the deuterated or alkylated products, respectively.

EXPERIMENTAL SECTION

Materials

The selected hydrocarbons for our study were either purchased from Aldrich Chemical Company or were synthesized by known procedures. The compounds chosen were bibenzyl, 1,1'-binaphthyl, 1,1,3-triphenylpropane, 9,10-diphenylanthracene and 9-benzyl-9,10-dihydroanthracene.

n-Heptane(Aldrich) was shaken with concentrated sulfuric acid and subsequently distilled. The purified solvent was stored in the presence of molecular sieve 5A. The other chemicals such as n-butyllithium (1.6 M solution in hexane), potassium tert-butoxide, hexamethylphosphoramide (99%), deuterium oxide (99.9 atom % D) and methyl alcohol-d (99.5 + atom % D) were used as received from the Aldrich Chemical Company.

Reaction Procedure

The base treatment of the pure compounds were carried out essentially as reported earlier for the coal systems.¹ n-Heptane (45-55 mL), potassium tert-butoxide (2.52 g, 22.5 mmoles) and n-butyllithium (15 mL 1.6 M solution in hexane, 24 mmoles) were added to a flame dried flask. The mixture was stirred at room temperature for 15 min, the pure compound (8-10 mmoles) was added and the solution was refluxed for 6 h. The color of the solution changed from dirty yellow to dark red or black on addition of the hydrocarbon. After 6 h of reflux the black reaction mixture was cooled and

dilute hydrochloric acid was added very slowly to the mixture at 0°C. The reaction was usually vigorous. The acidic reaction mixture was extracted with ether, washed with saturated sodium chloride solution and was dried over anhydrous magnesium sulfate. The ether extract was concentrated and analyzed by conventional GC and GC-MS procedures.

Synthesis of 9-benzyl-9,10-dihydroanthracene

The compound was synthesized by the method of Normant et al.⁴ 9,10-Dihydroanthracene (3.6 g, 20 mmole), hexamethylphosphoramide (15 mL) and tetrahydrofuran (45 mL) were added to a flame dried flask at -78°C under dinitrogen. The n-butyllithium reagent (16 mL, 24 mmole) was slowly added to the solution and the color of the solution turned dark red. After 2 h of stirring at -78°C, benzyl bromide (3.42 g, 20 mmole) was added to the anion over 15 minutes at -78°C and the mixture was slowly warmed to 0°C by replacing the dry-ice/acetone bath with an ice-water bath. After 12 h, the reaction mixture was acidified by dilute hydrochloric acid and the product was extracted into ether. The ether extract was washed and dried by the well established procedures. On removal of ether, yellowish white crystals appeared and on GC and GC/MS analysis, the crystals appeared to be the 9-benzyl-9,10-dihydroanthracene.

This crude product was further purified by recrystallization from petroleum ether to yield a pure compound, m.p. 113-114°C, reported⁴ m.p. 114-115°C.

Analysis of the Products

The products obtained after Lochmann's base treatments were identified by gas chromatographic analysis on a Perkin-Elmer Sigma 3B instrument using a 1.83 m x 0.32 cm column packed with 10% OV-101. For some reactions, the

products were also analyzed by GC-MS using Hewlett Packard 5970 series mass detector (70 ev) connected to 5890 gas chromatograph OV-101 capillary column, 15 m x 0.25 mm, 100-180°C/min). Some of the products were also analyzed by high resolution GC-MS using VG 70-250 instrument.

Results and Discussion

As mentioned earlier, the solubility of the coal samples increase quite significantly due to the Lochmann's base treatment. Hence it was proposed that carbon-carbon bond cleavage occurs with the Lochmann's base in refluxing heptane. In order to investigate the type of hydrocarbon structures undergoing bond cleavage these new studies were carried out on five pure compounds and the results are given in Table I.

The first two compounds, 1,1'-binaphthyl and bibenzyl were ideal examples of compounds linked through two sp^2 carbon atoms and two sp^3 carbon atoms respectively. As has been shown in the Table, 1,1'-binaphthyl does not decompose with the base and the starting material was recovered quantitatively.

Bibenzyl has four benzylic hydrogen atoms which are significantly acidic and may be abstracted by the base.

However, inasmuch as we recovered the starting material, 2, on acidification, we infer that structures like bibenzyl do not undergo carbon-carbon bond cleavage reactions with strong bases. It is pertinent to note that bibenzyls and substituted bibenzyls are known to undergo bond cleavage with the alkali metals.

Next, another aromatic hydrocarbon, 9,10-diphenylanthracene, which has considerable steric bulk due to the two peri phenyl groups, was treated with Lochmann's base.

3

Only about 10% of the starting material could be recovered at the end of the reaction. Decomposition of 3 was somewhat surprising since 1,1'-binaphthyl (1) did not undergo decomposition. Of the products that could be identified in the GC trace were benzene (71%) and two products with very similar retention times. The product is not 9-phenylanthracene and we are continuing our work to identify the products.

Finally, since we are interested in base promoted decomposition reactions of hydrocarbons, we selected two hydrocarbons which have acidic hydrogen atoms and also would lead on decomposition to stable carbanions. The first compound was 1,1,3-triphenylpropane (4).

Abstraction of the acidic benzylic hydrogen atoms should be facile, and the carbanions should give styrene-like products together with another stable benzylic carbanion upon decomposition. In fact when compound 4 was treated with Lochmann's base only 20% of the starting material was recovered. The products that could be identified in the GC, were toluene (23%) and diphenylmethane (24%).

Thus the hydrocarbon 4 does undergo fragmentation via two different routes A and B giving two different set of products. Although kinetically route B would be preferred, thermodynamically route A is more favorable. From the

ratio of toluene to diphenylmethane, the two decomposition products from two different routes, we conclude that almost 50% decomposition is via A and the remaining 50% via B. More work is underway to determine the preferred reaction pathway.

Finally, the last compound in our list, 9-benzyl-9,10-dihydroanthracene, 5, treated with Lochmann's base and the reaction mixture was quenched with an acid at the end of the reaction. About 76% of the starting material was obtained unchanged at the end of the reaction. Toluene and anthracene were identified as products of the reaction. No other materials was observed in the GC of the product mixture. Evidently, the strong base may abstract a benzylic hydrogen from the dihydroanthracene ring which undergoes vinyl-ogous β -elimination reaction.

CONCLUSION

Fase-catalyzed elimination reactions have been demonstrated for three hydrocarbons.

References

1. Stock, L. M., Chatterjee, K. and Ettinger, M. Eleventh Quarterly Report, April/May/June, 1990.
2. For a summary of the relevant literature, see (a) Grovenstein E. Jr., Bhatti, A. M., Quest, D. E., Segupta, D. and VanDerveer, D. J. Am. Chem. Soc., 1983, 105, 6290 and referenced therein; (b) Marcinow, Z., Hull, C. E. and Rabideau, P. W. J. Org. Chem., 1989, 54, 3602.
3. Chatterjee, K., 1990, Ph.D. Dissertation, The University of Chicago.
4. Cuvigny, T. and Normant, H. Bull. Soc. Chim. France, Sect. D., 19 65, 1872-81.

Table I. Reaction of Pure Compounds with Lochmann's base at 98°C for 6 hours followed by protonation.

Entry	Substrate	Percent substrate recovered at the end of the reaction	Comments
1.	1,1'-binaphthyl	100	-
2.	Bibenzyl	100	-
3.	9,10-diphenylanthracene	10	Unexpected bond cleavage was observed.
4.	1,1,3-Triphenylpropane	20	Toluene and diphenylmethane were the observed products. Carbanion formation followed by β -cleavage does take place.
5.	9,10-dibenzyl-9,10-dihydroanthracene.	76	Formation of benzylic anion followed by β -cleavage takes place thereby forming anthracene and toluene.

END

**DATE
FILMED**

10 / 27 / 92