

Societatea de Științe Matematice din România



GAZETA
MATEMATICĂ
SERIA B

11

publicație lunară pentru tineret
FONDATĂ ÎN ANUL 1895

Anul CXVIII, nr. 11/2013
ISSN 1584-9333

SUMARUL

Articole și Note matematice

- | | |
|--|-----|
| 1. Square Roots of Real 2×2 , by Nicolae Anghel | 489 |
| 2. Numere relativ prime având produsul pătrat perfect, de Dorel Miheț | 492 |

Pentru cercurile de elevi

- | | |
|--|-----|
| 3. Câteva probleme de tip Ramsey, de Vasile Pop | 494 |
|--|-----|

Examene și Concursuri

- | | |
|--|-----|
| 4. Concursul Interjudețean de Matematică și Informatică „Marian Țarină“, Ediția a XIII-a, Turda, 10-11 mai 2013, prezentare de Dorel I. Duca și Gheorghe Lobonț | 501 |
| 5. Concursul „Dimitrie Pompeiu“, Ediția a XIII-a, Botoșani, 17- 19 mai 2013, prezentare de Artur Bălăucă și Ioan Ciobănașu | 507 |

Probleme

- | | |
|--|------------|
| 6. Rezolvarea problemelor din Gazeta Matematică nr. 5/2013..... | 514 |
| 7. Probleme pentru examene naționale | 525 |
| 8. Probleme pentru ciclul primar (P:628 – P:637) | 527 |
| 9. Probleme pregătitoare pentru concursuri și olimpiade <ul style="list-style-type: none">• Probleme pentru gimnaziu (E:14561 – E:14574)• Probleme pentru liceu (26831 – 26844) | 528
530 |

Rubrica rezolvitorilor de probleme



© Toate drepturile privind reproducerea, parțială sau totală, sub orice formă, a materialelor publicate în Gazeta Matematică, sunt rezervate Societății de Științe Matematice din România.



Preț 9 lei

Colectia completa a Gazetei Matematice seria B, in format electronic este realizata de Intuitext (www.intuitext.ro)
in parteneriat cu Societatea de Științe Matematice din România.

GAZETA MATEMATICĂ

SERIA B

PUBLICAȚIE LUNARĂ PENTRU TINERET

Fondată în anul 1895

Anul CXVIII nr. 11

noiembrie 2013

ARTICOLE ȘI NOTE MATEMATICE

SQUARE ROOTS OF REAL 2×2 MATRICES

NICOLAE ANGHEL¹⁾

Abstract. In this note we investigate the real 2×2 matrices which admit real square roots.

Keywords: Matrix, Square Root

MSC : Primary: 15A23. Secondary: 11C20.

Only non-negative real numbers admit real square roots. Thinking of a real number as the simplest square matrix, a 1×1 matrix, an interesting question emerges. Which real $n \times n$ matrices, $n \geq 1$, admit real square roots? In other words, for which $A \in \mathcal{M}_n(\mathbb{R})$ is there an $S \in \mathcal{M}_n(\mathbb{R})$ such that $S^2 = A$?

In this short note we will give a complete answer to the above question in the case $n = 2$. As a corollary, we will also establish how many distinct real square roots a given 2×2 matrix has, and then proceed to describe them exactly.

Theorem. *For a given matrix $A \in \mathcal{M}_2(\mathbb{R})$ there are matrices $S \in \mathcal{M}_2(\mathbb{R})$ such that $S^2 = A$ if and only if $\det A \geq 0$ and, either $A = -\sqrt{\det A}I$ or $\text{tr}A + 2\sqrt{\det A} > 0$, where I is the 2×2 identity matrix. Obviously, in the latter case, $\text{tr}A + 2\sqrt{\det A} = 0$.*

Proof. Necessity. Assume that S is a real square root of A . From $S^2 = A$ we conclude that $(\det S)^2 = \det A$, so we must have $\det A \geq 0$. Any matrix $M \in \mathcal{M}_2(\mathbb{R})$ satisfies the Cayley-Hamilton equation, namely $M^2 - (\text{tr}M)M + (\det M)I = 0$. In particular, $S^2 - (\text{tr}S)S + (\det S)I = 0$ implies $A - (\text{tr}S)S + (\det S)I = 0$. There are two cases to consider:

¹⁾Professor, University of North Texas, Denton, TX

(1) $\text{tr}S \neq 0$. Since for a matrix $M \in \mathcal{M}_2(\mathbb{R})$ we have $\text{tr}(M^2) = (\text{tr}M)^2 - 2\det M$, by taking $M = S$ we have $\text{tr}A + 2\det S = (\text{tr}S)^2 > 0$. However, $\text{tr}A + 2|\det S| \geq \text{tr}A + 2\det S$, and so we get $\text{tr}A + 2|\det S| > 0$. Now $(\det S)^2 = \det A$ is equivalent to $|\det S| = \sqrt{\det A}$. In conclusion, $\text{tr}A + 2\sqrt{\det A} > 0$.

(2) $\text{tr}S = 0$. In this case, $A - (\text{tr}S)S + (\det S)I = 0$ becomes $A = -(\det S)I$. If $\det S < 0$, $\text{tr}A + 2\sqrt{\det A} = -4\det S > 0$, and there is nothing to prove. If $\det S \geq 0$, we have $\det S = \sqrt{\det A}$, and so $A = -\sqrt{\det A}I$. Clearly, then $\text{tr}A + 2\sqrt{\det A} = 0$. Therefore, there are no matrices A with $\text{tr}A + 2\sqrt{\det A} < 0$, which admit real square roots.

Sufficiency. If $\det A \geq 0$ and $\text{tr}A + 2\sqrt{\det A} > 0$, a direct calculation taking into account that $A^2 = (\text{tr}A)A - (\det A)I$ shows that

$$S := \frac{1}{\sqrt{\text{tr}A + 2\sqrt{\det A}}}(A + \sqrt{\det A}I)$$

is a real square root of A . Suitable equations, found in the necessity part of the proof, show that this is the only possible real square root of A with positive trace and non-negative determinant.

If $\det A \geq 0$ and $A = -\sqrt{\det A}I$, then it is easily seen that

$$S := \begin{pmatrix} 0 & 1 \\ -\sqrt{\det A} & 0 \end{pmatrix}$$

is a square root of A . □

Corollary. As the existence of real square roots goes, the following is true about any matrix $A \in \mathcal{M}_2(\mathbb{R})$:

(1) If $A \neq aI$, $a \in \mathbb{R}$, then A admits only finitely many real square roots, as follows:

(a) If $\det A > 0$ and $\text{tr}A - 2\sqrt{\det A} > 0$, there are exactly four distinct square roots, given by

$$S = \pm \frac{1}{\sqrt{\text{tr}A + 2\sqrt{\det A}}}(A + \sqrt{\det A}I)$$

or

$$S = \pm \frac{1}{\sqrt{\text{tr}A - 2\sqrt{\det A}}}(A - \sqrt{\det A}I).$$

(b) If $\det A < 0$, or $\det A \geq 0$ and $\text{tr}A + 2\sqrt{\det A} \leq 0$, there are no real square roots.

(c) Otherwise, there are exactly two distinct real square roots, given by $S = \pm \frac{1}{\sqrt{\text{tr}A + 2\sqrt{\det A}}}(A + \sqrt{\det A}I)$.

(2) If $A = aI$, $a \in \mathbb{R}$, then A admits infinitely many real square roots.

Regardless of a , the doubly infinite family $S = \begin{pmatrix} s & t \\ a-s^2 & -s \\ t & \end{pmatrix}$, $s, t \in \mathbb{R}$, $t \neq 0$, is in. If $a = 0$ we add to the above family the matrices $S = \begin{pmatrix} 0 & 0 \\ s & 0 \end{pmatrix}$, $s \in \mathbb{R}$, while if $a > 0$ we add the family $S = \begin{pmatrix} \pm\sqrt{a} & 0 \\ s & \mp\sqrt{a} \end{pmatrix}$, $s \in \mathbb{R}$, plus the two more matrices given by $S = \begin{pmatrix} \pm\sqrt{a} & 0 \\ 0 & \pm\sqrt{a} \end{pmatrix}$ (the signs correspond).

Proof. If we are in case (1)(a), a direct calculation, similar to that in the sufficiency part of the Theorem, shows that indeed the four proposed matrices are real square roots of A . Conversely, let S be a real square root of A . As in the proof of the Theorem, we then have $(\det S)^2 = \det A$, $(\text{tr}S)^2 = \text{tr}A + 2 \det S$, and $A - (\text{tr}S)S + (\det S)I = 0$.

If $\det S = \sqrt{\det A}$, then $\text{tr}S = \pm\sqrt{\text{tr}A + 2\sqrt{\det A}}$, and so

$$S = \pm \frac{1}{\sqrt{\text{tr}A + 2\sqrt{\det A}}} (A + \sqrt{\det A}I).$$

Similarly, if $\det S = -\sqrt{\det A}$ we get the other two matrices in the family of square roots.

The four matrices are distinct because, for instance, their traces are distinct:

$$\sqrt{\text{tr}A + 2\sqrt{\det A}} > \sqrt{\text{tr}A - 2\sqrt{\det A}} > -\sqrt{\text{tr}A - 2\sqrt{\det A}} > -\sqrt{\text{tr}A + 2\sqrt{\det A}}.$$

The case (1)(b) follows immediately from the Theorem, by negation, since $A \neq aI$, $a \in \mathbb{R}$.

In case (1)(c) „otherwise“ means after some „detective work“, $A \neq aI$, $a \in \mathbb{R}$ and in addition, $\det A > 0$ and $\text{tr}A + 2\sqrt{\det A} > 0$ and $\text{tr}A - 2\sqrt{\det A} \leq 0$, or $\det A = 0$ and $\text{tr}A > 0$. The claimed conclusion can then be reached as in (1)(b).

Finally, the case (2) is an easy ‘by hand’ calculation, given the simple structure of A .