Progress Report: DOE Environmental Management Science Grant

Date: October 15, 1997

Principal Investigator: David W. Ow

Proposal Title: "Molecular Genetics of Metal Detoxification: Prospects for Phytoremediation"

I. We proposed to characterize a number of fission yeast mutants that are hypersensitivity to cadmium and deficient in the production of metal-peptide complexes. For each of the mutants we sought to clone the gene responsible for the mutant phenotype and more importantly to define the gene function. Below, we summary the progress made thus far for each of the mutants.

Mutants that hypoproduce phytochelatins:

DS12: The gene has been cloned, but a full length cDNA remains to be isolated. We believe our longest clone is short at the 5' end by less than 100 bp. This gene encodes sulfite reductase and its function is needed for Pb-induced sulfide production, but not Cd-induced sulfide production. Since this enzyme acts upstream of cysteine biosynthesis, the likely reason that this mutant hypoproduces phytochelatins is that it fails to produce sufficient cysteine during Cd stress.

JS563 : The gene has been cloned and found to encode a sulfide dehydrogenase. In vitro, we found that the protein binds FAD, converts S^{2-} to S^0 while reducing quinone. The protein is membrane associated and has been localized to the mitochondria. Its likely function is to detoxify sulfide in the mitochondria resulting from cadmium-induced sulfide production. The sulfide electrons are likely used for the electron transport chain. Cells that have a defect in this enzyme cannot oxidize mitochondrial S^{2-} resulting in high toxic levels of S^{2-} during Cd stress. In addition, the high S^{2-} level precipitates Cd^{2+} to form CdS, and the lack of free Cd^{2+} fails to induce phytochelatin synthase activity to produce phytochelatin peptides.

JS282: The genomic clone that restores Cd tolerance to JS282 has been isolated. Surprisingly, this genomic clone when present in a multicopy vector in a wild type background causes hypersensitivity to Cd and selenium. The cDNA corresponding to the genomic clone has been isolated and its sequence is being determined.

JS246: A genomic clone that restores Cd tolerance has been isolated but its corresponding cDNA remains to be cloned. Wild type strains harboring the genomic clone become hypertolerant to cadmium. Extensive efforts are being made to find the cDNA clone, including the construction of new cDNA libraries.

Mutant that fails to accumulate a high molecular weight phytochelatin-cadmium-sulfide complex.

JS237: This mutant has a mRNA splicing defect in a gene homologous to the human Wiskott-Aldrich Syndrome protein (WASP). The human disease symptom is associated

with defects in cytoskeletal organization. Likewise, mutant JS237 exhibits aberrant arrangement of polymerized actin and the fission yeast WASP has been found to co-localize with cortical actin and to interact with actin associated proteins. The genomic clone complemented the defect associated with the polymerized actin organization, but did not fully restore Cd tolerance. Hence, we entertained the possibility of another gene encoded by the genomic clone. Indeed, we found that the mutation appears to affect more than the WASP mRNA. The data thus far suggest an upstream mRNA overlaps the WASP coding region. When the upstream DNA was sequenced, we found an open reading frame encoding a P-type ATPase which may be a cation transporter. Where this transporter resides within the cell and what role it plays in Cd tolerance are currently under investigation.

II. In addition to characterizing mutants that show a defect in metal tolerance, we also took the approach of isolating DNA clones that can confer hypertolerance to cadmium. A cDNA expression library was made in a fission yeast vector such that selection for clones that confer high level tolerance can be made. Some 40 clones were isolated and sequenced. These cDNAs reveal a variety of genes, including one that encodes a class I (animal-type) metallothionein protein. Hence, fission yeast also has a metallothionein which is not used in response to metal stress, but when overproduced through artificial means, can nonetheless function like animal metallothioneins. Other genes encode enzymes in glutathione biosynthesis and proteins that are involved in calcium binding, heat stress, protein synthesis or the generation of NAD(P)H. Some proteins show no sequence similarity to known proteins.

III. A cDNA library made from the mRNA of *Brassica juncea*, a metal-hyperaccumulating plant, was also expressed from the fission yeast expression vector. This library was transformed into fission yeast and cadmium hypertolerant clones were isolated. This resulted in the cloning of the *Brassica juncea* class II metallothionein genes. These genes are similar to those found in other plants, all of which are dissimilar to the animal class I genes, which as described above is also found in fission yeast. The cDNA library has predominantly short cDNAs, which may explain why the metallothionein genes are readily isolated. Current efforts are underway to construct a size-selected cDNA library that carries predominantly long inserts.