Radiological Conditions at the Southern Islands
of Rongelap Atoll

William L. Robison
Cynthia L. Conrado

RECEIVED
JUL 22 1996
OSTI

March 1996

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This report has been reproduced
directly from the best available copy.

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831
Prices available from (615) 576-8401, FTS 626-8401

Available to the public from the
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Rd.,
Springfield, VA 22161

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Radiological Conditions at the Southern Islands of
Rongelap Atoll

William L. Robison
Cynthia L. Conrado

Lawrence Livermore National Laboratory
Environmental Sciences Division

Report to DOE Environment, Safety, and Health
for Dr. Harry Pettengill, March 1992

March, 1996

TABLE OF CONTENTS

	Introductory Comments	01
Table 1	The number of food crops collected at the southern islands at Rongelap Atoll and the mean and median ^{137}Cs concentrations.....	02
Table 2	The number of indicator species collected at the southern islands at Rongelap Atoll and the mean and median ^{137}Cs concentrations.....	04
Table 3	The number of animal samples collected at the southern islands at Rongelap Atoll and the mean and median ^{137}Cs concentrations.....	06
Table 4	The number of soil samples collected at the southern islands at Rongelap Atoll.....	07
Table 5	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Erapuotsu Island (F32).....	08
Table 6	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Eniaetok Island (F33).....	08
Table 7	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Busch Island (F38).	09
Table 8	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Rongelap Island (F42).....	09
Table 9	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Arbar Island (F43).	10
Table 10	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Eniran Island (F46).	10

Table 11	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Tufa Island (F47).	11
Table 12	Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Borukka Island (F49).	11
Figure 1	Rongelap Island— ^{137}Cs distribution in soil.....	12
Figure 2	^{137}Cs in soil at southern islands of Rongelap Atoll.....	13
Appendix A	Radionuclide concentration summary in vegetation samples collected during the 1978 NMIRS and from 1986 through 1991 in the southern islands of Rongelap Atoll.	A-14
Table A-1	The ^{137}Cs radionuclide concentration summary for vegetation (food crops) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.....	A-15
Table A-2	The ^{137}Cs radionuclide concentration summary for vegetation (indicator species) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.	A-17
Appendix B	Radionuclide concentration summary in animal samples collected during the 1978 NMIRS and from 1986 through 1991 in the southern islands of Rongelap Atoll.	B-18
Table B-1	The ^{137}Cs radionuclide concentration summary for animals (food source) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.....	B-19
Appendix C	Radionuclide concentration summary of all soil profile samples collected during the 1978 NMIRS and from 1986 through 1991 in the southern islands of Rongelap Atoll....	C-20
Table C-1	Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Erapuotsu Island (F32).....	C-21

Table C-2	Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Eniaetok Island (F33).....	C-22
Table C-3	Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Busch Island (F38).....	C-23
Table C-4	Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trips in 1986 through 1989 on Rongelap Island (F42).	C-24
Table C-5	Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1991 on Arbar Island (F43).	C-25
Table C-6	Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Eniran Island (F46).....	C-26
Table C-7	Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Tufa Island (F47).	C-27
Table C-8	Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Borukka Island (F49).....	C-28
Table C-9	Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Eniaetok Island (F33).	C-29
Table C-10	Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Rongelap Island (F42).	C-30
Table C-11	Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Borukka Island (F49).	C-31
Table C-12	Plutonium 239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1989 on Eniaetok Island (F33).....	C-32

Table C-13	Plutonium 239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1986 on Rongelap Island (F42).	C-33
Table C-14	Plutonium 239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1989 on Borukka Island (F49).	C-34
Table C-15	Americium-241 radionuclide concentration summary for all soil profiles taken from our recent trip in 1991 on Erapuotsu Island (F32).	C-35
Table C-16	Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Eniaetok Island (F33).	C-36
Table C-17	Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Busch Island (F38).	C-37
Table C-18	Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trips in 1986 through 1989 on Rongelap Island (F42).	C-38
Table C-19	Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1991 on Arbar Island (F43).	C-39
Table C-20	Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Eniran Island (F46).	C-40
Table C-21	Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Tufa Island (F47).	C-41
Table C-22	Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Borukka Island (F49).	C-42
Appendix D	Maps of Islands.	D-43

INTRODUCTORY COMMENTS

The data presented in the following tables is the total available for each southern island; they include both the data from the 1978 Northern Marshall Island Radiological Survey (NMIRS) and trips to Rongelap Atoll from 1986 through 1991. There are additional samples that were taken at Rongelap Island in 1990 and 1991, and the data are unavailable for this report.

In one table we present the number of vegetation samples collected in the 1978 NMIRS and from 1986 through 1991. Again, the majority of the ^{137}Cs is from the 1986-1991 trips. We have not made additional analyses of $^{239+240}\text{Pu}$, ^{241}Am and ^{90}Sr because the concentrations are very low and these radionuclides contribute less than 5% of an already very small dose.

In another table we show the number of soil samples collected at each island in 1978 and the number collected since 1986. Most of the data are from 1986 through 1991. The major exception is ^{90}Sr where all of the data are from the 1978 NMIRS. We have done some additional Pu analyses of soils from Rongelap Eniaetok, and Borukka Island but none of the other southern islands. A significant amount of new data for ^{137}Cs and ^{241}Am have been generated from the samples collected from 1986 through 1991.

The data are presented in the form of summary tables, graphics, detailed appendices and aerial photographs of the islands with the sample locations marked. The identified sample locations from the 1978 NMIRS will be added later.

Table 1. The number of food crops collected at the southern islands at Rongelap Atoll and the mean and median ¹³⁷Cs concentrations.

Island Name	Island Code	Number of Samples		Cs-137 pCi/g wet wt.		
		Total	1978 NMIRS	1986-1991	median	mean
Drinking Coconut Meat						
Erapuotsu	32f	3	0	3	2.0	2.2
Eniaetok	33f	7	2	5	1.7	2.7
Busch	38f	5	0	5	1.1	1.1
Rongelap	42f	200	3	197	1.4	1.9
Arbar	43f	4	1	3	0.88	0.93
Eniran	46f	4	0	4	0.98	1.1
Tufa	47f	6	0	6	0.74	0.78
Borukka	49f	5	0	5	1.2	1.3
Drinking Coconut Juice						
Erapuotsu	32f	3	0	3	0.79	0.85
Eniaetok	33f	7	2	5	0.32	0.56
Busch	38f	5	0	5	0.28	0.32
Rongelap	42f	194	3	191	0.62	0.85
Arbar	43f	4	1	3	0.17	0.17
Eniran	46f	4	0	4	0.47	0.48
Tufa	47f	7	0	7	0.21	0.22
Borukka	49f	5	0	5	0.36	0.46
Copra Meat						
Eniaetok	33f	14	2	12	3.3	3.4
Rongelap	42f	72	14	58	2.6	3.4
Arbar	43f	1	0	1	0.82	0.82
Tufa	47f	1	0	1	1.1	1.1
Borukka	49f	6	1	5	2.0	2.9
Copra Juice						
Eniaetok	33f	14	2	12	2.4	2.5
Rongelap	42f	70	13	57	1.4	1.9
Arbar	43f	1	0	1	1.1	1.1
Tufa	47f	1	0	1	0.80	0.80
Borukka	49f	6	1	5	1.2	2.1

Table 1. The number of food crops collected at the southern islands at Rongelap Atoll and the mean and median ¹³⁷Cs concentrations (continued).

Island Name	Island Code	Number of Samples			Cs-137 pCi/g wet wt.	
		Total	1978 NMIRS	1986-1991	median	mean
<i>Pandanus</i>						
Erapuotsu	32f	3	0	3	2.0	2.4
Eniaetok	33f	15	2	13	2.2	4.0
Busch	38f	2	0	2	0.83	0.83
Rongelap	42f	78	16	62	5.9	7.6
Arbar	43f	4	2	2	1.4	3.1
Tufa	47f	2	0	2	4.0	4.0
Borukka	49f	9	4	5	1.0	1.4
<i>Breadfruit Meat</i>						
Rongelap	42f	30	1	29	3.0	3.2
<i>Tacca Meat</i>						
Busch	38f	1	0	1	2.3	2.3
Rongelap	42f	5	0	5	4.3	5.7
Tufa	47f	2	0	2	1.6	1.6
Borukka	49f	4	1	3	1.2	1.2
<i>Lime Meat</i>						
Rongelap	42f	9	0	9	1.5	1.6
<i>Squash Meat</i>						
Rongelap	42f	2	0	2	5.8	5.8

NOTE: Specific activity decay corrected to 1994.

Table 2. The number of indicator species collected at the southern islands at Rongelap Atoll and the mean and median ¹³⁷Cs concentrations.

Island Name	Island Code	Number of Samples			Cs-137 pCi/g wet wt.	
		Total	1978 NMIRS	1986-1991	median	mean
Morinda Fruit						
Eniaetok	33f	1	1	0	2.5	2.5
Rongelap	42f	2	0	2	1.6	1.6
Arbar	43f	1	0	1	0.47	0.47
Eniran	46f	3	0	3	0.56	0.57
Tufa	47f	1	0	1	0.74	0.74
Morinda Leaves						
Arbar	43f	1	0	1	0.38	0.38
Eniran	46f	3	0	3	0.22	0.31
Tufa	47f	1	0	1	0.66	0.66
Guettardia Leaves						
Rongelap	42f	1	0	1	2.9	2.9
Arbar	43f	3	0	3	0.38	0.67
Tufa	47f	1	0	1	0.25	0.25
Messerschmidia Leaves						
Rongelap	42f	2	0	2	0.49	0.49
Pisonia Leaves						
Tufa	47f	1	0	1	3.0	3.0
Pandanus Leaves						
Rongelap	42f	5	3	2	7.3	5.6
Scaevola Leaves						
Arbar	43f	4	0	4	0.27	0.32
Eniran	46	1	0	1	0.19	0.19
Tufa	47f	2	0	2	0.31	0.31
Borukka	49f	1	0	1	3.7	3.7

Table 2. The number of indicator species collected at the southern islands at Rongelap Atoll and the mean and median ^{137}Cs concentrations (continued).

Island Name	Island Code	Number of Samples			Cs-137 pCi/g wet wt.	
		Total	1978 NMIRS	1986-1991	median	mean
At-At Vine						
Tufa	47f	1	0	1	0.34	0.34
Tabero Fruit						
Rongelap	42f	3	0	3	2.0	1.8
Tabero Leaves						
Rongelap	42f	1	0	1	1.2	1.2
Coconut Fronds						
Rongelap	42f	16	0	16	1.1	2.4

NOTE: Specific activity decay corrected to 1994.

Table 3. The number of animal samples collected at the southern islands at Rongelap Atoll and the mean and median ¹³⁷Cs concentrations.

Island Name	Island Code	Number of Samples			Cs-137 pCi/g wet wt.	
		Total	1978 NMIRS	1986-1991	median	mean
Coconut Crab Muscle						
Erapuotsu	32f	1	0	1	4.9	4.9
Busch	38f	1	0	1	1.4	1.4
Arbar	43f	2	2	0	2.5	2.5
Eniran	46f	2	0	2	1.5	1.5
Tufa	47f	2	0	2	1.6	1.6
Borukka	49f	1	0	1	2.1	2.1
Coconut Crab Hepatopancreas						
Erapuotsu	32f	1	0	1	3.4	3.4
Busch	38f	1	0	1	0.98	0.98
Eniran	46f	2	0	2	1.1	1.1
Tufa	47f	2	0	2	1.1	1.1
Borukka	49f	1	0	1	1.9	1.9
Pig Heart						
Rongelap	42f	4	0	4	13	14
Pig Liver						
Rongelap	42f	6	2	4	5.3	5.6
Pig Muscle						
Rongelap	42f	7	2	5	13	14

NOTE: Specific activity decay corrected to 1994.

Table 4. The number of soil samples collected at the southern islands at Rongelap Atoll.

Number of Soil Samples for Cs-137 and Am-241					
Island Name	Island Code	1978		1986-1991	
		Profiles	0-5 cm.	Profiles	0-5 cm.
Erapuotsu	32f	0	0	3	3
Eniaetok	33f	6	6	15	15
Busch	38f	0	0	5	5
Rongelap	42f	28	27	90	267
Arbar	43f	7	6	7	7
Eniran	46f	0	0	7	6
Tufa	47f	0	0	8	8
Borukka	49f	3	3	10	10

Number of Soil Samples for Pu-239+240					
Island Name cm.	Island Code	1978		1986-1991	
		Profiles	0-5 cm.	Profiles	0-5
Erapuotsu	32f	0	0	0	0
Eniaetok	33f	6	6	0	15
Busch	38f	0	0	0	0
Rongelap	42f	20	18	0	59
Arbar	43f	0	0	0	0
Eniran	46f	0	0	0	0
Tufa	47f	0	0	0	0
Borukka	49f	3	3	0	10

Number of Soil Samples for Sr-90					
Island Name cm.	Island Code	1978		1986-1991	
		Profiles	0-5 cm.	Profiles	0-5
Erapuotsu	32f	0	0	0	0
Eniaetok	33f	6	6	0	0
Busch	38f	0	0	0	0
Rongelap	42f	21	20	0	0
Arbar	43f	0	0	0	0
Eniran	46f	0	0	0	0
Tufa	47f	0	0	0	0
Borukka	49f	3	3	0	0

Table 5. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Erapuotsu Island (F32).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	22	23	NDA	NDA	NDA	NDA	5.2	7.7
05-10	31	35	NDA	NDA	NDA	NDA	4.5	4.5
10-15	5	8.1	NDA	NDA	NDA	NDA	0.0	0.0
15-25	1.6	1.4	NDA	NDA	NDA	NDA	0.0	0.0
25-40	0.86	0.96	NDA	NDA	NDA	NDA	0.0	0.0
40-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0
00-05	22	23	NDA	NDA	NDA	NDA	5.2	7.7
00-10	26	29	NDA	NDA	NDA	NDA	4.4	4.4
00-15	18	22	NDA	NDA	NDA	NDA	0.0	0.0
00-25	11	14	NDA	NDA	NDA	NDA	0.0	0.0
00-40	7.4	9.0	NDA	NDA	NDA	NDA	0.0	0.0
00-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

NDA stands for no data available.

Table 6. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Eniaetok Island (F33).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	21	21	5.4	6.6	6.6	7.2	5.5	5.1
05-10	8.0	9.1	8.1	9.2	3.0	3.3	0.88	1.7
10-15	3.3	4.4	3.4	6.1	2.0	2.6	0.61	0.70
15-25	1.3	1.5	2.1	3.3	0.31	1.8	0.11	0.16
25-40	0.49	0.75	3.1	5.9	0.059	0.056	0.027	0.041
40-60	0.50	1.6	2.7	2.4	0.089	0.17	0.067	0.11
00-05	21	21	5.4	6.6	6.6	7.2	5.5	5.1
00-10	13	15	6.8	7.9	2.6	2.6	2.8	3.1
00-15	10	11	5.9	7.3	2.1	2.6	1.6	2.1
00-25	7.2	7.2	4.4	5.7	1.4	2.3	0.76	0.78
00-40	4.8	4.8	4.0	5.8	0.87	1.4	0.37	0.48
00-60	2.7	2.5	4.0	4.7	0.71	1.0	0.28	0.35

NOTE: Specific activity decay corrected to 1994.

Table 7. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Busch Island (F38).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	12	27	NDA	NDA	NDA	NDA	5.4	6.9
05-10	22	23	NDA	NDA	NDA	NDA	6.0	6.0
10-15	16	15	NDA	NDA	NDA	NDA	2.6	3.1
15-25	6.0	4.9	NDA	NDA	NDA	NDA	0.51	0.69
25-40	1.4	2.0	NDA	NDA	NDA	NDA	0.43	0.43
40-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0
00-05	12	27	NDA	NDA	NDA	NDA	5.4	6.9
00-10	23	25	NDA	NDA	NDA	NDA	3.5	3.5
00-15	22	22	NDA	NDA	NDA	NDA	3.5	3.5
00-25	15	15	NDA	NDA	NDA	NDA	2.4	2.4
00-40	9.9	10	NDA	NDA	NDA	NDA	1.3	1.3
00-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

NDA stands for no data available.

6

Table 8. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Rongelap Island (F42).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	14	14	5.2	4.8	2.8	3.4	1.6	2.0
05-10	6.5	7.4	3.3	5.5	1.0	2.1	0.68	1.3
10-15	2.9	4.0	3.1	4.1	0.50	0.92	0.48	0.61
15-25	1.2	1.8	2.2	3.1	0.20	0.27	0.18	0.31
25-40	0.36	0.60	1.4	1.8	0.090	0.13	0.046	0.079
40-60	0.19	0.36	0.0	0.0	0.0	0.0	0.020	0.071
00-05	14	14	5.2	4.8	2.8	3.4	1.6	2.0
00-10	9.4	11	4.1	5.2	1.8	2.6	1.7	2.2
00-15	7.5	8.8	3.7	4.8	1.4	2.0	1.4	1.6
00-25	5.1	6.0	3.6	4.2	1.2	1.4	0.87	1.0
00-40	3.5	3.9	2.9	3.3	0.82	0.90	0.32	0.38
00-60	2.4	2.7	0.0	0.0	0.0	0.0	0.18	0.25

NOTE: Specific activity decay corrected to 1994.

Table 9. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Arbar Island (F43).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	10	14	NDA	NDA	NDA	NDA	3.5	7.0
05-10	7.3	11	NDA	NDA	NDA	NDA	0.95	2.2
10-15	2.4	6.6	NDA	NDA	NDA	NDA	2.7	2.7
15-25	1.5	3.3	NDA	NDA	NDA	NDA	2.9	2.9
25-40	1.0	0.80	NDA	NDA	NDA	NDA	0.0	0.0
40-60	0.31	0.47	NDA	NDA	NDA	NDA	0.0	0.0
00-05	10	14	NDA	NDA	NDA	NDA	3.5	7.0
00-10	10	13	NDA	NDA	NDA	NDA	2.7	4.1
00-15	7.3	11	NDA	NDA	NDA	NDA	2.7	2.7
00-25	5.3	8.0	NDA	NDA	NDA	NDA	3.5	3.5
00-40	3.7	5.5	NDA	NDA	NDA	NDA	0.0	0.0
00-60	2.3	2.4	NDA	NDA	NDA	NDA	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

NDA stands for no data available.

Table 10. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Eniran Island (F46).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	13	13	NDA	NDA	NDA	NDA	5.0	4.8
05-10	8.3	9.8	NDA	NDA	NDA	NDA	1.8	1.8
10-15	3.3	3.9	NDA	NDA	NDA	NDA	0.60	0.60
15-25	2.0	2.4	NDA	NDA	NDA	NDA	0.0	0.0
25-40	0.34	0.91	NDA	NDA	NDA	NDA	0.0	0.0
40-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0
00-05	13	13	NDA	NDA	NDA	NDA	5.0	4.8
00-10	12	12	NDA	NDA	NDA	NDA	2.3	2.3
00-15	9.2	9.7	NDA	NDA	NDA	NDA	1.7	1.7
00-25	6.6	6.9	NDA	NDA	NDA	NDA	0.0	0.0
00-40	4.7	4.7	NDA	NDA	NDA	NDA	0.0	0.0
00-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

NDA stands for no data available.

Table 11. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Tufa Island (F47).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	18	24	NDA	NDA	NDA	NDA	3.6	4.5
05-10	21	22	NDA	NDA	NDA	NDA	2.1	5.2
10-15	8.4	11	NDA	NDA	NDA	NDA	1.4	1.3
15-25	3.0	4.4	NDA	NDA	NDA	NDA	0.20	0.38
25-40	1.1	1.9	NDA	NDA	NDA	NDA	0.0	0.0
40-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0
00-05	18	24	NDA	NDA	NDA	NDA	3.6	4.5
00-10	17	23	NDA	NDA	NDA	NDA	3.5	4.7
00-15	15	19	NDA	NDA	NDA	NDA	3.6	3.9
00-25	9.8	13	NDA	NDA	NDA	NDA	3.3	2.9
00-40	6.9	8.8	NDA	NDA	NDA	NDA	0.0	0.0
00-60	0.0	0.0	NDA	NDA	NDA	NDA	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

NDA stands for no data available.

Table 12. Average ^{137}Cs , ^{90}Sr , $^{239+240}\text{Pu}$, and ^{241}Am soil concentrations in pCi/g dry weight for soil profiles taken on Borukka Island (F49).

Soil depth (cm)	^{137}Cs		^{90}Sr		$^{239+240}\text{Pu}$		^{241}Am	
	Median	Mean	Median	Mean	Median	Mean	Median	Mean
00-05	22	25	12	12	5.3	6.9	3.2	4.5
05-10	13	15	9.3	7.6	0.66	0.93	0.88	2.6
10-15	7.2	9.1	6.2	5.6	0.25	0.29	0.61	0.76
15-25	2.6	4.3	2.4	3.0	0.18	0.16	0.20	0.26
25-40	0.84	1.4	2.5	2.5	0.069	0.069	0.035	0.092
40-60	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
00-05	22	25	12	12	5.3	6.9	3.2	4.5
00-10	17	20	10	9.6	4.3	4.1	2.2	3.3
00-15	14	17	9.0	8.3	2.9	2.8	1.7	2.0
00-25	9.5	12	6.0	6.2	1.8	1.7	0.82	1.3
00-40	6.1	7.8	5.2	5.2	1.3	1.3	0.48	0.46
00-60	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

NOTE: Specific activity decay corrected to 1994.

Figure 1. Rongelap Island— ^{137}Cs Distribution in Soil.

Figure 2. ^{137}Cs in soil at southern islands of Rongelap Atoll.

Appendix A

Radionuclide concentration summary in vegetation samples collected during the 1978 NMIRS and from 1986 through 1991 in the southern islands of Rongelap Atoll.

Table A-1. The ^{137}Cs radionuclide concentration summary for vegetation (food crops) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.

Island	Island		pCi g ⁻¹ wet wt					Mean	SD
name	code	N ^a	Minimum	Maximum	Median	Mean	SD	of logs	of logs
Drinking Coconut Meat									
Erapuotsu	32f	3	1.5 × 10 ⁰	3.1 × 10 ⁰	2.0 × 10 ⁰	2.2 × 10 ⁰	8.2 × 10 ⁻¹	7.4 × 10 ⁻¹	3.7 × 10 ⁻¹
Eniaetok	33f	7	1.7 × 10 ⁻¹	7.9 × 10 ⁰	1.7 × 10 ⁰	2.7 × 10 ⁰	2.9 × 10 ⁰	3.1 × 10 ⁻¹	1.4 × 10 ⁰
Busch	38f	5	5.6 × 10 ⁻¹	1.6 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	3.8 × 10 ⁻¹	3.6 × 10 ⁻²	3.9 × 10 ⁻¹
Rongelap	42f	200	2.9 × 10 ⁻¹	9.8 × 10 ⁰	1.4 × 10 ⁰	1.9 × 10 ⁰	1.4 × 10 ⁰	4.0 × 10 ⁻¹	6.9 × 10 ⁻¹
Arbar	43f	4	4.9 × 10 ⁻¹	1.5 × 10 ⁰	8.8 × 10 ⁻¹	9.3 × 10 ⁻¹	4.9 × 10 ⁻¹	-1.9 × 10 ⁻¹	5.7 × 10 ⁻¹
Eniran	46f	4	4.7 × 10 ⁻¹	1.9 × 10 ⁰	9.8 × 10 ⁻¹	1.1 × 10 ⁰	6.0 × 10 ⁻¹	-4.0 × 10 ⁻²	5.8 × 10 ⁻¹
Tufa	47f	6	3.7 × 10 ⁻¹	1.6 × 10 ⁰	7.4 × 10 ⁻¹	7.8 × 10 ⁻¹	4.2 × 10 ⁻¹	-3.6 × 10 ⁻¹	5.0 × 10 ⁻¹
Borukka	49f	5	6.3 × 10 ⁻¹	2.5 × 10 ⁰	1.2 × 10 ⁰	1.3 × 10 ⁰	7.4 × 10 ⁻¹	1.6 × 10 ⁻¹	5.5 × 10 ⁻¹
Drinking Coconut Juice									
Erapuotsu	32f	3	4.0 × 10 ⁻¹	1.4 × 10 ⁰	7.9 × 10 ⁻¹	8.5 × 10 ⁻¹	4.8 × 10 ⁻¹	-2.8 × 10 ⁻¹	6.2 × 10 ⁻¹
Eniaetok	33f	7	1.8 × 10 ⁻²	1.8 × 10 ⁰	3.2 × 10 ⁻¹	5.6 × 10 ⁻¹	6.1 × 10 ⁻¹	-1.2 × 10 ⁰	1.5 × 10 ⁰
Busch	38f	5	1.6 × 10 ⁻¹	4.5 × 10 ⁻¹	2.8 × 10 ⁻¹	3.2 × 10 ⁻¹	1.2 × 10 ⁻¹	-1.2 × 10 ⁰	4.1 × 10 ⁻¹
Rongelap	42f	194	9.2 × 10 ⁻²	3.7 × 10 ⁰	6.2 × 10 ⁻¹	8.5 × 10 ⁻¹	6.7 × 10 ⁻¹	-4.6 × 10 ⁻¹	8.0 × 10 ⁻¹
Arbar	43f	4	5.0 × 10 ⁻²	3.0 × 10 ⁻¹	1.7 × 10 ⁻¹	1.7 × 10 ⁻¹	1.1 × 10 ⁻¹	-2.0 × 10 ⁰	7.7 × 10 ⁻¹
Eniran	46f	4	1.9 × 10 ⁻¹	7.7 × 10 ⁻¹	4.7 × 10 ⁻¹	4.8 × 10 ⁻¹	2.5 × 10 ⁻¹	-8.6 × 10 ⁻¹	6.0 × 10 ⁻¹
Tufa	47f	7	7.9 × 10 ⁻²	3.5 × 10 ⁻¹	2.1 × 10 ⁻¹	2.2 × 10 ⁻¹	1.0 × 10 ⁻¹	-1.6 × 10 ⁰	5.6 × 10 ⁻¹
Borukka	49f	5	8.8 × 10 ⁻²	1.2 × 10 ⁰	3.6 × 10 ⁻¹	4.6 × 10 ⁻¹	4.2 × 10 ⁻¹	-1.1 × 10 ⁰	9.4 × 10 ⁻¹
Copra Meat									
Eniaetok	33f	14	6.1 × 10 ⁻¹	7.2 × 10 ⁰	3.3 × 10 ⁰	3.4 × 10 ⁰	1.9 × 10 ⁰	1.0 × 10 ⁰	7.0 × 10 ⁻¹
Rongelap	42f	72	8.4 × 10 ⁻¹	1.8 × 10 ¹	2.6 × 10 ⁰	3.4 × 10 ⁰	2.7 × 10 ⁰	1.0 × 10 ⁰	6.6 × 10 ⁻¹
Arbar	43f	1	8.2 × 10 ⁻¹	8.2 × 10 ⁻¹	8.2 × 10 ⁻¹	8.2 × 10 ⁻¹	0.0 × 10 ⁰	-1.9 × 10 ⁻¹	0.0 × 10 ⁰
Tufa	47f	1	1.1 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	0.0 × 10 ⁰	7.8 × 10 ⁻²	0.0 × 10 ⁰
Borukka	49f	6	9.5 × 10 ⁻¹	7.1 × 10 ⁰	2.0 × 10 ⁰	2.9 × 10 ⁰	2.4 × 10 ⁰	7.9 × 10 ⁻¹	7.9 × 10 ⁻¹
Copra Juice									
Eniaetok	33f	14	5.2 × 10 ⁻¹	5.6 × 10 ⁰	2.4 × 10 ⁰	2.5 × 10 ⁰	1.6 × 10 ⁰	6.5 × 10 ⁻¹	8.0 × 10 ⁻¹
Rongelap	42f	70	2.5 × 10 ⁻¹	6.4 × 10 ⁰	1.4 × 10 ⁰	1.9 × 10 ⁰	1.5 × 10 ⁰	3.4 × 10 ⁻¹	7.8 × 10 ⁻¹
Arbar	43f	1	1.1 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	0.0 × 10 ⁰	6.6 × 10 ⁻²	0.0 × 10 ⁰
Tufa	47f	1	8.0 × 10 ⁻¹	8.0 × 10 ⁻¹	8.0 × 10 ⁻¹	8.0 × 10 ⁻¹	0.0 × 10 ⁰	-2.2 × 10 ⁻¹	0.0 × 10 ⁰
Borukka	49f	6	3.6 × 10 ⁻¹	5.1 × 10 ⁰	1.2 × 10 ⁰	2.1 × 10 ⁰	2.0 × 10 ⁰	3.3 × 10 ⁻¹	1.0 × 10 ⁰
Pandanus									
Erapuotsu	32f	3	1.4 × 10 ⁰	4.0 × 10 ⁰	2.0 × 10 ⁰	2.4 × 10 ⁰	1.4 × 10 ⁰	7.9 × 10 ⁻¹	5.5 × 10 ⁻¹
Eniaetok	33f	15	5.3 × 10 ⁻¹	1.4 × 10 ¹	2.2 × 10 ⁰	4.0 × 10 ⁰	4.7 × 10 ⁰	8.0 × 10 ⁻¹	1.1 × 10 ⁰
Busch	38f	2	5.6 × 10 ⁻¹	1.1 × 10 ⁰	8.3 × 10 ⁻¹	8.3 × 10 ⁻¹	3.8 × 10 ⁻¹	-2.5 × 10 ⁻¹	4.8 × 10 ⁻¹
Rongelap	42f	78	5.1 × 10 ⁻¹	3.4 × 10 ¹	5.9 × 10 ⁰	7.6 × 10 ⁰	6.5 × 10 ⁰	1.7 × 10 ⁰	8.3 × 10 ⁻¹
Arbar	43f	4	5.7 × 10 ⁻¹	8.9 × 10 ⁰	1.4 × 10 ⁰	3.1 × 10 ⁰	3.9 × 10 ⁰	5.7 × 10 ⁻¹	1.2 × 10 ⁰
Tufa	47f	2	3.4 × 10 ⁰	4.7 × 10 ⁰	4.0 × 10 ⁰	4.0 × 10 ⁰	9.4 × 10 ⁻¹	1.4 × 10 ⁰	2.4 × 10 ⁻¹
Borukka	49f	9	2.8 × 10 ⁻¹	2.9 × 10 ⁰	1.0 × 10 ⁰	1.4 × 10 ⁰	9.5 × 10 ⁻¹	6.7 × 10 ⁻²	7.9 × 10 ⁻¹

Table A-1. The ^{137}Cs radionuclide concentration summary for vegetation (food crops) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991 (continued).

Island name	Island code	N ^a	pCi g ⁻¹ wet wt					Mean of logs	SD of logs	
			Minimum	Maximum	Median	Mean	SD			
Breadfruit Meat										
Rongelap	42f	30	1.0 × 10 ⁰	7.0 × 10 ⁰	3.0 × 10 ⁰	3.2 × 10 ⁰	1.5 × 10 ⁰	1.0 × 10 ⁰	4.9 × 10 ⁻¹	
Tacca Meat										
Busch	38f	1	2.3 × 10 ⁰	2.3 × 10 ⁰	2.3 × 10 ⁰	2.3 × 10 ⁰	0.0 × 10 ⁰	8.4 × 10 ⁻¹	0.0 × 10 ⁰	
Rongelap	42f	5	1.0 × 10 ⁰	1.5 × 10 ¹	4.3 × 10 ⁰	5.7 × 10 ⁰	5.4 × 10 ⁰	1.4 × 10 ⁰	9.6 × 10 ⁻¹	
Tufa	47f	2	5.2 × 10 ⁻¹	2.7 × 10 ⁰	1.6 × 10 ⁰	1.6 × 10 ⁰	1.5 × 10 ⁰	1.7 × 10 ⁻¹	1.2 × 10 ⁰	
Borukka	49f	4	5.2 × 10 ⁻¹	2.1 × 10 ⁰	1.2 × 10 ⁰	1.2 × 10 ⁰	7.5 × 10 ⁻¹	4.4 × 10 ⁻²	6.7 × 10 ⁻¹	
Lime Meat										
Rongelap	42f	9	1.2 × 10 ⁰	2.0 × 10 ⁰	1.5 × 10 ⁰	1.6 × 10 ⁰	2.5 × 10 ⁻¹	4.4 × 10 ⁻¹	1.6 × 10 ⁻¹	
Squash Meat										
Rongelap	42f	2	3.7 × 10 ⁰	7.8 × 10 ⁰	5.8 × 10 ⁰	5.8 × 10 ⁰	2.9 × 10 ⁰	1.7 × 10 ⁰	5.2 × 10 ⁻¹	

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table A-2. The ^{137}Cs radionuclide concentration summary for vegetation (indicator species) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.

through 1991.

Island name	Island code	N ^a	pCi g ⁻¹ wet wt.					Mean of logs	SD of logs	
			Minimum	Maximum	Median	Mean	SD			
Morinda Fruit										
Eniaetok	33f	1	2.5 × 10 ⁰	2.5 × 10 ⁰	2.5 × 10 ⁰	2.5 × 10 ⁰	0.0 × 10 ⁰	9.1 × 10 ⁻¹	0.0 × 10 ⁰	
Rongelap	42f	2	1.1 × 10 ⁰	2.1 × 10 ⁰	1.6 × 10 ⁰	1.6 × 10 ⁰	7.2 × 10 ⁻¹	4.4 × 10 ⁻¹	4.6 × 10 ⁻¹	
Arbar	43f	1	4.7 × 10 ⁻¹	4.7 × 10 ⁻¹	4.7 × 10 ⁻¹	4.7 × 10 ⁻¹	0.0 × 10 ⁰	-7.5 × 10 ⁻¹	0.0 × 10 ⁰	
Eniran	46f	3	2.8 × 10 ⁻¹	8.7 × 10 ⁻¹	5.6 × 10 ⁻¹	5.7 × 10 ⁻¹	3.0 × 10 ⁻¹	-6.7 × 10 ⁻¹	5.7 × 10 ⁻¹	
Tufa	47f	1	7.4 × 10 ⁻¹	7.4 × 10 ⁻¹	7.4 × 10 ⁻¹	7.4 × 10 ⁻¹	0.0 × 10 ⁰	-3.0 × 10 ⁻¹	0.0 × 10 ⁰	
Morinda Leaves										
Arbar	43f	1	3.8 × 10 ⁻¹	3.8 × 10 ⁻¹	3.8 × 10 ⁻¹	3.8 × 10 ⁻¹	0.0 × 10 ⁰	-9.7 × 10 ⁻¹	0.0 × 10 ⁰	
Eniran	46f	3	1.3 × 10 ⁻¹	5.6 × 10 ⁻¹	2.2 × 10 ⁻¹	3.1 × 10 ⁻¹	2.3 × 10 ⁻¹	-1.4 × 10 ⁰	7.4 × 10 ⁻¹	
Tufa	47f	1	6.6 × 10 ⁻¹	6.6 × 10 ⁻¹	6.6 × 10 ⁻¹	6.6 × 10 ⁻¹	0.0 × 10 ⁰	-4.2 × 10 ⁻¹	0.0 × 10 ⁰	
Guettardia Leaves										
Rongelap	42f	1	2.9 × 10 ⁰	2.9 × 10 ⁰	2.9 × 10 ⁰	2.9 × 10 ⁰	0.0 × 10 ⁰	1.1 × 10 ⁰	0.0 × 10 ⁰	
Arbar	43f	3	2.2 × 10 ⁻¹	1.4 × 10 ⁰	3.8 × 10 ⁻¹	6.7 × 10 ⁻¹	6.4 × 10 ⁻¹	-7.1 × 10 ⁻¹	9.5 × 10 ⁻¹	
Tufa	47f	1	2.5 × 10 ⁻¹	2.5 × 10 ⁻¹	2.5 × 10 ⁻¹	2.5 × 10 ⁻¹	0.0 × 10 ⁰	-1.4 × 10 ⁰	0.0 × 10 ⁰	
Messerschmidia Leaves										
Rongelap	42f	2	2.7 × 10 ⁻¹	7.1 × 10 ⁻¹	4.9 × 10 ⁻¹	4.9 × 10 ⁻¹	3.1 × 10 ⁻¹	-8.3 × 10 ⁻¹	6.9 × 10 ⁻¹	
Pisonia Leaves										
Tufa	47f	1	3.0 × 10 ⁰	3.0 × 10 ⁰	3.0 × 10 ⁰	3.0 × 10 ⁰	0.0 × 10 ⁰	1.1 × 10 ⁰	0.0 × 10 ⁰	
Pandanus Leaves										
Rongelap	42f	5	6.1 × 10 ⁻¹	9.3 × 10 ⁰	7.3 × 10 ⁰	5.6 × 10 ⁰	3.8 × 10 ⁰	1.4 × 10 ⁰	1.2 × 10 ⁰	
Scaevola Leaves										
Arbar	43f	4	1.7 × 10 ⁻¹	5.5 × 10 ⁻¹	2.7 × 10 ⁻¹	3.2 × 10 ⁻¹	1.7 × 10 ⁻¹	-1.3 × 10 ⁰	5.0 × 10 ⁻¹	
Eniran	46f	1	1.9 × 10 ⁻¹	1.9 × 10 ⁻¹	1.9 × 10 ⁻¹	1.9 × 10 ⁻¹	0.0 × 10 ⁰	-1.7 × 10 ⁰	0.0 × 10 ⁰	
Tufa	47f	2	2.9 × 10 ⁻¹	3.4 × 10 ⁻¹	3.1 × 10 ⁻¹	3.1 × 10 ⁻¹	3.4 × 10 ⁻²	-1.2 × 10 ⁰	1.1 × 10 ⁻¹	
Borukka	49f	1	3.7 × 10 ⁰	3.7 × 10 ⁰	3.7 × 10 ⁰	3.7 × 10 ⁰	0.0 × 10 ⁰	1.3 × 10 ⁰	0.0 × 10 ⁰	
At-At Vine										
Tufa	47f	1	3.4 × 10 ⁻¹	3.4 × 10 ⁻¹	3.4 × 10 ⁻¹	3.4 × 10 ⁻¹	0.0 × 10 ⁰	-1.1 × 10 ⁰	0.0 × 10 ⁰	
Tabero Fruit										
Rongelap	42f	3	1.4 × 10 ⁰	2.1 × 10 ⁰	2.0 × 10 ⁰	1.8 × 10 ⁰	3.8 × 10 ⁻¹	5.8 × 10 ⁻¹	2.3 × 10 ⁻¹	
Tabero Leaves										
Rongelap	42f	1	1.2 × 10 ⁰	1.2 × 10 ⁰	1.2 × 10 ⁰	1.2 × 10 ⁰	0.0 × 10 ⁰	2.1 × 10 ⁻¹	0.0 × 10 ⁰	
Coconut Fronds										
Rongelap	42f	16	2.7 × 10 ⁻¹	7.3 × 10 ⁰	1.1 × 10 ⁰	2.4 × 10 ⁰	2.4 × 10 ⁰	4.0 × 10 ⁻¹	9.9 × 10 ⁻¹	

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Appendix B

**Radionuclide concentration summary in animal samples
collected during the 1978 NMIRS and from 1986 through 1991
in the southern islands of Rongelap Atoll.**

Table B-1. The ^{137}Cs radionuclide concentration summary for animals (food source) collected from the southern islands of Rongelap Atoll during the 1978 NMIRS together with our recent trips in 1986 through 1991.

Island name	Island code	N ^a	pCi g ⁻¹ wet wt.					Mean of logs	SD of logs	
			Minimum	Maximum	Median	Mean	SD			
Coconut Crab Muscle										
Erapuotsu	32f	1	4.9 × 10 ⁰	4.9 × 10 ⁰	4.88 × 10 ⁰	4.9 × 10 ⁰	0.0 × 10 ⁰	1.6 × 10 ⁰	0.0 × 10 ⁰	
Busch	38f	1	1.4 × 10 ⁰	1.4 × 10 ⁰	1.39 × 10 ⁰	1.4 × 10 ⁰	0.0 × 10 ⁰	3.3 × 10 ⁻¹	0.0 × 10 ⁰	
Arbar	43f	2	2.3 × 10 ⁰	2.6 × 10 ⁰	2.5 × 10 ⁰	2.5 × 10 ⁰	1.9 × 10 ⁻¹	9.0 × 10 ⁻¹	7.8 × 10 ⁻²	
Eniran	46f	2	1.1 × 10 ⁰	2.0 × 10 ⁰	1.5 × 10 ⁰	1.5 × 10 ⁰	6.5 × 10 ⁻¹	3.8 × 10 ⁻¹	4.4 × 10 ⁻¹	
Tufa	47f	2	1.2 × 10 ⁰	2.0 × 10 ⁰	1.6 × 10 ⁰	1.6 × 10 ⁰	6.1 × 10 ⁻¹	4.2 × 10 ⁻¹	3.9 × 10 ⁻¹	
Borukka	49f	1	2.1 × 10 ⁰	2.1 × 10 ⁰	2.1 × 10 ⁰	2.1 × 10 ⁰	0.0 × 10 ⁰	7.3 × 10 ⁻¹	0.0 × 10 ⁰	
Coconut Crab Hepatopancreas										
Erapuotsu	32f	1	3.4 × 10 ⁰	3.4 × 10 ⁰	3.4 × 10 ⁰	3.4 × 10 ⁰	0.0 × 10 ⁰	1.2 × 10 ⁰	0.0 × 10 ⁰	
Busch	38f	1	9.8 × 10 ⁻¹	9.8 × 10 ⁻¹	9.8 × 10 ⁻¹	9.8 × 10 ⁻¹	0.0 × 10 ⁰	-2.3 × 10 ⁻²	0.0 × 10 ⁰	
Eniran	46f	2	9.6 × 10 ⁻¹	1.3 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	2.2 × 10 ⁻¹	1.1 × 10 ⁻¹	2.0 × 10 ⁻¹	
Tufa	47f	2	9.0 × 10 ⁻¹	1.3 × 10 ⁰	1.1 × 10 ⁰	1.1 × 10 ⁰	2.2 × 10 ⁻¹	7.0 × 10 ⁻²	2.5 × 10 ⁻¹	
Borukka	49f	1	1.9 × 10 ⁰	1.9 × 10 ⁰	1.9 × 10 ⁰	1.9 × 10 ⁰	0.0 × 10 ⁰	6.3 × 10 ⁻¹	0.0 × 10 ⁰	
Pig Heart										
Rongelap	42f	4	9.7 × 10 ⁰	2.2 × 10 ¹	1.3 × 10 ¹	1.4 × 10 ¹	5.6 × 10 ⁰	2.6 × 10 ⁰	3.8 × 10 ⁻¹	
Pig Liver										
Rongelap	42f	6	3.2 × 10 ⁰	8.5 × 10 ⁰	5.3 × 10 ⁰	5.6 × 10 ⁰	2.0 × 10 ⁰	1.7 × 10 ⁰	3.7 × 10 ⁻¹	
Pig Muscle										
Rongelap	42f	7	5.3 × 10 ⁰	2.9 × 10 ¹	1.3 × 10 ¹	1.4 × 10 ¹	7.8 × 10 ⁰	2.5 × 10 ⁰	5.6 × 10 ⁻¹	

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Appendix C

Radionuclide concentration summary of all soil profile samples collected during the 1978 NMIRS and from 1986 through 1991 in the southern islands of Rongelap Atoll.

Table C-1. Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Erapuotsu Island (F32).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	3	7.3×10^0	4.1×10^1	2.2×10^1	2.3×10^1	1.7×10^1	2.9×10^0	8.8×10^{-1}
05-10	3	3.0×10^1	4.3×10^1	3.1×10^1	3.5×10^1	7.4×10^0	3.5×10^0	2.0×10^{-1}
10-15	3	2.6×10^0	1.7×10^1	5.0×10^0	8.1×10^0	7.6×10^0	1.8×10^0	9.5×10^{-1}
15-25	3	7.0×10^{-1}	1.9×10^0	1.6×10^0	1.4×10^0	6.3×10^1	2.6×10^{-1}	5.4×10^{-1}
25-40	3	8.3×10^{-1}	1.2×10^0	8.6×10^{-1}	9.6×10^{-1}	1.9×10^{-1}	-5.8×10^{-2}	1.9×10^{-1}
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
C-21	0-05	3	7.3×10^0	4.1×10^1	2.2×10^1	2.3×10^1	2.9×10^0	8.8×10^{-1}
	0-10	3	1.9×10^1	4.2×10^1	2.6×10^1	2.9×10^1	3.3×10^0	4.1×10^{-1}
	0-15	3	1.4×10^1	3.4×10^1	1.8×10^1	2.2×10^1	3.0×10^0	4.5×10^{-1}
	0-25	3	9.1×10^0	2.1×10^1	1.1×10^1	1.4×10^1	2.6×10^0	4.3×10^{-1}
	0-40	3	6.1×10^0	1.3×10^1	7.4×10^0	9.0×10^0	2.1×10^0	4.1×10^{-1}
	0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-2. Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Eniaetok Island (F33).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	21	6.4×10^{-2}	5.7×10^1	2.1×10^1	2.1×10^1	1.7×10^1	2.3×10^0	1.8×10^0
05-10	21	2.8×10^{-2}	2.5×10^1	8.0×10^0	9.1×10^0	6.4×10^0	1.8×10^0	1.4×10^0
10-15	20	2.5×10^{-2}	1.4×10^1	3.3×10^0	4.4×10^0	3.4×10^0	1.1×10^0	1.3×10^0
15-25	21	4.3×10^{-2}	3.2×10^0	1.3×10^0	1.5×10^0	9.4×10^{-1}	7.6×10^{-2}	1.0×10^0
25-40	21	6.0×10^{-2}	2.4×10^0	4.9×10^{-1}	7.5×10^{-1}	6.1×10^{-1}	-6.5×10^{-1}	9.7×10^{-1}
40-60	6	2.3×10^{-1}	6.8×10^0	5.0×10^{-1}	1.6×10^0	2.6×10^0	-3.3×10^{-1}	1.2×10^0
0-05	21	6.4×10^{-2}	5.7×10^1	2.1×10^1	2.1×10^1	1.7×10^1	2.3×10^0	1.8×10^0
0-10	21	4.6×10^{-2}	3.5×10^1	1.3×10^1	1.5×10^1	1.0×10^1	2.2×10^0	1.5×10^0
0-15	20	4.0×10^{-2}	2.5×10^1	1.0×10^1	1.1×10^1	7.3×10^0	2.0×10^0	1.4×10^0
0-25	20	4.1×10^{-2}	1.5×10^1	7.2×10^0	7.2×10^0	4.5×10^0	1.6×10^0	1.3×10^0
0-40	20	5.0×10^{-2}	9.7×10^0	4.8×10^0	4.8×10^0	2.9×10^0	1.2×10^0	1.2×10^0
0-60	6	9.9×10^{-1}	4.6×10^0	2.7×10^0	2.5×10^0	1.4×10^0	7.9×10^{-1}	6.1×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-3. Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Busch Island (F38).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	5	2.2×10^0	8.5×10^1	1.2×10^1	2.7×10^1	3.4×10^1	2.6×10^0	1.4×10^0
05-10	5	1.1×10^1	3.4×10^1	2.2×10^1	2.3×10^1	1.1×10^1	3.0×10^0	5.3×10^{-1}
10-15	5	6.8×10^0	2.2×10^1	1.6×10^1	1.5×10^1	6.7×10^0	2.6×10^0	5.3×10^{-1}
15-25	5	7.5×10^{-1}	8.1×10^0	6.0×10^0	4.9×10^0	3.0×10^0	1.3×10^0	9.6×10^{-1}
25-40	5	3.1×10^{-1}	4.0×10^0	1.4×10^0	2.0×10^0	1.7×10^0	2.6×10^{-1}	1.1×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	5	2.2×10^0	8.5×10^1	1.2×10^1	2.7×10^1	3.4×10^1	2.6×10^0	1.4×10^0
0-10	5	6.8×10^0	5.3×10^1	2.3×10^1	2.5×10^1	1.9×10^1	3.0×10^0	8.4×10^{-1}
0-15	5	1.2×10^1	3.8×10^1	2.2×10^1	2.2×10^1	1.1×10^1	3.0×10^0	4.9×10^{-1}
0-25	5	9.5×10^0	2.3×10^1	1.5×10^1	1.5×10^1	5.4×10^0	2.7×10^0	3.5×10^{-1}
0-40	5	7.3×10^0	1.5×10^1	9.9×10^0	1.0×10^1	2.9×10^0	2.3×10^0	2.7×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-4. Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trips in 1986 through 1989 on Rongelap Island (F42).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	294	8.3×10^{-2}	5.7×10^1	1.4×10^1	1.4×10^1	9.6×10^0	2.3×10^0	1.1×10^0
05-10	114	6.9×10^{-2}	3.3×10^1	6.5×10^0	7.4×10^0	5.8×10^0	1.7×10^0	9.3×10^{-1}
10-15	114	2.0×10^{-1}	3.2×10^1	2.9×10^0	4.0×10^0	4.2×10^0	9.5×10^{-1}	9.8×10^{-1}
15-25	115	3.9×10^{-2}	1.2×10^1	1.2×10^0	1.8×10^0	1.8×10^0	7.2×10^{-2}	1.1×10^0
25-40	113	2.4×10^{-2}	3.0×10^0	3.6×10^{-1}	6.0×10^{-1}	6.6×10^{-1}	-1.0×10^0	1.0×10^0
40-60	93	2.9×10^{-2}	2.4×10^0	1.9×10^{-1}	3.6×10^{-1}	4.6×10^{-1}	-1.6×10^0	1.1×10^0
0-05	294	8.3×10^{-2}	5.7×10^1	1.4×10^1	1.4×10^1	9.6×10^0	2.3×10^0	1.1×10^0
0-10	113	8.5×10^{-2}	3.8×10^1	9.4×10^0	1.1×10^1	7.3×10^0	2.2×10^0	8.7×10^{-1}
0-15	111	1.2×10^{-1}	3.4×10^1	7.5×10^0	8.8×10^0	5.7×10^0	1.9×10^0	8.0×10^{-1}
0-25	110	1.7×10^{-1}	2.5×10^1	5.1×10^0	6.0×10^0	3.8×10^0	1.6×10^0	7.0×10^{-1}
0-40	108	1.3×10^{-1}	1.6×10^1	3.5×10^0	3.9×10^0	2.4×10^0	1.2×10^0	6.7×10^{-1}
0-60	88	1.4×10^{-1}	1.1×10^1	2.4×10^0	2.7×10^0	1.7×10^0	8.0×10^{-1}	6.5×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-5. Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1991 on Arbar Island (F43).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	13	4.0×10^0	3.8×10^1	1.0×10^1	1.4×10^1	1.1×10^1	2.4×10^0	6.7×10^{-1}
05-10	13	2.0×10^0	3.7×10^1	7.3×10^0	1.1×10^1	1.0×10^1	2.0×10^0	9.5×10^{-1}
10-15	13	6.9×10^{-1}	2.4×10^1	2.4×10^0	6.6×10^0	8.2×10^0	1.2×10^0	1.2×10^0
15-25	12	1.5×10^{-1}	1.6×10^1	1.5×10^0	3.3×10^0	4.9×10^0	3.8×10^{-1}	1.3×10^0
25-40	11	1.2×10^{-2}	1.8×10^0	1.0×10^0	8.0×10^{-1}	6.4×10^{-1}	-8.7×10^{-1}	1.6×10^0
40-60	4	5.9×10^{-2}	1.2×10^0	3.1×10^{-1}	4.7×10^{-1}	5.3×10^{-1}	-1.5×10^0	1.5×10^0
C-25	0-05	13	4.0×10^0	3.8×10^1	1.0×10^1	1.4×10^1	2.4×10^0	6.7×10^{-1}
	0-10	12	3.0×10^0	3.6×10^1	1.0×10^1	1.3×10^1	2.3×10^0	7.1×10^{-1}
	0-15	12	2.5×10^0	3.2×10^1	7.3×10^0	1.1×10^1	2.1×10^0	7.5×10^{-1}
	0-25	11	1.7×10^0	2.3×10^1	5.3×10^0	8.0×10^0	1.8×10^0	8.0×10^{-1}
	0-40	9	1.1×10^0	1.5×10^1	3.7×10^0	5.5×10^0	1.4×10^0	8.8×10^{-1}
	0-60	3	7.2×10^{-1}	4.2×10^0	2.3×10^0	2.4×10^0	6.5×10^{-1}	9.0×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-6. Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Eniran Island (F46).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	6	6.2×10^0	2.2×10^1	1.3×10^1	1.3×10^1	6.0×10^0	2.5×10^0	4.8×10^{-1}
05-10	7	1.7×10^0	2.3×10^1	8.3×10^0	9.8×10^0	7.3×10^0	2.0×10^0	8.6×10^{-1}
10-15	7	5.3×10^{-1}	7.3×10^0	3.3×10^0	3.9×10^0	2.5×10^0	1.1×10^0	9.1×10^{-1}
15-25	7	1.4×10^{-1}	6.1×10^0	2.0×10^0	2.4×10^0	2.3×10^0	3.4×10^{-1}	1.3×10^0
25-40	7	4.2×10^{-2}	2.5×10^0	3.4×10^{-1}	9.1×10^{-1}	1.0×10^0	-8.4×10^{-1}	1.5×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	6	6.2×10^0	2.2×10^1	1.3×10^1	1.3×10^1	6.0×10^0	2.5×10^0	4.8×10^{-1}
0-10	6	6.6×10^0	1.9×10^1	1.2×10^1	1.2×10^1	5.5×10^0	2.4×10^0	4.7×10^{-1}
0-15	6	5.5×10^0	1.5×10^1	9.2×10^0	9.7×10^0	3.9×10^0	2.2×10^0	4.1×10^{-1}
0-25	6	3.7×10^0	9.6×10^0	6.6×10^0	6.9×10^0	2.1×10^0	1.9×10^0	3.3×10^{-1}
0-40	6	2.4×10^0	6.1×10^0	4.7×10^0	4.7×10^0	1.4×10^0	1.5×10^0	3.4×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-7. Cesium-137 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Tufa Island (F47).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	8	6.5×10^0	6.1×10^1	1.8×10^1	2.4×10^1	1.8×10^1	2.9×10^0	7.0×10^{-1}
05-10	8	6.2×10^0	5.5×10^1	2.1×10^1	2.2×10^1	1.6×10^1	2.9×10^0	7.6×10^{-1}
10-15	8	2.0×10^0	3.1×10^1	8.4×10^0	1.1×10^1	9.5×10^0	2.0×10^0	9.3×10^{-1}
15-25	8	1.1×10^0	1.6×10^1	3.0×10^0	4.4×10^0	4.8×10^0	1.1×10^0	8.4×10^{-1}
25-40	8	2.5×10^{-1}	5.2×10^0	1.1×10^0	1.9×10^0	1.8×10^0	1.9×10^{-1}	1.1×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	8	6.5×10^0	6.1×10^1	1.8×10^1	2.4×10^1	1.8×10^1	2.9×10^0	7.0×10^{-1}
0-10	8	1.0×10^1	4.5×10^1	1.7×10^1	2.3×10^1	1.4×10^1	3.0×10^0	6.0×10^{-1}
0-15	8	7.9×10^0	3.6×10^1	1.5×10^1	1.9×10^1	1.1×10^1	2.8×10^0	6.1×10^{-1}
0-25	8	5.3×10^0	2.8×10^1	9.8×10^0	1.3×10^1	8.1×10^0	2.4×10^0	6.0×10^{-1}
0-40	8	3.5×10^0	2.0×10^1	6.9×10^0	8.8×10^0	5.6×10^0	2.0×10^0	6.1×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-8. Cesium-137 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Borukka Island (F49).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	13	6.7×10^0	5.1×10^1	2.2×10^1	2.5×10^1	1.2×10^1	3.1×10^0	5.4×10^{-1}
05-10	13	4.0×10^0	3.9×10^1	1.3×10^1	1.5×10^1	1.1×10^1	2.5×10^0	6.5×10^{-1}
10-15	13	2.7×10^0	2.5×10^1	7.2×10^0	9.1×10^0	6.6×10^0	2.0×10^0	7.1×10^{-1}
15-25	13	1.0×10^0	1.3×10^1	2.6×10^0	4.3×10^0	3.8×10^0	1.1×10^0	8.3×10^{-1}
25-40	13	1.6×10^{-1}	4.2×10^0	8.4×10^{-1}	1.4×10^0	1.3×10^0	-8.3×10^{-2}	1.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
C-28	0-05	13	6.7×10^0	5.1×10^1	2.2×10^1	2.5×10^1	3.1×10^0	5.4×10^{-1}
	0-10	13	1.3×10^1	4.3×10^1	1.7×10^1	2.0×10^1	3.0×10^0	3.6×10^{-1}
	0-15	13	9.4×10^0	3.7×10^1	1.4×10^1	1.7×10^1	2.7×10^0	3.8×10^{-1}
	0-25	13	6.1×10^0	2.8×10^1	9.5×10^0	1.2×10^1	2.4×10^0	4.1×10^{-1}
	0-40	13	3.9×10^0	1.9×10^1	6.1×10^0	7.8×10^0	2.0×10^0	4.3×10^{-1}
	0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-9. Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Eniaetok Island (F33).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	6	7.3×10^{-1}	1.5×10^1	5.4×10^0	6.6×10^0	5.5×10^0	1.5×10^0	1.2×10^0
05-10	6	2.9×10^0	2.0×10^1	8.1×10^0	9.2×10^0	6.0×10^0	2.0×10^0	6.9×10^{-1}
10-15	6	1.5×10^0	1.4×10^1	3.4×10^0	6.1×10^0	5.3×10^0	1.5×10^0	8.8×10^{-1}
15-25	6	5.3×10^{-1}	7.9×10^0	2.1×10^0	3.3×10^0	2.9×10^0	8.6×10^{-1}	9.7×10^{-1}
25-40	6	3.2×10^{-1}	2.3×10^1	3.1×10^0	5.9×10^0	8.4×10^0	1.0×10^0	1.4×10^0
40-60	6	2.4×10^{-1}	4.9×10^0	2.7×10^0	2.4×10^0	1.8×10^0	4.9×10^{-1}	1.2×10^0
0-05	6	7.3×10^{-1}	1.5×10^1	5.4×10^0	6.6×10^0	5.5×10^0	1.5×10^0	1.2×10^0
0-10	6	1.8×10^0	1.7×10^1	6.8×10^0	7.9×10^0	5.7×10^0	1.8×10^0	8.3×10^{-1}
0-15	6	1.7×10^0	1.6×10^1	5.9×10^0	7.3×10^0	4.8×10^0	1.8×10^0	7.3×10^{-1}
0-25	6	1.3×10^0	1.2×10^1	4.4×10^0	5.7×10^0	3.8×10^0	1.5×10^0	7.8×10^{-1}
0-40	6	9.0×10^{-1}	1.4×10^1	4.0×10^0	5.8×10^0	4.7×10^0	1.4×10^0	9.4×10^{-1}
0-60	6	6.8×10^{-1}	1.0×10^1	4.0×10^0	4.7×10^0	3.4×10^0	1.2×10^0	9.5×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-10. Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Rongelap Island (F42).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	20	1.2×10^{-1}	1.3×10^1	5.2×10^0	4.8×10^0	3.1×10^0	1.2×10^0	1.1×10^0
05-10	20	1.6×10^{-1}	1.6×10^1	3.3×10^0	5.5×10^0	5.0×10^0	1.2×10^0	1.2×10^0
10-15	20	2.4×10^{-2}	1.4×10^1	3.1×10^0	4.1×10^0	4.1×10^0	7.6×10^{-1}	1.5×10^0
15-25	20	3.9×10^{-1}	8.6×10^0	2.2×10^0	3.1×10^0	2.4×10^0	7.7×10^{-1}	9.3×10^{-1}
25-40	21	1.1×10^{-2}	7.5×10^0	1.4×10^0	1.8×10^0	1.7×10^0	-3.8×10^{-2}	1.5×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	20	1.2×10^{-1}	1.3×10^1	5.2×10^0	4.8×10^0	3.1×10^0	1.2×10^0	1.1×10^0
0-10	19	1.4×10^{-1}	1.5×10^1	4.1×10^0	5.2×10^0	3.9×10^0	1.3×10^0	1.1×10^0
0-15	18	2.8×10^{-1}	1.3×10^1	3.7×10^0	4.8×10^0	3.8×10^0	1.2×10^0	1.0×10^0
0-25	17	7.6×10^{-1}	9.9×10^0	3.6×10^0	4.2×10^0	3.1×10^0	1.1×10^0	8.5×10^{-1}
0-40	17	6.1×10^{-1}	7.4×10^0	2.9×10^0	3.3×10^0	2.2×10^0	9.2×10^{-1}	8.3×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-11. Strontium-90 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS on Borukka Island (F49).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	3	1.1×10^1	1.2×10^1	1.2×10^1	1.2×10^1	6.1×10^{-1}	2.5×10^0	5.4×10^{-2}
05-10	3	3.8×10^0	9.9×10^0	9.3×10^0	7.6×10^0	3.4×10^0	2.0×10^0	5.4×10^{-1}
10-15	3	3.4×10^0	7.3×10^0	6.2×10^0	5.6×10^0	2.0×10^0	1.7×10^0	4.0×10^{-1}
15-25	3	1.6×10^0	5.2×10^0	2.4×10^0	3.0×10^0	1.9×10^0	9.8×10^{-1}	6.1×10^{-1}
25-40	2	1.7×10^0	3.2×10^0	2.5×10^0	2.5×10^0	1.1×10^0	8.5×10^{-1}	4.5×10^{-1}
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	3	1.1×10^1	1.2×10^1	1.2×10^1	1.2×10^1	6.1×10^{-1}	2.5×10^0	5.4×10^{-2}
0-10	3	7.8×10^0	1.1×10^1	1.0×10^1	9.6×10^0	1.5×10^0	2.3×10^0	1.7×10^{-1}
0-15	3	6.4×10^0	9.5×10^0	9.0×10^0	8.3×10^0	1.7×10^0	2.1×10^0	2.2×10^{-1}
0-25	3	4.8×10^0	7.8×10^0	6.0×10^0	6.2×10^0	1.5×10^0	1.8×10^0	2.5×10^{-1}
0-40	2	4.4×10^0	6.1×10^0	5.2×10^0	5.2×10^0	1.2×10^0	1.6×10^0	2.3×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-12. Plutonium-239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1989 on Eniaetok Island (F33).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	21	2.3×10^{-2}	2.0×10^1	6.6×10^0	7.2×10^0	5.9×10^0	1.3×10^0	1.7×10^0
05-10	6	1.5×10^0	7.2×10^0	3.0×10^0	3.3×10^0	2.0×10^0	1.1×10^0	5.5×10^{-1}
10-15	6	2.3×10^{-1}	6.9×10^0	2.0×10^0	2.6×10^0	2.6×10^0	3.1×10^{-1}	1.4×10^0
15-25	6	4.5×10^{-3}	9.3×10^0	3.1×10^{-1}	1.8×10^0	3.7×10^0	-1.3×10^0	2.4×10^0
25-40	6	9.9×10^{-3}	8.7×10^{-2}	5.9×10^{-2}	5.6×10^{-2}	2.6×10^{-2}	-3.1×10^0	7.8×10^{-1}
40-60	6	1.6×10^{-2}	6.8×10^{-1}	8.9×10^{-2}	1.7×10^{-1}	2.5×10^{-1}	-2.5×10^0	1.3×10^0
0-05	21	2.3×10^{-2}	2.0×10^1	6.6×10^0	7.2×10^0	5.9×10^0	1.3×10^0	1.7×10^0
0-10	6	8.7×10^{-1}	5.9×10^0	2.6×10^0	2.6×10^0	1.8×10^0	7.7×10^{-1}	7.1×10^{-1}
0-15	6	6.5×10^{-1}	6.2×10^0	2.1×10^0	2.6×10^0	1.9×10^0	7.5×10^{-1}	7.3×10^{-1}
0-25	6	3.9×10^{-1}	7.5×10^0	1.4×10^0	2.3×10^0	2.6×10^0	4.2×10^{-1}	9.5×10^{-1}
0-40	6	2.5×10^{-1}	4.7×10^0	8.7×10^{-1}	1.4×10^0	1.6×10^0	-3.3×10^{-2}	9.4×10^{-1}
0-60	6	1.7×10^{-1}	3.2×10^0	7.1×10^{-1}	1.0×10^0	1.1×10^0	-3.6×10^{-1}	9.4×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-13. Plutonium-239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1986 on Rongelap Island (F42).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	77	2.7×10^{-1}	1.3×10^1	2.8×10^0	3.4×10^0	2.8×10^0	8.8×10^{-1}	9.3×10^{-1}
05-10	16	2.1×10^{-1}	1.0×10^1	1.0×10^0	2.1×10^0	2.5×10^0	2.0×10^{-1}	1.1×10^0
10-15	18	4.7×10^{-2}	3.5×10^0	5.0×10^{-1}	9.2×10^{-1}	9.8×10^{-1}	-7.0×10^{-1}	1.2×10^0
15-25	18	1.8×10^{-2}	1.1×10^0	2.0×10^{-1}	2.7×10^{-1}	2.6×10^{-1}	-1.8×10^0	1.1×10^0
5-40	19	5.8×10^{-4}	4.9×10^{-1}	9.0×10^{-2}	1.3×10^{-1}	1.3×10^{-1}	-2.8×10^0	1.7×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	77	2.7×10^{-1}	1.3×10^1	2.8×10^0	3.4×10^0	2.8×10^0	8.8×10^{-1}	9.3×10^{-1}
0-10	15	3.6×10^{-1}	6.2×10^0	1.8×10^0	2.6×10^0	2.1×10^0	5.4×10^{-1}	1.0×10^0
0-15	14	2.7×10^{-1}	5.2×10^0	1.4×10^0	2.0×10^0	1.6×10^0	3.0×10^{-1}	1.0×10^0
0-25	13	1.8×10^{-1}	3.5×10^0	1.2×10^0	1.4×10^0	1.1×10^0	-7.9×10^{-2}	1.0×10^0
0-40	13	1.1×10^{-1}	2.3×10^0	8.2×10^{-1}	9.0×10^{-1}	6.6×10^{-1}	-4.3×10^{-1}	9.4×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-14. Plutonium-239+240 radionuclide concentration summary for all soil profiles taken during the 1978 NMIRS together with our recent trip in 1989 on Borukka Island (F49).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	13	7.4×10^{-1}	1.6×10^1	5.3×10^0	6.9×10^0	4.0×10^0	1.7×10^0	7.4×10^{-1}
05-10	3	5.0×10^{-1}	1.6×10^0	6.6×10^{-1}	9.3×10^{-1}	6.1×10^{-1}	-2.1×10^{-1}	6.2×10^{-1}
10-15	3	2.4×10^{-1}	3.9×10^{-1}	2.5×10^{-1}	2.9×10^{-1}	8.8×10^{-2}	-1.3×10^0	2.8×10^{-1}
15-25	3	9.2×10^{-2}	1.9×10^{-1}	1.8×10^{-1}	1.6×10^{-1}	5.6×10^{-2}	-1.9×10^0	4.2×10^{-1}
25-40	2	4.8×10^{-2}	9.1×10^{-2}	6.9×10^{-2}	6.9×10^{-2}	3.0×10^{-2}	-2.7×10^0	4.5×10^{-1}
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	13	7.4×10^{-1}	1.6×10^1	5.3×10^0	6.9×10^0	4.0×10^0	1.7×10^0	7.4×10^{-1}
0-10	3	2.9×10^0	5.0×10^0	4.3×10^0	4.1×10^0	1.1×10^0	1.4×10^0	2.8×10^{-1}
0-15	3	2.1×10^0	3.4×10^0	2.9×10^0	2.8×10^0	6.8×10^{-1}	1.0×10^0	2.6×10^{-1}
0-25	3	1.3×10^0	2.1×10^0	1.8×10^0	1.7×10^0	3.9×10^{-1}	5.4×10^{-1}	2.4×10^{-1}
0-40	2	1.2×10^0	1.3×10^0	1.3×10^0	1.3×10^0	1.2×10^{-1}	2.2×10^{-1}	9.8×10^{-2}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-15. Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Erapuotsu Island (F32).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	3	3.2×10^0	1.5×10^1	5.2×10^0	7.7×10^0	6.1×10^0	1.8×10^0	7.8×10^{-1}
05-10	2	4.5×10^0	4.6×10^0	4.5×10^0	4.5×10^0	1.1×10^{-1}	1.5×10^0	2.3×10^{-2}
10-15	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
15-25	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
25-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	3	3.2×10^0	1.5×10^1	5.2×10^0	7.7×10^0	6.1×10^0	1.8×10^0	7.8×10^{-1}
0-10	2	3.9×10^0	4.8×10^0	4.4×10^0	4.4×10^0	6.6×10^{-1}	1.5×10^0	1.5×10^{-1}
0-15	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-25	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-16. Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Eniaetok Island (F33).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	18	2.4×10^{-1}	1.1×10^1	5.5×10^0	5.1×10^0	3.6×10^0	1.2×10^0	1.2×10^0
05-10	15	1.9×10^{-1}	8.2×10^0	8.8×10^{-1}	1.7×10^0	2.1×10^0	-3.9×10^{-2}	1.1×10^0
10-15	9	1.1×10^{-1}	2.0×10^0	6.1×10^{-1}	7.0×10^{-1}	6.2×10^{-1}	-7.7×10^{-1}	1.0×10^0
15-25	6	5.8×10^{-2}	4.7×10^{-1}	1.1×10^{-1}	1.6×10^{-1}	1.6×10^{-1}	-2.1×10^0	7.3×10^{-1}
25-40	6	1.5×10^{-2}	1.1×10^{-1}	2.7×10^{-2}	4.1×10^{-2}	3.5×10^{-2}	-3.4×10^0	6.7×10^{-1}
40-60	5	2.8×10^{-3}	3.4×10^{-1}	6.7×10^{-2}	1.1×10^{-1}	1.3×10^{-1}	-3.1×10^0	1.8×10^0
0-05	18	2.4×10^{-1}	1.1×10^1	5.5×10^0	5.1×10^0	3.6×10^0	1.2×10^0	1.2×10^0
0-10	15	3.9×10^{-1}	6.9×10^0	2.8×10^0	3.1×10^0	2.0×10^0	8.4×10^{-1}	8.8×10^{-1}
0-15	8	4.6×10^{-1}	4.9×10^0	1.6×10^0	2.1×10^0	1.7×10^0	3.9×10^{-1}	9.0×10^{-1}
0-25	6	3.2×10^{-1}	1.5×10^0	7.6×10^{-1}	7.8×10^{-1}	4.5×10^{-1}	-4.1×10^{-1}	6.3×10^{-1}
0-40	5	2.1×10^{-1}	9.2×10^{-1}	3.7×10^{-1}	4.8×10^{-1}	3.1×10^{-1}	-9.2×10^{-1}	6.7×10^{-1}
0-60	5	1.4×10^{-1}	6.4×10^{-1}	2.8×10^{-1}	3.5×10^{-1}	2.3×10^{-1}	-1.2×10^0	7.0×10^{-1}

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-17. Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Busch Island (F38).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	4	1.0×10^0	1.6×10^1	5.4×10^0	6.9×10^0	7.2×10^0	1.3×10^0	1.4×10^0
05-10	2	4.1×10^0	7.9×10^0	6.0×10^0	6.0×10^0	2.7×10^0	1.7×10^0	4.7×10^{-1}
10-15	3	2.1×10^0	4.5×10^0	2.6×10^0	3.1×10^0	1.2×10^0	1.1×10^0	3.8×10^{-1}
15-25	3	5.0×10^{-1}	1.1×10^0	5.1×10^{-1}	6.9×10^{-1}	3.1×10^{-1}	-4.4×10^{-1}	4.2×10^{-1}
25-40	1	4.3×10^{-1}	4.3×10^{-1}	4.3×10^{-1}	4.3×10^{-1}	0.0×10^0	-8.6×10^{-1}	0.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	4	1.0×10^0	1.6×10^1	5.4×10^0	6.9×10^0	7.2×10^0	1.3×10^0	1.4×10^0
0-10	2	2.6×10^0	4.5×10^0	3.5×10^0	3.5×10^0	1.3×10^0	1.2×10^0	3.8×10^{-1}
0-15	2	2.5×10^0	4.5×10^0	3.5×10^0	3.5×10^0	1.4×10^0	1.2×10^0	4.3×10^{-1}
0-25	2	1.9×10^0	2.9×10^0	2.4×10^0	2.4×10^0	7.0×10^{-1}	8.5×10^{-1}	3.0×10^{-1}
0-40	1	1.3×10^0	1.3×10^0	1.3×10^0	1.3×10^0	0.0×10^0	2.9×10^{-1}	0.0×10^0
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-18. Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trips in 1986 through 1989 on Rongelap Island (F42).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	267	4.0×10^{-2}	1.1×10^1	1.6×10^0	2.0×10^0	1.7×10^0	3.0×10^{-1}	9.7×10^{-1}
05-10	84	4.9×10^{-2}	7.2×10^0	6.8×10^{-1}	1.3×10^0	1.5×10^0	-2.8×10^{-1}	1.0×10^0
10-15	51	7.6×10^{-4}	2.6×10^0	4.8×10^{-1}	6.1×10^{-1}	5.5×10^{-1}	-1.1×10^0	1.4×10^0
15-25	35	7.5×10^{-3}	1.4×10^0	1.8×10^{-1}	3.1×10^{-1}	3.6×10^{-1}	-1.8×10^0	1.2×10^0
25-40	22	7.5×10^{-4}	4.4×10^{-1}	4.6×10^{-2}	7.9×10^{-2}	9.9×10^{-2}	-3.4×10^0	1.6×10^0
40-60	17	1.2×10^{-3}	4.0×10^{-1}	2.0×10^{-2}	7.1×10^{-2}	1.1×10^{-1}	-4.0×10^0	1.9×10^0
0-05	267	4.0×10^{-2}	1.1×10^1	1.6×10^0	2.0×10^0	1.7×10^0	3.0×10^{-1}	9.7×10^{-1}
0-10	78	4.4×10^{-2}	8.0×10^0	1.7×10^0	2.2×10^0	1.6×10^0	4.5×10^{-1}	9.2×10^{-1}
0-15	41	1.3×10^{-1}	5.6×10^0	1.4×10^0	1.6×10^0	1.3×10^0	1.8×10^{-1}	8.6×10^{-1}
0-25	21	9.4×10^{-2}	3.5×10^0	8.7×10^{-1}	1.0×10^0	9.4×10^{-1}	-4.0×10^{-1}	9.9×10^{-1}
0-40	10	6.5×10^{-2}	9.8×10^{-1}	3.2×10^{-1}	3.8×10^{-1}	2.8×10^{-1}	-1.3×10^0	8.9×10^{-1}
0-60	7	4.4×10^{-2}	7.3×10^{-1}	1.8×10^{-1}	2.5×10^{-1}	2.4×10^{-1}	-1.8×10^0	1.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-19. Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1991 on Arbar Island (F43).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	8	2.3×10^0	1.6×10^1	3.5×10^0	7.0×10^0	5.7×10^0	1.7×10^0	8.2×10^{-1}
05-10	6	3.5×10^{-1}	7.1×10^0	9.5×10^{-1}	2.2×10^0	2.7×10^0	1.8×10^{-1}	1.2×10^0
10-15	2	5.9×10^{-1}	4.7×10^0	2.7×10^0	2.7×10^0	2.9×10^0	5.1×10^{-1}	1.5×10^0
15-25	1	2.9×10^0	2.9×10^0	2.9×10^0	2.9×10^0	0.0×10^0	1.1×10^0	0.0×10^0
25-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	8	2.3×10^0	1.6×10^1	3.5×10^0	7.0×10^0	5.7×10^0	1.7×10^0	8.2×10^{-1}
0-10	6	1.5×10^0	9.8×10^0	2.7×10^0	4.1×10^0	3.4×10^0	1.2×10^0	7.8×10^{-1}
0-15	2	1.4×10^0	3.9×10^0	2.7×10^0	2.7×10^0	1.7×10^0	8.6×10^{-1}	7.0×10^{-1}
0-25	1	3.5×10^0	3.5×10^0	3.5×10^0	3.5×10^0	0.0×10^0	1.3×10^0	0.0×10^0
0-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-20. Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Eniran Island (F46).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	5	1.6×10^0	7.7×10^0	5.0×10^0	4.8×10^0	2.2×10^0	1.4×10^0	5.9×10^{-1}
05-10	2	1.2×10^0	2.4×10^0	1.8×10^0	1.8×10^0	9.0×10^{-1}	5.2×10^{-1}	5.2×10^{-1}
10-15	2	1.9×10^{-1}	1.0×10^0	6.0×10^{-1}	6.0×10^{-1}	5.8×10^{-1}	-8.4×10^{-1}	1.2×10^0
15-25	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
25-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	5	1.6×10^0	7.7×10^0	5.0×10^0	4.8×10^0	2.2×10^0	1.4×10^0	5.9×10^{-1}
0-10	2	2.0×10^0	2.5×10^0	2.3×10^0	2.3×10^0	3.4×10^{-1}	8.2×10^{-1}	1.5×10^{-1}
0-15	2	1.7×10^0	1.7×10^0	1.7×10^0	1.7×10^0	3.2×10^{-2}	5.4×10^{-1}	1.9×10^{-2}
0-25	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-21. Americium-241 radionuclide concentration summary for all soil profiles taken from our most recent trip in 1991 on Tufa Island (F47).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	8	1.8×10^0	9.4×10^0	3.6×10^0	4.5×10^0	2.6×10^0	1.4×10^0	5.6×10^{-1}
05-10	6	7.0×10^{-1}	1.6×10^1	2.1×10^0	5.2×10^0	6.0×10^0	1.1×10^0	1.2×10^0
10-15	4	1.6×10^{-1}	2.3×10^0	1.4×10^0	1.3×10^0	8.8×10^{-1}	-1.1×10^{-1}	1.2×10^0
15-25	4	1.7×10^{-1}	9.4×10^{-1}	2.0×10^{-1}	3.8×10^{-1}	3.8×10^{-1}	-1.3×10^0	8.2×10^{-1}
25-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	8	1.8×10^0	9.4×10^0	3.6×10^0	4.5×10^0	2.6×10^0	1.4×10^0	5.6×10^{-1}
0-10	6	1.9×10^0	9.6×10^0	3.5×10^0	4.7×10^0	3.2×10^0	1.3×10^0	6.8×10^{-1}
0-15	4	1.3×10^0	7.2×10^0	3.6×10^0	3.9×10^0	2.8×10^0	1.1×10^0	8.2×10^{-1}
0-25	3	8.5×10^{-1}	4.7×10^0	3.3×10^0	2.9×10^0	1.9×10^0	8.5×10^{-1}	9.0×10^{-1}
0-40	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Table C-22. Americium-241 radionuclide concentration summary for all soil profiles taken from the 1978 NMIRS together with our most recent trip in 1989 on Borukka Island (F49).

Soil depth (cm)	N ^a	pCi g ⁻¹ dry wt					Mean of logs	SD of logs
		Minimum	Maximum	Median	Mean	SD		
0-05	13	1.1×10^0	1.1×10^1	3.2×10^0	4.5×10^0	2.9×10^0	1.3×10^0	6.2×10^{-1}
05-10	9	1.6×10^{-1}	1.5×10^1	8.8×10^{-1}	2.6×10^0	4.8×10^0	5.3×10^{-2}	1.3×10^0
10-15	8	9.9×10^{-2}	2.3×10^0	6.1×10^{-1}	7.6×10^{-1}	7.4×10^{-1}	-7.6×10^{-1}	1.1×10^0
15-25	9	3.9×10^{-2}	8.8×10^{-1}	2.0×10^{-1}	2.6×10^{-1}	2.5×10^{-1}	-1.7×10^0	9.0×10^{-1}
25-40	5	1.6×10^{-2}	2.9×10^{-1}	3.5×10^{-2}	9.2×10^{-2}	1.2×10^{-1}	-3.0×10^0	1.2×10^0
40-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0
0-05	13	1.1×10^0	1.1×10^1	3.2×10^0	4.5×10^0	2.9×10^0	1.3×10^0	6.2×10^{-1}
0-10	9	1.3×10^0	8.1×10^0	2.2×10^0	3.3×10^0	2.2×10^0	1.0×10^0	6.0×10^{-1}
0-15	6	8.9×10^{-1}	3.4×10^0	1.7×10^0	2.0×10^0	1.0×10^0	5.6×10^{-1}	5.4×10^{-1}
0-25	5	5.9×10^{-1}	2.2×10^0	8.2×10^{-1}	1.3×10^0	7.4×10^{-1}	8.2×10^{-2}	6.0×10^{-1}
0-40	3	3.7×10^{-1}	5.2×10^{-1}	4.8×10^{-1}	4.6×10^{-1}	7.7×10^{-2}	-7.9×10^{-1}	1.8×10^{-1}
0-60	0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0	0.0×10^0

NOTE: Specific activity is decay corrected to 1994.

N^a stands for number of individual samples.

Appendix D

Maps of Islands

Thomas A. Lee
Estimate 100-00-00

Rongelap Atoll
Burck Island F-49