THE THERMAL DECOMPOSITION OF C₂H₅I

by

S. S. Kumaran, M.-C. Su,[‡] K. P. Lim, and J. V. Michael*

Chemistry Division, Argonne National Laboratory, Argonne, IL 60439, USA

Corresponding Author: **Dr. J. V. Michael**
D-193, Bldg 200
Argonne National Laboratory
Argonne, IL 60439
Phone: (708) 252-3171, Fax: (708) 252-4470
E-mail: Michael@ANLCHM.BITNET

(26th Symp. (Int'l) on Combust., Submitted Dec., 1995)

Desired Presentation: Oral
Preferred Publication: Proceedings
Category: React. Kinetics Code Numbers: 1.1, 1.5

	Text	Reference	Tables	Figures	Total
Count	3037 words	37 citations	4 pages	5 pages	
Equiv. Words	3037	747	800	1000	5584

(NOT PATENTABLE)

The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31-109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

[‡]Sabbatical Leave. Permanent address: Department of Chemistry, Butler University, Indianapolis, IN 46208.

This work was supported by the U. S. Department of Energy, Office of Basic Energy Sciences, Division of Chemical Sciences, under Contract No. W-31-109-Eng-38.

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

Abstract

The high temperature thermal dissociation of $\text{C}_2\text{H}_5\text{I}$ has been characterized in this study. Kinetics and overall yield experiments were performed over the temperature range, 946-2046 K, using the atomic resonance absorption spectrometric technique (ARAS) for the temporal detection of both product H- and I-atoms behind reflected shock waves. The $\text{C}_2\text{H}_5\text{I}$ decomposition proceeds by both C-I fission and HI elimination. Rate constants for the C-I fission process, measured over the temperature and density ranges, 946-1303 K and $0.82\text{--}4.4 \times 10^{18} \text{ cm}^{-3}$, respectively, can be well represented to within $\pm 37\%$ by the first-order expression: $k = 6.34 \times 10^9 \exp(-15894 \text{ K}/T) \text{ s}^{-1}$. Overall yield data for atomic product gave a branching ratio for C-I fission of (0.87 ± 0.11) suggesting that 13% of the reaction proceeds through molecular HI elimination. This conclusion is consistent with earlier studies that showed C-I fission to be the dominant dissociation channel.

The temperature and pressure dependences of the dissociation rate constants and the yield data have been theoretically described using three formulations of unimolecular rate theory. The best description was obtained with a full Master's equation analysis. However, all three calculations confirm that the HI elimination pathway is lower lying than the C-I fission process by $\sim 3 \text{ kcal mole}^{-1}$.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Introduction

Thermal chemical reactions often involve multiple reaction pathways. In both unimolecular and bimolecular reactions, the extent of branching into different product channels depends on the nature of the intermediate configurations and the thermochemistry. In unimolecular reactions, this is determined by the reaction thresholds and transition state properties for the respective pathways. There are not many reaction systems that delineate these features even though it has been suggested that understanding the kinetics in pyrolysis and combustion systems requires such detailed descriptions. [1] Before the development of unimolecular rate theory can continue to proceed, some unambiguous experimental observations on multi-channel dissociations are required. The effects of such quantities as the collisional energy transfer parameter and the fall-off behavior (as functions of both temperature and density) on the branching ratios would be particularly informative. Theoretical characterization would then be essential for a predictive application to unstudied systems. This of course is a strong need, for example, in the development of detailed kinetics in combustion systems. [1]

Recent experiments on CH_2Cl_2 [2] and COCl_2 [3] from this laboratory have shown these to be typical two-channel unimolecular dissociations involving both C-Cl bond fission and molecular elimination (HCl or Cl_2 , respectively). In this paper, the thermal decomposition kinetics of $\text{C}_2\text{H}_5\text{I}$ are studied using both H- and I-atom atomic resonance absorption spectrometric (ARAS) techniques. From thermodynamic considerations, the pathways available for thermal dissociation are C-I fission and molecular HI elimination. This should be contrasted with the other haloethanes, $\text{C}_2\text{H}_5\text{F}$, $\text{C}_2\text{H}_5\text{Cl}$, and $\text{C}_2\text{H}_5\text{Br}$, where HX elimination requires ~ 50 kcal mole $^{-1}$ whereas bond fission requires a much larger energy effectively eliminating C-X bond fission as a viable dissociation channel. In $\text{C}_2\text{H}_5\text{I}$, the C-I bond fission channel,

yields C_2H_5 which will undergo fast subsequent decomposition,

since $\Delta H_{0(1a)}^0 = 34.8 \text{ kcal mole}^{-1}$ is much lower than $\Delta H_{0(1)}^0$. The overall process is then $C_2H_5I = C_2H_4 + H + I$. C_2H_5I has in fact been used as a source of H-atoms for gas phase kinetics studies. [4,5]

The molecular elimination channel,

has been theoretically [6,7] shown to involve a 4-center transition state in contrast to the 3-center processes in CH_2Cl_2 and $COCl_2$. Both Herzler and Frank [4] and Mertens et al. [5] have assumed that reaction (2) was negligible. However, using the most recent value [8] for $\Delta_f H_{298K(C_2H_5I)}^0 = -1.72 \text{ kcal mole}^{-1}$ combined with the JANAF [9] recommendations for C_2H_4 and HI gives $\Delta H_{298K(2)}^0 = 20.6 \text{ kcal mole}^{-1}$. With $\Delta_f H_{298K(C_2H_5)}^0 = 28.7 \text{ kcal mole}^{-1}$ from Hanning-Lee et al., [10] $\Delta H_{298K(1)}^0 \cong 56 \text{ kcal mole}^{-1}$; i. e., $\sim 35 \text{ kcal mole}^{-1}$ higher. The reverse 4-center process, HI addition to C_2H_4 , has an estimated [6,11] insertion barrier of $\sim 30 \text{ kcal mole}^{-1}$, suggesting that the threshold energies for reactions (1) and (2) could be nearly isoenergetic.

There are earlier studies that have attempted to characterize the decomposition. R. A. Ogg, Jr. [12] studied the thermal system, HI with C_2H_5I , over small temperature (523-573 K) and pressure (145-177 Torr) ranges. Using calorimetric determination of [I] and with an assumed reaction mechanism, $k_1 = 1.82 \times 10^{13} \exp(-21638 \text{ K/T}) \text{ s}^{-1}$; i. e. reaction (1) was rationalized as being first-order. Reaction (2) was not even considered in this analysis. Sullivan [13] later reinterpreted the results of Ogg [12] and estimated $k_1 = 1.54 \times 10^{16} \exp(-26167 \text{ K/T}) \text{ s}^{-1}$. He further suggested that the C-I bond strength in C_2H_5I was

52.0 kcal mole⁻¹. Butler and Polanyi [14] were the first to postulate that both channels (1) and (2) could be thermally accessible in the dissociation of C₂H₅I. Yang and Conway [15] then studied the C₂H₅I pyrolysis at higher temperatures (660-794 K) using the toluene-carrier flow technique. They measured the rate constant for reaction (2) and concluded that it was first-order over the toluene pressure range, 2-18 Torr, and C₂H₅I pressure range, (0.6-40) x 10⁻³ Torr. They [15] reported $k_2 = 1.27 \times 10^{14} \exp(-26569 \text{ K}/T)$ at 14 Torr total pressure. From the extrapolated value, $k_1 = 4.50 \times 10^{13} \exp(-25160 \text{ K}/T)$, this work suggested that ~70% of the reaction occurs through reaction (1). It is clear that these earlier studies have not fully explored the C₂H₅I pyrolysis; i. e., the temperature and density dependences of the rate constants for (1) and (2).

A thorough study of the C₂H₅I decomposition requires kinetics measurements under chemical isolation conditions in order to unambiguously determine reaction rates and yields. The ARAS method combined with shock tube equipment is well suited for this purpose. The thermal dissociation of C₂H₅I was studied by observing the temporal formation of both H- and I-atoms behind reflected shock waves. The calibration for H- and I-atom ARAS detection techniques are completely independent and are detailed below. Because of the difference in the H-atom and I-atom ARAS detection sensitivities, separate set of experiments were carried out. However, these experiments were performed over nearly the same temperature ranges, 1080-2020 K (H-atom) and 946-2046 K (I-atom).

Experimental

The shock tube ARAS experiments were performed behind reflected shock waves. The experimental approach used here is similar to that described elsewhere. [16,17] Experiments were performed under highly dilute conditions so as to minimize secondary chemistry and thereby obtain chemical isolation.

Apparatus: The shock tube used here has already been described. [16] The 7 m circular tube (i.d. 9.74 cm) was separated from the driver by a thin aluminum diaphragm (4 mil unscored 1100-H18). The tube was routinely pumped to <10⁻⁸ Torr between experiments

by a packaged pump system (Edwards Vacuum Products, model CR100P). Shock waves were produced by manually breaking the diaphragm at a known driver pressure of high-purity He. Boundary layer corrected post reflected shock properties were calculated from incident shock wave velocities that were measured by eight equally spaced pressure transducers (PCB Piezotronics, Inc., model 113A2) located well down stream from the diaphragm. [18-20] Both pressure and absorption data were recorded using a Nicolet 4094C digital storage oscilloscope.

H-atom ARAS Detection: The ARAS technique was used to measure $[H]_t$. [21] The H-lamp was operated under a 2 Torr flow-pressure of UHP He with the microwave discharge power set to 40 W. H_2 and CH_4 impurities present in the He gave sufficient Ly_α signal through an O_2 (1 atm of dry air) gas filter. A discharge flow system containing high-purity H_2 in the optical path provided an atomic filter that was used to establish the fraction of non- Ly_α light. Line reversal was negligible under these conditions. The observed data were the time dependent transmittances, and from these, $[H]_t$ profiles were obtained from routine Ly_α line absorption calculations using the known absorption oscillator strength for a Doppler-broadened source at an equivalent lamp temperature of 480 K. [22,23] The Ly_α oscillator strength is *a priori* better known than those for any other spectral transition. Absolute $[H]$ determination with this ARAS detection method in reflected shock waves has already been experimentally validated. [21,24]

I-atom ARAS Detection: The ARAS technique for I-atom detection has been recently described. [25] The I-atom resonance lamp configuration (2 Torr flow of 0.1% CF_3I in UHP He, at 50 W microwave power) is similar to the Cl-atom lamp detailed elsewhere.²⁴ Lamp emission was observed with a solar blind EMR G14 PMT (quantum efficiency, ~3%) through a suprasil filter over a wavelength range of 165-185 nm without further spectral resolution. The absorption spectrum of I-atoms in this region corresponds to the transitions, $^2P_{3/2}$ and $^4P_{5/2,1/2} \leftarrow ^2P_{3/2}^0$ (principally $\lambda = 183.0$ nm). [26] A purge of dry N_2 between the PMT and the shock tube window eliminated O_2 from the beam path.

I-atom calibration using CF_3I as the thermal source of $[\text{I}]$ has been described. [25] In that study, varying CF_3I -Kr mixtures were shock heated at sufficiently high temperatures to completely dissociate CF_3I . The long time transmittance values correspond unambiguously to an I-atom concentration, $[\text{I}]_\infty$, equal to $[\text{CF}_3\text{I}]_0$. However, construction of an absorbance, $(\text{ABS})_\infty \equiv -\ln(I_\infty / I_0)$, against $[\text{I}]_\infty$ relationship (the curve-of-growth) requires a prior knowledge of the fraction of light that is non-resonant. This fraction was determined to be (0.49 ± 0.02) . [25] Long-time absorbance values, $(\text{ABS})_\infty$, corresponding to $[\text{I}]_\infty = [\text{CF}_3\text{I}]_0$, were calculated from the known non-resonant fraction of light, and the curve-of-growth was constructed. The curve-of-growth is linear for $[\text{I}] \leq 3 \times 10^{12} \text{ cm}^{-3}$ being best represented by the linear Beer's law expression,

$$(\text{ABS}) = 1.9215 \times 10^{-13} [\text{I}] \quad (3)$$

where $[\text{I}]$ is given in molecules cm^{-3} . The I-atom absorption cross-section is $\sigma = 1.933 \times 10^{-14} \text{ cm}^2$, and this is in good agreement with earlier determinations. [27,28] At higher concentrations the curve becomes increasingly non-linear.

Gases: High-purity He (99.995%) from Air Products and Chemicals was used as the driver gas. Kr (Scientific Grade, 99.997%) used as the diluent in the experimental mixtures was obtained from MG Industries. UHP Grade He (99.995%) from Airco was used as the diluent in the I-atom lamp mixture and for the H-atom resonance lamp. Also obtained from Airco was the high-purity H_2 (99.995%) for the atomic filter. CF_3I (99%) for the I-atom resonance lamp and $\text{C}_2\text{H}_5\text{I}$ (99%) were obtained from Aldrich Chemical Co. Inc. Reactant $\text{C}_2\text{H}_5\text{I}$ (99%) was further purified by bulb-to-bulb distillation with the middle third being retained for mixture preparation. All gas mixtures were stored in a glass vacuum line, and an MKS Baratron capacitance manometer was used for pressure measurements.

Results

In this study, two sets of shock tube ARAS experiments were carried out, and rate constants and yields were measured for both. Because reaction (1a) is much faster than (1) and secondary reactions are negligible at the $[\text{C}_2\text{H}_5\text{I}]_0$ used, all observed $[\text{H}]$ or $[\text{I}]$ is a result of reaction (1); i. e., $[\text{H}]_\infty$ or $[\text{I}]_\infty$ should unambiguously correspond to the yield from the C-I fission channel. This is confirmed by iterative kinetics calculations using the mechanism in Table I. Note that H-atom abstraction of either H or I from $\text{C}_2\text{H}_5\text{I}$ is not included in the mechanism because these rate constants are negligible and do not contribute to either $[\text{H}]_t$ or $[\text{I}]_t$ with the present dilute conditions. [30]

38 H-atom ARAS experiments at $T > 1220\text{K}$ were used to determine $[\text{H}]$ yields. From the transmittance value $T_\infty \equiv (I_\infty / I_0)$ (and corresponding absorbance value, $(\text{ABS})_\infty \equiv -\ln(T_\infty)$), $[\text{H}]_\infty$ was obtained through line absorption calculations [21-24] at the temperature of a given experiment. With the Table 1 mechanism, observed $[\text{H}]_\infty$ can be recovered by using a branching ratio, $k_1/(k_1+k_2)$, that is identical to the observed yield defined as $[\text{H}]_\infty/[\text{C}_2\text{H}_5\text{I}]_0$, showing that reaction (3) in Table I is negligible. The H-atom yield values are plotted in Fig. 1 as solid circles which give the least-squares average, $[\text{H}]_\infty/[\text{C}_2\text{H}_5\text{I}]_0 = (0.85 \pm 0.07)$, implying $k_2/k_1 = 0.18$. It is obvious that the molecular elimination channel, reaction (2), does have significance. Similar experiments were performed by observing I-atoms with the ARAS technique. $[\text{I}]_\infty$ was measured in 40 yield experiments over the temperature range 950-2050 K with the ARAS technique. [25] These experiments differ from the H-atom experiments in two ways: (i) the I-atom product from reaction (1) is directly observed and does not depend on the occurrence of reaction (1a), and; (ii) the $[\text{I}]$ detection range (i. e., $(1-8) \times 10^{12}$ molecule cm^{-3}) is ~ 10 times higher than that for $[\text{H}]$. The yield results are also displayed in Fig. 1 as open circles. This set gives $[\text{I}]_\infty/[\text{C}_2\text{H}_5\text{I}]_0 = (0.89 \pm 0.07)$, implying $k_2/k_1 = 0.12$, in agreement with the H-atom results. If the two sets are combined, the T-independent value, (0.87 ± 0.11) is obtained giving $k_2/k_1 \approx 0.15$.

Four H-atom rate constant measurements were made at $T < 1230$ K where the temporal formation of H-atoms was fully time resolved. The top panel of Fig. 2 shows a typical experimental record, and these transmittance data were converted to $[H]_t$ giving the bottom panel. With a fixed k_2/k_1 set by $[H]_\infty$, k_1 was adjusted to best fit the $[H]_t$ profile using the full Table 1 mechanism. This is illustrated in Fig. 2 (bottom panel) where the best fit to $[H]_t$ is shown as the line. Derived first-order rate constants for reaction (1) are given in Table 2 along with the corresponding second-order values. More extensive rate constant measurements were made with the I-atom ARAS detection method. A typical record is shown in Fig. 3. Using eqn. (3), transmittance data were converted to $[I]_t$ giving the profile displayed in the bottom panel. As before, $[I]_t$ was fitted with the Table 1 mechanism by adjusting k_1 and the ratio, k_2/k_1 . Even with somewhat higher $[C_2H_5I]_0$, the $[I]$ values are still not perturbed by abstraction (reaction (3) in Table 1). The solid line in the bottom panel of Fig. 3 illustrates the quality of fits used for deriving the first- and second-order rate constants listed in Table 2.

The results in Table 2 are plotted in Fig. 4 as first-order rate constants simply because we anticipate that this decomposition may be in the fall-off region. The linear least squares line shown in the figure is given by,

$$k_1 = 6.34 \times 10^9 \exp(-15894 \text{ K}/T) \text{ s}^{-1}, \quad (4)$$

for the T-range, 946-1303 K. The points shown in the figure are within $\pm 37\%$ of the line at the one standard deviation level.

Discussion

Since the threshold energies reactions (1) and (2) are within a few kcal mole⁻¹ of one another, the C_2H_5I decomposition might be a nearly degenerate two channel reaction system. Yang and Conway [15] showed that (2) is not negligible. However, the

subsequent experiments at higher temperature substantiate that (1) must dominate. [4,5] These conclusions are corroborated in the present work.

In order to fully characterize this decomposition, we have performed three types of unimolecular rate theory calculations that are similar to those in our earlier studies. [31-32] The first method uses the semi-empirical Troe formulation to rationalize the results while the second is a full RRKM calculation. [33] Both formulations are single channel type calculations and therefore disregard the effects of population depletion by the lower energy channel on the dissociation rate for the higher energy channel. The third method is also a full RRKM calculation (Program *UNIMOL* from Gilbert et al. [34]); however, it utilizes a full weak collision Master's equation analysis for the energy transfer processes. Here the mutual interferences of population effects for both channels are included in order to determine the total rate of depletion and the branching fractions into the product channels.

The models used and the results from the respective calculations are given in Table 3. All three methods conclude that the threshold energy for C-I fission is (52.0 ± 0.5) with molecular elimination being lower by ~ 3 kcal mole⁻¹. This suggests $\Delta_f H_{0K(C_2H_5I)}^0 = 5.0$ kcal mole⁻¹; i. e., 2.8 kcal mole⁻¹ higher than the Carson et al. [8] value. This new value and $E_{0(2)} = 48.9$ kcal mole⁻¹ suggests a potential energy barrier of 32.5 kcal mole⁻¹ for HI insertion into C₂H₄.

Excluding Lennard-Jones Kr-C₂H₅I collision parameters (calculated by methods outlined by Cambi et al. [37], implying $\sigma_{12} = 4.071$ Å and $\epsilon_{12} = 284.3$ K), the first two models in Table 3 involve adjusting 4 parameters, $E_{0(1)}$, $E_{0(2)}$, $\langle \Delta E_1 \rangle_{down}$, and $\langle \Delta E_2 \rangle_{down}$ in order to give good agreement for both the pressure dependent rate constants and the experimental yield, 0.87. A T-independent yield could be conserved (Fig. 1) if $\langle \Delta E_2 \rangle_{down}$ was substantially less than $\langle \Delta E_1 \rangle_{down}$ (see Table 3). However, the *UNIMOL* formulation successfully agreed with the data using a common energy transfer parameter, 410 cm⁻¹. The calculated values for the yields do however show a slight T-dependence as seen by the dashed line in Fig. 1. Clearly this slight variation is well within the data scatter. In all three

calculations, the predictions show unambiguously that both reactions (1) and (2) are strongly affected by pressure fall-off. This is undoubtedly due to the large number of degrees of freedom and, in particular, the values for the low lying frequencies in C_2H_5I . This justifies presenting the data as first-order in Fig. 4.

A comparison between the predicted density dependences from the *UNIMOL* calculation and experiment is shown in Fig. 5 where the present data in Table 2 have been plotted as second-order. The quality of fits with the first two methods is similar to that shown with this third method. Extrapolation of the third model to the conditions of Mertens et al. [5] predicts rate constants that are in excellent agreement with experiment. Lastly, our calculations show that the low temperature data of Yang and Conway [15] are not near the high pressure limit for reaction (2) as they had presupposed. Hence, ambiguities exist as to what collisional energy transfer parameter to use when toluene is the third body. With a high β_c , we predict a yield of ~ 0.8 to be compared to their rough estimate of 0.7, and the predicted apparent activation energy is ~ 51 whereas they report $52.8 \text{ kcal mole}^{-1}$.

Even though the single channel models can probably be used to assess the threshold energies for the two processes, rationalization of the data requires substantially different values for the two $\langle \Delta E \rangle_{down}$ values. We agree with Just [1] that weak collisional energy transfer for multi-channel dissociations should be considered with a unified model. In this instance, Master's equation analysis (Program *UNIMOL*) is clearly appropriate, and, as noted earlier, its application requires one less parameter to rationalize the data.

Acknowledgments

We thank B. Kirtman for providing us with a vibrational analysis for the HI elimination transition state. This work was supported by the U. S. Department of Energy, Office of Basic Energy Sciences, Division of Chemical Sciences, under Contract No. W-31-109-Eng-38.

References

- 1 Just, Th., *Twenty-Fifth Symposium (International) on Combustion*, The Combustion Institute, Pittsburgh, 1994, pp. 687-703.
- 2 Lim, K. P., and Michael, J. V., *Twenty-Fifth Symposium (International) on Combustion*, The Combustion Institute, Pittsburgh, 1994, pp. 809-816.
- 3 Lim, K. P., and Michael, J. V., *J. Phys. Chem.* 98: 211-215 (1994).
- 4 Herzler, J., and Frank, P., *Ber. Bunsenges. Phys. Chem.* 96: 1333-1338 (1992).
- 5 Mertens, J. D., Wooldridge, M. S., and Hanson, R. K., *Eastern States Section 1994 Technical Meeting, The Combustion Institute*, Pittsburgh, PA, paper presentation.
- 6 Toto, J. L., Pritchard, G. O., and Kirtman, B., *J. Phys. Chem.* 98: 8359-8370 (1994).
- 7 Wolfe, S., and Kim, C. K., *Israel J. Chem.* 33: 295-305 (1993).
- 8 Carson, A. S., Laye, P. G., Pedley, J. B., Welsby, A. M., Chickos, J. S., and Hosseini, S., *J. Chem. Thermodynamics* 26: 1103-1109 (1994).
- 9 Chase, M. W., Jr., Davies, C. A., Downey, J. R., Jr., Frurip, D. J., McDonald, R. A., and Syverud, A. N., *J. Phys. Chem. Ref. Data* 14: Supplement No. 1 (1985).
- 10 Hanning-Lee, M. A., Green, N. J. B., Pilling, M. J., and Robertson, S. H., *J. Phys. Chem.* 97: 860-870 (1993).
- 11 Benson, S. W., and O'Neal, H. E., *Kinetic Data on Gas Phase Unimolecular Reactions*, National Stand. Ref. Data Ser., U.S., Nat'l. Bur. Stand., Washington, D. C., 1970.
- 12 Ogg, R. A., Jr., *J. Am. Chem. Soc.* 56: 526-536 (1934).
- 13 Sullivan, J. H., *J. Phys. Chem.* 65: 722-727 (1961).
- 14 Butler, E. T., and Polanyi, M., *Trans. Faraday Soc.* 39: 19-36 (1943).
- 15 Yang, J.-H., and Conway, D. C., *J. Chem. Phys.* 43: 1296-1303 (1965).
- 16 Michael, J. V., *Prog. Energy Combust. Sci.* 18: 327-347 (1992).

-
- 17 Michael, J. V., in *Advances in Chemical Kinetics and Dynamics* (J. R. Barker, Ed.), JAI, Greenwich, 1992, Vol. 1, p. 47, for original references.
 - 18 Michael, J. V., and Sutherland, J. W., *Int. J. Chem. Kinet.* 18: 409-436 (1986).
 - 19 Michael, J. V., *J. Chem. Phys.* 90: 189-197 (1989).
 - 20 Michael, J. V., and Fisher, J. R., in *Seventeenth International Symposium on Shock Waves and Shock Tubes*, AIP Conference Proceedings 208 (Y. W. Kim, Ed.), American Institute of Physics, New York, 1990, p. 210.
 - 21 Lim, K. P., and Michael, J. V., *Twenty-Fifth Symposium (International) on Combustion*, The Combustion Institute, Pittsburgh, 1994, pp. 713-719, and references cited therein.
 - 22 Lynch, K. P., Schwab, T. C., and Michael, J. V., *Int. J. Chem. Kinet.* 8: 651-671 (1976).
 - 23 Maki, R. G., Michael, J. V., and Sutherland, J. W., *J. Phys. Chem.* 89: 4815-4821 (1985).
 - 24 Kumaran, S. S., Lim, K. P., and Michael, J. V., *J. Chem. Phys.* 101: 9487-9498 (1994).
 - 25 Kumaran, S. S., Su, M.-C., Lim, K. P., and Michael, J. V., *Chem. Phys. Lett.* 243: 59-63 (1995).
 - 26 Radzig, A. A., and Smirnov, B. M., *Reference Data on Atoms, Molecules, and Ions*, Springer Ser. Chem. Phys., Springer-Verlag, Heidelberg, 1985, Vol. 39.
 - 27 Takahashi, A., Inoue, A., and Inomata, T., *Twentieth International Symposium on Shock Waves*, Pasadena, CA, 1995, paper presentation.
 - 28 Braun-Unkoff, M., Frank, P., and Just, Th., *Ber. Bunsenges. Phys. Chem.* 94: 1417-1425 (1990).
 - 29 Baulch, D. L., Duxbury, J., Grant, S. J., and Montague, D. C., *J. Phys. Chem. Ref. Data* 10: Supplement No. 1, 1-1 (1981).

-
- 30 Westley, F., Frizzel, D. H., Herron, J. T., Hampson, R. F., and Mallard, W. G., *NIST Chemical Kinetics Database - ver. 5.0*, NIST Standard Reference Database 17, Gaithersburg, MD (1993).
- 31 Kumaran, S. S., Lim, K. P., Michael, J. V., and Wagner, A. F., *J. Phys. Chem.* 99: 8673-8680 (1995).
- 32 Kumaran, S. S., Su, M.-C., Lim, K. P., Michael, J. V., and Wagner, A. F., *J. Phys. Chem.*, in press.
- 33 Troe, J., *J. Chem. Phys.* 66: 4745-4757 (1977); *ibid.* 66: 4758-4775 (1977); Troe, J., *J. Phys. Chem.* 83: 114-??? (1979); Troe, J., *Ber. Bunsenges. Phys. Chem.* 87: 161-169 (1983); Gilbert, R. G., Luther, K., and Troe, J., *Ber. Bunsenges. Phys. Chem.* 87: 169-177 (1983).
- 34 Gilbert, R. G., Simth, S. C., and Jordan, M. J. T., *Program package UNIMOL program suite* (1993), School of Chemistry, Sydney University, NSW 2006, Australia, Gilbert_R@summer.chem.su.oz.au.
- 34 Herzberg, G., *Electronic Spectra of Polyatomic Molecules*, Van Nostrand, Princeton, 1966, p. 647.
- 36 Feng, Y., Niiranen, J. T., Bencsura, A., Knyazev, V. D., Gutman, D., and Tsang, W., *J. Phys. Chem.* 97: 871-880 (1993).
- 37 Cambi, R., Cappelletti, D., Liuti, G., and Pirani, F., *J. Chem. Phys.* 95: 1852-1861 (1991).

Table 1: Mechanism for C₂H₅I Pyrolysis.

1.	C ₂ H ₅ I + M → C ₂ H ₅ + I + M	k ₁ = to be fitted
1a.	C ₂ H ₅ → C ₂ H ₄ + H	k _{1a} = ∞
2.	C ₂ H ₅ I + M → C ₂ H ₄ + HI + M	k ₂ = to be fitted
3.	H + HI → H ₂ + I	k ₃ = 7.87 × 10 ⁻¹¹ exp(-330 K/T) ^a

^aRef. 29, units are cm³ molecule⁻¹ s⁻¹.

Table 2: Kinetics Data for the Thermal Dissociation of C₂H₅I.

P ₁ /(Torr)	^a M _s	^b p ₅ /(10 ¹⁸ cm ⁻³)	^c k _{1st} /(s ⁻¹)	^b T ₅ /(K)	^c k ₁ /(cm ³ molecule ⁻¹ s ⁻¹)
H-atom Results:					
X _{C₂H₅I} = 6.273×10 ⁻⁷					
8.41	2.197	1.410	9500	1230	6.74×10 ⁻¹⁵
X _{C₂H₅I} = 2.087×10 ⁻⁷					
15.91	2.054	2.464	1800	1100	7.30×10 ⁻¹⁶
15.89	2.029	2.416	1350	1080	5.59×10 ⁻¹⁶
15.86	2.177	2.633	12000	1217	4.56×10 ⁻¹⁵
I-atom Results:					
X _{C₂H₅I} = 1.247×10 ⁻⁶					
10.99	1.892	1.507	275	946	1.82×10 ⁻¹⁶
11.00	2.124	1.764	7443	1161	4.22×10 ⁻¹⁵
13.53	2.061	2.105	2665	1106	1.27×10 ⁻¹⁵
16.03	2.073	2.512	2290	1117	9.12×10 ⁻¹⁶
20.15	2.092	3.118	5300	1118	1.70×10 ⁻¹⁵
30.90	1.963	4.403	650	1000	1.48×10 ⁻¹⁶
10.98	2.113	1.749	9500	1150	5.43×10 ⁻¹⁵
10.95	1.974	1.595	825	1019	5.17×10 ⁻¹⁶
16.03	2.055	2.485	3200	1101	1.29×10 ⁻¹⁵
X _{C₂H₅I} = 9.976×10 ⁻⁷					
10.63	2.053	1.637	3800	1089	2.32×10 ⁻¹⁵
10.96	1.957	1.583	1300	1000	8.21×10 ⁻¹⁶
10.96	2.223	1.864	17000	1256	9.12×10 ⁻¹⁵
10.97	2.055	1.692	3900	1091	2.30×10 ⁻¹⁵
X _{C₂H₅I} = 2.199×10 ⁻⁶					
5.97	2.195	0.996	18000	1232	1.81×10 ⁻¹⁴
5.97	2.080	0.936	9000	1115	9.62×10 ⁻¹⁵
5.95	2.244	1.023	26240	1278	2.56×10 ⁻¹⁴
5.94	2.155	0.973	7500	1188	7.71×10 ⁻¹⁵
5.96	2.135	0.965	9000	1168	9.32×10 ⁻¹⁵

Table 2 (cont.)

$P_1/(\text{Torr})$	aM_s	$b\rho_5/(10^{18}\text{cm}^{-3})$	$k_{1st}/(\text{s}^{-1})$	$bT_5/(\text{K})$	$c k_1/(\text{cm}^3\text{molecule}^{-1}\text{s}^{-1})$
5.94	2.268	1.034	27000	1303	2.61×10^{-14}
5.95	2.043	0.911	3400	1080	3.73×10^{-15}
5.96	1.899	0.825	220	948	2.67×10^{-16}
5.88	1.976	0.858	2100	1021	2.45×10^{-15}
$X_{\text{C}_2\text{H}_5\text{I}} = 9.976 \times 10^{-7}$					
15.91	2.168	2.638	13250	1204	5.02×10^{-15}
15.91	2.067	2.492	4950	1108	1.99×10^{-15}
15.90	1.889	2.212	250	948	1.13×10^{-16}
15.95	1.959	2.324	1000	1013	4.30×10^{-16}
15.94	2.003	2.400	1800	1049	7.50×10^{-16}
15.99	2.041	2.457	3090	1087	1.26×10^{-15}
15.99	2.034	2.455	2825	1077	1.15×10^{-15}
15.86	2.112	2.541	7822	1154	3.08×10^{-15}
15.90	2.171	2.632	13300	1212	5.05×10^{-15}
15.87	1.970	2.331	1290	1023	5.53×10^{-16}

^aThe error in measuring the Mach number, M_s , is typically 0.5-1.0% at the one standard deviation level. ^bQuantities with subscript 5 refer to the thermodynamic state of the gas in the reflected shock region. ^cThe rate constants are derived as described in the text.

Table 3: Theoretical Calculations for the Dissociation Reaction: $C_2H_5I (+ M) \rightarrow$ Prods.

Species	Frequencies (cm ⁻¹)	Moments of Inertia (amu Å ²)	Source			
C ₂ H ₅ I	^a 3026, 2987(2), 2956, 2931, 1454(3), 1384, 1217(2), 1038, 939(2), 751, 520, 264	17.4, 169.7, 180.7, 3.20 (reduced int. rotor)	est. from ref. 35			
C ₂ H ₅	3112, 3033, 2987, 2920, 2842, 1440(3), 1366, 1186, 1175, 1138, 802, 540	5.1, 22.5, 24.2 1.12 (reduced int. rotor)	ref. 36			
C ₂ H ₄ -HI [‡]	^b 169, 211, 318, 598, 802, 1065, 1135, 1191, 1244, 1257, 1363, 1423, 1515, 2925, 2978, 3014, 3086	21.9, 263.4, 278.6	ref. 6			
Theoretical Fits:						
		<i>E</i> ₀ /kcal mole ⁻¹	⟨Δ <i>E</i> ⟩ _{down} /cm ⁻¹			
(1) C-I fission						
Troe semi-empirical		51.5	380			
RRKM		52.3	445			
Unimol		52.3	410			
(2) HI elimination						
Troe semi-empirical		48.0	119			
RRKM		50.0	182			
Unimol		48.7	410			
Theoretical Results (900-1400 K):						
^c log $\left(\frac{A_0}{\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}}\right)$		<i>n</i> ₀	<i>T</i> ₀	log $\left(\frac{A_\infty}{\text{s}^{-1}}\right)$	<i>T</i> _∞	
^d (1) C-I fission						
Troe semi-empirical		45.519	15.2	33799	15.210	25490
RRKM		44.985	15.0	33621	14.921	25986
UNIMOL		53.602	17.5	36599	14.899	25937
^e (2) HI elimination						
Troe semi-empirical		43.810	15.1	31615	14.450	25867
RRKM		42.284	14.5	32058	14.480	26960
UNIMOL		59.485	19.5	35793	14.461	26360

^aThese frequencies are in good agreement with our scaled STO-3G RHF-SCF *ab initio* calculations. ^bThese frequencies are from B. Kirtman and are based on Gaussian 94 *ab initio* calculations presented in ref. 6. In accordance with usual practice, they are scaled by 0.9. ^c $k_i = A_i T^{-n_i} \exp(-T_i/T)$ three parameter fits for limiting low- and high-pressure rate constants. ^dThe derived values of k_{-1}^∞ for each of the models from top to bottom are: 7.7, 5.5, and $(5.5 \pm 0.2) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. ^eThe derived values of k_{-2}^∞ for each of the models from top to bottom are: $2.78 \times 10^{-11} \exp(-18219 \text{ K}/T)$, $2.79 \times 10^{-11} \exp(-18978 \text{ K}/T)$, and $3.11 \times 10^{-11} \exp(-18533 \text{ K}/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

Figure Captions

- Figure 1: Plot of yield data, ● – H-atoms and ○ – I-atoms, for reaction (1) against T^{-1} for $948 \leq T \leq 2046$ K. The combined sets give the yield = 0.87 (solid line). The dashed line is the *UNIMOL* prediction (see text).
- Figure 2: The top panel shows a typical experimental H-atom ARAS signal, and the bottom panel shows the corresponding $[H]_t$ obtained from line absorption calculations (see text). The conditions are: $[C_2H_5I]_0 = 8.84 \times 10^{11} \text{ cm}^{-3}$ at $T = 1230$ K and $\rho = 1.410 \times 10^{18} \text{ cm}^{-3}$. The solid line is the calculated fit using the mechanism of Table 1 with $[H]_\infty = 0.71 [C_2H_5I]_0$ and $k_1 = 9500 \text{ s}^{-1}$.
- Figure 3: The top panel shows a typical experimental I-atom ARAS signal, and the bottom panel shows the corresponding $[I]_t$ obtained using eqn. (3). The conditions are: $[C_2H_5I]_0 = 2.62 \times 10^{12} \text{ cm}^{-3}$ at $T = 1106$ K and $\rho = 2.105 \times 10^{18} \text{ cm}^{-3}$. The solid line is the calculated fit using the mechanism of Table 1 with $[I]_\infty = 0.83 [C_2H_5I]_0$ and $k_1 = 2665 \text{ s}^{-1}$.
- Figure 4: Arrhenius plot of measured first-order rate constants for reaction (1), using H- and I-atom ARAS techniques. The solid line represents a linear-least-squares fit given by eqn. (4).
- Figure 5: Arrhenius plots of measured second-order rate constants for reaction (1) (Table 2): (▲) 6 Torr, (■) 11 Torr, and (●) 16 Torr loading pressure. The solid lines are *UNIMOL* predictions using the Table 3 model (see text).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5