MAY 1996

STRUCTURAL BIOLOGY RESEARCH

**AT THE
NATIONAL SYNCHROTRON LIGHT SOURCE**

BROOKHAVEN NATIONAL LABORATORY

**Associated Universities, Inc.
Upton, NY 11973-5000
Under Contract No. DE-AC02-76CH00016**

NATIONAL SYNCHROTRON LIGHT SOURCE

The world's largest facility for scientific research using x-rays and ultraviolet and infrared radiation is operated by the National Synchrotron Light Source Department. This year alone, a total of about 2,200 researchers from 360 institutions performed experiments at the world's largest source of synchrotron radiation. Researchers often work in collaboration with the Laboratory's staff scientists at the Light Source, conducting a wide range of innovative experiments in physics, chemistry, biology, materials science and various technologies.

CREATING A MAP OF LIFE

By defining the three-dimensional structures of biological macromolecules, scientists can create a map of life, a guide for exploring the biological and chemical interactions of the vast variety of molecules found in living organisms.

"Solving" the structure of a molecule—determining the exact location of each of its atoms—provides information on its function. Studies in structural biology may lead to new insights into how biological systems are formed and nourished, how they survive and grow, how they are damaged and die.

EXPLORING THE MOLECULES OF LIFE

At the National Synchrotron Light Source (NSLS), about 600 structural biologists from more than 100 institutions use an x-ray technique called protein crystallography at six dedicated beam lines to explore the molecules of life. They decipher the structures of proteins, complex combinations of amino acids that are an essential ingredient of every living cell.

Meanwhile, at numerous other beam lines, biologists use x-ray and ultraviolet light as probes to study proteins and other

biological structures, such as DNA, virus particles and carbohydrates, for obtaining data that is not accessible via protein crystallography.

To accommodate this large number of users, a new \$1.8 million structural biology wing was added to the NSLS during fiscal year 1995. The 6,300-square-foot addition contains eight new laboratories for preparation and storage of samples, important amenities for our many users.

A GROWING FIELD

The first crystallographic structure of a protein—that of myoglobin, which supplies oxygen to muscles—was solved in 1958 with the use of conventional laboratory x-rays. Since then, the field of structural biology has grown tremendously—and today, it is estimated that more than 100 structures are solved each month by scientists throughout the world.

Within the last decade, the development of synchrotron light sources and powerful computers has made solving macromolecular structures easier and faster. Also, the rapid growth of genetic-engineering technology makes it possible for scientists to make large amounts of synthetic proteins for this research. Previously, researchers were limited to studying proteins in the minute quantities obtained from natural sources.

SYNCHROTRON LIGHT: MULTIPLE BENEFITS

The NSLS provides x-rays that are 10,000 times more intense than those from a conventional x-ray source. This intensity results in higher resolution of data and images, which provides more detail, and allows scientists to study extremely small structures.

Another advantage of using a synchrotron source for this research is that the wavelength of the light can be fine-tuned to meet a scientist's needs. Thus, a three-dimensional structure can be determined from a single crystal, rather than from many crystals, which are hard to grow. Also, data collection at the NSLS is up to 50 times faster than it is using conventional light sources. With synchrotron x-rays, an entire series of data can be taken within a millisecond, while a protein crystal is in action, providing important information on its function.

FACETS OF DIFFRACTION

Biological molecules are studied using three basic protein crystallography methods at the NSLS: multiple isomorphous replacement (MIR), multiple-wavelength anomalous diffraction (MAD) and Laue diffraction. All involve x-ray diffraction, in which x-rays hit a protein crystal and scatter off its atoms in a pattern recorded by

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

detectors. The pattern relates to the placement of each atom in the molecule and can be deciphered by a computer.

While MIR involves using several crystals at a fixed wavelength, MAD requires only one crystal, which is exposed to many wavelengths of synchrotron light. Laue diffraction, which uses the full white synchrotron radiation spectrum, gives information on the dynamics of protein structure—how a crystal chemically changes over a short time-scale.

DESIGNING DRUGS AND ENZYMES

Many pharmaceutical companies do research at the NSLS to develop new drugs

for treating numerous diseases, from the common cold to AIDS.

In this “structure-based” drug design, researchers use protein crystallography techniques to design inhibitors, molecules that attach to enzymes in disease-causing agents and block them. Knowing the precise structure of a molecule and its inhibitor can help researchers develop more effective drugs that also produce fewer unwanted side effects.

Other companies use the NSLS to engineer enzymes for industrial purposes. For example, some chemical companies are looking for an enzyme that would be suitable for making nylon by environmentally benign fermentation processes, rather than

by using petroleum, which creates pollution.

A UNIQUE RESOURCE

New data on proteins and the other molecules of life discovered at the NSLS can be instantaneously shared with researchers, teachers and students who have access to the global Internet computer network.

Structural biologists working at the NSLS, along with researchers from all over the world, add to and obtain information from Brookhaven’s Protein Data Bank (PDB), an archive for three-dimensional structures of proteins, nucleic acids and other biological macromolecules. The PDB

Malcolm Capel (left), BNL, and Xiaodong Cheng, Cold Spring Harbor Laboratory, prepare to mount a crystal for data collection at beam line X12B.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MB

is a unique resource for the scientific community, as well as for others who wish to tap into its wealth of information.

Established in 1971, the PDB has grown considerably. Today, a user consults it almost every minute through the Internet, while others receive updated compact disks periodically. The number of structures stored has grown exponentially in recent years, with more than 100 now being added each month to the more than 4,000 already in the PDB. ■

STRUCTURAL BIOLOGY AT THE NSLS — PROGRESS AND PROMISE

From the toxin that causes cholera to DNA's proteins, the numerous and varied structural biology experiments conducted at the National Synchrotron Light Source (NSLS) help scientists understand the biological principles that govern life's complexity.

The following are a few examples of the rich diversity of research at the NSLS beam lines dedicated solely to protein crystallography, a technique in which x-ray probes are used to study the structure of a protein in its crystalline form. Proteins, made of any of numerous combinations of amino acids, are components of all living cells. Knowing a protein's structure can elucidate its function.

PROTEIN CRYSTALLOGRAPHY: A SAMPLING

BEAM LINE X4A

Using a technique called multiwavelength anomalous diffraction (MAD), researchers from the Howard Hughes Medical Institute at Columbia University and Texas A&M University have determined the structure of the tyrosine kinase domain, or region, of the human insulin receptor.

A receptor is a site on a cell's surface that combines with a hormone like insulin, or another biological substance, to produce a change in cell function. Tyrosine kinase is just one of hundreds of enzymes that help govern hormones.

The tyrosine kinase of this study is related to type II diabetes, which does not respond to insulin. In nondiabetics, the body activates the insulin receptor when the blood's sugar level is high. The receptor binds to insulin, signaling the tyrosine kinase to go to work and set off a series of biochemical changes that cause blood glucose to be converted to glycogen, which holds energy in reserve. In type II diabetics, however, the insulin receptor molecule is defective in its ability to sense the presence of insulin, and the extra blood sugar cannot be converted.

Structural details about tyrosine kinase may provide important clues on how to prevent type II diabetes. Further, since it is the first tyrosine kinase for which the three-dimensional structure has been determined, it is a crucial first step in understanding cellular signals and metabolic functioning.

BEAM LINE X8C

Using the technique known as isomorphous heavy atom replacement, Argonne National Laboratory researchers have solved the structure of the toxin that causes cholera, an infectious gastrointestinal disease that began in Asia and has spread to South America and Mexico.

The soluble toxin that the Argonne team investigated is deposited in the intestinal

Tyrosine kinase unit from the human insulin receptor. The areas numbered Y1158, Y1162, and Y1163 represent the enzyme's activation sites.

Cholera toxin. The darker, bottom part and the long chain extending to the top part are the parts of the protein that bind to the lining of the intestine. The top section is the part that invades the interior of intestine cells and stimulates them to secrete water, bringing on cholera's symptoms.

tract of cholera disease victims by the bacterium *Vibrio cholerae*. The toxin stimulates cells to secrete water, causing profuse diarrhea, then dehydration and, finally, collapse.

Knowing the structure of the toxin, researchers can now probe the chemistry of the disease process in order to develop a more effective vaccine. The one available now does not produce complete immunity against cholera.

BEAM LINE X12B

Several years ago, researchers from Brookhaven and Cold Spring Harbor Laboratory, working at X12C, determined the structure of the bacteriophage T7's lysozyme. A bacteriophage is a virus that infects bacteria, and the T7 strain has been commonly used by Brookhaven biologists

as a simple genetic model system, since it contains a single piece of DNA and only about 50 genes.

This year, because of the higher beam intensity and advanced instrumentation at X12B, a much higher-resolution structure for the lysozyme enzyme was achieved by Cold Spring Harbor scientists.

T7 lysozyme combines two functions in one small protein: It cuts open the bacterial cell wall, and it inhibits DNA transcription, or reading, by the T7 enzyme RNA polymerase. Transcription creates a messenger RNA molecule from a template DNA molecule, so that genetic information is transferred to messenger RNA and proteins can be made from those instructions.

Each lysozyme molecule appears to be able to perform only one of these functions at a time. Learning the structure of this

protein may provide new basic insights into the processes of gene regulation and DNA replication.

BEAM LINE X12C

Using x-ray diffraction techniques on this beam line, researchers from Brookhaven's Biology Department, Los Alamos National Laboratory, the Max Planck Institute for Medical Research and Rice University recently determined the structure of the oxygen-binding protein myoglobin in a transient state.

Myoglobin stores oxygen in the body's muscles and tissues. It is the first protein for which a crystal structure was determined, in 1958. Protein crystallographers have since studied this molecule extensively, compiling data on its structure, dynamics and function.

Phenyl ring (top center ring) of the amino acid phenylalanine-91 of the T7 lysozyme protein.

Three states of carbon monoxide (CO) bound in myoglobin. The grey depicts the tight complex between the two, in which the dumbbell-shaped CO molecule is bound to an iron atom. White represents myoglobin with no CO bound to it, and black is the transient state, trapped at 20 kelvins, with the CO molecule released but lingering nearby.

To simulate the action of an oxygen molecule as it binds to myoglobin, the researchers substituted the poisonous gas carbon monoxide (CO). Although CO binds more strongly than oxygen, the myoglobin molecule will release CO when it is illuminated with visible light. To determine the structure of the myoglobin-gas complex, the scientists bound carbon monoxide to myoglobin molecules in a crystal, then froze the crystal to near absolute zero, and finally, released the CO with a bright light.

The structure they found revealed the carbon monoxide molecule resting near the iron atom to which it was bound. This finding now can be compared to myoglobin's structure when CO is absent or tightly bound. The researchers observed details of the subtle shape change that accompanies binding and release of gas molecules, and their results provide crucial data on the dynamics of how a simple protein works.

BEAM LINE X25

Researchers from Rockefeller University and the Howard Hughes Medical Institute have determined the structure of a protein known as the TATA-box binding protein, or TBP, using isomorphous crystallography.

TBP is the DNA-binding subunit of the RNA polymerase enzyme. It positions the polymerase on the DNA to initiate gene transcription (described in the section on X12B). When TBP attaches itself to a strand of DNA, the DNA bends into a V-shape, thus helping to position the enzyme.

This study helped researchers investigate how TBP recognizes various DNA sequences and directs transcription from them. Examining TBP's structure may help researchers understand how individual genes are turned on or off in a cell in order to make specific proteins. Using this information, they may learn how to design pharmaceuticals to block or stimulate the production of these proteins, thus curtailing genetic diseases.

BEAM LINE X26C

Owned by the University of Chicago, X26C is the primary beam line for using

the white-beam Laue method, in which numerous x-ray diffraction intensities can be measured from a single crystal. This method enables researchers to take "molecular movies"—images of proteins as they change their structures during a chemical reaction.

A collaboration from Brookhaven, the Fred Hutchinson Cancer Research Center in Washington State, the University of California, Berkeley, and the Molecular Research Council in England used the Laue technique to solve the structures of two intermediate chemicals formed during the catalytic actions of the enzyme isocitrate dehydrogenase (IDH). These intermediate chemicals are involved in a two-step process that is part of ordinary respiration: oxidation and expulsion of a carbon dioxide molecule.

In the two-step reaction studied, researchers observed the transformation of isocitrate, a direct product of sugar metabolism, to oxalosuccinate, an intermediate that had never before been observed. They also recorded the conversion of oxalosuccinate to α -ketoglutarate. Each of these enzyme intermediates is catalyzed, or chemically changed, by a different portion of IDH. These data provide insight into fundamental metabolic processes.

In an effort to obtain the best protein crystals for data collection, researchers at X26C use x-ray topography techniques to find defects in protein crystals—in much the way that physicians use x-rays to find broken bones. A collaboration from Brookhaven, the University of Alabama at Huntsville and the University of Manchester in England applied x-ray topography to

protein crystallography for the first time to optimize crystal quality for experiments.

Crystals must be as perfect as possible in protein crystallography studies in order to give scientists the most precise structural data. Protein crystals were grown aboard a 1994 space shuttle mission and compared with those grown on earth. Using x-ray topography at X26C, the researchers found that protein crystals grown under constant temperature or in microgravity conditions found in space had fewer defects than those grown at slightly fluctuating temperatures or normal gravity conditions.

DIVERSE BIOLOGICAL RESEARCH

Complementing the protein crystallography research described above are vari-

ous structural biology experiments performed on noncrystalline materials. Here is a small sampling of these experiments:

BEAM LINE X1

Using the leading scanning x-ray microscope in the world, a collaborative team from the State University of New York at Stony Brook and Lawrence Livermore National Laboratory has mapped the distribution of DNA and protein in the human sperm nucleus. They have found that certain sperm structures, such as the segment that makes the first contact with an egg during fertilization, are particularly rich in protein. The next stage of the project is aimed at understanding the structural basis of certain types of male infertility.

BEAM LINE X9B

Researchers from the Albert Einstein College of Medicine and the University of Pennsylvania have used x-ray diffraction techniques on beam line X9B to analyze the structure of membrane proteins. The researchers have developed a novel method of attaching proteins to the surface of inorganic substrates in order to collect diffraction data. This creates a one-dimensional arrangement of the protein-membrane complexes, which is ideal for structural and functional studies.

BEAM LINE X11A

Researchers from North Carolina State University used the technique known as x-ray absorption spectroscopy to follow structural changes in ferritin, a protein that stores iron for use in hemoglobin and other iron-containing proteins. The technique allows researchers to look at how ferritin works in a solution that mimics the conditions in the human body, which is not possible using protein crystallography techniques. This basic research will help reveal how iron is used in the body and may be the pathway for designing better iron supplements.

BEAM LINES U9 AND U11

Structural biology experiments using ultraviolet light at these beam lines include two techniques: circular dichroism, which provides information on the structure of proteins, DNA, RNA and polysaccharides; and time-resolved fluorescence, which supplies data on molecular structure, photochemical reactions and interactions in nucleic acids and proteins.

For example, using time-resolved fluorescence, Brookhaven researchers are studying how the nucleic acid bases adenine and guanine respond to ultraviolet radiation. Such studies help us to understand the mechanisms by which the ultraviolet components of sunlight damage DNA and induce mutations and cancer. ■

TATA-box binding protein bound with a 14 base-pair DNA sequence. The unusual saddle-shaped protein induces a conformational change in DNA.

Cover Figure:

View of a specific complex between the transcription factor NF- κ B (p50 subunit) and DNA. The p50 protein (365 amino-acid residues in the form shown here) binds to DNA as a dimer. Its polypeptide chain (represented as a ribbon) folds into two domains (colored yellow and red). The amino-terminal domain (yellow) contains most of the segments that interact specifically with DNA; the carboxy-terminal domain (red) mediates dimerization. The DNA (blue) contains eleven base pairs. The molecular structure was determined by X-ray diffraction from crystals of the complex using NSLS beamlines X25 and X4A. Their adjustable wavelength monochromators and the very high intensity available at the X25 wiggler station were both essential for the success of the project. Collaborating institutions: X4A - Howard Hughes Medical Institute; X25 - Howard Hughes Medical Institute; the Department of Molecular and Cellular Biology and the Department of Chemistry, Harvard University. (Reference: C.W. Müller, F.A. Rey, M. Sodeoka, G.L. Verdine and S.C. Harrison. *Structure of the NF- κ B-p50 Homodimer Bound to DNA*. Nature, Vol. 373: 311-317; 26 January, 1995.)