CONF-961051-1

TITLE: A QUANTITATIVE TEST OF DIFFERENT MAGNETIC FIELD
MODELS USING CONJUNCTIONS BETWEEN DMSP AND
GEOSYNCHRONOUS ORBIT

AUTHOR(S): G. D. Reeves
L. A. Weiss
M. F. Thomsen
D. J. McComas

RECEIVED
MAR 13 1996
OSTI

SUBMITTED TO: Workshop on Radiation Belts
Brussels, Belgium
October 17-20, 1996

Los Alamos
NATIONAL LABORATORY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

A Quantitative Test of Different Magnetic Field Models Using Conjunctions Between DMSP and Geosynchronous Orbit

G. D. Reeves, L. A. Weiss, M. F. Thomsen, and D. J. McComas

Los Alamos National Laboratory, Mail Stop D436, Los Alamos, NM 87545, USA, reeves@lanl.gov

Abstract. We report here on a study which tests the magnetic field line mapping between geosynchronous orbit and the ionosphere. The mapping is determined both observationally and from five magnetospheric magnetic field models. The mapping is tested observationally by comparing electron energy spectra obtained by the Magnetospheric Plasma Analyzer (MPA) at geosynchronous orbit and by the DMSP spacecraft. Because the orbits are nearly perpendicular, in general, the spectra match well for only a few seconds providing a good determination of when DMSP crosses the geosynchronous drift shell. In this way the mapping between geosynchronous orbit and the ionosphere can be determined to better than one degree. We then compare the measured magnetic footpoints of geosynchronous orbit with the footpoints predicted by five magnetospheric field models: Tsyganenko-89, Tsyganenko-87, Tsyganenko-82, Oslen-Pfizer, and Hilmer-Voigt. Based on a set of over 100 measured magnetic conjunctions we find that, in general, there are significant differences between the mappings predicted by various magnetic field models but that there is no clear “winner” in predicting the observed mapping. We find that the range of magnetic latitudes at which we measure conjunctions is much broader than the range of latitudes which the models can accommodate. This lack of range is common to all magnetic field models tested. Although there are certainly cases where the models are not sufficiently stretched, we find that on average all magnetic field models tested are too stretched. This technique provides an excellent opportunity for testing future magnetic field models and for determining the appropriate parameterizations for those models.

Introduction

A crucial aspect of the modeling of the Earth’s radiation belts is modeling the Earth’s magnetic field. This study uses an observational criteria for determining the mapping between three low-altitude DMSP spacecraft and high-altitude geosynchronous spacecraft. It also compares the results of the measured mapping with the mapping predicted by several commonly used magnetic field models.

The current generation of empirical magnetic field models (including those of Tsyganenko) are statistical fits to single point measurements of the magnetic field measured by spacecraft in the magnetosphere. The models have commonly been tested by comparing model magnetic field vectors to the magnetic field measured by one or more spacecraft in the magnetosphere [e.g. *Tsyganenko*, 1989; *Fairfield*, 1991; *Peredo and Stern*, 1991; *Peredo et al.*, 1993; *Pulkkinen et al.*, 1994; *Thomsen et al.*, 1996]. However, one of the most common uses for magnetic field models is to trace the magnetic field lines to determine the magnetic connectivity between

two points in space or between a spacecraft and the ground and the *mapping* predicted by these empirical models has not been systematically tested. Tests of the magnetic field mapping are, in some ways, more sensitive because they integrate along the entire field line.

We apply this test of the field models to geosynchronous orbit for several reasons. Firstly, geosynchronous orbit is an inherently interesting region of space. It lies in the transition region between dipole-like and tail-like magnetic field lines and near the edge of the trapping region for energetic particles. It is sensitive to the effects of substorms and is a source region for the injection of particles into the ring current and radiation belts. Geosynchronous orbit is also heavily populated by satellites. The effects of the radiation environment on the operation of those satellites is a fundamentally important application of magnetospheric physics. Finally, there is a large and continuous database of plasma and energetic particle measurements from a series of Los Alamos instruments on geosynchronous satellites which make a large, systematic, and statistical study possible.

Finding Magnetic Conjunctions

The technique we use to establish magnetic conjunction between the low-altitude DMSP spacecraft and the geosynchronous satellites is to compare electron energy spectra on the two satellites as a function of time and to look for times when the spectra are very nearly identical. DMSP orbits at an altitude of approximately 850 km in a nearly polar, circular orbit with a period of approximately 90 minutes. Therefore DMSP crosses the geosynchronous L-shell approximately once every 23 minutes. The DMSP orbits are also sun-synchronous and therefore each DMSP satellite samples a nearly fixed region of local time.

The geosynchronous satellites orbit at a geocentric distance of $6.6 R_E$ with a period of 24 hours and therefore must pass through the local times sampled by the DMSP satellites. In this sense the orbits are perpendicular to each other and for each geosynchronous-DMSP satellite pair there are numerous possible conjunctions each day. To further increase the statistics we use data from two geosynchronous satellites (1989-046 and 1990-095) and from three DMSP satellites (DMSP F8, F9, and F10).

We define a "nominal conjunction" as a time when one DMSP and one geosynchronous satellite are within $\pm 10^\circ$ in magnetic longitude and when DMSP is between 50° and 80° magnetic latitude. During a nominal conjunction we examine the electron energy spectra from the SSJ/4 instrument on DMSP which measures precipitating electrons in 20 energy bins from 30 eV to 30 keV [Hardy *et al.*, 1984]. One complete energy spectrum is obtained every 1 second which provides excellent resolution in latitude. At the same time we examine electron energy spectra from the Magnetospheric Plasma Analyzer (MPA) instrument at geosynchronous orbit. The MPA is a spherical-sector electrostatic analyzer which measures electrons from about 1 eV to 40 keV [Bame *et al.*, 1993]. The MPA measures in a fan of 6 look directions and takes 24 azimuthal sectors in each 10-second spin of the spacecraft. The spacecraft spin axis points toward the center of the earth so excellent pitch angle coverage is obtained.

In comparing spectra it is important that several conditions be met. First, since DMSP measures only that portion of the distribution that is in the loss cone we use only the geosynchronous MPA spectrum that is most nearly field aligned (as described by Weiss *et al.* [1996] and Thomsen *et al.* [1996]). Second, our technique assumes that the magnetic field is unchanging in the time it takes DMSP to cross the geosynchronous field line so we limit analysis to times when the MPA spectrum is constant for several minutes. This criteria also assures that there are not strong variations in local time in the vicinity of the geosynchronous spacecraft. Finally, we also eliminate times when there is a field-aligned potential drop. This occurs naturally as a result of our spectral comparison because if DMSP is measuring an

Figure 1. An illustration of how good spectral matches are chosen. The top panel shows the RMS difference between the MPA and DMSP spectra as a function of time. The second panel shows MPA and DMSP spectra that met our matching criteria for this event. The bottom panel shows several DMSP spectra in this interval that did not match the MPA spectrum. The times of those spectra are shown with open circles in the top panel.

accelerated population and MPA is not then the spectra will not match. We assure this by adopting an extremely strict condition for spectral matching.

Figure 1 shows a typical spectral match and the criteria used to define it. In the top panel we plot the RMS difference between the spectrum measured by DMSP and by MPA for two minutes of a nominal conjunction. We define a "measured conjunction" as the times at which the RMS difference in the spectra falls below a threshold of 0.36. By examining a large number of spec-

tra we determined that this threshold is actually more selective than a visual examination of the spectra. Furthermore we require that a measured conjunction last for less than 30 seconds. This assures that the longitude range over which the spectra match is sufficiently small that we can distinguish between one magnetic field model and another. Finally, we also require that the RMS error does not fall below a second threshold of 0.42 for more than 60 seconds which assures that the minimum in the RMS is sharp and well-defined.

In this case our "measured conjunction" (as defined by the criteria above) lasted for 6 seconds. The middle panel in Figure 1 shows the MPA spectrum and DMSP spectra that satisfied our matching criteria. (We note that the energy spectra use the nominal calibrations of the instruments and are not normalized in any way.) In the bottom panel of Figure 1 we plot the same MPA spectrum along with 5 other DMSP spectra. The times of those spectra are marked with open circles on the RMS plot in the top panel. Clearly the spectral match for those times is significantly worse. We also point out that this is only a 2-minute portion of the DMSP crossing which occupies a very small portion of a typical DMSP spectrogram plot. Outside this 2-minute interval the DMSP spectrum did not even resemble the MPA spectrum which is what one would expect since during those times DMSP is mapping to very different parts of the magnetosphere.

This type of two-point spectral comparison has been used before to establish magnetic conjugacy [Sharp *et al.*, 1971; Mende and Shelly, 1976; Meng *et al.*, 1979; Lundin and Evans, 1985; Schumaker *et al.*, 1989; Mauk and Meng, 1991; Hones *et al.*, 1996]. What distinguishes this study from those earlier studies is that we have established an automated algorithm for identifying magnetic conjunctions and we have applied it to a set of satellites that have frequent conjunctions. This allows us to study the magnetic mapping from geosynchronous orbit to the ionosphere in a statistical manner.

Measured Conjunction Statistics

Using the technique described in the previous section we examined three months of data (March, September, and December, 1991) for nominal conjunctions between one of three DMSP satellites (F8, F9, and F10) and one of the two geosynchronous satellites (1989-045 and 1990-095). Out of over 3,500 nominal conjunctions we found 102 that satisfied our selection criteria. For each of the 102 conjunctions we identified the times of all spectra that met our criteria as well the single 1-second DMSP spectrum that best matched the MPA spectrum (as defined by the minimum RMS error).

Figure 2 shows the location of DMSP at the times when the best matched spectra were observed. Figure 2a shows DMSP's geographic location and Figure 2b shows its location in magnetic local time and magnetic latitude. Because we are using geosynchronous satel-

lites we are limited to the geographic longitudes those satellites sample. 1989-046 had footpoints near Alaska. 1990-095 has two clusters of footpoints – over western and central Russia – because it was moved in the middle of 1991. Likewise, because the DMSP satellites are sun-synchronous they sample only a limited range of local time. But, thanks to the rotation of the earth's dipole DMSP is able to sample about one half of the possible magnetic local times (as shown in Figure 2b).

It is apparent from Figure 2 that the footpoint of geosynchronous orbit generally lies in the auroral ionosphere. Most often it is in the region of diffuse aurora but it frequently lies in the region of discrete aurora. It is also apparent from the figure that the footpoint of geosynchronous orbit can be quite variable, spreading over more than 10° in magnetic latitude. In a related study Weiss *et al.*, [1996] investigate how well the measured location of the geosynchronous footpoint correlates with various magnetospheric indices such as Kp, AE, Dst, the local tilt of the field at geosynchronous orbit, and the equatorward edge of the auroral boundary. In this paper our emphasis is on evaluating how well various magnetic field models predict the location of the measured footpoint. An important point about Figure 2 is that no magnetic field models have been used to determine the magnetic footpoints of geosynchronous orbit. Therefore we have a completely model-independent data set of field line mappings which we can use to evaluate magnetospheric magnetic field models.

Comparison with Magnetic Field Models

We now compare the magnetic field mapping determined from the DMSP and MPA spectra with the magnetic field mapping predicted by various magnetic field models. The models we use are the Tsyganenko-89 [Tsyganenko, 1989], Tsyganenko-87 [Tsyganenko, 1987], Tsyganenko-82 [Tsyganenko and Usmanov, 1982], Olsen-Pfitzer [Olsen and Pfitzer, 1974], and Hilmer-Voigt [Hilmer and Voigt, 1995] magnetic field models. For each of these models we use the IGRF representation of the earth's internal field. This is very important for mapping studies because, as shown by Reeves and Weiss [1996], the deviation of the earth's field from a dipole can have a significant effect on the magnetic mapping from geosynchronous orbit to low altitudes.

Figure 3 illustrates the method we used to compare the footpoints predicted by the field models with the measured footpoint determined from DMSP and MPA spectra. This event is the same event shown in Figure 1. It was a conjunction between the DMSP F9 satellite and the geosynchronous satellite 1989-046. The best spectral match was recorded by DMSP at 08:44:30 UT when DMSP was at -59.78° magnetic latitude. DMSP was in the southern hemisphere moving poleward and during the 6-seconds of spectral match it moved only 0.4° in magnetic latitude.

An expanded plot of region of the conjunction is

Figure 2. The location of DMSP when it measured a magnetic conjunctions with a geosynchronous satellite. (A) The position of DMSP in geographic coordinates. Only northern hemisphere conjunctions are shown. (B) The position of DMSP in geomagnetic coordinates (magnetic latitude and magnetic local time). Conjunctions from both hemispheres are shown.

shown in the inset. Here we have plotted the location of the footpoint measured by DMSP with a square and the location of the footpoints predicted by the Tsyganenko-89 magnetic field model with circles. In both cases the footpoint is defined at an altitude of 100 km. The Tsyganenko-89 field model comes in five versions for integral values of the magnetic activity parameter K_p . For this event $K_p=3^-$ and the Tsyganenko-89 model predicted a footpoint which was 0.6° further poleward than the actual measured footpoint which is excellent agreement.

We calculated the difference in magnetic latitude of the footpoint for each of the five models used in this study and for each of the 102 conjunctions in our data set. We used each model "as advertised". In other words, for the Tsyganenko family of field models we used the actual K_p parameter for each conjunction to specify what version of the model to use. The Hilmer-Voigt model is specified by three parameters: Dst, the stand-off distance of the magnetopause (given by solar wind pressure), and the equatorward boundary of the auroral oval (given by DMSP electron precipitation signatures). For the Hilmer-Voigt model we again used the parameters that were appropriate for each event. The Olsen-Pfitzer model has no free parameters so the same model applies to all of our cases.

A histogram of the difference between the measured and model footpoints for each of the five models is plotted in Figure 4. The top panel shows the statistics for the Tsyganenko-89 model. Here, 32% of the model footpoints agreed with the measured footpoints to within

$\pm 1^\circ$, 65% were within $\pm 3^\circ$, and 83% were within $\pm 5^\circ$. Put another way, if you need to know the location of the footpoint of geosynchronous orbit to within 1° the Tsyganenko-89 model has a 32% probability of being correct. However, it also has a 17% chance of being off by more than 5° and the statistical uncertainty in the mapping is approximately 3° .

A fairly surprising result of this study is that, on average, all the field models tested perform about equally well – or equally poorly. The three generations of Tsyganenko magnetic field models contain various refinements and improvements but the changes in model did not improve the accuracy of the mapping from geosynchronous orbit to the ionosphere. This is in part because the measured footpoints have a much larger range of latitudes than is accommodated in the models. (Compare Figures 2 and 3.) However, the Tsyganenko models have a larger range than the Olsen-Pfitzer model (which has none) yet the Olsen-Pfitzer model does as good a job of predicting the location of the geosynchronous footpoint as any of the Tsyganenko models. This is no doubt in part because K_p is a poor parameter to use for determining the amount of stretching in the field. In fact *Hones et al.*, [1996] and *Weiss et al.* [1996] have shown that the K_p value needed for the Tsyganenko-89 model to reproduce the observations is completely uncorrelated with the actual K_p measured for an event. It may, then, be somewhat surprising that the Hilmer-Voigt model, which uses a set of parameters which might be expected to be better correlated with the measured footpoint of geosynchronous orbit, still does not perform significantly

Figure 3. Comparing the measured footpoint with the footpoint predicted by the Tsyganenko-89 model. The globe shows the location of DMSP when it measured a good spectral match with MPA. The inset shows the magnetic latitude and longitude of the measured footpoint with a bar for the location of DMSP during the entire spectral match and a square for the location of the best spectral match. The footpoints predicted by various Kp levels of the Tsyganenko-89 model are shown with circles. The actual Kp was 3⁻ which gives a difference for this event of only 0.6°.

better than its rivals.

Finally we note that the distributions are also not symmetric. The difference in magnetic latitude is calculated such that, regardless of what hemisphere DMSP was in, negative values represent cases where the measured footpoint was poleward of the model footpoint and positive values represent cases where the measured footpoint was equatorward of the model footpoint (for example the conjunction shown in Figure 3). The more stretched the field model is the further equatorward the model footpoint moves. (See Figure 3.) Therefore negative values mean that the model is too stretched compared to the observations and positive values mean that the model is not sufficiently stretched.

It is apparent from Figure 4 that all the models are, on average, too stretched. Our cases include a variety of types of magnetospheric conditions which include quiet times, growth phases, expansion phases, and recovery phases and a variety of activity levels from Kp=0 to Kp=9⁻. We note, however, that we have very few conjunctions in the midnight local time sector due to the limited range of local times sampled by the DMSP orbits. Therefore these results do not imply that the field models are too stretched compared to a growth phase field at midnight. Rather we suspect that in order to better represent the conditions at midnight the modelers may have made the models too stretched at other local times.

Figure 4. Histograms of the difference between the measured and model footpoints. All models share the same strengths and weaknesses as described in the text.

Conclusions

We have compiled a set of 102 conjunctions between one of three low-altitude DMSP satellites and one of two geosynchronous satellite using an algorithm that compares the field-aligned electron energy spectra measured at each location. Excellent spectral matches can be found for a subset of nominal conjunctions. Those conjunctions that meet our spectral matching criteria provide a sensitive and field model-independent determination of the magnetic footpoint of geosynchronous orbit.

We compared the measured footpoint obtained by the spectral matching technique with the footpoint predicted by the Tsyganenko-89, Tsyganenko-87, Tsyganenko-82, Olsen-Pfitzer, and Hilmer-Voigt field models. Surprisingly, we found that no one field model performed significantly better than any of the other models. The statistical uncertainty in the footpoint predicted by all of the field models tested was approximately $\pm 3^\circ$. Only about 25-30% of the time did the field model predict the conjunction to within $\pm 1^\circ$ and as much as 20% of the time the field model could be off by more than $\pm 5^\circ$.

We found that the footpoint of geosynchronous orbit varies over more than 10° of magnetic latitude. This is a larger range of latitudes than any of the field models tested can accommodate. This suggests that the next generation of magnetic field models should allow a greater range in the amount of stretching that they allow. However, we also found that all the field models were, on average, too stretched compared to the measured footpoints therefore adding more stretching to the models will only aggravate the problem.

This database of conjunctions is useful not only for evaluating the models that we have used here but can also be used to evaluate any magnetospheric magnetic field model. The database can be extended to include a larger number of cases and a larger number of seasons or magnetospheric conditions as needed. By using other low-altitude satellites it may be possible to get broader coverage in local time and by using other high-altitude satellites, such as CRRES or POLAR or CLUSTER it may be possible to systematically test other ranges of L. In addition to testing existing field models we are also using this database to help determine what magnetospheric parameters actually control the mapping between geosynchronous orbit and the ionosphere and therefore what measurements might be most useful for parameterizing future magnetospheric magnetic field models.

Acknowledgements. We are grateful to P. T. Newell for providing the DMSP data and to the US Department of Energy Office of Basic Energy Science for financial support.

References

- Bame, S. J., D. J. McComas, M. F. Thomsen, B. L. Barraclough, R. C. Elphic, J. P. Glore, J. T. Gosling, J. C. Chavez, E. P. Evans, and F. J. Wymer, Magnetospheric plasma analyzer for spacecraft with constrained resources, *Rev. Sci. Instrum.*, **64**, 1026, 1993.
- Fairfield, D. H., An evaluation of the Tsyganenko magnetic field model, *J. Geophys. Res.*, **96**, 1481, 1991.
- Hardy, D. A., L. K. Schmitt, M. S. Gussenhoven, F. J. Marshall, H. C. Yeh, T. L. Schumaker, A. Huber, and J. Pantazis, Precipitating electron and ion detectors SSJ/4 for the block 5D/flights 6-10 DMSP satellites: Calibration and data presentation, *Rep. AFGL-TR-84-0317*, Air Force Geophys. Lab., Hanscom AFB, Bedford Mass., 1984.
- Hilmer, R. V., and G.-H. Voigt, A magnetospheric magnetic field model with flexible current systems driven by independent physical parameters, *J. Geophys. Res.*, **100**, 5613, 1995.
- Hones, E. W., M. F. Thomsen, G. D. Reeves, L. A. Weiss, D. J. McComas, and P. T. Newell, Observational determination of magnetic connectivity of the geosynchronous region of the magnetosphere to the auroral oval, *J. Geophys. Res.*, *in press*, 1996.
- Lundin R., and D. S. Evans, Boundary layer plasmas as a source for high latitude, early afternoon auroral arcs, *Planet. Space Sci.*, **32**, 1389, 1985.
- Mauk, B. H., and C.-I. Meng, The aurora and middle atmospheric process, in *Auroral Physics*, edited by C.-I. Meng, M. J. Rycroft, and L. A. Frank, Cambridge Univ. Press, NY, 223, 1991.
- Mende, S. B., and E. G. Shelly, Coordinated ATS 5 electron flux and simultaneous auroral observations, *J. Geophys. Res.*, **81**, 97, 1976.
- Meng, C.-I., B. Mauk, and C. E. McIlwain, Electron precipitation of evening diffuse aurora and its conjugate electron fluxes near the magnetic equator, *J. Geophys. Res.*, **84**, 2545, 1979.
- Olsen, W. P., and K. A. Pfitzer, A quantitative model of the magnetospheric magnetic field, *J. Geophys. Res.*, **79**, 3739, 1974.
- Peredo, M. D., D. P. Stern, and N. A. Tsyganenko, Are existing magnetospheric models excessively stretched?, *J. Geophys. Res.*, **98**, 15,343, 1993.
- Peredo, M. D., and D. P. Stern, On the Position of the Near-Earth Neutral Sheet: A Comparison of Magnetic Model Predictions With Empirical Formulas, *J. Geophys. Res.*, **96**, 19,521, 1991.
- Pulkkinen, T. I., D. N. Baker, D. G. Mitchell, R. L. McPherson, C. Y. Huang, and L. A. Frank, Thin current sheets in the magnetotail during substorms: CDAW-6 revisited, *J. Geophys. Res.*, **99**, 5793, 1994.
- Reeves, G. D., and L. A. Weiss, Seven magnetic field models compared: Mapping from geosynchronous orbit to the auroral zone, *J. Geophys. Res.*, *submitted*, 1996.
- Schumaker, T. L., M. S. Gussenhoven, D. A. Hardy, and R. L. Carovillano, The relationship between diffuse auroral and plasma sheet electron distributions near local midnight, *J. Geophys. Res.*, **94**, 10,061, 1989.
- Sharp, R. D., D. L. Carr, R. G. Johnson, and E. G. Shelly, Coordinated auroral-electron observations from a synchronous and a polar satellite, *J. Geophys. Res.*, **76**, 7669, 1971.
- Thomsen, M. F., D. J. McComas, G. D. Reeves, and L. A. Weiss, An observational test of the Tsyganenko (T89a) model of the magnetospheric field, *J. Geophys. Res.*, *submitted*, 1996.
- Tsyganenko, N. A., A magnetospheric magnetic field model with a warped tail current sheet, *Planet. Space Sci.*, **37**, 5, 1989.
- Tsyganenko, N. A., Global quantitative models of the geomagnetic field in the cislunar magnetosphere for different disturbance levels, *Planet. Space Sci.*, **35**, 1347, 1987.
- Tsyganenko, N. A., and A. V. Usmanov, Determination of the magnetospheric current system parameters and development of experimental geomagnetic field models based on data from IMP and HEOS satellites, *Planet. Space Sci.*, **30**, 985, 1982.
- Weiss, L. A., M. F. Thomsen, G. D. Reeves, and D. J. McComas, An examination of the Tsyganenko (T89a) field model using a database of two-satellite magnetic conjunctions, *J. Geophys. Res.*, *submitted*, 1996.

D. J. McComas, G. D. Reeves, M. F. Thomsen, and L. A. Weiss, Los Alamos National Laboratory, Mail Stop D436, Los Alamos, NM 87545, USA, reeves@lanl.gov