CONF-9605149--2

ANL/CHM/CP-89998

Poster - "Ultrafast X" Optical
Society Mtg., San Diego, Calif.
May 28 - June 1, 1996

Factoring the Contribution of Through-Space and Through-Bond Interactions to Rates of

Photoinduced Electron Transfer in Donor-Spacer-Acceptor Molecules using Ultrafast

Transient Absorption Spectroscopy

David Gosztola¹, Bing Wang¹, Michael R. Wasielewski^{1,2}

¹Chemistry Division, Argonne National Laboratory, Argonne, IL 60439-4831, (708)252-6963

² Department of Chemistry, Northwestern University, Evanston, IL 60208-3113

Contributions from through-space and through-bond interactions to the electronic coupling matrix elements for photoinduced charge separation and recombination in linked donor-spacer-acceptor molecules were studied.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

The submitted manuscript has been authored by a contractor of the U.S. Government under contract No. W-31-109-ENG-38. Accordingly, the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Dlc

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

Factoring the Contribution of Through-Space and Through-Bond Interactions to Rates of Photoinduced Electron Transfer in Donor-Spacer-Acceptor Molecules using Ultrafast Transient Absorption Spectroscopy

David Gosztola¹, Bing Wang¹, Michael R. Wasielewski^{1,2}

¹Chemistry Division, Argonne National Laboratory, Argonne, IL 60439-4831, (708)252-6963

² Department of Chemistry, Northwestern University, Evanston, IL 60208-3113

It is well known that the rate of intramolecular electron transfer in linked donor-acceptor molecules is not only dependent upon the distance and orientation between the donor (D) and acceptor (A), but is also dependent upon the electronic properties of the intervening spacer (S) molecules between the redox centers[1]. If the distance between D and A is small enough to allow for direct overlap between the frontier orbitals of the donor and acceptor, electron transfer may occur by means of a through-space mechanism. When the D and A centers are too far apart for direct orbital overlap to be important, electron transfer may occur via a superexchange or through bond interaction. Both contributions to the total electronic coupling matrix element for electron transfer between the donor and acceptor depend on the distance and the orientation of the various molecular components. Thus, it is important to hold both the distance and geometry of the donor and acceptor fixed as the properties of the covalent spacer between them are varied. Achieving this degree of structural control in a D-S-A molecule is often a difficult task.

We present here a study of a series of linked D-S-A molecules. The molecules consisted of a 4-piperidinylnaphthalene-1,8-dicarboximide (ANI) electron donor and a N-(n-octyl)pyromellitimide (PI) electron acceptor attached to the 1,5- and 1,8-positions of either anthracene (ANC) or

dibenzobicyclo(2.2.2)octatriene

(DBO) spacers. For the 1,8-disubstituted

compounds (**5A** and **6A**), ANI and PI are held approximately cofacial with a center-to-center distance of 5.3 Å, whereas for the 1,5-disubstituted compounds (**5B** and **6B**), the center-to-center distance increases to 13.5 Å. These spacers position the ANI donor and PI acceptor at fixed distances relative to one another, provide the desired rigidity, and strongly limit through-bond electron transfer.

Fluorescence quantum yields (Φ_f), charge separation lifetimes (τ_{cs}) and charge recombination lifetimes (τ_{cr}) determined in toluene and THF are listed in the table below for the various compounds (compound **7** is simply the ANI chromophore with no electron acceptor). The fluorescence arises from the excited CT state of the ANI chromophore, which has a fluorescence quantum yield of 0.91 in toluene[2]. The fluorescence emission from **5a** and **6a** is strongly quenched in both toluene and THF, whereas **5b** and **6b** exhibit very little quenching in either solvent. The lifetimes were measured by monitoring the formation and decay kinetics of PI^- at 715 nm following 412 nm, 140 fs laser excitation. Figure 1 shows the transient absorption spectra for the four molecules 20 ps after excitation. The 715 nm PI^- band is present only in the top spectra proving that only in compounds **5a** and **6a** is electron transfer competitive with spontaneous decay of the ANI excited state.

A detailed analysis of the electronic coupling matrix element, V , using the data from the table as well as electrochemically determined redox potentials and reference compounds shows that charge separation in **5a** and **6a** in both toluene and THF occurs via a through-space

Cmpd	Solvent	Φ_f	τ_{cs} (ps)	τ_{cr} (ps)
7	Tol	0.91	--	8500
5a	Tol	<0.001	2.4	70, 250
5b	Tol	0.90	>8500	--
6a	Tol	.007	2.4	450
6b	Tol	0.90	>8500	--
7	THF	0.72	--	8500
5a	THF	<0.001	1.6	39
5b	THF	0.70	>8500	--
6a	THF	<0.001	1.8	43
6b	THF	0.70	>8500	--

mechanism[3]. Charge recombination in **5a** in toluene and **6a** in both toluene and THF is shown to also occur via a through-space mechanism. Charge recombination in **5a** in toluene however occurs via a mixture of through-space and through-bond mechanisms. The through-bond mechanism is thought to involve the intermediate ANI-ANC⁺-PI⁻.

Figure 1: Top: Transient absorption spectra of **5a**(—) and **6a**(-----) in toluene. Bottom: Transient absorption spectra of **5b**(—) and **6b**(-----) in toluene. All spectra were recorded 20 ps after excitation with a 140 fs pulse at 412 nm.

References

1. M. R. Wasielewski, *Chem. Rev.*, 92 (1992) 435-461. D. Gust, T. A. Moore, and A. L. Moore, *Acc. Chem. Res.*, 26 (1993) 198-205.
2. S. R. Greenfield, W. A. Svec, D. Gosztola, and M. R. Wasielewski, *J. Am. Chem. Soc.*, submitted (1995)
3. D. Gosztola, B. Wang, and M. R. Wasielewski, *J. Photochem. Photobiol.*, submitted (1995)

Acknowledgments

This work was supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, United States Department of Energy under contract No. W-31-109-Eng-38.