Final Report on LDRD Program

New XAFS Spectroscopic Investigations in the 1-2keV Region

Joe Wong, M. Fröba and E. Tamura
Lawrence Livermore National Laboratory

RECEIVED

JUL 08 1996

OSTI

March, 1996

This is an informal report intended primarily for internal or limited external distribution. The opinions and conclusions stated are those of the author and may or may not be those of the Laboratory.

Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract W-7405-ENG-48.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This report has been reproduced
directly from the best available copy.

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831
Prices available from (615) 576-8401, FTS 626-8401

Available to the public from the
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Rd.,
Springfield, VA 22161

UCRL

Final Report on LDRD Program

New XAFS Spectroscopic Investigations in the 1-2 keV Region

Tracking Code: 94-LW-062
(unclassified)

Joe Wong, M. Fröba and E. Tamura

Lawrence Livermore National Laboratory, University of California,

PO Box 808, Livermore, CA 94551

Chemistry and Materials Science

Overview - Until recently x-ray absorption fine structure (XAFS) measurements in the 1-2 keV region remained a challenging experimental task. This was primarily due to the lack of an adequate monochromator crystal that possessed both the required x-ray properties (large d-spacing, high resolution and reflectivity) and materials properties (ultra-high vacuum (UHV) compatibility, damage resistance in a synchrotron radiation beam, absence of constituent element absorption edges and stability, both thermal and mechanical). Traditionally, XAFS spectra in this photon energy range have been measured in a piece-wise fashion using a combination of monochromator crystals: beryl (1010) which covers from 780 to 1550 eV, quartz (1010) which covers 1480-1830 eV, InSb (111) which covers 1800-3700 eV and Ge(111) from 2000 eV upwards [1].

Very recently [2], we have an experimental breakthrough in XAFS spectroscopy in this soft x-ray region. This energy region is of great importance for materials and basic research since the K-edges of Na (1070 eV), Mg (1303 eV), Al (1559 eV) and Si (1839 eV), the L-edges of some 4p elements from Ga to Sr and the M-edges of the rare-earth elements fall within this energy window of the electromagnetic spectrum [3]. YB₆₆, a complex binary semiconducting yttrium boride having a cubic crystal structure with a lattice constant of 23.44 Å [4] has been singled out as a candidate monochromator material for synchrotron radiation in the 1-2 keV region [5,6]. There is no intrinsic absorption by the constituent elements in this region, which can adequately be dispersed by the (400) reflection having a 2d value of 11.76 Å (see Fig. 1). In terms of vacuum compatibility, resistance to radiation damage, thermal and mechanical stability, YB₆₆ satisfies all the material requirements for use as a monochromator in a synchrotron beam. In the past few years, LLNL in collaboration with a number of other research institutes [7], has pioneered the development of this unique man-made crystal for use as soft x-ray monochromator with synchrotron light sources for materials science studies. This effort led to an RD-100 Award in 1991[8], and a successful installation of YB₆₆ in January 1993 [2] as a double-crystal monochromator on the JUMBO beamline at SSRL.

Technical Accomplishments - The technical accomplishments are summarized as follows:
Experimental

- Observed direct effects of ligand electronegativity & coordination number on edge energy shift and whiteness intensity respectively in tetrahedrally coordinated Si-containing solids: Si,

SiC, Si₃N₄, SiO₂; and Al-containing compounds: Al₄C₃, AlN, Al₂O₃, AlF₃. Similar systematics were also observed for Mg compounds with ligands and coordination (see Fig. 2)

- Obtained first experimental evidence and elucidated the dependence of multiple scattering in both the Al and Si XANES on the Al-O-Si bond angle in a series of well characterized crystalline aluminosilicate sodalites of known Al-O-Si bond angle [9].
- Correlated intensities of bound-state transitions in the Al XANES spectra with the Al coordination in the Al₂SiO₅ polymorphic model compounds.
- Obtained correlation of disorder (inversion) in MgAl₂O₄ spinel with Mg K-edge XANES.
- Obtained correlation of width of Si K-edge whiteness with crystallinity in silica compounds.
- Obtained correlations of strength and shape of multiple scattering features with the degree of interactions between the host framework and guest molecules in silica sodalites.
- Identified dehydration sites in muscovite (mica) at the 6-fold coordinated Al octahedra.
- Completed Si K-edge measurements in orthosilicates (SiO₄ units), pyrosilicates (Si₂O₇ units), chain silicates (SiO₂)_n, layered silicates (SiO₃)_n and 3-dimensional network silicates.
- Completed experimental work on a series of transition metal (3d, 4d and 5d) disilicides.
- Obtained quality Br L-edge spectra to establish feasibility of L-edge investigations for the elements: Ga, Ge, As, Br,....., Rb, Sr which are important in environmental science.

Theoretical

- Successfully tested our computational code using a Fully Relativistic KKR Green Function Method, and obtained good agreement between the calculated and experimental spectra for both Cu and Al metal [10].
- Successfully developed an imbedded cluster approach to account for the long-range coulombic potential in ionic systems to compute XANES of MgO in good agreement with experiment.

Beamline Instrumentation

- Over the last 2 years of operation, established the long-term stability of YB₆₆ as an UHV compatible and high resolution monochromator in the 1-2 keV region
- Implemented a diamond window to isolate sample chamber from SPEAR UHV, thus enabling a wide range of powder specimens and model compounds to be measured under high vacuum conditions.
- Implemented a quick-change sample holder to maximize usage of beam time.
- Transpose data collection package from the original pdp 11/34 to a microVax.
- Successfully implemented a new Sc-doped YB₆₆ crystal pair with a higher (~25%) energy resolution. Thus we have now the world's (*only*) two pairs of working YB₆₆ monochromators on the JUMBO beamline at SSRL.

Our results in the past two years confirm not only an early prediction [5] for the use of YB₆₆ as synchrotron radiation-compatible soft x-ray monochromator, but also open up a new energy window for studying the bonding and structure of Mg-, Al- and Si-bearing materials *not readily possible* before [11,12]. Furthermore, with YB₆₆, Al EXAFS spectra can now be routinely recorded to 2060 eV, some 500 eV above its K-edge to give a much wider and usable EXAFS data range to $k = 11 \text{ \AA}^{-1}$. In the case of Mg-bearing materials, XAFS spectra may now be recorded from 1250 to 2000 eV, a range of 700 eV above the Mg edge to yield a k_{max} value to $\sim 13.5 \text{ \AA}^{-1}$ (see Fig. 3). This is compared with a much shorter k -range of $\sim 8 \text{ \AA}^{-1}$ available with beryl monochromator which limits data to just below the K-edge of its Al constituent [13].

References cited

1. Z. Hussain, E. Umbach, d.A. Shirley, J. Stohr, J. Feldnaus, Nucl. Instrum. Meth. 195, 115 (1982).
2. M. Rowen, Z.U. Rek, Joe Wong, T. Tanaka, G.N. George, I.J. Pickering, G.H. Via and G.E. Brown, Jr., Synchrotron Rad. News, Vol. 6(3), 1 (1993).
3. J.A. Bearden and A.F. Burr, Rev. Mod. Phys. 39, 125 (1967)
4. S.M. Richards and J.S. Kasper, Acta Cryst. B25, 237 (1969).
5. Joe Wong, W. L. Roth, B.W. Batterman, L.E. Berman, D.M. Pease, S. Heald and T. Barbee, Nucl. Instrum. Meth. 195, 133 (1982).
6. Joe Wong, G. Shimkaveg, W. Goldstein, M. Eckart, T. Tanaka, Z. Rek and H. Tompkins, Nucl. Instrum. Meth. A291, 243, (1990).
7. These include National Institute Research for Inorganic Materials (NIRIM), Tsukuba, Japan: SSR), Stanford; and Berlin Synchrotron Lab (BESSY), Berlin, Germany
8. Energy & Technology Review, UC-LLNL, Sept-Oct. issue (1991)
9. M. Fröba, Joe Wong, P. Behrens, P. Sieger, M. Rowen, t. Tanaka, Z. Rek and J. Felsche, Physica B, 208&209, 65 (1995).
10. E. Tamura, Phys. Rev. B 45, 3271 (1992).
11. G.E. Brown, Jr., F. Dikman and G. Waychunas, SSRL Activity Report 83/01 p.148, (1983).
12. T. Murata, T. Matsukawa and S. Naoe, Sol. State Comm. 66, 787, (1988).
13. J. Cerino, J. Stohr, N. Hower and R.Z. Bachrach, Nucl. Instrum. Meth. 172, 227 (1980).

Publications: resulted from this LDRD work:

1. Development of a YB₆₆ Soft X-ray Monochromator for Synchrotron Radiation, T. Tanaka, Y. Ishizawa, Joe Wong, Z. Rek, M. Rowen, F. Schaefer and B. Mueller, JJAP Series 10, Proc. of the 11th Internat. Sym. on Boron Compounds, 110-113, (1994).
2. New Opportunities in XAFS Investigations in the 1-2 keV Region, Joe Wong, G.N. George, I. J. Pickering, Z.Rek, M. Rowen, T. Tanaka, G.H. Via, B. DeVries D.E.W. Vaughan, and G.E. Brown, Jr., Solid State Comm. 92(7), 559-562, 1994.
3. Spectroscopic Investigations of Mesoporous M41S Silica phases, Michael Fröba, P. Behrens, Joe Wong, G. Engelhardt, Ch. Haggenmüller, G.van de Goor, M. Rowen, T. Tanaka, W. Schwieger, MRS Proc. 371, 99 (1995).
4. Correlation of multiple scattering features in XANES spectra of Al and Si K-edges to the Al-O-Si bond angle in aluminosilicate sodalites: An empirical study, M. Fröba, Joe Wong, P. Behrens, P. Sieger, M. Rowen, t. Tanaka, Z. Rek and J. Felsche, Physica B, 208&209, 65 (1995).
5. Al and Si K absorption edges of Al₂SiO₅ polymorphs using the new YB₆₆ soft x-ray monochromator M. Fröba, Joe Wong , M. Rowen, G.E. Brown, Jr., T. Tanaka and Z. Rek, Physica B, 208&209, 555 (1995).
6. New Aspects of 1-2 keV Spectroscopy, Joe Wong, Z. Rek, M. Rowen, T. Tanaka, F. Schäfers, B. Müller, G. N.; George, I.J. Pickering, G. Via, B. DeVries G. E. Brown, Jr., M. Fröba, Physica B, 208&209, 220 (1995)
7. XANES in Metals - Relativistic Effects and Core Hole Screening, E. Tamura, J. van Ek, M. Fröba and Joe Wong, Phys. Rev. Lett. 72(24), 4899 (1995).
8. Correlation of the growth parameter of YB₆₆ with defect structure and monochromator performance, Z.U. Rek, T Tanaka, M. Rowen and Joe Wong, in prep. for submission to J. Cryst. Growth
9. Anomalous scattering from {006} planes of YB₆₆ at the yttrium L-edges, T. Tanaka, M. Rowen, Z.U. Rek, Y. Kitajima, I. Higashi, Joe Wong, and Y. Ishizawa, in prep. for submission to J. Appl. Cryst.
10. YB₆₆: A New Soft X-ray Monochromator for Synchrotron Radiation. Part II Characterization, Joe Wong, M. Rowen, Z. U. Rek, T. Tanaka, F. Schäfers, B. Müller, in prep. for submission to Nucl. Instrum. Meth.

Soft X-ray Monochromators for Synchrotron Radiation

Figure 1. Soft X-ray monochromators for synchrotron radiation showing the unique features of YB_{66} in the 1-2 keV region. Positions of K absorption edges are marked by vertical arrows for the elements. Inset is a plot of FWHM of YB_{66} double crystal rocking curves as a function of photon energy under various beam conditions.

Figure 2. Normalized K-edge XANES spectra of selected Si, Al and Mg model compounds showing systematic edge shift and increase of whiteline intensity of the $1s \rightarrow 3p$ transition as a function of ligand type (electronegativity) and coordination number.

Figure 3. Experimental Mg K-edge XAFS spectrum of MgO and Al K-edge spectrum of Al_2O_3 illustrating the energy span of EXAFS data collectible with the YB₆₆ monochromator. The small panels are the corresponding phase-corrected (for the 1st oxygen shell) and k^3 -weighted Fourier transform of the extracted EXAFS signals.

L- & M-edge Spectroscopy in the 1-2 keV Region

Figure 4. List of elements with L and M absorption edges in the 1-2 keV (top). Normalized $L_{3,2}$ -edge XANES of Br in KBr and $KBrO_3$ showing richness of the spectral features with bonding and structure of Br in each of the material (bottom).

Technical Information Department • Lawrence Livermore National Laboratory
University of California • Livermore, California 94551