Letter to J. S. Kelly, AEC on the Project Sulky Concept

G. W. Johnson

RECEIVED

JAN 2 1 1997


OSTI

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

lin

June 29, 1964

MASTER



DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

UNIVERSITY OF CALIFORNIA

LAWRENCE RADIATION LABORATORY P. O. BOX 808 LIVERMORE, CALIFORNIA 94551

June 29, 1964

DFCLASSI ICATION

STAMP ON REVERSE.

Mr. John S. Kelly Director, Division of Peaceful Nuclear Explosives U. S. Atomic Energy Commission Washington 25, D. C.

Ref: COPK 63-33, Project Sulky Combut dated October 31, 1963 COPKF 63-4, Paraysis of Propert Sulky based December 27, 1963 COPKF 63-3, Typopsis of the Data and U. B. Astimates of the Distribution of radioactivity at long range from . Sulky

Dear Mr. Kelly:

As you recall, the 100-ton nuclear cratering experiment in hardrock, Sulky, was deferred when we requested permission to execute it a few months ago. After review of the experimental plant the uclear exception program, and the importance of nuclear excavation to the Ational objectives for an Isthmian canal, we are convinced the Sulky a statement should be executed as soon as practical.

There were several questions regarding and descetability of same a from Sulky at the national borders. While experience from the Pike ventice has been informative, no new definitive in formation has been obtained. However, it has been possible to modify the experimental plan to better document release of radioactivity from Sulky. We are convinced that Sulky will not be detected if it is fired under the wind conditions specified during the winter months; however, it is possil . USAN will not concur in this conclusion.

Details of both the changes in experime, al plan and Pike venting analysis are included.

We hereby request your approved to proceed with Project Sulky as originally requested with a read stang data of December 3, 1964. The radio-activity source measurements with slightly modified to improve measurement precision and the wind sector that the initial downwind direct a will fall between 135 and 165 degrees.

corrected by directions Section Sectin Section Section Section Section Section Section Section Section


GERALD W. JOHNSON

Associate Director for Plowshare.

Encl: Appear to Project Sulky Concept

An Evaluation of the Equivalent Fission Yield Vented from Pike

, ill right the contains information affecting the National-Defense of the United Sours within the meaning of the esplorage laws. Time 13 U. S. C. Secs. 793 and 794, the transmission or revelation of which in any manner to an unauthorized person is prohibited by law.



son is prohibited by law.

Brig. Gen. D. Crowson, DMA e copies

E. Shute, SAN

INTERNAL:

T. C. Merkle J. Gofman

J. Rosengren

C. W. Johnson E. Teller

A. C. Haussmann

R. Geockerman

J. Foster/D. Sewell

J. Reeves, MV00

John Kelly, DPNE

EXTERNAL:


:NOITUAIRIZIG

Mr. John S. Kelly

1961 '67 aung

of which in any internet of an unauthorized per-

This material and the National States within the National-Debate And Sect. 19 10 S. C., Sect. 195 and 35 an



I. INTRODUCTION

Since the preparation of the original concept letter additional experience with the detection and evaluation of the Pike vented fraction has improved our perspective on the problems of detectability at long-range and our confidence that most of the debris is deposited close in to the point of detonation.

The objectives of the experiment remain the same:

1. Experimental determination of the vented fraction.

These data are particularly important to the normalization of the theoretical relationships which have been derived to relate depth of burial to the vented fraction for a given medium. A modification of the experimental plan will be made to permit added emphasis on the determination of the airborne source strength within a mile or two of ground zero. These data will make it possible to speak more specifically to the question of the percentage of the total radioactivity carried to long range by the wind and its probability of detection at long range.

The USWB will be advised of this change in emphasis of the program, since it is particularly pertinent to the difference of opinion which exists between LRL and the USWB on the partition of the vented activity between that which readily falls out and that which reaches the ground only through convection and diffusion processes. The Pike results were insufficiently precise to shed any definitive information on the question. Several additional experiments will be required before the problem is well understood. An interpretation of the Pike data is included for reference.

Produce data on crater mechanics at a scaled depth greater than Dannyboy.

High-explosive experiments by their very nature are restricted to yields that are small compared to what is readily accomplished with

OFFICIAL USE ONLY

with nuclear explosives. Further, the data will always be suspect because of (a) the uncertainties in the application of scaling relationships and (b) the differences in the nature of the explosions conducted with HE and NE.


3. Measurement of physical effects of the explosion at intermediate and long range.

It is always possible that effects other than those attributable to radioactivity may be the limiting feature of some specific Plowshare application. In order to reduce the probability of there being surprises associated with the physical effects on future applications of nuclear explosives it is important to continue the accumulation of data which define the limits and the magnitude of such effects.

II. COMPARISON WITH PIKE

Pike was not a cratering shot nor was it a planned vent; as a result any attempt at comparison of Pike and Sulky must be restricted to comparing the history of the vented radioactive debris.

- 1. The mechanism of release of radioactive material from these two shots is quite different. Pike is more comparable to the release of material through a long, small diameter pipe than to a venting of material through overburden as it moves upward and outward during the formation of a crater. There is likely to be a considerable difference in particle size distribution. The evidence is that the vented fraction from Pike contained a larger percentage of its activity on small particles than would be expected for Sulky. The result in such a case would be that Sulky would deposit a larger fraction of the vented activity closer to ground zero, or to put it another way, that a smaller fraction of the vented activity should be carried to long distances.
- 2. The vented fraction on Sulky is expected to be a factor of 5 to 10 smaller than for Pike. Further, Sulky would be shot only under meteorological conditions which would provide the maximum possible residence time for the airborne debris within the boundaries of the United States. This was not true for Pike. The winds at shot time



were northerly, with minimum velocity and directional shear, and of appreciable speed. These are precisely opposite to the conditions which have been established for conduct of the Sulky experiment.

A careful choice of wind structure, combined with Sulky's smaller vented fraction, should reduce the probability of detection at any border to insignificance; in part because of the dilution with distance, and in part because of the small percentage probability of reaching either Mexico or the Pacific Coast if shooting is restricted to the 135 to 165 degree sector (see data included in Table I, as extracted from the USWB publication, "A Climatology of 700 mb Trajectories Originating at the Nevada Test Site," dated March 1964).

TABLE I
TRAJECTORY EXIT PROBABILITIES, Dec. Through Feb.*

TRAJECTORY EXIT	PROBABILITY %	MEDIAN NUMBER OF DAYS
West Canada	50	2.4
Est Canada	0	
Atlantic Ocean	50	4.0
Gulf of Mexico	0	
Mexico	0	
Pacific Ocean	0	

^{*}For trajectories originating at NTS in the 135 to 165 degree sector.

III. COST AND TIME ESTIMATE

No changes are anticipated. The modifications to the experimental plan as outlined above are believed to be within the original estimates.