**Cover Sheet for a Hanford
Historical Document
Released for Public Availability**

Released 1995

**Prepared for the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DECLASSIFIED

FINAL REPORT
PRODUCTION TEST 231-12
RECOVERY OF AMERICIUM

BY

G. V. PACKER
TECHNICAL SECTION
ENGINEERING DEPARTMENT

MARCH 2, 1954

Classification Cancelled and Changed To

DECLASSIFIED

By Authority of CG-PR-2

CH Henager, 7-31-92

By Jenni Malley 9-26-94

Verified By J E Saville 9-28-94

NOTUCNI

HANFORD ATOMIC PRODUCTS OPERATION
RICHLAND, WASHINGTON

GENERAL ELECTRIC

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DECLASSIFIED

MASTER

DECLASSIFIED

HW-30999

Technology - Hanford Processes

This document consists
of 16 pages

COPY 1 OF 1, SERIES MA

FINAL REPORT
PRODUCTION TEST 231-12
RECOVERY OF AMERICIUM

By

G. V. Packer

Plant Processes Unit
Separations Technology Sub-Section

March 2, 1954

HANFORD ATOMIC PRODUCTS OPERATION
RICHLAND, WASHINGTON

Operated for the Atomic Energy Commission by the
General Electric Company under Contract #W-31-109-Eng-52

[REDACTED]

[REDACTED]

[REDACTED]

Route To	Read By	Date	Route To	Read By	Date
Yellow Copy					
RECORD CENTER FILE					

RECEIVED
JUL 2 1956

300 AREA
CLASSIFIED FILES

DECLASSIFIED

DECLASSIFIED

-2-

HW-30999

Technology - Hanford Processes

INTERNAL DISTRIBUTION

COPY NUMBER

1	O. H. Greager - W. K. Woods
2	A. B. Greninger
3	F. W. Albaugh
4	V. R. Cooper
5	O. F. Hill
6	W. H. Reas
7	A. E. Smith - G. V. Packer
8	E. R. Irish
9	F. W. Woodfield
10	W. N. Mobley - O. F. Beaulieu
11	L. M. Knights - J. W. Jordan
12	J. E. Maider - T. W. Hauff
13	C. A. Priode
14	R. S. Bell - V. R. Chapman
15	C. T. Groswith
16	H. A. Moulthrop - L. M. Meeker
17	V. D. Donihee
18	A. H. Bushey - H. R. Schmidt
19	300 File
20	Yellow Copy

EXTERNAL DISTRIBUTION

COPY NUMBER

21	Armed Forces Special Weapons Project, Washington
22 - 23	Atomic Energy Commission, Washington
24	Chicago Patent Group
25 - 28	duPont Company, Augusta
29	duPont Company, Wilmington
30	Hanford Operations Office
31	Hanford Operations Office: Attn. J. I. Thomas
32 - 34	Knolls Atomic Power Laboratory
35	Patent Branch, Washington
36	USAF Headquarters
37 - 51	Technical Information Service, Oak Ridge

DECLASSIFIED

DECLASSIFIED

-3-

HW-30999

FINAL REPORT
PRODUCTION TEST 231-12
RECOVERY OF AMERICIUM

I. INTRODUCTION

In response to the need for isolating small quantities of americium for employment in research studies, principally at laboratories not connected with the Hanford Atomic Products Operation, a program was formulated ⁽¹⁾ ⁽²⁾ to accomplish this at minimum cost and disturbance to plutonium production. This report describes the operations employed and the results obtained in the isolation of americium from aged plutonium nitrate solution utilizing the Isolation Building process equipment.

II. OBJECTIVE

The objective of the production test was to establish a process for isolating americium from aged plutonium using existing equipment.

III. SUMMARY

A method for isolating Am^{241} from a mixture of plutonium isotopes using a peroxide precipitation nearly identical to that used to separate plutonium from lanthanum was developed for use in Hanford Atomic Products Operation plant equipment.

A total of 98 milligrams of Am^{241} were isolated from about 1600 grams of aged plutonium (as plutonium nitrate) and 950 milligrams of Am^{241} were isolated from about 6.4 Kg of aged plutonium (as metal scrap) using the isolation building equipment. Based on radiochemical analyses the yield ranged from 92 to 104 per cent.

The use of filter sticks to separate plutonium peroxide from the supernatant gives an acceptable Am^{241} concentration in the supernatant without further concentration. When the wash solution is added to the supernatant the total Am^{241} content is increased slightly, however, the concentration is reduced excessively.

DECLASSIFIED

DECLASSIFIED

-4-

HW-30999

IV. EXPERIMENTAL

The raw material for this production test was about 1600 grams of plutonium as dried plutonium nitrate aged 0.79 years to 1.06 years and 6400 grams of plutonium as metal scrap accumulated in storage between 1949 and 1953.

The concentrated and aged plutonium nitrate was flushed from the sample cans and processed in two can batches through a plutonium peroxide precipitation in the second cycle precipitator tank. The supernatant solution (including the wash supernatant) containing the Am^{241} was transferred to the supernatant collection tank where the excess hydrogen peroxide was thermally decomposed. The supernatant solutions of the 0.79 year aged plutonium were transferred to a glass lined evaporator in the Purification Building, concentrated from 91 to 13 liters and loaded into the shipping containers. Americium was isolated from the 1.06 year aged plutonium in the same manner except that the supernatant solution was not concentrated since it could be contained in the shipping vessel. The plutonium portions (peroxide cakes) were processed through the treatment which follows the normal second peroxide cycle.

The excess metal from samples taken in the fabrication building has been stored since 1949 pending recovery of the plutonium. In the spring of 1953 an americium recovery program was incorporated with the recovery of this aged metal using the plutonium peroxide precipitation process. The feed solution was prepared by dissolving the plutonium metal in hot nitric acid (70 per cent) and hydrofluoric acid (0.02M) and diluting with distilled water to the desired hydrogen-ion concentration (2.0M).

Instead of decanting off the supernatant solution, the peroxide slurry was filtered through the sintered stainless steel filters, which facilitated the removal of a greater volume fraction of solution from the cake than was possible by decanting.

DECLASSIFIED

DECLASSIFIED

-5-

HW-30999

It was advantageous, for this reason, to collect only the original americium - rich filtrate and exclude the wash filtrate solution, thereby eliminating the need for concentration of the solution containing the Am^{241} .

V. DISCUSSION

The extent of the batch recovery of americium from plutonium depends primarily upon the mechanical ability to transfer the solutions from one vessel to another without hold-up and upon thorough washing of the plutonium peroxide precipitate within chemical conditions compatible with low plutonium losses to the supernatant solution. The problem of solution hold-up was solved by following each transfer with flushes employing dilute nitric acid. It is not practical to remove all of the soluble americium by copiously washing the plutonium peroxide cake since this would dilute the americium excessively and result in solution volumes greater than those which can be economically concentrated. Based upon the relative solubility of the americium and the desire to keep the plutonium concentration and final combined supernatant volume low the normal three washes of plutonium peroxide cake (5.0 Kg each of 6.0 per cent nitric acid) were used for the first Am^{241} isolation runs. The combined supernatant-wash solution was concentrated in a glass-lined evaporator.

A summary of the Am^{241} isolation and concentration from the first eight cans (aged 0.79 years) of material processed is given in Tables I and II. The analytical results of these runs are presented in Tables III and IV to show the typical solution concentrations and significant impurities.

Additional Am^{241} was isolated from two sample cans of material aged 1.06 years. The americium²⁴¹ bearing supernatant solution was not concentrated because the solution volume was less than the capacity of the shipping container. This run is summarized in Table V.

DECLASSIFIED

DECLASSIFIED

-6-

HW-30999

Twenty typical americium by-product runs made during the 1953 plutonium recovery program are listed in Table VIII. These runs returned 6.4Kg of plutonium to the process and recovered 0.95% grams of Am^{241} (81 per cent yield of the Am^{241} radiochemically determined to be in the starting solutions).

The amounts of Am^{241} indicated in Tables I and V approximated the amounts which were calculated to be present based on the age of the plutonium and the pile exposure history of the original uranium metal. When uranium is discharged from a Hanford pile it contains some of the isotope, Pu^{241} , which is formed by a combination of nuclear reactions during the pile exposure. The amount is related to the MWD/T as shown in Figure I. ⁽⁵⁾ Plutonium-241 (14.2 year half-life) decays to Am^{241} by emitting a beta-particle. Thus when the uranium metal is stored 70 - 100 days before it is processed for plutonium isolation, 1 - 1.3 per cent of the Pu^{241} decays. The daughter isotope Am^{241} is removed with the waste and fission product solutions of the separations process, therefore, the plutonium that was set aside for this production test initially contained about 99 per cent of the Pu^{241} present in the uranium metal at the time of pile discharge. Figure II shows the relationship of Am^{241} growth per gram of Pu^{241} (present at start of aging) to the age of the plutonium since the last chemical separation, based on the half-lives 14.2 and 470 years of Pu^{241} and Am^{241} , respectively.

DECLASSIFIED

DECLASSIFIED

-7-

HW-30999

Calculating the amount of Am^{241} present in aged plutonium is illustrated below for the first run (see Tables I, III, and VI):

Amount of plutonium in starting solution	348 grams
Pile exposure of original uranium	394 MWD/Ton
Ratio: $\text{Pu}^{241}/\text{Pu}$ (from Figure I)	0.00134 grams/gram
Amount of Pu^{241}	$0.00134 \times 348 = 0.466$ grams
Age of plutonium since last chemical separation	0.79 years = 9.5 months
Ratio: $\text{Am}^{241}/\text{Pu}^{241}$ (from Figure II)	0.038 grams/gram
Calculated amount of Am^{241}	$0.038 \times 0.466 \times 1000 = 17.7$ mg
Amount of Am^{241} in the starting solution by radiochemical assay	20.7 mg

The actual amounts of Am^{241} are compared to the calculated amounts in Table VI. The actual values were used to back-calculate $\text{Pu}^{241}/\text{Pu}$ weight ratios for comparison to the curve of Figure I. The $\text{Pu}^{241}/\text{Pu}$ ratio for the first run based on the Am^{241} radiochemically determined to be the starting solution was computed as shown below:

Age of process starting solution	0.79 years = 9.5 months
Ratio: $\text{Am}^{241}/\text{Pu}^{241}$ (from Figure II)	0.038 grams/gram
Amount of Am^{241} in the starting solution	20.7 mg = 0.0207 gms
Amount of Pu^{241}	$0.0207/0.038 = 0.545$
Amount of plutonium in starting solution	348 grams
Ratio: $\text{Pu}^{241}/\text{Pu}$	$0.545/348 = 0.00156$ grams/g
Pile exposure of original uranium	394 MWD/ton
Ratio: $\text{Pu}^{241}/\text{Pu}$ (from Figure I)	0.00134 grams/gram

These and the calculations for the other four runs are given in Table VII.

GVP:ms

G. V. Packer

G. V. Packer

DECLASSIFIED

DECLASSIFIED

-8-

HW-30999

VI. REFERENCES

1. HW-16038, "Americium Separation from Plant Solutions", O. F. Hill to F. W. Albaugh, February 15, 1950.
2. HW-17518, "Discussions on Americium Processes", M. J. Szulinski and G. V. Packer to File, April 13, 1950.
3. HW-19119, "Laboratory Study of Americium Recovery Processes", C. F. Callis to File, October 1, 1950.
4. HW-17990, "Production Test 231-12 Recovery of Americium", M. J. Szulinski to File, June 2, 1950.
5. HW-26584, "The Isotopic Content and Specific Activity of Pile-Produced Plutonium", by E. M. Kinderman, H. R. Schmidt, G. J. Alkire, A. Chetham-Strode, and Roy Ko, January 12, 1953.

DECLASSIFIED

DECLASSIFIED

TABLE I
AMERICIUM MATERIAL BALANCE, ISOLATION BUILDING

Run No.	Metal History MWD/Ton	Am ²⁴¹ Total Weight, mg.			Material Balance Per Cent	Am ²⁴¹ Lost	
		Aged Starting Solution	Original+Wash Supernatant Solution	Concentrated Plutonium Cake Solution		In Cake Solution, Per Cent	Am ²⁴¹ Isolated, Per Cent
1	394	20.7		3.00			
2	335	18.4	35.9	1.88	104.3	12.5	92.0
3	337	16.3		0.98			
4	415	20.0	37.6	4.63	119.0	15.4	103.5
Total		75.4	73.5 (1)	10.49	111.4	13.9	

(1) The final combined solution of Am²⁴¹ contained approximately 3 grams of plutonium.

TABLE II
AMERICIUM MATERIAL BALANCE, PURIFICATION BUILDING

	<u>Input, mg.</u>	<u>Output, mg.</u>
Before Concentration (Combined analysis)	76.1	
After Concentration (for off-site)		57.6
After Concentration (for on-site)	---	4.0
Total	76.1	61.6
Material Balance, Per cent	81	
Unaccounted for loss, Per cent	19	

DECLASSIFIED

DECLASSIFIED

-10-

HW-30999

TABLE III
AMERICIUM RECOVERY ANALYTICAL RESULTS, ISOLATION BUILDING
FIRST EIGHT SAMPLE CANS

<u>Run No.</u>		<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>
Starting Solution	Am, g/l	2.71×10^{-3}	2.55×10^{-3}	1.96×10^{-3}	2.30×10^{-3}
	Pu, g/l	45.42	44.46	37.78	36.97
	H, g/l	1.07	1.14	1.55	1.00
	Fe, g/l	0.235	0.243	0.196	0.167
	Sp. Gr.	1.112	1.108	1.11	1.093
Volume, liters		7.6	7.2	8.3	8.7
Supernatant Solution (1)	Am, g/l	0.804×10^{-3}		0.718×10^{-3}	
	Pu, g/l	0.0863		0.0520	
	Fe, g/l	0.103		0.073	
	Sp. Gr.	---		1.052	
	Volume, liters	44.7		52.4	
Concentrated Plutonium Cake Solution	Am, g/l	2.35×10^{-3}	1.54×10^{-3}	0.76×10^{-3}	3.54×10^{-3}
	Pu, g/l	256.8	255.2	239.7	235.3
	Fe, g/l	0.507	0.483	0.378	0.350
	Sp. Gr.	1.673	1.681	1.659	1.654
	Volume, liters	1.28	1.22	1.29	1.31

(1) The supernatant solutions of runs 1 and 2 were combined and sampled; likewise, the supernatant solutions of runs 3 and 4 were combined and sampled.

DECLASSIFIED

DECLASSIFIED

-11-

HW-30999

TABLE IV
AMERICIUM RECOVERY ANALYTICAL RESULTS, PURIFICATION BUILDING
FIRST EIGHT SAMPLE CANS

Combined Supernatant Solution (Before Concentration)		Spectrochemical	Combined Supernatant Solution (After Concentration)	
			Chemical	
Am, g/l	0.836×10^{-3}	Ca, ppm of Pu	Am, g/l	4.74×10^{-3}
Pu, g/l	0.0954	Cr, ppm of Pu	Pu, g/l	0.6600
Sp. Gr.	1.045	La, ppm of Pu	Fe, g/l	1.064
Volume,		Na, ppm of Pu	Sp. Gr.	1.235
liters	91.0	Ni, ppm of Pu	Volume,	13.0
			liters	

DECLASSIFIED

DECLASSIFIED

TABLE V
AMERICIUM MATERIAL BALANCE, ISOLATION BUILDING

Run No.	Metal History MWD/Ton	Am ²⁴¹ Total Weight, mg.			Material Balance Per Cent	Am ²⁴¹ Lost	Am ²⁴¹ Isolated Per Cent
		Aged Starting Solution	Original+ Supernatant Solution	Wash Concentrated Plutonium Cake Solution		In Cake Solution, Per Cent	
5	372	26.5	24.2	2.2	99.6	8.3	91.5

TABLE VI
AM²⁴¹ CALCULATED TO BE IN THE STARTING SOLUTIONS

MWD/Ton	Pu, grams	Pu ²⁴¹ /Pu ⁽¹⁾ Wt. Ratio	Pu ²⁴¹ , grams	Age, months	Am ²⁴¹ /Pu ²⁴¹ Wt. Ratio ⁽²⁾	Am ²⁴¹ , mg	
						Calculated	Actual
394	348	0.00134	0.466	9.5	0.038	17.7	20.7
335	321	0.00097	0.311	9.5	0.038	11.8	18.4
337	315	0.00098	0.309	9.5	0.038	11.7	16.3
415	321	0.00149	0.478	9.5	0.038	18.2	20.0
372	332	0.00120	0.398	12.7	0.050	19.9	28.5

- (1) Obtained from Figure I.
(2) Obtained from Figure II.

DECLASSIFIED

DECLASSIFIED

TABLE VII
PLUTONIUM-241 CONTENT OF PLUTONIUM DERIVED FROM PILE
IRRADIATED URANIUM AT THE TIME OF LAST CHEMICAL SEPARATION

Run No.	Age of Process Starting Solution Months	Wt. Ratio Am ²⁴¹ /Pu ²⁴¹ from Figure II	Am ²⁴¹ in Process Starting Solution, Grams	Pu ²⁴¹ Present At Start of Aging	Total Pu Isotopes In Process Starting Solution, gms	Back- calculated Wt. Ratio Pu ²⁴¹ /Pu	MWD/ Ton	Fig. I Wt. Ratio Pu ²⁴¹ /Pu (Ref. 5)
1	9.5	0.038	0.0207	0.545	348	0.00156	394	0.00134
2	9.5	0.038	0.0184	0.484	321	0.00150	335	0.00097
3	9.5	0.038	0.0163	0.429	315	0.00136	337	0.00098
4	9.5	0.038	0.0200	0.526	321	0.00164	415	0.00149
5	12.7	0.05	0.0265	0.530	332	0.00160	372	0.00120

DECLASSIFIED

DECLASSIFIED

-14-

HW-30999

TABLE VIII
SOME TYPICAL AMERICIUM RECOVERY RUNS MADE DURING SPRING,
1953

Run No.	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Z-12-12-Z-16	11.0	310	53	56	20.0	4	48	1.3	91
Z-12-12-Z-17	9.5	330	54	41	17.0	3	47	0.9	87
Z-12-12-Z-18	18.6	270	36	49	27.5	4	30	1.5	83
Z-12-12-Z-19	14.5	340	53	54	15.0	4	44	1.2	83
Z-12-12-Z-20	10.4	330	42	46	10.0	3	35	0.9	83
Z-53-4-Z-25	9.8	300	63	73	9.0	4	41	1.3	65
Z-53-4-Z-26	11.4	340	47	63	14.0	10	43	2.9	92
Z-53-5-Z-1	12.3	300	64	75	11.5	2	46	0.7	72
Z-53-5-Z-5	12.2	350	72	76	21.0	2	63	0.6	87
Z-53-5-Z-6	13.9	390	71	73	13.5	2	54	0.5	77
Z-53-5-Z-8	10.9	330	66	65	10.0	2	45	0.6	69
Z-53-5-Z-9	10.3	260	51	64	13.0	7	44	2.7	86
Z-53-5-Z-14	11.8	360	73	82	12.0	5	56	1.4	77
Z-53-05-Z-15	12.5	340	82	82	13.5	8	72	2.4	88
Z-53-05-Z-18	11.1	340	87	86	12.0	4	65	1.2	75
Z-53-05-Z-23	18.0	310	59	64	21.0	9	50	2.9	84
Z-53-05-Z-24	20.7	360	64	67	22.0	1	51	2.8	80
Z-53-05-Z-26	19.3	280	54	59	22.0	2	46	0.7	86
Z-53-05-Z-27	19.6	270	38	36	20.5	2	34	0.7	89
Z-53-05-Z-28	19.2	300	48	48	21.0	2	36	0.7	75
Total, 20 runs		6410	1177	1259		80	950		

- (1) Starting solution volume, liters.
- (2) Starting solution Pu total grams.
- (3) Starting solution Am²⁴¹ total mg (radiochemical analysis).
- (4) Starting solution Am²⁴¹ total mg (calculated from history data).
- (5) Peroxide supernatant solution volume, liters.
- (6) Peroxide supernatant solution Pu total grams.
- (7) Peroxide supernatant solution Am²⁴¹ total mg.
- (8) Peroxide supernatant solution per cent Pu lost.
- (9) Peroxide supernatant solution per cent Am²⁴¹ recovered.

DECLASSIFIED

DECLASSIFIED

FIGURE I
PLUTONIUM-241 CONTENT OF PLUTONIUM
DERIVED FROM PILE-IRRADIATED URANIUM
AT THE TIME OF LAST CHEMICAL SEPARATION

DECLASSIFIED

DECLASSIFIED

-16-

HW-30999

FIGURE II
GRAMS OF AMERICIUM-241 AFTER AGING
ASSOCIATED WITH ONE GRAM
OF PLUTONIUM-241 AT START
OF AGING

DECLASSIFIED