DOE/OR/21383--T1

Computer Support to Run Models of the Atmosphere

Final Report for
Interagency Agreement DOE IAG DE-AI05-830R21383

between

National Aeronautics and Space Administration
Goddard Space Flight Center
Institute for Space Studies
2880 Broadway
New York, NY 10025

RECEIVED

OCT 16 1996

OSTI

and

U.S. Department of Energy
Office of Energy Research

Inez Fung

NASA Goddard Institute for Space Studies
2880 Broadway
New York, NY 10025

August 30 1996

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

Our research is focused on a better quantification of the variations in CO₂ exchanges between the atmosphere and biosphere and the factors responsible for these exchanges. The principal approach is to infer the variations in the exchanges from variations in the atmospheric CO₂ distribution.

The principal tool involves using a global three-dimensional tracer transport model to advect and convect CO₂ in the atmosphere. The tracer model we use was developed at the Goddard Institute for Space Studies (GISS) and is derived from the GISS atmospheric general circulation model. A special run of the GCM is made to save high-frequency winds and mixing statistics for the tracer model. We note the GISS tracer model is widely distributed, and forms the basis for the tracer models at Scripps Institution of Oceanography (Keeling et al., 1989), Max-Planck Institute, Hamburg (Heimann, 1995) and CSIRO, Australia (Enting, et al., 1993).

Source/sink distributions need to be specified for the tracer model. For some sources and sinks, e.g. fossil fuel combustion, the pattern and magnitudes can be specified with some confidence. For others, e.g. biospheric uptake, the patterns and magnitudes are the major unknowns in the carbon cycle. Where possible, we have compiled information about the "better-known" sources and sinks, and gridded the information on carbon exchanges into a format compatible with the atmospheric models. These source/sink patterns are distributed via CDIAC as DB1006. They form the basis for an international tracer model intercomparison (Law and Rayner, 1995).

To deduce the sought-after magnitudes and patterns of the net uptake by the biosphere and by the ocean, and to test hypotheses about "known" sources and sinks, we developed a new approach for the inverse problem. We first divide the biosphere and ocean into units (e.g. a vegetation type or the North Atlantic between 15N and 50N). For each unit, we assign a hypothetical source strength of 1 GtC/y. The tracer model is run for each unit. The resultant atmospheric CO₂ responses are then combined, with the source strength of each unit as those required to provide a best fit between

the simulated and observed atmospheric CO₂ variations.

Each tracer model experiment is integrated for a minimum of 3 years, so that the north-south gradient stabilizes for the specified source/sink. Typically, we divide the biosphere into 11 vegetation types, and the oceans into 15 regions. Together with the fossil fuel and landuse sources, an investigation of a single carbon budget scenario involves ~ 90 model years.

The tracer model experiments are carried out on the successive generations of computers at GISS. With the tremendous leaps in computational capability, the cpu requirement has dropped from 24 cpu hours/model year on the 1983 Amdahl computers to 5 cpu minutes/model year on the 1996 IBM 6000 RISC station.

References

Enting, I. G., C. M. Trudinger, R. J. Francey, and H. J. Granek (1993): Synthesis inversion of atmospheric CO₂ using the GISS tracer transport model. CSIRO DAR Tech. Report No. 29, 44 pp.

Keeling, C. D., S. C. Piper, M. Heimann (1989): A three-dimensional model of atmospheric CO₂ transport based on observed winds: 4: Mean annual gradients and interannual variations. In *Aspects of Climate Variability in the Pacific and Western Americas*, D. Peterson (ed.), Geophysical Monograph 55, American Geophysical Union.

Heimann, M. (1995): The global atmospheric tracer model TM2. Deutsches Klimarechenzentrum, Technical Report No. 10, 51 pp.

Law, R.M. and P.J. Rayner (1995): A comparison of modelled responses to prescribed CO₂ sources. CSIRO DAR Tech. Report No. 36.

The publications resulting from the IAG are listed below.

- Fung, I., K. Prentice, E. Matthews, J. Lerner and G. Russell (1983): A 3-D tracer model study of atmospheric carbon dioxide: Response to seasonal exchanges with the terrestrial biosphere. *J. Geophys. Res.*, 88, 1282- 1294.
- Emanuel, W.R., I. Fung, G.G. Killough, Jr., B. Moore III, and T.-H. Peng (1985): Modeling the global carbon cycle and changes in atmospheric carbon dioxide levels. Chapter 7 in *Atmospheric Carbon Dioxide and the Global Carbon Cycle*, J.R. Trabalka (ed), U.S. Department of Energy DOE/ER-0239, December, 1985.
- Tucker, C.J., I. Fung, C.D. Keeling and R.H. Gammon (1986): Relationship between atmospheric CO₂ variations and a satellite-derived vegetation index. *Nature*, 319, 195-198.
- Fung, I. (1986): Analysis of the seasonal and geographic patterns of atmospheric CO₂ distributions with a 3-D tracer model. In: *The Changing Carbon Cycle: A Global Analysis*, J.R. Trabalka and D.E. Reichle (eds), Springer-Verlag, New York.
- Heimann, M., C.D. Keeling and I. Fung (1986): Simulating the atmospheric carbon dioxide distribution with a three dimensional tracer model. In *The Changing Carbon: A Global Analysis*, J.R. Trabalka and D.E. Reichle (eds), Springer-Verlag, New York.
- Fung, I., C.J. Tucker and K.C. Prentice (1987): Application of advanced very high resolution radiometer vegetation index to study atmosphere- biosphere exchange of CO₂. *J. Geophys. Res.*, 92, 2999-3015.
- D'Arrigo, R., G. Jacoby and I. Fung (1987): The role of boreal forests in atmosphere-biosphere exchange of carbon dioxide. *Nature*, 329, 321-323.
- Fung, I.Y., C.J. Tucker and K.C. Prentice (1987): On the variability of atmospheric-biosphere exchange of CO₂. *Advances in Space Research*, 7, (11)175-(11)180.
- Fung, I. and J. John (1989): Interannual and longer-term changes of the terrestrial biosphere and their relationships to atmospheric CO₂ variations. In: *Proceedings of Third International Conference on Analysis and Evaluation of Atmospheric CO₂ Data Present and Past*, Environmental Pollution Monitoring and Research Programme No. 59, World Meteorological Organization.
- Tans, P.P., I. Fung and T. Takahashi (1990): Observational constraints on the global atmospheric CO₂ budget. *Science*, 247, 1431-1439.
- Fung, I. (1990): *An Earth Atlas*. An unpublished atlas compiled and produced to celebrate the 65th birthday of Professor Bert Bolin.
- Prentice, K. and I. Fung (1990): Bioclimatic simulations test the sensitivity of terrestrial carbon storage to perturbed climate. *Nature*, 346, 48-50.
- Takahashi, T., P. Tans and I. Fung (1992): Balancing the budget: Carbon dioxide sources and sinks, and the effects of industry. *Oceanus*, 35(1), 18-28.
- Bolin, B. and I. Fung (1992): The carbon cycle revisited. In: *Modeling the Earth System*, D. Ojima (ed.), pp. 151-164, OIES Global Change Institute Vol. 3, UCAR, Boulder, CO.
- Graedel, T.E., T.S. Bates, A.F. Bouwman, D. Cunnold, J. Dignon, I. Fung, D.J.

- Jacob, B.K. Lamb, J.A. Logan, G. Marland, P. Middleton, J.M. Pacyna, M. Placet and C. Veldt (1993): A compilation of inventories of emissions to the atmosphere. *Global Biogeochemical Cycles*, 7, 1-26.
- Fung, I. (1993): Models of oceanic and terrestrial sinks of anthropogenic CO₂: A review of the contemporary carbon cycle. In *The Biogeochemistry of Global Change: Radiative Trace Gases*, R.S. Oremland (editor), pp. 166-189, Chapman and Hall, NY.
- Dai, A.G. and I. Fung (1993): Can climate variability contribute to the "missing" CO₂ sink? *Global Biogeochemical Cycles*, 7, 599-609.
- Tans, P., I. Fung and I. Enting (1995): Storage versus flux budgets: The terrestrial Uptake of CO₂ during the 1980's. In *"Biotic Feedbacks in the Global Climatic System: Will the Warming Feed the Warming?"* G. Woodwell and F. Mackenzie (eds.), Oxford University Press. pp. 351-366.
- Fung, I. (1995) Perturbations to the biospheric carbon cycle: uncertainties in the estimates. In *"Biotic Feedbacks in the Global Climatic System: Will the Warming Feed the Warming?"* G. Woodwell and F. Mackenzie (eds.), Oxford University Press pp. 366-374.
- Fung, I. and T. Takahashi (1995): Assessment of factors affecting estimates of air-sea exchanges of CO₂ from pCO₂ gradients. In *"The Carbon Cycle"*, T. Wigley and D. Schimel (eds.) (in press).
- Friedlingstein, P., K. Prentice, I. Fung, J. John, and G. Brasseur (1995): Carbon-biosphere-climate interactions in the LGM climate. *J. Geophys. Res.*, 100, 7203-7221.
- Friedlingstein, P., I. Fung, E. Holland, J. John, G. Brasseur, D. Erickson, and D. Schimel (1995): On the contribution of biospheric CO₂ fertilization to the missing sink. *Global Biogeochemical Cycles*, 9, 541-556.
- Braziunas, T. F., I. Y. Fung and M. Stuiver (1995): The preindustrial atmospheric ¹⁴CO₂ latitudinal gradient as related to exchanges among atmospheric, oceanic and terrestrial reservoirs. *Global Biogeochemical Cycles*, 9, 565-584.
- Denning, A.S., I. Fung, and D. Randall (1995): Strong simulated meridional gradient of atmospheric CO₂ due to seasonal exchange with the terrestrial biota. *Nature*, 376, 240-242.