INCLUSIVE HADRONIC PRODUCTION OF THE B_c MESON VIA HEAVY QUARK FRAGMENTATION *

KINGMAN CHEUNG[†]

Center for Particle Physics, University of Texas at Austin, Austin TX 78712, U.S.A.

E-mail: cheung@utpapa.ph.utexas.edu

ABSTRACT

We summarize the studies on the hadronic production of S- and P-wave ($\bar{b}c$) mesons via direct fragmentation of the bottom antiquark as well as the Altarelli-Parisi induced gluon fragmentation.

The direct production of heavy mesons like J/ψ , Υ , and ($\bar{b}c$) mesons can provide very interesting tests for perturbative QCD. According to the potential model calculation¹, for ($\bar{b}c$) mesons the first two sets ($n = 1$ and $n = 2$) of S-wave states, the first ($n = 1$) and probably the entire second set ($n = 2$) of P-wave states, and the first set ($n = 1$) of D-wave states lie below the BD flavor threshold. Since the annihilation channel of excited ($\bar{b}c$) mesons is suppressed relative to the electromagnetic and hadronic transitions, the excited states below the BD threshold will cascade down into the ground state B_c via emission of photons and/or pions. Inclusive production of the B_c meson therefore includes the production of the $n = 1$ and $n = 2$ S-wave and P-wave states, and the $n = 1$ D-wave states. Here we do not include the D-wave contributions since they are expected to be very small.

Intuitively, the dominant production of ($\bar{b}c$) mesons at the large transverse momentum region should come from the direct fragmentation of the heavy $\bar{b}$ antiquark^{2,3}. We calculate the hadronic production of S- and P-wave ($\bar{b}c$) mesons using the fragmentation approach^{4,5,6}. The fragmentation approach essentially involves the factorization of the whole production process into the production of a high energy parton (a $\bar{b}$ antiquark or a gluon) and the fragmentation of this parton into various ($\bar{b}c$) states. The novel feature in our approach^{2,3} is that the relevant fragmentation functions at the heavy quark mass scale can be calculated in perturbative QCD. Let H denotes any ($\bar{b}c$) meson states. The differential cross section $d\sigma/dp_T$ versus the transverse momentum p_T of H is given by

$$\begin{aligned} \frac{d\sigma}{dp_T}(p\bar{p} \rightarrow H(p_T)X) &= \sum_{ij} \int dx_1 dx_2 dz f_{i/p}(x_1, \mu) f_{j/\bar{p}}(x_2, \mu) \left[\frac{d\hat{\sigma}}{dp_T}(ij \rightarrow \bar{b}(p_T/z)X, \mu) \right. \\ &\quad \left. \times D_{\bar{b} \rightarrow H}(z, \mu) + \frac{d\hat{\sigma}}{dp_T}(ij \rightarrow g(p_T/z)X, \mu) D_{g \rightarrow H}(z, \mu) \right]. \quad (1) \end{aligned}$$

*Talk presented at Beyond the Standard Model IV, Lake Tahoe, California (Dec 1994)

[†]Representing also Tzu Chiang Yuan, UC-Davis

MASTER

TS

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Figure 1: The differential cross section $d\sigma/p_T$ versus p_T of the $(\bar{b}c)$ meson (H) in various spin-orbital states with $n = 1$ at the Tevatron. The acceptance cuts are $p_T(H) > 6$ GeV and $|y(H)| < 1$.

For the production of $\bar{b}$ we include the subprocesses $gg \rightarrow b\bar{b}$, $g\bar{b} \rightarrow g\bar{b}$, and $q\bar{q} \rightarrow b\bar{b}$; while for the gluon g we include the subprocesses $gg \rightarrow gg$, $q\bar{q} \rightarrow gg$, and $gq(\bar{q}) \rightarrow gq(\bar{q})$. In Eq. (1), a common scale μ is chosen for parton distribution functions, parton-parton scattering, and fragmentation functions. We estimate the dependence on μ by varying the scale $\mu = (0.5 - 2)\mu_R$, where $\mu_R = \sqrt{p_T^2(\text{parton}) + m_b^2}$. This choice of scale avoids the large logarithms in the short-distance part $\hat{\sigma}$'s. However, logarithms of order μ_R/m_b have to be summed in the fragmentation functions, which is implemented by evolving the Altarelli-Parisi (AP) equations for the fragmentation functions^{4,5}. The initial conditions for the AP equations are the fragmentation functions that we can calculate by perturbative QCD at the initial scale μ_0 , which is of the order of the b -quark mass. At present, all the S-wave² and P-wave³ fragmentation functions for $\bar{b} \rightarrow (\bar{b}c)$ have been calculated to leading order in α_s . To obtain the fragmentation functions at an arbitrary scale greater than μ_0 , we numerically integrate the AP evolution equations.

Other details in inputs can be found in Ref. 6. We impose $p_T(H) > 6$ GeV and $|y(H)| < 1$ cuts on the $(\bar{b}c)$ state H . The p_T spectra for the $(\bar{b}c)$ state H with various spin-orbital quantum numbers are shown in Fig. 1 and Fig. 2 for $n = 1$ and $n = 2$, respectively. Thus, we can also obtain the inclusive production rate of B_c as a function of $p_T^{\min}(B_c)$ by integrating the p_T spectra. Table 1 gives the inclusive cross sections for the B_c meson at the Tevatron as a function of $p_T^{\min}(B_c)$, including $n = 1$ and $n = 2$ S- and P-wave state contributions. The variations versus the scale μ between $\mu_R/2$ and $2\mu_R$ are always within a factor of two, and are rather insensitive to changes in scale when $p_T^{\min}(B_c) \gtrsim 10$ GeV.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Figure 2: Same as Fig.1 for $n = 2$.

At the end of Run Ib at the Tevatron, the total accumulated luminosities can be up to 100–150 pb^{-1} or more. With $p_T > 6$ GeV, there are about 5×10^5 B_c^+ mesons. In the future, when the Main Injector is installed in the Run II, which can accumulate 1–2 fb^{-1} luminosity, there will be of order 10^7 B_c mesons. At the LHC there will be about 3×10^9 B_c mesons with $p_T > 10$ GeV at the assumed 100 fb^{-1} luminosity.

In conclusion, there should be enough signature events to confirm the existence of B_c at the Tevatron, and the LHC will be a copious source of B_c . This work was supported by US DOE-FG03-93ER40757.

1. E. Eichten and C. Quigg, Phys. Rev. **D49**, 5845 (1994).
2. E. Braaten, K. Cheung, and T. C. Yuan, Phys. Rev. **D48**, R5049 (1993).
3. T. C. Yuan, Phys. Rev. **D50**, 5664 (1994).
4. K. Cheung, Phys. Rev. Lett. **71**, 3413 (1993).

Table 1: Inclusive production cross sections for the B_c meson at the Tevatron including the contributions from all the S-wave and P-wave states below the BD threshold as a function of $p_T^{\min}(B_c)$. The acceptance cuts are $p_T(B_c) > 6$ GeV and $|y(B_c)| < 1$.

$p_T^{\min}$ (GeV)	σ (nb)		
	$\mu = \frac{1}{2}\mu_R$	$\mu = \mu_R$	$\mu = 2\mu_R$
6	2.81	5.43	6.93
10	0.87	1.16	1.22
15	0.26	0.29	0.26
20	0.098	0.097	0.083
30	0.021	0.018	0.014

5. K. Cheung and T. C. Yuan, Phys. Lett. **B325**, 481 (1994).
6. K. Cheung and T.C. Yuan, preprint UCD-95-4 and CPP-94-37 (Feb 1995).

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.