On the radial structure of fluctuations and turbulence induced flows

C. Hidalgo, M.A. Pedrosa, B. van Milligen, E. Sánchez, R. Balbín,
I. García-Cortés
Euratom-Ciemat, Madrid, Spain

G. Batanov, O. Fedyanin, N. Kharchev, Yu. Kholnov, S. V.
Shchepetov, K. Sarksyán, N.N. Skvortsova
General Physics Institute, Moscow, Russia

L. García
Carlos III University, Madrid, Spain

B. A. Carreras
Oak Ridge National Laboratory, Oak Ridge, USA

RECEIVED

OCT 27 1997

OSTI

Abstract

Comparative studies of the structure of turbulence have been carried out in stellarator plasmas with low (TJ-IU) and medium (L2-M) magnetic shear. In both devices, the value of the radial coherence of fluctuations is bursty. In the plasma edge region, the radial coherence is dominated by low frequency fluctuations (< 200 kHz) in TJ-IU, whereas in L2-M, the radial coherence is due to high frequency fluctuations (> 100 kHz). Numerical calculations of resistive ballooning turbulence show that in the non-linear phase, the correlation length is of the order of the width of the linear mode poloidal components, whereas in the linear phase, the correlation length is related to the width of the envelope. Toroidal coupling effects appear to be relevant in the nonlinear phase in the proximity of instability thresholds. The electrostatic Reynolds stress shows a radial gradient close to the velocity shear location in the TJ-IU torsatron, thus suggesting that this mechanism can drive poloidal flows.

Keywords:

stellarator, anomalous transport, flows, L-H transition

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

1 Introduction

A relevant issue for the question, whether transport exhibits a Bohm or gyro-Bohm scaling, is the understanding of the mechanisms that determine the radial correlation length of fluctuations. The closeness to turbulence thresholds [1], the type of instability driving the turbulence and the presence of sheared flows [2] are considered critical elements to determine the transition between different transport regimes as well as to account for a rapid change of transport coefficients [3].

In the framework of the physics of poloidal flows and fluctuations, it has been argued that the average poloidal flow profile can be modified by plasma fluctuations via radially varying Reynolds stress [4-5]. This mechanism might be involved in the L-H transition.

The goal of the present investigation is threefold. Firstly, to compare the statistical properties and the radial structure of fluctuations in the plasma boundary region of stellarator plasmas with low (TJ-IU) and medium (L2-M) magnetic shear. Secondly, the investigation of the radial structure of resistive ballooning turbulence using numerical calculations for both the linear and non-linear phase versus beta. Finally, to investigate flows driven by fluctuations (Reynolds stress) in the TJ-IU plasma boundary region.

2 Comparative studies of plasma turbulence in stellarator plasmas

Measurements of fluctuations and turbulent flux have been carried out in the plasma boundary region of the TJ-IU torsatron ($P_{ECRH}=200$ kW, $iota(0)\approx 0.23$, $R=0.6$ m, $-a\approx 0.1$ m, $-n_e\approx 0.5\times 10^{19}$ m $^{-3}$, $B_t=0.67$ T) [6] and of the L2-M stellarator ($P_{ECRH}=200$ kW, $iota(0)\approx 0.20$, $iota(a)\approx 1$, $R=1.0$ m, $-a\approx 0.11$ m, $-n_e\approx 1.5\times 10^{19}$ m $^{-3}$, $B_t=1.2$ T) [7]. Turbulence has been investigated using Langmuir probe arrays and signal processing tools described elsewhere [8].

In both devices, the ion saturation density current increases and the floating potential becomes more negative as the probe is inserted into

the plasma edge region. As observed in other devices, the radial electric field is sheared in the proximity of the (computed) location of the last closed flux surface (LCFS). In the L2-M stellarator the level of ion saturation current fluctuations is of the order $iI_s / I_s \approx 5\%$ for $r/r_{LCFS} < 0.8 - 0.9$ and about 20 % near the location of the LCFS. In the TJ-IU edge plasma region, the level of fluctuations is $iI_s / I_s \approx (15 - 20)\%$, which is about a factor of three higher than in L2-M.

In the TJ-IU torsatron, the value of the radial coherence of fluctuations decreases when increasing radial distance between probes. The value of the radial coherence associated with fluctuations and turbulent transport is intermittent, and the turbulent flux exhibits a smaller radial coherence than the one associated with fluctuations. The statistical properties of the probability distribution function of fluctuations in pressure gradients have been also investigated. To date, evidence has not been found for non-gaussian features in the plasma gradient fluctuation distribution that are related with the presence of critical gradients.

In the TJ-IU torsatron, the radial and poloidal frequency resolved coherence of fluctuations are quite similar (i.e., isotropic in terms of frequencies) and dominated by frequencies below 200 kHz (Fig. 1). However, in the L2-M stellarator, the radial and poloidal coherence structure of the fluctuations are different at the plasma edge, the radial coherence being dominated by high frequencies (larger than 100 kHz) (Fig. 2) and the poloidal coherence dominated by low frequencies (less than 100 kHz). In addition, in the TJ-IU torsatron, the apparent radial velocity of fluctuations is about 1 km/s, whereas in the L2-M stellarator, it is much higher (20 - 40 km/s). These results can be interpreted in terms of the influence of linear and non-linear coupling effects on turbulence [9].

3 Numerical calculations of resistive ballooning turbulence

The results of numerical calculations of resistive ballooning turbulence show that the correlation length in the nonlinear phase is of the order of the width of the linear mode poloidal components, as

opposed to the linear phase, where the correlation length is related to the width of the envelope. However, in low beta plasmas, toroidal coupling effects appear to be relevant in the nonlinear phase, and as a consequence the radial correlation increases [10]. A combined (experimental/theoretical) effort is in progress to investigate the statistical properties of the radial structure of fluctuations versus the proximity to the turbulent thresholds for different plasma instabilities (e.g., pressure gradient driven instabilities).

4 Flows driven fluctuations

The radial profile of the electrostatic Reynolds stress has been measured in the plasma boundary region of the TJ-IU torsatron using a multiple array of Langmuir probes. The spatial orientation of the Langmuir probe array provides a measurement of radial (iE_r) and poloidal (iE_θ) electric field fluctuations in a plasma volume smaller than the typical correlation volume of fluctuations. There is a significant reduction in the radial coherence of fluctuations in the vicinity of the velocity shear layer location, whereas the poloidal coherence of fluctuations and the root mean squared (rms) level of fluctuations change slightly in the region close to the shear location. The measured Reynolds stress (proportional to $\langle iE_\theta iE_r \rangle$) shows a radial variation around the velocity shear location, suggesting that such a mechanism can drive poloidal flows in the edge region of the TJ-IU torsatron (Fig. 3). On the contrary, the magnetic component of the Reynolds stress is negligible, due to the relative phase ($\approx \pi/2$) between poloidal and radial fluctuating components of the magnetic field [11]. Further studies are in progress to quantify the importance of Reynolds stress induced flow as a drive for poloidal rotation, a mechanism which might be involved in the L-H transition.

Acknowledgments

The authors would like to acknowledge stimulating discussions with the members of the European and U.S. Transport Task Force Turbulence Working Group.

References

- [1] X. Garbet and R. E. Waltz, Phys. Plasmas **3**, 1898 (1996).
- [2] K. Burrel, Phys. PLasmas **4**, 1499 (1997)
- [3] K. Gentle et al., Phys. Rev. Lett. **74**, 3620 (1995).
- [4] P.H. Diamond and Y. B. Kim, Phys. Fluids B **3**, 1626 (1991)
- [5] B. A. Carreras et. al., Phys. Fluids B **5**, 1491 (1993).
- [6] E. Ascasibar et al., in *Plasma Physics and Controlled Nuclear Fusion Research* (Proc. 15th Int. Conf. Seville, 1994), Vol 1, IAEA, Vienna (1995) p. 749.
- [7] V.V. Abrakov, D. K. Akulina et al., *Proceedings of the 10th International Conference on Stellarators*, EUR-CIEMAT **30**, 10 (1995).
- [8] M. A. Pedrosa et al., Proc. 23rd EPS Conf. on Controlled Fusion and Plasma Physics (Kiev) vol 20C, part II (1996) p 827.
- [9] G. M. Batanov et al., submitted for publication (1997).
- [10] L. García et al., Proc. 24rd EPS Conf. on Controlled Fusion and Plasma Physics (Berchtesgaden) (1997) (in press).
- [11] C. Hidalgo et al., submitted for publication (1997).

Figure Captions

Fig. 1 (a) Frequency spectrum and (b) radial coherence of fluctuations in the plasma edge region of the TJ-IU torsatron.

Fig. 2 (a) Frequency spectrum and (b) radial coherence of fluctuations in the plasma edge region of the L2-M stellarator.

Fig. 3 Radial profile of the electrostatic Reynolds stress in the plasma boundary region of the TJ-IU torsatron.

Fig. 1.- (a) Frequency spectrum and (b) radial coherence of fluctuations in the plasma edge region of the TJ-IU torsatron.

Fig. 2.- (a) Frequency spectrum and (b) radial coherence of fluctuations in the plasma edge region of the L2-M stellarator.

Fig. 3.- Radial profile of the electrostatic Reynolds stress in the plasma boundary region of the TJ-IU torsatron.

M98000679

DOE

⑱ DOE, XF

⑲ UC-900, DOE

19971202 076