2/17/86 W.B.

①

I-20494

DR-0957-1

UCRL-53574

Reaction of the Topopah Spring Tuff With J-13 Water at 120°C

V. M. Oversby

MASTER

July 18, 1984

Lawrence
Livermore
National
Laboratory

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Reaction of the Topopah Spring Tuff With J-13 Water at 120°C

V. M. Oversby

Manuscript date: July 18, 1984

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

LAWRENCE LIVERMORE LABORATORY
University of California • Livermore, California • 94550

Available from: National Technical Information Service • U.S. Department of Commerce
5285 Port Royal Road • Springfield, VA 22151 • \$8.50 per copy • (Microfiche \$4.50)

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *xb*

Contents

Abstract	1
Introduction	1
Experimental Procedure	2
Results and Discussion	7
Composition of the Caliche Component	7
Test Matrix Results at 120°C	18
Estimation of Steady-State Water Chemistry	27
Conclusions	28
Acknowledgments	28
References	29
Appendix A	A-1

Reaction of the Topopah Spring Tuff With J-13 Water at 120°C

Abstract

The Nevada Nuclear Waste Storage Investigations (NNWSI) Project is examining the suitability of the Topopah Spring Member of the Paintbrush Tuff for potential development as a high level nuclear waste repository. As part of the NNWSI Project, the Lawrence Livermore National Laboratory (LLNL) is responsible for the design and testing of waste packages suitable for use in the Topopah Spring tuff at Yucca Mountain. Definition of the physical and chemical environment of the waste package is part of that task. This report describes a series of hydrothermal experiments using crushed tuff from the Topopah Spring Member and natural ground water from well J-13. The purpose of these experiments is to define the changes in water chemistry that would result from temperature changes caused by emplacing high level nuclear waste in a repository in the Topopah Spring tuff.

Experiments were conducted at 120°C in Teflon-lined reaction vessels at four separate rock-to-water ratios and for reaction times up to 72 days. The composition of evaporite deposits contained in the pores of the surface-outcrop rock material used in these experiments is determined from solution compositions resulting from treatment of the rock before the start of the experiments. Results from the experiments at 120°C are compared with previous experimental results from hydrothermal reaction of the Topopah Spring tuff with J-13 water at 90 and 150°C.

The main conclusion that can be drawn from this work is that changes in the water chemistry due to heating of the rock-water system can be expected to be very minor. There is no significant source of anions (F^- , Cl^- , NO_3^- , or SO_4^{2-}) in the rock; solution anion compositions after reaction of pretreated rock with J-13 water differ very little from the starting compositions. The major changes in cations are an increase in silica to approximately the level of cristobalite solubility, supersaturation of aluminum followed by slow precipitation, and fairly rapid precipitation of calcium and magnesium due to the retrograde solubility of calcite. These results are in good agreement with those previously reported for reaction of the tuff with J-13 water at 90 and 150°C.

Introduction

Lawrence Livermore National Laboratory is developing designs for high level nuclear waste packages as part of the NNWSI Project. The potential repository site under study is located at Yucca Mountain in Nye County, Nevada; the reference repository horizon is in the Topopah Spring Member of the Paintbrush Tuff. At the proposed repository level the tuff is densely welded and devitrified. The rock consists predominantly of a fine-grained intergrown assemblage of quartz, cristobalite, and alkali feldspar, with a small proportion of phenocrysts. Phenocryst minerals are quartz, alkali feldspar, plagioclase, and biotite.

Performance objectives for high-level nuclear waste packages, as given in the Nuclear Regula-

tory Commission's regulations (10 CFR Part 60), require prediction of the corrosion performance of metals for up to 1000 years, and of the performance of waste forms for up to 10,000 years. Successful prediction of performance for these long time periods requires a thorough understanding of the physical and chemical environmental conditions to which the packages will be subjected. Definition of the waste package environment involves detailed description of the pre-emplacment (ambient) conditions at the proposed repository horizon and determination of the changes that will result after emplacement of waste packages.

Construction of a high-level waste repository can produce three types of changes in the repository and surrounding environment. First, mining

causes physical changes in the rock unit. Second, the radiation field produced by the waste can cause potential environmental changes. Third, the thermal output of the waste will heat the surroundings, causing physical and chemical changes in the repository rock and water systems. This report addresses the chemical changes caused by heating of repository rock with representative ground water.

The experiments were conducted using Topopah Spring tuff (Tpt) collected from Fran Ridge, several miles east of Yucca Mountain, and water from well J-13. The locations of the potential repository site at Yucca Mountain, Fran Ridge, and J-13 are shown in Fig. 1. The elevation is lower at J-13 than at Yucca Mountain, and the Tpt lies below the water table. The Tpt is the major producing horizon for J-13; thus, the water chemistry is probably close to the prevailing chemistry in the Tpt at Yucca Mountain. At Yucca Mountain, the Tpt lies well above the water table in the unsaturated zone. We have not yet obtained water samples from this region, but collection of samples is planned as part of the exploratory shaft testing program. Until samples of water are available from the unsaturated repository horizon, the water from well J-13 has been adopted as a reference water chemistry for NNWSI experimental work.

The expected environmental conditions in the unsaturated zone limit the temperature at which

liquid water will exist. For the expected case with no substantial sealing of fractures and pores in the host rock, the local atmospheric boiling point of unconfined water is approximately 95°C. Water contained in pores is held by capillary forces; this water might have a somewhat higher effective boiling point. To bracket the expected temperature range, and to provide data for geochemical modeling studies, hydrothermal experiments are being conducted over a range of temperatures. The experiments described in this report were conducted at 120°C. Results from experiments conducted at 90 and 150°C have already been reported (Oversby, 1984).

The experiments described here used crushed rock and Teflon-lined reaction vessels. The techniques used required quenching of solutions before separating the rock and water after reaction. Other experiments are in progress that use gold-bag rocking autoclaves, which allow sampling at temperature. Some experiments also in progress or recently completed use core wafer Tpt tuff, both from outcrop samples and from drill cores. The core wafer experiments allow identification of alteration products formed in the rock as well as identification of secondary phases formed by precipitation from solution. Results from these experiments will be the subject of separate reports.

Experimental Procedure

The water used in these experiments was obtained from well J-13 at the Nevada Test Site (shown in Fig. 1). Water sampling and storage procedures are described in Oversby (1984).

The rock samples were collected from an outcrop located at Fran Ridge (Fig. 1). A detailed description of the locality is given by Knauss (1984). Procedures used to prepare the rock for experiments are described by Oversby (1984). The rock material was less than 60 mesh size (250 micron) and was from split number A-3.

All experiments were conducted in Teflon capsules encased in metal containers. Standard Parr Acid Digestion Bombs (model number 4748) were used. They were cleaned at 120°C before use by the procedure given in Oversby (1984).

The material obtained from surface outcrops near Yucca Mountain contains readily soluble material that is not present in rocks protected by a significant amount of overlying rock (Oversby, 1983).

The soluble material can be easily removed by shaking the rock powder in water for two minutes, allowing the powder to settle, and decanting the solution. A further step of cooking overnight with J-13 water ensures substantially complete removal of the soluble material. The rock samples were pretreated using the room-temperature shaking step and an overnight cook in J-13 water at 120°C. The chemical compositions of these rinse solutions, which are discussed in the following section of this report, can be used to determine the composition of the soluble component. A detailed description of the pretreatment procedure is given in Oversby (1984).

At the start of the experiments J-13 water was weighed into the Teflon capsules containing pretreated rock. The amount of water used was kept constant at 48 g for all experiments; rock weights of 0.4, 0.8, 1.6, and 3.2 g were used. After water was added to the capsule, it was closed, placed in

Figure 1. Map showing the location of Yucca Mountain, Fran Ridge, and well J-13. The outcrop locality for rock samples used in this work is marked Tpt-FR. Tcfb-TS is the outcrop locality for Bullfrog tuff samples. Other localities marked are drill holes.

the outer metal container, and the assembly was placed in a Blue M oven that had been preheated to 120°C. The assembly remained in the oven for a reaction time of 1, 3, 6, 12, 24, 36, 48, 60, or 72 days.

At the end of the predetermined reaction time the assembly was removed from the oven, opened as soon as it was cool enough to handle (approximately 50°C), and the contents were allowed to cool to room temperature. The pH of the liquid was measured with a meter that had been calibrated using standard solutions of pH 7 and 10. Following pH measurement, the solution was filtered through #40 filter paper and then through a 0.1-micron filter. The filtered solution was split into four portions: (1) for anion analysis, stored in plastic, (2) for alkalinity, stored in glass, (3) and (4) acidified with two drops of 50% HNO₃ per 6 ml of sample, and stored in plastic for later analysis by inductively coupled plasma (ICP) spectrometry. The rock sample was recovered on the filter paper, dried in air, and transferred to a small storage bottle when dry.

Cation analyses were performed using an automated ICP spectrometer system. The system, detailed in Peck et al. (1979), uses a polychromator operated in the fixed-channel mode, giving simultaneous analysis for 28 elements. It is calibrated with standard solutions, and the operator employs a "check standard" that is run as a regular sample during each series of analyses. During the course of these experiments some difficulties were encountered with results for boron and potassium. These difficulties were identified through the use of control blank samples of J-13 water that are regularly submitted along with the hydrothermal test solutions. The control blanks of J-13 are given identification numbers that make them appear to the ICP operator to be part of the hydrothermal test set.

Table 1 gives a summary of results on J-13 control blank samples and one set of six J-13 analyses run by the ICP operator to confirm that the boron and potassium analysis problems had been corrected. These samples (Table 1b) were known by the operator to be J-13. The boron problem occurred in analyses performed on 8/4/83, and in all analyses during the period from December 1983 through March 1984. The boron content for control blanks run during this period was reported as below detection limits. The problem was subsequently identified as having been caused by incorrect data storage and reduction in the computer control system. This resulted in reported concentrations for boron that were systematically low by about 0.1 ppm. Since this is approximately the bo-

ron concentration in J-13, results for control blanks were reported as near or below the detection limit (i.e., close to zero).

The boron problem was corrected in May 1984, as can be seen in Table 1a. These analyses revealed a problem with potassium that had not previously been encountered. The potassium concentration of J-13 is about 5 ppm; the analyses of control blanks run on 5/17/84 gave approximately 3 ppm. In addition, results for most of the other elements are slightly off the expected values. These results were also affected by a problem with data storage and manipulation in the computer. This problem was corrected during June 1984. The J-13 analyses run on 6/13/84 give correct results for all elements (Table 1b).

Control blanks run on 6/22/84 give the expected results for J-13 except for No. 6. The results for most elements in this sample are erratic and indicate hardware problems during the analysis. This sample has been excluded from the data set used to obtain the average for J-13 samples run on 6/22/84.

A summary of all J-13 control blank analyses run on the polychromator ICP are given in Table 1d. The summary does not include the set run on 5/17/84 since these analyses are suspect for all elements. Details of analyses for the period from July 1983 through March 1984 are given in Oversby (1984).

During the course of these experiments a new monochromator ICP was brought on-line. To check on the precision and accuracy of the new instrument, a split of each of the solution samples for the rock-water matrix at 120°C was analyzed on both the new and old ICP units. A series of control blanks of J-13 water was submitted to the new ICP facility along with the solution samples. Data for the J-13 analyses are given in Table 1e; results for sample solutions are compared with those obtained on the polychromator ICP in Appendix A. Results for most elements on the monochromator ICP are slightly higher than those obtained from the polychromator unit. The exception is boron, which is lower in the new ICP results.

Anion analyses were performed using a Dionex Model 2100 Ion Chromatograph with a "fast run" AS4 separator column and an anion fiber suppressor. Detection limits are 0.1 ppm for Cl⁻ and F⁻ and 0.2 ppm for NO₃⁻ and SO₄⁻. Alkalinity measurements were made using a Technicon Autoanalyzer, an automated version of the standard total alkalinity titration. This method measures the sum of the OH⁻, HCO₃⁻, CO₃⁻, and any other titratable anions (such as H₃SiO₄⁻) in solu-

Table 1. Data for ICP analyses of J-13 water samples. Concentration in ppm.**a. Four J-13 water samples run on 5/17/84 — control blanks.**

No.	Al	B	Fe	Si	Ca	K	Mg	Na	Li
1	0.006	0.145	0.006	28.7	14.2	3.2	2.05	49.4	0.040
2	<0.002	0.145	0.003	28.7	14.2	2.8	2.06	49.6	0.039
3	<0.002	0.148	0.011	28.5	14.1	3.0	2.07	50.8	0.039
4	<0.002	0.146	<0.001	28.8	14.2	2.5	2.08	50.6	0.039
Average	0.002	0.146	0.005	28.7	14.2	2.9	2.07	50.1	0.039
Std dev.	0.003	0.001	0.004	0.11	0.04	0.3	0.011	0.61	0.0004

b. Six J-13 water samples run on 6/13/84, known by analyst to be J-13.

No.	Al	B	Fe	Si	Ca	K	Mg	Na	Li
1	bdl ^a	0.137	<0.002	27.3	12.7	5.05	1.92	44.2	0.041
2	bdl	0.141	0.002	27.4	12.8	4.94	1.92	44.8	0.042
3	bdl	0.137	0.002	27.2	12.7	5.17	1.91	44.6	0.042
4	bdl	0.137	0.003	27.4	12.8	5.30	1.92	45.1	0.042
5	bdl	0.139	0.012	27.3	12.8	4.56	1.92	45.0	0.042
6	bdl	0.137	0.006	27.2	12.8	5.34	1.92	44.6	0.042
Average		0.138	0.005	27.3	12.8	5.06	1.92	44.7	0.042
Std dev.		0.002	0.004	0.08	0.05	0.26	0.004	0.30	0.0004

c. Six J-13 water samples run on 6/22/84 — control blanks.

No.	Al	B	Fe	Si	Ca	K	Mg	Na	Li
1	0.022	0.135	0.012	26.6	12.5	4.94	1.91	44.0	0.042
2	0.021	0.138	0.010	26.6	12.5	4.60	1.90	44.0	0.041
3	0.017	0.138	0.014	26.8	12.7	5.28	1.92	44.9	0.043
4	0.021	0.138	0.026	26.6	12.6	5.37	1.92	44.7	0.043
5	0.012	0.134	0.013	26.5	12.7	4.41	1.91	44.9	0.042
6	0.032	0.140	0.069	27.3	12.3	4.02	1.86	43.3	0.041
Average	0.019	0.137	0.015	26.6	12.6	4.92	1.91	44.5	0.042
Std dev.	0.004	0.002	0.006	0.10	0.09	0.37	0.007	0.41	0.001

d. Average of all data for J-13 cation analyses, excluding 5/17/84.

Date	Al	B	Fe	Si	Ca	K	Mg	Na	Li
7/19/83	0.009	0.121	0.009	27.0	11.9	5.46	1.88	43.4	
8/4/83	0.003		0.002	26.2	13.5	5.19	2.07	44.9	
9/13/83	0.001	0.120	0.002	27.0	12.8	4.54	1.91	43.3	
10/24/83	0.013	0.125	0.008	26.9	13.3	4.89	1.93	43.6	
12/6/83	0.012		0.004	27.5	12.5	5.09		45.5	
12/16/83	0.012		<0.002	27.5	12.5	5.09		45.5	
Av2/15/84	0.040		0.015	26.2	12.2	4.73	1.90	41.2	
3/7/84	0.009		0.004	27.8	10.5	5.67	1.86	43.3	
Av3/7/84	0.011		0.004	27.0	13.0	5.51	1.92	43.4	
Av6/13/84	<0.002	0.138	0.005	27.3	12.8	5.10	1.92	44.7	0.042
Av6/22/84	0.019	0.137	0.015	26.6	12.6	4.92	1.91	44.5	0.042
Average	0.012	0.128	0.006	27.0	12.5	5.11	1.92	43.9	0.042
Std. dev.	0.010	0.008	0.005	0.49	0.77	0.32	0.06	1.19	

Table 1. (Continued)

e. Data for new ICP; four control blanks run on 3/27/84.

No.	Al	B	Fe	Si	Ca	Mg	Na
1	bdl ^a	0.11	0.028	27.5	13.8	2.11	51.3
2	bdl	0.10	0.026	27.0	13.3	2.11	46.8
3	bdl	0.11	0.030	26.7	13.8	2.08	48.7
4	bdl	0.12	0.025	26.7	13.9	2.10	46.5
Average	dl ^b = .028	0.11	0.027	27.0	13.7	2.10	48.3
Std dev.		0.007	0.002	0.33	0.23	0.01	1.91

^a bdl = below detection limit.^b dl = detection limit.

Table 2. Data for anion analyses of J-13 control blanks.

a. Anions measured by ion chromatography, ppm.

Date	F ⁻	Cl ⁻	NO ₃ ⁻	SO ₄ ²⁻
8/30/83	1.8	5.3	6.4	13.4
9/30/83	2.3	6.8	9.3	19.8
Av3/14/84	2.3	6.6	9.0	18.1
Av3/26/84	2.4	7.1	9.2	18.5
3/30/84	2.3	6.8	18.0	19.0
4/2/84	2.4	7.0	9.1	19.0
4/3/84	1.3	7.4	13.1	18.6
5/4/84	2.3	6.9	7.0	18.4
5/4/84	2.3	6.9	6.4	18.5
5/10/84	2.2	6.9	5.4	18.4
Average	2.2	6.8	9.3	18.2
Std dev.	0.33	0.53	3.57	1.65
Reaverage	2.2	6.9	9.6	18.7
New std dev	0.32	0.21	3.63	0.47

b. Alkalinity by Technicon Autoanalyzer.

Date	meq/l	HCO ₃ ⁻ (ppm)
7/27/83	2.29	140
8/5/83	2.13	130
5/2/83	2.16	132
1/11/84	2.46	150
3/13/84	2.10	128
Average	2.23	136
Std dev.	0.13	8.1

tions. For the pH range and silicon concentrations of samples resulting from these experiments, the contribution of silica species to alkalinity should be minor.

Table 2 gives a summary of results for anion and alkalinity measurements on control blank samples of J-13 water. Results obtained through 3/26/84 have been previously reported and discussed by Oversby (1984). Data for anion analyses

on 8/30/83 are systematically low and have been eliminated to obtain the average value for J-13 (labelled Reaverage in Table 2a). Analyses for nitrate were erratic during the period from 3/30/84 through 5/10/84. The reason for this problem is being investigated. Only four samples for which results are reported here were analyzed during 1984.

The control blank samples are run in the same sample sets as the solutions from the rock-water hydrothermal reaction experiments. The precision of the analytical results for the control blanks provides a good measure of the precision of the analytical results for the experimental solutions.

Results and Discussion

Composition of the Caliche Component

The rock samples used in the experiments were pretreated to remove the readily soluble salts that are associated with rock material collected from surface outcrops near Yucca Mountain. The composition of soluble salts removed from the Fran Ridge outcrop Topopah Spring tuff used in previous experiments was reported by Oversby (1984). Rock used in the present experiments was from a companion split of the same crushing and, as such, was expected to have the same quantity and type of soluble material. The rinse solution used to pretreat the 120°C samples was 48 ml of J-13 water in all cases. Table 3 gives the anion concentrations found in the room temperature rinsing step; data for anion concentrations from the overnight cooked rinse at 120°C are given in Table 4.

There is no indication that any fluoride or carbonate material is released to solution during either of the pretreatment rinsing steps. In fact, alkalinity decreases by approximately 10% during the room temperature rinse and remains essentially constant or decreases slightly during the cooked rinse.

The dominant anion components of the soluble material are nitrate and sulfate. There is also a chloride component, but it is present in much lower amounts. Anion concentrations in the rinse solutions are approximately linear with sample weight, as can be seen in Fig. 2, which plots the averages for chloride, nitrate, and sulfate concentrations from room temperature rinse solutions as a function of rock sample weight. This is the behavior to be expected if the anions are due to complete solution of a readily soluble phase or phases.

The quantity of anions released during the overnight cooked rinse is substantially smaller. A portion of these anions is undoubtedly due to retention of some of the room temperature rinse solution on the rock when that solution was removed by decantation.

The standard deviation quoted is that of the population; for a single analysis the appropriate value would be slightly higher, the difference amounting to a few percent of the value of the standard deviation in most cases.

The total amount of anions removed from the rock used in these experiments is slightly higher than the anions found on rock material from split A-4, which was used in the 90 and 150°C experiments (Oversby, 1984). The difference in amount is very small and, as will be seen below, correlates with a slightly greater amount of calcium, potassium, and sodium in the rinse solutions from split A-3 material.

Data for the pH of room temperature and 120°C rinse solutions are given in Table 5. The pH of the rinse solutions is systematically higher than that of J-13 water, and shows a positive correlation with rock sample weight. None of the anion components identified in the rinse solutions is a weak acid; therefore, the change in pH must be due to a component analyzed by ICP and reported as a cation.

Tables 6 through 13 give data for rinse solutions measured by ICP. The cooked rinse data are affected by reaction with the rock at 120°C, and therefore cannot be used to unambiguously identify the nature of the soluble component. Data for aluminum in the room temperature rinse solutions are slightly erratic for the 1.6- and 3.2-g samples; however, total aluminum release is small in all cases and is slightly less than that found for split A-4 (Oversby, 1984).

Boron release (Table 7) is linear with sample weight in both the room temperature and cooked rinse solutions. Boron can form a weak acid; this may be the cause of the pH change found in the rinse solutions. Boron was not a component of the soluble material associated with split A-4 (Oversby, 1984).

Iron concentrations (Table 8) are essentially constant at J-13 values for both the room temperature and cooked rinse solutions. Silicon (Table 9) is slightly lower than the J-13 concentration in the room temperature rinse, while it is higher in the cooked rinse solution due to reaction of the water with the rock at 120°C. Both of these results are the

Table 3. Anion concentrations in ppm from room temperature rinse solutions. Sample identification is final run time followed by sample weight in grams.

Sample ID	F ⁻	Cl ⁻	NO ₃ ⁻	SO ₄ ⁻²	Alkalinity	
					meq/l	HCO ₃ ⁻ (ppm)
J-13	2.2	6.9	9.6	18.7	2.23	136
1-0.4	2.4	8.9	19.3	28.0	2.03	124
3-0.4	2.2	8.9	18.8	21.9	2.04	124
6-0.4	2.2	8.4	18.5	25.8	2.10	128
12-0.4	2.2	8.3	19.4	25.1	1.99	121
24-0.4	2.2	8.5	19.0	25.5	2.01	123
36-0.4	2.4	8.7	19.5	26.5	2.10	128
48-0.4	2.2	7.9	19.3	28.4	1.95	119
60-0.4	2.1	8.2	16.4	25.0	2.09	127
72-0.4	2.3	8.5	19.9	26.9	2.02	123
Average	2.2	8.5	18.9	25.9	2.04	124
Std dev.	0.10	0.31	0.96	1.81	0.05	3
1-0.3	2.1	10.6	29.4	35.5	2.13	130
3-0.8	2.2	10.6	29.2	33.4	1.97	120
6-0.8	2.2	10.1	28.1	32.8	2.09	127
12-0.8	2.3	9.8	29.3	33.5	2.09	127
24-0.8	2.2	9.8	28.7	33.0	2.01	123
36-0.8	2.1	10.0	27.5	32.1	2.10	128
48-0.8	2.2	10.4	28.6	35.6	1.98	121
60-0.8	2.2	10.7	29.6	35.8	2.09	127
72-0.8	2.3	10.0	29.5	32.2	1.99	121
Average	2.2	10.2	28.9	33.8	2.05	125
Std dev.	0.07	0.34	0.67	1.40	0.06	4
1-1.6	2.2	13.4	48.0	46.7	1.99	121
3-1.6	2.3	13.4	48.1	45.8	2.06	126
6-1.6	2.1	12.4	44.6	43.9	2.13	130
12-1.6	2.3	12.9	49.3	46.5	1.94	118
24-1.6	2.3	12.8	48.0	45.2	2.09	127
36-1.6	2.1	12.8	44.1	43.5	2.05	125
48-1.6	2.2	13.1	45.9	46.1	2.08	127
60-1.6	2.2	13.4	48.3	47.1	2.10	128
72-1.6	2.3	12.7	45.8	45.5	2.11	129
Average	2.2	13.0	46.9	45.6	2.06	126
Std dev.	0.07	0.34	1.73	1.16	0.06	4
1-3.2	2.3	18.2	91.1	86.1	2.16	132
3-3.2	2.5	26.4	86.3	75.5	2.08	127
6-3.2	2.2	19.3	78.6	67.5	2.06	126
12-3.2	2.3	18.1	81.1	70.4	2.01	123
24-3.2	2.3	19.4	86.4	76.4	2.06	126
36-3.2	2.2	19.4	80.4	65.1	2.25	137
48-3.2	2.3	20.3	81.1	72.3	2.15	131
60-3.2	2.3	18.4	68.6	53.4	2.17	132
72-3.2	1.3	17.8		74.8	2.13	130
72-3.2B	2.3	17.1	78.1	67.3		
Average	2.2	19.4	81.3	73.9	2.12	129
Std dev.	0.31	2.49	6.03	6.51	0.07	4

Table 4. Anion concentrations in ppm from overnight cooked rinse solutions. Sample identification is final run time followed by sample weight in grams.

Sample ID	F ⁻	Cl ⁻	NO ₃ ⁻	SO ₄ ⁻	Alkalinity	
					meq/l	HCO ₃ (ppm)
J-13	2.2	6.9	9.6	18.7	2.23	136
1-0.4	2.3	7.9	11.5	20.9	2.05	125
3-0.4	2.2	7.1	5.1	18.4	2.06	126
6-0.4	2.2	7.0	10.5	18.7	2.08	127
12-0.4	2.3	7.3	10.9	19.6	2.11	129
24-0.4	2.3	7.1	10.8	18.9	2.09	127
36-0.4	2.3	7.2	11.1	19.5	2.08	127
48-0.4	2.3	8.1	11.6	21.3	2.05	125
60-0.4	2.0	6.2	9.4	16.8	2.17	132
72-0.4	2.2	7.2	10.7	19.5	2.18	133
Average	2.2	7.2	10.2	19.3	2.10	128
Std dev.	0.09	0.51	1.89	1.26	0.05	3
1-0.8	2.3	8.3	13.5	22.0	2.13	130
3-0.8	2.2	7.5	10.9	18.9	2.06	126
6-0.8	2.0	6.8	11.3	18.1	2.22	135
12-0.8	2.3	7.7	13.2	13.9	2.18	133
24-0.8	3.4	7.7	13.8	32.1	2.11	129
36-0.8	2.3	7.9	15.9	21.7	2.20	134
48-0.8	2.2	8.5	28.6	39.1	2.13	130
60-0.8	2.5	8.4	15.4	23.8	2.18	133
72-0.8	2.3	7.6	13.0	20.0	2.18	133
Average	2.4	7.8	15.0	23.4	2.15	131
Std dev.	0.38	0.70	5.03	7.39	0.05	3
1-1.6	2.3	9.0	17.0	24.0	2.13	130
3-1.6	2.3	8.3	14.1	21.5	2.07	126
6-1.6	2.4	8.6	17.4	23.2	2.14	131
12-1.6	2.3	8.6	16.9	23.0	2.19	134
24-1.6	2.3	8.3	16.9	22.7	2.19	134
36-1.6	2.5	8.7	18.4	24.4	2.15	131
48-1.6	2.2	8.9	16.4	23.4	2.22	135
60-1.6	2.3	8.1	16.0	22.3	2.18	133
72-1.6	2.5	8.8	18.3	24.0	2.23	136
Average	2.3	8.6	16.8	23.2	2.17	132
Std dev.	0.10	0.28	1.22	0.87	0.05	3
1-3.2	2.3	10.8	26.8	31.6	2.21	135
3-3.2	2.3	9.9	23.0	25.2	2.07	126
6-3.2	2.1	10.0	23.7	27.4	2.26	138
12-3.2	2.4	10.5	27.6	29.8	2.18	133
24-3.2	2.4	10.1	27.5	27.3	2.11	129
36-3.2	2.4	11.2	28.9	33.3	2.18	133
48-3.2	2.4	10.7	25.4	29.8	2.23	136
48-3.2B	2.3	10.4	25.9	26.7		
60-3.2	2.5	11.0	26.8	30.5	2.23	136
72-3.2	2.4	10.6	26.0	30.9	2.20	134
Average	2.4	10.5	26.2	29.3	2.19	133
Std dev.	0.10	0.41	1.49	2.39	0.06	3

Figure 2. Concentrations of anions in pretreatment room temperature rinse solutions (Table 3) plotted as a function of sample weight.

Table 5. pH of solutions from room temperature rinse and overnight-cooked rinse pretreatment. Initial J-13 pH = 7.6.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	8.21	8.28	8.37	8.40
3	8.14	8.12	8.20	8.23
6	8.04	8.18	8.25	8.29
12	8.13	8.23	8.34	8.40
24	8.12	8.24	8.33	8.35
36	nm ^a	nm	nm	nm
48	8.19	8.28	8.34	8.42
60	8.05	8.24	8.30	8.42
72	nm	nm	nm	nm
Average	8.13	8.22	8.30	8.36
Std dev.	0.06	0.05	0.06	0.07
Cooked rinse				
1	8.22	8.28	8.34	8.32
3	8.14	8.24	8.25	8.26
6	8.08	8.13	8.20	8.22
12	8.16	8.25	8.29	8.31
24	8.16	8.24	8.29	8.34
36	nm	nm	nm	nm
48	8.12	8.18	8.27	8.33
60	8.30	8.27	8.19	8.21
72	nm	nm	nm	nm
Average	8.17	8.23	8.26	8.28
Std dev.	0.07	0.05	0.05	0.05

^a Not measured.

Table 6. Aluminum concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 Al = 0.012 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	0.004	0.031	0.045	0.136
3	0.028	0.028	0.034	0.024
6	0.014	0.018	0.019	0.019
12	0.003	0.022	0.000	0.006
24	0.000	0.025	0.027	0.000
36	0.012	0.018	0.008	0.014
48	0.033	0.036	0.140	0.039
60	0.038	0.031	0.038	0.263
72	0.012	0.010	0.019	0.009
Average	0.016	0.024	0.037	0.057
Std dev.	0.013	0.008	0.039	0.083
Cooked rinse				
1	1.67	1.60	1.34	1.05
3	1.59	1.54	1.47	1.13
6	1.47	1.46	1.37	1.12
6R	1.49	1.48	1.40	1.14
12	1.57	1.62	1.30	1.08
24	1.48	1.53	1.26	1.01
36	1.48	1.54	1.30	0.96
36R	1.51	1.54	1.30	0.96
48	1.58	1.53	1.35	1.03
60	1.61	1.56	1.43	1.14
72	1.59	1.54	1.37	1.09
72R	1.62	1.54	1.38	1.09
Average	1.56	1.54	1.36	1.07
Std dev.	0.06	0.04	0.06	0.06

Table 7. Boron concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 B = 0.13 ppm.

Rinse sample no.	Weight (grams)				
	0.40	0.80	1.60	3.20	
1	0.118	0.120	0.133	0.159	Low
3	0.130	0.137	0.148	0.168	
6	0.102	0.095	0.102	0.115	
12	0.115	0.120	0.135	0.156	
24	0.118	0.122	0.132	0.163	Low
36	0.059	0.064	0.070	0.108	
48	0.119	0.127	0.140	0.162	
60	0.121	0.117	0.132	0.163	
72	0.046	0.050	0.057	0.074	Low
Average ^a	0.120	0.124	0.137	0.162	
Std dev. ^a	0.005	0.007	0.006	0.004	
Cooked rinse					
1	0.130	0.137	0.156	0.188	Low
3	0.132	0.141	0.167	0.213	
6	0.081	0.083	0.107	0.137	
6R	0.149	0.162	0.179	0.219	
12	0.129	0.151	0.154	0.197	Low
24	0.142	0.133	0.157	0.187	
36	0.056	0.064	0.081	0.121	
36R	0.150	0.158	0.177	0.227	
48	0.130	0.139	0.157	0.196	Low
60	0.131	0.136	0.156	0.196	
72	0.041	0.050	0.071	0.112	
72R	0.152	0.158	0.183	0.218	
Average ^a	0.138	0.146	0.165	0.205	
Std dev. ^a	0.009	0.011	0.011	0.014	

^a Excluding low values.

Table 8. Iron concentration in room temperature rinse solutions and overnight-cooked rinse solutions.
J-13 Fe = 0.006 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.28
1	0.024	0.046	0.088	0.049
3	0.018	0.019	0.049	0.040
6	0.004	0.003	0.001	0.002
12	0.003	0.007	0.006	0.004
24	0.010	0.012	0.005	0.005
36	0.003	0.005	0.004	0.004
48	0.056	0.019	0.024	0.033
60	0.022	0.007	0.004	0.015
72	0.004	0.003	0.006	0.001
Average	0.016	0.013	0.021	0.017
Std dev.	0.016	0.013	0.028	0.018
Cooked rinse				
1	0.015	0.021	0.045	0.031
3	0.029	0.034	0.031	0.045
6	bdl*	bdl	0.003	bdl
6R	0.003	0.005	0.015	0.006
12	0.001	0.010	0.002	0.003
24	0.017	0.010	0.014	0.012
36	0.002	0.002	0.002	0.002
36R	0.011	0.010	0.006	0.010
48	0.022	0.033	0.021	0.020
60	0.003	0.003	0.003	0.004
72	0.002	0.004	0.003	0.005
72R	0.024	0.011	0.010	0.016
Average	0.011	0.012	0.013	0.013
Std dev.	0.010	0.011	0.013	0.013

* bdl = below detection limits.

Table 9. Silicon concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 Si = 27.0 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	25.1	24.6	24.5	25.9
3	25.7	25.5	25.8	25.4
6	26.0	25.7	25.5	25.3
12	25.2	24.8	25.1	24.8
24	25.6	24.6	24.6	24.9
36	26.1	25.7	25.6	25.4
48	25.5	25.5	25.3	24.7
60	25.2	24.4	24.2	25.3
72	25.8	25.9	25.6	25.7
Average	25.6	25.2	25.1	25.3
Std dev.	0.34	0.55	0.54	0.38
Cooked rinse				
1	37.6	40.4	44.2	48.3
3	37.5	40.2	46.1	52.2
6	36.7	39.2	44.5	51.0
6R	36.5	39.1	44.4	51.2
12	36.9	40.9	42.8	50.4
24	37.7	37.8	43.2	46.3
36	36.9	40.0	44.1	48.9
36R	36.8	39.6	43.8	49.1
48	36.1	38.8	43.2	48.0
60	36.0	38.4	43.1	48.3
72	37.3	39.8	43.8	50.3
72R	36.9	39.5	43.6	50.8
Average	36.9	39.5	43.9	49.6
Std dev.	0.53	0.84	0.84	1.62

Table 10. Calcium concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 Ca = 12.5 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	16.0	19.5	28.6	45.2
3	17.0	21.2	29.7	46.0
6	16.4	19.4	27.4	43.2
12	15.4	20.2	28.5	42.1
24	17.1	19.3	27.5	45.1
36	16.1	19.6	27.9	41.8
48	15.5	19.6	27.1	41.4
60	15.6	19.3	27.2	43.3
72	16.0	21.1	27.6	44.8
Average	16.1	19.9	27.9	43.7
Std dev.	0.58	0.71	0.80	1.58
Cooked rinse				
1	10.6	10.8	10.8	11.1
3	11.1	10.8	10.3	11.1
6	11.0	10.9	11.2	10.5
6R	11.5	11.5	11.2	11.0
12	11.9	13.0	10.7	12.2
24	14.0	10.9	12.5	10.5
36	12.0	12.2	11.6	13.7
36R	12.4	12.3	12.1	14.4
48	10.0	10.9	10.1	10.4
60	11.0	11.3	10.5	11.1
72	12.1	11.7	11.9	11.4
72R	12.2	11.9	11.8	11.9
Average	11.6	11.5	11.2	11.6
Std dev.	0.99	0.69	0.73	1.21

Table 11. Magnesium concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 Mg = 1.92 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	1.79	1.93	2.16	2.40
3	1.83	1.89	2.10	2.37
6	1.79	1.87	2.02	2.33
12	1.76	1.82	1.97	2.23
24	1.81	1.84	1.98	2.26
36	1.82	1.86	2.01	2.30
48	1.84	1.88	2.02	2.33
60	1.78	1.78	1.92	2.29
72	1.82	1.88	2.04	2.34
Average	1.80	1.86	2.02	2.32
Std dev.	0.02	0.04	0.07	0.05
Cooked rinse				
1	0.29	0.26	0.32	0.31
3	0.19	0.21	0.18	0.26
6	0.27	0.27	0.25	0.23
6R	0.28	0.28	0.27	0.24
12	0.26	0.27	0.26	0.25
24	0.36	0.24	0.33	0.31
36	0.27	0.21	0.24	0.29
36R	0.26	0.21	0.24	0.29
48	0.27	0.30	0.26	0.26
60	0.27	0.24	0.21	0.23
72	0.24	0.24	0.23	0.20
72R	0.24	0.24	0.23	0.20
Average	0.27	0.25	0.25	0.25
Std dev.	0.04	0.03	0.04	0.04

Table 12. Potassium concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 K = 5.1 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	7.0	10.2	15.5	24.2
3	7.9	10.1	15.3	24.3
6	7.8	9.6	14.4	22.5
12	6.9	10.6	14.7	22.4
24	7.6	10.0	14.7	23.1
36	7.5	9.7	14.5	22.3
48	7.2	9.8	14.3	22.2
60	7.4	9.9	14.1	23.3
72	7.2	10.4	14.0	23.9
Average	7.4	10.0	14.6	23.1
Std dev.	0.33	0.30	0.48	0.79
Cooked rinse				
1	7.7	8.9	12.2	17.3
3	7.7	9.6	12.2	18.4
6	7.4	9.0	13.1	16.4
6R	7.5	8.5	11.8	15.4
12	7.4	10.3	11.4	16.8
24	8.6	9.0	12.9	15.6
36	7.7	9.8	12.4	17.4
36R	7.4	8.8	11.5	16.3
48	7.0	9.4	11.9	15.8
60	7.4	9.1	12.0	16.9
72	7.7	9.7	13.0	16.5
72R	7.3	9.3	11.4	15.9
Average	7.6	9.3	12.2	16.6
Std dev.	0.38	0.47	0.58	0.83

Table 13. Sodium concentration in room temperature rinse solutions and overnight-cooked rinse solutions. J-13 Na = 43.9 ppm.

Rinse sample no.	Weight (grams)			
	0.40	0.80	1.60	3.20
1	42.2	43.8	49.8	59.5
3	43.9	44.9	49.9	57.3
6	44.5	43.0	47.1	54.7
12	40.3	45.4	48.4	54.4
24	44.1	42.8	47.9	56.8
36	42.8	42.8	47.9	54.2
48	41.3	42.6	45.7	52.9
60	40.1	41.1	45.0	53.9
72	41.2	44.9	45.9	57.8
Average	42.3	43.5	47.5	55.7
Std dev.	1.6	1.3	1.6	2.1
Cooked rinse				
1	46.2	44.8	48.1	53.2
3	44.0	45.5	48.3	57.8
6	44.0	44.4	50.8	53.1
6R	47.3	49.1	52.5	58.0
12	44.2	50.4	44.9	53.7
24	52.1	43.4	51.5	48.6
36	44.9	46.8	47.5	54.3
36R	48.8	48.6	51.8	59.6
48	40.7	44.1	45.4	49.3
60	40.6	40.9	45.0	51.8
72	45.7	46.1	49.9	52.0
72R	48.0	49.4	52.2	57.3
Average	45.5	46.1	49.0	54.1
Std dev.	3.1	2.7	2.7	3.3

same as those obtained for split A-4 material (Oversby, 1984).

Calcium (Table 10), potassium (Table 12), and sodium (Table 13) concentrations show a linear increase with rock sample weight. Average values for the room temperature rinse solutions for these elements are plotted in Fig. 3 as a function of sample weight. Calcium is the dominant cation, with sodium and potassium being present in somewhat lesser amounts. The amounts found for split A-3 are slightly greater than those reported previously for split A-4 (Oversby, 1984); however, the differences are very small and the proportions of the three cations are essentially constant.

Data for magnesium (Table 11) show a slight decrease in concentration from the J-13 value for the 0.4- and 0.8-g samples and a slight increase for the 1.6- and 3.2-g samples. The cooked rinse data show substantial decreases in all cases due to the retrograde solubility of $MgCO_3$. The same trends were observed for split A-4 material (Oversby, 1984).

Charge balance calculations for the room temperature rinse solutions show that the equivalents of positive charge added to J-13 water by the K + Ca + Na components of the caliche material exactly balance the equivalents of negative charge added by the Cl + NO_3 + SO_4 for the 3.2-g samples. Negative charge exceeds positive charge by 12% for the 1.6-g samples and by 31% for the 0.8- and 0.4-g samples. For small sample weights, the differences between measured concentrations and J-13 values are small; therefore, a large part of the apparent lack of charge balance may be due to uncertainties in the analyses.

Test Matrix Results at 120°C

The test matrix used four sample weights: 0.4, 0.8, 1.6, and 3.2 g of crushed Topopah Spring tuff. The volume of J-13 water used was 48 ml in all cases. Blanks, which consisted of J-13 water without tuff, were run for 3, 12, 24, 36, and 60 days.

The large Parr bombs are too heavy for use with a roller device in the oven, so the tests were run in a static mode with the bombs placed upright. To prevent the rock powder from caking on the bottom of the Teflon capsule, each bomb was shaken by hand several times a day on Monday, Wednesday, and Friday during the reaction interval.

Tables 14 through 17 give data for anions measured by ion chromatography. Fluoride concentrations (Table 14) show a slight increase as a function of reaction time that is not correlated with sample weight. The blank solutions also show a similar in-

crease in fluoride. All fluoride concentration increases are attributed to fluoride release by the Teflon, which is a fluoro-carbon material. Chloride concentrations (Table 15) are similar to or slightly higher than J-13 chloride concentrations. Nitrate concentrations (Table 16) show behavior similar to chloride, while sulfate (Table 17) remains essentially at J-13 levels. These data show that there is little or no change in anion concentrations in the water as a result of reaction with the rock at 120°C. This is the same result as that obtained at 90 and 150°C (Oversby, 1984).

Figure 3. Concentrations of cations in pretreatment room temperature rinse solutions (Tables 10, 12, and 13) plotted as a function of sample weight.

Table 14. Fluoride concentration in ppm from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 fluoride concentration is 2.2 ppm.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	2.2	2.1	2.1	2.1	
3	2.2	2.3	2.2	2.4	2.2
6	2.2	2.1	2.1	2.5	
12	2.3	2.3	2.3	2.4	2.2
24	2.4	2.4	2.2	2.3	2.3
36	3.6	2.4	2.5	2.6	2.7
48	2.4	2.5	2.5	2.5	
60	2.7	2.8	2.5	2.6	2.5
72	2.5	2.8	2.8	2.6	

Table 15. Chloride concentration in ppm from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 chloride concentration is 6.9 ppm.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	8.0	7.6	7.7	7.8	
3	7.0	7.0	7.1	7.4	7.0
6	7.5	7.6	6.7	7.6	
12	7.0	7.2	7.4	7.7	7.1
24	7.4	7.4	7.3	8.1	7.1
36	8.5	7.1	7.5	7.9	7.3
48	6.9	7.1	7.3	7.9	
60	7.2	7.2	7.3	8.0	6.9
72	7.2	7.3	7.6	8.0	

Table 16. Nitrate concentration in ppm from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 nitrate concentration is 9.6 ppm.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	8.7	9.1	8.5	8.9	
3	9.1	9.2	8.8	10.4	9.3
6	9.4	9.1	9.0	10.8	
12	9.0	9.5	10.3	11.7	9.1
24	9.6	9.2	10.0	12.1	9.1
36	9.9	10.2	10.8	12.7	9.8
48	9.3	9.5	10.2	11.0	
60	9.6	9.9	10.5	13.2	9.4
72	9.9	10.0	10.2	12.4	

Table 17. Sulfate concentration in ppm from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 sulfate concentration is 18.7 ppm.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	19.1	19.3	20.4	19.3	
3	17.9	18.1	18.6	18.4	17.8
6	19.8	19.0	17.1	19.7	
12	18.0	17.9	18.1	19.2	18.2
24	17.8	14.4	17.9	19.2	17.8
36	17.8	18.2	18.7	19.7	19.5
48	17.8	18.1	18.1	19.1	
60	18.2	18.0	17.9	19.3	18.1
72	18.4	18.2	19.1	18.7	

Table 18a gives alkalinity of solutions in meq/l and Table 18b gives alkalinity calculated as ppm HCO_3^- . Alkalinity decreases in the blank solutions due to precipitation of magnesium and calcium carbonates. In addition, some carbon dioxide is probably lost to the Teflon capsule, with a resultant decrease in alkalinity (Knauss et al., 1983). Data for calcium, magnesium, and alkalinity for the blank solutions indicate that for all cases except the 3-day blank, the alkalinity is lower than

can be explained by precipitation of $(\text{Ca,Mg})\text{CO}_3$ alone. This supports the suggestion that some CO_2 is lost to the Teflon. Table 19 lists the pH values for solutions at the end of the reaction period. Most of the increase in pH is attributed to the loss of CO_2 to the Teflon; reactions in gold-bag rocking autoclaves do not show an increase in pH (Knauss et al., 1983).

Aluminum concentrations (Table 20) increase fairly rapidly and then show a slight decrease for

Table 18. Alkalinity measured from reaction of Topopah Spring tuff with J-13 water at 120°C.

a. Alkalinity in meq/l. Initial J-13 alkalinity is 2.23 meq/l.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	1.81	1.84	1.99	1.98	
3	1.86	1.87	2.00	1.78	2.02
6	1.77	1.86	1.86	1.91	
12	1.70	1.79	1.76	1.86	1.58
24	1.69	1.72	1.70	1.90	1.42
36	1.75	1.73	1.74	1.81	1.55
48	1.66	1.72	1.77	1.76	
60	1.78	1.72	1.80	1.85	1.37
72	1.72	1.71	1.78	1.84	

b. Alkalinity in ppm bicarbonate. Initial J-13 alkalinity is 136 ppm bicarbonate.

Run time (days)	Sample weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	110	112	121	121	
3	113	114	122	109	123
6	108	113	113	117	
12	104	109	107	113	96
24	103	105	104	116	87
36	107	106	106	110	95
48	101	105	108	107	
60	109	105	110	113	84
72	105	104	109	112	

Table 19. pH of solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 pH = 7.6.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	8.15	8.10	8.21	8.32	
3	7.93	8.02	8.16	8.24	7.96
6	8.02	8.02	8.20	8.29	
12	7.88	8.03	8.05	8.15	
24	8.14	8.38	8.35	8.41	8.30
36	8.38	8.59	8.57	8.56	8.44
48	8.34	8.43	8.80	8.78	
60	8.13	8.43	8.42	8.72	8.59
72	8.94	8.86	8.84	8.99	

Table 20. Concentration of aluminum in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Al = 0.012 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	1.29	1.90	1.85	1.52	
3	1.62	2.05	1.96	1.43	0.011
6	1.91	2.13	1.84	1.34	
6	1.84	1.98	1.78	1.31	Rerun ICP 6/22/84
12	1.74	1.96	1.69	1.24	0.019
24	1.78	2.10	1.69	1.17	0.016
36	1.77	2.12	1.73	1.17	0.015
48	1.68	1.95	1.68	1.18	
60	1.58	1.78	1.38	1.09	0.015
72	1.65	1.99	1.56	1.21	

longer reaction times. The slow decrease is most clearly seen in the data for the 1.6- and 3.2-g samples (Fig. 4). The line labelled "Rerun ICP 6/22/84" represents a reanalysis of the same solution that was originally analyzed on 8/4/83. Agreement of the results for aluminum for analyses made more than 10 months apart indicates that solution concentrations for aluminum are remarkably stable at room temperature for these samples. The behavior of aluminum at 120°C is intermediate between that seen at 90 and 150°C (Oversby, 1984).

Table 21 gives the boron concentrations for the solutions. Data for the 6-day samples were run on 8/4/83, at which time there was a problem with boron data reduction (see the "Experimental Procedures" section for details). Because boron should not precipitate from the solution, even during long term storage of the solutions, the samples were reanalyzed after the difficulties with boron data treatment were corrected. The line labelled "Rerun ICP 6/22/84" is the reanalysis of the same solution. The boron levels found on 6/22/84 are consistent with the pattern established by the other results and lend credence to our assumption

that boron should not precipitate from solution during storage at room temperature. There is a slight correlation of boron content with sample weight and reaction time at a given sample weight. The differences are small, especially as a function of reaction time, and may relate to incomplete removal of the soluble boron component rather than reaction of the major rock phases with the water.

Data for iron (Table 22) show that there is very little increase in iron in solution as a result of reaction of J-13 water with the tuff at 120°C. This is partly because of the limited amount of iron present in the rock, but it is mainly the result of the oxidizing conditions in the reaction system and the low solubility of iron under those conditions.

Data for silicon (Table 23) show an initial sharp increase followed by a further slow increase, as shown in Fig. 5, where the data for each sample weight are plotted as a function of reaction time. Data for the sample run after more than 10 months of storage at room temperature again show remarkable stability of solution concentrations over long storage periods. The same data are plotted in

Figure 4. Aluminum solution concentrations plotted as a function of reaction time.

Table 21. Concentration of boron in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 B = 0.13 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	0.116	0.116	0.117	0.119	
3	0.132	0.134	0.138	0.153	0.133
6	0.027	0.026	0.029	0.040	
6	0.140	0.137	0.145	0.155	Rerun ICP 6/22/84
12	0.130	0.129	0.137	0.157	0.126
24	0.133	0.139	0.138	0.155	0.118
36	0.127	0.123	0.133	0.156	0.119
48	0.131	0.137	0.144	0.166	
60	0.137	0.134	0.145	0.173	0.118
72	0.140	0.144	0.152	0.179	

Table 22. Concentration of iron in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Fe = 0.006 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	0.024	0.052	0.056	0.046	
3	0.061	0.043	0.050	0.052	0.118
6	0.007	0.006	0.007	0.005	
6	0.029	0.016	0.022	0.015	Rerun ICP 6/22/84
12	0.012	0.038	0.023	0.029	0.040
24	0.050	0.034	0.032	0.060	0.023
36	0.040	0.016	0.082	0.038	0.008
48	0.053	0.016	0.028	0.013	
60	0.024	0.049	0.024	0.022	0.463
72	0.031	0.044	0.028	0.035	

Table 23. Concentration of silicon in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Si = 27.0 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	33.4	34.8	37.9	40.4	
3	39.5	41.8	44.8	50.0	25.9
6	39.9	41.2	44.5	50.3	
6	40.1	41.5	44.9	51.5	Rerun ICP 6/22/84
12	44.9	45.1	49.3	57.3	24.8
24	48.1	50.6	54.8	63.4	24.0
36	50.4	50.2	58.3	68.0	24.6
48	51.7	52.7	59.8	70.5	
60	50.8	52.3	57.5	73.6	24.6
72	56.5	55.5	62.8	76.5	

Figure 5. Silicon solution concentrations plotted as a function of reaction time.

Fig. 6 as a function of a "normalized reaction parameter." Data for the 3.2-g sample are plotted in Fig. 6 vs the duration of the reaction (in days). Data for the 1.6-g sample are plotted vs the reaction time divided by two; this adjusts for the difference in the ratio of sample surface area to solution volume for the two samples. The normalized reaction parameter is calculated by multiplying the ratio (sample wt/3.2) times reaction time. If the reaction rate were dependent on the ratio of sample surface area to solution volume (SA/V), the data would define a single curve in the normalized reaction parameter plot. While there is some scatter in the data in Fig. 6, the dependence of reaction rate on SA/V is clear.

Data for 120°C silicon concentrations for the largest ratio of sample weight to solution volume are plotted in Fig. 7 along with data for reaction at 90 and 150°C for the same weight-to-volume ratio. As would be expected, the data at 120°C are intermediate between the results for the other two temperatures, and are somewhat closer to the 150°C values.

Results for calcium (Table 24) and magnesium (Table 25) show similar behavior, with an initial sharp decrease in concentration followed by a slow further decrease. After 72 days of reaction the concentration of calcium is approximately 3.5 ppm whereas the concentration of magnesium is about 0.1 ppm. These concentrations are both higher than those found at 150°C and lower than those found at 90°C, a fact that is consistent with the retrograde solubility of magnesium and calcium carbonate (Oversby, 1984). The data for the sample rerun after long storage show a slight decrease in solution concentrations of calcium; the reason for this is not known.

Potassium (Table 26) and sodium (Table 27) show no clear trend as a function of reaction time, but do show a slight positive correlation with sample weight. The correlation is more regular in the case of potassium than sodium. This may either be caused by the dissolution of some residual soluble material or by the rapid dissolution of a small amount of feldspar. The correlation with sample weight favors the former explanation. Potassium

Figure 6. Silicon concentrations plotted as a function of a normalized reaction parameter. See text for details.

Figure 7. Silicon concentration at 90, 120, and 150°C for constant sample weight to solution volume plotted as a function of reaction time.

concentrations are similar to those reported previously for 90 and 150°C, whereas sodium concentrations are somewhat higher (Oversby, 1984). Potassium data for the rerun samples with 0.4, 1.6, and 3.2 g of rock were run in a sample set that gave a low value of potassium in the J-13 control blank (No. 6, Table 1C). There is probably no significant

change in potassium in solution as a function of storage time. The 0.8-g rerun was run with a set of samples for which the control blank gave correct values for potassium and shows no significant difference from the first analysis. Sodium data for the rerun sample are within error limits of the first analysis.

Table 24. Concentration of calcium in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Ca = 12.5 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	6.71	7.08	8.84	9.41	
3	7.63	7.95	9.26	10.30	5.21
6	6.30	7.04	6.86	7.31	
6	5.97	6.42	6.50	7.00	Rerun ICP 6/22/84
12	6.22	5.86	6.76	7.03	3.76
24	5.29	4.95	4.59	4.96	2.43
36	4.17	3.75	3.97	4.19	2.05
48	4.62	4.95	3.94	4.11	
60	4.88	4.47	4.48	4.34	2.26
72	3.44	3.63	3.74	3.59	

Table 25. Concentration of magnesium in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Mg = 1.92 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	0.235	0.265	0.273	0.219	
3	0.272	0.213	0.231	0.235	0.596
6	0.102	0.095	0.080	0.071	
6	0.126	0.093	0.088	0.070	Rerun ICP 6/22/84
12	0.099	0.153	0.111	0.117	0.289
24	0.173	0.116	0.119	0.180	0.176
36	0.152	0.058	0.262	0.134	0.124
48	0.197	0.076	0.082	0.051	
60	0.115	0.181	0.097	0.079	0.180
72	0.096	0.138	0.095	0.109	

Table 26. Concentration of potassium in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 K = 5.1 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	5.6	6.4	7.6	9.3	
3	6.2	6.8	8.9	10.9	5.0
6	6.9	7.2	8.4	10.8	
6	6.1	5.9	7.8	9.5	Rerun ICP 6/22/84
12	6.3	6.8	8.2	10.4	4.5
24	6.6	7.1	8.1	9.5	4.5
36	6.0	6.9	7.5	8.8	4.3
48	6.3	7.3	7.8	8.9	
60	6.0	7.1	7.9	9.9	4.7
72	6.0	7.1	8.2	9.2	

Table 27. Concentration of sodium in solutions from reaction of Topopah Spring tuff with J-13 water at 120°C. Initial J-13 Na = 44 ppm.

Reaction time (days)	Weight (grams)				Blank
	0.4	0.8	1.6	3.2	
1	41.0	40.3	43.4	42.4	
3	45.9	46.2	51.3	52.3	46.2
6	43.3	44.7	44.5	46.2	
6	44.4	43.7	45.0	47.1	Rerun ICP 6/22/84
12	43.0	42.2	45.2	47.1	40.6
24	46.5	46.8	45.3	45.4	40.7
36	44.7	43.1	44.0	45.1	39.5
48	45.4	46.1	45.6	47.3	
60	42.7	44.1	44.6	51.1	40.9
72	46.5	44.7	47.0	50.4	

Table 28. Estimate of steady-state water chemistry for the system Topopah Spring Tuff — J-13 water.

Element	Concentration (ppm)		
	90°C ^a	120°C	150°C ^a
Al	0.4	1.2	1
B	0.1	0.1	0.1
Fe	0	0	0
Si	49	81	122
Ca	8	3.5	3
Mg	0.2	0.1	0.1
K	9	9	9
Na	40	45	40
F	2	2	2
Cl	7	7	7
NO ₃	9	9	9
SO ₄	18	18	18

^a From Oversby (1984).

Estimation of Steady-State Water Chemistry

None of the experiments reported in this paper reached steady-state solution concentrations. The matrix permeability of the Topopah Spring tuff is low and the downward infiltration rate of water at Yucca Mountain is also low. This suggests that water flow through the rock matrix material at the repository level would be slow enough to achieve steady-state solution compositions at the appropriate temperatures. Long term experiments using J-13 water and Topopah Spring tuff are in progress in Teflon reaction vessels and in gold-bag rocking autoclaves to determine the steady-state chemistry. Pending completion of those experiments, the data given above will be

used together with the solubility of cristobalite to predict the steady-state solution compositions.

Table 28 gives the estimated water chemistry at steady state for the system Topopah Spring tuff plus J-13 water. Data given in the sections on results showed that the only source of anions to change the water chemistry came from the caliche component, and not from the rock itself. Experiments using rock from drill cores have established that there is no caliche material associated with the Topopah Spring tuff where it lies below the region affected by surface runoff (Oversby, 1983). Thus, the estimates for anions are simply the original J-13 values, with a slight lowering in the case of sulfate to allow for some precipitation. Estimates for the expected composition at 90 and 150°C are from Oversby (1984).

Boron and iron concentrations are not expected to change significantly as the result of hydrothermal reaction of the Topopah Spring tuff and J-13 water. Potassium and sodium increase slightly, and appear to be essentially independent of temperature. Aluminum concentrations will increase above J-13 values, and presently exhibit a maximum value for the 120°C data set. Calcium and magnesium concentrations will decrease. Silicon

concentrations have been set at the cristobalite solubility level (calculated from data in Walther and Helgeson, 1977) for each temperature. Quartz is present in the rock and may control silicon solubility in the very long term; however, data obtained to date indicate that silica precipitation is very slow, and the kinetics of quartz formation may be too slow to control the solubility.

Conclusions

Reaction of the Topopah Spring Member of the Paintbrush Tuff with J-13 well water, a groundwater obtained from a locality at which the Topopah Spring tuff is an aquifer, shows that changes in water chemistry due to heating the rock and water at 120°C are limited to minor effects in all cases except silicon. Silicon concentrations increase as a result of dissolution of cristobalite and are expected to reach 81 ppm at 120°C at steady-state.

The surface outcrop samples used in these experiments contained a substantial amount of easily removed evaporite deposits on the rock surface. This type of material is not expected to be encountered at depth under Yucca Mountain (Oversby, 1983) and should be removed before the rock is used in experiments where the evaporite material might affect the results. A relatively simple pretreatment procedure is effective in removing the evaporite salts.

Anion data show that there is no detectable amount of anions in the rock at 120°C.

The pH of these experiments increased as a result of reaction. Data from comparable experiments conducted in gold-bag rocking autoclaves

do not show such an increase (Knauss et al., 1983). The increase in pH observed here is attributed to uptake of CO₂ by the Teflon reaction vessels. Alkalinity determinations for these experiments are expected to underestimate the alkalinity of the rock-water system for this reason.

Changes in concentration occurred for aluminum, potassium, sodium, magnesium, and calcium as a result of reaction. Magnesium and calcium decrease due to the precipitation of (Ca,Mg) CO₃ (Knauss et al., 1983). Aluminum increases to form a supersaturated condition followed by slow precipitation. The solubility of aluminum in this system appears to be controlled by gibbsite (Knauss et al., 1983). Potassium increases from 5 to 9 ppm, probably due to dissolution of alkali feldspar. There is no significant difference in potassium concentration as a function of temperature. Sodium increases are very small, and may be due either to feldspar dissolution or to incomplete removal of the soluble material.

The results of the experiments at 120°C are consistent with those obtained for the same rock-water system at 90 and 150°C (Oversby, 1984).

Acknowledgments

The success of these experiments is the result of the contributions of many individuals. Thanks to the care and effort of Jan Brown we made effective use of laboratory time and space, and had no instances where experiments had to be rerun. Cation analyses were performed by A. Langhorst for samples run on the polychromator and by T. Duewer for samples run on the monochromator system. Ion chromatography measurements were made by J. Lam, and alkalinity by R. Swanziger. Their careful measurements and attention to detail are greatly appreciated. Rock crushing, matrix execution, and pH measurements were done by Jan Brown. Without her efforts this work would not have been accomplished. T. Wolery provided the calculated solubility of cristobalite as a function of temperature. The manuscript was reviewed by L. Ballou, J. Delany, and K. Knauss; their time and effort are greatly appreciated.

REFERENCES

- Knauss, K. G. (1984), *Petrologic and Geochemical Characterization of the Topopah Spring Member of the Paintbrush Tuff: Outcrop Samples used in Waste Package Experiments*, Lawrence Livermore National Laboratory, Livermore, CA, UCRL-53558.
- Knauss, K. G., V. M. Oversby, and T. J. Wolery (1983), *Post Emplacement Environment of Waste Packages*, Lawrence Livermore National Laboratory, Livermore, CA, UCRL-89475.
- Oversby, V. M. (1983), *Performance Testing of Waste Forms in a Tuff Environment*, Lawrence Livermore National Laboratory, Livermore, CA, UCRL-90045.
- Oversby, V. M. (1984), *Reaction of the Topopah Spring Tuff with J-13 Well Water at 90°C and 150°C*, Lawrence Livermore National Laboratory, Livermore, CA, UCRL-53552.
- Peck, E. S., A. L. Langhorst, and D. W. O'Brien (1979), *Analyses of Natural Waters with an Automated Inductively Coupled Plasma Spectrometer System*, Lawrence Livermore National Laboratory, Livermore, CA, UCRL-81043.
- U.S. Nuclear Regulatory Commission (1983), 10 CFR Part 60, "Disposal of High-Level Radioactive Wastes in Geologic Repositories," Technical Criteria, Final Rule, *Federal Register* 48 (120).
- Walther, J. V. and H. C. Helgeson (1977), Calculation of the Thermodynamic Properties of Aqueous Silica and the Solubility of Quartz and its Polymorphs at High Temperatures and Pressures, *Amer. Jour. Sci.* 277, 1315-1351.

Appendix A

Tables A1 through A5 duplicate the polychrometer ICP data from Tables 20, 21, 23, 24, and 27. Splits of these solutions were also analyzed on the new monochrometer ICP. Results of those analyses are given in Tables A1 through A5, labelled as Nx. Agreement of results from the two instruments is generally quite good for aluminum and sodium. Data for boron show somewhat more variability than expected for the new instrument. Silicon data show a systematic difference between the instruments, with the new instrument giving higher concentrations. The amount of the difference increases as the silicon concentration increases. Data for J-13 control blanks gave the same silicon values for both instruments (27 ppm, Table 1). The results for calcium are somewhat erratic and do not show the expected level of agreement. Further work with the new instrument will be necessary before it can be used for routine analyses.

Table A1. Comparison of results from polychromator ICP (Table 20) with monochromator ICP (new instrument). Aluminum concentration in ppm.

Reaction time (days)	Weight (grams)				Blank
	0.40	0.80	1.60	3.20	
1	1.29	1.90	1.85	1.52	
N1	1.45	2.14	1.94	1.59	
3	1.62	2.05	1.96	1.43	0.011
N3	1.57	1.98	1.92	1.41	<0.03
6	1.91	2.13	1.84	1.34	
N6	1.91	2.15	1.78	1.29	
12	1.74	1.96	1.69	1.24	0.019
N12	1.74	1.94	1.68	1.24	<0.08
24	1.78	2.10	1.69	1.17	0.016
N24	1.73	2.05	1.66	1.10	<0.08
36	1.77	2.12	1.73	1.17	0.015
N36	1.74	2.17	1.76	1.21	<0.08
48	1.68	1.95	1.68	1.18	
N48		1.95	1.71		
60	1.58	1.78	1.38	1.09	0.015
N60		1.46	1.46	1.13	<0.08
72	1.65	1.99	1.56	1.21	
N72		2.09	1.58	1.27	

Table A2. Comparison of results from polychromator ICP (Table 21) with monochromator ICP (new instrument). Boron concentration in ppm.

Reaction time (days)	Weight (grams)				Blank
	0.40	0.80	1.60	3.20	
1	0.116	0.116	0.117	0.119	
N1	0.170	0.170	0.130	0.140	
3	0.132	0.134	0.138	0.153	0.133
N3	0.160	0.170	0.160	0.170	0.160
6	0.140	0.137	0.145	0.155	
N6	0.170	0.150	0.140	0.150	
12	0.130	0.129	0.137	0.157	0.126
N12	0.100	0.100	0.130	0.120	0.110
36	0.127	0.123	0.135	0.156	0.119
N36	0.110	0.120	0.130	0.140	0.110
48	0.131	0.137	0.144	0.166	
N48	0.110	0.130	0.130	0.190	
60	0.137	0.134	0.145	0.173	0.118
N60	0.180	0.170	0.170	0.200	0.140
72	0.140	0.144	0.152	0.179	
N72	0.130	0.130	0.130	0.150	

Table A3. Comparison of results from polychromator ICP (Table 23) with monochromator ICP (new instrument). Silicon concentration in ppm.

Reaction time (days)	Weight (grams)				Blank
	0.40	0.80	1.60	3.20	
1	33.4	34.8	37.9	40.4	
N1	37.0	39.2	40.3	43.5	
3	39.5	41.8	44.8	50.0	25.9
N3	39.0	41.1	43.9	53.8	28.3
6	39.9	41.2	44.5	50.3	
N6	43.0	45.2	47.9	60.6	
12	44.9	45.1	49.3	57.3	24.8
N12	45.8	46.2	48.9	65.4	27.8
24	48.1	50.6	54.8	63.4	24.0
N24	47.2	49.9	53.8	68.8	26.1
36	50.4	50.2	58.3	68.0	24.6
N36	51.8	54.1	60.6	72.2	24.9
48	51.7	52.7	59.8	70.5	
N48	57.6	59.1	67.4	82.8	
60	50.8	52.3	57.5	73.6	24.6
N60	58.0	66.6	66.6	82.7	26.8
72	56.5	55.5	62.8	76.5	
N72	62.4	63.9	70.8	88.0	

Table A4. Comparison of results from polychromator ICP (Table 24) with monochromator ICP (new instrument). Calcium concentration in ppm.

Reaction time (days)	Weight (grams)				Blank
	0.40	0.80	1.60	3.20	
1	6.71	7.08	8.84	9.41	
N1	7.00	7.70	9.30	11.00	
3	7.63	7.95	9.26	10.30	5.21
N3	6.45	7.10	7.90	8.52	4.10
6	6.30	7.04	6.86	7.31	
N6	6.00	6.80	7.40	7.50	
12	6.22	5.86	6.76	7.03	3.76
N12	5.71	5.31	6.10	5.97	2.80
24	5.29	4.95	4.59	4.96	2.43
N24	3.70	3.54	2.94	1.54	
36	4.17	3.75	3.97	4.19	2.05
N36	3.51	3.23	3.29	3.60	1.07
48	4.62	4.95	3.94	4.11	
N48	3.57	3.98	3.00	3.39	
60	4.88	4.47	4.48	4.34	2.26
N60	3.66	3.66	3.79	3.32	1.17
72	3.44	3.63	3.74	3.59	
N72	2.35	2.78	2.82	2.68	

Table A5. Comparison of results from polychromator ICP (Table 27) with monochromator ICP (new instrument). Sodium concentration in ppm.

Reaction time (days)	Weight (grams)				Blank
	0.40	0.80	1.60	3.20	
1	41.0	40.3	43.4	42.4	
N1	45.0	44.0	46.0	48.0	
3	45.9	46.2	51.3	52.3	46.2
N3	44.4	44.1	41.2	46.6	44.8
6	43.3	44.7	44.5	46.2	
N6	46.0	47.0	47.0	49.0	
12	43.0	42.2	45.2	47.1	40.6
N12	43.2	43.0	42.4	36.0	46.8
24	46.5	46.8	45.3	45.4	40.7
N24	43.4	41.1	39.5	44.8	39.8
36	44.7	43.1	44.0	45.1	39.5
N36	46.2	47.4	44.9	48.3	48.3
48	45.4	46.1	45.6	47.3	
N48	49.5	46.5	46.6	46.9	
60	42.7	44.1	44.6	51.1	40.9
N60	47.8	46.5	44.9	52.2	41.2
72	46.5	44.7	47.0	50.4	
N72	51.2	46.1	48.2	47.6	