LA-UR-973046

Approved for public release;
distribution is unlimited.

TITLE: MAGNETIC PHASE DIAGRAMS OF UNiGe

CONF-9708108-

RECEIVED

NOV 12 1997

OSTI

AUTHOR(S): H. Nakotte, LANSCE-12
I.H. Hagmusa, Van der Waals-Zeeman Inst., Univ of Amsterdam
J.C.P. Klaasse, Van der Waals-Zeeman Inst., Univ. of Amsterdam
M.S. Torikachvili, Dept. of Physics, San Diego State Univ.
A.H. Lacerda, NHMFL, Pulsed Field Facility
E. Bruck, Van der Waals-Zeeman Inst., Univ. of Amsterdam
F.R. de Boer, Van der Waal-Zeeman Inst., Univ of Amsterdam

SUBMITTED TO: International Conference on Research in High Magnetic Fields,
Sydney, Australia, August 4-6, 1997
Physica B

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Los Alamos
NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. The Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MAGNETIC PHASE DIAGRAMS OF UNiGe

H. Nakotte^a, I.H. Hagmusa^b, J.C.P. Klaasse^b, M.S. Torikachvili^c, A.H. Lacerda^d, E. Brück^b, K. Prokes^b and F.R. de Boer^b,

a: MLNSC, Los Alamos National Laboratory, Los Alamos NM 87545, USA

b: Van der Waals-Zeeman Institute, University of Amsterdam, 1018 XE Amsterdam, The Netherlands,

c: Department of Physics, San Diego State University, San Diego CA 92182, USA

d: NHMFL, Pulsed Field Facility, Los Alamos National Laboratory, Los Alamos NM 87545, USA

UNiGe undergoes two magnetic transitions in zero field. Here, the magnetic diagrams of UNiGe for $B \parallel b$ and $B \parallel c$ are reported. We performed temperatures scans of the magnetization in static magnetic fields up to 19.5T applied along the b and c axes. For both orientations 3 magnetic phases have been identified in the B-T diagrams. We confirmed the previously reported phase boundaries for $B \parallel c$, and in addition we determined the location of the phase boundaries for $B \parallel b$. We discuss a possible relationship of the two zero-field antiferromagnetic phases (commensurate: $T < 42\text{K}$; incommensurate: $42\text{K} < T < 50\text{K}$) and the field-induced phase, which, at low temperatures, occurs between 18 and 25T or 4 and 10T for $B \parallel b$ or $B \parallel c$, respectively. Finally, we discuss the field dependence of the electronic contribution γ to the specific heat for $B \parallel c$ up to 17.5T, and we find that its field dependence is similar to the one found in more itinerant uranium compounds.

Keywords: magnetic-phase diagram, antiferromagnetism, uranium compounds

UNiGe exhibits two antiferromagnetic transitions at about 50 and 42K in zero field [1]. Neutron-diffraction studies established that both zero-field phases are well described in terms of single- $\mathbf{q}$ non-collinear arrangement of the moments which form in two domains [2]. The magnetic phase below 42K is commensurate with $\mathbf{q}=(0,1/2,1/2)$, while the phase between 42 and 50K is incommensurate with $\mathbf{q}=(0,\delta,\delta)$ and $\delta \approx 0.35$. Bulk magnetic investigations of UNiGe revealed a highly anisotropic behaviour both in the magnetically-ordered and in the paramagnetic states [3]. At low temperatures, spin-flip transitions occur for fields applied along b and c axis. At 4.2K, two transitions are found in both the b -axis magnetization (~ 17 and ~ 25 T) and the c -axis magnetization (~ 4 and ~ 10 T). For fields applied along c , it was shown that a magnetic structure with $\mathbf{q}=(0,1/3,1/3)$ forms [1], while the second transition presumably leads to a forced-ferromagnetic alignment of the moments. On the other hand, it was found that the a -axis response is much weaker and cannot be aligned even in pulsed fields up to 38T [3].

For $B \parallel c$, the magnetic-phase boundaries of the two zero-field phases and the field-induced $(0,1/3,1/3)$ phase have been well established by previous magnetization [4], magnetoresistance [5] and specific-heat measurements [6]. For $B \parallel b$, on the other hand, far less work has been done and the studies have been restricted to field sweeps at fixed temperatures [4]. These studies had led to a relatively complex b -axis B-T diagram with 5 different magnetic phases. However, Purwanto et al. [7] pointed out that some of the 'phase lines' may be due to a misaligned crystallite, which was found in neutron-diffraction studies on the same sample. Therefore, it was argued that the intrinsic b -axis B-T phase diagram exhibits only 3 different magnetic phases similar to the c -axis B-T phase diagram. [7].

Here, we want to clarify the phase boundaries for $B \parallel b$ using a better-quality single crystal, which was used also previously in magnetoresistance studies for $B \parallel c$ [5]. We performed temperatures scans of the magnetic response in static magnetic fields up to 19.5T applied along the b and c axes. In addition, we studied the field dependence of the electronic contribution γ to the specific heat in magnetic fields up to 17.5T. The magnetization studies were performed in the 20T superconducting magnet at the Los-Alamos Pulse Field Facility of the National High Magnetic Field Laboratory; specific-heat studies in applied fields were done using the 17.5T superconducting magnet at the University of Amsterdam [8].

Fig. 1 shows the temperature dependence of the magnetization for various magnetic fields applied along the b and the c axis. For $B \parallel b$, the most prominent feature is the drop in the magnetization at the 42K transition. We find a smooth and continuous shift of this transition toward lower temperatures with increasing field until it disappears for $B > 18$ T. There is a second anomaly in the b -axis magnetization: It is a barely visible increase in the magnetic response around 50K (at 2T) which becomes slightly more pronounced, but for fields above 10T it changes sign and a weak drop in the magnetic response appears at similar temperatures. Taking the temperatures where the anomalies occur, we can construct the B-T phase diagram for $B \parallel b$ and $B \parallel c$. The results are shown in Fig. 2, where previous results have been included as well. It should be noted that we do not find any other obvious anomalies for the b -axis response, and this seems to confirm the picture proposed in ref. 7. Comparing the b -axis and c -axis magnetizations (Fig. 1) with the B-T phase diagrams (Fig. 2), the following picture emerges: For sufficiently low fields ($B < 10$ T for $B \parallel b$ and $B < 4$ T for $B \parallel c$), one enters the incommensurate $(0, \delta, \delta)$ phase, which causes the magnetic response to increase possibly suggesting changes in δ and/or domain

repopulation effects. For higher magnetic fields, but below the second spin-flip transition ($B \approx 25\text{T}$ for $B \parallel b$ and $B \approx 9.5\text{T}$ for $B \parallel c$), one enters the field-induced $(0,1/3,1/3)$ phase, which is accompanied by a slight drop in the magnetic response. A very similar behaviour in such intermediate fields for $B \parallel b$ and $B \parallel c$ may thereby indicate that the same $(0,1/3,1/3)$ phase forms for both field orientations. Entering the $(0,1/2,1/2)$ gives rise to a pronounced drop in the magnetic response. Application of fields in close proximity of the 'ferromagnetic' phase boundary (9T for $B \parallel c$) gives rise to a relatively complex behaviour, which likely can be attributed to domain effects.

Altogether, we believe that our magnetization results are consistent with a relationship between the 3 magnetic phases which has been proposed earlier [2]. The incommensurate phase is believed to consist out of ferromagnetic sheets with a $(++-)$ stacking with a spin-slip every 20th layer or so. The $(0,1/3,1/3)$ has the same $(++-)$ stacking of ferromagnetic sheets, but there is no spin-slip. The commensurate $(0,1/2,1/2)$ phase, on the other hand, has ferromagnetic sheets stacked in a $(+-)$ sequence.

Furthermore, we studied the field dependence of the electronic contribution γ to the specific heat. This provides some qualitative measure of Fermi-surface effects. For that, we measured c/T vs B for $B \parallel c$ at 850mK using a semi-adiabatic method. Data were taken with increasing field. As can be seen in Fig.3, γ is somewhat reduced at the fields where the spin-flip transitions ($\sim 3\text{T}$ and $\sim 11\text{T}$ at 850mK) occur. However, reduced values of γ at the first transition are present only in close proximity to the spin-flip transition. In fact, excluding the data points at 3 and 4T, one can fit all other values up to 11T to a quadratic behaviour (solid line in Fig. 3). Above the second transition, however, one finds almost linearly decreasing γ values up to the highest field applied (17.5T). This behaviour (quadratic increase below the transition, linear decrease above the transition) is surprisingly

similar to the findings for UNiAl [9], which one can consider as a moderately enhanced heavy-fermion system, whereas the $5f$ electrons in UNiGe are more localized. At present, we have very little idea how to interpret this observation. Furthermore, we find some evidence that γ is slightly enhanced ($\sim 2\text{-}3\text{mJ/K}^2\text{mol}$) between 3 and 9T after application of a field, but we did not study such hysteresis effects in details. Nevertheless, our results provide some evidence that the basic features of the Fermi surface in UNiGe survive the first spin-flip transition, while the nature of the Fermi surface seems to be modified at the second spin-flip transition.

In conclusion, we have confirmed the phase boundaries between the long-range ordered and the paramagnetic phases for $B \parallel c$, and we also established the location of the phase boundaries for $B \parallel b$. Finally, we found a field dependence of γ in UNiGe similar to the one found in more itinerant uranium compounds.

This work was supported in part by the division of Basic Energy Sciences of the U.S. Department of Energy and by the "Stichting voor Fundamenteel Onderzoek der Materie" (FOM).

references

- [1] V. Sechovsky, L. Havela, A. Purwanto, Allen C. Larson, R.A. Robinson, K. Prokes, H. Nakotte, E. Brück, F.R. de Boer, P. Svoboda, H. Maletta and M. Winkelmann; J. Alloys Comp. **213/214** (1994) 536
- [2] H. Nakotte, A. Purwanto, R.A. Robinson, Z. Tun, K. Prokes, A.C. Larson, L. Havela, V. Sechovsky, H. Maletta, E. Brück and F.R. de Boer, Phys. Rev. B **54** (1996) 7201.
- [3] L. Havela, V. Sechovsky, F.R. de Boer, E. Brück and H. Nakotte; Physica B **177** (1992) 159-63.
- [4] F.R. de Boer, K. Prokes, H. Nakotte, E. Brück, M. Hilbers, P. Svoboda, V. Sechovsky, L. Havela and H. Maletta; Physica B **201** (1994) 251
- [5] H. Nakotte, L. Havela, V. Sechovsky, A. Lacerda, A. Purwanto, K. Prokes, E. Brück,, F.R. de Boer and M.S. Torikachvili; in *Physical Phenomena at High Magnetic Fields II*, Singapore: World Scientific (1996) 209
- [6] I.H. Hagusma, J.C.P. Klaasse, E. Brück, K. Prokes, H. Nakotte and F.R. de Boer; J. Appl. Phys. **81** (1997) 4157.
- [7] A. Purwanto, V. Sechovsky, L. Havela, R.A. Robinson, H. Nakotte, Allen C. Larson, K. Prokes, E. Brück and F.R. de Boer; Phys. Rev. B **53** (1996) 758
- [8] J.C.P. Klaasse, Rev.Sci. Instrum. **68** (1997) 89

- [9] E. Brück, H. Nakotte, F.R. de Boer, P.F. de Chatel, H.P. van der Meulen, J.J.M. Franse, A.A. Menovsky, N.H. Kim-Ngan, L. Havela, V. Sechovsky, J.A.A.J. Perenboom, N.C. Tuan and J. Sebek; Phys. Rev. B **49** (1994) 8852

figure captions:

Fig. 1: Temperature dependence of the magnetization for various fields (a) applied along the b axis and (b) fields applied along the c axis. Note the presence of various anomalies (see text), which determine the magnetic phase boundaries.

Fig. 2: Magnetic phase diagram for fields (a) applied along the b axis and (b) applied along the c axis. The plots comprise the data from previous magnetization ($\times$)⁴, magnetoresistance (Δ)⁵ and present magnetization data ($\bullet$). The shaded area in (b) represents the region where hysteresis effects occur.

Fig. 3: Field dependence of the electronic contribution γ to the specific heat for fields applied along the c axis, determined by C/T values at 850mK. The arrows indicate the locations of metamagnetic transitions in the c -axis magnetization (~ 4 and ~ 11 T at 850mK). The error bars represent an estimate of the absolute error ($\sim 1\text{mJ/K}^2\text{mol}$), while the relative error is much smaller ($\sim 0.1\text{mJ/K}^2\text{mol}$). The solid line represents the fit of the data for $B < 10\text{T}$ (and excluding 3 and 4T) to a quadratic field dependence. Data have been taken with increasing field.

Fig. 1 H. Nakotte et al
Magnetic phase diagrams...

Fig. 2: H. Nakotte et al.
 Magnetic phase diagrams ...

Fig. 3: H. Nakotte et al.
Magnetic phase diagrams...