LA-UR- 97-4114

Approved for public release;
distribution is unlimited.

Title:

NUMERICAL SIMULATION OF EXPLOSIVE
VOLCANISM AND ITS EFFECTS ON THE
ATMOSPHERE

RECEIVED

FEB 02 1998

OSTI

Author(s):

Greg A. Valentine, EES-5
James E. Bossert, EES-8

Submitted to:

Final report on LDRD project submitted
to DOE Office of Scientific and
Technical Information (OSTI)

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

Los Alamos
NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. The Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible electronic image products. Images are produced from the best available original document.

Numerical Simulation of Explosive Volcanism and Its Effects on the Atmosphere

Greg A. Valentine*, James E. Bossert

Abstract

This is the final report of a one-year, Laboratory-Directed Research and Development (LDRD) project at the Los Alamos National Laboratory (LANL). The objective of this project was to begin work on combining two modeling approaches in order to advance the state-of-the-art in simulating and predicting explosive volcanic eruption dynamics and their effects. We began applying the CFDLIB family of codes for the near-field (high temperature, velocity, and particle concentration) region of an explosive eruption. We also applied the RAMS meteorological code to model the far-field dynamics of eruption clouds and ash fallout. Initial test runs were conducted in preparation for full-scale simulations that would eventually couple the two models for the most comprehensive volcano simulation tool to date. Eventual applications include aviation hazards, risk assessment, and extension to atmospheric collateral effects of conventional and nuclear weapons.

Background and Research Objectives

Explosive volcanism is one of the most dramatic of natural events. During these eruptions a mixture of hot particles and gas flow out of a volcanic vent at speeds of several 100 m/s. The mixture rises into the atmosphere and entrains air. Depending on the eruption conditions (velocity, gas content, particle size range, pressure, and vent size) and atmospheric conditions (cross winds, stratification), the mixture continues to rise to high altitudes or falls back to the ground, producing high-speed density currents that flow out across the landscape. Eruptions affect humans on a variety of time and space scales. Near an eruption site (within 1 - 100 km, the *near field*), eruptive effects are immediate and devastating. They include fast-moving (up to 300 m/s), hot, particle laden density currents, shock waves, and heavy fallout of ash from buoyant plumes. At intermediate distances (10s - 100s of km from eruption sites; *mesoscale*), plumes of ash leave damaging fallout deposits that choke machinery and agriculture and pose threats to aviation for the duration of an eruption (hours to weeks). At larger distances (*global scale*) the ash, gases, and aerosols that an eruption injects into the atmosphere can affect the global climate for several years after an eruption.

Los Alamos has a long-standing interest in volcanic and atmospheric processes due to a variety of programs, including geothermal energy, nuclear test containment, hazardous plume dispersion, and global climate change. Laboratory researchers lead the geophysical community in applying 2-D, time-dependent, multiphase hydrodynamic modeling to the near field dynamics of explosive eruptions (Wohletz et al., 1984; Valentine and Wohletz, 1989a,b; Valentine et al., 1991; Valentine et al., 1992). These approaches are now being used in the international geophysics community for both basic research and hazard mitigation (e.g., Dobran et al., 1993, 1994; Neri and Dobran, 1994; Kieffer, 1995), and have played a major role in establishing computational approaches for a range of magmatic processes (Bergantz, 1995). Since our early work the capability to compute near field dynamics has been superseded by the work of Dobran and colleagues.

Los Alamos is currently using and developing the most scientifically rigorous modeling systems for the simulation of atmospheric mesoscale processes (Bossert and Cotton, 1994a, 1994b; Margolin et al., 1995). These models have recently been used to examine dispersion processes in regions of highly complex terrain (Poulos and Bossert, 1995; Bossert and Poulos, 1995). To date, the eruption and atmosphere simulation capabilities of the Laboratory have not been combined.

The objectives of this proposal are two-fold. First, we will make a major advance in our ability to simulate the near field dynamics of explosive eruptions by applying the most sophisticated multiphase hydrodynamics codes available at the Laboratory -- the CFDLIB family of codes. The new capabilities will include 3-D, multiple particle sizes, and multicomponent gas. Second, we will combine the eruption modeling capability with our atmospheric transport capability. We will use the near field modeling to develop rigorous source terms for mesoscale transport of volcanic plumes and ash fallout. This work will constitute the next generation of eruption dynamics and atmospheric effects modeling.

Importance to LANL's Science and Technology Base and National R&D Needs

- *Forefront Science* -- Our previous work on the first generation of multiphase numerical eruption modeling was of extremely high visibility in the geophysical community. It introduced numerical simulation into a realm of geological sciences that had previously used only observation and simple theory. The work in this proposal will allow us to retain this high visibility and the programmatic funding that results from a strong technical reputation.

Weapons Effects -- The computational approaches and tools developed in this proposal will be directly applicable to modeling of nuclear and conventional weapons effects.

Weapons effects modeling in the past has been an important component of the weapons

program, but is now at risk due to lack of programmatic support. Our research will sharpen the appropriate modeling tools in a manner that maintains visibility and reputation in the research community and that addresses real problems.

- *Aviation Hazards* -- A direct and highly practical application of this work will be the mitigation of aviation hazards. Explosive volcanic regions such as the Pacific Rim, Indonesia, and the Mediterranean are also areas of heavy air traffic both for civilian and defense purposes. We are pursuing joint work with the Jet Propulsion Laboratory that would couple satellite observations of eruptions with real-time predictive modeling of plumes so that air traffic can be safely rerouted, particularly for flights over and around the Pacific Ocean

Scientific Approach and Accomplishments

Approach and accomplishments are as follows:

Theoretical underpinning. Comprehensive review of explosive volcanism modeling to date was conducted and used as part of the basis for a new multifield theoretical framework for describing eruption phenomena. The theoretical framework is based on conservation of mass, momentum, and energy for multicomponent gas with particles and droplets with a range of sizes. This was presented at an international conference in the form of a short course, and is being published as a chapter of a book that is tentatively titled "Physics of Explosive Eruptions."

Near-field dynamics. We began using the CFDLIB family of codes for near-field (high velocity, temperature, and particle loading regimes) dynamics of explosive eruptions. Much effort was put into learning the codes and their underlying theories, in order to be most efficient in applying them to volcanic problems. Results to date include simple gas explosions, and initial work on loading explosions with ash particles. These are essential first steps toward successful application to volcanic conditions.

Far-field dynamics. Numerical experiments with the regional meteorological code RAMS (Pielke, 1992) were conducted to determine what, if any, modifications would be necessary for the code to be able to model intense thermal buoyancy effects, ash transport, and fallout in explosive eruption clouds as they are transported downwind. Initial calculations for small-scale (but quite common) types of explosive eruptions were successfully conducted, and we determined that velocities of up to about 80 m/s can be computed without any modification to RAMS. We are exploring the adaptation of the

numerical models of precipitation and condensation in rain clouds to fallout and particle coalescence in volcanic plumes.

Because of the small amount and short duration of this grant (\$80K for 1 year), we have not yet obtained definitive new results. However, we have laid a strong foundation for gaining new levels of understanding and predictive capability for volcanic phenomena using these computer codes. We expect to be ready to present two papers at the next general congress of the International Association of Volcanology and Chemistry of the Earth's Interior (July, 1998).

Publications

Valentine, G.A., "Eruption Column Physics," in Physics of Explosive Eruptions (tentative book title), Elsevier Science Publisher, (in press, 1998).

References

- Bergantz, G.W., "Changing techniques and paradigms for the evaluation of magmatic processes" *Journal of Geophysical Research*, **100**, 17603-17613 (1995).
- Bossert J.E., Poulos G.S. (1995) "A numerical investigation of mechanisms affecting drainage flows in highly complex terrain," *Theoretical and Applied Climatology*, **52**, 119-134 (1995).
- Bossert, J.E., "Forecasting winds over the Pajarito Plateau: A preliminary investigation" LANL Report LA-UR-95-3907, 57 pp (1995).
- Bossert, J. E., "An investigation of flow regimes affecting the Mexico City region" *Journal of Applied Meteorology*, **36**, 119-140 (1997).
- Bossert, J. E., Poulos, G.S., "A numerical investigation of mechanisms affecting drainage flows in highly complex terrain," *Theoretical and Applied Climatology*, **52**, 119-134 (1995).
- Bossert, J. E., Cotton, W.R., "Regional-scale flows in mountainous terrain. Part I: A numerical and observational comparison," *Monthly Weather Review*, **122**, 1449-1471 (1994a).
- Bossert, J. E., Cotton, W.R., 1994, "Regional-scale flows in mountainous terrain. Part II: Simplified numerical experiments" *Monthly Weather Review*, **122**, 1472-1489 (1994b).
- Dobran, F., Neri, A., Macedonio, G., "Numerical simulation of collapsing volcanic columns," *Journal of Geophysical Research*, **98**, 4231-4259 (1993).
- Dobran, F., Neri, A., Todesco, M. (1994) "Assessing the pyroclastic flow hazard at Vesuvius," *Nature*, **367**, 551-554 (1994).
- Kieffer, S.W., "Numerical-models of caldera-scale volcanic eruptions on Earth, Venus, and Mars," *Science*, **269**, 1385-1391 (1995).
- Margolin, L., Reisner, J., Smolarkiewicz, P.J., "Application of the volume of fluid method to the advection-condensation problem," Submitted to *Monthly Weather Review* (1996).
- Neri, A., Dobran, F., "Influence of eruption parameters on the thermofluid dynamics of collapsing volcanic columns." *Journal of Geophysical Research*, **99**, 11833-11857 (1994).
- Pielke, R.A., W.R. Cotton, R.L. Walko, C.J. Tremback, W.A. Lyons, L.D. Grasso, M.E. Nicholls, M.D. Moran, D.A. Wesley, T.J. Lee, and J.H. Copeland, "A comprehensive meteorological modeling system -- RAMS," *Meteorology and Atmospheric Physics*, **49**, 69-91 (1992).

- Poulos, G.S., Bossert, J.E., "An observational and prognostic numerical investigation of complex terrain dispersion," *Journal of Applied Meteorology*, **34**, 650-669, (1995).
- Valentine, G.A., Wohletz, K.H., "Numerical models of Plinian eruption columns and pyroclastic flows," *Journal of Geophysical Research*, **94**, 1867-1887 (1989a).
- Valentine, G.A., Wohletz, K.H., "Environmental hazards of pyroclastic flows determined by numerical models," *Geology*, **17**, 641-644 (1989b).
- Valentine, G.A., Wohletz, K.H., Kieffer, S.W., "Sources of unsteady column dynamics in pyroclastic flow eruptions," *Journal of Geophysical Research*, **96**, 21887-21892 (1991).
- Valentine, G.A., Wohletz, K.H., Kieffer, S.W., "Effects of topography on facies and compositional zonation in caldera-related ignimbrites," *Geological Society of America Bulletin*, **104**, 154-165 (1992).
- Wohletz, K.H., McGetchin, T.R., Sandford, M.T., II, Jones, E.M., "Hydrodynamic aspects of caldera-forming eruptions: numerical experiments," *Journal of Geophysical Research*, **89**, 8269-8286 (1984).