

Thermochemistry of the Ethyl Radical and the C–H Bond Strength in Ethane

Paul Marshall

Department of Chemistry, University of North Texas, P.O. Box 305070, Denton, Texas 76203-5070

Received: April 12, 1999

CCSD(T) theory has been applied to the ethyl radical using aug-cc-pVDZ, cc-pVTZ, and cc-pVQZ basis sets, and extrapolated to the complete basis set limit. QCISD/6-311G(d,p) results were used to estimate unknown frequencies in C₂H₅, and zero-point energies were combined with a correction for core electron correlation to find a 0 K bond dissociation enthalpy of 416.0 kJ mol⁻¹ (422.5 kJ mol⁻¹ at 298 K). This corresponds to Δ_fH(C₂H₅) values of 131.5 and 120.5 kJ mol⁻¹ at 0 and 298 K, respectively. Uncertainties of ±2.1 kJ mol⁻¹ are assumed.

1. Introduction

The thermochemistry of the ethyl radical establishes the carbon–hydrogen bond dissociation enthalpy (BDE) in ethane



through the relation

$$\text{BDE} = \Delta_f H(\text{H}) + \Delta_f H(\text{C}_2\text{H}_5) - \Delta_f H(\text{C}_2\text{H}_6) \quad (2)$$


The enthalpies of formation of H and C₂H₆ are well-defined. The strength of the simplest primary C–H bond is of fundamental importance to chemistry, and there have been several experimental determinations, some of which are listed in Table 1. There has been controversy surrounding some of the kinetic determinations. McMillen and Golden¹ selected a value based on the rate constant *k*₃ for



combined with assumed parameters for the reverse rate constant *k*₋₃. That value was called into question by the work of Tsang² and Pacey and Wimalasena³ based on thermal decomposition of *n*-butane




Cao and Back⁴ considered the forward and reverse kinetics of the system



and also recommended a more positive Δ_fH(C₂H₅). Hanning-Lee et al.⁵ obtained the forward and reverse rate constants for the reaction



and thus the equilibrium constant *K*₆ = *k*₆/*k*₋₆ as a function of temperature. Ruscic et al.⁶ noted that some of these results were apparently inconsistent with the then current ionization potential (IP) of ethyl and value of Δ_fH(C₂H₅⁺), and redetermined the IP to obtain Δ_fH(C₂H₅). Seakins et al.⁷ and Nicovich et al.⁸ investigated the kinetics of




in both directions, and derived similar bond strengths via the equilibrium constant *K*₇. Recently, Dobis and Benson^{9,10} have criticized those studies of reaction 7, and measured *k*₇ values about an order of magnitude smaller. As Dobis and Benson noted,⁹ there is a similarly wide range in the literature values of *k*₋₇, which allows for a significant range in *K*₇ and therefore Δ_fH(C₂H₅).

The aim of the present work is to obtain the C₂H₅–H bond strength via high-level ab initio calculations using coupled cluster theory with basis sets of up to quadruple-ζ quality, extrapolated to the complete basis set limit. The results permit an independent assessment of the experimental determinations.

2. Methodology

The BDE could in principle be obtained directly via the energy change for reaction 1. A more accurate result is expected from the isodesmic process



where, because the number and type of each bond are conserved, errors arising from basis set incompleteness and incomplete accounting of electron correlation mostly cancel. The geometry and frequencies of each species were first calculated at the QCISD/6-311G(d,p) level of theory. Next, the geometries and energies were obtained using CCSD(T) theory with three basis sets, aug-cc-pVDZ, cc-pVTZ, and cc-pVQZ.^{11,12} Core electrons were excluded from the correlation treatments in all these calculations. To correct for this simplification, two energies at the final geometries were also obtained with a full double-ζ basis set augmented with polarization functions, D95(3df,2p),¹³ where the core electrons were included and then excluded in the CCSD(T) calculations. All computations were made with the Gaussian 94 suite of programs.¹⁴

3. Results and Discussion

The computed geometries for C₂H₅ and C₂H₆ are summarized in Figure 1 and Table 2. By comparison with the equilibrium (*r*_e) structures for C₂H₂ and C₂H₄, Peterson and Dunning¹⁵ found C–C and C–H bond length errors at the CCSD(T)/cc-pVQZ level of theory of around 0.003 Å (1 Å = 10⁻¹⁰ m), and an


TABLE 1: Some Values of Ethyl Thermochemistry and the C₂H₅-H BDE^a

$\Delta_f H_0$	$\Delta_f H_{298}$	BDE ₀	BDE ₂₉₈	method	ref
132.2 ± 2.4	108.4 ± 4.2	416.6 ± 2.4	410.9 ± 4.2	I + C ₂ H ₆ ⇌ HI + C ₂ H ₅	1
	119 ± 2		<i>n</i> -C ₄ H ₁₀ ⇌ 2 C ₂ H ₅	2,3	
	117		H + C ₂ H ₆ ⇌ H ₂ + C ₂ H ₅	4	
	120.2 ± 0.9		H + C ₂ H ₄ ⇌ C ₂ H ₅	5	
			C ₂ H ₅ → C ₂ H ₅ ⁺	6	
131.5 ± 2.1	121.9 ± 1.8	416.0 ± 2.1		HBr + C ₂ H ₅ ⇌ Br + C ₂ H ₆	7
	121.8 ± 2.1		423.8 ± 2.5	HBr + C ₂ H ₅ ⇌ Br + C ₂ H ₆	8
	118.8 ± 1.0		420.5 ± 1.3	HBr + C ₂ H ₅ ⇌ Br + C ₂ H ₆	9
	120.5 ± 2.1		422.5 ± 2.1	CCSD(T) theory	this work

^a In kJ mol⁻¹.**TABLE 2: CCSD(T) Geometries of C₂H₅ and C₂H₆^a**

parameter	ethyl (C _s symmetry)			ethane (D _{3d} symmetry)		
	aug-cc-pVDZ	cc-pVTZ	cc-pVQZ	aug-cc-pVDZ	cc-pVTZ	cc-pVQZ
CH _a ^b	1.112	1.099	1.098	1.106	1.092	1.091
CH _b ^b	1.106	1.092	1.091			
CH _c ^b	1.095	1.081	1.080			
CC ^b	1.506	1.492	1.489	1.540	1.529	1.526
CCH _a ^c	111.60	111.67	111.63	111.15	111.20	111.21
CCH _b ^c	111.44	111.52	111.57			
CCH _c ^c	120.78	120.74	120.80			
H _b CCH _a ^d	±119.42	±119.42	±119.36			
H _c CCH _a ^d	±85.06	±85.00	±85.44 ^e			

^a Atom labeling defined in Figure 1. ^b Bond length in 10⁻¹⁰ m. ^c Bond angle in degrees. ^d Dihedral angle in degrees. ^e Convergence to 0.5° accepted for this parameter, rather than default of 0.1°.

**Figure 1.** Structures of ethyl (C_s) and ethane (D_{3d}). Parameters given in Table 2.

error of 0.03° in the ethylene HCH angle. Presumably similar uncertainties apply here.

Frequency information is given in Table 3. Where available, experimental fundamental frequencies ν_0 were employed to calculate vibrational zero-point energy contributions. All the C₂H₆ modes are known,¹⁶ but in the case of C₂H₅, three of the 15 normal modes have not been observed.¹⁷ Figure 2 is a plot of experimental versus QCISD/6-311G(d,p) frequencies of ethyl, which have been discussed previously,^{18,19} and ethane, together with a linear fit constrained to pass through the origin. The root-mean-square deviation is 39 cm⁻¹ and the slope is 0.956. This scale factor, which mainly accounts for anharmonicity, was used here to predict ν_0 for the three missing modes for ethyl, at 145 (torsion), 780, and 1426 cm⁻¹.

Although the thermochemical tables of Gurvich et al.¹⁶ for C₂H₅ are based on a significant torsional barrier, there is evidence that the barrier is small. Cohen²⁰ used 0.71 kJ mol⁻¹, Suter and Ha²¹ obtained a zero-point energy corrected value of 0.31 kJ mol⁻¹ from MP2/6-31G(d,p) theory, East and Bunker²² found the same value from MP2/cc-pVTZ theory, and Tirtowidjojo et al.¹⁰ derived 0.56 kJ mol⁻¹ from fitting a sinusoidal potential to the QCISD/6-311G(d,p) harmonic frequency. In this

TABLE 3: QCISD/6-311G(d,p) Frequencies (in cm⁻¹) for Ethyl and Ethane, Scaled by 0.956

C ₂ H ₅	symmetry	C ₂ H ₆	symmetry
145	A''	311	A _{1u}
435	A'	795 (2) ^a	E _u
780	A''	981	A _{1g}
958	A'	1189 (2)	E _g
1030	A'	1364	A _{2u}
1161	A''	1392	A _{1g}
1362	A'	1455 (2)	E _g
1426	A'	1459 (2)	E _u
1440	A'	2918	A _{2u}
1442	A''	2921	A _{1g}
2871	A'	2976 (2)	E _g
2946	A'	2998 (2)	E _u
2987	A''		
3026	A'		
3124	A''		

^a Doubly degenerate.

work, QCISD(T)/6-311+G(3df,2p)//QCISD/6-311G(d,p) calculations at the minimum energy geometry and the transition state for internal rotation yield an electronic energy barrier of 0.30 kJ mol⁻¹. This falls to only 0.08 kJ mol⁻¹ after correcting for changes in zero-point energy. Recent high-resolution IR spectroscopy measurements²³ are consistent with a low rotational barrier, between 0.1 and 0.5 kJ mol⁻¹. For the purpose of calculating the temperature dependence of the enthalpy of ethyl, its torsion is therefore treated as free, with a symmetry number of 6 and a reduced moment of inertia, based on the CCSD(T)/cc-pVQZ geometry, of 2.10 × 10⁻⁴⁷ kg m². With use of the rigid rotor/harmonic oscillator approximation for the other motions of C₂H₅, H_{298,15}-H₀ is calculated to be 12.18 kJ mol⁻¹, which lies between the Gurvich et al.¹⁶ value of 11.81 kJ mol⁻¹ and the Thermodynamics Research Center tabulation of 12.30 kJ mol⁻¹.²⁴ On the same basis, the entropy S_{298,15} is found to be 247.7 J K⁻¹ mol⁻¹, which falls between the values employed by Nicovich et al.⁸ and Dobis and Benson,⁹ of 246.8 and 249.4 J K⁻¹ mol⁻¹, respectively.

The ab initio energies are listed in Table 4 and are used to

TABLE 4: Energies for Species in the Reaction $C_2H_6 + CH_3 \rightarrow C_2H_5 + CH_4^a$

calculation	$C_2H_6(^1A_{1g})/au$	$CH_3(^2A_2'')/au$	$C_2H_5(^2A')$ /au	$CH_4(^1A_1)/au$	$\Delta E/kJ mol^{-1}$
CCSD(T)/aug-cc-pVDZ	-79.597 970	-39.724 710	-78.931 486	-40.395 820	-12.15
CCSD(T)/cc-pVTZ	-79.674 445	-39.760 976	-79.002 506	-40.438 099	-13.61
CCSD(T)/cc-pVQZ	-79.698 718	-39.772 449	-79.025 724	-40.450 888	-14.30
CBS limit ^b					-14.90
CCSD(T)=full/D95(3df,2p)	-79.686 040	-39.765 936	-79.015 700	-40.442 112	-15.32
CCSD(T)=fc/D95(3df,2p)	-79.649 064	-39.748 112	-78.979 223	-40.423 535	-14.66
ZPE ^c	0.072 283	0.028 990	0.057 629	0.043 224	-1.10
ΔH_0					-16.67

^a In au, 1 au \approx 2625.3 kJ mol⁻¹. ^b Complete basis set limit. ^c Zero-point energy.




Figure 2. Comparison of calculated harmonic QCISD/6-311G(d,p) and observed fundamental vibrational frequencies for ethyl and ethane.

derive the energy change ΔE for reaction 8 at various levels of theory. The complete basis set limit was obtained by fitting the CCSD(T)/cc-pVnZ values to the functional form $\Delta E = \Delta E_\infty + A \exp(-Bn)$ and extrapolating to infinite n , i.e., to ΔE_∞ .²⁵ Two corrections were applied to this extrapolated ΔE . The first was for the neglect of core correlation, equal to $\Delta E[\text{CCSD(T)=full}] - \Delta E[\text{CCSD(T)=fc}]$ with the D95(3df,2p) basis set, which came to -0.67 kJ mol⁻¹. The second correction was for changes in vibrational zero-point energy, here set equal to $1/2 \sum h\nu_0$ where ν_0 is obtained from the experimental fundamental frequencies (except for the three missing modes for C_2H_5 discussed above). After these corrections, ΔE equals ΔH_0 for reaction 8, where

$$\Delta H_0 = \text{BDE}_0(C_2H_5-H) - \text{BDE}_0(CH_3-H) \quad (8)$$

Litorja and Ruscic²⁶ have recently measured $\text{BDE}_0(CH_3-H)$ to high precision, 432.64 ± 0.29 kJ mol⁻¹; hence $\text{BDE}_0(C_2H_5-H) = 415.96$ kJ mol⁻¹ is obtained. The present value of $H_{298.15} - H_0$ for C_2H_5 and tabulated values^{16,27} for C_2H_6 and H of 11.88 and 6.20 kJ mol⁻¹ imply $\text{BDE}_{298.15}(C_2H_5-H) = 422.46$ kJ mol⁻¹. These bond dissociation enthalpies can be combined with the enthalpies of formation of C_2H_6 and H, -68.38 and 216.04 kJ mol⁻¹ at 0 K and -84.00 and 218.00 kJ mol⁻¹ at 298.15 K, respectively,^{16,27} using eq 2, to find the $\Delta_f H(C_2H_5)$ values listed in Table 1.

The least certain input parameter is $\Delta_f H(C_2H_6)$ but the quoted uncertainty is only ± 0.4 kJ mol⁻¹.¹⁶ An earlier recommendation is 0.75 kJ mol⁻¹ more negative.²⁸ Because the zero-point energies mostly cancel for reaction 8, uncertainties in frequencies have little influence. Because the ΔE for reaction 8 is close to converged, the basis set extrapolation introduces little uncertainty: use of an alternative formula²⁵ $\Delta E = \Delta E_\infty + A/(l + 1/2)^4 + B/(l + 1/2)^6$ (where l is the highest angular

momentum function, 2, 3, and 4 here) leads to $\Delta E_\infty = -14.81$ kJ mol⁻¹ instead. In order to estimate the reliability of the derived $\Delta_f H(C_2H_5)$ values, it is noted that, using a similar computational approach applied to BDEs in the series CH_n and C_2H_n ($n \leq 4$), Peterson and Dunning¹⁵ found errors of up to 0.8 and 1.7 kJ mol⁻¹, respectively, by comparison with experiment. They expected accuracies of 2.1 kJ mol⁻¹ for two unknown C_2H_n BDEs, and this same error limit is used here. This may be somewhat pessimistic: the earlier work¹⁵ was not able to take advantage of the cancellations in isodesmic reactions, and there were changes in the C-C bond order.

The thermochemistry for reaction 8 can be compared to results from the Gaussian-x series of model chemistries. Ruscic et al.⁶ reported $\Delta H_0 = -13.0$ kJ mol⁻¹ from G1 theory. Use of G2 energies²⁹ in reaction 8 yields $\Delta H_0 = -13.7$ kJ mol⁻¹, and G3 data^{30,31} imply $\Delta H_0 = -12.9$ kJ mol⁻¹. The last calculation includes core electron correlation, like the present work. Those results are around 3–4 kJ mol⁻¹ more positive than our best calculation and would therefore lead to a similar deviation in $\Delta_f H(C_2H_5)$. We are applying Gx methods to the thermochemistry of larger alkyl radicals and the present results provide a calibration of the derived bond strengths.

Our computed thermochemistry may also be compared to the various experimental results in Table 1. Clearly it supports the newer measurements which are more positive than the results from iodination kinetics. There is accord within the uncertainty with the results of *n*-butane dissociation, and exceptionally close agreement with the results based on ionization of ethyl and on the kinetics of dissociation of ethyl radicals (combined with the reverse process, addition of H atoms to ethylene), to within 0.7 kJ mol⁻¹. The deviation of 3.5 kJ mol⁻¹ from the value based on H + C_2H_6 kinetics suggests there might be some moderate but unresolved discrepancies between forward and reverse rate constants in that system. The present results lie within the small range of thermochemistry based on HBr + $C_2H_5 \rightleftharpoons Br + C_2H_6$ kinetics. However, the narrowness of this range is fortuitous, because of the wide disparities in forward and reverse rate constants noted in the Introduction. Different selections lead to different equilibrium constants and a range of over 11 kJ mol⁻¹ in the derived $\Delta_f H(C_2H_5)$.⁹ The present work does not determine the rate constants, but does fix their ratio. In combination with tabulated data for the other reactants,^{16,27} our thermochemical data for C_2H_5 implies $K_7 = 4.6 \times 10^7$ at 298 K. This is rather close to the value of $(5.1 \pm 2.5) \times 10^7$ derived⁹ from flash photolysis measurements of k_7 and k_{-7} .^{7,8} The uncertainty assumed in $\Delta_f H(C_2H_5)$ implies about a factor of 2 uncertainty in the present K_7 at 298 K, which therefore accommodates some of the k_7 and k_{-7} results from VLPP and thermal measurements that yielded K_7 in the range $(1.2-3.0) \times 10^7$.⁹ With our computed K_7 , if an unambiguous determination of k_7 or k_{-7} can be made, then much of the uncertainty in the remaining quantity will also be resolved.

Conclusions

Coupled cluster theory has been combined with extrapolation of correlation consistent basis sets and a core electron correction to obtain the C_2H_5-H bond dissociation enthalpy at the complete basis set limit. The value is in good accord with several recent experiments, and constrains the ratio of forward and reverse kinetics for the $Br + C_2H_6$ system. The barrier to internal rotation in ethyl is found to be negligible, about 0.1 kJ mol^{-1} .

Acknowledgment. Dr. Branko Ruscic is thanked for valuable discussions. This work was supported by the R. A. Welch Foundation (Grant B-1174) and the UNT Faculty Research Fund, and computer facilities were provided in part by the ASC/MSRC supercomputing center and the Materials and Manufacturing Division, Air Force Research Laboratory, both at Wright-Patterson AFB, Ohio.

References and Notes

- (1) McMillen, D. F.; Golden, D. M. *Annu. Rev. Phys. Chem.* **1982**, *33*, 493.
- (2) Tsang, W. *Int. J. Chem. Kinet.* **1978**, *10*, 821.
- (3) Pacey, P. D.; Wimalasena, J. H. *J. Phys. Chem.* **1984**, *88*, 5657.
- (4) Cao, J.-R.; Back, M. H. *Int. J. Chem. Kinet.* **1984**, *16*, 961.
- (5) Hanning-Lee, M. A.; Green, N. J. B.; Pilling, M. J.; Robertson, S. H. *J. Phys. Chem.* **1993**, *97*, 860.
- (6) Ruscic, B.; Berkowitz, J.; Curtiss, L. A.; Pople, J. A. *J. Chem. Phys.* **1989**, *91*, 114.
- (7) Seakins, P. W.; Pilling, M. J.; Niiranen, J. T.; Gutman, D.; Krasnoperov, L. N. *J. Phys. Chem.* **1992**, *96*, 9847.
- (8) Nicovich, J. M.; van Dijk, C. A.; Kreutter, K. D.; Wine, P. J. *J. Phys. Chem.* **1991**, *95*, 9890.
- (9) Dobis, O.; Benson, S. W. *J. Phys. Chem. A* **1997**, *101*, 6030.
- (10) Benson, S. W.; Dobis, O. *J. Phys. Chem. A* **1998**, *102*, 5175.
- (11) Dunning, Jr., T. H. *J. Chem. Phys.* **1989**, *90*, 1007.
- (12) Kendall, R. A.; Dunning, Jr., T. H.; Harrison, R. J. *J. Chem. Phys.* **1992**, *96*, 6769.
- (13) Dunning, Jr., T. H.; Hay, P. J. In *Modern Theoretical Chemistry*; Schaefer, H. F., III, Ed.; Plenum: New York, 1976; pp 1–28.
- (14) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T. A.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. J. P.; Head-Gordon, M.; Gonzalez, C.; Pople, J. A. *GAUSSIAN 94*; Gaussian, Inc.: Pittsburgh, PA, 1995.
- (15) Peterson, K. A.; Dunning, Jr., T. H. *J. Chem. Phys.* **1997**, *106*, 4119.
- (16) Gurvich, L. V.; Veyts, I. V., Alcock, C. B., Eds. *Thermodynamic Properties of Individual Substances*, 4th ed.; Hemisphere: New York, 1992; Vol. 2.
- (17) Jacox, M. E. *J. Phys. Chem. Ref. Data* **1998**, *27*, 115.
- (18) Tirtowidjojo, M. M.; Colegrove, B. T.; Durant, J. L., Jr. *Ind. Eng. Chem. Res.* **1995**, *34*, 4202.
- (19) Hase, W. L.; Schlegel, H. B.; Balbyshev, V.; Page, M. *J. Phys. Chem.* **1996**, *100*, 5354.
- (20) Cohen, N. *J. Phys. Chem.* **1992**, *96*, 9052.
- (21) Suter, H. U.; Ha, T.-K. *Chem. Phys. Lett.* **1991**, *154*, 227.
- (22) East, A. L. L.; Bunker, P. R. *Chem. Phys. Lett.* **1998**, *282*, 49.
- (23) Sears, T. J.; Johnson, P. M.; Jin, P.; Oatis, S. *J. Chem. Phys.* **1996**, *104*, 781.
- (24) *TRC Thermodynamics Tables—Hydrocarbons*; Thermodynamics Research Center, Texas A & M University: Texas, 1983; table r-1930.
- (25) Martin, J. M. L. Calibration Study of Atomization Energies of Small Polyatomics. In *Computational Thermochemistry*; Irikura, K. K., Frurip, D. J., Eds.; ACS Symp. Ser. 677; American Chemical Society: Washington, DC, 1998; Chapter 12.
- (26) Litorja, M.; Ruscic, B. *J. Chem. Phys.* **1997**, *107*, 9852.
- (27) Gurvich, L. V.; Veyts, I. V., Alcock, C. B., Eds. *Thermodynamic Properties of Individual Substances*, 4th ed.; Hemisphere: New York, 1989; Vol. 1.
- (28) Wagman, D. D.; Evans, W. H.; Parker, V. B.; Schumm, R. H.; Halow, I.; Bailey, S. M.; Churney, K. L.; Nuttall, R. L. *J. Phys. Chem. Ref. Data* **1982**, *11*, Suppl. 2. *The NBS Tables of Chemical Thermodynamic Properties*.
- (29) Curtiss, L. A.; Raghavachari, K.; Trucks, G. W.; Pople, J. A. *J. Chem. Phys.* **1991**, *94*, 7221.
- (30) Curtiss, L. A.; Raghavachari, K.; Redfern, P. C.; Rassolov, V.; Pople, J. A. *J. Chem. Phys.* **1998**, *109*, 7764.
- (31) <http://chemistry.anl.gov/compmat/g3theory.htm>.