MIT/WHOI 95-20

DOE/OR/00033--T754

**Massachusetts Institute of Technology
Woods Hole Oceanographic Institution**

**Joint Program
in Oceanography/
Applied Ocean Science
and Engineering**

RECEIVED
JUL 11 1997
OSTI

DOCTORAL DISSERTATION

*Development and Application of the Mollusc
Arctica islandica as a Paleoceanographic Tool
for the North Atlantic Ocean*

by

Christopher R. Weidman

September 1995

MIT/WHOI

95-20

*Development and Application of the Mollusc
Arctica islandica as a Paleoceanographic Tool
for the North Atlantic Ocean*

by

Christopher R. Weidman

Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

and

Woods Hole Oceanographic Institution
Woods Hole, Massachusetts 02543

September 1995

MASTER

DOCTORAL DISSERTATION

Funding was provided by the National Science Foundation through Grant No. OCE9301015.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Reproduction in whole or in part is permitted for any purpose of the United States Government.
This thesis should be cited as: Christopher R. Weidman, 1995. Development and Application of the
Mollusc *Arctica islandica* as a Paleoceanographic Tool for the North Atlantic Ocean. Ph.D. Thesis.
MIT/WHOI, 95-20.

Approved for publication; distribution unlimited.

Approved for Distribution:

William B. Curry, Chair
Department of Geology and Geophysics

Marcia K. McNutt
MIT Director of Joint Program

John W. Farrington
WHOI Dean of Graduate Studies

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

**Portions of this document may be illegible
electronic image products. Images are
produced from the best available original
document.**

**Development and Application of the Mollusc *Arctica islandica*
as a Paleoceanographic Tool for the North Atlantic Ocean**

by

Christopher R. Weidman

B.S. , State University of New York College at Oneonta, New York
(1988)

Submitted in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

at the

Massachusetts Institute of Technology

and the

Woods Hole Oceanographic Institution

The Government reserves for itself and
September 1995 others acting on its behalf a royalty free,
nonexclusive, irrevocable, world-wide
license for governmental purposes to publish,
distribute, translate, duplicate, exhibit,
and perform any such data copyrighted by
the contractor.

Signature of Author.....

Joint Program in Marine Geology and Geophysics
Massachusetts Institute of Technology
Woods Hole Oceanographic Institution

Certified by.....

Glenn A. Jones
Thesis Supervisor

Accepted by.....

Deborah K. Smith
Chair, Joint Committee for Marine Geology and Geophysics
Massachusetts Institute of Technology
Woods Hole Oceanographic Institution

Development and Application of the Mollusc *Arctica islandica* as a Paleoceanographic Tool for the North Atlantic Ocean

by

Christopher R. Weidman

Submitted to the Massachusetts Institute of Technology and the Woods Hole Oceanographic Institution Joint Program in Oceanography/Applied Ocean Science and Engineering on August 10, 1995, in partial fulfillment of the requirements for a degree of Doctor of Philosophy.

ABSTRACT

Until now there has been no tool comparable to corals for reconstructing long term high-resolution geochemical time-series for the colder, higher-latitude oceans. In this thesis, the long-lived (+100 years) boreal mollusc (Bivalvia) *Arctica islandica* is shown to be practical for this purpose in the northern North Atlantic Ocean. The evidence, compiled here, demonstrates that the carbonate shell of this species faithfully records the ambient dissolved inorganic carbon's (DIC) radiocarbon ($\Delta^{14}\text{C}$) concentration and accurately reflects the ambient temperature in its stable oxygen isotope ($\delta^{18}\text{O}$) composition. However, the stable carbon isotope ($\delta^{13}\text{C}$) composition of the *A. islandica* shell likely is not a good recorder of ambient DIC $\delta^{13}\text{C}$, and likely responds to physiological controls. Four $\Delta^{14}\text{C}$ time histories are reconstructed from the annual bands of *A. islandica* shells for the higher-latitudes of the northern North Atlantic Ocean (from 41°N to 70°N). These ocean records show significant spatial and temporal differences in the evolution of the radiocarbon signal between the subpolar and subtropical regions and between eastern and western regions of the northern North Atlantic, which are attributed to regional differences in mixed-layer depth and the presence of deepwater sources. A 109-year interannual record of bottom temperatures for a location near the former Nantucket Lightship position has been reconstructed for the period 1875-1983 from the overlapped stable oxygen isotope composition of four *A. islandica* shells. This record's annual temperature anomalies (variation from the running ten-year mean) show significant positive correlation with regional bottom, sea surface and air temperature anomalies. However, the shell-derived bottom temperature record describes a century-long cooling ($\sim 1^\circ\text{C}$) in contrast to a century-long warming of regional sea surface temperatures of equal magnitude, indicating a long term divergence between surface and

bottom conditions. It is suggested that this contrast may be owed to a reduction in vertical mixing and increased seasonal stratification of shelf waters. This thesis fulfills the prophesy laid out nearly two decades ago by Thompson and Jones [1977] that *Arctica islandica* could someday be used to reconstruct past ocean history as "the *tree* of the North Atlantic".

Thesis Supervisor: Dr. Glenn A. Jones

Title: Associate Scientist, Woods Hole Oceanographic Institution

Acknowledgements

Every thesis owes its inspiration and much of its manifestation to others, animate and inanimate. Thanks are owed to Woods Hole for its incredible community of people whose vision and sharing have created a rich climate for learning and a wonderful place for my kids to grow up. I would also like to thank Nobska, Stony, and Gansett beaches, Eel Pond bridge, Community Hall, the Wall, Coffee at Pie in the Sky, the Martha's Vineyard Ferry, Surf Drive, Shining Sea Bike Path, and the strong westerlies which whistle about the MBL Library and make it hard to leave.

Glenn Jones, my thesis advisor, has provided strong guidance and encouragement throughout this project. I also thank him for his fountain of ideas, sense of the big picture, and his eye for detail. However, I want to thank him most for our late afternoon discussions when we lost track of time (apologies to our families).

I thank the members of my thesis committee for their time and advice: Dan McCorkle, Pat Lohmann, Ed Boyle, Steve Murawski, Dave Chapman, and Glen Gawarkiewicz. A special thanks is owed to K. C. Lohmann for one incredible and largely sleepless week that helped launch a good half of this thesis. I thank, also, some of my most important mentors and counselors, official and unofficial, over the years at WHOI and MIT: Dave Aubrey, John Southard, Ole Madsen, Scott Lehman, Bill Curry, Al Uchupi, and Graham Giese.

I owe more than I know to John, Jake, Abbie, Stella, Kathy, and Marcy, for not making the Education Office a bureaucracy and for covering for my wayward ways.

I thank all my Joint Program friends here at WHOI who have made the heavy loads bearable with much humor and camaraderie, and single out three examples: Gary Jaroslow for his companionship and fellow second incarnation perspective (Gary, schnapps or coffee?), Carl Friedrichs for his excellence and permanent optimism (Carl, how do you do this?), and Javier Escartin for his Catalan sense of the absurd (I can't #!*\$!! believe it). I thank my buddies at NOSAMS: Kathy, Ann, Bob, Susan, Liz, Dan, Karl, John, Greg and especially my officemate Al Gagnon for his friendly patience and ability to keep the mass spectrometers running.

I reserve my deepest thanks to my family. This has been a long road guys, and each of you has helped me every step of the way. To my mom and dad for their infectious love of science and chronic curiosity, where all of this began afterall. To Jean and to Bernie (who I miss very much) for their example of never too late and don't be afraid - just go ahead and do it. To my children, Dylan, Lailye and Kaya, of whom I am very proud for becoming independent, honest, and voracious consumers of life, whose lives have much to teach me. And finally, to Melissa Rose, the great love of my life and companion in this ever unfolding adventure, I whisper, "the tomatoes are ripe, babe, what's next ?"

Funding was partly provided by an ORAU Global Climate Change Fellowship.

CONTENTS

Abstract	2
Acknowledgements	4
Introduction	9
References	13

PART I. Development

Chapter	1.	A Shell-Derived Time History of Bomb- ¹⁴ C on Georges Bank and Its Labrador Sea Implications	17
		- Reprint - Removed for separate processing	
		Abstract	18
		Introduction	18
		Significance of Bomb- ¹⁴ C Studies	18
		Using Molluscs to Derive Ocean Radiocarbon Records	18
		Methods	19
		Results	19
		Discussion	21
		Comparison with North Atlantic Coral-Derived $\Delta^{14}\text{C}$ Records	21
		Georges Bank-Labrador Sea	23
		Georges Bank $\Delta^{14}\text{C}$ Budget	23
		Estimate of the Bomb- ¹⁴ C Time History for the Labrador Sea	24
		Gulf Stream and Antarctic Intermediate Water	25
		Ventilation/ ¹⁴ CO ₂ Uptake in the Labrador Sea	26
		Conclusions	27
		References	27

Chapter	2.	The Long-Lived Mollusc <i>Arctica islandica</i> : A New Paleoceanographic Tool for the Reconstruction of Bottom Temperatures for the Continental Shelves of the Northern North Atlantic Ocean	30
		Abstract	31
		Introduction	31
		Methods	32

Nantucket Shoals Lightship T and S Records	32
Predicted Aragonite $\delta^{18}\text{O}$ Record	34
Shell Sampling and Analysis	34
Results and Discussion	36
Conclusions	39
References	39
 Chapter 3. Stable Oxygen and Carbon Isotope Records from <i>Arctica islandica</i>	
Shells in a Monitored Laboratory Growth Experiment	41
Introduction	41
Methods	42
Laboratory Growth Experiment	42
Offshore Site Monitoring	43
Sample Preparation and Stable Isotope Analysis	44
Results	45
Water Column Properties	45
Temperature	45
Salinity	46
$\delta^{18}\text{O}$ -water	46
DIC $\delta^{13}\text{C}$	46
Pore Water DIC	47
Devil's Bridge Properties (natural cycle)	48
Shell Records	49
Ring Structure and Shell Growth	49
$\delta^{18}\text{O}$ -shell Record	50
$\delta^{13}\text{C}$ -shell Record	51
Discussion	52
$\delta^{18}\text{O}$ -aragonite	52
$\delta^{13}\text{C}$ -aragonite	57
Microhabitat / Pore Water $\delta^{13}\text{C}$	59
Physiological (Vital) Effects	60
Summary	63
Acknowledgements	65
References	66
Tables	69
Figures	73

PART II. Application

Chapter 4.	Time History of $\Delta^{14}\text{C}$ in the Surface North Atlantic Ocean	89
	Introduction	89
	General Circulation of North Atlantic and the $\Delta^{14}\text{C}$ Signal	91
	Coastal Environments and the $\Delta^{14}\text{C}$ Signal	93
	Methods	96
	Results	97
	Eastern North Atlantic $\Delta^{14}\text{C}$ Time Histories	97
	North Sea	97
	Iceland	98
	Norway	98
	Western North Atlantic $\Delta^{14}\text{C}$ Time Histories	99
	Bermuda	99
	Florida	99
	Georges Bank	100
	Discussion	100
	North Atlantic Pre-Bomb $\Delta^{14}\text{C}$	100
	Time History Comparison	100
	Comparison with Other Pre-bomb $\Delta^{14}\text{C}$ Measurements	101
	Suess Effect	107
	1950s Extrema: Norway and North Sea	108
	North Atlantic Post-Bomb $\Delta^{14}\text{C}$	110
	Time History Comparison	110
	Comparison with Other Post-bomb $\Delta^{14}\text{C}$ Measurements	113
	Determining the Offshore $\Delta^{14}\text{C}$ Signal	113
	North Sea	115
	Norway	116
	The Evolution of the Bomb- ^{14}C Signal in the North Atlantic	117
	Summary	119
	Acknowledgements	121
	References	122
	Tables	127
	Figures	132

Chapter 5.	A Century-Long Record of Bottom Temperatures Near Nantucket Shoals from Stable Oxygen Isotopic Analysis of <i>Arctica islandica</i> Shells	147
	Introduction	147
	Methods	148
	Location	148
	Sampling and Stable isotope Analysis	149
	Chronology	150
	Results	150
	Data	150
	Overlap Comparison and Chronology Check	151
	Band Width / Age Corrections	152
	T / $\delta^{18}\text{O}$ -aragonite and T / S / $\delta^{18}\text{O}$ -water Relationships	154
	The Combined Shell-Derived Temperature Records	156
	BTmean	157
	BTmax	158
	BTmin	158
	Discussion	159
	Nantucket Lightship Bottom Temperature Records	159
	Nantucket Lightship Sea Surface Temperature Records	160
	Brenton Reef Lightship and Boothbay Harbor SST Records	161
	Coastal Northeast U.S. Air Temperature Record	164
	Comparison of Detrended Records	164
	The Long-Term Trend Problem	165
	Conclusions	168
	Acknowledgements	171
	References	172
	Tables	174
	Figures	180
Conclusions		192
Appendix		199

INTRODUCTION

The carbon and oxygen isotopic compositions of marine waters and calcareous skeletons of marine organisms have been used to provide important evidence of physical and biological oceanographic processes. The carbon isotopes are important tracers of the global carbon cycle [Broecker, & Peng, 1982], being used to measure the ocean's uptake of atmospheric carbon dioxide [Suess, 1953][Revelle, & Suess, 1957][Craig, 1957][Broecker, Peng, Ostlund, & Stuvier, 1985][Druffel, & Suess, 1983][Quay, Tilbrook, & Wong, 1992], as well as the ocean's physical circulation [Broecker, Gerard, Ewing, & Heezen, 1960][Broecker, et al., 1982][Ostlund, Possnert, & Swift, 1987][Druffel, 1989][Toggweiler, Dixon, & Broecker, 1991]. The oxygen isotopes provide a handle on global and regional climate through the hydrological cycle [Shackleton, 1977][Fairbanks, & Matthews, 1978][Broecker, et al., 1982][Dettman, & Lohmann, 1993] and the dependence of calcium carbonate's isotopic composition on ambient temperature at the time of calcification [Epstein, Buchsbaum, Lowenstam, & Urey, 1953][Grossman, & Ku, 1986].

The most direct method of measuring the carbon and oxygen isotopic composition of seawater is to collect and analyze seawater samples. Intensive large ocean studies, such as GEOSECS (1970s), TTO (1980s), and WOCE (1990s), along with more limited seawater sampling efforts, provide an important synoptic perspective on the ocean's geochemical distribution. However, seawater measurements are snapshots of the ocean's isotopic properties and, unless taken in sequence, provide no information on the temporal variability of these properties, and no direct information on past conditions. Also, in the surface ocean where biologic and climatic processes have strong short term impacts on some of these isotopic properties, seawater measurements do not necessarily reflect the long term averages at these sites [Broecker, & Peng, 1980].

Biogenic marine carbonate records the ambient seawater's carbon and oxygen isotopic composition at the time the carbonate was precipitated, though this relationship is not always simple [Wefer, & Berger, 1991]. Sclerochronology, the study of sequentially-deposited calcareous growth structures, and isotopic analysis can be combined to reconstruct seasonal and annual records of past ocean conditions. Many marine organisms possess skeletal growth structures which enable the time of carbonate deposition to be determined quite accurately [Clark, 1974][Rhoads, & Lutz, 1980][Jones, 1983]. In the lower latitudes, seasonal and interannual isotope records have been reconstructed by

sampling and analyzing the annual bands of large stony corals and sclerosponges [Nozaki, Rye, Turekian, & Dodge, 1978][Druffel, & Linick, 1978][Druffel, 1981][Druffel, et al., 1983][Patzoid, 1984][Druffel, 1985][Druffel, 1987][Patzoid, Ristedt, & Wefer, 1987][Druffel, 1989][Chakraborty, Ramesh, & Krishnaswami, 1994]. In this way, banded-corals have yielded important information on the variability of past ocean conditions, as long term high-resolution monitors of marine geochemistry and temperature for the surface subtropical oceans.

These methods have also been applied to molluscs, which are not latitudinally constrained, and the results have been used: 1) to study aspects of molluscan life history, such as, shell growth as a function of metabolism, ontogeny, and environment [Rhoads, et al., 1980][Erlenkeuser, & Wefer, 1981][Jones, Williams, & Romanek, 1986][Krantz, Williams, & Jones, 1987][Romanek, Jones, Williams, Krantz, & Radke, 1987][Landman, Druffel, Cochran, Donahue, & Jull, 1988]; and 2) to reconstruct environmental records of the ocean, such as, modern seasonal and interannual variations in ocean circulation, temperature, salinity, and primary productivity [Killingsley, & Berger, 1979][Williams, Arthur, Jones, & Healy-Williams, 1982][Arthur, Williams, & Jones, 1983][Jones, Williams, & Arthur, 1983][Jones, Williams, Arthur, & Krantz, 1984], as well as similar conditions and properties in the ancient oceans [Epstein, et al., 1953][Krantz, 1990].

Despite its widespread application, inherent difficulties arise in reconstructing and interpreting isotopic records obtained from molluscs. Many mollusc species have very limited geographical and ecological ranges, which limit their use as paleoenvironmental recorders. The carbonate mass of most molluscs is small (relative to corals) which limits sampling resolution. Shell growth rates often slow with age, further reducing sampling resolution in the outer (more recent) bands of older specimens. Also, about 80% of studied marine bivalves are relatively short-lived (less than 20 years) [Heller, 1990] compared to the colonial life-span of stony corals (several hundred years), meaning that most molluscs are inappropriate for reconstructing long term high-resolution isotope records.

Periodicity of the mollusc growth bands must be established for each species because uncorrected sub-annual or episodic banding can create chronological uncertainty [Jones, 1981]. Like most organisms, bivalves respond to a host of external and internal influences, and so their shells are often imperfect integrators of ambient conditions. For example, shell growth often slows or ceases altogether under conditions of extreme cold or warmth, low nutrient or oxygen supply, high levels of sediment suspension or predation,

or owing to ontogenic or sexual stage [Lutz, & Rhoads, 1977][Rhoads, et al., 1980][Jones, et al., 1984][Harrington, 1989][Krantz, Jones, & Williams, 1989]. Finally, many calcareous marine organisms, including molluscs, form carbonate in disequilibrium with the isotopic composition of ambient seawater, and while unusual for the oxygen isotopes this is often the rule for carbon isotopes [Wefer, et al., 1991].

All of these problems constrain the approach to the selection and use of molluscs as tools for reconstructing isotopic time histories in the oceans, and as a consequence, no comparable tool has been developed for the colder, higher latitude oceans that parallels the paleoceanographic role that banded-corals have provided for the warmer subtropical surface oceans. In this thesis, the mollusc *Arctica islandica* has been selected for this purpose, because of its unique life history and because recent technological advances have created the opportunity to overcome the sampling problems formerly presented by slow growth, long-lived molluscs. Furthermore, *A. islandica* inhabits the northern North Atlantic Ocean, a region thought to play a critical role in the regulation of global climate as a primary source of deep water for the world's ocean [Warren, 1981][Stocker, & Broecker, 1992][Broecker, 1991] and as an important sink for atmospheric carbon dioxide [Baes, Bjorkstrom, & Mulholland, 1985][Sarmiento, & Sundquist, 1992][Broecker, & Peng, 1992].

Over the past decades, there has been a growing scientific interest in *A. islandica*, initially, stemming from this species potential value as a commercial fishery [Nichol, 1951][Loosanoff, 1953][Merrill, & Ropes, 1969]. Thompson, et al., [Thompson, Jones, & Dreibelbis, 1980] and Jones [Jones, 1980] were the first to recognize the potential of the bivalve *A. islandica* as a marine geochemical recorder, calling it the "tree of the north Atlantic shelf" [Thompson, et al., 1977], due to its great longevity, with individual ages commonly in excess of 100 years (with one documented specimen aged at 221 years old [Jones, 1983]). These authors noted that *A. islandica*, commonly known as the ocean quahog, is a large (~100 mm in radial length) infaunal species which inhabits the continental shelves and slopes on both sides of the North Atlantic Ocean, spanning a wide depth and latitude range, from 10 - 200 m, and from Cape Hatteras (~35°N) to the Barent's Sea (~70°N). Their studies, along with the subsequent work of others [Murawski, Ropes, & Serchuk, 1982][Turekian, Cochran, Nozaki, Thompson, & Jones, 1982][Ropes, Jones, Murawski, Serchuk, & Jerald, 1984], provided evidence supporting the annual nature of *A. islandica*'s growth bands.

While several studies have used the shells of *A. islandica* to estimate ocean $\Delta^{14}\text{C}$ [Erlenkeuser, 1976][Turekian, et al., 1982][Tanaka, Monaghan, & Turekian, 1990], none of these studies was able to take advantage of this species' longevity, because of the large sample mass requirements of conventional beta decay radiocarbon methods. Also, sampling methods used to obtain seasonal and interannual isotopic profiles from corals [Nozaki, et al., 1978][Druffel, 1989] and other faster growing but shorter-lived molluscs, such as *Spisula solidissima* [Jones, et al., 1983] and *Tridacna maxima* [Romanek, et al., 1987], have not been applicable to *A. islandica* because of its slow linear growth rate.

So, although many fundamental aspects of *A. islandica*'s life history have been established over the last three decades, its predicted potential as a paleoceanographic tool has, until now, gone largely unrealized. Two recent technological advances have been central in altering this situation: 1) the introduction of accelerator mass spectrometry (AMS), which has reduced the sample size requirements for radiocarbon analysis by 3 orders of magnitude [Linick, Damon, Donahue, & Jull, 1989]; and 2) the development of computer - aided microsampling equipment with micron-scale resolution which has provided the means to profile small growth structures in molluscs with precision [Dettman, et al., 1993]. Applying this new technology to *A. islandica*, the objectives of this thesis are: 1) the development and calibration of *A. islandica* as a new paleoceanographic tool for reconstructing oxygen and carbon isotope time-series; and 2) the demonstration that long term annual and seasonal time-histories of these isotopes can be reconstructed and used to investigate the temporal and spatial variability of the northern North Atlantic Ocean's temperature, physical circulation, and uptake of atmospheric $^{14}\text{CO}_2$. The first objective motivates chapters 1-3, which discuss the validity and accuracy of this new tool. The second objective motivates chapters 4-5 (and part of chapter 1) which discuss 1) the evolution of the radiocarbon signal in the North Atlantic during the period of anthropogenic influence based on the reconstruction of four high-latitude $\Delta^{14}\text{C}$ time-histories, and 2) a century-long interannual record of bottom temperatures on the northeastern U.S. continental shelf. Regarding the format of the thesis, each chapter has been designed as a separate and distinct paper, though all five chapters are intimately linked to the overall objective of this thesis and provide a context and foundation for each other.

References

- Arthur, M. A., D. F. Williams and D. S. Jones, Seasonal temperature-salinity changes and thermocline development in the mid-Atlantic Bight as recorded by the isotopic composition of bivalves, *Geology*, *11*, 655-659, 1983.
- Baes, C. F. J., A. Bjorkstrom and P. J. Mulholland, Uptake of carbon dioxide by the oceans, in: *Atmospheric Carbon Dioxide and the Global Carbon Cycle*, edited by J. R. Trabalka, pp. 81-111, U.S. Dept. of Energy, Oak Ridge, TN, 1985.
- Broecker, W. S., The great ocean conveyor, *Oceanogr.*, *4*, 79-89, 1991.
- Broecker, W. S., R. Gerard, M. Ewing and B. C. Heezen, Natural radiocarbon in the Atlantic Ocean, *J. Geophys. Res.*, *65*, 2903-2931, 1960.
- Broecker, W. S., and T. H. Peng, Seasonal variability in the C14/C12 ratio for surface ocean water, *Geophys. Res. Lett.*, *7*, 1020-1022, 1980.
- Broecker, W. S., and T. H. Peng, *Tracers in the Sea*, Lamont-Doherty Geological Observatory Publication, Columbia University, Palisades, N.Y., 1982.
- Broecker, W. S., T. H. Peng, G. Ostlund and M. Stuvier, The distribution of bomb radiocarbon in the ocean, *J. Geophys. Res.*, *90*, 6953-6970, 1985.
- Broecker, W. S., and T. S. Peng, Interhemispheric transport of carbon dioxide by ocean circulation, *Nature*, *356*, 587-589, 1992.
- Chakraborty, S., R. Ramesh and S. Krishnaswami, Air-sea exchange of CO₂ in the Gulf of Kutch, northern Arabian Sea based on bomb-carbon in corals and tree-rings, *Proc. Indian Acad. Sci. (Earth Planet. Sci.)*, *103*, 329-340, 1994.
- Clark, G. R., Growth lines in invertebrate skeletons, *Ann. Rev. Earth Planet. Sci.*, *2*, 77-99, 1974.
- Craig, H., The natural distribution of radiocarbon and the exchange time of carbon dioxide between atmosphere and sea, *Tellus*, *9*, 1-17, 1957.
- Dettman, D. L., and K. C. Lohmann, Seasonal change in Paleogene surface water $\delta^{18}\text{O}$: fresh-water bivalves of western North America, in *Climate Changes in Continental Isotopic Records*, *Geophys. Monogr. Ser.*, vol. 78, edited by P. Swart, K. C. Lohmann, J. McKenzie, and S. Savin, pp. 153-163, AGU, Washington, D. C., 1993.
- Druffel, E. M., Radiocarbon in annual coral rings from the eastern tropical Pacific, *Geophys. Res. Lett.*, *8*, 59-62, 1981.
- Druffel, E. M., Bomb radiocarbon in the Pacific: Annual and seasonal timescale variations, *J. Mar. Res.*, *45*, 667-698, 1987.
- Druffel, E. M., Decade time scale variability of ventilation in the North Atlantic: High-precision measurements of bomb radiocarbon in banded corals, *J. Geophys. Res.*, *94*, 3271-3285, 1989.

- Druffel, E. M., and T. W. Linick, Radiocarbon in annual coral rings, *Geophys. Res. Lett.*, 5, 913-916, 1978.
- Druffel, E. M., and H. E. Suess, On the radiocarbon record in banded corals: exchange parameters and net transport of $^{14}\text{CO}_2$ between atmosphere and surface ocean, *J. Geophys. Res.*, 88, 1271-1280, 1983.
- Druffel, E. R. M., Detection of El Nino and decade scale variations of sea surface temperature from banded coral records: Implications for the carbon dioxide cycle, in: *The Carbon Cycle and Atmospheric CO_2 : Natural Variations Archaean to Present*, *Geophys. Monogr. Ser.*, vol. 32, edited by E. T. Sundquist and W. S. Broecker, pp.111-122, AGU, Washington, D.C., 1985.
- Epstein, S., R. Buchsbaum, H. A. Lowenstam and H. C. Urey, Revised carbonate-water isotopic temperature scale, *Bull. Geol. Soc. Am.*, 64, 1315-1326, 1953.
- Erlenkeuser, H., C^{14} and C^{13} isotope concentration in modern marine mussels from sedimentary habitats, *Naturwis.*, 63, 338, 1976.
- Erlenkeuser, H., and G. Wefer, Seasonal growth of bivalves from Bermuda recorded in their O-18 profiles, *Proceedings of 4th International Coral Reef Symposium*, vol. 2, 643-648, Manila, 1981.
- Fairbanks, R. G., and R. K. Matthews, The marine oxygen isotope record in Pleistocene coral, Barbadoes, West Indies, *Quat. Res.*, 10, 181-196, 1978.
- Grossman, E. L., and T. Ku, Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects, *Chem. Geol.*, 59, 59-74, 1986.
- Harrington, R. J., Aspects of growth deceleration in bivalves: Clues to understanding the seasonal O-18 and C-13 record -- A comment on Krantz et al. (1987), *Palaeogeogr.*, *Palaeoclimatol.*, *Palaeoecol.*, 70, 399-403, 1989.
- Heller, J., Longevity in molluscs, *Malacol.*, 31, 259-295, 1990.
- Jones, D. S., Annual cycle of shell growth and reproduction in the bivalves-*Spisula Solidissima*-&-*Arctica islandica*-, PhD, Princeton University, 1980.
- Jones, D. S., Repeating layers in the molluscan shell are not always periodic, *J. Paleontol.*, 55, 1076-1082, 1981.
- Jones, D. S., Sclerochronology: reading the record of the molluscan shell, *Am. Scientist*, 71, 384-391, 1983.
- Jones, D. S., D. F. Williams and M. A. Arthur, Growth history and ecology of the Atlantic Surf Clam, *Spisula Solidissima* (Dillwyn), as revealed by stable isotopes and annual shell increments, *J. Mar. Biol. Ecol.*, 73, 225-242, 1983.
- Jones, D. S., D. F. Williams, M. A. Arthur and D. E. Krantz, Interpreting the Palaeoenvironmental, paleoclimatic and life history records in mollusc shells, *Geobios. Mem.*, 8, 333-339, 1984.
- Jones, D. S., D. F. Williams and C. S. Romanek, Life history of symbiont-bearing giant clams from stable isotope profiles, *Science*, 231, 46-48, 1986.

- Killingsley, J. S., and W. H. Berger, Stable isotopes in a mollusc shell: detection of upwelling events, *Science*, 205, 186-188, 1979.
- Krantz, D. E., Mollusk-isotope records of Plio-Pleistocene marine paleoclimate, U.S. Middle Atlantic Coastal Plain, *Palaaios*, 5, 317-335, 1990.
- Krantz, D. E., D. S. Jones and D. F. Williams, Reply to "Aspects of growth deceleration in bivalves: Clues to understanding the seasonal O-18 and C-13 record", *Palaeogeogr.*, *Palaeoclimatol.*, *Palaeoecol.*, 70, 403-407, 1989.
- Krantz, D. E., D. F. Williams and D. S. Jones, Ecological and paleoenvironmental information using stable isotope profiles from living and fossil molluscs, *Palaeogeogr.*, *Palaeoclimatol.*, *Palaeoecol.*, 58, 249-266, 1987.
- Landman, N. H., E. R. M. Druffel, J. K. Cochran, D. J. Donahue and A. J. T. Jull, Bomb-produced radiocarbon in the shell of the chambered nautilus: rate of growth and age at maturity, 89, 28-34, 1988.
- Linick, T. W., P. E. Damon, D. J. Donahue and A. J. T. Jull, Accelerator mass spectrometry: the new revolution in radiocarbon dating, *Quat. Internat.*, 1, 1-6, 1989.
- Loosanoff, V. L., Reproductive cycle in *Cyprina islandica*, *Biol. Bull. (Woods Hole)*, 104, 146-155, 1953.
- Lutz, R. A., and D. C. Rhoads, Anaerobiosis and a theory of growth line formation, *Science*, 198, 1222-1227, 1977.
- Merrill, A. S., and J. W. Ropes, The general distribution of the Surf Clam and Ocean Quahog, *Proc. Nat. Shellfish. Ass.*, 59, 40-45, 1969.
- Murawski, S. A., J. W. Ropes and F. M. Serchuk, Growth of the Ocean Quahog, *Arctica islandica*, in the Middle Atlantic Bight, *Fish. Bull.*, U.S., 80, 21-34, 1982.
- Nichol, D., Recent species of the veneroid pelecypod *Arctica*, *J. Wash. Acad. Sci.*, 41, 102-106, 1951.
- Nozaki, Y., D. M. Rye, K. K. Turekian and R. E. Dodge, C-13 and C-14 variations in a Bermuda coral, *Geophys. Res. Lett.*, 5, No.10, 825-828, 1978.
- Ostlund, H. G., G. Possnert and J. H. Swift, Ventilation rate of the deep Arctic Ocean from Carbon 14 data, *J. Geophys. Res.*, 92, 3769-3777, 1987.
- Patzoid, J., Growth rhythms recorded in stable isotopes and density bands in the reef coral *Porites lobata* (Cebu, Phillipines), *Coral Reefs*, 3, 78-90, 1984.
- Patzoid, J., H. Ristedt and G. Wefer, Rate of growth and longevity of a large colony of *Pentapora foliacea* (Bryozoa) recorded in their oxygen isotope profiles, *Mar. Biol.*, 96, 535-538, 1987.
- Quay, P. D., B. Tilbrook and C. S. Wong, Oceanic uptake of fossil fuel CO₂: carbon-13 evidence, *Science*, 256, 74-79, 1992.

- Revelle, R., and H. E. Suess, Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO₂ during the past decades, *Tellus*, 9, 18-27, 1957.
- Rhoads, D. C., and R. A. Lutz, Skeletal Growth of Aquatic Organisms, New York and London, Plenum Press, 1980.
- Romanek, C. S., D. S. Jones, D. F. Williams, D. E. Krantz and R. Radke, Stable isotopic investigation of physiological and environmental changes recorded in shell carbonate from the giant clam *Tridacna maxima*, *Mar. Biol.*, 94, 385-393, 1987.
- Ropes, J. W., D. S. Jones, S. A. Murawski, F. M. Serchuk and A. Jerald, Documentation of annual growth lines in Ocean Quahogs, *Arctica islandica* Linne, *Fish. Bull., U.S.*, 82, 1-19, 1984.
- Sarmiento, J. L., and E. T. Sundquist, Revised budget for the oceanic uptake of anthropogenic carbon dioxide, *Nature*, 356, 589-593, 1992.
- Shackleton, N., The oxygen isotope stratigraphic record of the late Pleistocene, *Phil. Trans. R. Soc. Lond.*, 280, 169-182, 1977.
- Stocker, T. F., and W. S. Broecker, NADW formation as a branch of the hydrological cycle, *EOS*, 73, No. 18, 202-203, 1992.
- Suess, H. E., Natural radiocarbon and the rate of exchange of CO₂ between the atmosphere and the sea, University of Chicago Press, Chicago, Ill., 1953.
- Tanaka, N., M. C. Monaghan and K. K. Turekian, C¹⁴ balance for the Gulf of Maine, Long Island Sound and the northern Middle Atlantic Bight: evidence for the extent of the Antarctic Intermediate Water contribution, *J. Mar. Res.*, 48, 75-87, 1990.
- Thompson, I., and D. S. Jones, The ocean quahog, *Arctica islandica*, 'tree' of the north Atlantic shelf, *Geol. Soc. of Am. Abstr. Programs*, 9, 1199, 1977.
- Thompson, I., D. S. Jones and Dreibelbis, Annual internal growth banding and life history of the Ocean quahog *Arctica islandica* (Mollusca: Bivalvia), *Mar. Bio.*, 57, 25-34, 1980.
- Toggweiler, J. R., K. Dixon and W. S. Broecker, The Peru upwelling and the ventilation of the South Pacific thermocline, *J. Geophys. Res.*, 96, 20,467-20,497, 1991.
- Turekian, K. K., D. Cochran, Y. Nozaki, I. Thompson and D. S. Jones, Determination of shell deposition rates of *Arctica islandica* from the New York Bight using natural ²²⁸Ra and ²²⁸Th and bomb-produced ¹⁴C, *Limnol. Oceanogr.*, 27, 737-741, 1982.
- Warren, B. A., *Evolution of Physical Oceanography*, MIT Press, Cambridge, Mass., 1981.
- Wefer, G., and W. H. Berger, Isotope paleontology: growth and composition of extant calcareous species, *Mar. Geol.*, 100, 207-248, 1991.
- Williams, D. F., M. A. Arthur, D. S. Jones and N. Healy-Williams, Seasonality and mean annual sea surface temperatures from isotopic and sclerochronological records, *Nature*, 296, 432-434, 1982.

Chapter 3

Stable Carbon and Oxygen Isotope Records from *Arctica islandica* Shells in a Monitored Laboratory Growth Experiment.

Introduction

Stable oxygen (^{18}O) and carbon (^{13}C) isotope analysis of the carbonate skeletons of marine organisms is an important means of reconstructing past ocean climate and carbon cycle conditions [Wefer, & Berger, 1991]. However, because these records are of biological origin, they can be biased by internal as well as external influences, and the former can vary among different phyla and species. Consequently, there is a prerequisite interest in verifying these skeletal isotope records with known ambient conditions. This can be done in the laboratory or in the field under natural conditions, both approaches having their advantages and disadvantages. In the previous chapter [Weidman, Jones, and Lohmann, 1994], a study was presented that compared a 12-year oxygen isotope record from a single shell with a predicted one derived from nearby lightship temperature and salinity records, an aragonite- $\delta^{18}\text{O}$ relationship from Grossman and Ku [Grossman, & Ku, 1986], and a $\delta^{18}\text{O}\text{-H}_2\text{O}$ / salinity relationship for the Gulf of Maine from Fairbanks [Fairbanks, 1982]. That study concluded that the *A. islandica*'s shell isotope record was in close agreement with the predicted one assuming an eight-month shell growth period. However, some limitations of that field study were that: 1) the lightship and shell growth sites while close (~15 km) (See Figure 1 in Chapter 2) were not identical allowing for possible differences in actual T and S properties, 2) the $\delta^{18}\text{O}$ -water was not measured directly but derived from salinity using a regional linear relationship, and finally, 3) no comparison was possible for the shell's $\delta^{13}\text{C}$ record as no contemporary time series of the

stable carbon isotope composition of the ambient dissolved inorganic carbon (DIC) exists for this shelf region. These deficiencies provide the motivation for the laboratory and monitored field study presented in this chapter, where *Arctica islandica* specimens were grown under closely monitored conditions in the laboratory and in several different substrates to compare the shells' carbon and oxygen isotope record with directly measured ambient properties. At the same time an offshore site was monitored, though less intensively, to record the natural variation of shelf water column properties.

Methods

Laboratory Growth Experiment:

Arctica islandica specimens representing a range of sizes/ages were captured live on June 4, 1993 from a location 5 km south of Moosepeak Light, Maine (44° 25' N, 67° 32' W) at 70 m depth (Figure 1). Water samples for salinity and isotopic analysis were taken and bottom temperatures were measured at this site at the time of capture. Specimens were transported to a laboratory at the National Marine Fisheries Service Aquarium at Woods Hole, where individual shell heights were measured with a micrometer, each shell was indelibly marked with an ID number, and placed into one of three bins submerged in a tank 1.2 m x 2.4 m in area with a water depth of 35 cm. Each of the three bins contained a different substrate, medium/coarse sand of ~0.5 mm diameter (15 cm deep), marine very fine sand/silt of ~ 0.06 mm diameter (15 cm deep), and no sediment at all. Unfiltered seawater from Woods Hole Harbor supplied the system which included a refrigerated recirculating loop to keep the tank's water temperature below 20°C during the warmest summer months. Tank inflow/outflow rate was maintained at about 20 l/min so that the residence time of the water in the tank was approximately 1 hour. This flow rate was chosen to exceed the maximum filtration rate determined for individual adult *A. islandica*

(0.12 l/ min) times the ~100 specimens that were placed in the tank [Winter, 1969][Kraus, Beal, Chapman, and McMartin, 1992].

Since Kraus, et al. [1992] have reported that handling strongly impacts shell growth rates for this species, the specimens, once placed in the bins, were not disturbed for the duration of the year-long calibration experiment. Tank temperature was recorded, almost daily, to the nearest tenth of a degree with a calibrated thermometer. Water samples were collected every two weeks for salinity and stable carbon and oxygen isotope analysis. During the warm summer months (July-September) when external seawater input was reduced and refrigerated recirculation increased, a commercially available invertebrate food supplement was added to the tank water once every two days.

In June 1994, just prior to harvesting the laboratory specimens, water column and pore water samples were collected from the two bins with the sand and mud substrate for carbon isotope analysis. In the case of the sand substrate this was accomplished using a syringe and extracting pore waters from three sediment depths of 1 cm, 3 cm, and 6 cm, respectively. In the case of the mud substrate this was done by coring the sediment with a 5 cm diameter plastic tube, removing three successively deep layers of mud at 1.4 cm intervals (total depth 0-4.2 cm), and centrifuging them to separate the pore water.

All laboratory specimens were harvested on the same day on June 10, 1994, their ID numbers verified, and frozen. Later, selected specimens were thawed out, their shell heights measured, and a subset selected for isotopic analysis.

Offshore Site Monitoring:

A location 5 km south of Gay Head (41° 19'N, 70° 50'W) and 34 m depth (Figure 1), referred to hereafter as the Devil's Bridge site, was revisited about once every month

and a half for one year to record bottom temperature and take water samples for salinity and carbon and oxygen isotope analysis. This site was selected because it is the ocean quahog habitat nearest to Woods Hole. Although, *A. islandica* specimens were gathered from this site as well, the population at this site was too old/large and its growth rate too slow to be useful for this high-resolution laboratory growth study.

Sample Preparation and Stable Isotope Analysis:

Carbonate for isotopic analysis was obtained from selected shells using the micro-sampling techniques described in the previous chapter [Weidman, *et al.*, 1994]. Oxygen and carbon isotopic analyses of the shell carbonate were performed on the VG Prism mass spectrometer with an automated on-line Isocarb extraction system at the Woods Hole Oceanographic Institution. For DIC carbon analysis, seawater was collected in 500 ml borosilicate glass bottles, poisoned with 0.2 ml of saturated HgCl_2 solution to cease any further biological activity in the sample, and sealed with greased glass stoppers for later analysis. For oxygen analysis, seawater was collected in 100 ml glass "salinity" bottles and sealed with airtight plastic screw caps for later analysis. CO_2 extraction of the bi-monthly collected water samples for carbon isotope analysis was carried out using standard stripping methods described in McNichol, *et al.* [McNichol, Jones, Hutton, & Gagnon, 1994]. Water samples collected for oxygen isotope analysis were prepared using a slight modification of methods first described in an unpublished report by T. Anderson [1983] at the University of Illinois and also later described by Allard [Allard, 1989] (see Appendix for details). These methods involve a freeze/thaw routine along with equilibrating an injected aliquot of CO_2 of known volume and isotopic composition with a seawater sample. Isotopic analyses of bi-monthly and offshore site waters were performed (in duplicate) on the VG Optima mass spectrometer at the Woods Hole Oceanographic Institution. CO_2 extraction of the pore waters and (simultaneously taken) ambient water column were carried out using methods developed by D. McCorkle for pore water $\delta^{13}\text{C}$ analysis [McCorkle,

Emerson, & Quay, 1985][McCorkle, Keigwin, Corliss, & Emerson, 1990][McCorkle, & Keigwin, 1994], and isotopic analyses of these samples were performed on the VG 602 mass spectrometer at the Woods Hole Oceanographic Institution. All analyses on all three mass-spectrometers were calibrated against the simultaneously run NBS/NIST standards (NBS-19 and VSMOW) and in-house standards calibrated to these NBS/NIST standards. The analytical precision of carbon and oxygen analyses, for carbonate and DIC samples is approximately $\pm 0.1\text{‰}$ (1σ) and for oxygen analyses of water samples is $\pm 0.06\text{‰}$ (rms of duplicate differences). For reporting stable oxygen and carbon isotope compositions, the standard notation is used where

$$\delta^{18}\text{O}_{\text{sample}} (\text{PDB}) = [({}^{18}\text{O}/{}^{16}\text{O}_{\text{sample}}/{}^{18}\text{O}/{}^{16}\text{O}_{\text{standard}}(\text{NBS-19}=-2.20\text{‰})) - 1] \times 1000$$

and

$$\delta^{13}\text{C}_{\text{sample}} (\text{PDB}) = [({}^{13}\text{C}/{}^{12}\text{C}_{\text{sample}}/{}^{13}\text{C}/{}^{12}\text{C}_{\text{standard}}(\text{NBS-19}=+1.95\text{‰})) - 1] \times 1000$$

and

$$\delta^{18}\text{O}_{\text{sample}} (\text{SMOW}) = [({}^{18}\text{O}/{}^{16}\text{O}_{\text{sample}}/{}^{18}\text{O}/{}^{16}\text{O}_{\text{standard}}(\text{VSMOW}=0.00\text{‰})) - 1] \times 1000$$

In this paper, Grossman and Ku's [1986] estimate of $\sim 0.2\text{‰}$ for the difference between SMOW and PDB scales is assumed.

Results

I. Water Column Properties

Temperature:

The tank water temperature record shows a full annual cycle ranging from 0° to 18°C with a mean of 10.8°C and standard deviation of $\pm 6.3^{\circ}$ (Figure 2a and Table 1). For convenience, the bi-monthly temperatures, which were measured at the same time as the water sample collections, are shown in Table 1. Bottom temperature at the Maine capture

site at the time of capture was 5.2°C (6/4/93). Captured specimens were introduced to the tank on June 5, 1993 (experiment day # 33), at which time laboratory water temperatures were already greater than 15°C. Water temperature reached a maximum in mid-September, dropped below 15°C in late October, reached a minimum in late January, and rose to just above 16°C at the experiment's termination on June 10, 1994.

Salinity:

Salinity in the laboratory tank ranged about 1‰ from 30.77‰ to 31.78‰ with a mean of 31.30‰ and a standard deviation of ± 0.28 ‰ (Figure 2b). Bottom salinity at the capture site at the time of capture was 31.52‰. Highest laboratory salinities were recorded in the fall and early winter and lowest in the early spring. Temperature and salinity display only a weak positive correlation ($r = +0.211$).

$\delta^{18}\text{O}$ -water:

The $\delta^{18}\text{O}$ (SMOW) of the tank water ranged about 0.4‰ from -1.59 to -1.22‰ with a mean of -1.44‰ and standard deviation of ± 0.08 ‰. Bottom $\delta^{18}\text{O}$ -water value at the capture site at the time of capture was -1.49‰ (Figure 2c). As expected, $\delta^{18}\text{O}$ -water values correlate positively with salinity ($r = +0.656$). The linear $\delta^{18}\text{O}$ -salinity relationship for the tank water is: $\delta^{18}\text{O}\text{-water} = 0.190 * S - 7.38$ (Standard Error of the Estimate (SEE) = 0.07‰). This is similar in slope and intercept to Fairbanks' [1982] $\delta^{18}\text{O}$ -salinity relationship for the Mid-Atlantic Bight (+0.258 and -9.14). The intercept value of ~ -7 ‰ indicates that the meteoric source of the fresh component in Woods Hole Harbor is local, based on Figure 1 in Fairbanks [1982].

$\delta^{13}\text{C}$ -DIC:

$\delta^{13}\text{C}$ (PDB) of the DIC ranged about 1‰ from -0.2‰ to +1.0‰ with a mean of +0.62‰ and standard deviation of ± 0.35 ‰ (Figure 2d). Bottom DIC $\delta^{13}\text{C}$ at the capture

site at the time of capture was 0.4‰ heavier than initial laboratory DIC. DIC $\delta^{13}\text{C}$ reached a minimum in mid-summer and a maximum in late spring, and so is negatively correlated with temperature ($r = -0.582$). This negative relationship may result from the dependence of the $\text{CO}_2\text{-HCO}_3$ equilibrium fractionation on temperature ($\sim -0.12\text{‰} / \text{C}^\circ$) [Deuser and Degens, 1967][Mook, 1986]. The correlation between temperature and DIC $\delta^{13}\text{C}$ data is weakened by the large $\delta^{13}\text{C}$ scatter at temperatures above 15°C . Deviation from equilibrium is likely for two reasons (Broecker and Peng, 1982): 1) carbon isotopic exchange is a rather slow process in seawater (~ 1 year for full equilibration of a 10 m layer) and seasonal temperature changes are generally too rapid to allow for complete equilibrium, and 2) marine biological cycles are accompanied by carbon isotopic fractionations which can move the isotopic composition of the local DIC reservoir away from equilibrium with the atmosphere. For example, phytoplankton blooms in the spring and early summer can result in a ^{13}C enrichment of the DIC as ^{12}C is preferentially taken up by the organisms.

Pore Water DIC:

The pore waters of the sand and mud substrates are very enriched (by $\sim 65\%$) in ΣCO_2 and very depleted in $\delta^{13}\text{C}$ (by $\sim 6\text{‰}$) relative to the ambient water column above the sediment surface (Table 2). A sharp gradient in ΣCO_2 and $\delta^{13}\text{C}$ properties occurs in the top 1 cm of the sediment column of both sand and mud substrates. These conditions are characteristic of natural marine sediments due to the breakdown of organic matter within the upper sediment layers (marine organic carbon $\delta^{13}\text{C} \sim -20\text{‰}$) [McCorkle, 1985]. Surprisingly, mud and sand pore waters show no significant differences in their DIC properties, despite a difference in permeability of $\sim 10^2$ based on grain size [Inman, 1963] and the consequent potential difference in pore water exchange with the overlying water column.

Devil's Bridge Properties (natural cycle):

Bottom temperature conditions at the Devil's Bridge site ranged from -0.8°C to 13.8°C with a mean 9.1°C and standard deviation of $\pm 5.3^{\circ}\text{C}$. Maximum temperatures were reached in late summer / early fall (Figure 3). Though no measurements were made during the winter months (Dec. - Feb.), temperatures could not have been much colder than -0.8°C recorded in early March, making that value a reasonable minimum for the year. In general, bottom temperatures at the Devil's Bridge site were cooler and had less range than in the laboratory tank. Salinity ranged 0.4‰ from 31.97‰ to 32.36‰ with a mean of 32.05‰ and standard deviation of $\pm 0.20\text{‰}$. Like the laboratory conditions, lowest salinity occurred in the summer and spring with the highest values in the fall.

The $\delta^{18}\text{O}$ (SMOW) of the bottom water ranged 0.25‰ from -1.12‰ to -1.37‰ with a mean of -1.21‰ and standard deviation of $\pm 0.10\text{‰}$. The $\delta^{18}\text{O}$ -water is correlated positively with salinity ($r = +0.760$). The linear $\delta^{18}\text{O}$ -salinity relationship for Devil's Bridge water is: $\delta^{18}\text{O}\text{-water} = 0.395 * S - 13.86$ ($\text{SEE} = 0.07\text{‰}$). This is similar in slope and intercept to Fairbanks' [1982] $\delta^{18}\text{O}$ -salinity relationship for the Gulf of Maine ($+0.421$ and -14.66), and suggests a higher-latitude source for its fresh component.

DIC $\delta^{13}\text{C}$ (PDB) ranged $\sim 0.7\text{‰}$ from $+0.71\text{‰}$ to $+1.39\text{‰}$ with a mean of $+1.02\text{‰}$ and standard deviation of $\pm 0.22\text{‰}$. Maximum $\delta^{13}\text{C}$ values were recorded in the first spring and early summer at the start of the monitoring program and reached a minimum in the early fall. $\delta^{13}\text{C}$ rebounded again by late fall but remained at a low level during the following spring. DIC $\delta^{13}\text{C}$ at the Devil's Bridge site is not significantly correlated to temperature ($r = +0.235$) and its trend is of the opposite sign from the laboratory DIC $\delta^{13}\text{C}$ record. However, both laboratory and offshore $\delta^{13}\text{C}$ profiles have minima when ambient temperatures are near their maxima, and both display maxima in the spring.

II. Shell Records

Stable carbon and oxygen isotope records (Table 4) were obtained from the shells of three young *A. islandica* specimens, each approximately the same size/age (24-27 mm, / 7-9 yrs-old), and each grown in a different substrate medium (sand, mud, no sediment) in the laboratory for one year. Laboratory shell growth was very limited in older specimens (see Figure 4 in chapter 2), making it difficult to obtain a seasonal sampling resolution in their shells. For this reason, in this study we have limited our analysis to shells within the specified size/age range, which as a group showed the most shell growth and afforded the greatest sampling resolution. Figure 4a-c displays each shell's carbon and oxygen isotope record separately.

Ring Structure and Shell Growth:

In Figure 4a-c, the vertical cross-hatched lines superimposed on the profiles represent the internal "rings", which are visible in radial cross-sections of the shells as thin translucent layers between wider more opaque growth increments. In chapters one and two these rings were shown to be annual in nature and deposited in the autumn when bottom temperatures are highest and $\delta^{18}\text{O}$ of the shell is most depleted. In the shell records shown here, the first $\sim 2/3$ of the isotope record is obtained from shell material that was deposited in the natural environment prior to capture, and the latter $\sim 1/3$ from material deposited in the laboratory setting over 1 year. The first two "rings" (from the left) are, thus, annual "rings", deposited in the fall of 1991 and 1992, respectively. The third "rings" are "disturbance rings", coincident with the start of the laboratory growth period (June 1993), and are, likely, related to the trauma experienced by the specimens at the time of their capture and/or subsequent sharp change in their ambient environment. The physical distance between the third and fourth rings is very small relative to the other bands, and in specimen qx102 it is just discernible. The fourth (or last) "rings" appear to have been

deposited during the autumn of 1993 as they lie near the $\delta^{18}\text{O}$ -shell minima. However, the fourth ring of specimen qx94 (sand) is positioned just after the maximum $\delta^{18}\text{O}$, though this is likely due to a placement error owing to the difficulty of visually gauging the actual position of the ring during micro-sampling. New radial shell growth for the three specimens averaged about 1.8 mm. Shell growth was greatest for the specimen grown in sand (2.15 mm), and least for the specimen grown in "no sediment" (1.39 mm). The specimen grown in mud had about average growth (1.83 mm).

$\delta^{18}\text{O}$ -shell record:

Between 37 and 57 aragonite powder samples were obtained from outer portions of each of the three shells, of which between 15 and 21 samples were obtained from the laboratory grown portion of each shell. So the isotope records can be compared more closely, the sample positions are normalized by setting the pre-laboratory (pre-capture) growth run from -1 to 0 and the laboratory growth run from 0 to +1 (Figure 5a-b). The oxygen isotope profiles of all three shells show strong similarities (Figure 5a). The pre-laboratory part of the $\delta^{18}\text{O}$ -shell records display similar shapes and approximately equal values for minima and maxima. In these portions of the profiles, $\delta^{18}\text{O}$ values range from about +1‰ to +2‰ with a mean of +1.4‰ (± 0.3 ‰). As described above, the minima occur at or near the annual "rings", and the maxima occur within a few samples after the annual "rings", so that the intra-band $\delta^{18}\text{O}$ profiles are largely characterized by negative trends. This is consistent with the seasonal $\delta^{18}\text{O}$ profiles from the Nantucket Lightship specimen of the previous chapter (see Figure 4 in chapter 2).

The laboratory-grown portions of the record begin at the third "ring" where $\delta^{18}\text{O}$ values are initially $\sim +1.2$ ‰ and are followed by a steep depletion. In this portion of the records, $\delta^{18}\text{O}$ values range from about 0‰ to +2‰ with a mean of +0.7‰ (± 0.7). The lab-grown segments exhibit a seasonal $\delta^{18}\text{O}$ range about twice that of the pre-lab segments.

Minima of about 0‰ occur within one or two samples after the disturbance ring. This sharp drop is consistent with the large temperature difference ($\Delta \sim 10^{\circ}\text{C}$) between the Maine capture site (5°C) and the laboratory tank (15°C). In all three shells, these first minima are confined within a very narrow sample space (1-2 samples), despite a prolonged period (5 months) of warm ($+15^{\circ}\text{C}$) tank temperatures, indicating very little shell growth during this time. These minima are followed by a sharp enrichment to maxima of $\sim +2\text{‰}$. A rather linear negative trend characterizes the remaining $\delta^{18}\text{O}$ data which reach terminal values of $\sim 0\text{‰}$. This negative trending $\delta^{18}\text{O}$ data comprises 60-85% of laboratory shell growth, and indicates that most of the new shell growth occurred during a few months in the winter/spring of 1994.

$\delta^{13}\text{C}$ -shell record:

Like the $\delta^{18}\text{O}$ profiles, the pre-laboratory portion of the $\delta^{13}\text{C}$ -shell profiles display very similar shapes and values for minima and maxima (Figure 5b). In this portion of the shell, $\delta^{13}\text{C}$ values range about 0.5‰ from about $+1.7\text{‰}$ to $+2.2\text{‰}$ with a mean of $+2.0\text{‰}$ ($\pm 0.2\text{‰}$). The annual $\delta^{13}\text{C}$ minima occur within a few samples after the annual $\delta^{18}\text{O}$ minima and before the next $\delta^{18}\text{O}$ maxima. This observation demonstrates that seasonal shell growth continues after annual "ring" deposition and that the samples from the region between the $\delta^{18}\text{O}$ minima and maxima are not simply a mixture of very enriched and depleted material divided by a hiatus in shell growth. If these samples were such a mixture of adjacent values, then no $\delta^{13}\text{C}$ minima could occur in this region. The annual $\delta^{13}\text{C}$ maxima occur near the latter part of each intra-band profile, but generally begin declining just before the annual "ring" is deposited, so that the intra-band $\delta^{13}\text{C}$ profile is dominated by a positive trend.

The $\delta^{13}\text{C}$ profiles of the lab-grown segments exhibit a much greater range and more variability between shells, than the pre-lab segments. In this portion of the shell, $\delta^{13}\text{C}$

values range $\sim 2.0\text{‰}$ from about $+0.2\text{‰}$ to $+2.2\text{‰}$ with a mean of $+1.5\text{‰}$ (± 0.5). Like the $\delta^{18}\text{O}$ data, the $\delta^{13}\text{C}$ profiles show a large depletion at the disturbance "rings" to minima between $+0.7\text{‰}$ and -0.3‰ . This ($\sim 2\text{‰}$) drop is consistent in sign, but larger than the difference in DIC $\delta^{13}\text{C}$ between the laboratory minimum and capture site ($\Delta \sim 1\text{‰}$). Like the pre-lab $\delta^{13}\text{C}$ profiles, the laboratory $\delta^{13}\text{C}$ minima occur near to, but after the $\delta^{18}\text{O}$ minima. The $\delta^{13}\text{C}$ minima are followed by an enrichment to maxima of $\sim +2.0\text{‰}$ that occur just prior to the end of each profile. The rate of this enrichment varies among the specimens; it is steep in the case of specimen qx94 (sand) and much less so in the case of qx156 (no sediment). The $\delta^{13}\text{C}$ profiles again display a depletion just before the profiles' terminations. The lab-grown segments show overall positive $\delta^{13}\text{C}$ trends, which is consistent in sign, but greater than the similar trend in the DIC $\delta^{13}\text{C}$.

Discussion

I. $\delta^{18}\text{O}$ -aragonite

The laboratory shell $\delta^{18}\text{O}$ profiles are superimposed onto a predicted $\delta^{18}\text{O}$ profile (Figure 6) for aragonite deposited in equilibrium with the laboratory tank water. This predicted profile was obtained by applying the tank's temperature and $\delta^{18}\text{O}$ -water data (Table 1) to the modified form of the Grossman and Ku (1986) paleotemperature equation (1a and 1b) (for mollusc data only) used in the previous chapter:

$$\delta^{18}\text{O}_{\text{aragonite(PDB)}} = 4.45 - 0.213 \cdot T (\text{C}^\circ) + \delta^{18}\text{O}_{\text{water(SMOW)}} \quad 1a)$$

$$T (\text{C}^\circ) = 20.86 - 4.69 \cdot (\delta^{18}\text{O}_{\text{aragonite(PDB)}} - \delta^{18}\text{O}_{\text{water(SMOW)}}) \quad 1b)$$

The predicted $\delta^{18}\text{O}$ -aragonite curve has a range of $\sim 4\text{‰}$ from -1.0‰ to $+3.0\text{‰}$ with a mean of 0.7‰ ($\pm 1.3\text{‰}$). The predicted range is about twice as large as that of the lab-grown shell $\delta^{18}\text{O}$ signals. In particular, the $\delta^{18}\text{O}$ -shell profiles do not become as enriched or as depleted as the predicted profile and neither are they in phase with each other, assuming continuous linear shell growth. The latter result alone implies that the shell growth rates must be variable. If the critical points (minima, maxima, and endpoints) of the predicted and shell records are assumed to coincide, then it becomes clear that most of the shell growth occurred during the last months of the experiment, and that little growth occurred during the first two-thirds of the experiment. This result may be due to a combination of physical shock associated with capture, resettling into the new laboratory environment, or the very warm tank temperatures during the first summer. As already mentioned, Kraus, et al. [1993], noted that handling the live *A. islandica* specimens during routine growth studies greatly reduces their growth rates, and so it is reasonable to suppose a period of convalescence after capture. If the validity of equation (1) for *A. islandica* is not assumed, and continuous (though non-linear) shell growth is assumed so that the entire temperature record is represented in the $\delta^{18}\text{O}$ -shell profiles, then a relationship can be derived from a fit of the tank temperature and shell's $\delta^{18}\text{O}$ maxima and minima:

$$T = 26.63 - 7.66^{*}(\delta^{18}\text{O}_{\text{aragonite(PDB)}} - \delta^{18}\text{O}_{\text{water(SMOW)}}) \quad 2)$$

This equation has roughly twice the negative slope as equation (1b). While this result cannot be justified as an equilibrium relationship based on thermodynamic arguments [O'Neil, Clayton, & Mayeda, 1969], it is perhaps possible that it is the result of some unknown non-linear "vital" effect. This seems unlikely owing to the preponderance of evidence which indicates that bivalves generally deposit their carbonate in oxygen isotopic equilibrium with the ambient environment [Wefer, et al., 1991]. However, a point in favor of the higher-slope relationship is that it accurately hindcasts the Maine capture site

temperature at the time of capture, whereas Grossman and Ku's relationship overestimates this value by $\sim 3^{\circ}\text{C}$. While this could be taken as evidence of a disequilibrium effect, it may be that the instantaneous properties measured at this site at the time of capture were not characteristic over the period of shell deposition for these samples, which may represent one to several weeks of shell growth. For instance, the discrepancy might be explained by the presence of fresher or lighter $\delta^{18}\text{O}$ -water conditions existing just prior to capture. Salinity data for this area (MARMAP station #141) do show a large ($\sim 1\text{‰}$) freshening of this magnitude which occurs in the spring, and dissipates rapidly by early summer. In addition, the source of this springtime freshening seems to be a seasonal pulse of coastal Scotian Shelf water into the Gulf of Maine, which itself has its origins in the Gulf of Saint Lawrence [Smith, 1983]. This higher latitude source should also carry a lighter $\delta^{18}\text{O}$ -water component for a given salinity [Fairbanks, 1982], and taken along with a 1‰ drop in salinity (equivalent to $\sim 0.5\text{‰}$ change in $\delta^{18}\text{O}$ -water) could account for the discrepancy between the predicted and observed values.

The more likely case, is that the *A. islandica* specimens deposited their aragonite in equilibrium with ambient conditions, and disparity between predicted $\delta^{18}\text{O}$ -aragonite and $\delta^{18}\text{O}$ -shell is the result of shell growth hiatuses. Shell growth cessation is common in bivalves during extremes in temperature, and has been reported in many species [Wilbur and Owen, 1964][Williams, Arthur, Jones, and Healy-Williams, 1982][Harrington, 1989][Wefer, et al., 1991]. More relevant, independent evidence exists from several studies for shell growth reduction or cessation in *A. islandica* during the coldest months [Jones, 1980a][Jones, 1980b][Lutz, Goodsell, Castagna, & Stickney, 1983][Kraus, Beal, Chapman, & McMartin, 1992]. In the previous chapter, the general validity of equation (1) was assumed, and bottom water T and S records from a nearby location along with Fairbanks' [1982] salinity- $\delta^{18}\text{O}$ -water relationship were used to predict a $\delta^{18}\text{O}$ -aragonite record. By comparing the shell and predicted records, a cold shutdown temperature of $\sim 6^{\circ}\text{C}$

($\pm 1^\circ$) (at monthly resolution) was implied. No warm-end shutdown temperature was noted as the shell record $\delta^{18}\text{O}$ minima corresponded with the predicted $\delta^{18}\text{O}$ minima. However, Mann [1982] noted that the distribution of *A. islandica* on the eastern U. S. continental shelf shows a shoreward limit that corresponds to $\sim 16^\circ\text{C}$ October (annual maximum) bottom isotherm. This suggests that temperatures greater than 16°C may be outside the normal temperature range of this species, and if so it would seem reasonable that they would respond poorly to temperatures exceeding this value.

Furthermore, in this chapter, the ambient temperature and $\delta^{18}\text{O}$ -water are specifically known throughout the period of calibration, which removes the uncertainty associated with these parameters present in the previous chapter. Assuming equilibrium deposition and applying the mean value of the lab-grown shells' $\delta^{18}\text{O}$ minima and maxima ($-0.15\text{‰} \pm 0.2\text{‰}$ and $+2.01\text{‰} \pm 0.3\text{‰}$, respectively) and the mean $\delta^{18}\text{O}$ -water ($-1.44\text{‰} \pm 0.08\text{‰}$) to equation (1), a mean cold and warm shutdown temperature are estimated to be $4.7^\circ\text{C} \pm 1.8^\circ$ and $14.8^\circ\text{C} \pm 1.3^\circ$, respectively. The cold shutdown temperature estimate of $\sim 5^\circ\text{C}$ is slightly lower, though not significantly different than the $\sim 6^\circ\text{C}$ estimate of the previous chapter. For the region near the Nantucket Lightship for the period from 1956-1971, bottom temperatures annually dropped $1^\circ\text{--}2^\circ\text{C}$ below the 5°C lower limit for about 3 to 4 months (January to April) on average, while the upper 15°C limit was exceeded only occasionally (twice by just 0.5°C in 14 years) and the mean annual maximum (October) bottom temperature for this period is fully 2°C below this cutoff.

Assuming that the shell growth hiatuses occur at the $\delta^{18}\text{O}$ -shell minima and maxima, then the $\delta^{18}\text{O}$ -shell profiles can be separated at these points and compared to different sections of the predicted $\delta^{18}\text{O}$ -aragonite curve (Figure 7). The growth periods of each of the shells can be determined by mapping the endpoints of each group onto the predicted curve. The data then fall into three groups. The first minima are arbitrarily

assigned to the second or fall group. The hiatus between the second and third groups is made near the $\delta^{18}\text{O}$ maxima but is also defined by a sharp increase in $\delta^{13}\text{C}$.

The first group consists of the first 1-2 points from each shell's laboratory growth and represents a mix of pre-laboratory growth and early laboratory growth. As the change from one temperature regime to the other ($\Delta \sim 10^\circ\text{C}$) was essentially instantaneous, the data indicating in-between temperature states must be the result of mixing between material deposited under cold pre-laboratory conditions and warm laboratory conditions. This material was deposited sometime during the first four months or 35% of the experimental period and yet represents only about 10% of shell growth.

The second group, which includes the first summer minima, consists of 2-4 data points from each shell and represents autumn and early winter growth. This growth period lasted about two months from October to December or about 20% of the experiment's duration and represents approximately the same percentage of growth. In a recent *A. islandica* study, Witbaard, et al. [Witbaard, Jenness, Van Der Borg, & Ganssen, 1994] suggest that seasonal shell growth stops at the annual ring and that the apparent fall growth in their $\delta^{18}\text{O}$ profile may simply represent a sampling artifact caused by mixing the shell's $\delta^{18}\text{O}$ minima and maxima. Although some smearing of the signal is inevitable during sampling, and already invoked here to explain the first group's "in-between" data, the $\delta^{13}\text{C}$ minima that also occur within the second group argue against a simple mixing between $\delta^{18}\text{O}$ minima and maxima, because these minima cannot be accounted for by a mixing of end members, and therefore indicate at least some real shell growth during this late autumn period.

The last group, or spring data, contain most of the shell data and represent most of the laboratory shell growth (70%). This growth period began in mid-March and lasted until

the experiment's termination in early June or approximately 20% of the experiment's duration. Figure 8 shows that the spring shell data, converted to temperature (using equation 1 and the mean $\delta^{18}\text{O}$ -water), closely follow the observed temperature curve indicating that shell growth is approximately linear over this time. The data from specimen qx94, grown in sand and having the most shell growth of the three shells, show the most deviation from the predicted curve, indicating a more variable shell growth rate for this faster growing individual.

II. $\delta^{13}\text{C}$ -aragonite

Currently, there is no consensus on the fractionation of aragonite deposited in equilibrium with the $\delta^{13}\text{C}$ -DIC of ambient waters. Figure 9 shows the results of applying the tank properties $\delta^{13}\text{C}$ -DIC and temperature to three recently published methods, all of which give disparate results. Mook [Mook, 1986] proposed the equation:

$$\delta^{13}\text{C-} \text{aragonite} = -4232/T \text{ (K}^\circ\text{)} + 16.90 + \delta^{13}\text{C-DIC} \quad 3a)$$

This relationship describes $\delta^{13}\text{C}$ -aragonite as a positive function of temperature, with an approximate linear slope $\sim +0.05/^\circ\text{C}$ in the range $0^\circ\text{-}20^\circ\text{C}$. Grossman and Ku (1986) using mollusc data from their own study proposed the equation:

$$\delta^{13}\text{C-} \text{aragonite} = 2.66 - 0.131 * T \text{ (C}^\circ\text{)} + \delta^{13}\text{C-DIC} \quad 3b)$$

The relationship shows $\delta^{13}\text{C}$ -aragonite as a negative function of temperature. Most recently, Romanek, et al, [Romanek, Grossman, & Morse, 1992] have performed inorganic precipitation experiments that show that $\delta^{13}\text{C}$ -aragonite has a constant offset of $\sim +2.7\text{‰}$ relative to ambient DIC with no significant temperature dependence. Hence, unlike

$\delta^{18}\text{O}$, there is no prior confidence in the predicted equilibrium curves as a baseline to compare the shell records.

The $\delta^{13}\text{C}$ of the tank DIC averages $+0.6\text{‰}$ while the lab-grown shell material averages $+1.6\text{‰}$, for a mean difference of 1.0‰ . This difference is much less than that expected from equilibrium using Romanek, et al.'s [1993] constant offset of $+2.7\text{‰}$. For the mean tank temperature of 10.8°C , it is also much less than the $+2.3\text{‰}$ predicted from Mook's equation (3a), but it is only a bit less than the $\sim +1.2\text{‰}$ predicted from Grossman and Ku's equation (3b).

The shell $\delta^{13}\text{C}$ profiles do show a general similarity to the trend of the $\delta^{13}\text{C}$ -DIC signal (Figure 10), in that minima occur during the first summer and become more enriched in the winter and spring. Assuming continuous shell growth, this could be taken as evidence that the shell $\delta^{13}\text{C}$ reflects the $\delta^{13}\text{C}$ variations in the ambient DIC. However, the greater amplitude of the shell $\delta^{13}\text{C}$ signals relative to the DIC signal suggests that the relationship between the $\delta^{13}\text{C}$ of the water column DIC and shell carbonate cannot be attributed to a simple offset from $\delta^{13}\text{C}$ -DIC.

Using the shell growth periods determined from fitting the $\delta^{18}\text{O}$ data onto the predicted $\delta^{18}\text{O}$ record, the paired shell $\delta^{13}\text{C}$ data can be assigned to a specific time of deposition (Figure 11). From this perspective, the variation in shell $\delta^{13}\text{C}$ bears little relation to the $\delta^{13}\text{C}$ -DIC profile. For example, $\delta^{13}\text{C}$ -DIC remains rather constant during the fall and spring growth periods, while shell $\delta^{13}\text{C}$ varies by about 1‰ . Also, while spring DIC is $\sim 0.2\text{‰}$ enriched in $\delta^{13}\text{C}$ relative to the previous fall, the spring shell $\delta^{13}\text{C}$ is approximately 1‰ enriched relative to the fall. However, the variation of shell data is similar to the variation predicted from Mook's [1986] positive temperature-dependent relationship (3a), although, the shell data are about 1‰ depleted relative to this predicted

curve. Also, the offset between the fall and spring shell data ($\sim 1\text{‰}$) is not reflected in the predicted values from Mook's relationship which differ by only $\sim 0.2\text{‰}$. Also not reflected in this predicted curve, is the shells' sharp depletion ($0.3\text{--}1.7\text{‰}$) in the late spring. Finally, the absence of a clear dependence on temperature is shown by the poor correlation between the shell (laboratory) $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ ($r = -0.178$) (Figure 12). These mixed results suggest that while the seasonal variation in *A. islandica* shell $\delta^{13}\text{C}$ may be controlled at times by a positive dependence on temperature, there seems to be no dependence at other times.

Microhabitat / Pore Water $\delta^{13}\text{C}$:

Studies have shown that the depleted $\delta^{13}\text{C}$ -DIC environment of marine sediments [McCorkle, *et al.*, 1985] can cause the tests of the infaunal benthic foraminifera to become depleted relative to tests of epifaunal species [Grossman, 1987][McCorkle, Keigwin, Corliss, and Emerson, 1990]. Analogously, Krantz, *et al.* [Krantz, Williams, & Jones, 1987] offered the hypothesis that the $\delta^{13}\text{C}$ composition of the shells of infaunal filter-feeding bivalves may be depleted relative to that of epifaunal species due to the uptake of pore water DIC. However, the $\delta^{13}\text{C}$ composition of *A. islandica* shells does not appear to be dependent on the $\delta^{13}\text{C}$ of ambient pore water DIC based on the results shown here. The mean $\delta^{13}\text{C}$ of the shell grown in the water column or "no sediment" is actually more depleted (qx156 $\delta^{13}\text{C} = +1.49\text{‰}$) than the shell grown in the mud (qx102 $\delta^{13}\text{C} = +1.77\text{‰}$) even though the latter specimen's ambient pore water $\delta^{13}\text{C}$ was depleted by $\sim 6\text{‰}$ relative to the overlying water column. The sand substrate's ambient pore water $\delta^{13}\text{C}$ was as depleted as the mud substrate's pore water, and yet the mean $\delta^{13}\text{C}$ of the shell grown in sand (qx94 = $+1.46\text{‰}$) is almost identical to that of the shell grown in the water column. These results indicate that the "microhabitat" analogy does not apply to the infaunal bivalve *A. islandica*, probably because this bivalve's internal fluid exchange with the overlying water column is rapid and overwhelms any external pore water influence. Based on studies that describe the significant influence of benthic megafauna on geochemical cycling in the

sediments [Aller, 1978][Aller and Yingst, 1985], it is perhaps more likely that infaunal filter-feeding bivalves, such as *A. islandica*, rather than being passive inhabitants of their sediment environment, may act instead as primary transport pathways of overlying waters to the sediments, thus enriching the local pore water $\delta^{13}\text{C}$ about the individual bivalves.

Physiological (Vital) Effects:

The strong similarity displayed among the shell's $\delta^{13}\text{C}$ profiles, especially the pre-laboratory growth portions, makes it difficult to discern whether the specimens are responding to environmental changes (external control) or a common physiological cycle (internal control). However, the absence of any consistent correspondence with ambient temperature and DIC variations and the greater contrast shown among the specimens' laboratory $\delta^{13}\text{C}$ profiles hints that individual physiology may play an important role, at least during periods of greater physical stress (e.g., handling, extreme laboratory conditions, reproductive activity). Stable carbon isotope studies involving the carbonate shells of marine organisms, almost unanimously, have discussed the role of vital effect or physiological controls in interpreting their $\delta^{13}\text{C}$ compositions, and most have concluded that it is a significant factor resulting in disequilibrium with ambient DIC (e.g., [Wefer and Berger, 1991]). The specific term "vital effect" is usually interpreted to mean the inclusion of metabolically-produced (internal respiration) CO_2 , which is usually isotopically lighter ($\sim -20\text{‰}$) than ambient DIC ($\sim 0\text{‰}$), into the carbonate shell of a marine organism causing this shell to become depleted relative to equilibrium [Grossman, 1987]. Tanaka, et al. [1986] have calculated that approximately 50% of the carbon in the shells of several coastal mollusc species is of metabolic origin. If we allow that the *A. islandica* shells are about 1‰ depleted relative to "equilibrium", based on the average difference with the predicted offsets ($0.2\text{--}1.7\text{‰}$) derived from the three published relationships, then these shells require only a 5% uptake of metabolic- CO_2 to cause this overall depletion. A fluctuation of $\sim \pm 5\%$ about this metabolic uptake value would also explain the seasonal $\delta^{13}\text{C}$ signal of $\sim 2\text{‰}$ observed

in the laboratory shell growth, though the pre-laboratory shell growth has a seasonal $\delta^{13}\text{C}$ signal only about 1/4 of this amplitude.

In this view, the shells' seasonal $\delta^{13}\text{C}$ variation suggests variable metabolic activity or at least a variation in the supply of metabolic carbon to the shell. As the shell becomes most depleted in the autumn and most enriched in the mid-spring this model would suggest greatest metabolic activity or supply in the autumn and least in the spring. However, it would be reasonable to expect that increased shell growth, such as occurred during the spring in the laboratory, would parallel increased tissue growth and metabolic activity, but the spring is precisely when the shells are most enriched - not depleted. Perhaps, increased fluid exchange with the ambient water accompanies growth and overwhelms the metabolic effect on shell composition, but in any case this seems to add a complication and makes a direct metabolic effect less likely. McConnaughey [1989a, 1989b] suggested that kinetic effects, such as those that might be associated with rapid rates of carbonate deposition, would tend to cause offsets in both $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ and result in increased correlation between them. He also noted that based on the equilibration rates for the various DIC ions, that kinetic effects would be least likely in organisms with high flushing rates, low metabolisms, and slow shell growth rates. All three conditions, would seem to apply to *A. islandica*, and as Figure 12 shows no obvious correlation between $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, a kinetic effect related to shell growth rate seems implausible. McConnaughey [1989a] also discussed an indirect metabolic effect that would lead to a positive correlation between a photosynthetic organism's skeletal enrichment and that organism's tissue growth. He explained that if tissue growth dominated the draw down on an organism's carbon reservoir in the fluid between the skeleton and the membrane, then the residual pool available for shell formation, could be enriched in ^{13}C as tissue growth would preferentially incorporate ^{12}C . However, this mechanism should not apply to non-photosynthetic organisms, such as *A. islandica*, because their tissues receive their carbon

from what they consume and supposedly do not draw on the ambient DIC. Nevertheless, it does seem that the *A. islandica* shell's pattern of springtime ^{13}C enrichment likely coincides with the period of increased tissue growth, for whatever reason.

Ontogenic depletion of shell $\delta^{13}\text{C}$ has been reported in many bivalves [Wefer and Berger, 1991] and we have observed it consistently in *A. islandica* shells. In general, evidence of long term ^{13}C (and ^{14}C) depletion in environmental records over the last century has attracted great interest because of the possibility that it might be tracking the uptake of CO_2 from the combustion of fossil-fuels (which are very depleted in ^{13}C) [Nozaki, Rye, Turekian, and Dodge, 1978][Quay, Tilbrook, and Wong, 1992]. However, owing to the absence of ambient records of DIC $\delta^{13}\text{C}$ in most of the molluscan studies, it has not been clear that these $\delta^{13}\text{C}$ depletions are of internal (ontogenic) or external (fossil-fuel) origin. Overlapped records from *A. islandica* shells of different ages and from the same location near position of the former Nantucket Lightship (Figure 13), show a $\delta^{13}\text{C}$ offset between them for the same time periods, demonstrating that these depletions are not the result of actual changes in the ambient environment and must be related to ontogeny. These records also show depletions of 1‰ or more during the first 15 years of growth. This corresponds approximately to the age when *A. islandica* reaches sexual maturity and also corresponds with a sharp reduction in growth rate [Thompson, Jones, and Ropes, 1980][Mann, 1982]. Again, these results appear consistent with a positive correlation between tissue growth and $\delta^{13}\text{C}$ enrichment, with a reduction in the rate of tissue growth as bivalves age paralleling a depletion in their shells' $\delta^{13}\text{C}$ composition. Alternately, increased metabolic activity associated with increased reproductive activity could also explain the ontogenic depletion trend. In either case, it is apparent that a physiological or vital effect plays a dominant role in the $\delta^{13}\text{C}$ composition of *A. islandica* shells.

Summary

$\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records were obtained from the aragonite shells of three *A. islandica* specimens grown in the laboratory in different substrates for 1 year, and compared with ambient conditions such as water temperature, salinity, $\delta^{18}\text{O}$ -water, water column DIC $\delta^{13}\text{C}$, and pore water DIC $\delta^{13}\text{C}$. Assuming *A. islandica*, like most bivalves, are depositing their shells in equilibrium with ambient conditions and using Grossman and Ku's [1986] aragonite paleotemperature equation, the shells' $\delta^{18}\text{O}$ record indicates that shell growth occurred when water temperatures were between 5°C and 15°C . No significant shell growth seemed to take place at higher or lower temperatures. Also, very little shell growth seems to have occurred during the first two-thirds of the experiment. Perhaps this is the result of a combination of physical shock associated with capture, resettling into a new environment, and excessively warm tank conditions during their first few months in the laboratory. Most (70%) of shell growth occurred in the last two or three spring months of the experiment. The springtime shell $\delta^{18}\text{O}$ data, converted to temperature, closely correspond to the observed temperature curve during this period, and indicate that shell growth was approximately linear for this period. These results suggest that *A. islandica* can be used to accurately reconstruct ambient temperatures within a 10°C range from 5° to 15°C .

The $\delta^{13}\text{C}$ signals of *A. islandica* shells do not appear to be in "equilibrium" with ambient conditions, though exactly what constitutes $\delta^{13}\text{C}$ equilibrium is still unsettled within the stable isotope community. Shell $\delta^{13}\text{C}$ averages about 1‰ heavier than ambient DIC, but is about 1‰ lighter than "equilibrium" based on three of the most recently published equilibrium relationships for aragonite. The shell's seasonal variation of $\sim 2\text{‰}$ cannot be attributed, primarily, to seasonal $\delta^{13}\text{C}$ -DIC variations, which remained rather constant during the fall and spring periods of shell growth. Temperature trends in the fall and spring do correlate positively with shell $\delta^{13}\text{C}$ trends as predicted by Mook's [1986]

equilibrium relationship. However, there are significant deviations from this positive correlation as demonstrated by the poor correlation between $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ and the sharp drop in shell $\delta^{13}\text{C}$ despite rising temperatures in the late spring. Pore water $\delta^{13}\text{C}$ or the "microhabitat effect" is shown to have no obvious influence on the $\delta^{13}\text{C}$ composition of *A. islandica* shells. An ontogenic $\delta^{13}\text{C}$ depletion trend noted in *A. islandica* shells during their first ~15 years of growth cannot be accounted for by variations in the external environment. As external controls have not been identified to account for the shell $\delta^{13}\text{C}$ variations, we suggest physiological controls.

This study reaffirms the conclusions of the previous chapter, that the $\delta^{18}\text{O}$ composition of *A. islandica*'s annual bands can be used to reconstruct most of the ambient temperature record within the range of temperatures this species is naturally likely to encounter, except for the coldest winter months. In contrast, this study has shown that the $\delta^{13}\text{C}$ composition of *A. islandica*'s annual bands 1) does not directly reflect ambient $\delta^{13}\text{C}$ -DIC and 2) displays seasonal and ontogenic variations that are more consistent with a physiological control, which, until further understood, precludes its use in reconstructing records of the ocean's carbon environment.

Acknowledgements:

Financial support for this work was provided by NOAA grant # NA36GPO291, Oak Ridge Associated Universities Global Climate Change Fellowship Program, and the WHOI Ocean Ventures Fund. I especially want to thank the personnel of the NOAA National Marine Fisheries Aquarium at Woods Hole for providing the laboratory space as well as their invaluable help in carrying out the experiment. I also want to thank Matthew Gould for skippering the *Mytilus* to and from the Devil's Bridge site on good days and bad, and Tuddy Erkhart of Jonesport, Maine for offering his boat, time, and friendly assistance in collecting the specimens for this experiment. Finally, I want to thank my son Dylan for filling in for me while I was away.

References:

- Allard, D. A., Masters Thesis, University of Rhode Island, 1989.
- Aller, R. C., Experimental studies of changes produced by deposit feeders on pore water, sediment, and overlying water chemistry, *Am. J. Sci.*, 278, 1185-1234, 1978.
- Aller, R. C., and J. Y. Yingst, Effects of the marine deposit-feeders *Heteromastus filiformis* (polychaeta), *Macoma balthica* (bivalvia) and *Tellina texana* (bivalvia) on averaged sedimentary solute transport, reaction, and microbial distributions, *J. Mar. Res.*, 43, 615-645, 1985.
- Broecker, W. S., and T. H. Peng, *Tracers in the Sea*, Lamont-Doherty Geological Observatory Publication, Columbia University, Palisades, N.Y., 1982.
- Deuser, W. G., and E. T. Degens, Carbon isotope fractionation in the system CO₂(gas)-CO₂(aqueous)-HCO₃-(aqueous), 215, 1033-1035, 1967.
- Fairbanks, R. G., The origin of Continental Shelf and Slope Water in the New York Bight and Gulf of Maine: evidence from H₂O-18/ H₂O-16 ratio measurements, *J. Geophys. Res.*, 87, 5796-5808, 1982.
- Grossman, E. L., and T. Ku, Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects, *Chem. Geol.*, 59, 59-74, 1986.
- Grossman, E. L., Stable isotopes in modern benthic foraminifera: A study of vital effect, *J. Foram. Res.*, 17, 48-61, 1987.
- Harrington, R. J., Aspects of growth deceleration in bivalves: Clues to understanding the seasonal O-18 and C-13 record -- A comment on Krantz et al. (1987), *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 70, 399-403, 1989.
- Inman, D. L., Sediments: physical properties and mechanics of sedimentation, in: *Submarine Geology*, edited by F. P. Shepard, pp. 101-151, Harper's Geoscience Series, Harper and Row, New York, 1963.
- Jones, D. S., Annual cycle of shell growth and reproduction in the bivalves-*Spisula solidissima*-&-*Arctica islandica*-, PhD, Princeton University, 1980a.
- Jones, D. S., Annual cycle of shell growth increment formation in two continental shelf bivalves and its paleoecologic significance, *Paleobio.*, 6, 331-340, 1980b.
- Krantz, D. E., D. F. Williams and D. S. Jones, Ecological and paleoenvironmental information using stable isotope profiles from living and fossil molluscs, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 58, 249-266, 1987.
- Kraus, M. G., B. F. Beal, S. R. Chapman and L. McMartin, A comparison of growth rates in *A. islandica* (Linnaeus, 1967) between field and laboratory populations, *J. Shellfish Res.*, 11, 289-294, 1992.

- Lutz, R. A., J. G. Goodsell, M. Castagna and A. P. Stickney, Growth of experimentally cultured ocean quahogs (*Arctica islandica* L.) in north temperate embayments, *J. World Maricult. Soc.*, 14, 185-190, 1983.
- Mann, R., The seasonal growth cycle of gonadal development in *Arctica islandica* from the southern New England Bight, *Fish. Bull.*, 80, 315-326, 1982.
- McConnaughey, T., C13 and O18 isotopic disequilibrium in biological carbonates: I. patterns, *Geochim. Cosmochim.*, 53, 151-162, 1989a.
- McConnaughey, T., C13 and O18 isotopic disequilibrium in biological carbonates: II. *In vitro* simulation of kinetic isotope effects, *Geochim. Cosmochim.*, 53, 163-171, 1989b.
- McCorkle, D. C., S. R. Emerson and P. D. Quay, Stable carbon isotopes in marine porewaters, *Earth Planet. Sci. Lett.*, 74, 13-26, 1985.
- McCorkle, D. C., and L. D. Keigwin, Depth profiles of $\delta^{13}\text{C}$ in bottom water and core-top *C. wuellerstorfi* on the Ontong-Java Plateau and Emperor Seamounts, *Paleocean.*, 9, 197-208, 1994.
- McCorkle, D. C., L. D. Keigwin, B. H. Corliss and S. R. Emerson, The influence of microhabitats on the carbon isotopic composition of deepsea benthic foraminifera, *Paleocean.*, 5, 161-185, 1990.
- McNichol, A. P., G. A. Jones, D. L. Hutton and A. R. Gagnon, The rapid preparation of seawater ΣCO_2 for radiocarbon analysis at the National Ocean Sciences AMS Facility, *Radiocarbon*, 36, 237-246, 1994.
- Mook, W. G., C13 in atmosphere CO_2 , *Neth. J. Sea Res.*, 20, 211-223, 1986.
- O'Neil, J. R., R. N. Clayton and T. K. Mayeda, Oxygen isotope fractionation in divalent metal carbonates, *J. Chem. Phys.*, 51, 5547-5558, 1969.
- Nozaki, Y., D. M. Rye, K. K. Turekian and R. E. Dodge, C-13 and C-14 variations in a Bermuda coral, *Geophys. Res. Lett.*, 5, No.10, 825-828, 1978.
- Quay, P. D., B. Tilbrook and C. S. Wong, Oceanic uptake of fossil fuel CO_2 : carbon-13 evidence, *Science*, 256, 74-79, 1992.
- Romanek, C. S., E. L. Grossman and J. W. Morse, Carbon isotopic fractionation in synthetic aragonite and calcite: effects of temperature and precipitation rate, *Geochim. Cosmochim.*, 56, 419-430, 1992.
- Smith, P. C., The mean and seasonal circulation off southwest Nova Scotia, *J. Phys. Oceanogr.*, 13, 1034-1054, 1983.
- Tanaka, N., M. C. Monaghan and D. M. Rye, Contribution of metabolic carbon to mollusc and barnacle shell carbonate, *Nature*, 320, 520-523, 1986.
- Thompson, I., D. S. Jones and J. W. Ropes, Advanced age for sexual maturity in the ocean quahog *Arctica islandica* (Mollusca: Bivalvia), *Mar. Biol.*, 57, 35-39, 1980.

- Weidman, C. R., G. A. Jones and K. C. Lohmann, The long-lived mollusc *Arctica islandica*: A new paleoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern North Atlantic Ocean, *J. Geophys. Res.*, **99**, 18,305-18,314, 1994.
- Wefer, G., and W. H. Berger, Isotope paleontology: growth and composition of extant calcareous species, *Mar. Geol.*, **100**, 207-248, 1991.
- Wilbur, K. M., and G. Owen, Growth, in: *Physiology of Mollusca*, edited by K. M. Wilbur and C. M. Yonge, pp. 211-242, Academic Press, New York, 1964.
- Williams, D. F., M. A. Arthur, D. S. Jones and N. Healy-Williams, Seasonality and mean annual sea surface temperatures from isotopic and sclerochronological records, *Nature*, **296**, 432-434, 1982.
- Winter, J. E., Über den Einfluss der Nahrungskonzentration und anderer Faktoren auf Filtrierleistung und Nahrungsausnutzung der Muscheln *Arctica islandica* und *Modiolus modiolus*, *Mar. Biol.*, **4**, 87-135, 1969.
- Witbaard, R., M. I. Jenness, K. Van Der Borg and G. Ganssen, Verification of annual growth increments in *Arctica islandica* L. from the North Sea by means of oxygen and carbon isotopes, *Neth. J. of Sea Res.*, **33**, 91-101, 1994.

Table 1: Ambient Water Properties-Initial Sites and Laboratory

Date	Experiment Day #	Temperature (C°)	Salinity (‰)	$\delta^{18}\text{O}$ (SMOW)	$\delta^{13}\text{C}$ (PDB)	
5/4/93	1	7.1	32.185	-1.13	1.22	Devil's Bridge
6/4/93	32	5.2	31.521	-1.49	1.10	Jonesport
5/5/93	2	12.5	31.290	-1.33	0.89	
6/3/93	31	15.8	31.286	-1.35	0.72	
6/18/93	46	15.9	31.264	-1.40	0.18	
7/1/93	59	15.2	31.222	-1.43	0.38	
7/15/93	73	15.9	31.121	-1.41	0.19	
7/28/93	86	16.2	31.190	-1.50	0.12	
8/16/93	105	16.3	31.286	-1.48	-0.21	
8/30/93	119	17.3	31.439	-1.48	0.54	
9/12/93	132	18.3	31.466	-1.39	0.08	
10/6/93	156	16.5	31.628	-1.47	0.35	
10/18/93	168	15.7	31.779	-1.33	0.68	
11/6/93	187	12.6	31.670	-1.22	0.76	
11/25/93	206	9.8	31.706	-1.40	0.67	
12/10/93	221	8.5	31.418	-1.43	0.54	
12/27/93	238	2.9	31.722	-1.45	0.71	
1/13/94	255	1.5	31.200	-1.49	-----	
1/27/94	269	-0.0	31.314	-1.47	0.96	
2/10/94	283	0.3	31.221	-1.49	0.99	
2/28/94	301	1.7	31.210	-1.49	0.61	
3/18/94	319	2.8	30.798	-1.58	0.92	
4/5/94	337	6.6	30.765	-1.59	0.92	
4/24/94	356	9.7	30.860	-1.52	0.99	
5/16/94	378	12.6	30.903	-1.47	0.80	
6/4/94	397	15.7	31.174	-1.37	0.95	
6/10/94	403	16.3	31.292	-1.34	0.82	
Mean		10.8	31.298	-1.44	0.62	
$\pm\sigma$		± 6.3	± 0.282	± 0.08	± 0.35	

Table 2: Ambient Pore Water and Water Column DIC Properties (Final State) 6/10/94

ID	Substrate	Depth (cm)	ΣCO_2 ($\mu\text{mol/ml}$)	$\delta^{13}\text{C}$ (‰)
BW-1	Water	0	2.011	+0.99
BW-2	Water	0	1.997	+0.97
BS1-1	Sand	1.0	3.204	-4.56
BS1-2	Sand	3.0	3.540	-5.46
BS1-3	Sand	6.0	3.373	-4.73
BS2-1	Sand	1.0	3.164	-5.05
BS2-2	Sand	3.0	3.385	-5.29
BS2-3	Sand	6.0	3.520	-4.36
BM1-1	Mud	0.0-1.4	2.944	-3.75
BM1-2	Mud	1.4-2.8	4.815	-8.51
BM2-1	Mud	0.0-1.4	2.410	-2.17
BM2-2	Mud	1.4-2.8	2.756	-3.88
BM2-3	Mud	2.8-5.6	3.269	-5.62

Figure Captions

Figure 1. Location Map

Figure 2. Laboratory Tank Water Properties: a) temperature; b) salinity; c) $\delta^{18}\text{O}$ -water; d) $\delta^{13}\text{C}$ -DIC.

Figure 3. Devil's Bridge Water Properties: a) temperature; b) salinity; c) $\delta^{18}\text{O}$ -water; d) $\delta^{13}\text{C}$ -DIC.

Figure 4. Individual shell oxygen (solid line) and carbon (dashed line) stable isotope profiles versus sample sequence. The double vertical lines represent the growth "ring" locations. The pre-laboratory growth / laboratory growth parts of the profiles are separated by the third "ring" from the left, with pre-laboratory growth to the left and laboratory growth to the right. a) specimen qx156 grown in no sediment, b) specimen qx94 grown in medium/coarse sand, c) specimen qx102 grown in mud (very fine sand/silt).

Figure 5. Comparative a) oxygen and b) carbon stable isotope profiles versus normalized shell growth. The "0" normalized growth position separates the pre-laboratory profile (-1 to 0) from the laboratory profile (0 to +1). The specimens' substrate types are indicated by solid line (no sediment), dashed line (sand), and dotted line (mud). Note that the pre-laboratory growth profiles display much less variation both individually and collectively than to the laboratory growth profiles.

Figure 6. Predicted laboratory $\delta^{18}\text{O}$ -aragonite profile (dashed) compared with $\delta^{18}\text{O}$ -shell profiles (solid). The profiles are plotted assuming constant continuous growth. The horizontal lines indicate the mean maximum and mean minimum $\delta^{18}\text{O}$ -shell values. Note that the amplitude of the $\delta^{18}\text{O}$ -shell profiles is approximately half that of the predicted profile, and suggests that either 1) the shells did not grow during the coldest and the warmest periods, or 2) that the specimen's were not depositing their aragonite in equilibrium. Also, note the profiles are not in phase indicating variable growth rates including, perhaps, periods of shell growth cessation.

Figure 7. Predicted $\delta^{18}\text{O}$ -aragonite profile and $\delta^{18}\text{O}$ -shell data. The $\delta^{18}\text{O}$ -shell data are plotted in time by matching up their maxima and minima with corresponding predicted $\delta^{18}\text{O}$ -aragonite values, and assuming periods of no shell growth. The double vertical lines represent the shell growth hiatuses. The substrate type of the $\delta^{18}\text{O}$ -shell data is indicated by: 1) 'o' no sediment; 2) '+' sand; and 3) '*' mud.

Figure 8. Spring tank temperature versus shell-derived temperature data for the same period. The shell-derived temperatures closely correspond to the actual ($\text{SEE} = \pm 0.95^\circ\text{C}$), and indicate that the shell growth is approximately linear during this period.

Figure 9. Tank $\delta^{13}\text{C}$ -DIC (solid) and three predicted $\delta^{13}\text{C}$ -aragonite profiles (dashed): 1) Romanek, et al.'s [1992] constant offset relationship (Ro); 2) Mook's [1986] positive temperature-dependent relationship (equation 3a) (Mo); 3) Grossman and Ku's [1986] negative temperature-dependent relationship (equation 3b) (Gr). Currently, there is no consensus on "equilibrium" fractionation between aragonite and ambient DIC.

Figure 10. Laboratory $\delta^{13}\text{C}$ -DIC profile (dashed) compared with $\delta^{13}\text{C}$ -shell profiles (solid). The profiles are plotted assuming constant continuous growth. The shells are more enriched than the DIC by about 1‰ on average and the shell profiles show some phase similarity to the DIC profile. However, the $\delta^{13}\text{C}$ -shell signals have a larger amplitude than the $\delta^{13}\text{C}$ -DIC signal. The $\delta^{13}\text{C}$ profile of the specimen grown in the water column (no sed) is actually lighter than that of the specimens grown in substrates whose pore waters are considerably lighter, indicating that pore water composition does not influence the shells' carbon isotope composition.

Figure 11. Laboratory $\delta^{13}\text{C}$ -DIC profile (solid), predicted $\delta^{13}\text{C}$ -aragonite profile (dashed) [Mook, 1986], and $\delta^{13}\text{C}$ -shell data. The $\delta^{13}\text{C}$ -shell data are plotted in time according to the same scheme used for their $\delta^{18}\text{O}$ -shell pairs in Figure 7. The vertical lines separate periods of growth and no growth. The substrate type of the $\delta^{13}\text{C}$ -shell data is indicated by: 1) 'o' no sediment; 2) '+' sand; and 3) '*' mud. Taken all together the fall shell and DIC data have about the same average value, while the spring shell data are about 1‰ heavier than the DIC. While the $\delta^{13}\text{C}$ -DIC remains rather constant during each growth period the $\delta^{13}\text{C}$ -shell shows variation. The trend of some of this variation is similar to Mook's [1986] predicted $\delta^{13}\text{C}$ -aragonite indicating a positive temperature dependence, however the sharp depletion trend near the shell profiles termination runs counter to this interpretation.

Figure 12. Laboratory shell $\delta^{13}\text{C}$ versus $\delta^{18}\text{O}$. No significant correlation exists between shell $\delta^{13}\text{C}$ versus $\delta^{18}\text{O}$.

Figure 13. Overlapped long term $\delta^{13}\text{C}$ records from two shells of different ages live-captured at the same time and at the same location on Nantucket Shoals. The first ~15 years of growth in each shell show a 1-1.5‰ depletion. The records overlap the years 1946-1956. The overlapping part of the record of the older specimen (solid line) averages about +1.8‰ and shows only a slight negative trend, while the overlapping part of the record of the younger specimen (dashed) averages about +2.3‰ and shows a steep negative trend. These results are interpreted as evidence of a strong physiological control on the $\delta^{13}\text{C}$ composition of *A. islandica* shells.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

Chapter 4

Time History of $\Delta^{14}\text{C}$ in the Surface North Atlantic Ocean

Introduction

The northern North Atlantic Ocean is thought to play a critical role in the regulation of global climate and ocean circulation [Broecker, 1991][Keigwin and Jones, 1994][Stocker and Broecker, 1994][Dickson and Brown, 1994]. This role is characterized by a large net meridional transport of heat from the lower-latitude ocean to the higher-latitude atmosphere and, as a result of this oceanic heat-loss to the atmosphere, the production of a large share of the world ocean's deep water and the warming of northern Europe's climate [Stommel, 1957][Worthington, 1970][Warren, 1981]. In addition, deepwater production provides a mechanism for sequestering anthropogenic CO_2 [Broecker, & Peng, 1992], and hence the character of this region's circulation provides important constraints on global climate changes associated with the "green-house" effect. In the last four decades, one of the most important techniques used to understand oceanographic processes has been the measurement of ocean radiocarbon ($\Delta^{14}\text{C}$). As a time-dependent tracer, $\Delta^{14}\text{C}$ provides oceanographers with a method of determining the structure and rates of the ocean's steady state physical circulation [Broecker, & Peng, 1982][Toggweiler, Dixon, & Bryan, 1989a] [Maier-Reimer, & Bacastow, 1990][Schlosser, et al., 1995] as well as the ocean's steady state absorption of atmospheric carbon dioxide [Revelle and Suess, 1957]. Although confounding steady state applications, nuclear bomb-produced ^{14}C has become a valuable tracer for measuring the ocean's transient uptake of atmospheric CO_2 , as well as a means of measuring water mass transport in the ocean [Broecker, et al., 1982][Quay, Stuiver, & Broecker, 1983][Toggweiler, Dixon, & Bryan, 1989b][Broecker, 1991]. In particular, global and regional ocean

sampling programs, such as GEOSECS (1970s), TTO (1980s), and WOCE (1990s), have provided snapshots of these tracers' ocean distribution. The temporal evolution of the ocean's bomb- ^{14}C signal has been more difficult to assess owing to the global scope and cost of seawater sampling programs. But since the late 1970s, beginning with Nozaki et al. [Nozaki, Rye, Turekian, & Dodge, 1978] and Druffel and Linick [Druffel, & Linick, 1978a], long, high-resolution time-series of surface ocean $\Delta^{14}\text{C}$ have been reconstructed from banded corals. These records have provided important pre-bomb (before 1958) information concerning the ocean's uptake of fossil-fuel produced CO_2 , as well as post-bomb information useful in studying the nature of upper ocean circulation [Toggweiler, Dixon, & Broecker, 1991] [Druffel, 1989]. However, these time-series have been limited to the warm subtropical oceans, the habitat of fast-growing banded-corals, and there is still a near absence of comparable records for the high-latitude oceans, and a corresponding lack of insight that radiocarbon time-series could provide for the globally important circulation processes operating in these regions.

Only recently has the reconstruction of high-resolution $\Delta^{14}\text{C}$ records for the high-latitude oceans become practical with the introduction of AMS-radiocarbon dating technology [Linick, Damon, Donahue, & Jull, 1989], which has enabled the analysis of very small samples (~10 mg) from the skeletal remains of cold ocean marine organisms, such as otoliths and the shells of slow-growing, long-lived molluscs. For example, Kalish [1993] employed AMS-dating of fish otoliths to reconstruct a 70-year radiocarbon time-history for New Zealand (39°S) coastal waters. Weidman and Jones [1993] used micro-sampling methods to extract carbonate samples from the shells of the long-lived mollusc *Arctica islandica* and AMS-analysis to reconstruct a 50-year radiocarbon time history for Georges Bank on the northeastern (41°N) U.S. continental shelf. Also, Jones and Weidman (in preparation) have similarly obtained a 70-year radiocarbon time-series from a long-lived mollusc collected on South Georgia Island shelf (55°S) within the Antarctic

Circumpolar Current. These studies have initiated the application of AMS-dating and micro-sampling methods to obtain cold ocean radiocarbon time-series. This paper builds on our earlier work on Georges Bank, by applying these techniques to obtain a set of radiocarbon time histories for the northeastern North Atlantic Ocean. These records are combined with Druffel's two coral-derived records from Bermuda and Florida [Druffel, 1989] to obtain a more complete picture of the evolution of the ^{14}C signal in the surface North Atlantic Ocean over the last century than has previously been possible.

General circulation of the North Atlantic and the surface $\Delta^{14}\text{C}$ signal:

The three new shell-derived time-histories presented here are all located in the eastern North Atlantic, while three previously published time histories, represented here, are from the western North Atlantic (Figure 1a). The three eastern North Atlantic shell collection sites are 1) southern North Sea ($54^{\circ} 24'\text{N}$, $6^{\circ} 30'\text{E}$) at 37 m depth, 2) Siglufjordur on the north coast of Iceland ($66^{\circ} 10'\text{N}$, $18^{\circ} 53'\text{W}$) at 22 m depth, and 3) Tromsøysund, on the northern coast of Norway ($69^{\circ} 38'\text{N}$, $18^{\circ} 57'\text{E}$) at 3 m depth. The three western North Atlantic shell and coral collection sites are 1) Georges Bank (41°N , 67°W) at 76 m depth [Weidman and Jones, 1993], 2) Florida (24°N , 81°W) at 4 m depth [Druffel, 1989], and 3) Bermuda (32°N , 65°W) at ~ 10 m depth [Druffel, 1989]. These coastal and shelf locations are adjacent to large-scale features of the North Atlantic's general surface circulation that will be described next, in broad terms, to provide a general oceanographic context for the radiocarbon time histories.

Although many quantitative aspects are still uncertain, the overall qualitative view of the North Atlantic's surface circulation is not very different from that proposed by Sverdrup, et al. [Sverdrup, Johnsen, & Fleming, 1942] (Figure 1a). In this simple view, two coupled basin-wide systems dominate the surface circulation in the North Atlantic described by an anti-cyclonic subtropical gyre and a cyclonic subpolar gyre [McCartney

and Talley, 1982][Schmitz, & McCartney, 1993]. In this paper, the term "subpolar gyre" refers to the greater subpolar gyre, in the manner of Dickson, et al. [1988] and Ellet [1993], which includes the Nordic Seas (e.g., the path of the Great Salinity Anomaly), as opposed to the lesser subpolar gyre, which refers only to the cyclonic recirculating flows of the Labrador Sea and Irminger Sea. The two systems are connected by a continuous northward surface flow along the western side of the subtropical gyre (Gulf Stream) and the south side of the subpolar gyre (North Atlantic Current). This northward surface flow is thought to largely balance the sub-surface southward flow of North Atlantic Deep Water (NADW), and together constitute the principal features of the North Atlantic's thermohaline circulation cell. The two North Atlantic surface gyres also can be described roughly by their mean temperature fields at 150 m depth [Robinson, Bauer, & Schroeder, 1979] (Figure 1b). The subtropical gyre region is defined by a zone with temperatures above 15°C, and the subpolar gyre region is defined by a zone with temperatures below 10°C. The zone between 10°C and 15°C, which incorporates much of the central eastern North Atlantic, has been described as a transition area between subpolar and subtropical waters [Krauss, 1986]. Another, subdivision can be made by separating the subpolar region into an eastern section with warmer temperatures between 5°-10°C and a western section with colder temperatures below 5°C. The eastern section includes the poleward flowing Atlantic waters of the Irminger Current on its west side and the North Atlantic Current and its extension the Norwegian Current on its east side. The western section includes the southward flowing polar waters of the East Greenland Current, the Arctic waters of the Greenland Sea, and the Labrador Sea. Lastly, due to strong seasonal cooling and increased cyclone frequency, the higher-latitude regions, especially in the west, experience much greater convective mixing than the lower-latitudes, expressed by a general deepening of the winter mixed-layer from south to north [Robinson, et al., 1979] (Figure 1c).

Broecker and Peng [1982] presented a latitudinal profile of surface $\Delta^{14}\text{C}$ for the Atlantic Ocean from GEOSECS and pre-bomb ocean water data (Chapter 1, Figure 1). The salient features of this profile are higher $\Delta^{14}\text{C}$ in the mid-latitudes and lower $\Delta^{14}\text{C}$ in the equatorial regions and the high-latitudes. The cause of the latitudinal $\Delta^{14}\text{C}$ variations are ascribed to ocean circulation processes, such as upwelling in the equatorial ocean which brings deep (low $\Delta^{14}\text{C}$) waters to the surface, subtropical convergence of surface (high $\Delta^{14}\text{C}$) waters, which have had a longer time to equilibrate with the atmosphere, and deep convective mixing in the high-latitudes which overturns the surface and deep layers and thus lowers the surface $\Delta^{14}\text{C}$.

Coastal environments and $\Delta^{14}\text{C}$ signal:

Within this ocean circulation framework, each of the six sites is also located within a shelf environment, and so its $\Delta^{14}\text{C}$ signal may differ from that of the adjacent ocean. The evolution of the surface ocean's $\Delta^{14}\text{C}$ concentration at any one location is a function of its initial value, the difference between the surface ocean and atmosphere $\Delta^{14}\text{C}$ concentrations, the CO_2 air-sea surface gas transfer rate (usually treated as a function of the regional mean wind speed), mixed-layer depth, vertical mixing/exchange rates with deeper ocean layers, and horizontal advection. These parameters have important ramifications for the $\Delta^{14}\text{C}$ signals of shelf waters. For example, mixed-layer depth is limited to the shelf depth, vertical exchange with deeper layers cannot occur, and the longer ocean water is isolated on the shelf, the more it will equilibrate with the atmosphere.

The German Bight region of the southern North Sea is primarily (90-95%) supplied by waters from the English Channel [Reid, Taylor, & Stephens, 1988], which itself is fed by surface waters from the central eastern North Atlantic [Pingree and Le Cann, 1989][Pingree, 1993]. The shelf pathway for waters from the English Channel to the German Bight site is shallow (< 50m), and waters at the North Sea site have a residence

time estimated to be ~ 4-6 months from the Dover Straits, as well as additional time in the shelf regions (English Channel and Celtic and Armorican Shelves) to the west [Reid, *et al.*, 1988]. These factors suggest a strong shelf influence on the North Sea's $\Delta^{14}\text{C}$ signal relative to that of its central eastern North Atlantic source.

Iceland is near the center of the greater subpolar gyre with southward flowing Arctic waters to its north and west and northward flowing Atlantic waters to its south and east. The northern Icelandic shelf waters are generally supplied by Atlantic water of Irminger Current origin (the north Icelandic Irminger Current), however considerable mixing with waters of Arctic origin occurs during the winter [Stefansson, 1962][Johannessen, 1986]. In addition, the Atlantic versus Arctic component can vary widely from year to year as the result of climatic variations [Stefansson, & Olafsson, 1991]. Although, the Siglufjordur site is relatively shallow (~20 m), it is on a narrow part of the shelf adjacent (~10 km) to a deep region (Skagafjardar-Djup) and its $\Delta^{14}\text{C}$ signal might therefore show only slight modification from that of the adjacent ocean.

The Tromsøysund, Norway site is adjacent to the offshore Norwegian Current which is the northern extension of the eastern limb of the subpolar gyre as we have defined it. However, this site is within a complex fjord system (~50 km inland) fed by shelf waters which are part of the Norwegian Coastal Current, a relatively fresh current extending northward from the Skaggeiat at the mouth of the Baltic Sea to the Barent's Sea [Johannessen, 1986][Gade, 1986]. In the northern Norwegian shelf region considerable mixing occurs between the offshore (Norwegian Current) and the inshore Norwegian Coastal Current waters [Rey, 1981]. Despite these complexities, tidal currents within Tromsøysund are strong and the hydrography (salinity and temperature) shows strong continuity and little modification from the shelf waters of the adjacent western coast [Normann, 1992]. These observations imply that the Tromsøysund $\Delta^{14}\text{C}$ record probably

reflects that of the shelf waters with little modification, but the influence of the coastal current (with its Baltic and runoff sources) may cause it to vary from that of the adjacent Norwegian Sea.

The Bermuda and Florida sites are both within the western region of the North Atlantic encompassed by the subtropical gyre. Druffel [1989] makes the reasonable assumption that the shallow waters of Bermuda's reef platform (~15 km at greatest extent) exchange rapidly with the surrounding Sargasso Sea surface. Thus, Bermuda's coral-derived $\Delta^{14}\text{C}$ signal is not expected to differ significantly from the adjacent oceanic $\Delta^{14}\text{C}$ signal, and should be representative of the mid-subtropical gyre region. The Florida coral was also obtained from a shallow reef platform and Druffel and Linick [1978] noted slightly lower salinity here than for the offshore Gulf Stream, and allowed for some slight Florida Bay influence. However, this site is located on a very narrow part of the shelf and only ~15 km from the center line of the Florida Current so that it is reasonable to assume, as Druffel [1989] did, that the Florida coral $\Delta^{14}\text{C}$ signal is essentially the same as that of the surface Gulf Stream.

The upstream source region for shelf water on Georges Bank is thought to be the Labrador Sea [Chapman, & Beardsley, 1989]. From estimates of the mean alongshelf velocity (~10 cm/s) the shelf water residence time (relative to the Labrador Sea) for water on Georges Bank is estimated to be about 1 year. Although the site where the shell was collected is relatively deep (76 m) and overlying waters in this region of Georges Bank are stratified during the warmer months of the year, complete vertical mixing occurs during the colder months of the year. It was argued in Weidman and Jones [1993] (Chapter 1) that the $\Delta^{14}\text{C}$ signal on Georges Bank is modified relative to the Labrador Sea $\Delta^{14}\text{C}$ signal due to enhanced atmospheric equilibration of the shelf waters.

Methods

All shells used in this study were obtained from live-captured specimens collected from dredge hauls. Shell samples were prepared by slicing each shell radially from the umbo through the ventral edge along the axis of maximum growth (see Chapter 1, Figure 3) using a slow-speed petrographic saw with twin diamond-edged blades. A shell slice ~2 mm thick was obtained this way and then epoxied onto a petrographic slide. The slide was mounted on an automated 3-axis stage and the outer shell layer of the slice was subsectioned into samples comprised of one or more annual bands with an ultra-thin (150 μm) diamond saw. Care was taken to extract only carbonate from the outer shell layer, and not from the adjacent inner nacreous layer which represents material deposited at a later time. Also, the periostracum (or outer organic layer) was removed, along with part of the outer prismatic layer just below the periostracum, in order to eliminate inclusion of surface contaminants or material affected by early diagenesis. The number of annual bands comprising a sample was constrained only by the minimum amount of aragonite required for a single AMS radiocarbon analysis (~10 mg). The shell of any one specimen contains a continuous record for the period of its life-span, however, since shell growth slows as the bivalve ages, sampling-resolution, and likewise temporal-resolution, is reduced in the later (more recent) stages of shell growth. Sometimes to overcome this handicap, samples from the inner, faster growth intervals from two or more shells of different ages were analyzed to produce a higher-temporal resolution record than otherwise would be obtainable from a single shell. Samples weighed from ~10-80 mg and integrated from 1 to 10 years (Table 1).

Samples were prepared for AMS analysis by first etching them in 10% HCl for 30 seconds to remove surface contaminants, followed by conversion to CO_2 in anhydrous phosphoric acid at 60°C (overnight), and finally conversion to graphite using a hydrogen

reduction and iron catalytic process [Gagnon and Jones, 1993]. Graphite targets were analyzed at the National Ocean Sciences AMS Facility at Woods Hole Oceanographic Institution, Woods Hole, Massachusetts. Analytical $\Delta^{14}\text{C}$ error for these samples ranged from $\pm 3.0\text{‰}$ to $\pm 13.5\text{‰}$ with a mean of $\pm 5.7\text{‰}$. All $\Delta^{14}\text{C}$ data are age-corrected to 1950 and normalized to $\delta^{13}\text{C}$ (PDB) value of -25‰ , according to methods outlined by Stuiver and Pollach [1977]. To correct for fractionation, $\delta^{13}\text{C}$ values were obtained either from sub-samples of the same material analyzed for $\Delta^{14}\text{C}$, interpolation with adjacent samples from within the same shell, or the mean $\delta^{13}\text{C}$ of other *A. islandica* shells in the region. *A. islandica* shell $\delta^{13}\text{C}$ values averaged $\sim +2\text{‰}$.

Results

The time histories for the eastern North Atlantic (the new results) are described first, but all six North Atlantic records (from 1940 on) are shown in Figure 2(a-f). Shell-derived data are also shown in Table 2(a-d). Coral-derived data are tabulated in Druffel [1989].

Eastern Atlantic $\Delta^{14}\text{C}$ Time Histories (New Results)

North Sea:

The German Bight $\Delta^{14}\text{C}$ record spans from 1948 to 1990 (Figure 2a, Table 1a) and is a composite of two overlapping shell records. This record has a minimum value of -83.3‰ in 1954, and a maximum value of $+239.3\text{‰}$ in 1967-1968, for an amplitude of 323‰ . The pre-bomb (before 1958) data have a mean value of -65.8‰ , and show a large depletion of $\sim 33\text{‰}$ between 1948 and 1954. The first definite bomb influence (significant rise above previous levels) in the shell record occurred in 1959. The steepest signal increase occurred between the years 1961 and 1966. The North Sea $\Delta^{14}\text{C}$ signal remained

near its maximum value for about 7 years until 1974-1975, and afterwards decayed by ~100‰ until the early 1980s, but has changed little since then.

Iceland:

The Siglufjörður $\Delta^{14}\text{C}$ time-history spans from 1872 to 1991 (Figure 2b and Table 1b), is composed of samples from three shells, and is the longest shell-derived $\Delta^{14}\text{C}$ record obtained thus far. This record has a minimum value of -59.5‰ in 1912-1921, and a maximum value of +56.9‰ in 1970-1971, for an amplitude of 116‰. The pre-bomb (before 1958) data have a mean value of -55.8‰, and show no significant trend. The first definite bomb influence in the shell record occurred in 1960. The steepest signal increase occurred between the years 1960 and 1964. After the peak in 1970-1971, the signal dropped to a local minimum in the mid-1970s, and rose to a second local maximum in the late 1970s. The signal declined slightly in the early 1980s and changed very little for the rest of the decade.

Norway:

The Tromsøysund $\Delta^{14}\text{C}$ time-history spans from 1940 to 1993 (Figure 2c and Table 1c) and is from a single shell. This record has a minimum value of -82.3‰ in 1956-1957, and a maximum value of +132.2‰ in 1969-1970, for an amplitude of 215‰. The pre-bomb (before 1958) data have a mean value of -60.9‰, and show a large depletion of ~36‰ between 1940 and 1956-1957. The first definite bomb influence in the shell record occurred in 1959-1960. The steepest signal increase occurred between the years 1957 and 1970. After the peak, the signal dropped sharply in the mid-1970s, but rose again to a second local maximum in the later 1970s. Between 1977 and 1988 the signal decayed by ~60‰, but has changed little since 1988.

Western North Atlantic $\Delta^{14}\text{C}$ Time Histories (Previously Published)

Bermuda:

Druffel's [1989] coral-derived Bermuda $\Delta^{14}\text{C}$ time history spans from 1950-1983 (Figure 2d). The Bermuda coral record has a minimum value of -54.8‰ in 1951, and a maximum value of $+159.2\text{‰}$ in 1977, for an amplitude of 214‰ . The pre-bomb (before 1958) data have a mean value of -46.7‰ , but this record does not extend far enough back to calculate a pre-bomb trend. However, Nozaki, et al. [1978] showed a depletion of $\sim 25\text{‰}$ between 1900 to 1950 from their coral-derived Bermuda record. The first definite bomb influence appears in the coral record in 1959. The steepest signal increase occurs between the years 1962 and 1964. Though the peak value occurs in 1977, the bomb-signal also shows an earlier local maximum of almost equal magnitude ($+157.6\text{‰}$) in 1973, and otherwise remains within $\sim 10\text{‰}$ of its maximum value for the decade from 1971 to 1981. The signal drops sharply in the early 1980s and the record ends in 1983, so no comparison can be made with the shell-derived records for the last decade.

Florida:

Druffel's [1989] coral-derived Florida $\Delta^{14}\text{C}$ time history spans from 1951-1983 (Figure 2e). The Florida coral record has a minimum value of -66.0‰ in 1952, and a maximum value of $+156.0\text{‰}$ in 1970, for an amplitude of 222‰ . The pre-bomb (before 1958) data have a mean value of -58.9‰ . This particular record also does not extend far enough back to calculate a pre-bomb trend, but Druffel and Linick [1978] published a longer record from this same site which showed a depletion of $\sim 12\text{‰}$ between 1900 and 1955. The first definite bomb influence occurred in 1958, and the steepest signal increase occurred between 1962 and 1965. The bomb signal remained near its maximum value for about 6 years, then dropped sharply in 1976, but changed little until 1983 when it dropped sharply again.

Georges Bank:

The Georges Bank $\Delta^{14}\text{C}$ time-history spans from 1939 to 1990 (Figure 2f and Table 1d) and is from a single shell. This record has a minimum value of -74.6‰ in 1951-1952, and a maximum value of $+77.0\text{‰}$ in 1973-1975, for an amplitude of 152‰ . The pre-bomb (before 1958) data have a mean value of -67.2‰ , and show a depletion of $\sim 13\text{‰}$ between 1939 and 1951-1952. The first definite bomb influence in the shell record occurred in 1959-1960. The steepest signal increase occurred between 1960 and 1965. The signal dropped off sharply after 1973-1975, but remained nearly constant throughout the 1980s.

Discussion

North Atlantic Pre-Bomb $\Delta^{14}\text{C}$

Time History Comparison:

Table 2 contains a summary of important parameters for each of the six $\Delta^{14}\text{C}$ time-histories. Pre-bomb (<1958) values range from -83‰ to -45‰ , and average -58‰ . Norway and North Sea sites have unusually low minima $< -80\text{‰}$ in the mid-1950s, which may be anomalous or represent unusual conditions (and will be discussed in a later section), and so the pre-bomb data for the period just from 1940 to 1951 (~ 1950) are also compared in Table 2. Bermuda and Georges Bank have the highest and lowest pre-bomb (~ 1950) values, respectively. This contrast is consistent with latitudinal position (see Chapter 1, Figure 1), Bermuda being in a region of low vertical exchange and converging surface flows, and Georges Bank having its source in the Labrador Sea where deep convective mixing occurs. The Florida pre-bomb (~ 1950) value is intermediate between Bermuda and Georges Bank. Druffel [1989] has explained this depletion in the Florida pre-bomb signal relative to Bermuda as evidence of an equatorial (upwelled) component in the

Florida Current. The eastern North Atlantic (~1950) pre-bomb mean values (Norway, Iceland, and the North Sea) are all approximately the same $\sim -55\text{‰}$, showing less regional contrast than western North Atlantic sites. Also, despite their higher-latitude location these sites are more enriched than the Georges Bank region.

Comparison with other pre-bomb $\Delta^{14}\text{C}$ measurements:

Pre-bomb ocean water $\Delta^{14}\text{C}$ data for the North Atlantic were obtained by Broecker, et al. [Broecker, Gerard, Ewing, & Heezen, 1960] (see Chapter 1, Figure 1) in the mid-1950s and their results show a North Atlantic mean surface water $\Delta^{14}\text{C}$ value of -49‰ ($n=18$), ranging from a high of -33‰ to a low of -65‰ . While that study noted no significant relationship between North Atlantic surface radiocarbon concentrations and specific hydrographic features, their sampling in the North Atlantic was limited to the subtropical gyre and no samples were taken north of 39°N . Fonselius and Ostlund [Fonselius, & Ostlund, 1959] obtained five pre-bomb (1957) surface ocean radiocarbon values averaging -43‰ for the northern North Atlantic above latitude 60°N . Results from the Denmark Strait ($n=3$) west of Iceland, averaged -48‰ , while two samples taken within the Norwegian Current averaged -35‰ . Broecker, et al. [1960] suggested a slight negative correction of about -3‰ to their own and Fonselius and Ostlund's data, owing to some initial bomb enrichment of the atmosphere by 1956. From the results of these two early ocean radiocarbon studies Broecker, et al., [1960] concluded that surface $\Delta^{14}\text{C}$ waters in the northern North Atlantic ($60^\circ\text{--}80^\circ\text{N}$) were more enriched ($+14\text{‰}$) than subtropical gyre ($15^\circ\text{--}40^\circ\text{N}$) surface waters, which fit broadly with the idea that the former were being replenished by the latter as part of the Atlantic's thermohaline circulation and by continued enrichment from contact with the atmosphere.

Druffel's [1989] Florida and Bermuda pre-bomb data average is -53‰ , which is effectively identical to Broecker, et al.'s [1960] average value of -52‰ for the subtropical

gyre region (using Broecker, et al.'s 3‰ correction). However, Fonselius and Ostlund's northern North Atlantic data are quite enriched (by ~15‰) compared to the shell-derived pre-bomb records for the four northern North Atlantic sites discussed here, which yield an average of -59‰. Pre-bomb $\Delta^{14}\text{C}$ data for the surface Nordic Sea regions also have been compiled from whole-shell analysis of selected museum shells [Mangerud, 1972][Mangerud and Gulliksen, 1975]. For the Norwegian Sea region, Mangerud and Gulliksen [1975] gave pre-anthropogenic means of -52‰, -57‰, and -61‰ for the southern and northern coasts of Norway and Spitzbergeb, respectively. They interpreted the northward depletion trend in their data as evidence of the increasing influence of Norwegian Current (lower $\Delta^{14}\text{C}$) and the decreasing influence of the northward flowing Norwegian Coastal Current (higher $\Delta^{14}\text{C}$), implying a relatively depleted core (< -60‰) for the Norwegian Current. Their Norwegian Sea values are in close agreement with this study's pre-1950 time history data for northeastern North Atlantic (North Sea, Norway, and Iceland). For the Greenland Sea and Denmark Strait region, these same studies showed shell-derived pre-bomb data with means of -69‰ and -68‰ from the east coast of Greenland (from Hjort, 1973) and from the southwest coast of Iceland (from Broecker and Olson, 1961), respectively. These values are more depleted than Iceland's north coast record (the nearest time history location), but are similar to the Georges Bank pre-bomb data (and our estimate for the Labrador Sea region in Chapter 1) as well as Broecker's [1979] estimate of -70‰ for the initial source of North Atlantic Deep Water (NADW).

It would appear then that Fonselius and Ostlund's [1959] ocean water data for the Nordic Sea regions are too enriched, perhaps because of bomb contamination, to represent mean pre-bomb surface conditions, and that the pre-bomb $\Delta^{14}\text{C}$ data from the shell records presented in this study are more in line with other regional pre-bomb data. In contrast, then to Broecker, et al.'s [1960] assessment, the shell pre-bomb results from this study show that the northern North Atlantic surface waters, on average, are more depleted than

subtropical gyre surface waters, and that the western part of this northern region is the most depleted region in the North Atlantic. While the difference between mid- and high latitude regions in the North Atlantic ($\Delta \sim 20\text{‰}$) is not nearly as great as in the South Atlantic and Pacific Oceans ($\Delta \sim 70\text{‰}$), the conclusion that no surface $\Delta^{14}\text{C}$ gradient exists between the subtropical and subpolar North Atlantic [Bard, 1988][Bard, et al., 1994] appears incorrect. The gradient disappears when the radiocarbon signature of the surface subpolar regions is blended out by combining data (including the Fonselius and Ostlund [1959] data) from the same latitudes, and thus masking the east/west asymmetry of the pre-bomb $\Delta^{14}\text{C}$ data. However, it is critical not to ignore this asymmetry because, as Bard, et al. [1994] have recently pointed out in their discussion of the Younger Dryas (~ 10 kyr B.P.) period, the presence and magnitude of the meridional surface $\Delta^{14}\text{C}$ gradient has important implications for the thermohaline circulation of the North Atlantic.

The pre-bomb north-south contrast in $\Delta^{14}\text{C}$ is at least partly explainable by the difference in winter mixed-layer depth and the consequent difference in surface residence time between these regions, which, in a one-dimensional sense, should allow the subtropical gyre waters to become more equilibrated with the atmosphere than the higher-latitude waters. However, because subpolar waters are thought to be largely replenished by surface waters of subtropical origin, and transport to these northern regions, via the North Atlantic Current system, should allow for further enrichment as a result of additional surface residence time, there is a need for a counterbalancing depleted source(s) to account for the lower $\Delta^{14}\text{C}$ of northern surface waters. In addition, the flushing times for the Nordic Seas and Labrador Sea basins are on the order of a decade or two [Broecker, 1979][Schlosser, et al., 1995][Bonisch and Schlosser, 1995], and so the $\Delta^{14}\text{C}$ depletion caused by radioactive decay is negligible ($\sim 1\text{--}3\text{‰}$) and should be more than compensated by the atmosphere-derived $\Delta^{14}\text{C}$ enrichment ($\sim 4\text{--}9\text{‰}$).

The particularly low values ($\sim -70\text{‰}$) of the western subpolar region highlight the problem and motivate the need for a significant deepwater source. Based on GEOSECS tritium data, Broecker [1979] calculated that the deep overflows from the Greenland and Norwegian Seas were contributing waters with an estimated pre-bomb-value of -68‰ to the NADW, and as mentioned, Mangerud and Gulliksen [1979] reported pre-anthropogenic shell values comparable to this for the Greenland Sea and Denmark Strait surface waters. Broecker [1979] further showed that, because of the short residence in the Nordic basins, the inflow $\Delta^{14}\text{C}$ concentration must approximately equal the outflow $\Delta^{14}\text{C}$ concentration. These combined results imply that the full water column within the subpolar gyre region south and west of Iceland may have carried a pre-bomb value of $\sim -70\text{‰}$. The problem is how to achieve a $\Delta^{14}\text{C}$ of -70‰ for this subpolar region if its only major sources are 1) equilibrated surface waters ($\sim -50\text{‰}$) of subtropical origin and 2) Polar waters of the East Greenland Current ($\sim -70\text{‰}$). Either, another depleted source(s) exists or subtropical surface waters do not contribute to the subpolar gyre. Since much evidence has been compiled to reject the latter option (i.e., [Lazier, 1973][*McCartney and Talley*, 1982][*McCartney and Talley*, 1984][*Schmitz and McCartney*, 1993]), a depleted source must be sought.

Two potential sources of the depleted waters are the intermediate and deep water reservoirs existing along the southern limits of the subpolar gyre that might be accessed by intense winter convection. Convection, here and northward, can extend to depths of 1000 m or more [*McCartney and Talley*, 1982]. Broecker [1979], argued that an important contributor to NADW and its $\Delta^{14}\text{C}$ composition was the underlying Antarctic Bottom Water (mid-latitude AABW: $\Delta^{14}\text{C} \sim -100\text{‰}$ at ~ 4000 m) which extends northward at least to mid-latitude in the North Atlantic. However, while he showed that the uptake of AABW quite successfully accounts for the downstream evolution of the NADW $\Delta^{14}\text{C}$ composition, it is more difficult to account for its presence in the surface subpolar waters. A second,

more viable potential source for the depleted component in northern North Atlantic surface waters is Antarctic Intermediate Water (mid-latitude AAIW: $\Delta^{14}\text{C} \sim -90\text{‰}$ at ~ 1000 m), which Tsuchiya [Tsuchiya, 1989] showed, through its silica signature, being transported well into the high-latitude (60°N) North Atlantic by the Gulf Stream-North Atlantic Current and extending along the subpolar front (Figure 3). McCartney and Talley [1982] have shown that the intense winter convection along the axis of the North Atlantic Current, Irminger Current, and the Labrador Sea allows for the formation of thick (500-1000 m) vertically homogeneous upper ocean water masses referred to as Subpolar Mode Water (SPMW). They showed that these upper ocean waters continue to evolve in temperature and salinity, increasing in density along their trajectories, and ultimately form the core waters of the Norwegian Current and Labrador Sea Water. These same processes potentially could mix surface subtropical waters with an initial (mid-latitude) value of -50‰ and underlying intermediate waters (AAIW) with an initial (mid-latitude) value -90‰ to form SPMW with a $\Delta^{14}\text{C}$ of -70‰ . In such a scheme, the contribution of Arctic water from the East Greenland Current with a similar value of -70‰ would not alter the $\Delta^{14}\text{C}$ composition of subpolar gyre water.

The presence of AAIW in the northern North Atlantic waters has been reported previously by others. Sverdrup, et al. [1942], in a simple schematic of North Atlantic intermediate water masses showed AAIW flowing northward as the underlying part of the western boundary current (Figure 188, p. 685). Broecker's [1979] paper, while focusing on the role of AABW, listed the very similar $\Delta^{14}\text{C}$, silica, and nitrate properties of Circumpolar intermediate waters, and did not preclude an important role for them in the make up of (upper) NADW. Broecker and Takahashi [1981] traced the influence of AAIW up to about 40°N in the central North Atlantic using GEOSECS results for ^{14}C , ^{226}Ra , silica, oxygen, nitrate, potential temperature, and salinity. Tanaka, et al. [1990], calculated that the Gulf of Maine $\Delta^{14}\text{C}$ budget required relatively equal inputs of AAIW (40%) and

Gulf Stream/ Sargasso Sea Water (60%). Following their arguments it was shown in chapter one [Weidman and Jones, 1993] that a 50:50 mixture of AAIW and Gulf Stream (or Sargasso Sea) Water could also account for the $\Delta^{14}\text{C}$ budget of the Labrador Sea. Maier-Reimer and Bacastow's [1990] pre-bomb $\Delta^{14}\text{C}$ tracer results from their ocean global circulation model (OGCM) show the northward penetration of an AAIW analog that results in northwestern North Atlantic surface values of -70‰ (Figure 4a). Alternately, it is the absence of this northward penetration of an AAIW analog in Toggweiler, et al.'s [1989a] OGCM pre-bomb $\Delta^{14}\text{C}$ results that allows for a more enriched (-40‰) northwestern North Atlantic (Figure 4b), which is at odds with the observed pre-bomb values in the western subpolar region. The introduction of this intermediate water component into the surface waters along the subpolar front about the south of the Labrador Sea and Iceland, would account for the lower radiocarbon concentrations of the northern North Atlantic waters in general, and the even more depleted conditions of the Labrador and Irminger Seas in particular.

A third possible source for depleted waters for the Labrador Sea region is Arctic waters from the Canadian Archipelago. Chapman and Beardsley [1989], suggested that continuity exists between the East Greenland, West Greenland, and Labrador Currents through westward flows across the Davis Strait (see Chapter 1, Figure 2). The coastal and inshore parts of these currents, being fresher and less dense than offshore Labrador Sea Water, remain somewhat distinct [Lazier, 1973] and therefore may carry the initial radiocarbon signature of the East Greenland Current (-70‰) relatively undiluted (in pre-bomb times) all the way to the Georges Bank region. Chapman and Beardsley [1989] also suggested that Arctic waters flowing southward through the Davis Strait are a major contributor (~2 Sv) to this flow, and Lazier [1973] indicated that waters from this same source (albeit through the Canadian Archipelago) form the major supply of low-salinity water to the surface of the Labrador Sea. This is significant in light of the fact that

Mangerud and Gulliksen [1975] reported three shell values from the southern coast of Ellesmere Island with a very depleted pre-bomb mean $\Delta^{14}\text{C}$ of -89‰ . However, this value is much lighter than the $\sim -50\text{‰}$ estimated by Ostlund et al. [1987], for pre-bomb Arctic surface waters of the Eurasian basin. Mangerud and Gulliksen [1975] suggest that the strong salinity stratification of the upper Arctic Ocean water effectively prevents significant exchange with the deeper waters (Canadian basin deep water $\Delta^{14}\text{C} \sim -105\text{‰}$) and explain their low Arctic shell values as either owed to a Pacific (Bering Strait) origin ($\Delta^{14}\text{C} < -100\text{‰}$) or a local depleted source. Granting these uncertainties, if the -89‰ value is typical of pre-bomb Arctic waters transported southward through the Canadian Archipelago, then these waters might be an important source of depleted waters for the Labrador Sea.

Suess Effect:

The "Suess effect" refers to the reservoir depletion caused by the dilution of atmospheric CO_2 from the combustion of (^{14}C -free) fossil fuels [Suess, 1953]. Tans, et al. [Tans, De Jong, & Mook, 1979] have shown a $\sim 20\text{‰}$ depletion in tree-ring $\Delta^{14}\text{C}$ from northern Europe between 1900 and 1950. The ocean's uptake of this atmospheric fossil-fuel CO_2 , has similarly caused a $\Delta^{14}\text{C}$ depletion in the ocean surface waters as recorded by corals [Druffel and Suess, 1983]. The magnitude of this depletion in the ocean over the same pre-bomb period averages $\sim 10\text{‰}$ or about half that of the atmosphere. However, as shown by Druffel and Suess [1983] the size of this ocean Suess effect varies widely from place to place owing to differing oceanographic processes, such as the rate of deep water renewal to the local surface waters and differences in regional air-sea gas exchange rates. For instance, the Suess effect derived from a Bermuda coral is $\sim 25\text{‰}$ [Nozaki, et al., 1978] or almost the same as Tans, et al.'s [1979] tree-ring value, while a Florida coral shows $\sim 11\text{‰}$ [Druffel and Linick, 1978] and a Galapagos coral measured only $\sim 6\text{‰}$ [Druffel, 1981]. Most of the ocean $\Delta^{14}\text{C}$ depletion exhibited in these coral records has

occurred since about 1930, so that if just the period from about 1940 to 1955 is considered, in order to be comparable with the shorter shell-derived records, most of the Suess effect should still be detectable in the northern North Atlantic records (shown in Table 2). For this shorter period, the magnitude of the Suess effect for Bermuda and Florida is approximately 18‰ and 11‰, respectively. The depletion for Georges Bank is ~13‰, or intermediate between that of Bermuda and Florida. In contrast, the Norway and North Sea records exhibit depletions greater than ~30‰, though this calculation is based on the very low values occurring at both sites in the mid-1950s. In contrast, the Iceland record spans almost the entire anthropogenic period, though with low-temporal resolution for the pre-bomb period, but surprisingly shows no significant depletion and perhaps even enrichment in the latter part of this signal. Corroborating this result, pre-bomb whole shell data from the southwest coast of Iceland [*Broecker and Olson*, 1961] do not show a Suess effect depletion either and actually indicate an 8‰ enrichment between the 1840s and the 1940s. Perhaps, convective and deep-water formation processes in the subpolar regions or episodic shifts between Arctic and Atlantic water masses on Iceland's north coast [*Stefansson and Olafsson*, 1991] result in ocean $\Delta^{14}\text{C}$ changes that overwhelm the Suess effect in this region. Taken together, these pre-bomb results display a great deal of variability concerning the size of the Suess effect in the North Atlantic, and suggest that reservoir corrections [*Mangerud and Gulliksen*, 1975][*Bard*, 1988] and regional fossil-fuel CO_2 uptake calculations based on a modelled 0-10‰ Suess effect for the surface North Atlantic possess a large uncertainty.

1950s Extrema: Norway and North Sea:

The very depleted minima ($< -80\text{‰}$) in the North Sea and Norway in the mid-1950s are difficult to explain, because there is no other evidence of a surface source this low in the eastern North Atlantic. Since, they represent only two data points and the North Sea value has a large precision error ($\pm 13\text{‰}$), it is possible that they are simply the result of analytical

error. However, these two points also roughly coincide with the rather low minimum in the Georges Bank record, and the 1950s is the period of maximum Suess effect. Tree-ring records from northern Europe do not indicate "clean-air" $\Delta^{14}\text{C}$ values much lower than -30‰ by 1950 [Tans, *et al.*, 1979], and atmospheric values of at least $\sim -80\text{‰}$ would be required to account for these extreme ocean minima. The Norway site is adjacent to a small city and within a fjord with steep mountains to the east, and perhaps it is possible that nearby fossil-fuel CO_2 emissions resulted in locally depleted atmospheric conditions. Meijer, *et al.* [in press] have recently shown a seasonal $\Delta^{14}\text{C}$ depletion of 40‰ for atmospheric CO_2 during the period 1974-1993 caused by local fossil-fuel burning at a sampling station near Groningen, Netherlands. Cain and Suess [1976] also showed that trees in New York City had values close to -90‰ in the 1940s. Further, two studies have shown that the highly industrialized Rhine Valley region of northern Germany displays atmospheric $\Delta^{14}\text{C}$ depletions greater than 100‰ relative to "clean air" during winter-time due to increased pollution and lower atmosphere inversions [Levin, Munnich, & Weiss, 1980][Segl, *et al.*, 1983]. However, this local pollution explanation seems unreasonable as a cause for the North Sea anomaly since that site is about 100 km offshore, and should not be subject to such a large local anthropogenic effect.

Unusually strong upwelling of intermediate waters along the subpolar front, perhaps in response to increased deepwater production, might have provided a source for such depleted values. Evidence of deep convective mixing in the central eastern North Atlantic to depths of ~ 1000 m has been reported and invoked to explain the ventilation of the intermediate depth waters of the eastern subtropical gyre region [Arhan, Colin de Verdiere, & Memery, 1994]. In the northwest Atlantic, Tanaka, *et al.* [1990] discussed the possibility of episodic upwelling of AAIW near the Labrador Sea and southwards along the shelf to explain unusually low pre-bomb $\Delta^{14}\text{C}$ values in Long Island Sound in the mid-1950s. Also, Dickson, *et al.* [1975, 1988] have reported significant climate reversals in the

northern North Atlantic, which could provide the mechanisms for unusual upwelling events and frontal shifts of subpolar and arctic water masses. The $\Delta^{14}\text{C}$ minima at Georges Bank, North Sea, and Norway differ in time from each other by the approximate water mass travel times between these locales estimated from the mean residual velocity of the subpolar gyre circulation ~ 1000 km/yr [Dickson, *et al.*, 1988]. If these anomalous observations are not merely coincidental and are unrelated to the Suess effect, it might suggest that unusual upwelling occurred in the northern North Atlantic in the early 1950s forming very depleted surface water which migrated about the greater subpolar gyre in the 1950s. Though, the Iceland record does not show a similar anomaly, its only 1950s data are a decadal average for 1942-1951 and the year 1958, and so it may simply be undetected. A higher resolution $\Delta^{14}\text{C}$ time-series for the 1950s from these same locations might help to resolve this issue.

North Atlantic Post-Bomb $\Delta^{14}\text{C}$

Time History Comparison:

The first certain appearance of bomb- ^{14}C (1959 ± 1 year) as well as the period of steepest ^{14}C signal increase (1960-1966) are nearly synchronous in all six time-histories. These results reflect the hemispheric homogeneity of the atmospheric bomb- ^{14}C signal, which reached a maximum in 1964-1965 [Levin, *et al.*, 1985] (Figure 5). However, large variations in signal amplitude exist among the different sites. The North Sea signal has the largest amplitude $\sim 300\text{‰}$. The Florida, Bermuda, and Norway signal amplitudes are intermediate $\sim 200\text{‰}$, while the Georges Bank and Iceland signals have the smallest amplitudes $\sim 110\text{--}150\text{‰}$. Phase differences exist regarding the timing of the signal maxima. The North Sea signal maximum occurs earliest about 1966-1968, only 1-3 years after the atmospheric signal peak, while Bermuda's broad peak is the most delayed, occurring in the mid- to late-1970s. The rest of the records display a greater phase similarity with their maxima occurring in the early 1970s. The decay of the bomb-signal, shown in Table 2 as a percentage reduction of the total bomb-signal amplitude by 1990, also varies among sites.

The North Sea record shows the largest signal decay, whereas the Iceland record exhibits only a slight decline over two decades.

In general, these observations are consistent with the broad-scale oceanographic conditions, of which perhaps the most important factor is the winter mixed-layer depth. To illustrate this, Figure 5 shows the effect of depth changes on an idealized mixed-layer ocean in response (solid lines) to the atmospheric bomb- ^{14}C signal (dashed line). The essential observation from this exercise is that shallower mixed layers equilibrate with the atmosphere more rapidly. Alternatively, increased gas exchange rates have the same proportional effect as decreased mixed-layer depth. As gas exchange rates increase northward over the North Atlantic as a function of increasing mean wind-speed [Toggweiler, *et al.*, 1989a] they should tend to cancel the effect of increased mixed-layer depth. However, in the North Atlantic gas-transfer rates vary by a factor of two (Figure 6), while the winter mixed-layer depth varies by a factor of 5 or more (Figure 1c), thus making mixed-layer depth the more critical parameter. Since subtropical regions (Bermuda and Florida) have shallower mixed-layers relative to subpolar regions (Georges Bank, Iceland), their $\Delta^{14}\text{C}$ signal amplitudes should tend to be greater. The North Sea and Norway records do not fit this generalization since both have rather large amplitudes and yet are near regions of deep mixing. However, this discrepancy serves as an indication of the shelf modification of these ocean $\Delta^{14}\text{C}$ signals [Weidman and Jones, 1993].

Shallower mixed-layers also should produce an earlier phase (maximum) and a more rapid signal decay (Figure 5). However, four of the profiles reach their peak at about the same time in the early 1970s, and a clear relation between mixed-layer depth and phase is not apparent in the data. This is partly because the broad peaks of the bomb signals, defined by a decade long period (from the late 1960s to the late 1970s) when $\Delta^{14}\text{C}$ fluctuated within $\sim 10\%$ of the maxima, make it difficult to determine precisely the phase of

these $\Delta^{14}\text{C}$ signals. Regional oceanographic processes, operating on interannual and decadal scales and resulting even in minor $\Delta^{14}\text{C}$ fluctuations could produce apparent phase shifts of several years. The North Sea record with its very early peak follows a shallow pattern, while Bermuda with its very late peak follows a deep pattern, which is opposite to what might be expected based on the mixed-layer depths of their source regions. The North Sea signal's early phase is again consistent with a strong shelf influence. Druffel [1989] explained Bermuda's delayed phase relative to Florida's as the result of deeper winter mixing in the Sargasso Sea. However, winter mixing at Bermuda is still much shallower than that taking place in any of the subpolar regions and yet Bermuda's peak occurs later, suggesting that the mixed-layer depth is not the sole factor in determining the phase of the bomb-signal. Druffel and Suess [1983], in discussing the Galapagos coral record, noted that the supply of intermediate or deep water to surface waters should cause a phase shift in the surface $\Delta^{14}\text{C}$ signal towards that of the atmospheric $\Delta^{14}\text{C}$ signal, as the depleted component (relatively unaffected by bomb input) would tend to drive down the concentration as soon as the atmospheric peak input passed. The advanced phase of the northern North Atlantic signals relative to Bermuda is also consistent with the other evidence of a deep depleted source for the northern surface waters. Signal decay is difficult to assess for the subtropical sites as both records end in 1983, but Druffel [1989] has explained the more rapid dropoff of the Florida signal relative to the Bermuda signal as evidence of the presence of upwelled waters in the Gulf Stream precursor. As seen in Table 2, the Georges Bank and Iceland signals decay at about half the rate (~20% drop relative to bomb-signal amplitude) as the North Sea and Norwegian signals (~35% drop relative to bomb-signal amplitude). This contrast could be the result of greater mixed-layer depths in the western subpolar source regions and/or greater shelf influence in the two eastern sites.

Comparison with Other Post-bomb $\Delta^{14}\text{C}$ Measurements:

Figure 7 shows GEOSECS [Stuiver and Ostlund, 1980] and TTO [Ostlund and Grall, 1987] North Atlantic surface data versus winter mixed-layer depth (using Robinson, et al. [1979]) at each GEOSECS and TTO site and Table 3 compares the regional mean GEOSECS and TTO values with the time-history data for years ~1972 and ~1981, and also shows the winter (February) mixed-layer depth for each region. These comparisons again demonstrate that the post-bomb surface $\Delta^{14}\text{C}$ in the North Atlantic during these periods can be seen largely as a function of mixed-layer depth, which is in line with Broecker and Peng's [1982] interpretation of the poleward depletion of surface $\Delta^{14}\text{C}$ in the North Atlantic. Possibly aliasing this relationship, however, is the presence of an intermediate/deep water component, whose $\Delta^{14}\text{C}$ composition would be relatively uninfluenced yet by bomb- ^{14}C and would thus dilute the bomb input in the surface waters. Table 3 also shows that the Florida, Bermuda, and Iceland data closely match their GEOSECS and TTO regional means, while the North Sea, Norway, and Georges Bank data are higher than their regional means. While, it cannot be certain that the regional means derived from GEOSECS and TTO data are representative of the ocean areas adjacent to the time-history sites, this comparison suggests that the former three records are not significantly modified by shelf influences, while the latter three records may have been altered by shelf influences.

Determining the Offshore $\Delta^{14}\text{C}$ Signal:

In chapter one [Weidman and Jones, 1993] it was shown that the Georges Bank's $\Delta^{14}\text{C}$ signal could be viewed as a shelf-modified Labrador Sea $\Delta^{14}\text{C}$ signal. The shelf effect can be summarized as the importation and isolation of ocean water on a shallow shelf, the essential effect of which is to enhance the equilibration of this water's $\Delta^{14}\text{C}$ concentration with the atmosphere relative to its ocean source. In this way the "shelf effect" results in the bomb-signal's amplification, an advancement of its phase towards that of the

atmosphere's signal. The resultant $\Delta^{14}\text{C}$ of shelf waters are thus a combination of the offshore $\Delta^{14}\text{C}$ which is presumably characteristic over a wide ocean region (such as the Labrador Sea or Norwegian Sea) and local shelf water depth and residence time. The ocean record can be estimated from an adjacent shelf record by inverting this relationship. A similar approach was used in chapter one to obtain a $\Delta^{14}\text{C}$ record for the Labrador Sea from the Georges Bank data, and is now used to reconstruct $\Delta^{14}\text{C}$ time histories for the central eastern North Atlantic and the Norwegian Sea. A simple model describes the relationship between a shelf site's $\Delta^{14}\text{C}$ signal and that of its offshore ocean source:

$$\Delta^{14}\text{C}_{\text{shelf}(+R)} = \Delta^{14}\text{C}_{\text{ocean}} + U_{\text{shelf}} * (\Delta^{14}\text{C}_{\text{atmosphere}} - \Delta^{14}\text{C}_{\text{ocean}}) \quad 1)$$

The shelf uptake rate $U_{\text{shelf}} = R * I / (h_{\text{shelf}} * \Sigma\text{CO}_2)$, where R is the shelf residence time, I is the air-sea CO_2 invasion rate, h_{shelf} is the mean shelf depth, and ΣCO_2 is total DIC (which is assumed here for simplicity to be the same for both ocean and shelf). Rewriting (1) to solve for the ocean signal we obtain:

$$\Delta^{14}\text{C}_{\text{ocean}(-R)} = (\Delta^{14}\text{C}_{\text{shelf}} - U_{\text{shelf}} * \Delta^{14}\text{C}_{\text{atmosphere}}) / (1 - U_{\text{shelf}}) \quad 2)$$

Given the shelf and atmospheric record, we can solve (2) by determining a value for U_{shelf} in two ways: 1) directly from estimates of the parameters R , I , h_{shelf} , and ΣCO_2 , and 2) indirectly by finding a U_{shelf} that fits the regional ocean GEOSECS and TTO data. The latter method can be expressed as

$$U_{\text{shelf}} = (\Delta^{14}\text{C}_{\text{shelf}} - \Delta^{14}\text{C}_{\text{GEOSECS or TTO}}) / (\Delta^{14}\text{C}_{\text{atmosphere}} - \Delta^{14}\text{C}_{\text{GEOSECS or TTO}}) \quad 3)$$

North Sea:

A determination of the uptake coefficient U_{shelf} for the North Sea region can be made directly. A residence time of 4-6 months has been calculated for the waters of English Channel origin in the German Bight region of the North Sea [Reid, *et al.*, 1988]. A quantitative estimate is not available for the residence time of waters of central eastern North Atlantic origin in the English Channel and Celtic shelf, but based on Pingree and Le Cann's [1989] study of residual currents ($\sim 1\text{-}3\text{ cm/s}$) on the Celtic and Armorican shelves an additional 8-12 month period does not seem unreasonable. This gives a total shelf residence time (from the central eastern North Atlantic to the German Bight) of 12-18 months or $R \sim 1.25\text{ year}$. A value of $20\text{ mol CO}_2\text{ m}^{-2}\text{ yr}^{-1}$ is assumed for the CO_2 invasion rate I , which is the approximate value that Broecker, *et al.* [1985] determined for the Atlantic Ocean. For h_{shelf} we assume an average value of 50 m. A value of 2 mol m^{-3} is used for ΣCO_2 . From these estimates we obtain $U_{\text{shelf}} \sim 0.25$ for the North Sea. Alternately, using the indirect method (3), the regional TTO surface value from the central eastern North Atlantic for 1981 of $\sim +109\text{‰}$, and an atmospheric value for 1981 of $\sim +240\text{‰}$ [Levin, *et al.*, 1985] we obtain an estimate of $U_{\text{shelf}} \sim 0.19$ for the North Sea.

Figure 8 shows several estimates of the $\Delta^{14}\text{C}$ signal in the North Sea using (2) the North Sea record, the atmospheric record, and several U_{shelf} values from 0.20 to 0.35. From this analysis a $\Delta^{14}\text{C}$ signal with a $U_{\text{shelf}} \sim 0.22$ is adopted for the shelf regions west of the German Bight (Table 4a). The estimated central eastern North Atlantic record peaks about 1974 with a maximum of $\sim +160\text{‰}$. The phase and amplitude of this signal are similar to that of the Bermuda signal, peaking in the mid-1970s, and showing a slow decay rate (drop of $\sim 15\%$ of total bomb-signal amplitude). The estimated pre-bomb record is dependent on our assumptions about the nature of the extreme 1950s minimum in the North Sea. Since no unusual atmospheric depletion is assumed in the model, then very depleted ocean waters for the 1950s are predicted. Alternately, if the 1950s ocean source water were

to be fixed at a low but reasonable value of -65‰ , then an atmospheric value of $\sim -140\text{‰}$ is still required to fit the North Sea minimum. While this level of depletion is similar to that found in the nearby industrialized Rhine valley, it seems too extreme to maintain over large areas of the English Channel and North Sea shelf, which makes the idea for an ocean source of the 1950s anomaly seem the more probable cause.

Norway:

The direct method cannot be used at present to determine U_{shelf} for the Norway site because an independent estimate of the shelf residence time (R) for water in Tromsøysund is not available, nor is there a quantitative estimate of relative mixing between the Norwegian Coastal Current and offshore Norwegian Current waters (which would affect our estimate of R). Using the indirect method (equation 3), the atmospheric values of $+465\text{‰}$ (1972) and $+240\text{‰}$ (1981), the Tromsøysund data for ~ 1972 and ~ 1981 (from Table 1c), and the regional GEOSECS and TTO data for the eastern subpolar region of $+54\text{‰}$ (1972) and $+63\text{‰}$ (1981), we obtain U_{shelf} values of 0.16 (1972) and 0.20 (1981).

Figure 9 shows the several estimates of the $\Delta^{14}\text{C}$ signal in the Norwegian Sea using (2) and the Tromsøysund record, the atmospheric record, and an array of U_{shelf} values from .10 to .20. From this analysis a $\Delta^{14}\text{C}$ signal with a $U_{\text{shelf}} \sim 0.15$ is adopted as the estimate for the $\Delta^{14}\text{C}$ signal of the Norwegian Sea (Table 4b). This $\Delta^{14}\text{C}$ signal has a similar, but slightly larger, amplitude than the Iceland signal, peaks in the mid-1970s, and slowly decays (drop of $\sim 16\%$ of total bomb-signal amplitude). A larger U_{shelf} value is not chosen, despite a better fit with GEOSECS and TTO data, because it produces a large local minimum in the mid-1960s bomb-curve which seems unrealistic since this is the time of peak atmospheric $\Delta^{14}\text{C}$, unless unusually deep mixing was occurring in the ocean source region. There is some evidence for this latter hypothesis. The mid-1960s was a time of unusual cooling in the northern North Atlantic [Dickson, *et al.*, 1975] and records from the

north coast of Iceland show a sharp change to a regime of colder springtime ocean temperatures ($\Delta \sim 4^{\circ}\text{C}$) and lower salinities ($\Delta \sim 0.5\text{‰}$) at this time [Stefansson and Olafsson, 1991]. Recently, Bonisch and Schlosser [1995] have shown results from a tracer and box modeling study of the Greenland/Norwegian Seas and Nansen Basin, which suggest that deep water production in this region was five times greater between the mid-1960s and 1980 than since that time. Also, Druffel [1989] has shown, from modeling results, that the upper ocean ventilation rates in the Sargasso Sea (mid-subtropical gyre) appeared to be at a 15-year maximum in the mid-1960s. Alternately, as the rate of change of atmospheric $\Delta^{14}\text{C}$ was greatest in the early to mid-1960s, our model and time-history may simply have too coarse a resolution to accurately reconstruct this sensitive part of the bomb-curve.

As in the North Sea case, the pre-bomb estimate for the offshore source region is dependent on our assumptions about the cause of the 1950s anomaly. As this site is quite shallow and within a few hundred meters of a small city, it is not unreasonable to suspect a strong anthropogenic (fossil-fuel combustion) effect. However, even if offshore waters are assumed to be as low -65‰ , a back estimate requires local atmosphere $\Delta^{14}\text{C}$ of $\sim -185\text{‰}$, which seems impossible to maintain over an area wide enough to affect the Tromsøysund surface water. This analysis adds weight to the alternative hypothesis of an offshore origin for the 1950s anomaly.

The Evolution of the bomb- ^{14}C signal in the North Atlantic:

In Figure 10, three estimated time histories for the Labrador Sea (from Chapter 1, Table 2), Norwegian Sea, and central eastern North Atlantic (derived from the Georges Bank, Norway, and North Sea records, respectively, and corrected for shelf effects) are combined with the time-histories for Iceland, Bermuda, and Florida, to present an overview of the evolution of the bomb- ^{14}C signal in the North Atlantic Ocean. A sharp

contrast in amplitudes is apparent between northern (subpolar) and southern (subtropical) regions with remarkable conformity within each region. Florida, Bermuda, and the central eastern North Atlantic have similar amplitudes of $\sim 220\text{‰}$ and maxima of $\sim +160\text{‰}$. These radiocarbon results add support to other evidence (i.e., McCartney and Talley, 1982) that suggest the surface waters in the transitional region south of the subpolar front, including those that feed the central eastern North Atlantic and the southern North Sea, generally share a more distinct connection with the subtropical gyre than with the subpolar gyre. The Bermuda and central eastern North Atlantic records also share a delayed phase relative to the Florida signal. Druffel [1989] has explained the phase difference between Bermuda and Florida as the result of deeper mixing in the Sargasso (~ 200 m) relative to the Florida Current (~ 100 m). However, this explanation still contains a discrepancy since both Florida and Bermuda have similar amplitudes which should also vary with mixed-layer depth. One solution to this problem is that the dilution of the atmospheric input resulting from a deeper mixed-layer in the Sargasso Sea is balanced by horizontal advection from converging surface waters.

The Labrador Sea, Iceland Sea, and Norwegian Sea have amplitudes of $\sim 120\text{‰}$ and maxima $\sim +60\text{‰}$. All reached their maxima in the mid-1970s and have decayed only by about 15% of their total bomb-signal amplitudes over 15 years (about 1/6 the atmospheric bomb-signal decay rate). These results are consistent with the deep winter mixed-layers (500-1000 m) of these northern areas that dilute the bomb- ^{14}C inventory and slow the ocean's equilibration with the atmosphere. The results, also, are consistent with a $\Delta^{14}\text{C}$ -depleted intermediate or deepwater contribution to the subpolar regions. The slow signal decay rates of the northern North Atlantic time histories might be partly owed to the sharply reduced convection and deep water production since ~ 1980 indicated by a number of multi-tracer studies (e.g., [Schlosser, *et al.*, 1991][Bonisch and Schlosser, 1995]). Also, because surface waters in these subpolar regions are, presently, more depleted than the

atmosphere, these regions should still be increasing their inventory of atmospheric bomb- ^{14}C or maintaining a balance with regional loss through deep water production. These results support Toggweiler, et al.'s [1989b] modeling results which predicted increasing inventories for northern North Atlantic and decreasing inventories for the subtropical North Atlantic during the 1980s. The estimated Iceland Sea and Labrador Sea bomb-signals are very similar, though the Labrador Seas pre-bomb data are $\sim 20\%$ lower, indicating a deepwater component (such as AAIW) in the Labrador Sea surface waters absent in the surface waters of the Iceland Sea. A small east-west difference is shown in the relative enrichment of the Norwegian Sea signal versus the Labrador Sea and Iceland Sea signals. This difference is in agreement with deeper mixing in the western section of the subpolar gyre and Nordic Seas, and the poleward advection of surface Atlantic water into the Norwegian Sea.

Summary

Three shell-derived $\Delta^{14}\text{C}$ time-histories from the eastern North Atlantic were combined with three previously published $\Delta^{14}\text{C}$ records from the western North Atlantic to investigate the temporal and spatial evolution of the $\Delta^{14}\text{C}$ in the surface North Atlantic Ocean. Pre-bomb conditions are more enriched in the subtropical regions than the subpolar regions, and most depleted in the Labrador Sea and most enriched in the Sargasso Sea. This contrast can be explained by 1) deeper convective mixing in the higher-latitude regions, and 2) a depleted source for the subpolar gyre waters of intermediate (AAIW) southern ocean origin, or possibly of Arctic origin via the Davis Strait. The high-latitude time-histories generally show pre-bomb depletions that are consistent with the Suess effect recorded by subtropical North Atlantic corals, but the magnitude of this effect varies greatly, and is much larger on the European shelf than in the western North Atlantic. Very low pre-bomb values ($< -80\%$) in the North Sea and Norway in the 1950s, cannot be

explained easily by a Suess effect depletion of the overlying atmosphere, and most likely are the result of unusually strong convection / upwelling in the northern North Atlantic during this time.

Initial appearance of the bomb- ^{14}C is synchronous in all North Atlantic $\Delta^{14}\text{C}$ records about 1959 (± 1 year). All North Atlantic $\Delta^{14}\text{C}$ records show a steep rise during the early to mid-1960s and reach maxima in the late 1960s to late 1970s. The North Sea record has the largest amplitude ($\sim 320\text{‰}$) and the Iceland record has the smallest ($\sim 120\text{‰}$). The bomb- ^{14}C signal amplitude and maxima are seen largely as a function of local and regional mixed-layer depth. The southern North Sea record's extremely large amplitude and advanced phase indicates enhanced equilibration owing to local shelf influence. By comparing regional GEOSECS and TTO surface $\Delta^{14}\text{C}$ data to the time history records, the degree of shelf influence can be recognized, corrected for, and used to derive estimated regional surface ocean time histories. The Georges Bank, Norway, and North Sea records have been corrected in this way to produce the first regional $\Delta^{14}\text{C}$ time-series for the Labrador Sea (done in chapter one), Norwegian Sea, and the central eastern North Atlantic. The "corrected" bomb- ^{14}C signals show a sharp contrast between northern (subpolar gyre) and southern (subtropical gyre) regions with remarkable conformity within each of these regions. Despite its high-latitude position, the southern North Sea is clearly more linked with subtropical gyre regions. Florida, Bermuda, and the central eastern North Atlantic have amplitudes and maxima that are double those of the Labrador Sea, Iceland Sea, and Norwegian Sea. These results are consistent with the deeper convection of the high-latitude regions as well as the likely contribution of an intermediate source to the subpolar gyre.

Despite the global climatic and oceanographic importance of high-latitude ocean regions, such as the northern North Atlantic, these areas still remain relatively devoid of radiocarbon data. This study has begun the process of closing this gap by taking advantage

of *A. islandica*'s longevity and wide geographic distribution to compile several radiocarbon time-histories throughout the northern North Atlantic Ocean. The results of this study demonstrate the utility of this new paleoceanographic tool and suggest that its further application combined with denser spatial coverage and higher-temporal resolution can be used to yield a more detailed history of North Atlantic circulation for the past century.

Acknowledgements

Financial support for this work was provided by NOAA grant # NA36GPO291, NSF grant # OCE9301015, Oak Ridge Associated Universities Global Climate Change Fellowship Program, and the WHOI Ocean Ventures Fund. All new radiocarbon analyses for this study were done at the National Ocean Sciences AMS Facility at WHOI. For critical help in collecting specimens, I would like to thank: Steve Murawski and Frank Almeida at the NOAA Northeast Center of National Marine Fisheries Service at Woods Hole for the North Sea specimens; Rindy Ostermann at WHOI, and Jon Olafsson and Hrafnkell Eiriksson, both at the Marine Research Institute in Reykjavik, Iceland for the Iceland specimens; and Jan Sundet of the Norwegian Institute of Fisheries and Aquaculture, Tromsø, Norway for the Norway specimens. I would also like to thank Bill Schmitz, Lynne Talley, and Mike McCartney for helping me to understand North Atlantic Ocean circulation.

References

- Arhan, M., A. Colin De Verdiere and L. Memery, The eastern boundary of the subtropical North Atlantic, *J. Phys. Ocean.*, 24, 1295-1316, 1994.
- Bard, E., Correction of accelerator mass spectrometry C14 ages measured in planktonic foraminifera: Paleoceanographic implications, *Paleoceanogr.*, 3, 635-645, 1988.
- Bard, E., M. Arnold, J. Mangerud, M. Paterne, L. Labeyrie, J. Duprat, M. A. Melieres, E. Sonstegaard and J. C. Duplessy, The North Atlantic atmosphere-sea surface ^{14}C gradient during the Younger Dryas climatic event, *Earth and Planet. Sci. Let.*, 126, 275-287, 1994.
- Bonisch, G., and P. Schlosser, Deep water formation and exchange rates in the Greenland/Norwegian Seas and the Eurasian Basin of the Arctic Ocean derived from trace balances, *Prog. Oceanog.*, 35, 29-52, 1995.
- Broecker, W. S., A revised estimate for the radiocarbon age of North Atlantic Deep Water, *J. Geophys. Res.*, 84, 3218-3226, 1979.
- Broecker, W. S., The great ocean conveyor, *Oceanogr.*, 4, 79-89, 1991.
- Broecker, W. S., R. Gerard, M. Ewing and B. C. Heezen, Natural radiocarbon in the Atlantic Ocean, *J. Geophys. Res.*, 65, 2903-2931, 1960.
- Broecker, W. S., and E. A. Olson, Lamont radiocarbon measurements VIII, *Radiocarbon*, 3, 176-204, 1961.
- Broecker, W. S., and T. H. Peng, *Tracers in the Sea*, Lamont-Doherty Geological Observatory Publication, Columbia University, Palisades, N.Y., 1982.
- Broecker, W. S., and T. S. Peng, Interhemispheric transport of carbon dioxide by ocean circulation, *Nature*, 356, 587-589, 1992.
- Broecker, W. S., and T. Takahashi, Hydrography of the central Atlantic-IV. Intermediate waters of Antarctic origin, *Deep-Sea Res.*, 28A, 177-193, 1981.
- Chapman, D. C., and R. C. Beardsley, On the origin of shelf water in the Middle Atlantic Bight, *J. Phys. Ocean.*, 19, 384-391, 1989.
- Dickson, R. P., and J. Brown, The production of North Atlantic Deep Water: sources, rates and pathways, *J. Geophys. Res.*, 99, 12,319-12,341, 1994.
- Dickson, R. R., H. H. Lamb, S.-A. Malmberg and J. M. Colebrook, Climatic reversal in northern North Atlantic, *Nature*, 256, 479-482, 1975.
- Dickson, R. R., J. Meincke, S. Malmberg and A. J. Lee, The "Great Salinity Anomaly" in the Northern North Atlantic 1968-1982, *Prog. Oceanog.*, 20, 103-151, 1988.

- Druffel, E. M., Decade time scale variability of ventilation in the North Atlantic: High-precision measurements of bomb radiocarbon in banded corals, *J. Geophys. Res.*, **94**, 3271-3285, 1989.
- Druffel, E. M., Radiocarbon in annual coral rings from the eastern tropical Pacific, *Geophys. Res. Lett.*, **8**, 59-62, 1981.
- Druffel, E. M., and T. W. Linick, Radiocarbon in annual coral rings, *Geophys. Res. Lett.*, **5**, 913-916, 1978.
- Druffel, E. M., and H. E. Suess, On the radiocarbon record in banded corals: exchange parameters and net transport of $^{14}\text{CO}_2$ between atmosphere and surface ocean, *J. Geophys. Res.*, **88**, 1271-1280, 1983.
- Ellett, D. J., The north-east Atlantic: A fan-assisted storage heater?, *Weather*, **48**, 118-126, 1993.
- Fonselius, S., and G. Ostlund, Natural radiocarbon measurements on the surface water from the North Atlantic and Arctic Sea, *Tellus*, **11**, 77-82, 1959.
- Gade, M., Features of fjord and ocean interaction, in: *The Nordic Seas*, edited by B. G. Hurdle, pp. 183-189, Springer-Verlag, New York, 1986.
- Gagnon, A. R., and G. A. Jones, AMS-graphite target production methods at the Woods Hole Oceanographic Institution between 1986-91, *Radiocarbon*, **35**, 301-310, 1993.
- Hjort, C., A sea correction for East Greenland, *Geologiska Foreningens i Stockholm Forhandlingar*, **95**, 132-134, 1973.
- Johannessen, O. M., Brief overview of the physical oceanography, in: *The Nordic Seas*, edited by B. G. Hurdle, pp. 103-127, Springer-Verlag, New York, 1986.
- Kalish, J. M., Pre- and Post-bomb radiocarbon in fish otoliths, *Earth Planet. Sci. Lett.*, **1993**.
- Krauss, W., The North Atlantic Current, *J. Geophys. Res.*, **91**, 5061-5074, 1986.
- Keigwin, L. D., and G. A. Jones, Western North Atlantic evidence for millennial-scale changes in ocean circulation and climate, *J. Geophys. Res.*, **35**, 12-397-12,410, 1994.
- Lazier, J. R. N., The renewal of the Labrador Sea water, *Deep-Sea Res.*, **20**, 341-353, 1973.
- Levin, I., B. Kromer, H. Schoch-Fischer, M. Bruns, M. Munnich, D. Berdau, J. C. Vogel and K. O. Munnich, 25 years of Tropospheric ^{14}C Observations in Central Europe, *Radiocarbon*, **27**, 1-19, 1985.
- Levin, I., K. O. Munnich and W. Weiss, The effect of anthropogenic CO_2 and ^{14}C sources on the distribution of ^{14}C in the atmosphere, *Radiocarbon*, **22**, 379-391, 1980.
- Linick, T. W., P. E. Damon, D. J. Donahue and A. J. T. Jull, Accelerator mass spectrometry: the new revolution in radiocarbon dating, *Quat. Internat.*, **1**, 1-6, 1989.

- Maier-Reimer, E., and R. Bacastow, Modelling of geochemical tracers in the ocean, in: *Climate-Ocean Interaction*, edited by M. E. Schlesinger, pp. 233-267, Kluwer Academic Publishers, Netherlands, 1990.
- Mangerud, J., Radiocarbon dating of marine shells, including a discussion of apparent age of Recent shells from Norway, *Boreas*, 10, 143-172, 1972.
- Mangerud, J., and S. Gulliksen, Apparent radiocarbon ages of Recent marine shells from Norway, Spitzbergen, and Arctic Canada, *Quatern. Res.*, 5, 263-273, 1975.
- McCartney, M. S., and L. D. Talley, The subpolar Mode Water of the North Atlantic Ocean, *J. Phys. Oceanogr.*, 12, 1169-1188, 1982.
- McCartney, M. S., and L. D. Talley, Warm-to-cold conversion in the northern North Atlantic Ocean, *J. Phys. Oceanogr.*, 14, 922-935, 1984.
- Meijer, H. A. J., J. V. D. Plicht, J. S. Gislefoss and R. Nydal, Long-time atmospheric ^{14}C and ^3H record near Groningen (Netherlands) and comparison with the Fruholmen (Norway) and Izana (Canary Islands) ^{14}C stations, *Radiocarbon*, in press.
- Normann, U., Havnmiljødata fra nord-norske fjorder 1991, Fiskeriforskning-Norges fiskerihøgskole, Tromsø, 1992.
- Nozaki, Y., D. M. Rye, K. K. Turekian and R. E. Dodge, C-13 and C-14 variations in a Bermuda coral, *Geophys. Res. Lett.*, 5, No.10, 825-828, 1978.
- Ostlund, H. G., and C. Grall, TTO: Tritium Laboratory Data Report, 16, University of Miami, 1987.
- Ostlund, H. G., G. Possnert and J. H. Swift, Ventilation rate of the deep Arctic Ocean from Carbon 14 data, *J. Geophys. Res.*, 92, 3769-3777, 1987.
- Pingree, R. D., Flow of surface waters to the west of the British Isles and in the Bay of Biscay, *Deep-Sea Res.*, 40, 369-388, 1993.
- Pingree, R. D., and B. Le Cann, Celtic and Armorican slope and shelf residual currents, *Prog. Oceanogr.*, 23, 303-338, 1989.
- Quay, P. D., M. Stuiver and W. S. Broecker, Upwelling rates for the equatorial Pacific Ocean derived from bomb ^{14}C distribution, *J. Mar. Res.*, 41, 769-792, 1983.
- Reid, P. C., A. H. Taylor and J. A. Stephens, The hydrography and hydrographic balances of the North Sea, in: *Pollution of the North Sea*, edited by W. Salomons, B. L. Bayne, E. K. Duursma and U. Forstner, pp. 3-19, Springer-Verlag, Heidelberg, Germany, 1988.
- Revelle, R., and H. E. Suess, Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO_2 during the past decades, *Tellus*, 9, 18-27, 1957.
- Rey, F., The development of the spring phytoplankton outburst at selected sites off the Norwegian coast, in: *The Norwegian Coastal Current*, edited by R. Sætre and M. Mork, pp. 649-680, The University of Bergen, Bergen, 1981.

- Robinson, M. K., R. A. Bauer and E. H. Schroeder, Atlas of North Atlantic - Indian Ocean Monthly Mean Temperatures and Mean Salinities of the Surface Ocean, Naval Oceanographic Office Reference Publication 18, Naval Oceanographic Office, 1979.
- Schlosser, P., G. Bonisch, B. Kromer, H. H. Loosi, R. Buhler, R. Bayer, G. Bonani and K. P. Koltermann, Mid-1980s distribution of tritium, ^3He , ^{14}C and ^{39}Ar in the Greenland/Norwegian Seas and the Nansen Basin of the Arctic Ocean, *Prog. Oceanog.*, 35, 1-28, 1995.
- Schlosser, P., G. Bonisch, M. Rhein and R. Bayer, Reduction of deep water formation in the Greenland Sea during the 1980s: evidence from tracer data, *Science*, 251, 1054-1056, 1991.
- Schmitz, W. J., and M. S. McCartney, On the North Atlantic circulation, *Rev. Geophys.*, 31, 29-49, 1993.
- Segl, M., I. Levin, H. Schoch-Fischer, M. Munnich, B. Kromer, J. Tschiersch and K. O. Munnich, Anthropogenic ^{14}C variations, *Radiocarbon*, 25, 583-592, 1983.
- Stefansson, U., *North Icelandic Waters*, Leiftur Ltd, Reykjavik, 269 p., 1962.
- Stefansson, U., and J. Olafsson, Nutrients and fertility of Icelandic waters, *J. Mar. Res. Inst. Reykjavik*, 12, 1-56, 1991.
- Stocker, T. F., and W. S. Broecker, Observation and Modeling of the North Atlantic Deep Water Formation and its Variability: Introduction, *J. Geophys. Res.*, 99, 12,317, 1994.
- Stommel, H., A survey of ocean current theory, *Deep-Sea Res.*, 4, 149-184, 1957.
- Stuiver, M., and H. G. Ostlund, GEOSECS Atlantic radiocarbon, *Radiocarbon*, 22, 1-24, 1980.
- Stuiver, M., and H. A. Polach, Discussion: Reporting of ^{14}C Data, *Radiocarbon*, 19, 355-363, 1977.
- Suess, H. E., Natural radiocarbon and the rate of exchange of CO_2 between the atmosphere and the sea, Proc. of the Conf. on Nuclear Processes in Geological Settings, University of Chicago Press, Chicago, Ill., 52-56, 1953.
- Sverdrup, H. V., M. W. Johnson and R. H. Fleming, *The Oceans*, Prentice-Hall, 1087p., 1942.
- Tanaka, N., M. C. Monaghan and K. K. Turekian, C^{14} balance for the Gulf of Maine, Long Island Sound and the northern Middle Atlantic Bight: evidence for the extent of the Antarctic Intermediate Water contribution, *J. Mar. Res.*, 48, 75-87, 1990.
- Tans, P. P., A. F. M. De Jong and W. G. Mook, Natural atmospheric ^{14}C variation and the Suess effect, *Nature*, 280, 826-828, 1979.
- Toggweiler, J. R., K. Dixon and W. S. Broecker, The Peru upwelling and the ventilation of the South Pacific thermocline, *J. Geophys. Res.*, 96, 20,467-20,497, 1991.

- Toggweiler, J. R., K. Dixon and K. Bryan, Simulations of radiocarbon in a coarse-resolution world ocean model, 1, steady state prebomb distributions, *J. Geophys. Res.*, 94, 8217-8242, 1989a.
- Toggweiler, J. R., K. Dixon and K. Bryan, Simulations of radiocarbon in a coarse-resolution world ocean model, 2, distributions of bomb-produced carbon 14, *J. Geophys. Res.*, 94, 8243-8264, 1989b.
- Tsuchiya, M., Circulation of the Antarctic Intermediate Water in the North Atlantic Ocean, *J. Mar. Res.*, 47, 747-755, 1989.
- Warren, B. A., *Evolution of Physical Oceanography*, MIT Press, Cambridge, Mass., 1981.
- Weidman, C. W., and G. A. Jones, A shell-derived C14 Time History on Georges Bank and its Labrador Sea Implications, *J. Geophys. Res.*, 98, 14,577-14588, 1993.
- Worthington, L. V., The Norwegian Sea as a mediterranean basin, *Deep-Sea Res.*, 17, 77-84, 1970.

Table 1. North Atlantic $\Delta^{14}\text{C}$ Time Histories derived from *Arctica islandica* shells.

A. North Sea-German Bight (54°N, 6°E) ~37 m depth

Year	$\Delta^{14}\text{C}$	$\pm\epsilon$	$\delta^{13}\text{C}^1$	Shell ID	Accession ID
1948*	-50.0	3.9	2.0	N-8A	OS-0093
1950*	-64.1	12.2	2.0	N-8A	OS-0094
1954*	-83.3	13.2	2.0	N-8A	rec #1018
1959*	-24.2	3.9	2.0	N-8A	OS-0090
1960	+9.5	7.8	2.0	N-8A II	OS-1351
1961-1963*	+26.2	4.3	2.0	N-8A	OS-0089
1963-1964	+116.6	9.1	2.0	N-8A II	OS-1349
1966	+234.2	8.5	2.0	N-1B	OS-0130
1967-1968	+239.3	13.3	2.0	N-8A II	OS-1350
1969	+222.8	7.6	2.0	N-1B	OS-0128
1969-1971*	+210.7	4.1	2.0	N-8A	OS-0091
1970	+228.4	5.0	2.0	N-1B	OS-0642
1971-1972	+216.7	3.8	2.0	N-8A II	OS-1424
1974	+220.4	6.2	2.0	N-1B	OS-0099
1974-1975*	+216.0	10.5	2.0	N-8A	OS-0088
1976	+186.7	3.7	2.0	N-8A II	OS-1400
1980-1981	+173.0	5.4	2.0	N-1B	OS-0097
1979-1983*	+133.7	13.5	2.0	N-8A	OS-0087
1982-1987	+137.2	6.1	2.0	N-1B	OS-0096
1989-1990	+126.4	7.0	2.0	N-1B	OS-0131

* these dates have been lagged by 7 years from those originally logged because of an error in logging these samples. Samples from N-8A II (adjacent slice from same shell) were later run to corroborate this 7-year shift.

1. $\delta^{13}\text{C}$ assumed from mean of the Iceland and Norway shells

B. Iceland-Siglufjodur (66°N, 19°W) 22m depth

Year	$\Delta^{14}\text{C}$	$\pm\epsilon$	$\delta^{13}\text{C}$	Shell ID	Accession ID
1872-1876	-55.0	3.5	2.6	I-1-1A	OS-2667
1892-1896	-53.8	3.3	2.5	I-1-1A	OS-2666
1912-1921	-59.5	3.5	1.9	I-1-1A	OS-2665
1942-1951	-54.8	3.8	1.8	I-1-1A	OS-2664
1958	-44.4	3.6	1.9	I-1-13B	OS-2736
1960	-22.3	5.3	1.8	I-1-13B	OS-2733
1964	+22.9	4.9	2.6	I-1-13B	OS-2731
1968-1969	+41.1	4.3	2.3	I-1-13B	OS-2734
1970-1971	+56.9	7.3	2.0	I-1-13B	OS-2730
1972-1973	+44.2	4.4	1.8	I-1-13B	OS-2735
1974-1975	+47.8	4.8	1.3	I-1-6B	OS-2672
1976	+40.8	4.1	1.7	I-1-6B	OS-2671
1974-1979	+53.3	4.8	1.7	I-1-13B	OS-2732
1979	+51.9	5.5	1.9	I-1-6B	OS-2669
1982-1983	+34.9	6.1	1.9	I-1-6B	OS-2673
1987	+43.7	4.4	1.8	I-1-6B	OS-2670
1991	+35.7	4.2	1.6	I-1-6B	OS-2668

Table 1. North Atlantic $\Delta^{14}\text{C}$ Time Histories derived from *Arctica islandica* shells.

C. Norway, Tromsø (70°N, 19°E) 3m depth

Year	$\Delta^{14}\text{C}$	$\pm\epsilon$	$\delta^{13}\text{C}$	Shell ID	Accession ID
1940	-44.5	3.0	2.5	T-2A	OS-3019
1945	-62.2	5.8	2.4	T-2A	OS-3018
1951-1952	-54.7	3.1	2.5	T-2A	OS-3017
1956-1957	-82.3	5.0	2.5	T-2A	OS-3016
1959-1960	-34.2	3.5	1.8	T-2A	OS-3015
1962-1963	+8.0	3.2	1.9	T-2A	OS-2865
1965-1966	+61.9	10.7	2.1	T-2A	rec #-6163
1969-1970	+132.2	4.3	1.8	T-2A	OS-2864
1973-1974	+105.3	3.8	2.1	T-2A	OS-2863
1976-1977	+122.3	3.3	1.8	T-2A	OS-2852
1979-1980	+100.0	5.4	1.7	T-2A	OS-2851
1982-1984	+97.0	4.0	1.7	T-2A	OS-3527
1988-1990	+62.4	4.0	1.5 ²	T-2A	OS-2738
1992-1993	+64.6	4.5	1.3	T-2A	OS-2737

2. $\delta^{13}\text{C}$ linearly interpolated from adjacent values

D. Georges Bank (41°N, 67°W) 76m depth

Year	$\Delta^{14}\text{C}$	$\pm\epsilon$	$\delta^{13}\text{C}^3$	Shell ID	Accession ID
1939	-61.3	6.2	2.3	GB61-7-1B	AA 6922
1942	-64.4	6.1	2.3	GB61-7-1B	AA 6923
1945-1946	-70.6	6.2	2.3	GB61-7-1B	AA 6924
1948-1949	-65.3	6.3	2.3	GB61-7-1B	AA 6925
1951-1952	-74.6	6.2	2.3	GB61-7-1B	AA 6927
1954-1955	-66.0	6.2	2.3	GB61-7-1B	AA 6928
1959-1960	-34.4	6.4	2.3	GB61-7-1B	AA 6929
1963-1965	+32.3	6.7	2.3	GB61-7-1B	AA 6930
1968-1970	+74.8	7.4	2.3	GB61-7-1B	AA 6932
1973-1975	+77.0	7.6	2.3	GB61-7-1B	AA 6933
1978-1980	+47.7	7.5	2.3	GB61-7-1B	AA 6934
1983-1985	+51.1	7.7	2.3	GB61-7-1B	AA 6935
1988-1990	+42.8	7.4	2.3	GB61-7-1B	AA 6936

3. $\delta^{13}\text{C}$ assumed from mean of 214 analysis from specimen from Nantucket Shoals

Table 2: North Atlantic ^{14}C Time History Summary

Location	(Lat., Long)	Pre-Bomb (<1958 / ~1950)	Pre-Bomb Minimum (Yr)	Post-Bomb Maximum (Yr)	Amplitude	Suess Effect ~1940+	Decay (% amplitude)
Florida	(25°N, 83°W)	-59 / -65	-66 (1952)	+156 (1970)	222	-11‰	--
Bermuda	(32°N, 64°W)	-47 / -49	-55 (1952)	+159 (1977)	214	-18‰	--
Georges Bank	(41°N, 67°W)	-75 / -70	-75 (1951-52)	+77 (1973-75)	152	-13‰	22
North Sea	(54°N, 6°E)	-66 / -57	-83 (1954)	+239 (1967-68)	322	-33‰	35
Iceland	(66°N, 19°W)	-56 / -55	-60 (1912-21)	+57 (1970-71)	117	~ 0‰	18
Norway	(70°N, 19°E)	-61 / -55	-82 (1956-57)	+132 (1969-70)	214	-36‰	32

Table 3: North Atlantic ^{14}C Time History and Ocean Data Summary

Location	~1972 (GEOSECS)	~1981(TTO)	TC Depth
Sub-Trop	139‰	131‰	208
Florida	155‰	128‰	
Bermuda	154‰	148‰	
Transitional	----	109‰	303
North Sea	218‰	153‰	
W. Sub-Pol	54‰	46‰	701
Georges Bank	76‰	49‰	
Iceland	44‰	43‰	
E. Sub-Pol	54‰	63‰	563
Norway	119‰	99‰	

Table 4a: Estimated $\Delta 14\text{C}$ Time History for the Central Eastern North Atlantic

Date	Atmosphere * $\Delta 14\text{C}$	Estimated Ocean $\Delta 14\text{C}$
1947	-30	-56
1949	-30	-74
1953	-30	-98
1958	+130	-68
1959	+227	-52
1960-1962	+266	-41
1962-1963	+537	-2
1965	+755	+87
1966-1967	+658	+121
1968	+564	+127
1968-1970	+543	+117
1969	+544	+139
1970-1971	+510	+134
1973	+429	+162
1973-1974	+415	+160
1975	+361	+138
1979-1980	+263	+129
1978-1982	+256	+109
1981-1986	+199	+120
1988-1989	+148	+120

Table 4b: Estimated $\Delta 14\text{C}$ Time History for the Norwegian Sea

Date	Atmosphere * $\Delta 14\text{C}$	Estimated Ocean $\Delta 14\text{C}$
1939	-30	-47
1944	-30	-68
1950-1951	-30	-59
1955-1956	+30	-102
1958-1959	+179	-72
1961-1962	+292	-42
1964-1965	+795	-67
1968-1969	+554	+58
1972-1973	+447	+45
1975-1976	+346	+83
1978-1979	+287	+67
1981-1983	+220	+75
1987-1989	+153	+46
1991-1992	+114	+56

* Atmospheric record is shown in Figure 3 and is a compilation of pre-bomb data from Tans, et al. [1979], and post-bomb data from Levin, et al. [1985] and Meijer, et al. [in press].

Figure Captions

Figure 1a. General surface circulation patterns of the surface North Atlantic Ocean (after Sverdrup, et al., 1942) and radiocarbon time history locations (Δ).

Figure 1b. Mean February water temperatures at 150 m depth (after Robinson, et al., 1979) and radiocarbon time history locations (Δ). These isotherm patterns vary very little throughout the year.

Figure 1c. Mean February depth to thermocline ($\sim$ annual maximum mixed-layer depth defined here by the depth where temperature is 1.1°C less than that of the surface) (after Robinson, et al., 1979) and radiocarbon time history locations (Δ).

Figure 2(a-f). Radiocarbon time histories for six locations in the North Atlantic Ocean. Eastern North Atlantic shell-derived records (new results): a) North Sea; b) Iceland; c) Norway. Western North Atlantic shell- and coral-derived records (previously published results): d) Bermuda (Druffel, 1989); e) Florida (Druffel, 1989); f) Georges Bank (Weidman and Jones, 1993).

Figure 3. Antarctic Intermediate Water Distribution in the North Atlantic: a) Silica ($\mu\text{mol/l}$) on the isopycnal whose depth is shown in b) Depth ($\times 100$ m) of AAIW core as identified by Tsushiya [1989] (from Tsushiya, 1989). Note that this surface outcrops in the Labrador and Irminger Seas. Comparing (b) with Figure 1c note that this core is accessible to convection almost everywhere north of the subpolar front.

Figure 4. a) Steady state OGCM $\Delta^{14}\text{C}$ results from a) Maier-Reimer and Bacastow, 1990, and b) Toggweiler, et al., 1989. The Maier-Reimer and Bacastow results reproduce a more realistic surface northern North Atlantic than Toggweiler's results owing to what appears to be the greater northward penetration of the former's AAIW analog (centered at ~ 1000 m at the equator).

Figure 5. Atmospheric bomb- ^{14}C signal (dashed line) modelled ocean bomb- ^{14}C signals (solid lines) in a simple mixed-layer ocean as a function of layer depth (or gas exchange rate). Shallower depths (or increased gas exchange rates) result in a reduced signal amplitude and earlier phase shift.

Figure 6. Distribution of wind speed dependent gas exchange rates ($\text{mol/m}^2/\text{yr}$) (from Toggweiler, et al., 1989a).

Figure 7. North Atlantic surface $\Delta^{14}\text{C}$ (GEOSECS (stars) and TTO (circles) data) versus February ($\sim$ annual maximum) mean depth to thermocline at those same locations. There is a strong correlation (GEOSECS: dashed line; TTO: solid line) between surface $\Delta^{14}\text{C}$ and winter mixed-layer depth.

Figure 8. Estimated central eastern North Atlantic $\Delta^{14}\text{C}$ signals (dashed lines) based on the North Sea ^{14}C time history (solid line) and back-corrected for shelf influence using various uptake coefficients from 0.20 to 0.35. The $\Delta^{14}\text{C}$ time history estimates are compared to the regional mean for the central eastern North Atlantic (solid circle) for ~ 1981 (TTO).

Figure 9. Estimated Norwegian Sea $\Delta^{14}\text{C}$ signals (dashed lines) based on the Tromsøysund $\Delta^{14}\text{C}$ time history (solid line) and back-corrected for shelf influence using various uptake coefficients from 0.10 to 0.20. The $\Delta^{14}\text{C}$ time history estimates are compared to the regional means for the Norwegian Sea (solid circles) for ~1972 (GEOSECS) and ~1981 (TTO).

Figure 10. Combined North Atlantic Ocean $\Delta^{14}\text{C}$ Time Histories (corrected for shelf effects). A strong signal contrast exists between subpolar and subtropical gyre influenced regions and close agreement among the signals within these regions. The subtropical gyre region signals have a much larger amplitude than the subpolar region signals.

Figure 1a

Figure 1c

Figure 2

Distribution of AAIW in the North Atlantic (from Tsushiya, 1989)

Figure 3

OGCM results (steady state) $\Delta^{14}\text{C}$ Distribution
(from Maier-Reimer and Bacastow, 1990)

Figure 4a

OGCM results (steady state) $\Delta^{14}\text{C}$ Distribution (from Toggweiler, et al., 1989)

Figure 4b

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Chapter 5

A Century-Long Record of Bottom Temperatures Near Nantucket Shoals from Stable Oxygen Isotopic Analysis of *Arctica islandica* Shells

Introduction

The method for deriving past ocean temperatures from the oxygen isotope composition of biogenic carbonates is a commonly used paleoceanographic technique with a four decade-long history [Urey, Lowenstam, Epstein, & McKinney, 1951][Epstein, Buchsbaum, Lowenstam, & Urey, 1953][Duplessy, 1978][Grossman, & Ku, 1986][Wefer, & Berger, 1991]. In particular, oxygen isotope studies of the shells of bivalved molluscs have been used to obtain information on shell growth banding and paleoenvironmental conditions [Rhoads, & Lutz, 1980][Arthur, Williams, & Jones, 1983][Jones, Williams, Arthur, & Krantz, 1984][Romanek, Jones, Williams, Krantz, & Radke, 1987][Krantz, Williams, & Jones, 1987]. However, because most bivalves have relatively short lives [Heller, 1990], isotope records generated from them have been limited to a decade or so in length, and the few long-lived species have slow shell growth rates, which has precluded the application of conventional sampling methods. The recent development of micro-sampling techniques [Dettman, & Lohmann, 1993] and low-sample weight mass spectrometry have advanced the potential of the field of stable isotope sclerochronology, by enabling the precise sampling of very small seasonal growth increments in long-lived bivalves. The specific methods as applied to the bivalve species *Arctica islandica* (longevity ~+200 years) were validated in an earlier study [Weidman, Jones, & Lohmann, 1994] which compared a 12-year oxygen isotope record in a single shell with bottom T and S data from the nearby Nantucket Lightship.

In this paper, a 109-year record (1875-1983) of bottom temperatures near the former location of the Nantucket Lightship has been compiled using stable oxygen isotopic analysis on the annual bands of shells of the bivalve mollusc *Arctica islandica*, more commonly known in this region as the ocean quahog. This record is the longest seasonal mollusc-derived paleotemperature record compiled to date and provides the first century-long seasonal record of bottom temperatures (from any source) on the North Atlantic continental shelves. Banded corals have been used previously to reconstruct ocean climate records for the warm sub-tropical regions (e.g., [Druffel, 1985][Aharon, 1991]), but only now, with this study, has reconstruction begun of century-length climate records for the colder high-latitude oceans, regions thought to be intimately linked with the regulation of global climate and ocean circulation. The ability to reconstruct these longer high-resolution records also provides a means of investigating historical climate events and monitoring synoptic oceanographic processes. To demonstrate this, the shell-derived climate record is compared to other historical air and sea surface data, enabling us to explore the relationship between atmosphere / sea surface temperatures and outer shelf bottom temperatures, and the temporal variability of this relationship from the intra-seasonal to the century-scale.

Methods

Location:

The century-long temperature record is derived from the oxygen isotope records of four *A. islandica* shells. All four specimens were live-captured from the same dredge haul during the 1992 NMFS Clam Survey (Leg II) in June, 1992 from station # 315 at 40° 28' N, 69° 29' W and 65 m depth (Figure 1). This location is approximately 9 km south of the former position of the Nantucket Lightship (40° 33' N, 69° 28' W and 59 m water depth). From lightship records [Weidman, *et al.*, 1994] and hydrographic survey data [Flagg,

1987], it is known that the water column near the capture site locale is isothermic from about November through March, and stratified during the warmer months of the year (see Chapter 2, Figure 2). For the period 1956-1957 and 1960-1971, when full water column temperature (T) and salinity (S) measurements were regularly performed at the lightship, bottom temperatures ranged from winter minima averaging 3.6°C ($\pm 0.9^{\circ}$) to autumn maxima averaging 12.9°C ($\pm 1.6^{\circ}$). Salinity averaged 32.72‰ ($\pm 0.37\text{‰}$) (see Chapter 2, Figure 3). The site is centered within a shelf water mass referred to as the "cold pool", which is a seasonally persistent subsurface band of relatively cold water situated between the 50 m and 80 m isobaths on the continental shelf, and extending from the southern flank of Georges Bank to the southern reaches of the Mid-Atlantic Bight [Houghton, Schlitz, Beardsley, Butman, & Chamberlin, 1982] (Figure 2).

Sampling and Stable Isotope Analysis:

The micro-sampling methods used to obtain carbonate samples from the annual growth bands of *A. islandica* shells have been detailed in earlier studies (Chapters 2 and 3). In Chapter 2 it was shown that shell growth did not occur during the coldest months of the year, and that the growth period corresponded to about an eight-month (May-December) growing season. For this reason, a target minimum resolution of eight samples per annual growth band was chosen so that each sample would represent approximately one month of shell growth (assuming a constant seasonal growth rate). Sampling generally began within the 3rd annual band, as the earlier (outer layer) shell material was usually too eroded to sample. The ontogenic decrease in seasonal shell growth currently limits the sampling to the inner (earliest) 20 to 50 annual bands of each shell. A longer record was thus reconstructed by overlapping the individual isotope records (with about a decade of overlap on average) from specimens of different ages.

The aragonite samples weighed $\sim 50 \mu\text{g}$ on average (ranging from 10 to $200 \mu\text{g}$), and were roasted (along with aragonite standards) in vacuo for 1 hour at 200°C to volatilize any organic matter. A VG Prism mass spectrometer with an on-line VG Isocarb extraction system was used for the analyses, which were performed at the Woods Hole Oceanographic Institution. The overall analytical precision reported for this instrument for oxygen isotopes, based on the daily analysis of four or more powdered National Institute of Standards and Technology carbonate standards, is $\sim \pm 0.1\text{‰}$ (1σ).

Chronology:

The specimen ages were determined by counting the internal bands in thin-sections made from radial slices of shell. In Table 1 are shown the specimen ages ($\pm$ uncertainty in years), number of bands sampled in each shell, and the estimated chronology of the sampled bands. Specimen ages range from 48 to 119 years-old. The banding in the two youngest shells was clear and unambiguous, but the two older shells had some growth disturbance features in their shells resulting in some uncertainty in their age estimates. Later, the shell chronologies were further checked by correlating the overlap section isotope signals and adjusting them, within the band count uncertainty range, only if it significantly improved the correlation (details discussed in next section). The specimen ages and chronologies shown in Table 1 are the adjusted and final ones.

Results

Data:

A total of 1579 samples were successfully analyzed from 139 bands obtained from the four shells, with sampling resolution ranging from 5 to 33 samples per growth band (~ 11.5 samples/band). For quantitative convenience, each growth year was defined as ending on the seasonal minimum $\delta^{18}\text{O}$ value and beginning on the sample following the

previous year's minimum $\delta^{18}\text{O}$ value. This method was synchronous with the visually identified growth bands, as in almost all years the seasonal $\delta^{18}\text{O}$ minimum occurred on or adjacent to the annual "ring" position (the "rings" being defined as the thin translucent layers which form the boundaries of each wider, opaque, annual growth increment). The data were then normalized to "monthly" values so that each year band was represented by eight values (our target minimum sampling resolution). Finally, as a simplification the "monthly" data are statistically arranged in three forms: 1) annual maximum (Omax); 2) annual minimum (Omin); and 3) annual mean (Omean).

Table 2 provides a statistical summary of each shell record and Figure 3 shows all the monthly data along with linear fits of their Omean, Omax, and Omin (slopes are summarized in Table 2). The mean ($\pm$ standard deviation) of Omean, Omax, and Omin are $+1.86\text{‰}$ ($\pm 0.18\text{‰}$), $+2.31\text{‰}$ ($\pm 0.21\text{‰}$), and $+1.32\text{‰}$ (0.25‰), respectively. The mean seasonal amplitude (Omax - Omin) is 0.99‰ . The Omin data show a positive (enriching) trend with age in all four shells. The trends of the Omean and Omax data vary consistently in sign with each other, but unlike the Omin data, do not trend in a consistent way in all shells. In each shell record, the seasonal amplitude (Omax - Omin) gets smaller with age, with the enrichment trend of Omin contributing the greatest part of this signal reduction. In each shell record, the Omin always has the largest standard deviation relative to Omax and Omean, although the difference among all three parameters is not large indicating a similar amount of interannual variation.

Overlap Comparison and Chronology Check:

The fidelity of the shell-derived $\delta^{18}\text{O}$ signals and relative chronologies was tested by comparing each of the three overlapped sections and performing standard correlation and lag tests. Figure 4 and Table 3 display the results of this comparison and correlation. The first (3B / 12B, $n = 5$) and third (28A / 4B, $n = 11$) overlaps had high and significant

correlations ($\alpha \sim .05$), without any adjustment of their relative (band-count) chronologies. Lag tests resulted in reduced correlation in all cases. The second (12B / 28A, $n = 14$) overlap showed a very poor correlation with the original (band-count) chronology (Omin: $r = -0.16$, Omean: $r = -0.256$, Omax: $r = +0.199$). However, a shift of +1 year (reducing 12B's estimated age by 1 year) significantly improved the fit, though the correlation is less significant ($\alpha \sim .10$) than that of the other two overlap sections. The lag test for the second overlap also identified two other equally significant fits, requiring shifts of -4 and -5 years (increasing the estimated age of 12B), but they were rejected since these shifts were beyond the uncertainty (± 1 year) range estimated for the shell. Also, since the first overlap (3B / 12B) showed a high correlation, their relative chronologies were assumed to be "locked", and so reduced the estimated age of specimen 3B by 1 year as well. As the chronologies of 3B and 12B are considered "locked", their combined uncertainty was estimated to be ± 1 year (the same as the uncertainty of 12B). These slight adjustments result in the "final" chronologies presented in the tables and figures.

Band Width/ Age Corrections:

In Figure 4 and Table 3, it appears that the greater the age difference between the overlapping shells, the larger the positive offset is between their Omin data. An alternate view is that the smaller the mean band widths of the older shells of the overlap pairs, the larger the offset. These alternate perspectives are obviously interrelated and may be explained as an age and/or sampling-precision effect. In the first case, it may be that there is an ontogenic shortening of the seasonal growth period or reduced autumn shell growth rates as a result, perhaps, of metabolic changes or increased reproductive activity, so that the warmest ambient temperatures are less represented in the shell's seasonal growth with age. This would account for a reduced seasonal $\delta^{18}\text{O}$ amplitude with age. This effect has been suggested for bivalves in general [Harrington, 1989] and for *Spisula solidissima* in particular [Williams, Arthur, Jones, and Healy-Williams, 1982].

In the second case, as the band widths decrease with shell age, there would be an increased tendency to mechanically smear the seasonal signal as the precision limits of the sampling process are approached. This mixing effect would be negligible when the band widths are large, but cause a small progressive reduction in signal amplitude as the band widths narrowed with shell age. This effect would be most pronounced in the Omin data, as the $\delta^{18}\text{O}$ minima (warmest temperatures of autumn) occur in the shell just prior to the $\delta^{18}\text{O}$ maxima (cold temperatures of early winter). The shells' $\delta^{18}\text{O}$ maxima would be less affected since the direction of sampling is chronological, and the bottom waters warm much more slowly in the spring than they cool in the autumn. Also, the Omean data would be little affected. Indeed, the offset differences for Omean and Omax are either small or not consistent with age/band width difference.

The age / band width effect can be corrected by calibrating the Omin overlap offsets with their age and band width differences. A linear fit between age differences and Omin offsets results in the age correction: Age Correction ($\Delta \delta^{18}\text{O}$) = $0.011 \text{ y}^{-1} + 0.143$. Since the 0-intercept is about 13 years, this relationship implies that an age correction need not be applied to data from bands which are less than 13 years-old. Interestingly, this is also the approximate mean age of sexual maturity for *A. islandica* [Mann, 1982].

If a band width effect is assumed instead, a similar bandwidth correction can also similarly derived: Band Width Correction ($\Delta \delta^{18}\text{O}$) = $-0.558 (\text{bandwidth (mm)}) + 0.204$. In this case, the 0-intercept is about 0.4 mm, and suggests that the correction should be applied to data from bands with widths smaller than this value. A more rigorous approach to correcting the sampling-precision effect can be done by using a mixing model, where each sample is treated as a combination of itself and a small amount of its neighbor. The

overlap offsets can be used to estimate the size of this mixing term, and the "true" sample values can be "unmixed" by inverting the model,

$$1) \delta^{18}\text{O}_{\text{unmixed}} = \delta^{18}\text{O}_{\text{mixed } 1} - [(MIX) * (\delta^{18}\text{O}_{\text{unmixed } 2} - \delta^{18}\text{O}_{\text{unmixed } 1})] / (RES + MIX)$$

where RES is the sample resolution (mm), MIX is the mixing term (mm), and unmixed 1 and 2 are adjacent samples (and their $\delta^{18}\text{O}$ difference is approximately the same as that between mixed samples). Using this last approach, we calculated a MIX $\sim 25 \mu\text{m}$ and used it to correct the Omin data for this sampling-precision effect (average correction $\sim 0.15\text{‰}$). No correction was applied to the Omean or Omax data.

T / $\delta^{18}\text{O}_{\text{aragonite}}$ and T / S / $\delta^{18}\text{O}_{\text{water}}$ Relationships:

As in chapters 2 and 3, a modified form of Grossman and Ku's [Grossman, *et al.*, 1986] aragonite- $\delta^{18}\text{O}$ equilibrium equation (for mollusc data only) is used to convert the shells' $\delta^{18}\text{O}_{\text{aragonite}}$ records to bottom temperature (T_b),

$$2) \quad T_b = 20.86 - 4.69 (\delta^{18}\text{O}_{\text{arag}} (\text{PDB}) - \delta^{18}\text{O}_{\text{water}} (\text{SMOW}))$$

The equation has been modified from the original to include the approximate difference ($\Delta \sim 0.2\text{‰}$) between CO_2 equilibrated with standard mean ocean water (SMOW) and CO_2 generated from the hydrolized carbonate standard Pee Dee belemnite (PDB). The standard notation for oxygen isotopic composition is used, where $\delta^{18}\text{O}_{\text{sample}} (\text{PDB}) = [^{18}\text{O}/^{16}\text{O}_{\text{sample}} / ^{18}\text{O}/^{16}\text{O}_{\text{standard}} (\text{PDB}) - 1] \times 1000$.

In equation 2, the term $\delta^{18}\text{O}_{\text{water}}$ is the isotopic composition of the ambient water, which can be estimated using Fairbanks' [Fairbanks, 1982] salinity- $\delta^{18}\text{O}_{\text{water}}$ relationship for the Gulf of Maine,

$$3) \quad \delta^{18}\text{O}_{\text{water}} (\text{SMOW}) = 0.421 (S) - 14.66$$

Rather than use a single mean value for our estimate of salinity (S), we use a variable one derived from a polynomial (2nd order) fit to T and S data for the period 1956-1971 (equation 4) when bottom data were recorded at the lightship (Figure 5). A quadratic relationship was chosen over a linear one because it captures the double maximum in the seasonal salinity cycle, the largest one at the autumn overturn when bottom temperatures are at a maximum and a smaller one in the late winter / early spring when bottom temperatures are coldest. This T / S function allows salinity (and thus $\delta^{18}\text{O}_{\text{water}}$) to be treated as a function of temperature,

$$4) \quad S = 0.018 T^2 - 0.277 T + 33.625 \quad (\text{SEE} = 0.322)$$

Substituting (4) into (3) to obtain

$$5) \quad \delta^{18}\text{O}_{\text{water}} (\text{SMOW}) = 0.008 T^2 - 0.117 T - 0.504$$

Substituting (5) into (2) to obtain

$$6) \quad T_b = 21.98 - [133.3 (\delta^{18}\text{O}_{\text{aragonite}}) - 43.1]^{0.5}$$

Note that (6) works only for $T_b < -22^\circ \text{C}$, which is well above recorded maximum bottom temperatures for this deep water area of the shelf. This relationship also suggests that as ambient waters in this region warm, $\delta^{18}\text{O}_{\text{aragonite}}$ will approach a minimum value $\sim +0.3\text{‰}$, because the ambient $\delta^{18}\text{O}_{\text{water}}$ would be getting heavier at the opposite rate from the equilibrium fractionation between aragonite and water. Hydrographically, this also implies that as bottom temperatures become $> 17^\circ \text{C}$ their S and $\delta^{18}\text{O}_{\text{water}}$ properties approach

those of slope water ($S > 34\text{‰}$ and $\delta^{18}\text{O}_{\text{water}} > +0.0\text{‰}$) [Fairbanks, 1982]. This result is supported by hydrographic data [Flagg, 1987] which shows that the nearest source of very warm bottom water is upper slope water, which can intrude landward along the outer shelf bottom during late summer and autumn.

The Combined Shell-Derived Temperature Records:

Equation (6) was used to convert the shell O_{mean} , O_{max} , and O_{min} data to bottom temperatures. The individual shell records were combined to form three 109-year annual temperature records from 1875 to 1983 (Table 4 and Figure 6). The overlap data are averaged to obtain a single temperature value for each year. We estimate no uncertainty in the chronology from 1924 to 1983 and a ± 1 year uncertainty for the period from 1875 to 1923. The shell-derived bottom temperatures, like the shell $\delta^{18}\text{O}$ data, are described by the three parameters: 1) annual mean (BTmean); 2) annual "monthly" maximum (BTmax); and an 3) annual "monthly" minimum (BTmin). Ten-year running means are also calculated and plotted in Figure 6 for each of the three records.

In chapter 2 [Weidman, *et al.*, 1994], it was argued that *A. islandica* shells' $\delta^{18}\text{O}$ composition can be used to estimate "monthly" bottom temperatures of ambient waters with an accuracy of $\pm 1.2^\circ\text{C}$, but since shell growth ceases during the winter at temperatures below $\sim 6^\circ\text{C}$, the coldest temperatures cannot be determined. In addition, in chapter 3, it was shown that shell growth seems to occur between ambient temperatures of 5° to 15°C , but not much above or below these cutoffs. As bottom temperatures at the Nantucket Lightship appear to rarely exceed 15°C , this upper limit should not usually constrain the shell's recording of annual maximum temperatures (BTmax). In contrast, BTmin and to some extent BTmean should tend to be biased towards warmer values than the corresponding statistics for ambient waters because of the cold end cutoff, which is $1\text{--}2^\circ\text{C}$ warmer than average minimum bottom temperatures in this region (see Chapter 2, Figure

3). However, the shell growth shutdown/turn on may not be entirely temperature driven, as suggested by the greater variability of the annual minimum temperature ($5.7^{\circ}\text{C} \pm 0.9^{\circ}\text{C}$) in this study's much larger data set than exhibited in the 12-year record ($6.0^{\circ}\text{C} \pm 0.6^{\circ}\text{C}$) discussed in chapter two (using equation 6 in this study to calculate both). The seasonal shell growth shutdown/turn on may be indirectly related to temperature through food or light availability and thus coupled to a particular time of year. For example, Figure 7 (from O'Reilly, Evans-Zetlin, and Busch, 1987] shows a time-series of seasonal primary production for the stratified region on the southern flank of Georges Bank. Note the large increase which occurs in May (spring phytoplankton bloom) and the sharp decline after November, so that the period of peak food availability (for a filter-feeder) is roughly coincident with *A. islandica*'s seasonal growth cycle.

BTmean:

BTmean values are not likely to be true annual means for the reasons discussed above, but they should provide a measure of the mean temperature during the shell growth season (~May-December). However, the mean of BTmean ($7.64^{\circ}\text{C} \pm 0.90^{\circ}\text{C}$) is very close to the Nantucket Lightship annual mean bottom temperature of ($7.58^{\circ}\text{C} \pm 0.50^{\circ}\text{C}$) for the period 1960-1971, but shows more variability. The coldest year of the mean shell record is 1977 (5.30°C) and the warmest is 1898 (9.88°C). Linear regression of the data produces a slope of $-0.010^{\circ}\text{C y}^{-1}$ ($\pm 0.003^{\circ}\text{C}$), indicating that mean annual bottom temperatures in the "cold pool" region south of Nantucket Shoals have cooled $\sim 1.1^{\circ}\text{C}$ over the past century. The mid-1960s and mid-1970s are periods of sustained cold bottom temperatures. Sharp coolings ($\sim -2^{\circ}\text{C y}^{-1}$) occur after some of BTmean's warmest periods near the years 1880, 1905, and 1920. The 10-year running mean exhibits two peak warm periods centered about 1900 and 1919, with a cold period between them, and cold periods in the 1880s, and after 1960 until the end of the record in the late 1970s. The 10-year running mean shows a flat trend from 1925 to 1960.

BTmax:

The BTmax record represents the annual maximum "monthly" bottom temperatures, which generally occur in this region at the autumn overturn in late October and November. The mean of BTmax ($11.47^{\circ}\text{C} \pm 1.69^{\circ}\text{C}$) is about 1°C lower than the Nantucket Lightship mean annual monthly maximum bottom temperature of ($12.59^{\circ}\text{C} \pm 1.37^{\circ}\text{C}$) for the period 1960-1971, and exhibits similar variability. The coldest year of the maximum record is 1976 (8.14°C) and the warmest is 1878 (15.60°C). Linear regression of the data produces a slope of $-0.027^{\circ}\text{C y}^{-1}$ ($\pm 0.005^{\circ}\text{C}$), indicating that the annual maximum "monthly" bottom temperatures at this site have cooled a surprising $\sim -2.9^{\circ}\text{C}$ over the past century. In particular, the mid-1960s and mid-1970s are the longest sustained cold periods of the century, with maximum bottom temperatures dropping about 4°C between warmer peaks in the early 1960s, late 1960s-early 1970s, and late 1970s. In the 1880s, there is a similar decade long reduction of about 4°C between peak warm annual maximum temperatures in the early 1880s and early 1890s, but in an absolute sense this "cold" period is warmer by about 1°C than a century later. Similar to the BTmean data, the 10-year running mean rises from the cold decade of the 1880s to two peak warm periods centered about 1900 and 1915 with a colder period between them. There follows a more or less continuous cooling until the end of the record in the late 1970s, with the most rapid reduction in maximum bottom temperatures occurring after 1945.

BTmin:

As discussed above, the BTmin record cannot represent the actual annual minimum "monthly" bottom temperature as the shells do not grow during the coldest months (when $T < \sim 5^{\circ}\text{--}6^{\circ}\text{C}$), and it is still unclear whether a shutdown/turn on temperature or time is the more important factor controlling the seasonal cessation and commencement of shell deposition. However, this record should at least reflect the rate of springtime warming if a shutdown temperature is operating, with years having more rapid rates of warming

reflected in warmer minima. If seasonal shell growth is more of a time-controlled phenomenon, then the interannual variations will reflect real differences in temperature at a particular time in the spring, and need not be constrained to temperatures greater than $\sim 5^{\circ}\text{--}6^{\circ}\text{C}$. The mean of BTmin ($5.65^{\circ}\text{C} \pm 0.90^{\circ}\text{C}$) is only slightly colder than the mean bottom temperature ($5.90^{\circ}\text{C} \pm 0.82^{\circ}\text{C}$) for the month of May at the Nantucket Lightship for the period 1960-1971, and exhibits the same variability. The coldest year of the minimum record is 1881 (3.48°C) and the warmest year is 1898 (7.34°C). Linear regression of the data produces a slope of $-0.004^{\circ}\text{C y}^{-1}$ ($\pm 0.003^{\circ}\text{C}$), indicating that the annual minimum ($\sim$ May) "monthly" bottom temperatures at this site have cooled only $\sim 0.4^{\circ}\text{C}$ over the past century. The 1880s and 1960s stand out as the most persistent coldest periods of the past century. The 10-year running mean shows a 1°C warming from the early 1880s to 1900 which is a characteristic of all three shell-derived bottom temperature records. The 10-year running mean shows almost no trend for the next 60 years from 1900 to about 1960, and then drops off sharply by $\sim 1^{\circ}\text{C}$ for the rest of the record.

Discussion

Nantucket Lightship Bottom Temperature Records:

A comparison of the shell-derived bottom temperature records with the Nantucket Lightship bottom temperature records for the continuous period 1960-1971 shows a strong qualitative and quantitative fit (Figure 8). The lightship data are similarly divided into the annual mean, annual maximum, and the mean May bottom temperature (representing a comparable "minimum"). The period 1960-1971 is particularly distinctive because of the extreme and sustained drop in bottom temperatures which occurred in the mid-1960s. The magnitude and phase of this low-temperature anomaly recorded at the lightship are reflected in all three shell-derived records. The means of each of the shell-derived records as

compared to the lightship records for the same period are: shell / lightship mean = 7.20°C / 7.58°C ; shell / lightship maximum = 10.37°C / 12.59°C ; shell / lightship "minimum" = 5.27°C / 5.90°C . The shell-derived temperatures are always colder, though the offset is small in the mean and minimum data, but a significant $\sim 2^{\circ}\text{C}$ colder in the maximum data. Also, the mean and minimum records are much better correlated ($r = +0.653$ and $+0.710$, respectively) than the maximum records ($r = +0.265$). The weaker fit of the maximum data may indicate that the annual maximum bottom temperature is a more spatially heterogeneous phenomenon on the outer shelf, as it is synchronous with the autumn overturn, a convective (vertical) process of short duration. The lower annual maximum is consistent with the greater depth of the shell site and its more centered position within the "cold pool" (see Figure 2), and also the autumn overturn may be delayed relative to the shallower lightship position allowing for an even colder annual maximum to be experienced at the deeper site.

Nantucket Lightship Sea Surface Temperature Records:

The Nantucket Lightship sea surface temperature (SST) records are more temporally extensive (1878-1971) than the bottom temperature (BT) records, but are less satisfactory to use as a comparison data set. This is true largely because the Nantucket lightships were relocated a number of times over the period from 1878 to 1971. In addition, the records are not continuous, having gaps as long as 12 years. However, SST records from the same 1960-1971 period when bottom temperatures were logged, offer insight into the relationship between bottom and surface temperatures at this outer shelf region. The SST data are arranged into annual means, maxima, and minima (using the May SSTs as the comparable minima) (Figure 8). Like the lightship and shell-derived bottom temperatures, the lightship SSTs show a large cooling in the mid-1960s. During this period, overall cooling at the surface has almost the same magnitude as the cooling at the bottom, but there are important differences between surface and bottom temperature

signals. For example, the surface and bottom lightship data are strongly correlated when comparing their May data ($r = +0.855$), while their annual mean and maximum data are more weakly correlated ($r = +0.533$ and $+0.338$, respectively). This result is consistent with the idea that stratification during summer isolates the bottom waters from surface heating. By the time overturn occurs in the autumn, the magnitude of the summer surface anomaly is often modified considerably. In fact, the offset between the lightship's annual maximum bottom and sea surface temperatures tends to be greatest during years when either the bottom maxima are coldest or surface maxima are warmest, indicating that the bottom and surface anomalies are often out of phase with each other. A lag test confirms this phase difference in the annual maximum signal by showing improved correlations when the bottom signal precedes the surface signal by 1 year or when the surface signal precedes the bottom signal by 2 years. These observations demonstrate that bottom and sea surface temperature anomalies can vary from each other significantly, even at the same location on the outer shelf.

Brenton Reef Lightship and Boothbay Harbor SST Records:

Two coastal SST time-series from the New England region are compared with the shell-derived bottom temperatures (Figure 9). The Boothbay Harbor, Maine ($\sim 43^{\circ} 45' \text{ N}$, $69^{\circ} 40' \text{ W}$) SST record from 1906 to 1995 [Lazzari, 1995] is the longest, most continuous SST record on the U. S. east coast, and can be compared to the later 78 years of the shell-derived record (1906-1983). The Brenton Reef Lightship ($\sim 41^{\circ} 20' \text{ N}$, $71^{\circ} 30' \text{ W}$) SST record from 1879 to 1941 [Bumpus, 1957] is one of longest and oldest continuous SST records on the southern New England coast, and can be compared with most of the earlier portion of the shell-derived record. Both records overlap a 36-year period from 1906 to ~ 1941 . For comparison, these records are arranged into annual mean, maximum, and April (comparable minimum) records, and 10-year running means are computed for each parameter. The mean April SSTs are used here instead of May SSTs as was done for the

Nantucket Lightship records, because significant warming of nearshore coastal waters occurs during May, making the April coastal SSTs more reflective of the spring offshore conditions. The Brenton Reef April and maximum records are several years shorter than the annual mean record because of missing monthly data for the late 1930s.

A basic contrast between the two coastal temperature records is that Brenton Reef has a mean about 2°C warmer than the Boothbay Harbor (10.4°C versus 8.3°C). A linear fit to the Brenton Reef records show that both the annual mean and April data have positive slopes (mean = $+0.005 \text{ y}^{-1} \pm 0.006$, April = $+0.016 \text{ y}^{-1} \pm 0.009$), while the annual maximum data have a slight negative slope (maximum = $-0.001 \text{ y}^{-1} \pm 0.007$). Similarly, a linear fit to the Boothbay Harbor records also show that the annual mean and April data have positive slopes (mean = $+0.017 \text{ y}^{-1} \pm 0.004$, April = $+0.013 \text{ y}^{-1} \pm 0.006$), while the annual maximum data have a negative slope (maximum = $-0.016 \text{ y}^{-1} \pm 0.004$). Therefore the coastal sea surface records indicate a mean warming of $\sim 0.3^\circ\text{C}$ ($\pm 0.4^\circ\text{C}$) for the period between 1879 and 1941 and $\sim 1.3^\circ\text{C}$ ($\pm 0.3^\circ\text{C}$) for the period between 1906 and 1983. The equally strong warming trends indicated for the month of April of $\sim 1^\circ\text{C}$ ($\pm 0.5^\circ\text{C}$) for each period contrast with the cooling indicated for the annual maximum temperatures, implying that the mean annual warming trend is primarily a winter/spring phenomenon and not a summer feature. The summer temperature records at Boothbay Harbor prior to 1941 may be artificially high due to an instrumentation change in that year (K. Drinkwater, pers. comm., 1995), so that the slope of the annual maximum data may be biased. Correlation of the 1906-1941 overlapped section of these two coastal SST records reflects a significant ($\alpha = .05$) coherence between the two records (annual mean: $r = +0.492$, annual maximum: $r = +0.323$, April: $r = +0.438$). However, these r coefficients also indicate a good deal of local variation, especially between the annual maximum SSTs, which demonstrates that coastal SST anomalies within the same general region can differ substantially.

The 10-year running means of the Brenton Reef records, trend flat or downward for the 40-year period from the early 1880s to the early 1920s, and afterwards rise to record warming in the mid-1930s. The 10-year running means of the Boothbay Harbor records, trend flat or downward for the 30-year period from about 1910 to the early 1940s, rise sharply to maxima in the early 1950s, and then decline steeply to minima in the mid-1960s. The Boothbay Harbor running mean also shows a rise again in the 1970s. An obvious contrast between the running means in the overlapping period is the absence of a significant warming in the mid-1930s in the Boothbay Harbor records.

A comparison of these two SST records with the shell-derived bottom temperature record, shows aspects of agreement and disagreement. Correlation between the SST and the shell BT records reveals a poor coherence between the data sets (r coefficients between Brenton Reef and shell: annual mean = -0.126, annual maximum = +0.009, and spring minimum = +0.327; and Boothbay Harbor and shell: annual mean = +0.031, annual maximum = +0.283, and spring minimum = +0.260). The overall warming ($\Delta \sim +1^\circ\text{C}$) displayed in the SST mean record is in contrast to the overall cooling ($\Delta \sim -1^\circ\text{C}$) displayed in the shell-derived BT mean record. The warming trend ($\Delta \sim +1^\circ\text{C}$) from 1880 to 1900 indicated in the shell BT records, is not apparent during the same period in the records from Brenton Reef. The warmest part of the shell BT records are centered between 1900 and 1920, while the warmest parts of the SST mean records are later during mid-century. The shell BT records do not show any significant warming trend in mid-century, neither the 1930s maximum of the Brenton Reef record nor the 1950s maximum of the Boothbay Harbor record. Common to the SST and shell BT records is a sharp cooling trend from the 1950s until the late 1960s. In contrast, the 1970s are a period of warming in the SST records, but continued cooling in the shell BT records. Another shared characteristic of both SST and shell-derived BT records is that their annual maximum data all trend downward indicating long-term cooling of seasonal maximum temperatures over the past

century. This result reaffirms the notion that whatever warming may have occurred on the northeastern U.S. continental shelf over the last century, it has not been a summer phenomenon.

Coastal Northeast U. S. Air Temperature Record:

A composite air temperature record for northeastern U.S. coastal locations for the period from 1901 to 1983 [*Boden, Kaiser, Sepanski & Stoss, 1994*](Figure 10) is compared to the coastal SST and shell-derived BT records. The air temperature record is divided into annual mean, summer (maximum), and spring (comparable minimum) values. Linear fits of these data show positive slopes for all three parameters (mean: $= +0.010^{\circ}\text{C y}^{-1} \pm 0.003^{\circ}\text{C}$; summer $+0.013^{\circ}\text{C y}^{-1} \pm 0.003^{\circ}\text{C}$; spring $= +0.008^{\circ}\text{C y}^{-1} \pm 0.004^{\circ}\text{C}$), indicating a regional mean atmospheric warming of $\sim +0.8^{\circ}\text{C}$ over the first 80 years of this century. Correlation of the air temperature records with the SST data generally show a good coherence (Brenton Reef, Boothbay Harbor: annual means: $r = +0.685, +0.581$; annual maximum: $r = +0.532, +0.029$; April: $r = +0.599, +0.541$) with the exception of the Boothbay Harbor maximum. The 10-year running means of the air temperatures are similar to those of the Boothbay Harbor SSTs, showing a warm peak in the 1950s and a cold period in the mid-1960s. Correlation of the air temperature records with the shell-derived BT data show a poor coherence (annual means: $r = +0.080$; annual maximum: $r = -0.115$; April: $r = +0.060$). From this perspective, the SSTs at the coastal sites clearly seem to reflect atmospheric forcing, but the shell-derived BTs do not, with the exception of the extreme mid-1960s cold period.

Comparison of Detrended Records:

The mid-1960s exception suggests that perhaps a large measure of the inconsistency between the surface data (air and sea surface) and shell-derived bottom data

is in the long-term trends. To investigate this hypothesis the data sets were detrended by subtracting out the 10-year running means and intercomparing their normalized residuals or deviations (residual / SD(residuals)). These deviation records display a much improved positive coherence between the shell-derived data and surface records (Figure 11). The correlation is most significantly ($\alpha \sim .01$) improved between the shell BT and Boothbay Harbor SST deviation data (mean: $r = +0.435$; maximum: $r = +0.211$; minimum: $r = +0.456$), in fact, being slightly more significant than the correlation between the Boothbay Harbor and Brenton Reef SST records. The correlation is still quite significant ($\alpha \sim .05$) with the atmospheric data (mean: $r = +0.236$; maximum: $r = +0.185$; minimum: $r = +0.245$), but less so with the Brenton Reef data ($\alpha \sim .10$) (mean: $r = +0.169$; maximum: $r = -0.030$; minimum $r = +0.429$). The annual mean and minimum (spring) data show the strongest coherence, while the annual maximum data remain only weakly correlated. These results show that much of the short-term (decade-scale) interannual variation in the bottom temperature record is in phase and proportional to the short-term SST and air temperature anomalies, and therefore suggests that the short term shell-derived bottom temperature anomaly pattern can be used to infer SST and air temperature anomalies in the region. However, this positive relationship seems to break down on longer time-scales.

The Long-Term Trend Problem:

The disparity between the long-term shell-derived bottom temperature and surface temperature (air and sea) trends poses an important question. Are these opposing temperature trends real or is the difference an artifact of our assumptions? The assumptions fall into two categories: unaccounted for 1) biological processes affecting the shells' oxygen isotope chemistry or 2) water mass changes affecting the $\delta^{18}\text{O}$ properties of the ambient waters. The overlapped shell isotope records provided a means of detecting differences between individual specimens and bias due to "vital" effects. Small offsets were observed in the seasonal minimum $\delta^{18}\text{O}$ data that appeared to be ontogenic in nature, and

would have the effect of deriving colder annual maximum temperatures by $\sim 1^{\circ}\text{C}$ in the later (more recent growth) portions of the longer shell records. However, the mean and minimum temperature data are unaffected and the maximum data have been corrected for these effects, and so a biological cause for the long-term trend difference is ruled out.

It is possible that as the result of undetected changes in shelf circulation over the last century, there has been a gradual enrichment of the ambient $\delta^{18}\text{O}_{\text{water}}$ of outer shelf waters, which is aliased in our isotope-derived record as an apparent long-term cooling trend. This effect would be difficult to sort out, because circulation changes large enough to affect the $\delta^{18}\text{O}_{\text{water}}$ properties of shelf water might just as easily affect its temperature properties as well. If an enrichment of ambient $\delta^{18}\text{O}_{\text{water}}$ has occurred, one way this could have happened is indicated by three shelf circulation studies [Fairbanks, 1982] [Chapman, Barth, Beardsley, & Fairbanks, 1986] [Chapman, & Beardsley, 1989]. These studies describe shelf water as essentially a two component mixture of fresher Scotian shelf water ($\delta^{18}\text{O}_{\text{water}} \sim -1.09\text{‰}$) and saltier Slope water ($\delta^{18}\text{O}_{\text{water}} \sim +0.75\text{‰}$), whose relative proportions can vary both spatially and temporally. Using their model, a 10% increase in the relative proportion of Slope water would cause a $+0.5\text{‰}$ salinity increase and a $+0.2\text{‰}$ $\delta^{18}\text{O}_{\text{water}}$ enrichment, or the equivalent of a 1°C cooling in temperature if unaccounted for in estimating shelf $\delta^{18}\text{O}_{\text{water}}$. However, if this mechanism is invoked as the cause of the long-term temperature trend difference between shell-derived BTs and SSTs; it simply shifts the problem, for it implies that shelf waters were a surprising $\sim 1\text{‰}$ fresher a century ago to account for the 2°C relative difference (1°C SST warming + 1°C BT cooling). This conclusion, while pointing up the uncertainty of the absolute temperature estimate from $\delta^{18}\text{O}$ data, would still motivate the need to understand how such a significant shelf circulation change occurred, and if that change was related to air and sea surface warming.

The other option is to assume that the shell-derived bottom temperature trends are real, and to look for oceanographic mechanisms to explain the contrast with coastal surface temperature trends. The simplest explanation may be that higher sea surface temperatures in the New England region lead directly or indirectly to greater stratification of adjacent deep shelf waters and increased volume and isolation of cold bottom waters on the outer shelf. This explanation has support in Houghton et al.'s [1983] study of the "cold pool" and its seasonal persistence in the Mid-Atlantic Bight. They suggest that the prime source of "cold pool" water is the large volume of very cold dense water formed in the winter over the Nantucket Shoals due to strong tidal mixing. As surface heating increases in the late spring and summer this same mechanism produces a large volume of warm, less dense water which then spreads out as a near-surface layer over the colder denser bottom waters. Further surface heating increases this stratification, which reduces vertical mixing and heat flux between the warm mixed-layer and the cold bottom waters. If surface temperatures were to rise, or seasonal warming were to begin earlier in the spring or end later in the fall, a consequence would be reduced vertical mixing, an increase in the initial volume of cold bottom water, and a persistence of cold bottom temperatures. Other indirect causes that have been associated with reduced vertical mixing and increased stratification are increased precipitation and local runoff at certain times of year (warm/wet versus cold/dry climate patterns), decreased wind stress, or changes in storm frequency and intensity patterns. In support of the latter of these, Dickson and Namias [1976] showed that decreased cyclone frequencies and intensities for the period 1948-1957 were coupled with warming along the U. S. eastern seaboard, while increased storminess for the period (1958-1969) was linked with a strong cooling trend in this region. These ideas on a cause for bottom cooling synchronous with surface warming, are purely speculative, but they demonstrate that oceanographic mechanisms, similar to those used to explain the existence of the "cold pool", may be operating to create this contrast.

Conclusions

A 109-year annual time-series (1875-1983) of bottom temperatures has been reconstructed for a location (at 65 m depth) near the former Nantucket Lightship from the oxygen isotope records of four shells of the long-lived mollusc species *Arctica islandica*. This site is centered within a seasonal hydrographic feature referred to as the "cold pool", and so these records provide a century-long thermal history of this outer continental shelf subsurface water mass. The method of establishing the chronology by annual band counting, and using the overlapped isotope records from several shells of different ages to reconstruct a longer record was validated by the strong correlations between the overlapped portions of the four shell records. The shell-derived bottom temperature record is presented as three annual records: annual mean, annual maximum, and annual minimum. The annual means represent the means of the shell growth seasons, average 7.6°C ($\pm 0.9^{\circ}\text{C}$), and indicate a century-long cooling of $\sim 1.1^{\circ}\text{C}$. The annual maxima represent bottom temperatures at about the time of the autumn overturn, average 12.6°C ($\pm 1.3^{\circ}\text{C}$), and indicate a large century-long reduction of $\sim 2.9^{\circ}\text{C}$ in the autumn bottom temperature. The annual minima represent the spring or May bottom temperatures, average 5.7°C ($\pm 0.9^{\circ}\text{C}$) and indicate a modest reduction of $\sim 0.4^{\circ}\text{C}$ in spring bottom temperatures over the last century. The shell records describe a $\sim 1^{\circ}\text{C}$ warming from a cold period in the 1880s to a record warm period centered about 1900. A cold period ~ 1905 -1910 is followed by another peak warming about 1915, after which there is general decline in temperatures for the rest of the record. A steeper decline in temperature occurs after 1960. All records indicate that the 1880s, 1960s, and 1970s were the three most prolonged, coldest periods of the past century.

Shell-derived bottom temperature records show a strong similarity with the Nantucket Lightship bottom temperature records for the period 1960-1971, reconfirming

that the shell-derived temperatures are accurately reflecting bottom temperatures on the outer shelf. The coherence between these records is strongest between the mean and minimum records and weakest between the maximum records. The latter result may indicate greater horizontal heterogeneity in bottom waters on the outer shelf at the autumn overturn. Long records of sea surface temperature from two coastal locations in New England and an air temperature record from the northeast U.S. coastal region show a strong positive correlation with each other, though there are significant local variations among these records at both short and long time-scales. Coastal air and sea surface records indicate a long-term $\sim 1^{\circ}\text{C}$ warming over the last century with peak warming occurring in mid-century, most apparent in their annual mean and spring records. The shell-derived bottom temperature records show a poor correlation with the surface temperature records. However, detrending the records of their 10-year running means, greatly improves the correlation between these records, and indicates that the short-term temperature anomaly patterns are similar, while the long-term temperature trends differ.

One primary contrast between the long-term trends is that the shell-derived records indicate a mean century-long cooling of $\sim 1^{\circ}\text{C}$, while mean air and sea surface records describe a century-long warming of equal magnitude. We have considered whether this contrast is a real temperature difference or an artifact. A biological or "vital effect" cause for this disparity has been rejected because the overlapping shell records provide a constraint on individual and ontogenic variation. It is possible that shelf water $\delta^{18}\text{O}$ has become more enriched over the past century, owing to changes in shelf water circulation. For example, a 10% increase in the amount of Slope water relative to Scotian shelf water would produce an apparent 1°C cooling if not accounted for. However, this explanation would also imply that the shelf waters have become $\sim 0.5\text{‰}$ saltier over the past century as well. While, this hypothesis may be correct, it merely shifts the problem, from one of a significant temperature change to one of a significant salinity change (or some of both) either of which

would have broad implications for shelf circulation. Finally, it seems equally possible that the shell-derived bottom temperature trends are real and of opposite sign to surface temperature trends. To explain this contrast, it is suggested that increased surface warming is directly or indirectly related to increased or prolonged stratification of shelf waters, which leads to a net reduction in vertical mixing and heat flux into the bottom waters of the "cold pool" and thus, an increase in the seasonal persistence of this continental hydrographic feature over the past century.

Acknowledgements

Financial support for this work was provided by NOAA grant # NA36GPO291 and Oak Ridge Associated Universities Global Climate Change Fellowship Program. All sampling and stable isotope analyses for this study were performed at the Woods Hole Oceanographic Institution. Specimen collections were carried out during the 1992 NMFS Clam Survey with the invaluable assistance of crew of the R/V Delaware II and the Survey Group at NOAA's Northeast Center of the National Marine Fisheries Service at Wood Hole.

References

- Arthur, M. A., D. F. Williams and D. S. Jones, Seasonal temperature-salinity changes and thermocline development in the mid-Atlantic Bight as recorded by the isotopic composition of bivalves, *Geology*, *11*, 655-659, 1983.
- Boden, T. A., D. P. Kaiser, R. J. Sepanski and F. W. Stoss, Trends '93: A compendium of data on global change, Carbon Dioxide Information Analysis Center, Environmental Sciences Division, Oak Ridge National Laboratory, ESD Publication No. 4195, 1994.
- Bumpus, D. F., Surface water temperatures along Atlantic and Gulf Coasts of the United States, 214, U.S. Fish and Wildlife Service, 1957.
- Chapman, D. C., J. A. Barth, R. C. Beardsley and R. G. Fairbanks, On the continuity of mean flow between the Scotian Shelf and the Middle Atlantic Bight, *J. Phys. Oceanog*, *16*, 758-772, 1986.
- Chapman, D. C., and R. C. Beardsley, On the origin of shelf water in the Middle Atlantic Bight, *J. Phys. Ocean.*, *19*, 384-391, 1989.
- Dettman, D. L., and K. C. Lohmann, Seasonal change in Paleogene surface water $\delta^{18}\text{O}$: fresh-water bivalves of western North America, in *Climate Changes in Continental Isotopic Records*, *Geophys. Monogr. Ser.*, vol. 78, edited by P. Swart, K. C. Lohmann, J. McKenzie, and S. Savin, pp. 153-163, AGU, Washington, D. C., 1993.
- Dickson, R. R., and J. Namias, North American influences on the circulation and climate of the North Atlantic sector, *Mon. Weather Rev.*, *104*, 1255-1265, 1976.
- Druffel, E. R. M., Detection of El Nino and decade scale variations of sea surface temperature from banded coral records: Implications for the carbon dioxide cycle, in: *The Carbon Cycle and Atmospheric CO₂: Natural Variations Archaeon to Present*, edited by E. T. Sundquist and W. S. Broecker, pp. Geophysical Monograph 32, American Geophysical Union, Washington, D.C., 1985.
- Duplessy, J. C., Isotope Studies, in: *Climate Change*, edited by J. Gribben, pp. 46-67, Cambridge University Press, New York, 1978.
- Epstein, S., R. Buchsbaum, H. A. Lowenstam and H. C. Urey, Revised carbonate-water isotopic temperature scale, *Bull. Geol. Soc. Am.*, *64*, 1315-1326, 1953.
- Fairbanks, R. G., The origin of Continental Shelf and Slope Water in the New York Bight and Gulf of Maine: evidence from $\text{H}_2\text{O}-18/\text{H}_2\text{O}-16$ ratio measurements, *J. Geophys. Res.*, *87*, 5796-5808, 1982.
- Flagg, C. N., Hydrographic structure and variability, in: *The Georges Bank Book*, edited by R. H. Backus, pp. 108-124, MIT Press, Cambridge, MA, 1987.
- Grossman, E. L., and T. Ku, Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects, *Chem. Geol.*, *59*, 59-74, 1986.
- Harrington, R. J., Aspects of growth deceleration in bivalves: Clues to understanding the seasonal O-18 and C-13 record -- A comment on Krantz et al. (1987), *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, *70*, 399-403, 1989.

- Heller, J., Longevity in molluscs, *Malacol.*, 31, 259-295, 1990.
- Houghton, R. W., R. Schlitz, R. C. Beardsley, B. Butman and J. L. Chamberlin, The Middle Atlantic Bight Cold Pool: evolution of the temperature structure during summer 1979, *J. Phys. Ocean.*, 12, 1019-1029, 1982.
- Jones, D. S., Annual cycle of shell growth and reproduction in the bivalves-*Spisula solidissima*-&-*Arctica islandica*-, PhD, Princeton University, 1980.
- Jones, D. S., D. F. Williams, M. A. Arthur and D. E. Krantz, Interpreting the Palaeoenvironmental, paleoclimatic and life history records in mollusc shells, *Geobios. Mem.*, 8, 333-339, 1984.
- Krantz, D. E., D. F. Williams and D. S. Jones, Ecological and paleoenvironmental information using stable isotope profiles from living and fossil molluscs, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 58, 249-266, 1987.
- Lazzari, M. A., Monthly and annual means of sea surface temperature, Boothbay Harbor, Maine, 1905 through 1994, Research Reference Document 95/1, Maine Department of Marine Resources, Environmental Monitoring Project at the Marine Sciences Laboratory, West Boothbay Harbor, Maine, 1995.
- Mann, R., The seasonal growth cycle of gonadal development in *Arctica islandica* from the southern New England Bight, *Fish. Bull.*, 80, 315-326, 1982.
- O'Reilly, J. E., C. Evans-Zetlin and D. A. Busch, Primary production, in: *The Georges Bank Book*, edited by Backus, pp. MIT Press, Cambridge, MA, 1987.
- Rhoads, D. C., and R. A. Lutz, Skeletal Growth of Aquatic Organisms, New York and London, Plenum Press, 1980.
- Romanek, C. S., D. S. Jones, D. F. Williams, D. E. Krantz and R. Radke, Stable isotopic investigation of physiological and environmental changes recorded in shell carbonate from the giant clam *Tridacna maxima*, *Mar. Biol.*, 94, 385-393, 1987.
- Urey, H. C., H. A. Lowenstam, S. Epstein and C. R. McKinney, Measurement of paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark, and Southeastern United States, *Bull. Geol. Soc. Am.*, 62, 399-416, 1951.
- Wefer, G., and W. H. Berger, Isotope paleontology: growth and composition of extant calcareous species, *Mar. Geol.*, 100, 207-248, 1991.
- Weidman, C. R., G. A. Jones and K. C. Lohmann, The long-lived mollusc *Arctica islandica*: A new paleoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern North Atlantic Ocean, *J. Geophys. Res.*, 99, 18,305-18,314, 1994.
- Williams, D. F., M. A. Arthur, D. S. Jones and N. Healy-Williams, Seasonality and mean annual sea surface temperatures from isotopic and sclerochronological records, *Nature*, 296, 432-434, 1982.

Table 1. Specimen Band Chronology Data

Specimen ID	Age ($\pm$ uncertainty) (years)	Bands sampled	Chronology (years)
CS92-315-3B	119 ($\pm$ 2)	45	1875 - 1919
CS92-315-12B	79 ($\pm$ 1)	23	1915 - 1937
CS92-315-28A	70 ($\pm$ 0)	33	1924 - 1956
CS92-315-4B	48 ($\pm$ 0)	38	1946 - 1983

Table 2: Shell $\delta^{18}\text{O}$ Summary

Specimen	Mean Mean (σ)	Mean Max (σ)	Mean Min (σ)	Mean Amp	Slope Mean	Slope Max	Slope Min	Slope Amp
3B	1.82 (.18)	2.30 (.25)	1.22 (.24)	1.08	-0.003	-0.007	+0.003	-0.010
12B	1.78 (.19)	2.31 (.17)	1.18 (.31)	1.13	+0.010	+0.006	+0.015	-0.009
28A	1.83 (.13)	2.23 (.16)	1.35 (.24)	0.88	-0.002	-0.004	+0.007	-0.011
4B	1.97 (.20)	2.40 (.22)	1.49 (.24)	0.90	+0.011	+0.010	+0.013	-0.003
Mean	1.86 (.18)	2.31 (.21)	1.32 (.25)	0.99	+0.004	+0.001	+0.010	-0.008

Table 3. Overlap Comparison

Overlap ID	Overlap Size (years)	Chronology (years)	Δ mean	Δ max	Δ min	r mean	r max	r min	Age Δ (years)	Band Width (mm) Young / Old
3B / 12B	5	1915 - 1919	+0.12	-0.02	+0.26	0.84	0.62	0.95	40	6.61 / 0.23
12B / 28A	14	1924 - 1937	-0.03	+0.08	-0.08	0.60	0.43	0.35	9	2.77 / 0.29
28A / 4B	11	1946 - 1956	0.00	-0.08	+0.15	0.74	0.62	0.80	23	3.61 / 0.27

Table 4. Combined Shell-Derived Bottom Temperature Record

Year	Mean	Max	Min
1875	6.46	11.53	4.06
1876	6.76	11.66	3.77
1877	7.57	14.52	4.86
1878	9.88	15.60	6.80
1879	8.78	11.81	6.41
1880	8.23	11.50	6.50
1881	6.37	10.60	3.48
1882	7.71	15.12	5.66
1883	6.94	12.16	4.06
1884	7.71	11.02	5.30
1885	7.20	10.87	4.90
1886	7.57	10.07	4.75
1887	6.67	10.61	4.59
1888	7.16	11.90	4.55
1889	8.18	11.74	6.89
1890	8.14	10.29	6.72
1891	8.04	13.96	6.72
1892	8.04	10.99	6.07
1893	7.39	13.03	5.34
1894	7.80	12.91	6.07
1895	7.66	10.99	5.10
1896	8.23	13.09	5.38
1897	9.56	14.77	6.89
1898	9.08	12.06	7.34
1899	8.09	11.97	6.07
1900	7.66	11.70	6.07
1901	8.78	14.27	6.76
1902	9.29	13.20	7.20
1903	9.45	14.31	7.30
1904	7.99	12.40	5.10
1905	6.85	10.15	4.55
1906	8.63	13.15	6.33
1907	6.94	9.74	5.26
1908	6.67	9.78	5.54
1909	6.07	13.16	4.48
1910	8.43	13.44	6.59
1911	8.18	12.22	6.67
1912	7.80	11.34	5.74
1913	7.71	13.30	6.33
1914	7.80	12.10	6.63
1915	8.52	13.50	6.31
1916	8.00	10.67	5.87
1917	8.19	15.35	5.06
1918	8.66	13.53	6.26
1919	9.54	14.43	6.93

Year	Mean	Max	Min
1920	8.23	10.11	6.12
1921	8.98	12.59	6.67
1922	6.67	9.50	5.14
1923	7.20	11.21	5.18
1924	7.69	11.25	5.96
1925	8.56	13.54	6.27
1926	7.23	10.11	4.89
1927	8.53	11.99	6.28
1928	7.45	10.62	5.65
1929	6.61	9.05	5.10
1930	8.40	14.45	5.85
1931	8.63	12.64	6.36
1932	7.58	12.87	5.75
1933	8.43	11.89	6.44
1934	7.48	10.24	5.40
1935	6.93	10.67	5.13
1936	8.29	12.74	6.08
1937	7.51	11.62	5.21
1938	7.62	12.45	5.66
1939	6.85	9.50	5.22
1940	7.57	11.86	5.22
1941	8.58	13.14	6.85
1942	8.63	11.09	7.20
1943	8.23	14.58	5.62
1944	7.80	12.39	5.34
1945	7.85	10.51	6.63
1946	8.14	12.19	5.77
1947	7.88	11.08	5.64
1948	7.16	10.80	5.73
1949	8.34	11.55	6.90
1950	6.74	9.57	5.09
1951	7.95	10.90	6.68
1952	8.09	10.68	6.13
1953	8.81	11.52	7.44
1954	7.60	9.74	6.15
1955	7.81	10.90	6.09
1956	7.95	10.74	6.31
1957	7.57	10.23	6.07
1958	6.72	11.31	4.82
1959	7.57	11.13	5.14
1960	8.88	11.85	6.20
1961	7.48	10.76	5.42
1962	7.95	12.99	5.74
1963	7.20	10.34	5.02
1964	6.37	8.83	4.71
1965	6.12	9.84	4.21
1966	6.24	8.87	4.90
1967	6.29	8.89	4.36
1968	7.07	9.48	4.55
1969	6.80	9.90	5.38

Year	Mean	Max	Min
1970	8.14	12.54	6.24
1971	7.90	10.12	6.54
1972	7.20	9.82	5.91
1973	6.54	8.49	5.87
1974	6.54	9.73	4.75
1975	6.20	10.11	4.44
1976	6.03	8.14	5.38
1977	5.30	8.75	3.81
1978	7.43	11.12	5.87
1979	6.37	8.58	4.59
1980	5.74	8.43	4.06
1981	5.99	9.53	3.92
1982	6.80	9.77	5.22
1983	7.43	11.48	4.75
Mean ($\pm 1\sigma$)	7.64 (± 0.90)	11.47 (± 1.69)	5.65 (± 0.90)
Minimum (Year)	5.30 (1977)	8.14 (1976)	3.48 (1881)
Maximum (Year)	9.88 (1878)	15.60 (1878)	7.34 (1898)

Figure Captions

Figure 1. Nantucket Shoals and Georges Bank. Specimen collection site is approximately nine kilometers south of the former position of the Nantucket Lightship.

Figure 2. Figures from Houghton, et al.'s [1982] Mid-Atlantic Bight (MAB) "Cold Pool" study showing the specimen collection site (solid triangle). This position is at approximately the same depth contour as the New England Shelf Experiment 79's (NSFE79) station N2, which is centered within the northern portion of the "Cold Pool".

Figure 3. "Monthly" $\delta^{18}\text{O}$ (PDB) record for each specimen plotted versus its estimated chronology. Linear trends are plotted for each record's Omean (annual mean), Omax (annual maximum), and Omin (annual minimum).

Figure 4. Overlap $\delta^{18}\text{O}$ record comparison. The last parts of the records for Omean, Omin, Omax of the older shells (solid line) are compared with the first parts of the records of the younger shells. These overlap pairs show significant correlation, though the Omin data in the first and third pair are offset, with the older shells' records being more enriched.

Figure 5. Bottom temperature versus bottom salinity (monthly means) at the Nantucket Lightship for the period 1956-1957 and 1960-1971. A quadratic fit to the data is used to model the double salinity maximum occurring when temperatures are coldest and warmest.

Figure 6. Combined shell-derived bottom temperature records, which include annual mean (BTmean), annual (autumn) maximum (BTmax), and annual (spring) minimum (BTmin). Running 10-year means (centered on the year) are plotted for each statistical record.

Figure 7. Seasonal primary production in the stratified region on the southern flank of Georges Bank. The dashed line is the moving average of seven adjacent samples (from O' Reilly, et al., 1987).

Figure 8. Comparison between the shell-derived bottom temperatures (solid lines) and the Nantucket Lightship bottom (dashed lines) and surface (dotted lines) temperatures.

Figure 9. Boothbay Harbor (solid lines) and Brenton Reef Lightship SST (dashed lines) records: annual mean, annual maximum, and April. 10-year running means are plotted for each record. (Boothbay data from Lazzari, 1995 and Brenton Reef Lightship data from Bumpus, 1957)

Figure 10. Regional (coastal northeast U.S. cities) air temperature records: annual mean, summer (maximum), and spring (comparable minimum). 10-year running means are plotted for each record. (data from Boden, et al., 1990)

Figure 11. Detrended sea surface temperature, air temperature, and shell-derived bottom temperature (deviation) Records. Deviation records = $[T - \text{running mean (10y)}] / \text{SD}(\text{residuals})$.

Figure 1

MAB Cold Pool Study (Figures from Houghton, R. W., et al., 1982)

Top: Transect Locations (note NSFE79)
 Middle: Monthly averaged T along NSFE79
 Bottom: Marmap BTs

Specimen collection position (▲)

Figure 2

Figure 3

— 3B
 - - - 12B

— 12B
 - - - 28A

— 28A
 - - - 4B

Figure 4

Figure 5

Figure 6

Figure 7

— SHELL BT
 - - - NTKT BT
 NTKT SST

Figure 8

----- Brenton Reef LS
mean: 1879-1941
max: 1879-1937
min: 1879-1935

———— Boothbay Harbor
mean, min, max: 1906-1983
(no 1949 data)

Figure 9

Air Temperatures for Coastal Northeast U.S.

Figure 10

Figure 11

CONCLUSIONS

The need for methods to reconstruct marine geochemical records of meaningful length with seasonal, annual, and decadal scale resolution for the cold, higher latitude oceans, has long been recognized. The recent availability of computer-aided micro-sampling equipment and micromass mass spectrometry now provides a way for long-lived, slow growth rate molluscs, such as *A. islandica*, to be used for this purpose. The primary objectives of this thesis have been: 1) the development and calibration of *A. islandica* as a new paleoceanographic tool for the mid- and high-latitudes of the North Atlantic Ocean; and 2) the demonstration that *A. islandica*-derived time histories of $\Delta^{14}\text{C}$, and $\delta^{18}\text{O}$ can be reconstructed for periods up to a century, and used to investigate the temporal and spatial variability of the northern North Atlantic's continental shelf bottom temperature, physical circulation, and uptake of atmospheric $^{14}\text{CO}_2$. In pursuit of these objectives, the first three chapters have described the developmental aspects and shown that the carbon and oxygen isotopic composition *A. islandica* shells can be used to reconstruct accurate time-series of ambient ocean radiocarbon and temperature. However, the stable carbon isotopic (^{13}C) composition of the *A. islandica* shell appears to be offset from "equilibrium" with ambient DIC and its seasonal and interannual variations seem to be largely of physiological origin, and therefore not as useful a recorder of environmental conditions. The last two chapters have focused on applications, describing 1) the evolution of the $\Delta^{14}\text{C}$ signal throughout the surface northern North Atlantic over the last 50 to 100 years, and 2) the first century-long record of continental shelf bottom temperatures for a higher-latitude (cold ocean) location.

In chapter one, a time history of $\Delta^{14}\text{C}$ on Georges Bank from 1939 to 1990 was derived from the annual growth bands of a single shell of the bivalve *A. islandica*. This was the first long-term ^{14}C time-series reconstructed for the North Atlantic outside the subtropical regions and demonstrated that long-lived molluscs can be used for high-latitude

$\Delta^{14}\text{C}$ reconstructions. The Georges Bank bomb- ^{14}C signal was shown to be in phase with, but to have a smaller amplitude than, two subtropical North Atlantic bomb- ^{14}C time histories derived from corals. The depleted pre-bomb conditions on Georges Bank imply a deepwater source, which was identified as originating in the Labrador Sea. A model based on continuity between the Labrador Sea and Georges Bank can account for the $\Delta^{14}\text{C}$ budget on Georges Bank and was also used to back calculate a $\Delta^{14}\text{C}$ time history for the Labrador Sea. This calculation prescribes a lower prebomb $\Delta^{14}\text{C}$ value for the surface Labrador Sea than previously estimated, suggesting greater penetration of the bomb signal and uptake of bomb-produced ^{14}C . These results strengthen the view that the Labrador Sea is a primary region of deepwater formation as well as a strong sink for atmospheric CO_2 . Using a mixing model to examine changes in ventilation and/or atmospheric $^{14}\text{CO}_2$ uptake in the Labrador Sea, it was suggested that there was greater vertical mixing in the mid-1960s in contrast to reduced vertical mixing in the early 1970s. These results correspond with independent observations of salinity changes in the North Atlantic and Labrador Sea, which are related to the Great Salinity Anomaly, thus indicating a possible linkage between deepwater formation and surface salinity in the northern North Atlantic.

Chapter two describes the first oxygen isotopic evidence confirming that *A. islandica* deposits its growth bands annually, and that the annual mark is deposited at or shortly after bottom temperatures reach their seasonal maximum, which occurs at the Nantucket Lightship location at the autumn overturn. Shell growth appears coincident with the eight warmest months of the year (May-December) and not during the four coldest months (January-April) based on an ambient transition temperature of $\sim 6^\circ\text{C}$. The salinity corrected shell $\delta^{18}\text{O}$ data exhibit a relationship to the observed bottom temperatures not significantly different from that predicted using Grossman and Ku's [1986] paleotemperature equations for aragonite (from mollusc data only). This evidence suggests that *A. islandica* deposits its aragonite shell in oxygen isotopic equilibrium with the

surrounding water and is an accurate ($\pm 1.2^{\circ}\text{C}$) monitor of local bottom temperatures during the May through December shell growth season. The consistent cold (shell growth) shutdown temperature observed in this study suggests that *A. islandica* shells have the potential to independently record the paleo- $\delta^{18}\text{O}$ (and paleosalinity) composition of its growth environment.

In chapter three, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records, obtained from the shells of *A. islandica* specimens grown in the laboratory in different substrates for one year, were compared with ambient conditions such as water temperature, salinity, $\delta^{18}\text{O}$ -water, water column DIC $\delta^{13}\text{C}$, and pore water DIC $\delta^{13}\text{C}$. Using Grossman and Ku's [1986] aragonite paleotemperature equation (from mollusc data only), the shell $\delta^{18}\text{O}$ records indicate that shell growth occurred when water temperatures were between 5°C and 15°C . Most of shell growth occurred in the last two or three spring months of the experiment, and the springtime shell $\delta^{18}\text{O}$ data closely correspond to the predicted $\delta^{18}\text{O}$ curve for this period. These results suggest that *A. islandica* can be used to accurately reconstruct ambient temperatures within a 10°C range from 5° to 15°C . The $\sim 5^{\circ}\text{C}$ cold shutdown value is not significantly lower than the 6°C value reported in chapter two, but the difference may also be the result of better constraint on the ambient $\delta^{18}\text{O}$ -water and the higher-sampling resolution in this laboratory experiment versus chapter two's Nantucket Lightship comparison study.

The $\delta^{13}\text{C}$ composition of *A. islandica* shells does not appear to be in equilibrium with ambient conditions, though exactly what constitutes $\delta^{13}\text{C}$ equilibrium is still unsettled within the stable isotope community. Shell $\delta^{13}\text{C}$ averages about 1‰ heavier than ambient DIC, but are from 0.2-1.7‰ lighter than "equilibrium" based on three of the most recently published equilibrium relationships. The shell's seasonal variation of ~ 2 ‰ cannot be attributed to ambient DIC $\delta^{13}\text{C}$ variations, which were nearly constant throughout the

period of shell growth. Temperature trends in the fall and spring correlate positively with shell $\delta^{13}\text{C}$ trends as predicted by Mook's [1986] equilibrium relationship, however there are large deviations from this correlation as well. Pore water $\delta^{13}\text{C}$ is shown to have no obvious effect on shell $\delta^{13}\text{C}$. This result does not support the hypothesis that $\delta^{13}\text{C}$ offsets between infaunal and epifaunal bivalves are the result of pore water influence. As external influences cannot fully explain the seasonal $\delta^{13}\text{C}$ signal a physiological effect is suspected which would account for seasonal $\delta^{13}\text{C}$ variation as a positive function of tissue growth. *A. islandica* shells show a $\sim 1\%$ depletion trend during their first 15 years of growth which is related to ontogeny (e.g., reduced tissue growth or the onset of reproductive activity) and not to changes in the external environment.

In chapter four, three shell-derived $\Delta^{14}\text{C}$ time-histories from the eastern North Atlantic are combined with three previously published $\Delta^{14}\text{C}$ records from the western North Atlantic to investigate the evolution of the bomb-signal in the surface North Atlantic Ocean. From these records, pre-bomb (~ 1950) surface North Atlantic $\Delta^{14}\text{C}$ values average -59% , and are most depleted in the Labrador Sea ($\sim -70\%$) and most enriched in the Sargasso Sea ($\sim -50\%$). This $\sim 20\%$ difference between subpolar and subtropical gyre pre-bomb surface waters is attributed to a substantial depleted source for the western subpolar regions, such as Antarctic Intermediate Water convectively mixing along the subpolar front and/or southward flowing polar surface waters from the Davis Strait region. In general, the pre-bomb records show depletion trends that are consistent with the Suess effect, but the magnitude of these depletions varies greatly and are largest on the European shelves, and negligible at Iceland. However, in the North Sea and Norway records, these steep trends are driven by very depleted $\Delta^{14}\text{C}$ values in the mid-1950s that cannot be explained easily by the Suess effect depletion of the overlying atmosphere, and may be the result of increased convection or upwelling in the northern North Atlantic in the early 1950s.

Initial appearance of the bomb- ^{14}C is synchronous in all North Atlantic $\Delta^{14}\text{C}$ records about 1959 (± 1 year). All North Atlantic $\Delta^{14}\text{C}$ records show a steep rise during the early to mid-1960s and reach maxima in the late 1960s to late 1970s. The North Sea record has the largest amplitude ($\sim 320\text{‰}$) and the Iceland record has the smallest ($\sim 120\text{‰}$). Shelf influence is suspected of causing the extremely large amplitude and advanced phase of the North Sea signal relative to other North Atlantic signals. Where shelf influence is considered significant, the adjacent regional ocean $\Delta^{14}\text{C}$ time-histories are corrected by inverting a simple atmospheric uptake model and inputting the shell-derived $\Delta^{14}\text{C}$ records, the shelf water depths, and shelf residence times, and fitting to GEOSECS and TTO surface $\Delta^{14}\text{C}$ data for each ocean region. The Georges Bank, Norway, and North Sea records have been corrected in this way to produce regional records for the Labrador Sea, Norwegian Sea, and the central eastern North Atlantic (Bay of Biscay).

The amplitudes of the "corrected" North Atlantic bomb- ^{14}C signals show a sharp contrast between northern (subpolar gyre) and southern (subtropical gyre) regions with remarkable conformity within each of these regions. Florida, Bermuda, and the central eastern North Atlantic have similar amplitudes of $\sim 220\text{‰}$ and maxima of $\sim +160\text{‰}$. The Labrador Sea, Iceland Sea, and Norwegian Sea have amplitudes of $\sim 120\text{‰}$ and maxima $\sim +60\text{‰}$. These results reflect the contrast between the depths of the winter mixed-layer in the northern and southern regions, but are also consistent with a depleted or deepwater contribution to the subpolar gyre. Also, because the subpolar gyre surface waters are currently more depleted than the atmosphere, these regions should be still increasing their inventory of atmospheric bomb- ^{14}C or maintaining a balance with regional loss through deep water production. A small east-west difference is shown in the relative enrichment of the Norwegian Sea signal versus the Labrador Sea and Iceland Sea signals. This difference is in agreement with deeper mixing in the western section of the subpolar gyre, and the poleward advection of surface Atlantic water into the Norwegian Sea.

In chapter five, a 109-year annual time-series (1875-1983) of bottom temperatures has been reconstructed from the oxygen isotope records of four *A. islandica* shells live-captured near the former position of the Nantucket Lightship at 65 m depth. This site is centered within the "cold pool", and so this time-series provides a century-long thermal history of this outer continental shelf hydrographic feature. The method of establishing the chronology by annual band counting, and using the overlapped isotope records from several shells of different ages to reconstruct a longer record was validated by significant positive correlations between the overlapped portions of the four shell records. The shell-derived bottom temperatures are presented as three annual records: annual mean, annual maximum, and annual minimum. The annual means represent the means of the shell growth seasons, average 7.6°C ($\pm 0.9^{\circ}\text{C}$), and indicate a century-long cooling of $\sim 1.1^{\circ}\text{C}$. The annual maxima represent bottom temperatures at about the time of the autumn overturn, average 11.5°C ($\pm 1.7^{\circ}\text{C}$), and indicate a large century-long cooling of $\sim 2.9^{\circ}\text{C}$. The annual minima represent the spring bottom temperatures, average 5.7°C ($\pm 0.9^{\circ}\text{C}$) and indicate a modest cooling of $\sim 0.4^{\circ}\text{C}$ over the last century. A sharp decline in temperature occurs in all three shell records after 1960, and all records indicate that the mid-1880s, mid-1960s, and mid-1970s were the most prolonged, coldest periods of the past century.

Shell-derived bottom temperature records show a strong similarity to and significant correlation with the Nantucket Lightship bottom temperature records for the period 1960-1971. The shell-derived bottom temperature records show a poor correlation with the coastal sea surface and air temperature records. However, the positive coherence between these records is greatly improved by comparing their detrended or residual (deviation from 10-year running mean) records. This indicates that the short-term anomaly patterns between the continental shelf bottom temperatures and coastal sea surface and air temperatures are similar, while their long-term temperature trends are dissimilar. The primary contrast

between the long-term trends is that the shell-derived records indicate a century-long cooling of $\sim 1^{\circ}\text{C}$, while regional air and coastal sea surface records describe a century-long warming of equal magnitude. It was considered whether this temperature contrast is real or an artifact. Biological or "vital effects" have been rejected as a cause of the disparity because the overlapping shell records provide a check on this type of variation. It was estimated that a 10% increase in the amount of Slope water relative to Scotian shelf water would produce $\sim +0.5\text{‰}$ salinity increase and a $\sim +0.2\text{‰}$ $\delta^{18}\text{O}$ -water increase, which would falsely indicate an apparent $\sim 1^{\circ}\text{C}$ cooling. While, this is a possibility, it merely shifts the problem, from one of a significant temperature change to one of a salinity change, either of which would have broad implications for shelf circulation. It seems equally possible that the shell-derived bottom temperature trends are real and of opposite sign to surface temperature trends. To explain this contrast, it was suggested that increased surface warming is related, directly or indirectly, to increased or prolonged stratification of shelf waters and a net reduction in vertical mixing and heat flux into the bottom waters of the "cold pool".

Arctica islandica is an abundant long-lived species on continental shelves of the northern North Atlantic Ocean. In this thesis, evidence has been compiled which demonstrates that important oceanographic problems can be addressed by utilizing the carbon and oxygen composition of *A. islandica* shells. Finally, this thesis fulfills the task laid out nearly two decades ago by Thompson and Jones [1977] that *Arctica islandica* could someday be used to reconstruct past ocean history as "the *tree* of the North Atlantic".

Appendix

CO₂-H₂O Equilibration Technique: Modified Anderson Method

Water samples collected for oxygen isotope analysis were prepared using a slight modification of methods first described in an unpublished report by T. Anderson [1983] at the University of Illinois, Urbana and also later described by D. Allard [1989] in his Master's Thesis at the University of Rhode Island. These methods essentially involve several freeze/thaw routines along with a standard technique consisting of equilibrating an injected aliquot of CO₂ of known volume and isotopic composition with a seawater sample. The methods produce high repeatability ($\pm 0.01\%$) and accuracy.

I. Water-CO₂ Equilibration:

1. Using a 5 ml (B-D) plastic syringe fitted to a 0.2 ml (Fisher Brand: length 30 cm, external diameter: 5.7 mm) pyrex pipette, several 3-4 ml of seawater from the sample storage bottle is drawn up into the syringe and flushed out as discard at least twice, before drawing up an equal volume to be transferred to the reaction vessel.

2. Before transferring the sample water to the reaction vessel, any water drops on the outside of the pyrex pipette are dried with a clean tissue.

3. Next the pipette is inserted into the neck of the reaction vessel. *Our standard reaction vessel for this procedure is a 20 cm long pyrex tubing (outside diameter: 9 mm; inside diameter: 7 mm) welded to a 10 cm long, 13 mm diameter pyrex test tube (with a rounded bottom). This lower 10 cm portion of the reaction vessel is referred to hereon as the bulb, and the upper 20 cm portion is referred to as the neck.* 1 ml of sample water is then transferred from the syringe to the base of the bulb and then the pipette is removed from the reaction vessel carefully. Care is taken not to get any sample water drops on the upper insides of the bulb or the neck of the reaction vessel.

4. Next, the reaction vessel, kept upright, is connected to the vacuum line.

5. Before evacuation, the base portion of the reaction vessel is submerged in a CO₂ slurry bath for 3 minutes to freeze the water sample.

6. The reaction vessel is then evacuated to a pressure of less than 10 mTorr. This takes about 1 minute on our vacuum line. *Anderson has suggested that this step can be repeated meaning the water sample in the reaction vessel can be thawed, refrozen, and re-evacuated. We have found that while this does remove some residual air locked in the ice, the practical effect on the isotopic composition is negligible, while it markedly increases the chances of cracking the reaction vessel. For these reasons we do not repeat this step.*

7. The CO₂ slurry bath is removed and replaced by a liquid N₂ bath.

8. An aliquot (90 μ mol) of CO₂ was then frozen into the reaction vessel. This takes about 3 minutes to complete.

9. Then the neck the reaction vessel is flame sealed, and allowed to thaw.

10. The sealed reaction vessel is then fully emerged in water bath with a temperature maintained at 25°C ($\pm 0.1^\circ\text{C}$). *For the water bath we have found that a large insulated picnic cooler filled with 25°C water, and then kept in a room whose temperature is maintained close to 25°C, provides a very satisfactory storage bath. With this low-budget setup, the water should be checked 2-3 times a day to make sure water temperature is maintained at 25°C ($\pm 0.1\%$).* Our simple bath has no vibrators to hasten equilibration, and we have found that at least two full days are required for complete equilibration, and to be cautious we have allowed samples to equilibrate for three days.

II. CO₂ Extraction.

- 1. The equilibrated sample vessel is removed from the water bath, and dried off immediately with a paper towel.**
- 2. The upper neck of the vessel is then scored on one side, and inserted into a tube cracker on the vacuum line.**
- 3. The bulb of the reaction vessel is submerged in a CO₂ slurry bath and frozen for 5 minutes.**
- 4. Meanwhile, the cracker chamber is evacuated and checked for leaks.**
- 5. The neck of the reaction vessel is cracked and sample CO₂ released.**
- 6. A liquid N₂ bath is placed onto the collection vessel (either valve or flame-seal type) and 5 minutes is allowed for the CO₂ sample to be transferred.**
- 7. The sample is pumped over for 15 seconds, then collection vessel is isolated (either by closing its valve or by flame sealing).**

Extraction is complete.

III. Further Procedure Notes

1. Calibration and Procedure Test

A series of runs with an in-house standard seawater have been carried to test the repeatability of these procedures. Also, this standard has been directly calibrated with V-SMOW. Results of these runs follow.

N-13 (in-house I)			N-17 (in-house II)			V-SMOW		DATE
	(PDB)	(SMOW)		(PDB)	(SMOW)	(PDB)	(SMOW)	
1	0.247	-1.321	6	0.235		1	1.561	
6	0.255	-1.313				2	1.575	
3	0.243	-1.325						
mean:	0.248	-1.320		0.235	-1.333	1.568	0.00	12/28-29/94
<hr/>								
14	0.247	-1.325				3	1.572	
16	0.261	-1.311						
20	0.223	-1.349						
mean:	0.243	-1.329				1.572	0.00	1/6-7/95
<hr/>								
15	0.304	-1.324				4	1.628	
17	0.310	-1.318						
22	0.322	-1.306						
mean:	0.312	-1.316				1.628	0.00	1/9/95
<hr/>								
25	0.191	-1.340	26	0.190		5	1.531	
26	0.224	-1.307						
27	0.195	-1.336						
mean:	0.203	-1.328		0.190	-1.341	1.531	0.00	1/10/95
<hr/>								
Mean:	-1.323 $\pm$ 0.013			-1.337				

Note: the V-SMOW value of $\sim +1.57\text{‰}$ relative to (PDB) is $\sim 1.85\text{‰}$ larger than the values of $\sim -0.22\text{‰}$ to -0.27‰ mentioned in the literature for this relative difference. This discrepancy is due to the fact that we have established our reference gas (PDB) value from NIST carbonate standards on a system where the phosphoric acid reaction temperature is 90°C rather than the classic 25°C . Based on Swart et al.'s [Swart, Burns, and Leder, 1991] paper this reaction temperature difference would produce $\sim 2.0\text{‰}$ offset, which is bit larger than the offset in our data. Another small portion of the difference is due to the very depleted (-17.0‰) sacrificial gas (our "Matt Gas") used for equilibration. This causes the sample gas (if near V-SMOW values) to be about 0.05‰ depleted relative to its actual value. Thus our V-SMOW value is actually $\sim +1.62\text{‰}$ (PDB- 90°) about 1.9‰ larger than PDB- 25° which moves it a bit closer to Swart, et al.'s predicted offset of $\sim 2\text{‰}$.

2. Figure 1 shows the effect of equilibration temperature on $\delta^{18}\text{O}$ -water. The slope of this line is $-0.165\text{‰}/^\circ\text{C}$. This seems to be a potentially important source of analytical error, as a tenth of a degree variation in equilibration temperature can account for most of the analytical error in our data.

3. Figure 2 shows the results of an equilibration completion test. A negative exponential fit to the data yields a half equilibration time of ~ 7 hours or 0.29 days. The exponential coefficient is -2.38 with time in days. For an initial water-gas isotopic difference of 16‰ , it takes approximately 3 days to reach equilibrium (within $\pm 0.01\text{‰}$ of the final value).

Equilibrium Temperature Data:

Temperature C°	(N-13: CO ₂) δ ¹⁸ O (VSMOW)
25	-1.32
49	-5.31
53	-5.94
59	-6.95

$$\delta^{18}\text{O}_{\text{gasT}} = -0.165 \cdot T + 4.135 + \delta^{18}\text{O}_{\text{w}25^\circ}$$

A-Figure 1

Equilibration Time Data:

Time (N-13: CO₂)
days $\delta^{18}\text{O}$ (VSMOW)

0	-17.23
1	-2.81
1	-2.84
3	-1.33
3	-1.32
3	-1.36
3	-1.33
9	-1.32

$$\delta^{18}\text{O}_{\text{gas}_t} = \delta^{18}\text{O}_{\text{water}} - (\delta^{18}\text{O}_{\text{water}} - \delta^{18}\text{O}_{\text{gas}_0}) * \exp^{-2.38t}$$

A-Figure 2