Appendix 2

Final Report DOE Subcontract Ocean Margins Program

BOTTOM BOUNDARY LAYER MEASUREMENTS IN OMP

Thomas F. Gross, Skidaway Inst. Oceanography

Albert J. Williams III, Woods Hole Oceanographic Institute

DOE/ER/62179--T1-Pt. 3

ABSTRACT:

The main role of the Benthic Acoustic Stress Sensor (BASS) tripods within the Ocean Margins Program experiments was to detect and quantify organic carbon rich particle transport off the shelf. This requires measures of the turbulent boundary layer flow and bed stress, the physical forcing of the particle transport, as well as the concentration and type of particles which are being transported. The BASS tripods were deployed at sites 17 and 26. Data from site 26 were recovered spanning three periods: Feb. 2 - April 6, May 13 - June 27, June 28 - Aug. 18. Site 17 was occupied Feb. 12 - April 11. The BASS tripods were arrayed with five BASS sensors measuring detailed velocity parameters within four meters of the seabed. Velocity time series indicate a usually weak tidal flow which produces small bed stress by itself. On the occasions when a strong flow, probably the Gulf Stream, crosses the area, the bed shear stress increases dramatically to as much as 10 dyne cm^{-2} . This is competent to move unconsolidated sediments in the area. Other instruments from the tripods include: two conductivity / temperature sensor pairs, five WetStar fluorometers, thermistors, transmissometer, optical backscatterance sensors and a pressure sensor.

INTRODUCTION:

The benthic boundary layer is the region where the mean water column flow is affected by seabed drag and decreases from the mean water column velocity to zero at the seabed. The rapid reduction of velocity is forced by turbulent shear stress, which may be described as the vertical flux of mean horizontal momentum. However the stress is forced by the barotropic and baroclinic pressure gradients and therefore is best described by this force balance, rather than simply by the mean velocity gradient. In addition the stress may be functionally partitioned by averaging time scales into mean momentum flux stress and bottom wave induced stress. The fast accelerations near the sea bed caused by surface gravity waves may create large near bed stress capable of suspending sediment, but, because wave motion averages to near zero mean velocity, this large stress has only an indirect effect the mean momentum flux or total particulate and sediment transport.

To describe these many faces of the benthic boundary layer detailed measurements of the 3-D velocity profile near the sea bed were made at several locations during the OMP experiment. The data are evaluated for:

- 1) Mean Stress. Derived from the slope of the logarithmic velocity profile.
- 2) Turbulent Stress and Energy. Obtained from cross-covariance of rapidly sampled turbulent velocity vectors.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible electronic image products. Images are produced from the best available original document.

- 3) Bottom wave motions. Usually the near bed wave motions may be described by average wave period, average wave velocity and average orbital excursion. These are all estimated from the Fourier decomposed velocity time series by identifying the peak wave frequency and assigning to it the wave motion band variance.
- 4) Particulate Response to Stress. Using direct measurements with OBS of particulate concentration at a few points above the seabed, a Rouse profile is calculated from the mean stress velocity scale, u^* , and the particle fall velocity. Total suspended load is then calculated as the depth integral of the Rouse profile and transport rates are calculated using both the concentration profile and velocity profile data.

DATA COLLECTED:

All instruments mounted on the BASS tripod were controlled and data logged to the central BASS 340 Megabyte data logger. The sampling scheme was devised to fill the 340 Megabyte data logger disk in the 103 days. A data record consists of all instrument data sampled at 2Hz for 133 seconds for 22876 bytes per record. A data record was collected every ten minutes starting 1 min after the top of the hour.

- 1.) Velocity Measurements. Five BASS sensors at 447.5 cmab, 257.5 cmab, 122.5 cmab, 62.5 cmab and 32.5 cmab.
- 2.) Wave Motion Measurements. Paro-Scientific pressure sensor at 400 cmab.
- 3.) Water Properties. 2 Seabird conductivity and temperature pairs 100 cmab, 400 cmab.
- 4.) Chlorophyll Measurements. Five WETStar Miniature Fluorometers 445 cmab, 258 cmab, 110 cmab, 35 cmab and 28 cmab.
- 5.) Particulate Measurements. Five Optical Backscatterance Sensors 445 cmab, 258 cmab, 110 cmab, 35 cmab and 28 cmab. One SeaTech Light Transmissometer, 25 cm pathlength at 400 cmab.

RESULTS:

The 40 hour low pass velocity vectors demonstrate the large eddy veering flows which move through the area (Fig. 1). On March 6, June 5 and August 14, large and persistent flows toward the NE, the direction which would be expected for Gulf Stream intrusions into the area were observed. These are the largest amplitude velocities observed and indicate that critical conditions for particle transport will be highly biased to NE transport. Other events may correlate with atmospheric wind events. However the passage of tropical storm Andrew on June 20 was only a ten hour event and thus is not evident in the low passed signal.

The second panel of Fig.1 shows samples of the velocity vector time series with the 40 hour low pass signal removed. Dominating the resultant signal is the tidal forcing which at 4 meters above the bed appears to be about 10 cm/s at the M2 frequency.

Figure 2 summarizes the analysis of turbulence parameters obtained from the logarithmic velocity profile technique. The shear velocity, U^* , covaries strongly with the low passed velocity. The tidal background U^* can be seen from the low amplitude 40 hour low

passed times (~March 20) to be only 1 cm/s. Bed sediments will be winnowed of all but particles large enough to avoid motion at this persistent level of stress. The large values of Z_o (> 5 cm) are outside the acceptable range of Nikuradse physical roughness. Very rough boundary layers (non-homogeneous on scales of meters, e.g. rocks, boulders and bottom undulations) exhibit this, but with scattered and low $R2$ also indicate violation of the log profile model. Some directional dependence may indicate local roughness problems of this type. During strong flow Z_o reduces to approximately 1 cm such as during the March 6 event. When $R2$, the correlation coefficient for the logarithmic profile fit, is less than 0.8 the log profile is not really present, and the estimates of U^* and Z_o are useless. (However differentiating profiles by $R2$ in the range of 0.90 - 1.0 does not indicate statistically different profile quality.) As expected large $R2$ occurs with large velocity and large U^* . Since sediment transport from bottom stress only occurs with large U^* the estimates with low $R2$ and low U^* can be safely ignored.

Figure 3 summarizes the turbulence quantities obtained from the ten minute records of 2 Hz sampled data. Average speed for ten minute intervals is plotted first for comparison to turbulent quantities. $Q2$, the turbulent velocity variance, within a 133 second record is an indicator of the magnitude of turbulence within the bottom boundary layer, but, may, at times be caused by the presence of wave motion near the sea bed. The correlation for Reynolds stress, $u'w'$, is expected to be negative for a well behaved boundary layer. Wave motions or upstream obstacles deflecting the current can cause positive values. Short averaging times of 133 seconds also allow a large statistical uncertainty (on order $\pm 50\%$) accounting for some positive values. The energy and stress vary together for the most part, however they do not always track with the mean velocity. While the mean velocity forces the turbulence, the effects of stratification and surface gravity waves can greatly alter the relationship.

CONCLUSIONS

Several anomalous factors make this data set interesting. On July 15-18 very large kinetic energy and stresses are observed with little speed increase. While on August 14 very large speeds have no signature in energy or stress. The June 5 event has large speed and possibly wave motion (or energy in the 5-30 second band), but the energy and stress do not grow until several days later. However in the same deployment (and therefore with the same surrounding topography) the event of May 14-16, with flow from the same direction and amplitude is nicely correlated with energy and stress.

Preliminary bottom transport estimates indicate that a few events dominate the transport of near bottom particles off the shelf. Sweeps, by what might be the Gulf Stream, are more important to sediment transport than are local storms, even hurricanes. Storm waves have a difficult time penetrating to the 75 meter depth. Even then the transport of particles depends upon a mean current, which may not be present during an atmospheric event.

40 hr low pass

Tidal band (2 hour-40 hour)