NITROGEN INTERACTIONS AND PHOTOSYNTHETIC RESPONSES TO CO₂: WORK PLAN FOR BIOCON EXPERIMENT/PHYSIOLOGICAL MEASUREMENTS AT CEDAR CREEK

In fulfillment of the subcontract
from the University of Minnesota to Brookhaven National Laboratory

D. Ellsworth, Brookhaven National Laboratory

Preface:

Nitrogen plays a critical role in photosynthetic function, which in turn can affect many ecosystem processes through its effects on plant growth and ecosystem carbon cycles. As a result of its central role in photosynthetic enzymes, leaf N status can affect the magnitude of photosynthetic enhancement by elevated CO₂. It is now well-recognized that species responses to elevated CO₂ may be different when species are grown in isolation vs. in a mixed community. Part of this effect may result from effects on leaf N itself as a result of species competition for N in N-limited ecosystems, and part of the effect is simply a function of dilution of leaf nutrients in leaves with greater carbohydrates as a result of elevated CO₂. However, photosynthetic efficiency itself may be affected if N-limited plants reallocate N within leaves away from carboxylation functions under elevated CO₂ compared to ambient plants (Drake et al. 1997). Thus while there is cause to believe that there are interactive effects of N and CO₂ on species photosynthetic physiology, there is little experimental data to support or reject this idea, particularly in realistic ecosystems under field conditions.

Biotic interactions, most notably the presence of N-fixing plants, can affect ecosystem N stocks and carbon cycling via effects of N on photosynthetic function (Chapin et al. 1997, Hooper and Vitousek 1997). If photosynthetic responses of leaves and ecosystems are largely mediated through canopy N, then biotic as well as inorganic N sources will affect the magnitude of these responses. Under elevated CO₂ there is evidence from the Swiss FACE experiment that growth and photosynthetic function are most responsive to CO₂ in species mixtures including N-fixers (Hebeis et al. 1997, Lüscher et al. 1998, S.P. Long, pers. comm.). However, in that experiment there were confounding management factors and species diversity effects *per se* could not be tested. Still, photosynthetic studies showed that CO₂-induced photosynthetic adjustments in the Swiss FACE experiment were greater under low N and in monocultures than in the mixture of grasses and the N-fixing species clover (S.P. Long, pers. comm.). Effects of species diversity and N-fixers in specific on plant CO₂ responses in interaction with N have important implications for predicting ecosystem responses to elevated CO₂ under a variety of site conditions, and may also temper management for mitigation of ecosystem CO₂ responses.

Objective: Determine Nitrogen and Biodiversity (N-fixer) effects on photosynthetic responses in interaction with elevated CO₂ in the BioCON FACE experiment.

Hypotheses:

- 1) The magnitude of photosynthetic responses to elevated CO₂ will vary in magnitude according to canopy N status.
- 2) Photosynthetic acclimation of species will be greatest for N-limited plants, and progressively lower for plants with progressively higher leaf N.
- 3) Species interactions with legumes will improve N status and hence the photosynthetic response to elevated CO₂.

My specific prediction is that the magnitude of the photosynthetic response to CO₂ will be greatest for C-3 plant species in high N x diversity plots followed by high N monoculture, low N x diversity+N fixer plots, and low N x diversity+N fixer plots (see Figure below).

Design:

The site is the BioCON FACE experiment (Cedar Creek Experiment E141) at the Cedar Creek Natural History Area. The overall experimental design for BioCON is described in the original proposal by Reich et al. and similar to that used in Tilman et al. (1997) except that species are grown in combinations of 1, 4, 9 and 16 species mixtures within high and low N sectors of ambient and elevated CO₂ FACE rings. The specific measurement design will involve temperature and light-controlled measurements of

photosynthetic responses to CO₂ (A-c_i curves) for each species in a variety of plots. Each species (see list below) will be measured in monoculture stands in high and low N plots under ambient and elevated CO₂. In addition, each species will be measured in 9-species polyculture stands containing no legumes (-N fixer) or containing at least 2 species of legumes (+N fixer). Thus the overall design is 4 C-3 species x 2 CO₂ levels x 2 N levels x 3 diversity levels (monoculture, diversity-N fixer, and diversity+N fixer) x 2 to 3 reps (depending on plot type). The measurement matrix yields a total of 128 A-c_i curves to be measured. In addition, both legume species (see list below) will be measured in monoculture at high and low N and CO₂ and one legume (probably *Lespedeza*) will be also measured in the 9 species diversity +N fixer plots (16 + 12 more A-c_i curves).

Three visits will be made to the site in 1998 (May, July and Aug.) for the purpose of conducting A-c_i curve measurements and conductance measurements under the Ellsworth/TECO project. Early season perennials will be measured in May 1998 as well as some of the species with later phenology. The majority of species will be measured in July except those that show early senescence. For the projected 156 A-c_i curves anticipated to be measured in the experimental design, roughly 25 full working days will be required and at least two gas exchange systems will need to be operated simultaneously, not including equipment problems and inclement weather considerations.

A-c_i curves will be analyzed using the biochemical model of photosynthesis (Farquhar model; see Long 1991, Wullschlegel 1993) to determine if adjustment of photosynthetic characteristics has occurred and the nature of these adjustments in all measured treatment combinations. Foliage will be analyzed for N concentration and SLA, and a subset of foliar samples will be run for $\delta^{13}\text{C}$ (to determine the c/c_a setpoint for photosynthesis) and $\delta^{15}\text{N}$ (to see if N transfer from legumes can be inferred). Data will be analyzed for effects of CO₂ on photosynthetic efficiency (initial slope of A-c_i curve as a function of N) under the treatment conditions, and for CO₂, N and legume diversity effects on total N as a determinant of photosynthesis under growth c_a and photosynthetic efficiency. The analysis of A-c_i curves as a function of the experimental treatment will yield dependencies of the initial slope (V_{cmax} from the Farquhar model) and A_{net} at a given c_a as a function of leaf N (see attached figure). To the extent that there are interactions among experimental variables, the effects of these variables on both A_{net} and N will not be additive.

LIST of species for experimentation:

Agropyron repens, C-3 grass (or *Koeleria cristata* ??)

Bromus inermis, C-3 grass

Achillea millefolium, C-3 Forb

Solidago rigida, C-3 Forb

Lespedeza capitata, Legume

Lupinus perennis, Legume

References

- Chapin, F.S. III, B.H. Walker, R.J. Hobbs, D.U. Hooper, J. H. Lawton, O. E. Sala, D. Tilman. 1997. Biotic control over the functioning of ecosystems. *Science* 277: 500
- Drake B.G., M. Gonzales-Meler and S.P. Long. 1997. More efficient plants: A consequence of rising atmospheric CO₂? *Annual Review of Plant Physiology and Plant Molecular Biology* 48: 609-639.
- Hebeisen, T., A. Lüscher, S. Zanetti, B.U. Fischer, U. Hartwig, M. Frehner, G.R. Hendrey, H. Blum and J. Nösberger. 1997. Growth response of *Trifolium repens* L. and *Lolium perennne* L. as monocultures and bi-species mixture to free air CO₂ enrichment and management. *Global Change Biology* 3: 149-60.
- Hooper D.U. and P.M. Vitousek. 1997. The effects of plant composition and diversity on ecosystem processes. *Science* 277:1302
- Long S.P. 1991. Modifications of the response of photosynthetic productivity to rising temperature by atmospheric CO₂ concentrations: has its importance been underestimated? *Plant, Cell and Environment* 14: 729-739.
- Lüscher, A., G.R. Hendrey and J. Nösberger. 1998. Long-term responsiveness to free-air CO₂ enrichment of functional types, species and genotypes of plants from fertile permanent grassland. *Oecologia* 113: 37-45.

- Tilman, D., J. Knops, D. Wedin, P. Reich, M. Ritchie, E. Siemann. 1997. The influence of functional diversity and composition on ecosystem processes. *Science* 277:1300
- Wedin, D.A. and D. Tilman. 1996. Influence of nitrogen loading and species composition on the carbon balance of grasslands. *Science* 274:1720.
- Wullschlegel S.D. 1993. Biochemical limitations to carbon assimilation in C₃ plants - A retrospective analysis. *Journal of Experimental Botany* 44: 907-920.

Figure: A family of A-c_i curves for a C-3 plant in the BioCON experiment illustrating the dependence of curve parameters on leaf N, which in turn is influenced by experimental treatment (+N, +N-fixer, +CO₂ and combinations thereof).