Evaluate Status of Pacific Lamprey in the Clearwater River Drainage, Idaho

Annual Report 2000

This Document should be cited as follows:

Cochnauer, Tim, Christopher Claire, "Evaluate Status of Pacific Lamprey in the Clearwater River Drainage, Idaho", 2000 Annual Report, Project No. 200002800, 28 electronic pages, (BPA Report DOE/BP-00000090-1)

Bonneville Power Administration
P.O. Box 3621
Portland, OR 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

**EVALUATE STATUS OF PACIFIC LAMPREY IN THE CLEARWATER RIVER
DRAINAGE, IDAHO**

ANNUAL REPORT 2000

Prepared by:
Tim Cochnauer, Regional Fishery Manager
Christopher Claire, Fishery Technician

Idaho Department of Fish and Game
Clearwater Region
1540 Warner
Lewiston, ID 83501

Prepared for:
U.S. Department of Energy
Bonneville Power Administration
Environmental Fish and Wildlife
P.O. Box 3621
Portland, OR 98203-3621

Project Number 2000-028-00
Contract Number 00000090-00001

TABLE OF CONTENTS

ABSTRACT	1
INTRODUCTION.....	2
PROJECT AREA	2
OBJECTIVES	3
METHODS	3
RESULTS	5
SUMMARY AND DISCUSSION	9
LITERATURE CITED.....	11
APPENDIX A.....	12
APPENDIX B	23

LIST OF TABLES

Table 1.	Habitat substrate classification for sampling sites in the South Fork Clearwater River Drainage.....	5
Table 2.	Habitat locations of Pacific lamprey juveniles in randomly sampled units in Red River and selected units in South Fork Clearwater River, ID, 2000.....	8

LIST OF FIGURES

Figure 1.	Geographic location of Pacific lamprey investigations in the South Fork Clearwater River, drainage, 2000.....	3
Figure 2.	Length frequency of Pacific lamprey (N=13) captured in Red River migrant trap, South Fork Clearwater River drainage, ID, 2000.....	6
Figure 3.	Length frequency of Pacific lamprey (N=154) captured by electroshocking in Red River, ID, 2000.....	7
Figure 4.	Length frequency of Pacific lamprey (N=93) captured by electroshocking in South Fork Clearwater River, ID, 2000.....	9

ABSTRACT

Recent decline of Pacific lamprey *Lampetra tridentata* adult migrants to the Snake River drainage has focused attention on the species. Adult returns in 1995-1999 were more than ten magnitudes less than returns in the early 1960's. Human activities in the Snake River and Clearwater River drainages have altered ecosystem habitat in the last 100 years and likely the productive potential of Pacific lamprey habitat. Logging, stream impoundment, road construction, grazing, mining, and community development have dominated habitat alteration in the Clearwater River system and Snake River corridor. Hydroelectric projects in the Snake River corridor impact juvenile Pacific lamprey outmigrants and returning adults. Juvenile lamprey outmigrants potentially pass through turbines, turbine bypass and collection systems, and spillway structures at lower Snake River hydroelectric dams. Clearwater River drainage hydroelectric facilities including the Pacific Power and Light Dam on the Clearwater River in Lewiston, Idaho, impacted Pacific lamprey populations, however, the degree of impact is unknown (1920's-early 1970's). Hydroelectric dam construction (Harpster Dam) on the South Fork of the Clearwater River resulted in obstructed salmonid passage in the mid-1900's. Habitat alterations in the Snake River basin and Clearwater River drainage have had numerous negative effects on salmon *Oncorhynchus spp.* and steelhead trout *O. mykiss* populations (wild fish), but the magnitude of impacts on lamprey productivity and survival is unknown. Thorough understanding of Pacific lamprey habitat use and life history processes is needed to facilitate management and restoration of the species. Through Bonneville Power Administration support, the Idaho Department of Fish and Game began investigation into the status of Pacific lamprey populations in Idaho's Clearwater River drainage in 2000. Trapping, electrofishing, and spawning ground redd surveys were used to determine where Pacific lamprey persist in the South Fork of the Clearwater River drainage. Habitat surveys evaluating juvenile habitat use were primarily conducted in the Red River subbasin. Red River subbasin resource manipulations have resulted in elevated stream sediment, stream destabilization, riparian canopy reduction, and water temperature extremes. A total of 262 juvenile Pacific lamprey were captured during the 2000 field season. Sampling in the Red River drainage yielded the largest number of Pacific lamprey juveniles. Preliminary findings indicate Pacific lamprey juveniles, while present, are not numerous or widely distributed. Age of juveniles captured was determined using length frequency.

INTRODUCTION

The Pacific lamprey *Lampetra tridentata* is facing the same migratory hazards and habitat degradation as other anadromous fish species in Idaho. Because this fish is not recognized as a sport or game fish species, little attention has been given to its status. Basic life history, distribution, and population status are urgently needed to fully understand this species and to begin intensive management before extinction occurs in Idaho.

The South Fork Clearwater River (SFCR) drainage of north central Idaho is an important study area as both Pacific lamprey ammocoetes and macrothemia have been captured in outmigrant smolt traps since 1992.

Understanding juvenile fish population composition, migrational behavior, and habitat needs will provide basic information to better manage Pacific lamprey. Without this knowledge, the opportunity for preservation of critical habitat may be lost. This project will add to our knowledge of Pacific lamprey and provide critical information to minimize future degradation of habitat.

PROJECT AREA

The Clearwater River drainage is located in north central Idaho and encompasses approximately 2.5 million hectares. The SFCR, one of the four major tributaries of the Clearwater River, encompasses 300,440 hectares. The upper SFCR watershed has several large meadow complexes with low stream gradients and fine substrates. The lower SFCR reaches are predominantly canyon confined and boulder substrate dominated. In 1910 Grangeville Electric Light and Power Company built a hydroelectric dam on the SFCR at rkm 32.0. In 1937 Avista Utilities, (formerly Washington Water Power), acquired the dam. Steelhead trout *O. mykiss* migration was possible, although limited, over the dam from 1935 to 1949. High flows destroyed the fishway in 1945 eliminating adult salmonid passage until the dam was removed in 1963. Adult Pacific lamprey passage may have occurred during this entire period as adult lamprey have the ability to climb above water surface levels (G. Starke, U.S. Army Corps of Engineers, personal communications). Pacific lamprey returns to the main Clearwater River in the period following removal of the SFCR dam could have provided recolonization stock for the SFCR drainage.

The current land ownership of the SFCR watershed is U.S. Forest Service (68%), private (28%), Nez Perce Tribe (0.9%), Bureau of Land Management (2%), and State of Idaho (0.7%). Land use activities range from predominantly forestry related in the upper SFCR to livestock pasture and grazing in lower reaches. Historically, mining was centered in the upper reaches. Extensive mining from the 1860's to the mid-1900's occurred in four SFCR tributaries, Crooked River, Red River, American River, and Newsome Creek. Generally mining in the watershed has impacted fish production in varying degrees through sedimentation and instream riparian degradation. Riparian

canopy removal in the meadow complexes of the upper SFCR watershed contributes to stream temperature extremes.

OBJECTIVES

1. Determine life history characteristics of Pacific lamprey ammocoetes and macrothemia in the Clearwater River drainage.
2. Determine habitat requirements of Pacific lamprey in the Clearwater River drainage.
3. Determine distribution of Pacific lamprey in SFCR drainage.

METHODS

Electroshocking with an ABP-2 electrofisher was used to capture fish in the stream channel. Any Pacific lamprey mortalities were to be retained for statolith banding age determination, following procedures described by Beamish and Medland (1988). Determination of habitat usage and distribution was focused in the Red River drainage. We segmented Red River into one-kilometer sections from its mouth upstream to the uppermost bridge crossing. We prioritized 41 sampling locations based on random selection of kilometer section and sampled the first 100 meters of each selected section. The habitat in each of the sampled sections was classified as to type (Table 1). The first riffle, pool, glide, etc., was electroshocked from its downstream boundary upstream without repeating in identical habitat types. Water depth, water velocity, and substrate were measured and recorded at the site of individual captures. Other selected sites in the SFCR drainage were also sampled by electroshocking.

Three downstream migrant traps currently operated by Idaho Department of Fish and Game (IDFG) in the SFCR drainage were used to monitor Pacific lamprey downstream movements. The Crooked River scoop trap (rkm 1.0) was operated from March 26 to October 31. A 1.22m diameter rotary screen trap on American River (rkm 3.0) was operated from March 26 to October 31. Another 1.22m diameter rotary screen trap on Red River (rkm 6.0) was operated from March 23 to October 31.

Captured Pacific lamprey juveniles were anesthetized, and total lengths and body weights were measured. Individuals were then recovered in fresh water and released near the site of capture. All Pacific lamprey juveniles (greater than 100 mm in length) captured by electroshocking were marked with fluorescent orange elastomer on the left side behind the gill openings. Outmigrant estimates past traps were made using Beamish and Levings (1991) trap-area fished methods.

Eight kilometers of Red River and its tributaries were surveyed for spawning adult Pacific lamprey from April 19 to July 7.

Figure 1. Geographic location of Pacific lamprey investigations in the S. F. Clearwater River drainage, 2000.

Table 1. Habitat substrate classification for sampling sites in the South Fork Clearwater River drainage.

Habitat Units	Substrate Classification
Falls	Substrate Type (mm)
	Large Boulder 512 +
Cascades	Small Boulder 256-512
	Cobble 64-256
Rapids	Coarse Gravel 16-64
Typical	Medium Gravel 8-16
Boulders	Fine Gravel 2-8
Bedrock	Course Sand .5-2
	Fine Sand 0.062-0.50
Riffles	Silt/Organic 0.004-0.062
Typical	
Pocket-water	
Glide	
Pools	
Lateral Scour Pool	
Straight Scour Pool	
Plunge Pool	
Dammed Pool	
Alcove	

Snake River juvenile lamprey passage information at Lower Granite project was obtained from Washington Department of Fish and Wildlife in 2000. Juvenile mortality samples of macrothamia and ammocoetes were collected. Bypass mortality, collection processes, and sampling procedures were observed at Lower Granite facility on five occasions from April 1 to July 1.

RESULTS

During 2000, no Pacific lamprey ammocoetes or macrothamia were captured in the Crooked River scoop or the American River rotary screen traps. Fifteen Pacific lamprey ammocoetes and one macrothamia were captured in the Red River rotary screen trap. The average total length of the ammocoetes was 125 mm and the length of the macrothamia was 145 mm (Figure 2).

Figure 2. Length frequency of Pacific lamprey (N=13) captured in the Red River migrant trap, South Fork Clearwater River drainage, ID, 2000

Because of the low numbers of Pacific lamprey captured at traps, no mark-recapture outmigrant estimates were performed. Based on trap-area fished, a total of 93 ammocoetes and 7 macrothalmia were estimated to have migrated past the Red River trap in 2000.

Because statolith banding examination for age determination requires sacrificing individuals, no age determinations were attempted for fish captured. Future age determination may include statolith banding examination. Age assessment, based on length frequency (Figures 2-4), suggests there may be four age classes in the fish sampled.

A total of 154 Pacific lamprey ammocoetes were captured by electroshocking in Red River. Juvenile lamprey were found in three sections of Red River up to rkm 3.0, however, no sections between rkm 3.0 and 8.0 were sampled due to the randomness of selection. Tributaries of Red River (Siegal Creek, Red Horse Creek, and S.F. Red River) were sampled, but no Pacific lamprey were found. The largest Pacific lamprey captured electroshocking measured 166 mm while the smallest (found in SFCR) was 47 mm (Figure 3). No macrothalmia were captured while electroshocking.

More Pacific lamprey ammocoetes were captured in lateral scour pool habitat than any other single habitat type, however, the highest Red River density was found in the alcove (non-random) habitat type (Table 2). Individuals were mostly found inhabiting sand/silt substrate behind boulders and were captured in water depths ranging from just over 1.0 meter to less than 0.1 meter.

Figure 3. Length frequency of Pacific lamprey (N=154) captured by electroshocking in Red River, ID, 2000.

Although numerous potential spawning sites were identified, no adult Pacific lamprey or redds were observed during 2000 redd surveys.

Non-random habitat surveys were conducted on potential juvenile lamprey habitat sites in the main stem SFCR, American River, and Newsome Creek. Five sites were sampled in the Newsome Creek drainage from rkm 0.1 to rkm 13.4, five in the American River drainage from rkm 0.1 to rkm 18.0, and one in the SFCR at rkm 44.3, 8.8 km upstream of the Mt. Idaho bridge. No lamprey were found sampling Newsome Creek and American River. Ninety-three lamprey were sampled in the SFCR site (Figure 4).

Table 2. Habitat locations of Pacific lamprey juveniles in randomly sampled units in Red River and selected units in South Fork Clearwater River, ID, 2000.

RED RIVER					
Habitat Type	Lamprey Captured	Total Area Fished (m²)	Total Time Fished(Min)	Fish/100m²	CPUE(Fish/Min.)
Lateral Scour Pool	99	1397.9	966	7.1	0.10
Riffle	2	662.6	325	0.3	0.01
Riffle w/Pockets	3	420.4	125	0.7	0.02
Rapids w/Boulders	4	175.1	140	2.3	0.03
Straight Scour Pool	25	211.3	170	11.8	0.15
Alcove (non-random)	19	7.5	20	253.3	0.95
TOTALS:	152	2874.8	1746	Na	Na
SOUTH FORK CLEARWATER RIVER					
Lateral Scour (along margin)	93	11.0	20	845.5	4.65

Figure 4. Length frequency of Pacific lamprey (N=93) captured by electroshocking in South Fork Clearwater River, ID, 2000.

SUMMARY AND DISCUSSION

It is not known if lamprey densities observed in 2000 reflect overall low population numbers in the SFCR drainage as the low juvenile densities would be expected following several years of a few spawning adults. Relatively high effort to catch rates were partially a result of random sampling in sections of the Red River drainage outside the expected potential habitat range. Potential juvenile habitat sampled in Newsome Creek and American River did not yield lamprey. It is possible distribution of lamprey in the SFCR basin is limited. However, further sampling needs to be completed before definitive population numbers and distribution can be determined. Length frequency distribution data indicated possibly four age classes present, assuming growth rates similar to those noted by Scott and Crossman (1973) for juveniles in native habitat. Age determination precision can only be increased with future sampling. This year's sampling (random units) found the greatest number of juveniles in lateral scour habitat type. Alcove habitat type, however, showed the highest total density per meter squared. Flow velocities are generally lower in alcoves than lateral scour pools providing energy conserving burrowing conditions for ammocoetes.

No Pacific lamprey juvenile mortalities were observed at outmigrant traps, but three mortalities were noted as a result of handling while electroshocking in the SFCR.

Juvenile lamprey passage information obtained from Washington Department of Fish and Wildlife at Lower Granite dam has not been fully reviewed. Preliminary findings indicate that the majority of juveniles pass the project between March 30 and July 1.

Both life stages (ammocoete and macrothemia) are present in Lower Granite samples. Rearing of some ammocoetes in the Snake River pool could account for both life stages being sampled at the project.

LITERATURE CITED

- Beamish, R. J., and C. D. Levings. 1991. Abundance and Freshwater Migrations of the Anadromous Parasitic Lamprey, *Lampetra tridentata*, in a Tributary of the Fraser River, British Columbia. Can. J. of Fish. and Aquatic Sci. 48: pp.1250-1263.
- Beamish, F. W., and T. E. Medland. 1988. Age Determination for Lampreys. Transactions of the American Fisheries Society. 117: pp.63-71.
- Scott, W. B., and E. J. Crossman. 1973. Freshwater Fishes of Canada. Fisheries Research Board of Canada Ottawa. Bulletin 184. pp. 42-45.

APPENDIX A

A1. Red River Pacific Lamprey Length and Weight

Date Electrofished	Km/ Reach	Habitat Type	Area Fished M ²	Time Fished (Min.)	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans/ Untrans.
7/28/00	01/01	RIP	420.4	125	3	3	135	3.8	LCO	U
7/28/00	01/01	RIP					122	2.9	LCO	U
7/28/00	01/01	RIP					136	4.1	LCO	U
8/12/00	01/01	LSP	78.0	46	3	3	127	3.6	LCO	U
8/12/00	01/01	LSP					115	2.7	LCO	U
8/12/00	01/01	LSP					145	5.0	LCO	U
8/12/00	01/01	RIF	102.7	70	2	2	159	6.6	LCO	U
8/12/00	01/01	RIF					134	4.1	LCO	U
8/12/00	01/01	RBB	na	na	2		--	--	--	U
8/12/00	01/01	RBB					--	--	--	U
8/12/00	00/09	ALC	7.5	20	19	19	102	1.4	LCO	U
8/12/00	00/09	ALC					150	5.5	LCO	U
8/12/00	00/09	ALC					150	5.7	LCO	U
8/12/00	00/09	ALC					116	2.6	LCO	U
8/12/00	00/09	ALC					151	5.6	LCO	U
8/12/00	00/09	ALC					140	4.7	LCO	U
8/12/00	00/09	ALC					117	2.9	LCO	U
8/12/00	00/09	ALC					105	1.7	LCO	U
8/12/00	00/09	ALC					124	3.1	LCO	U
8/12/00	00/09	ALC					80	1.2	LCO	U
8/12/00	00/09	ALC					151	5.8	LCO	U
8/12/00	00/09	ALC					156	6.7	LCO	U
8/12/00	00/09	ALC					150	5.2	LCO	U
8/12/00	00/09	ALC					148	6.0	LCO	U
8/12/00	00/09	ALC					147	5.6	LCO	U
8/12/00	00/09	ALC					157	7.4	LCO	U
8/12/00	00/09	ALC					139	4.2	LCO	U
8/12/00	00/09	ALC					119	3.1	LCO	U
8/12/00	00/09	ALC					128	3.4	LCO	U
8/13/00	03/04	LSP	150.0	175	45	45	146	5.8	LCO	U
8/13/00	03/04	LSP					150	5.7	LCO	U
8/13/00	03/04	LSP					152	5.5	LCO	U
8/13/00	03/04	LSP					162	7.0	LCO	U
8/13/00	03/04	LSP					127	4.3	LCO	U
8/13/00	03/04	LSP					135	4.2	LCO	U
8/13/00	03/04	LSP					127	3.7	LCO	U
8/13/00	03/04	LSP					125	3.7	LCO	U

*mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A1 (Con't). Red River Pacific Lamprey Length and Weight

Date	Km/ Electrofished	Habitat Type	Area Fished M ²	Time Fished (Min.)	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans/ Untrans.
8/13/00	03/04	LSP					141	4.9	LCO	U
8/13/00	03/04	LSP					159	6.1	LCO	U
8/13/00	03/04	LSP					157	6.7	LCO	U
8/13/00	03/04	LSP					152	5.9	LCO	U
8/13/00	03/04	LSP					156	6.5	LCO	U
8/13/00	03/04	LSP					147	5.7	LCO	U
8/13/00	03/04	LSP					148	5.8	LCO	U
8/13/00	03/04	LSP					144	5.5	LCO	U
8/13/00	03/04	LSP					146	5.7	LCO	U
8/13/00	03/04	LSP					159	6.4	LCO	U
8/13/00	03/04	LSP					159	6.4	LCO	U
8/13/00	03/04	LSP					154	6.2	LCO	U
8/13/00	03/04	LSP					156	6.8	LCO	U
8/13/00	03/04	LSP					155	6.3	LCO	U
8/13/00	03/04	LSP					145	5.6	LCO	U
8/13/00	03/04	LSP					130	3.9	LCO	U
8/13/00	03/04	LSP					134	4.1	LCO	U
8/13/00	03/04	LSP					165	7.3	LCO	U
8/13/00	03/04	LSP					152	6.1	LCO	U
8/13/00	03/04	LSP					95	1.7	LCO	U
8/13/00	03/04	LSP					147	5.6	LCO	U
8/13/00	03/04	LSP					152	5.5	LCO	U
8/13/00	03/04	LSP					155	5.6	LCO	U
8/13/00	03/04	LSP					166	7.6	LCO	U
8/13/00	03/04	LSP					155	6.5	LCO	U
8/13/00	03/04	LSP					166	10.9	LCO	U
8/13/00	03/04	LSP					160	6.9	LCO	U
8/13/00	03/04	LSP					161	6.8	LCO	U
8/13/00	03/04	LSP					108	1.9	LCO	U
8/13/00	03/04	LSP					121	3.2	LCO	U
8/13/00	03/04	LSP					97	1.8	LCO	U
8/13/00	03/04	LSP					110	2.5	LCO	U
8/13/00	03/04	LSP					99	1.8	LCO	U
8/13/00	03/04	LSP					98	1.6	LCO	U
8/13/00	03/04	LSP					91	1.3	LCO	U
8/13/00	03/04	LSP					107	2.0	LCO	U
8/13/00	03/04	LSP					92	1.3	LCO	U
8/14/00	03/04	LSP	152.3	135	51	50	155	5.9	LCO	U
8/14/00	03/04	LSP					154	6.0	LCO	U
8/14/00	03/04	LSP					157	6.6	LCO	U
8/14/00	03/04	LSP					162	8.0	LCO	U
8/14/00	03/04	LSP					153	6.6	LCO	U
8/14/00	03/04	LSP					155	5.7	LCO	U
8/14/00	03/04	LSP					99	1.8	LCO	U

*mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A1 (Con't). Red River Pacific Lamprey Length and Weight

Date	Km/	Habitat	Area	Time	Lamprey	Lamprey	Length	Weight	Recapture/	Trans/
Electrofished	Reach	Type	Fished	Fished	Captured	Measured	(mm)	(g)	Marked*	Untrans.
			M ²	(Min.)						
8/14/00	03/04	LSP					157	7.1	LCO	U
8/14/00	03/04	LSP					155	6.5	LCO	U
8/14/00	03/04	LSP					138	4.5	LCO	U
8/14/00	03/04	LSP					152	6.3	LCO	U
8/14/00	03/04	LSP					158	7.1	LCO	U
8/14/00	03/04	LSP					148	5.5	LCO	U
8/14/00	03/04	LSP					157	6.5	LCO	U
8/14/00	03/04	LSP					156	6.4	LCO	U
8/14/00	03/04	LSP					142	5.1	LCO	U
8/14/00	03/04	LSP					149	5.8	LCO	U
8/14/00	03/04	LSP					142	5.1	LCO	U
8/14/00	03/04	LSP					155	7.2	LCO	U
8/14/00	03/04	LSP					152	6.2	LCO	U
8/14/00	03/04	LSP					150	5.2	LCO	U
8/14/00	03/04	LSP					139	4.4	LCO	U
8/14/00	03/04	LSP					155	6.3	LCO	U
8/14/00	03/04	LSP					147	5.2	LCO	U
8/14/00	03/04	LSP					97	1.9	LCO	U
8/14/00	03/04	LSP					93	1.9	LCO	U
8/14/00	03/04	LSP					160	7.9	LCO	U
8/14/00	03/04	LSP					148	5.6	LCO	U
8/14/00	03/04	LSP					147	6.1	LCO	U
8/14/00	03/04	LSP					155	7.1	LCO	U
8/14/00	03/04	LSP					125	3.6	LCO	U
8/14/00	03/04	LSP					152	5.8	LCO	U
8/14/00	03/04	LSP					106	2.0	LCO	U
8/14/00	03/04	LSP					137	4.6	LCO	U
8/14/00	03/04	LSP					151	6.2	LCO	U
8/14/00	03/04	LSP					160	7.6	LCO	U
8/14/00	03/04	LSP					158	6.5	LCO	U
8/14/00	03/04	LSP					143	5.0	LCO	U
8/14/00	03/04	LSP					135	4.5	LCO	U
8/14/00	03/04	LSP					109	2.4	LCO	U
8/14/00	03/04	LSP					109	2.8	LCO	U
8/14/00	03/04	LSP					152	6.1	LCO	U
8/14/00	03/04	LSP					137	4.2	LCO	U
8/14/00	03/04	LSP					142	4.8	LCO	U
8/14/00	03/04	LSP					87	1.3	LCO	U
8/14/00	03/04	LSP					94	1.4	LCO	U
8/14/00	03/04	LSP					145	5.1	LCO	U
8/14/00	03/04	LSP					156	6.2	LCO	U
8/14/00	03/04	LSP					85	1.1	LCO	U
8/14/00	03/04	LSP					103	1.8	LCO	U
8/14/00	03/04	LSP					--	--	--	U

*mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A1 (Con't). Red River Pacific Lamprey Length and Weight

Date Electrofished	Km/ Reach	Habitat Type	Area	Time	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans/ Untrans.
			Fished M ²	Fished (Min.)						
8/16/00	03/04	RBB	128.0	100	4	4	155	6.4	LCO	U
8/16/00	03/04	RBB					156	6.2	LCO	U
8/16/00	03/04	RBB					159	7.3	LCO	U
8/16/00	03/04	RBB					157	6.1	LCO	U
9/9/00	03/04	SCP	199.9	170	25	25	136	3.8	LCO	U
9/9/00	03/04	SCP					143	5.3	LCO	U
9/9/00	03/04	SCP					152	6.0	LCO	U
9/9/00	03/04	SCP					147	5.7	LCO	U
9/9/00	03/04	SCP					162	6.9	LCO	U
9/9/00	03/04	SCP					165	7.1	LCO	U
9/9/00	03/04	SCP					155	5.7	LCO	U
9/9/00	03/04	SCP					153	5.9	LCO	U
9/9/00	03/04	SCP					144	4.9	LCO	U
9/9/00	03/04	SCP					161	6.5	LCO	U
9/9/00	03/04	SCP					132	3.7	LCO	U
9/9/00	03/04	SCP					150	6.1	LCO	U
9/9/00	03/04	SCP					150	5.3	LCO	U
9/9/00	03/04	SCP					104	2.1	LCO	U
9/9/00	03/04	SCP					101	2.0	LCO	U
9/9/00	03/04	SCP					154	5.7	LCO	U
9/9/00	03/04	SCP					153	5.5	LCO	U
9/9/00	03/04	SCP					148	5.8	LCO	U
9/9/00	03/04	SCP					165	7.4	LCO	U
9/9/00	03/04	SCP					151	6.3	LCO	U
9/9/00	03/04	SCP					157	5.7	LCO	U
9/9/00	03/04	SCP					152	6.4	LCO	U
9/9/00	03/04	SCP					135	4.3	LCO	U
9/9/00	03/04	SCP					140	4.4	LCO	U
9/9/00	03/04	SCP					92	1.5	LCO	U
TOTAL			1238.8	841	154	151				

*mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A2. South Fork Clearwater River Pacific Lamprey Length and Weight

Date Electrofished	Km/ Reach	Habitat Type	Area	Time	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans./ Untrans.
			Fished M ²	Fished (Min)						
9/10/00	44.3/na	LSP	11.0	20	93	91	57	0.5	--	U
9/10/00	44.3/na	LSP					70	0.7	LCO	U
9/10/00	44.3/na	LSP					70	0.7	--	U
9/10/00	44.3/na	LSP					66	0.7	LCO	U
9/10/00	44.3/na	LSP					52	0.5	--	U
9/10/00	44.3/na	LSP					67	0.5	--	U
9/10/00	44.3/na	LSP					62	0.4	--	U
9/10/00	44.3/na	LSP					65	0.5	--	U
9/10/00	44.3/na	LSP					81	1.1	LCO	U
9/10/00	44.3/na	LSP					79	0.9	LCO	U
9/10/00	44.3/na	LSP					70	0.8	--	U
9/10/00	44.3/na	LSP					75	0.7	--	U
9/10/00	44.3/na	LSP					55	0.3	--	U
9/10/00	44.3/na	LSP					77	0.8	--	U
9/10/00	44.3/na	LSP					71	--	--	U
9/10/00	44.3/na	LSP					61	0.6	--	U
9/10/00	44.3/na	LSP					78	0.8	--	U
9/10/00	44.3/na	LSP					81	1.1	LCO	U
9/10/00	44.3/na	LSP					75	0.8	--	U
9/10/00	44.3/na	LSP					81	1.2	LCO	U
9/10/00	44.3/na	LSP					115	1.6	LCO	U
9/10/00	44.3/na	LSP					116	2.3	LCO	U
9/10/00	44.3/na	LSP					63	0.4	--	U
9/10/00	44.3/na	LSP					61	0.4	--	U
9/10/00	44.3/na	LSP					83	1.0	--	U
9/10/00	44.3/na	LSP					77	0.9	--	U
9/10/00	44.3/na	LSP					60	0.5	--	U
9/10/00	44.3/na	LSP					65	0.5	--	U
9/10/00	44.3/na	LSP					100	1.6	LCO	U
9/10/00	44.3/na	LSP					103	2.4	LCO	U
9/10/00	44.3/na	LSP					79	1.0	--	U
9/10/00	44.3/na	LSP					71	0.7	--	U
9/10/00	44.3/na	LSP					120	2.7	LCO	U
9/10/00	44.3/na	LSP					73	0.8	--	U
9/10/00	44.3/na	LSP					63	0.5	--	U
9/10/00	44.3/na	LSP					75	0.8	--	U
9/10/00	44.3/na	LSP					67	0.6	--	U
9/10/00	44.3/na	LSP					57	0.4	--	U
9/10/00	44.3/na	LSP					69	0.7	--	U
9/10/00	44.3/na	LSP					78	0.8	--	U
9/10/00	44.3/na	LSP					59	0.4	--	U
9/10/00	44.3/na	LSP					75	0.8	--	U
9/10/00	44.3/na	LSP					79	0.9	--	U

Mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A2. (Con't). South Fork Clearwater River Pacific Lamprey Length and Weight

Date	Km/ Electrofished	Habitat Type	Area Fished M ²	Time Fished (Min)	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans./ Untrans.
9/10/00	44.3/na	LSP					75	0.9	--	U
9/10/00	44.3/na	LSP					47	0.2	--	U
9/10/00	44.3/na	LSP					77	0.8	--	U
9/10/00	44.3/na	LSP					124	3.0	LCO	U
9/10/00	44.3/na	LSP					86	1.2	LCO	U
9/10/00	44.3/na	LSP					75	0.8	--	U
9/10/00	44.3/na	LSP					89	1.2	LCO	U
9/10/00	44.3/na	LSP					89	1.3	--	U
9/10/00	44.3/na	LSP					87	1.3	--	U
9/10/00	44.3/na	LSP					135	4.0	--	U
9/10/00	44.3/na	LSP					119	2.8	LCO	U
9/10/00	44.3/na	LSP					92	1.3	LCO	U
9/10/00	44.3/na	LSP					66	0.7	--	U
9/10/00	44.3/na	LSP					75	0.7	--	U
9/10/00	44.3/na	LSP					80	1.0	--	U
9/10/00	44.3/na	LSP					65	0.5	--	U
9/10/00	44.3/na	LSP					59	0.5	--	U
9/10/00	44.3/na	LSP					74	0.7	LCO	U
9/10/00	44.3/na	LSP					73	0.7	--	U
9/10/00	44.3/na	LSP					62	0.5	--	U
9/10/00	44.3/na	LSP					82	1.0	--	U
9/10/00	44.3/na	LSP					83	1.3	LCO	U
9/10/00	44.3/na	LSP					116	2.6	--	U
9/10/00	44.3/na	LSP					92	1.3	LCO	U
9/10/00	44.3/na	LSP					107	2.1	--	U
9/10/00	44.3/na	LSP					116	2.4	LCO	U
9/10/00	44.3/na	LSP					129	3.6	LCO	U
9/10/00	44.3/na	LSP					79	1.1	--	U
9/10/00	44.3/na	LSP					93	1.4	LCO	U
9/10/00	44.3/na	LSP					61	0.5	--	U
9/10/00	44.3/na	LSP					77	0.9	--	U
9/10/00	44.3/na	LSP					85	1.0	--	U
9/10/00	44.3/na	LSP					65	0.5	--	U
9/10/00	44.3/na	LSP					75	0.8	--	U
9/10/00	44.3/na	LSP					68	0.7	--	U
9/10/00	44.3/na	LSP					80	1.1	LCO	U
9/10/00	44.3/na	LSP					95	1.6	LCO	U
9/10/00	44.3/na	LSP					115	2.5	--	U
9/10/00	44.3/na	LSP					102	1.7	--	U
9/10/00	44.3/na	LSP					84	1.1	--	U
9/10/00	44.3/na	LSP					121	2.9	--	U
9/10/00	44.3/na	LSP					115	2.6	--	U
9/10/00	44.3/na	LSP					90	1.2	--	U
9/10/00	44.3/na	LSP					70	0.7	--	U

Mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A2. (Con't). South Fork Clearwater River Pacific Lamprey Length and Weight

Date	Km/ Electrofished	Habitat Type	Area Fished M ²	Time Fished (Min)	Lamprey Captured	Lamprey Measured	Length (mm)	Weight (g)	Recapture/ Marked*	Trans./ Untrans.
9/10/00	44.3/na	LSP					76	0.9	--	U
9/10/00	44.3/na	LSP					62	0.5	--	U
9/10/00	44.3/na	LSP					92	1.3	--	U
9/10/00	44.3/na	LSP					88	1.3	--	U
9/10/00	44.3/na	LSP					--	--	--	U
9/10/00	44.3/na	LSP					--	--	--	U
TOTALS			11.0	20	93	91				

Mark = LCO = left center orange

Trans = transformed

Untrans = untransformed

A3. Red River Habitat Surveys for Electrofished Units

RED RIVER SURVEYS OF UNITS ELECTROFISHED FOR LAMPREY

Sample Personnel: Claire, Peterson

		UNIT MEASUREMENTS								
HABITAT UNIT (Km from Mouth)	DATE	TIME	LAMPREY CAPTURED	Length (m)	Slope %	Max Depth (m)	Channel AVG (m)	Wetted AVG (m)	Area (m²)	
LSP	LATERAL SCOUR POOL									
	1.033 - 1.045	27-Jul-00	9:30 AM	3	13	0.5	0.8	na	6	78.0
	26.9084 - 26.9258	29-Jul-00	11:30 AM	0	17.4	0.5	0.5	10.5	8.8	153.1
	26.9258 - 26.9428	29-Jul-00	12:10 PM	0	17	0.25	0.7	8.8	7.0	119.0
	26.9513 - 26.9713	29-Jul-00	3:25 PM	0	20	0.5	0.8	8.4	6.6	132.0
	20.63 - 20.719	30-Jul-00	8:00 AM	0	33	2	0.75	10.9	13.1	432.3
	3.4 - 3.43	10-Aug-00	9:25 AM	95	30	0.5	0.9	13.0	10.1	303.0
	33.982- 34.0115	25-Aug-00	8:40 AM	0	29.5	0.25	0.55	7.1	6.1	180.0
SCP	STRAIGHT SCOUR POOL									
	3.482 - 3.498	9-Sep-00	10:00 AM	25	16	0.25	0.65	16.2	14.1	199.9
RIF	RIFFLE									
	1.033 - 1.045	12-Aug-00	9:30 AM	2	13	0.5	0.8	12.2	7.9	102.7
	26.882 - 26.9084	30-Jul-00	8:10 AM	0	26.4	0.75	0.25	11.2	10.3	271.9
	26.9428 - 26.9513	1-Aug-00	8:15 AM	0	8.5	1	0.35	10.4	8.0	68.0
	20.6 - 20.63	10-Aug-00	8:00 AM	0	30	1	0.4	8.5	7.3	219.0
RIP	RIFFLE WITH POCKETS									
	1.0 - 1.033	28-Jul-00	8:25 AM	3	28.5	1.25	0.75	18.3	14.8	421.8
RBB	RAPIDS WITH BOULDERS									
	1.0641 - 1.0681	15-Aug-00	9:00 AM	0	4	0.75	0.7	13.9	11.7	46.8
	3.43 - 3.444	16-Aug-00	8:35 AM	4	14	3	0.55	12.1	9.2	128.8
ALC	ALCOVE									
	0.99	12-Aug-00	1:00 PM	19	2.2	0	0.4		3.4	7.5

A3 (Con't). Red River Habitat Surveys for Electrofished Units

RED RIVER SURVEYS OF UNITS ELECTROFISHED FOR LAMPREY

Sample Personnel: Claire, Peterson

HABITAT UNIT (Km from Mouth)	WATER VELOCITIES (m/sec)***						STREAM SHADE				TOTAL
	Substrate		60% from substrate			75%	LEFT BNK	CNTR UP	CNTR DWN	RIGHT BNK	
	25%	50%	75%	25%	50%						
LSP											
1.033 - 1.045	0.287	0.491	0.454	*	0.500	0.456	0.0%	0.0%	0.0%	7.5%	7.5%
26.9084 - 26.9258	0.217	0.271	0.232	*	*	*	24.0%	0.0%	0.0%	10.5%	34.5%
26.9258 - 26.9428	0.130	0.141	0.194	0.388	0.359	0.272	0.0%	0.0%	0.0%	3.5%	3.5%
26.9513 - 26.9713	0.246	0.298	0.136	0.219	0.348	0.220	0.0%	7.5%	0.0%	15.0%	22.5%
20.63 - 20.719	0.054	0.152	0.126	0.152	0.253	*	10.5%	0.0%	0.0%	4.5%	15.0%
3.4 - 3.43	0.028	0.201	0.181	0.032	0.343	*	25.0%	1.5%	4.5%	13.5%	44.5%
33.982- 34.0115	0.201	0.048	0.308	*	0.362	*	0.0%	0.0%	6.0%	4.5%	10.5%
SCP											
3.482 - 3.498	0.574	0.047	0.180	*	0.349	*	9%	1.50%	3%	22.50%	36%
RIF											
1.033 - 1.045	0.287	0.491	0.454	*	0.500	0.456	0.0%	0.0%	0.0%	7.5%	7.5%
26.882 - 26.9084	0.468	0.772	0.088	*	*	*	18.0%	0.0%	0.0%	6.0%	24.0%
26.9428 - 26.9513	0.313	0.617	0.455	*	*	*	0.0%	0.0%	0.0%	3.0%	3.0%
20.6 - 20.63	0.474	0.833	0.531	*	*	*	12.0%	1.5%	3.0%	1.5%	18.0%
RIP											
1.0 - 1.033	0.397	0.378	0.545	*	*	*	12.0%	1.5%	0.0%	25.0%	38.5%
RBB											
1.0641 - 1.0681	0.633	0.840	0.264	*	*	*	Not taken				
3.43 - 3.444	0.297	0.065	0.610	*	*	*	10.5%	3.0%	4.5%	10.5%	28.5%
Alcove											
0.99	**	**	**				0%	7.5%	10.5%	15.0%	33.0%

* Water too shallow for velocity to be taken

** Flow less than lower limit of standard speed rotor

*** Substrate = ~ 5 cm above substrate 60% = 60% up water column percentages from left bank facing upstream (25%, stream center, 75%)

A3 (Con't). Red River Habitat Surveys for Electrofished Units

RED RIVER SURVEYS OF UNITS ELECTROFISHED FOR LAMPREY

Sample Personnel: Claire, Peterson

HABITAT UNIT (Km from Mouth)	SUBSTRATE COMPOSITION %									
	LG. BLDR	SM. BLDR	COBBLE COBBLE	COARSE GRVL.	MED. GRVL	FINE GRVL	COARSE SAND	FINE SAND	SILT/ ORG.	BED ROCK
LSP										
1.033 - 1.045	20%	23%	14.5%	12%	5%	5%	10%	10%	0.5%	
26.9084 - 26.9258	1%	25%	28%	15%	8%	8%	8%	5%	2%	
26.9258 - 26.9428	2%	35%	20%	10%	5%	11%	10%	5%	2%	
26.9513 - 26.9713	2%	15%	27%	16%	8%	8%	10%	5%	9%	
20.63 - 20.719	2%	4%	33%	20%	10%	10%	10%	3%	8%	
3.4 - 3.43	33%	15%	8%	7%	5%	15%	8%	7%	2%	
33.982- 34.0115	0%	12%	40%	5%	10%	15%	6%	8%	4%	
SCP										
3.482 - 3.498	12%	25%	20%	19%	5%	5%	5%	5%	1%	3%
RIF										
1.033 - 1.045	20%	23%	14.5%	12%	5%	5%	10%	10%	0.5%	
26.882 - 26.9084	0%	10%	40%	18%	5%	5%	10%	10%	2%	
26.9428 - 26.9513	0%	20%	35%	21%	5%	8%	5%	5%	1%	
20.6 - 20.63	0.5%	3%	41%	20%	12%	8%	8%	7%	0.5%	
RIP										
1.0 - 1.033	25%	30%	17%	7%	5%	5%	8%	2%	1%	
RBB										
1.0641 - 1.0681	37.5%	27%	15%	5%	5%	5%	3%	2%	0.5%	
3.43 - 3.444	35%	20%	12%	8%	5%	8%	7%	3%	2%	
Alcove										
0.99	20%	10%	2%	2%	3%	5%	5%	35%	18%	

APPENDIX B

EXPENDITURES
(As of December 31, 2000)

Personnel	\$25,711
Personnel Benefits	9,528

Subtotal Personnel	\$35,239
--------------------	----------

Supplies	\$13,314
Travel	200
Subcontractor	5,000
Indirect Charge	11,019

Subtotal Operating	\$29,533
--------------------	----------

Total	\$64,772
-------	----------