are necessary to solve the TOV equations with this new EoS and, then, obtain the relevant astrophysical features solely in terms of α_s and the beta function.

The pressure, in units of the pressure of a free gas, is shown in Figure 1 as a function of the quark chemical potential. There, we show the results up to first order in α_s and the one which includes the $O(\alpha_s^2)$ contribution. It is clear from this plot that the interactions between quarks can not be ignored, even in this large- μ region. Moreover, contrary to the case of finite-temperature perturbation theory [17], the series is reasonably well-behaved. Then, perturbation theory seems to be applicable for the pressure in the range of μ considered, but it does not imply that it works well for other observables as pointed out recently by Rajagopal and Shuster [18]. The remarkable feature in this figure is the fact that the pressure vanishes for $\mu = 0.767 \text{ GeV}$, so that the star is self-bound and represents a new class of solution at ultrahigh densities. This fact is emphasized in Figure 2. There one can see that the result for the case $\bar{\mu} = \mu$ represents a new branch in the EoS and can not be matched with hadronic equations of state [19]. In fact, there is a big gap separating the two regions.

FIG. 1. Total pressure (in units of the free gas pressure) as a function of the chemical potential for the case $\bar{\mu} = \mu$. We show the result up to first order in α_s and the one which includes the second-order contribution.

FIG. 2. Equation of state for the case $\bar{\mu} = \mu$. The free gas and a hadronic equation of state are displayed for comparison.

Figure 3 shows the total mass as a function of the central energy density for the $\bar{\mu} = \mu$ case. As mentioned before, extrema in such a plot signal gravitational instability and define a maximum allowed total mass for a stable star. For the case under consideration, the maximum mass is of the order of $0.35M_\odot$ and the central energy density can reach values almost a hundred times greater than ϵ_0 . These results should be compared to the ones obtained by using the

MIT bag model equation of state, namely $M_{max} \approx 2M_{\odot}$ and $\epsilon_c^{max} \approx 8\epsilon_0$.

FIG. 3. Total mass (in solar mass units) as a function of the central energy density, for the case $\bar{\mu} = \mu$.

Figure 4 displays the mass-radius relation for the cases $\bar{\mu} = \mu$, $\bar{\mu} = 2\mu$ and $\bar{\mu} = 3\mu$. They should be compared to the usual results for the maximum mass, quoted above, and for the maximum radius, $R_{max} \approx 11km$. One can see the strong dependence on the choice of the scale $\bar{\mu}$. In fact, for higher values of $\bar{\mu}$, one can reach the region that reproduces the results obtained by using the usual approach.

FIG. 4. Mass-radius relation for the cases $\bar{\mu} = \mu$, $\bar{\mu} = 2\mu$ and $\bar{\mu} = 3\mu$.

IV. CONCLUSION AND FUTURE WORK

Although the patterns of the curves for the mass-radius relation presented in Figure 4 resemble those obtained for strange stars within the MIT bag model approach, the phenomenology encountered in our treatment is very different. The quark stars which result from perturbative QCD are roughly five times smaller, they are less massive ($M \approx 0.3 - 0.9 M_{\odot}$ as compared to $M \approx 2 M_{\odot}$ for strange stars), and much denser. They represent an entirely new class of ultradense stars. The strong dependence on the choice of the scale $\bar{\mu}$, which must be chosen in a way that minimizes the contribution of higher-order logarithmic corrections [7], is of course unpleasant and is under current investigation [13]. However, it does not diminish the spectrum of exciting possibilities opened by the results provided by perturbative QCD. The possible existence of quark stars could also have considerable impact on the study of cold quark matter in connection with color superconductivity [20]. They might be detected in micro-lensing experiments.

In fact, the MACHO project has reported micro-lensing events for the Large Magellanic Cloud [21], interestingly with mass ranges close to our calculated quark star masses of $M = 0.15\text{--}0.9M_{\odot}$.

It is a pleasure to thank Larry McLerran, Tony Rebhan, Robert Harlander, Jonathan Lenaghan and André Peshier for fruitful discussions and illuminating remarks. This work was partially supported by the U.S. Department of Energy under Contract No. DE-AC02-98CH10886 and by CNPq (Brazil) through a post-doctoral fellowship.

-
- [1] The reader is referred to N. K. Glendenning, *Compact Stars — Nuclear Physics, Particle Physics, and General Relativity* (Springer, New York, 2000) for a more detailed account and a more complete list of references.
 - [2] R. C. Tolman, Phys. Rev. **55**, 364 (1939); J. R. Oppenheimer and G. M. Volkoff, *ibid.*, p. 374.
 - [3] A. Hewish *et al.*, Nature **217**, 709 (1968).
 - [4] M. Gell-Mann and Y. Neeman, *The Eightfold Way* (Benjamin, New York, 1964).
 - [5] D. D. Ivanenko and D. F. Kurdgelaidze, Astrophysics **1**, 251 (1965).
 - [6] J. C. Collins and M. J. Perry, Phys. Rev. Lett. **34**, 1353 (1975).
 - [7] B. Freedman and L. McLerran, Phys. Rev. D **17**, 1109 (1978).
 - [8] E. Witten, Phys. Rev. D **30**, 272 (1984). See also A. R. Bodmer, Phys. Rev. D **4**, 1601 (1971).
 - [9] E. Farhi and R. L. Jaffe, Phys. Rev. D **30**, 2379 (1984).
 - [10] P. Haensel, J. L. Zdunik, and R. Schaeffer, Astron. Astrophys. **160**, 121 (1986). C. Alcock, E. Farhi, and A. Olinto, Astrophys. J. **310**, 261 (1986).
 - [11] S. Weinberg, *Gravitation and Cosmology — Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972).
 - [12] R. D. Pisarski and D. H. Rischke, Phys. Rev. D **61**, 051501(R) (2000).
 - [13] E. S. Fraga, R. D. Pisarski and J. Schaffner-Bielich, work in progress.
 - [14] V. Baluni, Phys. Rev. D **17**, 2092 (1978).
 - [15] J. P. Blaizot, E. Iancu, and A. Rebhan, hep-ph/0005003 (2000).
 - [16] D. Groom *et al.*, Eur. Phys. J. **C15**, 1 (2000).
 - [17] C.-X. Zhai and B. Kastening, Phys. Rev. D **52**, (1995). E. Braaten and A. Nieto, Phys. Rev. Lett. **76**, 1417 (1996). J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. Lett. **83**, 2139 (1999).
 - [18] K. Rajagopal and E. Shuster, hep-ph/0004074 (2000).
 - [19] J. Schaffner and I. N. Mishustin, Phys. Rev. C **53**, 1416 (1996).
 - [20] M. Alford, K. Rajagopal, and F. Wilczek, Phys. Lett. B **422**, 247 (1998); Nucl. Phys. B **537**, 443 (1999). R. Rapp, T. Schäfer, E. V. Shuryak, and M. Velkovsky, Phys. Rev. Lett. **81**, 53 (1998). D. Page, M. Prakash, J. M. Lattimer, and A. Steiner, Phys. Rev. Lett. **85**, 2048 (2000). M. Alford, J. Berges, and K. Rajagopal, Nucl. Phys. B **571**, 269 (2000). M. Alford, J. A. Bowers and K. Rajagopal, hep-ph/0008208. R. Rapp, E. Shuryak and I. Zahed, hep-ph/0008207.
 - [21] C. Alcock *et al.*, astro-ph/0001272 (2000).