NUCLEAR ENERGY RESEARCH INITIATIVE (NERI) PROGRAM
ANNUAL TECHNICAL PROGRESS REPORT

Project Title: Development and Validation of Temperature Dependent Thermal Neutron Scattering Laws for Applications and Safety Implications in Generation IV Reactor Designs
Grant Number: DE-FG03-02SF22514/A000 (NERI Project 2001-140)
Principal Investigator: Ayman I. Hawari
Institution(s): North Carolina State University (NCSSU), Oak Ridge National Laboratory (ORNL), and Instituto Balseiro (IB)
Report Period: 9/1/2001 – 8/31/2002 (Phase I)

The overall objectives of this project are to critically review the currently used thermal neutron scattering laws for various moderators as a function of temperature, select a well documented and representative set of experimental data sensitive to the neutron spectra to generate a data base of benchmarks, update models and models parameters by introducing new developments in thermalization theory and condensed matter physics into various computational approaches in establishing the scattering laws, benchmark the results against the experimental set, and in the case of graphite, perform a validation experiment by observing neutron slowing down as a function of temperatures equal to or greater than room temperature, and understand the implications of the obtained results on the ability to accurately determine the operating and safety characteristics (e.g., the moderator temperature coefficient of reactivity, buildup and depletion of transuranium elements) of given reactor designs, which include generation IV nuclear power systems, advanced research reactors and power reactors for space applications.

MILESTONES AND TASK STATUS IN PHASE 1

The tasks for this phase are being performed in parallel with some overlap. The following table lists the tasks for this phase:

An update of project schedule and task status

Milestone/Task Description	Planned Completion Date	Actual Completion Date
(a) Evaluate models and data used in current calculations of scattering law for the proposed materials	March 31, 2002	Completed
(b) Evaluate improved new models and data relevant to the thermalization problem in neutron and condensed matter physics	May 31, 2002	Completed (will be revisited as the project progresses)
(c) Start the implementation of new computational tools to calculate with new models and/or data	May 31, 2002	Completed
(d) Evaluate experimental data available to define a set of benchmarks	March 31, 2002	near completion (will continue as the project progresses)
(e) Start the development of models to compute experimental benchmarks	August 31, 2002	near completion (will continue as the project progresses)
(f) Begin designing the graphite benchmark experiments that will be performed at ORELA	May 31, 2002	Completed
(g) Begin preparation of thermal neutron scattering libraries	August 31, 2002	Completed

NARRATIVE OF PROGRESS

Phase 1 of this project achieved several major milestones: First, the methodologies that can be used to calculate the thermal neutron scattering kernels were studied and implemented. As a result, the tools that are needed to perform the required analysis have been setup, tested, and are available to perform calculations for the materials of interest to this work. Second, specific scattering kernels were generated for graphite, Be, and BeO, which allowed comparisons to be made to previous methodologies and data sets. Third, the graphite benchmark experiment is in advanced design stages and construction of its heating system is expected to begin within the first quarter of phase 2, while the experimental preparations and preliminary experimental runs are to be performed during the third and fourth quarters of phase 2.

Tasks (a), (b), and (c):

During this phase, the research performed at NCSU, ORNL, and IB introduced new data and methods that can be used for evaluating the thermal neutron scattering kernels for moderators/reflectors that are of interest in nuclear technology. Primarily, our studies focused on establishing the appropriate methodologies and tools for evaluating and developing the thermal neutron scattering kernels. From the start, it became clear to us that the most efficient approach would be to establish and validate the main tools that would be used for this purpose, which then can be used to study the various materials. As a result, the following capabilities have been developed by the members of this project:

- 1) At ORNL, both the NJOY99/LEAPR and the GASKET methodologies (computer codes) are currently implemented. Both of these codes are capable of calculating the scattering laws, which are then used by the THERMR and ACER modules of NJOY to produce the cross section libraries. In addition, both codes were modified to eliminate numerical problems that produced inconsistencies that, in certain cases, can result in erroneous values for the cross sections. Subsequently, the codes were used to calculate the scattering kernel for graphite. The figure below illustrates the excellent agreement between LEAPR (marked by +) and GASKET (solid line) methodologies as a result of the above analysis. The work also focused on investigating the graphite phonon spectra that represent the starting point of the calculation of scattering kernels. In this case, two phonon spectra were utilized: the Young-Koppel phonon spectrum (J. Chem. Phys., 42, 1, 357, 1965) and the spectrum based on the work of Nicklow, Wakabayashi, and Smith (Phys. Rev. B, 5, 12, 1972).

The scattering law for graphite as a function of momentum and energy transfer. A comparison between GASKET and LEAPR results

- 2) At IB, the work focused on the application of a synthetic approach to construct the frequency spectra and subsequently utilize them in the inelastic kernel calculations. In the case of the elastic component, the approach was to approximate coherent elastic scattering by an incoherent calculation. This was justified based on the fact that incoherent like behavior will dominate in systems such as nuclear reactors. Finally, a special version of the NJOY code is being created that will also contain in it the GASKET methodology, which will facilitate the production of the scattering kernel using either approach. In this work, calculations were performed for graphite, beryllium and beryllium oxide. The figure below shows the results of the calculation for Be in BeO.

Inelastic and elastic thermal neutron scattering cross sections for Be in BeO at 296 K

- 3) At NCSU, The NJOY99 code system was set up and tested using the basic data of Young-Koppel and Nicklow, Wakabayashi, and Smith. Furthermore, we studied the use of ab initio physics methods to calculate the input needed by NJOY. Namely the phonon spectra that control thermal neutron interaction in crystalline materials. In this case, we implemented the VASP-PHONON code system. Currently we are able to generate the phonon spectrum using this system and, subsequently, use it in the NJOY system to produce the thermal neutron scattering kernel. The figures below show the phonon spectra for graphite based on the Young-Koppel (YK), Nicklow-Wakabayashi-Smith (ORNL), and ab initio (PH) models, and the thermal neutron scattering kernels for graphite.

The graphite phonon spectra obtained using the 3 methods described in the text.

Inelastic thermal neutron scattering cross section for graphite at 296 K and 1200 K. The curve labeled PH is based on ab initio models and is expected to change once better models are developed.

Tasks (d) and (e):

The initial intention during this phase was to evaluate a body of benchmarks for validating the scattering kernels. However, as the project progressed, it became clear to us that this effort can be most efficient when coordinated with the scattering law evaluation and validation studies for a specific material. Therefore, at this stage our combined effort has identified appropriate benchmark-quality experimental data for both graphite (3 benchmarks) and light water (the NIST leakage experiment). As reported previously, computational models were set up for the graphite benchmarks, and work is underway for the light water experiment. In addition, extensive literature searches were performed to identify experiments within the last 30 years that may be used as benchmarks. Sorting of these references is currently underway. Therefore this step will continue during phase 2.

Task (f):

The initial intention during this phase was to perform preliminary designs of the graphite benchmark experiment during this phase of the project. However, as the work progressed, it became clear that a near final design is needed during this phase to ensure the timely construction of the heating system that is needed to perform the experiment. Therefore, a major time investment was made to achieve that objective. To date, our design shows that a graphite rectangular pile that is 75-cm x 75-cm x 75-cm can produce the required results. In addition, we plan to use Pu-239 fission detectors, which are capable of producing easily interpretable time signatures. Currently, our calculations show that, below 1 eV, deviations on the order of 5% (in the detector response) exist between the LEAPR based inelastic scattering data that we generated during this work, and the standard GASKET based data that is contained in the MCNP libraries (as distributed by RSICC). The results of the calculations were also discussed with the potential vendor that can supply the heating system. We expect to finalize this work during the first quarter of Phase 2.

The figures below show the time dependent Pu-239 detector response at different temperatures and the variation in that response due to the variation in the cross section (LEAPR vs. GASKET generated cross sections).

The time spectra at a distance of 0.25 m from the pile surface and for temperatures of 300 K and 800 K. The peaks labeled 1, 2, and 3 correspond to resonance groups and/or resonances in the Pu-239 fission cross-section at approximately 40-300 eV, 8-30 eV, and 0.3 eV respectively

A comparison between the time dependent Pu-239 detector responses as predicted based on the new LEAPR data, and the standard MCNP data at 296 K.

Task (g):

Currently, all collaborators on this project are capable of generating the thermal neutron scattering libraries using the NJOY methodology. Therefore, this task is being implemented as the final step in a process that begins with task (c). Also, as mentioned above, modifications at ORNL have been made to the LEAPR module to eliminate numerical problems, and the GASKET methodology was implemented. At IB, the GASKET methodology is currently being incorporated into LEAPR. At NCSU, the ability to calculate basic phonon spectra was implemented using the VASP-PHONON code system. Using the various tools, thermal neutron scattering kernels were generated for graphite, beryllium, and beryllium oxide.

COST PERFORMANCE

During this phase, Dr. Ayman I. Hawari (PI on this project) relocated from the University of Cincinnati (UC) to assume the position of Associate Professor of Nuclear Engineering and Nuclear Reactor Program Director in the Department of Nuclear Engineering at North Carolina State University (NCSU). In addition, the two graduate students associated with this project relocated from UC to NCSU. The process of transferring the NERI grant from UC to NCSU was initiated in late December 2001 / early January 2002, and was completed in May, 2002. Once the contract was transferred, we initiated the hiring of a research associate, who will join the project team at NCSU. The position was offered to Dr. Victor H. Gillett (of Institute Balseiro, Argentina) and he plans to join our group around November 1, 2002. In addition, we have made the necessary preparations to begin equipment purchases in relation to the experimental work that will be performed during the fourth quarter of phase 2, and phase 3. The discrepancy between the actual and planned spending levels can be entirely accounted for by taking these expenditures (including staff benefits and university overhead costs) into account. Therefore, although the spending at NCSU during phase 1 was below the expected limits, we expect to adjust for that during phase 2.

At ORNL the spending from the account continues at its regular levels. The figure below shows the cumulative cost of the project for both institutions.

Project cost performance at the end of each month at NCSU and ORNL

PUBLICATIONS

To date, this project has resulted in the following accepted publications:

- 1) Ayman I. Hawari, Iyad Al-Qasir, Bernard W. Wehring, Felix C. Difilippo, and John-Paul Renier, "INVESTIGATION OF NEUTRON THERMALIZATION IN GRAPHITE MODERATORS," accepted to ANS winter meeting, Washington DC, 2002.
- 2) Ayman I. Hawari, Tong Zhou, Bernard W. Wehring, Felix C. Difilippo, and John-Paul Renier, "EXPERIMENT FOR BENCHMARKING THE THERMAL NEUTRON SCATTERING LAW IN GRAPHITE AT THE ORELA FACILITY," accepted to ANS winter meeting, Washington DC, 2002.
- 3) Felix C. Difilippo, John-Paul Renier, and Ayman I. Hawari, "BENCHMARKS FOR THE SCATTERING KERNEL OF GRAPHITE," accepted to the PHYSOR meeting, Seoul, South Korea, 2002.