NUCLEAR DATA AND MEASUREMENTS SERIES

ANL/NDM-152

Technical note on:-

DISPERSION CONTRIBUTIONS TO NEUTRON REACTIONS

by

R. D. Lawson and A. B. Smith

April, 2001

ARGONNE NATIONAL LABORATORY, ARGONNE, ILLINOIS

Operated by THE UNIVERSITY OF CHICAGO

for the U. S. DEPARTMENT OF ENERGY

under Contract W-31-109-Eng-38

Argonne National Laboratory, with facilities in the states of Illinois and Idaho, is owned by the United States Government, and operated by the University of Chicago under provisions of a contract with the Department of Energy.

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government, nor The University of Chicago, nor any agency thereof, nor any of their employees or officers, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, Argonne National Laboratory, or The University of Chicago.

Available electronically at <http://www.doe.gov/bridge>

Available for a processing fee to U. S. Department of Energy and its contractors, in paper, from:

U. S. Department of Energy
Office of Scientific and TEchnical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
phone: (865) 576-8401
fax: (865) 576-5728
email: reports@adonis.osti.gov

Nuclear Data and Measurement Series

Reports in the Argonne National Laboratory Nuclear Data and Measurement Series present results of studies in the field of microscopic nuclear data. The primary objective of the series is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the compilation and evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication, but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

PUBLICATIONS IN THE ANL/NDM SERIES

A listing of recent issues in this series is given below. Issues and/or titles prior to ANL/NDM-130 can be obtained from the National Technical Information Service, U. S. Department of Commerce, National Technical Information Service, Technology Administration, Springfield, VA 22161, or by contacting the author of this report at the following address:-

Technology Development Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, IL 60439
USA

In addition, information on the ANL/NDM report Series can be found on the Internet at the following URL:
<http://www.td.anl.gov/reports/ANLNDMReports.html>. This Internet address contains complete textual and tabular information.

A. B. SMITH AND P. T. GUENTHER

Fast-Neutron Interaction with the Fission Product ^{103}Rh
ANL/NDM-130, September 1993

A. B. SMITH AND P. T. GUENTHER

Fast-Neutron Scattering from Vibrational Palladium Nuclei
ANL/NDM-131, October 1993

A. B. SMITH

Neutron Interaction with Doubly-Magic ^{40}Ca
ANL/NDM-132, December 1993

A. B. SMITH

Neutron Scattering at $Z = 50$:- Tin
ANL/NDM-133, September 1994

A. B. SMITH, S. CHIBA AND J. W. MEADOWS

An Evaluated Neutronic File for Elemental Zirconium
ANL/NDM-134, September 1994

A. B. SMITH

Neutron Scattering from Uranium and Thorium
ANL/NDM-135, February 1995

A. B. SMITH

Neutron Scattering and Models:- Iron
ANL/NDM-136, August 1995

A. B. SMITH

Neutron Scattering and Models:- Silver
ANL/NDM-137, July 1996

A. B. SMITH

Neutron Scattering and Models:- Chromium
ANL/NDM-138, June 1996

W. P. POENITZ AND S. E. AUMEIER

The Simultaneous Evaluation of the Standards and Other Cross Sections of Importance for Technology
ANL/NDM-139, September 1997

- J. T. Daly and D. L. Smith
A Compilation of Information on the $^{31}\text{P}(p,\gamma)^{32}\text{S}$ Reaction and Properties of Excited Levels in ^{32}S
 ANL/NDM-140, November 1997
- A. B. SMITH
Neutron Scattering and Models:- Titanium
 ANL/NDM-141, July 1997
- A. B. SMITH
Neutron Scattering and Models:- Molybdenum
 ANL/NDM-142, July 1999
- R. E. MILLER AND D. L. SMITH
A Compilation of Information on the $^{31}\text{P}(p,\gamma)^{33}\text{Cl}$ Reaction and Properties of Excited Levels in ^{33}Cl
 ANL/NDM-143, July 1997
- R. E. MILLER AND D. L. SMITH
A Compilation of Information on the $^{31}\text{P}(p,\gamma)^{28}\text{Si}$ Reaction and Properties of Excited Levels in ^{28}Si
 ANL/NDM-144, November 1977.
- R. D. LAWSON AND A. B. SMITH
Abares-A Neutron Spherical Optical-Statistical Model Code: A Users Manual
 ANL/NDM-145, June 1998.
- R. T. KLANN AND W. P. POENITZ
Non-destructive Assay of EBR-II Blanket Elements Using Resonance Transmission Analysis
 ANL/NDM-146, August 1998.
- A. B. SMITH
ELEMENTAL ABAREX -- A User's Manual
 ANL/NDM-147, June 1999.
- D. G. NABEREJNEV AND D. L. SMITH
A Method to Construct Covariance Files in ENDF/B Format for Criticality Safety Applications
 ANL/NDM-148, June 1999.
- A. B. Smith
Neutrons and Antimony:- Physical Measurements and Interpretations
 ANL/NDM-149, July 2000
- A. B. Smith
Neutrons and Antimony:- Neutronic Evaluations of ^{121}Sb and ^{123}Sb
 ANL/NDM-150, July 2000.
- A. B. Smith
Fast-neutrons Incident on Holmium
 ANL/NDM-151, December 2000.

Technical note on:-

DISPERSION CONTRIBUTIONS TO NEUTRON REACTIONS^{*}

by

R. D. Lawson and A. B. Smith
Argonne National Laboratory
Argonne, Illinois

April, 2001

Keywords:

Simple formulation of dispersion effects in neutron reactions.

^{*}This work supported by the United States Department of Energy
under contract W-31-109-Eng-38.

TABLE OF CONTENTS

Text -----	1
References -----	6
Table 1 -----	7
Figure 1 -----	9
Figure 2 -----	10
Figure 3 -----	11
Appendix, FORTRAN Source listing -----	12

Technical Note On:-

DISPERSION CONTRIBUTIONS TO NEUTRON REACTIONS

It is well known that there is a dispersion relationship linking the real and imaginary optical-model potentials and reflecting causality ([MNS86], [Hod94], [Sat83]). This relationship can be expressed as

$$J(E) = J_{hf}(E) + \frac{P}{\pi} \int_{-\infty}^{+\infty} \frac{J_w(E')}{E - E'} dE', \quad (1)$$

where $J(E)$ is the strength of the total real potential, $J_{hf}(E)$ is the strength of the local-equivalent Hartree-Fock interaction, $J_w(E)$ is the strength of the imaginary potential, and "P" denotes the principal-value integral (herein, unless otherwise stated, all potential strengths are expressed as volume-integrals per nucleon). The integral of Eq. 1 can be broken into two parts, a surface component denoted by $\Delta J_{sur}(E)$ and a volume component given by $\Delta J_{vol}(E)$, where

$$\Delta J_{sur}(E) = \frac{P}{\pi} \int_{-\infty}^{+\infty} \frac{J_{sur}(E')}{E - E'} dE' \quad (2)$$

and

$$\Delta J_{vol}(E) = \frac{P}{\pi} \int_{-\infty}^{+\infty} \frac{J_{vol}(E')}{E - E'} dE', \quad (3)$$

and where $J_{sur}(E')$ and $J_{vol}(E')$ are surface- and volume-imaginary potential strengths, respectively. Clearly, $J(E)$ then becomes

$$J(E) = J_{hf}(E) + \Delta J_{vol}(E) + \Delta J_{sur}(E) \quad (4)$$

and

$$J(E) = J_{eff}(E) + \Delta J_{sur}(E), \quad (5)$$

where $J_{eff}(E)$ is obviously the sum of the first two terms of Eq. 4. Thus, in order to self-consistently interpret neutron-reaction data at energies $\lesssim 30$ MeV one must have some way of evaluating $\Delta J_{sur}(E)$ and $\Delta J_{vol}(E)$. Complex approximations can be found in the literature, involving a number of assumptions. What is given here is an approach roughly outlined approximately a decade ago [LGS87]. It is

simple, effective, and has been used a number of times by the Argonne Group. With reasonable assumptions as to potential geometric forms, the concepts described here can be easily implemented using the spherical optical-model code ABAREX ([LS98], [Smi99]) and the coupled-channels-model code ECIS96 [Ray96]. It is a proven and useful method which is documented here to assure that it does not vanish into the mists of time.

In the following paragraphs the basic assumptions, the algebraic manipulations and the resulting formulas are set forth, and an example given. There follows a short FORTRAN program which will calculate the integrals of Eqs. 2 and 3, using reasonable formulations of the imaginary potential. The net qualitative effect of the dispersion relation on neutron-induced reactions at energies of $\approx < 30$ MeV is to add a fraction of both the surface- and volume-imaginary potentials to the real potential. The attached FORTRAN code will calculate both the total and surface fraction of these contributions as a function of energy. The ABAREX optical-model program will directly utilize the surface component in a wide range of neutron calculations using the keyword DISP" as discussed in refs. [LS98], and [Smi99]. The surface, volume and total integrals of Eqs. 1, 2 and 3 are also provided by the program and can be applied to a wide range of dispersive considerations.

The present procedure assumes the energy dependence of the imaginary potential strength illustrated in Fig. 1. It was assumed that the imaginary potential was symmetric about the Fermi Energy ($-E_f$) [JLM77], (herein the absolute value of the Fermi Energy is used). It was further assumed that from $-E_f \rightarrow 0$ the energy dependence of the surface-imaginary potential follows a parabolic behavior [JLM77], and concurrently the volume-imaginary potential is zero. From zero energy to some energy E_n the imaginary-potential strength was taken to be entirely a surface term and to increase linearly with energy as more channels open. Experimental analysis indicates that E_n is probably in the $20 \rightarrow 30$ MeV range, although some experimental interpretations extend the imaginary volume absorption to lower energies [WG86]. From E_n the surface-imaginary strength was assumed to fall linearly to zero at some energy E_v . E_v is uncertain but is reasonably in the range of $50 \rightarrow$ several-hundred MeV. Concurrently, the volume absorption rises linearly from zero at E_n to J_v at E_v . Beyond E_v the J_v was assumed to be constant to infinity. These assumptions are quite reasonable and supported by experimental evidence. However, other linear or non-linear representations can be used, if the numerical formalism is modified accordingly.

Consider the integral in Eq. 2 over the energy range x where $-2E_f \leq x \leq 0$ of Fig. 1. This range consists of sections "A" and "B" of Fig. 1. In this energy range the parabolic behavior of J_{sur} is given in Table 1. This behavior implies that the integral of Eq. 2 in regions A or B of Fig. 1 is proportional to $I(E)$ where

$$\begin{aligned} I(E) &= P \int \frac{(x+E_f)^2}{(x-E)} \cdot dx \\ &= P \int \frac{(x^2 + 2xE_f + E_f^2)}{(x-E)} \cdot dx \\ &= P \int \frac{x(x-E) + (x-E)(2E_f+E) + (E+E_f)^2}{(x-E)} \cdot dx. \end{aligned} \quad (5)$$

The only term in this evaluation that can lead to trouble is the logarithmic integral involving the term proportional to $(E+E_f)^2$ and this occurs when E lies in the range of the integration. In particular, consider the interval $-E_f \leq x \leq 0$. If E does not lie in this region then

$$P \int_{-E_f}^0 \frac{dx}{(x-E)} = \ln(x-E) \Big|_{-E_f}^0 = \ln \left[\frac{E}{(E+E_f)} \right]. \quad (6)$$

On the other hand, if E does lie in the integration range then

$$\begin{aligned} P \int_{-E_f}^0 \frac{dx}{(x-E)} &= \lim_{\epsilon \rightarrow 0} \left[\int_{-E_f}^{E-\epsilon} \frac{dx}{(x-E)} + \int_{E+\epsilon}^0 \frac{dx}{(x-E)} \right] \\ &= \lim_{\epsilon \rightarrow 0} \left[\ln \left(\frac{-\epsilon}{-E_f-E} \right) + \ln \left(\frac{-E}{-\epsilon} \right) \right] \\ &= \lim_{\epsilon \rightarrow 0} \ln \left[\left(\frac{\epsilon}{+E_f+E} \right) \left(\frac{-E}{-\epsilon} \right) \right] \\ &= \ln \left[\frac{-E}{E_f+E} \right]. \end{aligned} \quad (7)$$

But E , by supposition, is a negative number between 0 and $-E_f$ so that the argument of the logarithm is positive and the only effect of the principal-value condition is that for any value of E , either inside or outside the range of integration, this term becomes $\ln \left| \frac{E}{E+E_f} \right|$, where $|\dots|$ stands for the absolute value of the enclosed quantity. The value of this integral in

regions A and B of Fig. 1 is given by $\Delta J_{\text{sur}}(E)$ in Table 1.

The contributions to $\Delta J_{\text{sur}}(E)$ and $\Delta J_{\text{vol}}(E)$ in the other regions shown in Fig. 1 can be evaluated by similar methods, and the corresponding analytic expressions are given in Table 1.

In order to find the total contribution the imaginary potential makes to the real optical-model potential one must sum up the entries shown for the various regions, as given in Table 1. These include the various volume contributions, $\Delta J_{\text{vol}}(E)$. In particular, the volume contributions in the "G" and "H" regions are evaluated in the limit $c \rightarrow \infty$. Thus, "G" plus "H" are

$$\begin{aligned}\Delta J_{\text{vol}}(E)(\text{regions G+H}) &= \frac{J_v}{\pi} \lim_{c \rightarrow \infty} \left[\ln \left| \frac{c-E}{E_v-E} \right| + \ln \left| \frac{2E_f+E_v+E}{c+E} \right| \right] \\ &= \frac{J_v}{\pi} \lim_{c \rightarrow \infty} \left[\ln \left| \frac{c-E}{E_v-E} \right| \cdot \left| \frac{2E_f+E_v+E}{c+E} \right| \right] \\ &= \frac{J_v}{\pi} \ln \left| \frac{2E_f+E_v+E}{E_v-E} \right|. \quad (8)\end{aligned}$$

Therefore, $\Delta J_{\text{vol}}(E)$, Eq. 3, becomes

$$\begin{aligned}\Delta J_{\text{vol}}(E) &= \frac{J_v}{\pi(E_n-E_v)} \left[(E_n-E) \ln \left| \frac{E_v-E}{E_n-E} \right| + (2E_f+E_n+E) \ln \left| \frac{2E_f+E_n+E}{2E_f+E_v+E} \right| \right] \\ &+ \\ &\quad (E_n-E_v) \ln \left| \frac{2E_f+E_v+E}{E_v-E} \right| \quad (9)\end{aligned}$$

In practice, since J_{hf} in Eq. 1 is not well known, one has the option of assuming the real optical-model potential emerging from a fit to neutron reaction data is already the sum of $J_{\text{hf}}(E)$ and $\Delta J_{\text{vol}}(E)$. On the other hand, since the radius and diffuseness of the volume absorption may be different than those of the real potential, if one wants to be self consistent one should iterate on the problem.

One can calculate the total real surface potential, $\Delta J_{\text{sur}}(E)$ of Eq. 2, by summing the contributions to $\Delta J_{\text{sur}}(E)$ given in Sections A through F of Table 1. When the keyword "DISP", and associated ratio numerical values calculated by the program of the Appendix, is included in the input of the spherical optical-statistical model code ABAREX ([LS98],[Smi99]) the

contribution of $\Delta J_{\text{sur}}(E)$, discussed above, will automatically be calculated. ΔJ_{sur} and ΔJ_{vol} can also be employed in applications by using well known equations [Elt58] to convert strengths in volume-integrals-per-nucleon to potential magnitudes, assuming geometric shapes of the potentials (e.g. Saxon-Wood real form and Saxon-Wood-derivative imaginary form). The same results can be obtained with ABAREX and some other codes (e.g. ECIS96 [Ray96]) if the input parameters are suitably arranged.

It has been repeatedly shown that surface dispersive effects in neutron-induced reactions at lower energies (e.g., $\approx < 30$ MeV) have a substantive impact on the processes and should be included in the calculations. There is also a volume contribution due to dispersive effects as set forth in Eq. 3 and Table 1. It can be calculated as illustrated in the example given below. However, as pointed out above, it can not be experimentally separated from the energy dependence of the local-equivalent Hartree-Fock potential or from the possible energy dependence of the nucleon-nucleon force.

The appendix to this Note is a small FORTRAN program which will evaluate the above equations and derive the "DISP" ratio for ABAREX or other model calculations. Certainly, other methods of evaluating the equations of Table 1 are possible. In following such alternate courses attention must be given to the poles of the various equations of Table 1. The results obtained with the program of the appendix are illustrated in Figs. 2 and 3 using a representative experimentally-derived potential for ^{121}Sb as described in ref. [Smi00]. The illustrative calculations assumed a Saxon-Woods-derivative surface-imaginary potential strength given by

$$J_w(E) = 45.63 + 1.797 \cdot E \quad (\text{MeV} \cdot \text{fm}^3) \quad (10)$$

to an E_n of 25 MeV. Above 25 MeV it was assumed that the surface component of the imaginary potential linearly fell to zero at 60 MeV. Concurrently, the volume absorption strength linearly increased from zero at 25 MeV to $75 \text{ MeV} \cdot \text{fm}^3$ at 60 MeV and then remained constant to higher energies. This behavior is the same as that schematically illustrated in Fig. 1. The Fermi Energy was taken to be $|7.86|$ MeV. The ΔJ_{sur} of Eq. 2 and the ΔJ_{vol} of Eq. 3 are indicated in Fig. 2. ΔJ_{vol} of Fig. 2 illustrates its approximately linear energy dependence, as cited above. Using ΔJ_{tot} of Fig. 2, one can obtain the fraction of the total imaginary strength that is added to the real potential due to dispersion effects, as illustrated by the ΔJ_{tot} curve of Fig. 3. This contribution will include both a

surface and volume component. ΔJ_{sur} of Fig. 2 can be used to determine the fraction of the surface-imaginary strength that contributes to the real potential as a result of dispersion effects (value of the "DISP" input of ABAREX). That is shown by the ΔJ_{sur} curve of Fig. 3. The difference between ΔJ_{tot} and ΔJ_{sur} of Fig. 3 reflects the contribution of the volume imaginary effects, particularly at larger energies. Of course, the dispersive ratios ("DISP" of ABAREX) are defined where the respective potential strengths are non-zero, i.e. in the denominator of the ratio. Somewhat different results will be obtained with alternative initial assumptions and/or imaginary potential strengths. However, the qualitative behavior of the results is approximately the same at energies of ≤ 30 MeV. The relatively small quantitative differences will not greatly effect the subsequent model predictions. In principle, the calculational procedures should be pursued in an iterative manner; first determining the model, calculating the dispersive corrections, and then redetermining the model, etc., until the model parameters reasonably converge. In practice one iteration is usually sufficient for acceptable accuracy.

References

- [Elt58] Elton L., Rev. Mod. Phys. 30 (1958) 557.
- [Hod94] Hodgson P. E. The nucleon optical model (World Science, 1994).
- [JLM77] Jeukenne J., Lejeune A. and Mahaux C., Phys. Rev. C16 (1977) 80.
- [LGS87] Lawson R., Guenther P. and Smith A., Phys. Rev. C36 (1987) 1298.
- [LS98] Lawson R. and Smith A., Argonne National Laboratory Report ANL/NDM-145 (1998).
- [MNS86] Mahaux C., Ngo H. and Satchler G., Nucl. Phys. A449 (1986) 354.
- [Ray96] Raynal J., The coupled-channels code ECIS96, private communication 1996; see also CEA Report, CEA-N-2772 (1994).
- [Sat83] Satchler G., Direct nuclear reactions (Clarendon, Oxford, 1983)
- [Smi99] Smith A., Argonne National Laboratory Report ANL/NDM-147 (1999).
- [Smi00] Smith A., Argonne National Laboratory Report ANL/NDM-149 (1999).
- [WG86] Walter R. and Guss P., Proc. conf. on nucl. data for basic and applied sci. Eds. P. Young et al., vol.-2, p-277 (Gordon and Breach, New York (1986)).

Argonne National Laboratory
April, 2001

Table 1. Parameterizations of dispersion contributions in the eight energy regions of Fig. 1.

Integral, Region A

$$J_{\text{sur}}(x) = \frac{J_0}{E_f^2} (x + E_f)^2$$

$$\Delta J_{\text{sur}}(E) = \frac{J_0}{E_f^2} \frac{1}{\pi} \left[-\frac{E_f^2}{2} + E_f(2E_f + E) + (E + E_f)^2 \ln |E/(E_f + E)| \right]$$

$$J_{\text{vol}}(x) = 0$$

$$\Delta J_{\text{vol}}(E) = 0$$

Integral, Region B

$$J_{\text{sur}}(E) = \frac{J_0}{E_f^2} (x + E_f)^2$$

$$\Delta J_{\text{sur}}(E) = \frac{J_0}{E_f^2} \frac{1}{\pi} \left[\frac{E_f^2}{2} + E_f E + (E + E_f)^2 \ln |(E_f + E)/(2E_f + E)| \right]$$

$$J_{\text{vol}}(x) = 0$$

$$\Delta J_{\text{vol}}(E) = 0$$

Integral, Region C

$$J_{\text{sur}}(x) = \frac{1}{E_n} \left[J_0 E_n + (J_n - J_0)x \right]$$

$$\Delta J_{\text{sur}}(E) = \frac{1}{\pi} \left[(J_n - J_0) + \left(J_0 + \frac{(J_n - J_0)E}{E_n} \right) \ln \left| \frac{E - E_n}{E} \right| \right]$$

$$J_{\text{vol}}(x) = 0$$

$$\Delta J_{\text{vol}}(E) = 0$$

Integral, Region D

$$J_{\text{sur}}(x) = \frac{1}{E_n} \left[J_0 E_n + (J_0 - J_n)(x + 2E_f) \right]$$

$$\Delta J_{\text{sur}}(E) = \frac{1}{E_n} \frac{1}{\pi} \left[(J_0 - J_n)E_n + \left[J_0 E_n + (2E_f + E)(J_0 - J_n) \right] \ln \left| \frac{2E_f + E}{2E_f + E + E_n} \right| \right]$$

$$J_{\text{vol}}(x) = 0$$

$$\Delta J_{\text{vol}}(E) = 0$$

Integral, Region E

$$J_{\text{sur}}(x) = \frac{J_n}{E_n - E_v} (-E_v + x)$$

$$\Delta J_{\text{sur}}(E) = \frac{J_n}{(E_n - E_v)} \frac{1}{\pi} \left[(E_v - E_n) + (E - E_v) \ln \left| \frac{E_v - E}{E_n - E} \right| \right]$$

$$J_{\text{vol}}(x) = \frac{J_v}{E_n - E_v} (E_n - x)$$

$$\Delta J_{\text{vol}}(E) = \frac{J_v}{(E_n - E_v)} \frac{1}{\pi} \left[(E_n - E_v) + (E_n - E) \ln \left| \frac{E_v - E}{E_n - E} \right| \right]$$

Integral, Region F

$$J_{\text{sur}}(x) = \frac{J_n}{E_v - E_n} (2E_f + E_v + x)$$

$$\Delta J_{\text{sur}}(E) = \frac{J_n}{(E_v - E_n)} \frac{1}{\pi} \left[(E_v - E_n) + (2E_f + E_v + E) \ln \left| \frac{2E_f + E_n + E}{2E_f + E_v + E} \right| \right]$$

$$J_{\text{vol}}(x) = \frac{J_v}{E_n - E_v} (2E_f + E_n + x)$$

$$\Delta J_{\text{vol}}(E) = \frac{J_v}{(E_n - E_v)} \frac{1}{\pi} \left[(E_v - E_n) + (2E_f + E_n + E) \ln \left| \frac{2E_f + E_n + E}{2E_f + E_v + E} \right| \right]$$

Integral, Region G

$$J_{\text{sur}}(x) = 0$$

$$\Delta J_{\text{sur}}(E) = 0$$

$$J_{\text{vol}}(x) = J_v$$

$$\Delta J_{\text{vol}}(E) = \frac{J_v}{\pi} \lim_{c \rightarrow \infty} \left[\ln \left| \frac{c - E}{E_v - E} \right| \right]$$

Integral, Region H

$$J_{\text{sur}}(x) = 0$$

$$\Delta J_{\text{sur}}(E) = 0$$

$$J_{\text{vol}}(x) = J_v$$

$$\Delta J_{\text{vol}}(E) = \frac{J_v}{\pi} \lim_{c \rightarrow \infty} \left[\ln \left| \frac{2E_f + E_v + E}{c - E} \right| \right]$$

In the limit $c \rightarrow \infty$

$$\Delta J_{\text{vol}}(E) \text{ for regions G+H} = \frac{J_v}{\pi} \ln \left| \frac{2E_f + E_v + E}{E_v - E} \right|$$

Fig. 1. Schematic diagram of the imaginary-potential energy dependence, as described in the text. "0" symbols indicate surface components of the potential, and "+" symbols the volume components.

Fig. 2. Dispersive integral contributions as discussed in the text. ΔJ_{sur} , ΔJ_{vol} and ΔJ_{tot} refer to surface, volume and total dispersive contributions, as discussed in the text.

Fig. 3. Dispersive fraction as discussed in the text. ΔJ_{sur} and ΔJ_{tot} denote the surface and total dispersive-ratio contributions.

Appendix, FORTRAN source listing

```

CCCCC  CALCULATION OF DISPERSION CONTRIBUTIONS FROM R. LAWSON FORMULAS.      1
CCCCD  OUTPUT DIRECTLY APPLICABLE TO ABAREX.                                2
CCCCC  SIMPLE FORMULATION DESCRIBED IN TECHNICAL MEMO ABOVE.                3
CCCCC  PROVIDES TOTAL, SURFACE AND VOLUME INTEGRAL                          4
CCCCC  OUTPUT ON FILE "OUT".                                                5
CCCCC  RUNS ON SIMPLE PC WITH LAHEY COMPILER.                               6
CCCCC  ALSO RUNS ON VAX WORK STATION AND VAX CLUSTER.                       7
CCCCC  SHOULD RUN ON ANY MODEST FACILITY WITH A FORTRAN-77 COMPILER.        8
COMMON EF,E,EX,DELTA,EN,EV,FE
DIMENSION H(3,1500),X(4,1500)
C      DIMENSIONED FOR 500 MESH POINTS
C      IF MORE MESH POINTS ARE DESIRED ENLARGE THE ABOVE DIMENSION
C      STATEMENT AND INCREASE 'NUM' (BELOW) CORRESPONDINGLY.
NUM=1500
C      ALL ENERGIES ARE IN MEV. ALL STRENGTHS (J) ARE VOLUME-INTEGRALS-
C      PER-NUCLEON IN UNITS OF MEV-FM**3,
1      FORMAT(' EF, ABS VALUE OF FERMI ENERGY (MEV), ZERO EXIT, (F12.6)')
2      FORMAT(F12.6)
3      FORMAT(' EN OF FIG. 1 OF TECH. NOTE IN MEV, (F12.6)')
4      FORMAT(' EV OF FIG. 1 OF TECH. NOTE IN MEV, (F12.6)')
5      FORMAT(' JO OF FIG. 1 OF TECH. NOTE IN MEV-FM**3, (F12.6)')
6      FORMAT(' JN OF FIG. 1 OF TECH. NOTE IN MEV-FM**3, (F12.6)')
7      FORMAT(' JV OF FIG. 1 OF TECH. NOTE IN MEV-FM**3, (F12.6)')
8      FORMAT(' E-START OF CALCULATION IN MEV, (F12.6)')
9      FORMAT(' E-END OF CALCULATION IN MEV, (F12.6)')
C      E-START SHOULD BE LESS THEN E-END
10     FORMAT(' DE, ENERGY INCREMENT OF THE CALCULATION IN MEV, (F12.6)')
11     FORMAT(' TOTAL DISP ', 'EF,JO,JN,JV',4(2X,F8.3))
20     FORMAT(' SURFACE DISP ', 'EF,JO,JN,JV',4(2X,F8.3))
13     FORMAT(' TOTAL INTEGRAL')
15     FORMAT(10E10.3)
16     FORMAT(1H )
17     FORMAT(' DELTA FOR POLES IN MEV, TYPICALLY 0.05 (F12.6)')
18     FORMAT(' SURFACE INTEGRAL')
19     FORMAT(' VOLUME INTEGRAL')
OPEN(10,FILE='OUT',STATUS='NEW')
100    CONTINUE
NJ=1
WRITE(*,1)
READ(*,2)EF
IF(EF.EQ.0.0)stop
WRITE(*,3)
READ(*,2)EN
WRITE(*,4)
READ(*,2)EV
WRITE(*,5)
READ(*,2)WJO
WRITE(*,6)
READ(*,2)WJN
SL=(WJN-WJO)/EN
WRITE(*,7)
READ(*,2)WJV
WRITE(*,17)
READ(*,2)DELTA
WRITE(*,8)
READ(*,2)ESTART
WRITE(*,9)
READ(*,2)ESTOP
WRITE(*,10)
READ(*,2)DE
SLMAX=(WJV-WJN)/(EV-EN)
DO 200 JJ=1,NUM
E=ESTART+FLOAT(JJ-1)*DE
EE=ABS(E+EF)
IF(EE.LE.EF)FE=WJO*EE**2/EF**2
IF(EE.GT.EF.AND.EE.LE.(EN+EF))FE=WJO+SL*(EE-EF)

```

```

IF (EE.GT.(EN+EF).AND.EE.LE.(EV+EF)) FE=WJN+SLMAX*(EE-EF-EN)
IF (EE.GT.(EV+EF)) FE=WJV
IF (ABS (FE).LT.1.E-6) FE=1.E-6
CALL CHECK
A=(WJ0/((EF**2)*3.14159))*(-.5*EF**2+EF*(2.*EF+EX)+(EX+EF)**2)*
&ALOG (ABS (EX/(EF+EX)))
B=(WJ0/((EF**2)*3.14159))*(.5*EF**2+EF*EX+(EX+EF)**2)*
&ALOG (ABS ((EF+EX)/(2.*EF+EX)))
C=(1./3.14159)*((WJN-WJ0)+(WJ0+((WJN-WJ0)/EN)*EX)*
&ALOG (ABS ((EX-EN)/EX)))
D=(1./3.14159)*((WJ0-WJN)+(WJ0+((2.*EF+EX)*(WJ0-WJN))/EN)*ALOG
&(ABS ((2.*EF+EX)/(2.*EF+EX+EN))))
ESS=(WJN/(3.14159*(EN-EV)))*((EV-EN)+(EX-EV)*ALOG (ABS ((EV-EX)/
&(EN-EX))))
EVV=(WJV/(3.14159*(EN-EV)))*((EN-EV)+(EN-EX)*ALOG (ABS ((EV-EX)/
&(EN-EX))))
FSS=(WJN/(3.14159*(EV-EN)))*((EV-EN)+(2.*EF+EV+EX)*
&ALOG (ABS ((2.*EF+EN+EX)/(2.*EF+EV+EX))))
FVV=(WJV/(3.14159*(EN-EV)))*((EV-EN)+(2.*EF+EN+EX)*
&ALOG (ABS ((2.*EF+EN+EX)/(2.*EF+EV+EX))))
GH=(WJV/(3.14159))*ALOG (ABS ((2.*EF+EV+EX)/(EV-EX)))
FINT=A+B+C+D+ESS+EVV+FSS+FVV+GH
FINS=A+B+C+D+ESS+FSS
FINV=EVV+FVV+GH
DISPT=FINT/FE
IF (EX.GE.EV) GO TO 30
DISPS=FINS/FE
GO TO 31
30 DISPS=0.0
31 IF (E.GE.0.) GO TO 41
DISPT=0.0
DISPS=0.0
41 H(1,NJ)=EX
H(2,NJ)=DISPT
H(3,NJ)=DISPS
X(1,NJ)=EX
X(2,NJ)=FINT
X(3,NJ)=FINS
X(4,NJ)=FINV
IF (E.GT.ESTOP) GO TO 35
NJ=NJ+1
200 CONTINUE
35 NJ=NJ-1
WRITE(10,11) EF,WJ0,WJN,WJV
DO 40 J=1,NJ
IF (H(1,J).LE.0.) GO TO 40
WRITE(10,15) H(1,J),H(2,J)
40 CONTINUE
WRITE(10,16)
WRITE(10,20) EF,WJ0,WJN,WJV
DO 42 J=1,NJ
IF (H(1,J).LE.0.) GO TO 42
WRITE(10,15) H(1,J),H(3,J)
42 CONTINUE
WRITE(10,16)
WRITE(10,13)
DO 50 J=1,NJ
WRITE(10,15) X(1,J),X(2,J)
50 CONTINUE
WRITE(10,16)
WRITE(10,18)
DO 60 J=1,NJ
WRITE(10,15) X(1,J),X(3,J)
60 CONTINUE
WRITE(10,16)
WRITE(10,19)

```


	DO 70 J=1,NJ	133
	WRITE(10,15)X(1,J),X(4,J)	134
70	CONTINUE	135
	WRITE(10,16)	136
	GO TO 100	137
	END	138
CCCCC	CC	139
	SUBROUTINE CHECK	140
	COMMON EF,E,EX,DELTA,EN,EV,FE	141
C GENERAL		142
	IF(FE.GT.0.0)GO TO 5	143
6	FORMAT(' POTENTIAL=0 IS ILLEGAL')	144
	WRITE(*,6)	145
	STOP	146
5	EX=E	147
	IF(EN.LT.EV)GO TO 1	148
2	FORMAT(' ILLEGAL INPUT, STOP')	149
	WRITE(*,2)	150
	STOP	151
1	IF(EF.NE.0.0)GO TO 3	152
4	FORMAT(' ILLEGAL FERMI-E, STOP')	153
	WRITE(*,4)	154
	STOP	155
C A		156
3	IF(ABS(EF+E).LE.DELTA.AND.E.GE.-EF) EX=-(EF-DELTA)	157
	IF(ABS(EF+E).LT.DELTA.AND.E.LE.-EF) EX=-(EF+DELTA)	158
	IF(ABS(E).LE.DELTA.AND.E.GE.0.) EX=DELTA	159
	IF(ABS(E).LE.DELTA.AND.E.LT.0.) EX=-DELTA	160
C B		161
	IF(ABS(E+EF).LE.DELTA.AND.E.GT.-EF) EX=-EF+DELTA	162
	IF(ABS(E+EF).LE.DELTA.AND.E.LE.-EF) EX=-EF-DELTA	163
	IF(ABS(E+2.0*EF).LE.DELTA.AND.E.GT.-(2.*EF)) EX=-2.0*EF+DELTA	164
	IF(ABS(E+2.0*EF).LT.DELTA.AND.E.LE.-(2.*EF)) EX=-2.0*EF-DELTA	165
C C		166
	IF(ABS(E-EN).LE.DELTA.AND.E.GT.EN) EX=EN+DELTA	167
	IF(ABS(E-EN).LT.DELTA.AND.E.LE.EN) EX=EN-DELTA	168
	IF(ABS(E).LE.DELTA.AND.E.GE.0.) EX+=DELTA	169
	IF(ABS(E).LE.DELTA.AND.E.LT.0.) EX-=DELTA	170
C D		171
	IF(ABS(E+2.*EF).LE.DELTA.AND.E.GT.(-2.*EF)) EX=-2.*EF+DELTA	172
	IF(ABS(E+2.*EF).LT.DELTA.AND.E.LE.(-2.*EF)) EX=-2.*EF-DELTA	173
	IF(ABS(E+(2.*EF+EN)).LE.DELTA.AND.E.GT.(-(2.*EF+EN))) EX=	174
	&-(2.*EF+EN)+DELTA	175
	IF(ABS(E+(2.*EF+EN)).LT.DELTA.AND.E.LE.(-(2.*EF+EN))) EX=	176
	&-(2.*EF+EN)-DELTA	177
C E-S		178
	IF(ABS(EN-E).LE.DELTA.AND.E.GT.EN) EX=EN+DELTA	179
	IF(ABS(EN-E).LE.DELTA.AND.E.LE.EN) EX=EN-DELTA	180
	IF(ABS(E-EV).LE.DELTA.AND.E.GT.EV) EX=EV+DELTA	181
	IF(ABS(E-EV).LT.DELTA.AND.E.LE.EV) EX=EV-DELTA	182
C E-V		183
	IF(ABS(EN-E).LE.DELTA.AND.E.GT.EN) EX=EN+DELTA	184
	IF(ABS(EN-E).LE.DELTA.AND.E.LE.EN) EX=EN-DELTA	185
	IF(ABS(E-EV).LE.DELTA.AND.E.GT.EV) EX=EV+DELTA	186
	IF(ABS(E-EV).LE.DELTA.AND.E.LE.EV) EX=EV-DELTA	187
C F-S		188
	IF(ABS((2.*EF+EN)+E).LE.DELTA.AND.E.GT.-(2.*EF+EN)) EX=	189
	&-(2.*EF+EN)+DELTA	190
	IF(ABS((2.*EF+EN)+E).LE.DELTA.AND.E.LE.-(2.*EF+EN)) EX=	191
	&-(2.*EF+EN)-DELTA	192
	IF(ABS((2.*EF+EV)+E).LE.DELTA.AND.E.GT.-(2.*EF+EV)) EX=	193
	&-(2.*EF+EV)+DELTA	194
	IF(ABS((2.*EF+EV)+E).LT.DELTA.AND.E.LE.-(2.*EF+EV)) EX=	195
	&-(2.*EF+EV)-DELTA	196
C F-V		197
	IF(ABS((2.*EF+EN)+E).LE.DELTA.AND.E.GT.-(2.*EF+EN)) EX=	198

&-(2.*EF+EN)+DELTA	199
IF (ABS ((2.*EF+EN)+E) .LE. DELTA.AND.E.LE.-(2.*EF+EN)) EX=	200
&-(2.*EF+EN)-DELTA	201
IF (ABS ((2.*EF+EV)+E) .LE. DELTA.AND.E.GT.-(2.*EF+EV)) EX=	202
&-(2.*EF+EV)+DELTA	203
IF (ABS ((2.*EF+EV)+E) .LE. DELTA.AND.E.LE.-(2.*EF+EV)) EX=	204
&-(2.*EF+EV)-DELTA	205
C G+H	206
IF (ABS (-(2.*EF+EV)-E) .LE. DELTA.AND.E.GT.-(2.*EF+EV)) EX=	207
&-(2.*EF+EV)+DELTA	208
IF (ABS (-(2.*EF+EV)-E) .LE. DELTA.AND.E.LE.-(2.*EF+EV)) EX=	209
&-(2.*EF+EV)-DELTA	210
IF (ABS (EV-E) .LE. DELTA.AND.E.GT.EV) EX=EV+DELTA	211
IF (ABS (EV-E) .LT. DELTA.AND.E.LE.EV) EX=EV-DELTA	212
RETURN	213
END	214
	215