The engineered phytoremediation of ionic and methylmercury pollution

Subject line: DE-FG07-96-ER20257.EMSP 70054 Final Report

Principal Investigator: Richard B. Meagher, Department of Genetics, Life Sciences Building, Green Street, University of Georgia, Athens, GA 30602, phone: 706-542-1444, fax: 706-542-1387, email: Meagher@arches.uga.edu

Graduate students working on EMSP project

Student	Position	% Time Project	Current Status
Sarah Marshburn	Genetics Graduate Student with RBM	100%	Beginning Ph.D. 2002– Joining laboratory June, 2003.
Andrew Heaton	Ecology Graduate Student with RBM	100%	Ph.D. candidate, Finishing Dec., 2002
Anne Marie Zimeri	Genetics Graduate Student with RBM	100%	Ph.D. candidate, Finish 2003-2004
Raoufa Rahman	Graduate Student - Alexandria Univ., Alexandria, Egypt -RBM is Co-Advisor	100%	Ph.D. candidate – Finishing June, 2003

FINAL PROGRESS REPORT

RESEARCH OBJECTIVE

Our long-term objective is to enable highly productive plant species to extract, resist, detoxify, and/or sequester toxic organic and heavy metal pollutants (Meagher, 2000) applying scientific strategies and technologies from a rapidly developing field called phytoremediation. The phytoremediation of toxic elemental and organic pollutants requires the use relatively different approaches (Meagher, 2000). **Our current specific objectives are to use transgenic plants to control the chemical species, electrochemical state, and aboveground binding of mercury to a) prevent methylmercury from entering the food-chain, b) remove mercury from polluted sites, and c) hyperaccumulate mercury in aboveground tissues for later harvest.** Various parts of this strategy are being critically tested by examining different genes in model plants and field species and comparing the results to control plants as recently reviewed (Meagher et al., 2000; Rugh et al., 2000). A positive spin off of this work has been a strategy for the phytoremediation of arsenic (Dhankher et al., 2002).

RESEARCH PROJECT AND IMPLICATIONS

During the last grant period we focused our efforts on examining transgenic plant species that would be more useful along and in aquatic environments where methylmercury is a produced (e.g., rice and cottonwood) and expanding the genetic capabilities of model plants to hyperaccumulation. We completed many physiological experiments with transgenic tobacco, cottonwood and rice. MerA expressing tobacco (Heaton et al., in prep.), cottonwood (Che et al., submitted) and rice (Heaton et al., submitted) plants are extremely resistant to mercury even in sediments where it is quite toxic and kills wild-type control cottonwood plants.

Methylmercury (MeHg) produced by native bacteria at mercury-contaminated wetland sites is a far more serious problem than Hg(II). MeHg is inherently more toxic than Hg(II), is efficiently biomagnified by several orders of magnitude in the food chain, and poses the most immediate threat to animal populations. MeHg is responsible for the vast majority of cases of mercury poisoning from mercury contaminated fish. MeHg has been a major focus of our research for the last two years. Model plants, *Arabidopsis* and tobacco, have been transformed with a modified bacterial organomercurial lyase gene (*merB*) to degrade methylmercury and other forms of organic mercury (PMA) to the less toxic Hg(II) (Bizily et al., 1999). *Arabidopsis* plants expressing both *merA* and *merB* are resistant to even higher levels of MeHg and are capable of efficiently converting MeHg to Hg(0) at levels 1000 times faster than control plants (Bizily et al., 2000) as shown in the figure below. However, this research demonstrated conclusively that MerB activity is rate limiting in the coupled MerA/MerB catalyzed reaction. During the last year (2000/2001) we focused on improving the efficiency of MerB activity by targeting the enzyme to subcellular environments (Bizily et al., 2002). In a very significant finding, MerB enzyme targeted to the endoplasmic reticulum (ER) or through the ER to the cell wall processes methylmercury 10 to 20 times more efficiently than cytoplasmic MerB. Subcellular targeting of enzymes is an exciting new breakthrough for phytoremediation of toxic heavy metals or organics. Our work suggests that native macrophytes (e.g., trees, shrubs, grasses) can be engineered to thrive on and detoxify the most abundant forms of ionic and organic mercury at polluted sites (Meagher, 2000; Meagher et al., 2000).

During this last grant year we focused research in three new areas, each with the aim of developing plants that hyperaccumulate mercury and other metals. First, a postdoctoral fellow,

Yujing Li, isolated genes encoding the three enzymes in the phytochelatin synthesis pathway γ -glutamylcysteine synthetase (*E. coli*) (Li et al., submitted), glutathione synthetase (*E. coli*) (Li et al., in prep.) and phytochelatin synthetase (fission yeast, *Arabidopsis*, genes in collaboration with Dr. J. Schroeder) (Li et al., in preparation). They were cloned for constitutive plant expression (all tissues and organs) and for focused expression aboveground in leaves and stems. These constructs have been tested in transgenic *Arabidopsis* and show that plants expressing any one of these three enzymes are more resistant to mercury and arsenic than control plants. In addition, we isolated three specific monoclonal antibodies to monitor the plant expression levels of three of these proteins using a novel rapid immunological protocol developed in our laboratory (Li et al., 2001). Second, a graduate student isolated several of the diverse *Arabidopsis* metallothionein genes and demonstrated their differential binding specificities for Hg(II) and other metal ions (Zimeri et al., in prep).

Third, we examined three new promoter systems for expressing phytoremediating enzymes like MerA and MerB. We developed the ACT2pt (pt = promoter terminator) system, based on *Arabidopsis* actin gene ACT2 (Balish, in prep.). ACT2pt/*merA* expresses MerA protein at 5 times higher levels than the widely accepted 35Sp/*NOS* constitutive expression system. Dr. Balish has shown that ACT2pt/*merA* appears to be more consistent in being expressed in nearly all transgenic *Arabidopsis* and tobacco lines tested instead of only 60-70% of the 35Sp/*merA*/*NOS* lines. Similarly, we developed SRS1pt based on our previous work a decade ago on the light induced SRS1 soybean rubisco small subunit gene (Dhankher et al., 2002; Li et al., submitted). This new cassette expresses very high levels of a GUS reporter in leaves, but not roots. It will be used to express genes that assist with the hyperaccumulation of metals in above-ground tissue. A third plant promoter system was obtained from a European laboratory for root-specific expression of MerA and MerB, to be used as part of our mercury hyperaccumulation efforts (Heaton, in prep.). This tobacco promoter is active in roots, but further research is needed to see if it is expressed strongly and specifically enough for efficient expression of *merA* and detoxification of Hg(II) in roots.

PLANNED ACTIVITIES ON SUBMITTED RENEWAL

In order to advance this mercury phytoremediation strategy, our research has focused on the following Specific Aims and we have made the initial progress indicated on all six: (1) to

increase the transport of mercury to aboveground tissue through the root expression of MerA; (2) to identify small mercury binding peptides that enhance hyperaccumulation aboveground (initial results are positive with EC and MerP peptides); (3) to test the ability of multiple genes acting together to enhance resistance and hyperaccumulation (several new gene combinations have been quite successful); (4) to construct a simple molecular system for creating male/female sterility, allowing engineered grass, shrub, and tree species to be released indefinitely at contaminated sites (initial data are positive on model plants engineered with remediable nutrient-based sterility); (5) to test the ability of transgenic cottonwood and rice plants to detoxify ionic mercury and prevent methylmercury release from contaminated sediment (a significant reduction of mercury from sediment was demonstrated); and (6) to conduct field testing with transgenic plants for the remediation of mercury (the initial results of which demonstrate significant reductions in total mercury in a three month period). The results of these experiments will enable the phytoremediation of methyl- and ionic mercury by a wide spectrum of deep-rooted, fast-growing plants adapted to diverse environments.

INFORMATION ACCESS (PUBLICATIONS AND WEB SITE)

Bizily, S., Kim, R., Kandasamy, M., and Meagher, R.B. (2002). Subcellular targeting of methylmercury lyase enhances its specific activity for organic mercury detoxification in plants. *Plant Physiol.* in press.

Bizily, S., Rugh, C.L., and Meagher, R.B. (2000). Phytodetoxification of hazardous organomercurials by genetically engineered plants. *Nat. Biotechnol.* **18**, 213-217.

Bizily, S., Rugh, C.L., Summers, A.O., and Meagher, R.B. (1999). Phytoremediation of methylmercury pollution: *merB* expression in *Arabidopsis thaliana* confers resistance to organomercurials. *Proc Natl Acad Sci USA* **96**, 6808-6813.

Che, D.S., Meagher, R.B., Heaton, A.C.P., Lima, A., and Merkle, S.A. (submitted). Expression of mercuric ion reductase in eastern cottonwood confers mercuric ion reduction and resistance.

Dhankher, O.P., Li, Y., Rosen, B.P., Shi, J., Salt, D., Senecoff, J.F., Sashti, N.A., and Meagher, R.B. (2002). Engineering tolerance and hyperaccumulation of arsenic in plants by combining arsenate reductase and gamma-glutamylcysteine synthetase expression. *Nat Biotechnol* **20**, 1140-5.

- Heaton, A., Rugh, C., and Meagher, R.** (in prep.). Genetically engineered *Nicotiana tabacum* as a model for mercury phytoremediation. *Plant Phys.*
- Heaton, A., Rugh, C., Wang, J., Kim, R., and Meagher, R.** (submitted). Toward detoxifying mercury-polluted aquatic sediments using rice genetically-engineered for mercury resistance. *Environmental Toxicology and Chemistry*
- Li, Y., Kandasamy, M.K., and Meagher, R.B.** (2001). Rapid isolation of monoclonal antibodies: monitoring enzymes in the phytochelatin synthesis pathway. *Plant Physiol.* **127**, 711-719.
- Li, Y., Parkash Dhankher, O., and Meagher, R.** (in prep.). Engineering metal hyperaccumulation: II. Overexpression of glutathione synthetase in plants confers high level arsenic and mercury resistance.
- Li, Y., Parkash Dhankher, O., and Meagher, R.** (submitted). Engineering metal hyperaccumulation: I. Overexpression of γ -glutamylcysteine synthetase in plants confers high level arsenic and mercury tolerance.
- Li, Y., Parkash Dhankher, O., Schroeder, J., and Meagher, R.** (in preparation). Engineering metal hyperaccumulation: III. Leaf-specific expression of phytochelatin synthase in plants confers high level arsenic and mercury resistance.
- Meagher, R.B.** (2000). Phytoremediation of toxic elemental and organic pollutants. *Curr. Opin. Plant Biol.* **3**, 153-162.
- Meagher, R.B., Rugh, C.L., Kandasamy, M.K., Gragson, G., and Wang, N.J.** (2000). Engineered phytoremediation of mercury pollution in soil and water using bacterial genes. In: *Phytoremediation of Contaminated Soil and Water*, N. Terry, and G. Banuelos, eds (Boca Raton: Lewis Publishers), pp. 203-221.
- Rugh, C.L., Bizily, S.P., and Meagher, R.B.** (2000). Phytoremediation of environmental mercury pollution. In: *Phytoremediation of toxic metals: Using plants to clean-up the environment*, B. Ensley, and I. Raskin, eds (New York: Wiley and Sons), pp. 151-169.