PROJECT TITLE: Atmospheric Aerosol Source-Receptor Relationships: The Role Of Coal-Fired Power Plants

DOE AWARD #: DE-FC26-01NT41017

REPORT: Semi-Annual Technical Progress Report

REPORTING PERIOD: September 2004 – February 2005

ISSUED: April 2005

SUBMITTED TO: US Department of Energy
National Energy Technology Laboratory
P.O. Box 10940
626 Cochrans Mill Rd
Pittsburgh, PA 15236-0940
Attn: Bill Aljoe

PREPARED BY: Carnegie Mellon University
5000 Forbes Ave
Pittsburgh, PA 15213

PIs: Allen L. Robinson, Spyros N. Pandis, Cliff I. Davidson
(412) 268 – 3657; (412) 268 – 3348 (fax)
e-mail: alr@andrew.cmu.edu

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ABSTRACT

This report describes the technical progress made on the Pittsburgh Air Quality Study (PAQS) during the period of September 2004 through February 2005. Significant progress was made this project period on the analysis of ambient data, source apportionment, and deterministic modeling activities. The major experimental achievement this project period was the characterization of the mercury and fine particle emissions from two modern, large, commercial pulverized coal boilers. This testing completes the field work component of the Source Characterization Activity.

This report highlights results from mercury emission measurements made using a dilution sampler. The measurements clearly indicate that mercury is being transformed from an oxidized to an elemental state within the dilution. However, wall effects are significant making it difficult to determine whether or not these changes occur in the gas phase or due to some interaction with the sampler walls.

This report also presents results from an analysis that uses spherical aluminum silicate (SAS) particles as a marker for primary $PM_{2.5}$ emitted from coal combustion. Primary emissions from coal combustion contribute only a small fraction of the $PM_{2.5}$ mass (less than 1.5% in the summer and less than 3% in the winter) at the Pittsburgh site. Ambient SAS concentrations also appear to be reasonably spatially homogeneous. Finally, SAS emission factors measured at pilot-scale are consistent with measurements made at full-scale.

This report also presents results from applying the Unmix and PMF models to estimate the contribution of different sources to the $PM_{2.5}$ mass concentrations in Pittsburgh using aerosol composition information. Comparison of the two models shows similar source composition and contribution for five factors: crustal material, nitrate, an Fe, Mn, and Zn factor, specialty steel production, and a cadmium factor. PMF found several additional factors. Comparison between source contributions for the similar factors shows reasonable agreement between the two models. The sulfate factor shows the highest contribution to local $PM_{2.5}$ with an annual average contribution of approximately 28% (from PMF). The nitrate, crustal material, and primary OC and EC factors also show significant contributions on the order of 10-14%. The sulfate factor is affected by photochemistry and therefore shows maximum values in summer.

TABLE OF CONTENTS

Disclaimer	2
Abstract	3
Table of Contents	4
List of Figures	5
Excutive Summary	6
Experimental	7
Activity 1. Project Management	7
Activity 2. Ambient Monitoring	7
Activity 3. Source Characterization.....	7
Activity 4. Source Apportionment	7
Activity 5. Three-Dimensional Deterministic Modeling	8
Results and Discussion	8
1. Single Particle Characterization.....	8
4. Development and Evaluation of Measurement Methods.....	9
3. Effects of dilution on mercury speciation in coal-fired power plant plumes.....	10
4. Source Apportionment of Primary Coal Emissions with SAS.....	16
5. Source Apportionment using PMF and UNMIX	20
Conclusions.....	26
References.....	27
Presentations and Publications.....	27
Publications:.....	27
Presentations:	32

LIST OF FIGURES

Figure 1: Schematic of CMU Dynamic Dilution Sampler (Dilution Tunnel)	10
Figure 2: Time series of diluted and undiluted mercury concentration data taken on September 30	12
Figure 3: Flue gas and diluted sample mercury concentration. Note - Diluted samples have been corrected for tunnel blank and dilution ratio.	13
Figure 4: Ratios of diluted versus undiluted mercury concentrations. Diluted samples were blank and dilution corrected.....	14
Figure 5. Time series of SAS number concentration and PM _{2.5} mass measured during the July 2001 intensive.	16
Figure 6. Time series of SAS number concentrations measured at the CMU site and the NETL site for January 2001.....	17
Figure 7. SAS emission factors as a fraction of PM _{2.5} mass measured during pilot-scale and full-scale boiler testing.....	19
Figure 8. Estimate PM _{2.5} mass contributed by primary coal emissions to the Pittsburgh Supersite during the Summer 2001 intensive.	20
Figure 9. Unmix source compositions apportioned by PM _{2.5} mass. All ten factors are outputs of PMF, while only the first six factors are outputs of Unmix. Although the Unmix regional transport factor includes primary OC and EC, it is graphed for comparison with the PMF sulfate factor.	21
Figure 10. PMF and Unmix source contributions apportioned by PM _{2.5} mass. All ten factors are outputs of PMF, while only the first six factors are outputs of Unmix. Note that for comparison purposes, the Unmix regional transport factor is compared to the PMF sulfate plus primary OC and EC factors.	22
Figure 11. Monthly average Unmix source contributions. Height of the bars corresponds to the monthly average PM _{2.5} mass measured with a TEOM. The study average represents the average source contributions from July 11, 2001 through July 31, 2002. Unmix uses PM _{2.5} mass as a fitting species so the mass of PM _{2.5} unexplained by Unmix is less than 1%	23
Figure 12. Monthly average PMF source contributions. Height of the bars corresponds to the monthly average PM _{2.5} mass measured with a TEOM. The study average represents the average source contributions from July 11, 2001 through July 31, 2002. The unexplained mass is the difference between the monthly average PM _{2.5} mass and the sum of the monthly averaged source contributions from each factor.	24
Figure 13. Linear regression results of PMF and Unmix source contributions.....	25

EXECUTIVE SUMMARY

With support from the US Department of Energy and the US Environmental Protection Agency, Carnegie Mellon University is conducting detailed studies of the ambient particulate matter in the Pittsburgh, PA metropolitan area. The work includes ambient monitoring, source characterization, and modeling (statistical and deterministic) for source apportionment. The major objectives of the project include:

- To achieve advanced characterization of the PM in the Pittsburgh region. Measurements include the PM size, surface, volume, and mass distribution; chemical composition as a function of size and on a single particle basis; temporal and spatial variability.
- To obtain accurate current fingerprints of the major primary PM sources in the Pittsburgh region using traditional filter-based sampling and state-of-the-art techniques.
- To estimate the impact of the various sources (transportation, power plants, natural, etc.) on the PM concentrations in the area using both statistical and deterministic models.
- To quantify the responses of the PM characteristics to changes in these emissions in support of the emission control decision making in the area.
- To develop and evaluate current and next generation aerosol monitoring techniques for both regulatory applications and for determination of source-receptor relationships.

This document is the eighth semi-annual progress report for this project. During this project period significant progress was made on the analysis of ambient data, source apportionment, and deterministic modeling activities. Major achievements this project period and results described in this progress report include:

- Measurement of mercury and fine particle emissions from two full-scale pulverized coal boilers using a state-of-the-art dilution sampling system.
- Analysis of ambient and emission data for spherical aluminum silicate (SAS) particle concentrations to estimate the contribution of coal combustion to primary PM_{2.5} at the Pittsburgh site.
- Application of Unmix and PMF receptor models to apportion PM_{2.5} mass in the Pittsburgh area.
- Investigation of the effects of dilution on mercury oxidation state in coal-fired power plant plumes.

EXPERIMENTAL

This section provides an overview of the effort on various project activities. This project period the majority of the effort went into Activity 3 Source Characterization, Activity 4 Source Apportionment, and Activity 5 Three-Dimensional Modeling.

Activity 1. Project Management

During this project period additional data were submitted to EPA for inclusion in the Supersites Relational Database and to ATS for inclusion in the DOE-sponsored air quality database.

Activity 2. Ambient Monitoring

The purpose of this activity is to create an extensive database of ambient PM measurements for source apportionment, examination of aerosol processes, evaluation of instrumentation, and air quality model development and evaluation. Data collection is complete. Work has continued on the analysis of the ambient data set and selected results from this analysis are shown in the Results and Discussion section of this report.

Activity 3. Source Characterization

The purpose of this activity is to develop updated emission profiles for important source categories around Pittsburgh. Updated source profiles are being developed through a combination of source testing, fence line measurements, and analysis of highly time resolved data collected at the central site. These profiles are used in the source apportionment and deterministic modeling activities.

The major accomplishment in Activity 3 this project period was emission testing on two full scale coal-fired power plants. One test focused on effects of dilution on mercury partitioning on coal fired power plant plumes. The second test focused on measuring fine particle emission rates. The testing was done on modern, large (>400 MWe) pulverized coal fired boilers with low NO_x burners and electrostatic precipitators (ESP) for control of particle emissions. The mercury experiments were done on a boiler with an SCR unit for NO_x control. The particle characterization was performed downstream of the ESP. Neither plant had equipment for sulfur control. Both plants were operating on a blend of eastern bituminous and lower sulfur western sub-bituminous coal. This completes the field work component of the Activity 3. Selected results from the source characterization activities are shown in the Results and Discussion section of this report.

Activity 4. Source Apportionment

The purpose of this activity is to quantify the contribution of different sources to the fine PM_{2.5} levels in Pittsburgh. Significant effort was expended on the source

apportionment analysis this project period and selected results are shown in the Results and Discussion section.

Activity 5. Three-Dimensional Deterministic Modeling

The purpose of this activity is to evaluate the performance of the three-dimensional chemical transport model (PMCAMx) with air quality data collected by this and other projects. PMCAMx is a publicly available computer modeling system for the integrated assessment of photochemical and PM pollution. This CTM has been recently upgraded by the CMU team and ENVIRON to include state-of-the-art description of aerosol dynamics and thermodynamics, cloud chemistry, and wet removal processes. PMCAMx+ is the research version of the code and it includes the latest developments in Carnegie Mellon organic and inorganic aerosol and aqueous-phase chemistry modules. During this project period research focused on evaluating national inventories for primary organic carbon and elemental carbon used by the model.

RESULTS AND DISCUSSION

1. Single Particle Characterization

The University of California at Davis operated the RSMS-3, a single particle mass spectrometer, for twelve consecutive months (306 operation days) at the central monitoring site. This instrument provides semi-continuous, highly time resolved measurements of single particle size and composition for particulate pollution. The data are being used for single particle characterization, particle classification, source attribution, and correlating/combining with measurements from collocated instruments.

Approximately 236,000 single particle mass spectra were collected throughout the duration of the Pittsburgh Supersite experiment using RSMS-3, a third generation single particle mass spectrometer. The instrument was operated semi-continuously for 306 days, sampling particles with aerodynamic diameters in the range of 30 – 1100 nm and collecting both positive and negative ion spectra, particle size and time of detection for each particle measured. The entire data set has been fully processed and analyzed. Spectra have been clustered into 20 distinct particle classes based on the distribution of their positive ion mass peaks. Negative ion spectra were classified independently within each positive ion class. Frequency of occurrence versus particle size, month of the year and wind direction has also been calculated for the full data set, as well as within each class. Results indicate a rich array of multi-component ultrafine particles composed primarily of carbon and ammonium nitrate. Approximately 54% of all the particles measured fell into the carbonaceous ammonium nitrate (CAN) class. These particles were observed in all size bins and from most wind directions for the entirety of this study. Ubiquitous sources throughout the area, including vehicular emissions and secondary organic aerosol formation, are considered to be responsible for a larger fraction of these particles. In terms of particle number, metal containing aerosol dominated the remainder of the particle classes identified. These particles were rich in K⁺, Na⁺, Fe⁺, Pb⁺, and to

a lesser extent, Ga⁺ and Zn⁺. They tended to be smaller in size and were highly correlated with specific wind directions, facilitating the isolation of specific sources. (Bein et al. 2005)

4. Development and Evaluation of Measurement Methods

The University of Maryland has developed multi-element simultaneous GFAAS methods for determining 11 metals in three analytical groups (Group 1, predominately crustal elements, Al, Cu, Fe, Mn, and Cr; Group 2, tracers of coal and oil combustion as well as other anthropogenic sources, Se, As, Pb, and Ni; and Group 3, tracers of municipal incinerator aerosol - Zn and Cd), in ambient slurry samples collected at 30 minute intervals with the University of Maryland - Semicontinuous Elements in Aerosol Sampler, SEAS II. Addition of acid (0.2% v/v nitric acid) and ultrasonic treatment (15 min) was used to improve slurry stability and matrix composition. Wide calibration ranges were needed for analysis of the slurry samples. Improved calibration ranges were observed by using a mini carrier gas flow during the atomization stage. Compromise in analytical sensitivity varied as atomizer conditions for metals changed. Palladium (4 ug) and hydrogen (5% in Argon) were found to be effective modifiers for group one and two elements. A fast furnace temperature program was developed for Group 3. Zinc showed two-fold higher linearity in the method as compared with standard methods. Detection limits by the SEAS-II-GFAA approach were compared with traditional filter XRF, LA-ICPMS, and INAA techniques used in air quality studies. The efficacy of the analytical methods developed were applied to urban PM_{2.5} collected in Baltimore for use in a NIST fine-particulate Standard Reference material. Results obtained by the GFAA methods agreed well with the results obtained by NIST using Instrumental Neutron Activation Analysis. (Pancras 2005)

The University of California Davis coupled the Versatile Aerosol Concentration Enrichment System (VACES) to a single particle mass spectrometer (RSMS-3) to determine if the pre-concentration of particles induced by the VACES increases the hit rate of RSMS-3, as well as to monitor whether the VACES introduces any changes in the composition of the particles sampled. The goal was to increase the particle hit rate, and thus detection efficiency, of RSMS-3 in order to broaden its applicability to conditions other than polluted urban areas, for example cleaner rural sites or the stratosphere.

Results from the field evaluation indicate that VACES increased the RSMS-3 hit rate by 5-20 times at particle sizes ranging from 40 to 640 nm. VACES only enhances the hit rate by about a factor of 2 for large particle sizes because the RSMS-3 flow rates for these particles did not match the optimum operating condition of VACES. During the 3 days of measurements most of the particles were a mixture of carbonaceous material and ammonium nitrate with a variation across the spectrum from particles that were mostly carbonaceous to particles that were mostly ammonium nitrate. Both ambient and concentrated carbonaceous and ammonium nitrate composition distributions were indistinguishable with the RSMS-3, suggesting that VACES introduces an insignificant artifact for those particles. (Zhao et al. 2005)

3. Effects of dilution on mercury speciation in coal-fired power plant plumes

Dilution sampling was carried to investigate the changes in partitioning of mercury in plume of coal-fired power plants. Measurements were made on a 817 MW unit firing a blend of 60% powder river basin sub-bituminous coal and 40% mid-sulfur eastern bituminous coal. The boiler is wall-fired, with low-NO_x burners. The boiler has an ESP with SO₃ and NH₄ conditioning. An SCR is also used for NO_x control. Measurements were made over a period of 3 days downstream of the SCR but upstream of the ESP.

A dynamic dilution sampler was used to investigate changes in partitioning of mercury upon dilution (Figure 1). The system is designed to simulate atmospheric dilution and is constructed out of stainless steel. For the mercury testing all of the wetted components in contact with sample were coated with a 1200 Å thick layer of Silcosteel, applied by Restek Performance Coatings. In addition to the main dilution tunnel, a residence time chamber (RTC, not shown in Figure 1) was brought inline during some testing periods. With the RTC inline, the overall aging time of the diluted exhaust can be varied between 2 seconds and 5 minutes.

Figure 1: Schematic of CMU Dynamic Dilution Sampler (Dilution Tunnel)

Mercury sampling and speciation were via Mercury Continuous Emission Monitors (CEMs) operated by the University of North Dakota – Energy and Environmental Research Center (UND-EERC). A PS Analytical Sir Galahad CEM measured gas-phase total and elemental mercury concentrations inside the dilution sampler. These concentrations ranged from 0 to 250 ng/m³. As operated, the Sir Galahad collected sample for 5 minutes before analysis; analysis required 3 to 4 minutes and alternated between elemental and total mercury analyses. A Tekran Mercury Vapor Analyzer was used to sample the un-diluted flue gas. The Tekran measured concentrations in the range of 0 to 10 µg/m³ and alternately measured elemental mercury for 10 minutes and total mercury for 45 minutes; data points were recorded every 2 minutes in both modes.

Measurements were made over a three day period. On September 28th and on the morning of the 29th indicated significant losses of mercury inside the system. In an attempt to reduce these losses, the walls of the dilution tunnel and residence time chamber were heated to an outer surface temperature of 180 °C. Upon heating, mercury levels inside the tunnel dramatically increased presumably due to mercury desorption from the tunnel walls. This saturated the Sir Galahad analyzer, limiting data collection on the 29th. All of the experiments performed on September 30th were done with heated walls.

Blanks were measured at the beginning and end of each day of experiments by operating the system on filtered dilution air. Data collected from the dilution tunnel were blank corrected using a best estimate tunnel blank for the specific time period. The data are also dilution corrected using a period-average dilution ratio calculated using the background-corrected flue gas and dilution tunnel CO₂ concentrations. In subsequent discussion, the abbreviation DT refers to experiments performed using only the dilution tunnel, while DT + RTC refers to experiments using both the dilution tunnel and residence time chamber were online. The residence time in the two cases was approximately 2 seconds and 5 minutes, respectively.

Time series of the uncorrected (no blank or dilution correction) mercury measurements made on September 30 is shown in Figure 2. Data points in the top portion of the figure are the flue-gas mercury concentrations, which show a modest increase in total mercury concentrations over the course of the day.

The data in the lower portion of Figure 2 are mercury concentrations measured in the dilution tunnel, with different testing periods annotated. There are large changes in mercury concentrations inside the dilution sampler corresponding to changes in sampler operating conditions. For example, switching the mercury monitor from the end of the dilution tunnel to the residence time chamber at ~ 15:30 caused a large increase in measured concentrations. A constant dilution ratio of 30 was maintained throughout this period. The dilution ratio was increased to 100 at 17:00 after which there is a slow decay in measured total and elemental mercury concentrations inside this system. This decay does show a flattening trend, but sampling was stopped at approximately 20:00, when the exhaust sampling probe was detached from the dilution system and it was operated with filtered dilution air. At the end of the day, the blank level of elemental mercury is significantly higher than at the beginning of the day. Figure 2 shows the blank slowly decaying as the system is continually flushed with filtered dilution air overnight.

Figure 2: Time series of diluted and undiluted mercury concentration data taken on September 30

A summary of the data from each test condition is shown in Figure 3. Undiluted concentrations are an average of values from the Tekran CEM over the period when the tunnel was operated at a constant dilution ratio. The in-stack data indicate a clear predominance of oxidized mercury at this location in the plant. The total mercury concentration in the stack varied between 3 and $7.2 \mu\text{g}/\text{m}^3$. Elemental mercury ranged between 0.06 to $0.25 \mu\text{g}/\text{m}^3$. Oxidized mercury levels varied between 2.8 and $7.1 \mu\text{g}/\text{m}^3$ contributing over 90%, and generally closer to 98%, of the total mercury.

Data from the dilution sampler are shown dilution- and blank-corrected along with the flue-gas concentrations in Figure 3. It is clear that mass closure was very poor during the periods when the tunnel was unheated (on September 28 and 29). The mass balance improved somewhat when the heat-tape was added to the system (September 30). All samples except the first DT DR100 show a significant increase in dilution-corrected elemental mercury concentration relative to the flue-gas concentrations. This is a clear indicating that there is conversion of oxidized mercury to elemental mercury occurring inside the dilution sampler. The measurements made at the end of the residence time chamber show a large increase in elemental mercury and approach mass closure with the flue-gas samples.

Figure 3: Flue gas and diluted sample mercury concentration. Note - Diluted samples have been corrected for tunnel blank and dilution ratio.

Mass closure and mercury transformation can be seen more clearly in Figure 4, which plots the ratios of the diluted to undiluted total, elemental and oxidized mercury concentrations presented in Figure 3. Perfect mass balance closure is a ratio of 1. The three unheated data points show very poor mass closure, with a nearly complete loss of oxidized mercury in the dilution tunnel. During these experiments, on average, only 14% of the total mercury mass measured in the flue-gas is captured by measurements in the dilution tunnel. On the other hand, the dilution corrected elemental mercury concentrations measured inside the tunnel are higher by approximately a factor of two relative to that measured in the flue gas, indicating some reduction of elemental mercury in the dilution tunnel.

Heating the tunnel walls improved mass closure. The two experiments with only the heated dilution tunnel inline show an improving total mercury mass closure relative to the flue-gas measurements; 29% of mass was recovered at DR = 100 and 42% at DR = 30. There was also a significant increase in elemental mercury levels in the diluted exhaust; elemental mercury levels in the diluted gases were on the order of 10 times those in the un-diluted gas. These two experiments also indicate that the amount of reduction occurring in the dilution tunnel increases at the higher dilution ratio. The diluted measurements with DR = 30 showed 9 times more elemental mercury than the undiluted gas, while those at DR = 100 show 16 times more.

Improved mass closure was observed in measurements made at the end of the residence time chamber. This chamber added approximately 5 minutes aging time to the system before the mercury measurements. The inclusion of the residence time chamber improved total mercury mass closure significantly, to 65% and 79% at dilution ratios of 30 and 100, respectively. However, there is a concern about whether appropriate blank corrections were applied. As indicated in the time series in Figure 2, the tunnel blank levels of total and especially elemental mercury increased significantly during the course of the day. This suggests that variable adsorption and desorption of mercury is occurring and thus complicates blank-correction of dilution tunnel measurements. These experiments showed that nearly all the mercury at the end of the RTC was in elemental form, with dilution corrected elemental mercury concentrations being a factor of 70 to 85 higher than those under non-diluted conditions. Also shown is an almost total reduction of oxidized mercury within the dilution tunnel, with 5 and 11% of the initial concentrations of flue gas oxidized mercury found in the diluted sample. As in the experiments without the residence time chamber, a larger reduction effect is seen in the run at a higher dilution ratio, suggesting a connection between dilution ratio and the mercury transformation.

Figure 4: Ratios of diluted versus undiluted mercury concentrations. Diluted samples were blank and dilution corrected.

Although the data clearly show that oxidized mercury is being converted into an elemental state inside the dilution sampler, it is difficult to conclude whether or not these transformations are occurring on the tunnel walls or in the gas phase. However, there are

some trends that are consistent across all tests. First, there are significant losses of oxidized mercury to the tunnel walls, especially when they are not heated. The question then becomes whether the higher wall temperature reduces the absorption of mercury to the walls, or simply increasing the rate at which absorbed mercury desorbs. The fact that the mass closure improved over the course of the September 30th seems to suggest that the tunnel wall becomes 'saturated' with mercury, approaching an equilibrium state. However, the blanks of total mercury at the beginning and end of the testing on September 30th were approximately the same, while the elemental mercury blank was 4 times higher at the end of the day. This indicates that the tunnel background oxidized mercury level decreased over the course of the day and suggests that mercury was being released from the walls in elemental form during testing. If so, then some fraction of the observed transformation of oxidized to elemental mercury is occurring on the walls. Two distinct hypotheses emerge to viably explain the reduction of mercury in the tunnel:

1. Significant portions of the oxidized mercury in the flue gas are being reduced to its elemental form in the dilution tunnel in the gas phase. The magnitude of this effect increases with increasing dilution ratio and residence time – given sufficient time in the tunnel the vast majority of oxidized mercury is reduced.
2. The reduction of oxidized mercury is occurring on the heated dilution tunnel walls. Oxidized mercury is adsorbing onto the walls and surface reactions leads to its reduction to elemental form and subsequent release into the gas stream. Heating the walls increases the rate at which adsorbed mercury is released to the gas stream and influences the measurements of elemental mercury made in the system.

When the heat-tape was initially applied and mercury levels in the tunnel initially spiked (but before the analyzer saturated), the total mercury level peaked much more quickly (50 minutes earlier) than the elemental level. The implication is that the thermally-driven desorption was largely out-gassing oxidized mercury into the dilution tunnel. Since the CEM signal was saturated, nothing can be said about the relative levels of elemental and total mercury. However, this behavior suggests that reduction is not strictly occurring on the tunnel walls and lends some support to the first hypothesis.

Arriving at firm conclusions based on this data is complicated by the lack of a complete tunnel characterization during this short test period. Tunnel background mercury levels before and during tests are difficult to quantify because only two tunnel blanks -at the beginning and end of testing- were collected during the major day of data collection. In this analysis blanks from the beginning of the day was applied to data taken without the RTC online, while the blank from the end of the day was applied to the last two tests with the RTC installed. The blank levels in the tunnel at intermediate points are unknown. Attempts at interpreting the data are further complicated by the fact that our only view of the transient response of the tunnel to external heating was truncated due to instrument issues.

In future experiments, tunnel blanks taken in between test points would allow a better characterization of the transformation occurring inside the dilution tunnel. If further experiments at elevated tunnel wall temperatures are conducted, quantifying the transient response of mercury speciation in the tunnel to changes in tunnel surface temperature would be helpful. In addition, other methods for reducing wall losses might be developed. Some possibilities are applying a more inert surface coating to the tunnel, designing a tunnel that uses a sheath flow to minimize contact between the sample and the tunnel walls or running the tunnel at other surface temperatures. Development of such a sampler would be best accomplished on a laboratory or pilot-facility scale.

4. Source Apportionment of Primary Coal Emissions with SAS

RJ Lee Group continued analyzing samples collected on polycarbonate filters to determine ambient spherical aluminum silicate (SAS) particle number concentrations on both source and ambient filters. SAS particles are thought to be a unique tracer for coal-fire power plant formed from the high temperature processing of fly ash (Fisher et al. 1978; Webber et al. 1985; Eatough et al. 1996).

CCSEM methods were used to measure the size and obtain the elemental composition for 2500 individual particles from each ambient and source sample. The maximum and minimum diameter measured during the analysis was used to calculate the volume of each particle and each particle was assigned a density based on a common oxide in proportion to the elements present, determined by the EDS analysis. Images and spectra collected during the analysis were then reviewed for each individual particle to distinguish SAS from non-spherical material.

Figure 5. Time series of SAS number concentration and PM_{2.5} mass measured during the July 2001 intensive.

Time series of SAS number concentration and PM_{2.5} mass are shown in Figure 5 for the July 2001 intensive. At times the two parameters are correlated, for example

concentrations of both peaked in early August. However, in middle of July there was another spike in PM_{2.5} mass concentration while SAS concentrations remain low. SAS concentrations like those of other PM parameters vary episodically presumably in response to meteorological conditions.

Figure 6. Time series of SAS number concentrations measured at the CMU site and the NETL site for January 2002.

Figure 6 shows time series of SAS number concentrations measured at the CMU site and the NETL site for January 2002. SAS concentrations in the winter are comparable to those in the summer. There is also reasonable in the SAS concentrations measured at the two sites on 4 of the 5 days when data are available at both sites. On the 1/5/02 SAS concentrations at the CMU site were significantly higher than at the NETL site. An SO₂ plume impacted the CMU site for much of the day.

The SAS number concentrations data shown in Figures 5 and 6 can be used to derive an upper bound on the contribution of primary emissions from coal-fired power plants to ambient PM_{2.5} levels in Pittsburgh. The estimate is an upper bound because it assumes that coal boilers are the only source of SAS particles. To estimate the contribution of primary coal emissions to ambient PM_{2.5} we need a SAS emission factor for coal combustion -- number of SAS particles/μg primary PM_{2.5} emissions. The contribution of primary coal-fired power plant emissions to the ambient PM_{2.5} was determined by dividing the total number of SAS per sample collection volume by the total number of SAS per μg of coal emissions.

SAS emission factors were determined based on CCSEM analysis filters collected during source tests. Data are available from both pilot-scale and full scale testing. All filter samples were collected after particulate control devices. Samples were collected

with the Carnegie Mellon University dilution sampling system. Pilot scale measurements were made using the Combustion and Environmental Research Facility (CERF) at the Department of Energy National Energy Technology Laboratory. The CERF is a pulverized-coal combustor designed to simulate the time-temperature history of a commercial coal boiler. At full load it consumes 20 kg of pulverized coal per hour, roughly 150 kW when burning a typical US bituminous coal. Coal is injected through a swirl-stabilized burner at the top of a 3-m tall and 45-cm diameter refractory lined combustion zone. Combustion products then flow into a horizontal convective section, through two flue gas coolers, heat-traced piping, and into a bag house. The filter samples considered here were collected after the bag house. Measurements at pilot-scale were made while firing a range of common utility and industrial fuels representing: Prater Creek Coal is an eastern bituminous coal with low sulfur and ash content; Black Thunder Coal is a low-sulfur, high-calcium sub-bituminous coal from the Powder River Basin; and Bailey Mine Coal is a high-sulfur, Pittsburgh seam bituminous coal.

The full-scale measurements were made on large, modern pulverized coal boilers burning blends of eastern bituminous and western sub-bituminous coals. On boiler #1 the measurements were made upstream of the particle control device and on boiler #2 they were made downstream of the particle control device.

SAS emission factors are summarized in Figure 7 and Table 1. The emission factors vary by a factor of 7 with the emission factor for Pittsburgh #8 coal being the lowest and the emission factor for the bituminous/sub-bituminous blend fired in Boiler #2 being the highest. The measurements at pilot-scale are comparable to those made at full scale. The most direct appropriate comparison is between the pilot-scale measurements made while firing Powder River Basin coal (PRB, a western sub-bituminous coal) and the full-scale data since the blends considered at full scale were predominantly western sub-bituminous. The emission factor measured while firing the PRB falls in between the emission factors measured at full-scale. The data suggest that there may be some rank dependence in the emission rate of SAS with the SAS emission rate of the lower rank sub-bituminous coals being higher than that of the higher rank eastern bituminous coals.

Table 1. SAS emission factors (number of SAS particles/ μg primary $\text{PM}_{2.5}$ emissions) for different fuels.

Test	SAS/ μg
<u>Pilot-Scale Combustor</u>	
Prater Creek Coal (Test 1)	2.16E+05
Prater Creek Coal (Test 2)	2.14E+05
Pittsburgh #8 Coal	1.25E+05
Powder River Basin	5.81E+05
Prater Creek Coal-Wood Blend	3.16E+05
<u>Full-Scale Boiler</u>	
Boiler #1	4.40E+05
Boiler #2	7.11E+05

Figure 7. SAS emission factors as a fraction of PM_{2.5} mass measured during pilot-scale and full-scale boiler testing.

Predictions of the PM_{2.5} mass contributed by primary emission from coal combustion for the Summer 2001 Intensive are shown in Figure 8. Using the Pittsburgh #8 coal emission factor produces the highest estimate and the Boiler #2 profile produces the lowest with the monthly-average coal contribution for July 2001 varying between 0.09 to 0.55 $\mu\text{g}/\text{m}^3$ of PM_{2.5} (0.25% to 1.6% of the ambient monthly average PM_{2.5} mass). The monthly average contributions for January 2002 are essentially the same on a concentration basis, 0.08 to 0.58 $\mu\text{g}/\text{m}^3$ of PM_{2.5}, but larger on a relative basis, 0.5% to 3.1% of ambient PM_{2.5} mass, because of the lower ambient levels during the wintertime. These estimates are consistent with expectations that primary coal emissions are a small contributor to ambient fine particulate matter mass.

Figure 8. Estimate PM_{2.5} mass contributed by primary coal emissions to the Pittsburgh Supersite during the Summer 2001 intensive.

5. Source Apportionment using PMF and UNMIX

An objective of the Pittsburgh Air Quality Study was to determine the major sources of PM_{2.5} in the Pittsburgh region. Daily 24-hour averaged filter-based data were collected for 13 months, starting in July 2001, including sulfate and nitrate data from IC analysis, trace element data from ICP-MS analysis, and organic and elemental carbon from the thermal optical transmittance (TOT) method and the NIOSH thermal evolution protocol. These data were used in two source-receptor models, Unmix and PMF.

The Unmix model created a six-factor solution with the source compositions shown in Figure 9 and source contributions shown in Figure 10. Some combinations of the species considered resulted in a solution that was not feasible. The combination of species that provided the best solution included Ca, Ti, Cr, Mn, Fe, Zn, Mo, Cd, sulfate, nitrate, OC, and EC. Three parameters designed to evaluate the model results are the minimum R², the signal-to-noise ratio, and the strength. The R² value is related to the proportion of variance of each species explained by the factors. For all species, the minimum R² value is recommended to be greater than 0.8. For the selected species, the minimum R² value was 0.86. The minimum signal-to-noise ratio is the smallest estimated signal-to-noise ratio for any of the factors in the model, recommended to be greater than 2. A value of 2.32 was obtained using this dataset. The strength is a measure of the confidence in the model. Strength is recommended to be greater than 3, but with some datasets this is unachievable and thus a strength less than 3 may still be acceptable (Henry, personal communication). For this dataset, it was impossible to find a

combination of species yielding a strength greater than 3, and the final solution had a strength of 1.41.

Figure 9. Unmix source compositions apportioned by $PM_{2.5}$ mass. All ten factors are outputs of PMF, while only the first six factors are outputs of Unmix. Although the Unmix regional transport factor includes primary OC and EC, it is graphed for comparison with the PMF sulfate factor.

Figure 10. PMF and Unmix source contributions apportioned by $\text{PM}_{2.5}$ mass. All ten factors are outputs of PMF, while only the first six factors are outputs of Unmix. Note that for comparison purposes, the Unmix regional transport factor is compared to the PMF sulfate plus primary OC and EC factors.

Species for which the ambient data correlated strongly with the source contributions (correlation coefficient greater than 0.7) allow determination of the source types. The six factors in the model have been designated a crustal material factor, a regional transport factor, a nitrate factor, an Fe, Mn, and Zn factor, a specialty steel production factor, and a cadmium factor. Descriptions of the nature of the factors, such as their contributions to $PM_{2.5}$ mass on a seasonal basis as well as a yearly average, are described below. To determine the mass contribution to $PM_{2.5}$, the total $PM_{2.5}$ mass was included as a species in the model, and the calculated source compositions and contributions were normalized by the Unmix-apportioned $PM_{2.5}$ mass. Because $PM_{2.5}$ is used as a fitting species, together these six factors account for all of the $PM_{2.5}$ mass. Figure 11 shows the Unmix source contributions normalized by the Unmix-apportioned $PM_{2.5}$ mass, averaged monthly and for the entire study.

Figure 11. Monthly average Unmix source contributions. Height of the bars corresponds to the monthly average $PM_{2.5}$ mass measured with a TEOM. The study average represents the average source contributions from July 11, 2001 through July 31, 2002. Unmix uses $PM_{2.5}$ mass as a fitting species so the mass of $PM_{2.5}$ unexplained by Unmix is less than 1%

PMF allows inclusion of more species in the model due to the consideration of uncertainties that enables handling of missing and below detection limit data. Species included in the PMF solution are $PM_{2.5}$ sulfate, nitrate, OC, EC, Mg, K, Ca, Ti, V, Cr, Mn, Fe, Ni, Cu, Zn, Ga, As, Se, Mo, Cd, Ba, and Pb. We can identify a tracer species for each factor based on the source compositions shown in Figure 9. However, a better indication is the correlation of the species ambient concentration with the PMF-modeled

source contribution. A correlation greater than 0.7 is a good indication of a tracer species. Based on the tracer species for each factor, the factors were defined as crustal material (Ca and Ti tracers), sulfate, nitrate, an Fe, Zn, and Mn factor, specialty steel (Mo and Cr tracers), cadmium, coal combustion (Ga tracer), lead, selenium, and primary OC and EC (OC and EC tracers).

A comparison of compositions and contributions for factors found by both Unmix and PMF are shown in Figures 9 and 10. These source categories include crustal material, sulfate, nitrate, steel production, specialty steel, and cadmium. The factors not found by Unmix but found by PMF are described below, and their source compositions and contributions are also shown in Figures 9 and 10. Figure 12 shows the average PMF source contributions apportioned by average $PM_{2.5}$ mass concentration. A total of 22% of the $PM_{2.5}$ measured with the TEOM is not apportioned to any source by PMF. This missing mass could be explained by species not included in the model, such as particulate ammonium, or the presence of water in the particles that was measured as $PM_{2.5}$. If all of the sulfate is assumed to be ammonium sulfate, the missing mass fraction decreases to 13%. Assuming that all of the nitrate is ammonium nitrate as well decreases the missing mass fraction to 9%.

Figure 12. Monthly average PMF source contributions. Height of the bars corresponds to the monthly average $PM_{2.5}$ mass measured with a TEOM. The study average represents the average source contributions from July 11, 2001 through July 31, 2002. The unexplained mass is the difference between the monthly average $PM_{2.5}$ mass and the sum of the monthly averaged source contributions from each factor.

Agreement between the two models is generally quite good, both in composition of sources and source contribution trends. However, there are a few significant differences. While source contributions track well for the two models, showing similar trends in

concentration with time, the magnitude of the contribution does not agree for some factors. Figure 13 shows the results of a linear regression of the Unmix source contributions against the same factors found by PMF. For this comparison, the PMF sulfate and primary OC and EC factors were added together for comparison with the Unmix regional transport factor. The PMF sulfate and primary OC and EC factors were added together for comparison with the Unmix regional transport factor.

The R^2 values for all factors are reasonable and statistically significant, ranging from 0.64 for crustal material to 0.99 for the cadmium factor. The slope of the regression line, however, ranges from 0.41 for the Fe, Mn, and Zn factor to 6.2 for the specialty steel factor. The crustal material and the cadmium factors are within 20% in source contribution magnitude, suggesting that results are robust for these factors. PMF apportions more mass to the specialty steel factor due to the inclusion of 2% of the sulfate mass. PMF apportions less mass to the sulfate and primary OC and EC factors, the nitrate factor, and the Fe, Mn, and Zn factor. The apportionment of less mass to the sulfate and primary OC and EC factors by PMF as compared to the Unmix regional transport factor is likely due to Unmix fitting the model to total $PM_{2.5}$, while PMF has a significant fraction of unexplained mass. For the nitrate and Fe, Zn, and Mn factors, the difference is due to the apportionment of OC and EC. Unmix apportions 17% of the OC mass and 17% of the EC mass to the nitrate factor whereas PMF apportions 7% OC mass and 5% EC mass to the nitrate factor. For the Fe, Mn, and Zn factor, the apportionment is 12% of the OC mass and 18% of the EC mass explained by Unmix, but only 4% of the OC mass and 7% of the EC mass is explained by the same PMF factor. Results from previous comparisons of PMF and Unmix show similar conclusions: convergence for some factors but poor agreement for others (Poirot et al., 2001; Maykut et al., 2003).

Figure 13. Linear regression results of PMF and Unmix source contributions.

In comparing Figures 11 and 12, the average source contributions as a percent of average $PM_{2.5}$ mass are within a few percent for the two models for all factors, with the

exception of the Unmix regional transport factor (68%) and the sulfate and primary OC and EC factor for PMF (total 40%). Unmix apportions all the mass, while on average PMF apportions only 78% of the mass, so some discrepancy is expected. PMF is more effective at discerning between primary and secondary OC; Unmix does not distinguish between the two and therefore can only give a large factor that is general regionally transported material and is not very informative from a policy-making perspective.

CONCLUSIONS

Significant progress was made this project period on the analysis of ambient data, source characterization, and source apportionment activities. This report highlights results from mercury emission measurements made using a dilution sampler. The measurements clearly indicate that mercury is being transformed from an oxidized to an elemental state within the dilution. Because of the effect of wall losses it is not clear whether or not this transformation occurs in the gas phase or due to some interaction with the sampler walls.

This report also presents results from an analysis that uses SAS as a marker for primary PM_{2.5} emitted from coal combustion. Primary emissions from coal combustion contribute only a small fraction of the PM_{2.5} mass at the Pittsburgh site. Ambient SAS concentrations also appear to be reasonably spatially homogeneous. Finally, SAS emission factors measured at pilot-scale appear consistent with measurements made at full-scale.

The Unmix and PMF models have been used to estimate the contribution of different sources to the PM_{2.5} mass concentrations in Pittsburgh. Comparison of the two models shows similar source composition and contribution for five factors: crustal material, nitrate, an Fe, Mn, and Zn factor, specialty steel production, and a cadmium factor. PMF found several additional factors: coal combustion, a lead factor, and a selenium factor assumed to be related to coal combustion. The PMF model found a sulfate factor separate from the OC and EC associated with primary emissions, while Unmix grouped these three species together into a single factor. Comparison between source contributions for the similar factors shows reasonable agreement between the two models. The sulfate factor shows the highest contribution to local PM_{2.5} with an annual average contribution of approximately 28% (from PMF). The nitrate, crustal material, and primary OC and EC factors also show significant contributions on the order of 10-14%. The sulfate factor is affected by photochemistry and therefore shows maximum values in summer. The nitrate factor is temperature sensitive due to the volatility of nitrate; maximum values of particulate nitrate occur in winter. The crustal material and vehicle sources somewhat more constant contributions throughout the year. The remaining factors contribute on a smaller scale and are defined by plume events, with peaks in concentration distinctly higher than average concentration.

REFERENCES

- Bein, K.J., Y. Zhao, A.S. Wexler and M.V. Johnston (2005), Speciation of size-resolved individual ultrafine particles in Pittsburgh, Pennsylvania, *J. Geophys. Res.*, 110, D07S05, doi:10.1029/2004JD004708.
- Eatough, D. J., Eatough, M. and Eatough, N. L. (1996). "Apportionment of sulfur oxides at Canyonlands during the winter of 1990 .3. Source apportionment of SO_x and sulfate and the conversion of SO₂ to sulfate in the Green River basin." *Atmospheric Environment* 30(2): 295-308.
- Fisher, G. L., Prentice, B. A., Silberman, D., Ondov, J. M., Biermann, A. H., Ragaini, R. C. and McFarland, A. R. (1978). "Physical and Morphological-Studies of Size-Classified Coal Fly-Ash." *Environmental Science & Technology* 12(4): 447-451.
- Maykut, N.; Lewtas, J.; Kim, E.; Larson, T. Source apportionment of PM_{2.5} at an urban IMPROVE site in Seattle, Washington. *Environ. Sci. Technol.* 2003, 37, 5135-5142.
- Poirot, R. L.; Wishinski, P. R.; Hopke, P. K.; Polisar, A. V. Comparative application of multiple receptor methods to identify aerosol sources in northern Vermont. *Environ. Sci. Technol.* 2001, 35, 4622-4636.
- Webber, J. S., Dutkiewicz, V. A. and Husain, L. (1985). "Identification of Submicrometer Coal Fly-Ash in a High-Sulfate Episode at Whiteface Mountain, New-York." *Atmospheric Environment* 19(2): 285-292.
- Zhao, Y., K.J. Bein, A.S. Wexler, C. Misra, P.M. Fine and C. Sioutas (2005), Field evaluation of the versatile aerosol concentration enrichment system (VACES) particle concentrator coupled to the rapid single-particle mass spectrometer (RSMS-3), *J. Geophys. Res.*, 110, D07S02, doi:10.1029/2004JD004644.

PRESENTATIONS AND PUBLICATIONS

Publications:

1. A. Khlystov, C. O. Stanier, S. Takahama, and S. N. Pandis (2005) Water content of ambient aerosol during the Pittsburgh Air Quality Study, *J. Geophys. Res.*, 110 (D7): Art. No. D07S10.
2. A. Khlystov, C. Stanier, and S.N. Pandis, "An algorithm for combining electrical mobility and aerodynamic size distributions data when measuring ambient aerosol," *Aerosol Science and Technology*, 38(S1), 229-238 (2004).
3. A. Khlystov, Q. Zhang, J. L. Jimenez, C. Stanier, S. N. Pandis, M. R. Canagaratna, P. Fine, C. Misra, and C. Sioutas (2005) In-situ concentration of semi-volatile aerosol using water-condensation technology, *J. Geophys. Res.*, submitted.

4. A. Polidori, Turpin, B., Lim H.J., Totten, L., Davidson, C., "Polarity and Molecular Weight/Carbon Weight of the Pittsburgh Organic Aerosol," *Aerosol Sci. Technol.* in preparation.
5. A. Polidori, Turpin, B., Lim, H.J., Cabada, J.C., Subramanian, R., Robinson, A., "Secondary Organic Aerosol Formation during the Pittsburgh Air Quality Study (PAQS)," *Aerosol Sci. Technol.* in preparation.
6. A.E. Wittig N. Anderson, A.Y. Khlystov, S.N. Pandis, C. Davidson, and A.L. Robinson, "Pittsburgh Air Quality Study overview," *Atmospheric Environment*, 38 (20), 3107-3125 (2004).
7. A.E. Wittig, S. Takahama, A.Y. Khlystov, S.N. Pandis, S. Hering, B. Kirby, and C. Davidson, "Semi-continuous PM_{2.5} inorganic composition measurements during the Pittsburgh Air Quality Study," *Atmospheric Environment*, 38 (20), 3201-3213 (2004).
8. A.L. Robinson, Donahue, N. M. and Rogge, W. F. (2005). "Photochemical oxidation and changes in molecular composition of organic aerosol in the regional context." *Journal of Geophysical Research-Atmospheres* In preparation.
9. A.P. Grieshop, Lipsky, E. M. and Robinson, A. L. (2005). "Measurements of In Use Vehicle Fine Particle Emissions Made in a Highway Tunnel." *Aerosol Science & Technology* in preparation.
10. Bein, K.J., Y. Zhao, A.S. Wexler and M.V. Johnston, Speciation of size-resolved individual ultrafine particles in Pittsburgh, Pennsylvania, *J. Geophys. Res.*, doi:10.1029/2004JD004708, in review.
11. C.O. Stanier, A. Khlystov, and S.N. Pandis, "Chemical processes and long-range transport of aerosols: Insights from the Pittsburgh Air Quality Study," in *Long Range Transport of Air Pollution*, Kluwer (2002).
12. C.O. Stanier, A. Khlystov, W.R. Chan, M. Mandiro, and S.N. Pandis, "A method for the in situ measurement of fine aerosol water content of ambient aerosols: The Dry-Ambient Aerosol Size Spectrometer (DAASS)," *Aerosol Science and Technology*, 38(S1), 215-228 (2004).
13. C.O. Stanier, A.Y. Khlystov, and S.N. Pandis, "Ambient aerosol size distributions and number concentrations measured during the Pittsburgh Air Quality Study (PAQS)," *Atmospheric Environment*, 38 (20), 3275-3284 (2004).
14. C.O. Stanier, A.Y. Khlystov, and S.N. Pandis, "Nucleation events during the Pittsburgh Air Quality Study: Description and relation to key meteorological, gas phase, and aerosol parameters," *Aerosol Science and Technology*, 38(S1), 253-264 (2004).

15. D. Vayenas, S. Takahama, C. Davidson, and S. N. Pandis (2005) Simulation of the thermodynamics and removal processes in the sulfate-ammonia-nitric acid system: Implications for PM_{2.5} control strategies, *J. Geophys. Res.*, 110 (D7): Art. No. D07S14.
16. D.B. Kane, J.J. Wang, K. Frost, M.V. Johnston, (2002) "Detection of Negative Ions from Individual Ultrafine Particles", *Analytical Chemistry* 74, 2092-2096.
17. D.B. Millet, N. M. Donahue, S. N. Pandis, A. Polidori, C. O. Stanier, B. J. Turpin, and A. H. Goldstein (2005) Partitioning VOCs and organic aerosols into primary and secondary sources: Results from the Pittsburgh Air Quality Study, *J. Geophys. Res.*, 110 (D7): Art. No. D07S07.
18. E. A. Weitkamp, E.M. Lipsky, P. Pancreas, J. Ondov, A. Polidori, B.J. Turpin, A.L. Robinson, "Fine Particle Emission Profile for a Large Coke Production Facility Based on Highly Time Resolved Fence Line Measurements," *Atmospheric Environment*, submitted.
19. E.M. Lipsky and A.L. Robinson, "Effects of Dilution on Fine Particle Mass and Partitioning of Semi-volatile Organics in Diesel Exhaust and Wood Smoke," *Environmental Science & Technology*, submitted.
20. E.M. Lipsky, A.L. Robinson, "Design and evaluation of a portable dilution sampling system for measuring fine particle emissions from combustion systems," *Aerosol Science & Technology*, in press.
21. G.A. Lithgow, A.L. Robinson, and S.G. Buckley, "Ambient measurements of metal-containing PM_{2.5} in an urban environment using laser-induced breakdown spectroscopy," *Atmospheric Environment*, 38 (20), 3319-3328 (2004).
22. J. Hybl, G. Lithgow, and S.G. Buckley, (2003) "Laser-Induced Breakdown Spectroscopy Detection of Biological Material," *Applied Spectroscopy* 57(10) pp 1207-1215
23. J.C. Cabada S. Rees, S. Takahama, A. Khlystov, S.N. Pandis, C.I. Davidson, and A.L. Robinson, "Mass size distributions and size resolved chemical composition of fine particulate matter at the Pittsburgh Supersite," *Atmospheric Environment*, 38 (20), 3127-3141 (2004).
24. J.C. Cabada, A. Khlystov, A.E. Wittig, C. Pilinis, and S.N. Pandis, "Light scattering by fine particles during the Pittsburgh Air Quality Study: Measurements and modeling," *Journal of Geophysical Research--Atmospheres*, 109 (D16), Art. No. D16S03 (2004).
25. J.C. Cabada, S.N. Pandis, and A.L. Robinson, "Sources of atmospheric carbonaceous particulate matter in Pittsburgh, Pennsylvania," *Journal of the Air & Waste Management Association*, 52, 732-741 (2002).

26. J.C. Cabada, S.N. Pandis, R. Subramanian, A.L. Robinson, A. Polidori, and B. Turpin, "Estimating the secondary organic aerosol contribution to PM_{2.5} using the EC tracer method," *Aerosol Science and Technology*, 38(S1), 140-155 (2004).
27. J.P. Pancras, Ondov, J. M., Gazula, S., Moore, J., Tuch, T., (2005). Highly-Time Resolved Metals Measurements with SEAS-II at Four U.S. Supersites. *Aerosol Sci. Technology*, in preparation.
28. J.P. Pancras, Ondov, J. M., Zeisler, R., (2005) Multielement Electrothermal AAS Determination of Eleven Marker Elements in Fine Ambient Aerosol Slurry Samples Collected with SEAS-II, *Analytica Chimica Acta*, in press.
29. K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston (2005), Speciation of size-resolved individual ultrafine particles in Pittsburgh, Pennsylvania, *J. Geophys. Res.*, 110, D07S05, doi:10.1029/2004JD004708.
30. K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky and A.L. Robinson, A hypothesis for the missing source of ultrafine calcium particles in the atmosphere, *Atm. Environ.*, in preparation.
31. K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, Source sampling experiments using a single particle mass spectrometer during the Pittsburgh Air Quality Study: source characterization and quantitative estimates of source contribution, *J. Geophys. Res.*, in preparation.
32. K.J. Bein, Y. Zhao, A.S. Wexler, N.J. Pekney, C.I. Davidson, M.P. Tolocka, M.V. Johnston and G. Evans, Characterization of smoke plumes from Canadian forest fires detected at three separate locations, Pittsburgh, Baltimore and Toronto, using single particle mass spectrometry, *Atm. Environ.*, in preparation.
33. L. Zhou, E. Kim, P. K. Hopke, C. Stanier, and S.N. Pandis (2005) "Mining airborne particulate size distribution data by positive matrix factorization," *J. Geophys. Res.*, 110 (D7): Art. No. D07S19.
34. L. Zhou, E. Kim, P.K. Hopke, C.O. Stanier, and S. Pandis, "The advanced factor analysis on Pittsburgh particle size-distribution data," *Aerosol Science and Technology*, 38(S1), 118-132 (2004).
35. L. Zhou, P. K. Hopke, C. O. Stanier, S. N. Pandis, J. M. Ondov, and J. P. Pancras (2005) Investigation of the relationship between chemical composition and size distribution of airborne particles by Partial Least Square (PLS) and Positive Matrix Factorization (PMF), *J. Geophys. Res.*, 110 (D7): Art. No. D07S18.
36. L. Zhou, W. Liu, and P.K. Hopke, (2004) "Comparison of Two Trajectory Based Models for Locating Particle Sources for Two Rural New York Site", *Atmospheric Environment* 38 (13): 1955-1963.

37. L.M. Zhou, P.K. Hopke, P. Paatero, J.M. Ondov, J.P. Pancras, N.J. Pekney, and C.I. Davidson, "Advanced factor analysis for multiple time resolution aerosol composition data," *Atmospheric Environment*, 38 (29), 4909-4920 (2004).
38. N.J. Pekney, C.I. Davidson, A.L. Robinson, L. Zhou, Hopke, P.K. D.J. Eatough "Identification of Major Sources of PM_{2.5} in Pittsburgh using PMF and Unmix," *Aerosol Science & Technology*, submitted.
39. N.J. Pekney, K.J. Bein, C.I. Davidson, A.S. Wexler and M.V. Johnston, Identification of sources of atmospheric PM at the Pittsburgh Supersite: RSMS III and filter-based Positive Matrix Factorization, *Atm. Environ.*, in preparation.
40. Q. Zhang, C. O. Stanier, M. R. Canagaratna, J. T. Jayne, D. R. Worsnop, S. N. Pandis, and J. L. Jimenez (2004) Insights into the chemistry of new particle formation and growth events in Pittsburgh based on aerosol mass spectrometry, *Environ. Sci. Tech.*, 38 (18): 4797-4809.
41. Q. Zhang, M. R. Canagaratna, J. T. Jayne, D. R. Worsnop, and J. L. Jimenez (2005) Time and size-resolved chemical composition of submicron particles in Pittsburgh: Implications for aerosol sources and processes, *J. Geophys. Res.*, 110 (D7): Art. No. D07S09.
42. R. Subramanian, A.Y. Khlystov, J.C. Cabada, and A.L. Robinson, "Positive and negative artifacts in particulate organic carbon measurements with denuded and undenuded sampler configurations," *Aerosol Science and Technology*, 38(S1), 27-48 (2004).
43. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Selection of Source Profiles for Chemical Mass Balance Modeling using Organic Molecular Markers." *Environmental Science & Technology* Under review.
44. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Source Apportionment of Gasoline and Diesel Vehicles using the Chemical Mass Balance Model and Molecular Markers." *Environmental Science & Technology* In preparation.
45. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Source Apportionment of Primary Organic Aerosol in Pittsburgh, PA using organic molecular markers." *Environmental Science & Technology* In preparation.
46. R. Subramanian, Khlystov, A. Y. and Robinson, A. L. (2005). "Measurement of elemental carbon using the Thermal-Optical Transmittance Technique: Fundamental assumptions and effect of peak inert-mode temperature." *Aerosol Science and Technology*, In preparation.

47. R.R. Anderson, W. K. Modey, D. V. Martello, L. J. Lucas, C. I. Davidson, and D. J. Eatough (2005) Apportionment of ambient primary and secondary pollutants during a 2001 summer study in Pittsburgh using EPA UNMIX, *J. Geophys. Res.*, submitted.
48. S. Takahama, A.E. Wittig, D.V. Vayenas, C.I. Davidson, and S.N. Pandis, "Modeling the diurnal variation of nitrate during the Pittsburgh Air Quality Study," *Journal of Geophysical Research--Atmospheres*, 109 (D16), Art. No. D16S06 (2004).
49. S.L. Rees, A.L. Robinson, A. Khlystov, C.O. Stanier, and S.N. Pandis, "Mass balance closure and the Federal Reference Method for PM_{2.5} in Pittsburgh, Pennsylvania," *Atmospheric Environment*, 38 (20), 3305-3318 (2004).
50. S.N. Pandis, "Estimates of diesel and other emissions: Overview of the Supersite program," in *Improving Estimates of Diesel and Other Emissions for Epidemiological Studies*, HEI Communication 10, Health Effects Institute, Boston, MA, (2003).
51. S.S. Park, Pancras, P., Ondov, J. M., Robinson, A. 2005, "Application of the Pseudo-Deterministic Multivariate Receptor Model to resolve Power Plant Influences on Pittsburgh Air Quality." *Atmos. Environ.*, in preparation.
52. T.M. Gaydos, C.O. Stanier CO, S.N. Pandis (2005) "Modeling of in situ ultrafine atmospheric particle formation in the eastern United States," *J. Geophys. Res.*, 110 (D7): Art. No. D07S12.
53. W. Tang, T. Raymond, B. Wittig, C. Davidson, S.N. Pandis, A. Robinson, and K. Crist, (2004) "Spatial variations of PM_{2.5} during the Pittsburgh Air Quality Study," *Aerosol Science and Technology*, 38(S2): 80-90.
54. W.K. Modey, D.J. Eatough, R.R. Anderson, D.V. Martello, S. Takahama, L.L. Lucas, and C.I. Davidson, "Ambient fine particulate concentrations and chemical composition at two sampling sites in metropolitan Pittsburgh: a 2001 intensive summer study," *Atmospheric Environment*, 38 (20), 3165-3178 (2004).
55. Y. Zhao, K. J. Bein, A. S. Wexler, C. Misra, P. M. Fine, and C. Sioutas (2005) Field evaluation of the VACES particle concentrator coupled to the RSMS-3 single particle mass spectrometer, *J. Geophys. Res.*, 110 (D7): Art. No. D07S02.

Presentations:

1. "Investigation of nucleation bursts in the Pittsburgh air quality study", 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (C. O. Stanier, A. Y. Khlystov, and S. N. Pandis).
2. "Monitoring of water content of ambient aerosol during the Pittsburgh Air Quality Study" 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (A. Y. Khlystov, C. O. Stanier, D. Vayenas, and S. N. Pandis).

3. Performance of the Aerodynamic Particle Sizer 3320 during the Pittsburgh Air Quality Study (PAQS)” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (A. Khlystov, C. Stanier, and S. N. Pandis).
4. “Sulfate-ammonia-nitric acid interactions in an urban area” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (S. Takahama, A. Khlystov, B. Wittig, S. V. Hering, C. Davidson, A. Robinson, and S. N. Pandis).
5. “Sampling artifacts during measurement of ambient carbonaceous aerosol” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (R. Subramanian, A. Y. Khlystov, J. C. Cabada, S. N. Pandis, and A. L. Robinson).
6. “Formation and properties of regional aerosol: Some insights from the Pittsburgh Air Quality Study”, NASA-GSFC, Greenbelt MD, May 2002,(C. Stanier, A. Khlystov, S. Rees, J. Cabada, A. Robinson, C. Davidson, and S. N. Pandis)
7. “Seasonal composition of PM_{2.5} and performance of the Federal Reference Method in Pittsburgh”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (S. L. Rees, S. Takahama, A. L. Robinson, A. Khlystov, and S. N. Pandis).
8. “Continuous measurements of ammonia, sulfate, and nitrate in Pittsburgh: Implications for PM_{2.5} control strategies”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (B. Wittig, A. Khlystov, S. Takahama, C. Davidson, A. Robinson, S. Hering, and S. N. Pandis).
9. “The contribution of long-range transport and secondary organic aerosol to PM_{2.5} in Pittsburgh”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (J. C. Cabada, R. Subramanian, S. N. Pandis, A. L. Robinson, W. Tang, N. J. Anderson, T. Raymond, and C. I. Davidson).
10. “The Dry-Ambient Size Spectrometer: A new technique for the automatic on-line measurement of the atmospheric aerosol water size distribution”, Annual Meeting of American Geophysical Union, San Francisco, December 2001 (A. Khlystov, C. O. Stanier, S. N. Pandis).
11. “The July 2001 intensive of the Pittsburgh Air Quality Study”, Annual Meeting of AAAR, Portland, Oregon, October 2001 (C. I. Davidson, A. L. Robinson, and A. Khlystov, S. N. Pandis).
12. “Sources of atmospheric carbonaceous particulate matter in Pittsburgh”, Annual Meeting of AAAR, Portland, Oregon, October 2001 (J. Cabada, S. N. Pandis and A. L. Robinson).
13. “Automated measurements of dry and wet ambient aerosol distributions”, Annual Meeting of AAAR, Portland, Oregon, October 2001 (A. Y. Khlystov, W. R. Chan, C. O. Stanier, M. Mandiro, and S. N. Pandis)

14. "Continuous measurements of ammonia and ammonium in ambient air", Annual Meeting of AAAR, Portland, Oregon, October 2001 (A. Khlystov, J. Sauser, R. Otjes, and S. N. Pandis).
15. The contribution of secondary organic aerosol to PM_{2.5} concentrations in Pittsburgh, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (J. C. Cabada, S. N. Pandis, A. L. Robinson, R. Subramanian, A. Polidori, and B. Turpin).
16. Preliminary results from the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (S. N. Pandis, C. I. Davidson, A. L. Robinson, and A. Y. Khlystov)
17. Monitoring of water content of ambient aerosol during the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (A. Y. Khlystov, C. O. Stanier, D. Vayenas, and S. N. Pandis)
18. Investigation of nucleation bursts during the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (C. O. Stanier, A. Y. Khlystov, B. Wittig, S. N. Pandis, Y. Zhou, K. Bein, A. S. Wexler, C. Misra, and C. Sioutas)
19. Atmospheric particulate matter: Physics, chemistry, and Chemical Transport Models, PM AAAR 2003, Pittsburgh PA March 2003 (B. Koo, K. Fahey, T. Gaydos, and S. N. Pandis)
20. Secondary organic aerosol contribution to carbonaceous PM_{2.5} concentrations in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (J. C. Cabada, S. N. Pandis, B. Wittig, A. Robinson, R. Subramanian, A. Polidori, and B. J. Turpin)
21. Using ultrafine concentrators to increase the hit rates of single particle mass spectrometers, PM AAAR 2003, Pittsburgh PA March 2003 (Y. Zhao, K. J. Bein, A. S. Wexler, C. Misra, P. M. Fine, and C. Sioutas)
22. PM_{2.5} Federal Reference Method performance relative to mass balance closure, PM AAAR 2003, Pittsburgh PA March 2003 (S. L. Rees, A. L. Robinson, A. Khlystov, C. O. Stanier, and S. N. Pandis)
23. Examining the assumptions behind elemental carbon measurements using the thermal-optical transmittance technique, PM AAAR 2003, Pittsburgh PA March 2003 (R. Subramanian, A. Y. Khlystov, and A. L. Robinson)
24. Spatial variations of PM_{2.5} during intensive sampling of PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (W. Tang, C. I. Davidson, T. R. Raymond, S. N. Pandis, B. Wittig, A. Khlystov, and A. L. Robinson)
25. Fenceline sampling adjacent to a large coke production facility in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (E. A. Weitkamp, E. Lipsky, A. Robinson, N. Anderson, H. Leifeste, R. Subramanian, J. Cabada, A. Khlystov, C.

- Stanier, L. Lucas, S. Takahama, B. Wittig, C. Davidson, S. Pandis, A. Polidori, H. J. Lim, B. Turpin, P. Pancras, and J. Ondov)
26. In-use vehicle emissions source characterization study: Squirrel Hill tunnel Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (E. M. Lipsky, A. Robinson, N. Anderson, H. Leifeste, R. Subramanian, J. Cabada, S. Rees, A. Khlystov, C. Stanier, L. Lucas, S. Takahama, B. Wittig, C. Davidson, S. N. Pandis, A. Polidori, H. J. Lim, and B. Turpin)
 27. Water content of ambient aerosol during PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (A. Khlystov, C. Stanier, and S. N. Pandis)
 28. Diurnal and seasonal trends in outdoor particle size distributions measured at urban and rural locations during PAQS (C. Stanier, A. Khlystov, and S. N. Pandis)
 29. Mass and chemically resolved size compositions of fine particulate matter at the Pittsburgh Supersite, PM AAAR 2003, Pittsburgh PA March 2003, (J. C. Cabada, S. N. Pandis, S. Rees, S. Takahama, A. Khlystov, A. L. Robinson, and C. I. Davidson)
 30. Simulation of the atmospheric aerosol size/composition distribution in a three-dimensional chemical transport model, PM AAAR 2003, Pittsburgh PA March 2003 (T. M. Gaydos, K. M. Fahey, B. Koo, and S. N. Pandis)
 31. Application of PMCAMx to the South Coast Air Basin and the Eastern United States, PM AAAR 2003, Pittsburgh PA March 2003 (B. Koo, K. M. Fahey, T. M. Gaydos, and S. N. Pandis)
 32. Principal component analysis of trace elements in PM_{2.5} in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (N. J. Anderson, C. I. Davidson, S. N. Pandis, A. Robinson, and A. Khlystov)
 33. Source apportionment using particle size distribution data from PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (L. Zhou, E. Kim, P. K. Hopke, C. Stanier, and S. N. Pandis)
 34. Highly time-resolved measurements of elemental composition at the Baltimore, St. Louis, and Pittsburgh Supersites using the UM High Frequency Aerosol Slurry Sampler: Unprecedented resolution of the sources of primary atmospheric aerosol PM AAAR 2003, Pittsburgh PA March 2003 (J. M. Ondov, J. Pancras, S. Gazula, M. Yu, J. Turner, A. Robinson, S. N. Pandis, N. D. Poor, and R. K. Stevens)
 35. "Size Resolved Chemical Classification of Dual Polarity Single-Ultrafine-Particle Mass Spectrometry Data Collected During Pittsburgh Supersite Experiment", K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.

36. "Dynamic Data Classification Using a Component-Weighted Similarity Algorithm", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.
37. "Diurnal Variations of Ultra-fine Particles in Pittsburgh Measured by a Rapid Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.
38. "Rapid Single Particle Mass Spectrometry and EPA Supersite Experiments", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, NYU Institute of Environmental Medicine, Tuxedo, NY, December 2002.
39. "Speciation of Size-Resolved Individual Ultrafine Particles in Pittsburgh, Pennsylvania", *K.J. Bein, Y. Zhao, A.S. Wexler, D.A. Lake, D.B. Kane, M.P. Tolocka, K.P. Rhoads, M.V. Johnston and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
40. L. Zhou, E. Kim, and P.K. Hopke Analysis of the Particle Size Distribution Data Collected During Pittsburgh Air Quality Study, Presented to 21st Annual Conference of the American Association for Aerosol Research, Charlotte, NC, October 7-11, 2002.
41. L. Zhou, E. Kim, P.K. Hopke, C. Stanier, and S. Pandis Source Apportionment Using Particle Size Distribution Data from the Pittsburgh Air Quality Study (PAQS), presented to 2003 AAAR PM Meeting on Particulate Matter: Atmospheric Sciences, Exposure and the Fourth Colloquium on PM and Human Health, Pittsburgh, PA, March 30 to April 4, 2003.
42. L. Zhou, P.K. Hopke, and W. Liu, Comparison of three back trajectory based models--PSCF, SQTBA, residence-time weighted concentration model for identifying particle sources for Potsdam and Stockton, New York, presented to 21st Annual Conference of the American Association for Aerosol Research, Anaheim, CA, October 20-24, 2003.
43. L. Zhou, P.K. Hopke, and P. Paatero, Advanced Factor Analysis for Aerosol Composition Data with Various Temporal Resolutions, presented to 21st Annual Conference of the American Association for Aerosol Research, Anaheim, CA, October 20-24, 2003.
44. S.G. Buckley, F. Ferioli, G.A. Lithgow, "Combustion System Analysis Using Laser-Induced Breakdown Spectroscopy," Paper 403, 22nd International Congress on Applications of Lasers and Electro-Optics, Jacksonville, FL, October 13-16, 2003.

45. G.A. Lithgow, A.L. Robinson, and S.G. Buckley, "Ambient Measurements of Inorganic Species in an Urban Environment Using LIBS," Second International Conference on Laser-Induced Plasma Spectroscopy, Orlando, FL, September 24 – 27, 2002.
46. G.A. Lithgow, A.L. Robinson, S.G. Buckley, "Ambient Particle Measurements in an Urban Environment Using Laser-Induced Breakdown Spectroscopy (LIBS)," Paper 11E4, American Association of Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
47. Ondov, J. M. (2001). Sources of Metals Influencing Air Quality at the Pittsburgh Supersite: Data collected with the UM High-Frequency Aerosol Slurry Sampler.
48. Ondov, J. M. (2002). Unprecedented Source Apportionment with the University of Maryland Semi-Continuous Elements in Aerosol System Presented at the Regional AQ Modeling & Data Analysis Meeting, sponsored by MARAMA, OTC, NESCAUM and MANE-VU, January 23-24, 2002, Baltimore, MD.
49. Ondov, J. M., (2003) Short-Term Concentrations of Metals SEAS via SEAS and Selected Variables. Presented at the Mid-Atlantic Region Air Monitoring Association Conference in Baltimore, Jan 21-22, at the Harbor Court Hotel.
50. Ondov, J. M., Pancras, J. P., Gazula, S., Moore, J. A., Park, S. S., Chang, Y. C., Squib, K., Powel, J., Mitkus, R., Turner, J., Yu, M.N.S., Robinson, A., Pandis, S., Davidson, C. (2003). Identification of Sources From High-Frequency Elements Measurements at 4 Supersites. Semi-annual EPA Supersites meeting in Atlanta, Jan 22-23.
51. Ondov, J. M. (2003). High-frequency metals measurements at the Pittsburgh Supersite and Coke Plant site with the University of Maryland SEAS. Seminar presented at the Pittsburgh Supersite Project workshop, March 5, Carnegie Mellon University.
52. Ondov, J. M. (2003) Highlights of SEAS Metals Data for the Pittsburgh Supersite. Presented at the Pittsburgh Air Quality Study Meeting, Pittsburgh, PA, April 4,5, 2003.
53. Ondov, J. M. (2003) New Paradigm for Air Pollution Control: Pseudo Deterministic Receptor Modeling of Highly Time Resolved Ambient Aerosol Composition Data Derived from the UMCP SEAS. Seminar presented to the Source Apportionment Group, US EPA, Research Triangle Park, November 12.
54. Ondov, J. M., Emission Rates of Pollutants from Stationary Sources Using Highly Time Resolved Ambient Measurements and a New Pseudo Deterministic Hybrid Receptor Model. Presented at the EPA Center for Hazardous Substances in Urban Environments, Research Program Internal Workshop, January 5, 2004.

55. Ondov, J. M., New Pseudo Deterministic Model for Individual Source Apportionment Using Highly Time Resolved Data. Presented at the MARAMA MANE-VU Science Meeting, Baltimore, MD, January 27-29, 2004.
56. Park, S. S., Pancras, J. P., Gazula, S., Ondov, J. M. (2002) Sources of Elemental Aerosol Constituents in Pittsburgh Using Positive Matrix Factorization of Highly Time-resolved Data. Presented at the American Association of Aerosol Research meeting, 21st Annual AAAR Conference October 7-11, Charlotte.
57. Ondov, J. M., Pancras, J. P., Gazula, S., Yu, M. N. S., Turner, J., Robinson, A., Pandis, S., Stevens, R. K., Poor (2003). Highly Time-Resolved Measurements of Elemental Composition at the Baltimore, St. Louis, Pittsburgh, and Tampa Supersites Using the UM High-Frequency Aerosol Slurry Sampler: Unprecedented Resolution of the Sources of Primary Atmospheric Aerosol. To be presented at the Association of Aerosol Research meeting, Particulate Matter: Atmospheric Sciences, Exposure, and the Fourth Colloquium on PM and Human Health, March, Pittsburgh.
58. Ondov, J. M., Pancras, J. P., Park, S. S., Poor, N., Turner, J. R., Yu, M., Lipsky, E., Weitkamp, E., Robinson, A. (2003) PM emission rates from highly time-resolved ambient concentration measurements. Presented at the October meeting of the American Society for Aerosol Research, San Diego.
59. Ondov, J. M., Poor, N. (2003) Emission Inventory Development through highly-time-resolved ambient sampling. NARSTO Workshop on Innovative Methods for Emission-Inventory Development and Evaluation, University of Texas, Austin; October 14-17, 2003.
60. G.A. Lithgow and S.G. Buckley, "Detection and Classification of Biological Aerosols Using Laser-Induced Breakdown Spectroscopy," Paper 10B4, American Association of Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
61. S.G. Buckley "Laser-Induced Breakdown Spectroscopy for Detection of Biological Aerosols - Potential and Perspective," PITTCO 2005, Orlando FL, Feb. 27-Mar. 4, 2005. (invited)
62. S.G. Buckley "LIBS as a Combustion and Aerosol Diagnostic," PacificChem 2005, Honolulu, HI, Dec 15-20, 2005. (invited)
63. Cabada, J.C., Pandis, S.N., Robinson, A.L., Davidson, C.I., Polidori, A., Turpin, B.J., and Subramanian, R., "The contribution of Secondary Organic Aerosol to PM_{2.5} in Pittsburgh," Presented at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
64. Offenberg, J.H., Polidori, A., Porcja, R., Turpin, B.J., "Functional group composition by size and polarity in Pittsburgh, PA and insights into aerosol

- processing,” Presented at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
65. Polidori, A., Turpin, B.J., Lim, H.J., Robinson, A., Subramanian, R., Cabada, J.C., “Semi-continuous organic particulate matter measurements during Pittsburgh Air Quality Study (PAQS),” Poster presentation at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
 66. Subramanian, R., Robinson, A.L., Cabada, J.C., Pandis, S.N., Wittig, B., Polidori, A., Turpin, B.J., Hering, S.V., Modey, W.K., Eatough, D.J., “Intercomparison of ambient carbonaceous aerosol samplers used during the Pittsburgh Air Quality Study,” Poster presentation at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
 67. Cabada, J.C., Pandis, S.N., Robinson, A.L., Davidson, C.I., Polidori, A., Turpin, B.J., and Subramanian, R., “The contribution of Secondary Organic Aerosol to PM_{2.5} in Pittsburgh,” Platform Presentation at the Annual Meeting of the American Institute of Chemical Engineers, Indianapolis, IN, November 2002.
 68. Millet, D.B., Donahue, N.M., Polidori, A., Stanier, C.O., Turpin, B.J., Goldstein, A.H., “VOC-Aerosol Relationships at the Pittsburgh Air Quality Study”, Presented at the Berkeley Atmospheric Science Center symposium, Berkeley, CA, September 2003.
 69. Polidori, A., Turpin, B., Lim, H.J., Cabada, J.C., Subramanian, R., Robinson, A., “Secondary Organic Aerosol Formation During the Pittsburgh Air Quality Study (PAQS)”, Presented at the Environmental Sciences Graduate Student Association (ESGSA), New Brunswick, NJ, June 2004.
 70. Polidori, A., Turpin, B.J., Lim, H.J., Totten L., Davidson, C., “Characterization of the Organic Fraction of Atmospheric Aerosols” Platform Presentation at the American Association for Aerosol Research Annual Conference, Atlanta, GA, October 2004.
 71. Robinson, A.L., Subramanian, R., Gaydos, T., Pandis, S.N., Bernardo-Bricker, A., Rogge, W.F., Polidori, A., Turpin, B.J., Clarke, L., Hernandez, M., “Synthesis of Source Apportionment Estimates of Organic Aerosol in the Pittsburgh Region” Platform Presentation at the American Association for Aerosol Research Annual Conference, Atlanta, GA, October 2004.
 72. Polidori, A., Turpin, B., Lim, H.J., Subramanian, R., Robinson, A., Pandis, S., Cabada, J.C., “Local and Regional Secondary Organic Aerosol Formation: Insights from a Year at Pittsburgh and Comparisons with Los Angeles and Atlanta,” Platform Presentation at the American Association for Aerosol Research “Supersite” Conference, Atlanta, GA, February 2005.

73. Weitkamp, E., Lipsky, E., Robinson, A., Polidori, A., Turpin, B., Pancras, P., Ondov, J., Bernado-Bricker, A., Vasquez, O., Rogge, W., "Fine Particle Emission Profile for a Large Coke Production," Poster presentation at the American Association for Aerosol Research "Supersite" Conference, Atlanta, GA, February 2005.
74. "Source Sampling and Characterization Using a Single Particle Mass Spectrometer during the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
75. "Size-Resolved Chemical Classification of Single Particle Mass Spectrometry Data Collected during the Pittsburgh Supersite Experiment: Source Attribution", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
76. "Laboratory Experiments Examining Ultrafine Particle Production by Re-breathing of Road Dust through a Diesel Engine", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky and A.L. Robinson, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
77. "Detection of a Contaminating Plume during a Roadway Tunnel Source Sampling Experiment Using a Single Particle Mass Spectrometer", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
78. "Detection and Characterization of a Smoke Plume from Canadian Forest Fires during the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler, N.J. Pekney, C.I. Davidson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
79. "Field Evaluation of the VACES Particle Concentrator Coupled to the RSMS-3 Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, C. Misra, P.M. Fine and C. Sioutas, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
80. "Identification of Sources of Atmospheric PM at the Pittsburgh Supersite: RSMS-III and Filter-based Positive Matrix Factorization", *N.J. Pekney, K.J. Bein, C.I. Davidson, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
81. "Laboratory Experiments Examining Ultrafine Particle Production by Re-breathing of Road Dust through a Diesel Engine", *K.J. Bein, Y. Zhao, A.S.

- Wexler, E. Lipsky and A.L. Robinson, American Association for Aerosol Research Annual Conference, Atlanta, GA, October 4-8, 2004.
82. "Measurements of Nitrate Particles in Pittsburgh, PA, Using Rapid Single Particle Mass Spectrometry", *Y. Zhao, K.J. Bein, A.S. Wexler, M.P. Tolocka and M.V. Johnston, American Association for Aerosol Research Annual Conference, Atlanta, GA, October 4-8, 2004.
 83. "Size Resolved Chemical Classification of Dual Polarity Single-Ultrafine-Particle Mass Spectrometry Data Collected During the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
 84. "Dynamic Data Classification Using a Component-Weighted Similarity Algorithm", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
 85. "Diurnal Variations of Ultra-fine Particles in Pittsburgh Measured by a Rapid Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
 86. "Experimental Validation of a High Transmission Rate Inlet for Ultra-fine Single Particle Mass Spectrometers", *Y. Zhao, K.J. Bein, P. Middha and A.S. Wexler, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
 87. "Synthesis of Research on In-situ Particle Nucleation in Western Pennsylvania: Which Hypotheses for Nuclei Formation and Growth are Consistent with Field Observations, Mass Spectrometry, and Modeling?.", *C. Stanier, T. Gaydos, A.Y. Khlystov, Q. Zhang, M. Caragaratna, J. Jayne, D. Worsnop, J. Jimenez, D. Millet, A. Goldstein, K.J. Bein, Y. Zhao, A.S. Wexler and S.N. Pandis, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
 88. "Speciation of Size-Resolved Individual Ultrafine Particles in Pittsburgh, Pennsylvania", *K.J. Bein, Y. Zhao, A.S. Wexler, D.A. Lake, D.B. Kane, M.P. Tolocka, K.P. Rhoads, M.V. Johnston and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
 89. "Using Ultrafine Concentrators to Increase the Hit Rate of Single Particle Mass Spectrometers", *Y. Zhao, K.J. Bein, A.S. Wexler, C. Misra, P.M. Fine and C. Sioutas, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.

90. "Investigation of Urban Aerosols at the Baltimore Particulate Matter Supersite by Single Particle Mass Spectrometry", *D.A. Lake, M.P. Tolocka, M.V. Johnston, K.J. Bein, Y. Zhao, A.S. Wexler, K.P. Rhoads, and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
91. "Investigation of Nucleation Bursts during the Pittsburgh Air Quality Study, *C. Stanier, A.Y. Khlystov, B. Wittig, S.N. Pandis, K.J. Bein, Y. Zhao, A.S. Wexler, C. Misra and C. Sioutas, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
92. G. Casuccio*, T. Lersch, R.J. Lee, A. Robinson, E. Weitkamp, and D. Martello, "Estimating of the Impact of Primary Coal Fired Boiler Emissions to PM_{2.5} And PM₁₀ Using Computer Controlled Scanning Electron Microscopy Data," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005.
93. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson, and M.V. Johnston, "Source Sampling and Characterization Using a Single Particle Mass Spectrometer During the Pittsburgh Supersite Experiment," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005.
94. M. Hernandez*, L. Clarke, L. Angenent, and A. Robinson, "Primary Biopolymer Associations with Fine Particulate Matter," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005.
95. S.S. Park*, J.P. Pancras, J.M. Ondov, A. Robinson, and C. Davidson, "Application of the UM Multivariate Pseudo-Deterministic Receptor Model to Resolve Power Plant Influences on Air Quality at the CMU Supersite," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005.
96. R. Subramanian*, A. Robinson, and A. Khlystov, "OC/EC Analysis with Thermal-Optical Methods: Effects of Temperature Protocol and Non-Carbonaceous Compounds," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005.
97. A. Wittig*, A. Robinson, and S. Pandis, "Continuous PM_{2.5} Mass Measurements at the Pittsburgh Air Quality Study Supersite," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7-11, 2005. (poster)
98. W. Tang, T. Raymond, B. Wittig, C. Davidson*, S. Pandis, A. Robinson, and K. Crist, "Spatial Variations of PM_{2.5} in the Pittsburgh Region," American

Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.

99. A. Polidori*, B. Turpin, H.J. Lim, R. Subramanian, A. Robinson, S. Pandis, and J.C. Cabada, “Local and Regional Secondary Organic Aerosol Formation: Insights from a Year at Pittsburgh and Comparison with Los Angeles and Atlanta,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
100. T. Lane, R. Pinder, M. Shrivastava, S.N. Pandis, and A.L. Robinson*, “Evaluation of the Primary Organic Carbon Emission Inventory for the Eastern United States,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
101. O. Sevimoglu*, W.F. Rogge, A. Bernardo, A. Robinson and R. Subramanian, “Fine Particulate Abrasion Products from Leaf Surfaces of Urban Plants: Comparison Between Los Angeles and Pittsburgh,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
102. A. Bernardo-Bricker*, W.F. Rogge, O. Sevimoglu, A. Robinson, E. Lipsky, and A. Grieshop, “Source Profiles for Organic PM_{2.5} for Diesel Trucks and Gasoline Vehicles Determined for the Squirrel Hill Tunnel in Pittsburgh, PA,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
103. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, and A.L. Robinson, “Laboratory Experiments Examining Ultrafine Particle Production by Re-Breathing of Road Dust Through a Diesel Engine,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
104. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson, and M.V. Johnston, “Detection of a Contaminating Plume During a Roadway Tunnel Source Sampling Experiment Using a Single Particle Mass Spectrometer,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
105. A.L. Robinson*, S.L. Rees, A. Khlystov, C.O. Stanier, and S.N. Pandis, “Mass Balance Closure and the Federal Reference Method for PM_{2.5} in Pittsburgh, Pennsylvania,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.

106. R. Subramanian*, A. Robinson, A. Bernardo-Bricker, and W.F. Rogge, "Organic Carbon Mass Balance and Source Apportionment of Primary Organic Carbon in the Pittsburgh Region Using Molecular Markers," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
107. A.L. Robinson*, E.M. Lipsky, N. Pekney, W.F. Rogge, A. Bernardo-Bricker, and O. Sevimglu, "Fine Particle Emission Profile for Road Dust in Pittsburgh, Pennsylvania," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
108. E. Weitkamp*, E. Lipsky, A. Robinson, A. Polidori, B. Turpin, P. Pancras, J. Ondov, A. Bernardo-Bricker, O.R. Vasquez, and W.F. Rogge, "Fine Particle Emission Profile for a Large Coke Production, American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
109. A. Grieshop*, E. Lipsky, and A. Robinson, "Fuel-Based Particulate Matter and Gaseous Emission Factors Determined from Vehicles in Pittsburgh, PA's Squirrel Hill Tunnel," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
110. A.L. Robinson*, N.M. Donahue, K. Huff Hartz, A. Sage and E. Weitkamp, "Field and Laboratory Experiments Examining the Stability of Organic Molecular Markers Used for Source Apportionment," American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
111. A.L. Robinson*, N.M. Donahue, A.M. Sage, K.E. Huff Hartz, and E.A. Weitkamp, "Field and Laboratory Experiments Examining the Stability of Organic Molecular Markers Used for Source Apportionment," 2004 AGU Fall Meeting, San Francisco, CA, December 13–17, 2004.
112. A.L. Robinson*, R. Subramanian, A. Bernardo-Bricker, and W.F. Rogge, "Source Apportionment of Primary Organic Carbon in the Pittsburgh Region using Molecular Markers and Different Receptor Models," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
113. K.E. Huff Hartz*, E.A. Weitkamp, A.M. Sage, A.A. Presto, A.L. Robinson, and N.M. Donahue, "Kinetics of Atmospheric Processing of Organic Particulate Matter: A Relative Rates Approach," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)

114. E.A. Weitkamp*, K.E. Huff Hartz, A.M. Sage, A.L. Robinson, N.M. Donahue, W.F. Rogge, and A. Bernardo-Bricker, "Evaluation of the Oxidation Kinetics of Molecular Markers Used for Source Apportionment of Primary Organic Aerosol," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
115. A.L. Robinson*, R. Subramanian, T. Gaydos, S.N. Pandis, A. Bernardo-Bricker, W.F. Rogge, A. Polidori, B.J. Turpin; L. Clarke, and M. Hernandez, "Synthesis of Source Apportionment Estimates of Organic Aerosol in the Pittsburgh Region," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
116. E. M. Lipsky* and A.L. Robinson, "Effects of Dilution Ratio and Residence Time on the Partitioning of Semi-Volatile Organic Carbon in Emissions from a Wood Stove And Diesel Engine," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
117. K.J. Bein*, Y. Zhao, A.S. Wexler, E.M. Lipsky, and A.L. Robinson, "Laboratory Experiments Examining Ultrafine Particle Production by Rebreathing of Road Dust Through a Diesel Engine," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
118. A.M. Sage*, K.E. Huff Hartz, E.A. Weitkamp, A.L. Robinson, and N.M. Donahue "Laboratory Measurement of Heterogeneous Oxidation Kinetics of Organic Aerosols," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
119. A.P. Grieshop*, E.M. Lipsky, and A.L. Robinson, "Fuel-Based Particulate Matter and Gaseous Emission Factors determined from Vehicles in Pittsburgh, PA's Squirrel Hill Tunnel," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
120. N.M. Donahue* and A.L. Robinson, "Theoretical, In Situ, and Laboratory Constraints on Organic Aerosol Oxidation," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
121. C. P. Rose*, C. Torrey, V. Aleven, A. Robinson, C. Wu, and K. Forbus, "CycleTalk: Towards a Dialogue Agent that Guides Design with an Articulate Simulator," ITS 2004 7th International Conference on Intelligent Tutoring Systems, Maceió, Alagoas, Brazil August, 30–September 03, 2004.
122. A.L. Robinson*, "Are Organic Molecular Markers Used for Source Apportionment Stable?," presented at Telluride Workshop on Atmospheric Aerosol Aging, Telluride, CO, August 9–13, 2004. (invited)

123. A.L. Robinson*, "Comparison of Emission- and Receptor-based Model Predictions of Primary Organic Aerosol in Pittsburgh, PA," presented at LADCO Academic Community Meeting, Chicago, IL, June 11, 2004. (invited)
124. A.L. Robinson*, "Sources of Organic Particulate Matter in Pittsburgh, PA," presented at Mid-Atlantic Regional Air Management Association Science Meeting (MANE-VU/MARAMA 2004), Baltimore, MD, January 27–29, 2004 (invited).