PROJECT TITLE: Atmospheric Aerosol Source-Receptor Relationships: The Role Of Coal-Fired Power Plants

DOE AWARD #: DE-FC26-01NT41017

REPORT: Semi-Annual Technical Progress Report

REPORTING PERIOD: March 2005 – August 2005

ISSUED: December 2005

SUBMITTED TO: US Department of Energy
National Energy Technology Laboratory
P.O. Box 10940
626 Cochrans Mill Rd
Pittsburgh, PA 15236-0940
Attn: Bill Aljoe

PREPARED BY: Carnegie Mellon University
5000 Forbes Ave
Pittsburgh, PA 15213

PIs: Allen L. Robinson, Spyros N. Pandis, Cliff I. Davidson
(412) 268 – 3657; (412) 268 – 3348 (fax)
e-mail: alr@andrew.cmu.edu

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ABSTRACT

This report describes the technical progress made on the Pittsburgh Air Quality Study (PAQS) during the period of March 2005 through August 2005. Significant progress was made this project period on the source characterization, source apportionment, and deterministic modeling activities.

This report highlights new data on road dust, vegetative detritus and motor vehicle emissions. For example, the results show significant differences in the composition in urban and rural road dust. A comparison of the organic of the fine particulate matter in the tunnel with the ambient provides clear evidence of the significant contribution of vehicle emissions to ambient PM. The source profiles developed from this work are being used by the source-receptor modeling activities.

The report presents results on the spatial distribution of PMF-factors. The results can be grouped into three different categories: regional sources, local sources, or potentially both regional and local sources. Examples of the regional sources are the sulfate and selenium PMF-factors which most likely-represent coal fired power plants. Examples of local sources are the specialty steel and lead factors. There is reasonable correspondence between these apportionments and data from the EPA TRI and AIRS emission inventories.

Detailed comparisons between PMCAMx predictions and measurements by the STN and IMPROVE measurements in the Eastern US are presented. Comparisons were made for the major aerosol components and PM_{2.5} mass in July 2001, October 2001, January 2002, and April 2002. The results are encouraging with average fraction biases for most species less than 0.25. The improvement of the model performance during the last two years was mainly due to the comparison of the model predictions with the continuous measurements in the Pittsburgh Supersite. Major improvements have included the descriptions: of ammonia emissions (CMU inventory), night time nitrate chemistry, EC emissions and their diurnal variation, and nitric acid dry removal.

TABLE OF CONTENTS

Disclaimer	2
Abstract	3
Table of Contents	4
List of Figures	5
Executive Summary	6
Experimental	7
Activity 1. Project Management	7
Activity 2. Ambient Monitoring	7
Activity 3. Source Characterization	7
Activity 4. Source Apportionment	7
Activity 5. Three-Dimensional Deterministic Modeling	7
Results and Discussion	8
Development of a fingerprint for urban and rural road dust in Pittsburgh, PA	8
Development of a source profile for leaf detritus in Pittsburgh, PA	13
Organic composition of vehicle emissions in the Squirrel Hill Tunnel	16
Spatial source apportionment	20
Evaluation of PMCAMx predictions	26
Conclusions	31
References	32
Presentations and Publications	33
Publications:	33
Presentations:	38

LIST OF FIGURES

Figure 1. Elemental composition of urban and rural road dust samples.....	9
Figure 2. Organic composition of urban and rural road dust by compound class.	10
Figure 4. n-alkanoic acids as a function of carbon number.	11
Figure 5. PAHs in urban and rural road dust.	11
Figure 6. Motor vehicle markers in urban and rural road dust.	12
Figure 7. Road dust steroids and resin acids.....	13
Figure 8. n-Alkanes in leaf sample as a function of carbon number.	15
Figure 9. n-Alkanoic acids in leaf sample as a function of carbon number.....	15
Figure 10. Average fuel-based emissions of sum of hopanes for the three different sampling periods.	18
Figure 11. Comparison of distribution of hopanes, steranes and PAHs in tunnel emissions (bars) with ambient concentrations (horizontal symbols).....	19
Figure 12. Comparison of distribution of n-alkylcyclohexanes and n-alkanes in tunnel emissions (height of bars & right hand axis) with ambient concentrations (horizontal symbols & left hand axis).	20
Figure 13: Comparison of predicted PM _{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for July 2001. Also shown are the 1:1 and ±30% lines.	27
Figure 14: Comparison of predicted PM _{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for October 2001. Also shown are the 1:1 and ±30% lines.	28
Figure 15: Comparison of predicted PM _{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for January 2002. Also shown are the 1:1 and ±30% lines.	29
Figure 16: Comparison of predicted PM _{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for April 2002. Also shown are the 1:1 and ±30% lines.	30

EXECUTIVE SUMMARY

With support from the US Department of Energy and the US Environmental Protection Agency, Carnegie Mellon University is conducting detailed studies of the ambient particulate matter in the Pittsburgh, PA metropolitan area. The work includes ambient monitoring, source characterization, and modeling (statistical and deterministic) for source apportionment. The major objectives of the project include:

- To achieve advanced characterization of the PM in the Pittsburgh region. Measurements include the PM size, surface, volume, and mass distribution; chemical composition as a function of size and on a single particle basis; temporal and spatial variability.
- To obtain accurate current fingerprints of the major primary PM sources in the Pittsburgh region using traditional filter-based sampling and state-of-the-art techniques.
- To estimate the impact of the various sources (transportation, power plants, natural, etc.) on the PM concentrations in the area using both statistical and deterministic models.
- To quantify the responses of the PM characteristics to changes in these emissions in support of the emission control decision making in the area.
- To develop and evaluate current and next generation aerosol monitoring techniques for both regulatory applications and for determination of source-receptor relationships.

This document is the ninth semi-annual progress report for this project. During this project period significant progress was made on the source characterization, source apportionment, and deterministic modeling activities. Major achievements this project period and results described in this progress report include:

- Development of fingerprints for urban and rural road dust in Pittsburgh, PA.
- Development of fingerprint for vegetative detritus based on a composite sample of major tree types in the Pittsburgh region.
- Analysis of vehicle emission measured in tunnel to develop an aggregate, fleet-average emission profiles for motor vehicles in the Pittsburgh area.
- Comparison of PMCAMx+ predictions for PM_{2.5} mass and major species to data from the STN and IMPROVE networks in the Eastern US for July 2001, October 2001, January 2002, and April 2002.

EXPERIMENTAL

This section provides an overview of the effort on various project activities. This project period the majority of the effort went into Activity 3 Source Characterization, Activity 4 Source Apportionment, and Activity 5 Three-Dimensional Modeling.

Activity 1. Project Management

During this project period additional data were submitted to NARSTO for permanent archive and EPA for inclusion in the Supersites Relational Database.

Activity 2. Ambient Monitoring

The purpose of this activity is to create an extensive database of ambient PM measurements for source apportionment, examination of aerosol processes, evaluation of instrumentation, and air quality model development and evaluation. Data collection is complete. Work has continued on the analysis of the ambient data set.

Activity 3. Source Characterization

The purpose of this activity is to develop updated emission profiles for important source categories around Pittsburgh. Updated source profiles are being developed through a combination of source testing, fence line measurements, and analysis of highly time resolved data collected at the central site. These profiles are used in the source apportionment and deterministic modeling activities. Work this project period has focused on analysis of results. A paper on the fence line measurements made adjacent to a metallurgical coke production facility was accepted for publication (Weitkamp et al. 2005). A paper was also submitted on measurements made in the highway tunnel (Grieshop et al. 2005). Finally, analysis was completed on the road dust and vegetative detritus samples. Selected results from this work are presented in the Results and Discussion section.

Activity 4. Source Apportionment

The purpose of this activity is to quantify the contribution of different sources to the fine PM_{2.5} levels in Pittsburgh. Results on the spatial distribution of different source contributions are described in the Results and Discussion section.

Activity 5. Three-Dimensional Deterministic Modeling

The purpose of this activity is to evaluate the performance of the three-dimensional chemical transport model (PMCAMx) with air quality data collected by this and other projects. PMCAMx is a publicly available computer modeling system for the integrated assessment of photochemical and PM pollution. This CTM has been recently upgraded

by the CMU team and ENVIRON to include state-of-the-art description of aerosol dynamics and thermodynamics, cloud chemistry, and wet removal processes. PMCAMx+ is the research version of the code and it includes the latest developments in Carnegie Mellon organic and inorganic aerosol and aqueous-phase chemistry modules. During this project period work focused on evaluating model performance for the October 2001, January 2002, and April 2002 periods.

RESULTS AND DISCUSSION

Development of a fingerprint for urban and rural road dust in Pittsburgh, PA

Vehicle traffic contributes particulate matter from emissions from tailpipes but by re-suspending road dust. To develop a chemical fingerprint for road dust in the Pittsburgh region, paved road dust samples were collected from 5 rural and 6 urban/suburban sites in the summer of 2003 using a vacuum sampler constructed out of Teflon and stainless steel. Samples were dried in an oven at 100 °C and then passed through a 37 µm sieve to remove large particles. The samples were combined to create a composite urban and rural dust sample.

The sieved road dust was then re-suspended in a glass vessel using HEPA and activated carbon cleaned air. The filter samples were analyzed for: PM_{2.5} mass via gravimetric analysis, OC & EC: Thermal-Optical with Sunset Instrument and NIOSH protocol, elemental composition (acid digestion followed by ICP-MS), and organic speciation followed by GC-MS. Multiple samples were collected and analyzed for each dust sample to verify measurement repeatability.

Figure 1 shows the elemental composition of the PM_{2.5} mass of the urban and rural road dust samples. The largest constituents of the road dust include organic carbon (OC), calcium, aluminum and iron. Assuming an organic mass to organic carbon ratio of 1.8, the quantified species contribute 43% and 46% of the rural and urban road dust PM_{2.5} mass, respectively. Major components not analyzed for include silicon and oxygen.

Comparing the urban and rural samples indicates that the urban road dust is enriched in metals associated anthropogenic sources; notably Fe, Zn, Cu, Pb, Cr, Ni, Mo, Sb. The rural road dust is enriched in elements associated with soil dust; notably Ca and Mn. The urban road dust is modestly enriched in carbon.

GC-MS analysis was performed on solvent extracted samples to quantitatively determine contribution of more than 100 individual organic species to the road dust. Figure 2 shows the contribution of different compound classes to the road dust OC. In total, the identified compounds contribute 0.35% and 0.94% of the OC mass in the rural and urban road dust samples, respectively.

For most of the compound classes, the urban road dust is significantly enriched compared to the rural sample. For example, the contribution of n-alkanes to the urban

sample is almost a factor of 4 greater than in the rural sample. The notable exception to this is the enrichment of steroids (cholesterol and beta-sitosterol) in the rural sample.

Figure 1. Elemental composition of urban and rural road dust samples.

Figures 3 and 4 present the distribution of n-alkanes and n-alkanoic acids as a function of carbon number. These two classes contribute the majority of the identified organic mass. The distribution of these species is similar to that measured in a road dust sample collected in Los Angeles (Rogge et al., 1993). For the n-alkanes, the urban samples are enriched across the entire range of carbon numbers. The distribution of lower molecular n-alkanes (C-19-C-25) is characteristic for vehicle exhaust. The highest concentrations are observed in higher molecular weight odd n-alkanes, which is a known signature for vegetative detritus. However, higher even n-alkanes also contribute a significant fraction of organic road dust mass indicating the importance of non-biological sources of high molecular weight n-alkanes.

Figure 4 shows the distribution of n-alkanoic acids with carbon number. For the n-alkanoic acids, stearic and palmitic acids are the dominant homologues in both the urban and rural sample and a strong even-to-odd distribution is prevalent. This pattern in high molecular weight n-alkanoic acids is a characteristic of vegetative detritus. Unlike the n-alkanes, the urban samples are only enriched at certain carbon numbers, notably C-16, C-18 and C-20. Little enrichment in the urban sample is observed in the higher molecular weight n-alkanoic acids.

Figure 2. Organic composition of urban and rural road dust by compound class.

Figure 3. n-Alkanes as a function of carbon number.

Figure 5 compares polycyclic aromatic hydrocarbon concentrations in the urban and rural road dust samples. PAHs are emitted by a number of combustion sources including gasoline powered vehicles, natural gas and wood combustion. Coke production is thought to be the dominant source of PAH in the Pittsburgh metropolitan area.

Figure 4. n-alkanoic acids as a function of carbon number.

Figure 5. PAHs in urban and rural road dust.

The urban road dust sample is significantly enriched in PAH relative to the rural sample. This enrichment underscores the influence of anthropogenic activities on road dust composition in urban environments. For higher molecular weight PAH (C-18 or larger), the ratio of urban to rural PAH is 4.1 ± 0.5 .

Figure 6 compares motor vehicle marker concentrations in urban and rural road dust. Hopanes and steranes are commonly used as markers for diesel and gasoline motor vehicles. Benzothiazole is a marker for tire wear aerosol. Alkylcyclohexanes have been identified in vehicle exhaust.

All of these markers are substantially enriched in the urban road dust sample consistent with the much higher vehicular traffic on urban roads. It is interesting to

compare the enrichment of these molecular markers to elemental carbon (EC). Motor vehicles and diesel vehicles in particular are the dominant source of EC in urban environments. The urban to rural ratio of EC in road dust is 1.25 (Figure 2) versus 2.4 for hopanes and 4.3 for steranes.

Figure 6. Motor vehicle markers in urban and rural road dust.

Figure 7 compares urban and rural road dust steroid and dehydroabietic acid levels. Dehydroabietic acid is a tracer of soft wood combustion and cholesterol is a marker for meat cooking emissions. Beta-sitosterol is a plant derived sterol, also known as a phytosterol. Beta-sitosterol is measured in soft-wood smoke and is also found in plant abrasion products.

The two steroids are the only quantified compounds that were enriched in the rural road dust relative to the urban road dust. We interpret this as the rural road dust being enriched in material of biological origin.

Figure 7. Road dust steroids and resin acids.

Development of a source profile for leaf detritus in Pittsburgh, PA

Due to wind-induced mechanical shear and the rubbing motion of leaves against each other, an unspecified amount of fine particulate epicuticular wax protrusions and leaf deposits are released to the atmosphere that have been identified in urban and rural ambient PM_{2.5} samples. Given the findings from PAQS regarding the potentially significant contribution of cellulose to ambient PM_{2.5} it is important to quantify the contribution of these waxes. In order to determine the contributions of these waxy epicuticular fine particles to the atmospheric fine particle burden, source profile have to be available for the sites of interest.

Up to now, only one source profile has been available for leaf samples collected in Los Angeles (Rogge et al. 1993). As part of PAQS, we have developed an additional source profile for organic constituents associated with fine particulate leaf surface abrasion products has been generated for Pittsburgh, PA

Green leaves from 11 common tree species that are characteristic for the Pittsburgh area were harvested during September of 2003 and composited according to the tree distribution for that area. The distribution of leaves is given in Table 1. This tree distribution is generally representative of forests that are described as a “Mixed Oak Forest” or “Appalachian Oak Forest”. The leaves were collected from the Schenley Park, a park with 500 acres that is situated in Pittsburgh. The park is influenced by human activities, including picnic areas equipped with barbeque places.

To simulate a windblown generation process that dislodges waxy leaf surface protrusions, the leaf composite was placed in a clean Teflon bag and mechanically agitated while passing purified air through the bag, similar to the procedure used for the Los Angeles leaf samples. The fine waxy protrusions ($dp \leq 2.5 \mu m$) shed from the leaf surfaces were extracted and analyzed using gas chromatography/mass spectrometry. Individual organic compounds were identified included: n-Alkanes, iso- and anteiso-

alkanes, n-alkanoic acids, n-alkenoic acids, alkanals, n-alkanones, steroids, pentacyclic triterpenoids, phenolic type compounds, and others. Trace amounts of PAH were identified and quantified as well.

The filter samples were subsequently extracted with methylene chloride. Prior to extraction, known amounts of a suite consisting of 7 perdeuterated n-alkanes were spiked onto the filter samples. The amount added was based on the organic carbon (OC) content of the samples as determined by OC/EC analysis.

The fine particulate leaf surface abrasion products were extracted successively four times by mild ultrasonic agitation with methylene chloride. The total sample extract was combined and reduced to about 100 μ L using rotary evaporation followed by gentle high-purity N₂-stream evaporation. Next, the sample extracts were treated with diazomethane, in order to convert carboxylic acids to their methyl ester analogues. Sample extracts were then analyzed using an Agilent 6890 Gas Chromatography equipped with a 30 m long DB-5MS column, coupled to an Agilent 5973 Mass Selective Detector.

Table 1. Distribution of leaves in composite sample.

Pittsburgh Vegetation Sample Composite		Fraction of Mass in Composite (%)
Ailanthus	<i>Simaroubaceae</i>	7.3
White Ash	<i>Fraxinus Americana</i>	6.3
Ohio Buckeye	<i>Aesculus Glabra</i>	1.0
Wild Red Cherry	<i>Prunus Pensylvanica</i>	2.2
Black Oak	<i>Quercus Velutina</i>	13.4
American Elm	<i>Ulmus Americana</i>	5.0
Scarlet Hawthorne	<i>Crataegus laevigata</i>	4.0
Norway Maple	<i>Acer Platanoides</i>	51.4
Common Locust	<i>Gleditsia triacanthos</i>	5.4
Pin Oak	<i>Quercus Palustris</i>	1.5
Slippery Elm	<i>Ulmus Rubra</i>	2.5

Figure 8 shows that the higher molecular n-alkanes associated with leaf waxes show a pronounced preference for odd carbon numbered n-alkanes in the Pittsburgh samples. Waxy n-alkanes concentration levels for the Pittsburgh leaf abrasion sample is somewhat lower than found for Los Angeles (Rogge et al. 1993), possibly a result of the different plant distribution as well as differences in climate between the two locations.

Figure 8. n-Alkanes in leaf sample as a function of carbon number.

Figure 9. n-Alkanoic acids in leaf sample as a function of carbon number.

Figure 9 indicates that the n-alkanoic acids in leaf surface waxy protrusions show a strong preference for even carbon numbered n-alkanoic acids for carbon number greater than C14. Comparing the source profiles for leaf surface abrasion products from green leaves, it can be seen that the concentrations for higher molecular weight ($\geq C24$) even carbon numbered n-alkanoic acids is about 2-3 times higher at Pittsburgh than measured in Los Angeles. A more than 4-fold increase in the mass concentration of higher molecular weight n-alkanoic acids ($\geq C24$) is observed in dead leaf abrasion products. Possible explanations include oxidation of n-alkanols and n-alkanals to form n-alkanoic

acids in the dead leaf material and/or the effect of environmental growth conditions (e.g., temperature) or leaf age on wax composition.

In the Pittsburgh fine particulate leaf surface sample, no n-alkanols were identified. Consequently, no comparison for these compound class is possible between this two locations. Also, only the major n-alkanals could be identified in the Pittsburgh sample and these show somewhat lower concentrations than determined for Los Angeles.

Because ambient particulate matter is deposited onto surfaces, including leaf surfaces, it is not surprising to find as well PAHs. Fluoranthene and pyrene show about 10 times higher concentrations in the leaf surface abrasion products in the Pittsburgh sample when compared to Los Angeles samples. This is not surprising, considering the substantially higher ambient PAHs concentrations in Pittsburgh than typically found in Los Angeles.

Organic composition of vehicle emissions in the Squirrel Hill Tunnel

Interstate-376 in Pittsburgh passes through the Squirrel Hill tunnel. The 4,225 ft long twin-bore tunnel carries both and non-commercial traffic. Sampling was performed in the west bound bore which has a constant 2.5 percent grade so the vehicles are under load.

As part of the sampling twenty PM2.5 Filter/PUF pairs were collected for organic speciation from the tunnel and nearby ambient sites. These samples were collected in 2002 during November 12-15 (Tuesday-Friday), and November 18-21 (Monday-Thursday). The sampler was set up above the ceiling of the tunnel at approximately 12 feet above the road and approximately 100 feet from the exit-end of the tunnel. Sampling was performed using a Tisch Environmental, Inc. Model TE-1000 PUF sampler operated at a flow rate of 145 lpm. A PM2.5 cut was obtained with a URG cyclone installed upstream of the Tisch sampler.

Traffic count and speed data were gathered via a RTMS (Remote Traffic Microwave Sensor) operated for the Pennsylvania Department of Transportation (PennDOT). Fleet composition was determined from PennDOT video recorded at the entrance and exit of the tunnel's west-bound tube. Vehicles were manually classified as light-duty (LD) and heavy-duty (HD) based on visual inspection of video. Heavy-duty vehicles include tractor-trucks, large single-unit trucks and buses. City buses did not regularly pass through the tunnel. While this approach separates vehicles by size and not fuel-type, the assumption that most large vehicles are diesel-powered and smaller ones gasoline-powered is reasonable. Less than 5% of heavy-duty vehicles with more than three axles in the U.S. fleet are gasoline-powered (VIUS 2002). The portion of light duty vehicles powered by diesel fuel is similarly small; around 1% of vehicles in US households were diesel powered (EIA 1994) and at most 15% of light-duty trucks are diesel powered (VIUS 2002). Therefore, we assume that all of the heavy duty vehicles are diesels (HDDV) and light duty vehicles (LDV) are gasoline powered. Traffic composition is summarized in Table 2.

Table 2. Summary of traffic data for organic speciation sampling periods.

Date	Time Period	Traffic Density (vehicles/hour)	Average speed (mph)	Average Truck	Period Identification
				Fraction (% fuel)	
Nov. 15	00:00 – 06:00	867	58	29%	High-Truck
Nov. 18	00:00 – 06:00	944	56	28%	High-Truck
Nov. 19	00:00 – 06:00	923	58	33%	High-Truck
Nov. 12	0:700-09:00	3766	30	13%	Low-Speed
Nov. 13	0:700-09:00	3877	29	13%	Low-Speed
Nov. 20	0:700-09:00	4058	29	9%	Low-Speed
Nov. 15	10:00-16:30	3312	48	17%	High-Speed
Nov. 21	10:00-16:30	3152	41	21%	High-Speed

These samples were analyzed by GC-MS for n-alkanes, iso- and anteiso-alkanes, n-alkanoic acids, n-alkenoic acids, aliphatic dicarboxylic acids, aromatic polycarboxylic acids, alkylcyclohexanes, resin acids, levoglucosan, phenolic type compounds, sterols, thiazoles, hopanes, steranes, PAHs, oxy-PAHs, N-hetero-PAHs, S-hetero-PAHs, and others. This involved spiking each filter with a known amount of an internal standard mix consisting of a suite of seven perdeuterated n-alkanes (C12, C16, C20, C24, C28, C32 and C36) to correct for losses incurred throughout the analytical procedure. Extraction was then performed with dichloromethane using mild ultrasonication for the filters, and a Repeated Compression Extraction Apparatus (RCEA) for the PUFs. Both extracts were combined and rotavaporated to reduce the solvent volume to about 2 ml. The volume was further reduced to ~200µl using a gentle stream of ultrapure N₂. The extracts were methylated using freshly prepared diazomethane in order to convert fatty acids to their methyl esters analogs prior to GC/MS analysis.

GC/MS analysis of derivatized extracts was carried out on a GC model HP-6890 coupled to a MSD model HP-5973 (Agilent Technologies). The GC was operated in the splitless mode and was equipped with a capillary column DB-5MS. Temperature programming of the GC consisted of the following steps: isothermal at 60°C for 2 min, temperature ramp of 6 °C/min up to 300°C, and isothermal hold for 18 min. The injector temperature was set at 300 °C and the GC/MS interface temperature at 285 °C. Helium was used as carrier gas with a flow rate of 1.2 ml/min. The injected sample volume was 1µl. The MSD was operated in the electron impact mode at 70 eV electron energy, scan range 50-550 amu. The GC/MS data was acquired and processed using a Agilent ChemStation.

The concentrations of the 96 organic compounds identified in the PM_{2.5} of the tunnel samples were corrected by subtracting their ambient concentration levels corresponding to the days when the tunnel samples were collected. These corrected concentrations were then converted to fuel-based emission factors by multiplying the compounds concentrations by the conversion factor reported in the table above. The multipliers to convert concentrations to their equivalent emission factors were calculated for each sample by Grieshop et. al. (2005), based on the background-corrected CO₂ and CO

concentrations measured in the tunnel, and the carbon fraction in the diesel fuel and gasoline, weighted by the fleet composition.

Figure 10 shows the emission factors for the sum of seven individual hopanes, calculated as an average of the samples collected at each of the time periods considered. The emission factors for the “High-Truck”, “Low-Speed” and “High-Speed” periods are very similar. The emission profiles for the steranes for the same three periods show the same trend, and the ratio of total hopanes to total steranes was found to be 1.67 ± 0.07 $\mu\text{g/Kg}$. Likewise, our findings show that the emission factors profiles for n-alkanes (C19-C32), and n-alkylcyclohexanes (C17-C28), are as well very similar for all three time-periods. This indicates that the composition of these organic compounds in the fleet emissions from gasoline and diesel powered vehicles is preserved in different driving conditions.

Figure 10. Average fuel-based emissions of sum of hopanes for the three different sampling periods.

Figures 11 - 12 illustrates that the vehicular signature can be clearly recognized in the ambient air samples. Starting with the hopanes and steranes, the ambient concentrations profile and the tunnel emission factors profile show great similarity. The same similarity is observed for the PAHs, particularly for the higher molecular weight PAHs ranging from benzo[a]anthracene to coronene. For the ambient n-alkanes concentration profiles, Figure 12 clearly shows the contribution of the leaf surface abrasion particulates in the ambient samples by the characteristic predominance of odd-to-even carbon number n-alkanes.

Figure 11. Comparison of distribution of hopanes, steranes and PAHs in tunnel emissions (bars) with ambient concentrations (horizontal symbols).

Figure 12. Comparison of distribution of n-alkylcyclohexanes and n-alkanes in tunnel emissions (height of bars & right hand axis) with ambient concentrations (horizontal symbols & left hand axis).

Spatial source apportionment

Ambient PM_{2.5} composition data in Pittsburgh, PA have been used with Positive Matrix Factorization (PMF) to determine the major sources of PM_{2.5} sampled (Pekney et al. 2005). We have now used the potential source contribution function (PSCF) with the PMF-modeled source contributions to estimate the spatial locations of the sources. We have also used the Conditional Probability Function (CPF) to calculate the probability that a source is located within a particular wind direction sector.

The Conditional Probability Function (CPF) calculates the probability that a source is located within a particular wind direction sector, $\Delta\Theta$, or:

$$CPF = \frac{m_{\Delta\Theta}}{n_{\Delta\Theta}} \quad (1)$$

where $n_{\Delta\Theta}$ is the number of times that the wind passed through direction sector $\Delta\Theta$, and $m_{\Delta\Theta}$ is the number of times that the source contribution peaked while the wind passed through sector $\Delta\Theta$ (Ashbaugh 1985). To use CPF with the Pittsburgh Supersite data, the 24-hour averaged source contribution data have been applied to all 15-minute wind direction averages measured at the site for each date. All time periods of wind speed less than 1 m/s have been removed from the dataset. To calculate $m_{\Delta\Theta}$, the highest 25% of

source contribution concentrations were counted. CPF is useful in determining the direction of a source from a receptor site; however, it cannot determine the actual location of the source.

PSCF calculates the probability that a source is located at latitude i and longitude j . The basis of PSCF is that if a source is located at (i,j) , an air parcel back trajectory passing through that location indicates that material from the source can be collected and transported along the trajectory to the receptor site. PSCF solves

$$PSCF = \frac{m_{ij}}{n_{ij}} \quad (2)$$

where n_{ij} is the number of times that the trajectories passed through the cell (i,j) and m_{ij} is the number of times that a source concentration was high when the trajectories passed through the cell (i,j) . The criteria for determining m_{ij} were either the 75th or 90th percentile highest source contributions, depending on the structure of the source contributions for each factor. Therefore, a cell (i,j) with a PSCF value close to one indicates a probable source location. A probability field is constructed for the domain covered by the back trajectories such that locations from which transport to the receptor site results in high concentrations of a source material are most likely locations of sources of that material.

The Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model was used to calculate air mass back-trajectories in 1-hour sequential increments ending at the receptor site at 40.44°N latitude and 79.94°W longitude (Draxler 1999). HYSPLIT was run four times a day for every day of the study, or every six hours from July 11, 2001 – July 31, 2002. A starting height of 500m was used. Six-hour back trajectories were selected because this time is sufficient to determine probable locations of both local and regional sources, and increasing the time of trajectories also increases the uncertainty, resulting in unrealistic source locations.

The trajectory domain extends from 35°–50° north latitude and 75°–90° west longitude, and is divided into $0.1^\circ \times 0.1^\circ$ cells, resulting in 22,500 total cells. The maximum n_{ij} is 1570, although many cells on the outer edges of the grid have n_{ij} less than 10. These outer-edge cells could have misleading large PSCF values if a trajectory passed through them during a source contribution peak, resulting in mistaken attribution of a source to an area. To reduce the effect of spurious large ratios in grid cells with low n_{ij} , a weighting function W_{ij} is multiplied by the PSCF value:

$$W_{ij} = \begin{cases} 1.0 & 9 < n_{ij} \\ 0.9 & 8 < n_{ij} \leq 9 \\ 0.8 & 7 < n_{ij} \leq 8 \\ 0.7 & 6 < n_{ij} \leq 7 \\ 0.6 & 5 < n_{ij} \leq 6 \\ 0.5 & 4 < n_{ij} \leq 5 \\ 0.4 & 3 < n_{ij} \leq 4 \\ 0.3 & 2 < n_{ij} \leq 3 \\ 0.2 & 1 < n_{ij} \leq 2 \\ 0.1 & n_{ij} \leq 1 \end{cases} \quad (3)$$

The results from the PSCF and CPF analysis are summarized in Table 3. PMF-modeled factor names are given with their tracer species, annual average contribution to PM_{2.5} mass concentrations in Pittsburgh, source regions as determined by the PSCF and CPF analyses, and possible sources as identified by the tracer species and the TRI. The PSCF and CPF results for the PMF factors presented can be grouped into three different categories: regional sources, local sources, or potentially both regional and local sources. Examples of the regional sources are the sulfate and selenium PMF-factors which most likely-represent coal fired power plants. For these factors, the PSCF results show most probable source locations in agreement with TRI or AIRSDatabase source locations while the CPF results show a slightly offset most probable direction that does not agree well with the PSCF results.

Examples of local sources are the specialty steel and lead factors. For these factors the CPF results show clear most probable directions that agree well with both the PSCF results, which show a clustering of most probable locations near the site and the known locations of molybdenum, chromium, and lead sources by TRI. Although the cadmium factor is assumed to also represent a local source or sources, the CPF results were inconclusive, with no direction showing a strongly significant probability. The gallium-rich and Fe, Mn, and Zn factors could represent either local or regional sources, or both.

The gallium-rich factor PSCF and CPF results show very good agreement but as this source cannot be identified, a conclusion as to its being regional or local cannot be drawn. And while the Fe, Mn, and Zn factor PSCF and CPF results are similar to results for the sulfate and selenium factors, the difference that there are known local Mn and Zn sources is the direction indicated as most probable by CPF. Therefore, this factor can be considered both regional and local. In conclusion, PSCF appears to better predict locations of regional sources while CPF is useful in pinpointing directions of local sources.

PMF, CPF and PSCF are powerful tools for determining the major sources of particulate pollution in an urban area like Pittsburgh, but each model has its limitations. Both models rely on counting statistics and therefore require large amounts of data such

that results are statistically significant. PMF cannot distinguish between sources in the same area that impact the receptor location. This may lead to a mixing of different types of sources that makes it difficult to determine the apportionment of source contributions to PM_{2.5} mass concentrations. CPF can provide the probability that a source is located in a particular direction, but cannot provide any information as to the proximity of the source. PSCF, however, provides probability that a source is located in a particular grid cell, but is dependent upon the accuracy of the HYSPLIT model. HYSPLIT trajectory accuracy decreases with increasing trajectory time, and effects of the boundary layer are not considered.

In an urban area like the Pittsburgh, where there are many industrial activities in and near the city as well as in the surrounding heavily industrialized region, clearly identifying source locations using any of these models is difficult. High concentrations of some pollutant species can occur for a wide range of wind directions. Furthermore, mixing height as well as wind direction can affect airborne concentrations of the tracer species, although only the latter is used in these models. When 24-hour data are used, source plumes that occur on the order of minutes are masked due to the variability of wind direction over a day. Despite these limitations, probable locations are determined for several of the modeled sources that have been shown to correlate with the locations of known sources as reported to the TRI. This study demonstrated that CPF and PSCF aided in determining PM source locations to significantly improve source apportionment study.

Table 3. Summary of sources identification from PMF, CPF, and PSCF analysis. Annual average contribution each source makes to fine particle mass is given as percent of annual average PM_{2.5} mass.

PMF-Factor (% of PM_{2.5})	Tracer Species	Possible source direction from site, CPF	Possible source areas and direction from site, PSCF	Sources and industry types as identified by the TRI except where indicated
Sulfate (33)	Sulfate	SE	Ohio River Valley: SW	Electric services, petroleum and coal products (from AIRS Database, EPA 1999)
Crustal Material (12)	Ca, Ti	NE, SW	No most probable location	Wind-blown soil crust and road dust (Watson et al. 2001)
Vehicle Emissions and Road Dust (11)	OC, EC, hopanes, PAH's, n-alkenes	NNE, SE	No most probable location	Highways and roads (Schauer et al. 1996)
Nitrate (8)	Nitrate	NNE, SW	No most probable location	Vehicles, electric services (Watson et al. 2001)
Wood Combustion, Vegetative Detritus, Cooking (4)	OC, EC, resin acids, levoglucosan, syringols, n-alkenes	No probability of direction > 0.35	No most probable location	Wildfires, residential wood burning and cooking, restaurants (Simoneit 1989, Simoneit et al. 1993, 1999)
Gallium-rich source (3)	Ga (and some As, Cu, Ni, and V)	NW	Western PA and NE Ohio: NW	Ni and Cu: Electric services, blast furnaces and steel mills, electrometallurgical products, steel and copper foundries, metalworking and manufacturing As and V: Electric services
Steel Production (2)	Fe, Zn, Mn	SE	Ohio River Valley: SW	Zn and Mn: Blast furnaces and steel mills, foundries, electrometallurgical products, electric services, and primary and secondary non-ferrous metals

Table 3 (cont.)

PMF-Factor (% of PM_{2.5})	Tracer Species	Possible source direction from site, CPF	Possible source areas and direction from site, PSCF	Sources and industry types as identified by the TRI except where indicated
Lead (2)	Pb	N	Western PA: N and S	Electric services, blast furnaces and steel mills, copper foundries, metal sanitary ware, refuse systems, hydraulic cement, inorganic pigments and paints, pressed and blown glass and glassware, custom compound purchased resins, electrometallurgical products, storage batteries, primary and secondary non-ferrous metals, and steel pipe and tubes
Specialty Steel Production and Processing (1)	Cr, Mo	NNE, ESE	Local: N, E, and S	Mo: Blast furnaces and steel mills, primary metal products, secondary nonferrous metals, electrometallurgical products, industrial inorganic chemicals, steel foundries, primary and secondary non-ferrous metals and metal products. Cr: source types listed for Mo plus electric services, cold finishing of steel shapes, steel pipe and tubes, fabricated pipe and fittings, rolling mill machinery, special dies, tools, jigs, and fixtures, fabricated metal products, metal coating and allied services, fabricated plate work (boiler shops), and nonferrous rolling and drawing.
Cadmium (1)	Cd	No probability of direction > 0.25	Tri-state region: NW, SW, S	Primary and secondary non-ferrous metals, pressed and blown glass and glassware, custom compound purchased resins, refuse systems, and inorganic pigments, paints, and allied products
Selenium (1)	Se	SE	Ohio River Valley: SW, and to a lesser extent NE	Primarily electric services

Evaluation of PMCAMx predictions

PMCAMx+ is applied in the eastern United States during the four seasons of the year (July 2001, October 2001, January 2002, April 2002). The first days of each simulation have been excluded in order to limit the effect the initial conditions have on the results. For boundary conditions, low background concentrations are used for all species. The aerosol module includes ten size sections varying in size from 40 nm to 10 μ m, with the equilibrium approach being used to model aerosol condensation and evaporation. The modeling domain covers a 3492x3240 km region in the eastern United States with 36x36 km grid resolution and fourteen vertical layers totaling 6 km. Inputs to the model include horizontal wind components, temperature, pressure, water vapor, vertical diffusivity, clouds, and rainfall, all created using the meteorological model MM5 (Grell et al., 1995) by the Lake Michigan Air Directors Consortium (LADCO).

The emission inventory used is Midwest Regional Planning Organization's Base E inventory, which was put together and supplied by LADCO [LADCO, 2003]. This inventory (including primary carbonaceous material) is based on the U.S. EPA's 1999 National Emissions Inventory (version 2.0) [U.S. EPA, 2001]. Spatial and temporal improvements have been made to develop the Base E emissions inventory: the inventory has improvements to the temporal ammonia emissions made at Carnegie Mellon University, uses MOBILE6 for vehicular sources [U.S. EPA, 2003], BIOME3 for biogenic emissions [Wilkinson and Janssen, 2001], and has reduced dust emissions. Three different representative days are available: an average weekday, Saturday, and Sunday. There are two types of emissions, elevated point sources and gridded sources, which include mobile sources, area/non-road mobile sources, and biogenic sources.

Scatter plots comparing model predictions to daily measurements made at AIRS and IMPROVE network sites for July 2001, October 2001, January 2002 and April 2002 are shown in Figures 13-16. Data are shown for overall PM_{2.5} mass and major aerosol species including sulfate, nitrate, ammonium, organic matter, and elemental carbon. The average fractional bias of the model predictions for each season for these different species is shown in Figure 17.

The current performance of PMCAMx in the Eastern U.S. for the major aerosol components and PM_{2.5} during all seasons is encouraging. The improvement of the model performance during the last two years was mainly due to the comparison of the model predictions with the continuous measurements in the Pittsburgh Supersite. Major improvements have included the descriptions: of ammonia emissions (CMU inventory), night time nitrate chemistry, EC emissions and their diurnal variation, and nitric acid dry removal. The most significant remain issues include: improvement of meteorological fields and wet deposition, ammonia emissions in October, primary organic PM emissions, and description of SOA formation.

Figure 13: Comparison of predicted PM_{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for July 2001. Also shown are the 1:1 and $\pm 30\%$ lines.

Figure 14: Comparison of predicted PM_{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for October 2001. Also shown are the 1:1 and $\pm 30\%$ lines.

Figure 15: Comparison of predicted PM_{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for January 2002. Also shown are the 1:1 and $\pm 30\%$ lines.

Figure 16: Comparison of predicted PM_{2.5} mass, sulfate, nitrate, ammonium, organic mass, and elemental carbon to data from the AIRS and IMPROVE Networks for April 2002. Also shown are the 1:1 and $\pm 30\%$ lines.

Figure 17. Average fractional bias of PMCAMx+ predictions for different seasons. A positive fractional bias indicates the model is overpredicting the observed concentrations. For example, as shown in Figure 14, the model significantly overpredicts ammonium concentrations in October 2001.

CONCLUSIONS

Significant progress was made this project period on the source characterization, source apportionment and model evaluation activities. On the source characterization front, this report highlights new data on road dust, vegetative detritus and motor vehicles. The results show significant differences in the composition in urban and rural road dust. Also there are clear differences in the organic composition of vegetative detritus in Pittsburgh compared to previous results obtained in Los Angeles. This is not unexpected given the differences in climate and plant species. Finally, a comparison of the organic of the fine particulate matter in the tunnel with the ambient provides clear evidence of the significant contribution of vehicle emissions to ambient PM. The source profiles developed from this work are being used by the source-receptor modeling activities.

We presented results from source apportionment analysis of the spatial distribution of PMF-factors. The PSCF and CPF results for the PMF-factors presented can be grouped into three different categories: regional sources, local sources, or potentially both regional and local sources. Examples of the regional sources are the sulfate and selenium PMF-factors which most likely-represent coal fired power plants. Examples of local sources are the specialty steel and lead factors. The nitrate, vehicle emissions and road dust, wood combustion, vegetative detritus and cooking, and crustal material factor CPF and PSCF results were inconclusive as sources of these factors exist in all directions from the site and therefore one would not expect a clear probability field in any one direction.

The current performance of PMCAMx in the Eastern U.S. for the major aerosol components and PM_{2.5} during all seasons is encouraging. The improvement of the model performance during the last two years was mainly due to the comparison of the model predictions with the continuous measurements in the Pittsburgh Supersite. Major

improvements have included the descriptions: of ammonia emissions (CMU inventory), night time nitrate chemistry, EC emissions and their diurnal variation, and nitric acid dry removal. The most significant remain issues include: improvement of meteorological fields and wet deposition, ammonia emissions in October, primary organic PM emissions, and description of SOA formation. We are now using the model to investigate response of ambient PM_{2.5} concentrations to changes in emissions.

REFERENCES

EIA (1994). Residential Transportation Energy Consumption Survey, U.S. Energy Information Administration.

Grieshop, A. P., Lipsky, E. M., Pekney, N. J., Takahama, S. and Robinson, A. L. (2005). "Diesel and Gasoline Vehicle Gaseous and Fine Particle Emission Factors Measured in a Highway Tunnel in Pittsburgh, PA." *Atmospheric Environment* Submitted.

Pekney, N. J., Davidson, C. I., Robinson, A. L., Zhou, L., Hopke, P. K. and Eatough, D. (2005). "Identification of Major Sources of PM_{2.5} in Pittsburgh using PMF and UNMIX." *Aerosol Science & Technology* in press.

Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R. and Simoneit, B. R. T. (1993). "Sources of Fine Organic Aerosol .4. Particulate Abrasion Products From Leaf Surfaces of Urban Plants." *Environmental Science & Technology* 27(13): 2700-2711.

VIUS (2002). Vehicle Inventory and Use Survey, U.S. Census Bureau.

Weitkamp, E. A., Lipsky, E. M., Pancras, P., Ondov, J., Polidori, A., Turpin, B. J. and Robinson, A. L. (2005). "Fine Particle Emission Profile for a Large Coke Production Facility based on Highly Time Resolved Fence Line Measurements." *Atmospheric Environment* 39: 6719-6733.

PRESENTATIONS AND PUBLICATIONS

Publications:

1. A. Khlystov, C. O. Stanier, S. Takahama, and S. N. Pandis (2005) Water content of ambient aerosol during the Pittsburgh Air Quality Study, *J. Geophys. Res.*, 110 (D7): Art. No. D07S10.
2. A. Khlystov, C. Stanier, and S.N. Pandis, "An algorithm for combining electrical mobility and aerodynamic size distributions data when measuring ambient aerosol," *Aerosol Science and Technology*, 38(S1), 229-238 (2004).
3. A. Khlystov, Q. Zhang, J. L. Jimenez, C. Stanier, S. N. Pandis, M. R. Canagaratna, P. Fine, C. Misra, and C. Sioutas (2005) In-situ concentration of semi-volatile aerosol using water-condensation technology, *J. Geophys. Res.*, submitted.
4. A. Polidori, Turpin, B., Lim H.J., Totten, L., Davidson, C., "Polarity and Molecular Weight/Carbon Weight of the Pittsburgh Organic Aerosol," *Aerosol Sci. Technol.* in preparation.
5. A. Polidori, Turpin, B., Lim, H.J., Cabada, J.C., Subramanian, R., Robinson, A., "Secondary Organic Aerosol Formation during the Pittsburgh Air Quality Study (PAQS)," *Aerosol Sci. Technol.* in preparation.
6. A.E. Wittig N. Anderson, A.Y. Khlystov, S.N. Pandis, C. Davidson, and A.L. Robinson, "Pittsburgh Air Quality Study overview," *Atmospheric Environment*, 38 (20), 3107-3125 (2004).
7. A.E. Wittig, S. Takahama, A.Y. Khlystov, S.N. Pandis, S. Hering, B. Kirby, and C. Davidson, "Semi-continuous PM_{2.5} inorganic composition measurements during the Pittsburgh Air Quality Study," *Atmospheric Environment*, 38 (20), 3201-3213 (2004).
8. A.L. Robinson, Donahue, N. M. and Rogge, W. F. (2005). "Photochemical oxidation and changes in molecular composition of organic aerosol in the regional context." *Journal of Geophysical Research-Atmospheres* In preparation.
9. A.P. Grieshop, Lipsky, E. M. and Robinson, A. L. (2005). "Measurements of In Use Vehicle Fine Particle Emissions Made in a Highway Tunnel." *Aerosol Science & Technology* in preparation.
10. Bein, K.J., Y. Zhao, A.S. Wexler and M.V. Johnston, Speciation of size-resolved individual ultrafine particles in Pittsburgh, Pennsylvania, *J. Geophys. Res.*, doi:10.1029/2004JD004708, in review.

11. C.O. Stanier, A. Khlystov, and S.N. Pandis, "Chemical processes and long-range transport of aerosols: Insights from the Pittsburgh Air Quality Study," in *Long Range Transport of Air Pollution*, Kluwer (2002).
12. C.O. Stanier, A. Khlystov, W.R. Chan, M. Mandiro, and S.N. Pandis, "A method for the in situ measurement of fine aerosol water content of ambient aerosols: The Dry-Ambient Aerosol Size Spectrometer (DAASS)," *Aerosol Science and Technology*, 38(S1), 215-228 (2004).
13. C.O. Stanier, A.Y. Khlystov, and S.N. Pandis, "Ambient aerosol size distributions and number concentrations measured during the Pittsburgh Air Quality Study (PAQS)," *Atmospheric Environment*, 38 (20), 3275-3284 (2004).
14. C.O. Stanier, A.Y. Khlystov, and S.N. Pandis, "Nucleation events during the Pittsburgh Air Quality Study: Description and relation to key meteorological, gas phase, and aerosol parameters," *Aerosol Science and Technology*, 38(S1), 253-264 (2004).
15. D. Vayenas, S. Takahama, C. Davidson, and S. N. Pandis (2005) Simulation of the thermodynamics and removal processes in the sulfate-ammonia-nitric acid system: Implications for PM_{2.5} control strategies, *J. Geophys. Res.*, 110 (D7): Art. No. D07S14.
16. D.B. Kane, J.J. Wang, K. Frost, M.V. Johnston, (2002) "Detection of Negative Ions from Individual Ultrafine Particles", *Analytical Chemistry* 74, 2092-2096.
17. D.B. Millet, N. M. Donahue, S. N. Pandis, A. Polidori, C. O. Stanier, B. J. Turpin, and A. H. Goldstein (2005) Partitioning VOCs and organic aerosols into primary and secondary sources: Results from the Pittsburgh Air Quality Study, *J. Geophys. Res.*, 110 (D7): Art. No. D07S07.
18. E. A. Weitkamp, E.M. Lipsky, P. Pancreas, J. Ondov, A. Polidori, B.J. Turpin, A.L. Robinson, "Fine Particle Emission Profile for a Large Coke Production Facility Based on Highly Time Resolved Fence Line Measurements," *Atmospheric Environment*, submitted.
19. E.M. Lipsky and A.L. Robinson, "Effects of Dilution on Fine Particle Mass and Partitioning of Semi-volatile Organics in Diesel Exhaust and Wood Smoke," *Environmental Science & Technology*, submitted.
20. E.M. Lipsky, A.L. Robinson, "Design and evaluation of a portable dilution sampling system for measuring fine particle emissions from combustion systems," *Aerosol Science & Technology*, in press.
21. G.A. Lithgow, A.L. Robinson, and S.G. Buckley, "Ambient measurements of metal-containing PM_{2.5} in an urban environment using laser-induced breakdown spectroscopy," *Atmospheric Environment*, 38 (20), 3319-3328 (2004).

22. J. Hybl, G. Lithgow, and S.G. Buckley, (2003) "Laser-Induced Breakdown Spectroscopy Detection of Biological Material," *Applied Spectroscopy* 57(10) pp 1207-1215
23. J.C. Cabada S. Rees, S. Takahama, A. Khlystov, S.N. Pandis, C.I. Davidson, and A.L. Robinson, "Mass size distributions and size resolved chemical composition of fine particulate matter at the Pittsburgh Supersite," *Atmospheric Environment*, 38 (20), 3127-3141 (2004).
24. J.C. Cabada, A. Khlystov, A.E. Wittig, C. Pilinis, and S.N. Pandis, "Light scattering by fine particles during the Pittsburgh Air Quality Study: Measurements and modeling," *Journal of Geophysical Research--Atmospheres*, 109 (D16), Art. No. D16S03 (2004).
25. J.C. Cabada, S.N. Pandis, and A.L. Robinson, "Sources of atmospheric carbonaceous particulate matter in Pittsburgh, Pennsylvania," *Journal of the Air & Waste Management Association*, 52, 732-741 (2002).
26. J.C. Cabada, S.N. Pandis, R. Subramanian, A.L. Robinson, A. Polidori, and B. Turpin, "Estimating the secondary organic aerosol contribution to PM_{2.5} using the EC tracer method," *Aerosol Science and Technology*, 38(S1), 140-155 (2004).
27. J.P. Pancras, Ondov, J. M., Gazula, S., Moore, J., Tuch, T., (2005). Highly-Time Resolved Metals Measurements with SEAS-II at Four U.S. Supersites. *Aerosol Sci. Technology*, in preparation.
28. J.P. Pancras, Ondov, J. M., Zeisler, R., (2005) Multielement Electrothermal AAS Determination of Eleven Marker Elements in Fine Ambient Aerosol Slurry Samples Collected with SEAS-II, *Analytica Chimica Acta*, in press.
29. K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston (2005), Speciation of size-resolved individual ultrafine particles in Pittsburgh, Pennsylvania, *J. Geophys. Res.*, 110, D07S05, doi:10.1029/2004JD004708.
30. K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky and A.L. Robinson, A hypothesis for the missing source of ultrafine calcium particles in the atmosphere, *Atm. Environ.*, in preparation.
31. K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, Source sampling experiments using a single particle mass spectrometer during the Pittsburgh Air Quality Study: source characterization and quantitative estimates of source contribution, *J. Geophys. Res.*, in preparation.
32. K.J. Bein, Y. Zhao, A.S. Wexler, N.J. Pekney, C.I. Davidson, M.P. Tolocka, M.V. Johnston and G. Evans, Characterization of smoke plumes from Canadian forest fires detected at three separate locations, Pittsburgh, Baltimore and Toronto, using single particle mass spectrometry, *Atm. Environ.*, in preparation.

33. L. Zhou, E. Kim, P. K. Hopke, C. Stanier, and S.N. Pandis (2005) "Mining airborne particulate size distribution data by positive matrix factorization," *J. Geophys. Res.*, 110 (D7): Art. No. D07S19.
34. L. Zhou, E. Kim, P.K. Hopke, C.O. Stanier, and S. Pandis, "The advanced factor analysis on Pittsburgh particle size-distribution data," *Aerosol Science and Technology*, 38(S1), 118-132 (2004).
35. L. Zhou, P. K. Hopke, C. O. Stanier, S. N. Pandis, J. M. Ondov, and J. P. Pancras (2005) Investigation of the relationship between chemical composition and size distribution of airborne particles by Partial Least Square (PLS) and Positive Matrix Factorization (PMF), *J. Geophys. Res.*, 110 (D7): Art. No. D07S18.
36. L. Zhou, W. Liu, and P.K. Hopke, (2004) "Comparison of Two Trajectory Based Models for Locating Particle Sources for Two Rural New York Site", *Atmospheric Environment* 38 (13): 1955-1963.
37. L.M. Zhou, P.K. Hopke, P. Paatero, J.M. Ondov, J.P. Pancras, N.J. Pekney, and C.I. Davidson, "Advanced factor analysis for multiple time resolution aerosol composition data," *Atmospheric Environment*, 38 (29), 4909-4920 (2004).
38. N.J. Pekney, C.I. Davidson, A.L. Robinson, L. Zhou, Hopke, P.K. D.J. Eatough "Identification of Major Sources of PM_{2.5} in Pittsburgh using PMF and Unmix," *Aerosol Science & Technology*, submitted.
39. N.J. Pekney, K.J. Bein, C.I. Davidson, A.S. Wexler and M.V. Johnston, Identification of sources of atmospheric PM at the Pittsburgh Supersite: RSMS III and filter-based Positive Matrix Factorization, *Atm. Environ.*, in preparation.
40. Q. Zhang, C. O. Stanier, M. R. Canagaratna, J. T Jayne, D. R. Worsnop, S. N. Pandis, and J. L. Jimenez (2004) Insights into the chemistry of new particle formation and growth events in Pittsburgh based on aerosol mass spectrometry, *Environ. Sci. Tech.*, 38 (18): 4797-4809.
41. Q. Zhang, M. R. Canagaratna, J. T. Jayne, D. R. Worsnop, and J. L. Jimenez (2005) Time and size-resolved chemical composition of submicron particles in Pittsburgh: Implications for aerosol sources and processes, *J. Geophys. Res.*, 110 (D7): Art. No. D07S09.
42. R. Subramanian, A.Y. Khlystov, J.C. Cabada, and A.L. Robinson, "Positive and negative artifacts in particulate organic carbon measurements with denuded and undenuded sampler configurations," *Aerosol Science and Technology*, 38(S1), 27-48 (2004).
43. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Selection of Source Profiles for Chemical Mass Balance Modeling

using Organic Molecular Markers." Environmental Science & Technology Under review.

44. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Source Apportionment of Gasoline and Diesel Vehicles using the Chemical Mass Balance Model and Molecular Markers." Environmental Science & Technology In preparation.
45. R. Subramanian, Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F. and Robinson, A. L. (2005). "Source Apportionment of Primary Organic Aerosol in Pittsburgh, PA using organic molecular markers." Environmental Science & Technology In preparation.
46. R. Subramanian, Khlystov, A. Y. and Robinson, A. L. (2005). "Measurement of elemental carbon using the Thermal-Optical Transmittance Technique: Fundamental assumptions and effect of peak inert-mode temperature." *Aerosol Science and Technology*, In preparation.
47. R.R. Anderson, W. K. Modey, D. V. Martello, L. J. Lucas, C. I. Davidson, and D. J. Eatough (2005) Apportionment of ambient primary and secondary pollutants during a 2001 summer study in Pittsburgh using EPA UNMIX, *J. Geophys. Res.*, submitted.
48. S. Takahama, A.E. Wittig, D.V. Vayenas, C.I. Davidson, and S.N. Pandis, "Modeling the diurnal variation of nitrate during the Pittsburgh Air Quality Study," *Journal of Geophysical Research--Atmospheres*, 109 (D16), Art. No. D16S06 (2004).
49. S.L. Rees, A.L. Robinson, A. Khlystov, C.O. Stanier, and S.N. Pandis, "Mass balance closure and the Federal Reference Method for PM_{2.5} in Pittsburgh, Pennsylvania," *Atmospheric Environment*, 38 (20), 3305-3318 (2004).
50. S.N. Pandis, "Estimates of diesel and other emissions: Overview of the Supersite program," in *Improving Estimates of Diesel and Other Emissions for Epidemiological Studies*, HEI Communication 10, Health Effects Institute, Boston, MA, (2003).
51. S.S. Park, Pancras, P., Ondov, J. M., Robinson, A. 2005, "Application of the Pseudo-Deterministic Multivariate Receptor Model to resolve Power Plant Influences on Pittsburgh Air Quality." *Atmos. Environ.*, in preparation.
52. T.M. Gaydos, C.O. Stanier CO, S.N. Pandis (2005) "Modeling of in situ ultrafine atmospheric particle formation in the eastern United States," *J. Geophys. Res.*, 110 (D7): Art. No. D07S12.
53. W. Tang, T. Raymond, B. Wittig, C. Davidson, S.N. Pandis, A. Robinson, and K. Crist, (2004) "Spatial variations of PM_{2.5} during the Pittsburgh Air Quality Study," *Aerosol Science and Technology*, 38(S2): 80-90.
54. W.K. Modey, D.J. Eatough, R.R. Anderson, D.V. Martello, S. Takahama, L.L. Lucas, and C.I. Davidson, "Ambient fine particulate concentrations and chemical

composition at two sampling sites in metropolitan Pittsburgh: a 2001 intensive summer study,” *Atmospheric Environment*, 38 (20), 3165-3178 (2004).

55. Y. Zhao, K. J. Bein, A. S. Wexler, C. Misra, P. M. Fine, and C. Sioutas (2005) Field evaluation of the VACES particle concentrator coupled to the RSMS-3 single particle mass spectrometer, *J. Geophys. Res.*, 110 (D7): Art. No. D07S02.

Presentations:

1. “Investigation of nucleation bursts in the Pittsburgh air quality study”, 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (C. O. Stanier, A. Y. Khlystov, and S. N. Pandis).
2. “Monitoring of water content of ambient aerosol during the Pittsburgh Air Quality Study” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (A. Y. Khlystov, C. O. Stanier, D. Vayenas, and S. N. Pandis).
3. Performance of the Aerodynamic Particle Sizer 3320 during the Pittsburgh Air Quality Study (PAQS)” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (A. Khlystov, C. Stanier, and S. N. Pandis).
4. “Sulfate-ammonia-nitric acid interactions in an urban area” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (S. Takahama, A. Khlystov, B. Wittig, S. V. Hering, C. Davidson, A. Robinson, and S. N. Pandis).
5. “Sampling artifacts during measurement of ambient carbonaceous aerosol” 6th International Aerosol Conference, Taipei, Taiwan, September 2002 (R. Subramanian, A. Y. Khlystov, J. C. Cabada, S. N. Pandis, and A. L. Robinson).
6. “Formation and properties of regional aerosol: Some insights from the Pittsburgh Air Quality Study”, NASA-GSFC, Greenbelt MD, May 2002, (C. Stanier, A. Khlystov, S. Rees, J. Cabada, A. Robinson, C. Davidson, and S. N. Pandis)
7. “Seasonal composition of PM_{2.5} and performance of the Federal Reference Method in Pittsburgh”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (S. L. Rees, S. Takahama, A. L. Robinson, A. Khlystov, and S. N. Pandis).
8. “Continuous measurements of ammonia, sulfate, and nitrate in Pittsburgh: Implications for PM_{2.5} control strategies”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (B. Wittig, A. Khlystov, S. Takahama, C. Davidson, A. Robinson, S. Hering, and S. N. Pandis).
9. “The contribution of long-range transport and secondary organic aerosol to PM_{2.5} in Pittsburgh”, PM_{2.5} and Electric Power Generation, Pittsburgh, April 2002 (J. C. Cabada, R. Subramanian, S. N. Pandis, A. L. Robinson, W. Tang, N. J. Anderson, T. Raymond, and C. I. Davidson).

10. "The Dry-Ambient Size Spectrometer: A new technique for the automatic on-line measurement of the atmospheric aerosol water size distribution", Annual Meeting of American Geophysical Union, San Francisco, December 2001 (A. Khlystov, C. O. Stanier, S. N. Pandis).
11. "The July 2001 intensive of the Pittsburgh Air Quality Study", Annual Meeting of AAAR, Portland, Oregon, October 2001 (C. I. Davidson, A. L. Robinson, and A. Khlystov, S. N. Pandis).
12. "Sources of atmospheric carbonaceous particulate matter in Pittsburgh", Annual Meeting of AAAR, Portland, Oregon, October 2001 (J. Cabada, S. N. Pandis and A. L. Robinson).
13. "Automated measurements of dry and wet ambient aerosol distributions", Annual Meeting of AAAR, Portland, Oregon, October 2001 (A. Y. Khlystov, W. R. Chan, C. O. Stanier, M. Mandiro, and S. N. Pandis)
14. "Continuous measurements of ammonia and ammonium in ambient air", Annual Meeting of AAAR, Portland, Oregon, October 2001 (A. Khlystov, J. Sauser, R. Otjes, and S. N. Pandis).
15. The contribution of secondary organic aerosol to PM_{2.5} concentrations in Pittsburgh, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (J. C. Cabada, S. N. Pandis, A. L. Robinson, R. Subramanian, A. Polidori, and B. Turpin).
16. Preliminary results from the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (S. N. Pandis, C. I. Davidson, A. L. Robinson, and A. Y. Khlystov)
17. Monitoring of water content of ambient aerosol during the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (A. Y. Khlystov, C. O. Stanier, D. Vayenas, and S. N. Pandis)
18. Investigation of nucleation bursts during the Pittsburgh Air Quality Study, AGU Fall Meeting 2002, San Francisco CA Dec. 2002 (C. O. Stanier, A. Y. Khlystov, B. Wittig, S. N. Pandis, Y. Zhou, K. Bein, A. S. Wexler, C. Misra, and C. Sioutas)
19. Atmospheric particulate matter: Physics, chemistry, and Chemical Transport Models, PM AAAR 2003, Pittsburgh PA March 2003 (B. Koo, K. Fahey, T. Gaydos, and S. N. Pandis)
20. Secondary organic aerosol contribution to carbonaceous PM_{2.5} concentrations in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (J. C. Cabada, S. N. Pandis, B. Wittig, A. Robinson, R. Subramanian, A. Polidori, and B. J. Turpin)

21. Using ultrafine concentrators to increase the hit rates of single particle mass spectrometers, PM AAAR 2003, Pittsburgh PA March 2003 (Y. Zhao, K. J. Bein, A. S. Wexler, C. Misra, P. M. Fine, and C. Sioutas)
22. PM_{2.5} Federal Reference Method performance relative to mass balance closure, PM AAAR 2003, Pittsburgh PA March 2003 (S. L. Rees, A. L. Robinson, A. Khlystov, C. O. Stanier, and S. N. Pandis)
23. Examining the assumptions behind elemental carbon measurements using the thermal-optical transmittance technique, PM AAAR 2003, Pittsburgh PA March 2003 (R. Subramanian, A. Y. Khlystov, and A. L. Robinson)
24. Spatial variations of PM_{2.5} during intensive sampling of PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (W. Tang, C. I. Davidson, T. R. Raymond, S. N. Pandis, B. Wittig, A. Khlystov, and A. L. Robinson)
25. Fenceline sampling adjacent to a large coke production facility in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (E. A. Weitkamp, E. Lipsky, A. Robinson, N. Anderson, H. Leifeste, R. Subramanian, J. Cabada, A. Khlystov, C. Stanier, L. Lucas, S. Takahama, B. Wittig, C. Davidson, S. Pandis, A. Polidori, H. J. Lim, B. Turpin, P. Pancras, and J. Ondov)
26. In-use vehicle emissions source characterization study: Squirrel Hill tunnel Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (E. M. Lipsky, A. Robinson, N. Anderson, H. Leifeste, R. Subramanian, J. Cabada, S. Rees, A. Khlystov, C. Stanier, L. Lucas, S. Takahama, B. Wittig, C. Davidson, S. N. Pandis, A. Polidori, H. J. Lim, and B. Turpin)
27. Water content of ambient aerosol during PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (A. Khlystov, C. Stanier, and S. N. Pandis)
28. Diurnal and seasonal trends in outdoor particle size distributions measured at urban and rural locations during PAQS (C. Stanier, A. Khlystov, and S. N. Pandis)
29. Mass and chemically resolved size compositions of fine particulate matter at the Pittsburgh Supersite, PM AAAR 2003, Pittsburgh PA March 2003, (J. C. Cabada, S. N. Pandis, S. Rees, S. Takahama, A. Khlystov, A. L. Robinson, and C. I. Davidson)
30. Simulation of the atmospheric aerosol size/composition distribution in a three-dimensional chemical transport model, PM AAAR 2003, Pittsburgh PA March 2003 (T. M. Gaydos, K. M. Fahey, B. Koo, and S. N. Pandis)
31. Application of PMCAMx to the South Coast Air Basin and the Eastern United States, PM AAAR 2003, Pittsburgh PA March 2003 (B. Koo, K. M. Fahey, T. M. Gaydos, and S. N. Pandis)

32. Principal component analysis of trace elements in PM_{2.5} in Pittsburgh, PM AAAR 2003, Pittsburgh PA March 2003 (N. J. Anderson, C. I. Davidson, S. N. Pandis, A. Robinson, and A. Khlystov)
33. Source apportionment using particle size distribution data from PAQS, PM AAAR 2003, Pittsburgh PA March 2003 (L. Zhou, E. Kim, P. K. Hopke, C. Stanier, and S. N. Pandis)
34. Highly time-resolved measurements of elemental composition at the Baltimore, St. Louis, and Pittsburgh Supersites using the UM High Frequency Aerosol Slurry Sampler: Unprecedented resolution of the sources of primary atmospheric aerosol PM AAAR 2003, Pittsburgh PA March 2003 (J. M. Ondov, J. Pancras, S. Gazula, M. Yu, J. Turner, A. Robinson, S. N. Pandis, N. D. Poor, and R. K. Stevens)
35. "Size Resolved Chemical Classification of Dual Polarity Single-Ultrafine-Particle Mass Spectrometry Data Collected During Pittsburgh Supersite Experiment", K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.
36. "Dynamic Data Classification Using a Component-Weighted Similarity Algorithm", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.
37. "Diurnal Variations of Ultra-fine Particles in Pittsburgh Measured by a Rapid Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 2003.
38. "Rapid Single Particle Mass Spectrometry and EPA Supersite Experiments", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, NYU Institute of Environmental Medicine, Tuxedo, NY, December 2002.
39. "Speciation of Size-Resolved Individual Ultrafine Particles in Pittsburgh, Pennsylvania", *K.J. Bein, Y. Zhao, A.S. Wexler, D.A. Lake, D.B. Kane, M.P. Tolocka, K.P. Rhoads, M.V. Johnston and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
40. L. Zhou, E. Kim, and P.K. Hopke Analysis of the Particle Size Distribution Data Collected During Pittsburgh Air Quality Study, Presented to 21st Annual Conference of the American Association for Aerosol Research, Charlotte, NC, October 7-11, 2002.
41. L. Zhou, E. Kim, P.K. Hopke, C. Stanier, and S. Pandis Source Apportionment Using Particle Size Distribution Data from the Pittsburgh Air Quality Study (PAQS), presented to 2003 AAAR PM Meeting on Particulate Matter:

Atmospheric Sciences, Exposure and the Fourth Colloquium on PM and Human Health, Pittsburgh, PA, March 30 to April 4, 2003.

42. L. Zhou, P.K. Hopke, and W. Liu, Comparison of three back trajectory based models--PSCF, SQTBA, residence-time weighted concentration model for identifying particle sources for Potsdam and Stockton, New York, presented to 21st Annual Conference of the American Association for Aerosol Research, Anaheim, CA, October 20-24, 2003.
43. L. Zhou, P.K. Hopke, and P. Paatero, Advanced Factor Analysis for Aerosol Composition Data with Various Temporal Resolutions, presented to 21st Annual Conference of the American Association for Aerosol Research, Anaheim, CA, October 20-24, 2003.
44. S.G. Buckley, F. Ferioli, G.A. Lithgow, "Combustion System Analysis Using Laser-Induced Breakdown Spectroscopy," Paper 403, 22nd International Congress on Applications of Lasers and Electro-Optics, Jacksonville, FL, October 13-16, 2003.
45. G.A. Lithgow, A.L. Robinson, and S.G. Buckley, "Ambient Measurements of Inorganic Species in an Urban Environment Using LIBS," Second International Conference on Laser-Induced Plasma Spectroscopy, Orlando, FL, September 24 – 27, 2002.
46. G.A. Lithgow, A.L. Robinson, S.G. Buckley, "Ambient Particle Measurements in an Urban Environment Using Laser-Induced Breakdown Spectroscopy (LIBS)," Paper 11E4, American Association of Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
47. Ondov, J. M. (2001). Sources of Metals Influencing Air Quality at the Pittsburgh Supersite: Data collected with the UM High-Frequency Aerosol Slurry Sampler.
48. Ondov, J. M. (2002). Unprecedented Source Apportionment with the University of Maryland Semi-Continuous Elements in Aerosol System Presented at the Regional AQ Modeling & Data Analysis Meeting, sponsored by MARAMA, OTC, NESCAUM and MANE-VU, January 23-24, 2002, Baltimore, MD.
49. Ondov, J. M., (2003) Short-Term Concentrations of Metals SEAS via SEAS and Selected Variables. Presented at the Mid-Atlantic Region Air Monitoring Association Conference in Baltimore, Jan 21-22, at the Harbor Court Hotel.
50. Ondov, J. M., Pancras, J. P., Gazula, S., Moore, J. A., Park, S. S., Chang, Y. C., Squib, K., Powel, J., Mitkus, R., Turner, J., Yu, M.N.S., Robinson, A., Pandis, S., Davidson, C. (2003). Identification of Sources From High-Frequency Elements Measurements at 4 Supersites. Semi-annual EPA Supersites meeting in Atlanta, Jan 22-23.

51. Ondov, J. M. (2003). High-frequency metals measurements at the Pittsburgh Supersite and Coke Plant site with the University of Maryland SEAS. Seminar presented at the Pittsburgh Supersite Project workshop, March 5, Carnegie Mellon University.
52. Ondov, J. M. (2003) Highlights of SEAS Metals Data for the Pittsburgh Supersite. Presented at the Pittsburgh Air Quality Study Meeting, Pittsburgh, PA, April 4,5, 2003.
53. Ondov, J. M. (2003) New Paradigm for Air Pollution Control: Pseudo Deterministic Receptor Modeling of Highly Time Resolved Ambient Aerosol Composition Data Derived from the UMCP SEAS. Seminar presented to the Source Apportionment Group, US EPA, Research Triangle Park, November 12.
54. Ondov, J. M., Emission Rates of Pollutants from Stationary Sources Using Highly Time Resolved Ambient Measurements and a New Pseudo Deterministic Hybrid Receptor Model. Presented at the EPA Center for Hazardous Substances in Urban Environments, Research Program Internal Workshop, January 5, 2004.
55. Ondov, J. M., New Pseudo Deterministic Model for Individual Source Apportionment Using Highly Time Resolved Data. Presented at the MARAMA MANE-VU Science Meeting, Baltimore, MD, January 27-29, 2004.
56. Park, S. S., Pancras, J. P., Gazula, S., Ondov, J. M. (2002) Sources of Elemental Aerosol Constituents in Pittsburgh Using Positive Matrix Factorization of Highly Time-resolved Data. Presented at the American Association of Aerosol Research meeting, 21st Annual AAAR Conference October 7-11, Charlotte.
57. Ondov, J. M., Pancras, J. P., Gazula, S., Yu, M. N. S., Turner, J., Robinson, A., Pandis, S., Stevens, R. K., Poor (2003). Highly Time-Resolved Measurements of Elemental Composition at the Baltimore, St. Louis, Pittsburgh, and Tampa Supersites Using the UM High-Frequency Aerosol Slurry Sampler: Unprecedented Resolution of the Sources of Primary Atmospheric Aerosol. To be presented at the Association of Aerosol Research meeting, Particulate Matter: Atmospheric Sciences, Exposure, and the Fourth Colloquium on PM and Human Health, March, Pittsburgh.
58. Ondov, J. M., Pancras, J. P., Park, S. S., Poor, N., Turner, J. R., Yu, M., Lipsky, E., Weitkamp, E., Robinson, A. (2003) PM emission rates from highly time-resolved ambient concentration measurements. Presented at the October meeting of the American Society for Aerosol Research, San Diego.
59. Ondov, J. M., Poor, N. (2003) Emission Inventory Development through highly-time-resolved ambient sampling. NARSTO Workshop on Innovative Methods for Emission-Inventory Development and Evaluation, University of Texas, Austin; October 14-17, 2003.

60. G.A. Lithgow and S.G. Buckley, "Detection and Classification of Biological Aerosols Using Laser-Induced Breakdown Spectroscopy," Paper 10B4, American Association of Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
61. S.G. Buckley "Laser-Induced Breakdown Spectroscopy for Detection of Biological Aerosols - Potential and Perspective," PITTCON 2005, Orlando FL, Feb. 27-Mar. 4, 2005. (invited)
62. S.G. Buckley "LIBS as a Combustion and Aerosol Diagnostic," PacificChem 2005, Honolulu, HI, Dec 15-20, 2005. (invited)
63. Cabada, J.C., Pandis, S.N., Robinson, A.L., Davidson, C.I., Polidori, A., Turpin, B.J., and Subramanian, R., "The contribution of Secondary Organic Aerosol to PM_{2.5} in Pittsburgh," Presented at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
64. Offenberg, J.H., Polidori, A., Porcja, R., Turpin, B.J., "Functional group composition by size and polarity in Pittsburgh, PA and insights into aerosol processing," Presented at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
65. Polidori, A., Turpin, B.J., Lim, H.J., Robinson, A., Subramanian, R., Cabada, J.C., "Semi-continuous organic particulate matter measurements during Pittsburgh Air Quality Study (PAQS)," Poster presentation at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
66. Subramanian, R., Robinson, A.L., Cabada, J.C., Pandis, S.N., Wittig, B., Polidori, A., Turpin, B.J., Hering, S.V., Modey, W.K., Eatough, D.J., "Intercomparison of ambient carbonaceous aerosol samplers used during the Pittsburgh Air Quality Study," Poster presentation at the American Association for Aerosol Research Annual Conference, Charlotte, NC, October 2002.
67. Cabada, J.C., Pandis, S.N., Robinson, A.L., Davidson, C.I., Polidori, A., Turpin, B.J., and Subramanian, R., "The contribution of Secondary Organic Aerosol to PM_{2.5} in Pittsburgh," Platform Presentation at the Annual Meeting of the American Institute of Chemical Engineers, Indianapolis, IN, November 2002.
68. Millet, D.B., Donahue, N.M., Polidori, A., Stanier, C.O., Turpin, B.J., Goldstein, A.H., "VOC-Aerosol Relationships at the Pittsburgh Air Quality Study", Presented at the Berkeley Atmospheric Science Center symposium, Berkeley, CA, September 2003.
69. Polidori, A., Turpin, B., Lim, H.J., Cabada, J.C., Subramanian, R., Robinson, A., "Secondary Organic Aerosol Formation During the Pittsburgh Air Quality Study (PAQS)", Presented at the Environmental Sciences Graduate Student Association (ESGSA), New Brunswick, NJ, June 2004.

70. Polidori, A., Turpin, B.J., Lim, H.J., Totten L., Davidson, C, "Characterization of the Organic Fraction of Atmospheric Aerosols" Platform Presentation at the American Association for Aerosol Research Annual Conference, Atlanta, GA, October 2004.
71. Robinson, A.L., Subramanian, R., Gaydos, T., Pandis, S.N., Bernardo-Bricker, A., Rogge, W.F., Polidori, A., Turpin, B.J., Clarke, L., Hernandez, M., "Synthesis of Source Apportionment Estimates of Organic Aerosol in the Pittsburgh Region" Platform Presentation at the American Association for Aerosol Research Annual Conference, Atlanta, GA, October 2004.
72. Polidori, A., Turpin, B., Lim, H.J., Subramanian, R., Robinson, A., Pandis, S., Cabada, J.C., "Local and Regional Secondary Organic Aerosol Formation: Insights from a Year at Pittsburgh and Comparisons with Los Angeles and Atlanta," Platform Presentation at the American Association for Aerosol Research "Supersite" Conference, Atlanta, GA, February 2005.
73. Weitkamp, E., Lipsky, E., Robinson, A., Polidori, A., Turpin, B., Pancras, P., Ondov, J., Bernado-Bricker, A., Vasquez, O., Rogge, W., "Fine Particle Emission Profile for a Large Coke Production," Poster presentation at the American Association for Aerosol Research "Supersite" Conference, Atlanta, GA, February 2005.
74. "Source Sampling and Characterization Using a Single Particle Mass Spectrometer during the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
75. "Size-Resolved Chemical Classification of Single Particle Mass Spectrometry Data Collected during the Pittsburgh Supersite Experiment: Source Attribution", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
76. "Laboratory Experiments Examining Ultrafine Particle Production by Re-breathing of Road Dust through a Diesel Engine", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky and A.L. Robinson, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
77. "Detection of a Contaminating Plume during a Roadway Tunnel Source Sampling Experiment Using a Single Particle Mass Spectrometer", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.

78. "Detection and Characterization of a Smoke Plume from Canadian Forest Fires during the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler, N.J. Pekney, C.I. Davidson and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
79. "Field Evaluation of the VACES Particle Concentrator Coupled to the RSMS-3 Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, C. Misra, P.M. Fine and C. Sioutas, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
80. "Identification of Sources of Atmospheric PM at the Pittsburgh Supersite: RSMS-III and Filter-based Positive Matrix Factorization", *N.J. Pekney, K.J. Bein, C.I. Davidson, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research International Specialty Conference, Atlanta, GA, February 7-11, 2005.
81. "Laboratory Experiments Examining Ultrafine Particle Production by Re-breathing of Road Dust through a Diesel Engine", *K.J. Bein, Y. Zhao, A.S. Wexler, E. Lipsky and A.L. Robinson, American Association for Aerosol Research Annual Conference, Atlanta, GA, October 4-8, 2004.
82. "Measurements of Nitrate Particles in Pittsburgh, PA, Using Rapid Single Particle Mass Spectrometry", *Y. Zhao, K.J. Bein, A.S. Wexler, M.P. Tolocka and M.V. Johnston, American Association for Aerosol Research Annual Conference, Atlanta, GA, October 4-8, 2004.
83. "Size Resolved Chemical Classification of Dual Polarity Single-Ultrafine-Particle Mass Spectrometry Data Collected During the Pittsburgh Supersite Experiment", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
84. "Dynamic Data Classification Using a Component-Weighted Similarity Algorithm", *K.J. Bein, Y. Zhao, A.S. Wexler and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
85. "Diurnal Variations of Ultra-fine Particles in Pittsburgh Measured by a Rapid Single Particle Mass Spectrometer", *Y. Zhao, K.J. Bein, A.S. Wexler, and M.V. Johnston, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
86. "Experimental Validation of a High Transmission Rate Inlet for Ultra-fine Single Particle Mass Spectrometers", *Y. Zhao, K.J. Bein, P. Middha and A.S. Wexler, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
87. "Synthesis of Research on In-situ Particle Nucleation in Western Pennsylvania: Which Hypotheses for Nuclei Formation and Growth are Consistent with Field

- Observations, Mass Spectrometry, and Modeling?.”, *C. Stanier, T. Gaydos, A.Y. Khlystov, Q. Zhang, M. Caragaratna, J. Jayne, D. Worsnop, J. Jimenez, D. Millet, A. Goldstein, K.J. Bein, Y. Zhao, A.S. Wexler and S.N. Pandis, American Association for Aerosol Research Annual Conference, Anaheim, CA, October 20-24, 2003.
88. “Speciation of Size-Resolved Individual Ultrafine Particles in Pittsburgh, Pennsylvania”, *K.J. Bein, Y. Zhao, A.S. Wexler, D.A. Lake, D.B. Kane, M.P. Tolocka, K.P. Rhoads, M.V. Johnston and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
 89. “Using Ultrafine Concentrators to Increase the Hit Rate of Single Particle Mass Spectrometers”, *Y. Zhao, K.J. Bein, A.S. Wexler, C. Misra, P.M. Fine and C. Sioutas, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
 90. “Investigation of Urban Aerosols at the Baltimore Particulate Matter Supersite by Single Particle Mass Spectrometry”, *D.A. Lake, M.P. Tolocka, M.V. Johnston, K.J. Bein, Y. Zhao, A.S. Wexler, K.P. Rhoads, and D.J. Phares, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
 91. “Investigation of Nucleation Bursts during the Pittsburgh Air Quality Study, *C. Stanier, A.Y. Khlystov, B. Wittig, S.N. Pandis, K.J. Bein, Y. Zhao, A.S. Wexler, C. Misra and C. Sioutas, Association for Aerosol Research Annual Conference, Charlotte, NC, October 7-11, 2002.
 92. G. Casuccio*, T. Lersch, R.J. Lee, A. Robinson, E. Weitkamp, and D. Martello, “Estimating of the Impact of Primary Coal Fired Boiler Emissions to PM_{2.5} And PM₁₀ Using Computer Controlled Scanning Electron Microscopy Data,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
 93. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson, and M.V. Johnston, “Source Sampling and Characterization Using a Single Particle Mass Spectrometer During the Pittsburgh Supersite Experiment,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
 94. M. Hernandez*, L. Clarke, L. Angenent, and A. Robinson, “Primary Biopolymer Associations with Fine Particulate Matter,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
 95. S.S. Park*, J.P. Pancras, J.M. Ondov, A. Robinson, and C. Davidson, “Application of the UM Multivariate Pseudo-Deterministic Receptor Model to Resolve Power Plant Influences on Air Quality at the CMU Supersite,” American Association of

Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.

96. R. Subramanian*, A. Robinson, and A. Khlystov, “OC/EC Analysis with Thermal-Optical Methods: Effects of Temperature Protocol and Non-Carbonaceous Compounds,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
97. A. Wittig*, A. Robinson, and S. Pandis, “Continuous PM_{2.5} Mass Measurements at the Pittsburgh Air Quality Study Supersite,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
98. W. Tang, T. Raymond, B. Wittig, C. Davidson*, S. Pandis, A. Robinson, and K. Crist, “Spatial Variations of PM_{2.5} in the Pittsburgh Region,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
99. A. Polidori*, B. Turpin, H.J. Lim, R. Subramanian, A. Robinson, S. Pandis, and J.C. Cabada, “Local and Regional Secondary Organic Aerosol Formation: Insights from a Year at Pittsburgh and Comparison with Los Angeles and Atlanta,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
100. T. Lane, R. Pinder, M. Shrivastava, S.N. Pandis, and A.L. Robinson*, “Evaluation of the Primary Organic Carbon Emission Inventory for the Eastern United States,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
101. O. Sevimoglu*, W.F. Rogge, A. Bernardo, A. Robinson and R. Subramanian, “Fine Particulate Abrasion Products from Leaf Surfaces of Urban Plants: Comparison Between Los Angeles and Pittsburgh,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
102. A. Bernardo-Bricker*, W.F. Rogge, O. Sevimoglu, A. Robinson, E. Lipsky, and A. Grieshop, “Source Profiles for Organic PM_{2.5} for Diesel Trucks and Gasoline Vehicles Determined for the Squirrel Hill Tunnel in Pittsburgh, PA,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
103. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, and A.L. Robinson, “Laboratory Experiments Examining Ultrafine Particle Production by Re-Breathing of Road

- Dust Through a Diesel Engine,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
104. K.J. Bein*, Y. Zhao, A.S. Wexler, E. Lipsky, A.L. Robinson, and M.V. Johnston, “Detection of a Contaminating Plume During a Roadway Tunnel Source Sampling Experiment Using a Single Particle Mass Spectrometer,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
 105. A.L. Robinson*, S.L. Rees, A. Khlystov, C.O. Stanier, and S.N. Pandis, “Mass Balance Closure and the Federal Reference Method for PM_{2.5} in Pittsburgh, Pennsylvania,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
 106. R. Subramanian*, A. Robinson, A. Bernardo-Bricker, and W.F. Rogge, “Organic Carbon Mass Balance and Source Apportionment of Primary Organic Carbon in the Pittsburgh Region Using Molecular Markers,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.
 107. A.L. Robinson*, E.M. Lipsky, N. Pekney, W.F. Rogge, A. Bernardo-Bricker, and O. Sevimoglu, “Fine Particle Emission Profile for Road Dust in Pittsburgh, Pennsylvania,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
 108. E. Weitkamp*, E. Lipsky, A. Robinson, A. Polidori, B. Turpin, P. Pancras, J. Ondov, A. Bernardo-Bricker, O.R. Vasquez, and W.F. Rogge, “Fine Particle Emission Profile for a Large Coke Production, American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
 109. A. Grieshop*, E. Lipsky, and A. Robinson, “Fuel-Based Particulate Matter and Gaseous Emission Factors Determined from Vehicles in Pittsburgh, PA’s Squirrel Hill Tunnel,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005. (poster)
 110. A.L. Robinson*, N.M. Donahue, K. Huff Hartz, A. Sage and E. Weitkamp, “Field and Laboratory Experiments Examining the Stability of Organic Molecular Markers Used for Source Apportionment,” American Association of Aerosol Research (AAAR) Specialty Conference: Particulate Matter, Supersites Program & Related Studies, Atlanta, GA, February 7–11, 2005.

111. A.L. Robinson*, N.M. Donahue, A.M. Sage, K.E. Huff Hartz, and E.A. Weitkamp, "Field and Laboratory Experiments Examining the Stability of Organic Molecular Markers Used for Source Apportionment," 2004 AGU Fall Meeting, San Francisco, CA, December 13–17, 2004.
112. A.L. Robinson*, R. Subramanian, A. Bernardo-Bricker, and W.F. Rogge, "Source Apportionment of Primary Organic Carbon in the Pittsburgh Region using Molecular Markers and Different Receptor Models," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
113. K.E. Huff Hartz*, E.A. Weitkamp, A.M. Sage, A.A. Presto, A.L. Robinson, and N.M. Donahue, "Kinetics of Atmospheric Processing of Organic Particulate Matter: A Relative Rates Approach," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
114. E.A. Weitkamp*, K.E. Huff Hartz, A.M. Sage, A.L. Robinson, N.M. Donahue, W.F. Rogge, and A. Bernardo-Bricker, "Evaluation of the Oxidation Kinetics of Molecular Markers Used for Source Apportionment of Primary Organic Aerosol," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
115. A.L. Robinson*, R. Subramanian, T. Gaydos, S.N. Pandis, A. Bernardo-Bricker, W.F. Rogge, A. Polidori, B.J. Turpin, L. Clarke, and M. Hernandez, "Synthesis of Source Apportionment Estimates of Organic Aerosol in the Pittsburgh Region," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
116. E. M. Lipsky* and A.L. Robinson, "Effects of Dilution Ratio and Residence Time on the Partitioning of Semi-Volatile Organic Carbon in Emissions from a Wood Stove And Diesel Engine," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
117. K.J. Bein*, Y. Zhao, A.S. Wexler, E.M. Lipsky, and A.L. Robinson, "Laboratory Experiments Examining Ultrafine Particle Production by Rebreathing of Road Dust Through a Diesel Engine," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
118. A.M. Sage*, K.E. Huff Hartz, E.A. Weitkamp, A.L. Robinson, and N.M. Donahue, "Laboratory Measurement of Heterogeneous Oxidation Kinetics of Organic Aerosols," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)
119. A.P. Grieshop*, E.M. Lipsky, and A.L. Robinson, "Fuel-Based Particulate Matter and Gaseous Emission Factors determined from Vehicles in Pittsburgh, PA's Squirrel Hill Tunnel," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004. (poster)

120. N.M. Donahue* and A.L. Robinson, "Theoretical, In Situ, and Laboratory Constraints on Organic Aerosol Oxidation," American Association of Aerosol Research 23rd Annual Conference (AAAR 2004), Atlanta, GA, October 4–8, 2004.
121. A.L. Robinson*, "Are Organic Molecular Markers Used for Source Apportionment Stable?," presented at Telluride Workshop on Atmospheric Aerosol Aging, Telluride, CO, August 9–13, 2004. (invited)
122. A.L. Robinson*, "Comparison of Emission- and Receptor-based Model Predictions of Primary Organic Aerosol in Pittsburgh, PA," presented at LADCO Academic Community Meeting, Chicago, IL, June 11, 2004. (invited)
123. A.L. Robinson*, "Sources of Organic Particulate Matter in Pittsburgh, PA," presented at Mid-Atlantic Regional Air Management Association Science Meeting (MANE-VU/MARAMA 2004), Baltimore, MD, January 27–29, 2004 (invited).