



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division


Submitted to Physical Review D

LIMITS ON NEUTRINO OSCILLATIONS FROM MUON-DECAY NEUTRINOS

P. Némethy, R.L. Burman, D.R.F. Cochran, J. Duclos,
J.S. Frank, C.K. Hargrove, V.W. Hughes, H. Kaspar,
U. Moser, R.P. Redwine, and S.E. Willis

June 1980

RECEIVED
LAWRENCE
BERKELEY LABORATORY
DEC 3 1980
LIBRARY AND
DOCUMENTS SECTION



LBL-11104cs

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Limits on Neutrino Oscillations from Muon-decay Neutrinos

P. Némethy, R.L. Burman, D.R.F. Cochran, J. Duclos, J.S. Frank,
C.K. Hargrove, V.W. Hughes, H. Kaspar, U. Moser, R.P. Redwine
and S.E. Willis^(a)

Yale University, New Haven, CT 06530, Los Alamos Scientific Laboratory,
Los Alamos, NM 87545, Lawrence Berkeley Laboratory, Berkeley, CA 94720,
Massachusetts Institute of Technology, Cambridge, MA 02139, National
Research Council, Ottawa, Ontario K1A 0R6, Canada, Centre d'Etudes
Nucléaires de Saclay, F-91190 Gif-sur-Yvette, France, Swiss Institute
for Nuclear Research, CH-5234 Villigen, Switzerland and University of
Berne, CH-3012 Berne, Switzerland.

Abstract: No evidence for neutrino oscillations is seen in our experi-
ment which observed neutrinos from muon-decays at rest.
Upper limits on oscillation parameters are presented for
neutrino mixing of the kind $\nu_e \leftrightarrow \nu_\mu$ and also of the kind
 $\nu_e \leftrightarrow \nu_i$, $i \neq \mu$.

In a recent Letter¹, we pointed out that our neutrino experiment on
the nature of muon conservation also provides an upper limit on neutrino
oscillations. Here we present a more detailed analysis of this result.

Neutrino oscillations, first proposed by B. Pontecorvo² and by Z.
Mako, et al³, are of considerable interest in the light of gauge
theories with broken lepton flavor symmetry. Experimental upper limits
on neutrino oscillations have been reported by E. Bellotti et al.,⁴ and
by J. Blietschau et al.⁵; F. Reines et al. have reported evidence for
neutrino instability.⁶

In our analysis we make use of our previously published evidence¹
that muon conservation is an additive law. We also make the simplifying
assumption that oscillations occur between only two neutrino states.
Neutrino mixing is then described by a 2 x 2 matrix and the oscillations
depend on two parameters, the mixing angle θ , and the mass difference
 $\Delta = (m_1^2 - m_2^2)$ between neutrino mass eigenstates. The oscillation

(a) Present address: Fermilab, Batavia, IL 60510

probability for neutrinos of momentum p at a distance D from the source is given by

$$P(\nu_a \rightarrow \nu_b) = 0.5 \sin^2 2\theta (1 - \cos \frac{D\Delta}{2p}) . \quad (1)$$

In the experiment we utilized a six-ton water Cerenkov counter to observe ν_e and $\bar{\nu}_e$ from the decay chain $\pi^+ \rightarrow \mu^+ \nu_\mu$ (at rest) and $\mu^+ \rightarrow e^+ \nu_e \bar{\nu}_\mu$ (at rest) by the charged current reactions $\bar{\nu}_e p \rightarrow n e^+$ (in H_2O) and $\nu_e d \rightarrow p p e^-$ or $\bar{\nu}_e d \rightarrow n n e^+$ (in D_2O). The neutrino source was the Clinton P. Anderson Meson Physics Facility (LAMPF) beam stop at a mean distance of 9 m from the detector. For details, see Ref. 1. The results⁷ are

$$R = \bar{\nu}_e / \mu^+ \text{ decay} = 0.00 \pm 0.06 \quad (2)$$

and

$$R' = \nu_e / \mu^+ \text{ decay} = 1.09 \pm \begin{matrix} 0.37 \\ 0.41 \end{matrix} \quad (3)$$

where we have added (in quadrature) a $\pm 10\%$ uncertainty in neutrino flux and a $+25\%/-10\%$ uncertainty⁸ in the neutrino deuteron cross-section calculation of J.S. O'Connell⁹ to our experimental error in R' .

Our null result (2) for R is a direct upper limit on $\nu_e \leftrightarrow \nu_\mu$ oscillations producing $\bar{\nu}_e$ from the $\bar{\nu}_\mu$ in the muon decay. To evaluate this limit we weight the muon-decay $\bar{\nu}_\mu$ spectrum by the E^2 dependence of the cross section and by the oscillation probability (1) averaged over the finite detector size (1.8 m) to obtain a predicted spectrum shape and normalization for any combination of the oscillation parameters Δ and θ . After folding in the experimental resolution we fit these spectra to our observed spectrum of H_2O events (Ref. 1, Fig. 2) above our energy cutoff of 25 MeV, to obtain the 68% and 90% confidence level upper limits on Δ , as a function of the mixing parameter, $\sin^2 2\theta$, shown in Fig. 1.

Our heavy water measurement (3) does not distinguish electron neutrinos and electron antineutrinos. Since two muon neutrinos are produced for every electron neutrino in the $\pi-\mu-e$ decay sequence,

$\nu_\mu \leftrightarrow \nu_e$ oscillations would increase R' . However, our water measurement (2), which yielded Fig. 1, is a far more sensitive test for oscillations of this kind and limits their contribution to R' to a negligible level.

In the absence of $\nu_e \leftrightarrow \nu_\mu$ oscillations, ν_e can still disappear by oscillations of the kind $\nu_e \leftrightarrow \nu_i$, $i \neq \mu$, (e.g., $\nu_e \leftrightarrow \nu_\tau$), thus decreasing R' . Therefore, our observation (3) of R' at full strength puts a limit, albeit much weaker because of the big error bars, on such oscillations. For any combination of Δ and θ , we fit the expected spectrum of the original ν_e events, less those that have changed into ν_i , to our observed spectrum (Ref. 1, Fig. 1) of D_2O events (above 25 MeV) with the folding procedure described above. We obtain the 68% and 90% confidence level upper limits on Δ , as a function of $\sin^2 2\theta$, shown in Fig. 2.

We note that the curves of Fig. 1 and Fig. 2 are not asymptotic. The limits oscillate with Δ , dramatically in the case of $\nu_e \leftrightarrow \nu_i$ ($i \neq \mu$). Fig. 3 and Fig. 4 show the large Δ behavior of the limits for both cases.

We conclude that our experiment does not show evidence for neutrino oscillations at the levels of sensitivity indicated in the figures.

This research was supported in part by the U.S. Department of Energy under Contracts No. EY-76-C-3075, No. W-7405-ENG-36 and No. W-7405-ENG-48.

References

1. S.E. Willis, V.W. Hughes, P. Némethy, R.L. Burman, D.R.F. Cochran, J.S. Frank, R.P. Redwine, J. Duclos, H. Kaspar, C.K.Hargrove and U. Moser, Phys. Rev. Lett. 44, 522 (1980).
2. B. Pontecorvo, Zh. Eksp. Teor. Phys. 33, 549 (1957) transl. in Soviet Phys. JETP 6, 429 (1958); B. Pontecorvo, Zh. Eksp. Teor. Fiz. 53, 1725 (1967) transl. in Soviet Phys. JETP 26, 984 (1968).
3. Z. Mako et al., Prog. Teor. Phys. 28, 247 (1962).
4. E. Bellotti et al., Nuovo Cimento Lett. 17, 533 (1976).
5. J. Blietschau et al., Nucl. Phys. B133, 205 (1978).
6. F. Reines et al., Univ. of California at Irvine preprint (1980).
7. After correcting an error in the expected $\bar{\nu}_e p$ cross section in Ref. 1.
8. J.S. O'Connell, private communication.
9. J.S. O'Connell, Los Alamos Scientific Laboratory Report No. LA-5175-MA (1973), unpublished.

Figure Captions

- Fig. 1 Upper limit on $\nu_e \leftrightarrow \nu_\mu$ from H₂O data.
- Fig. 2 Upper limit on $\nu_e \leftrightarrow \nu_i$ ($i \neq \mu$) from D₂O data.
- Fig. 3 Large Δ behavior of $\nu_e \leftrightarrow \nu_\mu$ limit. The allowed region is to the left of the curves.
- Fig. 4 Large Δ behavior of $\nu_e \leftrightarrow \nu_i$ ($i \neq \mu$) limit. The allowed region is to the left of the curves.




Fig. 1





Fig. 2



9-80

3854A3

Fig. 3




Fig. 4