Derived Concentration Guideline Levels for Argonne National Laboratory's Building 330 Area

Environmental Science Division

About Argonne National Laboratory

Argonne is a U.S. Department of Energy laboratory managed by UChicago Argonne, LLC under contract DE-AC02-06CH11357. The Laboratory's main facility is outside Chicago, at 9700 South Cass Avenue, Argonne, Illinois 60439. For information about Argonne and its pioneering science and technology programs, see www.anl.gov.

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor UChicago Argonne, LLC, nor any of their employees or officers, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of document authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, Argonne National Laboratory, or UChicago Argonne, LLC.

Derived Concentration Guideline Levels for Argonne National Laboratory's Building 330 Area

prepared by
S. Kamboj and C. Yu
Environmental Science Division, Argonne National Laboratory

March 18, 2011

CONTENTS

1	INTRODUCTION	1
1.1	Site Description.....	1
1.2	Building History.....	1
1.3	Derivation of Guideline Levels.....	1
2	SCENARIO DESCRIPTION	3
2.1	Land Use Considerations	3
2.2	Exposure Scenarios	3
2.3	Pathways and Key Parameters	4
3	DOSE/SOURCE CONCENTRATION RATIOS.....	8
3.1	Probabilistic Analysis	8
3.2	Deterministic Analysis.....	13
4	DERIVED CONCENTRATION GUIDELINE LEVELS	22
4.1	DCGLs Based on Probabilistic Analysis	22
4.2	DCGLs Based on Deterministic Analysis.....	22
4.3	Summary DCGLs for Building 330 Area	22
5	REFERENCES	25
	APPENDIX A: Parameters Used for Probabilistic Analysis	26
	APPENDIX B: Parameters Used for Deterministic Analysis.....	51
	APPENDIX C: Detailed Probabilistic Analysis Results	69

TABLES

1	Summary of Applicable Exposure Pathways for Different Scenarios Considered for Argonne's Building 330 Area	6
2	List of Key Parameters for Applicable Exposure Pathways.....	7
3	Total Peak DSR Percentiles of Am-241 for Different Exposure Scenarios for the Building 330 Area	10

TABLES (Cont.)

4	Total Peak DSR Percentiles of Ba-133 for Different Exposure Scenarios for the Building 330 Area	11
5	Total Peak DSR of C-14 for Different Exposure Scenarios for the Building 330 Area.....	12
6	Total Peak DSR Percentiles of Cs-137 for Different Exposure Scenarios for the Building 330 Area	13
7	Total Peak DSR Percentiles of Pu-238 for Different Exposure Scenarios for the Building 330 Area	14
8	Total Peak DSR Percentiles of Pu-239 for Different Exposure Scenarios for the Building 330 Area	15
9	Total Peak DSR Percentiles of Sr-90 for Different Exposure Scenarios for the Building 330 Area	16
10	Total Peak DSR Percentiles of Tc-99 for Different Exposure Scenarios for the Building 330 Area	17
11	DSRs at Peak of the Means, Mean of the Peaks, and 95th Percentile of the Peaks for Different Scenarios	18
12	Maximum DSRs for the Industrial Scenario for the Building 330 Area	19
13	Maximum DSRs for the Offsite Resident Scenario for the Building 330 Area.....	19
14	Maximum DSRs for the Recreational Scenario for the Building 330 Area	20
15	Maximum DSRs for the Onsite Resident Scenario for the Building 330 Area	20
16	Total DSRs for Different Scenarios for the Building 330 Area.....	21
17	DCGLs Using Probabilistic Analysis at a 25 mrem/yr Dose Limit for the Building 330 Area	23
18	DCGLs Using Deterministic Analysis at the 25 mrem/yr Dose Limit for Building 330 Area	24
19	Summary of DCGLs for the Building 330 Area	24

TABLES (Cont.)

A.1	Parameter Values/Distributions Used in the Analysis for Different Scenarios	27
A.2	Parameter Distributions Used in the Analysis – Contaminated Zone and Unsaturated Zone K_d Values.....	36
A.3	Parameter Distributions Used in the Analysis – Saturated Zone K_d Values	37
A.4	Parameter Distributions Used in the Analysis for the Plant Transfer Factor.....	38
A.5	Parameter Distribution Used in the Analysis for the Meat Transfer Factor	39
A.6	Parameter Distribution Used in the Analysis for the Milk Transfer Factor.....	40
A.7	Parameter Value/Distribution Used in the Analysis for the Fish and Crustacea Transfer Factor	41
A.8	Notations Used in Input Parameter Table 1	42
A.9	Parameter Correlations for Probabilistic Analyses in Different Scenarios.....	43
A.10	Additional Input Parameters for RESRAD-OFFSITE Analysis.....	44
B.1	Parameter Values Used for Deterministic Analysis of Different Scenarios	52
B.2	Parameter Values Used in the Deterministic Analysis for K_d , Plant, Meat, and Milk Transfer Factors.....	60
B.3	Parameter Value/Distribution Used in the Analysis for the Fish and Crustacea Transfer Factor	61
B.4	Additional Input Parameters for RESRAD-OFFSITE Analysis.....	62
C.1	Peak DSR Percentiles for Different Exposure Pathways for Am-241 in Industrial Use Scenario	69
C.2	Peak DSR Percentiles for Different Exposure Pathways for Ba-133 in Industrial Use Scenario	70
C.3	Peak DSR Percentiles for Different Exposure Pathways for C-14 in Industrial Use Scenario	71
C.4	Peak DSR Percentiles for Different Exposure Pathways for Cs-137 in Industrial Use Scenario	72

TABLES (Cont.)

C.5	Peak DSR Percentiles for Different Exposure Pathways for Pu-238 in Industrial Use Scenario	73
C.6	Peak DSR Percentiles for Different Exposure Pathways for Pu-239 in Industrial Use Scenario	74
C.7	Peak DSR Percentiles for Different Exposure Pathways for Sr-90 in Industrial Use Scenario	75
C.8	Peak DSR Percentiles for Different Exposure Pathways for Tc-99 in Industrial Use Scenario	76
C.9	Peak DSR Percentiles for Different Exposure Pathways for Am-241 in Offsite Resident Scenario	77
C.10	Peak DSR Percentiles for Different Exposure Pathways for Ba-133 in Offsite Resident Scenario	78
C.11	Peak DSR Percentiles for Different Exposure Pathways for C-14 in Offsite Resident Scenario	79
C.12	Peak DSR Percentiles for Different Exposure Pathways for Cs-137 in Offsite Resident Scenario	80
C.13	Peak DSR Percentiles for Different Exposure Pathways for Pu-238 in Offsite Resident Scenario	81
C.14	Peak DSR Percentiles for Different Exposure Pathways for Pu-239 in Offsite Resident Scenario	82
C.15	Peak DSR Percentiles or Different Exposure Pathways for Sr-90 in Offsite Resident Scenario	83
C.16	Peak DSR Percentiles for Different Exposure Pathways for Tc-99 in Offsite Resident Scenario	84
C.17	Peak DSR Percentiles for Different Exposure Pathways for Am-241 in Recreational Use Scenario	85
C.18	Peak DSR Percentiles for Different Exposure Pathways for Ba-133 in Recreational Use Scenario	86

TABLES (Cont.)

C.19	Peak DSR Percentiles for Different Exposure Pathways for C-14 in Recreational Use Scenario	87
C.20	Peak DSR Percentiles for Different Exposure Pathways for Cs-137 in Recreational Use Scenario	88
C.21	Peak DSR Percentiles for Different Exposure Pathways for Pu-238 in Recreational Use Scenario	89
C.22	Peak DSR Percentiles for Different Exposure Pathways for Pu-239 in Recreational Use Scenario	90
C.23	Peak DSR Percentiles for Different Exposure Pathways for Sr-90 in Recreational Use Scenario	91
C.24	Peak DSR Percentiles for Different Exposure Pathways for Tc-99 in Recreational Use Scenario	92
C.25	Peak DSR Percentiles for Different Exposure Pathways for Am-241 in Onsite Resident Scenario	93
C.26	Peak DSR Percentiles for Different Exposure Pathways for Ba-133 in Onsite Resident Scenario	94
C.27	Peak DSR Percentiles for Different Exposure Pathways for C-14 in Onsite Resident Scenario	95
C.28	Peak DSR Percentiles for Different Exposure Pathways for Cs-137 in Onsite Resident Scenario	96
C.29	Peak DSR Percentiles for Different Exposure Pathways for Pu-238 in Onsite Resident Scenario	97
C.30	Peak DSR Percentiles for Different Exposure Pathways for Pu-239 in Onsite Resident Scenario	98
C.31	Peak DSR Percentiles for Different Exposure Pathways for Sr-90 in Onsite Resident Scenario	99
C.32	Peak DSR Percentiles for Different Exposure Pathways for Tc-99 in Onsite Resident Scenario	100

1 INTRODUCTION

The derived concentration guideline level (DCGL) is the allowable residual radionuclide concentration that can remain in soil after remediation of the site without radiological restrictions on the use of the site. It is sometimes called the single radionuclide soil guideline or the soil cleanup criteria. This report documents the methodology, scenarios, and parameters used in the analysis to support establishing radionuclide DCGLs for Argonne National Laboratory's Building 330 area.

1.1 SITE DESCRIPTION

Argonne National Laboratory (Argonne or ANL) is located 43 km southwest of downtown Chicago in DuPage County, Illinois. The site occupies 607 ha and is mostly surrounded by the Waterfall Glen Forest Preserve, which is used mainly as a recreational area. The Argonne site contains a number of small ponds and streams. The primary drainage on the site is through Sawmill Creek that runs through the site and enters the Des Plaines River. Some southernmost parts of the site drain directly into the Des Plaines River.

1.2 BUILDING HISTORY

Building 330, the former Chicago Pile 5 (CP-5) Reactor facility, was designed for neutron beam, neutron flux, gamma ray, and biological studies in addition to serving as a reactor training facility. The reactor became operational in 1954 and was shut down in 1979. The decommissioning of the reactor and supporting facilities started in 1991 and was completed by the U.S. Department of Energy (DOE) Office of Environmental Management (EM) in late 2000. A final release status survey under DOE Order 5400.5 was completed in 2000 by Argonne Health Physics (ANL-HP) with confirmatory surveys done by the Oak Ridge Institute for Science and Education (ORISE). The building was released for industrial reuse at that time, but still contained significant quantities of tritium in the foundations and soils under the existing concrete slabs. Building 330 was reclassified as below Hazard Category 3 (i.e., no longer considered a nuclear facility). Since it was shut down, the building remained vacant and had deteriorated over time. It was demolished in 2010, and the building's debris disposed of as low-level radioactive waste.

1.3 DERIVATION OF GUIDELINE LEVELS

The contaminants of concern identified for Building 330 are Am-241, Ba-133, C-14, Cs-137, Pu-238, Pu-239, Sr-90, and Tc-99 (Gerbig 2011; Matton 2011). The soil guidelines for these radionuclides are calculated using a dose constraint of 25 mrem/yr for all scenarios. It is assumed that the current industrial use of the site will continue with an institutional control period of 100 years. All land use scenarios are described in Section 2. The soil guidelines are derived for all scenarios, and the results are presented in Section 4. The final DCGLs will be determined by

DOE after considering other factors such as cost-benefit analysis, as low as reasonably achievable (ALARA) policy, and the applicability to other portions of Argonne site.

2 SCENARIO DESCRIPTION

2.1 LAND USE CONSIDERATIONS

The current land use for the site after it is remediated is industrial use with institutional control. However, after the institutional control period ends, the site could be rezoned for other purposes such as recreational use or residential use, or continued as an industrial use site. In developing DCGLs for the Building 330 area's residual soil radioactivity, four potential exposure scenarios are considered to cover all likely future land uses within a 1,000-year time frame. However, if the peak dose were to occur at a later time, the time frame for the analysis would be increased to 10,000 years. The four scenarios vary with respect to the type of land use and food source consumed. Two scenarios (industrial worker onsite and surface water use offsite) capture the dose to different receptors for the current land use conditions. Another two scenarios (recreational use and subsistence farming) capture the potential dose to different receptors after the institutional control period has ended. The industrial use scenario after the institutional control has ended will result in lower doses compared to the current industrial use because the peak dose in the time frame of 1–10,000 years is used and, due to radioactive decay of some short-lived radionuclides that exist onsite, the dose would be less after the institutional control period of 100 years. Therefore, the industrial scenario after the institutional control period has ended is not considered in this analysis. During the institutional control period, it is possible that contaminants may migrate to offsite locations by either surface runoff or wind erosion. Therefore, offsite residents may be exposed to onsite contamination during and after the institutional control period. This scenario is included in the DCGL analysis.

2.2 EXPOSURE SCENARIOS

There are 4 scenarios considered in the derivation of soil guidelines (DCGLs) for the Building 330 area. Two scenarios (Scenario A and B) are current use scenarios and another 2 scenarios (Scenario C and D) are future use scenarios. These 4 scenarios are described in detail in this section. The pathways and parameters considered in these 4 scenarios are discussed in Section 2.3.

Scenario A (the current use onsite worker scenario) assumes continued industrial use of the site. Under this scenario, it is assumed that a receptor would work onsite either indoors or outdoors (a probability distribution representative of an onsite worker will be assigned to the time fraction spent onsite). It is also assumed that the worker will not ingest contaminated water, plant, fish, meat, or milk obtained from the site. It was assumed that the dose to the worker was only from the residual contamination in the soil in the vicinity of Building 330.

Scenario B (the current use offsite resident scenario) assumes the use of surface water by a hypothetical offsite resident. Under this scenario, it is assumed that the contamination from surface runoff mixes with the storm water drainage and discharges to the Sawmill Creek. A hypothetical offsite resident uses this surface water for irrigation and feeding livestock. However, a public water supply is available in the area, and surface water is not used for

drinking purposes. The exposure pathways considered are ingestion of plant food grown with the irrigation water from the Sawmill Creek and ingestion of meat and milk from livestock that were fed with water from Sawmill Creek. The fish consumed by the resident are also caught from Sawmill Creek. It should be noted that currently very few people fish in Sawmill Creek or use the water for any purposes. The exposure to external radiation, inhalation of dust and radon, and incidental ingestion of soil are considered for this offsite resident scenario.

Scenario C (the likely future use scenario) assumes recreational use of the site. This scenario assumes that, after the institutional control period, the current forest preserve area bordering the site is expanded to include the Building 330 area. A hypothetical person (recreationist) camps on the remediated area for two weeks per year and is involved in recreational activities such as hiking, skiing, biking, and horseback riding. During this time period, it is assumed that all drinking water used by the hypothetical individual is drawn from the surface water (pond) that captures water from the surface runoff adjacent to the remediated area. Also the individual ingests fish taken from that pond. The site has clayey soil of low permeability and poor drainage; therefore, a pond could be easily built that can capture surface water from runoff. The recreationist does not ingest plant food growing in the area and also does not ingest milk from the livestock or wild animal raised in the area. However, part of the meat diet is from the livestock or wild animal raised in the area.

Scenario D (possible but unlikely future use scenario) assumes residential use of the site after the institutional control period. For this scenario, it is assumed that after the institutional control of the site, the industrial activities would end and the site is re-zoned for residential use. It is assumed that the site is used by a resident farmer. The resident farmer in this situation would be exposed to direct external radiation; internal radiation from inhalation of contaminated dust; internal radiation from inhalation of radon and its progeny; and internal radiation from ingestion of water, plant foods, meat, milk, and fish and incidental ingestion of soil. All water used by the resident farmer for drinking, irrigation, and household usage is drawn from a deep well adjacent to the remediated area. The individual ingests fish caught from a nearby pond. A shallow well scenario is not considered due to the site hydrogeological characteristics (Patton et al. 1990). Although a perched water table may exist onsite, the water yield of this groundwater system would be low and not capable of sustaining a resident farmer.

2.3 PATHWAYS AND KEY PARAMETERS

Potential radiation doses resulting from multiple exposure pathways are considered in this analysis for all exposure scenarios. These pathways include:

1. Direct exposure to external radiation from remediated soil material,
2. Internal radiation from inhalation of contaminated dust,
3. Internal radiation from inhalation of radon and its progeny,

4. Internal radiation from ingestion of plant foods grown in the remediated area and irrigated with an onsite water source,
5. Internal radiation from ingestion of meat from livestock raised onsite and fed with fodder grown in the remediated area and irrigated with water drawn from an onsite well or pond (the water ingested by livestock is also drawn from an onsite water source),
6. Internal radiation from ingestion of milk from livestock raised onsite and fed with fodder grown in the remediated area and irrigated with onsite well water or pond water (the water ingested by milk cows is also drawn from an onsite water source),
7. Internal radiation from ingestion of fish from a pond located downgradient from the decontaminated area,
8. Internal radiation from incidental ingestion of onsite soil,
9. Internal radiation from drinking water from an onsite water source (groundwater or surface water),
10. Internal radiation from ingestion of plant foods grown offsite and irrigated with contaminated surface water,
11. Internal radiation from ingestion of meat from livestock raised offsite and fed with fodder grown offsite but irrigated with contaminated creek water (the water ingested by livestock is also drawn from the creek),
12. Internal radiation from ingestion of milk from livestock raised offsite and fed with fodder grown offsite but irrigated with contaminated creek water (the water ingested by milk cow is also drawn from the creek), and
13. Ingestion of fish from an offsite surface water source.

Table 1 summarizes the applicable exposure pathways for all scenarios, and Table 2 lists the key parameters for the applicable exposure pathways.

TABLE 1 Summary of Applicable Exposure Pathways for Different Scenarios Considered for Argonne's Building 330 Area

Exposure Pathway	Applicable Pathways			
	Current Use		Future Use	
	Scenario A ^a	Scenario B ^b	Scenario C ^c	Scenario D ^d
Direct external gamma exposure	Yes	Yes	Yes	Yes
Inhalation of dust	Yes	Yes	Yes	Yes
Inhalation of radon and its progeny	Yes	Yes	Yes	Yes
Ingestion of soil	Yes	Yes	Yes	Yes
Ingestion of plant foods grown onsite	No	No	No	Yes
Ingestion of meat from livestock raised onsite	No	No	Yes	Yes
Ingestion of milk from livestock raised onsite	No	No	No	Yes
Ingestion of fish from an onsite pond	No	No	Yes	Yes
Ingestion of water from a downgradient well	No	No	No	Yes
Ingestion of water from a nearby surface water source	No	No	Yes	No
Ingestion of plant foods grown offsite	No	Yes	No	No
Ingestion of meat from livestock raised offsite	No	Yes	No	No
Ingestion of milk from livestock raised offsite	No	Yes	No	No
Ingestion of fish from surface water source offsite	No	Yes	No	No

^a Industrial worker (current use onsite worker scenario): no consumption of plant food, meat, milk, water, and fish.

^b Offsite resident (current use offsite resident scenario): the water used for irrigation and feeding livestock as well as the fish ingested are taken from an offsite surface water body.

^c Recreationist (likely future use scenario): no consumption of plant food and milk.

^d Onsite resident (possible but unlikely future use scenario): all onsite pathways are included.

TABLE 2 List of Key Parameters for Applicable Exposure Pathways

Exposure Pathway	Key Parameters
Direct external gamma exposure	Time fraction spent onsite and external gamma shielding factor
Inhalation of dust	Inhalation rate, time fraction spent onsite, mass loading for inhalation, and indoor dust filtration factor
Inhalation of radon and its progeny	Time fraction spent onsite, radon diffusion coefficient and emanation coefficient
Ingestion of soil	Soil ingestion rate and time spent onsite
Ingestion of plant foods grown onsite	Plant transfer factor, plant ingestion rate, water dilution factor, and release rate from the source
Ingestion of meat from livestock raised onsite	Plant transfer factor, meat transfer factor, meat ingestion rate, water dilution factor, and release rate from the source
Ingestion of milk from livestock raised onsite	Plant transfer factor, milk transfer factor, milk ingestion rate, water dilution factor, and release rate from the source
Ingestion of fish from a nearby pond	Fish bioaccumulation factor, aquatic food contaminated fraction, surface water dilution factor, and release rate from the source
Ingestion of water from a downgradient well	Water ingestion rate, infiltration rate, K_d values (or leach rate) for contaminants, and hydrogeological parameters for the site
Ingestion of water from a nearby surface water source	Water ingestion rate, surface water dilution factor, and release rate from the source
Ingestion of plant foods grown offsite	Plant transfer factor, plant ingestion rate, surface water dilution factor, and release rate from the source
Ingestion of meat from livestock raised offsite	Plant transfer factor, meat transfer factor, meat ingestion rate, surface water dilution factor, and release rate from the source
Ingestion of milk from livestock raised offsite	Plant transfer factor, milk transfer factor, milk ingestion rate, surface water dilution factor, and release rate from the source
Ingestion of fish from surface water source offsite	Fish bioaccumulation factor, aquatic food contaminated fraction, surface water dilution factor, and release rate from the source

3 DOSE/SOURCE CONCENTRATION RATIOS

Each scenario discussed in Section 2 was translated from its potential transport and environmental pathways into a specific set of parameter values. The RESRAD (onsite) and RESRAD-OFFSITE codes include many input parameters that can be classified as metabolic, behavioral, and physical parameters. The metabolic parameter represents a metabolic characteristic of the potential receptor, and its value is independent of the scenario but may be different for different population groups. Behavioral parameter values depend on the receptor's behavior in the scenario (e.g., parameter values for a recreationist scenario could be different from those for a subsistence farmer scenario). The physical parameters are source- and site-specific, and their values do not change for a different group of receptors. The analysis was done for an adult member of the population. For the behavioral parameters, scenario-specific, or mean, or median values were used. For physical parameters, site-specific values (or distributions), whenever available, were used. If site-specific values were not available, RESRAD (onsite) default values were used. For probabilistic analysis, distributions from the NUREG/CR-6697 report (NRC 2000) were used.

Radionuclide dose/source (concentration) ratios (DSR) were calculated from both probabilistic and deterministic analyses. Deterministic analysis uses a single value for each input parameter, resulting in a single dose output value. The probabilistic analysis uses parameter distributions to identify the variability and uncertainty in dose estimates resulting from variability and uncertainty in the input parameters. A probability distribution is specified for each input parameter with distribution. A model is run repeatedly, using different values for each input parameter with distribution for each run. Instead of giving a single model output as in a deterministic run, probabilistic analysis produces a set of outputs that are equal in number to the number of realizations selected (realizations are equal to the number of observations times number of repetitions).

The RESRAD (onsite) computer code, version 6.5 (Yu et al. 2001), was used for analyzing the industrial, recreational use, and onsite resident scenarios. RESRAD-OFFSITE computer code, version 2.6 (Yu et al. 2007), was used for analyzing the offsite resident scenario. The time frame considered in this analysis was 1,000 years. However, if the peak dose occurred at later time, the time frame for the analysis was increased to 10,000 years. Radioactive decay and ingrowth were considered in deriving dose/source concentration ratios. The various parameters used in the codes for probabilistic analysis are listed in the Appendix A. The various parameters used in the codes for deterministic analysis are listed in the Appendix B.

3.1 PROBABILISTIC ANALYSIS

The details of parameter selection are presented in Appendix A, where Table A.1 lists all input parameter distributions/values used in the probabilistic dose analysis for different scenarios. Some of the parameters used in the analysis were radionuclide/element-specific (e.g., K_d values, transfer factors) and required multiple values because the analysis was

conducted for multiple radionuclides; therefore, the radionuclide- and element-specific parameters used in the analysis are listed in separate tables.

Parameter distributions for K_d values are listed in Tables A.2 and A.3. Parameter distributions for plant transfer factors, meat transfer factors, milk transfer factors, and fish and crustacean transfer factors are listed in Tables A.3–A.7, respectively. The distribution function notations used in the RESRAD (onsite) input parameter table (Table A.1) are listed in Table A.8.

Some of the input parameters are correlated. In cases for which a clear relationship exists between parameters, strong correlations were used as input to ensure proper pairing. These include the pairs for bulk density and total porosity (correlation coefficient of -0.99), bulk density and effective porosity (correlation coefficient of -0.99), and total porosity and effective porosity (correlation coefficient of 0.99). Table A.9 lists the correlations used in the analysis. A few parameter values and distributions that are scenario-specific and distinguish the scenarios from one another (such as intake rates and time spent onsite) are specified for each scenario. For those parameters not specified with a single value, the default probabilistic distribution was used. This results in a dose probability distribution (presented as the cumulative distribution function, CDF). RESRAD-OFFSITE code requires many more parameters compared to the RESRAD (onsite) code; the additional input parameters for the RESRAD-OFFSITE analysis are listed in Table A.10.

Since the peak dose depends linearly on radionuclide concentration, the analysis was done for unit concentration of each radionuclide for four scenarios. For this analysis, 4,500 input sets (1,500 observations and 3 repetitions) were generated. For each set of sampled parameter values, peak dose in the time interval 0–1,000 years was calculated. Cumulative probabilities were estimated from the resulting dose distribution for each scenario.

Tables 3–10 provide total peak dose-to-source ratio (DSR) percentiles (mrem/yr per pCi/g) for different exposure scenarios. Also listed in Tables 3–10 are the mean of the peaks and peak of the means DSR. The mean of the peaks DSR is always greater than or equal to the peak of the means DSR. However, if the peak DSRs occurred at the same time for all input sets analyzed, then there would not be any difference in the mean of the peaks and peak of the means DSR. Peak DSR from 4,500 realizations were analyzed, and the percentiles for different exposure pathways are provided in Appendix C. Water-independent (WI) and water-dependent (WD) pathways are distinguished by the extensions WI and WD, respectively. Tables C.1–C.8 show the peak DSR percentiles for different exposure pathways for individual radionuclides in the industrial use scenario. For Ba-133, Cs-137, and Sr-90, external exposure was the dominant pathway. For Am-241 and Tc-99, external exposure and soil ingestion were the dominant pathways. For C-14, only inhalation was the dominant pathway. For Pu-238 and Pu-239, soil ingestion and inhalation were two dominant pathways.

Tables C.9–C.16 show the peak DSR percentiles for different exposure pathways for individual radionuclides in the offsite resident scenario. For all radionuclides except Ba-133, water-dependent pathways were dominant, and for Ba-133, the water-independent pathway was dominant. For Am-241, Pu-238, and Pu-239, fish and plant ingestion were dominant pathways. For Ba-133, the external exposure pathway was dominant. For C-14, fish ingestion resulted in

TABLE 3 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Am-241 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	1.47E-03	2.16E-03	1.81E-03	2.96E-02
10%	4.06E-03	2.69E-03	1.96E-03	3.61E-02
15%	6.76E-03	3.32E-03	2.05E-03	4.05E-02
20%	9.42E-03	4.17E-03	2.11E-03	4.48E-02
25%	1.11E-02	4.91E-03	2.16E-03	4.91E-02
30%	1.21E-02	5.62E-03	2.22E-03	5.33E-02
35%	1.27E-02	6.21E-03	2.26E-03	5.75E-02
40%	1.33E-02	6.66E-03	2.31E-03	6.29E-02
45%	1.37E-02	6.92E-03	2.36E-03	6.81E-02
50%	1.43E-02	7.12E-03	2.40E-03	7.46E-02
55%	1.47E-02	7.31E-03	2.44E-03	8.03E-02
60%	1.52E-02	7.49E-03	2.49E-03	8.78E-02
65%	1.57E-02	7.69E-03	2.54E-03	9.62E-02
70%	1.62E-02	7.95E-03	2.58E-03	1.07E-01
75%	1.68E-02	8.26E-03	2.64E-03	1.20E-01
80%	1.75E-02	8.66E-03	2.71E-03	1.37E-01
85%	1.84E-02	9.24E-03	2.79E-03	1.61E-01
90%	1.97E-02	1.00E-02	2.89E-03	1.96E-01
95%	2.19E-02	1.19E-02	3.05E-03	2.68E-01
Mean of the peaks	1.35E-02	6.98E-03	2.40E-03	1.05E-01
Peak of the means	1.35E-02	6.87E-03	2.40E-03	1.01E-01

most dose. For Cs-137, fish, meat, and milk ingestion resulted in most dose. For Sr-90, fish, meat, and plant ingestion resulted in most dose. For Tc-99, plant, fish, and milk ingestion resulted in most dose.

Tables C.17–C.24 show the peak DSR percentiles for different exposure pathways for individual radionuclides in the recreational use scenario. For C-14 and Tc-99, water-dependent pathways were dominant and for others (Am-241, Ba-133, Cs-137, Pu-238, Pu-239, and Sr-90), water-independent pathways were dominant. For Am-241, Pu-238, and Pu-239, meat ingestion, soil ingestion, and inhalation were dominant pathways. For Ba-133, external exposure was dominant. For C-14 and Tc-99, aquatic food ingestion resulted in most dose. For Cs-137, meat ingestion and external exposure resulted in most dose. For Sr-90, meat ingestion resulted in most dose.

Tables C.25–C.32 show the peak DSR percentiles for different exposure pathways for individual radionuclides in the onsite resident scenario. For C-14 and Tc-99, water-dependent pathways were dominant, and for others (Am-241, Ba-133, Cs-137, Pu-238, Pu-239, and Sr-90), water-independent pathways were dominant. For Am-241, plant ingestion, external exposure, and soil ingestion were dominant. For Pu-238 and Pu-239, plant ingestion, soil ingestion, and inhalation were dominant pathways. For Ba-133, external exposure was dominant. For C-14, aquatic food ingestion resulted in most dose. For Cs-137, external exposure and meat ingestion

TABLE 4 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Ba-133 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	3.96E-02	6.66E-05	2.31E-07	1.95E-06
10%	1.16E-01	7.01E-05	1.27E-05	9.88E-05
15%	1.93E-01	7.37E-05	4.04E-05	2.71E-04
20%	2.69E-01	7.64E-05	6.11E-05	3.68E-04
25%	3.40E-01	7.84E-05	7.58E-05	4.19E-04
30%	3.61E-01	8.00E-05	8.46E-05	4.60E-04
35%	3.74E-01	8.12E-05	9.04E-05	4.92E-04
40%	3.88E-01	8.22E-05	9.37E-05	5.27E-04
45%	4.01E-01	8.31E-05	9.62E-05	5.65E-04
50%	4.14E-01	8.41E-05	9.77E-05	6.05E-04
55%	4.24E-01	8.49E-05	9.88E-05	6.43E-04
60%	4.33E-01	8.58E-05	9.95E-05	6.79E-04
65%	4.41E-01	8.68E-05	1.00E-04	7.21E-04
70%	4.50E-01	8.78E-05	1.00E-04	7.74E-04
75%	4.60E-01	8.87E-05	1.01E-04	8.30E-04
80%	4.84E-01	8.99E-05	1.01E-04	9.02E-04
85%	5.09E-01	9.12E-05	1.01E-04	9.97E-04
90%	5.34E-01	9.30E-05	1.01E-04	1.11E-03
95%	5.70E-01	9.59E-05	1.01E-04	1.31E-03
Mean of the peaks	3.80E-01	8.32E-05	8.04E-05	6.32E-04
Peak of the means	3.80E-01	8.06E-05	8.04E-05	6.32E-04

resulted in most dose. For Sr-90, plant and meat ingestion resulted in most dose. For Tc-99, water and plant ingestion resulted in most dose.

Table 3 shows the total peak DSR percentiles of Am-241 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The future onsite resident (after the institutional control period of 100 years) would receive the highest dose.

Table 4 shows the total peak DSR percentiles of Ba-133 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The industrial worker (current use scenario) would receive the highest dose.

Table 5 shows the total peak DSR percentiles of C-14 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The current offsite resident (current use scenario) would receive the highest dose.

Table 6 shows the total peak DSR percentiles of Cs-137 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The industrial worker (current use scenario) would receive the highest dose.

TABLE 5 Total Peak DSR Percentiles (mrem/yr per pCi/g) of C-14 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	2.22E-06	2.48E-04	0.00E+00	0.00E+00
10%	5.82E-06	3.76E-04	0.00E+00	0.00E+00
15%	1.02E-05	4.97E-04	0.00E+00	0.00E+00
20%	1.34E-05	6.40E-04	0.00E+00	0.00E+00
25%	1.53E-05	8.00E-04	0.00E+00	0.00E+00
30%	1.68E-05	9.86E-04	0.00E+00	0.00E+00
35%	1.79E-05	1.21E-03	0.00E+00	0.00E+00
40%	1.90E-05	1.47E-03	0.00E+00	0.00E+00
45%	2.02E-05	1.85E-03	0.00E+00	0.00E+00
50%	2.13E-05	2.36E-03	0.00E+00	0.00E+00
55%	2.23E-05	2.96E-03	0.00E+00	0.00E+00
60%	2.34E-05	3.89E-03	0.00E+00	0.00E+00
65%	2.45E-05	5.35E-03	0.00E+00	0.00E+00
70%	2.56E-05	7.94E-03	1.77E-17	0.00E+00
75%	2.70E-05	1.27E-02	4.24E-03	0.00E+00
80%	2.87E-05	2.31E-02	1.55E-02	0.00E+00
85%	3.05E-05	4.19E-02	5.45E-02	1.01E-03
90%	3.32E-05	8.49E-02	1.17E-01	3.45E-03
95%	3.73E-05	2.27E-01	2.50E-01	1.05E-02
Mean of the peaks	2.10E-05	5.95E-02	4.61E-02	3.36E-03
Peak of the means	2.10E-05	1.50E-02	1.58E-02	3.54E-03

Table 7 shows the total peak DSR percentiles of Pu-238 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The future onsite resident (after the institutional control period of 100 years) would receive the highest dose.

Table 8 shows the total peak DSR percentiles of Pu-239 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The future onsite resident (after the institutional control period of 100 years) would receive the highest dose.

Table 9 shows the total peak DSR percentiles of Sr-90 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The future onsite resident (after the institutional control period of 100 years) would receive the highest dose.

Table 10 shows the total peak DSR percentiles of Tc-99 for different exposure scenarios considered in deriving DCGLs for the Building 330 area at Argonne. The future onsite resident (after the institutional control period of 100 years) would receive the highest dose.

Table 11 shows the peak of the mean, mean of the peak, and 95th percentile peak DSR of each radionuclide for different scenarios. These values were used in deriving DCGL values. The scenario that results in most dose for and individual radionuclide is highlighted.

TABLE 6 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Cs-137 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	7.12E-02	2.69E-03	1.27E-02	7.02E-02
10%	2.09E-01	3.10E-03	1.33E-02	7.93E-02
15%	3.47E-01	3.45E-03	1.37E-02	8.78E-02
20%	4.85E-01	3.76E-03	1.42E-02	9.41E-02
25%	6.22E-01	4.08E-03	1.46E-02	1.00E-01
30%	6.46E-01	4.39E-03	1.51E-02	1.07E-01
35%	6.70E-01	4.71E-03	1.55E-02	1.14E-01
40%	6.93E-01	5.05E-03	1.60E-02	1.21E-01
45%	7.17E-01	5.41E-03	1.65E-02	1.28E-01
50%	7.41E-01	5.76E-03	1.71E-02	1.36E-01
55%	7.56E-01	6.13E-03	1.77E-02	1.43E-01
60%	7.71E-01	6.54E-03	1.83E-02	1.52E-01
65%	7.87E-01	7.07E-03	1.92E-02	1.61E-01
70%	8.02E-01	7.62E-03	2.00E-02	1.71E-01
75%	8.18E-01	8.38E-03	2.11E-02	1.83E-01
80%	8.62E-01	9.14E-03	2.25E-02	1.99E-01
85%	9.06E-01	1.03E-02	2.45E-02	2.23E-01
90%	9.51E-01	1.18E-02	2.76E-02	2.56E-01
95%	1.01E+00	1.49E-02	3.49E-02	3.17E-01
Mean of the peaks	6.78E-01	6.87E-03	1.95E-02	1.59E-01
Peak of the means	6.78E-01	6.88E-03	1.95E-02	1.59E-01

3.2 DETERMINISTIC ANALYSIS

The details of parameter selection for the deterministic analysis are presented in Appendix B, where Table B.1 lists all input parameter values used in the deterministic dose analysis for different scenarios. Some of the parameters used in the analysis were radionuclide/element-specific (e.g., K_d values, transfer factors) and required multiple values because analysis was conducted for multiple radionuclides. The parameter values used for these radionuclide- and element-specific analyses are listed in separate tables.

Parameter values for K_d , plant transfer factors, meat transfer factors, and milk transfer factors are listed in Tables B.2. Parameter values for fish and crustacean transfer factors are listed in Table B.3. RESRAD-OFFSITE code requires many more parameters compared to RESRAD (onsite) code; the additional input parameters for RESRAD-OFFSITE analysis are listed in Table B.4.

Table 12 lists the calculated maximum dose/source concentration ratios (DSRs) (mrem/yr per pCi/g) for different exposure pathways for the industrial use scenario. RESRAD (onsite) version 6.5 was used in the analysis. For Ag-108m, Ba-133, Co-60, Cs-137, Eu-152, Eu-154, Eu-155, Np-237, Sr-90, U-235, and U-238, external exposure was the dominant exposure pathway. For Am-241 and Pu-241, external exposure, inhalation, and soil ingestion pathways all

TABLE 7 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Pu-238 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	6.12E-04	8.84E-04	6.07E-04	1.42E-02
10%	1.65E-03	1.18E-03	6.87E-04	1.74E-02
15%	2.51E-03	1.37E-03	7.47E-04	2.00E-02
20%	3.24E-03	1.51E-03	8.00E-04	2.22E-02
25%	3.85E-03	1.65E-03	8.37E-04	2.48E-02
30%	4.36E-03	1.75E-03	8.72E-04	2.75E-02
35%	4.81E-03	1.88E-03	9.09E-04	3.03E-02
40%	5.21E-03	2.02E-03	9.42E-04	3.33E-02
45%	5.56E-03	2.15E-03	9.83E-04	3.65E-02
50%	5.92E-03	2.29E-03	1.02E-03	4.05E-02
55%	6.27E-03	2.47E-03	1.06E-03	4.43E-02
60%	6.65E-03	2.65E-03	1.11E-03	4.90E-02
65%	7.07E-03	2.89E-03	1.15E-03	5.36E-02
70%	7.54E-03	3.17E-03	1.20E-03	5.97E-02
75%	7.97E-03	3.49E-03	1.25E-03	6.82E-02
80%	8.46E-03	3.91E-03	1.32E-03	7.85E-02
85%	9.12E-03	4.62E-03	1.40E-03	9.26E-02
90%	1.00E-02	5.69E-03	1.49E-03	1.13E-01
95%	1.13E-02	8.18E-03	1.61E-03	1.56E-01
Mean of the peaks	5.98E-03	3.13E-03	1.06E-03	5.71E-02
Peak of the means	5.98E-03	3.12E-03	1.06E-03	5.71E-02

contributed to the dose. For C-14, only inhalation was the dominant exposure pathway. For I-129 and Tc-99, soil ingestion and external exposure were dominant exposure pathways. For Ni-59 and Ni-63, soil ingestion was the dominant exposure pathway. For Pu-238, Pu-239, and Pu-240, inhalation and soil ingestion were two dominant exposure pathways. For U-234, radon inhalation at later time contributed significantly to the dose. For most radionuclides, the maximum dose/source ratio would occur immediately following remedial action. For Pu-241 and U-234, the maximum dose/source concentration ratio would occur at a later time due to buildup of progeny. To check if the peak dose would occur after 1,000 years, the time frame for the analysis was extended to 10,000 years. Table 12 also lists the time of maximum DSR for each radionuclide.

Table 13 lists the calculated maximum DSRs for different exposure pathways for the offsite resident scenario. RESRAD-OFFSITE version 2.6 was used for this analysis. For most radionuclides, water-dependent pathways were dominant. For Ba-133, the external exposure pathway was the dominant exposure pathway. For Am-241, Pu-238, and Pu-239, fish and plant ingestion were dominant exposure pathways. For C-14, aquatic food ingestion resulted in the most dose. For Cs-137, fish, meat, and milk ingestion resulted in the most dose. For Sr-90, fish, meat, milk, and plant ingestion resulted in the most dose. For Tc-99, plant, fish, and milk ingestion resulted in the most dose. To check if the peak dose would occur after 1,000 years, the

TABLE 8 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Pu-239 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	6.73E-04	1.04E-03	1.46E-03	3.39E-02
10%	1.81E-03	1.39E-03	1.66E-03	4.18E-02
15%	2.76E-03	1.64E-03	1.80E-03	4.76E-02
20%	3.57E-03	1.82E-03	1.92E-03	5.33E-02
25%	4.24E-03	1.97E-03	2.02E-03	5.98E-02
30%	4.79E-03	2.13E-03	2.10E-03	6.62E-02
35%	5.27E-03	2.29E-03	2.19E-03	7.26E-02
40%	5.72E-03	2.46E-03	2.27E-03	7.99E-02
45%	6.11E-03	2.60E-03	2.36E-03	8.80E-02
50%	6.51E-03	2.77E-03	2.45E-03	9.69E-02
55%	6.89E-03	2.96E-03	2.54E-03	1.07E-01
60%	7.30E-03	3.20E-03	2.65E-03	1.18E-01
65%	7.76E-03	3.44E-03	2.75E-03	1.30E-01
70%	8.27E-03	3.77E-03	2.87E-03	1.44E-01
75%	8.75E-03	4.15E-03	3.02E-03	1.64E-01
80%	9.28E-03	4.70E-03	3.18E-03	1.88E-01
85%	1.00E-02	5.39E-03	3.36E-03	2.23E-01
90%	1.10E-02	6.51E-03	3.56E-03	2.70E-01
95%	1.24E-02	9.05E-03	3.87E-03	3.73E-01
Mean of the peaks	6.56E-03	3.67E-03	2.54E-03	1.37E-01
Peak of the means	6.56E-03	3.55E-03	2.54E-03	1.37E-01

time frame for the analysis was extended to 10,000 years. Table 13 also lists the time of maximum DSR for each radionuclide.

Table 14 lists the maximum DSRs for different exposure pathways for the recreational use scenario. RESRAD (onsite) version 6.5 was used in the analysis. The calculations were done after the institutional control period of 100 years. Maximum DSR would occur at 100 years (immediately after the institutional control period) for all radionuclides. For C-14 and Tc-99, the maximum dose is from water-dependent pathways (such as food ingestion irrigated by contaminated water, water ingestion, and fish ingestion). To check if the peak dose would occur after 1,000 years, the time frame for the analysis was extended to 10,000 years. Table 14 also lists the time of maximum DSR for each radionuclide.

Table 15 lists the maximum DSRs for different exposure pathways for the onsite resident scenario. RESRAD (onsite) version 6.5 was used in the analysis. The calculations were done after the institutional control period of 100 years. Maximum DSRs would occur at 100 years (immediately after the institutional control period) for all radionuclides. For C-14 and Tc-99, the maximum dose is from water-dependent pathways (such as ingestion of food irrigated by contaminated water, water ingestion, and fish ingestion). To check if the peak dose would occur after 1,000 years, the time frame for the analysis was extended to 10,000 years. Table 15 also lists the time of maximum DSR for each radionuclide.

TABLE 9 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Sr-90 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	9.62E-04	8.35E-04	1.18E-04	4.42E-03
10%	2.84E-03	9.28E-04	9.18E-04	2.97E-02
15%	4.65E-03	1.01E-03	1.64E-03	5.30E-02
20%	6.54E-03	1.07E-03	2.31E-03	7.31E-02
25%	8.25E-03	1.14E-03	2.88E-03	9.77E-02
30%	8.72E-03	1.22E-03	3.51E-03	1.18E-01
35%	9.02E-03	1.28E-03	4.23E-03	1.44E-01
40%	9.35E-03	1.36E-03	4.93E-03	1.70E-01
45%	9.66E-03	1.43E-03	5.69E-03	2.04E-01
50%	9.94E-03	1.52E-03	6.56E-03	2.35E-01
55%	1.02E-02	1.60E-03	7.69E-03	2.79E-01
60%	1.04E-02	1.70E-03	9.24E-03	3.21E-01
65%	1.06E-02	1.82E-03	1.08E-02	3.72E-01
70%	1.09E-02	1.95E-03	1.28E-02	4.25E-01
75%	1.12E-02	2.13E-03	1.54E-02	5.03E-01
80%	1.16E-02	2.33E-03	1.86E-02	6.07E-01
85%	1.22E-02	2.62E-03	2.34E-02	7.50E-01
90%	1.29E-02	3.09E-03	3.09E-02	9.71E-01
95%	1.38E-02	3.99E-03	4.75E-02	1.44E+00
Mean of the peaks	9.15E-03	1.86E-03	1.35E-02	4.17E-01
Peak of the means	9.15E-03	1.70E-03	1.35E-02	4.17E-01

Table 16 shows the peak DSR of each radionuclide in different scenarios. These values were used in deriving DCGL values. The scenario that results in most dose for individual radionuclide is highlighted.

TABLE 10 Total Peak DSR Percentiles (mrem/yr per pCi/g) of Tc-99 for Different Exposure Scenarios for the Building 330 Area

Percentile	Industrial Use	Offsite Resident	Recreational Use	Onsite Resident
5%	3.44E-06	1.03E-04	5.78E-06	1.60E-05
10%	9.83E-06	2.50E-04	1.69E-05	4.02E-05
15%	1.62E-05	3.90E-04	3.34E-05	8.08E-05
20%	2.27E-05	5.63E-04	5.84E-05	1.58E-04
25%	2.65E-05	7.84E-04	9.76E-05	5.40E-04
30%	2.87E-05	1.02E-03	1.50E-04	1.68E-03
35%	3.04E-05	1.29E-03	2.16E-04	3.40E-03
40%	3.18E-05	1.53E-03	3.41E-04	5.71E-03
45%	3.30E-05	1.79E-03	4.72E-04	8.61E-03
50%	3.41E-05	2.03E-03	6.48E-04	1.29E-02
55%	3.54E-05	2.31E-03	8.73E-04	1.95E-02
60%	3.65E-05	2.59E-03	1.17E-03	2.79E-02
65%	3.77E-05	2.86E-03	1.50E-03	3.98E-02
70%	3.89E-05	3.18E-03	1.93E-03	5.62E-02
75%	4.02E-05	3.54E-03	2.49E-03	7.78E-02
80%	4.18E-05	4.02E-03	3.31E-03	1.03E-01
85%	4.38E-05	4.62E-03	4.36E-03	1.36E-01
90%	4.70E-05	5.50E-03	6.31E-03	1.83E-01
95%	5.18E-05	7.06E-03	1.07E-02	2.54E-01
Mean of the peaks	3.21E-05	2.59E-03	2.56E-03	5.68E-02
Peak of the means	3.21E-05	1.06E-03	9.62E-04	5.41E-02

TABLE 11 DSRs (mrem/yr per pCi/g) at Peak of the Means, Mean of the Peaks, and 95th Percentile of the Peaks for Different Scenarios

Nuclide	Industrial Use			Offsite Resident			Recreational Use			Onsite Resident		
	Peak of the Means	Mean of the Peaks	95th% of the Peaks	Peak of the Means	Mean of the Peaks	95th% of the Peaks	Peak of the Means	Mean of the Peaks	95th% of the Peaks	Peak of the Means	Mean of the Peaks	95th% of the Peaks
Am-241	1.35E-02	1.35E-02	2.19E-02	6.87E-03	6.98E-03	1.19E-02	2.40E-03	2.40E-03	3.05E-03	1.05E-01	1.01E-01	2.68E-01
Ba-133	3.80E-01	3.80E-01	5.70E-01	8.06E-05	8.32E-05	9.59E-05	8.04E-05	8.04E-05	1.01E-04	6.32E-04	6.32E-04	1.31E-03
C-14	2.10E-05	2.10E-05	3.73E-05	1.50E-02	5.95E-02	2.27E-01	1.58E-02	4.61E-02	2.50E-01	3.54E-03	3.36E-03	1.05E-02
Cs-137	6.78E-01	6.78E-01	1.01E+00	6.88E-03	6.87E-03	1.49E-02	1.95E-02	1.95E-02	3.49E-02	1.59E-01	1.59E-01	3.17E-01
Pu-238	5.98E-03	5.98E-03	1.13E-02	3.12E-03	3.13E-03	8.18E-03	1.06E-03	1.06E-03	1.61E-03	5.71E-02	5.71E-02	1.56E-01
Pu-239	6.56E-03	6.56E-03	1.24E-02	3.55E-03	3.67E-03	9.05E-03	2.54E-03	2.54E-03	3.87E-03	1.37E-01	1.37E-01	3.73E-01
Sr-90	9.15E-03	9.15E-03	1.38E-02	1.70E-03	1.86E-03	3.99E-03	1.35E-02	1.35E-02	4.75E-02	4.17E-01	4.17E-01	1.44E+00
Tc-99	3.21E-05	3.21E-05	5.18E-05	1.06E-03	2.59E-03	7.06E-03	9.62E-04	2.56E-03	1.07E-02	5.41E-02	5.68E-02	2.54E-01

TABLE 12 Maximum DSRs for the Industrial (Current Use) Scenario for the Building 330 Area

Nuclide	Maximum Dose/Source Concentration Ratio ^a (mrem/yr)/(pCi/g) for the Industrial Use Scenario				Time of Maximum DSR (yr)
	External Exposure	Inhalation	Radon	Ingestion of Soil	
Am-241	8.23E-03	6.60E-03	0.00E+00	6.16E-03	0.00E+00
Ba-133	3.81E-01	6.66E-07	0.00E+00	4.48E-05	0.00E+00
C-14	3.66E-07	1.78E-05	0.00E+00	2.65E-06	0.00E+00
Cs-137	6.76E-01	2.65E-06	0.00E+00	3.96E-04	0.00E+00
Pu-238	2.62E-05	7.54E-03	2.35E-13	7.06E-03	0.00E+00
Pu-239	5.74E-05	8.26E-03	0.00E+00	7.71E-03	0.00E+00
Sr-90	8.69E-03	1.10E-05	0.00E+00	9.38E-04	0.00E+00
Tc-99	2.23E-05	8.41E-07	0.00E+00	1.86E-05	0.00E+00

^a All values are reported to three significant figures. The maximum dose/source concentration ratio would occur at time zero (immediately following remedial action) for all radionuclides.

TABLE 13 Maximum DSRs for the Offsite Resident (Current Use) Scenario for the Building 330 Area

Nuclide	Maximum Dose/Source Concentration Ratio ^a (mrem/yr)/(pCi/g) for Offsite Resident Scenario								Time of Maximum DSR (yr)
	External Exposure	Inhalation	Radon	Plant Ingestion	Meat Ingestion	Milk Ingestion	Soil Ingestion	Fish Ingestion	
Am-241	1.09E-06	6.57E-06	0.00E+00	1.32E-03	1.05E-05	1.56E-06	9.18E-07	6.69E-03	5.86
Ba-133 ^b	7.85E-05	4.26E-10	0.00E+00	6.92E-06	2.19E-07	2.03E-06	5.58E-09	6.62E-06	6.84
C-14	3.98E-10	0.00E+00	0.00E+00	8.02E-05	4.52E-05	4.57E-05	3.54E-09	3.38E-02	152
Cs-137	2.20E-04	2.43E-09	0.00E+00	8.76E-05	4.07E-04	4.00E-04	3.74E-08	4.39E-03	3.91
Pu-238	2.79E-09	7.29E-06	1.99E-15	1.53E-03	2.44E-05	9.05E-07	8.70E-07	1.84E-03	4.88
Pu-239	1.10E-06	7.75E-06	0.00E+00	1.87E-03	3.66E-05	1.10E-06	1.74E-04	1.92E-03	1118
Sr-90	2.89E-06	9.73E-09	0.00E+00	2.74E-04	2.82E-04	2.43E-04	1.11E-07	3.94E-04	4.88
Tc-99	1.44E-07	2.02E-14	0.00E+00	2.70E-03	2.49E-05	7.19E-04	1.43E-07	5.74E-04	75.2

^a All values are reported to three significant figures.

^b For Ba-133, the dose is from both water-independent and water-dependent pathways.

TABLE 14 Maximum DSRs for the Recreational (Likely Future Land Use) Scenario for the Building 330 Area

Maximum Dose/Source Concentration Ratio ^a (mrem/yr)/(pCi/g) for Recreational Use Scenario								Time of Maximum DSR (yr)
Nuclide	External Exposure	Inhalation	Radon	Water Ingestion	Meat Ingestion	Soil Ingestion	Fish Ingestion	
Am-241	1.18E-03	1.16E-03	0.00E+00	0.00E+00	3.05E-04	8.81E-04	0.00E+00	1.00E+02
Ba-133	9.80E-05	2.10E-10	0.00E+00	0.00E+00	1.73E-08	1.15E-08	0.00E+00	1.00E+02
C-14 ^b	0.00E+00	0.00E+00	0.00E+00	4.24E-17	7.52E-16	0.00E+00	6.58E-13	1.00E+02
Cs-137	1.12E-02	5.38E-08	0.00E+00	0.00E+00	2.46E-03	6.55E-06	0.00E+00	1.00E+02
Pu-238	2.00E-06	7.05E-04	1.50E-13	0.00E+00	3.72E-04	5.38E-04	0.00E+00	1.00E+02
Pu-239	9.59E-06	1.70E-03	0.00E+00	0.00E+00	8.92E-04	1.29E-03	0.00E+00	1.00E+02
Sr-90	1.16E-04	1.80E-07	0.00E+00	0.00E+00	5.02E-03	1.26E-05	0.00E+00	1.00E+02
Tc-99 ^b	2.58E-12	1.20E-13	0.00E+00	3.05E-06	5.70E-09	2.15E-12	1.90E-05	1.00E+02

^a All values are reported to three significant figures. The maximum dose/source concentration ratio would occur at 100 years (immediately after the institutional control period) for all radionuclides.

^b For C-14 and Tc-99, the maximum dose is from water dependent pathways (such as ingestion of food irrigated by contaminated water, water ingestion, and fish ingestion).

TABLE 15 Maximum DSRs for the Onsite Resident (Unlikely Future Land Use) Scenario for the Building 330 Area

Maximum Dose/Source Concentration Ratio ^a (mrem/yr)/(pCi/g) for On-site Resident Scenario										Time of Maximum DSR (yr)
Nuclide	External Exposure	Inhalation	Radon	Water Ingestion	Plant Ingestion	Meat Ingestion	Milk Ingestion	Soil Ingestion	Fish Ingestion	
Am-241	1.72E-02	6.46E-03	0.00E+00	0.00E+00	5.47E-02	5.63E-04	3.21E-05	1.76E-02	0.00E+00	1.00E+02
Ba-133	1.43E-03	1.17E-09	0.00E+00	0.00E+00	3.57E-06	4.35E-08	1.46E-07	2.30E-07	0.00E+00	1.00E+02
C-14 ^b	0.00E+00	0.00E+00	0.00E+00	4.63E-14	6.49E-15	3.98E-15	6.24E-15	0.00E+00	3.45E-13	1.00E+02
Cs-137	1.64E-01	3.00E-07	0.00E+00	0.00E+00	1.63E-02	1.42E-02	4.65E-03	1.31E-04	0.00E+00	1.00E+02
Pu-238	2.92E-05	3.93E-03	1.75E-07	0.00E+00	3.34E-02	6.87E-04	1.04E-05	1.07E-02	0.00E+00	1.00E+02
Pu-239	1.40E-04	9.45E-03	0.00E+00	0.00E+00	8.00E-02	1.65E-03	2.35E-05	2.57E-02	0.00E+00	1.00E+02
Sr-90	1.70E-03	1.01E-06	0.00E+00	0.00E+00	2.34E-01	4.72E-02	1.40E-02	2.51E-04	0.00E+00	1.00E+02
Tc-99 ^b	3.78E-11	6.66E-13	0.00E+00	3.26E-03	2.89E-04	2.48E-06	6.42E-05	4.29E-11	8.77E-06	1.00E+02

^a All values are reported to three significant figures. The maximum dose/source concentration ratio would occur at 100 years (immediately after the institutional control period) for all radionuclides.

^b For C-14 and Tc-99, the maximum dose is from water dependent pathways (such as food ingestion irrigated by contaminated water, water ingestion, and fish ingestion).

TABLE 16 Total DSRs for Different Scenarios for the Building 330 Area

Total Dose/Source Concentration Ratio ^a (mrem/yr)/(pCi/g) for Different Scenarios				
Nuclide	Industrial Use ^b	Offsite Resident ^b	Onsite Resident ^c	Recreational Use ^d
Am-241	2.10E-02	8.03E-03	9.66E-02	3.52E-03
Ba-133	3.81E-01	9.44E-05	1.44E-03	9.80E-05
C-14	2.08E-05	3.40E-02	4.08E-13	6.59E-13
Cs-137	6.77E-01	5.51E-03	1.99E-01	1.37E-02
Pu-238	1.46E-02	3.41E-03	4.88E-02	1.62E-03
Pu-239	1.60E-02	4.02E-03	1.17E-01	3.89E-03
Sr-90	9.63E-03	1.20E-03	2.97E-01	5.14E-03
Tc-99	4.17E-05	4.02E-03	3.63E-03	2.21E-05

^a All values are reported to three significant figures.

^b Current use scenario.

^c Future unlikely use scenario.

^d Future likely use scenario.

4 DERIVED CONCENTRATION GUIDELINE LEVELS

The derived concentration guideline level (DCGL) is the concentration of residual radioactive material that can remain in the decontaminated area and still allow use of the area without radiological restrictions. Given a dose limit of H_{EL} for an individual, the derived concentration guideline level (DCGL) for an individual radionuclide (i) can be calculated as

$$DCGL_i \text{ (pCi/g)} = H_{EL} \text{ (mrem/yr)} / DSR_i \text{ (mrem/yr per pCi/g)}$$

where DSR_i is the total dose to source concentration ratio. The total DSRs for probabilistic analysis at peak of the means, mean of the peaks, and the 95th percentile of the peaks are listed in Table 11; for deterministic analysis, they are listed in Table 16 for different scenarios. The dose limit, H_{EL} , used in calculating DCGLs is 25 mrem/yr.

4.1 DCGLS BASED ON PROBABILISTIC ANALYSIS

Table 17 shows the DCGLs derived from different scenarios at mean and 95th percentile of the peak DSRs based on a 25 mrem/yr dose. For some radionuclides, the current land use scenarios (industrial worker or offsite resident scenarios) resulted in the most restrictive DCGLs, and for others, the future land use (onsite resident) scenario resulted in the most restrictive DCGLs. The most restrictive DCGLs are highlighted in Table 17 and listed in the last column under the heading “Most Restrictive” for peak of the means, mean of the peaks, and 95th percentile of the peak DSRs.

4.2 DCGLS BASED ON DETERMINISTIC ANALYSIS

Table 18 shows the DCGLs derived from a deterministic analysis of different scenarios based on a 25 mrem/yr dose limit. For some radionuclides, the current land use scenarios (industrial worker or offsite resident scenarios) resulted in the most restrictive DCGLs and for others, the future land use (onsite resident) scenario resulted in the most restrictive DCGLs. The most restrictive DCGL is highlighted in Table 18 and listed in the last column under the heading “Most Restrictive.”

4.3 SUMMARY DCGLS FOR BUILDING 330 AREA

Table 19 compares the DCGLs derived based on probabilistic analysis at peak of the means, mean of the peaks, 95th percentile of the peak DSRs, and the DCGLs derived from deterministic analysis for Building 330 area at Argonne. Also listed in the last column of Table 19 are the proposed DCGLs for the eight radionuclides analyzed. These proposed DCGLs are selected by considering both probabilistic and deterministic analyses results, and they are reported to two (2) significant figures.

TABLE 17 DCGLs (pCi/g) Using Probabilistic Analysis at a 25 mrem/yr Dose Limit for the Building 330 Area

Nuclide	Industrial Use			Offsite Resident			Recreational Use			Onsite Resident			Most Restrictive		
	Peak of the Means	Mean of the Peaks	95th of the Peaks	Peak of the Means	Mean of the Peaks	95th of the Peaks	Peak of the Means	Mean of the Peaks	95th of the Peaks	Peak of the Means	Mean of the Peaks	95th of the Peaks	Peak of the Means	Mean of the Peaks	95th of the Peaks
Am-241	1.85E+03	1.85E+03	1.14E+03	3.64E+04	3.58E+03	2.10E+03	1.04E+04	1.04E+04	8.20E+03	2.38E+02	2.48E+02	9.33E+01	2.38E+02	2.48E+02	9.33E+01
Ba-133	6.58E+01	6.58E+01	4.39E+01	3.10E+05	3.00E+05	2.61E+05	3.11E+05	3.11E+05	2.48E+05	3.96E+04	3.96E+04	1.91E+04	6.58E+01	6.58E+01	4.39E+01
C-14	1.19E+06	1.19E+06	6.70E+05	1.67E+03	4.20E+02	1.10E+02	1.58E+03	5.42E+02	1.00E+02	7.06E+03	1.05E+04	2.38E+03	1.58E+03	4.20E+02	1.00E+02
Cs-137	3.69E+01	3.69E+01	2.48E+01	3.63E+03	3.64E+03	1.68E+03	1.28E+03	1.28E+03	7.16E+02	1.57E+02	1.57E+02	7.89E+01	3.69E+01	3.69E+01	2.48E+01
Pu-238	4.18E+03	4.18E+03	2.21E+03	8.01E+03	7.99E+03	3.06E+03	2.36E+04	2.36E+04	1.55E+04	4.38E+02	4.38E+02	1.60E+02	4.38E+02	4.38E+02	1.60E+02
Pu-239	3.81E+03	3.81E+03	2.02E+03	7.04E+03	6.81E+03	2.76E+03	9.84E+03	9.84E+03	6.46E+03	1.82E+02	1.82E+02	6.70E+01	1.82E+02	1.82E+02	6.70E+01
Sr-90	2.73E+03	2.73E+03	1.81E+03	1.47E+04	1.34E+04	6.27E+03	1.85E+03	1.85E+03	5.26E+02	6.00E+01	6.00E+01	1.74E+01	6.00E+01	6.00E+01	1.74E+01
Tc-99	7.79E+05	7.79E+05	4.83E+05	2.36E+04	9.65E+03	3.54E+03	2.60E+04	9.77E+03	2.34E+03	4.62E+02	4.40E+02	9.84E+01	4.62E+02	4.40E+02	9.84E+01

TABLE 18 DCGLs Using Deterministic Analysis at the 25 mrem/yr Dose Limit for Building 330 Area

DCGLs (pCi/g) from Deterministic Analysis at the 25 mrem/yr Dose Limit from Different Scenarios					
Nuclide	Current use		Future Use		
	Industrial	Offsite Resident	Onsite Resident	Recreational	Most Restrictive
Am-241	1.19E+03	3.11E+03	2.59E+02	7.10E+03	2.59E+02
Ba-133	6.56E+01	2.65E+05	1.74E+04	2.55E+05	6.56E+01
C-14	1.20E+06	7.36E+02	NA ^a	NA ^a	7.36E+02
Cs-137	3.70E+01	4.54E+03	1.26E+02	1.83E+03	3.70E+01
Pu-238	1.71E+03	7.33E+03	5.13E+02	1.55E+04	5.13E+02
Pu-239	1.56E+03	6.23E+03	2.14E+02	6.44E+03	2.14E+02
Sr-90	2.60E+03	2.09E+04	8.42E+01	4.86E+03	8.42E+01
Tc-99	5.99E+05	6.22E+03	6.89E+03	1.13E+06	6.22E+03

^a DCGLs at specific activity limit.

TABLE 19 Summary of DCGLs for the Building 330 Area

Probabilistic DCGLs (pCi/g)					
Nuclide	At 95th Percentile of the Peak DSRs	At Mean of the Peak DSRs	At Peak of the Mean DSRs	Deterministic DCGLs (pCi/g)	Proposed ^a DCGLs (pCi/g)
Am-241	9.33E+01	2.38E+02	2.48E+02	2.59E+02	2.4E+02
Ba-133	4.39E+01	6.58E+01	6.58E+01	6.56E+01	6.5E+01
C-14	1.00E+02	4.20E+02	5.42E+02	7.36E+02	5.4E+02
Cs-137	2.48E+01	3.69E+01	3.69E+01	3.70E+01	3.6E+01
Pu-238	1.60E+02	4.38E+02	4.38E+02	5.13E+02	4.3E+02
Pu-239	6.70E+01	1.82E+02	1.82E+02	2.14E+02	1.8E+02
Sr-90	1.74E+01	6.00E+01	6.00E+01	8.42E+01	6.0E+01
Tc-99	9.84E+01	4.40E+02	4.62E+02	6.22E+03	4.6E+02

^a The proposed DCGLs are rounded to 2 significant figures.

The derived concentration guidelines listed in Table 19 are for a large homogenously contaminated area. When implementing the derived concentration guideline levels for decontamination of a site, the sum of fractions rule applies. That is, the summation of the radionuclide concentrations S_i remaining onsite for radioanuclides i divided by their $DCGL_i$ should not be greater than unity; that is,

$$\sum_i S_i / DCGL_i \leq 1 .$$

5 REFERENCES

Gerbig, C., 2011, *Characterization Sample & Analysis Plan E Wing Building 330 Argonne National Laboratory*, prepared by DeNuke Contracting Services, Inc., Oak Ridge, Tenn., revised March 13.

Matton, P.B., 2011, personal communication between P.B. Matton, Argonne National Laboratory, and C. Yu, Argonne National Laboratory, Subject: "Bldg. 330 Soil Characterization Final Report," March 15.

NRC (U.S. Nuclear Regulatory Commission), 2000, *Development of Probabilistic RESRAD 6.0 and RESRAD-BUILD 3.0 Computer Codes*, NUREG/CR-6697 and ANL/EAD/TM-98, prepared by C. Yu et al., Argonne National Laboratory, Argonne, Ill., for U.S. Nuclear Regulatory Commission, Washington, DC, Nov.

Patton, T.L., R.H. Pearl, and S.Y. Tsai, 1990, *Hydrological conditions at the 800 Area at Argonne National Laboratory*, ANL/EAIS/TM-29, Argonne National Laboratory, Argonne, Ill., August.

Yu, C., et al., 2007, *User's Manual for RESRAD-OFFSITE Version 2*, ANL/EVS/TM/07-1, Argonne National Laboratory, Argonne, Ill.

Yu, C., et al., 2001, *User's Manual for RESRAD Version 6*, ANL/EAD-4, Argonne National Laboratory, Argonne, Ill.

APPENDIX A:
PARAMETERS USED FOR PROBABILISTIC ANALYSIS

Tables A.1–A.10 list all input parameter distributions or parameter values used in the probabilistic dose analysis for different exposure scenarios.

TABLE A.1 Parameter Values/Distributions Used in the Analysis for Different Scenarios^a

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Title					
Title	Scenario dependent	Scenario dependent	Scenario dependent	Scenario dependent	Scenario definition
Dose factor library	ICRP-60	ICRP-60	ICRP-60	ICRP-60	ICRP-60 methodology based DCFs (external from FGR-12 [Eckerman 1993] and internal from ICRP-72 [ICRP 1996])
Cut-off half-life (180 d or 30 d)	30 d	30 d	30 d	30 d	RESRAD default
Number of points (32, 64, 128, 256, 512, 1,024)	32	32	32	32	RESRAD default
Linear spacing/log spacing	Log spacing	Log spacing	Log spacing	Log spacing	RESRAD default
Maximum no. of points for dose	17	17	17	17	RESRAD default
Maximum no. of points for risk	1	1	1	1	Using a smaller integration point will shorten the calculation time
Use line draw character (yes/no)	Yes	Yes	Yes	Yes	RESRAD default
Find peak pathway dose (yes/no)	Yes	Yes	Yes	Yes	
Save all files after each run (yes/no)	Yes	Yes	Yes	Yes	
Time integrated probabilistic risk (yes/no)	No	No	No	No	Dose-to-source ratio (DSR) is calculated
Calculation Parameters					
Basic radiation dose limit (mrem/year)	25	25	25	25	Not used in DSR calculation
Times for calculation (years)	1, 3, 10, 30, 100, 300, 1,000	1, 3, 10, 30, 100, 300, 1,000	100, 300, 1,000	100, 300, 1,000	Up to the time horizon for dose calculation
Source					
Nuclide concentration (pCi/g)	1	1	1	1	Dose-source ratios calculated
Transport Factors					
Distribution coefficient for all zones (cm ³ /g)	Tables A.2 and A.3	Tables A.2 and A.3	Tables A.2 and A.3	Table A.2 and A.3	Distribution from Sheppard and Thibault (1990)
Number of unsaturated zones	1	1	1	1	RESRAD default
Time since placement of material (y)	0	0	0	0	RESRAD default
Groundwater concentration (pCi/L)	0	0	0	0	RESRAD default
Leach rate (1/y)	0	0	0	0	RESRAD default

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Solubility limit (mol/L)	0	0	0	0	RESRAD default
Use plant/soil ration (check box)	No	No	No	No	RESRAD default
Transfer Factors					
Plant transfer factor (wet-plant weight concentration, pCi/g/dry soil weight concentration, pCi/g)	NR ^d	Table A.4	Table A.4	NR	Distribution from NUREG/CR-6697 (NRC 2000)
Meat transfer factor (concentration in meat, pCi/g/rate of intake, pCi/d)	NR	Table A.5	Table A.5	Table A.5	Distribution from NUREG/CR-6697 (NRC 2000)
Milk transfer factor (concentration in milk, pCi/L/rate of intake, pCi/d)	NR	Table A.6	Table A.6	NR	Distribution from NUREG/CR-6697 (NRC 2000)
Fish transfer factor (concentration in fish, pCi/kg/concentration in water, pCi/L)	NR	Table A.7	Table A.7	Table A.7	Distribution from NUREG/CR-6697 (NRC 2000)
Crustacea transfer factor (concentration in crustacea, pCi/kg/concentration in water, pCi/L)	NR	Table A.7	Table A.7	Table A.7	RESRAD default
Contaminated Zone Parameters					
Area of contaminated zone (m ²)	10,000	10,000	10,000	10,000	Kamboj 2011; Matton 2011
Thickness of contaminated zone (m)	1	1	1	1	Kamboj 2011; Matton 2011
Length parallel to aquifer flow (m)	100	100	100	100	RESRAD default
Does the initial contamination penetrate the water table?	No	No	No	No	RESRAD default
Cover and Contaminated Zone Hydrological Data					
Cover depth (m)	0	0	0	0	No cover layer assumed
Density of cover material (g/cm ³)	NR	NR	NR	NR	Not required when cover depth equals zero
Cover erosion rate (m/y)	NR	NR	NR	NR	Not required when cover depth equals zero
Density of primary contaminated zone (g/cm ³)	TN ^{e,f} (1.696, 0.1855, 0.001, 0.999)	TN ^{e,f} (1.696, 0.1855, 0.001, 0.999)	TN ^{e,f} (1.696, 0.1855, 0.001, 0.999)	TN ^{e,f} (1.696, 0.1855, 0.001, 0.999)	Site-specific soil type (silty clay – Golchert et al. 2010) distribution

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Contaminated zone erosion rate (m/y)	1.65E-4	1.65E-4 ^g	1.65E-4	1.65E-4	To match the soil erosion of 1 ton/acre/yr in Illinois (NRCS 2010)
Contaminated zone total porosity	TN ^f (0.36, 0.07, 0.001, 0.999)	TN ^f (0.36, 0.07, 0.001, 0.999)	TN ^f (0.36, 0.07, 0.001, 0.999)	TN ^f (0.36, 0.07, 0.001, 0.999)	Site-specific soil type (silty clay – Golchert et al. 2010) distribution
Contaminated zone field capacity	0.2	0.2	0.2	0.2	RESRAD default
Contaminated zone hydraulic conductivity (m/y)	0.32	0.32	0.32	0.32	Site-specific (Quinn 2010)
Contaminated zone b parameter	TLN ^f (2.29, 0.259, 0.001, 0.999)	TLN ^f (2.29, 0.259, 0.001, 0.999)	TLN ^f (2.29, 0.259, 0.001, 0.999)	TLN ^f (2.29, 0.259, 0.001, 0.999)	Site-specific soil type (silty clay – Golchert et al. 2010) distribution
Humidity in air (g/m ³)	8	8	8	8	Used only for H-3
Evapotranspiration coefficient	U(0.5, 0.75)	U(0.5, 0.75)	U(0.5, 0.75)	U(0.5, 0.75)	Distribution from NUREG/CR-6697 (NRC 2000)
Wind speed (m/s)	4.6	NR	4.6	4.6	Site-specific value for Chicago (NCDC 2010)
Precipitation rate (m/y)	0.9177	0.9177	0.9177	0.9177	Site-specific value from Golchert et al. 2010 ^b
Irrigation rate (m/y)	0.0	0.1	0.1	0.0	Site-specific (ANL 1996)
Irrigation mode (overhead/ditch)	NR	NR	Overhead	NR	RESRAD default
Runoff coefficient	0.2	0.2	0.2	0.2	Site-specific for woodland (Yu et al. 2001, page E-7)
Watershed area for nearby stream or pond (m ²)	1,000,000	NR	1,000,000	1,000,000	RESRAD default
Accuracy for water/soil computation	0.001	NR	0.001	0.001	RESRAD default
Saturated Zone Hydrological Data					
Density of saturated zone (g/cm ³)	NR	TN(1.52, 0.230, 0.001, 0.999)	TN(1.52, 0.230, 0.001, 0.999)	TN(1.52, 0.230, 0.001, 0.999)	Generic soil type distribution
Saturated zone effective porosity	NR	TN(0.355, 0.0906, 0.001, 0.999)	TN(0.355, 0.0906, 0.001, 0.999)	TN(0.355, 0.0906, 0.001, 0.999)	Generic soil type distribution
Saturated zone total porosity	NR	TN(0.425, 0.0867, 0.001, 0.999)	TN(0.425, 0.0867, 0.001, 0.999)	TN(0.425, 0.0867, 0.001, 0.999)	Generic soil type distribution
Saturated zone field capacity	NR	0.2	0.2	0.2	
Saturated zone hydraulic conductivity (m/y)	NR	947	947	947	Site-specific (ANL 1996)

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Saturated zone b parameter	NR	NR	BLN(1.06, 0.66, 0.5, 30)	BLN(1.06, 0.66, 0.5, 30)	Generic soil type distribution
Saturated zone hydraulic gradient	NR	U(0.007, 0.023)	U(0.007, 0.023)	U(0.007, 0.023)	Site-specific (Patton et al. 1990)
Water table drop rate (m/y)	NR	NR	0.001	0.001	RESRAD default
Well pump intake depth (m below water table)	NR	NR	5	5	Site-specific
Model for water transportation (nondispersion/mass-balance)	NR	NR	Nondispersion	Nondispersion	RESRAD default
Well pumping rate (m ³ /y)	NR	NR	250	250	RESRAD default
Unsaturated zone parameters					
Thickness of unsaturated zone (m)	NR	13.9	13.9	13.9	Site-specific (48 ft – 1 m) (Quinn 2010)
Density of unsaturated zone (g/cm ³)	NR	TN ^f (1.696, 0.1855, 0.001, 0.999)	TN ^f (1.696, 0.1855, 0.001, 0.999)	TN ^f (1.696, 0.1855, 0.001, 0.999)	Site-specific soil type (silty clay) distribution
Unsaturated zone effective porosity	NR	TN ^f (0.289, 0.0735, 0.001, 0.999)	TN ^f (0.289, 0.0735, 0.001, 0.999)	TN ^f (0.289, 0.0735, 0.001, 0.999)	Site-specific soil type (silty clay) distribution
Unsaturated zone total porosity	NR	TN ^f (0.36, 0.07, 0.001, 0.999)	TN ^f (0.36, 0.07, 0.001, 0.999)	TN ^f (0.36, 0.07, 0.001, 0.999)	Site-specific soil type (silty clay) distribution
Unsaturated zone field capacity	NR	0.2	0.2	0.2	RESRAD default
Unsaturated zone hydraulic conductivity (m/y)	NR	0.32	0.32	0.32	Site-specific (Quinn 2010)
Unsaturated zone b parameter	NR	TLN ^f (2.29, 0.259, 0.001, 0.999)	TLN ^f (2.29, 0.259, 0.001, 0.999)	TLN ^f (2.29, 0.259, 0.001, 0.999)	Site-specific soil type (silty clay) distribution
Occupancy, Inhalation, and External Gamma Parameters					
Inhalation rate (m ³ /y)	11,400	8400	8400	14,000	Scenario-specific
Mass loading for inhalation (g/m ³)	Empirical	Empirical	Empirical	Empirical	NRC 2000
Exposure duration (y)	25	30	30	30	Not used in DSR calculations
Indoor dust filtration factor	U(0.15, 0.95)	U(0.15, 0.95)	U(0.15, 0.95)	U(0.15, 0.95)	Distribution from NRC 2000
External gamma shielding factor	NR	BLN(-1.3, 0.59, 0.044, 1)	BLN(-1.3, 0.59, 0.044, 1)	NR	Distribution from NRC 2000

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Indoor time fraction	0/empirical	NR	0.6833	0	Scenario-specific (EPA exposure factor) for U-234 industrial scenario used empirical (because of radon inhalation)
Outdoor time fraction	Empirical/0	NR	0.0833	0.0384	Scenario-specific (EPA exposure factor)
Shape of contaminated zone (circular/noncircular)	Circular	Circular	Circular	Circular	RESRAD default
Ingestion Pathway Dietary Data					
Fruit, vegetable and grain consumption (kg/y)	NR	160	160	NR	Distribution from NRC 2000
Leafy vegetable consumption (kg/y)	NR	14	14	NR	Distribution from NRC 2000
Milk consumption (L/y)	NR	92	92	NR	Distribution from NRC 2000
Meat and poultry consumption (kg/y)	NR	63	63	T(7.0, 19.0, 61)	Scenario-specific
Fish consumption (kg/y)	NR	5.4	5.4	T(4.7, 12, 57)	Scenario-specific
Other sea food consumption (kg/y)	NR	0.9	0.9	NR	RESRAD default
Soil ingestion (g/y)	T(0, 18.3, 36.5)	T(0, 18.3, 36.5)	T(0, 18.3, 36.5)	T(0, 18.3, 36.5)	Distribution from NRC 2000
Drinking water intake (L/y)	NR	510	510	1.4L/d ^g 14 = 19.6 L	Distribution from NRC 2000 and scenario-specific for recreation use
Drinking water contaminated fraction	NR	0	1	1	RESRAD default and for offsite receptor scenario water is not contaminated
Household water contaminated fraction	1	1	1	NR	RESRAD default
Livestock water contaminated fraction	NR	1	1	1	RESRAD default
Irrigation water contaminated fraction	NR	1	1	1	RESRAD default
Aquatic food contaminated fraction	NR	0.5	0.5	0.5	Distribution from NRC 2000
Plant food contaminated fraction	NR	-1	-1	NR	Calculated by RESRAD from area factor
Meat contaminated fraction	NR	-1	-1	1	Scenario-specific
Milk contaminated fraction	NR	-1	-1	NR	Calculated by RESRAD from area factor

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Ingestion Pathway, Nondietary Data					
Livestock fodder intake for meat (kg/d)	NR	NR	68	10.6	RESRAD default or scenario-specific
Livestock fodder intake for milk (kg/d)	NR	NR	55	NR	RESRAD default
Livestock water intake for meat (L/d)	NR	50	50	6.4	RESRAD default or scenario-specific
Livestock water intake for milk (L/d)	NR	160	160	NR	RESRAD default
Livestock intake of soil for meat (kg/d)	NR	NR	0.5	0.5	RESRAD default
Livestock intake of soil for milk (kg/d)	NR	NR	0.5	NR	RESRAD default
Mass loading for foliar deposition (g/m ³)	NR	NR	0.0001	0.0001	RESRAD default
Depth of soil mixing layer (m)	T(0.0, 0.15, 0.6)	T(0.0, 0.15, 0.6)	T(0.0, 0.15, 0.6)	T(0.0, 0.15, 0.6)	Distribution from NRC 2000
Depth of roots (m)	NR	U(0.3, 1.0)	U(0.3, 1.0)	U(0.3, 1.0)	Upper bound equal to depth of contamination
Drinking water fraction from groundwater source	NR	0	1	0	Scenario-specific
Household water fraction from groundwater source	NR	0	1	NR	Scenario-specific
Livestock water fraction from groundwater source	NR	0	1	0	RESRAD default for resident farmer scenario and 0 for recreational use scenario because surface water is used by livestock
Irrigation water fraction from groundwater source	NR	0	1	0	RESRAD default
Plant Factors					
Wet weight crop yield for non-leafy vegetables (kg/m ²)	NR	TLN(0.56, 0.48, 0.001, 0.999)	TLN(0.56, 0.48, 0.001, 0.999)	NR	Distribution from NRC 2000
Length of growing season for non-leafy vegetables (y)	NR	0.17	0.17	NR	RESRAD default
Translocation factor for non-leafy vegetables	NR	0.1	0.1	NR	RESRAD default
Weathering removal constant (1/y)	NR	T(5.1, 18, 84)	T(5.1, 18, 84)	T(5.1, 18, 84)	Distribution from NRC 2000
Wet foliar interception fraction for non-leafy vegetables	NR	0.25	0.25	NR	RESRAD default

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Dry foliar interception fraction for non-leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Wet weight crop yield for leafy vegetables (kg/m ²)	NR	1.5	1.5	NR	RESRAD default
Length of growing season for leafy vegetables (y)	NR	0.25	0.25	NR	RESRAD default
Translocation factor for leafy vegetables	NR	1	1	NR	RESRAD default
Wet foliar interception fraction for leafy vegetables	NR	T(0.06, 0.67, 0.95)	T(0.06, 0.67, 0.95)	NR	Distribution from NRC 2000
Dry foliar interception fraction for leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Wet weight crop yield for fodder (kg/m ²)	NR	NR	1.1	1.1	RESRAD default
Length of growing season for fodder (y)	NR	NR	0.08	0.08	RESRAD default
Translocation factor for fodder	NR	NR	1	1	RESRAD default
Wet foliar interception fraction for fodder	NR	NR	0.25	0.25	RESRAD default
Dry foliar interception fraction for fodder	NR	NR	0.25	0.25	RESRAD default
Storage-Times-Before-Use Data					
Storage time for fruits, non-leafy vegetables and grain (d)	NR	14	14	NR	RESRAD default
Storage time for leafy vegetables (d)	NR	1	1	NR	RESRAD default
Storage time for milk (d)	NR	1	1	NR	RESRAD default
Storage time for meat (d)	NR	NR	20	182.5	RESRAD default for resident farmer and scenario-specific for recreational use
Storage time for fish (d)	NR	7	7	0	RESRAD default and scenario-specific for recreational use
Storage time for crustacea and mollusks (d)	NR	7	7	0	RESRAD default and scenario-specific for recreational use
Storage time for well water (d)	NR	1	1	NR	RESRAD default for resident farmer and not required for recreational use
Storage time for surface water (d)	NR	1	1	0	RESRAD default for resident farmer and 0 for recreational use

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Storage time for livestock fodder (d)	NR	NR	45	0	RESRAD default for resident farmer and scenario-specific for recreational use
Radon Data					
Cover total porosity	NR	NR	NR	NR	No cover
Cover volumetric water content	NR	NR	NR	NR	No cover
Cover radon diffusion coefficient	NR	NR	NR	NR	No cover
Total porosity of the house or building foundation	0.1	0.1	0.1	0.1	RESRAD default
Volumetric water content of the foundation	0.03	0.03	0.03	0.03	RESRAD default
Diffusion coefficient for radon gas in foundation material (m ² /s)	3E-07	3E-07	3E-07	3E-07	RESRAD default
Diffusion coefficient for radon gas in contaminated zone soil (m ² /s)	2E-06	2E-06	2E-06	2E-06	For site-specific soil type
Emanating power of radon-222	0.28	0.28	0.28	0.28	For site-specific soil type (Yu et al. 1993)
Radon vertical dimension of mixing (m)	2.0	2.0	2.0	2.0	RESRAD default
Average building air exchange rate (h ⁻¹)	0.5	0.5	0.5	0.5	RESRAD default
Height of building (room) (m)	2.5	2.5	2.5	2.5	RESRAD default
Building indoor area factor	0	0	0	0	RESRAD default
Bulk density of house or building foundation (g/cm ³)	2.4	2.4	2.4	2.4	RESRAD default
Thickness of house or building foundation (m)	0.15	0.15	0.15	0.15	RESRAD default
Building depth below ground surface (m)	-1	-1	-1	-1	RESRAD default
Carbon-14 Data					
C-12 concentration in local water (g/cm ³)	2E-5	2E-5	2E-5	2E-5	RESRAD default
C-12 concentration in contaminated soil (g/g)	0.03	0.03	0.03	0.03	RESRAD default
Fraction of vegetation carbon absorbed from soil	0.02	0.02	0.02	0.02	RESRAD default
Fraction of vegetation carbon absorbed from air	0.98	0.98	0.98	0.98	RESRAD default
Thickness of evasion layer of C-14 in soil (m)	0.3	0.3	0.3	0.3	RESRAD default
C-14 evasion flux rate from soil (sec ⁻¹)	7E-7	7E-7	7E-7	7E-7	RESRAD default

TABLE A.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
C-12 evasion flux rate from soil (sec ⁻¹)	1E-10	1E-10	1E-10	1E-10	RESRAD default
Grain fraction in beef cattle feed	0.8	0.8	0.8	0.8	RESRAD default
Grain fraction in milk cow feed	0.2	0.2	0.2	0.2	RESRAD default

^a RESRAD-OFFSITE code is used for analyzing offsite resident scenario, and RESRAD code is used for other scenarios. Extra parameters required for RESRAD-OFFSITE code are listed in Table A.10.

^b For offsite resident and residential farmer scenarios, the parameter values for leafy vegetable consumption, meat and poultry consumption, and fish consumption are RESRAD defaults. Parameter values for inhalation rate, indoor and outdoor time fractions, livestock fodder intake for meat, livestock water intake for meat, livestock intake of soil, plant food contaminated fraction, and meat contaminated fractions are RESRAD defaults and are calculated based on the contaminated area; storage time of meat, fish, surface water, and livestock fodder are RESRAD defaults.

^c For the recreational use scenario, it is assumed that the individual would be involved in moderate activity onsite; therefore, an inhalation rate of moderate activity (1.6 m³/h) is assumed (EPA 1997). The parameter value for meat and poultry consumption is based on the assumption that a recreationist kills a deer and brings the catch home and uses its meat over a year; the distribution is based on the game consumption statistics among households that hunt. For fish consumption, it is assumed that a recreationist catches a fish and consumes it the same day; distribution is based on the home-caught fish consumption statistics among households that fish. The parameter values for indoor and outdoor time fractions are from scenario description. All meat consumed is assumed to be contaminated. For the storage time of meat, since the meat is consumed over a year, an average storage time of 182.5 days is assumed. For the storage time of fish, since the fish is consumed on the same day, a storage time of zero days is assumed. Since the game animal uses fresh forage and surface water, a storage time of zero days is assumed for both parameters.

^d NR – value is not required for this scenario.

^e See Table A.8 for the notation used for the type of distribution.

^f It is assumed that the soil type is silty clay at the Building 330 area (Golchert et al. 2010) and distributions representative of silty clay soil are used.

^g Cover and management are changed to 0.0506 to get this erosion rate in RESRAD-OFFSITE code.

**TABLE A.2 Parameter Distributions
Used in the Analysis – Contaminated
Zone and Unsaturated Zone K_d
Values**

K_d (cm ³ /g) Distribution ^a			
Lognormal Distribution Parameters			
Element	μ	σ	$\exp(\mu)$
Ac	7.8	3.22 ^b	2,400
Am	9.0	2.6	8,400
Ba ^c	6.33	3.22	561
C	0.8	0.8 ^d	1
Cs	7.5	1.6	1,900
Np	4.0	3.8	55
Pa	7.9	3.22 ^b	2,700
Pb	6.3	2.76 ^b	550
Po	8.0	1.68 ^b	3,000
Pu	8.5	2.1	5,100
Ra	9.1	1.3	9,100
Sr	4.7	2.0	110
Tc	0.2	0.06	1
Th	8.6	2.6	5,800
U	7.3	2.9	1,600

Source: Sheppard and Thibault (1990)
unless noted otherwise.

^a Except for Ba, the distributions are for clayey soil, which is the site-specific soil type for the contaminated zone and unsaturated zone.

^b Value not available from Sheppard and Thibault (1990). The value used is default for generic soil type.

^c Distribution for clayey soil is not available for Ba; therefore, the distribution for generic soil is used.

^d Value not available from Sheppard and Thibault (1990). The value used is default for sandy soil type.

**TABLE A.3 Parameter Distributions
Used in the Analysis – Saturated
Zone K_d Values**

K_d (cm ³ /g) Distribution ^a			
Lognormal Distribution Parameters			
Element	μ	σ	$\exp(\mu)$
Ac	6.1	3.22 ^b	450
Am	7.6	2.6	1,900
Ba ^c	6.33	3.22	561
C	1.1	0.8	5
Cs	5.6	2.5	280
Np	1.4	1.7	5
Pa	6.3	3.22 ^b	550
Pb	5.6	2.76 ^b	270
Po	5.0	1.6	150
Pu	6.3	1.7	550
Ra	6.2	3.2	500
Sr	2.6	1.6	15
Tc	-2.0	1.8	0.1
Th	8.0	2.1	3,200
U	3.5	3.2	35

Source: Sheppard and Thibault (1990)
unless noted otherwise.

- ^a Except for Ba, the distributions are for sandy soil, which is assumed to be the soil type for the saturated zone.
- ^b Value not available from Sheppard and Thibault (1990). The value used is default for generic soil type.
- ^c Distribution for sandy soil is not available for Ba; therefore, the distribution for generic soil is used.

**TABLE A.4 Parameter Distributions Used
in the Analysis for the Plant Transfer
Factor**

Plant Transfer Factor (wet plant concentration, pCi/g/dry soil concentration, pCi/g) Distribution			
Lognormal Distribution Parameters			
Element	μ	σ	$\exp(\mu)$
Ac	-6.91	1.1	1.0×10^{-3}
Am	-6.91	0.9	1.0×10^{-3}
Ba	-4.61	1.1	1.0×10^{-2}
C	-0.36	0.9	7.0×10^{-1}
Cs	-3.22	1.0	4.0×10^{-2}
Np	-3.91	0.9	2.0×10^{-2}
Pa	-4.61	1.1	1.0×10^{-2}
Pb	-5.52	0.9	4.0×10^{-3}
Po	-6.9	0.9	1.0×10^{-3}
Pu	-6.91	0.9	1.0×10^{-3}
Ra	-3.22	0.9	4.0×10^{-2}
Sr	-1.20	1.0	3.0×10^{-1}
Tc	1.61	0.9	5.0×10^0
Th	-6.91	0.9	1.0×10^{-3}
U	-6.21	0.9	2.0×10^{-3}

Source: NRC 2000.

**TABLE A.5 Parameter Distribution
Used in the Analysis for the Meat
Transfer Factor**

Meat Transfer Factor (concentration in meat, pCi/g/rate of intake, pCi/d) Distribution			
Lognormal Distribution Parameters			
Element	μ	σ	$\exp(\mu)$
Ac	-6.91	1.1	1.0×10^{-3}
Am	-9.90	0.2	5.0×10^{-5}
Ba	-8.52	0.9	2.0×10^{-4}
C	-3.47	1	3.1×10^{-2}
Cs	-3.00	0.4	5.0×10^{-2}
Np	-6.91	0.7	1.0×10^{-3}
Pa	-12.21	1.0	5.0×10^{-6}
Pb	-7.13	0.7	8.0×10^{-4}
Po	-5.30	0.7	5.0×10^{-3}
Pu	-9.21	0.2	1.0×10^{-4}
Ra	-6.91	0.7	1.0×10^{-3}
Sr	-4.61	0.4	1.0×10^{-2}
Tc	-9.21	0.7	1.0×10^{-4}
Th	-9.21	1.0	1.0×10^{-4}
U	-7.13	0.7	8.0×10^{-4}

Source: NRC 2000.

**TABLE A.6 Parameter Distribution
Used in the Analysis for the Milk
Transfer Factor**

Milk Transfer Factor (concentration in milk, pCi/L/rate of intake, pCi/d) Distribution			
Lognormal Distribution Parameters			
Element	μ	σ	$\exp(\mu)$
Ac	-13.12	0.9	2.0×10^{-6}
Am	-13.12	0.7	2.0×10^{-6}
Ba	-7.6	0.7	5.0×10^{-4}
C	-4.4	0.9	1.2×10^{-2}
Cs	-4.61	0.5	1.0×10^{-2}
Np	-11.51	0.7	1.0×10^{-5}
Pa	-12.21	0.9	5.0×10^{-6}
Pb	-8.11	0.9	3.0×10^{-4}
Po	-7.82	0.7	4.0×10^{-4}
Pu	-13.82	0.5	1.0×10^{-6}
Ra	-6.91	0.5	1.0×10^{-3}
Sr	-6.21	0.5	2.0×10^{-3}
Tc	-6.91	0.7	1.0×10^{-3}
Th	-12.21	0.9	5.0×10^{-6}
U	-7.82	0.6	4.0×10^{-4}

Source: NRC 2000.

TABLE A.7 Parameter Value/Distribution Used in the Analysis for the Fish and Crustacea Transfer Factor

Element	Crustacea Transfer Factor (concentration in crustacea, pCi/Kg/ concentration in water, pCi/L)	Fish Transfer Factor (concentration in fish, pCi/Kg/concentration in water, pCi/L) Distributions		
		Lognormal Distribution Parameters		
		μ	σ	$\exp(\mu)$
Ac	1,000	2.7	1.1	15
Am	1,000	3.4	1.1	30
Ba	200	1.4	1.1	4
C	9,100	10.8	1.1	50,000
Cs	100	7.6	0.7	2,000
Np	400	3.4	1.1	30
Pa	110	2.3	1.1	10
Pb	100	5.7	1.1	300
Po	20,000	4.6	1.1	100
Pu	100	3.4	1.1	30
Ra	250	3.9	1.1	50
Sr	100	4.1	1.1	60
Tc	5	3.0	1.1	20
Th	500	4.6	1.1	100
U	60	2.3	1.1	10

Source: NRC 2000.

**TABLE A.8 Notations Used in Input
Parameter Table 1**

Notation	Type of Distribution
$TN(\mu, \sigma, a, b)^a$	Truncated Normal
$TLN(\mu, \sigma, a, b)^b$	Truncated Lognormal-N
$BLN(\mu, \sigma, a, b)^c$	Bounded Lognormal-N
$U(a, b)^d$	Uniform
$LU(a, b)^e$	Log Uniform
$T(a, c, b)^f$	Triangular
Continuous linear ^g	Empirical
Continuous log ^h	Empirical

^a For truncated normal distribution, μ is mean, σ is standard deviation, a is lower quantile, and b is upper quantile of the distribution.

^b For truncated lognormal-N distribution, μ is mean and σ is standard deviation of underlying normal distribution; and a is lower quantile, and b is upper quantile of the distribution.

^c For bounded lognormal-N distribution, μ is mean and σ is standard deviation of underlying normal distribution; and a is minimum, and b is maximum of the distribution.

^d For uniform distribution, a and b parameters represent the minimum and maximum values, respectively.

^e For log uniform distribution, a and b parameters represent the minimum and maximum values, respectively.

^f For triangular distribution, a is minimum, c is mode, and b is maximum of the distribution.

^g The continuous linear distribution is defined by the user for the N points, and the distribution is assumed to vary linearly between the cumulative distribution functions specified by the user for the N points.

^h The continuous log distribution is defined by the user for the N points, and the distribution is assumed to vary logarithmically between the cumulative distribution functions specified by the user for the N points.

TABLE A.9 Parameter Correlations for Probabilistic Analyses in Different Scenarios

Parameter 1	Parameter 2	Correlation Coefficient	Comments
Contaminated zone soil density	Contaminated zone total porosity	−0.99	The two parameters are strongly negatively correlated
Unsaturated zone soil density	Unsaturated zone total porosity	−0.99	The two parameters are strongly negatively correlated
Unsaturated zone soil density	Unsaturated zone effective porosity	−0.99	The two parameters are strongly negatively correlated
Saturated zone soil density	Saturated zone total porosity	−0.99	The two parameters are strongly negatively correlated
Saturated zone soil density	Saturated zone effective porosity	−0.99	The two parameters are strongly negatively correlated
Unsaturated zone total porosity	Unsaturated zone effective porosity	0.99	A correlation of 0.99 provides satisfactory pairing of sampling data
Saturated zone total porosity	Saturated zone effective porosity	0.99	A correlation of 0.99 provides satisfactory pairing of sampling data

TABLE A.10 Additional Input Parameters for RESRAD-OFFSITE Analysis

Parameter	Value	Comment
Site Layout^a		
X dimension of primary contamination (m)	100	Total contaminated area = 10,000 m ²
Y dimension of primary contamination (m)	100	
Smaller X coordinate of the fruit, grain, non-leafy vegetables plot (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the fruit, grain, non-leafy vegetables plot (m)	465.6	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller Y coordinate of the fruit, grain, non-leafy vegetables plot (m)	434	Closest distance to the site boundary
Larger Y coordinate of the fruit, grain, non-leafy vegetables plot (m)	466	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller X coordinate of the leafy vegetables plot (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the leafy vegetables plot (m)	465.6	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller Y coordinate of the leafy vegetables plot (m)	468	Leafy vegetable plot is assumed to be next to the nonleafy vegetable plot
Larger Y coordinate of the leafy vegetables plot (m)	500	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller X coordinate of the pasture, silage growing area (m)	450	Close to the vegetable plots
Larger X coordinate of the pasture, silage growing area (m)	550	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller Y coordinate of the pasture, silage growing area (m)	500	Close to the vegetable plots
Larger Y coordinate of the pasture, silage growing area (m)	600	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller X coordinate of the grain fields (m)	500	Close to the vegetable plots
Larger X coordinate of the grain fields (m)	600	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller Y coordinate of the grain fields (m)	600	Close to the vegetable plots
Larger Y coordinate of the grain fields (m)	700	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller X coordinate of the dwelling site (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the dwelling site (m)	465.6	The dwelling site is assumed to be 31.6 m × 31.6 m and have an area of 1,000 m ²
Smaller Y coordinate of the dwelling site (m)	400	Closest distance to the site boundary
Larger Y coordinate of the dwelling site (m)	432	The dwelling site is assumed to be 31.6 m × 31.6 m and have an area of 1,000 m ²
Smaller X coordinate of the surface water body	526.5	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Larger X coordinate of the surface water body	826.5	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas

TABLE A.10 (Cont.)

Parameter	Value	Comment
Smaller Y coordinate of the surface water body	356.8	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Larger Y coordinate of the surface water body	656.8	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Source Release and Deposition Velocity		
Deposition velocity (m/s)	0.001	For all radionuclides except I-129 (0.01)
Distribution Coefficients		
Sediment in surface water body (cm ³ /g)	Table A.2	
Fruit, grain, non-leafy vegetable fields (cm ³ /g)	Table A.2	
Leafy vegetable fields (cm ³ /g)	Table A.2	
Pasture, silage growing areas (cm ³ /g)	Table A.2	
Livestock feed grain fields (cm ³ /g)	Table A.2	
Offsite dwelling site (cm ³ /g)	Table A.2	
Transfer Factors		
Fruit, grain, nonleafy vegetable transfer factor	Table A.3	
Leafy vegetable transfer factor	Table A.3	
Pasture and silage transfer factor	Table A.3	
Livestock feed grain transfer factor	Table A.3	
Storage Time		
Storage time for pasture and silage (d)	1	Default
Storage time for livestock feed grain (d)	45	Default
Primary Contamination		
Deposition velocity of dust (m/s)	0.001	Default
Rainfall and runoff factor	160	Default
Slope-length-steepness factor	0.4	Default
Cover and management factor	0.0506	Adjusted to match site-specific erosion rate
Support practice factor	1	Default
Contaminated Zone		
Soil erodibility factor of contaminated zone	0.4	Default
Clean Cover		
Soil erodibility factor of clean cover	NR	No cover
Volumetric water content of clean cover	NR	No cover
Agriculture/Livestock Feed Growing/Offsite Dwelling Area Parameters		
Fraction of area directly over primary contamination for all fields	0	Receptor offsite

TABLE A.10 (Cont.)

Parameter	Value	Comment
Irrigation applied per year for all fields (m/y)	0.1	Site specific
Evapotranspiration coefficient for all fields	U(0.5, 0.75)	Distribution
Runoff coefficient for all fields	U(0.1, 0.8)	Distribution
Depth of soil mixing layer or plow layer for all fields (m)	See Table A.1	
Volumetric water content for all fields	0.3	Default
Dry bulk density of soil for all fields (g/cm ³)	TN ^d (1.696, 0.1855, 0.001, 0.999)	Site-specific soil type (silty clay – Golchert et al. 2010) distribution from NRC 2000
Soil erodibility factor for all fields	0.4	Default
Slope-length-steepness factor for all fields	0.4	Default
Cover and management factor for all fields	0.0506	Adjusted to match site-specific erosion rate
Support practice factor for all fields	1	Default
Atmospheric Transport		
Release height (m)	1	Default
Release heat flux (cal/s)	0	Default
Anemometer height (m)	10	Default
Ambient temperature (K)	285	Default
AM atmospheric mixing height (m)	400	Default
PM atmospheric mixing height (m)	1,600	Default
Dispersion model coefficients	Pasquill-Gifford	Default
Wind speed terrain	Rural	Default
Elevation of off-site location, relative to ground level at primary contamination, for all fields (m)	0	Default
Grid spacing for areal integration (m)	10	Default
Joint frequency of wind speed and stability class for a 16-sector wind rose	Actual values from Chicago Midway	Site-specific
Unsaturated Zone Parameters		
Unsaturated zone longitudinal dispersivity (m)	0.1	Default
Saturated Zone Hydrological Data		
Thickness of saturated zone (m)	18.3	Default
Saturated zone longitudinal dispersivity to surface water body (m)	10	Default
Saturated zone horizontal lateral dispersivity to surface water body (m)	1	Default
Disperse vertically	Check box	Yes
Saturated zone vertical lateral dispersivity to surface water body (m)	.06	Default
Depth of aquifer contributing to surface water body (m)	10	Default

TABLE A.10 (Cont.)

Parameter	Value	Comment
Surface Water Body		
Sediment deliver ratio	1	Default
Volume of surface water body (m ³)	150,000	Default
Mean residence time of water in surface water body (y)	1	Default
Groundwater Transport Parameters		
Distance from downgradient edge of contamination to well in the direction parallel to aquifer flow (m)	–55	Site-specific
Distance from downgradient edge of contamination to surface water body in the direction parallel to aquifer flow (m)	256.8	Site-specific
Distance from downgradient edge of contamination to well in the direction perpendicular to aquifer flow (m)	–424	Site-specific
Distance from downgradient edge of contamination to the right edge of surface water body in the direction perpendicular to aquifer flow (m)	–150	Default
Distance from downgradient edge of contamination to the left edge of the surface water body in the direction perpendicular to aquifer flow (m)	150	Default
Main subzones in saturated zone	1	Default
Main subzones in each partially saturated zone	1	Default
Nuclide-specific retardation in all subzones, longitudinal dispersion in all but the subzone of transformation?	Yes	Default
Water Use		
Quantity of water consumed by an individual (L/yr)	See Table A.1	Household water not contaminated
Number of household individuals consuming and using water	4	Default
Quantity of water for use indoors of dwelling per individual (L/d)	225	Household water not contaminated
Quantity of water for beef cattle (L/d)	50	Default
Number of beef cattle	2	Default
Quantity of water for dairy cows (L/d)	160	Default
Number of dairy cows	2	Default
Well pumping rate (m ³ /yr)	250	RESRAD default
Ingestion Rates		
Drinking water intake from affected area	0	Drinking water not contaminated
Fish intake from affected area	0.5	Default
Crustacea and mollusks intake from affected area	0.5	Default
Fruit, grain, non-leafy vegetables fraction from affected area	0.5	Default
Leafy vegetables fraction from affected area	0.5	Default
Meat fraction from affected area	0.5	50% of the meat consumed is from contaminated area

TABLE A.10 (Cont.)

Parameter	Value	Comment
Milk fraction from affected area	0.5	50% of the meat consumed is from contaminated area
Livestock intake		
Pasture and silage intake for beef cattle (kg/d)	14	Default
Grain intake for beef cattle (kg/d)	54	Default
Soil from pasture and silage intake for beef cattle (kg/d)	0.1	Default
Soil from grain intake for beef cattle (kg/d)	0.4	Default
Pasture and silage intake for dairy cows (kg/d)	44	Default
Grain intake for dairy cows (kg/d)	11	Default
Soil from pasture and silage intake for dairy cows (kg/d)	0.4	Default
Soil from grain intake for dairy cows (kg/d)	0.1	Default
Livestock Feed Factors (for Pasture and Silage, Grain)		
Wet weight crop yield (kg/m ²)	1.1, 0.7	Default
Duration of growing season (yr)	0.08, 0.17	Default
Foliage to food transfer coefficient	1, 0.1	Default
Weathering removal constant	20, 20	Default
Foliar interception factor for irrigation	0.25, 0.25	Default
Foliar interception factor for dust deposition	0.25, 0.25	Default
Root depth (m)	See Table A.1	Distribution from NUREG/CR-6697 (NRC 2000)
Occupancy Factors		
Indoor time fraction on primary contamination	0	
Outdoor time fraction on primary contamination	0	
Indoor time fraction on offsite dwelling site	0.5	
Outdoor time fraction on offsite dwelling site	0.1	
Time fraction in fruit, grain, and non-leafy vegetable fields	0.1	
Time fraction in leafy vegetable fields	0.1	
Time fraction in pasture and silage fields	0.1	
Time fraction in livestock grain fields	0.1	

^a All the areas (dwelling, agricultural areas) are assumed to be less than 350 m away from the contaminated area. This is a conservative assumption because the distance to the site boundary from the Building 330 area is greater than 350 m.

REFERENCES

- ANL (Argonne National Laboratory), 1996, *Interim Action Documentation Report: Decontamination and Demolition of the 317 Area South Vaults*, Argonne National Laboratory, Argonne Ill., May 31.
- Eckerman, K.F., and J.C. Ryman, 1993, External Exposure to Radionuclides in Air, Water, and Soil, Exposure to Dose Coefficients for General Application, Based on the 1987 Federal Radiation Protection Guidance, EPA 402-R-93-076, Federal Guidance Report No. 12, prepared by Oak Ridge National Laboratory, Oak Ridge, Tenn., for U.S. Environmental Protection Agency, Office of Radiation and Indoor Air, Washington, D.C.
- EPA (U.S. Environmental Protection Agency), 1997, *Exposure Factor Handbook*, EPA/600/P-95/002Fa, Office of Research and Development, National Center for Environmental Assessment, Washington, D.C.
- Golchert, N.W., T.M. Davis, and L.P. Moos, 2010, *Argonne Site Environmental Report for Calendar Year 2009*, ANL-10/02, Argonne National Laboratory, Argonne, Ill., September.
- ICRP (International Commission on Radiological Protection), 1991, *1990 Recommendations of the International Commission on Radiological Protection*, ICRP Publication 60, Ann. ICRP 21(1-3), Pergamon Press, Oxford, England.
- ICRP, 1996, Age-Dependent Doses to Members of the Public from Intake of Radionuclides: Part 5, Compilation of Ingestion and Inhalation Coefficients, ICRP Publication 72, Ann. ICRP 26(1), Pergamon Press, Oxford, England.
- Kamboj, S., 2011, personal communication between S. Kamboj, Argonne, and R.L. Johnson, Argonne, Subject: "Bldg. 310 RESRAD Analysis Project Status," January 7.
- Matton, P.B., 2011, personal communication between P.B. Matton, Argonne, and R.J. Johnson, Argonne, Subject: "Bldg. 330 D&D Project-RESRAD, DCGLs, Clean-up Parameters," January 14.
- NCDC (National Climatic Data Center), 2010, "Average Wind Speed." Available at <http://lwf.ncdc.noaa.gov/oa/climate/online/ccd/avgwind.html>.
- NRC (U.S. Nuclear Regulatory Commission), 2000, *Development of Probabilistic RESRAD 6.0 and RESRAD-BUILD 3.0 Computer Codes*, NUREG/CR-6697 and ANL/EAD/TM-98, prepared by C. Yu et al., Argonne National Laboratory, Argonne, Ill., for U.S. Nuclear Regulatory Commission, Washington, D.C., Nov.
- NRCS (Natural Resources Conservation Service), 2010, "Report Cards Are Out – Illinois NRI Gets Gold Star!," news release, May 25.

Patton, T.L., R.H. Pearl, and S.Y. Tsai, 1990, *Hydrological Conditions at the 800 Area at Argonne National Laboratory*, ANL/EAIS/TM-29, Argonne National Laboratory, Argonne, Ill., August.

Quinn, J., 2010, personal communication between J. Quinn, Argonne, and S. Kamboj, Argonne, November 12.

Sheppard, M.I. and D.H. Thibault, 1990, "Default Soil Solid/Liquid Partition Coefficients, Kds, for Four Major Soil Types: A Compendium," *Health Physics* 59(4):471–482.

Yu, C., et al., 1993, *Data Collection Handbook to Support Modeling the Impacts of Radioactive Material in Soil*, ANL/EAIS-8, Argonne National Laboratory, Argonne, Ill., April.

Yu, C., et al., 2001, *User's Manual for RESRAD Version 6*, ANL/EAD-4, Argonne National Laboratory, Argonne, Ill.

APPENDIX B:
PARAMETERS USED FOR DETERMINISTIC ANALYSIS

Tables B.1–B.4 list all input parameter distributions or parameter values used in the deterministic dose analysis for different user scenarios.

TABLE B.1 Parameter Values Used for Deterministic Analysis of Different Scenarios^a

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Title					
Title	Scenario dependent	Scenario dependent	Scenario dependent	Scenario dependent	Scenario definition
Dose factor library	ICRP-60	ICRP-60	ICRP-60	ICRP-60	ICRP-60 methodology based DCFs (external from FGR-12 and internal from ICRP 1996)
Cut-off half-life (180 d or 30 d)	30 d	30 d	30 d	30 d	RESRAD default
Number of points (32, 64, 128, 256, 512, 1024)	32	32	32	32	RESRAD default
Linear spacing/log spacing	Log Spacing	Log Spacing	Log Spacing	Log Spacing	RESRAD default
Maximum no. of points for dose	17	17	17	17	RESRAD default
Maximum no. of points for risk	1	1	1	1	Using a smaller integration point will shorten the calculation time
Use line draw character (yes/no)	Yes	Yes	Yes	Yes	RESRAD default
Find peak pathway dose (yes/no)	Yes	Yes	Yes	Yes	
Save all files after each run (yes/no)	Yes	Yes	Yes	Yes	
Time integrated probabilistic risk (yes/no)	No	No	No	No	Dose-to-source ratio (DSR) is calculated
Calculation Parameters					
Basic radiation dose limit (mrem/year)	25	25	25	25	Not used in DSR calculation
Times for calculation (years)	1, 3, 10, 30, 100, 300,1,000	1, 2, 3, 4, 5, 6, 7, 8, 970	100, 300, 1,000	100, 300, 1,000	Up to the time horizon for dose calculation
Source					
Nuclide concentration (pCi/g)	1	1	1	1	Dose-source ratios calculated
Transport Factors					
Distribution coefficient for all zones (cm ³ /g)	Table B.2	Table B.2	Table B.2	Table B.2	Site-specific soil type
Number of unsaturated zones	1	1	1	1	RESRAD default
Time since placement of material (y)	0	0	0	0	RESRAD default
Groundwater concentration (pCi/L)	0	0	0	0	RESRAD default
Leach rate (1/y)	0	0	0	0	RESRAD default
Solubility limit (mol/L)	0	0	0	0	RESRAD default
Use plant/soil ration (check box)	No	No	No	No	RESRAD default

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Transfer Factors					
Plant transfer factor (wet-plant weight concentration, pCi/g/dry soil weight concentration, pCi/g)	NR ^d	Table B.2	Table B.2	NR	RESRAD default
Meat transfer factor (concentration in meat, pCi/g/rate of intake, pCi/d)	NR	Table B.2	Table B.2	Table B.2	RESRAD default
Milk transfer factor (concentration in milk, pCi/L/rate of intake, pCi/d)	NR	Table B.2	Table B.2	NR	RESRAD default
Fish transfer factor (concentration in fish, pCi/Kg/concentration in water, pCi/L)	NR	Table B.3	Table B.3	Table B.3	RESRAD default
Crustacea transfer factor (concentration in crustacea, pCi/kg/concentration in water, pCi/L)	NR	Table B.3	Table B.3	Table B.3	RESRAD default
Contaminated Zone Parameters					
Area of contaminated zone (m ²)	10,000	10,000	10,000	10,000	Matton 2011; Kamboj 2011
Thickness of contaminated zone (m)	1	1	1	1	Matton 2011; Kamboj 2011
Length parallel to aquifer flow (m)	100	100	100	100	RESRAD default
Does the initial contamination penetrate the water table?	No	No	No	No	RESRAD default
Cover and Contaminated Zone Hydrological Data					
Cover depth (m)	0	0	0	0	No cover layer assumed
Density of cover material (g/cm ³)	NR	NR	NR	NR	Not required when cover depth equals zero
Cover erosion rate (m/y)	NR	NR	NR	NR	Not required when cover depth equals zero
Density of primary contaminated zone (g/cm ³)	1.76	1.76	1.76	1.76	Site-specific (ANL 1996)
Contaminated zone erosion rate (m/y)	1.65E-4	1.65E-4 ^e	1.65E-4	1.65E-4	To match the soil erosion of 1 ton/acre/yr at Illinois (NRCS 2010)
Contaminated zone total porosity	0.3	0.3	0.3	0.3	Site-specific (ANL 1996)
Contaminated zone field capacity	0.2	0.2	0.2	0.2	RESRAD default

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Contaminated zone hydraulic conductivity (m/y)	0.32	0.32	0.32	0.32	Site-specific (Quinn 2010)
Contaminated zone b parameter	10.4	10.4	10.4	10.4	Site-specific soil type (silty clay) value from Yu et al. 1993 ^f
Humidity in air (g/m ³)	8	8	8	8	Used only for H-3
Evapotranspiration coefficient	0.65	0.65	0.65	0.65	Site-specific
Wind speed (m/s)	4.6	NR	4.6	4.6	Site-specific value for Chicago
Precipitation rate (m/y)	0.9177	0.9177	0.9177	0.9177	Site-specific value from Golchert et al. 1010 ^b
Irrigation rate (m/y)	0.0	0.1	0.1	0.0	Site-specific (ANL 1996)
Irrigation mode (overhead/ditch)	NR	NR	Overhead	NR	RESRAD default
Runoff coefficient	0.2	0.2	0.2	0.2	Site-specific for woodland (Yu et al. 2001, page E-7)
Watershed area for nearby stream or pond (m ²)	1,000,000	NR	1,000,000	1,000,000	RESRAD default
Accuracy for water/soil computation	0.001	NR	0.001	0.001	RESRAD default
Saturated Zone Hydrological Data					
Density of saturated zone (g/cm ³)	NR	1.76	1.76	1.76	Site-specific (ANL 1996)
Saturated zone effective porosity	NR	0.14	0.14	0.14	Site-specific
Saturated zone total porosity	NR	0.3	0.3	0.3	Site-specific (ANL 1996)
Saturated zone field capacity	NR	0.14	0.14	0.14	Site-specific
Saturated zone hydraulic conductivity (m/y)	NR	947	947	947	Site-specific (ANL 1996)
Saturated zone b parameter	NR	4.38	4.38	4.38	Site-specific soil type
Saturated zone hydraulic gradient	NR	.015	.015	.015	Site-specific (Patton et al. 1990)
Water table drop rate (m/y)	NR	NR	0.001	0.001	RESRAD default
Well pump intake depth (m below water table)	NR	NR	5	5	Site-specific
Model for water transportation (nondispersion/mass-balance)	NR	NR	Nondispersion	Nondispersion	RESRAD default
Well pumping rate (m ³ /y)	NR	NR	250	250	RESRAD default
Unsaturated Zone Parameters					
Thickness of unsaturated zone (m)	NR	13.9	13.9	13.9	Site-specific (48 ft – 1 m) (Quinn 2010) or scenario-specific
Density of unsaturated zone (g/cm ³)	NR	1.76	1.76	1.76	Site-specific (ANL 1996)
Unsaturated zone effective porosity	NR	0.2	0.2	0.2	Site-specific (ANL 1996)

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Unsaturated zone total porosity	NR	0.3	0.3	0.3	Site-specific (ANL 1996)
Unsaturated zone field capacity	NR	0.2	0.2	0.2	RESRAD default
Unsaturated zone hydraulic conductivity (m/y)	NR	0.32	0.32	0.32	Site-specific (Quinn 2010)
Unsaturated zone b parameter	NR	10.4	10.4	10.4	Site-specific soil type (silty clay) value from Yu et al. 1993
Occupancy, Inhalation, and External Gamma Parameters					
Inhalation rate (m ³ /y)	11,400	8,400	8,400	14,000	Scenario-specific
Mass loading for inhalation (g/m ³)	1E-4	1E-4	1E-4	1E-4	RESRAD default
Exposure duration (y)	25	30	30	30	Not used in DSR calculations
Indoor dust filtration factor	0.4	0.4	0.4	0.4	RESRAD default
External gamma shielding factor	NR	0.7	0.7	NR	RESRAD default
Indoor time fraction	0/0.2283	NR	0.6833	0	Scenario-specific
Outdoor time fraction	0.2283/0	NR	0.0833	0.0384	Scenario-specific
Shape of contaminated zone (circular/noncircular)	Circular	Circular	Circular	Circular	RESRAD default
Ingestion Pathway Dietary Data					
Fruit, vegetable and grain consumption (kg/y)	NR	160	160	NR	RESRAD default
Leafy vegetable consumption (kg/y)	NR	14	14	NR	RESRAD default
Milk consumption (L/y)	NR	92	92	NR	RESRAD default
Meat and poultry consumption (kg/y)	NR	63	63	19	RESRAD default and scenario-specific for recreational use
Fish consumption (kg/y)	NR	5.4	5.4	12	RESRAD default and scenario-specific for recreational use
Other sea food consumption (kg/y)	NR	0.9	0.9	NR	RESRAD default
Soil ingestion (g/y)	36.5	36.5	36.5	36.5	RESRAD default
Drinking water intake (L/y)	NR	510	510	1.4L/d ^g 14 = 19.6 L	RESRAD default and scenario-specific for recreational use
Drinking water contaminated fraction	NR	0	1	1	RESRAD default and for offsite receptor scenario, water is not contaminated
Household water contaminated fraction	1	1	1	NR	RESRAD default

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Livestock water contaminated fraction	NR	1	1	1	RESRAD default
Irrigation water contaminated fraction	NR	1	1	1	RESRAD default
Aquatic food contaminated fraction	NR	0.5	0.5	0.5	Distribution from NRC 2000
Plant food contaminated fraction	NR	–1	–1	NR	Calculated by RESRAD from area factor
Meat contaminated fraction	NR	–1	–1	1	Scenario specific
Milk contaminated fraction	NR	–1	–1	NR	Calculated by RESRAD from area factor
Ingestion Pathway, Nondietary Data					
Livestock fodder intake for meat (kg/d)	NR	NR	68	10.6	RESRAD default or scenario-specific
Livestock fodder intake for milk (kg/d)	NR	NR	55	NR	RESRAD default
Livestock water intake for meat (L/d)	NR	50	50	6.4	Or scenario-specific
Livestock water intake for milk (L/d)	NR	160	160	NR	RESRAD default
Livestock intake of soil for meat (kg/d)	NR	NR	0.5	0.5	RESRAD default
Livestock intake of soil for milk (kg/d)	NR	NR	0.5	NR	RESRAD default
Mass loading for foliar deposition (g/m ³)	NR	NR	0.0001	0.0001	RESRAD default
Depth of soil mixing layer (m)	0.15	0.15	0.15	0.15	RESRAD default
Depth of roots (m)	NR	0.9	0.9	0.9	RESRAD default
Drinking water fraction from groundwater source	NR	0	1	0	Scenario-specific
Household water fraction from groundwater Source	NR	0	1	NR	Scenario-specific
Livestock water fraction from groundwater source	NR	0	1	0	RESRAD default for resident farmer scenario and 0 for recreational use scenario because surface water is used by livestock
Irrigation water fraction from groundwater source	NR	0	1	0	RESRAD default
Plant Factors					
Wet weight crop yield for non-leafy vegetables (kg/m ²)	NR	0.7	0.7	NR	RESRAD default

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Length of growing season for non-leafy vegetables (y)	NR	0.17	0.17	NR	RESRAD default
Translocation factor for non-leafy vegetables	NR	0.1	0.1	NR	RESRAD default
Weathering removal constant (1/y)	NR	20	20	20	Distribution from NRC 2000
Wet foliar interception fraction for non-leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Dry foliar interception fraction for non-leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Wet weight crop yield for leafy vegetables (kg/m ²)	NR	1.5	1.5	NR	RESRAD default
Length of growing season for leafy vegetables (y)	NR	0.25	0.25	NR	RESRAD default
Translocation factor for leafy vegetables	NR	1	1	NR	RESRAD default
Wet foliar interception fraction for leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Dry foliar interception fraction for leafy vegetables	NR	0.25	0.25	NR	RESRAD default
Wet weight crop yield for fodder (kg/m ²)	NR	NR	1.1	1.1	RESRAD default
Length of growing season for fodder (y)	NR	NR	0.08	0.08	RESRAD default
Translocation factor for fodder	NR	NR	1	1	RESRAD default
Wet foliar interception fraction for fodder	NR	NR	0.25	0.25	RESRAD default
Dry foliar interception fraction for fodder	NR	NR	0.25	0.25	RESRAD default
Storage-Times-Before-Use Data					
Storage time for fruits, non-leafy vegetables and grain (d)	NR	14	14	NR	RESRAD default
Storage time for leafy vegetables (d)	NR	1	1	NR	RESRAD default
Storage time for milk (d)	NR	1	1	NR	RESRAD default
Storage time for meat (d)	NR	NR	20	182.5	RESRAD default for resident farmer scenario and scenario-specific for recreational use
Storage time for fish (d)	NR	7	7	0	RESRAD default and scenario-specific for recreational use

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Storage time for crustacea and mollusks (d)	NR	7	7	0	RESRAD default and scenario-specific for recreational use
Storage time for well water (d)	NR	1	1	NR	RESRAD default for resident farmer scenario and not required for recreational use scenario
Storage time for surface water (d)	NR	1	1	0	RESRAD default for resident farmer scenario and 0 for recreational use scenario
Storage time for livestock fodder (d)	NR	NR	45	0	RESRAD default for resident farmer scenario and scenario-specific for recreational use scenario
Radon Data^g					
Cover total porosity	NR	NR	NR	NR	No cover
Cover volumetric water content	NR	NR	NR	NR	No cover
Cover radon diffusion coefficient	NR	NR	NR	NR	No cover
Total porosity of the house or building foundation	0.1	0.1	0.1	0.1	RESRAD default
Volumetric water content of the foundation	0.03	0.03	0.03	0.03	RESRAD default
Diffusion coefficient for radon gas in foundation material (m ² /s)	3E-07	3E-07	3E-07	3E-07	RESRAD default
Diffusion coefficient for radon gas in contaminated zone soil (m ² /s)	2E-06	2E-06	2E-06	2E-06	RESRAD default (the value is representative of silty clay soil type)
Emanating power of radon-222	0.28	0.28	0.28	0.28	Clay soil type (Yu et al. 1993)
Radon vertical dimension of mixing (m)	2.0	2.0	2.0	2.0	RESRAD default
Average building air exchange rate (h ⁻¹)	0.5	0.5	0.5	0.5	RESRAD default
Height of building (room) (m)	2.5	2.5	2.5	2.5	RESRAD default
Building indoor area factor	0	0	0	0	RESRAD default
Bulk density of house or building foundation (g/cm ³)	2.4	2.4	2.4	2.4	RESRAD default
Thickness of house or building foundation (m)	0.15	0.15	0.15	0.15	RESRAD default
Building depth below ground surface (m)	-1	-1	-1	-1	RESRAD default

TABLE B.1 (Cont.)

Input Parameters	Current Use Scenarios		Future Use Scenarios		Comments
	Industrial Worker	Offsite Resident ^b	Resident Farmer ^b	Recreationist ^c	
Carbon-14 Data^g					
C-12 concentration in local water (g/cm ³)	2E-5	2E-5	2E-5	2E-5	RESRAD default
C-12 concentration in contaminated soil (g/g)	0.03	0.03	0.03	0.03	RESRAD default
Fraction of vegetation carbon absorbed from soil	0.02	0.02	0.02	0.02	RESRAD default
Fraction of vegetation carbon absorbed from air	0.98	0.98	0.98	0.98	RESRAD default
Thickness of evasion layer of C-14 in soil (m)	0.3	0.3	0.3	0.3	RESRAD default
C-14 evasion flux rate from soil (sec ⁻¹)	7E-7	7E-7	7E-7	7E-7	RESRAD default
C-12 evasion flux rate from soil (sec ⁻¹)	1E-10	1E-10	1E-10	1E-10	RESRAD default
Grain fraction in beef cattle feed	0.8	0.8	0.8	0.8	RESRAD default
Grain fraction in milk cow feed	0.2	0.2	0.2	0.2	RESRAD default

^a RESRAD-OFFSITE code is used for analyzing offsite resident scenario, and RESRAD (onsite) code is used for other scenarios. Extra parameters required for RESRAD-OFFSITE code are listed in Table B-4.

^b For offsite resident and residential farmer scenarios, the parameter values for food consumption are RESRAD defaults. Parameter values for inhalation rate, indoor and outdoor time fractions, are scenario-specific. The parameter values for livestock fodder intake for meat, livestock water intake for meat, livestock intake of soil, plant food contaminated fraction, and meat contaminated fractions are code defaults and are calculated based on the contaminated area. The parameter values for storage time of meat, fish, surface water, and livestock fodder are RESRAD defaults.

^c For the recreational use scenario, it is assumed that the individual would be involved in moderate activity onsite; therefore, an inhalation rate of moderate activity (1.6 m³/h) is assumed (EPA 1997). The parameter value for meat and poultry consumption is based on the assumption that a recreationist kills a deer and brings the catch home and uses its meat over a year; the value is based on the game consumption statistics among households that hunt. For fish consumption, it is assumed that a recreationist catches a fish and consumes it the same day; value is based on the home-caught fish consumption statistics among households that fish. The parameter values for indoor and outdoor time fractions are from scenario description. All the meat consumed is assumed to be contaminated. For the storage time of meat, since the meat is consumed over a year, an average storage time of 182.5 days is assumed. For the storage time of fish, since the fish is consumed on the same day, a storage time of zero days is assumed. Since the game animal uses fresh forage and surface water, a storage time of zero days is assumed for both parameters.

^d NR – value is not required for this scenario.

^e Cover and management are changed to 0.0506 to get site-specific erosion rate in RESRAD-OFFSITE areas.

^f It is assumed that the soil type is silty clay at the Argonne site, and values representative of silty clay soil are used.

^g All the parameters for radon (except radon emanation rate) and C-14 are kept at code default values.

TABLE B.2 Parameter Values Used in the Deterministic Analysis for K_d , Plant, Meat, and Milk Transfer Factors

Element	K_d (cm ³ /g)		Plant Transfer Factor (pCi/kg)/(pCi/kg)	Meat Transfer Factor (pCi/kg)/(pCi/d)	Milk Transfer Factor (pCi/L)/(pCi/d)
	Saturated Zone ^a	Contaminated and Unsaturated Zone ^b			
Ac	450	2,400	2.5×10^{-3}	2.0×10^{-5}	2.0×10^{-5}
Am	1,900	8,400	1.0×10^{-3}	5.0×10^{-5}	2.0×10^{-6}
Ba	560	560	5.0×10^{-3}	2.0×10^{-4}	5.0×10^{-4}
C	5	1	7.0×10^{-1}	3.1×10^{-2}	1.2×10^{-2}
Cs	280	1,900	4.0×10^{-2}	3.0×10^{-2}	8.0×10^{-3}
Np	5	55	2.0×10^{-2}	1.0×10^{-3}	5.0×10^{-6}
Pa	550	2,700	1.0×10^{-2}	5.0×10^{-3}	5.0×10^{-6}
Pb	270	550	1.0×10^{-2}	8.0×10^{-4}	3.0×10^{-4}
Po	150	3,000	1.0×10^{-3}	5.0×10^{-3}	3.4×10^{-4}
Pu	550	5,100	1.0×10^{-3}	1.0×10^{-4}	1.0×10^{-6}
Ra	500	9,100	4.0×10^{-2}	1.0×10^{-3}	1.0×10^{-3}
Sr	15	110	3.0×10^{-1}	8.0×10^{-3}	2.0×10^{-3}
Tc	0.1	1	5.0×10^0	1.0×10^{-4}	1.0×10^{-3}
Th	3,200	5,800	1.0×10^{-3}	1.0×10^{-4}	5.0×10^{-6}
U	35	1,600	2.5×10^{-3}	3.4×10^{-4}	6.0×10^{-4}

^a Based on sandy soil type from RESRAD data collection handbook or RESRAD default values for saturated zone.

^b Based on clay soil type from RESRAD data collection handbook or RESRAD default values for contaminated and unsaturated zone.

TABLE B.3 Parameter Value/Distribution Used in the Analysis for the Fish and Crustacea Transfer Factor

Element	Crustacea Transfer Factor (concentration in crustacea, pCi/Kg/ concentration in water, pCi/L)	Fish Transfer Factor (concentration in crustacea, pCi/Kg/ concentration in water, pCi/L)
Ac	1,000	15
Am	1,000	30
Ba	200	4
C	9,100	50,000
Cs	100	2,000
Np	400	30
Pa	110	10
Pb	100	300
Po	20,000	100
Pu	100	30
Ra	250	50
Sr	100	60
Tc	5	20
Th	500	100
U	60	10

TABLE B.4 Additional Input Parameters for RESRAD-OFFSITE Analysis

Parameter	Value	Comment
Site Layout^a		
X dimension of primary contamination (m)	100	Total contaminated area = 10,000 m ²
Y dimension of primary contamination (m)	100	Total contaminated area = 10,000 m ²
Smaller X coordinate of the fruit, grain, non-leafy vegetables plot (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the fruit, grain, non-leafy vegetables plot (m)	466	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller Y coordinate of the fruit, grain, non-leafy vegetables plot (m)	434	Closest distance to the site boundary
Larger Y coordinate of the fruit, grain, non-leafy vegetables plot (m)	466	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller X coordinate of the leafy vegetables plot (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the leafy vegetables plot (m)	465.6	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller Y coordinate of the leafy vegetables plot (m)	468	Leafy vegetable plot is assumed to be next to the nonleafy vegetable plot
Larger Y coordinate of the leafy vegetables plot (m)	500	Assume the vegetable plot is 31.6 m × 31.6 m, with an area of 1,000 m ²
Smaller X coordinate of the pasture, silage growing area (m)	450	Close to the vegetable plots
Larger X coordinate of the pasture, silage growing area (m)	550	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller Y coordinate of the pasture, silage growing area (m)	500	Close to the vegetable plots
Larger Y coordinate of the pasture, silage growing area (m)	600	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller X coordinate of the grain fields (m)	500	Close to the vegetable plots.
Larger X coordinate of the grain fields (m)	600	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller Y coordinate of the grain fields (m)	600	Close to the vegetable plots
Larger Y coordinate of the grain fields (m)	700	Assume the livestock feed area is 100 m × 100 m, with an area of 10,000 m ²
Smaller X coordinate of the dwelling site (m)	434.4	Closest distance to the site boundary
Larger X coordinate of the dwelling site (m)	465.6	The dwelling site is assumed to be 31.6 m × 31.6 m and have an area of 1,000 m ²
Smaller Y coordinate of the dwelling site (m)	400	Closest distance to the site boundary
Larger Y coordinate of the dwelling site (m)	432	The dwelling site is assumed to be 31.6 m × 31.6 m and have an area of 1,000 m ²
Smaller X coordinate of the surface water body	526.5	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Larger X coordinate of the surface water body	826.5	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas

TABLE B.4 (Cont.)

Parameter	Value	Comment
Smaller Y coordinate of the surface water body	356.8	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Larger Y coordinate of the surface water body	656.8	The surface water body is assumed to be close to the dwelling site and the vegetable and livestock feed areas
Source Release and Deposition Velocity		
Deposition velocity (m/s)	0.001	For all radionuclides except I-129 (0.01)
Distribution Coefficients		
Sediment in surface water body (cm ³ /g)	Table B.2	
Fruit, grain, nonleafy vegetable fields (cm ³ /g)	Table B.2	
Leafy vegetable fields (cm ³ /g)	Table B.2	
Pasture, silage growing areas (cm ³ /g)	Table B.2	
Livestock feed grain fields (cm ³ /g)	Table B.2	
Off-site dwelling site (cm ³ /g)	Table B.2	
Transfer Factors		
Fruit, grain, non-leafy vegetable transfer factor	Table B.2	
Leafy vegetable transfer factor	Table B.2	
Pasture and silage transfer factor	Table B.2	
Livestock feed grain transfer factor	Table B.2	
Storage Time		
Storage time for pasture and silage (d)	1	Default
Storage time for livestock feed grain (d)	45	Default
Primary Contamination		
Deposition velocity of dust (m/s)	0.001	Default
Rainfall and runoff factor	160	Default
Slope-length-steepness factor	0.4	Default
Cover and management factor	0.0506	Adjusted to match site-specific erosion rate
Support practice factor	1	Default
Contaminated Zone		
Soil erodibility factor of contaminated zone	0.4	Default
Clean Cover		
Soil erodibility factor of clean cover	NR	No cover
Volumetric water content of clean cover	NR	No cover
Agriculture/Livestock Feed Growing/Offsite Dwelling Area Parameters		
Fraction of area directly over primary contamination for all fields	0	Receptor offsite
Irrigation applied per year for all fields (m/y)	0.1	Site specific

TABLE B.4 (Cont.)

Parameter	Value	Comment
Evapotranspiration coefficient for all fields	Table B.1	
Runoff coefficient for all fields	Table B.1	
Depth of soil mixing layer or plow layer for all fields (m)	Table B.1	
Volumetric water content for all fields	0.3	Default
Dry bulk density of soil for all fields (g/cm ³)	Table B.1	
Soil erodibility factor for all fields	0.4	Default
Slope-length-steepness factor for all fields	0.4	Default
Cover and management factor for all fields	0.0488	Adjusted to match site-specific erosion rate
Support practice factor for all fields	1	Default
Atmospheric Transport		
Release height (m)	1	Default
Release heat flux (cal/s)	0	Default
Anemometer height (m)	10	Default
Ambient temperature (K)	285	Default
AM atmospheric mixing height (m)	400	Default
PM atmospheric mixing height (m)	1,600	Default
Dispersion model coefficients	Pasquill-Gifford	Default
Wind speed terrain	Rural	Default
Elevation of offsite location relative to ground level at primary contamination for all fields (m)	0	Default
Grid spacing for areal integration (m)	10	Default
Joint frequency of wind speed and stability class for a 16-sector wind rose	Actual values from Chicago Midway	Site-specific
Unsaturated Zone Parameters		
Unsaturated zone longitudinal dispersivity (m)	0.1	Default
Saturated Zone Hydrological Data		
Thickness of saturated zone (m)	18.3	Site-specific
Saturated zone longitudinal dispersivity to surface water body (m)	10	Default
Saturated zone horizontal lateral dispersivity to surface water body (m)	1	Default
Disperse vertically	Check box	Yes
Saturated zone vertical lateral dispersivity to surface water body (m)	0.06	Default
Depth of aquifer contributing to surface water body (m)	10	Default
Surface Water Body		
Sediment deliver ratio	1	Default
Volume of surface water body (m ³)	150,000	Default

TABLE B.4 (Cont.)

Parameter	Value	Comment
Mean residence time of water in surface water body (y)	1	Default
Groundwater Transport Parameters		
Distance from downgradient edge of contamination to well in the direction parallel to aquifer flow (m)	–1	Site-specific
Distance from downgradient edge of contamination to surface water body in the direction parallel to aquifer flow (m)	256.8	Site-specific
Distance from downgradient edge of contamination to well in the direction perpendicular to aquifer flow (m)	–424	Site-specific
Distance from downgradient edge of contamination to the right edge of surface water body in the direction perpendicular to aquifer flow (m)	–150	Default
Distance from downgradient edge of contamination to the left edge of the surface water body in the direction perpendicular to aquifer flow (m)	150	Default
Main subzones in saturated zone	1	Default
Main subzones in each partially saturated zone	1	Default
Nuclide-specific retardation in all subzones, longitudinal dispersion in all but the subzone of transformation?	Yes	Default
Water Use		
Quantity of water consumed by an individual (L/yr)	Table B.1	Household water not contaminated
Number of household individuals consuming and using water	4	Default
Quantity of water for use indoors of dwelling per individual (L/d)	225	Household water not contaminated
Quantity of water for beef cattle (L/d)	50	Default
Number of beef cattle	2	Default
Quantity of water for dairy cows (L/d)	160	Default
Number of dairy cows	2	Default
Well pumping rate (m ³ /yr)	NR	RESRAD default
Ingestion Rates		
Drinking water intake from affected area	0	Drinking water not contaminated
Fish intake from affected area	0.5	Default
Crustacea and mollusks intake from affected area	0.5	Default
Fruit, grain, non-leafy vegetables fraction from affected area	0.5	Default
Leafy vegetables fraction from affected area	0.5	Default
Meat fraction from affected area	0.5	50% of the meat consumed is from contaminated area
Milk fraction from affected area	0.5	50% of the meat consumed is from contaminated area

TABLE B.4 (Cont.)

Parameter	Value	Comment
Livestock Intake		
Pasture and silage intake for beef cattle (kg/d)	14	Default
Grain intake for beef cattle (kg/d)	54	Default
Soil from pasture and silage intake for beef cattle (kg/d)	0.1	Default
Soil from grain intake for beef cattle (kg/d)	0.4	Default
Pasture and silage intake for dairy cows (kg/d)	44	Default
Grain intake for dairy cows (kg/d)	11	Default
Soil from pasture and silage intake for dairy cows (kg/d)	0.4	Default
Soil from grain intake for dairy cows (kg/d)	0.1	Default
Livestock Feed Factors (for Pasture and Silage, Grain)		
Wet weight crop yield (kg/m ²)	1.1, 0.7	Default
Duration of growing season (yr)	0.08, 0.17	Default
Foliage to food transfer coefficient	1, 0.1	Default
Weathering removal constant	20, 20	Default
Foliar interception factor for irrigation	0.25, 0.25	Default
Foliar interception factor for dust deposition	0.25, 0.25	Default
Root depth (m)	Table B.1	
Occupancy Factors		
Indoor time fraction on primary contamination	0	Default
Outdoor time fraction on primary contamination	0	Default
Indoor time fraction on offsite dwelling site	0.5	Default
Outdoor time fraction on offsite dwelling site	0.1	Default
Time fraction in fruit, grain, and non-leafy vegetable fields	0.1	Default
Time fraction in leafy vegetable fields	0.1	Default
Time fraction in pasture and silage fields	0.1	Default
Time fraction in livestock grain fields	0.1	Default

^a All the areas (dwelling, agricultural areas) are assumed to be less than 350 m away from the contaminated area. This is a conservative assumption because the distance to the site boundary from the Building 330 area is greater than 350 m.

REFERENCES

- ANL (Argonne National Laboratory), 1996, Interim Action Documentation Report: Decontamination and Demolition of the 317 Area South Vaults, Argonne National Laboratory, Argonne Ill., May 31.
- Eckerman, K.F., and J.C. Ryman, 1993, External Exposure to Radionuclides in Air, Water, and Soil, Exposure to Dose Coefficients for General Application, Based on the 1987 Federal Radiation Protection Guidance, EPA 402-R-93-076, Federal Guidance Report No. 12, prepared by Oak Ridge National Laboratory, Oak Ridge, Tenn., for U.S. Environmental Protection Agency, Office of Radiation and Indoor Air, Washington, D.C.
- EPA (U.S. Environmental Protection Agency), 1997, *Exposure Factor Handbook*, EPA/600/P-95/002Fa, Office of Research and Development, National Center for Environmental Assessment, Washington, D.C.
- Golchert, N.W., T.M. Davis, and L.P. Moos, 2010, *Argonne Site Environmental Report for Calendar Year 2009*, ANL-10/02, Argonne National Laboratory, Argonne, Ill., September.
- ICRP (International Commission on Radiological Protection), 1991, *1990 Recommendations of the International Commission on Radiological Protection*, ICRP Publication 60, Ann. ICRP 21(1-3), Pergamon Press, Oxford, England.
- ICRP, 1996, Age-Dependent Doses to Members of the Public from Intake of Radionuclides: Part 5, Compilation of Ingestion and Inhalation Coefficients, ICRP Publication 72, Ann. ICRP 26(1), Pergamon Press, Oxford, England.
- Kamboj, S., 2011, personal communication between S. Kamboj, Argonne, and R.L. Johnson, Argonne, Subject: "Bldg. 310 RESRAD Analysis Project Status," January 7.
- Matton, P.B., 2011, personal communication between P.B. Matton, Argonne, and R.J. Johnson, Argonne, Subject: "Bldg. 330 D&D Project-RESRAD, DCGLs, Clean-up Parameters," January 14.
- NRC (U.S. Nuclear Regulatory Commission), 2000, *Development of Probabilistic RESRAD 6.0 and RESRAD-BUILD 3.0 Computer Codes*, NUREG/CR-6697 and ANL/EAD/TM-98, prepared by C. Yu et al., Argonne National Laboratory, Argonne, Ill., for U.S. Nuclear Regulatory Commission, Washington, D.C., Nov.
- NRCS (Natural Resources Conservation Service), 2010, "Report Cards Are Out – Illinois NRI Gets Gold Star!," news release, May 25.
- Patton, T.L., R.H. Pearl, and S.Y. Tsai, 1990, *Hydrological Conditions at the 800 Area at Argonne National Laboratory*, ANL/EAIS/TM-29, Argonne National Laboratory, Argonne, Ill., August.

Quinn, J., 2010, personal communication between J. Quinn (Argonne) and S. Kamboj, Argonne, November 12.

Yu, C., et al., 1993, *Data Collection Handbook to Support Modeling the Impacts of Radioactive Material in Soil*, ANL/EAIS-8, Argonne National Laboratory, Argonne, Ill., April.

Yu, C., et al., 2001, *User's Manual for RESRAD Version 6*, ANL/EAD-4, Argonne National Laboratory, Argonne, Ill.

APPENDIX C:

DETAILED PROBABILISTIC ANALYSIS RESULTS

Tables C.1–C.8 show the peak DSR percentiles (mrem/yr per pCi/g) for different exposure pathways for the industrial use scenario. RESRAD (onsite) version 6.5 was used in the analysis. For Ba-133, Cs-137, and Sr-90, external exposure was the dominant exposure pathway. For Am-241 and Tc-99, soil ingestion and external exposure were the dominant exposure pathways. For C-14, only inhalation was the dominant exposure pathway. For Pu-238 and Pu-239 soil ingestion and inhalation were two dominant exposure pathways.

TABLE C.1 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Am-241 in Industrial Use Scenario

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	1.47E-03	8.66E-04	1.86E-04	2.76E-04
10%	4.06E-03	2.54E-03	4.46E-04	6.60E-04
15%	6.76E-03	4.23E-03	6.94E-04	1.00E-03
20%	9.42E-03	5.89E-03	9.35E-04	1.33E-03
25%	1.11E-02	7.57E-03	1.14E-03	1.65E-03
30%	1.21E-02	7.86E-03	1.30E-03	1.95E-03
35%	1.27E-02	8.15E-03	1.44E-03	2.27E-03
40%	1.33E-02	8.43E-03	1.57E-03	2.53E-03
45%	1.37E-02	8.72E-03	1.70E-03	2.78E-03
50%	1.43E-02	9.01E-03	1.82E-03	3.02E-03
55%	1.47E-02	9.20E-03	1.97E-03	3.26E-03
60%	1.52E-02	9.39E-03	2.14E-03	3.50E-03
65%	1.57E-02	9.58E-03	2.34E-03	3.73E-03
70%	1.62E-02	9.76E-03	2.53E-03	4.00E-03
75%	1.68E-02	9.95E-03	2.78E-03	4.31E-03
80%	1.75E-02	1.05E-02	3.07E-03	4.64E-03
85%	1.84E-02	1.10E-02	3.46E-03	5.04E-03
90%	1.97E-02	1.16E-02	3.99E-03	5.53E-03
95%	2.19E-02	1.23E-02	4.87E-03	6.27E-03
Mean	1.35E-02	8.26E-03	2.11E-03	3.09E-03
Min	1.08E-04	7.56E-05	1.14E-06	5.60E-06
Max	4.08E-02	1.71E-02	1.65E-02	1.15E-02

**TABLE C.2 Peak DSR Percentiles (mrem/yr per pCi/g) for
Different Exposure Pathways for Ba-133 in Industrial Use
Scenario**

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	3.96E-02	3.96E-02	1.87E-08	2.00E-06
10%	1.16E-01	1.16E-01	4.45E-08	4.74E-06
15%	1.93E-01	1.93E-01	6.96E-08	7.27E-06
20%	2.69E-01	2.69E-01	9.28E-08	9.57E-06
25%	3.40E-01	3.40E-01	1.14E-07	1.20E-05
30%	3.61E-01	3.61E-01	1.31E-07	1.41E-05
35%	3.74E-01	3.74E-01	1.45E-07	1.63E-05
40%	3.88E-01	3.88E-01	1.58E-07	1.82E-05
45%	4.01E-01	4.01E-01	1.71E-07	2.01E-05
50%	4.14E-01	4.14E-01	1.83E-07	2.18E-05
55%	4.24E-01	4.24E-01	1.98E-07	2.35E-05
60%	4.33E-01	4.33E-01	2.15E-07	2.53E-05
65%	4.41E-01	4.41E-01	2.34E-07	2.69E-05
70%	4.50E-01	4.50E-01	2.54E-07	2.89E-05
75%	4.60E-01	4.60E-01	2.80E-07	3.11E-05
80%	4.84E-01	4.84E-01	3.09E-07	3.35E-05
85%	5.09E-01	5.09E-01	3.47E-07	3.63E-05
90%	5.34E-01	5.34E-01	4.02E-07	4.00E-05
95%	5.70E-01	5.70E-01	4.90E-07	4.55E-05
Mean	3.80E-01	3.80E-01	2.12E-07	2.23E-05
Min	3.50E-03	3.50E-03	1.15E-10	4.07E-08
Max	7.91E-01	7.91E-01	1.66E-06	8.39E-05

**TABLE C.3 Peak DSR Percentiles (mrem/yr per pCi/g) for
Different Exposure Pathways for C-14 in Industrial Use
Scenario**

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	2.22E-06	3.91E-08	1.94E-06	1.19E-07
10%	5.82E-06	1.14E-07	5.28E-06	2.85E-07
15%	1.02E-05	1.89E-07	9.27E-06	4.35E-07
20%	1.34E-05	2.64E-07	1.20E-05	5.79E-07
25%	1.53E-05	3.37E-07	1.38E-05	7.13E-07
30%	1.68E-05	3.51E-07	1.50E-05	8.45E-07
35%	1.79E-05	3.65E-07	1.62E-05	9.74E-07
40%	1.90E-05	3.78E-07	1.72E-05	1.10E-06
45%	2.02E-05	3.90E-07	1.84E-05	1.20E-06
50%	2.13E-05	4.02E-07	1.93E-05	1.31E-06
55%	2.23E-05	4.12E-07	2.03E-05	1.41E-06
60%	2.34E-05	4.21E-07	2.14E-05	1.52E-06
65%	2.45E-05	4.28E-07	2.24E-05	1.61E-06
70%	2.56E-05	4.37E-07	2.36E-05	1.73E-06
75%	2.70E-05	4.47E-07	2.50E-05	1.86E-06
80%	2.87E-05	4.69E-07	2.64E-05	2.00E-06
85%	3.05E-05	4.93E-07	2.83E-05	2.18E-06
90%	3.32E-05	5.18E-07	3.11E-05	2.39E-06
95%	3.73E-05	5.54E-07	3.50E-05	2.71E-06
Mean	2.10E-05	3.69E-07	1.93E-05	1.34E-06
Min	1.45E-07	3.37E-09	1.35E-07	2.41E-09
Max	7.84E-05	7.68E-07	7.40E-05	5.03E-06

**TABLE C.4 Peak DSR Percentiles (mrem/yr per pCi/g) for
Different Exposure Pathways for Cs-137 in Industrial Use
Scenario**

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	7.12E-02	7.11E-02	7.46E-08	1.77E-05
10%	2.09E-01	2.08E-01	1.79E-07	4.24E-05
15%	3.47E-01	3.47E-01	2.79E-07	6.46E-05
20%	4.85E-01	4.85E-01	3.76E-07	8.58E-05
25%	6.22E-01	6.22E-01	4.58E-07	1.06E-04
30%	6.46E-01	6.46E-01	5.23E-07	1.26E-04
35%	6.70E-01	6.69E-01	5.77E-07	1.46E-04
40%	6.93E-01	6.93E-01	6.30E-07	1.63E-04
45%	7.17E-01	7.17E-01	6.84E-07	1.79E-04
50%	7.41E-01	7.40E-01	7.32E-07	1.94E-04
55%	7.56E-01	7.56E-01	7.90E-07	2.09E-04
60%	7.71E-01	7.71E-01	8.60E-07	2.25E-04
65%	7.87E-01	7.87E-01	9.37E-07	2.40E-04
70%	8.02E-01	8.02E-01	1.01E-06	2.58E-04
75%	8.18E-01	8.17E-01	1.12E-06	2.78E-04
80%	8.62E-01	8.62E-01	1.23E-06	2.98E-04
85%	9.06E-01	9.06E-01	1.39E-06	3.24E-04
90%	9.51E-01	9.51E-01	1.60E-06	3.56E-04
95%	1.01E+00	1.01E+00	1.96E-06	4.03E-04
Mean	6.78E-01	6.78E-01	8.46E-07	1.99E-04
Min	6.21E-03	6.20E-03	4.58E-10	3.60E-07
Max	1.40E+00	1.40E+00	6.61E-06	7.42E-04

TABLE C.5 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-238 in Industrial Use Scenario

Percentile	All Pathways	External	Inhalation w/o Rn	Radon (WI)	Soil Ingestion
5%	6.12E-04	2.76E-06	2.13E-04	1.01E-11	3.16E-04
10%	1.65E-03	8.08E-06	5.09E-04	2.50E-11	7.56E-04
15%	2.51E-03	1.35E-05	7.94E-04	4.50E-11	1.15E-03
20%	3.24E-03	1.88E-05	1.07E-03	7.21E-11	1.53E-03
25%	3.85E-03	2.41E-05	1.30E-03	9.43E-11	1.89E-03
30%	4.36E-03	2.50E-05	1.49E-03	1.19E-10	2.24E-03
35%	4.81E-03	2.60E-05	1.64E-03	1.41E-10	2.60E-03
40%	5.21E-03	2.69E-05	1.79E-03	1.62E-10	2.91E-03
45%	5.56E-03	2.78E-05	1.95E-03	1.81E-10	3.19E-03
50%	5.92E-03	2.87E-05	2.08E-03	2.03E-10	3.46E-03
55%	6.27E-03	2.93E-05	2.25E-03	2.24E-10	3.73E-03
60%	6.65E-03	2.99E-05	2.45E-03	2.48E-10	4.02E-03
65%	7.07E-03	3.05E-05	2.67E-03	2.74E-10	4.27E-03
70%	7.54E-03	3.11E-05	2.89E-03	3.05E-10	4.58E-03
75%	7.97E-03	3.17E-05	3.18E-03	3.37E-10	4.95E-03
80%	8.46E-03	3.34E-05	3.52E-03	3.76E-10	5.32E-03
85%	9.12E-03	3.52E-05	3.95E-03	4.23E-10	5.77E-03
90%	1.00E-02	3.69E-05	4.57E-03	4.90E-10	6.33E-03
95%	1.13E-02	3.93E-05	5.57E-03	6.08E-10	7.19E-03
Mean	5.98E-03	2.63E-05	2.41E-03	2.41E-10	3.54E-03
Min	2.74E-05	2.41E-07	1.30E-06	9.48E-14	6.42E-06
Max	2.76E-02	5.44E-05	1.88E-02	2.03E-09	1.32E-02

**TABLE C.6 Peak DSR Percentiles (mrem/yr per pCi/g) for
Different Exposure Pathways for Pu-239 in Industrial Use
Scenario**

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	6.73E-04	6.04E-06	2.33E-04	3.45E-04
10%	1.81E-03	1.77E-05	5.58E-04	8.25E-04
15%	2.76E-03	2.94E-05	8.69E-04	1.26E-03
20%	3.57E-03	4.12E-05	1.17E-03	1.67E-03
25%	4.24E-03	5.28E-05	1.43E-03	2.06E-03
30%	4.79E-03	5.48E-05	1.63E-03	2.44E-03
35%	5.27E-03	5.68E-05	1.80E-03	2.84E-03
40%	5.72E-03	5.88E-05	1.97E-03	3.17E-03
45%	6.11E-03	6.08E-05	2.13E-03	3.48E-03
50%	6.51E-03	6.28E-05	2.28E-03	3.78E-03
55%	6.89E-03	6.41E-05	2.46E-03	4.07E-03
60%	7.30E-03	6.54E-05	2.68E-03	4.39E-03
65%	7.76E-03	6.67E-05	2.92E-03	4.66E-03
70%	8.27E-03	6.81E-05	3.16E-03	5.01E-03
75%	8.75E-03	6.94E-05	3.49E-03	5.40E-03
80%	9.28E-03	7.31E-05	3.85E-03	5.81E-03
85%	1.00E-02	7.69E-05	4.32E-03	6.30E-03
90%	1.10E-02	8.07E-05	5.00E-03	6.91E-03
95%	1.24E-02	8.59E-05	6.10E-03	7.85E-03
Mean	6.56E-03	5.75E-05	2.64E-03	3.87E-03
Min	3.02E-05	5.27E-07	1.43E-06	7.00E-06
Max	3.02E-02	1.19E-04	2.06E-02	1.44E-02

**TABLE C.7 Peak DSR Percentiles (mrem/yr per pCi/g) for
Different Exposure Pathways for Sr-90 in Industrial Use
Scenario**

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	9.62E-04	9.03E-04	3.08E-07	4.18E-05
10%	2.84E-03	2.67E-03	7.35E-07	1.00E-04
15%	4.65E-03	4.42E-03	1.15E-06	1.53E-04
20%	6.54E-03	6.18E-03	1.55E-06	2.02E-04
25%	8.25E-03	7.91E-03	1.89E-06	2.51E-04
30%	8.72E-03	8.27E-03	2.16E-06	2.96E-04
35%	9.02E-03	8.58E-03	2.38E-06	3.44E-04
40%	9.35E-03	8.86E-03	2.60E-06	3.84E-04
45%	9.66E-03	9.17E-03	2.82E-06	4.22E-04
50%	9.94E-03	9.47E-03	3.03E-06	4.57E-04
55%	1.02E-02	9.68E-03	3.26E-06	4.94E-04
60%	1.04E-02	9.87E-03	3.55E-06	5.32E-04
65%	1.06E-02	1.01E-02	3.88E-06	5.66E-04
70%	1.09E-02	1.03E-02	4.20E-06	6.07E-04
75%	1.12E-02	1.05E-02	4.63E-06	6.54E-04
80%	1.16E-02	1.10E-02	5.11E-06	7.04E-04
85%	1.22E-02	1.16E-02	5.71E-06	7.62E-04
90%	1.29E-02	1.22E-02	6.63E-06	8.34E-04
95%	1.38E-02	1.30E-02	8.09E-06	9.52E-04
Mean	9.15E-03	8.68E-03	3.49E-06	4.69E-04
Min	7.80E-05	7.57E-05	1.90E-09	8.52E-07
Max	1.96E-02	1.80E-02	2.74E-05	1.76E-03

TABLE C.8 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Tc-99 in Industrial Use Scenario

Percentile	All Pathways	External	Inhalation w/o Rn	Soil Ingestion
5%	3.44E-06	2.35E-06	2.37E-08	8.40E-07
10%	9.83E-06	6.91E-06	5.70E-08	2.00E-06
15%	1.62E-05	1.15E-05	8.88E-08	3.05E-06
20%	2.27E-05	1.61E-05	1.19E-07	4.06E-06
25%	2.65E-05	2.05E-05	1.46E-07	5.00E-06
30%	2.87E-05	2.14E-05	1.67E-07	5.90E-06
35%	3.04E-05	2.22E-05	1.84E-07	6.85E-06
40%	3.18E-05	2.30E-05	2.02E-07	7.71E-06
45%	3.30E-05	2.38E-05	2.17E-07	8.44E-06
50%	3.41E-05	2.45E-05	2.34E-07	9.17E-06
55%	3.54E-05	2.51E-05	2.52E-07	9.88E-06
60%	3.65E-05	2.56E-05	2.75E-07	1.06E-05
65%	3.77E-05	2.61E-05	2.99E-07	1.13E-05
70%	3.89E-05	2.66E-05	3.25E-07	1.21E-05
75%	4.02E-05	2.73E-05	3.58E-07	1.31E-05
80%	4.18E-05	2.85E-05	3.94E-07	1.41E-05
85%	4.38E-05	3.00E-05	4.42E-07	1.53E-05
90%	4.70E-05	3.16E-05	5.10E-07	1.67E-05
95%	5.18E-05	3.39E-05	6.23E-07	1.90E-05
Mean	3.21E-05	2.25E-05	2.70E-07	9.37E-06
Min	2.52E-07	2.01E-07	1.50E-10	1.72E-08
Max	8.21E-05	4.69E-05	2.13E-06	3.57E-05

Tables C.9–C.16 show the peak DSR percentiles (mrem/yr per pCi/g) for different exposure pathways for individual radionuclides in the offsite resident scenario. RESRAD-OFFSITE version 2.6 was used in the analysis. For all radionuclides except Ba-133, water-dependent pathways were dominant, and for Ba-133, the water-independent pathway was dominant. For Am-241, Pu-238, and Pu-239, fish and plant ingestion were dominant exposure pathways. For Ba-133, the external exposure pathway was dominant. For C-14, aquatic food ingestion resulted in most dose. For Cs-137, fish, meat, and milk ingestion resulted in most dose. For Sr-90, fish, meat, and plant ingestion resulted in most dose. For Tc-99, plant, fish, and milk ingestion resulted in most dose.

TABLE C.9 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Am-241 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	2.16E-03	2.72E-06	1.99E-03	1.21E-04	2.90E-06	3.07E-07	9.24E-07	9.69E-08	6.63E-06	5.17E-09	3.51E-11	1.72E-12	2.15E-10
10%	2.69E-03	5.72E-06	2.47E-03	1.62E-04	3.68E-06	4.41E-07	1.95E-06	9.70E-08	6.64E-06	6.08E-09	3.86E-11	2.24E-12	3.08E-10
15%	3.32E-03	8.77E-06	3.07E-03	2.00E-04	4.50E-06	5.44E-07	2.96E-06	9.70E-08	6.64E-06	6.76E-09	4.15E-11	2.66E-12	3.89E-10
20%	4.17E-03	1.18E-05	3.85E-03	2.35E-04	5.50E-06	6.53E-07	4.01E-06	9.70E-08	6.64E-06	7.40E-09	4.41E-11	3.02E-12	4.80E-10
25%	4.91E-03	1.46E-05	4.56E-03	2.68E-04	6.61E-06	7.47E-07	4.96E-06	9.70E-08	6.64E-06	8.02E-09	4.63E-11	3.43E-12	5.47E-10
30%	5.62E-03	1.69E-05	5.21E-03	2.97E-04	7.37E-06	8.52E-07	5.92E-06	9.70E-08	6.64E-06	8.68E-09	4.88E-11	3.80E-12	6.16E-10
35%	6.21E-03	1.94E-05	5.75E-03	3.30E-04	8.04E-06	9.51E-07	6.97E-06	9.70E-08	6.64E-06	9.27E-09	5.07E-11	4.21E-12	6.85E-10
40%	6.66E-03	2.19E-05	6.14E-03	3.58E-04	8.62E-06	1.05E-06	8.13E-06	9.70E-08	6.64E-06	9.83E-09	5.30E-11	4.63E-12	7.48E-10
45%	6.92E-03	2.41E-05	6.40E-03	3.86E-04	9.11E-06	1.17E-06	9.27E-06	9.70E-08	6.64E-06	1.05E-08	5.53E-11	5.07E-12	8.09E-10
50%	7.12E-03	2.62E-05	6.57E-03	4.15E-04	9.62E-06	1.29E-06	1.03E-05	9.70E-08	6.64E-06	1.12E-08	5.78E-11	5.59E-12	8.71E-10
55%	7.31E-03	2.78E-05	6.72E-03	4.50E-04	1.01E-05	1.45E-06	1.14E-05	9.70E-08	6.64E-06	1.19E-08	6.03E-11	6.10E-12	9.31E-10
60%	7.49E-03	2.93E-05	6.88E-03	4.88E-04	1.06E-05	1.61E-06	1.27E-05	9.70E-08	6.64E-06	1.27E-08	6.32E-11	6.76E-12	9.95E-10
65%	7.69E-03	3.07E-05	7.09E-03	5.27E-04	1.10E-05	1.77E-06	1.40E-05	9.70E-08	6.64E-06	1.35E-08	6.62E-11	7.40E-12	1.06E-09
70%	7.95E-03	3.22E-05	7.30E-03	5.64E-04	1.14E-05	1.95E-06	1.52E-05	9.70E-08	6.64E-06	1.45E-08	7.00E-11	8.06E-12	1.13E-09
75%	8.26E-03	3.33E-05	7.58E-03	6.18E-04	1.19E-05	2.22E-06	1.66E-05	9.70E-08	6.64E-06	1.54E-08	7.43E-11	9.02E-12	1.20E-09
80%	8.66E-03	3.47E-05	8.01E-03	6.75E-04	1.24E-05	2.50E-06	1.81E-05	9.70E-08	6.64E-06	1.67E-08	7.91E-11	1.01E-11	1.28E-09
85%	9.24E-03	3.61E-05	8.59E-03	7.46E-04	1.30E-05	2.89E-06	1.99E-05	9.70E-08	6.64E-06	1.86E-08	8.56E-11	1.16E-11	1.39E-09
90%	1.00E-02	3.78E-05	9.36E-03	8.68E-04	1.38E-05	3.42E-06	2.22E-05	9.70E-08	6.64E-06	2.12E-08	9.50E-11	1.37E-11	1.50E-09
95%	1.19E-02	4.02E-05	1.13E-02	1.06E-03	1.49E-05	4.56E-06	2.57E-05	9.70E-08	6.64E-06	2.56E-08	1.12E-10	1.79E-11	1.69E-09
Mean	6.98E-03	2.38E-05	6.49E-03	4.80E-04	9.24E-06	1.72E-06	1.14E-05	9.70E-08	6.64E-06	1.28E-08	6.37E-11	7.17E-12	8.95E-10
Min	1.49E-03	1.90E-07	1.42E-03	3.37E-05	1.64E-06	5.33E-08	4.81E-08	9.46E-08	6.47E-06	1.97E-09	2.27E-11	6.44E-13	1.13E-11
Max	1.08E-01	4.98E-05	1.08E-01	3.70E-03	2.14E-05	1.26E-05	3.97E-05	9.70E-08	6.64E-06	2.13E-07	3.07E-10	7.49E-11	2.43E-09

TABLE C.10 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Ba-133 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	6.66E-05	1.17E-05	3.91E-06	1.17E-06	5.06E-08	6.46E-07	5.64E-10	6.15E-05	6.52E-10	3.29E-11	2.41E-13	2.64E-12	4.45E-14
10%	7.01E-05	1.73E-05	5.54E-06	1.48E-06	7.05E-08	8.64E-07	8.27E-10	6.26E-05	6.64E-10	3.80E-11	3.40E-13	3.40E-12	6.42E-14
15%	7.37E-05	2.23E-05	6.49E-06	1.68E-06	8.99E-08	1.05E-06	1.08E-09	6.29E-05	6.67E-10	4.27E-11	4.20E-13	4.03E-12	8.32E-14
20%	7.64E-05	2.68E-05	7.15E-06	1.87E-06	1.07E-07	1.20E-06	1.34E-09	6.30E-05	6.69E-10	4.74E-11	5.01E-13	4.60E-12	1.01E-13
25%	7.84E-05	3.03E-05	7.44E-06	2.05E-06	1.26E-07	1.38E-06	1.58E-09	6.31E-05	6.70E-10	5.10E-11	5.87E-13	5.18E-12	1.17E-13
30%	8.00E-05	3.30E-05	7.56E-06	2.22E-06	1.47E-07	1.54E-06	1.83E-09	6.31E-05	6.70E-10	5.51E-11	6.74E-13	5.72E-12	1.30E-13
35%	8.12E-05	3.53E-05	7.65E-06	2.39E-06	1.68E-07	1.70E-06	2.05E-09	6.31E-05	6.70E-10	5.91E-11	7.70E-13	6.32E-12	1.44E-13
40%	8.22E-05	3.70E-05	7.75E-06	2.57E-06	1.90E-07	1.87E-06	2.27E-09	6.32E-05	6.70E-10	6.33E-11	8.68E-13	6.91E-12	1.58E-13
45%	8.31E-05	3.85E-05	7.85E-06	2.75E-06	2.15E-07	2.06E-06	2.49E-09	6.32E-05	6.70E-10	6.77E-11	9.75E-13	7.55E-12	1.71E-13
50%	8.41E-05	3.98E-05	7.97E-06	2.94E-06	2.41E-07	2.27E-06	2.71E-09	6.32E-05	6.70E-10	7.24E-11	1.10E-12	8.22E-12	1.85E-13
55%	8.49E-05	4.10E-05	8.08E-06	3.16E-06	2.74E-07	2.49E-06	2.91E-09	6.32E-05	6.70E-10	7.72E-11	1.23E-12	8.99E-12	1.99E-13
60%	8.58E-05	4.21E-05	8.23E-06	3.37E-06	3.09E-07	2.75E-06	3.16E-09	6.32E-05	6.70E-10	8.19E-11	1.38E-12	9.79E-12	2.11E-13
65%	8.68E-05	4.31E-05	8.40E-06	3.61E-06	3.50E-07	3.03E-06	3.38E-09	6.32E-05	6.70E-10	8.69E-11	1.56E-12	1.07E-11	2.25E-13
70%	8.78E-05	4.43E-05	8.61E-06	3.88E-06	4.01E-07	3.34E-06	3.62E-09	6.32E-05	6.70E-10	9.35E-11	1.78E-12	1.18E-11	2.38E-13
75%	8.87E-05	4.55E-05	8.86E-06	4.18E-06	4.59E-07	3.73E-06	3.90E-09	6.32E-05	6.70E-10	1.01E-10	2.05E-12	1.31E-11	2.55E-13
80%	8.99E-05	4.69E-05	9.20E-06	4.54E-06	5.33E-07	4.16E-06	4.19E-09	6.32E-05	6.70E-10	1.09E-10	2.38E-12	1.47E-11	2.73E-13
85%	9.12E-05	4.84E-05	9.67E-06	4.97E-06	6.49E-07	4.76E-06	4.54E-09	6.32E-05	6.70E-10	1.20E-10	2.85E-12	1.68E-11	2.94E-13
90%	9.30E-05	5.04E-05	1.05E-05	5.60E-06	8.00E-07	5.63E-06	5.05E-09	6.32E-05	6.70E-10	1.37E-10	3.60E-12	1.98E-11	3.23E-13
95%	9.59E-05	5.32E-05	1.23E-05	6.60E-06	1.12E-06	7.40E-06	5.76E-09	6.32E-05	6.70E-10	1.64E-10	4.98E-12	2.55E-11	3.61E-13
Mean	8.32E-05	3.70E-05	8.21E-06	3.32E-06	3.72E-07	2.91E-06	2.88E-09	6.28E-05	6.66E-10	8.21E-11	1.67E-12	1.04E-11	1.91E-13
Min	4.88E-05	2.07E-06	1.92E-06	3.07E-07	5.42E-09	1.67E-07	2.96E-11	4.32E-05	4.58E-10	1.17E-11	6.53E-14	1.02E-12	3.77E-15
Max	4.40E-04	3.61E-04	9.33E-05	1.83E-05	4.10E-06	2.09E-05	4.70E-08	6.32E-05	6.70E-10	9.38E-10	2.13E-11	7.58E-11	1.13E-12

TABLE C.11 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for C-14 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	2.48E-04	4.57E-12	2.43E-04	4.02E-07	4.44E-07	1.03E-06	1.40E-11	1.27E-15	9.49E-07	2.40E-11	1.54E-11	2.95E-11	6.03E-16
10%	3.76E-04	6.03E-12	3.66E-04	5.31E-07	5.01E-07	1.06E-06	2.03E-11	1.31E-15	9.95E-07	3.04E-11	1.68E-11	3.07E-11	8.64E-16
15%	4.97E-04	7.71E-12	4.86E-04	7.40E-07	5.82E-07	1.09E-06	2.71E-11	1.34E-15	1.03E-06	3.72E-11	1.82E-11	3.15E-11	1.14E-15
20%	6.40E-04	9.77E-12	6.27E-04	1.04E-06	6.71E-07	1.12E-06	3.53E-11	1.38E-15	1.05E-06	4.49E-11	1.97E-11	3.23E-11	1.38E-15
25%	8.00E-04	1.19E-11	7.79E-04	1.36E-06	7.97E-07	1.15E-06	4.42E-11	1.42E-15	1.07E-06	5.40E-11	2.14E-11	3.29E-11	1.61E-15
30%	9.86E-04	1.38E-11	9.68E-04	1.71E-06	9.40E-07	1.18E-06	5.63E-11	1.47E-15	1.09E-06	6.51E-11	2.33E-11	3.36E-11	1.91E-15
35%	1.21E-03	1.57E-11	1.19E-03	2.07E-06	1.15E-06	1.23E-06	6.97E-11	1.52E-15	1.11E-06	8.00E-11	2.53E-11	3.43E-11	2.25E-15
40%	1.47E-03	1.75E-11	1.44E-03	2.55E-06	1.40E-06	1.29E-06	8.51E-11	1.57E-15	1.12E-06	9.45E-11	2.76E-11	3.50E-11	2.61E-15
45%	1.85E-03	2.04E-11	1.80E-03	3.21E-06	1.70E-06	1.39E-06	1.06E-10	1.63E-15	1.14E-06	1.08E-10	3.03E-11	3.56E-11	2.97E-15
50%	2.36E-03	3.00E-11	2.32E-03	4.26E-06	2.32E-06	1.69E-06	1.40E-10	1.66E-15	1.16E-06	1.19E-10	3.32E-11	3.63E-11	3.39E-15
55%	2.96E-03	6.93E-11	2.89E-03	6.15E-06	3.42E-06	2.99E-06	2.55E-10	1.69E-15	1.17E-06	1.34E-10	3.63E-11	3.71E-11	3.86E-15
60%	3.89E-03	1.46E-10	3.79E-03	1.07E-05	6.52E-06	5.74E-06	5.35E-10	1.72E-15	1.19E-06	1.49E-10	3.99E-11	3.79E-11	4.37E-15
65%	5.35E-03	2.68E-10	5.19E-03	1.94E-05	1.21E-05	1.16E-05	1.07E-09	1.74E-15	1.20E-06	1.65E-10	4.42E-11	3.87E-11	4.88E-15
70%	7.94E-03	5.22E-10	7.56E-03	3.54E-05	2.12E-05	2.25E-05	2.06E-09	1.77E-15	1.22E-06	1.84E-10	4.92E-11	3.97E-11	5.42E-15
75%	1.27E-02	9.98E-10	1.23E-02	7.13E-05	3.86E-05	4.26E-05	3.76E-09	1.79E-15	1.24E-06	2.11E-10	5.44E-11	4.09E-11	5.97E-15
80%	2.31E-02	1.74E-09	2.24E-02	1.37E-04	7.48E-05	7.08E-05	6.97E-09	1.82E-15	1.26E-06	2.39E-10	6.16E-11	4.26E-11	6.60E-15
85%	4.19E-02	3.28E-09	4.00E-02	3.03E-04	1.62E-04	1.33E-04	1.30E-08	1.85E-15	1.29E-06	2.76E-10	7.10E-11	4.45E-11	7.32E-15
90%	8.49E-02	6.40E-09	8.30E-02	6.75E-04	3.80E-04	2.92E-04	2.53E-08	1.88E-15	1.32E-06	3.42E-10	8.41E-11	4.76E-11	8.23E-15
95%	2.27E-01	1.40E-08	2.26E-01	2.26E-03	1.20E-03	6.93E-04	5.96E-08	1.94E-15	1.36E-06	4.79E-10	1.11E-10	5.36E-11	9.64E-15
Mean	5.95E-02	2.68E-09	5.89E-02	4.88E-04	2.43E-04	1.21E-04	1.16E-08	1.62E-15	1.16E-06	1.68E-10	4.59E-11	3.93E-11	4.09E-15
Min	4.91E-05	2.14E-12	4.66E-05	1.92E-07	3.40E-07	9.64E-07	4.29E-13	1.16E-15	7.82E-07	9.76E-12	1.22E-11	2.74E-11	1.81E-17
Max	9.51E+00	1.51E-07	9.51E+00	5.68E-02	3.86E-02	9.22E-03	6.30E-07	4.26E-15	1.55E-06	5.22E-09	1.39E-09	1.76E-09	8.85E-14

TABLE C.12 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Cs-137 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	2.69E-03	1.81E-04	1.31E-03	1.72E-05	3.22E-04	2.16E-04	2.45E-08	1.76E-04	2.67E-09	4.75E-10	1.47E-09	7.21E-10	1.61E-12
10%	3.10E-03	2.10E-04	1.71E-03	2.08E-05	3.76E-04	2.56E-04	3.48E-08	1.76E-04	2.67E-09	5.72E-10	1.72E-09	8.52E-10	2.27E-12
15%	3.45E-03	2.22E-04	2.03E-03	2.34E-05	4.18E-04	2.88E-04	4.39E-08	1.76E-04	2.67E-09	6.43E-10	1.94E-09	9.60E-10	2.84E-12
20%	3.76E-03	2.32E-04	2.31E-03	2.57E-05	4.56E-04	3.17E-04	5.04E-08	1.76E-04	2.67E-09	7.22E-10	2.14E-09	1.06E-09	3.27E-12
25%	4.08E-03	2.38E-04	2.62E-03	2.77E-05	4.91E-04	3.44E-04	5.70E-08	1.76E-04	2.67E-09	7.98E-10	2.32E-09	1.16E-09	3.66E-12
30%	4.39E-03	2.44E-04	2.90E-03	2.97E-05	5.25E-04	3.70E-04	6.26E-08	1.76E-04	2.67E-09	8.74E-10	2.50E-09	1.24E-09	3.99E-12
35%	4.71E-03	2.49E-04	3.19E-03	3.19E-05	5.54E-04	3.97E-04	6.81E-08	1.76E-04	2.67E-09	9.42E-10	2.70E-09	1.33E-09	4.31E-12
40%	5.05E-03	2.54E-04	3.50E-03	3.38E-05	5.87E-04	4.22E-04	7.31E-08	1.76E-04	2.67E-09	1.03E-09	2.92E-09	1.42E-09	4.59E-12
45%	5.41E-03	2.58E-04	3.87E-03	3.61E-05	6.22E-04	4.51E-04	7.80E-08	1.76E-04	2.67E-09	1.10E-09	3.14E-09	1.52E-09	4.88E-12
50%	5.76E-03	2.62E-04	4.21E-03	3.86E-05	6.55E-04	4.75E-04	8.29E-08	1.76E-04	2.67E-09	1.19E-09	3.37E-09	1.62E-09	5.16E-12
55%	6.13E-03	2.66E-04	4.60E-03	4.11E-05	6.92E-04	5.07E-04	8.75E-08	1.76E-04	2.67E-09	1.31E-09	3.63E-09	1.74E-09	5.48E-12
60%	6.54E-03	2.70E-04	5.04E-03	4.41E-05	7.28E-04	5.40E-04	9.25E-08	1.76E-04	2.67E-09	1.42E-09	3.94E-09	1.86E-09	5.75E-12
65%	7.07E-03	2.75E-04	5.50E-03	4.70E-05	7.73E-04	5.73E-04	9.84E-08	1.76E-04	2.67E-09	1.57E-09	4.25E-09	1.99E-09	6.04E-12
70%	7.62E-03	2.79E-04	6.08E-03	5.09E-05	8.19E-04	6.15E-04	1.04E-07	1.76E-04	2.67E-09	1.72E-09	4.63E-09	2.15E-09	6.42E-12
75%	8.38E-03	2.84E-04	6.75E-03	5.56E-05	8.73E-04	6.61E-04	1.11E-07	1.76E-04	2.67E-09	1.94E-09	5.08E-09	2.33E-09	6.81E-12
80%	9.14E-03	2.90E-04	7.60E-03	6.12E-05	9.39E-04	7.20E-04	1.18E-07	1.76E-04	2.67E-09	2.26E-09	5.67E-09	2.58E-09	7.21E-12
85%	1.03E-02	2.97E-04	8.70E-03	6.83E-05	1.02E-03	7.90E-04	1.27E-07	1.76E-04	2.67E-09	2.75E-09	6.57E-09	2.87E-09	7.73E-12
90%	1.18E-02	3.06E-04	1.03E-02	7.85E-05	1.14E-03	8.84E-04	1.38E-07	1.76E-04	2.67E-09	3.54E-09	8.06E-09	3.36E-09	8.42E-12
95%	1.49E-02	3.17E-04	1.34E-02	1.01E-04	1.31E-03	1.06E-03	1.53E-07	1.76E-04	2.67E-09	5.16E-09	1.14E-08	4.31E-09	9.31E-12
Mean	6.87E-03	2.58E-04	5.38E-03	4.74E-05	7.16E-04	5.35E-04	8.56E-08	1.76E-04	2.67E-09	1.84E-09	4.55E-09	1.99E-09	5.30E-12
Min	9.86E-04	4.90E-05	3.63E-04	6.18E-06	1.06E-04	6.61E-05	1.36E-09	1.75E-04	2.65E-09	2.00E-10	7.65E-10	3.55E-10	8.30E-14
Max	5.79E-02	3.91E-04	5.56E-02	8.40E-04	2.60E-03	2.20E-03	3.60E-07	1.76E-04	2.67E-09	5.23E-08	7.55E-08	3.10E-08	2.08E-11

TABLE C.13 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-238 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Radon-WD	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Radon	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	8.84E-04	1.40E-08	6.03E-04	1.33E-10	1.82E-04	9.05E-06	2.99E-07	7.24E-07	1.11E-10	7.58E-06	2.25E-10	4.95E-09	6.17E-11	1.26E-12	7.45E-11
10%	1.18E-03	1.64E-08	7.77E-04	3.25E-10	2.39E-04	1.24E-05	3.75E-07	1.11E-06	1.11E-10	7.58E-06	5.65E-10	5.85E-09	6.64E-11	1.47E-12	1.09E-10
15%	1.37E-03	1.77E-08	8.74E-04	4.88E-10	2.80E-04	1.49E-05	4.45E-07	1.46E-06	1.26E-10	7.58E-06	8.38E-10	6.61E-09	6.98E-11	1.65E-12	1.35E-10
20%	1.51E-03	1.86E-08	9.73E-04	6.19E-10	3.16E-04	1.68E-05	5.05E-07	1.85E-06	1.62E-10	7.58E-06	1.07E-09	7.28E-09	7.27E-11	1.81E-12	1.55E-10
25%	1.65E-03	1.95E-08	1.06E-03	7.08E-10	3.46E-04	1.81E-05	5.59E-07	2.17E-06	1.88E-10	7.58E-06	1.21E-09	7.85E-09	7.52E-11	1.95E-12	1.76E-10
30%	1.75E-03	2.04E-08	1.17E-03	7.74E-10	3.72E-04	1.92E-05	6.05E-07	2.46E-06	2.06E-10	7.58E-06	1.32E-09	8.45E-09	7.74E-11	2.09E-12	1.94E-10
35%	1.88E-03	2.12E-08	1.27E-03	8.17E-10	4.05E-04	2.01E-05	6.56E-07	2.72E-06	2.18E-10	7.58E-06	1.39E-09	9.01E-09	7.97E-11	2.23E-12	2.09E-10
40%	2.02E-03	2.21E-08	1.38E-03	8.45E-10	4.36E-04	2.10E-05	7.09E-07	3.04E-06	2.27E-10	7.58E-06	1.44E-09	9.70E-09	8.19E-11	2.37E-12	2.26E-10
45%	2.15E-03	2.29E-08	1.50E-03	8.68E-10	4.67E-04	2.20E-05	7.61E-07	3.32E-06	2.33E-10	7.58E-06	1.49E-09	1.03E-08	8.42E-11	2.52E-12	2.40E-10
50%	2.29E-03	2.37E-08	1.66E-03	8.84E-10	5.01E-04	2.27E-05	8.16E-07	3.58E-06	2.38E-10	7.58E-06	1.54E-09	1.09E-08	8.65E-11	2.67E-12	2.57E-10
55%	2.47E-03	2.44E-08	1.80E-03	8.96E-10	5.34E-04	2.35E-05	8.65E-07	3.87E-06	2.42E-10	7.58E-06	1.59E-09	1.17E-08	8.87E-11	2.83E-12	2.72E-10
60%	2.65E-03	2.50E-08	1.99E-03	9.05E-10	5.77E-04	2.42E-05	9.35E-07	4.14E-06	2.45E-10	7.58E-06	1.62E-09	1.25E-08	9.13E-11	3.01E-12	2.86E-10
65%	2.89E-03	2.57E-08	2.21E-03	9.11E-10	6.26E-04	2.51E-05	1.01E-06	4.40E-06	2.47E-10	7.58E-06	1.66E-09	1.33E-08	9.40E-11	3.20E-12	3.00E-10
70%	3.17E-03	2.64E-08	2.46E-03	9.16E-10	6.76E-04	2.60E-05	1.08E-06	4.74E-06	2.49E-10	7.58E-06	1.70E-09	1.43E-08	9.68E-11	3.42E-12	3.18E-10
75%	3.49E-03	2.73E-08	2.80E-03	9.20E-10	7.30E-04	2.69E-05	1.17E-06	5.07E-06	2.50E-10	7.58E-06	1.74E-09	1.55E-08	1.00E-10	3.66E-12	3.37E-10
80%	3.91E-03	2.83E-08	3.22E-03	9.25E-10	7.94E-04	2.80E-05	1.27E-06	5.41E-06	2.51E-10	7.58E-06	1.79E-09	1.69E-08	1.04E-10	3.97E-12	3.57E-10
85%	4.62E-03	2.93E-08	3.92E-03	9.36E-10	8.76E-04	2.93E-05	1.40E-06	5.88E-06	2.52E-10	7.58E-06	1.84E-09	1.87E-08	1.08E-10	4.35E-12	3.84E-10
90%	5.69E-03	3.08E-08	5.02E-03	9.61E-10	9.87E-04	3.09E-05	1.58E-06	6.45E-06	2.53E-10	7.58E-06	1.90E-09	2.09E-08	1.14E-10	4.89E-12	4.15E-10
95%	8.18E-03	3.31E-08	7.38E-03	1.01E-09	1.18E-03	3.34E-05	1.92E-06	7.34E-06	2.54E-10	7.58E-06	1.99E-09	2.52E-08	1.24E-10	5.80E-12	4.59E-10
Mean	3.13E-03	2.36E-08	2.53E-03	7.69E-10	5.72E-04	2.23E-05	9.19E-07	3.72E-06	2.12E-10	7.58E-06	1.40E-09	1.24E-08	8.92E-11	2.98E-12	2.60E-10
Min	3.37E-04	2.72E-09	2.13E-04	4.46E-12	6.07E-05	4.24E-06	8.48E-08	3.08E-08	1.10E-10	7.51E-06	6.34E-12	2.05E-09	4.61E-11	6.27E-13	6.05E-12
Max	4.32E-02	6.72E-08	4.25E-02	1.25E-09	3.09E-03	4.83E-05	3.72E-06	1.21E-05	2.56E-10	7.58E-06	2.37E-09	5.38E-08	3.31E-10	1.24E-11	8.37E-10

TABLE C.14 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-239 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	1.04E-03	1.50E-07	6.68E-04	2.32E-04	1.03E-05	3.22E-07	7.83E-06	6.93E-09	8.30E-06	9.77E-09	1.49E-10	2.60E-12	7.67E-10
10%	1.39E-03	2.33E-07	8.83E-04	3.29E-04	1.44E-05	4.31E-07	1.30E-05	6.93E-09	8.30E-06	1.18E-08	1.89E-10	3.26E-12	1.16E-09
15%	1.64E-03	2.97E-07	9.99E-04	3.93E-04	1.80E-05	5.08E-07	1.83E-05	6.93E-09	8.30E-06	1.35E-08	2.18E-10	3.77E-12	1.53E-09
20%	1.82E-03	3.55E-07	1.10E-03	4.48E-04	2.09E-05	5.77E-07	2.26E-05	6.93E-09	8.30E-06	1.50E-08	2.39E-10	4.22E-12	1.84E-09
25%	1.97E-03	4.16E-07	1.20E-03	5.02E-04	2.26E-05	6.45E-07	2.69E-05	6.93E-09	8.30E-06	1.65E-08	2.58E-10	4.65E-12	2.19E-09
30%	2.13E-03	4.71E-07	1.30E-03	5.49E-04	2.42E-05	7.14E-07	3.18E-05	6.93E-09	8.30E-06	1.80E-08	2.75E-10	5.04E-12	2.54E-09
35%	2.29E-03	5.31E-07	1.43E-03	5.91E-04	2.56E-05	7.71E-07	3.61E-05	6.93E-09	8.30E-06	1.95E-08	2.90E-10	5.45E-12	2.85E-09
40%	2.46E-03	5.83E-07	1.55E-03	6.37E-04	2.68E-05	8.25E-07	4.12E-05	6.93E-09	8.30E-06	2.09E-08	3.06E-10	5.85E-12	3.14E-09
45%	2.60E-03	6.41E-07	1.70E-03	6.85E-04	2.81E-05	8.92E-07	4.61E-05	6.93E-09	8.30E-06	2.21E-08	3.22E-10	6.30E-12	3.49E-09
50%	2.77E-03	6.93E-07	1.87E-03	7.38E-04	2.92E-05	9.60E-07	5.11E-05	6.93E-09	8.30E-06	2.37E-08	3.39E-10	6.80E-12	3.80E-09
55%	2.96E-03	7.39E-07	2.03E-03	7.86E-04	3.02E-05	1.02E-06	5.63E-05	6.93E-09	8.30E-06	2.54E-08	3.54E-10	7.33E-12	4.12E-09
60%	3.20E-03	7.77E-07	2.27E-03	8.44E-04	3.14E-05	1.09E-06	6.13E-05	6.93E-09	8.30E-06	2.76E-08	3.71E-10	7.93E-12	4.43E-09
65%	3.44E-03	8.17E-07	2.51E-03	9.06E-04	3.26E-05	1.17E-06	6.74E-05	6.93E-09	8.30E-06	2.98E-08	3.89E-10	8.53E-12	4.76E-09
70%	3.77E-03	8.54E-07	2.82E-03	9.71E-04	3.40E-05	1.26E-06	7.31E-05	6.93E-09	8.30E-06	3.22E-08	4.09E-10	9.21E-12	5.12E-09
75%	4.15E-03	8.89E-07	3.24E-03	1.05E-03	3.55E-05	1.35E-06	7.94E-05	6.93E-09	8.30E-06	3.50E-08	4.33E-10	1.01E-11	5.53E-09
80%	4.70E-03	9.24E-07	3.80E-03	1.14E-03	3.71E-05	1.47E-06	8.66E-05	6.93E-09	8.30E-06	3.86E-08	4.63E-10	1.11E-11	5.96E-09
85%	5.39E-03	9.59E-07	4.51E-03	1.25E-03	3.91E-05	1.63E-06	9.55E-05	6.93E-09	8.30E-06	4.34E-08	4.95E-10	1.21E-11	6.44E-09
90%	6.51E-03	1.00E-06	5.64E-03	1.42E-03	4.17E-05	1.86E-06	1.07E-04	6.93E-09	8.30E-06	5.15E-08	5.34E-10	1.39E-11	7.07E-09
95%	9.05E-03	1.06E-06	8.27E-03	1.75E-03	4.55E-05	2.21E-06	1.23E-04	6.93E-09	8.30E-06	6.64E-08	6.03E-10	1.65E-11	8.01E-09
Mean	3.67E-03	6.50E-07	2.87E-03	8.28E-04	2.89E-05	1.07E-06	5.61E-05	6.93E-09	8.30E-06	2.91E-08	3.53E-10	7.88E-12	3.99E-09
Min	3.32E-04	9.91E-09	1.95E-04	5.53E-05	4.34E-06	1.00E-07	3.31E-07	6.84E-09	8.19E-06	3.50E-09	5.98E-11	9.87E-13	8.48E-11
Max	7.18E-02	1.30E-06	7.04E-02	3.56E-03	6.86E-05	4.76E-06	1.94E-04	6.93E-09	8.30E-06	2.35E-07	1.00E-09	3.56E-11	1.25E-08

TABLE C.15 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Sr-90 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	8.35E-04	9.80E-07	1.34E-04	7.27E-05	2.05E-04	1.20E-04	3.24E-08	2.09E-06	1.08E-08	2.46E-09	2.34E-09	7.32E-10	1.83E-12
10%	9.28E-04	1.27E-06	1.59E-04	8.92E-05	2.48E-04	1.43E-04	4.71E-08	2.11E-06	1.09E-08	3.20E-09	3.13E-09	9.52E-10	2.71E-12
15%	1.01E-03	1.50E-06	1.84E-04	1.02E-04	2.80E-04	1.62E-04	5.90E-08	2.12E-06	1.10E-08	3.87E-09	3.97E-09	1.15E-09	3.36E-12
20%	1.07E-03	1.71E-06	2.06E-04	1.14E-04	3.10E-04	1.78E-04	7.00E-08	2.12E-06	1.10E-08	4.46E-09	4.77E-09	1.37E-09	4.02E-12
25%	1.14E-03	1.91E-06	2.33E-04	1.26E-04	3.38E-04	1.93E-04	8.01E-08	2.12E-06	1.10E-08	5.15E-09	5.58E-09	1.59E-09	4.64E-12
30%	1.22E-03	2.08E-06	2.59E-04	1.39E-04	3.65E-04	2.07E-04	9.03E-08	2.13E-06	1.10E-08	5.87E-09	6.35E-09	1.85E-09	5.21E-12
35%	1.28E-03	2.23E-06	2.86E-04	1.52E-04	3.92E-04	2.22E-04	1.01E-07	2.13E-06	1.10E-08	6.71E-09	7.26E-09	2.08E-09	5.84E-12
40%	1.36E-03	2.38E-06	3.18E-04	1.69E-04	4.23E-04	2.37E-04	1.10E-07	2.13E-06	1.10E-08	7.58E-09	8.23E-09	2.38E-09	6.42E-12
45%	1.43E-03	2.51E-06	3.54E-04	1.86E-04	4.53E-04	2.52E-04	1.20E-07	2.13E-06	1.10E-08	8.72E-09	9.24E-09	2.70E-09	6.99E-12
50%	1.52E-03	2.61E-06	3.92E-04	2.02E-04	4.87E-04	2.69E-04	1.30E-07	2.13E-06	1.10E-08	9.82E-09	1.05E-08	3.04E-09	7.62E-12
55%	1.60E-03	2.71E-06	4.40E-04	2.24E-04	5.24E-04	2.88E-04	1.41E-07	2.13E-06	1.10E-08	1.12E-08	1.18E-08	3.52E-09	8.29E-12
60%	1.70E-03	2.81E-06	4.92E-04	2.48E-04	5.68E-04	3.06E-04	1.52E-07	2.13E-06	1.10E-08	1.28E-08	1.35E-08	3.99E-09	8.90E-12
65%	1.82E-03	2.90E-06	5.56E-04	2.76E-04	6.23E-04	3.30E-04	1.63E-07	2.13E-06	1.10E-08	1.44E-08	1.54E-08	4.53E-09	9.57E-12
70%	1.95E-03	3.00E-06	6.34E-04	3.12E-04	6.80E-04	3.55E-04	1.74E-07	2.13E-06	1.10E-08	1.66E-08	1.78E-08	5.27E-09	1.02E-11
75%	2.13E-03	3.11E-06	7.31E-04	3.57E-04	7.65E-04	3.84E-04	1.87E-07	2.13E-06	1.10E-08	1.96E-08	2.08E-08	6.18E-09	1.10E-11
80%	2.33E-03	3.21E-06	8.67E-04	4.30E-04	8.66E-04	4.20E-04	2.02E-07	2.13E-06	1.10E-08	2.37E-08	2.44E-08	7.40E-09	1.19E-11
85%	2.62E-03	3.34E-06	1.06E-03	5.19E-04	1.02E-03	4.67E-04	2.21E-07	2.13E-06	1.10E-08	2.95E-08	3.08E-08	9.06E-09	1.30E-11
90%	3.09E-03	3.49E-06	1.35E-03	6.71E-04	1.28E-03	5.46E-04	2.45E-07	2.13E-06	1.10E-08	3.86E-08	4.00E-08	1.22E-08	1.44E-11
95%	3.99E-03	3.70E-06	1.95E-03	1.01E-03	1.89E-03	6.77E-04	2.82E-07	2.13E-06	1.10E-08	5.93E-08	6.28E-08	1.87E-08	1.65E-11
Mean	1.86E-03	2.50E-06	6.49E-04	3.34E-04	7.18E-04	3.23E-04	1.40E-07	2.12E-06	1.10E-08	1.80E-08	1.93E-08	5.70E-09	8.17E-12
Min	3.60E-04	3.22E-07	7.43E-05	2.32E-05	8.90E-05	4.80E-05	2.25E-09	1.69E-06	8.78E-09	6.76E-10	5.37E-10	2.42E-10	1.31E-13
Max	3.53E-02	1.19E-05	1.21E-02	7.99E-03	3.15E-02	8.94E-03	1.01E-06	2.13E-06	1.10E-08	4.79E-07	1.08E-06	4.14E-07	4.31E-11

TABLE C.16 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Tc-99 in Offsite Resident Scenario

Percentile	Total	External-WD	Fish	Plant-WD	Meat-WD	Milk-WD	Soil-WD	External-WI	Inhalation	Plant-WI	Meat-WI	Milk-WI	Soil-WI
5%	1.03E-04	4.68E-09	9.77E-06	4.88E-05	6.78E-07	1.78E-05	1.65E-09	1.59E-09	8.28E-10	1.12E-10	4.95E-13	7.63E-12	4.35E-15
10%	2.50E-04	9.03E-09	2.18E-05	1.21E-04	1.76E-06	3.97E-05	3.65E-09	1.59E-09	8.31E-10	1.39E-10	6.94E-13	1.07E-11	6.23E-15
15%	3.90E-04	1.39E-08	3.59E-05	1.99E-04	2.87E-06	6.69E-05	5.68E-09	1.60E-09	8.33E-10	1.65E-10	8.80E-13	1.30E-11	7.58E-15
20%	5.63E-04	2.02E-08	5.29E-05	2.85E-04	4.16E-06	9.71E-05	8.34E-09	1.60E-09	8.35E-10	1.89E-10	1.04E-12	1.53E-11	8.68E-15
25%	7.84E-04	2.74E-08	7.27E-05	4.02E-04	5.66E-06	1.34E-04	1.12E-08	1.61E-09	8.37E-10	2.10E-10	1.21E-12	1.79E-11	9.70E-15
30%	1.02E-03	3.77E-08	9.33E-05	5.15E-04	7.54E-06	1.73E-04	1.45E-08	1.61E-09	8.38E-10	2.39E-10	1.39E-12	2.07E-11	1.05E-14
35%	1.29E-03	4.93E-08	1.20E-04	6.41E-04	9.16E-06	2.18E-04	1.84E-08	1.61E-09	8.40E-10	2.65E-10	1.59E-12	2.36E-11	1.15E-14
40%	1.53E-03	6.12E-08	1.47E-04	7.81E-04	1.10E-05	2.59E-04	2.28E-08	1.61E-09	8.42E-10	2.92E-10	1.81E-12	2.66E-11	1.23E-14
45%	1.79E-03	7.24E-08	1.74E-04	8.92E-04	1.30E-05	3.06E-04	2.77E-08	1.62E-09	8.44E-10	3.20E-10	2.06E-12	3.02E-11	1.31E-14
50%	2.03E-03	8.20E-08	2.07E-04	1.02E-03	1.53E-05	3.58E-04	3.21E-08	1.62E-09	8.45E-10	3.51E-10	2.28E-12	3.43E-11	1.38E-14
55%	2.31E-03	8.91E-08	2.44E-04	1.16E-03	1.76E-05	4.14E-04	3.71E-08	1.62E-09	8.47E-10	3.86E-10	2.57E-12	3.89E-11	1.46E-14
60%	2.59E-03	9.60E-08	2.97E-04	1.33E-03	2.01E-05	4.77E-04	4.22E-08	1.63E-09	8.49E-10	4.31E-10	2.92E-12	4.32E-11	1.54E-14
65%	2.86E-03	1.03E-07	3.59E-04	1.51E-03	2.32E-05	5.43E-04	4.78E-08	1.63E-09	8.51E-10	4.80E-10	3.36E-12	4.92E-11	1.63E-14
70%	3.18E-03	1.10E-07	4.24E-04	1.72E-03	2.72E-05	6.25E-04	5.34E-08	1.63E-09	8.52E-10	5.39E-10	3.86E-12	5.64E-11	1.73E-14
75%	3.54E-03	1.17E-07	5.18E-04	1.98E-03	3.19E-05	7.27E-04	5.96E-08	1.64E-09	8.54E-10	6.05E-10	4.47E-12	6.46E-11	1.82E-14
80%	4.02E-03	1.25E-07	6.50E-04	2.30E-03	3.81E-05	8.45E-04	6.67E-08	1.64E-09	8.56E-10	6.92E-10	5.31E-12	7.54E-11	1.95E-14
85%	4.62E-03	1.33E-07	8.48E-04	2.67E-03	4.60E-05	1.02E-03	7.49E-08	1.64E-09	8.57E-10	8.23E-10	6.37E-12	9.32E-11	2.07E-14
90%	5.50E-03	1.46E-07	1.19E-03	3.35E-03	5.97E-05	1.24E-03	8.47E-08	1.65E-09	8.59E-10	1.03E-09	8.02E-12	1.19E-10	2.22E-14
95%	7.06E-03	1.63E-07	1.82E-03	4.70E-03	8.63E-05	1.75E-03	1.00E-07	1.65E-09	8.61E-10	1.39E-09	1.18E-11	1.71E-10	2.49E-14
Mean	2.59E-03	7.80E-08	4.92E-04	1.53E-03	2.57E-05	5.48E-04	3.91E-08	1.62E-09	8.45E-10	5.01E-10	3.70E-12	5.54E-11	1.41E-14
Min	3.93E-06	3.55E-10	1.76E-07	1.74E-06	1.17E-08	5.68E-07	1.99E-11	1.56E-09	8.12E-10	3.54E-11	8.99E-14	1.49E-12	1.35E-16
Max	4.25E-02	2.36E-07	4.07E-02	2.62E-02	6.81E-04	1.36E-02	1.93E-07	1.67E-09	8.69E-10	5.23E-09	7.73E-11	2.54E-09	3.76E-14

Tables C.17–C.24 show the peak DSR percentiles (mrem/yr per pCi/g) for different exposure pathways for individual radionuclides in the recreational use scenario. RESRAD (onsite) version 6.5 was used in the analysis. For C-14 and Tc-99, water-dependent pathways were dominant, and for others (Am-241, Ba-133, Cs-137, Pu-238, Pu-239, and Sr-90), water-independent pathways were dominant. For Am-241, Pu-238, and Pu-239, meat ingestion, soil ingestion, and inhalation were dominant exposure pathways. For Ba-133, external exposure was dominant. For C-14 and Tc-99, aquatic food ingestion resulted in most dose. For Cs-137, meat ingestion and external exposure resulted in most dose. For Sr-90, meat ingestion resulted in most dose.

TABLE C.17 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Am-241 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	1.00E-03	1.18E-04	1.78E-04	1.27E-04	0.00E+00	0.00E+00	0.00E+00	1.81E-03
10%	1.11E-03	1.63E-04	2.18E-04	1.84E-04	0.00E+00	0.00E+00	0.00E+00	1.96E-03
15%	1.14E-03	1.92E-04	2.51E-04	2.27E-04	0.00E+00	0.00E+00	0.00E+00	2.05E-03
20%	1.16E-03	2.12E-04	2.78E-04	2.64E-04	0.00E+00	0.00E+00	0.00E+00	2.11E-03
25%	1.17E-03	2.28E-04	3.01E-04	2.95E-04	0.00E+00	0.00E+00	0.00E+00	2.16E-03
30%	1.17E-03	2.47E-04	3.24E-04	3.25E-04	0.00E+00	0.00E+00	0.00E+00	2.22E-03
35%	1.17E-03	2.63E-04	3.49E-04	3.54E-04	0.00E+00	0.00E+00	0.00E+00	2.26E-03
40%	1.17E-03	2.80E-04	3.73E-04	3.81E-04	0.00E+00	0.00E+00	0.00E+00	2.31E-03
45%	1.18E-03	2.96E-04	3.99E-04	4.06E-04	0.00E+00	0.00E+00	0.00E+00	2.36E-03
50%	1.18E-03	3.13E-04	4.31E-04	4.29E-04	0.00E+00	0.00E+00	0.00E+00	2.40E-03
55%	1.18E-03	3.34E-04	4.60E-04	4.50E-04	0.00E+00	0.00E+00	0.00E+00	2.44E-03
60%	1.18E-03	3.59E-04	4.93E-04	4.73E-04	0.00E+00	0.00E+00	0.00E+00	2.49E-03
65%	1.18E-03	3.86E-04	5.27E-04	4.99E-04	0.00E+00	0.00E+00	0.00E+00	2.54E-03
70%	1.18E-03	4.12E-04	5.63E-04	5.27E-04	0.00E+00	0.00E+00	0.00E+00	2.58E-03
75%	1.18E-03	4.45E-04	6.03E-04	5.59E-04	0.00E+00	0.00E+00	0.00E+00	2.64E-03
80%	1.18E-03	4.88E-04	6.47E-04	5.92E-04	0.00E+00	0.00E+00	0.00E+00	2.71E-03
85%	1.18E-03	5.47E-04	6.98E-04	6.27E-04	5.86E-11	1.68E-09	1.23E-12	2.79E-03
90%	1.18E-03	6.13E-04	7.66E-04	6.72E-04	3.34E-09	9.40E-08	6.80E-11	2.89E-03
95%	1.18E-03	7.26E-04	8.80E-04	7.33E-04	4.30E-08	1.23E-06	9.07E-10	3.05E-03
Mean	1.14E-03	3.59E-04	4.67E-04	4.28E-04	6.01E-09	2.76E-07	1.81E-10	2.40E-03
Min	2.58E-08	2.78E-10	5.15E-09	5.46E-10	0.00E+00	0.00E+00	0.00E+00	5.24E-08
Max	1.18E-03	2.55E-03	1.46E-03	8.80E-04	2.24E-07	4.41E-05	1.58E-08	4.99E-03

TABLE C.18 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Ba-133 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	2.04E-07	1.38E-13	5.50E-11	1.10E-11	0.00E+00	0.00E+00	0.00E+00	2.31E-07
10%	1.27E-05	6.12E-12	2.11E-09	4.93E-10	0.00E+00	0.00E+00	0.00E+00	1.27E-05
15%	4.04E-05	1.74E-11	4.67E-09	1.35E-09	0.00E+00	0.00E+00	0.00E+00	4.04E-05
20%	6.10E-05	2.41E-11	6.91E-09	2.02E-09	0.00E+00	0.00E+00	0.00E+00	6.11E-05
25%	7.57E-05	3.04E-11	9.15E-09	2.61E-09	0.00E+00	0.00E+00	0.00E+00	7.58E-05
30%	8.46E-05	3.52E-11	1.12E-08	3.12E-09	0.00E+00	0.00E+00	0.00E+00	8.46E-05
35%	9.03E-05	3.88E-11	1.34E-08	3.59E-09	0.00E+00	0.00E+00	0.00E+00	9.04E-05
40%	9.36E-05	4.26E-11	1.60E-08	4.01E-09	0.00E+00	0.00E+00	0.00E+00	9.37E-05
45%	9.61E-05	4.62E-11	1.89E-08	4.40E-09	0.00E+00	0.00E+00	0.00E+00	9.62E-05
50%	9.76E-05	4.99E-11	2.18E-08	4.83E-09	0.00E+00	0.00E+00	0.00E+00	9.77E-05
55%	9.87E-05	5.35E-11	2.48E-08	5.20E-09	0.00E+00	0.00E+00	0.00E+00	9.88E-05
60%	9.95E-05	5.70E-11	2.88E-08	5.58E-09	0.00E+00	0.00E+00	0.00E+00	9.95E-05
65%	1.00E-04	6.20E-11	3.37E-08	5.97E-09	0.00E+00	0.00E+00	0.00E+00	1.00E-04
70%	1.00E-04	6.76E-11	3.97E-08	6.38E-09	0.00E+00	0.00E+00	0.00E+00	1.00E-04
75%	1.01E-04	7.35E-11	4.71E-08	6.79E-09	0.00E+00	0.00E+00	0.00E+00	1.01E-04
80%	1.01E-04	8.02E-11	5.66E-08	7.21E-09	0.00E+00	0.00E+00	0.00E+00	1.01E-04
85%	1.01E-04	8.97E-11	7.08E-08	7.81E-09	0.00E+00	0.00E+00	0.00E+00	1.01E-04
90%	1.01E-04	1.04E-10	9.39E-08	8.47E-09	0.00E+00	0.00E+00	0.00E+00	1.01E-04
95%	1.01E-04	1.25E-10	1.44E-07	9.30E-09	0.00E+00	0.00E+00	0.00E+00	1.01E-04
Mean	8.03E-05	5.49E-11	4.16E-08	4.73E-09	5.71E-11	3.77E-10	2.55E-13	8.04E-05
Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max	1.01E-04	4.74E-10	2.23E-06	1.17E-08	7.26E-08	4.96E-07	3.21E-10	1.02E-04

TABLE C.19 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for C-14 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
10%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
15%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
20%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
25%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
30%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
35%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
40%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
45%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
50%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
55%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
60%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
65%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
70%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.24E-21	1.71E-17	9.54E-22	1.77E-17
75%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.58E-07	4.23E-03	1.88E-07	4.24E-03
80%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.02E-07	1.55E-02	3.84E-07	1.55E-02
85%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.02E-07	5.45E-02	6.66E-07	5.45E-02
90%	6.11E-08	3.03E-06	6.00E-02	1.62E-07	1.48E-06	1.17E-01	1.17E-06	1.17E-01
95%	6.19E-08	3.90E-06	9.22E-02	2.56E-07	2.81E-06	2.50E-01	2.26E-06	2.50E-01
Mean	8.40E-09	5.08E-07	1.20E-02	3.04E-08	4.66E-07	4.61E-02	3.97E-07	4.61E-02
Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max	6.26E-08	9.13E-06	3.04E-01	4.30E-07	1.53E-05	1.14E+01	2.02E-05	1.14E+01

TABLE C.20 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Cs-137 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	9.79E-03	5.80E-09	1.90E-03	9.80E-07	0.00E+00	0.00E+00	0.00E+00	1.27E-02
10%	1.04E-02	7.82E-09	2.38E-03	1.42E-06	0.00E+00	0.00E+00	0.00E+00	1.33E-02
15%	1.07E-02	8.99E-09	2.84E-03	1.73E-06	0.00E+00	0.00E+00	0.00E+00	1.37E-02
20%	1.09E-02	9.87E-09	3.23E-03	2.01E-06	0.00E+00	0.00E+00	0.00E+00	1.42E-02
25%	1.10E-02	1.07E-08	3.67E-03	2.23E-06	0.00E+00	0.00E+00	0.00E+00	1.46E-02
30%	1.10E-02	1.15E-08	4.11E-03	2.46E-06	0.00E+00	0.00E+00	0.00E+00	1.51E-02
35%	1.11E-02	1.23E-08	4.51E-03	2.65E-06	0.00E+00	0.00E+00	0.00E+00	1.55E-02
40%	1.11E-02	1.31E-08	5.03E-03	2.85E-06	0.00E+00	0.00E+00	0.00E+00	1.60E-02
45%	1.11E-02	1.38E-08	5.52E-03	3.03E-06	0.00E+00	0.00E+00	0.00E+00	1.65E-02
50%	1.12E-02	1.46E-08	6.09E-03	3.21E-06	0.00E+00	0.00E+00	0.00E+00	1.71E-02
55%	1.12E-02	1.56E-08	6.61E-03	3.37E-06	0.00E+00	0.00E+00	0.00E+00	1.77E-02
60%	1.12E-02	1.67E-08	7.32E-03	3.55E-06	0.00E+00	0.00E+00	0.00E+00	1.83E-02
65%	1.12E-02	1.79E-08	8.15E-03	3.75E-06	0.00E+00	0.00E+00	0.00E+00	1.92E-02
70%	1.12E-02	1.92E-08	9.00E-03	3.94E-06	0.00E+00	0.00E+00	0.00E+00	2.00E-02
75%	1.13E-02	2.07E-08	1.01E-02	4.17E-06	0.00E+00	0.00E+00	0.00E+00	2.11E-02
80%	1.13E-02	2.26E-08	1.16E-02	4.40E-06	0.00E+00	0.00E+00	0.00E+00	2.25E-02
85%	1.13E-02	2.55E-08	1.35E-02	4.67E-06	0.00E+00	0.00E+00	0.00E+00	2.45E-02
90%	1.13E-02	2.87E-08	1.65E-02	5.00E-06	0.00E+00	0.00E+00	0.00E+00	2.76E-02
95%	1.13E-02	3.35E-08	2.38E-02	5.43E-06	0.00E+00	0.00E+00	0.00E+00	3.49E-02
Mean	1.09E-02	1.68E-08	8.62E-03	3.20E-06	0.00E+00	0.00E+00	0.00E+00	1.95E-02
Min	1.69E-04	3.90E-11	4.22E-05	4.19E-08	0.00E+00	0.00E+00	0.00E+00	2.11E-04
Max	1.13E-02	1.19E-07	1.80E-01	6.50E-06	0.00E+00	0.00E+00	0.00E+00	1.91E-01

TABLE C.21 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-238 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Radon (WI)	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	1.78E-06	7.65E-05	6.21E-14	2.25E-04	7.97E-05	0.00E+00	0.00E+00	0.00E+00	6.07E-04
10%	1.90E-06	1.02E-04	7.85E-14	2.74E-04	1.15E-04	0.00E+00	0.00E+00	0.00E+00	6.87E-04
15%	1.94E-06	1.18E-04	9.13E-14	3.07E-04	1.43E-04	0.00E+00	0.00E+00	0.00E+00	7.47E-04
20%	1.96E-06	1.30E-04	1.01E-13	3.42E-04	1.65E-04	0.00E+00	0.00E+00	0.00E+00	8.00E-04
25%	1.97E-06	1.40E-04	1.10E-13	3.73E-04	1.84E-04	0.00E+00	0.00E+00	0.00E+00	8.37E-04
30%	1.98E-06	1.51E-04	1.19E-13	4.03E-04	2.02E-04	0.00E+00	0.00E+00	0.00E+00	8.72E-04
35%	1.99E-06	1.61E-04	1.29E-13	4.33E-04	2.19E-04	0.00E+00	0.00E+00	0.00E+00	9.09E-04
40%	1.99E-06	1.72E-04	1.39E-13	4.64E-04	2.35E-04	0.00E+00	0.00E+00	0.00E+00	9.42E-04
45%	2.00E-06	1.81E-04	1.49E-13	4.95E-04	2.51E-04	0.00E+00	0.00E+00	0.00E+00	9.83E-04
50%	2.00E-06	1.91E-04	1.59E-13	5.23E-04	2.64E-04	0.00E+00	0.00E+00	0.00E+00	1.02E-03
55%	2.00E-06	2.03E-04	1.69E-13	5.61E-04	2.77E-04	0.00E+00	0.00E+00	0.00E+00	1.06E-03
60%	2.00E-06	2.19E-04	1.81E-13	5.98E-04	2.92E-04	0.00E+00	0.00E+00	0.00E+00	1.11E-03
65%	2.00E-06	2.35E-04	1.91E-13	6.40E-04	3.06E-04	0.00E+00	0.00E+00	0.00E+00	1.15E-03
70%	2.00E-06	2.51E-04	2.06E-13	6.86E-04	3.23E-04	0.00E+00	0.00E+00	0.00E+00	1.20E-03
75%	2.00E-06	2.71E-04	2.22E-13	7.29E-04	3.41E-04	0.00E+00	0.00E+00	0.00E+00	1.25E-03
80%	2.00E-06	2.97E-04	2.42E-13	7.91E-04	3.61E-04	0.00E+00	0.00E+00	0.00E+00	1.32E-03
85%	2.00E-06	3.34E-04	2.66E-13	8.60E-04	3.83E-04	0.00E+00	0.00E+00	0.00E+00	1.40E-03
90%	2.00E-06	3.78E-04	3.06E-13	9.42E-04	4.11E-04	0.00E+00	0.00E+00	0.00E+00	1.49E-03
95%	2.00E-06	4.44E-04	3.63E-13	1.07E-03	4.47E-04	0.00E+00	0.00E+00	0.00E+00	1.61E-03
Mean	1.95E-06	2.20E-04	1.78E-13	5.73E-04	2.63E-04	1.47E-10	1.02E-09	1.56E-12	1.06E-03
Min	3.28E-10	1.23E-09	2.67E-15	9.70E-09	1.89E-09	0.00E+00	0.00E+00	0.00E+00	1.32E-08
Max	2.00E-06	1.55E-03	1.03E-12	1.95E-03	5.32E-04	2.86E-07	2.84E-06	2.77E-09	2.49E-03

TABLE C.22 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-239 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	8.56E-06	1.81E-04	5.41E-04	1.91E-04	0.00E+00	0.00E+00	0.00E+00	1.46E-03
10%	9.10E-06	2.48E-04	6.58E-04	2.77E-04	0.00E+00	0.00E+00	0.00E+00	1.66E-03
15%	9.32E-06	2.84E-04	7.39E-04	3.41E-04	0.00E+00	0.00E+00	0.00E+00	1.80E-03
20%	9.41E-06	3.14E-04	8.17E-04	3.94E-04	0.00E+00	0.00E+00	0.00E+00	1.92E-03
25%	9.48E-06	3.39E-04	8.91E-04	4.42E-04	0.00E+00	0.00E+00	0.00E+00	2.02E-03
30%	9.51E-06	3.66E-04	9.61E-04	4.83E-04	0.00E+00	0.00E+00	0.00E+00	2.10E-03
35%	9.54E-06	3.89E-04	1.03E-03	5.24E-04	0.00E+00	0.00E+00	0.00E+00	2.19E-03
40%	9.56E-06	4.13E-04	1.11E-03	5.63E-04	0.00E+00	0.00E+00	0.00E+00	2.27E-03
45%	9.58E-06	4.36E-04	1.18E-03	5.96E-04	0.00E+00	0.00E+00	0.00E+00	2.36E-03
50%	9.59E-06	4.61E-04	1.26E-03	6.31E-04	0.00E+00	0.00E+00	0.00E+00	2.45E-03
55%	9.60E-06	4.94E-04	1.35E-03	6.64E-04	0.00E+00	0.00E+00	0.00E+00	2.54E-03
60%	9.60E-06	5.28E-04	1.43E-03	7.00E-04	0.00E+00	0.00E+00	0.00E+00	2.65E-03
65%	9.61E-06	5.66E-04	1.53E-03	7.36E-04	0.00E+00	0.00E+00	0.00E+00	2.75E-03
70%	9.61E-06	6.06E-04	1.64E-03	7.72E-04	0.00E+00	0.00E+00	0.00E+00	2.87E-03
75%	9.62E-06	6.52E-04	1.75E-03	8.16E-04	0.00E+00	0.00E+00	0.00E+00	3.02E-03
80%	9.62E-06	7.13E-04	1.90E-03	8.65E-04	0.00E+00	0.00E+00	0.00E+00	3.18E-03
85%	9.62E-06	8.00E-04	2.06E-03	9.22E-04	0.00E+00	0.00E+00	0.00E+00	3.36E-03
90%	9.62E-06	9.02E-04	2.26E-03	9.87E-04	0.00E+00	0.00E+00	0.00E+00	3.56E-03
95%	9.62E-06	1.06E-03	2.56E-03	1.08E-03	0.00E+00	0.00E+00	0.00E+00	3.87E-03
Mean	9.38E-06	5.30E-04	1.37E-03	6.31E-04	1.82E-14	2.24E-13	3.38E-16	2.54E-03
Min	6.72E-08	1.24E-06	1.23E-05	2.42E-06	0.00E+00	0.00E+00	0.00E+00	1.82E-05
Max	9.62E-06	3.74E-03	4.63E-03	1.29E-03	2.78E-11	4.04E-10	7.19E-13	6.03E-03

TABLE C.23 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Sr-90 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	1.57E-06	6.82E-10	1.07E-04	8.55E-08	0.00E+00	0.00E+00	0.00E+00	1.18E-04
10%	1.49E-05	5.84E-09	8.34E-04	5.86E-07	0.00E+00	0.00E+00	0.00E+00	9.18E-04
15%	3.59E-05	1.19E-08	1.57E-03	1.24E-06	0.00E+00	0.00E+00	0.00E+00	1.64E-03
20%	5.40E-05	1.75E-08	2.21E-03	1.87E-06	0.00E+00	0.00E+00	0.00E+00	2.31E-03
25%	7.09E-05	2.30E-08	2.78E-03	2.45E-06	0.00E+00	0.00E+00	0.00E+00	2.88E-03
30%	8.41E-05	2.77E-08	3.40E-03	3.04E-06	0.00E+00	0.00E+00	0.00E+00	3.51E-03
35%	9.46E-05	3.21E-08	4.11E-03	3.60E-06	0.00E+00	0.00E+00	0.00E+00	4.23E-03
40%	1.03E-04	3.53E-08	4.78E-03	4.14E-06	0.00E+00	0.00E+00	0.00E+00	4.93E-03
45%	1.09E-04	3.88E-08	5.55E-03	4.62E-06	0.00E+00	0.00E+00	0.00E+00	5.69E-03
50%	1.14E-04	4.22E-08	6.44E-03	5.14E-06	0.00E+00	0.00E+00	0.00E+00	6.56E-03
55%	1.19E-04	4.57E-08	7.55E-03	5.55E-06	0.00E+00	0.00E+00	0.00E+00	7.69E-03
60%	1.22E-04	4.94E-08	9.10E-03	6.03E-06	0.00E+00	0.00E+00	0.00E+00	9.24E-03
65%	1.25E-04	5.36E-08	1.07E-02	6.52E-06	0.00E+00	0.00E+00	0.00E+00	1.08E-02
70%	1.28E-04	5.82E-08	1.27E-02	7.00E-06	0.00E+00	0.00E+00	0.00E+00	1.28E-02
75%	1.30E-04	6.41E-08	1.52E-02	7.50E-06	0.00E+00	0.00E+00	0.00E+00	1.54E-02
80%	1.31E-04	7.04E-08	1.85E-02	8.06E-06	0.00E+00	0.00E+00	0.00E+00	1.86E-02
85%	1.33E-04	7.87E-08	2.33E-02	8.76E-06	0.00E+00	0.00E+00	0.00E+00	2.34E-02
90%	1.33E-04	8.97E-08	3.08E-02	9.72E-06	0.00E+00	0.00E+00	0.00E+00	3.09E-02
95%	1.34E-04	1.13E-07	4.74E-02	1.07E-05	0.00E+00	0.00E+00	0.00E+00	4.75E-02
Mean	9.53E-05	4.72E-08	1.34E-02	5.15E-06	2.21E-07	1.35E-05	5.66E-08	1.35E-02
Min	0.00E+00	0.00E+00	4.49E-29	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.49E-29
Max	1.35E-04	4.56E-07	5.62E-01	1.43E-05	1.01E-04	2.47E-02	2.84E-05	5.62E-01

TABLE C.24 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Tc-99 in Recreational Use Scenario

Percentile	External	Inhalation w/o Rn	Meat Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Meat Ingestion (WD)	All Pathways
5%	1.05E-13	1.14E-15	6.12E-12	3.41E-14	7.79E-07	4.17E-06	1.41E-09	5.78E-06
10%	2.34E-13	2.81E-15	1.57E-11	8.11E-14	1.76E-06	1.30E-05	4.07E-09	1.69E-05
15%	4.25E-13	5.20E-15	3.34E-11	1.58E-13	3.16E-06	2.78E-05	7.40E-09	3.34E-05
20%	7.26E-13	9.21E-15	6.30E-11	2.67E-13	4.95E-06	4.95E-05	1.23E-08	5.84E-05
25%	1.24E-12	1.62E-14	1.12E-10	4.43E-13	7.74E-06	8.31E-05	1.93E-08	9.76E-05
30%	1.97E-12	2.61E-14	1.88E-10	7.09E-13	1.17E-05	1.25E-04	3.05E-08	1.50E-04
35%	3.16E-12	4.17E-14	3.04E-10	1.17E-12	1.78E-05	1.80E-04	4.51E-08	2.16E-04
40%	5.01E-12	6.51E-14	5.03E-10	1.87E-12	2.68E-05	2.79E-04	6.95E-08	3.41E-04
45%	8.06E-12	1.05E-13	9.03E-10	3.06E-12	3.89E-05	3.97E-04	9.58E-08	4.72E-04
50%	1.36E-11	1.80E-13	1.63E-09	5.11E-12	5.52E-05	5.40E-04	1.30E-07	6.48E-04
55%	2.45E-11	3.38E-13	3.26E-09	9.62E-12	7.81E-05	7.46E-04	1.71E-07	8.73E-04
60%	5.72E-11	8.07E-13	1.14E-08	2.25E-11	1.06E-04	1.02E-03	2.28E-07	1.17E-03
65%	3.76E-10	6.00E-12	1.67E-07	1.50E-10	1.41E-04	1.33E-03	2.88E-07	1.50E-03
70%	3.75E-06	2.93E-08	7.50E-05	8.08E-07	1.79E-04	1.76E-03	3.52E-07	1.93E-03
75%	3.77E-06	3.75E-08	1.44E-04	1.18E-06	2.11E-04	2.30E-03	4.30E-07	2.49E-03
80%	3.79E-06	4.50E-08	2.21E-04	1.47E-06	2.33E-04	3.10E-03	5.34E-07	3.31E-03
85%	3.80E-06	5.33E-08	3.27E-04	1.69E-06	2.50E-04	4.16E-03	6.64E-07	4.36E-03
90%	3.81E-06	6.56E-08	4.93E-04	1.96E-06	2.65E-04	6.14E-03	8.70E-07	6.31E-03
95%	3.82E-06	8.47E-08	8.44E-04	2.32E-06	2.81E-04	1.05E-02	1.25E-06	1.07E-02
Mean	1.30E-06	1.94E-08	1.62E-04	5.45E-07	1.03E-04	2.46E-03	3.29E-07	2.56E-03
Min	1.71E-15	1.79E-17	5.18E-14	8.70E-16	0.00E+00	0.00E+00	0.00E+00	1.18E-09
Max	3.86E-06	2.43E-07	1.15E-02	3.12E-06	3.60E-04	1.28E-01	7.23E-06	1.29E-01

Tables C.25–C.32 show the peak DSR percentiles (mrem/yr per pCi/g) for different exposure pathways for individual radionuclides in the onsite resident scenario. RESRAD (onsite) version 6.5 was used in the analysis. For C-14 and Tc-99, water-dependent pathways were dominant, and for others (Am-241, Ba-133, Cs-137, Pu-238, Pu-239, and Sr-90), water-independent pathways were dominant. For Am-241, plant ingestion, external exposure, and soil ingestion were dominant. For Ba-133, external exposure was dominant. For C-14, aquatic food ingestion resulted in most dose. For Cs-137, external exposure and meat ingestion resulted in most dose. For Pu-238 and Pu-239, plant ingestion, soil ingestion, and inhalation were dominant exposure pathways. For Sr-90, plant and meat ingestion resulted in most dose. For Tc-99, water and plant ingestion resulted in most dose.

TABLE C.25 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Am-241 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	4.45E-03	6.67E-04	1.13E-02	3.75E-04	9.85E-06	2.50E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.96E-02
10%	5.00E-03	9.09E-04	1.59E-02	4.14E-04	1.28E-05	3.66E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.61E-02
15%	5.44E-03	1.06E-03	2.00E-02	4.42E-04	1.54E-05	4.54E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.05E-02
20%	5.84E-03	1.20E-03	2.40E-02	4.64E-04	1.78E-05	5.29E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.48E-02
25%	6.20E-03	1.36E-03	2.81E-02	4.86E-04	2.01E-05	5.96E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.91E-02
30%	6.53E-03	1.53E-03	3.24E-02	5.04E-04	2.24E-05	6.55E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.33E-02
35%	6.87E-03	1.70E-03	3.67E-02	5.23E-04	2.46E-05	7.10E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.75E-02
40%	7.24E-03	1.85E-03	4.16E-02	5.40E-04	2.70E-05	7.61E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.29E-02
45%	7.61E-03	2.01E-03	4.68E-02	5.55E-04	2.96E-05	8.10E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.81E-02
50%	8.00E-03	2.18E-03	5.24E-02	5.72E-04	3.22E-05	8.57E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.46E-02
55%	8.41E-03	2.35E-03	5.92E-02	5.88E-04	3.51E-05	9.00E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.03E-02
60%	8.86E-03	2.55E-03	6.64E-02	6.06E-04	3.88E-05	9.47E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.78E-02
65%	9.40E-03	2.76E-03	7.46E-02	6.28E-04	4.24E-05	9.98E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.62E-02
70%	9.98E-03	2.99E-03	8.51E-02	6.49E-04	4.73E-05	1.05E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.07E-01
75%	1.06E-02	3.26E-03	9.80E-02	6.77E-04	5.19E-05	1.11E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.20E-01
80%	1.14E-02	3.59E-03	1.14E-01	7.03E-04	5.92E-05	1.18E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.37E-01
85%	1.25E-02	4.08E-03	1.38E-01	7.44E-04	6.79E-05	1.25E-02	1.95E-07	2.20E-09	7.47E-09	1.23E-09	3.58E-11	1.61E-01
90%	1.40E-02	4.73E-03	1.72E-01	8.04E-04	8.06E-05	1.34E-02	5.21E-06	6.92E-08	2.01E-07	3.60E-08	9.94E-10	1.96E-01
95%	1.64E-02	5.87E-03	2.37E-01	8.98E-04	1.05E-04	1.46E-02	6.35E-05	7.90E-07	2.46E-06	4.25E-07	1.04E-08	2.68E-01
Mean	8.83E-03	2.57E-03	8.04E-02	5.95E-04	4.16E-05	8.54E-03	4.51E-03	1.13E-04	1.25E-04	1.48E-06	1.55E-07	1.05E-01
Min	1.40E-11	5.49E-12	2.30E-10	2.22E-12	1.35E-13	2.85E-11	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.81E-10
Max	2.37E-02	1.96E-02	2.19E+00	2.34E-03	4.19E-04	1.97E-02	4.68E+00	1.18E-01	9.48E-02	1.08E-03	1.04E-04	4.89E+00

TABLE C.26 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Ba-133 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	1.72E-06	8.51E-13	1.52E-08	1.46E-10	4.88E-10	2.74E-10	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.95E-06
10%	9.61E-05	4.34E-11	4.90E-07	5.21E-09	2.25E-08	1.21E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.88E-05
15%	2.63E-04	9.94E-11	1.05E-06	1.02E-08	4.50E-08	2.63E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.71E-04
20%	3.61E-04	1.45E-10	1.56E-06	1.52E-08	6.20E-08	3.90E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.68E-04
25%	4.06E-04	1.78E-10	2.05E-06	2.00E-08	7.94E-08	5.07E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.19E-04
30%	4.46E-04	2.05E-10	2.64E-06	2.53E-08	9.47E-08	6.11E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.60E-04
35%	4.82E-04	2.35E-10	3.21E-06	3.08E-08	1.11E-07	7.10E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.92E-04
40%	5.15E-04	2.69E-10	3.90E-06	3.59E-08	1.28E-07	8.14E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.27E-04
45%	5.53E-04	3.02E-10	4.59E-06	4.36E-08	1.47E-07	8.88E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.65E-04
50%	5.92E-04	3.32E-10	5.42E-06	5.20E-08	1.67E-07	9.73E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.05E-04
55%	6.28E-04	3.66E-10	6.30E-06	6.11E-08	1.91E-07	1.05E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.43E-04
60%	6.70E-04	4.01E-10	7.43E-06	7.01E-08	2.22E-07	1.12E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.79E-04
65%	7.06E-04	4.38E-10	8.59E-06	8.32E-08	2.51E-07	1.19E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.21E-04
70%	7.58E-04	4.84E-10	1.01E-05	1.00E-07	2.88E-07	1.27E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.74E-04
75%	8.14E-04	5.36E-10	1.21E-05	1.21E-07	3.36E-07	1.36E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.30E-04
80%	8.88E-04	5.94E-10	1.48E-05	1.46E-07	4.04E-07	1.45E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.02E-04
85%	9.78E-04	6.69E-10	1.84E-05	1.84E-07	4.89E-07	1.56E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.97E-04
90%	1.10E-03	7.99E-10	2.44E-05	2.49E-07	6.34E-07	1.69E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.11E-03
95%	1.29E-03	9.99E-10	3.83E-05	3.96E-07	9.55E-07	1.85E-07	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.31E-03
Mean	6.21E-04	3.94E-10	1.08E-05	1.10E-07	3.00E-07	9.45E-08	5.76E-08	2.98E-10	2.22E-09	2.03E-10	8.31E-10	6.32E-04
Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max	2.00E-03	3.58E-09	5.15E-04	8.98E-06	1.36E-05	2.32E-07	5.56E-05	3.98E-07	2.15E-06	3.51E-07	1.05E-06	2.04E-03

TABLE C.27 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for C-14 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
10%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
15%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
20%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
25%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
30%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
35%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
40%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
45%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
50%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
55%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
60%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
65%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
70%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
75%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
80%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
85%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.77E-04	5.95E-04	1.94E-05	8.11E-06	2.26E-05	1.01E-03
90%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.70E-04	2.46E-03	6.22E-05	2.52E-05	7.21E-05	3.45E-03
95%	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.21E-03	7.98E-03	1.35E-04	5.33E-05	1.51E-04	1.05E-02
Mean	7.48E-10	5.07E-08	3.01E-04	1.17E-04	4.00E-05	5.17E-09	1.91E-04	3.11E-03	2.19E-05	8.52E-06	2.40E-05	3.36E-03
Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max	1.02E-06	5.13E-05	3.02E-01	1.17E-01	4.01E-02	7.72E-06	1.88E-02	3.00E+00	1.78E-03	7.99E-04	2.35E-03	3.00E+00

TABLE C.28 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Cs-137 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	4.31E-02	3.22E-08	3.03E-03	6.08E-03	1.47E-03	2.00E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.02E-02
10%	4.80E-02	4.36E-08	4.37E-03	7.97E-03	1.93E-03	2.81E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.93E-02
15%	5.20E-02	5.04E-08	5.60E-03	9.57E-03	2.37E-03	3.44E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.78E-02
20%	5.55E-02	5.70E-08	6.83E-03	1.12E-02	2.76E-03	3.98E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.41E-02
25%	5.90E-02	6.43E-08	8.07E-03	1.29E-02	3.15E-03	4.46E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.00E-01
30%	6.23E-02	7.29E-08	9.35E-03	1.45E-02	3.61E-03	4.89E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.07E-01
35%	6.55E-02	8.04E-08	1.07E-02	1.65E-02	4.07E-03	5.31E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.14E-01
40%	6.89E-02	8.63E-08	1.22E-02	1.84E-02	4.60E-03	5.67E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.21E-01
45%	7.22E-02	9.40E-08	1.40E-02	2.07E-02	5.12E-03	6.05E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.28E-01
50%	7.64E-02	1.02E-07	1.58E-02	2.33E-02	5.79E-03	6.37E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.36E-01
55%	8.02E-02	1.10E-07	1.79E-02	2.62E-02	6.49E-03	6.73E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.43E-01
60%	8.46E-02	1.19E-07	2.03E-02	2.94E-02	7.34E-03	7.06E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.52E-01
65%	8.95E-02	1.29E-07	2.32E-02	3.39E-02	8.31E-03	7.47E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.61E-01
70%	9.48E-02	1.39E-07	2.68E-02	3.83E-02	9.43E-03	7.84E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.71E-01
75%	1.01E-01	1.51E-07	3.11E-02	4.40E-02	1.09E-02	8.29E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.83E-01
80%	1.09E-01	1.66E-07	3.66E-02	5.17E-02	1.31E-02	8.79E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.99E-01
85%	1.19E-01	1.89E-07	4.45E-02	6.23E-02	1.58E-02	9.33E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.23E-01
90%	1.33E-01	2.21E-07	5.65E-02	8.16E-02	2.01E-02	9.98E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.56E-01
95%	1.55E-01	2.73E-07	8.11E-02	1.19E-01	2.90E-02	1.09E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.17E-01
Mean	8.42E-02	1.20E-07	2.60E-02	3.87E-02	9.63E-03	6.39E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.59E-01
Min	2.59E-03	2.95E-10	5.07E-04	1.41E-03	2.54E-04	2.00E-06	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.56E-03
Max	2.23E-01	9.13E-07	7.17E-01	1.21E+00	4.52E-01	1.29E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.45E+00

TABLE C.29 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-238 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Radon (WI)	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Radon (WD)	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	7.67E-06	4.19E-04	8.08E-08	7.18E-03	4.69E-04	4.78E-06	1.61E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.42E-02
10%	8.58E-06	5.69E-04	9.58E-08	9.90E-03	5.13E-04	5.71E-06	2.28E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.74E-02
15%	9.34E-06	6.59E-04	1.05E-07	1.25E-02	5.41E-04	6.46E-06	2.82E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.00E-02
20%	9.94E-06	7.47E-04	1.11E-07	1.49E-02	5.68E-04	7.05E-06	3.29E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.22E-02
25%	1.05E-05	8.42E-04	1.16E-07	1.74E-02	5.93E-04	7.63E-06	3.67E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.48E-02
30%	1.11E-05	9.55E-04	1.21E-07	2.01E-02	6.13E-04	8.10E-06	4.03E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.75E-02
35%	1.18E-05	1.04E-03	1.27E-07	2.28E-02	6.33E-04	8.62E-06	4.37E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.03E-02
40%	1.24E-05	1.13E-03	1.31E-07	2.57E-02	6.54E-04	9.17E-06	4.68E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.33E-02
45%	1.30E-05	1.23E-03	1.37E-07	2.89E-02	6.77E-04	9.76E-06	4.98E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.65E-02
50%	1.36E-05	1.33E-03	1.41E-07	3.25E-02	6.97E-04	1.03E-05	5.26E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.05E-02
55%	1.44E-05	1.44E-03	1.45E-07	3.65E-02	7.18E-04	1.10E-05	5.52E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.43E-02
60%	1.52E-05	1.56E-03	1.50E-07	4.09E-02	7.39E-04	1.16E-05	5.82E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.90E-02
65%	1.60E-05	1.69E-03	1.56E-07	4.60E-02	7.65E-04	1.24E-05	6.12E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.36E-02
70%	1.70E-05	1.83E-03	1.62E-07	5.23E-02	7.93E-04	1.32E-05	6.47E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.97E-02
75%	1.82E-05	1.99E-03	1.70E-07	6.03E-02	8.22E-04	1.42E-05	6.81E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.82E-02
80%	1.96E-05	2.19E-03	1.77E-07	7.03E-02	8.63E-04	1.55E-05	7.21E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.85E-02
85%	2.14E-05	2.49E-03	1.87E-07	8.47E-02	9.10E-04	1.70E-05	7.66E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.26E-02
90%	2.37E-05	2.90E-03	1.99E-07	1.05E-01	9.80E-04	1.94E-05	8.21E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.13E-01
95%	2.76E-05	3.60E-03	2.21E-07	1.48E-01	1.08E-03	2.31E-05	8.92E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.56E-01
Mean	1.51E-05	1.58E-03	1.44E-07	4.95E-02	7.28E-04	1.17E-05	5.25E-03	1.36E-07	9.78E-10	9.88E-17	5.22E-09	1.12E-09	1.41E-09	5.71E-02
Min	1.73E-09	4.82E-09	4.67E-09	1.46E-06	8.59E-08	3.59E-08	3.21E-08	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.65E-06
Max	3.97E-05	1.19E-02	3.51E-07	8.99E-01	3.63E-03	8.14E-05	1.07E-02	2.67E-04	3.52E-06	2.86E-13	1.03E-05	3.01E-06	3.26E-06	9.12E-01

TABLE C.30 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Pu-239 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	3.69E-05	1.00E-03	1.68E-02	1.12E-03	1.05E-05	3.86E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.39E-02
10%	4.13E-05	1.37E-03	2.38E-02	1.23E-03	1.27E-05	5.50E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.18E-02
15%	4.47E-05	1.59E-03	3.00E-02	1.30E-03	1.44E-05	6.80E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.76E-02
20%	4.78E-05	1.80E-03	3.57E-02	1.36E-03	1.58E-05	7.91E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.33E-02
25%	5.07E-05	2.03E-03	4.18E-02	1.42E-03	1.71E-05	8.80E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.98E-02
30%	5.34E-05	2.30E-03	4.80E-02	1.47E-03	1.83E-05	9.74E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.62E-02
35%	5.65E-05	2.51E-03	5.44E-02	1.52E-03	1.96E-05	1.05E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.26E-02
40%	5.93E-05	2.74E-03	6.15E-02	1.57E-03	2.08E-05	1.12E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.99E-02
45%	6.24E-05	2.97E-03	7.00E-02	1.62E-03	2.23E-05	1.19E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.80E-02
50%	6.55E-05	3.20E-03	7.83E-02	1.67E-03	2.37E-05	1.26E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.69E-02
55%	6.89E-05	3.46E-03	8.76E-02	1.72E-03	2.51E-05	1.32E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.07E-01
60%	7.26E-05	3.75E-03	9.92E-02	1.77E-03	2.66E-05	1.39E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.18E-01
65%	7.67E-05	4.06E-03	1.11E-01	1.84E-03	2.85E-05	1.47E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.30E-01
70%	8.16E-05	4.39E-03	1.26E-01	1.91E-03	3.05E-05	1.55E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.44E-01
75%	8.70E-05	4.78E-03	1.45E-01	1.98E-03	3.28E-05	1.63E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.64E-01
80%	9.35E-05	5.26E-03	1.69E-01	2.06E-03	3.57E-05	1.73E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.88E-01
85%	1.02E-04	6.00E-03	2.03E-01	2.18E-03	3.96E-05	1.84E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.23E-01
90%	1.14E-04	6.97E-03	2.51E-01	2.35E-03	4.52E-05	1.97E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.70E-01
95%	1.33E-04	8.61E-03	3.52E-01	2.59E-03	5.44E-05	2.13E-02	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.73E-01
Mean	7.23E-05	3.79E-03	1.19E-01	1.74E-03	2.69E-05	1.26E-02	6.47E-11	1.72E-13	2.49E-12	2.39E-13	3.84E-13	1.37E-01
Min	7.59E-07	9.27E-06	2.85E-04	1.19E-05	4.10E-07	1.46E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.63E-04
Max	1.91E-04	2.87E-02	2.15E+00	8.69E-03	1.91E-04	2.57E-02	1.61E-07	3.34E-10	6.19E-09	5.22E-10	5.42E-10	2.18E+00

TABLE C.31 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Sr-90 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	1.35E-05	5.27E-09	3.13E-03	8.29E-04	1.96E-04	1.79E-06	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.42E-03
10%	1.05E-04	3.67E-08	2.06E-02	5.19E-03	1.17E-03	1.27E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.97E-02
15%	2.37E-04	7.46E-08	3.82E-02	9.20E-03	2.26E-03	2.57E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.30E-02
20%	3.58E-04	1.06E-07	5.33E-02	1.34E-02	3.15E-03	3.85E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.31E-02
25%	4.44E-04	1.37E-07	7.15E-02	1.75E-02	4.14E-03	5.10E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.77E-02
30%	5.01E-04	1.69E-07	8.81E-02	2.17E-02	5.06E-03	6.22E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.18E-01
35%	5.53E-04	1.91E-07	1.06E-01	2.65E-02	6.18E-03	7.26E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.44E-01
40%	6.02E-04	2.17E-07	1.27E-01	3.17E-02	7.39E-03	8.34E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.70E-01
45%	6.48E-04	2.45E-07	1.51E-01	3.79E-02	8.78E-03	9.28E-05	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.04E-01
50%	6.92E-04	2.77E-07	1.74E-01	4.39E-02	1.04E-02	1.02E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.35E-01
55%	7.48E-04	3.07E-07	2.07E-01	5.06E-02	1.24E-02	1.10E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.79E-01
60%	7.99E-04	3.40E-07	2.37E-01	5.92E-02	1.46E-02	1.21E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.21E-01
65%	8.54E-04	3.74E-07	2.77E-01	6.97E-02	1.69E-02	1.30E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.72E-01
70%	9.23E-04	4.16E-07	3.21E-01	8.25E-02	2.00E-02	1.39E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.25E-01
75%	9.90E-04	4.63E-07	3.75E-01	9.87E-02	2.40E-02	1.49E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.03E-01
80%	1.08E-03	5.15E-07	4.55E-01	1.19E-01	2.92E-02	1.61E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.07E-01
85%	1.18E-03	5.83E-07	5.69E-01	1.50E-01	3.67E-02	1.76E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.50E-01
90%	1.33E-03	6.92E-07	7.29E-01	1.94E-01	4.76E-02	1.93E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.71E-01
95%	1.59E-03	8.96E-07	1.06E+00	2.99E-01	7.58E-02	2.15E-04	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.44E+00
Mean	7.37E-04	3.39E-07	3.11E-01	8.42E-02	2.08E-02	1.03E-04	2.75E-04	3.89E-06	1.15E-05	2.19E-05	1.15E-05	4.17E-01
Min	0.00E+00	0.00E+00	3.30E-29	8.02E-30	2.42E-30	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.34E-29
Max	2.57E-03	2.91E-06	5.45E+00	1.95E+00	4.71E-01	2.85E-04	1.93E-01	3.40E-03	7.74E-03	1.77E-02	8.91E-03	7.38E+00

TABLE C.32 Peak DSR Percentiles (mrem/yr per pCi/g) for Different Exposure Pathways for Tc-99 in Onsite Resident Scenario

Percentile	External	Inhalation w/o Rn	Plant Ingestion (WI)	Meat Ingestion (WI)	Milk Ingestion (WI)	Soil Ingestion	Water Ingestion	Aquatic Foods	Plant Ingestion (WD)	Meat Ingestion (WD)	Milk Ingestion (WD)	All Pathways
5%	6.58E-13	7.02E-15	2.00E-08	4.44E-11	5.21E-10	6.19E-13	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.60E-05
10%	1.56E-12	1.75E-14	4.79E-08	1.12E-10	1.26E-09	1.56E-12	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.02E-05
15%	2.89E-12	3.48E-14	9.25E-08	2.25E-10	2.57E-09	3.09E-12	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.08E-05
20%	5.07E-12	5.84E-14	1.58E-07	3.97E-10	4.45E-09	5.25E-12	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.58E-04
25%	7.82E-12	9.38E-14	2.69E-07	6.72E-10	7.29E-09	8.72E-12	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.40E-04
30%	1.26E-11	1.47E-13	4.40E-07	1.10E-09	1.26E-08	1.30E-11	1.41E-03	1.85E-06	1.27E-04	8.90E-07	2.34E-05	1.68E-03
35%	1.88E-11	2.36E-13	7.17E-07	1.71E-09	2.06E-08	2.02E-11	2.91E-03	5.19E-06	2.69E-04	2.00E-06	5.30E-05	3.40E-03
40%	2.97E-11	3.68E-13	1.13E-06	2.62E-09	3.40E-08	3.07E-11	5.03E-03	9.42E-06	4.57E-04	3.57E-06	8.96E-05	5.71E-03
45%	4.53E-11	5.64E-13	1.71E-06	4.10E-09	5.20E-08	4.63E-11	7.51E-03	1.48E-05	7.23E-04	5.83E-06	1.41E-04	8.61E-03
50%	7.16E-11	8.30E-13	2.66E-06	6.57E-09	7.82E-08	7.10E-11	1.13E-02	2.29E-05	1.11E-03	9.07E-06	2.11E-04	1.29E-02
55%	1.11E-10	1.32E-12	4.27E-06	1.07E-08	1.26E-07	1.13E-10	1.70E-02	3.39E-05	1.59E-03	1.32E-05	3.23E-04	1.95E-02
60%	1.70E-10	2.01E-12	6.58E-06	1.75E-08	2.02E-07	1.71E-10	2.43E-02	5.10E-05	2.33E-03	1.86E-05	4.81E-04	2.79E-02
65%	2.69E-10	3.34E-12	1.04E-05	2.68E-08	3.16E-07	2.71E-10	3.52E-02	7.30E-05	3.34E-03	2.56E-05	7.03E-04	3.98E-02
70%	4.23E-10	5.28E-12	1.64E-05	4.20E-08	5.09E-07	4.32E-10	4.95E-02	1.04E-04	4.72E-03	3.60E-05	9.62E-04	5.62E-02
75%	6.36E-10	8.24E-12	2.55E-05	6.56E-08	7.92E-07	7.00E-10	6.82E-02	1.57E-04	6.45E-03	5.07E-05	1.33E-03	7.78E-02
80%	1.00E-09	1.24E-11	4.15E-05	1.10E-07	1.29E-06	1.08E-09	8.97E-02	2.24E-04	8.70E-03	6.87E-05	1.82E-03	1.03E-01
85%	1.58E-09	1.97E-11	6.62E-05	1.78E-07	2.17E-06	1.71E-09	1.20E-01	3.31E-04	1.15E-02	9.34E-05	2.52E-03	1.36E-01
90%	2.77E-09	3.49E-11	1.16E-04	3.40E-07	3.80E-06	2.95E-09	1.61E-01	5.07E-04	1.55E-02	1.36E-04	3.72E-03	1.83E-01
95%	5.48E-09	7.28E-11	2.89E-04	8.67E-07	1.01E-05	6.14E-09	2.25E-01	9.41E-04	2.35E-02	2.22E-04	6.02E-03	2.54E-01
Mean	5.13E-07	7.46E-09	3.53E-03	1.11E-05	1.20E-04	5.76E-07	4.98E-02	2.08E-04	5.39E-03	4.77E-05	1.25E-03	5.68E-02
Min	1.08E-14	1.67E-16	3.36E-10	6.82E-13	1.34E-11	5.25E-15	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.07E-07
Max	5.64E-05	1.82E-06	4.67E-01	3.90E-03	2.44E-02	5.69E-05	5.53E-01	1.38E-02	1.60E-01	1.58E-03	3.84E-02	6.38E-01

Environmental Science Division

Argonne National Laboratory

9700 South Cass Avenue, Bldg. 240

Argonne, IL 60439-4847

www.anl.gov

U.S. DEPARTMENT OF
ENERGY

Argonne National Laboratory is a U.S. Department of Energy
laboratory managed by UChicago Argonne, LLC