

---

# Princeton Plasma Physics Laboratory

---

PPPL-

PPPL-



Prepared for the U.S. Department of Energy under Contract DE-AC02-09CH11466.

# Princeton Plasma Physics Laboratory

## Report Disclaimers

---

### Full Legal Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

### Trademark Disclaimer

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors.

---

## PPPL Report Availability

### Princeton Plasma Physics Laboratory:

<http://www.pppl.gov/techreports.cfm>

### Office of Scientific and Technical Information (OSTI):

<http://www.osti.gov/bridge>

---

### Related Links:

[U.S. Department of Energy](#)

[Office of Scientific and Technical Information](#)

[Fusion Links](#)

# Self-Similar Nonlinear Dynamical Solutions for One-Component Nonneutral Plasma in a Time-Dependent Linear Focusing Field

Hong Qin<sup>1,2</sup> and Ronald C. Davidson<sup>1</sup>

<sup>1</sup>*Plasma Physics Laboratory, Princeton University, Princeton, NJ 08543*

<sup>2</sup>*Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China*

## Abstract

In a linear trap confining a one-component nonneutral plasma, the external focusing force is a linear function of the configuration coordinates and/or the velocity coordinates. Linear traps include the classical Paul trap and the Penning trap, as well as the newly proposed rotating-radio-frequency traps and the Möbius accelerator. This paper describes a class of self-similar nonlinear solutions of nonneutral plasma in general time-dependent linear focusing devices, with self-consistent electrostatic field. This class of nonlinear solutions includes many known solutions as special cases.

PACS numbers: 52.27.Jt, 37.10.Ty

Nonneutral plasmas are often confined in traps with external focusing fields, such as the Paul trap [1, 2] and the Penning trap [3–8]. For most of the often-used traps, the external focusing forces are linear functions of the configuration coordinates and/or the velocity coordinates. For example, in a Paul trap the external focusing force in the transverse direction is proportional to the transverse displacement from the trap axis, whereas in a Penning trap the transverse focusing force is proportional to the transverse velocity. (Strictly speaking, this linearity is valid near the axis for real devices.) We call these types of traps *linear* traps. The strength of the external focusing field is generally allowed to vary with time. In many cases, such as the Paul trap, it is necessary to have time-dependent focusing fields to provide transverse confinement. Obviously, the quadrupole and solenoidal focusing lattices in particle accelerators [9] are also linear focusing devices. Recently, new types of traps, such as rotating-radio-frequency ion traps [10], and new types of focusing lattices, such as the Möbius accelerator [11], have been proposed. The main feature of these new devices is that the focusing force components in different directions are linearly coupled, which offers advantages in terms of stability and focusing strength over standard traps and focusing lattices. Yet, they all fit into the category of *linear* focusing devices. In this paper, we describe a class of self-similar nonlinear dynamical solutions of nonneutral plasmas in general linear focusing devices, with self-consistent electrostatic potential generated by the oscillating one-component plasma. The starting point of the present study is the set of macroscopic fluid equations with self-consistent electric field, which model the nonlinear dynamics of nonneutral plasmas. The class of nonlinear dynamical solutions admitted by the fluid equations includes many of the known modes in linear focusing systems as special cases, such as the well-known transverse envelope oscillations of a charged particle bunch in focusing lattices of accelerators and store rings, and the equilibrium solution of a cold nonneutral plasma in a time-independent Penning trap. It also includes new collective oscillation modes that haven't been reported before. As an example, a nonlinear collective oscillating mode in a time-dependent Penning trap is identified.

Collective dynamics of nonneutral plasmas are of considerable practical importance [2, 12–20]. Previous approaches for investigating collective dynamics of a nonneutral plasma typically first find an equilibrium solution, then analyze the evolution of linear perturbations relative to the equilibrium. An excellent example is the linear eigenmode analysis by Dubin [13, 14, 16, 17] and Bollinger [2]. Dubin [14] also developed a class of self-similar solution for

the penning trap [Detailed discussion is given after Eq.(11). Compared with the classical studies, the analysis presented here examines a new class of nonlinear modes that are applicable to time-dependent linear focusing devices, including Paul traps, and Penning traps with time-dependent confining magnetic field  $B_0(t)\mathbf{e}_z$ , where there exists no quasi-steady equilibrium state ( $\partial/\partial t = 0$ ) for the plasma. It also applies to the periodic focusing lattice with quadrupole and solenoidal magnets in accelerators and storage rings. In this sense, the results presented here are intended to extend our fundamental understanding of time-dependent linear focusing devices by a considerable amount. It is a new result that many of the fundamental modes in these important devices can be united through the theoretical development reported here.

We model the dynamics of a one-component nonneutral plasma in an applied linear focusing field including the effects of the self-generated electrostatic field,  $\mathbf{E} = -\nabla\varphi$ , by the following set of macroscopic fluid equations:

$$\frac{\partial n}{\partial t} + \nabla \cdot (n\mathbf{v}) = 0, \quad (1)$$

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} + \frac{q}{m} \nabla \varphi + \frac{\nabla P}{mn} + \kappa_1(t) \cdot \mathbf{x} + \kappa_2(t) \cdot \mathbf{v} = \mathbf{0}, \quad (2)$$

$$\left( \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla \right) \left( \frac{P}{n^\gamma} \right) = 0, \quad (3)$$

$$\nabla^2 \varphi = -4\pi qn, \quad (4)$$

where  $-\kappa_1(t) \cdot \mathbf{x}$  is the applied focusing force proportional to the displacement  $\mathbf{x}$ , and  $-\kappa_2(t) \cdot \mathbf{v}$  is the focusing force proportional to the average flow velocity  $\mathbf{v}$ . Here,  $q$  and  $m$  are the particle charge and mass, respectively,  $n(\mathbf{x}, t)$  is the particle number density,  $\mathbf{v}(\mathbf{x}, t)$  is the average flow velocity, and  $\varphi(\mathbf{x}, t)$  is the space-charge potential generated by the charged particles. The conducting boundaries are assumed to be far away, *i.e.*,  $|\mathbf{x}_w| \rightarrow \infty$ . The time-dependent tensors  $\kappa_1(t)$  and  $\kappa_2(t)$  include all of the known linear focusing forces as special cases. In a Paul trap, the focusing coefficients are

$$\kappa_1(t) = \text{Diag}[\kappa_{xx}(t), \kappa_{yy}(t), \kappa_{zz}(t)], \quad \kappa_2(t) = 0, \quad (5)$$

Here,  $\text{Diag}[a, b, c]$  denotes the  $3 \times 3$  diagonal matrix with diagonal components  $a$ ,  $b$ , and  $c$ . For a standard Paul trap, the transverse field is a quadrupole potential, and  $\kappa_{xx}(t) = -\kappa_{yy}(t)$  are the (oscillatory) transverse focusing coefficients, and the focusing coefficient  $\kappa_{zz}(t) > 0$  provides longitudinal confinement of the particles in the  $z$ -direction. [In Eq.(5), we do not

generally require that  $\kappa_{xx}(t) = -\kappa_{yy}(t)$ . The meaning of Paul trap here is a bit more general than the standard convention.] In a Penning trap, transverse confinement is provided by a uniform axial magnetic field  $B_0(t) \mathbf{e}_z$ , and  $\kappa_1(t)$  and  $\kappa_2(t)$  are given by

$$\kappa_1(t) = \begin{pmatrix} -\frac{1}{2}\omega_z^2(t) & -\Omega'(t) & 0 \\ \Omega'(t) & -\frac{1}{2}\omega_z^2(t) & 0 \\ 0 & 0 & \omega_z^2(t) \end{pmatrix}, \quad \kappa_2(t) = \begin{pmatrix} 0 & -2\Omega(t) & 0 \\ 2\Omega(t) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (6)$$

where  $\Omega(t) = qB_0(t)/2mc$  is the Larmor frequency, and  $\omega_z^2(t) > 0$  is the focusing coefficient in the longitudinal direction. The term  $\Omega'(t)$  denotes the time derivative of  $\Omega(t)$ , representing the force due to the inductive electric field when  $B_0(t)$  varies with time. As a simple model, the pressure  $P(\mathbf{x}, t)$  is taken to be a scalar, and the energy balance equation for the fluid is assumed to have the polytropic form in Eq. (3), where  $\gamma$  is the polytropic index. We emphasize that the energy equation adopted in the present analysis is a simple theoretical model to allow analytical progress, as discussed in detail by Dubin [14] and Amiranashvili and Stenflo [21]. The physics conclusions obtained in this paper are not sensitive to this choice of model. To model the moment transport and heating due to viscous effects, we can include a viscous force  $-\mu [\nabla^2 \mathbf{v} - \frac{1}{3} \nabla \nabla \cdot \mathbf{v}] / mn$  on the left-hand side of Eq. (2), and a viscous heating term  $2(\gamma - 1)\mu [e_{ij}e_{ij} - \frac{1}{3}(\nabla \cdot \mathbf{v})^2] / \rho^\gamma$  on the right-hand side of Eq. (3). Here,  $\mu$  is the viscosity coefficient and  $e_{ij} \equiv (\partial v_i / \partial x_j + \partial v_j / \partial x_i) / 2$ . It can be shown that the self-similar solution ansatz also applies to the system of equations with viscosity. However, the dynamical equations for  $P(t) / n^\gamma$  will be different from Eqs. (14) and (15) because of viscous heating effects.

The class of nonlinear collective dynamical solutions of the one-component nonneutral plasma are specified by the following solution structures admitted by the system of fluid-Poisson equations (1)-(4). The density  $n(\mathbf{x}, t)$  is taken to be uniform inside an ellipsoid and zero outside. The shape and orientation of the ellipsoid depend on time  $t$ , and are determined from (see Fig. 1)

$$S(\mathbf{x}, t) = D_{ij}(t) x_i x_j < 1. \quad (7)$$

Inside the ellipsoid, the field quantities are assumed to be of the form

$$n(\mathbf{x}, t) = n(t), \quad v_i(\mathbf{x}, t) = v_{ij}(t) x_j, \quad P(\mathbf{x}, t) = p_0(t) - p_{ij}(t) x_i x_j. \quad (8)$$




Figure 1: The density  $n(\mathbf{x}, t)$  is uniform inside the ellipsoid and zero outside. The shape and orientation of the ellipsoid depend on time, and are determined from  $S(\mathbf{x}, t) = D_{ij}(t) x_i x_j < 1$ .

Here,  $x_i$  ( $i = 1, 2, 3$ ) denotes the three configuration coordinates of the displacement vector  $\mathbf{x}$ , and  $v_i$  ( $i = 1, 2, 3$ ) denotes the three components of the flow velocity vector  $\mathbf{v}$ . There is an implicit summation over the repeated indices in Eq. (8). Equations (7) and (8) specify a particular space-time structure of the dynamical solutions. The velocity vector  $v_i$  is a linear function of the displacement vector  $x_i$ , and the coefficient is a time-dependent tensor  $v_{ij}(t)$ . The pressure  $P$  is given by a time-dependent function  $p_0(t)$  plus a quadratic function of the displacement vector, specified by the symmetric tensor  $p_{ij}(t)$ . We note that the solution requires the density to be constant inside the bunch, and the pressure falls off parabolically towards the edge. As discussed by Dubin [14] and Amiranashvili and Stenflo [21], this is only an approximate model. Dubin [14] chose a special parabolic shape for the pressure such that its mean value corresponds to  $nT$ . Amiranashvili and Stenflo [21] chose a different parabolic shape to obtain the best fit to the particle-in-cell simulation. They showed that a parabolic pressure profile with constant density is a reasonable model for small plasmas.

The ellipsoid  $S(\mathbf{x}, t) = D_{ij}(t) x_i x_j < 1$  is determined self-consistently by the velocity field according to

$$\frac{dS}{dt} = \frac{\partial S}{\partial t} + \mathbf{v} \cdot \frac{\partial S}{\partial \mathbf{x}} = 0. \quad (9)$$

In terms of the matrices  $v$  and  $D$ , Eq. (9) can be expressed as

$$D' + v^T D + Dv = 0, \text{ or } D'_{ij} + v_{li} D_{lj} + D_{il} v_{lj} = 0, \quad (10)$$

where  $v^T$  denotes the transpose of  $v$ . If  $D_{ij}$  is initially symmetric and positive-definite, then the solution for  $D_{ij}(t)$  determined by Eq. (10) is symmetric and positive-definite at all subsequent times. This is because  $v_{li} D_{lj} + D_{il} v_{lj}$  is symmetric, and  $D_{ij}$  cannot cross the boundary  $|D_{ij}| = 0$ , which corresponds to infinitely large density and pressure. In addition, if the initial conditions are chosen such that the pressure  $P(\mathbf{x}, t)$  vanishes at the boundary of the ellipsoid  $S(\mathbf{x}, t) = 1$  at  $t = 0$ , then Eqs. (3) and (9) guarantee that the pressure vanishes at the boundary at all time, *i.e.*,  $P(\mathbf{x}, t)|_{S(\mathbf{x}, t)=1} = 0$  for  $t \geq 0$ .

For given  $D_{ij}$  and total number of charged particles  $N$ , the solution to Poisson's equation (4) with boundary condition of  $\varphi(|\mathbf{x}| \rightarrow \infty) = 0$  is given by [9]

$$\varphi = -\frac{3Nq}{4} \int_0^\infty \frac{ds}{\sqrt{(\lambda_1^2 + s)(\lambda_2^2 + s)(\lambda_3^2 + s)}} \left( 1 - \frac{X^2}{\lambda_1^2 + s} - \frac{Y^2}{\lambda_2^2 + s} - \frac{Z^2}{\lambda_3^2 + s} \right), \quad (11)$$

Here  $\lambda_1^{-2}, \lambda_2^{-2}, \lambda_3^{-2}$  are the three eigenvalues of  $D_{ij}$ , and  $(X, Y, Z)^T = Q^{-1}(x, y, z)^T$  denotes rotated coordinates. The orthogonal matrix  $Q$  is constructed from the three eigenvectors  $\alpha_1, \alpha_2, \alpha_3$  of  $D_{ij}$  as  $Q = (\alpha_1, \alpha_2, \alpha_3)$ . Note that  $\lambda_i$  and  $Q$  are uniquely determined by the matrix  $D_{ij}$ .

The self-similar solution has the form of homogeneous deformations, and the solution ansatz is similar to that constructed by Dubin [14] for a time-independent Penning trap. The fact that homogeneous deformations of flows lead to exact solutions of hydrodynamic equation system is well-known. Such solutions can be traced back to the 18th century and were first obtained for gravitating fluids by Chandrasekhar [22]. In plasma physics, the same ansatz has been used in magnetohydrodynamics, starting from the early work of Kulikovskiy [23]. Similar solutions have been reported for plasma expansion using kinetic equations [24], and an interpretation in terms of Lagrangian variables was given by Amiranashvili, Yu, and Stenflo [25].

Assuming the solution ansatz in Eq. (8), we find that the spatial dependence in the fluid equations (1)-(4) drops out, and the system reduce to a set of ordinary differential equations (ODEs) for the density  $n(t)$ , velocity matrix  $v_{ij}(t)$ , pressure matrix  $p_{ij}(t)$  and  $p_0(t)$  given

by

$$n'(t) + nTr(v) = 0, \quad (12)$$

$$v' + vv - \frac{q}{m}QEQ^{-1} + \kappa_1 + \kappa_2v - \frac{2p}{mn} = 0, \quad (13)$$

$$p' + v^T p + pv - \gamma p \frac{n'}{n} = 0, \quad (14)$$

$$\left(\frac{p_0}{n^\gamma}\right)' = 0. \quad (15)$$

Here,  $Tr(v)$  denotes the trace of  $v$ , and  $E$  is the matrix representation of the self-electric field, which expresses the electric field in  $(X, Y, Z)$  coordinates as  $E_{ij}X_j$ . From Eq. (11), we obtain

$$E_{ij} = \text{Diag}(E_1, E_2, E_3),$$

$$E_1 = \frac{3Nq}{2\lambda_1^3}G\left(\frac{\lambda_2}{\lambda_1}, \frac{\lambda_3}{\lambda_1}\right), E_2 = \frac{3Nq}{2\lambda_2^3}G\left(\frac{\lambda_1}{\lambda_2}, \frac{\lambda_3}{\lambda_2}\right), E_3 = \frac{3Nq}{2\lambda_3^3}G\left(\frac{\lambda_1}{\lambda_3}, \frac{\lambda_2}{\lambda_3}\right),$$

$$G(u, v) \equiv \int_0^\infty \frac{ds}{\sqrt{(1+s)(u^2+s)(v^2+s)}}. \quad (16)$$

Note that the space-charge force  $qQEQ^{-1}/m$  in Eq. (13) is uniquely determined by the matrix  $D_{ij}$ .

Equations (10), and (12)-(14) form a system of ODEs for  $D_{ij}$ ,  $n$ ,  $v_{ij}$ , and  $p_{ij}$ , which determines a class of nonlinear dynamical solutions of the one-component plasma in the external focusing field. We do not treat Eq. (15) as an independent dynamical equation, because Eq. (15) can be integrated directly to give  $p_0(t)/n^\gamma(t) = \text{const}$ . Eq. (15) determines how the internal energy changes in response to density variations. In general, this class of solutions has twenty-two time-dependent variables. If the pressure matrix  $p_{ij}$  and density  $n$  are solved for in terms of  $v$  and  $D$ , and we do not count them as independent fields, then the system has fifteen time-dependent variables.

For certain forms of applied focusing fields, Eqs. (10) and (12)-(14) admit solutions with reduced dimensions. For example, in the time-dependent Paul trap described by Eq. (5), the system obviously admits solutions with  $Q = I$ , where  $I$  is the identity matrix, and diagonal solutions with  $v_{ij}$ ,  $p_{ij}$ , and  $D_{ij}$ ,

$$v_{ij} = \text{Diag}[u_1, u_2, u_3], p_{ij} = \text{Diag}[p_1, p_2, p_3], D_{ij} = \text{Diag}[D_1, D_2, D_3]. \quad (17)$$

The ODE system for  $n$ ,  $u_i$ ,  $p_i$  and  $D_i$  is then given by

$$n'(t) + n \sum_{i=1}^3 u_i = 0, \quad (18)$$

$$u_i' + u_i^2 - \frac{q}{m} E_i + \kappa_{ii} - \frac{2p_i}{mn} = 0, \quad (19)$$

$$p_i' + 2u_i p_i - \gamma p_i \frac{n'}{n} = 0, \quad (20)$$

$$D_i' + 2u_i D_i = 0. \quad (21)$$

In the above equations, there is no summation over a repeated index  $i$ .

The dynamics in three dimensions are coupled through the space-charge potential and density. From Eq. (21), we obtain  $u_i = \lambda_i'/\lambda_i$ , where  $\lambda_i = 1/\sqrt{D_i}$ . Then Eq. (19) reduces to an equation for  $\lambda_i$ ,

$$\lambda_i'' + \kappa_{ii} \lambda_i - \frac{q}{m} E_i \lambda_i - \frac{2p_i \lambda_i}{mn} = 0. \quad (22)$$

Equation (22) has a similar form to the familiar envelope equation for charged particle beams in a periodic focusing lattice. To see this, let's consider the special case where  $\gamma = 2$ ,  $\kappa_{xx} = \kappa_{yy} = \kappa_r$ , and the beam cross-section is circular, *i.e.*,  $\lambda_1 = \lambda_2 = r_b$ . We further assume that the ellipsoid is very long, *i.e.*,  $\lambda_1 = \lambda_2 = r_b \ll \lambda_3$ , and the focusing in the  $z$ -direction is sufficiently weak that the charge bunch is uniform in the  $z$ -direction over a scale-length comparable to  $r_b$ . Then  $E_1 = E_2 = 3Nq/(\lambda_3 2r_b^2)$  [9], where  $N = 4\pi n r_b^2 \lambda_3 / 3 = \text{const.}$  is the total number of particles in the long charge bunch. From Eq. (18), we obtain  $n = N_l / \pi r_b^2$ , where  $N_l \equiv 3N/4\lambda_3$  represents the line density. Equation (20) can be integrated to give  $p_i \lambda_i^2 n^{-2} = \text{const.}$  Making use of Eq. (15), we obtain the pressure solution as

$$P = p_0(t) \left( 1 - \frac{x^2}{\lambda_1^2} - \frac{y^2}{\lambda_2^2} - \frac{z^2}{\lambda_3^2} \right), \quad (23)$$

which vanishes on the plasma boundary. The equations for  $r_b = \lambda_1 = \lambda_2$  then becomes

$$r_b'' + \kappa_r r_b - \frac{K_b}{r_b^2} - \frac{\varepsilon^2}{r_b^3} = 0. \quad (24)$$

Here,  $K_b \equiv 2N_l q^2 / m$  and  $\varepsilon^2 \equiv 2p_r r_b^2 n^{-2} N_l / m$  represent the self-field perveance and the transverse emittance-squared. Note that the emittance is constructed from the constant of the motion  $p_r r_b^2 n^{-2}$ , and the line density  $N_l$ , which is approximately constant for a long charge bunch. The envelope equation (24) is identical to Eq. (6.60) in Ref. [9].

Another interesting example is the time-dependent Penning trap given by Eq. (6). It is a well-known fact that in a Penning trap, the single-particle transverse equations of motion

transform to uncoupled linear oscillator equations in a frame rotating with the instantaneous Larmor frequency  $\Omega(t)$  [26]. The rotation matrix is given by

$$R(\theta) = \begin{pmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \theta = -\int_0^t \Omega(t) dt.$$

Here, we show that the macroscopic fluid dynamics also enjoys this desirable transformation property. The density  $n$ , velocity matrix  $v_{ij}$ , pressure matrix  $p_{ij}$ , and shape matrix  $D_{ij}$  in the laboratory frame are transformed to their counterparts  $\bar{n}$ ,  $\bar{v}_{ij}$ ,  $\bar{p}_{ij}$ ,  $\bar{D}_{ij}$  in the rotating frame as

$$n = \bar{n}, \quad D = \bar{D}R(\theta), \quad p = R(-\theta)\bar{p}R(\theta), \quad (25)$$

$$v = R(-\theta)\bar{v}R(-\theta) + \Omega \frac{dR(\theta)}{d\theta} R(\theta), \quad (26)$$

Substituting Eqs. (25) and (26) into Eqs. (1) and (4), we find that the  $\bar{n}$ ,  $\bar{v}_{ij}$ ,  $\bar{p}_{ij}$ , and  $\bar{D}_{ij}$  satisfy the following equations in the rotating frame

$$\bar{n}'(t) + \bar{n}Tr(\bar{v}) = 0, \quad (27)$$

$$\bar{v}' + \bar{v}\bar{v} - \frac{q}{m}\bar{Q}\bar{E}\bar{Q}^{-1} + \bar{\kappa}_1 - \frac{2\bar{p}}{m\bar{n}} = 0, \quad (28)$$

$$\bar{p}' + \bar{v}^T\bar{p} + \bar{p}\bar{v} - \gamma\bar{p}\frac{\bar{n}'}{\bar{n}} = 0, \quad (29)$$

$$\bar{D}' + \bar{v}^T\bar{D} + \bar{D}\bar{v} = 0. \quad (30)$$

Here, the transformed focusing matrix  $\bar{\kappa}_1$  is diagonal,

$$\bar{\kappa}_1 = \text{Diag}[\Omega^2(t) - \frac{1}{2}\omega_z^2(t), \Omega^2(t) - \frac{1}{2}\omega_z^2(t), \omega_z^2(t)], \quad (31)$$

and there is no  $\bar{\kappa}_2$  term in the rotating frame. This is similar in form to the case of a Paul trap in the laboratory frame. The difference is that the (1, 1) and the (2, 2) components of  $\bar{\kappa}_{1ij}$  are the same, whereas in the case of a standard Paul trap,  $\kappa_{xx} = -\kappa_{yy}$ . Because of this, Eqs. (27)-(30) admit diagonal solutions of the form in Eq. (17) with  $\bar{Q} = I$ ,  $\bar{p}_1 = \bar{p}_2 = p_r$ , and  $\bar{D}_1 = \bar{D}_2 = 1/r_b^2$ . Equations (27) and (29) can be integrated to give  $p_r r_b^2/n^\gamma = \text{const.}$  and  $p_z z_b^2/n^\gamma = \text{const.}$ , or equivalently,  $p_r r_b^{2+2\gamma} z_b^\gamma = \text{const.}$  and  $p_z r_b^{2\gamma} z_b^{2+\gamma} = \text{const.}$ , where use is made of  $n^\gamma r_b^{2\gamma} z_b^\gamma = (3N/4\pi)^\gamma = \text{const.}$ . The corresponding nonlinear envelope equations

for  $r_b$  and  $z_b$  are given by

$$r_b'' + \left( \Omega^2 - \frac{1}{2} \omega_z^2(t) \right) r_b - \frac{3Nq}{2r_b^2} G \left( 1, \frac{z_b}{r_b} \right) - \frac{\varepsilon_{r,\gamma}^2}{r_b^{2\gamma-1} z_b^{\gamma-1}} = 0, \quad (32)$$

$$z_b'' + \omega_z^2 z_b - \frac{3Nq}{2z_b^2} G \left( \frac{r_b}{z_b}, \frac{r_b}{z_b} \right) - \frac{\varepsilon_{z,\gamma}^2}{r_b^{2\gamma-2} z_b^\gamma} = 0, \quad (33)$$

where  $\varepsilon_{r,\gamma}^2 \equiv 8\pi p_r r_b^{2\gamma+2} z_b^\gamma / 3mN$  and  $\varepsilon_{z,\gamma}^2 \equiv 8\pi p_z r_b^{2\gamma} z_b^{2+\gamma} / 3mN$  are two constants of the motion. Here,  $p_z = \bar{p}_3$  and  $\bar{D}_3 = 1/z_b^2$ . For a time-dependent Penning trap,  $\Omega$  and  $\omega_z$  are time-dependent, and the nonlinear dynamical solutions are described by Eqs. (32) and (33).

It can be shown that Eq. (32) reduces exactly to Eq. (24) when  $\gamma = 2$  and the transverse focusing is weak and the plasma ellipsoid is long, *i.e.*,  $z_b \gg r_b$ .

When  $\Omega$  and  $\omega_z$  are time-independent, Eqs. (27)-(30) possess another class of stationary ( $\partial/\partial t = 0$ ) solutions with  $\bar{Q} = I$ ,  $\bar{D} = \text{Diag}[1/r_b^2, 1/r_b^2, 1/z_b^2] = \text{const.}$ ,  $\bar{p} = \text{Diag}[p_r, p_r, p_z] = \text{const.}$ , and

$$v = \begin{pmatrix} 0 & \omega_r & 0 \\ -\omega_r & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \text{const.} \quad (34)$$

In this case, Eqs. (27)-(30) reduce to

$$\omega_r^2 + \frac{qE_r}{m} - \Omega^2 + \frac{1}{2} \omega_z^2(t) + \frac{2p_r}{mn} = 0, \quad (35)$$

$$\frac{qE_z}{m} - \omega_z^2 + \frac{2p_z}{mn} = 0. \quad (36)$$

When the nonneutral plasma ellipsoid is long in the  $z$ -direction and the transverse pressure  $p_r$  is negligibly small, Eq. (35) recovers to the well-known equilibrium radial force-balance equation for a cold nonneutral plasma column in a Penning trap (in the un-rotated laboratory frame) [26] as a special case with  $qE_r/m = 3Nq^2/2z_b r_b^2 m = \omega_p^2/2$ . For other types of linear focusing devices, such as rotating-radio-frequency traps and the Möbius accelerator, the nonlinear dynamical solutions described by Eqs. (8)-(14) can be calculated in a straightforward manner, using the particular  $\kappa_1$  and  $\kappa_2$  for each trap. Because of page limitations, these applications will be presented in a future paper.

In conclusion, a class of nonlinear dynamical solutions of a one-component nonneutral plasma admitted by the macroscopic warm-fluid equations with self-consistent electrostatic potential has been identified. The space-time structure of the solutions has the self-similar form specified in Eqs. (7) and (8). It includes many of the known modes in linear focusing

systems as special cases, as well as new nonlinear collective oscillation modes which haven't been reported before.

This research was supported by the U.S. Department of Energy (DE-AC02-09CH111466) and the National Natural Science Foundation of China (NSFC-11075162).

- 
- [1] W. Paul and H. Steinwede, *Zeitschrift für Naturforschung* **8A**, 448 (1953).
  - [2] J. Bollinger, D. Heinzen, F. Moore, W. Itano, and DJ, *Physical Review A* **48**, 525 (1993).
  - [3] F. M. Penning, *Physica (Utrecht)* **3**, 873 (1936).
  - [4] J. Malmberg and T. O'Neil, *Physical Review Letters* **39**, 1333 (1977).
  - [5] T. O'Neil, *Physics of Fluids* **23**, 2216 (1980).
  - [6] J. Bollinger and D. Wineland, *Physical Review Letters* **53**, 348 (1984).
  - [7] C. Driscoll, J. Malmberg, and K. Fine, *Physical Review Letters* **60**, 1290 (1988).
  - [8] J. Fajans and L. Friedland, *American Journal of Physics* **69**, 1096 (2001).
  - [9] R. C. Davidson and H. Qin, *Physics of Intense Charged Particle Beams in High Energy Accelerators* (Imperial College Press and World Scientific, 2001).
  - [10] T. Hasegawa, M. Jensen, and J. Bollinger, *Physical Review A* **71**, 1 (2005).
  - [11] R. Talman, *Physical Review Letters* **74**, 1590 (1995).
  - [12] L. Turner, *Physics of Fluids* **30**, 3196 (1987).
  - [13] D. Dubin, *Physical Review Letters* **66**, 2076 (1991).
  - [14] D. Dubin, *Physics of Fluids B* **5**, 295 (1993).
  - [15] L. Turner and J. M. Finn, *Physics of Plasmas* **2**, 1378 (1995).
  - [16] D. Dubin and T. O'Neil, *Reviews of Modern Physics* **71**, 87 (1999).
  - [17] D. Dubin, *Physics of Plasmas* **12**, 042107 (2005).
  - [18] N. D. Naumov, *Plasma Physics Reports* **30**, 269 (2004).
  - [19] A. R. Karimov, L. Stenflo, and M. Y. Yu, *Physics of Plasmas* **16**, 102303 (2009).
  - [20] A. Olivetti, J. Barré, B. Marcos, F. Bouchet, and R. Kaiser, *Physical Review Letters* **103**, 1 (2009).
  - [21] S. Amiranashvili and L. Stenflo, *Physical Review E* **67** (2003).
  - [22] S. Chandrasekhar, *Ellipsoidal Figures of Equilibrium* (Yale University Press, 1968).
  - [23] A. G. Kulikovskiy, *Doklady Akad. Nauk. SSSR N.S.* **117**, 199 (1957).

- [24] D. Dorozhkina and V. Semenov, *Physical Review Letters* **81**, 2691 (1998).
- [25] S. Amiranashvili, M. Y. Yu, and L. Stenflo, *Physics of Plasmas* **10**, 1239 (2003).
- [26] R. C. Davidson, *Physics of Nonneutral Plasmas* (Imperial College Press and World Scientific, 2001), p. 41, 2nd ed.



The Princeton Plasma Physics Laboratory is operated  
by Princeton University under contract  
with the U.S. Department of Energy.

Information Services  
Princeton Plasma Physics Laboratory  
P.O. Box 451  
Princeton, NJ 08543

Phone: 609-243-2245  
Fax: 609-243-2751  
e-mail: [pppl\\_info@pppl.gov](mailto:pppl_info@pppl.gov)  
Internet Address: <http://www.pppl.gov>