Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review Letters

JETS OF NUCLEAR MATTER FROM HIGH ENERGY HEAVY
ION COLLISIONS

H. Stöcker, L.P. Csernai, G. Graebner, G. Buchwald,
H. Kruse, R.Y. Cusson, J.A. Maruhn, and W. Greiner

February 1981

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 6 1981

LIBRARY AND
DOCUMENTS SECTION

LBL-11774 Rev. c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Jets of Nuclear Matter from High Energy
Heavy Ion Collisions*

H. Stöcker

Gesellschaft für Schwerionenforschung (GSI)
D6100 Darmstadt, Germany, and
Nuclear Science Division, Lawrence Berkeley Laboratory
University of California, Berkeley, CA 94720, U.S.A.

and

L. P. Csernai,** G. Graebner, G. Buchwald, H. Kruse,†
R. Y. Cusson,†† J. A. Maruhn, and W. Greiner
Institut für Theoretische Physik
D6000 Frankfurt am Main, Germany

Abstract: The nuclear fluid dynamical model with final thermal breakup is used to study the reactions $^{20}\text{Ne} + ^{238}\text{U}$ and $^{40}\text{Ar} + ^{40}\text{Ca}$ at $E_{\text{LAB}} = 390 \text{ MeV/n}$. The calculated double differential cross sections $d^2\sigma/d\Omega dE$ are in agreement with recent experimental data. However, it is shown that the azimuthal dependence of the triple differential distributions $d^3\sigma/dE d\cos\theta d\phi$, to be obtained from 4π exclusive experiments with single event analysis, can yield considerably deeper insight into the collision process and allow for snapshots of the reactions. Strongly correlated jets of nuclear matter are predicted.

*This work was supported by Bundesministerium für Forschung und Technologie (BMFT), Deutscher Akademischer Austausch Dienst (DAAD) and the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract W-7405-ENG-48.

**On leave from CRIP, Budapest, Hungary

† Present address: Gesellschaft für Schwerionenforschung, Darmstadt, W. Germany

††Permanent address: Physics Department, Duke University, Durham, North Carolina

Recent measurements of proton cross sections with high associated multiplicities [1] provide further evidence for predominant sideways emission of fragments [2] from high-energy heavy ion collisions. Several features of the data [1] are in qualitative agreement with three-dimensional (3D) nuclear fluid dynamical (NFD) calculations [3]. This is of great importance as it might indicate the presence of strong compression effects in nuclear collisions inherent in the NFD model [2-6].

Here we analyze the reactions $^{20}\text{Ne} + ^{238}\text{U}$ and $^{40}\text{Ar} + ^{40}\text{Ca}$ at $E_{\text{LAB}} = 390 \text{ MeV/n}$ [1]. We emphasize the great importance of 4π exclusive measurements for the understanding of the basic reaction mechanisms. Jets of nuclear matter are predicted. The NFD model we use includes [6] compressibility, an effective density dependent n-n interaction, Fermi pressure, internal excitations, a realistic treatment of the nuclear binding and surface properties, and the final thermal breakup [7-9]. Nuclear viscosity and thermal conductivity [8-11] have been neglected as in all previous 3D calculations because of numerical expenditure; the thermal energy is produced by shock heating [3-6]. Azimuthally (ϕ) averaged double differential particle cross sections $d^2\sigma/d\Omega dE$ and ϕ -dependent triple differential distributions $d^3\sigma/d\cos\theta d\phi dE$ are calculated boosting the Maxwell Boltzmann momentum distribution in each fluid element by the corresponding collective flow velocity into the laboratory [8-9]. This is done in a late stage in the reaction, so that the final distributions depend only negligibly on the exact value of the breakup density $\rho_{\text{BU}} \sim 0.5 \rho_0$ [7-9].

In order to estimate the differences between light and heavy fragment production, distributions for free nucleons (p,n) are also calculated [9]. This is done by shifting down the Fermi distributions of the nucleons in each fluid element according to the corresponding binding energy and evaporating the high momentum tails having internal energy $\epsilon \geq mc^2$ only [9]. These prescriptions lead to reasonable agreement with calculations assuming chemical equilibrium to describe the breakup into various light nuclei [12].

Figure 1 shows contours of the calculated double differential invariant nucleon cross section in the rapidity-transverse momentum plane $y_{||}/p_T$ together with the multiplicity selected observed proton inclusive cross sections [1] for the reaction ^{20}Ne (393 MeV/n) + ^{238}U . The theoretical results for $b = 4$ fm are consistent with the high multiplicity selected experimental data with an estimated mean $\bar{b} \approx 4$ fm [1]. In particular, the "emission pattern" [1], i.e., the positions of the maximum cross section at given p_T , and its shift with p_T agrees rather well (dashed lines). In agreement with the data [1], the calculated angular distributions of low energy particles ($E_{\text{Kin}} \sim 10$ MeV) show a broad maximum at large angles ($\theta_{\text{LAB}} \approx 75^\circ$), which shifts forward with increasing nucleon energy while the height decreases. A detailed quantitative comparison is presently inhibited, since the data result from a complicated folding of impact parameter dependent multiplicity distributions [1]. This can not easily be incorporated into the theoretical calculations, e.g., by simply assuming a sharp

cut-off impact parameter $b = 4.5$ fm [5]. The qualitative features of the calculated ϕ averaged contour plots, however, do not change dramatically with impact parameter, once violent collisions with $b < 6$ fm are selected. The emission pattern shifts only towards slightly larger angles when b is decreased. This means, unfortunately, that ϕ averaged double differential cross sections are of limited value for obtaining information on the reaction dynamics and on the nuclear equation of state [5,6]. Therefore, we next consider whether the azimuthal dependence of the differential cross sections, to be obtained from 4π exclusive experiments with single event analysis [13-16], can provide more specific dynamical information.

Figure 2 shows the triple differential cross sections $d^3\sigma/d\cos\theta d\phi dE$ in the scattering plane, i.e., the $y_{||}/p_T$ plane at $\phi = 0^\circ/180^\circ$, for the reaction $^{20}\text{Ne}(393 \text{ MeV/n}) + ^{238}\text{U}$ at various impact parameters b . For head-on collisions, $b = 0$ fm, the two maxima at $p_T/m \sim 0.1-0.2$ indicate the azimuthally symmetric large angle sideways emission of matter with temperatures $T < 10$ MeV, caused by the strong shockwave [2,3]. At intermediate impact parameters, a considerable azimuthal asymmetry appears. A strong maximum at smaller transverse and longitudinal velocities indicates the presence of a large chunk of cold, slowly moving matter, namely the target residue at $\phi = 180^\circ$. A flat local maximum in the projectile hemisphere ($\phi = 0^\circ$) at larger p_T and $y_{||}$ reflects some sideways deflected fragments of the beam particles. The spread of the maxima in ϕ depends strongly on b ; for intermediate b it is on the order of $\Delta\phi \sim 40^\circ$. The

apparent large collective transverse and longitudinal momentum transfer (the bounce off process [3,6,13,15]) results from the high pressure in the "participant" head shock zone, pushing the nuclear residues apart to opposite directions ($\Delta\phi \sim 180^\circ$). This process is of great importance, as it intimately connects the momentum transfer to be observed in bounce off events with the quantity of central interest, namely the nuclear equation of state $P(\rho, T)$ [17]. At large impact parameters ($b > 6$ fm) the invariant cross sections peak more closely to the initial projectile and target momenta. Maxima at finite p_T are found even in the azimuthally averaged particle cross sections.

The symmetric system ^{40}Ar (388 MeV/n) + ^{40}Ca shows a similarly strong forward-backward peaked distribution at large b . The ϕ -averaged double differential invariant nucleon cross sections exhibit a structureless "fireball" distribution at smaller impact parameters ($b \sim 2$ fm), i.e. the contour lines in the $y_{\parallel}/p_T$ plane are circles centered around y_{cm} . However, we observe in the triple differential cross sections again a symmetric two jet structure at finite p_T being superimposed on the broad thermal "fireball" background. The connection between the two jet maxima in the scattering plane defines the jet axis, which rotates around y_{cm} as a function of impact parameter b . The angle, $\theta_{\text{jet}}(b)$, of the jet axis with respect to the beam direction is given in Table 1. It is directly connected to the "emission pattern" [1] in the cm frame, thus giving the direction of the main fluid flow.

The mean momentum perpendicular to the jet axis may serve as a measure for the temperature in the system. The distance between the maxima, i.e. the mean momentum along the jet axis, will be influenced by the transport properties of the matter. For example, a large viscosity slows down the motion in the jet direction. Thus an event shape analysis [21] in terms of thrust and sphericity might be appropriate for 4π exclusive experiments.

For head-on collisions of equal nuclei, the compression in the shock zone is maximized, and most of the matter participates in the strong compression. The jet patterns give way to an azimuthally symmetric disk of nuclear matter, expanding towards 90° in the cm system [18]. It eventually results in doughnut-shaped triple differential cross sections symmetric around the beam axis. The strong collective transverse matter flow [18] with large mean velocity $p_T/m \sim 0.4$ is caused by the high pressure in the shock region, in analogy to the intermediate impact parameters. Remnants from the squeeze-out can still be seen at small, but finite impact parameters, $b \lesssim 2$ fm, thus giving rise to additional out-of-plane jet structures at $\theta_{cm} \sim 90^\circ$, $\phi \sim 90^\circ$ (four jet events [19]).

The "chemical" composition in various regions of phase space is another observable of great importance, which may yield information on the nuclear equation of state [20]. Since the temperature in the shock zone is much higher than the temperatures in the projectile and target remnants, we expect the predominant emission of free nucleons from the "fireball". Hence, the jet structure is much more pronounced for bound

nuclei (e.g. α , ^{12}C). The reason for that is twofold: clusters are heavier and thus have a smaller thermal velocity than nucleons at the same temperature, so that their distributions are not broadened as much by thermal effects. In addition, they are produced preferentially in the "cold" parts of the matter distribution, which for the bounce off is just the central part of the projectile and target residues. This effect is seen in Fig. 2, $b = 4$ fm, comparing the distribution of all particles with the free nucleon distribution. The latter is less structured and resembles a large fraction of the thermal background in the particle distribution. The distribution of the bound nuclei may be roughly estimated by subtracting the free nucleon distribution from that of all particles. One is thus led to the conclusion that the collective effects should be observable most clearly in, e.g., the α particle distributions. It might be useful to note that first indications for predominant sideways emission were observed using α -particle detection [2] and that recent data on deuteron and triton emission [15] show sharper sideways maxima than the protons [1]. Therefore, a detailed calculation of the cluster formation, e.g. in a chemical equilibrium model [12], is required for a quantitative comparison with future 4π exclusive experiments.

In conclusion, we have shown that triple differential particle cross sections offer a unique tool for the investigation of the complicated reaction dynamics in high-energy heavy ion collisions. Exclusive 4π experiments with single event analysis and special emphasis on production and correlations of

the emitted fragments can provide snapshots of bulk motion, mass and temperature distributions and energy flux in violent nuclear collisions.

We gratefully acknowledge fruitful discussions with M. Gyulassy, H.H. Gutbrod, S. Nagamiya, A. Poskanzer, H. Pugh, A. Sandoval, R. Stock, I. Tanihata, and K. Wolf.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Dept. of Energy under Contract W-7405-ENG-48.

References

- 1) R. Stock, et al., Phys. Rev. Lett. 44 (1980) 1243.
- 2) H.G. Baumgardt, E. Schopper, J. Physics G5 (1979) L231;
H.G. Baumgardt, et al., Z. Physik A237 (1975) 359.
- 3) H. Stöcker, J.A. Maruhn, W. Greiner, Phys. Rev. Lett. 44
(1980) 725.
- 4) A.A. Amsden, F.H. Harlow, J.R. Nix, Phys. Rev. C15 (1977)
2059;
F.H. Harlow, A.A. Amsden, J.R. Nix, J. Comp. Phys. 20
(1976) 119;
A.A. Amsden, G.T. Bertsch, F.H. Harlow, J.R. Nix, Phys.
Rev. Lett. 35 (1975) 905.
- 5) G.F. Bertsch, A.A. Amsden, Phys. Rev. C18 (1978) 1293;
J.R. Nix, D. Strottman, LA-UR-80-2068, to be published.
- 6) H. Stöcker, J.A. Maruhn, W. Greiner, Z. Physik A293 (1979)
173, and Prog. Part. Nucl. Phys. 4 (1980) 133.
- 7) P. Danielewicz, Nucl. Phys. A314 (1979) 465;
J. Bondorf, S. Garpman, J. Zimanyi, Nucl. Phys. A296
(1978) 320;
A. Sierk, J.R. Nix, Phys. Rev. C22 (1980) 1920.
- 8) L.P. Csernai, H.W. Barz, Z. Physik A296 (1980) 173.
- 9) G. Buchwald, et al., University Frankfurt Preprint 42-1980.
- 10) H. Stöcker, J.A. Maruhn, W. Greiner, Phys. Lett. 81B
(1979) 303.

- 11) H.H. Tang, C. Y. Wong, Phys. Rev. C21 (1980) 1846;
C.Y. Wong, J.A. Maruhn, T.S. Welton, Nucl. Phys. A253
(1975) 469.
- 12) L.P. Csernai, P. Subramanian, G. Buchwald, G. Graebner,
J.A. Maruhn, W. Greiner, H. Stöcker, ITPUF-38-80, to be
published.;
J. Kapusta and D. Strottman, LA-UR-80-1831, to be
published.
- 13) H.H. Gutbrod, Proc. Symp. Rel. Heavy Ions, Darmstadt,
March 1978 (GSI-P5-1978 p. 124);
W.G. Meyer, H.H. Gutbrod, Ch. Lukner, A. Sandoval,
Phys. Rev. C22 (1980) 179.
- 14) A.I. Warwick, et al., to be published.
- 15) H.H. Gutbrod, LBL-11123, Proc. Sem. High Energy Nucl.
Int., K. Nakai (ed.), Hakone, Japan, July 1980, p. 93.
- 16) R. Renfordt, et al., to be published.
- 17) H. Stöcker and B. Müller, GSI-Preprint 80-4, to be
published;
H. Stöcker, M. Gyulassy and J. Boguta, LBL-12095, to be
published.
- 18) W. Scheid, H. Müller, W. Greiner, Phys. Rev. Lett. 32
(1974) 741;
J. Hofmann, W. Scheid, W. Greiner, Nuovo Cim. 33A (1976)
343;
H. Stöcker, R. Y. Cusson, J.A. Maruhn, W. Greiner,
Z. Physik A294 (1980) 125 and A290 (1979) 297;
P. Siemens and J. Rasmussen, Phys. Rev. Lett. 42 (1979) 880.

- 19) H. Stöcker, invited talk at the Int. Conf. Extreme States
in Nucl. Syst., Dresden, February 1980, GSI Preprint 80-9.
- 20) H. Stöcker, W. Greiner, W. Scheid, Z. Physik A286 (1978)
121;
P. J. Siemens, J. I. Kapusta, Phys. Rev. Lett. 43 (1979)
1486.
- 21) For a review see G. Wolf, DESY 80/13, to be published.

Table 1: The jet angle, θ_{jet} , relative to the beam axis as a function of impact parameter at $E_{\text{LAB}} = 390 \text{ MeV/n}$

b[fm]	0	2	3	4	6	8
Ne + U	90°	68°	--	45°	26°	7°
Ar + Ca	90°	44°	28°	17°	2°	--

Figure Captions

Fig. 1) Contour diagrams in the plane of transverse momentum p_T in units of $p_T/(m_N c)$ and rapidity $y_{||} = 1/2 \ln [(E + p_{||})/(E - p_{||})]$ for the reaction $^{20}\text{Ne} + ^{238}\text{U}$ at $E_{\text{LAB}} = 393 \text{ MeV/n}$.

Compare the calculated invariant nucleon cross section $\frac{1}{p} d^2\sigma/d\Omega dE$ [$\text{n/MeV}^2 \text{ sr c}^{-1}$] at $b = 4 \text{ fm}$ with the measured proton cross sections for peripheral collisions (low associated multiplicities M , thin lines) and "central" collisions with an estimated mean $\bar{b} \approx 4 \text{ fm}$ (high M , thick lines) [1]. Note that in the upper and lower part of the figure, one and five lines respectively are drawn for each decade of cross section.

Fig. 2) Azimuthally dependent triple differential particle cross sections $\frac{1}{p} d^3\sigma/dE d\cos\theta d\phi$ [$\text{n/MeV}^2 \text{ sr c}^{-1}$] at various impact parameters for the same reaction as above. Shaded areas indicate flat local maxima. The lower right frame shows the distribution of free nucleons at $b = 4 \text{ fm}$ to be compared with the distribution of all particles in the lower left frame.

Ne $\rightarrow$ U

$E_{LAB} = 393$ MeV/n

XBL 8011-7415

Fig. 1

XBL 8010-12672

Fig. 2