WHAT IS THE EMITTANCE OF THE INJECTED BEAM?

G. Parzen

January 10, 1988

1

The usual Statement ^{is} ~~is~~ that the ~~the~~ emittance of the injected beam is $\epsilon_x = \epsilon_y = 10$ (normalized) for heavy ions in RHIC.

Tracking studies requires a more precise statement. Tracking requires that a 4-dimensional surface be specified ~~in~~ in x, x', y, y' space that contains 95% of the beam. The tracking studies can then investigate the stability of the particles inside this surface.

A simple way to specify this surface is by

$$\epsilon_T = \epsilon_x(x, x') + \epsilon_y(y, y') = C,$$

where C is a constant chosen so that this surface contains 95% of the beam. One reason for using this expression is that ϵ_T is ^{very} roughly a constant of the motion.

It will be argued below that the proper choice of C is

$$C = 10 (5/3) = 16.7$$

corresponding to the statement that $\epsilon_x = \epsilon_y = 10$.

Assuming for the moment that $C = 16.7$ is the proper choice of C , then in tracking studies where runs ~~are~~ done with $\epsilon_x = \epsilon_y$ then the proper starting emittance is

$$\epsilon_x = \epsilon_y = 8.33$$

If runs are done with $\epsilon_y = 0$, then the starting ϵ_x is

$$\epsilon_x = 16.7$$

$$\epsilon_y = 0$$

These two points are on the surface $\epsilon_T = \text{constant}$ that contains 95% of the beam.

I assume that the statement $\epsilon_x = \epsilon_y = 10$ means that for the projection of the particles on the x, x' plane, ~~the largest~~ 95% of the particles have an ϵ_x which is smaller than $\epsilon_x = 10$, and a similar statement applies to the y, y' plane.

I assume that the distribution $p(x, x', y, y')$ is gaussian with the form

$$p(x, x', y, y') \sim \exp\left(-(\epsilon_x(x, x') + \epsilon_y(y, y'))/\bar{\epsilon}\right) \quad (1)$$

The projection on the xx' plane has the distribution

$$p(x, x') = \int dy dy' p(x, x', y, y') \sim \exp\left(-\epsilon_x(x, x')/\bar{\epsilon}\right) \quad (2)$$

In order for 95% of the particle to ~~smaller~~ have an ϵ_x which is smaller than $\epsilon_x = 10$ then

$$\bar{\epsilon} = 10/3 \quad (3)$$

The fraction of the particles that have a total emittance, $\epsilon_T = \epsilon_x + \epsilon_y$, which is smaller than $\bar{\epsilon}_T$ is given by

$$F(\epsilon_T) = 1 - \exp(-\epsilon_T/\bar{\epsilon}) (1 + \epsilon_T/\bar{\epsilon}). \quad (4)$$

This may be derived from Eq (1) for $\rho(x, x', y, y')$. The choice ϵ_T that includes 95% of the particles is $\epsilon_T = 5 \bar{\epsilon}$ (the actual answer is closer to $4.8 \bar{\epsilon}$).

Thus the ϵ_T that contains 95% of the particles is

$$\epsilon_T = 5 \bar{\epsilon}$$

$$\epsilon_T = 5 \left(\frac{10}{3} \right)$$

$$\epsilon_T = 16.67$$

I wish to thank Harald Hahn
for his suggestion ~~of~~ for the
meaning of the statement that
 $e_x = 10$.