Nuclear Chemistry Research
at
Michigan State University

Annual Report
for
Calendar Year 1969
[AEC Contract No. AT(11-1)-1779]

Wm. C. McHarris, Principal Investigator

LEGAL NOTICE

THIS report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission
A. Makes any warranty or representation, expressed or implied with respect to the accuracy, completeness, or usefulness of the information disclosed in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights, or
B. Assume any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.
As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepared, furnished, or provided access to, any information pursuant to the employment or contract with the Commission, or his subcontract with such contractor.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

N.B. The material contained herein consists of an annual report to the U. S. Atomic Energy Commission on research performed under contract AT(11-1)-1779 and is not to be construed as publication in the open literature. Many of the results are complete, but others are likely to undergo change before final publication takes place. Before quoting from this report, you are requested to obtain permission from the author of the section in question, who will be very happy to supply you with the very latest information we have on the particular subject.

Contents

Preface	v
<i>Part I: EXPERIMENTAL</i>	
1. The Electron Capture Decay of Pb^{200} - R. Doebler	1
2. Search for an Isomeric State in Pb^{200} - R. Doebler	12
3. The Decay of Pb^{201} - R. Doebler	15
4. The Decay of Bi^{205} and States in Pb^{205} - K. Kosanke	24
5. The Electron Capture Decay of 12-Hour Bi^{204} - J. Cross	33
6. The Electron Capture Decay of Gd^{149} - R. Eppley	41
7. The $M4$ Isomer of Gd^{145} - R. Eppley	51
8. The Peculiar Decay of Gd^{145g} - R. Eppley	55
9. Gd^{144} and Gd^{143} - R. Eppley	65
10. Decays of the Two Isomers of Sm^{141} - R. Eppley	67
11. A Search for Dy^{147m} - R. Eppley	74
12. Decay Studies on Tb^{151} - R. Firestone	77
13. Radiations from the Decay of Yb^{167} - R. Goles	79
14. As^{74} Decay and Forbidden Zero-Phonon Transitions - W. McHarris	85
15. The Decays of Fe^{53g} and Fe^{53m} - J. Black	93
16. The Decay of 38.4-Min Zn^{63} - G. Giesler	103
17. The Decay of 9.3-Hour Zn^{62} - G. Giesler	109
18. The α Decay of Es^{252} - W. McHarris	113
19. The (p,t) Reaction on Pr^{141} - R. Goles	116
20. In-Beam Study of the Decay of Sc^{40} to Ca^{40} - J. Black	123
21. The Possibility of β -Delayed α 's from Ga^{62} or Ga^{63} - G. Giesler	124
22. On-Line γ -Ray Spectroscopy - R. Warner	126
<i>Part II: THEORETICAL</i>	
23. Simple Shell-Model Calculations on Three-Quasiparticle States in Pr^{139} - W. McHarris	128
24. $M4$ Transition Probabilities in Nuclei below $N=82$ - R. Eppley	133

Part III: TECHNIQUES AND INSTRUMENTATION

25. Status of the PDP-9 Computer - W. McHarris	137
26. Multianalyzer Programs for the PDP-9 Computer - O. McHarris	140
27. Auxiliary Programs - R. Eppley	141
28. Cyclotron On-Line Mass Separation - K. Kosanke	147
29. Rabbit -- Pneumatic Target System - K. Kosanke and J. Black	153
30. Slow Pulsing of the MSU Cyclotron Beam - J. Black	156
31. Megachannel γ -Ray Spectra - R. Doebler	159
32. Spectrum Analysis Routines - J. Cross and G. Giesler	163
33. A New Method for Plotting Spectra - R. Eppley	169
34. Some Interesting Compton-Scattering Problems Encountered in Ge(Li)-Ge(Li) Coincidence Experiments - G. Giesler	171
35. Calibration of γ -Ray Standards - R. Doebler	175
36. Progress with the Compton-Suppression NaI(Tl) Annulus - K. Kosanke	178
37. Measurement of Low-Energy Conversion Electrons with the MSU $\pi\sqrt{2}$ Iron-Free Spectrometer - B. Johnston and K. Kosanke	183
38. A Szilard-Chalmers Separation for Isomer-Ratio Enhancements - R. Goles	190
39. Preparation of Thin Pr^{141} Target Foils - R. Goles	193
40. A Single-Crystal Ge(Li) Conversion-Coefficient Spectrometer - R. Doebler	195
Personnel	203
Publications Involving Members of the Nuclear Chemistry Group	204

Preface

The year 1969 marks the completion of the first four years of nuclear chemistry research at Michigan State University. A type of research that did not exist here prior to 1965 has by now attracted a large, enthusiastic, and successful group of students. The extent of this can be seen by comparing the increase in both size and sophistication of this year's Annual Report over that of last year's (COO-1779-4).

Our nuclear chemistry research program is centered around the MSU Sector-Focused Cyclotron in the Cyclotron Laboratory, part of the Department of Physics. We have had a very close relationship with the nuclear physicists there, indeed, the best Chemistry-Physics cooperation of which I am aware. Even our primary financial support has come from them, and, although this is an Annual Report to the Chemistry Division, we must state that our major support to date has come from (in decreasing order):

- 1) The MSU Cyclotron Laboratory, under their NSF grant,
- 2) Michigan State University, especially the Department of Chemistry,
- and 3) Our Nuclear Chemistry contract from the AEC.

Thus, this Annual Report should be considered as supplementing and complementing the Cyclotron Laboratory's Annual Report to the NSF, for without their aid very few of the accomplishments reported herein could have been realized.

The reports that follow cover a wide variety of topics and also exhibit a wide variation in their degree of polish and completeness. Some are condensations of papers already in print, some half-finished, some ready for publication, and some even speculative. However, they should give one an overall view of the types of research in which we are working. Nuclear spectroscopy, especially off-line β - γ spectroscopy, still accounts for the major portion, but we have been branching out into many other fields as well. The nominal cutoff date for the reports was 1 November 1969, but since that time we have completed many other experiments that could not be included. For example, we now have a number of good spectra from the $\text{Pr}^{141}(p,t)\text{Pr}^{139}$ reaction and are well on our way toward a complete angular distribution -- slow, because the cross-section is so low that roughly 8 hr is required for each angle. Also, the in-beam γ -ray spectroscopy is well underway and has produced worthwhile re-

sults on some of the Xe isotopes. Most of the projects reported on are not simply isolated experiments but fit into programs of larger overall scope -- an idea of some of these can be obtained by glancing at the Ph.D. thesis topics on p. 203.

One of the problems arising out of our close cooperation with the nuclear physics groups is that it often becomes quite difficult to isolate what "nuclear chemistry has done" from what "nuclear physics has done". One of the most charming aspects of the lab as a whole is that it has not become rigidly compartmentalized, so the amount of interplay and cross-talk between and among individuals and groups is maximal. Thus, when writing a specifically "nuclear chemistry" Annual Report, there is always the danger of taking more credit than one should from the work of others. On some reports we have specifically mentioned the contributions of others outside our group, but such contributions are really much greater and more widespread than can always be acknowledged. We have worked most closely with Dr. W. H. Kelly and his group, consisting of Dr. D. B. Beery (now at Manchester College, North Manchester, Ind.), R. R. Todd, W. B. Chaffee, and S. M. Mustafa -- indeed, you will find one or more of their names on all of our publications and one or more of our names on all of their publications. But for the fact that he is a "physicist" and we are "chemists", rather than all natural philosophers together, he would likely be co-principal investigator on this Annual Report. We also thank the faculty at the Cyclotron Laboratory, all of whom have contributed directly or indirectly to this Report: T. Arnette, S. Austin, W. Benenson, H. Blosser, G. Crawley, A. Galonsky, C. Gruhn, M. Gordon, E. Kashy, H. McManus, B. Freedom, R. Sherr (visiting from Princeton), and B. H. Wildenthal. Special thanks also go to H. Hilbert, the Chief Engineer, R. deForest, the Computer Engineer, and C. Merritt, for the aid in computer operations.

The success of our nuclear chemistry program rests on many foundations: free and ample access to a superior cyclotron, good back-up equipment, a cooperative Physics Department, a supporting Chemistry Department which has decided that nuclear chemistry is at MSU to stay. In the last analysis, however, the prime reason for success rests in the number and quality of our graduate students. In these days when there is a certain reluctance in speculating as to the future of certain sciences and some justified criticism of turning

out too many overtrained scientists, I think the words *quality* and *superior* must not be hid under a veil of false modesty but brought out into the open. If sheer numbers were the main factor, our nuclear chemistry group could easily have become two or three times its present size several years ago. The present graduate students, however, have all proven themselves to be more than capable of becoming much more than competent scientists, and herein lies the real justification for our group. MSU and specifically nuclear chemistry at MSU has proven itself able to attract students, excellent students who are willing to put in the very long hours necessary for a rather severe and demanding experimental discipline, who operate the cyclotron themselves and construct detectors and equipment in addition to gathering and interpreting results, and who are gaining the necessary theoretical as well as experimental knowledge not, perhaps, to become strict theoreticians but to decide that there are many times when they can do a better job of analyzing their own experiments and relating these to physical reality than can someone else. I believe that there is demand for such people and ample justification for the support of programs that train and educate such people -- and that such programs should not take a back seat to purely technological and/or military feats.

The typing of this report was done by Mrs. I. Samra and the cover design, by K. Kosanka. For a closer scrutiny of the cover, you might check p. 151.

East Lansing, Mich.

WCM

17 December 1969

1. The Electron Capture Decay of Pb^{200}

R. Doebler

The decay of 21-h Pb^{200} has been reported on previously [1]; however, during the past year we have revised and added to our data on this isotope.

Pb^{200} was initially produced using 37-MeV protons on natural thallium foil to induce the $\text{Tl}^{203}(p,4n)\text{Pb}^{200}$ reaction. However, with He^3 beams from the MSU sector-focused cyclotron coming into routine use, we switched to the reaction $\text{Hg}^{202}(\text{He}^3,5n)\text{Pb}^{200}$ using a 45-MeV He^3 beam on a target of natural HgO . This reaction has the advantage of reducing the amount of Pb^{203} contamination by a factor of about three. The samples were aged from 1 to 3 days and the lead activity chemically separated from the Tl and Hg contaminants. To reduce the contamination from 26-h Tl^{200} , which was constantly growing in, a new sample, free of Tl, was prepared every hour. This procedure was followed for all spectra taken.

Figure 1 shows the γ -ray singles spectrum obtained in 7 hours using a 2.5% efficient $\text{Ge}(\text{Li})$ detector with a resolution of 2.3-keV FWHM on the 1332-keV γ -ray of Co^{60} . Nineteen γ -transitions belonging to the decay of Pb^{200} were found, having the energies and intensities listed in table 1. Of these, the 155.29-, 193.39-, 348.23-, 377.92-, and 525.54-keV transitions had not been detected in earlier studies [2-4]. The 155.29- and 193.39-keV transitions have only been tentatively assigned to the decay of Pb^{200} .

The results of integral and anti-coincidence experiments using an 8"x8" $\text{NaI}(\text{Tl})$ annulus and a 7-cm³ $\text{Ge}(\text{Li})$ detector (figures 2 and 3) show that all γ -rays are involved in strong coincidence except the 450.6-, 525.5-, and the 605.4-keV transitions. These are therefore assumed to be unfed ground state transitions; although, assuming our tentative placement of the 155.29-keV transition is correct, the 450.6-keV level may be slightly fed from the 605.4-keV level. The results of these experiments are given in table 1.

The placement of the transitions in the level scheme was based on data obtained from a two-dimensional coincidence experiment of 2048x

2048 channels using two large volume Ge(Li) detectors. The two-dimensional coincidence experiment and some results from our experiment on Pb^{200} are described in ref. [5]. A summary of the coincidence data obtained from this experiment is given in table 2. In addition to its usefulness in helping to place the transitions in the decay scheme, the two-dimensional coincidence experiment also aided in resolving the doublet peak composed of the 289.24- and 289.92-keV transitions [5].

The conversion coefficients for most of the transitions were determined using γ -ray intensities from our singles runs and the conversion electron data of Wierhed [2]. In order to normalize the two sets of data we assumed that the 147-keV γ -transition from the 0- first excited state to the 2- ground state [6] was a pure $E2$ transition and that the theoretical conversion coefficients of Hager and Seltzer [7] are good approximations for this $E2$ transition. Table 3 contains the transition data along with the multipolarity assignments we propose for those transitions where both conversion electron and γ -ray intensities are available. In only one of these transitions is there a significant amount of $M1, E2$ mixing, and this is for the 257.2-keV transition, which is $\sim 60 \pm 15\%$ $E2$. This lack of $E2$ admixture would seem to indicate that the levels in Tl^{200} populated in the decay of Pb^{200} are relatively "pure" single-particle states with little contribution from collective states.

From the conversion coefficients calculated in table 3 and the theoretical conversion coefficients of Hager and Seltzer where experimental values were not available, the total transition intensities as well as the K -conversion intensities were calculated. From the measured K x-ray intensity, K -conversion intensity, and the K fluorescent yield, the total β -feeding relative intensity to the ground state of the daughter was determined to be $< 0.5\%$, corresponding to a $\log ft > 8$. The total β -feeding relative intensity for decay to each excited daughter state was then calculated using the total transition intensities assuming feeding to the ground state to be zero. $\log ft$ values based on these relative intensities were calculated; these appear in the decay scheme shown in figure 4.

The $\log ft$ values for the excited states in Tl^{200} range from

6.1 to 8.0 indicating that these β -transitions are either allowed or first-forbidden transitions. Based on the multipolarity assignments and the 2- ground state, it appears that all the beta transitions to the excited states are first forbidden. Assuming these beta transitions to be first forbidden and using the multipolarity assignments we were able to assign spins and parities to all the levels in Tl^{200} populated in the decay of Pb^{200} . These are given in the decay scheme along with the total transition intensities in percent.

References

- [1] R. Doeblner, 1968 Annual Report, Nuclear Chemistry Group M.S.U. (COO-1779-4).
- [2] B. Wirhed and C. J. Herrlander, Ark. Fys. 23, 355 (1962).
- [3] K. E. Bergkvist, I. Bergström, C. J. Herrlander, S. Hultberg, H. Slätis, E. Sokolowski, A. H. Wapstra, and T. Wielding, Phil. Mag. 46, 65 (1955).
- [4] T. R. Gerholm, Ark. Fys. 11, 55 (1956).
- [5] For a discussion of the two-dimensional gamma-gamma coincidence experiment, see Sec. 30 of this annual report.
- [6] B. Aström, B. Johansson, and I. Bergström, Ark. Fys. 12, 205 (1957).
- [7] R. S. Hager and E. C. Seltzer, Nucl. Data, 4A, 1 (1968).

Table 1
Energies and Relative Intensities of Gamma Rays
from the Decay of Pb^{200}

Measured energies (keV)	Relative intensities		
	Singles	Integral $\gamma\gamma$ coincidence	Anticoincidence
K x-rays	3156 $\pm$ 350	-	-
109.54 $\pm$ 0.04	14.5 $\pm$ 2.0	370	12
142.28 $\pm$ 0.03	95.1 $\pm$ 5.0	4900	55
147.63 $\pm$ 0.03	1133 $\pm$ 30	21,500	1020
155.29 $\pm$ 0.1	1.4 $\pm$ 0.5	-	-
161.32 $\pm$ 0.04	9.1 $\pm$ 1.0	650	6.6
193.39 $\pm$ 0.1	1.0 $\pm$ 0.4	-	-
235.62 $\pm$ 0.04	129 $\pm$ 4.0	7000	81
257.19 $\pm$ 0.03	134 $\pm$ 4.0	5100	100
268.36 $\pm$ 0.03	119 $\pm$ 5.0	6000	80
289.24 $\pm$ 0.15	32 $\pm$ 10	1950	70
289.92 $\pm$ 0.15	51.6 $\pm$ 10		
302.93 $\pm$ 0.05	5.0 $\pm$ 1.0	125	3.5
315.60 $\pm$ 0.08	6.7 $\pm$ 1.0	260	4.5
348.23 $\pm$ 0.08	4.8 $\pm$ 1.5	130	-
377.92 $\pm$ 0.05	0.8 $\pm$ 0.3	9.7	-
450.56 $\pm$ 0.05	$\approx$ 100	$\approx$ 100	$\approx$ 100
457.80 $\pm$ 0.07	3.5 $\pm$ 0.6	24	2.3
525.54 $\pm$ 0.06	12.6 $\pm$ 1.0	-	12
605.44 $\pm$ 0.06	16.9 $\pm$ 1.2	23	19

Table 2

Results of γ - γ Coincidence Study Using 2-Dimensional Analysis

Gated Energy (keV)	Energies of γ -rays in Coincidence with Gate		
	Strong	Weak	Very Weak
109.5	147,268	235	193
142.3	147,235, 315	161	-
147.6	109,142,235, 268,303,315 457	161,348,378	155
161.3	289.24	147.6	-
235.6	289.92,142 147,257	109	-
257.2	268,348,235	193,315	-
268.4	109,147,257	-	-
289.2	161,235,315	-	-
289.9			
302.9	147	193	-
315.6	142,147,289	257	-
348.2	147,257	-	-
450.5	-	-	155

Table 3
Transition Data for Pb²⁰⁰

Energy (keV)	Photon intensity	Conversion intensity		Experimental conversion coefficient	Theoretical conversion coefficient	Multiple order
109.5	14.5	K	22	5.2	5.7	M1
		L _I	4.2			
		L _{II}	0.42			M1
142.3	95.1	K	69	2.5	2.7	M1
		L _I (L _{II})	14.2	0.51	0.48	M1
		M _I	3.4	0.12	0.095	M1
147.6	1133	K	100	0.30	0.34	E2
		L _I (L _{II})	137	0.41	0.43	E2
		L _{III}	90	0.27	0.27	E2
		M _{II} M _{III}	63	0.19	0.19	E2
161.3	9.06	K	4.5	1.7	1.9	M1
235.6	129	K	26.7	0.70	0.65	M1
		L _I (L _{II})	4.6	0.12	0.11	M1
		M _I	1.35	0.036	0.023	M1
257.2	134	K	10.1	0.26	0.51	M1
					0.091	E2
		L _I (L _{II})	2.8	0.071	0.092	M1
					0.048	E2
		L _{III}	0.5	0.013	0.00063	M1
			0.019	E2		
268.4	119	K	18.4	0.53	0.46	M1
		L _I (L _{II})	3.1	0.089	0.077	M1
		M _I	0.65	0.018	0.016	M1
289.2	32	K	3.7	0.39	0.37	M1
289.9	51.6	K	5.4	0.35	0.37	M1
302.9	5.03	K	0.70	0.47	0.32	M1
		L _I (L _{II})	0.12	0.081	0.057	M1

Table 3. continued

315.6	6.69	K	0.68	0.35	0.29	$M1$
		$L_I(L_{II})$	0.20	0.10	0.049	$M1$
450.6	100	K	4.4	0.15	0.11	$M1$
		$L_I(L_{II})$	0.76	0.026	0.019	$M1$
		M	0.16	0.0054	0.0044	$M1$
457.8	3.49	K	0.40	0.39	0.11	$M1$
605.3	16.9	K	0.24	0.046	0.051	$M1$
		L	0.04	0.008	0.009	$M1$

Fig. 1-1. γ -ray singles spectrum of Pb^{200} obtained in 7 h with a 2.5% Ge(Li) detector. A chemical separation to remove Tl was performed each hour. Peaks not labeled come primarily from Pb^{201} and Tl^{200} .

Fig. 1-2. Pb^{200} integral coincidence spectrum. All γ -rays above the x rays were included in the gate. Only Pb^{200} γ -rays are labelled.

Fig. 1-3. Pb^{200} anticoincidence spectrum. Only Pb^{200} γ -rays are labelled; the others result from other Pb isotopes or from Tl^{200} .

Fig. 1-4. ^{200}Pb decay scheme.

2. Search for an Isomeric State in Pb^{200}

R. Doebler

The decay schemes of the neutron deficient lead isotopes are of great interest because these isotopes lie near the doubly magic shell of 82 protons and 126 neutrons and therefore provide good experimental tests of predictions made by many particle shell model calculations. They are also of interest because all of the odd-mass and several of the even-mass isotopes exist in isomeric forms. These isomeric states arise from the presence of low-lying single-particle $i_{13/2}$ states either alone, as in the case of the odd-mass isomers, or in combination with other single-particle wave functions to give high-spin states which are lowered in energy so they lie in a region of much lower spin states, as in the case of Pb^{202m} , Pb^{204m} , and Pb^{206m} .

Figure 1 shows the basic aspects of the even-mass lead isomers. From the apparent systematic behavior of these isomers we expected to find an isomeric state in Pb^{200} at about 2150 keV. This isomeric state has been the object of previous investigations [1,2] with no positive results. Bergkvist, et al., [2] set an upper limit of one hour on the half-life of Pb^{200m} .

We began our search for Pb^{200m} late in 1967. Our first attempts involved bombarding Tl^{203} with protons to induce the reaction $\text{Tl}^{203}(p,4n)\text{Pb}^{200m}$. This type of reaction has been used to produce the other isomers in good yield. Bombardments were made at proton energies of 31, 36, and 40 MeV. Four 1024-channel Ge(Li) γ -ray spectra containing the results of numerous bombardments were obtained at each energy, each sample being counted consecutively for 15-minute periods. These experiments failed to produce any evidence of a new isomer and we set an upper limit of ≈ 1 minute on the half life of any new activity.

We next attempted to populate an isomeric state in Pb^{200} from the decay of Bi^{200} . This procedure was similar to that used successfully in our study of 6.1-second Pb^{203m} [3]. 35-minute Bi^{200} has an estimated

electron capture decay energy of 6.5 MeV and a ground state spin of 7 and should therefore populate high spin states in Pb^{200} . Bi^{200} was produced by bombarding separated isotope Tl^{203} with 50-MeV τ 's to induce the reaction $\text{Tl}^{203}(\tau, 6n)\text{Bi}^{200}$. The Bi^{200} was chemically separated from the target and loaded onto an anion-exchange column. The Pb^{200} activity was eluted with 0.3M HCl periodically and counted using a 7-cm³ Ge(Li) detector. The rate of elution was varied for several different runs, the fastest rate being several drops every 5 seconds. However, we failed to detect any evidence of an isomeric state in Pb^{200} longer than ≈ 1 second in these experiments.

We are now in the process of performing delayed coincidence experiments on Bi^{200} decay in the millisecond range similar to those used to measure the 4-msec half-life of Pb^{205m} [4].

References

- [1] E. K. Hyde, I. Perlman, and G. T. Seaborg, *The Nuclear Properties of the Heavy Elements*, Prentice Hall, Inc., Englewood Cliffs, New Jersey (1964).
- [2] K. E. Bergkvist, I. Bergström, G. J. Herrlander, S. Hultberg, H. Slätis, E. Sokolowski, A. H. Wapstra, and T. Wielding, *Phil. Mag.* 46, 65 (1955).
- [3] R. E. Doebler, Wm. C. McHarris, and C. R. Gruhn, *Nucl. Phys.* A120, 489 (1968).
- [4] I. Bergström, E.C.O. Bonacalya, M. Perez, A. Jech, and P. Thieberger, *Nucl. Inst. and Meth.* 8, 151 (1960).

Fig. 2-1. Systematics and decay schemes of the known even-even Pb isomeric states.

3. The Decay of Pb^{201}

R. Doebler

As part of our investigation into the decay systematics of the neutron deficient lead isotopes, we have recently initiated a study of the decay of 9.4-h Pb^{201} to Tl^{201} . Although this isotope had been extensively studied using $\text{NaI}(\text{Tl})$ scintillation and magnetic conversion electron spectrometers [1-4], the decay scheme was far from being complete.

Pb^{201} was produced using 27-MeV protons to induce the $\text{Tl}^{203}(p,3n)\text{Pb}^{201}$ reaction on natural thallium foil. The samples were aged from 12 to 24 hours to allow the 3.62-h Pb^{202m} and 67-min Pb^{204m} activities to decay away. The lead activity was then chemically separated from the Tl and Hg contaminants.

Figure 1 shows a typical γ -ray singles spectrum obtained using a 7-cm³ $\text{Ge}(\text{Li})$ detector with a resolution of 3.0-keV FWHM on the 1332-keV γ -ray of Co^{60} . To reduce contamination from 73-h Tl^{201} which was constantly growing in, a new sample, free of Tl, was prepared every four hours. Although our investigation has not been completed, we have observed 46 γ -ray transitions belonging to the decay of Pb^{201} . These are listed in table 1 along with their relative intensities.

The results of our anticoincidence and integral coincidence experiments using an 8"x8" $\text{NaI}(\text{Tl})$ annulus in conjunction with a 7-cm³ $\text{Ge}(\text{Li})$ detector are also summarized in table 1. From these experiments it appears that the γ -rays of energies 1098.52, 1157.49, 1238.82, 1277.11, 1401.30, 1445.83, 1479.85, 1617.2, and 1672.02 keV are predominantly unfed ground state transitions, while the 692.41- and 331.15-keV transitions are partially fed ground state transitions. The placement of most of the γ -rays in the decay scheme was aided greatly by a two-dimensional γ - γ coincidence experiment of 4096x4096 channels using two large volume $\text{Ge}(\text{Li})$ detectors [5]. The results of this experiment are summarized in table 2. Based on these results and sum and difference relationships, we have proposed additional energy levels in Tl^{201} at 1134.81-, 1330.28-,

and 1420.01-keV.

Using the conversion electron data of Aasa et al. [4] and the γ -ray intensities from our singles runs, the conversion coefficients for 22 transitions were calculated. The two sets of data were normalized using the measured K -conversion coefficients of Pettersson et al. [3] for the 331-, 361-, and 585-keV transitions. The resulting conversion coefficients are listed in table 3 along with the theoretical conversion coefficients for $M1$ and $E2$ multipolarities obtained by interpolation from the tables of Sliv and Band [6]. It is worth noting that out of eighteen possible $M1$ transitions only six have a significant amount of $E2$ mixing. This lack of $M1, E2$ mixing has also been noted in the decay of Pb^{200} [7].

The decay scheme as presently known is shown in figure 2. The spin and parity assignments are based on our transition multipolarities and assume a $\frac{1}{2}^+$ ground state for Tl^{201} . They are in agreement with the assignments of Aasa, et al., [4] for those levels where they made assignments. While Pb^{201} decays primarily by electron capture, Anderson, et al. [8], have reported some β^+ decay. In our two-dimensional analysis we gated on the weak 511-keV peak, which we assumed arose from β^+ annihilation, and observed weak coincidences with transitions de-exiting the 1157.49-keV level. Based on this observation we propose some β^+ population of the 1157.49-keV level.

References

- [1] A. H. Wapstra, D. Maeder, G. J. Nijgh, and L. Th. M. Ornstein,
Physica 20, 169 (1954).
- [2] K. E. Bergqvist, I. Bergström, C. J. Herrlander, S. Hultberg,
H. Stätis, E. Sokolowski, A. H. Wapstra, and T. Wiedling,
Phil. Mag. 46, 61,65 (1955).
- [3] B. G. Pettersson, T. R. Gerholm, Z. Grabowski, and B. van Nooijen,
Nucl. Phys. 24, 196 (1961).
- [4] E. Aasa, T. Sundström, O. Bergman, J. Lindskog, and K. Sevier,
Arkiv Fysik 27, 133 (1964).
- [5] See section 30 of this Annual Report.
- [6] L. A. Sliv and I. M. Band, in *Alpha-, Beta-, and Gamma-Ray
Spectroscopy*, ed. by K. Siegbahn, North-Holland Pub. Co.,
Amsterdam (1965).
- [7] See section 1 of this Annual Report.
- [8] G. Anderson, E. Arberman, and B. Jung, Ark. Fys. 11, 297 (1957).

Table 1

Energies and Relative Intensities of Gamma-Rays from the Decay of Pb²⁰¹

Measured energies (keV)	Relative Intensities		
	Singles	Anticoincidence	Integral Coincidence
231.8 ± 0.2	4.7 ± 1	-	12.3
241.0 ± 0.1	14 ± 2	-	22
285.15 ± 0.1	13 ± 3	-	14
331.15 ± 0.06	5200 ± 400	3730	2780
345.02 ± 0.08	10 ± 2	-	33
361.25 ± 0.06	680 ± 60	167	1060
381.52 ± 0.1	14.5 ± 2	2.0	33
394.80 ± 0.1	17 ± 5	1.8	40
406.00 ± 0.1	140 ± 25	-	210
465.02 ± 0.1	21 ± 4	-	40
510.8 ± 0.2	3.0 ± 1	-	19
514.3 ± 0.2	10 ± 2	-	27
541.0 ± 0.2	14 ± 2	-	24
546.40 ± 0.1	16 ± 3	-	26
584.60 ± 0.08	224 ± 20	35	364
597.69 ± 0.08	22 ± 2	2.3	33
637.87 ± 0.08	20 ± 2	3.8	30
692.41 ± 0.08	270 ± 20	163	178
708.85 ± 0.10	47 ± 4	7.6	84
753.30 ± 0.20	6.9 ± 2	-	20
767.25 ± 0.08	207 ± 30	60	226
787.30 ± 0.20	40 ± 10	10	-
803.66 ± 0.07	94 ± 10	11	146
826.24 ± 0.20	147 ± 15	40	186
907.67 ± 0.08	380 ± 30	116	382
945.95 ± 0.08	480 ± 40	146	≅ 480
999.23 ± 0.07	40 ± 5	8.4	60
1070.04 ± 0.08	73 ± 8	20	69
1088.82 ± 0.10	56 ± 6	18	58
1098.52 ± 0.07	118 ± 10	112	-
1114.75 ± 0.15	9 ± 2	2	-
1148.75 ± 0.08	49 ± 5	14	49
1157.49 ± 0.08	7.1 ± 2	7	-
1238.82 ± 0.07	72 ± 8	71	-
1277.11 ± 0.07	≅ 100	≅ 100	-
1286.35 ± 0.09	3.5 ± 1	1.5	-
1308.32 ± 0.08	32 ± 4	12	33
1340.88 ± 0.09	27 ± 4	7	28
1401.30 ± 0.08	7.8 ± 2	7	-
1424.11 ± 0.09	6.0 ± 2	1.8	-
1434.10 ± 0.40	0.65 ± 0.2	-	-
1445.83 ± 0.30	2.5 ± 1	2.0	-
1479.85 ± 0.15	9.7 ± 2	10	-
1617.2 ± 0.4	1.0 ± 0.4	1.3	-
1672.02 ± 0.3	1.3 ± 0.8	1.2	-
1764.7 ± 1.0	1.6 ± 0.8	0.25	-

Table 2

Results of γ - γ Coincidence Study Using 2-Dimensional Analysis

Gated Energy (keV)	Energies of γ -rays in coincidence with gate		
	Strong	Weak	Very Weak
232	331,361	585,692 767,787	395,1098
241	331,907	361,585	692,803,946,1070, 1238
285	331	803	361,597,767,946
331	285,345,361, 406,585,767 803,826,907 946,999,1070, 1089,1149,1309, 1343	381,395,709, 787,1115	465,1424
345	331,803		826
361	331,406,585, 709,787	546,596 638,946	465
381	331,361,767	406	692
395	331,361	946	
406	331,361,692		
511	331	826	465,1157
541	331	361	546
546	331,361	692	
585	331,361,692	-	-
597	285,331,361	585	692
638	331,361	-	-
692	331,406,585	597,708	546,787
709	331,361	692	-
767	331	381,541	285,395
787	331,361	-	-
803	331,345,285	-	-
826	331,511	-	-
907	331,241	-	-

Table 2 continued

946	331	-	232
999	331	341	-
1070	331	-	-
1089	331	-	-
1098	-	381	-
1115	-	331	-
1149	331	-	-
1239	-	241	-
1277	-	-	232
1309	331	-	-
1341	331	-	-

Table 3. Transition Data for Pb^{201}

Energy keV	Photon intensity	Conversion intensity	Experimental conversion coefficient	Theoretical conversion coefficients		Multipole order
				M1	E2	
285	13	7.5	0.34	0.37	0.20	M1
331	5200	1000	≈ 0.113	0.25	0.051	M1+E2
345	10	7.4	0.44	0.22	0.047	M1
361	680	240	≈ 0.21	0.20	0.042	M1
381	14.5	4.2	0.17	0.15	0.037	M1
406	140	33	0.14	0.14	0.032	M1
546	16	1.9	0.07	0.066	0.017	M1
585	224	21	0.055	0.060	0.015	M1
692	270	5.0	0.011	0.040	0.011	E2
708	47	4.0	0.05	0.039	0.010	M1
767	207	9	0.026	0.029	0.0083	M1
803	94	1.3	0.008	0.026	0.0076	E2
820	45	0.7	0.009	0.025	0.0075	E2
826	147	2.8	0.011	0.024	0.0074	E2+M1
907	380	10.7	0.016	0.019	0.0062	M1
946	480	12	0.015	0.017	0.0056	M1
1070	73	0.8	0.0065	0.012	0.0045	E2+M1
1098	118	1.2	0.006	0.012	0.0043	E2+M1
1115	9	0.20	0.013	0.011	0.0042	M1
1148	49	0.6	0.007	0.010	0.0040	M1+E2
1239	72	0.7	0.006	0.0084	0.0035	M1+E2
1309	32	0.6	0.011	0.0072	0.0031	M1

Fig. 3-1. Pb²⁰¹ singles γ -ray spectrum taken with a 7-cm³ Ge(Li) detector. A chemical separation was performed every four hours during this run to remove the Tl²⁰¹ that grows in.

Fig. 3-1. Decay scheme of Pb^{201} as presently known from our experiments.

4. The Decay of Bi^{205} and States in Pb^{205}

K. Kosanke

Previous internal conversion studies [1,2] have reported 84 transitions in the decay of Bi^{205} , but with one or more electron intensities [3-5] for only 38 transitions. Of these 84 transitions, we have confirmed 70 directly and have added 26 new transitions, 19 definite, and 7 probable (also several additional possible transitions). Of the 84 transitions originally reported, γ -intensities were reported for only 15 single transitions and 8 collections of transitions. We report transition intensities for 95 transitions and upper limits on 11 of the transitions previously reported but not observed in this study. The current status of the γ -ray and internal conversion studies is listed in table 1 along with multipolarity assignments.

A recent γ -ray spectrum collected using a 2.3-keV (at 1332 keV) $\text{Ge}(\text{Li})$ detector with an efficiency of 2.5% has suggested that much of the remaining uncertainties in intensity of weak peaks and peaks that have been stripped [6,7] will be eliminated shortly.

A series of β^+ feeding spectra were collected using the 2.5% detector mentioned above and an 8"x8" split $\text{NaI}(\text{Tl})$ annulus [2]. The $\text{Ge}(\text{Li})$ detector was placed just below the tunnel on the outside of the annulus and run in triple coincidence (511 keV-511 keV-any) with the halves of the annulus. The results of these runs are summarized in table 2. Corrections to the data for β^+ feeding to the 703.4-keV level in decreasing order of importance are: 703.4-keV coincidence with transitions that can pair produce in the annulus (corrected using the 1770-keV $\rightarrow$ 570-keV coincidence in Bi^{207}), 703.4-keV coincidences with legitimate transitions falling in the 511-keV gates of the annulus (corrected using the 1043-keV $\rightarrow$ 550-keV coincidence of Bi^{205} assuming 0% β^+ feeding), feeding of the 703.4-keV level from higher β^+ fed levels (10% of the feeding to 988-keV level in Bi^{205} goes to the 703.4-keV level). Corrections to the data for β^+ feeding to the 987.5-keV level, in decreasing order of importance are: 987.5-keV coincidences with legitimate transitions falling in the 511-keV gates of the annulus (corrected

using the 1043-keV $\rightarrow$ 550-keV coincidence in Bi²⁰⁵ assuming 0% β^+ feeding, and completely chance triple coincidences which are not very likely. It is possible that there is additional β^+ feeding to a level(s) at $\approx$ 775 keV, but the data are as yet inconclusive [?].

A two dimensional Ge(Li)-Ge(Li) megachannel coincidence experiment [9] was performed. The data have not been completely analyzed; however, approximately half the transitions observed in singles, including several of the new transitions, have been placed in a partial decay scheme (see figure 1) on the basis of the coincidence spectra. Two previously tentative levels [1] at 1593.5 and 1776.0 keV have been conclusively established, and additional tentative levels at 2124.6 and 2488.0 keV are proposed.

Up to this point the various shell-model calculations on states in Pb²⁰⁵ have not been very successful. Part of this has come about because they have been based on incomplete and often erroneous level schemes. After our level scheme is completed (within a few months, depending on the analysis of the remaining two-dimensional data), we hope to be able to make more meaningful comparisons with the various calculations and, if necessary, work on some of them ourselves.

References

- [1] S. H. Vegors, Jr., R. L. Heath, and D. G. Proctor, Nucl. Phys. 48, 230 (1963).
- [2] K. E. Bergkvist and R. Stockendal, Ark. Fys. 27, 339 (1964).
- [3] M. Schmorak, R. Stockendal, J. S. McDonell, and I. Bergström, Nucl. Phys. 2, 193 (1956).
- [4] R. Stockendal and S. Hultberg, Ark. Fys. 15, 33 (1959).
- [5] R. Stockendal, Ark. Fys. 17, 553 (1960).
- [6] Michigan State University, Nuclear Chemistry Annual Report, 1968, p. 30 (COO-1779-4).
- [7] For further discussion and spectra see "Annulus Progress" section 35 in this report.
- [8] C. F. Perdriat, J. H. Brunner, and H. J. Leisi, Helv. Phys. Acta 35, 175 (1962).
- [9] For a discussion of 2-dimensional analysis technique see "Megachannel Analysis" section 31 in this report.

Table 1. Current Status of the Decay of Bi²⁰⁵

Transition Energy		γ-intensity		β intensity K line Ref. 2,3,4 or 5	Multipolarity
This Work	Ref. 1	This Work ^a	Ref. 1		
-	2.3	-	-	-	-
-	26.2	-	-	[5]	M2[5] ^b
-	45.4	-	-	[5]	-
-	90.0	-	-	[5]	E1[5] ^b
115.2	115.1	.29±.03	-	[5]	M1
122.7±1.	122.9	.02±.01	-	-	-
128.9±1.	128.7	.02±.01	-	-	-
185.1	185.1	.31	2.7	40[5]	M1[E2]
205.8	205.8	.11	-	-	-
221.2	221.2	.08	-	-	-
236.0±.2	236.0	.22	-	8[3]	-
237.1±1.	237.2		-		-
-	248.9	<.05	-	-	-
260.6	260.5	4.00	5.7±1.5	160[5]	M1
262.9	262.9	1.2 ±.1		46[5]	M2
282.3	282.3	1.7 ±.2	6.0±1.	41[3]	M1
284.2	284.2	5.5		160[3]	M1
296.4±1.	297.2	.05±.2	-	-	-
310.4	310.4	.34	-	4.3[5]	E3
313.1	313.0	.17	-	4.9[5]	M1
349.6	349.6	1.6	2.7±.7	28[3]	M1
361.5	361.6	.16	-	-	-
-	421.8	<.1	-	-	-
493.7	493.8	1.2	3.5±1.	11[3]	-
498.6	498.8	.62	-	-	-
511.4±.3	511.7	3.1 ±.2	6.3±1.5	22[5]	M1
549.8	550.1	8.9	39.0±5.	6.3[4]	E1
561.6	561.7	.14		-	-
570.9	570.7	13.1		74[3]	M1
574.0	574.1	2.0		-	-
576.4	576.3	.86		-	-
579.8	580.0	16.8		24[3]	E2

Table 1 continued

592.3	-	.32	-	-	-
626.7	627.0	1.8	6.0±.8	6.7[3]	M1
-	634.4	<0.1	-	-	-
688.5	688.7	.62	-	-	-
703.4	703.4	100.	100.	100.	E2
-	710.8	<.1	-	-	-
-	712.6	<.1	-	-	-
717.4	717.5	.84	-	-	-
720.6	720.5	.32	-	-	-
723.6	723.5	.49	-	-	-
729.5	729.5	.18	-	-	-
745.1	745.0	1.8	13.0±3.	2.1[3]	M1+E2
759.1	759.0	3.8		3.1[3]	E2
761.4	761.2	2.3		2.0[3]	E2
770.9	-	.16	-	-	-
780.9	780.8	1.7	-	-	-
787.9	-	.32	-	-	-
795.6	795.6	.40	-	-	-
800.8±.3	-	.61	-	-	-
806.9±.3	806.1	.64	-	-	-
813.8	813.8	1.5	-	-	-
828.2 ^d	828.1	1.2	-	-	-
852.8	-	.16	-	-	-
860.1	859.9	1.4	5.5±2.	-	-
871.8	872.0	1.3		-	-
890.0	890.0	2.1	-	-	-
894.5	894.9	1.9	-	-	-
901.8	-	.42	-	-	-
910.8	910.9	5.4	10.6±2.	9.8[3]	M1
-	913.4	<.25	-	-	-
921.7	-	.18	-	-	-
931.7	-	.13	-	-	-
950.7	-	1.1	-	-	-
971.4	971.9	.89	-	-	-
978.0	-	.23	-	-	-
987.5	987.8	55.3	55.6	27.5[3]	E2

Table 1 continued

1001.9	1002.4	1.8	-	2.1[3]	M1
1013.6 $\pm$.3	1013.8 ^c	.23	-	2.1[2]	M4
1014.6 $\pm$.3	1014.4 ^c	3.2	-	3.3[2]	M1
1031.2	-	.20	-	-	-
1037.4 $\pm$.3	-	.67	-	-	-
1040.1 $\pm$.5	-	.27	-	-	-
1043.7	1043.7	25.8	28.0 $\pm$ 5.	32.5[3]	M1
-	1048.2	<.1	-	-	-
1060.7	-	.26 $\pm$.1	-	-	-
-	1062.7	<.25	-	-	-
1065.8	-	.35 $\pm$.1	-	-	-
1072.3	1072.4	.99	-	-	-
1107.5	-	.34 $\pm$.05	-	-	-
-	1128.1	<.1	-	-	-
1190.0	1190.0	8.1	7.8 $\pm$ 1.5	7.6[3]	M1
1199.8	1199.8	.61	-	-	-
1208.7	1208.3	1.9	-	-	-
1216.3	-	.26	-	-	-
1264.5	1264.7	.88	-	-	-
1276.0	-	.22	-	-	-
1283.7	-	.17	-	-	-
1351.6 $\pm$.5	1351.7	3.25 $\pm$.7	$\approx$ 3.	1.1[3]	M1+E2
1439.3	1439.0	.54 $\pm$.15	-	-	-
1500.2 ^d	1500.0	1.13	4.5 $\pm$ 1.5	1502.5	-
1521.6	-	.68		0.4[3]	-
1551.0	1552.2	4.7		-	-
1563.3	-	.84		-	-
1577.4	-	.47 $\pm$.1	-	-	-
1592.9	-	.32 $\pm$.1	-	-	-
1614.5	1615.4	8.5	7.5 $\pm$ 2.	3.1[3]	M1
1619.4	-	1.1	-	-	-
1717.5 $\pm$ 2.	1721.0	.13 $\pm$.1	-	-	-
1764.5	1766.1	124.	115. $\pm$ 10.	43.6[3]	M1
-	1768.9	<4.		-	-
1776.0	1777.7	14.2		5.2[3]	M1

Table 1 continued

1818.2	-	.18	-	-	-
1861.9	1863.9	23.6	} 36. 5.	2.0[3]	E1
1903.6	1905.9	9.4		0.6[3]	E1
-	1925.1	<.01		-	-
1966.0	-	.04	-	-	-
2565.2	2567.8	.19	-	.029[4]	E3
2606.8	2610.0	.11	-	.016[4]	E3

^aUncertainties listed in energies and intensities found in this work represent uncertainties in the data (i.e., for weak peaks, or for peaks that have been striped from interfering peaks) and should be taken to be in addition to errors of the magnitude of 0.025% on energies and 10% on intensities.

^bMultipolarity assigned on basis of electron intensities only.

^cEnergies from Ref. [2].

^dUnresolved doublet.

Table 2. β^+ feedings in the decay of Bi^{205} .

Energy of Level (keV)	Predicted $\epsilon_K/\beta^+ \rightarrow$ % feeding			This Work	Ref. [8]
703.4	80 ^a	$\rightarrow$	.18%	$.072 \pm .02\%$ ^b	$.058 \pm .01\%$
987.5	300	$\rightarrow$	.018%	$.008 \pm .004\%$ ^c	-
1043.7	500	$\rightarrow$	.006%	<.002	-

^aUsing graphs reproduced in: *Table of Isotopes*, Lederer, Hollander, and Perlman, 6th ed (1968).

^bError limits are rather generous and may be reduced when decay scheme is complete.

^cThe magnitude of the corrections on the feeding to this level is considerably smaller than for feeding to the 703-keV level, accordingly the smaller error limits.

Fig. 4-1. Preliminary decay scheme of ^{205}Bi , showing the present status of our progress on this nuclide. Roughly half of the known transitions are placed with reasonable assurance in this level scheme.

5. The Electron Capture Decay of 12-hour Bi^{204}

J. Cross

The electron capture decay of Bi^{204} ($t_{1/2}=11.6$ h) has long been of considerable interest, since the product of the decay, Pb^{204} was the first even-even nucleus found to have a long-lived metastable state (67 min-2186 keV level). In addition $_{82}\text{Pb}^{204}$ lies at the closed shell of protons ($N=82$) and only four neutrons removed from the $N=126$ closed neutron shell; thus it should be of great importance to compare the theoretical shell model calculations with experimental results.

Previous investigators [1-4] have reported a total of 67 γ -ray transitions, found predominantly by conversion electron spectroscopy. Only 21 of these transitions have been placed in the decay scheme (fig.1); however a few other conclusions have been reached based upon existing data:

- 1) The first and second levels lie at 899 and 1275 keV respectively and have spins and parities of $2+$ and $4+$.
- 2) All γ -ray cascades pass through the 899-keV, $2+$ level with little or no decay going directly to the ground state.
- 3) About 10% of the Bi^{204} decay passes through the 9- isomeric state.
- 4) A large number of populated levels lie above 2000 keV.

Most other investigators have avoided this isotope since the large number of transitions and resulting thousands of sums is staggering. While an absolutely complete decay scheme is almost beyond question, we are in the process of extending the range and scope of the γ -transitions and decay scheme considerably.

The Bi^{204} was produced with excellent yield by bombarding Pb^{206} separated isotope with 29.5-MeV protons from the MSU Sector-Focused Cyclotron to induce the $\text{Pb}^{206}(p,3n)\text{Bi}^{204}$ reaction. The only contaminant of consequence is Bi^{205} ($t_{1/2}=15.3$ d) produced by the $\text{Pb}^{206}(p,2n)$ reaction. However the excitation functions for Pb^{206} with protons indicate that at 29 MeV the ratio

of Bi^{204} to Bi^{205} should be about 3:1. This, coupled with the large difference in $t_{1/2}$ makes the production extremely efficient, and the sample remains adequately active for about 48-60 h after bombardment. We have performed chemical separations (ion-exchanger and nitrate precipitation), but we conclude that at present no chemical separation is required.

The object of our present investigation is three-fold:

- 1) preparation of improved Bi^{204} decay scheme
- 2) preparation of improved Bi^{203} decay scheme
- 3) shell model calculations involving comparison with experimental results.

At present we are completing the energy and intensity calculations of the statistically significant peaks in Bi^{204} . Table 1 lists the energy values and relative intensities (uncorrected for detector efficiency at this time) for these γ -rays. Figure 2 shows a typical γ -ray spectrum obtained with a Ge(Li) detector having a resolution of 2.4 keV at 1332 keV of Co^{60} . The resulting spectra were analyzed by MOIRAE and SAMPO (see section on *Data Analysis Routines*). The major calibration peaks were reproducible to ± 0.10 keV, the 0-1000 keV region to about ± 0.15 keV, the 1000-2000 keV region to about ± 0.3 keV, and from 2000 keV up to about ± 0.50 keV. The intensities of course are much less accurate, with most medium to large peaks falling in the ± 5 -15% region and small peaks up to about $\pm 22\%$.

Using the NaI(Tl) annulus in our laboratory a 511-511- γ triple coincidence was performed to detect the double escape peaks present. By gating on the 511-511 pulses only double escape γ -rays are recorded. More of this technique will be discussed in the section on the annulus later in this annual report.

Our recent work involved a 4 day, 7 million event Ge(Li)-Ge(Li) coincidence experiment using a computerized recording program already used for Nd^{139m} and Pb^{200} [5]. This involves using the Sigma 7 computer to store coincident events from two 4096-channel ADC's on magnetic tape - the data being retrieved and sorted by computer at a later date. This coincidence run should give us the data required to produce a much more accurate decay

sceme, possibly completed as early as December 1969.

Not only is the Bi^{204} study important in itself but also as a prelude to the Bi^{203} study lately proposed in our laboratory. Since Bi^{203} and Bi^{204} have very similar $t_{1/2}$'s it is impractical to attempt any separation of these isotopes. Once Bi^{204} is known however, Bi^{203} can be prepared in the presence of Bi^{204} easily and consequently analyzed.

Bibliography

- [1] A. R. Fritsch and J. M. Hollander, J. Inorg. Nucl. Chem 6, 165 (1958).
- [2] A. R. Fritsch, Univ. Calif. Rad. Lab. Report UCRL-3452, June 1956.
- [3] R. Stockendal, T. Novakov, B. Johansson, and M. Schmorak, Ark. Fys. 14, 65 (1968).
- [4] R. Stockendal, Ark. Fys. 17, 579 (1960).
- [5] Mich. State Univ. Nuclear Chemistry Annual Report, 1968, COO-1779-4.

Table 1. γ -Ray Data for Bi²⁰⁴

Energy (keV)	Photon Intensity*	Energy (keV)	Photon Intensity*
374.74	100.0000	473.50	0.1190
59.28	0.4390	477.75	0.1722
80.40	-	501.76	0.7630
96.54	0.1350	510.67	0.3649
100.74	0.5790	522.22	0.5090
109.38	0.1890	532.72	1.0323
140.80	4.0525	543.37	0.2247
147.36	0.5222	548.74	0.2632
153.52	0.5255	585.02	0.2121
164.92	0.4585	595.13	0.2568
169.83	1.0513	597.83	0.2823
176.17	4.2732	604.73	0.1487
212.70	1.0830	611.88	0.1699
216.11	4.1750	631.88	0.0516
219.46	6.7967	654.88	0.0769
222.15	3.0433	661.55	1.5521
227.46	0.6388	663.43	0.4771
240.40	0.7795	670.70	6.1540
248.91	5.3050	683.39	0.1233
257.52	0.1758	690.75	0.5024
286.32	0.5971	703.91	0.2298
289.25	5.6658	709.13	0.7648
291.35	1.8720	710.48	0.7518
304.45	0.2485	718.43	0.4509
320.85	0.2437	725.15	0.4976
331.07	0.5706	736.07	0.3546
340.59	0.1799	745.23	0.3182
405.82	0.3359	753.78	0.5063
412.24	0.3435	765.37	0.2430
421.55	1.1974	771.31	0.1870
432.53	0.0580	791.16	1.5557
438.46	0.8303	821.13	0.2656
440.34	2.6350	823.05	0.2271
447.08	0.4760	827.62	0.2211
455.92	0.1329	831.95	0.4038
461.70	0.1169	834.10	0.4370
468.23	0.4805	841.06	0.1042

*Relative values -- efficiency curve not yet ready. Also, these are working numbers taken directly from computer output, so the last two significant figures should be disregarded.

Table 1 continued

847.16	0.3787	1232.92	0.1182
881.05	0.1707	1240.04	0.0398
899.15	38.7857	1259.08	0.1120
911.84	9.4400	1261.71	0.0350
918.26	4.0288	1274.81	0.6927
924.16	0.0269	1290.61	0.0482
934.13	0.1327	1299.21	0.0289
941.44	0.0535	1328.22	0.0910
950.33	0.0578	1334.53	0.0716
958.77	0.0680	1347.93	0.0529
964.32	0.1881	1353.35	0.2015
979.45	1.5100	1373.68	0.1047
983.98	20.1757	1380.01	0.0499
990.34	0.3841	1383.62	0.0348
1014.19	0.0259	1414.74	0.2180
1021.93	0.0618	1446.12	0.0185
1027.59	0.0239	1447.52	0.0461
1037.34	0.1256	1453.67	-
1043.63	0.4421	1466.36	0.0750
1049.04	0.0626	1468.82	0.0803
1056.55	0.0575	1475.08	0.0176
1060.17	0.1631	1487.78	0.0430
1064.22	0.3098	1495.39	0.0107
1092.10	0.0305	1517.53	0.0721
1095.08	0.0632	1524.05	0.1711
1102.18	0.1802	1536.54	0.0816
1111.27	0.4799	1569.19	0.0273
1127.50	0.0823	1572.86	0.0435
1132.89	0.2506	1589.45	0.0606
1139.77	0.1802	1607.19	0.0398
1146.54	0.0511	1639.38	0.0620
1157.66	0.1673	1645.62	0.1102
1167.01	0.0621	1652.00	0.1020
1181.24	0.0187	1654.87	0.1019
1198.98	0.0657	1659.07	0.0360
1203.84	0.5593	1665.67	0.0242
1211.74	0.8604	1669.28	0.0169

Table 1 continued

1675.07	0.0139	2100.62	0.0060
1685.87	0.0268	2137.57	0.0037
1689.12	0.1040	2169.44	0.0055
1700.14	0.0451	2172.21	0.0039
1703.32	0.3600	2176.93	0.0064
1709.85	0.0166	2183.68	0.0044
1715.22	0.0246	2251.63	0.0047
1731.53	0.1119	2263.38	0.0374
1749.82	0.0477	2279.37	0.0046
1755.29	0.2075	2312.89	0.0082
1760.97	0.0324	2325.95	0.0047
1778.49	0.0454	2433.32	0.0033
1780.33	0.0550	2450.72	0.0056
1796.91	0.0174	2472.58	0.0048
1803.95	0.0100	2475.63	0.0030
1815.53	0.0263	2517.74	0.0222
1818.18	0.0836	2566.14	0.0097
1826.55	0.0906	2655.12	0.0044
1836.59	0.0109	2668.24	0.0020
1851.00	0.0133	2680.93	0.0363
1856.86	0.0424	2686.82	0.0294
1916.43	0.0078	2721.15	0.0032
1925.87	0.0884	2758.80	0.0091
1941.27	0.1222	2765.26	0.0016
1956.74	0.0581	2794.38	0.0014
1958.10	0.0600	2802.07	0.0017
1964.77	0.0562	2837.33	0.0219
1896.31	0.2076	2854.92	0.0018
1907.23	0.0249	2864.65	0.0010
1988.00	0.0045	2898.01	0.0013
2009.56	0.0105	2947.99	0.0007
2014.09	0.0069	2955.55	0.0017
2048.24	0.0194	2976.89	0.0018
2045.99	0.0105	2990.49	0.0004
2064.19	0.0046	3011.36	0.0023
2084.22	0.0050	3063.00	0.0005
2092.56	0.0050		

PRESENT DECAY SCHEME Bi^{204}

Fig. 5-1. The presently known, very limited decay scheme of Bi^{204} . We have already placed many more levels with a good degree of confidence, but, considering the large number of γ -rays, the final decay scheme should be quite complex.

Fig. 5-2. Bi²⁰⁴ singles γ -ray spectrum taken with a 2.5% Ge(Li) detector having a resolution in this run of about 2.4 keV.

6. The Electron-Capture Decay of Gd^{149}

R. Eppley

Further work has been carried out on the electron-capture decay of Gd^{149} during the past year. As a result several changes have been made in the decay scheme since last year's report [1].

Gd^{149} was produced with the MSU cyclotron by the $Gd^{151}(p,3n)$ reaction using proton beam energies of 25-28 MeV. For this work, natural Bu_2O_3 targets produced the desired activity cleanly enough such that no chemical separations were necessary. Twenty-three γ -rays have been attributed to Gd^{149} decay, with another γ -ray being questionable. Table 1 lists these γ -rays and their relative intensities. Figure 1 is the singles spectrum taken with a 7-cm³ Ge(Li) detector manufactured in this laboratory.

There has been some confusion as to whether the peaks with energies of 963, 993, and 1013 keV should be included in the Gd^{149} decay scheme. These peaks are evident in the spectrum (see figure 1), particularly right after bombardment, but appear to decay with different half-lives than the Gd^{149} .

A new peak has been found at 430.3 keV. This peak is visible in the coincidence spectrum that is gated on the 600-keV region. This is shown in figure 2. This energy fits as a transition from the 1097.3-keV level to the 666.2-keV level.

There is another significant change in the new, modernized decay scheme. Other authors [2,3] have reported a 459.9-keV gamma feeding the 496.4-keV (11/2-) level. In our attempts to confirm this placement, both prompt and delayed coincidence runs were performed. The prompt spectrum, gated on the 396.5-keV γ is shown in figure 3. The delayed spectrum, gated on the 149.6-keV peak which is in prompt coincidence with the 346.5-keV peak, is shown in figure 4. For these spectra a 7-cm³ Ge(Li) detector was used to record the spectra and a 3"x3" NaI(Tl) detector provided the coincidence signals. The resolving time for the coincidence circuit was of the order of 150 ns. The delayed spectrum was obtained by adding 0.5 μ s delay to the spectrum side of the coincidence circuit. This set-up allowed

us to see transitions feeding the 496.4-keV level ($t_{1/2} = 2.3\mu s$). As can be seen in figure 4, the 459.9-keV γ is not present. Thus, our conclusion is that this γ -ray does not proceed to the 496.4-keV level. The 459.9-keV γ is present in prompt integral-coincidence spectra and absent in anti-coincidence spectra, suggesting that it is indeed in prompt coincidence. By placing this transition as proceeding to ground from a level of the same energy, we can account for all the experimental data. The 478.7-keV transition is placed from the 939.1-keV level to the 459.9-keV level. This arrangement is consistent with the coincidence data, energy sums, and transition intensity data.

The experimental internal conversion coefficients were calculated by use of the electron intensities reported by Harmatz and Handley [4]. The photon and electron intensities were normalized by assuming the 346.5-keV transition to be a pure $M2$, with $\alpha_K = 0.20$. The theoretical conversion coefficients are interpolated values from the tables of Hager and Seltzer [5]. Figure 5 shows the theoretical conversion coefficient curves for the K -shell and for the various multipolarities. Superimposed on these curves are the experimentally measured K -shell coefficients. The deduced transition multipolarities are given in Table 1.

The decay scheme presented in figure 6, is based on coincidence and anti-coincidence runs, intensity balances for each level, and sums and differences checks. We are presently writing up the Gd^{149} work and the interpretation of the states in Eu^{149} . This should be complete and submitted to *The Physical Review* before the Christmas vacation. Preprints may be had by asking for "R. E. Eppley, Wm. C. McHarris, and W. H. Kelly, 'The Electron-Capture Decay of Gd^{149} '".

Table 1. Transition Data for Gd^{149}

Energy	Photon Intensity	K-Conversion Intensity ^{a,b,c}	Total Intensity	Exp. α_K	Multipole Order
K x-rays	-	-	-	-	-
149.6±0.2	233.4 ±10.0	124.9	376.6	0.54	M1
214.5±0.6	0.81±0.1	-	0.84	-	(E1)
252.3±0.7	1.08±0.25	0.14	1.23	0.130	M1
260.5±0.3	5.79±0.40	0.15	5.92	2.59 10 ⁻²	E1
272.0±0.2	14.60±0.6	1.75	16.3	1.20 10 ⁻¹	M1
298.5±0.2	126.7 ± 10	10.75	138.1	8.48 10 ⁻²	M1
346.5±0.3	100. ±4	20.0	123.6	0.20	M2
405.5±0.7	3.68±1.46	0.033	3.71	8.97 10 ⁻³	E1
430.3	0.33±0.05	-	0.33	-	(E1)
459.9±0.3	2.42±0.19	0.073	2.49	3.02 10 ⁻²	M1
478.7±0.3	0.95±0.1	0.011	0.96	1.16 10 ⁻²	E2
496.4±0.3	7.21±0.35	0.238	7.49	3.30 10 ⁻²	E3
516.4±0.3	11.05±1.5	0.213	11.3	1.93 10 ⁻²	M1
534.2±0.3	13.20±0.55	0.256	13.5	1.94 10 ⁻²	M1
645.2±0.3	5.93±0.5	0.018	5.94	3.04 10 ⁻³	E1
663.3±0.7	1.09±0.2	0.016	1.10	1.47 10 ⁻²	(M1)
666.2±0.7	3.93±0.6	0.026	3.96	6.62 10 ⁻³	E2
748.2±0.3	34.6 ±4.0	0.071	34.7	2.05 10 ⁻³	E1
788.6±0.3	29.6 ±3.0	0.15	29.8	5.08 10 ⁻³	M1+E2
812.4±0.5	0.55±0.13	-	0.55	-	(M1)
863 1?	0.32±0.10	w	e	e	e
875.8±0.4	0.90±0.11	0.0041	0.91	4.56 10 ⁻³	M1
933.3±0.	2.21±0.5	-	2.22	-	(M1)
939.1±0.	8.99±1.4	0.044	9.03	4.89 10 ⁻³	(M1)
947.7±0.5	3.67±0.64	0.0062	3.68	1.69 10 ⁻³	E2?

^a Intensities from ref. 4.

^b Intensities renormalized such that the α_K (346.5 keV) $\cong$ 0.20.

^c Errors in the relative intensities are reported as being 10% for the most intense peaks and increasing for the weaker ones.

^d For the purpose of arriving at total intensity values, theoretical L and M conversion coefficients were used for the indicated multipolarities. Interpolated values from Hager and Seltzer.

^e Not included in decay scheme.

References

- [1] COO-1779-4, Annual Report for Calendar Year 1968, Nuclear Chemistry Research at Michigan State University.
- [2] H. J. Prask, J. J. Reidy, E. G. Funk, Jr., and J. W. Mihelich, Nucl. Phys. 36, 441 (1962).
- [3] J. M. Jaklevic, E. G. Funk, and J. W. Mihelich, Nucl. Phys. 84, 618 (1966).
- [4] B. Harmatz and T. H. Handley, Nucl. Phys. 81, 481 (1966).
- [5] R. S. Hager and E. C. Seltzer, Nucl. Data, 4A, 1 (1968).

Fig. 6-1. Gd^{149} singles γ -ray spectrum taken with a 7-cm³ Ge(Li) detector.

Fig. 6-2. Gd^{149} coincidence spectrum in which a NaI(Tl) detector was set on the 600-keV region. The very weak 430-keV γ is enhanced greatly in this spectrum.

Fig. 6-3. Gd^{149} coincidence spectrum that aided in determining those γ -rays connected with the isomeric state at 496.4 keV in Eu^{149} . With a prompt gate on the 346.5-keV γ , this spectrum emphasizes the transitions following that state.

Fig. 6-4. Gd^{149} delayed coincidence spectrum that aided in determining which γ -rays feed the isomeric state at 496.4 keV in Eu^{149} . A 0.5- μsec delay was used in the spectrum side of the circuit.

Fig. 6-5. Experimental and theoretical conversion coefficients for the transitions in Eu^{149} following Gd^{149} decay.

7. The $M4$ Isomer of Gd^{145}

R. Eppley

During the past year work has been done to determine the K and L conversion coefficients for Gd^{145m} more precisely than before. As reported in our 1968 Annual Report, where we first reported the discovery of this isomer, the metastable state has an energy of 721.4 ± 0.4 keV and a half-life of 85 ± 3 sec.

Our measurements of the conversion coefficients involved the simultaneous counting of the γ -rays and the conversion electrons from the isomeric transition. The first measurements, as reported previously, were carried out with a 7-cm³ Ge(Li) detector for the γ -ray intensity and a plastic scintillator (Pilot B, 1-1/4" x 1/2") for the conversion-electron intensity. With this set-up the K conversion coefficient was determined to be approximately 0.1.

Our most recent measurements have utilized the same Ge(Li) detector in conjunction with a Si(Li) detector for the conversion-electron measurements. This detector, 1000 μ in thickness, is run at methanol-dry ice temperature with a bias of 200 V. With this detector it is possible to separate the K and L conversion lines. The Gd^{145m+g} electron spectrum is shown in fig. 1, the γ -ray spectrum in fig. 2. In the γ -ray spectrum the unlabelled peaks are transition in Eu^{145} from Gd^{145g} decay.

The results from these intensity measurements are given in Table 1. From the I_{eK}/I_{γ} comparison the transition is unambiguously an $M4$. The I_{eK}/I_{eL} ratio places it as an $M3$ or an $M4$. For the transition to be an $M3$ there would have to be another odd-parity state in the vicinity (assuming one of the involved states to be the $h_{11/2}$ state). From shell-model theory as well as the systematics of the $N=81$ nuclei, this is probably not the case. Therefore, we suggest the transition to be an $M4$ transition from the $h_{11/2}$ state to the $d_{3/2}$ state. As can be seen in the next section of this Annual Report, the $d_{3/2}$ state is probably not the ground state of Gd^{145} , but it lies quite close to the ground state. For more detail on the systematics

and calculations of $N=81$ isomeric transitions, see section 24 of this Annual Report.

Table 1.
Conversion Coefficients for the Gd^{145} Isomeric Transition.

	Experimental	Theoretical ^a			
		E3	E4	M3	M4
I_{e_K}/I_γ	0.120±0.015	0.0109	0.0241	0.0540	0.118
I_{e_K}/I_{e_L}	5.41	3.50	3.53	5.77	4.88

^aR. S. Hager and E. C. Seltzer, Nucl. Data A4, 1 (1968)

Fig. 7-1. Electron spectrum showing conversion lines from the 721.4-keV M_4 isomeric transition in Gd^{145} . This spectrum was taken with a 1000-v Si(Li) detector mounted in a dip-stick cryostat with a 0.1-mil Havar window.

Fig. 7-2. γ -ray spectrum from 34.8-sec Gd^{145m} . The 721.4-keV γ is the $M4$ isomeric transition. The unlabeled γ 's all result from the decay of 21.8-min Gd^{145g} .

8. The Peculiar Decay of Gd^{145g}

R. Eppley

In the past year most of our experimental work on Gd^{145g} has been completed, but some work still remains to be finished on data analysis.

All of our experimental work this past year has been based on the preparation of Gd^{145} by the $Sm^{144}(\tau, 2n)Gd^{145}$ reaction. The target was Sm_2O_3 that was enriched to 95.10% in Sm^{144} . τ beams of 20 MeV were used for the bombardments ($Q = -10.6$ MeV). A variety of experiments have been carried out, including both integral coincidence and anti-coincidence γ -ray measurements, two-dimensional coincidence measurements, 511-511 keV coincidence measurements, and half-life measurements.

For these measurements two γ -ray detectors were employed. One is a 7-cm³ Ge(Li) detector with a resolution of 2.9 keV as measured with Co^{60} . The other detector, also Ge(Li), has a resolution of 2.2 keV and an efficiency of 2.5%. The singles γ -ray spectrum is shown in figure 1. Because of the short half-life of Gd^{145} there is not enough time to carry out a chemical separation, so impurities must be picked out both by energy values and by relative half-lives. Figure 2 is the plot of the spectrum resulting from requiring a 511-511 coincidence in the gating detectors. For this run an 8"x8" NaI(Tl) split-ring annulus was used. Each half of the annulus was used to set a 511-keV window. In order to obtain a gate pulse, both halves of the annulus were required to "see" a 511-keV γ -ray within a resolving time of 100 ns. In this manner the resulting coincidence spectrum should be of transitions either proceeding directly to ground or through cascades to ground that depopulate levels fed by the positron decay of the Gd^{145g} .

Figure 3 shows the spectrum of an integral coincidence run. Again, the coincidence resolving time was 100 ns. The gating pulses were supplied by one-half of the above-mentioned annulus, and the 7-cm³ detector was used for the coincidence spectrum.

Table 1 is a list of the γ -ray energies and intensities which

have been attributed to the decay of Gd^{145g} . This table is still somewhat incomplete, pending the final analysis of the data.

The half-life of the Gd^{145g} decay has been measured with the help of our Sigma-7 computer. This was accomplished by using the program GEORGE. This program allows us to take data through a 50-MHz ADC (maximum of 8192 channels) and to have a "live" display on an 11" scope at the same time. This program allows us to dump the spectrum onto a disk at precise intervals which are determined at the beginning of the run. This dumping process takes significantly less than one second and does not interrupt the counting for any longer than this time. The spectra can then be punched on cards as they are removed from the disk, thereby making the start and stop times of the data acquisition independent of the card punching time. A pulser peak is included in each spectrum so that the dead-time can be accounted for and the appropriate corrections can be applied to the data. There is a great advantage to being able to acquire half-life data for the entire spectrum simultaneously. In this manner the half-life for any or all of the peaks in the spectrum can be measured.

The peaks at 1757.8 keV and 1880.6 keV were used for the present half-life determination. Forty spectra were obtained, taken consecutively, each one representing a two-minute time span. After background subtraction and dead time corrections, the points were least-squared to a straight line on a semi-log plot of counts/time vs time. From an average of these calculations the half-life was determined to be 21.8 ± 0.6 min.

At first glance, Gd^{145} has unusual decay properties. First, there is some evidence that the ground state spin is $1/2$, although other odd-neutron nuclei just below $N=82$ all apparently have a ground state spin of $3/2$. Second, the majority of the transitions are to the two high lying levels of 1757.8 keV and 1880.6 keV. This is indeed unusual in the face of seemingly good single-particle levels in the daughter which would furnish higher decay energies.

Eu^{145} levels have also been investigated by K. S. Toth, et al. [7].

They determined level spins by the $\text{Sm}^{144}(\tau, d)\text{Eu}^{145}$ reaction as well as by looking at Gd^{145} decay. Figure 4 shows the tentative level structure for Eu^{145} as determined in our investigation. This is compared with the level scheme proposed by Toth, et al. Toth lists a level at 716 keV with a spin of $11/2^-$. While Eu^{145} should indeed have a state with this spin, we have not seen evidence for it, as would be expected, considering the low spin of Gd^{145} . There is, of course, a γ -ray transition at 721.4 keV due to the isomeric transition in Gd^{145} .

One explanation for the peculiar behavior of the decay of Gd^{145} is to postulate an intermediate three-quasiparticle configuration. This type of mechanism is depicted in figure 5. The levels shown are not drawn to any energy scale and are for pictorial purposes only. In order for this mechanism to be feasible, two assumptions must be made as to level placement. First, it is necessary for the $h_{11/2}$ proton level to have dropped to the vicinity of the $d_{5/2}$ level. In this manner the ground state wave function for Eu^{145} could have a $11/2^-$ component. Second, the $h_{9/2}$ neutron level from the next oscillator shell must lie at an attainable energy relative to the usual shell model levels in the $50 \leq N \leq 82$ region. We have estimated that this level should be in the vicinity of 2 MeV, thus making the second assumption a reasonable one.

With these assumptions the decay could proceed as

$$\begin{aligned} & |(\pi h_{11/2})^2 (\nu s_{1/2})^{-1}\rangle \\ & \rightarrow |(\pi h_{11/2})^1 (\nu h_{9/2})^1 (\nu s_{1/2})^{-1}\rangle \\ & \rightarrow |(\pi h_{11/2})^1\rangle. \end{aligned}$$

By this formalism we do not mean to imply a ground state wave function simply of the form $|\pi h_{11/2}\rangle^n$, but only that a small portion of the wave function has this character. Assuming the ground state of Gd^{145} to have spin $1/2$, a transition to the lower lying Eu^{145} levels would be several times forbidden by strict shell model selection rules. The transitions would be hindered to the point where the (relatively small) $\pi h_{11/2}$ admixture would govern the decay rate.

Other nuclei in this region have also been shown to decay to three-particle states; for further details see section 22 of this Annual Report.

Reference

- [1] K. S. Toth, E. Newman, B. H. Wildenthal, R. L. Auble, R. M. Gaedke, and M. F. Roche, International Conference on Radioactivity in Nuclear Spectroscopy, Vanderbilt Univ., Aug 11-15, 1969 (p. 7).

Table 1. γ -ray Data for Gd^{145g}

Energy ^{a,b}	Singles Intensity
329.5 $\pm$ 0.2	100.00
511.01	-
736.0 $\pm$ 0.2 (1757.8 D.E.)	135.38
808.5 $\pm$ 0.2	353.58
858.9 $\pm$ 0.3 (1880.6 D.E.)	222.91
893.2	16.55
952.3	85.93
1041.9 $\pm$ 0.2	425.48
1072.0 $\pm$ 0.4	119.74
1247.3 (1757.8 S.E.)	29.72
1273.8	37.07
1369.7 (1880.6 S.E.)	57.85
1473.0 (2493.7 D.E.)	45.46
1567.4	39.82
1600.0	88.50
1620.3 (2641.5 D.E.)	103.78
1719.6	49.79
1757.8	1599.8
1815.9	40.28
1845.3	20.16
1880.6 $\pm$ 0.4	1521.4
1891.2	9.35
2070.1	11.77
2079.1	13.42
2451.7	21.71
2493.7	61.40
2580.4	37.38
2621.7	17.58
2641.5	91.58
2779.2	10.26
3543.6	6.22

^aThis is a preliminary set of energy values.

^bThe energies which are assigned error limits are the ones used to determine the remainder of the energy values in the table.

Fig. 8-1. Singles γ -ray spectrum of 21.8-min Gd^{145g} taken with a 7-cm³ Ge(Li) detector; repeated bombardments were required to obtain this spectrum.

Fig. 8-2. Gd^{145} pair spectrum. The source was placed inside a NaI(Tl) split annulus, each side of which was gated on the 511-keV region; thus, β^+ feeding, double-escape peaks, and transitions fed by either appear in the spectrum.

Fig. 8-3. Gd^{145g} integral coincidence spectrum, using one-half of a NaI(Tl) split annulus and a 7-cm³ Ge(Li) detector.

TENTATIVE LEVEL SCHEME FOR Eu^{145}

-63-

Fig. 8-4. Tentative decay scheme of Gd^{145g} . In (a) we show the decay scheme obtained from our work. This is compared with the levels from ref. [2], which are shown in (b).

Eu^{145}
 $63\text{Eu}82$

Fig. 8-5. Schematic representation of transitions between states in Gd^{145} and Eu^{145} , showing how the ground-state configuration of Gd^{145} might force its decay to go into three-particle states in Eu^{145} .

9. Gd¹⁴⁴ and Gd¹⁴³

R. Eppley

Recently we have begun the study of Gd¹⁴⁴ and Gd¹⁴³. These isotopes are produced with the MSU cyclotron by the Sm¹⁴⁴(α ,3n)Gd¹⁴⁴ ($Q = -20.2$ MeV) and Sm¹⁴⁴(α ,4n)Gd¹⁴³ ($Q = -31.7$ MeV) reactions, respectively.

The first experiments carried out were a series of reactions run with beam energies ranging from 25 MeV to 50 MeV in 5-MeV steps. In this manner we could observe which peaks appeared together at the various energies. At each energy a series of timed spectra were taken such that rough half-lives could be determined. Above 35 MeV the Gd¹⁴⁵ peaks were gone. In the 35-MeV to 50-MeV spectra peaks were present whose half-lives ranged from 2.3 min to 6.0 min. Several were grouped with an average half-life of about 5.1 min. We are investigating whether these peaks are daughters or grand-daughters of the Gd¹⁴⁴ decay. Keller and Münzel [1] have reported a half-life for Gd¹⁴⁴ of 4.5 min with no visible γ -rays. They report γ -rays in Sm¹⁴⁴ of 2.48 MeV, 1.66 MeV, and 0.82 MeV from the decay of 10.5 sec Eu¹⁴⁴. Due to the state of transient equilibrium existing between Gd¹⁴⁴ and Eu¹⁴⁴, the Sm¹⁴⁴ peaks would appear to decay with the same half-life as Gd¹⁴⁴. This will make the assignment of the peaks a difficult proposition, as the half-lives are too short to permit chemical separation.

The half-lives of the peaks in the 2-3 minute range possibly belong to the decay Gd¹⁴³. A complicating factor is that Malan, Münzel and Pfennig [2] report the half-life of Eu¹⁴³ as being 2.3 min. This means that, assuming a 2.5 min half-life for Gd¹⁴³, by 5 min after the end of the bombardment both Gd¹⁴³ and Eu¹⁴³ would be decaying at more or less the same rate.

Other evidence to be sought for the production of Gd¹⁴⁴ and Gd¹⁴³ will be in the study of the various decay products. Confirmation of known transitions will be evidence that the desired Gd activities were indeed initially produced.

References

- [1] K. A. Keller and H. Münzel, Radiochim. Acta 9, 176 (1968).
- [2] H. P. Malan, H. Münzel, and G. Pfennig, Radiochim. Acta 5, 25 (1966).

10. Decays of the Two Isomers of Sm^{141}

R. Eppley (in cooperation with R. Todd of the Cyclotron Laboratory and with R. Warner)

We are investigating the decay of Sm^{141} which we produce by the $\text{Nd}^{142}(\tau, 4n)\text{Sm}^{141}$ reaction. The target material is Nd_2O_3 enriched to 99% in Nd^{142} . A beam energy of 40 MeV is used ($Q = -27.5$ MeV). We have also produced the activity (inadvertently) by the $\text{Sm}^{144}(\tau, \alpha 2n)\text{Sm}^{141}$ reaction.

The decay of Sm^{141} is being investigated in an effort to determine whether or not the daughter nucleus, Pm^{141} , exhibits a level structure expected of three-quasi-particle coupled levels such are found in Pr^{139} [1].

Several authors have reported on investigations of this nuclear species. Table 1 is a summary of some of the results. The results of the half-life measurements for Sm^{141g} are in substantial agreement with our own measured value of 22.1 min as discussed later. The half-life of Sm^{141g} has only been estimated based on systematics of the region and results predicted by the single particle model. As described later we have found a 10-minute component in the spectrum which may be due to Sm^{141g} decay.

Based on a least-squares fit of $M4$ levels, which have been measured in other $N=79$ nuclei, we arrive at an energy of 171 keV for the expected $h_{11/2}$ level. A level at this energy would mean a very small transition intensity which is $M4$ in nature. The reported value of <1% would be correct and thus make difficult the observation of a peak at this energy. We have looked for this transition both in γ -ray spectra and in electron spectra. A large internal conversion coefficient would characterize such a transition and consequently the conversion peak would be prominent for even a fairly weak transition. Our electron spectra show no such conversion peak. This could mean, among other things, that the transition is indeed very weak and/or there is another level of intermediate spin which has come between the $h_{11/2}$ and $d_{3/2}$ levels.

Our half-life measurements were carried out with help from our Sigma-7 computer. Data acquisition was accomplished with GEORGE, a data

taking routine which is programmed to dump the spectrum onto the computer disc at predetermined intervals, erase memory, and begin taking more data. This dumping process takes less than one second to complete so that a negligible error is introduced into the data due to indeterminateness of knowing the starting and stopping time for each interval. A pulser peak is included in each spectrum for determining the necessary dead-time correction of each spectrum. By using this method any or all of the peaks in the species under investigation can be used for half-life determinations. In this manner we have determined the Sm^{141m} half-life to be 22.1 ± 0.5 min.

Figure 1 shows the spectrum collected from several targets, no one of which was counted for more than 30 minutes. A list of energies and relative intensities, which are tentatively assigned to the decay of Sm^{141g+m} is given in table 2. These values are subject to change as more of our data is analyzed.

Thus far we have taken 2-dimensional coincidence, anti-coincidence, 511 keV-511 keV coincidence, prompt and delayed integral coincidence, and electron spectra for this species.

Figure 2 shows the delayed-gate integral spectrum. The 511-keV peak is an anomaly due to the detector arrangement. This has been demonstrated by its absence on subsequent spectra with different detector geometry. This spectrum was taken with a 0.5- μ s gate delay and coincidence resolving time of 100 ns. The enhanced peaks are at 197, 432, and 628 keV. By virtue of the gate delay, these peaks should be from a delayed level in the daughter nucleus. It seems clear, from systematics of the region, that this isomeric level is the $\pi h_{11/2}$. The 432-keV γ -ray has been measured to decay with the 22.1-minute half-life, thus confirming that the 22.1-minute activity is indeed due to the isomeric level in Sm^{141} . These results are summarized in figure 3, which is a representation of the partial decay scheme. The $h_{11/2}$ level in Pm^{141} has an estimated half-life of 150 ns. This value is based on the 40-ns value for this level as measured in Pr^{139} and relating the two nuclides through the single-particle equations for an $M2$ transition.

Figure 4 is the 197 keV gated coincidence spectrum. This is an example of a spectrum taken with the 2-dimensional coincidence set-up. For

this run two Li(Ge) detectors and associated coincidence electronics were used in conjunction with task EVENT in the Sigma-7 computer. The fast coincidence resolving time was on the order of 100 ns.

All evidence points to the 197 keV transition being from the first excited state in Pm^{141} . The enhanced peaks in the spectrum then must be in fast coincidence with this transition, thus giving some idea of the ground state de-excitations of Sm^{141} .

Table 1
Published Data for Sm^{141}

Authors	Half-life for Sm^{141m}	Half-life ^a for Sm^{141g}	% M4 ^a Transition	E_γ for ^a M4
[2]	21.5 min	<10 min	-	215 keV
[3]	23.5 min	≈2 min	<1%	≈200 keV

^aEstimated Values

References

- [1] Wm. C. McHarris, D. B. Beery, and W. H. Kelly, phys. Rev. Lett. 22, 1191 (1969); see also section 23 of this Annual Report.
- [2] R. Arl't, B. Baier, G. Muziol, L. K. Peker, G. Pfrepper, Kh. Shtrusnyi, and D. Khristov, JINR, Dubna (USSR), Report 21 (1967).
- [3] H. J. Bleyl, H. Münzel, and G. Pfinning, Radiochim. Acta 8, 200 (1967).

Fig. 10-1. Singles γ -ray spectrum from combined Sm^{141m+g} taken with a 2.5%, 2.3-keV resolution Ge(Li) detector.

Fig. 10-2. Sm^{141m} delayed gate integral coincidence spectrum, demonstrating the existence of an $11/2^-$ "isomeric" state in Pm^{141} at 629 keV.

PARTIAL DECAY SCHEME FOR Sm^{141}

Fig. 10-3. Partial decay scheme for the Sm^{141} system. We have evidence for many more levels, including some that appear to be three-particle states in Pm^{141} (cf. section 23 of this Annual Report).

Fig. 10-4. Slice taken from the Sm¹⁴¹ megachannel, 2-dimensional spectrum. A 2.0% Ge(Li) detector was gated on the 197.8-keV γ , which deexcites the first excited state of Pm¹⁴¹, and the coincident spectrum seen by a 2.5% Ge(Li) detector is shown here.

11. A Search for Dy^{147m}

R. Eppley

Two days were spent this summer at Yale University in the search for Dy^{147m}. This search was initiated after the observation of an M4 isomer in Gd¹⁴⁵. Both of these nuclei possess 81 neutrons and should exhibit similar energy levels. M4 isomers have been experimentally observed in N=81 nuclei extending from Te¹³³ (Z=52) through Gd¹⁴⁵ (Z=64); consequently, Dy¹⁴⁷ (Z=66) is the next logical extension for investigation. The Heavy Ion Linear Accelerator was used in an effort to produce the Nd¹⁴²(C¹²,7n) Dy^{147m} reaction.

Since no observation of Dy¹⁴⁷ has been reported in the literature, we had no readily available information as to what to look for in the resulting spectra. The best handle should lie with the supposed M4 isomeric transition. The half-life and transition energy were estimated by extrapolating the measured M4 data for N=81 nuclei. This was accomplished by fitting the experimental data to a least-squared quadratic curve and extrapolating to Dy¹⁴⁷. In this manner the extrapolated value for the M4 transition energy in Dy¹⁴⁷ was 604±16 keV and the square of the M4 matrix element, $|M_p|^2$, was 3.01. Using these values in the Moszkowski equation for a single-particle, M4 transition, the half-life came out to be 5 minutes (assuming a total internal conversion coefficient to be 0.321). It should be added that the $|M_p|^2$ values used in the least-square process were obtained by use of the same Moszkowski formula. This then was our starting point for the search for Dy^{147m}.

The following are the calculated Q-values for the 6n, 7n, and 8n reactions:

<u>Reaction</u>	<u>Q-Value (calc)</u>
Nd ¹⁴² (C ¹² ,6n)Dy ¹⁴⁸	-67.0 MeV
Nd ¹⁴² (C ¹² ,7n)Dy ¹⁴⁷	-78.9 MeV
Nd ¹⁴² (C ¹² ,8n)Dy ¹⁴⁶	-89.1 MeV

Consequently, our first attempts utilized a C^{12} beam of 90 MeV.

Two detectors were used in this search, a 20-cm³ Ge(Li) γ -ray detector and a 1000 μ Si(Li) electron detector. The γ -spectra were collected on a PDP-8/I computer using 2048 channels and the electron spectra were collected on a RIDL 400-channel analyzer.

The first attempt, using a 90-MeV beam, yielded no 5-min activity in the 500-800 keV region. In addition there was no evidence for a conversion electron peak in the electron spectrum, as should be the case for an $M4$ transition in this energy range.

With apparently negative results at 90 MeV, we made bombardments at 80, 100, 110, and 120 MeV. In no instance did we observe conversion electron peaks that could be attributed by Dy^{147m} .

It is interesting that we did observe the 1757.8-keV and 1880.6-keV peaks from the decay of Gd^{145} in some of the γ -ray spectra, as well as the 721.4-keV $M4$ peak of Gd^{145m} on both the γ -ray and electron spectra. The reaction producing this Gd^{145} activity was probably the $Nd^{142}(C^{12}, \alpha 5n)Gd^{145}$ reaction.

In view of these results, several possibilities can be listed as contributing to the negative experiment.

- 1) The $M4$ transition may have a significantly shorter half-life than our predicted value of 5 minutes. If this is the case we would not have observed the transition, due to the necessary 2-3 minute lapse between the end of the bombardment and the start of the counting procedure.
- 2) The $M4$ transition energy may be significantly different than the predicted value. The electron detector is useful only in the range of 300-1000 keV. However, it seems unlikely that the transition energy would fall outside this range.
- 3) The decay from the isomeric level may be predominantly positron or electron capture decay which would make the $M4$ a weak transition and one that could possibly be overlooked.

- 4) Assuming the $M4$ transition to be an $h_{11/2}$ to $d_{3/2}$ transition, it is possible that a state with an intermediate spin drops between these two states, thus allowing the decay to proceed step-wise with transitions of lower multipolarity. In this event no isomeric transition would be present.
- 5) There is apparently a high probability of competing reactions. From observation we were able to confirm competition from the $(C^{12}, \alpha n)$ reaction. Undoubtedly, there are other $(C^{12}, \alpha n)$ as well as $(C^{12}, p n)$ reactions which complicate the picture and possibly leave the desired reaction as a minor product.

Several of the targets were counted on a Ge(Li) γ -detector several days after the bombardments in order to determine which decay products were present in an effort to determine which mass chains were involved. In this manner we hope to obtain added evidence as to whether any of the observed transitions in the original spectra were indeed from Dy^{147} . Further comment will have to await the further analysis of the mass of data which has been accumulated.

12. Decay Studies on Tb¹⁵¹

R. Firestone

During 1952-1963 several papers [1-11] were published on the decay of Tb¹⁵¹, but in recent years little further work has continued on this subject. Of considerable early interest was the small (0.0005%) α -decay branch, which has been quite thoroughly studied [1-4]. Preliminary decay scheme work was done by several authors [5-11] using conversion electron and NaI(Tl) techniques.

Tb¹⁵¹ can be produced at the Michigan State University Sector-Focused Cyclotron by bombardment of separated isotope Eu¹⁵¹ with 20-MeV τ 's or 40-MeV α 's by Eu¹⁵¹(τ ,3n)Tb¹⁵¹ and Eu¹⁵¹(α ,4n)Tb¹⁵¹ reactions, respectively. This process is greatly complicated by the fact that production of Tb¹⁵¹ necessarily implies Tb¹⁵² as a by-product. Both isotopes have 18-h half-lives, so the complete spectrum of Tb¹⁵² must be identified before the Tb¹⁵¹ can be attempted. Little mention is made in the literature of how or whether this problem was solved, leading to some doubt as to the validity of ascribing certain γ -rays to the correct isotopes.

Initial work in obtaining γ -ray spectrum for Tb¹⁵¹ using a 7 cm³ Ge(Li) detector indicates a more complicated situation than evidenced by previous work. Unfortunately, the complexity of these spectrum has made classification of γ -rays nearly impossible, so further work on obtaining a good spectrum of Tb¹⁵² is currently in progress. It is expected that a knowledge of Tb¹⁵² spectrum will allow extrication of the peaks belonging to Tb¹⁵¹.

References

- [1] J. O. Rasmussen, Jr., S. G. Thompson, and A. Ghiorso, Phys. Rev. 89, 33 (1963).
- [2] K. S. Toth and J. O. Rasmussen, Phys. Rev. 109, 121 (1958).
- [3] R. D. Macfarlane and D. W. Seegmiller, Nucl. Phys. 53, 449 (1963).
- [4] K. Ya. Gromov, I. Mahunka, M. Mahunka, and T. Fenyés, Izv. Akad. Nauk. S.S.S.R., Ser. Fiz. 29, 194 (1965).
- [5] J. W. Mihelich, B. Harmatz, and T. H. Handley, Phys. Rev. 108, 989 (1957).
- [6] N. M. Anton'eva, A. A. Bashilov, B. S. Dzhelepov, and B. K. Preobrazhenskii, Doklady Akad. Nauk S.S.S.R. 119, 241 (1958).
- [7] V. I. Baranovskii, A. N. Murin, V. N. Pokrovskii, and I. A. Iutlandov, Izv. Akad. Nauk. S.S.S.R., Ser. Fiz. 22, 808 (1958).
- [8] K. S. Toth, S. Björnholm, M. H. Jørgensen, O. B. Nielsen, O. Skilbreid, and R. Svanheden, J. Inorg. Nucl. Chem., 14, 1 (1960).
- [9] A. T. Strigachev, L. S. Novikov, A. A. Sorokin, V. A. Khalkin, N. V. Tsvetkova, and V. S. Shpinel, Izv. Akad. Nauk. S.S.S.R., Ser. Fiz. 25, 813 (1961).
- [10] B. Harmatz, T. H. Handley, and J. W. Mihelich, Phys. Rev. 128, 1186 (1962).
- [11] A. T. Strigachev, A. A. Sorokin and V. S. Shpinel', Izv. Akad. Nauk. S.S.S.R., Ser. Fiz. 26, 252 (1961).

13. Radiations from the Decay of Yb^{167}

R. Goles

The present work represents a continuation of the study of the γ -ray cascades accompanying the electron capture, positron decay of Yb^{167} . A number of studies [1-5] have contributed to the now well-known ground state rotational band-level scheme of Tm^{167} . Recently, P. Paris [6] has conducted a γ -ray study of this decay with both $\text{Ge}(\text{Li})$ and $\text{Si}(\text{Li})$ detectors in an attempt to extend the level scheme of Tm^{167} . Significant new data have now been obtained regarding the decay system through use of high resolution high efficiency $\text{Ge}(\text{Li})$ detectors.

Samples of 18.5-min Yb^{167} were produced by bombarding the oxide of naturally occurring Tm^{169} with 23.5-MeV protons accelerated by the Michigan State University Sector-Focused Cyclotron. Samples were bombarded for approximately 3 min with average currents of 2 μA , and the resulting activities were counted for periods not exceeding 3 half-lives.

The γ -ray spectrum of Tm^{167} below 500 keV contain only those photopeaks corresponding to transitions between the already well-established ground-state rotational levels and has appeared in a previous publication [6]. A typical γ -ray spectrum associated with the decay of Yb^{167} in the energy region from 500 to 2000 keV as obtained in this lab appears in figures 1a and 1b. One unusual aspect of the decay of Yb^{167} , as a glance at figure 1b will verify, is the large number of closely spaced, high energy, low intensity γ rays occurring in the 1300- to 1500-keV energy interval. Since the decay energy of Yb^{167} is only ≈ 1.9 MeV, the high density of these high energy photopeaks indicates a significant feeding of high lying rotational band levels whose decay character appears to have a definite interband nature.

The energies of the lines found in our γ -ray spectrum are listed in Table 1. The detector used in this energy calibration was a newly acquired $\text{Ge}(\text{Li})$ detector with an efficiency of $\approx 2\frac{1}{2}\%$ and an energy resolu-

tion of 2.1 keV (FWHM) for the 1332-keV Co^{60} peak. However, an efficiency curve has not yet been completed for this new detector, and, as a result, relative intensity data will not be presented here.

The energy measurements of the prominent γ -rays accompanying the decay of Yb^{167} were made by counting the Yb^{167} sources simultaneously with several well-known calibration sources. A list of the calibration sources and the corresponding transitions used appear in Table 2. In determining the energy curve of a given spectrum, a least-squares fit of the photopeak centroids of the calibration transitions was made to a quadratic equation after background had been subtracted from under these peaks. The background correction for each peak was made by fitting a cubic equation to several channels adjacent to both sides of the peak and then subtracting. The energies of the lower intensity transitions, which were obscured by the calibration standards, were then determined similarly by using the stronger Yb^{167} γ -rays as secondary standards.

Attempts at finding a consistent correlation between the above reported γ -rays and the well-established ground state rotational band of Tm^{167} have not as yet been successful. Both coincidence and anti-coincidence experiments using an 8x8 in. NaI(Tl) split annulus [7] in conjunction with a Ge(Li) detector have been thwarted by the combined effects of the low efficiency of the Ge(Li) detector used and the low intensity characteristics of the high energy transitions. However, with the newly acquired 21% efficiency Ge(Li) detector, renewed efforts in this area will be forthcoming.

In addition, within the next year a series of experiments involving the (p,t) reaction on Tm^{169} will be conducted in order to begin a systematic examination of this reaction in terms of α -decay theory [8]. As a by-product of this investigation, the resultant particle spectra and the angular distribution of states will be exceedingly valuable in extending the presently limited decay scheme of Yb^{167} .

References

- [1] T. H. Handley and E. L. Olson, Phys. Rev. 94, 968 (1954).
- [2] B. Harwitz, T. H. Handley, and J. W. Michelich, Phys. Rev. 114, 1082 (1959).
- [3] R. G. Wilson and M. L. Pool, Phys. Rev. 120, 1296 (1960).
- [4] K. Ya. Gromov, A. S. Danagulyan, A. T. Strigachev, and V. S. Shpinel, U.S.S.R. Yad. Fiz., 1137 (1965).
- [5] T. Tamivä, Nucl. Phys. 62, 305 (1965).
- [6] P. Paris, J. J. Phys. (Paris) 28, 388 (1967).
- [7] R. L. Auble, D. B. Beery, G. Berzins, L. M. Beyer, R. C. Etherton, W. H. Kelly, and Wm. C. McHarris, Nucl. Instr. Methods 51, 61 (1967).
- [8] H. J. Mang and J. O. Rasmussen, Dan. Mat.-Fys. Skr. 2, No. 3 (1962); J. K. Pogenburg, Ph.D. Thesis (Berkeley), UCRL-16187 (1965).
- [9] C. M. Lederer, J. M. Hollander, and I. Perlman, *Table of Isotopes* (John Wiley and Sons, New York, 1966) 6th. Ed.
- [10] S. M. Brahmavar, J. H. Hamilton, A. V. Ramayya, E. F. Zangar, and C. E. Bemis, Jr., Nucl. Phys. A125, No. 1 (1969).

Table 1. Energies of γ -Rays Greater than 511 keV Present
in the Decay of Yb^{167}

Measured γ -ray energy (keV)	Measured γ -ray energy (keV)	Measured γ -ray energy (keV)	Measured γ -ray energy (keV)
541.54 $\pm$ 0.3	923.46 $\pm$ 0.3	1288.00 $\pm$ 0.1	1455.12 $\pm$ 0.1
547.57 $\pm$ 0.2	933.51 $\pm$ 0.3	1289.92 $\pm$ 1.0	1460.65 $\pm$ 0.4
665.05 $\pm$ 0.5	936.44 $\pm$ 0.3	1304.76 $\pm$ 0.2	1464.69 $\pm$ 0.3
672.13 $\pm$ 0.3	970.83 $\pm$ 0.4	1306.79 $\pm$ 0.2	1481.04 $\pm$ 0.3
680.44 $\pm$ 0.3	998.38 $\pm$ 0.1	1320.48 $\pm$ 0.2	1486.46 $\pm$ 0.3
687.07 $\pm$ 0.2	1008.97 $\pm$ 0.2	1321.74 $\pm$ 0.2	1487.61 $\pm$ 0.1
688.58 $\pm$ 0.4	1022.81 $\pm$ 0.2	1332.79 $\pm$ 0.1	1498.09 $\pm$ 0.2
694.05 $\pm$ 0.4	1026.23 $\pm$ 0.2	1336.68 $\pm$ 0.3	1508.96 $\pm$ 0.5
695.58 $\pm$ 0.4	1037.02 $\pm$ 0.1	1339.94 $\pm$ 0.3	1511.89 $\pm$ 0.1
697.28 $\pm$ 0.4	1048.68 $\pm$ 0.2	1342.32 $\pm$ 0.3	1516.67 $\pm$ 0.3
707.67 $\pm$ 0.4	1050.36 $\pm$ 0.2	1355.27 $\pm$ 0.3	1517.99 $\pm$ 0.3
719.71 $\pm$ 0.4	1067.56 $\pm$ 0.3	1358.31 $\pm$ 0.4	1525.45 $\pm$ 0.2
733.13 $\pm$ 0.3	1069.52 $\pm$ 0.3	1361.54 $\pm$ 0.2	1533.54 $\pm$ 0.2
791.56 $\pm$ 0.2	1110.31 $\pm$ 0.2	1365.53 $\pm$ 0.5	1537.64 $\pm$ 0.2
794.19 $\pm$ 0.5	1113.50 $\pm$ 0.4	1369.88 $\pm$ 0.1	1570.47 $\pm$ 0.1
815.92 $\pm$ 0.3	1137.12 $\pm$ 0.5	1385.06 $\pm$ 0.2	1587.18 $\pm$ 0.1
829.34 $\pm$ 0.2	1139.56 $\pm$ 0.2	1393.12 $\pm$ 0.2	1619.04 $\pm$ 0.3
832.88 $\pm$ 0.2	1213.22 $\pm$ 0.1	1401.92 $\pm$ 0.3	1619.65 $\pm$ 0.3
846.17 $\pm$ 0.2	1217.07 $\pm$ 0.1	1410.41 $\pm$ 0.3	1631.82 $\pm$ 0.2
903.32 $\pm$ 0.3	1234.64 $\pm$ 0.1	1427.68 $\pm$ 0.2	1643.33 $\pm$ 0.3
905.33 $\pm$ 0.3	1241.89 $\pm$ 0.1	1433.46 $\pm$ 0.4	1644.19 $\pm$ 0.3
920.29 $\pm$ 0.1	1284.91 $\pm$ 0.5	1438.39 $\pm$ 0.1	

Table 2. γ -Ray Calibration Standards

Source	γ -ray energy (keV)	Reference
Na ²²	1274.53 $\pm$ 0.10	9
Sc ⁴⁶	889.18 $\pm$ 0.10 1120.41 $\pm$ 0.10	9
Co ⁶⁰	1173.226 $\pm$ 0.040 1132.483 $\pm$ 0.046	9
Zn ⁶⁵	1115.44 $\pm$ 0.01	9
Y ⁸⁸	897.96 $\pm$ 0.10 1836.08 $\pm$ 0.07	9
Ag ^{110m}	657.71 $\pm$ 0.03 706.68 $\pm$ 0.04 763.88 $\pm$ 0.04 884.67 $\pm$ 0.04 1475.73 $\pm$ 0.04 1504.90 $\pm$ 0.08 1562.22 $\pm$ 0.06	10
Ta ¹⁸²	1121.28 $\pm$ 0.12 1221.42 $\pm$ 0.10	9
Bi ²⁰⁷	569.63 $\pm$ 0.08 1063.58 $\pm$ 0.06 1769.71 $\pm$ 0.13	9

Fig. 13-1. Yb¹⁶⁷ spectrum showing the γ -rays with energies greater than 511 keV. This spectrum was taken with our 2.5% detector.

14. As⁷⁴ Decay and Forbidden Zero-Phonon Transitions.

W. McHarris

(in cooperation with J. Van Hise, Tri-State College,
and C. Papperillo, Notre Dame)

Sources of 18-d As⁷⁴ were prepared by three methods for use in this study: Ge⁷⁴(d,2n)As⁷⁴ (using the MIT cyclotron), As⁷⁵(γ,n)As⁷⁴ (at Livermore), and Ge⁷⁴(p,n)As⁷⁴ (using the MSU sector-focused cyclotron). After chemical purification, the sources were counted on a number of detectors at a number of different laboratories. A "typical" spectrum taken with the Ge(Li) detector system at Livermore is shown in fig. 1, and the list of γ-ray energies and intensities is given in Table 1. These energies and intensities are a composite of those obtained at MSU, Andrews University (while JVH was a faculty member there), Notre Dame, and Livermore; the agreement among the different laboratories was found to be excellent. In addition to singles spectra, various Compton-suppressed spectra were taken at MSU and Livermore (cf. section 36 of this Annual Report), and pair spectra were taken at MSU. A portion of the MSU Compton-suppressed spectrum that shows the existence of a weak 734.2-keV γ is shown in fig. 2.

The whole point of re-studying the decay of this nuclide, which had previously been studied by a considerable number of groups [1], was to look for the possible existence of a very weak 278.6-keV transition between the 1482.86-keV level and the 1204.31-keV level in Ge⁷⁴. These levels are presumably the 0+ and 2+ members of the two-phonon triplet, and according to straightforward theory of a vibrational nucleus [2], such a transition should be absolutely forbidden. However, recently such a transition was found to occur between the 4+ and 2+ states of a similar triplet in Ni⁶⁰, populated by the decay of Co⁶⁰ [3]. There the ratio of the observed transition rate to the single-particle rate was given a lower limit of 0.174. Our decay scheme for As⁷⁴ is shown in fig. 3, where it can be seen that we did not observe the analogous transition. The assignments given in fig. 3 are given on the basis of our data and also draw liberally on already published data, including reactions studies.

Focusing our attention on the states in Ge^{74} rather than Se^{74} , we give the following brief resume of our findings. The collective nature of the first excited state is apparently clear through Coulomb-excitation work [4], where it was interpreted as a single-phonon quadrupole vibrational state. It is not unreasonable to interpret the $2+$, $4+$ and $0+$ states next highest at 1204, 1464, and 1483 keV as the two-phonon triplet. That Ge^{74} is indeed not a pure harmonic vibrator is easily seen by the existence of the 1204-keV transition down to the ground state. Our search for the existence of the 278.6-keV transition, had it been successful, would have been further evidence of the failure of the simple vibrator model to explain this nucleus.

Using the Compton-suppressed spectra, we could place an upper limit on the 278.6-keV transition of $7.4 \times 10^{-3} \%$ of the intensity of the 595.896-keV transition. Of real interest, of course, is to compare the rate (its upper limit in this case) of the 278.6-keV transition with that of a model prediction. As the half-life of the 1482.86-keV state has not been measured, we must resort to a more approximate route if any estimate of the 278.6-keV rate is to be made. A survey of medium weight nuclei reveals that the transition from the $0+$, $2+$, or $4+$ members of the two-phonon triplets to the $2+$ single-phonon state tend to be ≈ 20 times as fast as the single-particle $E2$ rate. If we then assume the present 886.96-keV γ to be similarly enhanced by a factor of about 20 over the single-particle rate, we can place an upper limit on the ratio of experimental to single-particle ratios for the 278.6-keV transition by simply comparing their energies raised to the fifth power (for quadrupole transitions). When this is done, we find that

$$\lambda_{\text{exp}} / \lambda_{\text{single-particle}} \leq 1114,$$

which does not really say much about its degree of forbiddenness or non-forbiddenness.

Clearly As^{74} is not too useful for a search for a zero-phonon transition because of the weak population of the $0+$ state at 1483 keV and the even weaker population of the 1464-keV state, from which another zero-phonon transition could arise. This overcomes the favorable part, which is the relatively wide spacing between the $2+$ member of the triplet and the other

two members.

While no model predicts quantitatively the level spacings of Ce^{74} , the vibrator model does much better than any sort of rotator model. While addition of a cubic term to the oscillator potential does not relieve the degeneracy of the $0+$ and $4+$ members of the two-phonon triplet, a quartic term does remove this degeneracy. Both the cubic and quartic terms added to the oscillator potential as calculated by Belyaev and Zelevenskii [5] and Brink [6] does not give any improvement over the potential including only the quartic term. The best approximation to experiment yet is the oscillator plus quartic potential of Ferreira, Alcaraz, and Navarro [7]. And the γ -unstable rotator of Davydov and Chaban, while predicting $0+$, $2+$, and $4+$ states, places these at 1027 ($2+$), 1248 ($4+$), and 1922 ($0+$) keV, clearly not a satisfactory situation [8].

References

- [1] J. H. Hamilton, F. E. Coffman, A. V. Ramayya, and N. S. Johnson, Nucl. Phys. A132, 254 (1969), and references to previous work contained therein.
- [2] D. A. Bromley and J. Weneser, Comments Nucl. Particle Phys. 1, 75 (1967).
- [3] J. R. Van Hise and D. C. Camp, Phys. Rev. Lett. 23, 1248 (1969).
- [4] R. L. Robinson, P. H. Stelson, F. K. McGowan, J. L. C. Ford, and W. T. Milner, Nucl. Phys. 74, 281 (1965).
- [5] S. T. Belyaev and V. G. Zelevinskii, Izv. Akad. Nauk SSSR, Ser. Fiz. 28, 127 (1964).
- [6] D. M. Brink, Argonne National Laboratory report ANL-6797 (1963).
- [7] P. L. Ferreira, J. A. C. Alcaras, and V. C. A. Navarro, Phys. Rev. 136, B1243 (1964).
- [8] A. S. Davydov and A. A. Chaban, Nucl. Phys. 20, 499 (1960).

Table 1. Ag^{74} γ -Ray Energies and Intensities.

E_{γ} (keV)	I_{γ}
595.896 $\pm$ 0.08	100
608.39 $\pm$ 0.08	0.92
634.78 $\pm$ 0.08	24.8
715.33 $\pm$ 0.2	0.014
734.2 $\pm$ 0.3	0.0059
867.0 $\pm$ 0.7	0.0088
886.96 $\pm$ 0.1	0.043
993.46 $\pm$ 0.08	0.035
1101.1 $\pm$ 0.2	0.010
1204.34 $\pm$ 0.08	0.46
1269.0 $\pm$ 0.6	0.0031
1602.09 $\pm$ 0.5	0.012
2198.1 $\pm$ 0.3	0.029

Fig. 14-2. As^{74} Compton-suppressed spectrum taken at MSU by WCM. Only the portion showing the new 734.2-keV γ is shown.

Fig. 14-3. As^{74} decay scheme.

✓ 15. The Decays of Fe^{53g} and Fe^{53m}

J. Black

Iron-53, an 8.5-min nuclide decaying by both positron emission and electron capture, was first produced by a $Cr(\alpha, n)$ reaction on naturally occurring Cr [1]. Consequent investigations have been made using a number of methods, including scattering and (p, α) reactions [2-5]. Since modern techniques of γ -ray spectroscopy with semiconductor detectors have made it possible to study short-lived nuclides with greater ease and good resolution, it was decided to reinvestigate this decay.

The γ -rays emitted by this nuclide were examined with high-resolution Ge(Li) detectors. In addition, the positron spectrum was investigated with a Si(Li) surface-barrier detector. A precise energy determination was made for the 377.9-keV γ , together with the observation of two additional γ -rays attributable to the decay of Fe^{53g} and four γ -rays attributable to Fe^{53m} . The positron spectrum taken with the Si(Li) detector was examined in singles mode and also in coincidence with the 377.9-keV γ . The results of the γ -ray and positron studies were used to establish decay schemes for both Fe^{53} isomers (fig. 1).

Sources were produced by the reaction $Mn^{55}(p, 3n)Fe^{53}$, a relatively clean and unexplored method of production. Mn metal powder (99.94%) was bombarded with 34-MeV protons from the MSU Cyclotron. Typically, 0.5-g targets were bombarded with a 2- μ A beam for 3 min. To remove unwanted contaminants and insure that only the Fe^{53} was being counted, a chemical separation was carried out after each bombardment. The source was dissolved in 8.0-N HCl and the iron extracted with isopropyl ether. After washing with acid, the iron was extracted with distilled water, and a small portion of this was used for counting. For the positron work, a small portion was carefully dried on a thin foil holder and then counted with the Si(Li) detector. Most counting periods were for roughly 24 h, so multiple bombardments and separations were performed -- material from each bombardment being counted for ≈ 20 min.

A 7-cm³, five-sided coaxial Ge(Li) detector manufactured in this laboratory was used to determine the energies and intensities of the γ -rays.

Typical resolution obtained with this detector was 2.9 keV FWHM for the 1332.48-keV γ from Co^{60} . In addition, a 2.5% coaxial Ge(Li) detector having a resolution of 2.1 keV and a peak-to-Compton ratio of 16.5/1 was used for energy determinations. The energies of the γ -rays were measured by counting the Fe^{53g+m} sources simultaneously with a number of well-known calibration sources. A background correction was made for each peak by fitting a linear equation to channels above and below the peak and then subtracting. The centroids of unknown peaks were determined in a similar manner and the corresponding energies calculated from the calibration curve. The energies of the weaker Fe^{53g+m} γ -rays were verified by using the established stronger γ -rays as internal standards. Unwanted activities with different half-lives (prevailing even after the chemical separation) were detected by obtaining spectra periodically as the sources aged. Four γ -rays in the region 500-2500 keV were present in the Fe^{53g} spectra even after careful chemical separation. These peaks all had half-lives of 2-3 min, and they were identified as belonging to Fe^{53m} . A recent investigation by Eskola [6] has resulted in a decay scheme for 2.58-min Fe^{53m} . Our results for energies and half-lives confirm the results reported in his work. We have also observed additional weak γ -rays after the chemical separation and with apparently the Fe^{53g} half-life, so we attribute these to Fe^{53g} decay. A singles γ -ray spectrum taken with the 2.5% detector is shown in fig. 2. A list of energies and relative intensities for the γ -rays from the two isomers is given in Tables 1 and 2.

In addition to the singles data, our investigation has included the following γ -ray studies using Ge(Li) detectors in conjunction with an 8"x8" NaI(Tl) split annulus: anticoincidence, integral coincidence, and coincidence with the 377.9-keV γ . Most of these experiments were carried out without performing a chemical separation on the target. In addition, no specific effort was made to optimize the 2.5-min Fe^{53m} . Although these runs were of sufficient length (≈ 24 h) to produce reasonable statistics, few positive results were obtained. The overwhelming abundance of the 377.9- and 511-keV γ 's makes the detection of the very weak, higher-energy peaks rather difficult.

The anticoincidence spectrum revealed practically complete suppression of the 701.0-, 1011.5-, 1328.1-, and 2339.6-keV γ 's. The spectrum of γ -rays in coincidence with the 377.9-keV γ showed evidence for the 1011.5-

-keV γ , although no trace of other Fe^{53g+m} γ -rays was found. The integral coincidence experiment showed all γ -rays assigned to either isomer except for the 1619.9-keV transition attributed to Fe^{53g} . (This spectrum, however, did not cover energies above 2500 keV and would not include the 2750.7-keV γ assigned to Fe^{53g} .)

The Si(Li) surface-barrier detector used for positron detection was mounted in a dipstick cryostat and had an area of 200 mm² and a thickness of 1000 μ . In order to reduce noise the cryostat dipstick was immersed in a dry ice-methanol bath. The detector itself was covered by an evacuated Al can with a 0.10-mil Havar window for admission of the positrons. After the chemical separation, a drop of the solution was dried on a thin backing and placed in an appropriate holder on the detector can. Each sample was counted for approximately 20 min, using an ORTEC 109A preamplifier, a Canberra 1417 amplifier, and a Nuclear Data ND-160 analyzer. The spectra were calibrated by using the conversion electrons of Cs^{137} and Bi^{207} . Using the data-analysis task MOIRAE, a first-order background fit was made to the conversion-electron peaks of these standards and their centroids were determined. A polynomial fit of energy to channel number was then made using the program DATA IN. Using the tables of NBS Applied Math Series #13 [7], the corrected Fermi ordinates were obtained and Fermi plots prepared. Fig. 3 shows a Fermi plot of an Fe^{53} beta spectrum. The end-points of the three resolved beta groups are given in Table 3.

It would appear quite reasonable that if the 2.8-MeV group feeds the ground state, then the 2.4-MeV group feeds the first excited state at 377.9 keV. To examine this possibility, a beta-gamma coincidence experiment was carried out. A 3"x3" NaI(Tl) detector was used to gate on the 377.9-keV γ , while the Si(Li) detector was used to observe the positrons in coincidence with this γ -ray. The Fermi plot of the resultant beta spectrum revealed a highest end-point group at 2.4 MeV, 0.4 MeV less than the most energetic group in the ungated spectrum. Thus, the beta-gamma coincidence experiment confirms the earlier hypothesis that the 2.4-MeV group feeds the 377.9-keV level.

The decay schemes of Fe^{53g} and Fe^{53m} resulting from this study are shown in fig.1. Excited state energies are given in keV. The Fe^{53g} state probably at 2750.7 keV has not been reported previously. In addition, a pre-

cise energy determination was made on the rather well-known state at 377.9 keV. The state at 1619.9 keV is in agreement with other recent work [8]. Although reported earlier [5, 8], no evidence was found in this study to support a level in the vicinity of 1290 keV. Some evidence was seen for levels at higher energies, but they have not been verified by sufficient experimentation to warrant their inclusion in the decay scheme. The half-life of Fe^{53m} was taken from the work of Darnedde [9]. The level at 2339.6 keV was observed from its full-energy photopeak in addition to the double-escape and cascade peaks. The growth in Fe^{53g} activity as the isomer decayed showed indeed that the decay occurs through the ground state.

Examination of the single-particle estimates for a state with an energy of 3040.6 keV and a half-life of 2.5 min reveals that it must have a spin difference of at least five units between it and the ground state (e.g., $M5$ and $E5$ transitions would have half-lives of only 4.9 and 2.4 sec, respectively, and higher multipolarities might approach the observed half-life). Since the ground-state spin is known to be $7/2$, one infers that the isomeric state must have a spin of at least $17/2$. If the isomeric state is assumed to be a three-particle configuration, then the predicted spin value is $19/2$. The 701.0-keV transition from the isomeric state to the 2339.6-keV state is most likely an $E4$ or $M4$. Single-particle estimates of half-lives for such transitions are 0.65 and 2.08 min, respectively. Comparison with the known half-life would indicate that the $M4$ multipolarity is the correct assignment, but one cannot rule out the $E4$ as a possibility. Nevertheless, the transition is certainly accompanied by a spin change of four units. This fixes the spin of the 2339.6-keV state as $11/2$. On the basis of the observed γ -ray branching ratios, Eskola [6] has assigned the first excited state a spin of $9/2$.

In shell-model terms, the configuration of Fe^{53} is $(f_{7/2})^{-3}$, with one neutron hole and two proton holes in the $N=Z=28$ doubly closed shell. With the assumption of an inert Ca^{40} core, the possible states of the remaining pure $f_{7/2}$ configuration in the jj -coupling scheme and with seniority $v=3$ are $1/2^-$, $3/2^-$, $(5/2^-)^2$, $(7/2^-)^2$, $(9/2^-)^2$, $(11/2^-)^2$, $(13/2^-)^2$, $15/2^-$, $17/2^-$, and $19/2^-$ [10]. This three-particle multiplet would normally be degenerate in a central field, but splitting will occur because of residual interactions. McCullen, et al. [11], have made shell-model calculations for nuclei in the

$1f_{7/2}$ shell, taking the residual interaction to be a two-body interaction, of which the strength is determined from the observed levels of the configuration $(1f_{7/2})^2$ in Sc^{42} and Ca^{42} . The results of these calculations agree well with the experimental results for the lower-lying excited states in Fe^{53} . They predict the large spin gap below the $19/2$ state, which is responsible for the isomer, and the predicted energies are quite good for the $11/2$ and $9/2$ states.

The assignment of $7/2^-$ for the spin and parity of the ground state of Mn^{53} is in agreement with shell-model predictions and also with experimental evidence [12]. In shell-model terms the ground state is predicted to be an $(f_{7/2})^{-1}$ neutron state. The excited state at 377.9 keV is probably $5/2^-$, in agreement with other odd-mass nuclei in the $f_{7/2}$ shell region. This first excited state would presumably have a $(f_{5/2})^{-1}(f_{7/2})^{-2}$ configuration. Theoretically, the second excited state should be $3/2^-$ [13], which would also be consistent with other odd-mass nuclides in this region. The configuration would be $(p_{3/2})^{-1}(f_{7/2})^{-2}$.

There is a good possibility that three-particle isomers similar to Fe^{53m} may exist in isotopes such as Sc^{43} , Ti^{43} , and Co^{53} . The large decay energy of the Ti^{43} and Co^{53} , and the possibility of the decay of their $19/2^-$ state by superallowed positron decay to $19/2^-$ states in Sc^{43} and Fe^{53} , leads to the prediction that the half-lives would be shorter than 1 sec. However, decay of the $19/2^-$ state in Sc^{43} should be similar to Fe^{53m} . The failure of earlier experiments [6] to detect any of these isomers can be attributed primarily to the short half-lives involved and the experimental difficulties they entail. Therefore, we are now preparing an investigation using the recently-constructed Rabbit [14] system at the MSU Cyclotron that will permit the observation of isomers with much shorter half-lives; with pulsed-beam techniques half-lives as short as milliseconds should be observable [15].

References

- [1] L. N. Ridenour and W. J. Henderson, Phys. Rev. 52, 889 (1937).
- [2] S. E. Arnell and S. Sterner, Ark. Fys. 25, 423 (1964).
- [3] P. H. Vuister, Nucl. Phys. 83, 593 (1966).
- [4] S. Sterner, L. Jonsson, and S. E. Arnell, Ark. Fys. 31, 567 (1966).
- [5] J. O. Juliano, C. W. Kocher, T. D. Nainen, and A. C. G. Mitchell, Phys. Rev. 113, 602 (1959).
- [6] K. Eskola, Ann. Acad. Sci. Fenn. AVI, No. 261, p. 45 (1967).
- [7] *Tables for the Analysis of Beta Spectra*, National Bureau of Standards, Applied Math. Series, No. 13 (1951).
- [8] M. T. McEllistrem, K. W. Jones, and D. M. Sheppard, Bull. Am. Phys. Soc. 13, 1426 (1968).
- [9] I. Dervedde, Z. Phys. 216, 103 (1968).
- [10] B. H. Flowers, Proc. Roy. Soc. A212, 248 (1952).
- [11] J. D. McCullen, B. P. Bayman, L. Zamick, Phys. Rev. 134, B515 (1964).
- [12] W. Dobrowolaki, R. V. Jones, and C. D. Jaffries, Phys. Rev. 104, 1378 (1956).
- [13] A. deShalit, *Selected Topics in Nuclear Theory*, ed. by F. Janouch, I.A.E.A., Vienna, 1963.
- [14] Section 29 of this Annual Report.
- [15] Sections 22 and 30 of this Annual Report.

Table 1.
Fe^{53m} γ-Rays.

E_{γ} (keV)	Relative Intensity
701.0	100
1011.5	80
1328.1	88
2339.6	23

Table 2.
Fe^{53g} γ -Rays.

E_{γ} (keV)	Relative Intensity
377.9	41
1619.9	0.50
2750.7	0.30

Table 3.
Fe⁵³ β^+ Groups.

Group No.	Energy (MeV)
1	2.80 $\pm$ 0.10
2	2.39 $\pm$ 0.10
3	1.71 $\pm$ 0.15

PROPOSED DECAY SCHEMES OF ^{53}Fe & $^{53\text{m}}\text{Fe}$

Fig. 15-1.

Fig. 15-2. Fe^{53g+m} singles γ -ray spectrum taken with a 7-cm³ Ge(Li) detector. The sources used had been chemically purified, and many repeated bombardments were required to produce this spectrum, which represents some 17 hr of counting time.

Fig. 15-3. Fermi-Kurie plot of the positron spectrum from Fe^{53g} . The spectrum was taken with a 1000 μ Si(Li) detector.

16. The Decay of 38.4-Min Zn^{63} .

G. Giesler

In the last three years several authors [1-3] have reported data on Zn^{63} obtained by using Ge(Li) detectors. Only the most recent one [3], however, reports any coincidence work. For the coincidence detectors, they used a Ge(Li)-NaI(Tl) arrangement, gating the NaI(Tl) detectors on broad regions centered about 511, 669.6, and 961.9 keV. This is the sole coincidence work published on Zn^{63} decay.

As preparation for the studies of Ga^{63} , Ga^{62} , and other possible delayed α -emitters in this region, we also examined the decay of Zn^{63} . In addition to singles spectra, we also obtained two-dimensional coincidence data using the system described in section 31 of this Annual Report. The Ge(Li) detectors were 2.5 and 2.0% efficient. Fig. 1 shows a singles spectrum. The highest energy peak seen is at 3044.4 keV. Higher energy peaks were seen in other spectra, but they were removed with chemical separation. This spectrum is, in general, similar to those previously published. In fig. 2 we show the results of the two-dimensional coincidence experiments. Figs. 2A and 2E show the 2.5% and 2.0% integral coincidence spectra, respectively. Normally we set gates on the y axis and used the x axis as the display axis (the x axis has better resolution), but the results were substantiated by using the x axis as gate and y axis as display. The 450-keV γ is in coincidence with the 962-keV γ , as shown by figs. 2B and 2F. In 2B, gating on the 450-keV γ increases the intensity of the 962-keV γ above that of the 670-keV γ , whereas everywhere else the 670-keV γ is the more intense. By gating on the 962-keV γ , fig. 2F, the 450-keV γ stands out much more strongly than anywhere else, including singles. The 511-keV gate indicates the states that are fed by β^+ emission. In figs. 2D and 2I, respectively, the 670- and 1412-keV gates indicate no coincidence other than with the 511-keV $\gamma^{\pm}$ and its Compton. Only the 450-keV γ and $\gamma^{\pm}$ are in coincidence with the 962-keV γ . The 1040-keV peak, which in singles looks like a small Compton edge, is in coincidence with the $\gamma^{\pm}$ and the 670-keV γ . The 1392-keV γ , fig. 2H, appears to be only in coincidence with the 670-keV γ and is not β^+ fed. With experiments

such as this two-dimensional analysis, we have come up with the preliminary decay scheme shown in fig. 3. Energies are placed only on those γ -rays in cascades -- the others all appear to be ground-state transitions. A few weak γ -rays remain to be placed, so further work is in progress to place these and, more importantly, to interpret the decay scheme, i.e., try to determine the structures of the states that have been found.

References:

- [1] P. Holmberg, E. Spung, and A. Lundán, Soc. Sci. Fenn., Comm. Phys. Math. 32, No. 12 (1966).
- [2] D. DeFrenne, M. Dorikens, L. Dorikens-Vanpraet, and J. Demuynck, Nucl. Phys. A103, 203 (1967).
- [3] I. Borchert, Z. Phys. 223, 473 (1969).

Fig. 16-1. Zn⁶³ singles γ -ray spectrum taken with our 2.5% Ge(Li) detector, using a chemically purified source.

Fig. 16-2(a). Slices from a two-dimensional, megachannel coincidence spectrum for Zn^{63} . In the top spectrum of this fig. and of part (b) are shown the integral coincidence spectra, and in the lower portions are shown the various selected slices.

Fig. 16-2(b).

Fig. 16-3. ^{63}Zn decay scheme. We are reasonably confident about the energies of the levels, but the assignments of quantum numbers are still in progress.

17. The Decay of 9.3-Hour Zn^{62} .

G. Giesler

Within the last few years there has been a flurry of activity in studying the decay of Zn^{62} . Four authors [1-4] have reported their studies using Ge(Li) detectors. Within this set of results, only those of Bakhru [3] do not fall in close agreement with the others. He reports a transition at 682 keV and does not see one reported at 638 keV by the others. Based on these differences, he proposed a decay scheme somewhat different from the one proposed by the other authors. None saw any transitions of higher energies, although most had looked for them.

Our only reason for entering this already crowded domain was to recheck the consistency of the various results. We had decided that Ga^{62} was a good candidate for β -delayed α 's (see section 21 of this Annual Report), so it was necessary to check on activities in the $A=62$ chain quite thoroughly. Thus, we decided to look for higher-energy transitions in Zn^{62} decay.

A typical γ -ray spectrum is shown in fig. 1. Below 700 keV there is good agreement with the results of the previous authors. Eight transitions are seen above 700 keV, three of which come from the Cu^{62} daughter. Also present is a peak from Zn^{65} produced from the Cu^{65} in the natural Cu target foil. This is the only possible contaminant, since all other contaminants are removed by ion-exchange separations plus allowing the source to decay for several hours before counting. Two peaks, those from K^{40} and Tl^{208} , are present from room background.

There are two peaks remaining, those at 1279 and 1430 keV. Many experiments have been performed and prove that they are indeed from the Zn^{62} decay chain. The possibility of their occurring from the decay of Cu^{62} is ruled out since there is no combination of levels (well-known from many reaction studies) that has an energy difference close to 1279 keV, and only one combination, 3751 keV (3-) to 2336 (4+), that is close to the 1430-keV transition. The latter can be ruled out since the 1+ ground state of Cu^{62} lies only 3939 keV above the 0+ Ni^{62} ground state. These two transitions fit better to the 1285- and 1432-keV states seen by Morrison [5] in $\text{Ni}^{62}(\tau, d)$ scattering. At

present, we have determined the intensities of the 1279- and 1430-keV γ 's to be about 1×10^{-4} and 7×10^{-4} that of the 556-keV γ , respectively. Based on this, we propose the decay scheme in fig. 2. Further work is now being performed to characterize the transitions better.

References:

- [1] K. I. Roulston, B. H. Becker, and R. A. Brown, Phys. Lett. 24B, 93 (1967).
- [2] S. Antman, H. Pettersson, and A. Suarez, Nucl. Phys. A94, 289 (1967).
- [3] H. Bakhru, Phys. Rev. 169, 889 (1968).
- [4] B. J. Hoffman and D. G. Sarantites, Phys. Rev. 177, 1640 (1969).
- [5] G. Morrison, private communication (1969).

Fig. 17-1. γ -ray spectrum of chemically-purified Zn^{62} taken with a 7-cm³ Ge(Li) detector.

Fig. 17-2. ^{62}Zn decay scheme.

✓
18. The α Decay of Es^{252}

W. McHarris

(in cooperation with I. Ahmad and P. Fields, from
 Argonne National Laboratory)

Our studies of the α decay of Es^{252} to states in Bk^{248} (of particular interest because it crosses the $N=152$ subshell) appear to be more or less complete, but there are some open questions remaining concerning its ϵ decay (20%) to Cf^{252} . As a result, near the end of 1968 a fresh source was produced by bombarding a $\approx 150\text{-}\mu\text{g}$ Cf^{252} target with $\approx 14\text{-MeV}$ deuterons in the Argonne Cyclotron for about 5 days. Care was taken to prevent the presence of any Es^{254} , whose α groups would dominate the middle-energy portion of the Es^{252} α spectrum, and the source was set aside to allow the unavoidable 20-d Es^{253} [from a $\text{Cf}^{252}(d,p)\text{Cf}^{253}\xrightarrow{\beta^-}\text{Es}^{253}$ sequence] to decay away. Thus, in early 1970 we plan another look at Es^{252} decay, directed primarily at the ϵ -decay side, but also reexamining the α -decay side with improved α - and γ -ray detectors.

Because our subsequent studies may well produce slight alterations in energies and intensities, we do not present our formal α -decay scheme here; several interesting features in it, nevertheless, are worth comment. The ground state of Es^{252} appears to be a $7+$ state -- a triplet coupling of a $7/2+$ [633+] proton with a $7/2+$ [613+] neutron, the same as the ground state of Es^{254} [1]. However, whereas the favored decay of the latter could populate a $7+$ band starting at 86 keV in Bk^{250} , because of the $N=152$ gap the corresponding band in Bk^{248} starts at 593 keV. The 6.055-MeV favored α group as a result accounts for only $1.0\pm 0.1\%$ of the total α decay, and a 5.990-MeV group (mostly $l=2?$) presumably to the $8+$ member of the band at 658 keV is only $0.06\pm 0.03\%$ abundant. The lower-lying states in Bk^{248} all involve neutron states from below $N=152$, and its presumed ground-state band is an $8-$ band composed of a triplet coupling of the $7/2+$ [633+] proton with a $9/2-$ [734+] neutron. The $9-$ and $10-$ members of this band lie at 70.8 1.0 and 151.5 1.0 keV, respectively. The three α groups populating this band have energies (and intensities) of 6.638 ± 0.003 ($82.0\pm 1.0\%$),

6.568 (13.2±0.4%), and 6.489 keV (2.1±0.1%).

Characterizing this 8- "ground-state" band has involved a higher amount of relying on systematics and calculations than usual. First of all, it is not really clear that it is the true ground state. There appear to be two Bk²⁴⁸ isomers, a 2- state that decays by both β^- and ϵ decay, and a long-lived isomer that was identified through mass spectrometry and which could be the 8- state [2]. Searches have been made to detect small feedings between the two, but to date there have been no definitive results. Second, the rotational spacings of the members of this band are not what one would normally expect for a K=8 band. Calculation of the rotational constant, $\hbar^2/2\mathcal{J}$, from the simple $I(I+1)$ equation leads to a value of 3.93 keV (based on the 8-to-9 spacing), much smaller than encountered even in odd-odd nuclei in this region. Similarly, if one predicts the apparent K from the (9-10)/(8-9) spacings, its value comes out more like 6 than 8. This implies a large *positive* B in the rotational-energy equation,

$$E_I = E_0 + \frac{\hbar^2}{2\mathcal{J}} I(I+1) + BI^2(I+1)^2.$$

These apparent anomalies can be explained, indeed, would be expected, when one considers the Coriolis coupling and distortion inherent in the proton and neutron states involved. The Coriolis operator (cf. ref. [3] for a derivation and discussion of these results) has the form,

$$- \frac{\hbar^2}{2\sigma\mathcal{J}} j_{\pm} I_{\mp},$$

producing matrix elements of the form,

$$- \frac{\hbar}{2\sigma\mathcal{J}} [I(I+1) - K(K\pm 1)]^{1/2} [j(j+1) - \Omega(\Omega\pm 1)]^{1/2}.$$

Thus, it can be seen that large I 's contribute greatly to the distortion, and so do large j 's (even though the j 's become mixed in these deformed states, the j from the original spherical state always remains as a large component). The 7/2+[633+] proton state originates from the $i_{13/2}$ spherical state and is known to cause severe band distortion in Bk²⁵⁰ [1]; also, the 9/2-[734+] neutron state originates from the $j_{15/2}$ spherical state. Conditions are right, therefore, in the 8- band in Bk²⁴⁸ for very large Coriolis distortions. When calculating these,

we used the experimental values of distortions produced by these states in neighboring odd-mass nuclei, and the results duplicate the 8- band distortion quite well.

The other handle on identifying this band correctly comes from the calculated α transition probabilities to the three states seen in the band. These are simply a function of Clebsch-Gordan coefficients,

$$\frac{1}{HF} = \sum_l \frac{1}{HF_l} (I_i \pm K_f - K_i \ K_i | I_i \pm I_f \ K_f)^2 ,$$

where the HF_l (hindrance factors for given l waves) can be determined by normalizing the observed intensities to two states. We have tried other spins, for example the fairly reasonable 5- and 6+ possibilities for Es^{252} , but no lower spins seem to be able to duplicate the observed intensities.

All the α hindrance factors are abnormally small, in fact, and we are presently looking at how mixing can affect other states. There are several other high-lying states that receive fairly large amounts of α population. Because the 8- band lies so much lower than the 7+, it might be possible, say, to find another 7+ state, this time an octupole vibration based on the 8- band, lying fairly close to the 593-keV 7+ band. The possibilities for mixing would be enormous, so we expect a rather substantial amount of information to come out of the studies on this nucleus when they are completed.

References

- [1] Wm. C. McHarris, F. S. Stephens, F. Asaro, and I. Perlman, Phys. Rev. 144, 1031 (1966).
- [2] J. A. Milsted, A. M. Friedman, and C. M. Stevens, Nucl. Phys. 71, 299 (1965).
- [3] Wm. C. McHarris, Ph.D. thesis, University of California, Berkeley, Lawrence Radiation Laboratory Report UCRL-11784 (1965).

19. The (p,t) Reaction on Pr^{141}

R. Goles

The decay properties of Nd^{139} have been reported in recent studies [1-5]. In the latest of these investigations, a relatively complete decay scheme was proposed on the basis of the γ -rays associated with this decay [5]. In addition, probable nucleon configurations for important transitions between Nd^{139} and Pr^{139} have been postulated in order to explain some of the unusual systematics involved in the decay of the isomeric $11/2^-$ state [6]. The latest decay schemes [5] for both the ground and isomeric species of Nd^{139} are given in fig. 1. Because of the large amount of information already available on the states in Pr^{139} , and because of the significantly different interpretation given to the high-lying states, we decided to investigate the states in this nucleus more thoroughly by the (p,t) reaction on 100% abundant Pr^{141} . In this manner we could characterize those states not reached by the beta decay of the Nd^{139} isomers and could perform meaningful shell-model calculations on these states.

In the present work, $\approx 300 \mu\text{g}/\text{cm}^2$ Pr^{141} targets on $25 \mu\text{g}/\text{cm}^2$ carbon backings were bombarded with 500-nA beams of 40-MeV protons accelerated by the MSU Sector-Focused Cyclotron. The beam resolution as calculated from the measured magnetic fields and slit apertures is commonly better than 40 keV. Details of this transport and beam analysis system have been discussed previously [7]. A dE/dx vs E counter telescope, which consisted of two $\text{Si}(\text{Li})$ detectors, was used to identify and also to measure the energies of the outgoing particles.

The particle-identification system chosen for these experiments depends on the difference in energy lost in the ΔE detector for particles of different mass and charge but the same kinetic energy. The signals from the ΔE and E detectors were summed (called the Σ signal) at the detector telescope, and all three pulses (ΔE , E , and Σ) were passed through charge sensitive pre-amplifiers and sent to the data acquisition area. Fig. 2 shows the summing circuit. Using the electronics set-up shown in fig. 3, a slow coincidence was required between the ΔE and E signals. The latter two signals were then sent to a Northern Scientific dual 8192 ADC. The Cyclotron Laboratory Sigma 7 computer and the acquisition task TOOTSIE [8] were used to analyze the digital

signals received from the pair of coupled ADC's.

The computer code TOOTSIE stores the two-dimensional E - ΔE spectrum and can display it on an oscilloscope. Because of the difference in energy loss (ΔE signal) of particles of different charge and mass for the same total energy (E signal), the different particles seen by the detector telescope fall into distinctly different and nonoverlapping bands in the above display. Figure 4 shows just such a typical two-dimensional display. The code then allows gate lines to be drawn about the desired particle band or bands in the form of polynomial fits to designated points. These gate lines are then used to route all E signals falling within a given set to any of four 8192-channel memory blocks where the total particle energy spectrum is stored.

Because of the very low cross-sections involved in plucking two neutrons out of the closed shell of Pr^{139} , a complete angular distribution at 40 MeV has not yet been completed. But the preliminary results are very reassuring in that numerous states in Pr^{139} are populated, so by next year we hope to have extensive information about states in Pr^{139} .

References:

- [1] K. Ya. Gromov, A. S. Danagulyan, A. T. Strigachev, and V. S. Shpinel', *Izv. Akad. Nauk SSSR, ser. fiz.* 27, 1357 (1963).
- [2] J. Gilat and W. J. Tretyl, LRL Report UCRL-17299, p. 20 (1967).
- [3] J. Lange, KFK Report, summarized in J. Lange, H. Münzel, and I. Leitel, *Radiochim. Acta* 8, 123 (1967).
- [4] K. Ya. Gromov, A. S. Danagulyan, L. N. Nikityuk, V. V. Murav'eva, A. A. Sorokin, M. Z. Shtal', and V. S. Shpinel', *Zh. Eksperim. i Teor. Fiz.* 47, 1644 (1964).
- [5] D. B. Beery, W. H. Kelly, and Wm. C. McHarris, *Phys. Rev.*, scheduled for 20 Nov 1969 issue.
- [6] Wm. C. McHarris, D. B. Beery, and W. H. Kelly, *Phys. Rev. Lett.* 22, 1191 (1969).
- [7] G. Mackenzie, E. Kashy, M. M. Gordon, and H. G. Blosser, *I.E.E.E. Trans. Nucl. Sci.* 450, NS-14, No. 3 (1967).
- [8] D. Bayer and W. Benenson, private communication (1969).

Fig. 19-1(b) The decay scheme of Nd^{139g} , taken from ref. [5].

SUMMING CIRCUIT

Fig. 19-2. Block diagram of the summing circuit of the ΔE and E detectors preceding the preamplifiers.

EXPERIMENTAL ELECTRONICS

Fig. 19-3. Block diagram of the particle-identification electronics used to obtain (p, t) spectra.

Fig. 19-4. Two-dimensional contour display of the lines one obtains in a ΔE vs total E (Σ) particle-identification experiment. By drawing lines (fit to polynomials) enclosing the region of interest, one can obtain a digital selection of the desired events.

20. In-Beam Study of the Decay of Sc^{40} to Ca^{40}

J. Black

The bulk of the available information on the levels in Ca^{40} has been obtained through scattering experiments [1-5]. Although some earlier work on Sc^{40} decay has been performed [6] the rather short half-life of Sc^{40} has prevented detailed γ -ray investigations of the levels in Ca^{40} . However, with the advent of slow-pulsing techniques for the MSU Cyclotron, together with relatively good on-line detection systems, it is now possible to study the gamma decay of Sc^{40} to obtain the levels in Ca^{40} with some degree of confidence. Isotopically separated Ca^{40} in the form of the carbonate, will be placed in a goniometer which incorporates a precision positioning device and target changer. Beam optics before this instrument produce a well-focused proton beam at the target position, which together with a shielded beam dump, provides a relatively low background condition for counting. A (p,n) reaction using 20-MeV protons will be used to produce the 0.179-second Sc^{40} . It is anticipated that the beam will be pulsed on for 500 milliseconds, then allowing 100 milliseconds for the electronics to recover from the beam burst, and finally counting the sample for approximately 700 milliseconds. Our initial bombardments should have been completed before this Annual Report gets printed.

References

- [1] C. M. Braams, Phys. Rev. 101, 1764 (1956).
- [2] P. M. Endt and C. Van Der Leun, Nucl. Phys. A105, 1 (1967).
- [3] J. R. Erekine, Phys. Rev. 149, 854 (1966).
- [4] W. S. Gray, R. A. Kenetick, and J. J. Kraushaar, Nucl. Phys. 67, 542 (1965).
- [5] A. Springer and B. G. Harvey, Phys. Letters 14, 116 (1965).
- [6] E. Kashy and J. L. Snelgrove, Phys. Rev. 172, 1124 (1968), and references therein.

21. The Possibility of β -Delayed α 's from Ga^{62} or Ga^{63}

G. Giesler

Delayed particle emission has been known for many years. In fission, nuclei are produced but are so far from β stability that the β decay leads to neutron unstable levels, which then emit neutrons rather than γ -rays to move toward stability. Some examples of this are $\text{I}^{137-139}$ and Sb^{135} . Light nuclei commonly decay by β -delayed proton emission. The wide level spacing and large decay energies present there increase the probability of delayed particle emission. Also present to a small extent is β -delayed α emission. It has only been observed in the β decay of nuclei of the type $N=Z\pm 2$, where Z is odd decaying to an $N=Z$ even-even nuclei. Presently, only 8 nuclei have been observed to produce β -delayed α 's: Li^8 , B^8 , B^{12} , N^{12} , N^{16} , Na^{20} , Al^{24} , and Cl^{32} . The longest observed β half-life is the 7.4 sec one of N^{16} .

In the article reporting the discovery of 33-sec Ga^{63} , Nurmi and Fink [1] reported that their purpose was to look for delayed alphas, but were unable to detect any. Their estimate of the possibility of β -delayed α emission was based on the mass tables calculated by Myers and Swiatecki [2]. The α binding energy of Zn^{63} based on these calculations is 1.738 MeV and that based on experimental data of MTW [3] is 3.49 MeV; these can be compared with a Ga^{63} β -decay energy of ≈ 5.5 MeV. This difference in α -binding energies could mean the difference between observing or not observing the β -delayed α 's.

By going to the as of yet unreported Ga^{62} , one obtains the more favorable situation of 10.7 MeV estimated β -decay energy vs 1.162 MeV (calculated [2]) or 3.33 MeV (experimental [3]) α -binding energies. Ga^{62} , however, has a correspondingly higher Q value for its production and a shorter half-life. F6nges and Rupp [4] report that, based on the Myers-Swiatecki nuclear mass-tables and work by Tassgapers and Nurmi [5], the heaviest delayed α -active isotope of gallium is Ga^{65} with an α -decay energy of 4.42-MeV.

Because of the short half-lives involved, progress on this has

been slow. Ga^{63} has been produced, but the presence of delayed alphas is inconclusive. Work on the Ga^{62} is also in progress.

References

- [1] M. Nurmia and R. Fink, Phys. Lett. 14, 136 (1965).
- [2] J. Mattauch, W. Thiele, and A. Wapstra, Nucl. Phys. 67, 1,32,73 (1965).
- [3] T. Fénges and B. Rupp, ATOMKI Kozlem. 10, 11C (1968).
- [4] W. Myers and W. Swiatecki, UCTL-11980 (1965).
- [5] R. Taagepera and M. Nurmia, Ann. Acad. Sci. Fennicae, A VI, No. 78, 1 (1961).

22. On-Line γ -Ray Spectroscopy

R. Warner

A program of γ -ray spectroscopy on-line with the MSU cyclotron has been initiated in cooperation with Professor W. H. Kelly in the Physics Department. The techniques currently being used are similar to those first described by Yamazaki and Ewan [1], with modifications to take advantage of the Cyclotron Laboratory's excellent on-line computing facility and recent developments in commercially available timing circuitry [ORTEC Model 453 Constant Fraction Timing Discrimination]. The computer software used is called TOOTSIE and was written by D. L. Bayer of the Physics Department. Although TOOTSIE was originally written for particle identification, it is sufficiently flexible for general multiparameter data acquisition and is used unmodified for the energy vs time experiments in progress.

The MSU Cyclotron Laboratory is uniquely well suited for reaction γ -ray spectroscopy in other ways as well: A. The time-width of the beam bunch from the cyclotron has been measured to be as narrow as 0.1 nsec; therefore, the time resolution of the detectors used will determine the lower limit of lifetimes which can be measured directly. The upper limit for such measurements will soon be extended without limit by a beam pulsing system currently being perfected. B. A precision goniometer [2] which was originally designed for charged particle experiments is ideal for measuring γ -ray angular distributions. Preliminary measurements of the neutron flux in the vicinity of the goniometer show the target-out/target-in ratio to be only 5%, and minor modifications are expected to reduce the neutron and γ -ray backgrounds further.

Two experiments are in progress: A search is under way for a short-lived isomer in Xe^{128} . Four other $N=74$ nuclei are already known [3] to contain isomeric states, and the current work should establish whether the same microscopic configuration is responsible for all of these states. A study of the relative sensitivity of planar-drift and five-sided drift $\text{Ge}(\text{Li})$ detectors to damage by fast neutrons is proceeding in cooperation with Nuclear Diodes, the manufacturer of the detectors.

A comparative study of population by (α, xn) and (τ, xn) reactions will be initiated soon, and γ -ray spectroscopy with (heavy ion, xn) reactions will be done as soon as heavy ion beams are available from the cyclotron.

References

- [1] T. Yamazaki and G. T. Ewan, Nucl. Instr. Methods 62, 101 (1968).
- [2] K. M. Thompson and C. R. Gruhn, Bull. Am. Phys. Soc. 13, 701 (1968).
- [3] P. Ward, F. S. Stephens and R. M. Diamond, *Nuclear Chemistry Division Annual Report, 1968*, UCRL-18667, 54 (1969). R. A. Warner, Ph.D. Thesis, University of California, Davis (1969).
R. A. Warner and J. E. Draper, Phys. Rev. (to be published).

23. Simple Shell-Model Calculations on Three-Quasiparticle States in Pr^{139}

W. McHarris

(in cooperation with R. Muthukrishnan, now at North Texas State University, and A. Kromminga, Calvin College)

During the past year we discovered a rather startling phenomenon in the decay of $11/2^- \text{Nd}^{139m}$. Its ϵ decay preferentially populates six high-lying, high-spin, odd-parity states in Pr^{139} . We interpreted this as the structure of Nd^{139m} being peculiarly suited for directing its decay into a multiplet of three-quasiparticle states [1]. The transition can be characterized as

$$(\pi d_{5/2})^2 (\nu d_{3/2})^{-2} (\nu h_{11/2})^{-1} \rightarrow (\pi d_{5/2}) (\nu d_{3/2})^{-1} (\nu h_{11/2})^{-1}.$$

This decay scheme can be seen in fig. 19-1 of this Annual Report. A symbolic shell-model representation of the important transitions between various states in Nd^{139} and Pr^{139} is reproduced in fig. 1.

The states in this proposed three-quasiparticle multiplet are important not only because of the peculiar decay itself, but primarily because they afford a wealth of information about *specific, well-defined* states in a nucleus near 2 MeV that normally would be available only for states near the ground state. For example, the states deexcite by many apparently enhanced γ -ray transition to other states in the multiplet, but the transitions out of the multiplet appear to be very highly hindered -- there are many instances where $M1$ or $E2$ transitions in the neighborhood of 100 keV are much faster than $E1$'s of several MeV. Thus, the β decay itself should give much information about the major components of the wave functions of these states, and so also should the enhanced γ -ray transitions. Once the states are fairly well characterized, the retarded transitions to states lower in Pr^{139} should enable us to learn something about the small admixtures in the states.

With this in mind, very simple-minded shell-model calculations were undertaken to determine what sort of success one could obtain when using an extremely truncated basis space. For calculation of negative-parity states in Pr^{139} , only two configurations were considered, $(\nu d_{3/2})^{-2} (\pi h_{11/2})$ (the con-

figuration of the 11/2- "isomeric" state at 821.9 keV) and $(\nu h_{11/2})^{-1}(\nu d_{3/2})^{-1} \times (\pi d_{5/2})$ (presumably the major component in the three-quasiparticle multiplet). Such a configuration would produce the following possible states: 3/2, 5/2(2), 7/2(4), 9/2(5), 11/2(6), 13/2(5), 15/2(4), 17/2(2), and 19/2, all with negative parity. As the states populated by Nd^{139m} ϵ decay are most likely 11/2- or 9/2-, we concentrate on these in what follows. The unperturbed energies were taken to be

$$E_0(\nu d_{3/2})^{-2}(\pi h_{11/2}) = 1.1 \text{ MeV (from Pr}^{141} [2])$$

and

$$E_0(\nu h_{11/2})^{-1}(\nu d_{3/2})^{-1}(\pi d_{5/2}) = 746 \text{ keV (from Ce}^{139} [3]).$$

The interaction used was that proposed by True [4],

$$v = V_0 e^{-r^2/r_0^2} (W + MP^x + BP^y + HP^xP^y),$$

where P^x is the space exchange, P^y is the spin exchange, and the other values are $V_0 = -48.75$, $r_0 = 1.85$ fm, $W = 0.42$, $M = 0.42$, $B = 0.08$, $H = 0.08$, and $\hbar\omega = 41/A^{1/3}$ (for $A = 139$). The first results are shown in fig. 2, where the energies are compared with the experimentally determined energies. It is quite heartening that, of the possible states, two 11/2- and one 9/2- states lie high enough so that their population is either unlikely or impossible, leaving only eight possible states to be populated. One of these is the 821.9-keV 11/2- "isomeric" state and one could be the state (11/2-) observed at 1396.9 keV, which our experimental information did not allow us to characterize very completely [5] -- both states should be primarily of the configuration, $(\nu d_{3/2})^{-2} \times (\pi h_{11/2})$. The other six, three 9/2- and three 11/2-, lie at approximately the correct energies and consist primarily of the $(\nu h_{11/2})^{-1}(\nu d_{3/2})^{-1}(\pi d_{5/2})$ configuration, so they well could be good approximations to the experimentally-observed three-quasiparticle states.

Of course, energy agreements are not really terribly good tests of shell-model calculations, and one must try the transition probabilities as well. Our first results on calculating electromagnetic transition probabilities have been less successful, and it appears that we shall have to include other configurations, especially the configuration, $(\nu h_{11/2})^{-1}(\nu d_{3/2})^{-1}(\pi d_{5/2})^2(\pi g_{7/2})^{-1}$. That this configuration is indeed important can be seen, for example, from the

transition between the 2174.3- and the 113.8-keV states. However, inclusion of this three-hole, two-particle configuration complicates the picture considerably, and results have not been obtained as of yet. Similarly, calculation of the β transition probabilities quantitatively also requires the Nd^{139m} state to be the three-hole, two-particle configuration, depending on how inert the $d_{3/2}$ neutron pair (holes) really is.

At present the calculations are being continued, for the amount of concrete information available from this nucleus and from certain of its neighbors [1] should provide a severe work-out of shell-model calculations. Perhaps the most gratifying part of the investigation is that one can obtain reasonable results with such an oversimplified configuration space.

References

- [1] Wm. C. McHarris, D. B. Beery, and W. H. Kelly, Phys. Rev. Letters 22, 1191 (1969).
- [2] H. Baer, private communication (1968).
- [3] D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev., scheduled for the 20 Nov 1969 issue.
- [4] W. W. True, Phys. Rev. 130, 1530 (1963).
- [5] D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev., scheduled for the 20 Nov 1969 issue.

TRANSITIONS FOLLOWING THE DECAY OF $\text{Nd}^{139\text{m}}$

Fig. 22-1. Symbolic shell-model representation of some important transitions between states in Nd^{139} and/or Pr^{139} .

Fig. 22-2. Comparison of our shell-model calculations with the observed state energies. The 9/2- and 11/2- states are offset in the calculated column for purposes of clarity. The 820-keV 11/2- state was used to normalize the energies. The states marked with *** contain the largest components of the $(\nu d_{5/2})^{-2}(\pi h_{11/2})$ configuration, although there is a fair amount of mixing and fragmentation of this configuration. The other states contain only small amounts of this configuration and consist primarily of $(\pi d_{5/2})(\nu d_{3/2})^{-1}(\nu h_{11/2})^{-1}$.

23. M4 Transition-Probabilities in Nuclei below N=82

R. Eppley

Since our first observation [1] of Gd^{145m} , an $M4$ transition ($\nu h_{11/2} \rightarrow \nu d_{3/2}$), we have been looking for possible explanations for the deviations of the experimental matrix elements from the predictions of the single-particle shell model. We are presently interested in both the $N=81$ and the $N=79$ isotones that have experimentally measured $M4$ transitions. The energies and transition probability ratios are shown in fig. 1 for the $N=81$ series.

These $M4$ transitions, particularly the $N=81$ series, should be fairly pure single-particle transitions. The $11/2^-$ state is of high spin and is the opposite parity from the nearby levels; thus, configuration mixing of this state should indeed be small. However, the $3/2^+$ state could have small admixtures, which might account for the deviations from the shell-model predictions.

Gupta and Lawson [2] looked at this question of configuration mixing for $M4$ transitions. They assumed a wave function of the form $\phi_0 + \sum_i \alpha_i \phi_i$, where ϕ_0 is the wave function for the pure configuration and ϕ_i are the wave functions for the admixed ones. This is a good representation when the admixtures are small as they should be in this case. Their conclusion was that the experimental deviations probably cannot be accounted for solely by configuration mixing.

Kotajima and Morinaga [3] come to the same conclusion, that the effect cannot be due only to configuration mixing. They mention that better agreement should be reached by using a realistic potential in the radial matrix elements.

The latest published calculations are by Jansen, Morinaga, and Signorini [3]. In addition to confirming our experimental observation of the Gd^{145m} isomer, they made calculations comparing the experimental matrix elements of the $N=81$ $M4$ isomers with theoretical estimates corrected for configuration mixing and using "realistic" radial matrix elements.

The radial matrix elements are given by

$$|M_r|^2 = \left| \langle R_p(r) \left| \left(\frac{r}{R_0} \right)^{L+1} \right| R_i(r) \rangle \right|^2,$$

where $R_0 = r_0 A^{1/3}$. This was solved by solving the radial part of the Schrödinger equation using the potential

$$U(r) = V_{ws}(r) + V_{so}(r),$$

where V_{ws} is the Wood-Saxon potential and V_{so} is the spin-orbit potential. Values given by Horie and Oda [5] were used to correct for configuration mixing. This approach is demonstrated to remove a large part of the Z dependence from the matrix elements. However, the matrix elements are still smaller than the theoretical values.

For our part then, we are exploring another approach to the problem in terms of the quasi-particle formalism. We may be able to show that these transitions are hindered, relative to the single particle estimates, because of varying admixtures of a particle-hole component in the matrix elements. This component will have a lower probability than hole-hole or particle-particle components, thus hindering the transitions. The quasi-particle formalism is described in terms of a system of nucleons subject to a nuclear force including a pairing force. A nuclear ground state equation for such a system can be written after the fashion of the superconductivity equation given by Bardeen, Cooper, and Schrieffer [6],

$$\phi_{\text{BCS}} = \prod_v (U_v + V_v a_v^\dagger a_v^\dagger) |0\rangle.$$

This describes a state having a probability amplitude U_v^2 that the level v is occupied and an amplitude V_v^2 that level v is unoccupied. $a_v^\dagger$ and a_v are creation operators which create particles in the level v when operating on the vacuum state, $|0\rangle$. From this equation it is seen that particles are always created in pairs in the same level (one particle time reversed). This resulting even-even system is then considered as a quasi-particle vacuum state, denoted by $|\tilde{0}\rangle$. All quasi-particle excitations can be expressed relative to this vacuum by operating on the quasi-particle (even-even) vacuum state as follows.

$$\psi_{\text{quasi-particle}} = a_v^\dagger |\tilde{0}\rangle,$$

where $\alpha_v^\dagger$ is a quasi-particle creation operator and is of the form $\alpha_v^\dagger = U_v \alpha_v^\dagger - V_v \alpha_{\bar{v}}$.

When a single quasi-particle is formed, this amounts to an alteration in the occupation of one state. Thus, the quasi-particle is a mixture of a particle in the state v and a hole in state $\bar{v}$. In this sense the quasi-particle representation is a configuration mixing representation.

In this manner it should be possible to describe the $N=81$ $M4$ transitions in the quasi-particle formalism. The matrix elements can be written in the form $\langle \psi_{h11/2} | M4 | \psi_{d3/2} \rangle$, where the wave functions are written as ψ quasi-particle mentioned above. One of the problems of this method is to determine the values V_v^2 and U_v^2 for the levels in question. One of the primary means of experimentally determining these quantities is by the (d,p) and (d,t) reactions. The intensity of the reaction depends on the probability that a neutron is in the state v , therefore is a measure of V_v^2 and U_v^2 . We are presently investigating the research done with these reactions for the $X_{81\pm1}^{A\pm1}$ nuclei in this region, where X represents an odd-mass nucleus with 81 neutrons and exhibits the $M4$ isomerism. Since there are seven members presently in the $N=81$ set, there are a total of 14 reactions from which it would be desirable to have information. Of these reactions eight are with a stable target material and thus practicable.

References

- [1] R. E. Eppley, Wm. C. McHarris, and W. H. Kelly, Bull. Am. Phys. Soc. 13, 1467 (1968).
- [2] K. K. Gupta and R. D. Lawson, Phys. Rev. 114, 326 (1959).
- [3] K. Kotajima and H. Morinaga, Nucl. Phys. 16, 231 (1960).
- [4] G. Jansen, H. Morinaga, and C. Signorini, Nucl. Phys. A128, 247 (1969).
- [5] H. Horie and T. Oda, Progr. Theor. Phys. 32, 65 (1964).
- [6] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. 108, 1175 (1957).

Fig. 23-1. Systematics of the $N=81$ $M4$ isomeric transitions. At top the energy differences between the $h_{11/2}$ and $d_{3/2}$ states are given, and at the bottom the reduced transition probabilities (radial matrix elements squared).

25. Status of the PDP-9 Computer

W. McHarris

The PDP-9 computer, which was purchased in 1968 to serve our group primarily as a multi-purpose, multichannel analyzer, has remained in almost constant use during the past year. In fact, we have recently been forced to go to a sign-up procedure in order to insure that there be a smooth flow between data collection experiments, data reduction, and systems programming. The original configuration, consisting of a basic 8192 18-bit word machine, with teletype and fast paper tape punch (50 cps) and optical reader (300 cps), plus a type NH14B nuclear interface to a Northern Scientific type NS-620 dual 12-bit ADC and a 34C interface to a Tektronix RM503 oscilloscope, was found to be more than adequate for a great variety of pulse-height experiments and applications of varying complexity, but the I/O was found to be quite cumbersome. This was especially true when it was necessary to punch out many spectra in a short time, as when working with short half-lives.

This spring the computer has been improved by the addition of a type TC02 DECTape control and two TU55 DECTape transports, which have speeded up the I/O considerably. The nuclear interface was also updated from a type NH14B to an NH14C, thereby implementing the "list" mode as well as the accumulator (AC) and add-one-to-memory (MB+1) modes. Programs for operation of the list mode are now in preparation (see next section), and we hope to be able to run a significant number of our 2-d megachannel spectra (section 31) on the PDP-9 in the future. As an aid to spectrum analysis, we have acquired a Tektronix type 611 storage scope, and a type VP09 interface has been ordered to drive this scope. The VP09 comes with the same device numbers as the existing 34C interface, so we plan to change the device numbers of the VP09 ourselves, allowing the simultaneous operation of both scopes. A switch will be used to restore the original device numbers when needed to use the DEC-furnished I/O software for the VP09.

The PDP-9 has been reasonably trouble-free, with the significant exception of the fast paper tape punch. This accounted for three-fourths

of the down time and service calls. Even after being sent back to the factory and being rebuilt, this punch continued to prove troublesome, so it was finally replaced with another unit that appears to operate satisfactorily.

As mentioned in several places elsewhere in this Annual Report, our spectrum analysis routines have proven to be fairly severe bottlenecks. This is especially true with the scope programs on the Cyclotron Laboratory Sigma-7 computer, for the display takes up enough of the central processor time that it can slow down other functions of the time-shared computer. Thus, those operating the cyclotron are often loath to have such a display operating while they are taking data. Another bottleneck is the paper tape link between the two computers: block binary output from the PDP-9 is read into the Sigma 7 through a teletype reader. The PDP-9 now reads the IBM-format tapes from all the fixed-wired analyzers in the laboratory now and converts the tapes to block binary for the Sigma 7, so this bottleneck is emphasized. We are now planning to interface the two computers, and it is hoped that early next year we can transfer data between them in blocks. This will allow us to make much more use of the PDP-9 for data analysis, relieving the overloaded Sigma 7 of some of the more routine programs.

The next two sections in this Annual Report give some idea of the types of data collection programs and auxiliary programs that are now in use on the PDP-9. In fig. 1 an idea of the physical layout is shown.

Fig. 25-1 (next page). Photograph of the PDP-9 with a sampling of its associated equipment, including its operators, who double as cyclotron operators and as nuclear chemistry graduate students. Seated: Richard Eppley and Ray Doeblner. Standing: Greg Giesler, Ken Kosanke, Jerry Black, Richard Firestone, Jim Cross, and Ron Goles.

26. Multianalyzer Programs for the PDP-9 Computer

O. McHarris

Further progress has been made with the PDP-9 multianalyzer programs so as to enable this computer to carry a larger burden of our data acquisition and analysis. The programs furnished by DEC still form the heart of our multianalyzer software, but fairly extensive modifications have been made or are in progress to streamline them. Among these are the following:

1) Modification of the one-dimensional programs so as to allow one to choose at will the size of a spectrum and the number of spectra taken at a given time through routing. This is especially useful for following short-lived activities, and it will be incorporated, with computer interrupts, into our rabbit hardware (cf. § 29).

2) Enable the computer to read various paper-tape formats on call. At present the PDP-9 reads the IBM (odd parity) formats from all the various analyzers in the laboratory and converts this to block binary, which can be read by the Sigma-7. It can also read various irregular ASCII formats, including the one in use at the Yale HIAC (§ 11). This rather circuitous path is necessary because of the slow paper tape input on the Sigma 7 -- about 10 min to read a 4096-channel block binary tape vs over an hour for the IBM format.

3) Alteration of our overflow routines to allow more than the $2^{18}-1$ maximum number of events per channel. The overflow can either be typed out on the teletype, whereupon the computer resets and continues to collect data as before, or it can signal dumping the spectrum onto DECTape, whereupon the reset starts the next spectrum from a cleared memory. The latter is useful when the overflow starts to occur frequently, as when the low-energy Compton background approaches maximum. An "ad hoc" double-precision modification, in which one can set aside extra words for double-precision channels only on the (not necessarily previously predicted) channels that overflow is still in progress.

4) Modification of the two-dimensional program to allow use of the listing mode for megachannel analysis. With only DECTapes this is still "mini-megachannel" analysis but should improve with IBM tape or interfacing to the Sigma-7.

5) Improvements of the data-analysis routines to approximate the program MOIRAE (§ 32), which is one of our major bottlenecks. Implementation of this awaits the larger Tektronix scope interface, promised early in Feb.

27. Auxiliary Programs

R. Eppley

While the PDP-9 computer has been indispensable for data acquisition, it has also proven to be very useful in performing various routine types of numerical calculations. The FORTRAN programming capability of the PDP-9 has been used for this purpose.

All of the programs are stored on DECTape and can be run under the keyboard monitor system. Each is brought into core by the appropriate TTY command. Directions for the use of each program is included at the beginning of each program listing.

The following is a short description of some of the available programs:

1. ARITH: This program is for carrying out simple numerical calculations. It allows the PDP-9 to be used as an electronic calculator. Presently it is capable of performing addition, subtraction, multiplication, division, square, square root, cube, cube root, common logs, natural logs, antilogs, and exponential calculations. Each type of calculation is specified by a numerical code. The code and the numbers to be manipulated are entered on the TTY. The answer is presented as print-out on the TTY.
2. INTEN: This program is for the calculation and renormalization of γ -ray intensities. If peak areas are input (such as from MOIRAE: see discussion in Data Analysis section) the appropriate detector efficiency correction is applied to each of the peak areas in order to arrive at the correct transition intensity. Transition intensities can then be renormalized to other transitions without having to re-enter the numbers on the TTY. Intensity ratios can be renormalized in the same manner without applying the detector efficiency correction.

3. FERM3: For the calculation of Fermi functions. This program was developed to be an aid in the construction of Fermi-Kurie plots for electron and positron spectra. Rather than being a program for the extrapolation of values from published tables of Fermi functions, this program, using a power series method, calculates the function needed at each point of the spectrum.

The Fermi function can be written as [1]

$$f(Z, n) = n^{2+2s} e^{\pm \pi \delta} |\Gamma(1+s+i\delta)|^2$$

where Z = atomic number

n = electron momentum

$$s = \sqrt{1-\gamma^2} - 1$$

$$\gamma = Z/137.0$$

$$\gamma = Z\sqrt{1+n^2}/137.0 \quad n$$

The difficulty in the numerical computation is in evaluating the complex gamma function. It is of the form $|\Gamma(x+iy)|^2$ where $x = 1+s$ and $y = \delta$

This expression is evaluated by use of the expansion [2]

$$\left| \frac{\Gamma(x+iy)}{\Gamma(x)} \right|^2 = \prod_{n=0}^{\infty} [1+y^2/(x+n)^2]^{-1}$$

also,

$$\Gamma(x) = 1/\prod_{k=1}^{\infty} C_k x^k$$

where the coefficients C_1 through C_{26} are given in reference [2]. The function that we need is then constructed from the product of those two series.

$$|\Gamma(x+iy)|^2 = \left(\prod_{k=1}^{\infty} C_k x^k \right) \left(\prod_{n=0}^{\infty} [1+y^2/(x+n)^2]^{-1} \right)$$

The number of terms needed varies depending on the values of Z and n input.

4. DECAY: This program is a computational aid for the construction of

electron capture decay schemes. For a decay scheme of any complexity there are numerous time consuming calculations for which a change of any of the numbers means that the whole procedure must be repeated. Thus, this program can save a large amount of time. The program calculates electron capture ratios, K -capture intensities, and total capture intensities including capture to the ground state. It also calculates the percent decay to each level and the $\log ft$ values for each of the electron capture transitions.

5. PROB: This program is for the calculation of transition probabilities and half-lives of γ -ray transitions based on the single particle equations of Moszkowski. The only input information needed is the type of transition (neutron or proton), the transition energy and the value of the statistical factor. The output consists of the transition probability and half-life for the transitions $E1$ through $E5$ and $M1$ through $M5$.
6. PLYFT: This is a general polynomial least-squares program. "Best" curves can be calculated for polynomials up through degree 10. The output consists of a reduced chi square, the coefficient for each term of the polynomial and the calculation of any additional points specified in the input information.

The next two programs are used for the interpolation of the theoretical internal conversion coefficients from the published table.

7. ICC: This is a general program which will interpolate the conversion coefficient for any energy in the range of the table by providing a least-squared polynomial fit of a specified order to the log-log plot of the tabular values. It is

generally the case that a polynomial fit over a large region (say 50 keV-1500 keV) is quite unsatisfactory. However energies above about 150 keV can usually be fit quite well by a quadratic fit. Sometimes trial and error is necessary for a specific case. All energies of a given multipole order which fall within the energy limits of the "best" curve can be calculated at one time.

8. INTERP This is a program based on the theory of splines and adapted from the program given by Hagen and Seltzer [3]. With this method, the entire energy region given in the table can be covered at one time. This is accomplished by a piecewise fitting of the tabulated points to a cubic equation. As stated by Hager and Seltzer, this program minimizes the integral of the mean-square-difference of the second derivative of the trial curve and that of the curve which it approximates. This program does not work in some energy regions since there is not always a sufficient number of points to satisfy the equation.

9. MEL This program is for the calculation of both experimental and theoretical γ -ray half-lives and nuclear matrix elements using the extreme-single-particle model equations. The experimental half-life is corrected for internal conversion (which the program interpolates from the theoretical tabular data) to arrive at the γ -ray half-life. As an example, the square of the $M4$ matrix element is calculated from the equation

$$|M4|^2 = 1.16 \times 10^6 \times r_0^{-6} \times A^{-2} \times E_{\gamma}^{-9} \times S^{-1} \times T_{\frac{1}{2}}^{-1} \\ \times \left(1 + \alpha_K \frac{(1 + \alpha_L + \alpha_M + \dots)}{\alpha_K} \right)^{-1}$$

where:

$$r_0 = R_0 A^{1/3}$$

A = mass number
 E_γ = gamma transition energy
 S = statistical factor
 $T_{1/2}$ = experimental half-life
 α_i = conversion coefficients

With these results, comparisons can be made between the experimental results and shell model theory.

10. VC: This program is for the calculation of vector coupling coefficients. This is accomplished by use of the general equation as given by Judd [4].
11. RACAH: This program is for the calculation of Racah coefficients utilizing the general formula as given by Judd [4].
12. DP: For the calculation of the activities present at a given time for a chain of radioactive products. It is useful to know at what time, after a cyclotron bombardment, which decay products will cause measurable interference in the γ -ray spectrum of the isotope being studied. This program solves the Bateman equations with the initial conditions being $t = 0$; $N_1 = N_1^0$, $N_2^0 = N_3^0 = \dots N_n^0 = 0$. Thus, the number of atoms of the n th member of the chain, at time t , is found from the general equation

$$N_n(t) = C_1 e^{-\lambda_1 t} + C_2 e^{-\lambda_2 t} + \dots + C_n e^{-\lambda_n t}$$
 The program output gives $N_n(t)$ and the activity of component n , $(A_n(t))$, vs time for each species in the chain for the time interval and increments specified in the input.
13. TRIGA This program is for the calculation of the expected activity of samples submitted for irradiation with the MSU TRIGA reactor. The calculation assumes a 10 mg target and a neutron flux of 2×10^{12} neutrons/cm². sec. The input consists of the

mass number of each stable isotope for the element, the half-lives of each single-neutron capture species, the isotopic abundance of each stable species, all pertinent neutron capture cross-sections and the bombardment times which are to be used. The output gives the bombardment time vs activity at the end of the bombardment for each of the active species.

References

- [1] *Tables for the Analysis of Beta Spectra*, NBS Applied Mathematics Series 13.
- [2] *Handbook of Mathematical Functions*, Ed. M. Abramowitz and I. Stegun, Dover Publications, Inc., New York (1965).
- [3] R. S. Hager and E. C. Seltzer, *Nuclear Data*, Volume 4, Number 1 and 2, February 1968, Section A.
- [4] Brian R. Judd, *Operator Techniques in Atomic Spectroscopy*, McGraw-Hill, New York (1963).

28. Cyclotron On-Line Mass Separation

K. Kosanke

(in cooperation with H. Hilbert of the
Cyclotron Laboratory)

Historically, the study of nuclides farther and farther from β -stability has been hindered by several nuclear properties. The first is the rapidly diminishing half-lives of the reaction products. This effectively eliminates the possibility of doing a chemical separation on the target and generally forces one to use some sort of pulsed cyclotron beam technique, with counting experiments conducted in the beam area. The second is the overlapping of excitation functions. This results in the production of whole collections of different nuclei in production reactions, a very serious complication in view of the increasing complexities in the decay of nuclides far from β -stability.

A system providing, on a continuous basis and in times considerably less than one second, reaction products that have been isotopically separated would be an especially powerful technique in combating these problems. We are constructing such a system. It is essentially just combining of a couple of proven techniques, helium thermalization and jet transport of nuclear recoils [1] and electrostatic quadrupole mass filtering [2].

The on-line mass separation system under construction and its operation are shown schematically in figure 1. A collimated cyclotron beam enters the apparatus, striking a thin target in an atmosphere of helium (≈ 2 atm pressure). Those nuclei near the back of the target, interacting with the beam, will be recoiled out of the target into the helium. Information on some possible cyclotron beams, nuclear reactions, and recoil energies is listed in Table 1. The recoils are then thermalized ($E_T \approx 3\text{eV}$ [3]) by collisions with the helium atoms. If conditions are met such that laminar flow is set up in the capillary tube [1] the nuclear recoils will be swept out of the system with the helium at approximately sonic velocity. The capillary is run through the cyclotron vault wall into a relatively low background area. Then before the recoils leave the capillary they are ionized by a discharge through the helium. As the recoils and helium

leave the capillary they will diverge, because their velocity drops and because of space charge effects. However, the helium will diverge much more rapidly (because of its low mass) than will the nuclear recoils. This property is then used to skim off most of the helium such that the quadrupole will be run at a pressure of $\approx 10^{-5}$ torr. The ionized recoils are reaccelerated slightly ($E_p \approx 40\text{eV}$) and focussed into the quadrupole assembly [4]. Only those recoils of the proper e/m (axial velocity independent) will have stable orbits through the electrostatic quadrupole [2]. As the recoils exit the quadrupole assembly they are either drawn into a particle multiplier for setting up the instrument or drawn onto a mylar film and collected for nuclear counting experiments. The physical location of the equipment in the cyclotron experimental areas is shown in figure 2.

The e/m separated isotopes from the quadrupole can be studied using any of several techniques: γ -ray spectrum analysis using solid state detectors, internal conversion electron spectrum analysis using solid state detectors, the MSU $\pi\sqrt{2}$ beta spectrometer or eventually using the MSU six-sector beta spectrometer on-line, or α and β decay using solid state detectors. Also, many of the product nuclides would lie close enough to the line of particle instability that decays such as beta-delayed proton emission [5] or α emission [6] might be observed.

Sometimes the identification of mass number will not be enough to identify a species uniquely, such cases as when the reactions (τ, pxn) and $(\tau, (x+1)n)$ are competing reactions. In these instances the MSU six-sector beta spectrometer could be used to determine atomic number by looking at K -conversion electron energies.

Clearly this collection of equipment (MSU sector-focused, isochronous cyclotron, on-line electrostatic isotope separator, MSU six-sector beta spectrometer) is uniquely well suited for studying nuclides far from the line of beta stability.

References

- [1] R.D. MacFarlane, R. A. Gouch, N. S. Oakey, and D. F. Torgerson,
ORO-3820-2 (1969); Nucl. Inst. and Meth., to be published.
- [2] W. Paul, H. P. Reinhard, and U. von Zahn, Z. Phys. 152, 143
(1958).
- [3] K. Valli and E.K. Hyde, Phys. Rev. 176, 1377 (1968).
- [4] Extranuclear Labs. Model 324-9 with 3/4" quadrupole rods and a
D₂ head.
- [5] P.L. Reeder, A.M. Poskanzer, R.A. Esterlaund, and R. McPherson,
Phys. Rev., 147, 781 (1966).
- [6] See paper 21 in this report.

Table 1

Some of the Considerations Necessary in the Design
of Experiments Using the Isotope Separation System.

MSU Cyclotron beams ^a	Maximum cyc. Beam currents ^b	Ag ¹⁰⁹ Target		Pb ²⁰⁸ Target	
		Possible reaction	Recoil energy ^c	Possible reaction	Recoil energy ^c
30 MeV p	~20 μ A	(p,4n)	0.28 MeV	(p,4n)	0.15 MeV
50 MeV p	~20 μ A	(p,7n)	0.48 MeV	(p,6n)	0.25 MeV
50 MeV He ³	~3 μ A	(He ³ ,9n)	1.45 MeV	(He ³ ,9n)	0.74 MeV
76 MeV He ³	~10 μ A	(He ³ ,12n)	2.28 MeV	(He ³ ,11n)	1.14 MeV

^aIt is expected that the MSU cyclotron will also be producing C¹², N¹⁴,
and O¹⁶ beams shortly.

^bIndications are that the beam currents used will be limited by our
ability to cool our targets and not by the cyclotron itself.

^cAssumes isotropic emission of neutrons and no energy degradation in the
target material.

Fig. 28-1. Block and schematic diagrams of the on-line isotope separator.

29. Rabbit---Pneumatic Target System

K. Kosanke and J. Black

In the study of moderately short lived nuclides, it is undesirable to use pulsed cyclotron beam techniques (with their inherent problems background and radiation damage to detectors) if it is possible to remove the target for counting before the ratio of parent activity to interfering activities drops too low. As a result, a pneumatic target system (rabbit) has been developed in an attempt to study these short-lived β -decay nuclides ($t_{1/2} \geq 1$ sec.) and, further, to make more efficient use of cyclotron time in general.

The target holder (rabbit) and the cyclotron beam end of the system (rabbit hutch) are shown in figure 1. The rabbit is 3" long and 2" in diameter with the target carried in the central open space. The target is both air cooled directly and water cooled by conduction through the holder. The rabbit is reproducibly oriented ($\pm 3^\circ$) with respect to the cyclotron beam. The

mechanism is to rotate the rabbit by a jet of air on a set of fins near its base (dark portion near rabbit's base) then to stop it with the attraction between a small magnet mounted in the top portion of the rabbit and an electromagnet in the rabbit hutch.

The rabbit is propelled through a cellulose acetate piping (with bends made of poly vinyl chloride pipe) by setting up a pressure difference across the

rabbit. We are using ≈ 20 pounds pressure on one side and a partial vacuum, created by a large commercial vacuum cleaner, on the other side. At the end of its path, the rabbit is stopped with an air cushion, created by closing a solenoid valve to the partial vacuum source when the rabbit (and its small

magnet) passes a sensitive coil around the pipe.

Preliminary tests of the completed system indicate that we will be able to bring targets out into a counting area in times t , $1.0 \text{ sec} \leq t < 2.5 \text{ sec}$. The cost of the system is ~\$500.00 plus machine shop time.

Fig. 29-1. (next page) Photographs of the rabbit and a trial rabbit built to withstand considerable shock (both in top photograph) and of the rabbit hutch, at -45° out of the first beam-analysis magnet of the transport system.

30. Slow Pulsing of the MSU Cyclotron Beam

J. Black (in conjunction with P. Sigg of the Cyclotron Laboratory)

The MSU Sector-Focused Cyclotron has the capacity to operate in both continuous (i.e., continuous in a macroscopic sense, ignoring the natural RF bunching) and pulsed modes. The pulsed mode may be obtained in two ways: First, the beam may be disrupted by deflection plates during the first turn, such that beam is achieved only when power is applied to these plates. Second, the RF itself may be pulsed. Both of these methods have been used successfully with the MSU Cyclotron.

The present pulsing system, employing the deflection plate method, is used primarily for fast pulsing, i.e., making use of RF bunching, yielding pulses in the nsec region. Earlier in the development of the machine, a slow pulsing system existed which pulsed the RF to produce beam pulses in the milli-second to 10 second range. However, the advent of rather sophisticated logic and protection circuits in the RF system itself now prohibits the use of this simple RF pulsing. To circumvent this difficulty, an amplifier has been designed which will enable the slow pulsing unit to power the first turn deflection plates. This method is quite simple and should prove much easier to use than the RF pulsing technique. This system will provide beam-on times in the range of 0.01 to 24 seconds, and off times of 0.01 to 100 seconds.

Another approach to the slow pulsing problem utilizes the same amplifier and deflection plates as above, but makes use of a multivibrator clock module designed by Dr. Peter Miller of the MSU Cyclotron Laboratory. This module can provide beam-on (or beam-off) times ranging from 0.4 to 4 seconds. An inhibit feature may be used to delay the counting for a period after the beam is turned off. In addition, routing signals are provided which may be used to obtain up to seven spectra - each of which contains only the counts for a specific time interval after the counting begins (figure 1). For example, one might want a spectrum for each of seven 0.4- second intervals following a bombardment. This would be very useful in determining half-lives and contaminants for various short-lived species. Provision is also made for a spectrum

to be taken during the beam-on period if desired.

A detailed, on-line gamma-ray spectroscopy study of Sc^{40} and other short-lived isotopes is now being initiated, using the above slow pulsing techniques (cf. section 20 of this Annual Report).

LOGIC PULSES FOR SLOW PULSING AVAILABLE FROM CLOCK MODULE

RANGES

MASTER CLOCK (1/2 CYCLE) 0.4 → 4.0 SEC.

ROUTING PULSES .04 → 2.0 SECOND

STEP FUNCTION FORMED FROM ROUTING PULSES

Fig. 30-1.

31. Megachannel γ -Ray Spectra

R. Doebler

We have previously reported on a two-dimensional γ -ray coincidence system in which we used a 3"x3" NaI(Tl) detector with a 7-cm³ Ge(Li) detector [1]. In the past year many changes have been made in our system, the most significant being the replacement of the above detectors with two large volume Ge(Li) detectors.

Figure 1 shows in flowsheet form the apparatus used in a typical two-dimensional coincidence experiment. When two photons are detected in coincidence, the linear gates are opened, allowing the output from detector 1 to go to the x -ADC while the output of detector 2 goes to the y -ADC. With the x - and y -ADC's operating in the synchronous mode, the signals going into the two ADC's must arrive within 1 μ sec of each other in order to be recorded as a coincidence event. The 8192-channel ADC's are interfaced to the Cyclotron Laboratory's Sigma 7 computer, and the program TASK EVENT reads the data from both ADC's through its multipurpose interface. The events are then stored in a dedicated buffer in the computer until the buffer fills, whereupon the contents of the buffer are written on magnetic tape. The two coincident γ -rays are recorded by channel number in one word of memory in the computer and on the magnetic tape. Since the events are written to tape, this system cannot be used for count rates greater than ≈ 2000 coincidence events per second.

The data on the magnetic tape are later recovered off-line using another program. This program reads each event off the tape and separates the x and y addresses. It will then use up to 10 digital gates per pass of the tape to define coincidence spectra. Each gate can be either an "integral coincidence", that is, an open gate, or a selected gate of a few channels' width containing a peak of interest. The gates can be set on either the x or y address and background subtraction can be performed. The resulting one-dimensional coincidence spectra are then punched on cards for later plotting and data analysis.

Figure 2 shows some results from a two-dimensional γ - γ coincidence

experiment of 2048x2048 channels on Pb^{200} decay [2]. The x -side detector was a 2.5% efficient $\text{Ce}(\text{Li})$ detector with a resolution of 2.3 keV on the 1332-keV peak of Co^{60} , and the y -side had a 2.0% efficient detector with a resolution of 3.6 keV. The detectors were mounted at 90° to each other with a graded lead absorber bisecting the angle as in figure 1 to reduce Compton scattering between the detectors [3]. Figure 2a shows the y -side integral coincidence spectrum, while figure 2b shows a part of the x -side integral coincidence spectrum obtained in an 18-hour period. Figures 2c and 2d show a part of the x -side spectra obtained from gating on the 161-keV and 235-keV regions respectively of the y -side. The gated regions are shown in figure 2a by the crosshatching. The background subtraction subroutine in the recovery program was used to compensate for coincidence events involving the underlying Compton background.

This example was chosen because it illustrates a very useful technique for resolving doublets that differ only slightly in energy. In the decay of Pb^{200} two of the states populated lie at 289.24 and 289.92 keV, and both of these states decay at least partially to the ground state, giving rise to a doublet differing by only 0.68 keV. These two states, moreover are fed from above by different γ -rays; the 235-keV γ feeds the higher state and the 161-keV γ feeds the lower state. That this is indeed the case can be seen in figure 2c and 2d, where the centroids of the 289.7-keV doublet can be seen to shift, depending on which γ -ray is used for the gate.

Our two-dimensional γ -ray system has been used extensively during the past year and with the recent replacement of the 2.0% detector with a 3.63% efficient detector having a resolution of 1.9-keV it will be even more valuable for the very complicated decay schemes presently being investigated.

References

- [1] R. Doebler, 1968 Annual Report, Nuclear Chemistry Group Michigan State University (COO-1779-4).
- [2] See section 1 of this annual report.
- [3] See section 3 of this annual report.

Fig. 31-1. Block diagram of the electronics used for collecting megachannel γ -ray spectra with the Sigma-7 computer.

Fig. 31-2. Portions of a megachannel spectrum for Pb^{200} . A. γ -side (poorer resolution) integral coincidences. B. x -side integral coincidence showing the region near the 289-keV doublet on an expanded scale. C. x -side in coincidence with the 161-keV region. D. x -side in coincidence with the 235-keV region.

32. Spectrum Analysis Routines

J. Cross and G. Giesler

Two of the spectrum analysis routines used in the MSU cyclotron laboratory are SAMPO (developed by J. Routti, University of California, Berkeley) and MOIRAE (developed by R. Au and G. Berzins at Michigan State University).

SAMPO is a computerized data analysis routine written by J. Routti and adapted to the MSU cyclotron's Sigma 7 computer by T. Arnette and C. Merritt. Control may be maintained either via a scope and light pen and/or Fortran control cards. Some of the varied uses of SAMPO are:

- 1) to produce Calcomp or printer plots of the spectrum
- 2) to calculate energies
- 3) to calculate areas and relative intensities
- 4) and to calculate half-life data.

At present this routine is utilized as a separate program, that is without scope control. SAMPO's mathematical evaluation involves an initial shape parameter calibration over the region to be analyzed. Strong, well-resolved singlet peaks should be used in this calibration. The program fits a Gaussian curve to the upper portion of the peak and leading and trailing exponentials to the base of the peak. The shape parameters are stored and a linear interpolation used to get the parameters for any other peak under consideration. Once the shape parameters have been calculated for any given spectrum they may be reread directly into the program via control cards for subsequent runs, thus considerably shortening future calculation times. Once the shape calibration has been established, the energy calibration may be calculated. The energy values may be fit to linear or higher order polynomials at the operator's discretion. Similarly efficiency calibrations may be made by specifying a number of well-spaced peaks and their corrected relative intensities.

The actual analysis portion of SAMPO may be performed under two basic modes:

- 1) Automatic - in which SAMPO subroutines search out all statistically significant peaks, evaluate suitable fitting intervals and fit the

peak using the shape, energy, and efficiency calibration data.

- 2) controlled - in which the operator directly describes a fitting interval, the number and location of peaks to be found in that interval, the fitting then being made under these described conditions.

Figure 1 shows a sample output of a fitted peak, with the following values being returned:

- 1) exact centroid (and error)
- 2) energy (and error)
- 3) area-counts (and error in percent)
- 4) intensity-counts (and error in percent)

Because of other excellent data analysis routines in our laboratory SAMPO has only recently been put to much advantage. The foremost advantage found to date has been in reducing time required to strip multiplets in complex spectra. SAMPO allows us to strip all the possible multiplets in a single spectrum in the time previously required to strip a single peak by hand. An example of the usefulness of the program is seen in figure 1, depicting the 212-222 keV quartet seen in the Bi^{204} spectrum earlier in this report (section 5). The quartet was stripped by indicating the end points used for background and the approximate center channel of each peak in the multiplet. The dots are actual data points, the + 's are SAMPO fitted points, and * 's are points where the two are identical. Note that the fit is quite excellent. For a set of 4 separate Bi^{204} runs the energy range was about 0.35 keV on each peak and the intensities range within 11%.

MOIRAE is a program that uses a live-display scope and sense switches to perform spectrum analysis. After data are either read in from cards or transferred from a data acquisition program, the switches control the display, log or linear, expansion and shifting of the axes, and the various computational routines. Figure 2 shows the display of a typical spectrum. The information at the top gives the counts in the channel and channel position of the long pointer, the number of cycles in the log display, the run number, the routine presently in use, and the order of the background fit. The Compton edge in the center of the display is shown expanded in figures 3 and 4. In figure 3, a

4th order fit to the background determined by a set of points is shown. Switches are used to move the tall pointer to the position of points in the spectrum used to calculate the background. The short lines indicate the position of points that have been accepted. Either a set of individual points or regions of points can be used to calculate the background. Figure 4 shows the result of subtracting the background, along with the original spectrum. From this one selects the peak limits, and whether to use the full raw peak or the portion of the peak with counts greater than one-third the maximum to calculate the centroid. The program then finds the centroid, area, sum of raw data and background, and square root of that sum. This information can be punched on cards, and, along with the background and raw data, plotted on a Calcomp plotter, and printed. After accumulating all the analyses one chooses, he can then have the computer calculate a calibration curve using a least-squares fit to a quadratic equation and calculate intensities from the area. Also all statistical quantities such as standard deviation of peak energies from the curve are calculated.

Naturally, each of these routines has distinct advantages and disadvantages. MOIRAE is particularly useful since it allows immediate operator control over such parameters as background order, background and points and interval, peak endpoints, etc. SAMPO, on the otherhand, has the distinct advantage of being able to fit multiplet peaks accurately, which MOIRAE is unable to do presently. On the other hand, SAMPO is handicapped by a low order (second) background fit which makes it difficult to fit peaks accurately in some regions of the spectra. MOIRAE has the added disadvantage of tying up the display scope of the Sigma 7 for hours at a time. To alleviate this difficulty somewhat, we are presently adapting the analysis programs for the PDP-9 computer so that they can perform the same functions as MOIRAE.

Figs. 32-2,3,4 (overside). Photographs of the oscilloscope displays from the spectrum analysis program MOIRAE, as run on the Sigma-7 computer.

Fig. 32-2. Display of a portion of the Bi^{204} γ -ray spectrum including the quartet of fig. 32-1. A fifth-order fit has been made to part of this spectrum and can be seen as the intense line following the Compton edge in the middle, under the quartet in question.

Fig. 32-3. Expanded view of a fourth-order fit having just been made. The long pointer is moved by means of switches to select points in the background for calculating the fit. The short pointers record those points selected.

Fig. 32-4. Display of the peaks after the calculated fourth-order background has been subtracted.

33. A New Method for Plotting Spectra

R. Eppley

Recently we have adopted the method of presenting γ -ray spectra as the square root of the counts/channel vs the channel number. An example of this is shown in figure 1-A, the spectrum of Sm^{147} presented as a square-root plot. Figure 1-B is the same spectrum plotted as a semi-log plot and figure 1-C is the same spectrum plotted linearly.

As can be seen from these plots, the square-root plot retains the good features of both of the other plots, as well as having additional good features of its own. The square-root plot is similar to the semi-log plot in that the peaks with low intensity are still emphasized much more than on a linear plot. At the same time the more intense peaks are still very prominent as on a linear plot. In addition the square-root plot automatically and directly gives the statistical error in the count of each channel. Thus, the plot, as well as being a representation of the spectrum, is also a plot of the error of each channel.

Fig. 33-1. Comparison of square-root plot with linear and semi-log plots.

34. Some Interesting Compton-Scattering Problems Encountered
in Ge(Li)-Ge(Li) Coincidence Experiments

G. Giesler

With the recent availability of large volume, high-resolution Ge(Li) detectors, we have been able to perform high-resolution 2-dimensional coincidence experiments more or less routinely, as described in section 31. With all their advantages, such experiments also present several problems. One is the detection of Compton scattering. In one of these Ge(Li) detectors about 90% of the γ -rays incident are Compton scattered, mostly in the backward direction. Since many samples studied have a very short half-life, the angle between detectors is often close to 180° to increase the effective geometry. As a result, the back scattered Compton γ -rays are captured in the second detector within the resolving time of the system. This produces a coincidence count even though the sample may produce only one γ -ray. In order to study this effect more carefully, a series of experiments was performed using a Cs^{137} sample as the source in a 2-dimensional coincidence setup.

Figure 1 shows the effect of angle between detectors and the effect of using an absorber. At 180° and no absorber, the Compton scattering detected is much more sharply focused in the Compton edge and backscatter peaks than at 90° , as well as having about 4.4 times as many counts overall. By putting a $\frac{1}{4}$ " graded lead absorber with a $\frac{1}{4}$ " diameter hole for the sample between the detectors, the Compton edge and backscatter peaks are eliminated from the 90° position, and greatly reduced for the 180° position.

In order to study this more fully, several gates were set on the counts from one detector to determine what is in coincidence with each region of the spectrum. Figure 2 shows the two integral-coincidence spectra along with the selected gates. The dip in the Compton background, denoted C in figure 2F, is seen also at other angles. Comparing the spectrum from that gate with those of 2B and 2D, the gates on either side, we see that the Compton edge has decreased in intensity and broadened significantly. Also in going from 2B to 2C to 2D, a peak appears between the Compton edge and the full energy peak, moving to lower energies as the gates go to higher energies.

The sum of the energy of this peak and that of the gate is that of the full energy peak. This new peak is caused by a γ -ray emitted from the source Compton scattering in the first detector; the Compton γ -ray released then Compton scatters in the second detector and the new Compton γ -ray completely captured in first detector, all well within the system resolving time.

As expected, gating on the backscatter peak produces a strong Compton-edge peak, figure 2B, and gating on the Compton edge produces a strong backscatter peak along with a higher Compton background below the backscatter peak, figure 2H. The gate on the background between the backscatter and Compton edge peaks, figure 26, produces a typical Cs^{137} spectrum with slightly emphasized backscatter, difference, and Compton edge peaks. The valley between the Compton edge and full energy peaks also produces a typical Cs^{137} spectrum with only a slight accent on the backscatter peak figure 2I. Since the full energy peak could only come from chance coincidence, it produces the normal Cs^{137} spectrum as seen in figure 2J.

Similar results have been obtained at different relative detector angles, and with the use of an absorber, though the peaks were not so pronounced. Therefore, when new peaks appear in a 2-dimensional $\text{Ge(Li)}\text{-Ge(Li)}$ coincidence experiment, one must be careful, since they are not necessarily from transitions in the isotope, but from scattering between detectors.

Cs^{137} COINCIDENCE

Fig. 34-1. Spectra demonstrating the effects of angles between the detectors and of absorbers on the Compton-scattered events. Left: no absorber; Right: graded lead absorber with small hole for the sample.

Fig. 34-2. Spectra demonstrating the effects of moving the gates to various position on the scattered spectra. At the top left and right are the x and y integral spectra, respectively. The other spectra are from the x side in coincidence with the designated y gate, as shown in the y integral coincidence.

35. Calibration of γ -Ray Standards

R. Doebler

With the development of high resolution Ge(Li) detectors it has become possible to measure γ -ray energies with a precision of better than 100 eV. This does not present any significant problem for low energy (100 to 1000 keV) γ -ray spectroscopy, where a great number of energy standards are available which are known precisely to tens of eV's. However, for γ -ray energies greater than 1 MeV there are significantly fewer γ -ray standards with this precision. We have, therefore, recently undertaken the task of remeasuring the energies of some common standards listed in the 1967 edition of the *Table of Isotopes* [1] in the region 1000 to 2000 keV.

The calibration standards used in these determinations are listed in Table 1. The values are the average of those compiled by Marion [2] and I. Gunnink, et al. [3], except for Co^{56} , where the energies of Gunnink, et al., were used exclusively.

The Ge(Li) detectors used had resolutions of 2.1- and 2.3-keV for the 1332-keV peak of Co^{60} and efficiencies of 3.6% and 2.5%, respectively. The spectra were taken with both a Northern Scientific 4096-channel ADC interfaced to a PDP-9 computer and a Northern Scientific 8192-channel ADC interfaced to an SDS Sigma-7 computer.

Energy determinations were made by counting the "unknown" simultaneously with several of the standards listed in Table 1. The centroid channels of the peaks in a spectrum were determined following subtraction of a 3rd order least-squares fitted background. From the centroid channels of the standards a least-squares quadratic calibration curve was calculated and the energies of the "unknown" γ -rays computed from this curve.

We have, so far, completed only the energy determination of selected γ -rays in Ag^{110m} and Ta^{182} . The Ag^{110m} and Ta^{182} sources were made at the MSU Triga reactor, as were many of the calibration standards listed in Table 1.

The results of our work are summarized in Table 2 along with the results obtained in previous studies.

Table 1. Calibration Standards

Isotope	Energies (keV)		
	Marion [2]	Gunnink [3]	Values Used
Mn ⁵⁴	834.81±0.03	834.84±0.05	834.83±0.04
Y ⁸⁸	898.04±0.04	898.02	898.03±0.04
Co ⁶⁰	1173.23±0.04	1173.23±0.04	1173.23±0.04
Na ²²	1274.55±0.04	1274.55	1274.55±0.04
Co ⁶⁰	1332.49±0.04	1332.51±0.02	1332.50±0.03
Na ²⁴	1368.53±0.04	-	1368.53±0.04
K ⁴⁰	1460.75±0.06	-	1460.75±0.06
Na ²⁴	1731.91±0.12	1732.13±0.06	1732.08±0.06
Y ⁸⁸	1836.13±0.04	1836.13±0.05	1836.13±0.05
Na ²⁴	2753.92±0.12	2754.14±0.06	2754.09±0.06
Co ⁵⁶	846.76±0.05	846.78±0.06	846.78±0.06
	1037.97±0.07	1037.85±0.07	1037.85±0.07
	1175.26±0.13	1175.08±0.07	1175.08±0.07
	1238.34±0.09	1238.29±0.04	1238.29±0.04
	1360.35±0.09	1360.22±0.04	1360.22±0.04
	1771.57±0.10	1771.33±0.06	1771.33±0.06
	2015.49±0.20	2015.33±0.07	2015.33±0.07
	2035.03±0.12	2034.90±0.06	2034.90±0.06
	2598.80±0.12	2598.52±0.05	2598.52±0.05

Table 2. Energy Calibration Results for Ag^{110m} and Ta^{182}

Ag^{110m} :		
Energies in keV		
Brahmavar et al. [4]	Legrand et al. [5]	Present Work
884.67 $\pm$ 0.04	884.66 $\pm$ 0.04	884.68 $\pm$ 0.04
937.48 $\pm$ 0.04	937.46 $\pm$ 0.06	937.48 $\pm$ 0.04
1384.22 $\pm$ 0.04	1384.35 $\pm$ 0.06	1384.26 $\pm$ 0.05
1475.73 $\pm$ 0.04	1475.81 $\pm$ 0.08	1475.76 $\pm$ 0.07
1504.90 $\pm$ 0.08	1505.11 $\pm$ 0.14	1505.01 $\pm$ 0.07
1562.22 $\pm$ 0.06	1562.37 $\pm$ 0.12	1562.35 $\pm$ 0.08
Ta^{182} :		
	Table of Isotopes [1]	Present Work
	1121.28 $\pm$ 0.12	1121.31 $\pm$ 0.04
	1189.03 $\pm$ 0.20	1189.06 $\pm$ 0.04
	1221.42 $\pm$ 0.10	1221.42 $\pm$ 0.04

References

- [1] C. M. Leder, J. M. Hollander and I. Perlman, *Table of Isotopes*, John Wiley and Sons, New York 1967, 6th. Ed.
- [2] J. B. Marion, *Nucl. Data*, A4, 301 (1968).
- [3] R. Gunnink, R. A. Meyer, J. B. Niday, and R. P. Anderson, *Nucl. Instr. Methods* 65, 26 (1968).
- [4] S. M. Brahmavar, J. H. Hamilton, A. V. Ramayys, E. P. F. Zganjar, C. E. Bemis, Jr., *Nucl. Phys.* A125, 217 (1969).
- [5] J. Legrand and J. P. Boulanger, *Compt. Rend.* 265, 697 (1967).

36. Progress with the Compton-Suppression

NaI(Tl) Annulus

K. Kosanke

At a time when large ($\geq 2.5\%$) Ge(Li) γ -ray detectors are allowing megachannel analysis [1] to begin to become commonplace, a $8'' \times 8''$ NaI(Tl) split annulus [2] has remained a primary tool in our decay studies. In fact, these same larger Ge(Li) detectors used in conjunction with the annulus have expanded its usefulness. This report discusses three of the experiments which have become more or less routine in our decay studies: Triple coincidence experiments, anti-Compton experiments, and anticoincidence experiments.

A triple γ -coincidence [511 keV-511 keV (at 180°)-any] run was made on Bi^{205} in an attempt to refine previously reported [3] β^+ feeding data. The experimental set up and β^+ results are shown below; also see fig. 1 for spectra (for further discussions see section 4 of this report).

β^+ feeding to Pb^{205} :

703.4 keV level $\rightarrow$ 0.072 $\pm$ 0.02%

987.5 keV level $\rightarrow$ 0.008 $\pm$ 0.004%

1043. keV level $\rightarrow$ $< 0.002\%$

^a Photomultipliers

^b Halves of annulus

^c Source and ≈ 20 mils of Pb

foil shielding

^d 2.5% Ge(Li) detector

An anti-Compton [≥ 90 keV- ≥ 90 keV]-anti any] run was made on Bi^{205} in an attempt to reduce uncertainties in intensity for some of the particularly weak transitions appearing in singles by suppressing the Compton distributions. The experimental set up is shown below and spectra are in

fig. 2. Although the data have not been completely analyzed, it is apparent that intensity uncertainties will be reduced. Also, several additional weak peaks were seen and the possibility that some are transitions in Pb^{205} is being investigated.

An anticoincidence $\{(\geq 90 \text{ keV} - \geq 90 \text{ keV} - \geq 90 \text{ keV}) - \text{anti any}\}$ run was made on Bi^{205} using an additional 3"x3" NaI(Tl) detector in the tunnel of the annulus opposite the Ge(Li) detector in an attempt to verify some of the new assignments made in our partially completed decay scheme for Bi^{205} . The experimental set up is shown below and spectra in fig. 2.

- a 2.5% Ge(Li) detector
- b Photomultipliers
- c Halves of annulus
- d Graded Pb collimator
- e 3"x3" NaI(Tl) detector
- f Bi^{205} source

Again, these data have not been completely analyzed. However, the data do support the placement of the 890.0-keV γ in the new decay scheme ($1593.5 \rightarrow 703.4 \text{ keV}$ rather than $890.0 (7/2) \rightarrow 0.0 \text{ keV}$ [4] and tends to confirm the existence of a level at 1776.0 keV. Further this run suggest that the 828.2-keV transition may be feeding the isomeric state at 1013.8 keV.

References

- [1] For further discussions see Megachannel Analysis, section 31 of this report
- [2] R. L. Auble, D. B. Beery, G. Berzins, L. M. Beyer, R. C. Etherton, W. H. Kelly, and Wm. C. McHarris Nucl. Instr. and Meth. 51, 61 (1967).
- [3] D. F. Peppard, G. W. Mason, and S. W. Moline, J. Inorg. Nucl. Chem. 5, 141 (1957).
- [4] S. H. Vegors, Jr., R. L. Heath and D. G. Proctor, Nucl. Phys. 48, 230 (1963).

Fig. 36-1. Bi²⁰⁵ pair spectrum, demonstrating the use of our 8x8" NaI(Tl) annulus for measuring weak β^+ feedings.

Fig. 36-2. Spectra demonstrating the difference between anti-Compton (source external) and anticoincidence (source internal) configurations.

37. Measurement of Low-Energy Conversion Electrons with the MSU $\pi\sqrt{2}$ Iron-Free Spectrometer

B. Johnston† and K. Kosanke

Complementary to the study of the β -decay of Bi^{205} described earlier in this report [54], an attempt is under way to measure directly the energy and relative intensity of the 2.3-keV transition in Pb^{205} . This is an $E2$ transition from the first excited state to the ground state ($1/2^- \rightarrow 5/2^-$). Although well documented from observations of a collection of doublets separated by ≈ 2.3 keV [1] in the γ -ray and conversion electron spectra and from observed trends of the $1/2^-$ and $5/2^-$ states [2] in the neutron deficient Pb isotopes, the transition has never been observed directly. Our best estimate, as determined by stripping the moderately intense 260.6- 262.9-keV doublet (using a Ge(Li) detector with 2.3 keV resolution FWHM for the 1332.5-keV transition in Co^{60}) is for a 2.33 ± 0.02 keV transition and an intensity of 11.6% relative to the 703-keV transition (i.e., $\approx 2\%$ of all transitions) based on transitions intensities of the observed doublet.

The MSU $\pi\sqrt{2}$ β -ray spectrometer [3] is being used to measure the low-energy electron spectrum. In brief, it is a 30-cm iron-free, double-focusing, β -ray spectrometer with ultimate resolving power of about 0.045%. A post-focusing acceleration technique has been developed which extends the low energy range of the spectrometer down to less than 1 keV [4] (see Fig. 1). The acceleration is accomplished by means of a 5 to 7 kV electric potential at the exit slit of the spectrometer. This raises the energy of the electrons to a point where they can be detected with a conventional thin-window G. M. tube at an efficiency of nearly 100%. A negative 300-V retardation bias is maintained just before the high voltage cell to repel very low energy free electrons that would cause spurious counts in the detector.

The idea of this experiment is to observe the 2.33-keV transition directly by measuring the resulting internal conversion spectrum. However, this spectrum will be superimposed on the M -Auger spectrum. Thus, consideration must be given to the fact that the M -Auger lines could obscure the conversion spectrum.

Rough estimates indicate that there will be $\sim 1\frac{1}{2}$ M -shell vacancies per decay and that the majority of these vacancies will be concentrated in the M_4 and M_5 subshells. Since the M -Auger intensity is distributed over many transitions, any individual transition will occur for only a few percent of the total decays, with the M_4 and M_5 transitions being the most significant. It appears that conversion intensities are not going to be too much smaller than most of the individual M -Auger intensities. Thus, the conversion lines should be visible if they do not lie too near an Auger line.

To check this the energies for all the Auger lines between 1.415 and 2.250 keV have been calculated (see fig. 2) and a summary of possible interfering lines is given in Table 1. For the conversion spectrum the uncertainty in the transition energy enters (± 20 eV) and only the relative positions of the conversion lines can be predicted with any accuracy. However, limits have been set as to the absolute positions of the conversion lines in the Auger spectrum (See Table 1 and fig. 2).

The most prominent conversion lines are expected to be the N_2 and N_3 lines, with the rest being somewhat weaker. There are seven Auger lines in the region where the N_2 conversion line is expected. Even though all of these originate from an initial M_4 or M_5 vacancy, none is expected to be intense, since the two electrons involved in the transition are in different shells, or from the same shell but with l values differing by 2 or more [5]. Further, if the transition energy is slightly higher than 2.33 keV, the conversion line will fall in a fairly open spot of the Auger spectrum. Thus, there is a good chance that the N_2 conversion line can be observed and identified.

The situation is not quite so favorable for the N_3 line, as 13 Auger lines fall in the region where it is expected. In addition, the $M_4-N_4N_4$ line should be one of the more intense M -Auger lines. Nevertheless, if another conversion line is identified there is a good chance that the N_3 line can also be identified.

Although the N_1 conversion should be weak, its position falls in a rather open spot of the Auger spectrum. There are three lines in this region and only one, the $M_1-N_1O_1$ is expected to be of much consequence. There

is a fair chance that this conversion line can be observed if it has any intensity at all.

A pessimistic view is held for the chances of observing any of the other lines without first positively identifying other conversion lines. The N_4 and N_5 lines would be very weak and located in the vicinity of 6 Auger lines, one of which ($N_5-N_4N_6$) may be quite intense. Likewise, the O conversion will be weak and located near several Auger lines.

We have now begun this experiment but have as yet only incomplete spectra.

Footnote and References

- [1] The doublets observed are 260.6-262.9 keV, 574.0-576.4 keV, and 759.1-761.4 keV.
- [2] E. K. Hyde, I. Perlman, and G. T. Seaborg, *Nuclear Properties of the Heavy Elements*, Vol. II. (Prentice-Hall, 1964).
- [3] L. J. Velinsky, Ph.D. Thesis, MSU, (1964).
- [4] R. J. Krisciokaitis, Ph.D. Thesis, MSU, (1965).
- [5] Slatis and Rockbeager, *Arkiv Fysik*, preprint (1969) and S. K. Haynes, Private Communication.

Table 1. Auger and Coster-Kronig Lines within the Limits
for the N Conversion Lines

Auger and Coster-Kronig Lines		N Conversion Lines	
Identification	Energy (keV)	Energy (keV)	Identification
		1.416)	
$M_5-N_1O_1$	1.436	1.436	N_1
$M_5-N_1N_6$	1.437		
$M_5-N_1N_7$	1.442		
		1.456)	
		1.546)	
$M_5-N_1O_4$	1.565		
$M_5-N_2O_1$	1.566	1.566	N_2
$M_5-N_1O_5$	1.567		
$M_5-N_2N_6$	1.567		
$M_5-N_2N_7$	1.572		
$M_4-N_1O_2$	1.581		
$M_5-N_1P_1$	1.584		
		1.586)	
		1.665)	
$M_4-N_1O_4$	1.667		
$M_4-N_2O_1$	1.668		
$M_4-N_1O_5$	1.669		
$M_4-N_2N_6$	1.669		
$M_4-N_2N_7$	1.674		
$M_5-N_3O_1$	1.685	1.685	N_3
$M_4-N_1P_1$	1.686		
$M_5-N_3N_6$	1.686		
$M_4-N_1P_2$	1.690		
$M_5-N_3N_7$	1.691		
$M_5-N_2O_4$	1.695		
$M_5-N_2O_5$	1.697		
$M_4-N_4N_4$	1.701		
		1.705)	

Table 1. continued

		1.875		
$M_2-N_1N_2$	1.875			
$M_3-N_2N_5$	1.876		1.895	N_4
$M_5-N_4O_1$	1.895			
$M_5-N_1N_6$	1.896			
		1.897		
$M_5-N_4N_7$	1.901			
$L_1-L_3N_1$	1.909			
		1.915		
			1.917	N_3
		1.937		

Note: The energy limits placed on the conversion lines reflect the uncertainty in the energy of the 2.33 ± 0.02 keV transitions.

POST FOCUSING ACCELERATION CELL AND VOLTAGE CONNECTIONS

Fig. 37-1. Postfocusing acceleration cell and voltage connections for measuring low-energy electrons with the MSU $\pi/2$ spectrometer.

38. A Szilard-Chalmers Separation for
Isomer-Ratio Enhancements

R. Goles

A number of studies of the decay of Nd^{139} have contributed to the now relatively complete decay schemes of Nd^{139m} and Nd^{139g} [1-5]. Primary among these was the high-resolution γ -ray study by Beery, et al. [5], whose decay scheme for both the ground and metastable states appear in fig. 19-1 of this Annual Report.

In determining the ground-state decay scheme, Beery, et al., produced Nd^{139g} sources by bombarding Pr_2O_3 targets with 29-MeV protons for 245 sec in order to optimize the ground state over the metastable state. Experiments were carried out immediately upon concluding each bombardment, and the γ -rays resulting specifically from Nd^{139g} decay were followed as their intensities dropped from their initial values to those when transient equilibrium with Nd^{139m} was achieved. However, most of the γ -ray intensity, even this soon after the bombardments, originates from the 5.5-hr Nd^{139m} decay, for some 88% of all β -decay from 30-min Nd^{139g} proceeds directly to the ground state of Pr^{139} . A typical spectrum of Nd^{139g} obtained by Beery, et al., appears in fig. 1. Since the possibility exists that many low-intensity transitions following the 30-min Nd^{139g} decay could be masked by the 5.5-hr Nd^{139m} background activity, the aim of the present work is to attempt to rid the Nd^{139g} activity of its metastable background through the use of a Szilard-Chalmers reaction.

The procedure used has been described previously [6,7] and, in brief, consists of producing quinoline solutions of phthalocyanine complexes of Nd^{139} after transient equilibrium has been achieved. It was hoped that the highly converted nature of the $M4$ isomeric transition between the metastable and ground states would be so disruptive, in terms of electronic disturbances produced, as to break the organometallic bonds holding the Nd^{139} atom in the complex and thereby allow the ground state to be separated from the metastable species through aqueous extraction. Of course, the decay of both the ground and metastable states of Nd^{139} would also be expected to produce free Pr^{139} ,

but the spectrum and the decay scheme of this daughter has previously been well-characterized by Beery, et al. [8], and would not be expected to interfere in the energy region that is of primary interest in Nd^{139g} decay.

The results to date, although decidedly positive, have not succeeded in enhancing the ratio of ground to metastable activity over that obtained through the conventional counting techniques described earlier. The presence of the metastable species in the aqueous extract has been found to result, in part, from radiation damage suffered by the organometallic complexes as a result of too high a specific activity in the organic phase. The combined effects of smaller amounts of activity, larger volumes of organic solvent, and the addition of carrier have increased the total ground state yield, but it still remains significantly below the enhancement level desired.

Since the Szilard-Chalmers reactions are generally very specific as to the conditions under which they work well, a systematic search will be directed toward determining the destabilizing conditions responsible for limiting the desired level of separation in the above chemical system. Paralleling this approach, an investigation will be conducted for new systems involving inorganic metallic complexes as described in ref. [9].

References

- [1] K. Ya. Gromov, A. S. Danagulyan, A. T. Strigachev, and V. S. Shpinel', *Izv. Akad. Nauk SSSR, ser. fiz.* 27, 1357 (1963).
- [2] J. Gilat and W. J. Tretyl, LRL Report UCRL-17299, p. 20 (1967).
- [3] J. Lange, KFK Report, summarized in J. Lange, H. Münzel, and I. Leitzl, *Radiochim. Acta* 8, 123 (1967).
- [4] K. Ya. Gromov, A. S. Danagulyan, L. N. Nikityuk, V. V. Murav'eva, A. A. Sorokin, M. Z. Shtal', and V. S. Shpinel', *Zh. Eksperim. i Teor. Fiz.* 47, 1644 (1964).
- [5] D. B. Beery, W. H. Kelly, and Wm. C. McHarris, *Phys. Rev.*, scheduled for the 20 Nov 1969 issue.
- [6] W. Herr, *Z. Naturforsch.* 9a, 180 (1954).
- [7] T. Stenström and B. Jung, *Radiochim. Acta*, 4, 3 (1965).
- [8] D. B. Beery, W. H. Kelly, and Wm. C. McHarris, *Phys. Rev.*, scheduled for the 20 Nov 1969 issue.
- [9] H. A. C. McKay, in *Prog. Nucl. Phys.* (O. Frisch, ed.), Vol. I., Pergamon Press (1950).

Fig. 38-1. Example of the low-energy portion of a Nd^{139m} spectrum, demonstrating the background problems from the more intense Nd^{139m} γ -rays.

39. Preparation of Thin Pr^{141} Target Foils

R. Goles

In conjunction with experiments concerning the (p,t) reaction on Pr^{141} , thin ($0.1\text{--}1.0\text{ mg/cm}^2$) Pr^{141} foils have been prepared by vacuum evaporation. The vacuum evaporation apparatus used consisted of an 8 cfm mechanical pump, a 200 l/sec diffusion pump with a liquid nitrogen cooled baffle, and a evaporation chamber consisting of a 18 cm $\phi \times 32$ cm high glass bell jar. High voltage as well as high current base plate feed throughs allows the conventional resistance method of heating to be converted easily to that of an electron bombardment system, which is especially important where high temperature work is required or where precious isotopic material is involved. Realistic pressures obtainable by the above system are of the order of $\sim 10^{-7}$ torr.

The foils themselves were prepared by evaporating 100% abundant Pr^{141} metal from a 5 mil tantalum boat onto a $20\text{ }\mu\text{g/cm}^2$ carbon backing contained on a glass slide. Having a much higher Q value for (p,t) reactions than Pr^{141} , the carbon backings were used only on a convenient releasing agent for the deposited Pr^{141} foils. In order to insure a good uniformity of the foils, the carbon backings were always placed 4-6 in. above the boat.

Since the oxidation of Pr occurs readily in air, the bell jar was always let up to atmospheric pressure with the inert gas argon. The foils were then floated free of the glass slide, and mounted with all possible haste and stored under vacuum.

Under vacuums produced by normal roughing conditions, the useful lifetime of targets of $\sim 300\text{ }\mu\text{g/cm}^2$ thickness were found to be of the order of two weeks. However, Pr^{141} targets of indefinite lifetimes have been produced by sandwiching the foil between thin films ($\sim 5\text{ }\mu\text{g/cm}$) of the plastic, formvar. The thin films of formvar required are produced when a 0.1% solution of formvar in ethylene chloride is added dropwise to a container of water. The plastic film forms on the water surface and is usually carefully removed with a thin wire ring. The method is somewhat tedious but very useful.

Measurements of target thicknesses are conveniently obtained through the use of an α -gauge. The α -gauge consist of a well-collimated mono-energetic α beam, a Si(Li) semiconductor detector, and a position adjustable target rack, which is indispensable in determining target uniformities. With this α -gauge, target thicknesses can easily be determined within 10%.

The target preparation program will be extended in the near future to include the neighboring elements. Tb, Ho, and Tm.

40. A Single-Crystal Ge(Li) Conversion-Coefficient Spectrometer*

R. Doebler

(with R. R. Todd and C. R. Gruhn's group of the Cyclotron Lab.)

Lithium-drifted germanium detectors have been used mainly as γ -ray detectors for a number of years. Recently, several experimenters have noted the presence of conversion electrons in low-energy x-ray spectra observed with thin-windowed Ge(Li) detectors [1]. The possibility of using a small Ge(Li) detector in a thin windowed mount to measure γ -rays and electron intensities simultaneously and hence serve as a single-crystal conversion coefficient spectrometer was proposed by C. R. Gruhn and Wm. C. McHarris.

There are several advantages to such a system, principally that it uses fewer electronics and only one analyzer. Also, a single efficiency curve can be made for the single crystal Ge(Li) spectrometer, one that contains information about both γ -ray and electron intensities. Finally, the single-crystal spectrometer is potentially a powerful tool for coincidence experiments involving another detector, because it would mean the difference between a double or triple coincidence or sequential double coincidence experiments. Disadvantages of such a device are source strength, i.e., the detectors are necessarily small, so strong sources are required, and the straggling experienced by the electrons in passing through the window, especially for lower energies. This straggling makes it very difficult to separate the *L* and *M* electron peaks because of their small energy difference. There also appears to be no single optimum size of detector to cover all energy regions.

The Ge(Li) detector used in these experiments was manufactured in this laboratory by C. Maggiore. It is a planar drift device with a drift depth of ≈ 7 mm and an active volume of ≈ 0.4 cm³. It was mounted in a conventional dipstick cryostat with a window of 0.25 mil Havar foil. The orientation of the detector and source is indicated in fig. 1. The detector has a resolution of 1.8-keV FWHM for the 661.6-keV γ -ray of Cs¹³⁷.

The relative efficiencies were found to depend sensitively on source shape and position. To insure uniform reproducibility, the sources were

vaporized through 3-mm diameter masks onto 0.25-mil Pt foil backings. These were mounted on Al rings which fit into the slot shown in Fig. 1. The efficiency of the detector was calibrated by using the standards listed in Table 1, whose conversion coefficients had been determined precisely. An efficiency curve was also obtained using theoretical conversion coefficients from the table of R. S. Hager and E. C. Seltzer [2]. The intensities of the electron peaks were determined by first taking a spectrum of both electrons and γ -rays, as shown for Cs^{137} in fig. 2a. Then an Al absorber was placed between the source and the detector to stop the electrons, and a spectrum taken which contained only the γ -rays, as shown in fig. 2b. By overlapping the spectra and subtracting, the conversion electron intensities were easily obtained.

The efficiency curve for the present detector is shown in fig. 3. It takes the form of the relative efficiencies for γ -rays and electrons, i.e.,

$$C = \frac{D(\gamma)}{D(e^-)},$$

vs the γ -ray energy, E_γ . To obtain the conversion coefficient of a transition, one merely obtains an experimental intensity ratio for the γ -rays and electrons and multiplies this by the corresponding ordinate from fig. 3, i.e.,

$$\alpha = C \frac{A_{e^-}}{A_\gamma}.$$

Because the efficiency for electrons varies very slowly, the above efficiency curve is very nearly the same as the efficiency curve for γ -rays alone. The latter curve is presented in Fig. 3 with an offset scale for comparison.

An example of the use of the detector is shown in fig. 4, where spectra for Pb^{204m} are shown. The difference between α_K for the 899.1-keV $E2$ and the 912.0-keV $E5$ transition is striking. Our measured values for these are given in Table 2, where they are compared with the theoretical values of ref. [2]. The agreement is quite good, as is the agreement with previously measured values [3]. We also include the K/L ratio for the $E5$ transition, but one cannot extract a reliable K/L ratio for the $E2$ because

of the complexity of this portion of the spectrum.

Our experiments show that a small Ge(Li) detector with a thin window to allow the simultaneous counting of γ -rays and conversion electrons shows considerable promise as a convenient piece of equipment for measuring conversion coefficients from spectra that are not too complex.

Table 1. Conversion Coefficient Calibration Standards.

Nuclide	γ -energy	Multi-polarity	α_K experimental	Ref.	α_K theoretical ^a
Sn ¹¹³	391.7	M4	$4.38 \pm 0.08 \times 10^{-1}$	4	4.48×10^{-1}
Cs ¹³⁴	795.8	E2	$2.46 \pm 0.30 \times 10^{-3}$	5	2.58×10^{-3}
Cs ¹³⁴	1365	E2	$6.8 \pm 0.5 \times 10^{-4}$	5	8.19×10^{-4}
Cs ¹³⁷	661.595	M4	$8.94 \pm 0.10 \times 10^{-2}$	6	9.28×10^{-2}
Bi ²⁰⁷	569.6	E2	$1.56 \pm 0.07 \times 10^{-2}$	7	1.59×10^{-2}
Bi ²⁰⁷	1064	M4	$9.0 \pm 0.9 \times 10^{-2}$	7	9.70×10^{-2}

^aref. 8.

Table 2. Pb^{204m} Conversion Data.

Transition Energy (keV)	Multi-polarity	α_K experimental	α_K theoretical ^a	K/L experimental	K/L theoretical ^a
899.1	E2	0.0072 ± 0.0022	0.0065	---	---
912.0	E5	0.0545 ± 0.0045	0.046	1.66 ± 0.25	1.73

^aref. 8.

References

- [1] R. H. Pehl, F. S. Goulding, D. A. Landis, M. Lenziinger, Nucl. Instr. Methods, 59, 45 (1968).
- [2] R. S. Hager and E. C. Seltzer, Nucl. Data A4, 1 (1968).
- [3] C. J. Herrlander, R. Stockendal, J. A. McDoneil, and I. Bergström, Nucl. Phys. 1, 643 (1956).
- [4] J. R. Hamilton, in C. M. Lederer, J. M. Hollander, and I. Perlman, *Table of Isotopes* (J. Wiley and Sons, New York, 1967) 6th Ed.
- [5] R. A. Brown and G. T. Ewan, Nucl. Phys. 68, 325 (1965); W. Van Wyngaarden and R. D. Conner, Can. J. Phys. 42, 42 (1964).
- [6] J. S. Merritt and J. C. V. Taylor, Anal. Chem. 37, 351 (1965).
- [7] K. Siegbahn, Nucl. Phys. A93, 63 (1967).

*This paper is a condensation of a journal article, C. R. Gruhn, R. R. Todd, C. J. Maggiore, W. H. Kelly, R. E. Doebler, and Wm. C. McHarris, Nucl. Instr. Methods 75, 109 (1969).

Fig. 40-1. Small Ge(Li) detector mounted in a dipstick cryostat having a 0.25-mil Havar foil window to permit the detection of γ -rays and conversion electrons simultaneously.

Fig. 40-2. Spectra demonstrating the use of absorbers to select γ -rays and electrons separately. Top: Cs^{137} spectrum taken directly through the Havar window. Bottom: Cs^{137} spectra taken through an Al absorber to remove electrons.

Fig. 40-3. Efficiency curves. The solid curve represents the composite γ -ray-electron efficiencies. Triangles are calculated from theoretical conversion coefficients, $\times$'s from experimental ones. The dashed curve shows a (displaced) γ -ray efficiency curve.

Fig. 40-4. Pb^{204m} spectra, demonstrating the use of the conversion-coefficient spectrometer.

PERSONNEL

Principal Investigator:

Wm. C. McHarris, Assoc. Prof. of Chemistry and Physics, Ph.D. (Nuclear Chemistry), University of California, Berkeley, 1965.

Research Associate:

Ray A. Warner, Ph. D. (Nuclear Physics), University of California, Davis, 1969.

Graduate Students:

Jerry N. Black, B. S. (Chemistry), W. Va. Wesleyan, 1965. [Ph. D. Thesis Topic, "Pulsed In-Beam Studies of the Decays of Sc^{40} and Other Very Short-Lived Species"].

James B. Cross, B. S. (Chemistry), University of Dayton, 1968. [Ph. D. Thesis Topic, "The Decays of Bi^{204} and Bi^{203} , with Shell-Model Calculations on the States in the Pb Daughters"].

Raymond E. Doebler, B. A. (Chemistry), Cornell University, 1965. [Ph. D. Thesis Topic, "Systematics and Decays of Very Light Pb Isotopes"].

Richard E. Eppley, B. A. (Chemistry), North Central College (Ill.), Jan. 1966. [Ph. D. Thesis Topic, "Decays of Neutron-Deficient Gd Isotopes and Systematics of Nuclei Near the $N=82$ Shell"].

Richard B. Firestone, B. S. (Chemistry), University of Michigan, 1967.

Gregg C. Giesler, B. S. (Chemistry), University of Illinois, 1966. [Ph. D. Thesis Topic, "Very Short-Lived and Neutron-Deficient Ga and Zn Isotopes -- Delayed Alpha Emission"].

Ronald W. Gales, B. S. (Chemistry), University of Illinois, 1966. [Ph.D. Thesis Topic, "The (p,t) Reaction on Rare-Earth Isotopes"].

Kenneth L. Kosanke, B. S. (Chemistry), Wisconsin State University, White-water, 1966. [Ph.D. Thesis Topic, "On-Line Mass Separation with an Electrostatic Quadrupole Mass Filter"].

Scientific Aide (unpaid):

Orilla S. McHarris, Ph.D. (Physics), University of California, Berkeley, 1968.

Publications Involving Members of the
Nuclear Chemistry Group

I. Papers in Print during the Calendar Year 1969

1. "The States of ^{117}Sn ", D. B. Beery, G. Berzins, W. B. Chaffee, W. H. Kelly, and Wm. C. McHarris, Nucl. Phys. A123, 649 (1969).
2. "Three-Quasiparticle Multiplet in Pr^{139} ", Wm. C. McHarris, D. B. Beery, and W. H. Kelly, Phys. Rev. Letters 22, 1191 (1969).
3. "Neutron Deficient Members of the $A=139$ Decay Chain: I. 5.5-Hour Nd^{139m} and 30-Minute Nd^{139g} ", D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev., scheduled for the 20 November 1969 issue.
4. "Neutron Deficient Members of the $A=139$ Decay Chain: II. 4.5-Hour Pr^{139} ", D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev., scheduled for the 20 November 1969 issue.
5. "A Single Crystal $\text{Ge}(\text{Li})$ Conversion-Coefficient Spectrometer", C. R. Gruhn, R. R. Todd, C. J. Maggiore, W. H. Kelly, R. E. Doebler, and Wm. C. McHarris, Nucl. Instr. Methods, 75, 109 (1969).
6. "Three-Quasiparticle Multiplets in Nuclei Below $N=82$ ", D. B. Beery, R. Muthukrishnan, W. H. Kelly, and Wm. C. McHarris, International Conference on the Properties of Nuclear States, Montreal, 25-30 August 1969, Contributions, p. 203 (1969), Les Presses de la Université de Montréal (1969).
7. "A $\text{NaI}(\text{Tl})$ Split Annulus for Coincidence, Anticoincidence, Triple Coincidence, and Pair Spectrometry", W. H. Kelly and Wm. C. McHarris, International Conference on Radioactivity in Nuclear Spectroscopy, Vanderbilt University, 11-15 August 1969, to be published in the proceedings, Vol. II (contributed papers).
8. "Data Acquisition from Simultaneous Experiments Using the MSU Sigma-7 Computer", Wm. C. McHarris, R. F. Au, D. L. Bayer, W. Benenson, R. A. DeForest, W. H. Kelly, and W. E. Merritt, International Conference on Radioactivity in Nuclear Spectroscopy, Vanderbilt University, 11-15 August 1969, to be published in the proceedings, Vol. I (invited papers).

II. Papers Submitted to Journals or to Be Submitted by the End of Calendar Year 1969

9. "The Electron Capture Decay of Gd^{149} ", R. E. Eppley, Wm. C. McHarris, and W. H. Kelly, Phys. Rev.
10. "Decay of ^{74}As ", J. R. Van Hise, Wm. C. McHarris, C. Papperillo, E. G. Funk, and J. W. Mihelich, Nucl. Phys.
11. "The Population of States in Ce^{140} by Pr^{140} Decay", D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev.
12. "Decays of Ba^{131m} , Ba^{133m} , and Ce^{137g+m} ", D. B. Beery, W. H. Kelly, and Wm. C. McHarris, Phys. Rev.
13. "Compton Scattering Problems in Ge(Li)-Ge(Li) Coincidence Experiments", G. C. Giesler, Wm. C. McHarris, and W. H. Kelly, Nucl. Instr. Methods.
14. "States in Tl^{200} Populated by the Electron-Capture Decay of Pb^{200} ", R. E. Doebler, Wm. C. McHarris, and W. H. Kelly, Phys. Rev.
15. "The $M4$ Isomer Gd^{145m} ", R. E. Eppley, Wm. C. McHarris, and W. H. Kelly, Phys. Rev.

III. Abstracts and Papers Presented at Meetings

1. " γ Ray Studies of the Decay of 30-min. Nd^{139g} ", D. B. Beery, Wm. C. McHarris, and W. H. Kelly, American Physical Society 1969 Winter Meeting in New York, 3-6 Feb 1969, paper AE12, Bull. Am. Phys. Soc. 14, 20 (1969).
2. "Nuclear Chemistry Research with Heavy Ions", Wm. C. McHarris, ANL-AUA Regional Conference on Discussion of a Midwest Accelerator for Nuclear Chemistry and Physics, Chicago, 11 Jan 1969, Argonne Physics Division Report PHY-1969A, p. 77.
3. "The Michigan State University 50-MeV Sector-Focused Cyclotron" (Invited Paper), Symposium on "Major National Nuclear Chemistry Facilities", American Chemical Society Minneapolis Meeting, 13-18 Apr 1969, paper NUCL .
4. "A Three-Quasiparticle Interpretation of the High-Lying Negative-Parity States in Pr^{139} ", Wm. C. McHarris, D. B. Beery, and W. H. Kelly, American Physical Society 1969 Spring Meeting in Washington, 28 Apr-1 May 1969, paper HF10, Bull. Am. Phys. Soc. 14, 627 (1969).
5. "The Population of States in Ce^{140} by Pr^{140} Decay", D. B. Beery,

- W. H. Kelly, and Wm. C. McHarris, American Physical Society 1969 Spring Meeting in Washington, 28 Apr-1 May 1969, paper EF11, Bull. Am. Phys. Soc. 14, 570 (1969).
6. "Decays of Ba^{131m} , Ba^{133m} , and Ce^{137g+m} , W. H. Kelly, D. B. Beery, and Wm. C. McHarris, American Physical Society 1969 Spring Meeting in Washington, 28 Apr-1 May 1969, paper EF8, Bull. Am. Phys. Soc. 14, 569 (1969).
7. "The Decay Scheme of Zn^{63} ", G. C. Giesler, Wm. C. McHarris, and W. H. Kelly, American Physical Society 1969 Spring Meeting in Washington, 28 Apr-1 May 1969, paper DF8, Bull. Am. Phys. Soc. 14, 551 (1969).
8. "Data Acquisition from Simultaneous Experiments Using the MSU Sigma-7 Computer" (Invited Paper), cf. paper #8 in the list above.