Department of Applied Mathematics and Computer Science

School of Engineering and Applied Science

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from, the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

An Algorithm for Solving
Integer Linear Programming Problems

by

Leon Cooper

Washington University

This research was performed under AEC

Research Contract No. AT(11-1)-1493

Report No. COO-1493-31

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED
leg

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Preface

The work described in this report was performed and written in 1963-64. For that reason, I have not changed its original form. It was never formally presented as a report or paper. I am issuing it now for several reasons. First, a paper was presented by H. Eto, "A Simplicies Decomposition Algorithm for Integer Linear Programming" at the June, 1969 meeting of the Operations Research Society in Denver, which has substantially the same content as this paper. Mr. Eto read an early version of this work but does not acknowledge my name or work. His algorithm is based on my early paper. Second, it is now realized that this work represents one of a large class of related algorithms which are currently being developed and which will be presented in subsequent reports and publications by the present author. Third, Dr. R.A. Cook developed some of these ideas successfully into a computational method. He explored one particular method of avoiding having to solve large sets of linear inequalities. This work is reported in his doctoral dissertation, "An Algorithm for Integer Linear Programming", Washington University, 1966. Unfortunately, Dr. Cook has not seen fit to publish this work. However, a report has been issued containing most of this work, Report AM-6J-2, May, 1965, "An Algorithm for Integer Linear Programming", by R.A. Cook and L. Cooper.

An Algorithm For Solving Integer LP Problems

I. Introduction

This paper is concerned with solving linear programming problems in integers. Whereas non-integer linear programming theory makes use of the properties of convex sets, i.e., the fact that all feasible solutions to a linear programming problem are members of a convex set and conversely, all members of this set are feasible solutions, this principle can not be used directly, if at all, to solve the integer programming problem. The reason for this is that the set of all feasible solutions to this problem does not, strictly speaking, constitute a convex set. A simple example as shown in Figure 1 in 2-space illustrates this point. Only the lattice points in the shaded area are feasible integer points. It is clear that this collection of points is not a convex set. Even if one wished to consider the convex hull of these lattice points (the shaded area in Figure 1) it is not at all a simple matter to obtain a knowledge of new constraints which would define the convex hull of these lattice points. Even if this is done and simplex theory applied there may still be problems of application.

The two main proposals for solving integer linear programming problems are those of Gomory [1] and Land and Doig [2]. The method of Gomory starts with the simplex solution. If this solution is not an integer solution then a new constraint may be added according to rules developed by

Figure I

Gomory. This process is repeated until an optimal integer solution is obtained.

In practice, Gomory's algorithm, embodied in computer codes, has had only moderate success. Many people have reported even small problems (e.g., a matrix of size 4×12) which were taken off a computer with no solution after several thousand iterations. However, many other problems have been solved.

Land and Doig's algorithm also starts with the simplex solution. If it is not an integer solution, one of the variables is then made integer by using the notions of parametric programming to determine the integer values in the range of feasible values of this variable. The variable is set at the best value with reference to the objective function and then this process is repeated for a second variable. This is done again for the next best integer value of the first variable and the second variable. From particular combinations of values of these two variables we proceed to make a third variable an integer, a fourth variable, etc. This is a laborious process and not much computational experience has been reported with this algorithm. However, given enough time, it would undoubtedly find a solution. It can also be used easily with mixed integer programming problems.

In the remainder of this paper a new algorithm is proposed for the solution of integer linear programming problems. It is quite different in spirit and detail from either the Gomory algorithm or the algorithm of Land and Doig.

II. General Description

The problem we wish to solve can be stated as the following maximization problem:

$$\begin{aligned} \text{Max } Z &= \bar{c} \bar{x} \\ \text{subject to: } A \bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \\ \bar{x} &\text{ integral} \end{aligned} \tag{1}$$

We assume that A is an $m \times n$ matrix consisting of integer elements, $\bar{b}$ is an m -vector with integer components and $\bar{x}$ is an n -vector with integer components. We assume further that $\bar{c}$ is an n -vector with integer components. The vectors are usually considered to be points in an n -dimensional Euclidean space E_n . We assume further that (1) has a finite maximum.

If we solve the usual linear programming problem, i.e., the problem given by (1) with the integrality restriction suppressed, (hereafter denoted by (1)'), we will obtain a solution $\bar{x} = \bar{x}_m$ with a corresponding value of the objective function $z = z_m$. We will assume that in this solution of the non-integer linear programming problem there has been no significant loss in precision in the computation of z_m . If desired the simplex calculation could be done in terms of rational fractions, although this is probably not necessary.

From the integrality of $\bar{c}$, it follows that the value z_0 of the objective function for the optimal solution to (1) must be an integer, and

that $z_0 \leq z_m$. Thus, if the problem has a solution, it must be that the optimal value of $\bar{c} \bar{x}$ is one of the values $\lfloor z_m \rfloor, \lfloor z_m \rfloor - 1, \dots, \lfloor z_m \rfloor - i, \dots$, where $\lfloor a \rfloor$ denotes the greatest integer $\leq a$.

The central idea of the algorithm is to form, at each iteration, a constraint set such that any lattice point in the set is automatically a solution to the problem, and then to find such a lattice point, if it exists. At any iteration, this set of constraints consists of the original set given by (1)' and one additional equality, which has the form

$$\bar{c} \bar{x} = Z_i^* \quad (2)$$

where

$$Z_i^* = \lfloor z_m \rfloor - (i-1) \quad i=1, 2, \dots \quad (3)$$

and i represents the i^{th} iteration. If, for $i=1$, the feasible region defined by (1)' and (2) contains a lattice point, then clearly it represents the optimal solution. If the region contains no lattice point, then the required value for $\bar{c} \bar{x}$ is decreased by 1 and the argument repeated, and so on.

As illustrated in Figure II, the above consists of moving the optimal hyperplane into the feasible region of (1)' by discrete amounts until it intersects a lattice point. Point C represents the optimal solution to (1)'. Since point D, the optimal solution to (1), is not an extreme point of the constraint set described by (1)' and (2), the simplex algorithm cannot be used to find it.

The i^{th} iteration thus consists of solving the following problem:
find a lattice point $\bar{x}$ such that

Figure II

$$\begin{aligned} A\bar{x} &\leq \bar{b} \\ \bar{c}\bar{x} &= z_i^* \\ \bar{x} &\geq \bar{0} \end{aligned}$$

(4)

This is done by first solving

$$\bar{c}\bar{x} = z_i^*$$

(2)

as a linear diophantine equation.

After the general solution to the linear diophantine equation is obtained and substituted into

$$\begin{aligned} A\bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \end{aligned}$$

(5)

the inequalities (5) can be examined to determine whether or not they possess an integer solution. If (5) does possess an integer solution it is automatically optimal. If it does not, we move to the next iteration, i.e., penetrate more deeply into the convex polyhedron of feasible solutions and repeat the process. This entire process will now be described in more detail.

Equation (2) can be written:

$$C_1 x_1 + C_2 x_2 + \dots + C_n x_n = z_i^*$$

(6)

Equation (6) is to be treated and solved as a linear diophantine equation.

The general theory of such equations is very old and well known. (See [4]).

If the greatest common divisor of $c_1, c_2, \dots, c_n$ (symbolized $D(c_1, c_2, \dots, c_n)$) does not divide z_1^* , then there are no solutions to (6). Hence if (6) possesses no integer solutions at any iterative stage, we can immediately move on to the next iteration. If $D(c_1, c_2, \dots, c_n)$ does divide z_1^* , we must now determine if any of the solutions satisfy the conditions expressed by inequalities (5), i.e., whether there are any non-negative solutions in the feasible region of the optimal hyperplane (2).

The general solution to a linear diophantine equation in n variables can be written in the following form (See [4]):

$$x_j = \sum_{k=1}^{n-1} d_{jk} \beta_k + \gamma_j \quad j=1, 2, \dots, n \quad (7)$$

where d_{jk} and γ_j are known integer constants and the β_k are arbitrary integer variables. It is a relatively simple computational matter to solve (6) and obtain the solution (7) and we have written computer codes to determine the d_{jk} and γ_j of the solution (7).

After the equations (7) have been obtained they are substituted into (5), the original constraint set of the integer linear programming problem. First we write (7) in matrix form as:

$$\bar{X} = D\bar{\beta} + \bar{\gamma} \quad (8)$$

where: $\bar{x}$ is an n -vector of the x_j

D is an $n \times (n-1)$ matrix of d_{jk}

$\bar{\beta}$ is an $(n-1)$ - vector of the β_k

$\bar{\gamma}$ is an n -vector of the γ_j

If we now substitute (8) into (5) we have:

$$\begin{aligned} A\bar{x} &= A[D\bar{\beta} + \bar{\gamma}] \leq \bar{b} \\ \bar{x} &= D\bar{\beta} + \bar{\gamma} \geq \bar{0} \end{aligned} \quad (9)$$

If we now define:

$$\begin{aligned} P &= \begin{bmatrix} -AD \\ D \end{bmatrix} \\ \bar{f} &= \begin{bmatrix} A\bar{\gamma} - \bar{b} \\ -\bar{\gamma} \end{bmatrix} \end{aligned} \quad (10)$$

where P is an $(m+n) \times (n-1)$ matrix

$\bar{f}$ is an $(m+n)$ - vector

then inequalities (9) can be written:

$$P\bar{\beta} \geq \bar{f} \quad (11)$$

The major problem at each iteration is to determine whether or not inequalities (11), do or do not have any integer solutions, since any integer solution to (11) is optimal.

Solving a large set of linear inequalities such as (11) can be a tedious business. Hence the crux of the computational problem will be to find relatively rapid, efficient methods which may possibly take advantage of the special structure of the inequalities. Some possibilities are presently being investigated. The computational problem, in general, is discussed in a subsequent section. It should be pointed out, however, that there is a general procedure of elimination that, if all else fails, can be used to solve a set of inequalities such as (11).

Fourier [5] and more recently Motzkin [6] have both given consideration to the general problem of the solution of linear inequalities. The method proposed here is based on these ideas. Motzkin [6] has given a detailed theoretical analysis of the conditions for existence of solutions to linear inequalities. The elimination method referred to here is as follows:

It is desired to solve

$$P\bar{\beta} \geq \bar{f} \quad (11)$$

which is a set of $m+n$ inequalities in $n-1$ variables, the β_k . Rewriting (11) in component form, we have

$$\sum_{k=1}^{n-1} p_{jk} \beta_k - f_j \geq 0 \quad j=1,2,\dots,m+n \quad (12)$$

Let J_1 (J_2) be the subset of the set of integers $j=1,2,\dots,m+n$ corresponding to those inequalities of (12) which contain a negative (positive) coefficient of β_1 . In other words, J_1 contains those j such that p_{j1} is negative;

J_2 those for which p_{j1} is positive. From these inequalities we can write:

$$\beta_1 \leq \sum_{k=2}^{n-1} p'_{jk} \beta_k - f'_j \quad \text{for } j \in J_1 \quad (13)$$

$$\beta_1 \geq \sum_{k=2}^{n-1} p'_{jk} \beta_k - f'_j \quad \text{for } j \in J_2 \quad (14)$$

where

$$\begin{aligned} p'_{jk} &= -p_{jk}/p_{j1} & k=2,3,\dots,n-1 \\ f'_j &= f_j/p_{j1} & j \in J_1 \cup J_2 \end{aligned}$$

From (13) and (14) we can obtain a new set of inequalities of the form

$$\sum_{k=2}^{n-1} p'_{j_1 k} \beta_k - f'_{j_1} \geq \sum_{k=2}^{n-1} p'_{j_2 k} \beta_k - f'_{j_2} \quad j_1 \in J_1 \text{ and } j_2 \in J_2$$

or

$$\sum_{k=2}^{n-1} (p'_{j_1 k} - p'_{j_2 k}) \beta_k - (f'_{j_1} - f'_{j_2}) \geq 0 \quad j_1 \in J_1 \text{ and } j_2 \in J_2 \quad (15)$$

If the set J_1 has n_1 elements and the set J_2 has n_2 elements, then there will be $N_1 = n_1 n_2$ inequalities of the form (15). In general, the maximum number

of new inequalities which can be generated is $\leq \left(\frac{m+n}{2}\right)^2$; i.e., $N_1 \leq \left(\frac{m+n}{2}\right)^2$. Normally we would expect N_1 to be quite a bit less than this, however, since not all inequalities (12) need contain β_1 (p_{j1} might be zero), and since the maximum would only be approached if n_1 and n_2 were nearly equal.

The inequality set (15) has the same form as (12) except that it has one fewer variable, since β_1 has been eliminated. This elimination process can now be repeated with β_2, β_3 , etc., until finally, one arrives at a set of N_{n-2} inequalities involving a single variable β_{n-1} . If this set is consistent, then it may have one or more integer β_{n-1} for a solution. If it is not consistent, the original set of inequalities (11) has no integer solutions. If a β_{n-1} has been found, the back solution for $\beta_{n-2}, \dots, \beta_1$ is easily carried out by substitution and the question of consistency determined at each stage.

It is easily shown by induction that an upper bound for the number of inequalities N_{n-2} involving β_{n-1} , at the last stage of the elimination process, is given by

$$N_{n-2} \leq 4 \left(\frac{m+n}{4} \right)^{2^{n-2}} \quad (16)$$

The problem to be solved at each iteration has now been discussed. If an integer solution is encountered at any stage, it is optimal. The only remaining point is the finiteness of the algorithm.

It is theoretically possible that the convex set of points described by

$$\begin{aligned} A\bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \end{aligned} \quad (5)$$

contains no lattice points. This could occur either because the set is too "small", or more likely because some of the constraints were actually equalities which have no integer solutions. In any case, this could be determined if necessary by solving (5) in exactly the same way as described previously. If $\bar{x}_L$ is an integer solution, then clearly the stated problem has a solution, and

$$Z_L = \bar{c}\bar{x}_L \leq Z_0 \leq Z_M = \lfloor Z_m \rfloor \quad (17)$$

That is, only integer values between z_L and z_M need be tried as values for z_i^* .

Alternatively, a lower bound z_L for trial values of z_i^* could be obtained by solving the corresponding minimization problem

$$\begin{aligned} \text{Min } Z &= \bar{c}\bar{x} \\ \text{subject to: } A\bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \end{aligned} \quad (18)$$

If z_ℓ represents the solution to (18), then certainly

$$Z_L = \lceil z_\ell \rceil \leq Z_0 \leq Z_M = \lfloor Z_m \rfloor \quad (19)$$

where $\lceil a \rceil$ denotes the least integer $\geq a$.

However, it is possible that (18) has an unbounded minimum. In this case rather than resort to (17), it may be preferable to impose some arbitrary lower bound, which could be dictated by the particular problem involved.

III. Summary of the Algorithm.

1. We seek to solve the problem

$$\text{Max } z = \bar{c}\bar{x}, A\bar{x} \leq \bar{b}, \bar{x} \geq \bar{0}, \bar{x} \text{ integral} \quad (\text{A})$$

(The matrix A, vectors $\bar{b}$, $\bar{c}$ are integral)

2. Solve problem (A), without the integer restriction on $\bar{x}$, by a linear programming algorithm. Let the optimal value of the objective function be z_m . Also determine z_l as $\min z = \bar{c}\bar{x}$ subject to $A\bar{x} \leq \bar{b}$, $\bar{x} \geq \bar{0}$, or by some other method as described previously. Let $z_M = \lfloor z_m \rfloor$ and $z_L = \lceil z_l \rceil$.
3. If $\bar{x}_m$, the solution yielding z_m , is integral we are done.
If not, we go to 4.
4. The typical iterative calculation now begins: On the i^{th} iteration, let $z_i^* = z_M - (i-1)$, $i=1,2,\dots$ We now consider the problem:

Find an integer solution $\bar{x}$, if any exists, to :

$$\begin{aligned} A\bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \\ \bar{c}\bar{x} &= z_i^* \end{aligned} \quad (\text{B})$$

5. The set of inequalities, (B) is solved by first solving $\bar{c} \bar{x} = z_i^*$ as a linear diophantine equation, substituting into

$$\begin{aligned} A\bar{x} &\leq \bar{b} \\ \bar{x} &\geq \bar{0} \end{aligned}$$

and determining integer solutions, if any.

6. If (B) has at least one integer solution we are done. If not, go to 7.
7. Increase i by 1. If $z_i^* \geq z_L$, we return to 4. and continue the calculation. If $z_i^* < z_L$, then there is no integer solution to (A).

IV. Numerical Example

Consider the following problem:

$$\begin{aligned} \text{Max } z &= 3x_1 + 4x_2 + x_3 + 7x_4 \\ \text{Subject to : } 8x_1 + 3x_2 + 4x_3 + x_4 &\leq 7 \\ 2x_1 + 6x_2 + x_3 + 5x_4 &\leq 3 \\ x_1 + 4x_2 + 5x_3 + 2x_4 &\leq 8 \\ x_j &\geq 0, \text{ integer } j=1,2,3,4 \end{aligned} \tag{20}$$

The optimal solution to the non-integer problem, found by simplex, is

$$\bar{x} = (0.842, 0, 0, 0.2632)$$

$$z_m = 4.368$$

Thus, $z_M = \lfloor 4.368 \rfloor = 4$. Clearly $z_L = 0$ is a lower bound.

We now attempt to find solutions, if any exist, to the following:

$$8x_1 + 3x_2 + 4x_3 + x_4 \leq 7 \quad (21)$$

$$2x_1 + 6x_2 + x_3 + 5x_4 \leq 3 \quad (22)$$

$$x_1 + 4x_2 + 5x_3 + 2x_4 \leq 8 \quad (23)$$

$$3x_1 + 4x_2 + x_3 + 7x_4 = 4 \quad (24)$$

$$x_1 \geq 0 \quad (25)$$

$$x_2 \geq 0 \quad (26)$$

$$x_3 \geq 0 \quad (27)$$

$$x_4 \geq 0 \quad (28)$$

Equation (24) is solved as a diophantine equation:

$$3x_1 + 4x_2 + x_3 + 7x_4 = 4$$

$$x_3 = 4 - 3x_1 - 4x_2 - 7x_4$$

$$\therefore x_1 = \beta_1$$

$$x_2 = \beta_2$$

$$x_3 = 4 - 3\beta_1 - 4\beta_2 - 7\beta_3$$

$$x_4 = \beta_3$$

(29)

where $\beta_1, \beta_2, \beta_3$ are arbitrary integers.

We now write (21) - (23), (25) - (28) in terms of β_j as follows, by substituting from (29):

$$4\beta_1 + 13\beta_2 + 27\beta_3 - 9 \geq 0 \quad (30)$$

$$\beta_1 - 2\beta_2 + 2\beta_3 - 1 \geq 0 \quad (31)$$

$$14\beta_1 + 16\beta_2 + 33\beta_3 - 12 \geq 0 \quad (32)$$

$$\beta_1 \geq 0 \quad (33)$$

$$\beta_2 \geq 0 \quad (34)$$

$$-3\beta_1 - 4\beta_2 - 7\beta_3 + 4 \geq 0 \quad (35)$$

$$\beta_3 \geq 0 \quad (36)$$

β_1 can be eliminated from (30) - (33), (35) giving rise to the following set (adding in (34) and (36)):

$$23\beta_2 + 53\beta_3 - 11 \geq 0 \quad (37)$$

$$-10\beta_2 - \beta_3 + 1 \geq 0 \quad (38)$$

$$-8\beta_2 + \beta_3 + 20 \geq 0 \quad (39)$$

$$-4\beta_2 - 7\beta_3 + 4 \geq 0 \quad (40)$$

$$\beta_2 \geq 0 \quad (41)$$

$$\beta_3 \geq 0 \quad (42)$$

β_2 can be eliminated from (37) - (41), which together with (42) gives the following:

$$507\beta_3 - 87 \geq 0 \quad (43)$$

$$-\beta_3 + 1 \geq 0 \quad (44)$$

$$447\beta_3 + 372 \geq 0 \quad (45)$$

$$\beta_3 + 20 \geq 0 \quad (46)$$

$$51\beta_3 + 48 \geq 0 \quad (47)$$

$$-7\beta_3 + 4 \geq 0 \quad (48)$$

$$\beta_3 \geq 0 \quad (49)$$

Examining (43) - (49) successively, we obtain:

$$\begin{array}{ll}
 \beta_3 \geq \frac{87}{507} & \therefore \beta_3 \geq 1 \\
 \beta_3 \leq 1 & \therefore 1 \leq \beta_3 \leq 1 \text{ or } \beta_3 = 1 \\
 \beta_3 \geq -\frac{372}{447} & \therefore \beta_3 = 1 \\
 \beta_3 \geq -20 & \therefore \beta_3 = 1 \\
 \beta_3 \geq -\frac{48}{51} & \therefore \beta_3 = 1 \\
 \beta_3 \leq \frac{4}{7} & \therefore \beta_3 \neq 1, \text{ which is a contradiction.}
 \end{array}$$

It is therefore clear that there are no integer solutions to (21) - (28).

We would now continue the process and at the next iteration, attempt to resolve (21) - (28) with equation (24) replaced by:

$$3x_1 + 4x_2 + x_3 + 7x_4 = 3 = z_M - 1$$

This problem would also prove to have no integer solution, as would the following iteration with $z_3^* = 2$.

At the 4th iteration $z_4^* = 1$, and we attempt to solve (21) - (28)

with (24) replaced by

$$3x_1 + 4x_2 + x_3 + 7x_4 = 1 \quad (24)^*$$

Solving (24)*, we obtain

$$\begin{array}{ll}
 x_1 = \beta_1 & \\
 x_2 = \beta_2 & \\
 x_3 = 1 - 3\beta_1 - 4\beta_2 - 7\beta_3 & (29)^* \\
 x_4 = \beta_3 &
 \end{array}$$

and corresponding to (30) - (36) we get by substitution

$$4\beta_1 + 13\beta_2 + 27\beta_3 + 3 \geq 0 \quad (30)^*$$

$$\beta_1 - 2\beta_2 + 2\beta_3 + 2 \geq 0 \quad (31)^*$$

$$14\beta_1 + 16\beta_2 + 33\beta_3 + 3 \geq 0 \quad (32)^*$$

$$\beta_1 \geq 0 \quad (33)^*$$

$$\beta_2 \geq 0 \quad (34)^*$$

$$-3\beta_1 - 4\beta_2 - 7\beta_3 + 1 \geq 0 \quad (35)^*$$

$$\beta_3 \geq 0 \quad (36)^*$$

Eliminating β_1 gives:

$$23\beta_2 + 53\beta_3 + 13 \geq 0 \quad (37)^*$$

$$-10\beta_2 - \beta_3 + 7 \geq 0 \quad (38)^*$$

$$-8\beta_2 + \beta_3 + 23 \geq 0 \quad (39)^*$$

$$-4\beta_2 - 7\beta_3 + 1 \geq 0 \quad (40)^*$$

$$\beta_2 \geq 0 \quad (41)^*$$

$$\beta_3 \geq 0 \quad (42)^*$$

Eliminating β_2 we obtain a set corresponding to (43) - (49):

$$507\beta_3 + 291 \geq 0 \quad (43)^*$$

$$-\beta_3 + 7 \geq 0 \quad (44)^*$$

$$447\beta_3 + 633 \geq 0 \quad (45)^*$$

$$\beta_3 + 23 \geq 0 \quad (46)^*$$

$$51\beta_3 + 75 \geq 0 \quad (47)^*$$

$$-7\beta_3 + 1 \geq 0 \quad (48)^*$$

$$\beta_3 \geq 0 \quad (49)^*$$

Examining (43)* - (49)* we obtain successively:

$$\beta_3 \geq -\frac{291}{507}$$

$$\therefore \beta_3 \geq 0$$

$$\beta_3 \leq 7$$

$$\therefore 0 \leq \beta_3 \leq 7$$

$$\beta_3 \geq -\frac{633}{447}$$

$$\therefore 0 \leq \beta_3 \leq 7$$

$$\beta_3 \geq -23$$

$$\therefore 0 \leq \beta_3 \leq 7$$

$$\beta_3 \geq -\frac{75}{51}$$

$$\therefore 0 \leq \beta_3 \leq 7$$

$$\beta_3 \leq \frac{1}{7}$$

$$\therefore 0 \leq \beta_3 \leq 0 \quad \text{or} \quad \beta_3 = 0$$

Substituting $\beta_3 = 0$ into (37)* - (42)* we find that $\beta_2 = 0$. Substituting for β_2 and β_3 into (30)* - (36)* we find that $\beta_1 = 0$. Therefore from equations (29)* we find that:

$$x_1 = 0$$

$$x_2 = 0$$

$$x_3 = 1$$

$$x_4 = 0$$

is the optimal integer solution to (20).

V. Computational Considerations

The extent to which the algorithm proposed in this paper can be applied in practice to large integer linear programming problems which might be encountered in industrial and military practice has yet to be determined in detail. Studies are now under way to examine the feasibility of employing the algorithm for large problems. The major computational problems which are seen to exist are as follows:

- 1) Performing fixed point arithmetic on a digital computer with large numbers
- 2) Solving the sets of linear inequalities at each step of the iterative process.

Let us briefly examine each of these problems.

The problem of performing fixed point arithmetic and handling rational fractions is not a problem in any theoretical sense. For example, in a digital computer with variable word length and a very large fast access memory, the calculations could be easily performed. In a fixed word length computer each numerator and denominator could be represented by a single word and sub-routines written to perform rational fraction arithmetic. If the numbers became too large for a single word, in principle, it appears that two or more words could be used for each numerator and denominator. The extent to which this is practical and is required, is currently being explored.

The second problem, of solving large sets of linear inequalities, can also be handled in principle, as the second part of this paper indicates. However, the practicality of solving the large sets of inequalities that the

algorithm generates, by the general process of elimination that is described, has to be examined by performing calculations with actual problems, which is now being undertaken.

First, it should be noted that even if no better general way of solving these inequalities can be found, in many problems it is unnecessary to go through the complete elimination process described. At each iteration we wish to know if there is no solution or, if there is one, to find it as rapidly as possible. The following principles can be used and they can be incorporated in a computational code.

1. Any of the constraints of the original integer linear programming problems that are strict equalities can be solved simultaneously with the adjoined equality constraint as a set of linear diophantine equations thus reducing the total number of inequalities that need to be solved, and also the number of variables, since each equation can be used to eliminate one variable.
2. The set of inequalities (11) can be scanned for any obvious contradictions. If any occur, the calculation can proceed to the next iteration.
3. At any point in any iterative stage the uncovering of a contradiction implies that the set of inequalities has no solution. Hence, we can proceed to the next iteration.
4. The transition from one iteration to the next can be greatly simplified, since many of the calculations are the same at each iteration.

The general elimination process should not be as forbidding as the upper bound calculation indicates. Computer codes are now being written to explore this matter.

VI. Summary

A new algorithm has been described to solve integer linear programming problems which is based on the idea of solving as a linear diophantine equation a constraint parallel to the objective function hyperplane and then ascertaining if the original constraint inequalities and the linear diophantine solution have an integer solution in common. Some computational considerations which may affect the applicability of the algorithm to problems with many variables have also been briefly discussed.

References

1. Gomory, R.E., "Outline of an Algorithm for Integer Solutions to Linear Programs," Bulletin of The American Mathematical Society, LXIV pp. 275-8, (1958).
2. Land, A.H. and Doig, A.G., "An Automatic Method of Solving Discrete Programming Problems," Econometrica, XXVIII, No. 3, pp. 497-521, (1960).
3. Hadley, G. "Linear Programming," Addison-Wesley, 1962.
4. Cahen, Eugene. "Theorie des Nombres", Vol. I. Paris, A. Hermann 1914-1924.
5. Fourier, J.B.J., "Solution d'une question particuliere du Calcul des inequalities," 1826 and extracts from "Histoire de l'academie", 1823, 1824, Oeuvres II, pp. 317-328.
6. Motzkin, T.S., "Beitrage zur Theorie der Linearen Ungleichungen," Jersua'lem (1936) (Doctoral Dissertation, Basel).