FEB 11 1964

AROD # 14289

FINAL REPORT

AROD Grants DA-AROD-31-124-C117 and G534

"Research in Fluorine Chemistry"

June 1, 1961, to February 28, 1963

Department of Chemistry, University of Wisconsin, Madison, Wisconsin

Personnel:

Principal Investigator: John L. Margrave

Post-doctoral and Graduate Student Personnel: V. V. Dadape,
M. A. Frisch and A. S. Kana'an

Undergraduate Assistants: C. K. Barker, B. Bork, J. D. Borkowski,
R. G. Czerepinski and D. C. Lewis

The following projects were conducted in whole, or in part, with funds supplied under this AROD grant. Other support came through grants from the U. S. AEC, the NSF and the Wisconsin Alumni Research Foundation.

I. Synthesis of Tetrafluorochlorates

Early work which claimed to have produced MF_{1+x} and similar compounds by reaction of MCl or MBr or MI with F_2 has been shown to be in error. Actually, with KCl one produces $KClF_4$ and similarly with Rb and Cs halides.¹

1. (a) M. Silverthorn, L. Asprey and J. Margrave, J. Am. Chem. Soc. 83, 2955 (1961).
- (b) M. Silverthorn, M. S. Thesis, University of Wisconsin (1960).

Further studies of this system were conducted by R. Czerepinski who attempted the direct reaction

A 60-70% yield was obtained but high-purity $CsClF_4(s)$ could not be isolated.

The work has been summarized in the B. S. thesis of Mr. Czerepinski.²

2. R. G. Czerepinski, B. S. Thesis, University of Wisconsin, June, 1962.

Facsimile Price \$ 1.10

Microfilm Price \$.80

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

PATENT CLEARANCE OBTAINED. REFERENCE TO
THE PUBLIC IS APPROVED. PROCEEDINGS
ARE ON FILE IN THE RECEIVING SECTION.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

1. Reactions of SiH_4 with Oxides and Nitrides

A nickel system for handling SiH_4 and ClH_3 was constructed and utilized for studies of oxides and nitrides following the method of Katz and associates at the Argonne National Laboratory. Studies of IN samples were carried out which showed that the samples were essentially $\text{H}_{1.00}\text{N}_{1.00}$ but the experimental reproducibility was not better than $\pm 1\%$. Samples of SiO_2 were examined and it was established as 1.19 ± 0.01 .

1. The Properties of SiF_2 and H_2

Available data on SiF_2 and H_2 were reviewed and two publications resulted which established reasonable ranges for the thermodynamic stabilities of these molecules ^{3,4} Some quantitative properties of reduced fluorides

3. D. Pease, A. Farnen and J. Margrave, J. Phys. Chem. 66, 1200 (1962).
4. J. Margrave, J. Phys. Chem. 66, 1209 (1962).

of Si were sought

- (1) through a study of the transpiration weight loss of Si when exposed to a flow of SiH_4 according to the reaction

- (2) through observations of emission and absorption spectra of such systems, and

- (3) through observation of mass spectra of such systems.

The first two parts of this goal were covered by the Ph. D. thesis of A. S. Kazan who found that there is definitely a polymeric species of SiF_2 as the major species as the monomer. Pressure dependence studies support $x > 2$ and, by comparison with CF_2 , one predicts the existence of a

cyclic tetramer, Si_4F_8 .⁵ No new optical spectra were obtained.

5. (a) A. S. Konaun, Ph. D. Thesis, University of Wisconsin (1963).
(b) A. S. Konaun and J. L. Margrave, accepted for publication, Inorganic Chemistry (1964).
-

Mass spectrometric work concerned the products of the reaction between CaF_2 and Si in a Knudsen cell at high temperatures. $\text{SiF}(g)$ and $\text{SiF}_2(g)$ were definitely produced and their stabilities were established. For $\text{SiF}(g)$ one finds $\Delta H_{298}^\circ = -3 \pm 6$ Kcal/mole and for $\text{SiF}_2(g)$, one finds $\Delta H_{298}^\circ = -139 \pm 2$ Kcal/mole.⁶ A similar mass spectrometric study

6. T. C. Ehlert, Ph. D. Thesis, University of Wisconsin (1963).
-

of the $\text{Ge} + \text{CaF}_2$ reaction established $\Delta H_{298}^\circ [\text{GeF}(g)] = -7.4 \pm 5$ Kcal/mole and $\Delta H_{298}^\circ [\text{GeF}_2(g)] = -19 \pm 20$ Kcal/mole.

Studies of BF_2 , CF_2 and AlF_2 were not successful and further work is still in progress at the Rice University, Houston, Texas, under the extension of this grant.

IV. Thermodynamic Properties of Fluorine Compounds

A fluorine bomb calorimeter was operated in a collaborative project at the Argonne National Laboratory and the heats of formation of BF_3 , SiF_4 and HF were established.^{7,8,9} In addition, the heats of formation

7. S. Wise, H. Feder, W. Hubbard and J. Margrave, J. Phys. Chem. 65, 2157 (1961).
8. (a) S. Wise, H. Feder, W. Hubbard and J. Margrave, J. Phys. Chem. 66, 381 (1962).
(b) S. Wise, J. Margrave, H. Feder, W. Hubbard, J. Phys. Chem. 67, 815 (1963).
9. H. Feder, W. Hubbard, S. Wise and J. Margrave, J. Phys. Chem. 67, 1148 (1963).
-

of SiO_2^8 and BN^{10} were determined.

10. S. S. Wise, Ph. D. Thesis, University of Wisconsin (1963).

V. Miscellaneous Work

Several efforts were made to synthesize AlOF but none were successful. High pressures were used in an effort to get crystalline AlF but no definitive material was prepared. Samples of molybdenum oxychlorides were prepared but will have to be characterized by further work since there appear to be several oxychlorides.

Fluorine compounds (CF_4 , SiF_4 , CF_2Cl_2 , etc.) were passed through a plasma torch but no new compounds were synthesized. When CCl_4 was passed through a plasma torch, C_6Cl_6 was produced.¹¹

11. A. S. Kanaan and J. L. Margrave, Int. Sci. and Tech., No. 8, 75 (1962).