OCT 29 1963

UCRL-10841

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

CONF-271-1

MASTER

OLD AND NEW PROBLEMS IN THE FIELD OF VAPOR-LIQUID EQUILIBRIA

Otto Redlich

July 16, 1963

Facsimile Price \$ 1.10

Microfilm Price \$.88

Available from the
Office of Technical Services
Department of Commerce
Washington 25, D. C.

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:
A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Abstract of a Lecture to be delivered at the 13th Canadian Chemical Engineering Conference, Montreal, October 23, 1963.

Old and New Problems in the Field of Vapor-Liquid Equilibria

Otto Redlich

Department of Chemical Engineering, and Inorganic Materials Research Division,
Lawrence Radiation Laboratory, University of California, Berkeley, California.

The practical advantage of a systematic representation of vapor-liquid equilibria by means of activity coefficients rests, of course, on the fact that the results are loosened from the restriction to the original experimental conditions and can be extended, more or less reliably, to different conditions to multicomponent mixtures, and even to different substances.

We derive activity coefficients from the equilibrium condition that the chemical potential or partial molal free energy of a substance has the same value in the liquid and the vapor. Notwithstanding formal analogies, we have to treat the two phases differently, for the simple reason that we must assume the chemical potentials of the components in the gas to be known while the potentials in the liquid, or the activity coefficients, are the unknowns in the problem. The most common offense against the requirement of introducing a correct value of the free energy in the vapor phase is, of course, disregard of appreciable deviations from the laws of the perfect gaseous

mixture. Somewhat subtler, though essentially the same mistake, is the disregard of a change of the molecular state, such as the dimerization of formic acid, in the vapor state. The usual approximations for fugacity coefficients cannot correctly represent such chemical changes. Activity coefficients, on the other hand, as the unknowns in the calculation, stand on their own feet without any regard to the molecular state, provided only that the molecular weight in the dilute solution has been correctly introduced.

The phase rule shows that we overdetermine the unknowns of a binary mixture if we measure (for good experimental reasons) the mole fractions x_1 and y_1 in both phases as functions of temperature and pressure. The ensuing identical relation between the activity coefficients of the two components, essentially the relation of Duhem and Margules, has been a favorite subject of discussion in recent years. I wish to present my apologies because Mr. A. T. Kister and I¹ are in a way guilty of having started this discussion. By way of excuse it may be mentioned that in 1946 consistency criteria were quite useful, in view of the poor state of the available experimental material. Moreover, we said quite briefly (though with rigorous accuracy) what some later authors² said in extensive discussions. Perhaps we could have avoided some unjustified criticism³ regarding supposed restrictions of the area condition if we had expanded our discussion. We thought, however, at that time as we do today, that consistency criteria should be used but not much talked about; the experimenter should use this tool to control and improve his methods.

Automatic computation can be very useful if handled sensibly. The real question is how much one should leave to critical judgment in a problem in which functional relationships are quite complicated and experimental errors have widely varying influence. So far as I can see, the first correct completely mechanized procedure was recently suggested by Techo⁴. He smoothes the immediately observed quantities P and y_1 , introducing the equation of Duhem and Margules as a condition, and represents the results by orthogonal polynomials. Since he can assign equal weights to the

3

directly observed values, he avoids the usual pitfall of wrong weighting. Techo's procedure has the intrinsic advantage and disadvantage of a strictly mechanical method: It does not depend on any whim of the calculator and it does not utilize his experience and critical judgment. Moreover, it does not depend on, or utilize, any specific model.

Though often neglected, deviations of the vapor from a perfect gaseous mixture are frequently quite important. A few types of equations of state⁶⁻¹² have survived a large number of competitors. Each of these relations has important advantages, which require a special discussion. None is satisfactory in every respect so that it could be used exclusively.

The critical state of gaseous mixtures is of appreciable theoretical and practical interest. The critical lines of a binary mixture (T and P as functions of y_1) can be derived from an equation of state but the calculations are extremely involved¹³. The explicit representation of the limiting slopes¹⁴ of the critical lines in their endpoints (i.e., for the pure components) presents some advantage. Further progress will depend on improvements of the equations of state.

The relation between interaction in non-electrolyte solutions and the chemical nature of the components presents an old question. A condition of a satisfactory answer appears to be the further quantitative development of simple models for association, interassociation and compound formation¹⁵. As soon as these "chemical" influences are accounted for or whenever they are absent, one may expect to represent the remaining "physical" interaction by means of two parameters for any binary. Hildebrand's solubility parameters furnish an approximation for the interaction¹⁶. A group-interaction model¹⁷ and similar semi-empirical relations¹⁸ are practically useful and promise more elucidation in the future.

A considerable part of this review is based on work done at Shell Development Company, Emeryville, California. The author feels a profound obligation to the late Mr. A. Lacombe and the late Dr. R. W. Millar for inviting him to start this work. He has greatly enjoyed the cooperation of Mr. A. T. Kister, Dr. J. N. S. Kwong, Mr. C. E. Turnquist, Mr. C. M. Cable, Mr. J. M. Monger, Dr. M. N. Papadopoulos and Mr. A. K. Dunlop. He has greatly appreciated the opportunity to use unpublished data and results of Dr. C. L. Dunn, Mr. G. J. Pierotti, Dr. C. Black, Dr. C. H. Deal, Mr. E. L. Derr and others. He has also enjoyed the cooperation of Mr. F. J. Ackerman and Mr. R. D. Gunn at the University of California.

Bibliography

1. O. Redlich and A. T. Kister, Ind. Eng. Chem. 40, 341 (1948).
2. For instance, N. V. Ibl and B. F. Dodge, Chem. Eng. Sci. 2, 120 (1953).
3. J. S. Rowlinson, "Liquids and Liquid Mixtures," London, Butterworth (1959).
4. R. Tcho, Chem. Eng. Sci. 18, 27 (1963).
5. O. Redlich and J. N. S. Kwong, Chem. Revs. 44, 233 (1949).
6. Lewis-Randall-Pitzer-Brewer, "Thermodynamics," App. 1. New York: McGraw-Hill Book Company, 1961.
7. L. Riedel, Chem. Eng. Techn. 26, 83, 259, 679 (1954); 27, 209, 475 (1955); 28, 557 (1956).
8. O. Redlich and A. K. Dunlop, Chem. Eng. Progr. Symposium Series 52, 95 (1963).
Unpublished work by F. J. Ackerman and R. D. Gunn.
9. M. Benedict, G. B. Webb and L. C. Rubin, J. Chem. Phys. 8, 334 (1942); 10, 747 (1942).
10. C. Black, Ind. Eng. Chem. 50, 391 (1958).
11. J. J. Martin and Yu-Chun Hou, A.I.Ch.E. Journal 1, 142 (1955).
12. J. M. Prausnitz and R. D. Gunn, A.I.Ch.E. Journal 4, 430 (1958).
- J. M. Prausnitz and W. B. Carter, A.I.Ch.E. Journal 6, 611 (1960). J. M. Prausnitz and R. Anderson, A.I.Ch.E. Journal 7, 96 (1961).
13. H. J. Löffler, Chem. Ing. Techn. 34, 79 (1962).
14. O. Redlich and A. T. Kister, J. Chem. Phys. 36, 2002 (1962). O. Redlich and F. J. Ackerman, J. Chem. Phys. 38, 2740 (1963).

15. O. Redlich and A. T. Kister, J. Chem. Phys. 15, 849 (1947). G. Scatchard, Chem. Revs. 44, 7 (1949). J. Prigogine, V. Mathot and A. Desmyter, Bull. soc. chim. Belges 58, 547 (1949).
16. J. M. Prausnitz, W. C. Edmister and K. C. Chao, A.I.Ch.E. Journal 6, 214 (1960).
17. O. Redlich, E. L. Derr and G. J. Pierotti, J. Amer. Chem. Soc. 81, 2283 (1959). M. N. Papadopoulos and E. L. Derr, J. Amer. Chem. Soc. 81, 2283 (1959). G. M. Wilton and C. H. Deal, Ind. Eng. Chem. Fundamentals 1, No. 1, 20 (1962).
18. G. J. Pierotti, C. A. Deal and E. L. Derr, Ind. Eng. Chem. 51, 95 (1959).

6