

MASTER

E2/M1 Multipole Mixing Ratios in the
 "Spherical" Nuclei Te^{124} , Te^{126} , and Xe^{126*} †

Z. W. Grabowski, K. S. Krane[†] and R. M. Steffen

Department of Physics, Purdue University

Lafayette, Indiana 47907

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

ABSTRACT

E2/M1 mixing ratios of the 709 keV, 714 keV and 723 keV transitions in Te^{124} , the 754 keV transition in Te^{126} , and the 491 keV transition in Xe^{126} were measured using the γ - γ directional correlation technique. In addition, the measurements on the 646 keV-603^{keV}, 1691 keV-603 keV and 2091 keV-603 keV γ - γ cascades in Te^{124} are in agreement with pure E2 character of 646 keV transition and with pure E1 character of the 1691 keV and 2091 keV transitions in Te^{124} . Coaxial Ge(Li) and NaI(Tl) scintillation detectors were used in different combinations depending on the energy resolution required for a particular measurement. The results for the mixing ratios are $\delta_{709} = + 0.04^{+0.03}_{-0.05}$, $\delta_{714} = + 0.98 \pm 0.19$ and $\delta_{723} = - 3.4 \pm 0.1$ for the transitions in Te^{124} , $\delta_{754} = - 5.5^{+0.4}_{-0.3}$ for the $2^{+1} \rightarrow 2^{+}$ transition in Te^{126} , and $\delta_{491} = + 27^{+30}_{-9}$ for the $2^{+1} \rightarrow 2^{+}$ transition in Xe^{126} . There is an indication that the mixing ratios for the $2^{+1} \rightarrow 2^{+}$ transitions in even-even tellurium isotopes are negative and the mixing ratios of similar transitions in even-even Xe isotopes have positive values.

La 96

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

I. INTRODUCTION

The structure and the nature of the higher excited states of the even-even "spherical" nuclei in the mass region $100 \leq A \leq 150$ are not well understood. The first few excited states of these nuclei are interpreted as vibrational levels. However, the existence of cross-over transitions from the two-phonon 2^{+1} state to the ground state and the observation of finite quadrupole moments of the 2^{+} one - phonon states in many of these nuclei indicate that a description of these states in terms of harmonic oscillations of a spherical nuclear surface is not adequate. Furthermore, the existence of appreciable M1 components in the $2^{+1} \rightarrow 2^{+}$ transitions cannot be understood on the basis of a purely vibrational model. Clearly a more detailed microscopic description of the excited states of spherical nuclei is required before the relative strength of the M1 radiation components in gamma transitions of spherical nuclei can be understood. The interplay between the vibrational modes and the two-quasiparticle states is probably of major importance and should be considered in detail. Theoretical work in the region of intermediate nuclei is in progress¹. In the meantime, the present experimental work was undertaken in order to provide accurate experimental data on E2/M1 mixing ratios in order to test such calculations if and when they become available. In an earlier paper the E2/M1 multipole mixing ratios of the gamma transitions in the "spherical" Cd¹¹⁰ nucleus have been discussed². The present work describes an investigation of the E2/M1 multipole mixing ratios in the "spherical" nuclei Te¹²⁴, Te¹²⁶ and Xe¹²⁶.

II. THE LEVEL STRUCTURE OF THE Te^{124} , Te^{126} AND Xe^{126} NUCLEI

The levels of Te^{124} have recently been reinvestigated by Auer et al.³ and Meyer et al.⁴ from the decay of Sb^{124} and by Ragaini et al.⁵ from the decay of I^{124} . The lower excited states of Te^{124} (Fig. 1) exhibit a "vibrational" pattern which is expected for this "spherical" nucleus (proton number 52 and neutron number differing by only 10 from a closed neutron shell). The measurements of Warner and Draper⁶ show however, that the 4^+ level at 1249 keV could also be considered as a member of the ground-state quasi-rotational band. The E2/M1 mixing ratio of the 723 keV transition which proceeds between the two-phonon 2^{+1} and the one-phonon 2^+ level (in the vibrational picture), has been measured with conflicting results⁷⁻¹². The three lowest excited states of Te^{126} show clearly a vibrational character. Little is known about the higher excited states which have been investigated through nuclear reactions.

The level structure of Te^{126} shows similar interesting features. Whereas the lowest three excited states seem to conform to the vibrational picture, the higher excited states seem to be members of a rotational band up to 8^+ as found in $(\alpha, 2n)$ reactions¹³.

In the β - decay of I^{126} only the "vibrational" type levels of Te^{126} and Xe^{126} are populated. The decay scheme of I^{126} is shown in Fig. 2. The energy and intensity values are given as determined in the present investigation. They are in excellent agreement with the values of Singh and Taylor¹⁴. A weak gamma transition of (2046 ± 2) keV was observed in this laboratory which seems to correspond to the 2044.4 keV transition as reported in reference 14. The existence of a 1377.7 keV

transition¹⁴ could not be confirmed because of the interfering 1376 keV transition due to the I¹²⁴ impurity.

III. EXPERIMENTAL PROCEDURE

The directional correlation measurements were performed using both NaI(Tl) - Ge(Li) and Ge(Li) - Ge(Li) detector combinations. The former employed a 3 x 3 inch NaI(Tl) scintillator in conjunction with a coaxial Ge(Li) detector (ORTEC, 25 cc). A conventional (fast - slow) coincidence apparatus was employed having a resolving time of approximately 35 nsec. The Ge(Li) - Ge(Li) apparatus consisted of two coaxial detectors (ORTEC, 30 cc); the coincidence analysis was performed using a time-to-amplitude-converter system having an effective resolving time of approximately 50 nsec.

Radioactive Sb¹²⁴ was obtained from New England Nuclear Corporation in HCl solution. The solution was concentrated as needed and the liquid was placed inside a cylindrical glass ampoule 1.5 mm in diameter. The length of the liquid column varied from 5 - 8 mm.

The I¹²⁶ activity was produced as a result of (α , n) reactions on naturally-occurring antimony (57% Sb¹²¹, 43% Sb¹²³). Spectroscopically pure Sb metal powder was bombarded with 25 MeV alpha particles at the Argonne National Laboratory 60 - inch cyclotron. The irradiated powder was packed into an ampoule 3 mm in diameter and 10 mm in length. A considerable amount of I¹²⁴ activity resulted from this method of preparation, but the high resolution of the Ge(Li) detectors made it possible to easily correct for any effects on the measurement of the gamma rays from I¹²⁴.

Gamma ray spectra observed with Ge(Li) detectors from the Sb^{124} and I^{126} sources are shown in Figs. 3 and 4, respectively. The use of the solid-state detectors made possible a separation of the 723 keV transition from the 709 keV and 714 keV transitions in the Sb^{124} . A detail of this portion of the gamma spectrum is shown in Fig. 5. The separation of these lines is significant in attempts to measure the 723 keV-603 keV directional correlation, since both the 709 keV and 714 keV transitions are in coincidence with the 603 keV transition. In addition, as shown in Fig. 4, the 491 keV transition in I^{126} was resolved from the 511 keV positron-annihilation peak.

The various directional correlations were measured by determining the coincidence counting rates at seven different angles between 90° and 270° in the case of the NaI(Tl) - Ge(Li) measurements, and at nine different angles in the case of the Ge(Li) - Ge(Li) measurements. The counting rates were corrected for chance coincidences and for effects of competing cascades, and the directional correlation coefficients were extracted by fitting the measured counting rates to a function of the form:

$$W(\theta) = A'_0 + A'_{22} P_2(\cos \theta) + A'_{44} P_4(\cos \theta)$$

where

$$A'_{kk} = A_{kk} Q_{kk} G_{kk}$$

The correlation coefficients A_{kk} are defined in terms of the mixing ratios δ of Krane and Steffen². The geometrical correction factors

Q_{kk} were applied to correct for the finite solid angles of the detectors¹⁵. The perturbation factors G_{kk} were assumed to be equal to unity, the short lifetimes of the states involved in this measurement making external perturbations negligible.

IV RESULTS

In cases where it was feasible, directional correlations were measured for a given cascade using both NaI(Tl) - Ge(Li) and Ge(Li) - Ge(Li) configurations, in order to obtain a consistent set of results.

A summary of the present investigation of the Te^{124} gamma rays is given Table I. Results of some of the more recent measurements by other authors^{are} presented for comparison. Good agreement was obtained between the NaI(Tl) - Ge(Li) and Ge(Li) - Ge(Li) results of the present work in all cases. The low efficiency of the Ge(Li) detector for high-energy gamma rays made measurement of the 2091 keV - 603 keV directional correlation impossible for the Ge(Li) - Ge(Li) configuration. The resolution necessary for cascades involving the 709 keV and 714 keV transitions necessitated the use of the Ge(Li) - Ge(Li) system only.

As a check on the results involving the 709 keV and 714 keV transitions, the 1-3 directional correlation was measured between the 603 keV transition and the 709 keV-714 keV doublet. The results of this measurement were (with A_{00} normalized to unity):

$$A_{22} = 0.070 \pm 0.009$$

$$A_{44} = 0.052 \pm 0.015$$

Assuming the composition of the doublet to be 38% of the 709 keV and 62% of the 714 keV transitions, the directional correlation coefficients calculated on the basis of the results presented in Table I would be $A_{22} = 0.102$ and $A_{44} = -0.009$. The discrepancy between the calculated and measured values may be accounted for by considering the window accepting the doublet to include also a small amount (15%) of the 723 keV transition, as well as a portion (25%) due to the Compton background events primarily from the 1691 keV transition. It should be noted that neither of these corrections are necessary for the analyses of the 714 keV - 723 keV and 709 keV - 646 keV correlations.

Results for the directional correlation coefficients obtained from measurements on the I^{126} decay are shown in Table II. The large anisotropy characteristic of the 511 keV positron annihilation radiation caused significant problems in the measurements of the 491 keV - 389 keV directional correlation using the NaI(Tl) detector. Even with the NaI detector accepting the 389 keV transition (since the 491 keV transition cannot be resolved from the 511 keV line), the wide window widths necessary for the NaI detector resulted in the acceptance of a substantial amount of Compton background from the 511 keV radiation, with its corresponding large anisotropy. Thus it was necessary to use the Ge(Li) - Ge(Li) combination to measure this particular correlation.

To investigate the effects of the Compton background due to the positron annihilation radiation, the single-channel analyzer windows of the Ge(Li) detectors accepting the 389 keV and 491 keV peaks were moved so as to accept only the Compton background in an energy region either

above or below the peak. Investigations made with various combinations of window settings yielded a set of results which was reasonably consistent, particularly in view of the large statistical uncertainty due to the low counting rates. The directional correlation involving the Compton background showed the expected sharp peak at 180° . Such a correlation does not lend itself well to a Legendre polynomial analysis, and hence the measured coincidence counting rates of the 491 keV - 389 keV correlation were corrected for these effects before the correlation coefficients were extracted.

No such problems arose in the measurement of the 754 keV - 667 keV correlation, since both gamma rays lie above 511 keV, and hence this measurement could be performed using both detector combinations.

The mixing ratio for electric quadrupole-magnetic dipole admixture is defined in terms of the electromagnetic field multipole operators $\vec{A}_L(\pi)$ as²

$$\delta(\gamma_n) = \frac{\langle I_{n+1} \parallel \vec{j}_N \vec{A}_2^{(E)} \parallel I_n \rangle}{\langle I_{n+1} \parallel \vec{j}_N \vec{A}_1^{(M)} \parallel I_n \rangle}$$

or, in terms of Bohr-Mottelson²⁰ multipole matrix elements, as

$$\begin{aligned} \delta(\gamma_n) &= k_n \frac{\sqrt{3}}{10} \frac{\langle I_{n+1} \parallel \mathcal{M}(E2) \parallel I_n \rangle}{\langle I_{n+1} \parallel \mathcal{M}(M1) \parallel I_n \rangle} \\ &= k_n \frac{\sqrt{3}}{10} \Delta(\gamma_n) \end{aligned}$$

where the latter equation is evaluated in natural units ($\hbar = m_e = c = 1$).

In Table III are presented the mixing ratios $\delta(\gamma_n)$ and $\Delta(\gamma_n)$ derived

from the directional correlation coefficients given in Tables I and II.

V. DISCUSSION

(a) $\text{Sb}^{124} \rightarrow \text{Te}^{124}$

The results of the present investigation for the 646 keV - 603 keV cascade are consistent with the $4^+ - 2^+ - 0^+$ spin assignments, with results overlapping the expected values $A_{22} = 0.102$, $A_{44} = 0.009$. It is interesting to note, however, that there is some disagreement regarding the spin assignment of the 1249 keV level. Although the 4^+ assignment would certainly be preferred on the basis of the systematics of two-phonon levels in "vibrational" even-even nuclei, the directional correlation measurement alone cannot distinguish between an $I = 4$ assignment, with the 646 keV transition being pure E2, and $I = 3$ assignment, with the 646 keV transition being mixed E2/M1 having $\delta \approx 0.25$. Values of spin of the 1249 keV level lower than 3 and higher than 4 are excluded on the basis of the decay characteristics. The evidence from measurements of the K - conversion coefficients is contradictory; results based on measurements of conversion coefficients from the decay of Sb^{124} indicate a pure E2 character for the 646 keV transition^{5,21}, but the value obtained⁵ from the decay of I^{124} to Te^{124} is in agreement with M1 character for this radiation. In addition, the results of Dorikens-Vanpraet et al.¹¹ based on γ - γ measurements indicate a spin assignment of $I = 3$ for the 1249 keV level. At the present time however, it must be assumed that the $I = 4$ assignment is correct, on the basis of the systematics of even-even nuclei in this mass region.

The absence of direct feeding to the 1249 keV level from the $I^{124} (I=2^-)$ decay strongly supports this conclusion.

The results of various recent measurements of the mixing ratio of the 723 keV transition are in reasonably good agreement with the present results. Measurement of $\beta - \gamma$ directional correlations by Raghavan, Grabowski, and Steffen⁷ indicated the range of values $-2.8 \leq \delta(723) \leq -1.0$. Measurements of $\beta - \gamma - \gamma$ directional correlations by Glaubman and Oberholtzer¹⁰ have yielded $\delta(723) = -4.1 \pm 0.6$, and the $\gamma - \gamma$ measurements of Dorikens-Vanpraet et al.¹¹ and Stelson¹² have yielded $\delta(723) = -3.4 \pm 0.6$, respectively. While this latter result agrees precisely with the present value, the recent result, $\delta(723) = -7.5 \pm 2.0$, obtained in the experiment with oriented Sb^{124} nuclei by Sites and Steyert²² is at variance with it.


The results of the present work for the 1691 keV-603 keV and 2091 keV - 603 keV directional correlations are in agreement with the results for $3^- - 2^+ - 0^+$ cascades, with both the 1691 keV and 2091 keV transitions being of essentially E1 character ($|\delta| < 2 \times 10^{-2}$). The K-convention coefficients measurements confirm almost pure E1 character of these transitions^{5,21}.

The spin assignments for the 1958 keV and 2039 keV levels have been made on the basis of decay systematics⁴, rather than on the basis of previous directional correlation measurements. Considering the excitation energy only one is tempted to classify these levels as three phonon excitation (in the vibrational picture). The branching ratios of these states to the one-and two-phonon levels indicate some collective vibrational nature, with the transitions to the two-phonon levels

enhanced somewhat over those to the one-phonon level. It is to be expected that quasiparticle effects may be present at these high energies; such effects would tend to reduce the collective nature of the levels. Since the quasiparticle admixtures in these levels would be expected to be considerably greater than that in the 1326 keV level, it would be expected that the nature of the 709 keV and 714 keV transitions would be considerably less collective than that of the 723 keV transition. Based on the spin assignment of 2^+ and 4^+ for the 2039 keV and 1958 keV levels, this interpretation would be consistent with the E2/M1 mixing ratios measured in the present work, i.e. with both transitions showing substantial M1 admixtures.

It should be noted that on the basis of the directional correlation measurements involving the 709 keV and 714 keV transitions, the spin assignment of 3^+ for either the 1958 keV or the 2039 keV levels may not be excluded. An assignment of 3^+ for the 1958 keV level would be consistent with the directional correlation for a $3^+ - 4^+ - 2^+$ cascade with $\delta(709) \approx -0.45$, while a 3^+ assignment for the 2039 keV level would result in $\delta(714) \approx -0.3$ for the $3^+ - 2^+ - 2^+$ directional correlation. Using our value for the δ_{723} and Sliv's conversion coefficients²³ one can calculate the K-conversion coefficient for the 723 keV transition in Te^{124} as $\alpha_K(723) = 2.7 \times 10^{-3}$. The agreement between this calculated value and the measurements reported in Ref. 5 and 21 is excellent. The experimentally measured K-conversion coefficient for the 714 keV gamma transition^{5,21} when used with our $\delta(714)$ measurement strongly supports the 2^+ assignment for the 2039 keV level. However, the basis

of the measured conversion coefficients and our mixing ratio data it is not possible to decide between the 3^+ and 4^+ spin-parity assignment for the 1958 keV state in Te^{124} .



The results of the present investigation for the $2^{+1} - 2^+ - 0^+$ cascade in Te^{126} show agreement between the NaI(Tl) - Ge(Li) and Ge(Li) - Ge(Li) results; in addition the agreement with previous values as measured by Sakai et al. is good¹⁹. However, in previous investigations^{18,19} of Xe^{126} using NaI(Tl) detectors unambiguous values for the $2^{+1} - 2^+$ mixing ratio were not obtained due to the problems associated with the positron annihilation radiation. Such investigations have generally indicated large values of δ , results which are not inconsistent with those of the present work.

It is of interest to note the variations in values of δ in this mass region. The $2^{+1} - 2^+$ mixing ratio in Te^{122} has been measured¹⁸ to be $\delta = -3.7 \pm 0.3$. This latter result, together with those of the present investigation for Te^{124} and Te^{126} indicate that the values for the mixing ratios in all three of these even-even tellurium isotopes are negative. However, results for the $2^{+1} - 2^+$ transitions in Xe^{128} ($\delta = +6.4 \pm 1.5$)¹⁸ and for Xe^{132} ($\delta \approx +5$)²⁴ indicate a trend toward positive mixing ratios in even-even isotopes. At present, no explanation can be offered for this contrast. Tamura and Yoshida²⁵ have computed that the $(g\ 7/2)^2$ two quasi-proton state occurs at 1.36 MeV in Te and at 1.60 MeV in Xe; it thus would be expected to have a substantial influence on the properties of the 2^{+1} state in $\text{Te}(E_{2^{+1}} \approx 1.3$

MeV), but not nearly as much influence in the case of the 2^{+1} state of even-even nuclides of Xe ($E_{2^{+1}} < 1$ MeV in Xe^{126,128}; $E_{2^{+1}} \approx 1.3$ MeV in Xe¹³²).

* Work supported by the U.S. Atomic Energy Commission under contract AT(11-1) - 1746 (Chicago Operations Office).

‡ A preliminary report of this investigation has been presented at the International Conference on Angular Correlations in Delft (The Netherlands), August 17-21, 1970.

† Present address: Los Alamos Scientific Laboratory, Los Alamos, New Mexico.

1. B. Sørensen, Nucl. Phys. A142, 411 (1970); K. Kumar (private communication).
2. K. S. Krane and R. M. Steffen, Phys. Rev. C2, 724 (1970).
3. I. P. Auer, J. J. Reidy, and M. L. Wiedenbeck, Nucl. Phys. A124, 199 (1969).
4. R. P. Meyer, W. B. Walters and R. C. Ragaini, Nucl. Phys. A127, 595 (1969).
5. R. C. Ragaini, W. B. Walters, A. A. Noyes and R. A. Meyer, Phys. Rev. 187, 1721 (1969).
6. R. A. Warner and J. E. Draper, Phys. Rev. C1, 1069 (1970).
7. R. S. Raghavan, Z. W. Grabowski and R. M. Steffen, Phys. Rev. 139, 1 (1965).
8. H. Paul, Phys. Rev. 121, 1175 (1961).
9. T. Lindqvist and I. Marklund, Nucl. Phys. 4, 189 (1957).
10. M. J. Glaubman and J. D. Oberholtzer, Phys. Rev. 135, B1313 (1964).
11. L. Dorikens-Vanpraet, J. Demuyneck and M. Dorikens, Nucl. Phys. 73, 539 (1965).
12. P. H. Stelson, Phys. Rev. 157, 1098 (1967).
13. H. Morinaga and N. L. Lark, Nucl. Phys. 67, 315 (1965).
14. B. Singh and H. W. Taylor, Nucl. Phys. A147, 12 (1970).
15. D. C. Camp and A. L. Van Lehn, Nucl. Instr. and Math. 76, 192 (1969);
M. J. L. Yates, Alpha-Beta-and Gamma-Ray Spectroscopy (Ed. K. Sigbahn,
North-Holland Publishing Co., Amsterdam, 1965) p. 1691.
16. M. R. Meder and F. E. Durham, Nucl. Phys. A97, 78 (1967).
17. C. Weitkamp, Nucl. Phys. 43, 57 (1963).

18. I. Asplund, L. G. Strömberg and T. Wiedling, Ark. Fysik 18, 65 (1961).
19. M. Sakai, H. Ikegami, T. Yamazaki and K. Sugiyama, J. Phys. Soc. Japan 14, 983 (1959).
20. A. Bohr and B. R. Mottelson, Nuclear Structure, Vol. I (W. A. Benjamin, Inc., New York, 1969), p. 381.
21. E. P. Grigorev, A. V. Zolotavin, V. O. Sergeev and M. I. Sovtsov, Izv. Akad. Nauk SSSR, Ser. Fiz. 32, 773 (1968).
22. J. R. Sites and W. A. Steyert, Nucl. Phys. (To be published).
23. L. A. Sliv and M. I. Band in Alpha-Beta and Gamma-Ray Spectroscopy (Ed. K. Siegbahn, North-Holland Publishing Co., Amsterdam, 1965) p. 1639.
24. K. S. Krane and R. M. Steffen, Phys. Rev. C3, (1970).
25. T. Tamura and H. Yoshida, Nucl. Phys. 30, 579 (1962).

TABLE I

Results for the Directional Correlation Coefficients of the Te¹²⁴ Gamma Rays

Cascade (E in KeV)	Spin Sequence	A ₂₂	A ₄₄	Reference
646 - 603	4 ⁺ -2 ⁺ -0 ⁺	0.12 ± 0.04	0.0 ± 0.05	a
		0.099 ± 0.005	0.007 ± 0.008	Present, NaI(Tl)-Ge(Li)
		0.121 ± 0.025	0.010 ± 0.030	Present, Ge(Li)-Ge(Li)
714 - 723	2 ⁺ -2 ⁺ -2 ⁺	0.220 ± 0.015	-0.026 ± 0.030	Present, Ge(Li)-Ge(Li)
709 - 646	4 ⁺ -4 ⁺ -2 ⁺	0.187 ± 0.017	0.011 ± 0.033	Present, Ge(Li)-Ge(Li)
723 - 603	2 ⁺ -2 ⁺ -0 ⁺	0.14 ± 0.03	0.26 ± 0.04	a
		0.152 ± 0.008	0.304 ± 0.012	Present, NaI(Tl)-Ge(Li)
		0.130 ± 0.015	0.295 ± 0.020	Present, Ge(Li)-Ge(Li)
1691 - 603	3 ⁻ -2 ⁺ -0 ⁺	-0.063 ± 0.004	-0.009 ± 0.006	b
		-0.072 ± 0.005	-0.007 ± 0.005	c
		-0.099 ± 0.026	0.03 ± 0.03	a
		-0.060 ± 0.006	0.000 ± 0.008	Present, NaI(Tl)-Ge(Li)
		-0.064 ± 0.020	-0.005 ± 0.025	Present, Ge(Li)-Ge(Li)
2091 - 603	3 ⁻ -2 ⁺ -0 ⁺	-0.058 ± 0.005	0.001 ± 0.005	d
		-0.052 ± 0.012	0.037 ± 0.018	b
		-0.069 ± 0.015	0.004 ± 0.020	Present, NaI(Tl)-Ge(Li)

a Reference 12

b Reference 11

c Reference 16

d Reference 17

TABLE II

Results for the Directional Correlation Coefficients of Gamma Rays from the Decay of I^{126} ✓

Isotope	Cascade (E in keV)	Spin Sequence	A_{22}	A_{44}	Reference
Te^{126}	754 - 667	$2^+ - 2^+ - 0^+$	0.01 ± 0.02	0.35 ± 0.04	a
			0.059 ± 0.009	0.268 ± 0.014	b
			0.062 ± 0.009	0.330 ± 0.015	Present, NaI(Tl)-Ge(Li)
			0.052 ± 0.028	0.291 ± 0.035	Present, Ge(Li)-Ge(Li)
Xe^{126}	491 - 389	$2^+ - 2^+ - 0^+$	-0.03	0.29	a
			-0.071 ± 0.025	0.294 ± 0.041	b
			-0.103 ± 0.012	$\pm .341 \pm 0.030$	Present, Ge(Li)-Ge(Li)

a Reference 18

b Reference 19

TABLE III

E2/M1 Mixing Ratios of Gamma Transitions in Te^{124} , Te^{126} , and Xe^{126}

Isotope	Transition E(keV)	Spins	δ (γ)	Δ (γ) (natural units)
Te^{124}	723	$2^{+1}-2^{+}$	- 3.4 ± 0.1	- 13.9 ± 0.4
	714	$2^{+2}-2^{+1}$	+ 0.98 ± 0.19	+ 4.1 ± 1.3
	709	$4^{+1}-4^{+}$	+ $0.04^{+0.03}_{-0.05}$	+ $0.17^{+0.13}_{-0.21}$
Te^{126}	754	$2^{+1}-2^{+}$	- $5.5^{+0.4}_{-0.3}$	- $21.6^{+1.6}_{-1.2}$
Xe^{126}	491	$2^{+1}-2^{+}$	+ 27^{+30}_{-9}	+ 162^{+180}_{-54}

FIGURE CAPTIONS

Fig. 1. The decay of Sb^{124} (60d.) to levels of Te^{124} (Ref. 3).

Fig. 2. The decay of I^{126} (13d.) to levels of Te^{126} and Xe^{126} .

Fig. 3. Gamma-ray spectrum from the decay of Sb^{124} recorded with 30 cc. Ge(Li) detector.

Fig. 4. Gamma-ray spectrum from the decay of I^{126} recorded with 30 cc. Ge(Li) detector.

Fig. 5. Ge(Li) gamma-ray spectrum of Sb^{124} in the 700 keV energy region.







Fig 1 ; Z.W GRABOWSKI et al.

Fig. 2: Z.W. Graessner et al.








Fig 5
ZUB
Coannuli at all

