SUMMARY OF
SIGNIFICANT
RESULTS IN—

Mineral resources

Water resources

Engineering geology
and hydrology

Regional geology

Principles and processes

Laboratory and
field methods

Geographic surveys
and mapping

Management of resources
on public lands

Investigations in
other countries

OF—

Investigations in
progress

Reports published
fiscal year 1965

Operating agencies

Geological Survey offices

2nd SET
ENGINEERING LIBRARY
California Institute of Technology

GEOLOGICAL SURVEY RESEARCH 1965

geol

Chapter A

ENGINEERING LIBRARY
California Institute of Technology

metadc304147

GEOLOGICAL SURVEY RESEARCH 1965

WILLIAM T. PECORA, *Director*

GEOLOGICAL SURVEY PROFESSIONAL PAPER 525

Significant results of investigations for fiscal year 1965, accompanied by short papers in the fields of geology, hydrology, and related sciences. Published separately as Chapters A, B, C, and D.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON: 1965

UNITED STATES DEPARTMENT OF THE INTERIOR

STEWART L. UDALL, Secretary

GEOLOGICAL SURVEY

William T. Pecora, Director

GEOLOGICAL SURVEY RESEARCH 1965

Chapter A

GEOLOGICAL SURVEY PROFESSIONAL PAPER 525-A

A summary of recent significant scientific and economic results accompanied by a list of publications released in fiscal 1965, a list of geologic and hydrologic investigations in progress, and a report on the status of topographic mapping.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON: 1965

UNITED STATES DEPARTMENT OF THE INTERIOR

STEWART L. UDALL, Secretary

GEOLOGICAL SURVEY

William T. Pecora, Director

FOREWORD

"Geological Survey Research 1965" is the sixth annual review of the economic and scientific work of the U.S. Geological Survey. As in previous years the purpose of the volume is to make available promptly to the public the highlights of Survey investigations. This year the volume consists of 4 chapters (A through D) of Professional Paper 525. Chapter A contains a summary of significant results, and the remaining chapters are made up of collections of short technical papers.

Many of the results summarized in Chapter A are discussed in greater detail in the short papers or in reports listed in "Publications in Fiscal Year 1965," beginning on page A259. The tables of contents for chapters B through D are listed on pages A253–A258.

Numerous Federal, State, county, and municipal agencies listed on pages A205–A210 cooperated financially with the Geological Survey during fiscal 1965 and have contributed significantly to the results reported here. They are identified where appropriate in the short technical papers, but generally are not identified in the brief statements in chapter A.

Many individuals on the staff of the Geological Survey have contributed to "Geological Survey Research 1965." Reference is made to only a few. George Phair, Geologic Division, was responsible for organizing and assembling chapter A and for critical review of papers in chapters B–D, assisted by Frank W. Trainer, Water Resources Division. Marston S. Chase, Publications Division, was in charge of production aspects of the series, assisted by Jesse R. Upperco in technical editing, and William H. Elliott and James R. Hamilton in planning and preparing illustrations.

The volume for next year, "Geological Survey Research 1966," will be published as chapters of Professional Paper 550. Previous volumes are listed below, with their series designations.

Geological Survey Research 1960—Prof. Paper 400
Geological Survey Research 1961—Prof. Paper 424
Geological Survey Research 1962—Prof. Paper 450
Geological Survey Research 1963—Prof. Paper 475
Geological Survey Research 1964—Prof. Paper 501

WILLIAM T. PECORA,
Director.

CONTENTS

	Page		Page
Foreword	iii	Investigations of natural resources—Continued	
Investigations of natural resources	A1	Water resources—Continued	
Mineral resources	1	Midcontinent area—Continued	
Resource compilation	2	Kentucky	A29
Heavy metals	3	Tennessee	30
Iron	3	Alabama	30
Manganese	4	Mississippi	30
Molybdenum	4	Louisiana	31
Copper	4	Rocky Mountain area	32
Lead-zinc	5	Interstate studies	32
Mercury	5	Montana	32
Antimony	6	North Dakota	33
Gold and silver	6	Wyoming	33
Laboratory studies of sulfide ore minerals	7	South Dakota	34
Light metals and industrial minerals	8	Nebraska	34
Evaporites	8	Utah	34
Phosphate	9	Colorado	35
Clay and lightweight aggregate	10	Kansas	35
Beryllium	10	Arizona	36
Zeolites	11	New Mexico	36
Radioactive minerals	12	Oklahoma	36
Uranium	12	Texas	37
Thorium	13	Pacific coast area	37
Minor elements	14	Alaska	38
Organic fuels	14	Pacific Northwest	38
Coal	14	Nevada	39
Petroleum and natural gas	15	California	40
Oil shale and humate deposits	15	Hawaii and Guam	41
Techniques of mineral exploration	15	Management of natural resources on the public	
Water resources	17	land	42
Office of Water-Data Coordination	18	Classification of mineral lands	42
Atlantic coast area	18	Waterpower classification	43
Interstate studies	19	Supervision of prospecting, development, and	
New England	21	recovery of minerals	43
New York	21	Geology and hydrology applied to engineering and	
New Jersey	22	public health	45
Maryland	22	Investigations related to nuclear energy	45
West Virginia	22	Site evaluation	45
South Carolina	23	Plowshare program	46
Georgia	23	Geologic and hydrologic effects of nuclear	
Florida	23	explosions	46
Caribbean region	24	Basic geologic studies	48
Midcontinent area	25	Disposal of radioactive wastes	50
Interstate studies	25	Transport of radionuclides	50
Minnesota	26	Studies at nuclear-energy facilities	51
Wisconsin	26	Fate of radioactive liquid effluent	51
Michigan	27	Flow of air in tuff	52
Ohio	27	Water-contamination studies	52
Indiana	28	Detergents and pesticides in water supplies	53
Missouri	28	Laboratory studies	53
Arkansas	28	Field studies	53

	Page		Page
Geology and hydrology applied to engineering and public health—Continued		Regional geology—Continued	
Water-contamination studies—Continued		Basin and Range region—Continued	
Acid mine water	A53	Emplacement and age of intrusive rocks	A93
Industrial effects	54	Cenozoic volcanism	94
Brine and saline water	54	Geochemical exploration	94
Distribution of minor elements as related to public health	56	Geophysical studies	95
Problems in engineering geology and hydrology	57	Columbia Plateau and Snake River Plain	95
Land subsidence	58	Oregon	95
Evaporation suppression	58	Idaho	95
Floods	59	Pacific coast region	96
Outstanding floods of 1964-65	59	Washington	96
Flood frequency	60	Oregon	97
Flood mapping	60	California	97
Effects of the 1964 Alaska earthquake	61	Klamath Mountains and Northern Coast Ranges	97
Investigations of geologic effects	61	Southern Coast Ranges	98
Investigations of hydrologic effects	64	Sierra Nevada	99
Regional geology	67	San Joaquin Valley	99
Intermediate-scale geologic maps	68	Alaska	100
Maps of large regions	68	Northern Alaska	100
Coastal plains	69	West-central Alaska	102
Atlantic Coastal Plain	69	East-central Alaska	103
Gulf Coastal Plain and Mississippi embayment	71	Southwestern Alaska	103
New England and eastern New York	72	Southern Alaska	104
Eastern New York	72	Southeastern Alaska	104
Rhode Island	72	Puerto Rico	105
Massachusetts	72	Antarctica	106
Connecticut	74	Geologic and hydrologic investigations in other countries	109
Maine	74	Summary by countries	111
Appalachian region	75	Afghanistan	111
Structural and stratigraphic studies	75	Bolivia	111
Geophysical studies	78	Brazil	111
Eastern plateaus	78	Chile	112
Pennsylvania	78	Colombia	112
Kentucky	79	Ghana	112
Tennessee	81	Indonesia	113
Shield area and Upper Mississippi Valley	81	Nigeria	113
Michigan	81	Pakistan	113
Wisconsin	81	Panama	114
Minnesota	82	Philippine Islands	114
Nebraska	82	Saudi Arabia	115
Interior Highlands and eastern plains	82	Thailand	116
Northern Rocky Mountains and plains	83	International cooperation within the region of the Central Treaty Organization	116
Washington	83	Cooperative short-term topical investigations abroad	117
Idaho	83	Australia	117
Montana	84	Greenland	117
Wyoming	85	Iceland	117
North and South Dakota	86	Japan	117
Southern Rocky Mountains and plains	86	Leeward Islands	118
Geology of Precambrian rocks	86	Sino-Soviet geologic studies	118
Stratigraphy and structural geology	87	Study of Irazú Volcano, Costa Rica	118
Geology of volcanic and hypabyssal intrusive rocks	88	Investigations of principles and processes	121
Geophysical investigations	89	Paleontology	121
Quaternary geology	89	Foraminifera	121
Colorado Plateau	90	Lower invertebrates	122
Geologic studies	90	Mollusca	123
Geophysical studies	91	Arthropoda	124
Basin and Range region	92	Vertebrates	125
Stratigraphy and structural geology	92	Paleobotany	125
Nevada and eastern California	92	Marine geology and hydrology	127
Utah	93	Atlantic Continental Shelf and Slope	127
Arizona	93		

	Page		Page
Investigations of principles and processes—Continued		Investigations of principles and processes—Continued	
Marine geology and hydrology—Continued		Isotope and nuclear studies	A161
Marine geologic environment and processes	A128	Geochronology	161
Fresh-water-salt-water mixing	128	Radioactive disequilibrium	163
Water movement in estuaries	129	Stable-isotope investigations	164
Biohydrologic studies of estuaries	129	Lead	164
Oceanic islands	129	Oxygen and tritium	164
Astrogeology	130	Hydraulic and hydrologic studies	165
Earth-based lunar investigations	130	Surface water	165
Theoretical analyses of lunar phenomena	132	Ground water	168
Instrumentation and techniques for		Relations between ground water and surface	
land-based lunar study	132	water	172
Terrestrial and experimental impact and		Soil moisture and evapotranspiration	173
cratering phenomena	133	Sedimentation	175
Field studies of impact structures	133	Erosion	175
Laboratory studies of impact-related		Transportation	175
phenomena	133	Variability of sediment loads in rivers	176
Cosmochemistry and petrology	134	Deposition	176
Unmanned lunar exploration studies	136	Limnology	177
Ranger television investigations	136	Geomorphology	177
Surveyor television investigations	136	Plant ecology	178
Lunar Orbiter investigations	137	Glaciology	180
Lunar terrain analysis	137	Permafrost	181
Studies related to manned lunar exploration	137	Analytical methods	181
Astronaut training	137	Analytical chemistry	181
Investigations in support of manned lunar		Optical spectroscopy	183
exploration	138	X-ray fluorescence analysis	183
Geophysical investigations	139	Analysis of water	184
Studies of the crust and upper mantle	139	Hydrologic measurements and instrumentation	186
Regional seismic-refraction studies	139	Topographic surveys and mapping	189
Cooperative experiments in seismology	141	Mapping accomplishments	189
Regional gravity studies	142	Mapping in Antarctica	197
Geothermal investigations	142	Research and development	200
Theoretical and experimental geophysics	143	Establishment of test site for mapping	
Rock magnetism	143	research	200
Physical properties of minerals and rocks	145	Field surveys	200
Aerial infrared surveys	146	Photogrammetry	201
Solid-state studies	146	National Atlas	203
Field studies of thermoluminescence	146	Cooperating agencies for fiscal year 1965	205
Laboratory studies of thermoluminescence	146	Federal agencies	205
Magnetic-susceptibility measurements	146	State, county, and municipal agencies	205
Geochemistry, mineralogy, and petrology	147	U.S. Geological Survey offices	211
Field studies in petrology and geochemistry	147	Main centers	211
Silicic plutonic rocks	147	Public inquiries offices	211
Mafic and ultramafic rocks	148	Selected field offices in the United States and	
Silicic volcanic rocks	149	Puerto Rico	211
Metamorphic rocks	149	Conservation Division	211
Thermal water	150	Geologic Division	212
Diagenetic rocks	150	Topographic Division	212
Mineralogic studies and crystal chemistry	151	Water Resources Division	213
Experimental geochemistry	153	Offices in other countries	215
Basic stability relationships	153	Geologic Division	215
Studies of magmatic processes	153	Water Resources Division	215
Studies of metamorphic processes	154	Investigations in progress in the Geologic, Water	
Solubilities and exchange reactions	154	Resources, and Conservation Divisions	217
Studies of fluid inclusions in minerals	155	How to order U.S. Geological Survey publications	251
Geochemical data	156	Contents of Geological Survey Research 1965,	
Geochemistry of water	157	Chapters B, C, and D	253
Studies of atmospheric precipitation	157	Publications in fiscal year 1965	259
Studies of streams and lakes	158	Indexes	323
Ground-water studies	158	Publications index	323
Chemical equilibrium studies	159	Subject index	353
Thermal waters	159	Investigator index	371
Investigations at the Hawaiian Volcano			
Observatory	159		

ILLUSTRATIONS

	Page
FIGURE 1. Index map of the United States, showing areal subdivisions used in discussion of water resources	A19
2. Percentage of total United States mineral production that comes from Federal land	42
3. Index map of the conterminous United States, showing boundaries of regions referred to in discussion of regional geology	67
4. Index map of Alaska, showing boundaries of regions referred to in discussion of Alaska geology	101
5. Index map of Antarctica	106
6, 7. Views of erupting Irazú Volcano	118, 119
8. Sketch map of Costa Rica and its volcanoes	119
9. Index map of the Moon, showing status of geologic mapping	130
10. Progress of 7½- and 15-minute quadrangle topographic mapping	190
11. Status of 7½- and 15-minute quadrangle topographic mapping	191
12. Status of revision of large-scale topographic mapping	192
13. Status of 1:250,000-scale topographic mapping	193
14. Status of State topographic maps	194
15. Status of 1:1,000,000-scale topographic mapping	196
16. Plan for optical aerial mapping photography of Antarctica	198
17. Index map of Antarctica, showing status of topographic mapping	199

GEOLOGICAL SURVEY RESEARCH 1965

INVESTIGATIONS OF NATURAL RESOURCES

The U.S. Geological Survey as created by Act of Congress in 1879 was charged with the "classification of the public lands," and with the "examination of the geological structure, mineral resources and products of the national domain." A scanning of the highlights of the Geological Survey's program reported here will make it plain that resource studies, broadly interpreted, provided the keynote and the impetus for the greater part of the Survey's research in 1965 as it has each year since the agency's inception.

Although categories necessarily overlap, the separate sections of this summary may be said to be concerned in varying degrees with the "*what?*" "*where?*" and "*why?*" surrounding the natural occurrences of the Nation's mineral and water resources. In essence this first section attempts to answer the question "*what* are the distinguishing characteristics and economic importance of individual occurrences of specific commodities?" The section entitled "Regional Geology" is concerned with the total geologic environment, and attempts to answer the question "*where* within that environment are useful resources likely to be found?" The section entitled "Investigations of Principles and Processes" brings modern field and laboratory tools to bear upon the genetic processes that gave birth to the rocks and minerals of the earth's crust, and in so doing concentrated ores, fuels, and waters; it attempts to answer the question "*why* did these useful concentrations form where they did?"

MINERAL RESOURCES

Minerals and fuels, the fundamental ingredients of the American industrial economy, have been prime targets of the U.S. Geological Survey's work from its establishment in 1879. Developments in 1965 pinpoint some of the urgent resource problems currently brought into sharp focus by the demands of

a growing population and economy, and illustrate some of the measures being taken to solve them. Among the problems are scarcities that are developing among a few key metals as domestic consumption outstrips production and depletes world supplies. Silver, mercury, and gold are prime examples; industrial consumption of gold (to say nothing of monetary needs) now exceeds domestic production. Appalachia, an area depressed in part by the consequences of an overspecialized exploitation of its resource base, represents another kind of resource problem. The establishment of the Wilderness system illustrates still another: the competition of a growing population for diverse resources on lands where the exploitation of one seems to prevent the use of others.

Discovery of new supplies constitutes one of the prime solutions to the scarcity problem, and the Survey's work contributes importantly in this area by providing knowledge and methods that aid in prospecting for concealed deposits. For example, in May the Newmont Mining Corp. opened a gold deposit at Carlin, Nev., found by drilling beneath a thrust plate to test favorable environments identified by R. J. Roberts.¹ It promises to be the second largest gold mine in the United States. New rapid methods of analysis for gold, silver, and mercury, developed respectively by H. W. Lakin and H. M. Nakagawa (p. C168-C171),² by Nakagawa and Lakin (p. C172-C175), and by W. W. Vaughn, W. G. Cramer, and W. N. Sharp (p. B151-B154) should aid in prospecting for deposits of these metals. Identification of silver-bearing black calcite (D. F.

¹ R. J. Roberts, 1957, Major mineral belts in Nevada [abs.]: 1957 Pacific Southwest Mineral Industry Conf., Reno, Nev., Abstracts, p. 4.

R. J. Roberts, P. E. Hotz, James Gilluly, and H. G. Ferguson, 1958, Paleozoic rocks in north-central Nevada: Am. Assoc. Petroleum Geologists Bull., v. 42, no. 12, p. 2813-2857.

R. J. Roberts, 1960, Alinement of mining districts in north-central Nevada: Art. 9 in U.S. Geol. Survey Prof. Paper 400-B, p. B17-B19.

² See "Contents of Geological Survey Research 1965, Chapters B, C, and D" (p. A253-A258). This report is listed in the contents for chapter C (p. A255-A256).

Hewett and others, 1-65)³ points to a kind of silver deposit that has not been much prospected previously and suggests that silver, like uranium, may be found to occur in significant quantities in other types of deposits besides those mined in the past.

As a part of the Appalachia program, the U.S. Geological Survey, in cooperation with the U.S. Bureau of Mines and various State agencies, has begun studies aimed ultimately at developing a complex of local mineral resources that may serve as a base for new industries within Appalachia. Following initial emphasis on compilation and analysis of available resource data, work will focus on field studies to develop leads for prospecting.

The Wilderness Act directs the Geological Survey and Bureau of Mines to undertake surveys of Wilderness Areas to determine their mineral values. Primitive and other areas to be considered for inclusion in the system are also to be examined. The purpose of these surveys is to provide an objective description of potential resources that others can use in recommendations and decisions concerning the classification of the areas. Work has begun in seven of the Primitive Areas.

Salient results of the mineral-resource studies in 1965 are summarized in the following pages.

RESOURCE COMPILATION

Current knowledge of the mineral and water resources of Colorado, Idaho, Nevada, and New Mexico has been summarized in four reports prepared jointly by the U.S. Geological Survey and various State agencies. These reports describe known mineral deposits, and in addition make available the syntheses of much recent work pointing toward areas favorable for further exploration.

Resources of Colorado

The report on Colorado (U.S. Geol. Survey, 12-64) points out that major metallic resources have come largely from a narrow 250-mile-long mineralized belt extending diagonally across the State from the La Plata mineral district northeastward to the Jamestown district at the front of the Rockies in Boulder County. Throughout its length the Colorado Mineral Belt is characterized by bodies of Tertiary intrusive igneous rocks, many of which are related to concentrations of ore deposits. Favorable source rocks and possible reservoirs of petroleum and natural gas are described, and areas are indicated that have not yet been adequately tested.

³ See "Publications in Fiscal Year 1965" (p. A259-A322). This report is the first citation for 1965 under entries for D. F. Hewett, and is listed on p. A285.

Resources of Idaho

The report on Idaho (U.S. Geol. Survey, 16-64) notes that the Idaho batholith clearly constitutes a tungsten province and thus offers a promising field for prospecting and a reasonable chance for new discoveries. Other metallic mineral deposits in the central part of the State, particularly the productive lead-silver deposits, have long been regarded as related to the Idaho batholith. There is, however, ample evidence that many of these deposits stem from an early Tertiary epoch of mineralization, that is in turn associated with intrusions of a later age than the batholith. The search for new deposits should be guided by these associations.

Resources of Nevada

Most of the metallic deposits in Nevada, the summary volume (U.S. Geol. Survey, 17-64) concludes, are related to intrusive rocks, generally granodiorite to quartz monzonite in composition. The mining districts are not randomly distributed; many of the deposits are aligned along belts that trend northwestward, east-west, and northeastward across the State. More than 95 percent of the past production of copper, lead, zinc, gold, and silver has come from these belts. The distribution of deposits along the belts is not accidental but reflects fundamental stratigraphic and structural controls. These belts should be the best places to look for new ore deposits.

Resources of New Mexico

In New Mexico a similar beltlike distribution of the major metallic deposits is reported (U.S. Geol. Survey, 7-65). This belt extends from the southwestern corner to the north-central part of the State. Favorable targets within this zone are described. Energy resources in New Mexico are very large, and there are excellent chances of discovering new oil and gas fields.

Mineral fuels of eastern Montana

The future growth potential of eastern Montana's economy is based on unusually abundant resources of mineral fuels, according to a joint study of the U.S. Geological Survey and the U.S. Bureau of Mines (U.S. Geol. Survey, 6-65). One-eighth of the coal resources of the Nation, some 215 billion tons, is in eastern Montana; of this amount, 128 billion tons are of subbituminous rank and 87 billion tons are lignite. About a third of the subbituminous coal is in beds more than 10 feet thick, and about 3 percent of it lies in beds less than 120 feet below the surface, most of which can be recovered by open-pit mining methods. Thermal power plants are the

largest consumer of this stripping coal. Eastern Montana produces significant quantities of petroleum and natural gas, and many areas outside the known fields still are relatively untested. Oil has been produced from 17 known formations and gas from several different zones. The present density of testing outside known oil fields is only about 1 well for every 55 square miles.

Domestic resources of mercury analyzed

Domestic mercury resources and economic trends of the mercury industry have been analyzed by E. H. Bailey and R. M. Smith (4-64). After a number of years of declining prices, most of the domestic producers have terminated exploration and closed most of the mines. Domestic production in 1963 provided less than a fourth of the national consumption. However, the price of mercury advanced dramatically from \$180 a flask in 1963, to more than \$500 a flask by the end of 1964. This study shows that potential resources exist and that with resumption of exploration, the discovery of new deposits and expansion of production can be expected.

Domestic resources of silver evaluated

An analysis of domestic silver resources and economic trends has been made by T. H. Kiilsgaard (1-64). Because silver occurs in a variety of ore deposits in association with other metals, an economic analysis of silver production also must consider the production of the other metals, particularly the base metals lead and zinc. Significant changes in demand for these associated metals have direct impact on the output of silver. The increasing demand for silver for industrial usage has stimulated the search for new deposits by new exploratory techniques (p. C172-C175).

World thorium resources

In reviewing geology and resources of thorium in the United States and the world, J. C. Olson and W. C. Overstreet (1-64) point out that if moderately rich resources of thorium in the United States, expressed as tons per square mile in areas of igneous and high-grade metamorphic rock, are representative of worldwide distribution, corresponding world resources should be at least 3.6 million tons.

HEAVY METALS

IRON

Northwestern Adirondack Mountains, N.Y.

In a comprehensive report on the ore deposits of the St. Lawrence County magnetite district, New

York, B. F. Leonard and A. F. Buddington (1-64) describe two principal types of hypogene iron oxide deposits: magnetite deposits in skarn or marble, and magnetite deposits, with or without hematite, in microcline granite-gneiss. The ore bodies are interpreted as high-temperature replacement deposits genetically related to a "younger granite" series that intruded metasedimentary rocks of the Grenville Series. Earlier the metasediments had been intruded by anorthosite, gabbro, and quartz syenite. Renewed exploration and study of this district during and following World War II led to the discovery, under private and Government auspices, of 23 promising new deposits, as compared with 8 that were known from outcrops at least as early as 1840. None had been discovered in the intervening century. Magnetic surveys led to this success; among the finds was the first deposit located in North America by the then-new airborne magnetometer.

Michigan

Recent ground magnetometer surveys by J. E. Gair in the Palmer district, Marquette County, Mich., delimited two areas of anomalously high magnetic readings caused by iron-formation that may be of economic importance. The larger of the two anomalies lies at the east end of the district in secs. 26 and 27, T. 47 N., R. 26 W. and is as much as 27,000 gammas above background. The total extent of the area registering 3,000 gammas or more above background is about 2,800 feet by 1,200 feet. A smaller anomaly is centered near the northwest corner of the village of Palmer at the west end of the district. Maximum readings over this anomaly are about 3,500 gammas above background.

W. C. Prinz reports that some of the Ironwood Iron-Formation in the western half of T. 47 N., R. 44 W., in the East Gogebic Range, Mich., contains sufficient iron to make it of potential economic importance. Results of his recent geologic and magnetic survey, combined with data from old diamond drill holes, show that the middle few hundred feet of the formation is magnetite-rich and contains more than 35 percent iron for a strike length of at least 3 miles. Beneficiation tests were run on one sample of this rock by the Institute of Mineral Research, Michigan Technological University, but a concentrate of acceptable grade could not be made because the magnetite is poorly liberated. However, this sample represents only the uppermost 35 feet of the magnetic unit; the beneficiation properties of the remainder of the unit are unknown because of the lack of exposures. Thin sections from one drill hole

that penetrated the lower part of the magnetic unit showed coarse-grained magnetite for an interval of about 250 feet, suggesting that the lower part may be more amenable to concentration than the upper part.

Elongate and arcuate magnetic anomalies as high as 5,000 gammas in amplitude are prominent features of the southern half of the aeromagnetic map of part of Gogebic and Ontonagan Counties, Mich. (G. R. Boynton and J. L. Vargo, 5-64). From the continuity and broad bends in these anomalies, J. E. Case, C. E. Fritts, and W. C. Prinz infer that the anomalies represent folded magnetic units (locally iron-formation), presumably of Animikie age. If this interpretation is correct, the anomalies of higher amplitude merit further investigation as potential iron-ore sources.

MANGANESE

Significance of zoning in manganese veins

In the third of a series of comprehensive papers on the manganese oxides, D. F. Hewett (1-64) presents data that further support his earlier conclusion that many of the veins of manganese oxide in the Western States are of hypogene origin. He concludes that these veins formed near the surface where hot waters from great depth, containing manganese in the manganous state, met the zone of ground water that carried free oxygen. His data also indicate a genetic relation among several of the epithermal deposits in which manganese oxide veins nearest the surface will be succeeded in depth (1) by veins with abundant barite, (2) by those with abundant fluorite, (3) by veins of the epithermal gold-silver type, and finally (4) by veins of the epithermal base-metal type. If the above hypothesis is correct, Hewett notes that holes drilled to reach depths 500 to 1,000 feet below existing workings on manganese oxide veins may strike veins of manganese carbonate and silicate with barite, fluorite, and sulfide minerals that carry enough silver and gold to warrant exploitation.

Permian manganese nodules resemble deep-sea types

Manganese nodules that occur in rock of Permian age about 15 miles southwest of Dillon, Mont., are reported by R. A. Gulbrandsen to be remarkably similar to modern deep-sea manganese nodules. The nodules are in an approximately 2-foot-thick sandstone bed of the Franson Member of the Park City Formation and are located only about 3 feet below the base of the Retort Phosphatic Shale Member of the Phosphoria Formation. They tend to be spheri-

cal in shape with irregular surfaces; most are 2-4 centimeters in diameter. Cross sections of the nodules show many concentric rings of variable width. Semiquantitative spectrographic analyses of one nodule show the presence of 1.5 percent nickel and 0.2 percent cobalt, amounts of these elements that are characteristic of modern manganese nodules. Content of some elements differs markedly, however; for example, 0.02 percent copper in the nodule is unusually low, and 5 percent zinc is high. The nodules are considered to be of syngenetic origin like their modern-day counterparts.

MOLYBDENUM

New occurrences of potential economic importance in Colorado

Two mineralized areas containing molybdenum in amounts sufficient to justify further exploration have been delimited during the course of areal mapping in the Sawatch Range, according to Maurice Brock and Fred Barker. The minerals molybdenite, beryl, pyrite, and fluorite form an equilibrium assemblage in pegmatites, greisens, and high-temperature veins that represent the close of igneous activity in the Twin Lakes batholith. Field evidence indicates little or no time lapse between the magmatic pegmatites and the postmagmatic greisen and vein zones. This mineral assemblage is confined to an area of about 1 square mile in the batholith and is spatially related to small plutons of a porphyry that contains conspicuous quartz phenocrysts. Adjoining the Twin Lakes batholith is an extensive hydrothermally altered area containing molybdenite in a stockwork of closely spaced silicified fractures. Knowledge gained from a study of this deposit resulted in discovery of another similar molybdenite deposit in the vicinity of Tincup. This mineralized rock is limited to an area of less than a square mile, as indicated by molybdenum analyses of stream sediments in the area of limited outcrop.

COPPER

Structural study results in major discovery in Arizona

Recent diamond drilling in the Pima district, Arizona, by the Banner and Anaconda Mining Cos., has disclosed the presence of a concealed copper deposit of major commercial significance. This discovery is a result of an exploration program that was based in large part on an interpretation of the geologic structure of the Pima area made by J. R. Cooper, of the U.S. Geological Survey.⁴ Cooper pos-

⁴ J. R. Cooper, 1960, Some geologic features of the Pima mining district, Pima County, Arizona: U.S. Geol. Survey Bull. 1112-C, p. 63-103.

tulated that a large plate of rock in the northeastern part of the Pima district, which contains the Mission, Pima, and Palo Verde mines, was displaced about 6½ miles to the north-northwest along a nearly horizontal fault called the San Xavier thrust. The Anaconda Mining Co. is sinking a shaft to test drilling results and obtain large samples of ore for metallurgical tests, in preparation for open-pit mining. This deposit is believed by Cooper to be the roots of the displaced Mission, Pima, and Palo Verde ore bodies.

A smaller concealed deposit, discovered by Bear Creek Mining Co. about 4 miles northwest of Anaconda's deposit, is now held by Continental Materials Corp. which is sinking a shaft in preparation for underground mining. The Continental Materials ore body is 1½ miles from any previously known deposit; it is evidently in a slice of rock that was dragged into the San Xavier thrust zone. Other deposits of the same type probably occur within other parts of the thrust zone but may lie as much as 1,000 feet or more below the surface.

Geology of San Manuel copper deposit described

S. C. Creasey (1-65) described the geologic setting of the San Manuel area in Pinal County, Ariz., and, in a section prepared jointly with J. D. Pelletier, he related the copper deposit to the intrusive rocks, structure, and alteration halos. San Manuel is one of the major porphyry copper deposits of the country; its discovery resulted in part from earlier studies by the U.S. Geological Survey and the U.S. Bureau of Mines. Copper mineralization is closely associated with zones of potassium enrichment (probably in Late Cretaceous time) of Precambrian quartz monzonite and Cretaceous(?) granodiorite porphyry. In contrast, zones of argillized rock have abundant pyrite but very little copper.

LEAD-ZINC

Drill-core data on East Tennessee zinc district analyzed

Statistical treatment of drill-core data from the West New Market area in the East Tennessee zinc district confirms previous qualitative evaluation of the relationship of the amount of sphalerite to the stratigraphic thicknesses of limestone and dolomite units in the Kingsport Formation. In a preliminary report Helmuth Wedow, Jr., and J. R. Marie (p. B17-B22) show that the data are compatible with development of the ore-localizing breccia zones by thinning of soluble limestone beds and collapse of less soluble dolomite, with concomitant dilatance of

the dolomite. The dolomite thickens about a third of a foot for every foot of thinning of the underlying limestone, and zinc abundance increases logarithmically to the inverse thickness relation.

Computer analysis aids in exploration for base metals

A computer program has been used for trend-surface analysis of data on vertical stratigraphic thickness and metal-abundance in the Mascot-Jefferson City zinc district, Tennessee. The program has shown that simple geometric surfaces of the 2d and 3d degree, fitted by the least-squares method, account for a significant portion of the total variability of the selected data sets, according to current studies by Helmuth Wedow, Jr. In the case of widely spaced reconnaissance drilling, the trends of these surfaces point to areas of greater probability for favorable ore-bearing structures. Where drill holes are closely spaced to establish the limits of a particular ore body, the 2d-degree surface reveals the elliptical paraboloid habit of the collapse structure, with the long axis of the ellipse indicating the probable trend of the fracture system that localizes the structure.

Geology and ore possibilities in Antler Peak quadrangle, Nevada

R. J. Roberts (4-64) has described the stratigraphy and structure in a key area of the Antler Peak quadrangle where thrust faults of great magnitude superposed a western, eugeosynclinal sedimentary facies over an eastern, miogeosynclinal facies. Roberts and D. C. Arnold (2-65) have described the copper-gold, gold-silver, lead-zinc-silver, and silver and antimony deposits of the quadrangle, related their distribution to the intrusive rocks and structure, and made a number of suggestions for further prospecting. They report that the Duval Corp. has already had some success in recent exploration.

MERCURY

New association for mercury at Yellow Pine district, Idaho, reported

B. F. Leonard (p. B23-B28) reports that appreciable quantities of mercury as cinnabar occur in a stibnite deposit in the Yellow Pine district, central Idaho. Deposits of both antimony and cinnabar have been mined in this district, but this is the only deposit yet known in which the two metals occur together. The deposit is in a silicified zone associated with granite porphyry dikes that intrude the Idaho batholith. The amount of cinnabar is variable; specimens of antimony ore containing conspicuous cinnabar

bar contain about 0.5 to 1.0 percent mercury. Other antimony veins in the district should be tested for mercury.

Composition of natural waters closely associated with quicksilver and copper-lead-zinc-silver deposits

The thermal and mineral fluids now most closely associated with mercury deposits are characterized by relatively high content of Na^{+1} , HCO_3^{-1} , boron, and NH_4^{+1} ; by modest content of sulfide and total dissolved constituents; and by nearly neutral pH, according to D. E. White. The widely held theory of alkaline sulfide transport is not excluded completely by present evidence; other possible transport mechanisms more compatible with the mineralogy and presently associated fluids include the bisulfide, ammonia, and organic complexes, and thermal instability of HgS in sulfide-bearing water and vapor at temperatures about 150°C . Magma and thermally affected sedimentary and metamorphic rocks may all be sources of mercury, with proportions differing from place to place.

In the same study D. E. White also concludes that the rare fluids convincingly related to base-metal-silver deposits are probably brines of high salinity containing Na, Ca, and Cl. These brines generally may be sulfur-deficient, and effective ore formation may depend on some supplementary source of sulfur. Highly saline brines seldom if ever reach the surface in undiluted form because of their density: even when hot, they are much above the density of normal surface waters. Brines of the Na-Ca-Cl type may form in at least three ways: (1) Late-stage magmatic processes, (2) heating of connate waters after deep burial in sedimentary rocks, and (3) reaction of meteoric water + evaporates + rocks in a geothermal environment. Each may be a potent solvent for available base metals and silver.

Association of mercury and ammonia

The association of mercury and ammonia was confirmed by E. H. Pampeyan and R. C. Erd in the Palo Alto quadrangle, California, where the ammonium feldspar buddingtonite, previously recognized only in the Sulfur Bank quicksilver mine in Lake County, northern California, occurs with cinnabar and other metallic minerals in a dike which cuts Eocene sedimentary rocks. Two other ammonia-bearing minerals have been tentatively identified from this dike but are not yet fully described. Although the ammonia alteration has affected the enclosing rocks, the metallic mineralization is confined to the dike rock. At another locality in the quadrangle, ammonia is present in the glass shards of a Miocene diatomace-

ous tuff. The source of ammonia and the time of its introduction are uncertain, but they are presumed to be related to Miocene volcanic activity.

(For use of mercury as an indicator of ores of various types see the section "Techniques of Mineral Exploration.")

ANTIMONY

Ore-bearing fault zone extension noted in central Idaho

Mapping by B. F. Leonard has shown that the Meadow Creek fault zone, which localizes the antimony, gold, and tungsten ore bodies formerly exploited at the Yellow Pine and Meadow Creek mines, continues 2 miles southward from Stibnite, Idaho, to meet the Big Chief silicified zone. The 2-mile extension is concealed by glacial deposits but is indicated by Leonard's regional mapping of the area. Apparent right-lateral horizontal displacement along the Meadow Creek-Big Chief zone is estimated to be a mile or more. Shearing along this major structure probably controlled the development of an adjoining zone of fault blocks and of a nearby belt of northeast-trending fractures occupied by the Little Pistol dike swarm.

GOLD AND SILVER

USGS studies lead to major gold discovery near Carlin, Nev.

Mapping in north-central Nevada by a number of U.S. Geological Survey workers over a period of many years has recently been summarized in several major publications, which, through the guidance they have provided to private industry, have led to discovery of at least one important new ore deposit: the Carlin mine, near Carlin, Nev., opened in May 1965, by the Newmont Mining Corp. The target area was first suggested in a report by R. J. Roberts, P. E. Hotz, James Gilluly, and H. G. Ferguson,⁵ and more specifically in another article by Roberts, in "Geological Survey Research 1960".⁶ Newmont acquired claims in 1961, found gold in 1962, and presently estimates reserves of open-pit ore at 11 million tons containing 0.32 ounces of gold per ton. At its installed capacity of 2,000 tons per day, the anticipated annual yield of more than 200,000 ounces makes this mine the second largest gold producer in the United States, as well as the first major gold discovery in Nevada since 1902.

⁵ R. J. Roberts, P. E. Hotz, James Gilluly, and H. G. Ferguson, 1958, Paleozoic rocks of north-central Nevada: *Am. Assoc. Petroleum Geologists Bull.*, v. 42, p. 2813-2857.

⁶ R. J. Roberts, 1960, Alinement of mining districts in north-central Nevada: *Art. 9 in U.S. Geol. Survey Prof. Paper 400-B*, p. B17-B19.

Finely divided gold found in Pinyon and Harebell Formations, Wyoming

In northwestern Wyoming the Pinyon Conglomerate of Paleocene age and conglomerate in the Harebell Formation of Late Cretaceous age have been found to carry finely divided gold. In the course of a study of lead isotopes in gold, J. C. Antweiler tested more than 40 samples for gold content, using the analytical method developed by H. W. Lakin and H. M. Nakagawa (p. C168-C171) and found gold in both formations in values ranging from a few cents to a dollar per cubic yard. The Pinyon Conglomerate has long been known to contain gold, but gold has never before been reported in the Harebell Formation. Present evidence strongly indicates that both formations are major sources of gold in gravels of the Upper Snake River and its Wyoming tributaries. Little attention has been given to these formations because the finely divided river gold has seldom been profitably recovered, and the conglomerates themselves are very difficult to prospect.

Black calcite as potential source of silver

D. F. Hewett, Arthur Radtke, and Charles Taylor (1-65) noted that black calcite is widespread in many manganese deposits of the West. They found the black color to be due to a few percent of dispersed grains of a manganese oxide. Spectrographic analyses of the black residues remaining after acid decomposition of the calcite matrix in seven samples from districts in Utah, Nevada, Arizona, and New Mexico revealed, in addition to major amounts of manganese, measurable silver ranging up to 5 percent, or 1,500 ounces per ton. The host for the silver in the highest grade sample was found by electron-microprobe study to be a crystalline silver manganate. These results, when considered in the light of the well-known association of silver with manganese minerals in mining districts throughout the West, suggest that deposits containing black calcite should be given special attention in the urgent search for silver ore now underway.

LABORATORY STUDIES OF SULFIDE ORE MINERALS

Effects of minor elements in system Fe-S

P. B. Barton, Jr., has shown experimentally that the addition to pyrite-pyrrhotite charges of sufficient quantities of Mo, Pb, Zn, and Bi to form independent phases of each metal does not measurably affect the temperature of the reaction pyrite = pyrrhotite + sulfur liquid, in the presence of vapor. As noted by other workers, enough Cu to stabilize chalcopyrite

does lower the temperature of the reaction a few degrees. Some workers had attributed large effects to Pb and Mo, but Barton's findings refute these statements.

Barton has also studied the effect of a large number of elements on the (102) *d*-spacing of pyrrhotite; his data show that $d_{(102)}$ may be used safely to determine the metal : sulfur ratio of natural pyrrhotites, except for those unusually rich in Mn, Co, or Ni. Cu, in amount sufficient to form chalcopyrite or cubanite, influences the pyrrhotite $d_{(102)}$ spacing only above 550° C.

Metallogenetic grid constructed

P. B. Barton, Jr., and Priestley Toulmin 3d, have constructed a "metallogenetic grid" for ore minerals, analogous to Bowen's well-known petrogenetic grid for silicate and carbonate rocks. The metallogenetic grid describes the stability fields of sulfide mineral assemblages in terms of sulfur fugacity (f_{S_2}) and temperature. Data for the grid were taken from the literature and from Barton's and Toulmin's experimental work. Supplementing older determinations in the systems Fe-S, Ni-S, and Cu-Fe-S, by Barton and Toulmin, Barton in the past year has determined 4 equilibria in the system Fe-As-S, 5 in Fe-Sb-S, and 1 in Cu-Fe-S. As more data accumulate, the grid should prove increasingly useful in interpreting the conditions of origin of sulfide ore bodies.

Ideality and nonideality in sulfide minerals

Determination of activity-composition relations in solid solutions in the systems Fe-S (pyrrhotite), Fe-Zn-S (sphalerite), and Cu-Fe-S (chalcopyrite with Cu : Fe = 1 : 1), by P. B. Barton, Jr., and Priestley Toulmin 3d, suggests the general rule that substitutional sulfide solid solutions behave thermodynamically as nearly ideal solid solutions, while defect-structure solid solutions depart very strongly from ideality. This generalization, which is still being tested, may be of considerable significance in predicting thermodynamic behavior of minerals of variable composition.

Calculation of activities of elements in binary sulfides

Motoaki Sato and C. L. Christ have applied electrochemical methods to phase-equilibria studies of sulfide minerals. The general equation for a binary sulfide is:

$$E_{M_2S_j} = E^\circ_{M_2S_j} + \frac{RT}{4jF} \ln \left[\frac{(a_M^j)^2_{aq}}{(a_{S^{2-}})^j_{aq}} \cdot \frac{(a_S)^j M_2 S_j}{(a_M)^2 M_2 S_j} \right],$$

where E equals the electromotive force (EMF in volts) for a given cell and F equals Faradays. From

these results the activities of the elements in a binary sulfide can be determined. Two samples of covellite, CuS, from Butte, Mont., indicated sulfur activities of $10^{-2.07}$ and $10^{-1.63}$. Such large variations in EMF result from small variations in chemical composition. Calculated variations are: Ag₂S, 208; CuS, 61; Cu₂S, 386; PbS, 480; SnS, 427. Measured are Ag₂S, 235; and PbS, 483. The EMF is strongly temperature dependent, and hence is a valuable tool in the construction of phase diagrams. Ag₂S and PbS equilibrated at 200°C were assembled into an EMF cell and produced zero EMF at $200^\circ \pm 5^\circ\text{C}$. For natural sulfide pairs, the method could yield minimum temperatures of recrystallization.

Inversion of high-pressure tetragonal CuS

B. J. Skinner demonstrated that high-pressure tetragonal CuS (density, 5.932 ± 0.006 grams per cubic centimeter) inverts to chalcocite (density, 5.840 ± 0.010 g/cc) through a zero-order reaction. The reaction is temperature dependent, decreasing from 25°C to 100°C, at which temperature there is no inversion. These results indicate that tetragonal Cu₂S may be stable over a very small temperature range at 1 atmosphere pressure.

Covellite and djurleite found to coexist in nature

E. H. Roseboom found coexisting covellite and djurleite in natural specimens, contrary to the equilibrium synthetic system which implies that the two minerals should react to form either digenite-djurleite or digenite-covellite assemblages. A sample of covellite-digenite-djurleite from the Jumbo mine, Kennecott, Alaska, reacts to form djurleite + digenite in 24 hours at 95°C. At 60°C the reaction is two-thirds complete after 4 months. Roseboom suggests that the covellite formed as a supergene alteration of original djurleite.

Unquenchable high-temperature phases in system Cu-Ag-S

In studies of the system Cu-Ag-S, B. J. Skinner found unquenchable high-temperature phases which vary over wide ranges in composition. X-ray powder diffraction data suggest that the cation disordered structure proposed by Buerger and Wuensch⁷ for the high-temperature hexagonal polymorph of Cu₂S is also true for the phases found in Cu-Ag-S. These phases are structures with disordered cations that are in a liquid-like state set in arrays of sulfur atoms which are hexagonal closest packed, cubic closest packed, and body-centered cubic.

⁷ M. J. Buerger, and B. J. Wuensch, 1963, Distribution of atoms in high chalcocite, Cu₂S. *Science*, v. 141, p. 276-277.

LIGHT METALS AND INDUSTRIAL MINERALS

EVAPORITES

Bromine as a salinity indicator, Paradox Basin, Utah

O. B. Raup has used the bromine content of halite to estimate the salinity of the environment in which four halite units were formed in the upper part of the Paradox Member of the Hermosa Formation (Pennsylvanian), near Cane Creek, Utah. If further studies of other beds are equally successful, bromine may become a useful indicator pointing to horizons of high salinity, at which exploration for potash deposits may be fruitful. Samples of halite were collected from drill core at 2-foot intervals through the total thickness of each salt bed. The bromine content, as determined by X-ray fluorescence, ranges from 0.008 to 0.035 weight percent. The results show a systematic increase in bromine content, and thus in salinity, from the bottom to the top of each of the salt beds. The regularity of this increase indicates that each of the salt beds was deposited during a continuous period, and it can be inferred that there was neither a major change in depositional conditions nor subsequent metamorphism of the salt.

Borate deposits in Death Valley, Calif., lie along magnetic trough

The alignment of the main Tertiary borate deposits in the Furnace Creek area of Death Valley, Calif., mapped by J. F. McAllister, coincides with a conspicuous trough of low magnetic intensity shown plunging southeastward on an aeromagnetic map by G. E. Andreasen and F. A. Petrafeso.⁸ This relationship extends for 16 miles. The quantity of higher grade ore in exposed deposits increases, in a general way, with decrease of magnetic intensity to the southeast. That a regional structural element controls the locations of borate deposits is suggested further by a Bouguer gravity anomaly map covering an area from Death Valley to the Colorado River. The variations caused by low-density Cenozoic material have been removed from the map so that the gravity contours as shown reflect density contrasts within the basement rocks.⁹ A major zone of gravity highs parallel to Death Valley and another along the Colorado River are connected by a lesser zone of gravity highs curving southward. The connecting zone contains the arc of borate deposits from Fur-

⁸ G. E. Andreasen and F. A. Petrafeso, 1963, Aeromagnetic map of the east-central part of the Death Valley National Monument, Inyo County, California; U.S. Geol. Survey Geophys. Inv. Map GP-428.

⁹ W. H. Diment, S. W. Stewart, and J. C. Roller, 1961, Crustal structure from the Nevada Test Site to Kingman, Arizona, from seismic and gravity observations: *Jour. Geophys. Research*, v. 66, p. 201-214.

nace Creek across Amargosa Valley to Calleville Wash and White Basin east of Las Vegas, Nev.

Borates in California older than previously thought

W. C. Smith reports that vertebrate fossils discovered in beds that overlie the main sodium borate ore body at Kramer, Calif., throw new light on the age of the borate-bearing section at Kramer, heretofore assigned to the upper or Pliocene(?) part of the Tropic Group. R. H. Tedford finds that the fossils are early middle Miocene in age, and thus the borate section is substantially older than previously supposed. Definitive evidence of the age of the colemanite-bearing shales at East Kramer is still lacking, but from a few mollusks found in drill core D. W. Taylor suggests a late Cenozoic age.

Borate resources of California evaluated

A review of resource data by W. C. Smith indicates that the California borax industry can sustain production at recent rates (exceeding 300,000 tons of B_2O_3 per year) for about a hundred years. Kramer and Searles Lake account for about half of these reserves. In addition, some deposits of calcium-bearing borate, now worked on a small scale, are potentially important sources of boron products. Their estimated total reserves probably exceed 5 million tons of B_2O_3 , equal to about 15 years of current production.

PHOSPHATE

Occurrence of phosphate in Beaufort County, N.C.

The phosphorite unit of Beaufort County, N.C., which has been named the Pungo River Formation by J. O. Kimrey, was deposited in an east-trending basin, open to the sea on the northeast. The formation thickens from 20 to about 100 feet from west to east, but the thickening is due to intercalation of barren clay and dolomite beds. Phosphate sand makes up 50–75 percent of the Pungo River Formation in the economic part of the district. J. B. Cathcart finds that the phosphate mineral is a carbonate fluorapatite containing excess CO_2 and fluorine; most samples have between 28 and 31 percent P_2O_5 . The zeolite mineral clinoptilolite is widespread in the Pungo River Formation, but absent in the overlying Yorktown and underlying Castle Hayne Formations; it may be useful as a means of correlation.

Large reserves of phosphate in north Florida and south Georgia

Phosphorite now being explored by several companies in north Florida and adjacent parts of Georgia represents some scores of millions of tons of reserves. J. B. Cathcart notes that, although the age

of this phosphorite is unknown, its geology is reminiscent of the land-pebble field of south Florida. The base of the phosphorite is probably unconformable with the underlying Hawthorn Formation, and the overburden is only a thin Pleistocene(?) sand. Reserves are all in a concentrate of about 32 percent P_2O_5 ; the pebble fraction is minor in amount and too low in grade to be economic.

Enrichment of Bone Valley Formation in two stages documented

A comprehensive review of the genesis of the Bone Valley Formation in Florida by Z. S. Altschuler, J. B. Cathcart, and E. J. Young (1–64) documents the development of a major economic phosphorite through a two-stage secondary enrichment of low-grade, highly dispersed phosphate originally present in the limestones and dolomites of the Hawthorn Formation. Enrichment was first effected by residual accumulation and by solution and reprecipitation of apatite during subaerial weathering of the Hawthorn Formation. Marine transgression terminated the first stage of enrichment and initiated the second. The concentrate produced by earlier weathering was reworked, further enriched, and redeposited as extensive well-bedded deposits during marine recycling. In the process the P_2O_5 content of the individual pebbles and nodules was substantially upgraded.

Phosphate in Mississippian and Permian rocks of Utah

In the Morgan 15-minute quadrangle, in north-central Utah, R. J. Hite has mapped 3 miles of outcrop of the phosphate-bearing Park City Formation of Permian age and also a zone of phosphatic rock of Mississippian age. The area is about 6 miles southeast of Huntsville, Utah, in a faulted and otherwise structurally complex region. Two phosphorite units in the Park City Formation, sampled by E. M. Schell, are of economic interest: the lower one is 8.4 feet thick and averages 26.39 percent P_2O_5 ; the upper one is 5.5 feet thick and averages 28.69 percent P_2O_5 . The best zone of phosphate in the Mississippian rocks, as sampled by Schell, is a unit 1.2 feet thick that averages 27.79 percent P_2O_5 .

Phosphate in the Maquoketa Shale, Iowa, deserves study

C. E. Brown has collected samples from the zone of phosphatic nodules, pellets, and depauperate fossils in the lower few feet of the Maquoketa Shale near Dubuque, Iowa, and in adjacent parts of southwestern Wisconsin and northwestern Illinois. The few samples thus far analyzed show that the lower 3 feet of the shale near Dubuque contains as much as

17 percent P_2O_5 . Furthermore, a sample of 8.5 feet of dolomitic beds 10 to 15 feet above the base of the shale also contains 17 percent P_2O_5 . The lower part of the Maquoketa Shale is widely exposed, and the possibility that local topographic conditions have somewhere resulted in residual concentration of phosphatic material is worth examining.

CLAY AND LIGHTWEIGHT AGGREGATE

Clay in eastern Washington, its composition and utilization

X-ray fluorescence studies by J. W. Hosterman indicate a large variation in the alumina content of the clay deposits in Spokane County, Wash. Residual clay derived from basalt has an average of 22 percent Al_2O_3 , and a few samples have as much as 35 percent. The Latah Formation averages only about 20 percent, but one sample has 39 percent Al_2O_3 . Residual clay from igneous and metamorphic rocks also averages about 20 percent, but no individual sample exceeds 30 percent. These three types of clay are composed of either kaolinite or endellite, and the alumina content could be raised substantially if the nonclay minerals were removed by beneficiation. Clays in the Palouse Formation and in Pleistocene lake beds contain large amounts of montmorillonite and as a result contain less Al_2O_3 .

Tests made in collaboration with the staff of the Norris, Tenn., Research Laboratory of the U.S. Bureau of Mines indicate that some of the clay is suitable for making refractory products. Pyrometric-cone equivalent tests show that residual clay derived from the older igneous rocks would make high-heat-duty refractory products (P.C.E. 31-33), and if the clay were beneficiated, it would make super-heat-duty refractory products (P.C.E. above 33). The clays of the Latah Formation would make intermediate-heat-duty refractory products (P.C.E. 26-31), and the residual clay derived from basalt would make low-heat-duty refractory products (P.C.E. below 26).

Slate from Martinsburg Shale as a potential source of lightweight aggregate

Five large samples of slate from the Martinsburg Shale (Ordovician) collected by A. A. Drake, Jr., from outcrops near the Delaware River in New Jersey and Pennsylvania were fired in a rotary kiln by M. V. Denny and the late H. P. Hamlin, of the U.S. Bureau of Mines (p. D156-D162). The material formed bloated products comparable to presently used commercial material in strength, lightness, and low water absorption. The results suggest that most

if not all of the claystone of the formation would be suitable for lightweight aggregate.

BERYLLIUM

Binary distribution of minor elements at Spor Mountain, Utah

At the Roadside No. 5 beryllium deposit near Spor Mountain, Utah, one group of elements including Li and Sn is distributed evenly and widely at certain stratigraphic horizons, whereas another group of elements including Be, Zn, Pb, and Tl appears to be concentrated close to certain faults as well as along particular stratigraphic horizons. This binary distribution of elements suggests to D. R. Shawe a multiple-source hypothesis in which Li and related elements accumulated early in tuffaceous lake sediments, and Be and related elements were introduced by hot waters following faulting.

Unique mineral association of bertrandite near Gold Hill, Utah

W. R. Griffiths has called attention to several unusual characteristics of the recently discovered large bertrandite deposits near Gold Hill, Utah. These are in quartz monzonite host rocks, lacking the calcareous materials of the host rocks present in other nonpegmatitic beryllium districts, yet the deposits themselves contain much calcite. Furthermore, the deposits lack fluorite, and thus the beryllium cannot have been carried during mineralization as a complex fluoride, but may have been carried as a complex carbonate instead. This mechanism is reflected in the small suite of accessory elements in the deposits.

Beryllium belt in New Mexico and Arizona worth exploring

About a dozen occurrences of beryllium minerals fall in a well-defined belt extending from Socorro County, N. Mex., southwest into the southeast part of Arizona. They mark a province in which D. R. Shawe finds geologic characteristics that suggest that the region should be explored for volcanic-associated deposits similar to those at Spor Mountain, Utah. Like the Utah belt, the New Mexico-Arizona belt contains numerous fluorspar deposits, silicic volcanic rocks with unusually high beryllium content, and topaz-bearing rhyolites associated with thick sections of volcanic rocks containing permeable water-laid tuff. At one Tertiary volcanic locality the beryllium mineral bertrandite has been detected.

Beryllium at Lake George, Colo., related to Redskin Granite

In the Lake George beryllium area, Colorado, C. C. Hawley has found that the fine- to medium-grained

Redskin Granite, a late member of the Pikes Peak magma series, forms small stocks on the southwest flank of the Pikes Peak batholith, near a large bend in the batholith contact. Beryllium deposits are associated with these stocks. Spectrographic and chemical analyses show that the earlier, coarse-grained Pikes Peake Granite of the batholith is characterized by Ba, Sr, rare earths, and Zr. The finer grained Redskin Granite contains more of such elements as Be, Li, Sn, and Rb, which are also found in the beryllium deposits of the area. Thus it seems that the processes which enriched these elements in late members of the Pikes Peak magma series ultimately contributed Be and related elements to solutions that formed the beryllium deposits of the Lake George area.

Average beryllium content of large intrusions in Colorado

From analyses of 120 representative samples by gamma activation, W. N. Sharp has calculated an average value of 7.7 parts per million of beryllium in the Mount Antero leuco-granite stock and 5.6 ppm of beryllium for the Pikes Peak granite batholith in Colorado. The few analyses from samples that contain beryllium minerals have more than 30 ppm of beryllium. In granite containing less than 30 ppm, the beryllium is probably held in feldspar and mica.

Preliminary analyses of a rich blue, strongly pleochroic beryl from Mount Antero show it to have 6 to 8 percent Fe_2O_3 . This amount of iron, if verified by additional analyses, is most unusual for beryl.

Beryllium in thorite veins, Wet Mountains, Colo.

Samples from two thorite veins, collected by D. R. Shawe from the Wet Mountains, Colo., contain 0.02 to 0.2 percent beryllium. These lie within a few thousand feet of small bodies of Tertiary rhyolite intrusives that contain unusually high amounts of Be (6–16 parts per million), and may be genetically related to the veins.

Localization and zonation of beryllium, Lost River, Alaska

The Rapid River fault, according to C. L. Sainsbury, is the most important ore localizer for beryllium-fluorite deposits in the Lost River area, Alaska. The fault is mineralized for almost half of its extent over a distance of 8 miles. A strong zonation exists among the beryllium-fluorite deposits: the beryllium mineral helvite occurs nearest the granite in banded fluorite-magnetite skarn, succeeded by a zone of beryllium-fluorite-tin-sulfide veins in which chrysoberyl is the main beryllium mineral, and then by

beryllium-fluorite veins containing chrysoberyl, fluorite, euclase, hematite, and little or no tin or sulfide minerals.

ZEOLITES

Diagenesis of zeolites in Mohave Desert, Calif.

A conspicuous bedded tuff near the middle of the Ricardo Formation of Pliocene age in the El Paso Mountains of southern California has afforded R. A. Sheppard and A. J. Gude 3d, the opportunity to study the stages of diagenetic alteration from fresh rhyolitic glass to complete replacement of vitric material by clinoptilolite, opal, and montmorillonite (p. D44–D47). Tuff above an elevation of 3,000 feet generally is fresh, but the same tuff below 3,000 feet is altered. The tuff dips 10° to 20° northwest, and zeolitic alteration probably was accomplished by subsurface water after the Ricardo Formation had been tilted. Solution and hydrolysis of rhyolitic glass by the downward-moving subsurface water would increase the pH and salinity of the water, thereby providing a chemical environment more favorable for the formation of zeolites.

In altered rhyolitic tuffs of the Barstow Formation, zeolites that have been identified include analcime, chabazite, clinoptilolite, erionite, mordenite, and phillipsite. These are in nearly monomineralic beds or associated with other zeolites, clay minerals, opal, and potash feldspar. Chabazite is of particular interest because it is only rarely reported from sedimentary environments.

Preliminary study of tuffs from the Pleistocene Lake Tecopa indicates that vitric material is preserved only in the vicinity of fresh-water inlets. Elsewhere the tuffs are altered to zeolites and potash feldspar.

Zeolite makes up Chalk Cliffs at Mount Princeton, Colo.

Leonhardite, a calcium zeolite, has been found by W. N. Sharp to form the white chalky matrix surrounding fractured and crushed fragments of Mount Princeton Quartz Monzonite in the Chalk Cliffs at the site of the Mount Princeton hot springs on the west side of the Arkansas River valley near Buena Vista, Colo.

Zeolite replaces tuff in McMullen County, Tex.

Extensive deposits of white claylike altered tuff in McMullen County, Tex., consist largely of clinoptilolite associated with opal, montmorillonite, and a trace of quartz. D. H. Eargle regards this as probably the basal part of the Jackson Group (Eocene). The clinoptilolite was recognized from X-ray analyses by Dr. Shoichiro Hayashi, visiting Japanese geologist, and the abundance of it and other constit-

uents was determined by A. D. Weeks and M. L. Lindberg.

RADIOACTIVE MINERALS

URANIUM

Deposits in White Canyon and Elk Ridge areas, Utah, described

The uranium deposits close to the base of the Chinle Formation in the adjoining White Canyon and Elk Ridge areas, San Juan County, Utah, are described by R. E. Thaden, A. F. Trites, Jr., and T. L. Finnell (1-64) and by R. Q. Lewis and R. H. Campbell (1-65). In the White Canyon area the deposits are in sandstone and some interbedded mudstone that fill channels cut into the underlying Moenkopi Formation. In the Elk Ridge area most of the deposits are in similar bodies of rock that rest on the Moenkopi rather than in channels cut into it; ore does not occur in sandstone lenses of the Chinle where such lenses are separated from the Moenkopi by mudstone. Lewis and Campbell suggest that ore was deposited from solutions which migrated mainly through siltstone in the upper part of the Moenkopi and entered the host rocks at places where they were in contact with the Moenkopi.

The deposits of both areas are in a broad belt which extends eastward from the Colorado River at least to the Comb Ridge monocline and may extend into the adjoining Abajo Mountains area where, however, the Chinle Formation is buried by younger rocks (I. J. Witkind, 1-64).

Prelaccolithic age of uranium deposits in Abajo Mountains, Utah

The known uranium deposits in the Abajo Mountains area, Utah, are in the Salt Wash Sandstone Member of the Morrison Formation. Because one of these deposits is displaced by small faults, I. J. Witkind (1-64) suggests that the deposits were emplaced before the sedimentary rocks were domed and faulted by intrusion of the igneous rocks that form the laccoliths of the Abajo Mountains.

Uranium-bearing pipe formed by solution and collapse of limestone

Features observed by C. G. Bowles at the uranium-bearing pipelike structure at the Orphan mine on the South Rim of the Grand Canyon, Ariz., and at other pipes in the eastern part of the Grand Canyon area strongly suggest that the pipes originated prior to cutting of the canyon as the result of collapse of caverns formed by solution of the Redwall Limestone of Mississippian age. When the caverns collapsed

the pipes were formed by stoping of overlying rocks as high stratigraphically as the Coconino Sandstone of Permian age.

Map of Moab quadrangle locates uranium deposits

The geologic setting and distribution of uranium deposits in the important Lisbon Valley area, Utah, and the Uravan mineral belt, Colorado, are shown on a map of the Moab quadrangle covering 1° of latitude and 2° of longitude at a scale of 1:250,000 (P. L. Williams, 1-64). (For further details, see the section "Regional Geology, Colorado Plateau.")

Uranium deposits in Powder River Basin, Wyo.

Uranium deposits in the Powder River Basin, Wyo., are in sandstone mainly in the Wasatch Formation of Eocene age (W. N. Sharp and others, 1-64; W. N. Sharp and A. B. Gibbons¹⁰). The deposits are in sandstone which is relatively coarse grained and forms lenses elongate in a northerly direction. Much of the sandstone is drab yellowish gray, but the deposits are confined to a zone within which some lenses and parts of others are colored red to pink. The zone of partly red rocks is nearly 70 miles long and aligned with the subsurface axis of the basin. Most of the uranium is concentrated in drab sandstone immediately adjacent to the edge of the red sandstone, and the principal deposits occur very near the general boundary of the zone of partly red sandstone. The uranium-bearing rock forms various shaped bodies that conform to the irregular shape of the color contact. The intimate relation between uranium deposits and the edge of red sandstone suggests that both features are results of the same sequence of events.

Crescent-shaped ore bodies in Gas Hills and Crooks Gap areas, Wyoming

From study in the Gas Hills uranium district, Fremont and Natrona Counties, Wyo., F. C. Armstrong infers that the principal ore bodies are all along or near the same irregular sinuous contact between altered and unaltered rock that was once continuous throughout the district.

Known uranium deposits are distributed along three subparallel zones, the west, central, and east ore-trends, which are oriented west of north. The ore bodies are sinuously elongate in the direction of the ore trends, and at many places are irregularly crescent shaped in cross section. Altered rock occurs on the concave side of the crescent, and unaltered rock on the convex. Characteristically, thick high-

¹⁰ W. N. Sharp and A. B. Gibbons, 1964, Geology and uranium deposits of the southern part of the Powder River Basin, Wyoming: U.S. Geol. Survey Bull. 1147-D, p. D1-D60.

grade ore occupies the body of the crescent, and thin lower grade ore occurs as tabular bodies in the greatly extended lower horn of the crescent (lower limb ore). The crescents are convex southwestward in the west trend and northeastward in the central trend. Drilling shows that between these two ore trends is a thick prism of altered rock that is bounded, as though by parentheses, by crescent-shaped ore bodies. The orientation of the crescents in the two trends and the continuity of altered rock between them, strongly suggest that the north ends of the trends were once connected along a segment of contact that has been removed by erosion.

A corollary of this hypothesis is that only the thin lower-limb mineralized zone is present, beneath altered rock, in the area between the west and central trends. Locally this lower limb may be of sufficient thickness and grade to form small- to moderate-sized ore bodies. The same line of reasoning suggests that the south ends of the central and east ore trends are connected southward beneath cover. The area between the trends would be occupied by unmineralized and unaltered rock between convexly opposed crescents.

F. C. Armstrong also reports that an ore body of crescent-shaped cross section has been recognized by company geologists in the Reserve shaft mine of Continental Materials Corp. in the Crooks Gap area, Fremont County, Wyo. Uranium ore bodies of this form have been previously recognized and are common in the Shirley Basin and Powder River Basin areas. The similarity in shapes of ore bodies and in relative positions of altered rock, ore, and unaltered rock in all four areas suggest that the uranium deposits were all formed in the same manner.

Environments of roll ore bodies of uranium

In a review of information about discordant curved layers of uranium ore in sedimentary rocks, commonly called 'rolls, D. R. Shawe and H. C. Granger (1-65) suggest that environmental differences permit the distinction of two types. Rolls in the Colorado Plateau region are surrounded by a wide halo of reduced altered rock. They were probably formed before or during maximum burial of enclosing strata prior to major structural deformation. Rolls in the Tertiary basins of Wyoming are bounded on one side by relatively oxidized rock and on the other by relatively reduced rock. They were probably formed near the surface following deformation, uplift, and denudation.

Uranium deposits in Gulf Coastal Plain, Tex.

D. H. Eargle has noted that relatively unoxidized

uranium ore in the Galen mine, Karnes County, Tex., adjoins a tongue of bleached sand similar in color to altered sand adjacent to ore in some deposits in Wyoming. Coffinite, not previously known in Texas deposits, has been identified as one of the ore minerals by A. D. Weeks. Molybdenum and selenium minerals occur in 1 to 2 feet of sand overlying uraniferous sand.

Uranium deposits in Front Range Colo., fit zoning pattern

Uranium deposits in the Front Range, Colo., described by P. K. Sims and D. M. Sheridan (3-64), appear to be zonally arranged with respect to the Colorado mineral belt, as indicated by their distribution and size, and by systematic differences in mineralogy, structure, and texture of veins between deposits within the belt and those outside of it. The largest and most valuable ore bodies are in epithermal deposits marginal to the belt.

THORIUM

Thorium-bearing veins and rocks in New Mexico

Some of 45 feldspar-rich bodies in the southern Caballo Mountains, Sierra County, N. Mex., studied by M. H. Staatz, J. W. Adams, and N. M. Conklin (p. D48-D51) contain thorite. Uranophane and bastnaesite, a rare-earth mineral, are sparsely present in one body. The bodies are irregular in shape and range from a few feet to 300 feet in length.

Thorium-bearing veins, which occur in Dakota Sandstone and phonolite near Loughlin Peak, Colfax County, N. Mex., are reported by M. H. Staatz to contain a varied mineral assemblage which includes plumbogummite, zircon, xenotime, lepidocrocite, goethite, hematite, jarosite, quartz, and a rhabdophane-like thorium-bearing mineral. One vein from 1 to 1.5 feet thick and at least 600 feet long has an average grade of about 0.4 percent thorium.

Wet Mountains thorium area, Colorado, extended

Discovery by Q. D. Singewald of a complex pattern of felsite dikes mineralized with thorium in the southeast part of the Rosita quadrangle, Custer County, Colo., and of anomalous radioactivity along veins in the west half of T. 19 S., R. 71 W., Fremont County, greatly enlarges the areas in the vicinity of the Wet Mountains where rocks are known or suspected to be mineralized by thorium. The dikes in the Rosita quadrangle cut Precambrian rocks but are absent in areas of Tertiary volcanic rocks. The radioactivity of the dikes ranges from 2 to 3 times background where they are unaltered to many times background where they are veined by later minerals. In places the radioactivity exceeds 5.0 milliroentgens

per hour, the highest reading on a scintillation detector.

MINOR ELEMENTS

Use of absorption spectra to identify rare earths

The detection of rare-earth elements by their absorption spectra, although a method long known, has modern practical application in the investigation of rare-earth-bearing minerals. Recently J. W. Adams (2-65) has described in detail the absorption spectra of certain rare-earth minerals as determined by the hand spectroscope and microspectroscope. By the use of this convenient method many rare-earth minerals may be recognized in the field or laboratory and a distinction commonly can be made between dominantly cerium- or yttrium-group species.

Rare-earth-rich shear zone traced in Wet Mountains, Colo.

As part of his study of thorium resources, M. H. Staatz has traced a shear zone containing discontinuous rare-earth- and thorium-bearing veins for more than 1,500 feet in the Road Gulch area of the northern Wet Mountains, Colo. Selected samples from the veins contain as much as 19 percent rare earths and 1.1 percent thorium. Minerals that make up the veins include dolomite, siderite, calcite, quartz, hematite, goethite, barite, rutile, chlorite, fluorite, bastnaesite, monazite, thorite, zircon, and fergusonite.

ORGANIC FUELS

Studies that deal directly with the discovery, delineation, evaluation, and interpretation of coal, petroleum and natural gas, oil shale, and humate deposits are reported in this section. Related investigations that are concerned primarily with the geology and (or) geophysics of potential areas of production of such fuels are reported under other headings.

COAL

Rocky Mountains region

Projection of current trends in the use of coal for the production of coke and electricity has led Paul Averitt (1-65) to conclude that coal mining in the Rocky Mountains region will increase about 50 percent by 1970. An upward trend in mining began in 1958 and 1959 and terminated a postwar downtrend.

North-central New Mexico

A zone of subbituminous coking coal in the nearly flat lying Upper Cretaceous and Paleocene Raton Formation has been discovered and mapped by C. L. Pillmore (1-65, 2-65) in the Catskill NE and SE

quadrangles, Colfax County, north-central New Mexico. This coal zone crops out near the crests of flat ridges at an altitude of about 7,200 feet in a 16-square-mile area near the confluence of the Vermejo River and Gachupin Canyon, about 25 miles west of Raton, N. Mex. The overburden is generally less than 100 feet and commonly less than 50 feet thick. The zone averages about 7 feet in thickness and generally contains about 48 inches of coal. Near the mouth of Ancho Canyon in the eastern part of its outcrop area, however, the zone exceeds 15 feet in thickness and contains more than 100 inches of coal in beds as much as 30 inches thick.

Southwestern Oregon

The correlation of bituminous coal beds in the Tyee Formation of middle Eocene age between the Eden Ridge coal field and Squaw Basin, its southward extension, in Coos County, southwestern Oregon, is reevaluated by R. G. Wayland (1-65). On the basis of new core-hole data, brief field study, and photogeologic interpretation, the known coal beds in Squaw Basin are concluded to be of limited lateral extent and older than those at Eden Ridge. Squaw Basin is regarded as a possible source of strippable coal.

Western Arkansas

Original coal reserves in the Lower Hartshorne coal bed of the McAlester Formation of Pennsylvanian age in the Scranton, New Blaine, and Greenwood quadrangles, Logan and Sebastian Counties, western Arkansas, have been revised by B. R. Haley and T. A. Hendricks on the basis of new geologic data. Original reserves are estimated to be about 83 million tons of coal in the Scranton and New Blaine quadrangles and about 316 million tons of coal in the Greenwood quadrangle. These more reliable, new estimates differ by only 20 percent or less from previous estimates that were used in compiling the total coal reserves of the State (B. R. Haley ¹¹).

Eastern Pennsylvania

The original coal reserves in Pennsylvanian rocks of the Western Middle anthracite field of eastern Pennsylvania totalled more than 6.7 billion tons in beds exceeding 14 inches in thickness, according to a recent appraisal by H. H. Arndt. About 1.8 billion tons of anthracite and semianthracite had been mined or lost through mining operations as of January 1, 1963. Of the remaining 4.9 billion tons of reserves, about 2.8 billion tons are categorized as probable recoverable reserves.

¹¹ B. R. Haley, 1960, Coal resources of Arkansas: U.S. Geol. Survey Bull. 1072-P, p. 795-831.

Northwestern Alaska

Low-volatile bituminous coal in steeply dipping rocks of Early Mississippian age on the Lisburne Peninsula, northwestern Alaska, is described and evaluated by I. L. Tailleux (p. B34-B38). The coal is found in 13 seams 2½ feet or more thick and has an aggregate thickness of 70 feet. Analyses (on an as-received basis) of representative samples indicate that the coal is composed of 13-21 percent volatile matter, 68-80 percent fixed carbon, and 3-16 percent ash. The heating value of the coal ranges from 11,800 to 14,300 Btu. The coal has an important potential as a local source of fuel for the village of Point Hope, only 30 miles from the closest seams.

Eastern Montana

Recent information on the coal resources of eastern Montana is given in the section "Resource Compilation" under the side heading "Mineral Fuels of Eastern Montana."

PETROLEUM AND NATURAL GAS

Navajo Reservation, northeastern Arizona

The lower third of the Cutler Formation of Permian age probably includes marine rocks in the subsurface of northeastern Arizona, between the Monument and De Chelly upwarps. Subsurface correlation by R. B. O'Sullivan suggests that this part of the Cutler is continuous with part of the Supai Formation of Pennsylvanian and Permian age, which has yielded major shows of oil and gas in central Arizona. Although only a single slight show of oil and gas has been recorded to date from the lower third of the Cutler, a large area of northeastern Arizona that is underlain by this unit may have an oil- and gas-producing potential.

Denver Basin, northeastern Colorado

Porosity rather than structure may have localized accumulations of oil and gas in the Upper Cretaceous "D" sand of the Denver Basin, south of Fort Morgan, Colo. Unconnected, lenticular zones in the "D" sand that have greater effective porosity than lithologically similar enclosing beds of the same sand have been recognized by H. L. Cullins. These lenticular zones generally are about one mile long, half a mile wide, and 8 to 14 feet thick. Within a single oil and gas field, apparently anomalous gas pools that are structurally lower than oil pools, and oil or gas pools that are structurally lower than non-productive areas probably are controlled by these zones. Recognition and exploration of these porous zones may increase substantially the oil- and gas-producing potential of the Denver Basin.

OIL SHALE AND HUMATE DEPOSITS

Oil shale, northern Alaska

New data obtained by H. A. Tourtelot and I. L. Tailleux from recently collected samples may help interpret the possible origin of rich oil shale in northern Alaska. Samples that yield about 50 to 140 gallons of oil per ton consist almost entirely of a sporelike microfossil *Tasmanites*, identified by J. M. Schopf, at two localities (I. L. Tailleux,¹² p. D132, No. 2 and 5). Quartz and barite are the only accessory minerals in these samples. Semiquantitative spectrographic analyses of the ash of three samples show 0.1-3.0 percent barium, 150-700 parts per million of molybdenum, and 100-500 ppm of vanadium.

Shallow humate deposits, northwestern Florida

Widespread lenticular layers of coastal and near-coastal sands in northwestern Florida have been found to contain organic matter that is 95 to 99 percent humate. This humate, which is entirely and readily soluble in slightly alkaline water, has an average composition of 55 percent carbon, 4.4 percent hydrogen, 38.5 percent oxygen, 1.4 percent nitrogen, and 0.7 percent sulfur. According to estimates by V. E. Swanson and J. G. Palacas, 0.1 to 1 million tons of humate per square mile is present at a depth of less than 35 feet along a 3-mile-wide coastal strip between Pensacola and Apalachicola in northwestern Florida. The total content of easily soluble organic matter in this area of 300 to 500 square miles is estimated to be about 100 million tons.

TECHNIQUES OF MINERAL EXPLORATION

Tellurium and mercury as indicators of ore

In the Cripple Creek district, Colorado, a telluride-gold district, tellurium is widespread but is most concentrated in areas that have been mined for gold. According to G. B. Gott and J. H. McCarthy, Jr., mercury, tellurium, and silver seem to be associated with the Cripple Creek gold deposits, and the presence of any 2 of these 3 elements is indicative of higher than normal concentrations of gold. By way of contrast, tellurium concentrations in the vicinity of the Comstock lode are lower than expected, according to D. F. Davidson and H. W. Lakin, and seem to show no pattern of distribution that is strongly related to the silver deposits.

Mercury seems to be associated with all of the epithermal metal deposits studied to date by G. B. Gott and J. H. McCarthy, Jr. These include poly-

¹² I. L. Tailleux, 1964, Rich oil shale from northern Alaska: Art. 148 in U.S. Geol. Survey Prof. Paper 475-D, p. D131-D133.

metallic deposits in the Ely, Nev., and Park City, Utah, areas; silver deposits at Hamilton, Nev., and the gold deposits at Cripple Creek, Colo.

According to J. L. Jolly and A. V. Heyl (1-65) mercury anomalies may indicate concealed base-metal deposits. They report that 11 sphalerite samples from the central Kentucky and central Tennessee mineral districts contain from 1 to 10,000 parts per million of mercury. This is the first known occurrence of mercury in sphalerite east of the Mississippi River. The mercury probably occurs in the sphalerite lattice rather than as a discrete element, and its presence strongly suggests deposition by slightly alkaline solutions at a temperature range of 70° to 100°C. Mercury-bearing geochemical halos above concealed deposits of base metals are suggested by selected samples from wallrock suites in the central Kentucky district and in the Austinville, Va., district. At the Faircloth mine, Woodford County, Ky., the mercury content of the wallrock ranges from 0.17 ppm in dolomite adjacent to the vein minerals to 0.01 ppm in unaltered limestone. At the Ivanhoe mine in the Austinville district, a wallrock suite, consisting of partly altered carbonate rock in and near ore, contains 1.5 to 3.5 ppm of mercury in contrast to 0.03 ppm of mercury in normal or ordinary limestone. The presence of higher than background amounts of mercury in sphalerite and wallrock of the eastern and midwestern United States base-metal districts suggests the possibility of using mercury as a prospecting guide in these areas. The mercury-bearing sphalerite samples also contain 0.01 to 0.1 percent germanium and 1 to 2 percent cadmium.

Tellurium and mercury as indicators of "productive" jasperoid

Ninety-three samples of jasperoid from 22 areas, including 14 mining districts, were analyzed for tellurium by H. W. Lakin and for mercury by J. H. McCarthy, Jr., using sensitive methods capable of detecting 4 parts of tellurium and 1 part of mercury per hundred million. These samples had previously been classified by T. G. Lovering and J. C. Hamilton¹³ into "productive" and "nonproductive" types.

Tellurium content of these samples ranged from 200 to 0.04 parts per million, and mercury content from 90 to 0.02 ppm. In order of decreasing affinity, tellurium showed a statistically significant association with arsenic, bismuth, silver, antimony, lead, copper,

mercury, and barium. Mercury, in order of decreasing affinity, showed a statistically significant association with silver, lead, copper, tellurium, zinc, antimony, arsenic, bismuth, and molybdenum. Tellurium in concentrations of > 1 ppm and mercury in concentrations > 4 ppm were found to be indicators of jasperoid of the productive type.

Trace metals associated with gold deposits

As part of a study of geochemical halos, R. L. Erickson has learned that the low-grade gold deposits of north-central Nevada have a characteristic trace-metal suite of mercury, arsenic, and antimony. Small but anomalous amounts of tungsten are also common. Mercury seems to be most directly associated with gold, whereas arsenic and antimony may form halos around the most intense gold mineralization. The detection of anomalous amounts of these metals in soils and rocks is a useful exploration tool in the search for concealed gold ore.

Selenium as an indicator of uranium in Shirley Basin, Wyo.

Analyses of samples collected across roll-type bodies of uranium ore in the Shirley Basin, Wyo., show, according to E. N. Harshman, that selenium is notably concentrated at the contact of oxidized barren sandstone with ore-bearing sandstone. Because selenium is most readily moved in slightly alkaline relatively oxidized solutions and is precipitated by a decrease in pH (H. W. Lakin,¹⁴ p. 11-12), the distribution of selenium-rich rock suggests that slightly acidic conditions prevailed at the sites of deposition.

Accessory minerals in igneous rocks as indicators of ore

G. J. Neuerburg is studying patterns of distribution of minor accessory minerals in igneous bodies in the attempt to trace ore materials from the presumed source, through palimpsest channels of movement, toward areas of possible deposition. The distribution of accessory sulfides and magnetite, but not zircon, definitely appears to be related to skarn deposits of tungsten peripheral to the Asgood quartz monzonite stock, Humboldt County, Nev. Of the 15 deposits actually worked, 9 are adjacent to the alleged points of exit of the ore fluids. The principal objective of this mineralogic approach as a prospecting tool is to locate areas of higher-than-average ore potential, which could then be prospected intensively by a variety of other methods.

¹³ T. G. Lovering and J. C. Hamilton, 1962, Criteria for the recognition of jasperoid associated with sulfide ore: Art 63 in U.S. Geol. Survey Prof. Paper 450-C, p. C9-C11.

¹⁴ H. W. Lakin, 1961, Geochemistry of selenium in relation to agriculture, in M. S. Anderson, and others, Selenium in agriculture: U.S. Dept. Agriculture, Agriculture Handbook 200, p. 3-12.

Zinc-rich manganese-iron oxides cause spurious anomalies

On the basis of geochemical prospecting surveys of streams in Maine, F. C. Canney, E. V. Post, and W. H. Dennen report that some heavy-metal anomalies are apparently not related to mineral deposits but are derived, instead, from the zinc-rich nodules and coatings of manganese-iron oxides found to be plentiful in many stream courses. Sediment samples from streams draining mineralized areas, however, have above-average heavy metal: manganese ratios; thus this ratio appears to be a useful criterion for distinguishing between significant and nonsignificant anomalies.

New rapid analytical methods for gold and silver developed

New colorimetric procedures for the determination of silver and gold in geologic materials have been developed which should greatly aid in prospecting for those two elements. In the method reported by H. M. Nakagawa and H. W. Lakin for silver (p. C172-C175) the sample is digested with nitric acid and the silver is extracted with triisooctyl thiophosphate (TOTP) in benzene and stripped from the organic phase with dilute hydrochloric acid. Silver is then measured indirectly by its catalytic action on the persulfate oxidation of manganous ion to permanganate. The lower limit of sensitivity of the method is 0.01 parts per million. About 80 determinations can be made per man-day. In the method reported by Lakin and Nakagawa for gold (p. C168-C171), the gold in the sample is dissolved with sodium bromate and hydrobromic acid and then is extracted from dilute hydrobromic acid with ethyl ether. The ether is evaporated to dryness and the residue is taken up in water after which the solution is buffered and then extracted with a solution of 4,4'-bis (dimethylamino) thiobenzophenone (TMK) in isoamyl alcohol, forming a red gold-TMK complex. Absorbance of the complex is proportional to the amount of gold present. Thallium when present in relatively large amounts is an interference. As little as 0.02 ppm of gold can be determined, and about 25 analyses can be made per man-day.

Gamma-ray spectrometer used to locate zones of alteration

A truck-mounted gamma-ray spectrometer has been used by R. M. Moxham, R. S. Foote, and C. M. Bunker¹⁵ to make in situ analyses of uranium,

¹⁵ R. M. Moxham, R. S. Foote, and C. M. Bunker, 1965, Gamma-ray spectrometer studies of hydrothermally altered rocks: *Econ. Geology*, v. 60, no. 4, p. 653-671.

thorium, and potassium in hydrothermally altered zones associated with copper and copper-lead-zinc deposits in central Arizona. Equating the field spectrometric data with chemical analyses of samples collected at the spectrometer stations gave standard errors of 0.6 percent potassium, 3 parts per million of thorium, and 10 ppm of uranium for the outcrop analyses with the spectrometer. Potassium in the more intensely altered zones is about twice that in unaltered areas. There is no corresponding increase in thorium, so a higher potassium:thorium ratio also results from alteration; an abnormally high uranium content, locally in secular disequilibrium, was measured at the Bagdad, Ariz., porphyry copper deposit. The amount of potassium introduced by hydrothermal alteration is thought sufficient in quantity to be generally detectable by either surface or aerial spectrometry; spectrometric detection of anomalous uranium is probably marginal, owing to secular disequilibrium. The potassium:thorium ratio may provide a means of distinguishing, in such surveys, between hydrothermally altered high-potassium zones and unaltered extrinsically high-potassium intrusives.

Geobotanical prospecting in Alaska

H. T. Shacklette has found that many bryophyte (mosses and liverworts) species are associated in Alaska with concentrations of minerals, but only one species, *Gymnocolea acutiloba*, appears to function as an indicator on a regional scale. This species was found to grow only on substrata having a high copper content. Another species, *Cephalozia bicuspidata*, seems to be a local indicator of mineral deposits; it grew on a galena and sphalerite ore vein that contained 12.9 percent lead and 34.0 percent zinc, whereas other species at the site could not tolerate such high concentrations of these elements. Bryophytes appear to be indifferent to the concentrations of mercury, antimony, arsenic, limonite, nickel, and chromium that occur in rock outcrops in Alaska.

WATER RESOURCES

The U.S. Geological Survey investigates the occurrence, availability, and quality of surface and underground waters and the sediment discharge of streams. A hydrologic-data network which extends throughout the country provides continuing series of several types of basic data. During 1965, discharge and water-level data for surface waters were collected at about 8,000 stream-gaging stations and

about 800 lake-level stations. Continuous or periodic measurements of ground-water levels were made in about 16,000 observation wells. The quality of surface waters was monitored at about 1,670 stations; chemical, temperature, and sediment data were collected.

These basic hydrologic data are compiled and published in the following Water-Supply Paper series of the Geological Survey:

- "Surface Water Supply of the United States,"
- "Ground-Water Levels in the United States,"
- "Quality of Surface Waters of the United States," and
- "Quality of Surface Waters for Irrigation, Western United States."

The Surface Water Supply series, formerly published annually, is to be published at 5-year intervals beginning with the period 1961-65. Each report is in 16 parts: 14, determined by drainage basins, for the 48 conterminous States, and 2 for Alaska and Hawaii. Interim annual reports by States are being published. The reports on water quality are published annually, but conversion to 5-year publication, with interim annual reports, is anticipated. The water-quality records are grouped in terms of the same drainage basins used for the reports on surface-water supply. The reports on ground-water levels, also published at 5-year intervals, are in 6 parts which represent geographical sections of the country. In addition to these reports, a series of Water-Supply Papers is published which describes the magnitude and frequency of floods for the entire country, by drainage-basin areas, and another series describes notable floods each year.

Areal investigations of water resources are made largely in cooperation with State, local, or Federal agencies listed on page A205. These studies include the various aspects of the geologic and hydrologic environment that relate to the occurrence and movement of water on the surface and underground. Such studies of water resources stress the evaluation of sources of supply, chemical and physical composition, computation of the quantity available for use, description of the direction and rate of movement, evaluation of fluctuations in flow, and determination of disposition of the supply as use, waste, or outflow.

Diversified water-resources investigations are in progress in nearly every State. These fall into two general categories: "area" and "systems" studies. Area studies cover investigations of specific hydrologic problems within an area, generally comprising a political subdivision—the problems of a munici-

pality, a county, or a State. Systems studies, on the other hand, are investigations of the hydrologic environment of natural units such as a river basin or isolated valley or a major aquifer, whose area may include a number of political subdivisions. The purpose of these investigations is to determine the effect on the hydrologic system of changes in any part of it; for example, to predict how use of ground water in one municipality may influence stream-flow in another part of a river system.

Investigations stressing the economic aspects of water as a resource are treated in the following section under four areas (fig. 1), which correspond to the administrative subdivisions of the Water Resources Division.

OFFICE OF WATER-DATA COORDINATION

An Office of Water-Data Coordination has been established in the Water Resources Division of the U.S. Geological Survey to implement Circular A-67 of the Bureau of the Budget. This circular establishes guidelines for coordinating Federal activities in collecting water data and for developing a national plan to acquire these data. It assigns these coordinating and planning functions to the Department of the Interior, which in turn has assigned this responsibility to the Geological Survey.

Circular A-67 stresses the importance of participation in the planning of these activities by all Federal agencies concerned with the collection and use of water data. It identifies three principal functions of coordination:

- (1) Design and operation of a national network for the collection of water data.
- (2) Coordination of national-network and specialized water-data activities.
- (3) Maintenance of a central catalog of information on national-network and specialized water data, and on Federal activities being planned or conducted to acquire such data.

Among the chief purposes of Circular A-67 are assurance that (a) water information flows freely to those who require it for the effective development and management of water resources; (b) data are collected in a timely, effective, and economical fashion; and (c) information is readily accessible to all users of water information from a single focal point.

ATLANTIC COAST AREA

Moderate to plentiful supplies of water are commonly available in the Eastern States. Water-

FIGURE 1.—Index map of the United States, showing areal subdivisions used in discussion of water resources.

resources investigations of the U.S. Geological Survey in the Atlantic coast area have as their objective the description and interpretation of the occurrence of water, with emphasis on problems which affect water use. These problems relate to temporary excesses caused by floods and deficiencies due to drought such as that which persisted through the fiscal year in the Northeastern States; to natural variations in supply which reflect local and regional differences in rocks and soils, vegetation, and climate; and to natural quality of the water and changes in quality brought about, directly or indirectly, as a result of our use of this resource.

The results of project investigations summarized below illustrate the variety of studies in progress in this region of diverse geologic and hydrologic environments.

INTERSTATE STUDIES

Hydrologic atlas of Appalachia

Water will be an important resource in the proposed program for economic development of the Appalachian Region, according to W. J. Schneider

and others (1-65), because the combination of availability of water and accessibility by highways will largely determine the locations of future industrial areas. However, water problems already present—floods, acid mine drainage, and stream pollution—will become more serious with the further economic development unless remedial measures are taken. Maps in the new report by Schneider and others, published in 1965, show that an average of 150 billion gallons of water leaves the region each day as streamflow. Based on national averages this is enough to meet all water needs for more than six times the present population of the region. This flow, however, is extremely variable from place to place and year to year, resulting in floods and water shortages throughout the region. Ground water is also available in Appalachia for development, with small to moderate amounts available almost everywhere. Large supplies of up to 42 million gallons per day can be obtained from wells in some broad valleys in northern Pennsylvania and in the upper Ohio River Valley.

Ground-water resources of upper Ohio River basin

Abundant supplies of ground water are available for future development within the Allegheny River basin of New York and Pennsylvania, according to a reconnaissance survey of ground-water conditions by Morris Deutsch and J. C. Wallace. The north-western part of the basin is particularly well suited for large-scale ground-water development because it contains aquifers of highly permeable glacial outwash sand and gravel that are underlain by water-yielding sandstones of the Pocono Formation. Large supplies of ground water are also available in glacial and alluvial sediments along the entire course of the Allegheny River from its source to Pittsburgh, Pa., and in the valleys of tributary streams draining the unglaciated area upstream from Tionesta, Pa. South of the Clarion River, serious quality problems brought about by coal mining and interaquifer leakage of salt water will impede future large-scale ground-water developments.

In the Monongahela River drainage basin of West Virginia, Pennsylvania, and Maryland, a reconnaissance survey of ground-water conditions by P. R. Jordan and G. D. Dove showed that moderately large supplies of ground water are available from sandstones of the Pottsville and Allegheny Formations. In the industrial areas along the lower Monongahela and Youghiogheny Rivers near Pittsburgh, Pa., moderately large supplies of ground water can be pumped from the alluvium underlying the flood plains. Unfavorable ground-water conditions preclude large-scale development of ground water in the western third of the basin. New sources of large ground-water supplies in the central third of the basin could be found by extensive test drilling to locate areas of high yield from the lower part of the Conemaugh Formation, and to locate tracts outside their outcrop areas where the Pottsville and Allegheny Formations contain fresh water.

A reconnaissance survey of ground-water conditions in the Kanawha and Little Kanawha River drainage basins of North Carolina, Virginia, and West Virginia by G. D. Dove and J. C. Wallace revealed that the greatest potential for further development of ground water is in the alluvium along the lower reaches of the major streams. Care will be needed in the development of this source to prevent contamination of the water by industrial wastes from the rivers. The study also showed that moderately large ground-water supplies can be obtained from sandstones of the Pottsville and Allegheny Formations in a wide band across the central part of the Kanawha River basin.

Large ground-water supplies in Connecticut River basin

A map of the availability of ground water in the Connecticut River basin in New Hampshire and Vermont, being prepared by D. J. Cederstrom and A. L. Hodges, Jr., working in cooperation with the U.S. Army Corps of Engineers, shows where supplies of ground water of 1 to 5 million gallons per day can be obtained from tributary valleys, and where larger amounts are available at places along the Connecticut River itself. Yields greater than 1 mgd have been reported from individual deep wells. Some shallow gravelly deposits have very high permeability; yields greater than 200 gallons per minute have been obtained from wells less than 40 feet deep. Deep alluvial fill has been found in many tributary valleys. At Woodstock, Vt., for instance, a well which extends 140 feet below present stream level does not reach bedrock. These observations, which suggest that streams in upper New England formerly had a base level considerably below that of the present, indicate that thick water-bearing deposits are more common in these stream valleys than has previously been realized.

Hardness of ground water in Susquehanna River basin

P. R. Seaber and E. F. Hollyday, working in cooperation with the U.S. Army Corps of Engineers, (2-64) have related the hardness of ground water to the lithology of glacial drift, the direction of glacial movement, the topographic setting, and the history of the glacial deposits in the Susquehanna River basin in New York and Pennsylvania. Four distinct areas of ground-water hardness occur in the basin. The hardness is highest in areas underlain by drift containing many limestone erratics and is lowest in areas of few or no such erratics. The hardness data substantiate previous mapping of surficial deposits in the basin.

Natural sand filters protect ground water in Florida and Georgia

Many of the solution cavities and sinkholes in recharge areas of the productive Tertiary and Quaternary limestones in Florida and Georgia are filled with unconsolidated permeable deposits of Pleistocene and Recent age. V. T. Stringfield finds that much of the recharge to the limestones is through these permeable deposits, which are chiefly silica sand and which serve as filters by removing undesirable turbidity and contaminants. Presence of the unconsolidated deposits in solution cavities in the Miami area of Florida also retards the lateral movement of the water in the limestones, and thus helps maintain a fresh-water head and prevent salt-water encroachment.

NEW ENGLAND

Seismic studies aid delineation of potential aquifers

S. J. Pollock reports that seismic investigations in the Assabet River basin, in northeastern Massachusetts, have made possible the delineation of sites which appear to be most favorable for ground-water development. The two most promising areas are in the towns of Hudson and Stow, where beds of saturated sand and gravel appear to be as thick as 50–70 feet. It is estimated that properly constructed wells in these areas would yield up to several hundred gallons per minute; elsewhere in the basin, wells are likely to yield less than 150 gpm.

Ground water in Ten Mile and North Taunton River basins, Massachusetts

More than half the area of the Ten Mile and North Taunton basins, in southeastern Massachusetts, is underlain by alluvium and outwash deposits which are generally less than 100 feet thick, according to J. R. Williams. Most municipal and industrial water supplies are obtained from gravel in these deposits which, in places, yield as much as 1,000 gallons per minute to wells. Industrial and domestic wells in sedimentary rocks of Pennsylvanian age generally yield $\frac{1}{2}$ –20 gpm, but a few yield as much as 200 gpm.

Aquifers in Housatonic River basin, Massachusetts

Studies of the Housatonic River basin in western Massachusetts, by R. F. Norvitch, indicate that ice-contact deposits along the valley walls are locally good sources of ground water. Channel-fill deposits generally have a better ground-water potential in the upper reaches of the tributaries than in the main valley. Because of its cavernous nature the Stockbridge Limestone, which is the bedrock along part of the western side of the basin, may prove to be a highly productive aquifer.

Ground water in southern New Hampshire

Recent studies by J. M. Weigle indicate that a total sustained ground-water yield of 10 million gallons per day may be possible from glacial sand and gravel at places in the towns of Winchester, Swanzey, and Keene, in southwestern New Hampshire. Several times that much may be available where infiltration can be induced from the Ashuelot River and its tributaries. Glacial-lake deposits as thick as 100 feet underlie parts of the Keene basin, however; where the deposits are fine grained their potential yield to wells is considerably lower. At some places in the Ashuelot River basin the yields of individual wells may exceed 1 mgd, but probably most individual wells will yield not more than 0.1

mgd. Preliminary investigation of the Salem-Plaistow area, in southeastern New Hampshire, indicates that ground-water supplies sufficient for domestic needs can be obtained for many years in most towns in that area.

NEW YORK

Low flow of streams in Genesee River basin

Data collected at more than 500 sites in September and October 1964 by B. K. Gilbert, J. C. Kammerer, and F. J. Keller show that the flow of most unregulated streams in the 2,500-square-mile Genesee River basin, in western New York, is between 0.01 and 0.30 cubic feet per second per square mile of drainage area during month-long periods of little or no autumn rainfall. These low flows, which are exceeded more than 95 percent of the time, contrast with average annual flows estimated to be between 0.9 and 1.2 cfs per square mile. Preliminary interpretation suggests that dry-season flow of small streams draining mainly shale and glacial till is less than 0.1 cfs per square mile, whereas higher dry-weather flow characterizes small streams draining mainly glacial deposits of sand and gravel.

Discharge measurements made during the same period indicated that Spring Creek at Mumford, in the northern part of the basin, had a flow more than 40–100 times that of any other small stream in the basin. This stream is fed by limestone springs which evidently are recharged by precipitation over a much larger area than the drainage area (about 1 square mile) defined on topographic maps.

Streams lose flow above waterfalls in central New York

Streams in central New York have been found by I. H. Kantrowitz to lose flow in the reaches immediately above waterfalls. As much as 25 percent of the base-flow discharge of the streams moves under the falls. Drilled wells which tap carbonate-rock aquifers near the waterfalls yield up to 1,400 gallons per minute. The loss of streamflow and the high well yields are attributed to widening of vertical joints in the carbonate rocks near the falls by stream water charged with atmospheric carbon dioxide.

Downward ground-water flow in wells in Niagara County

Temperature profiles, periscope surveys, and water levels in wells studied by F. W. Trainer indicate that downward flow of water between fractures or groups of fractures occurs in many wells in flat-lying Lockport Dolomite in Niagara County, western New York. In areas of numerous and closely spaced wells the downward flow appears to have important

implications for recharge of deeper water-bearing zones in the dolomite, which in turn will lead to modification of the chemical quality, and local contamination, of water in the deeper zones. Under natural conditions the water in the deeper zones is more highly mineralized than that in the upper part of the dolomite.

NEW JERSEY

Ramapo River recharges an alluvial aquifer

Measurements of flow in the Ramapo River, analyzed by John Vecchioli, indicate that the stream is influent along part of its reach in New Jersey. All tributary inflows and the discharge at several places along the main channel were determined under conditions of high, intermediate, low, and very low flow during 1964. The observed seepage loss to an alluvial aquifer ranged from a few to several cubic feet per second, or a large part of the total pumpage from the aquifer (average, about 6 cfs, or 4 million gallons per day).

MARYLAND

Water resources in Patuxent River basin

Study of the Patuxent River basin by Deric O'Bryan and J. W. Crooks reveals that of the 1,960 million gallons per day of water brought to the Patuxent River basin from precipitation, about 64 percent, 1,260 mgd, is lost through evapotranspiration. The remainder, 700 mgd, is discharged from the basin as streamflow. The average flow from the basin ranges from about 630 to 780 mgd. Approximately 1 mgd of the total runoff is contributed by artesian aquifers in the lower part of the basin. Streamflows of 95-percent duration range from 0.03 to 0.15 mgd per square mile. Up to 40 mgd of water is transported out of the basin as public water supply for the Washington, D.C., metropolitan area.

Most of the water in the basin, except for tidal flow, can be used for public supplies and industrial purposes after minor treatment. The water generally is dilute in the Piedmont section of the basin and more highly mineralized in the coastal plain. Objectionable concentrations of iron are found in some ground-water supplies. Restrictions on waste disposal to streams are imposed by the State of Maryland, but objectionable levels of pollution exist in some small reaches of streams. The effect is not cumulative, however, and decreases generally downstream.

Preliminary estimates indicate that about 130,000 tons of sediment is discharged from the basin each year. Average sediment yield in undisturbed areas

of the Piedmont section of the basin is about 150 tons per square mile per year. Sediment yield in the coastal plain averages less than 100 tons per square mile per year. Small tributaries draining areas undergoing active urbanization yield sediment at rates up to 50,000 tons per square mile per year.

Water resources of Dorchester and Talbot Counties

F. K. Mack, R. A. Gardner, and W. E. Webb report that ground water provides most water supplies in Dorchester and Talbot Counties, in eastern Maryland. Most wells tap aquifers of Eocene or younger age at depths less than 600 feet, but a few are completed in aquifers of Late Cretaceous age at depths of about 1,000 feet. Fresh surface-water resources are limited because major streams of the two counties are tidal estuaries of the Chesapeake Bay. Studies of the specific conductance and the chloride content of water in the Choptank and Nanticoke Rivers, at times of lowest runoff, reveal that saline water moves several miles farther upstream than had been indicated by earlier studies.

Buried valley near Salisbury a source of ground water

D. H. Boggess and H. J. Hansen have mapped a 2-mile section of a deep buried channel that is located midway between Salisbury, in eastern Maryland, and the Delaware State line. A series of test holes across this paleochannel indicates that it trends east-west, extends about 200 feet below mean sea level (240 ft below land surface), and ranges in width from about 1,800 feet at the eastern end to about 4,800 feet at the western end. The channel is thought to have been incised into the Miocene surface, and refilled, during late Pleistocene time. The saturated sediments in the paleochannel appear to be a major potential source of good-quality ground water. Specific capacities of wells exceeding 50 gallons per minute per foot of drawdown are not uncommon, and well yields in the range of 2,000–5,000 gpm appear feasible.

WEST VIRGINIA

Water resources in Monongahela River basin

Studies in the Monongahela River basin by E. A. Friel and B. M. Wilmoth reveal that average annual runoff ranges from 0.75 million gallons per day per square mile in the northwestern part of the basin to 2 mgd per sq mi along the southern and southeastern parts of the basin. Over the entire basin, the lowest yearly runoff generally amounts to a little more than half the mean annual runoff. The median 7-day low flow, based on records for the period 1916–61, ranges from less than 0.013 mgd per sq mi in the western

part of the basin to more than 0.13 mgd per sq mi in the headwaters of the Blackwater River. The annual water loss (difference between precipitation and surface runoff) ranges from about 17 to 30 inches.

More than 1,500 square miles in the central part of the basin has good potential for the development of moderately large supplies of ground water. Present development of the aquifers is small; in large areas they are untapped. Small supplies can be obtained in the outcrop areas of all rock units. Most shortage problems occur in the western part of the basin, on hills and ridges underlain by shale. Areas of shallow saline ground water, in which the brine occurs in part naturally and in part as a result of gas-well operations, have been delineated, mostly beneath major stream valleys in the western part of the basin. The effects of coal mining on ground water and on the low flow of streams are relatively small except locally near mines, where low flows are increased and the quality of the water has deteriorated.

SOUTH CAROLINA

Ground-water system beneath coastal-plain sediments

An artesian system defined by G. E. Siple consists of fractures in crystalline rock that lies beneath 1,000 feet of coastal-plain sediments. The upper confining layer is a bed of saprolite 40 to 60 feet thick. Natural discharge is probably to streams in the boundary region of the Coastal Plain and Piedmont provinces where the confining bed has been partially or completely removed. The dissolved-helium content of water from the rock wells appears to vary directly with the proximity of the well to these discharge areas.

GEORGIA

Zone of low permeability in Glynn County

Water-level behavior resulting from occasional reduction in industrial pumping indicates a zone of low permeability between the Brunswick Peninsula and St. Simons and Jekyll Islands, in southeastern Georgia. This zone, as delineated by D. O. Gregg, was further substantiated by a map showing water-level declines from October 1962 to December 1963 caused by an increase of 36.7 million gallons per day in industrial pumping.

Ground water in Rockdale County

Ground-water studies by M. J. McCollum show that Rockdale County, in north-central Georgia, is underlain by weathered metamorphic rocks of Precambrian (?) age that have been intruded by granite

and by pegmatite and diabase dikes. Ground water in these rocks is more readily available where weathering is pronounced. The quality of the water is directly related to the composition of the rocks in which it occurs; thus, the water of best quality occurs in the Lithonia Gneiss of Watson (1902)¹⁶ and that with the highest iron concentration occurs in amphibolitic gneiss.

Lower part of aquifer in Cook County poorly productive

C. W. Sever reports that the drilling of a 870-foot test well at Adel, in Cook County, south-central Georgia, disclosed that the lower part of the principal artesian aquifer system between depths of about 440 and 850 feet consists of dense dolomitic limestone that yields practically no water.

Ground water in Floyd and Polk Counties

A ground-water investigation in the Paleozoic area of Georgia by Charles Cressler shows that in Polk County, in northwestern Georgia, yields of more than 1,000 gallons per minute can be obtained from wells in limestone. Limestone is less abundant in adjacent Floyd County, and ground-water conditions are less favorable, but yields up to 500 gpm are probable in some places. Eleven unused springs in the 2 counties discharge a total of more than 28 million gallons per day. The discharge of individual springs ranges from 0.5 to 15 mgd. The spring water is moderately hard and has a low iron content, making it suitable for many industrial uses.

FLORIDA

Ground-water circulation in Marion County, Fla.

Study of the quality of water in large springs in Marion County and vicinity, in north-central Florida, suggests to F. N. Visher that most of the circulation of ground water is through large solution openings in the upper few tens of feet of the carbonate rocks. The remainder of the circulation is through evaporite beds deep in the Floridan aquifer, where the solution of salt and other evaporites has produced a zone of very high permeability. The rest of the carbonate-rock sequence is less permeable and, except in local areas, is undergoing little solution at this time. In discharge areas, waters ascending from the evaporite beds mix with the water in the overlying carbonate rocks, resulting in a deterioration of the quality of the water in these areas. Water from evaporite beds is thought to be the principal source of dissolved material in poor-quality waters in much of the State.

¹⁶ T. L. Watson, 1902, A preliminary report on a part of the granites and gneisses of Georgia: Georgia Geol. Survey Bull. 9 A, p. 54, 125-127.

Water budget of mid-Gulf Coast basins

Preliminary estimates by R. N. Cherry of evapotranspiration, surface runoff, and ground-water runoff from the mid-Gulf Coast basins in the west-central coastal area are 35, 15, and 5 inches, respectively. The Floridan aquifer, which is composed of about 1,000 feet of limestone and dolomite ranging in age from Eocene to Miocene, underlies the basin. In 1963, St. Petersburg pumped about 22 million gallons per day from this aquifer.

Floridan aquifer tested at Key Largo

Information obtained by H. J. McCoy from a 1,070-foot well at the north end of Key Largo, drilled into the Floridan aquifer, the principal artesian aquifer of Florida, showed that the piezometric surface was 38 feet above mean sea level (or more than 15 feet higher than the level at which it had previously been). The water, although brackish, was relatively fresh, considering the distance offshore. The artesian water is being blended with fresher water piped from the Florida mainland and is used for golf-course irrigation. The economic feasibility of using conversion units to produce potable water from the brackish water in quantity, to supply the Florida Keys, is under study by the Florida Aqueduct Commission.

Aquifers in Myakka River basin

B. F. Joyner, Horace Sutcliffe, Jr., and J. D. Warren report that ground-water supplies in the Myakka River basin in southwestern Florida are obtained from the Hawthorn Formation and the Tampa, Suwannee, and Avon Park Limestones. The Hawthorn (youngest) Formation generally yields 50 gallons per minute or less of water that is of good quality except for fluoride, which ranges in concentration from 1.5 to 3 parts per million. The Tampa Limestone yields up to 1,000 gpm of water of good quality except that the fluoride locally exceeds 2.0 ppm. Wells which tap the Suwannee Limestone yield up to 1,000 gpm of water having dissolved-solids concentration exceeding 800 ppm and a sulfate concentration exceeding 500 ppm; those in the Avon Park Limestone yield 1,000–2,500 gpm of water having a dissolved-solids concentration of 850–2,000 ppm and a sulfate concentration of 500–1,500 ppm.

Water levels recover 200 feet near Panama City

J. B. Foster reports that water levels in the Floridan aquifer near Panama City, in western Florida, recovered a maximum of about 200 feet during 1964 as a result of abandonment of an industrial well

field. The well field had been in operation for 34 years, and the rate of pumping during the 5 years preceding 1964 was more than 11 million gallons per day.

Underseepage of Lake Okeechobee dikes caused by excavations

Studies by F. W. Meyer around the southern half of Lake Okeechobee indicate that seepage beneath the Hoover Dike is significant in the area between Clewiston and Moore Haven. The seepage is related to the excavation of borrow canals on either side of the dike. Increases in seepage can be expected all along the southern half of Lake Okeechobee if drainage canals in the adjacent agricultural areas are excavated parallel to and adjacent to the dike.

Wide range in flow and dissolved solids of Shark River Slough

Preliminary hydrologic studies of the Shark River Slough, in southern Florida, by B. F. Hartwell, Howard Klein, and F. B. Joyner (1–64) revealed wide ranges, from wet to dry years, in surface runoff and in dissolved solids. During the 10-year period prior to October 1962 median monthly flows ranged from 21 to 1,020 cubic feet per second; the average total yearly flow was about 250,000 acre-feet. The content of dissolved solids (about 500 parts per million) in areas of fresh ground water is more than three times greater during dry periods than during wet periods. The increase during dry weather, which is chiefly in calcium and carbonate, is thought to reflect lesser dilution of the ground water by recharge during dry periods than during wet ones.

CARIBBEAN REGION

Ground water overdeveloped in Ponce area, Puerto Rico

Study of the Ponce area, on the south coast of Puerto Rico, by N. E. McClymonds shows that the ground water is overdeveloped. Sustained pumping of wells as far as 2 miles from the coast has caused water levels to decline 10 to 20 feet below sea level in the major aquifer, a cavernous limestone. A layer of poorly permeable alluvium about 50 feet thick, overlying the limestone, seems to be holding sea water from encroaching inland to the area of depressed water level except southwest of Ponce, where there is evidence of encroachment. By the end of 1964, a drought year, water levels in many wells were down to the level of the pumps.

Surface-water supplies are extensively developed in the Ponce area, but considerable additional development is possible if storage reservoirs can be provided.

Water resources of Virgin Islands

D. G. Jordan and O. J. Cosner found from test drilling in the Virgin Islands National Park, St. John, that 1,000–15,000 gallons per day of fresh ground water are available from fractured volcanic tuffs and breccias. In coastal areas of the island, however, the threat of sea-water encroachment will impede development of ground-water supplies.

Small but significant quantities of fresh water have been discovered in St. Thomas as the result of test drilling. Three areas—on the north slopes of the Island, in Tutu Valley, and at the race track—warrant development of small public supplies. The total minimum yield from these areas is estimated to be 0.8 million gallons per day.

Development of ground water for public supplies on the Island of St. Croix is limited largely to alluvial and limestone aquifers in an area of about 6 square miles along the south-central coast. These aquifers will produce several million gallons per day of potable water, but they generally contain brackish water at depths of 40 to 100 feet below sea level. In areas underlain by fractured volcanic rock, supplies of from 1–10 gpm generally can be obtained. In nearly a third of the island, ground water is saline or is not present in usable quantities.

Surface-water runoff in the Virgin Islands appears to average less than 5 percent of the annual rainfall, or about 2 inches. Runoff from even large storms, much less than would be expected in an area where the slope of the land surface generally exceeds 200 feet in 1,000 feet, reflects the permeable rocks and deposits at and near the land surface.

MIDCONTINENT AREA

In the midcontinent area, water-resources investigations of the U.S. Geological Survey are directed toward solution of problems associated with distribution and increasing development of the available water. These studies include (1) appraisal of the available water resources in terms of location, quantity, and quality; (2) investigation of specific problems to provide hydrologically sound guidelines for their solution; and (3) facilitation of water-management practices by improving our understanding of the hydrologic environment.

In a few studies, previously unknown ground-water sources have been discovered that add significantly to the available supply. For example, fresh ground water has been found at depths as great as 2,600 feet below sea level in Pointe Coupee Parish, La. Other studies bear on declining ground-water

levels, deficient streamflow, and local overdevelopment of water sources. Thus, the need for an alternate and supplemental supply for the city of New Orleans has led to exploration and confirmation of large reserves of ground water in deep aquifers underlying Lake Pontchartrain. New alltime low ground-water levels in Madison and Shelby Counties, Tenn., followed increase in pumpage. Continuing deficiency of streamflow in part of the area emphasizes the need for water-shortage facilities and coordinated use of ground- and surface-water resources.

Significant results of these and other studies are reported below.

INTERSTATE STUDIES

Aquifers correlated in Louisiana and Texas

As a part of the compilation and analysis of data for the construction of an analog model of the aquifers in the Gulf Coast area of southeastern Texas and southwestern Louisiana, A. N. Turcan, Jr., Chabot Kilburn, and Roger Baker remapped the boundary between the Evangeline (Pliocene) and Chicot (Pleistocene) aquifers and correlated them between Houston, Tex., and the Atchafalaya River in Louisiana. In this interstate area the name "Jasper aquifer" has been proposed for deeper sediments of Miocene age that form an aquifer which is a mappable hydrologic unit, is composed of alternating beds of sand and clay, and contains fresh water in parts of the project area.

Water resources of Mississippi embayment region

R. L. Hosman, T. W. Lambert, and A. T. Long find that Tertiary aquifers are important sources of ground-water supplies in a total area of about 75,000 square miles, or about three-fourths of the entire Mississippi embayment. As a group, these aquifers are capable of sustaining much larger withdrawals than the present rate, which is in excess of 500 million gallons per day. Some of the aquifers are not tapped by wells, in large areas where they are known to contain usable water, because supplies are available from shallower aquifers. They will become increasingly important in these areas when eventually they are utilized to avoid or delay depletion of the shallower supplies. Most of the aquifers are in the Wilcox and Claiborne Groups, and the largest total withdrawal is from Claiborne aquifers. The most extensively developed is the Sparta Sand, which yields about 25 percent of the total withdrawals from Tertiary aquifers. The piezometric surface in the Sparta Sand shows several large

cones of depression caused by large withdrawals. A few of these cones have coalesced, and others seem to be coalescing and will probably form one large depression eventually. Use of deeper aquifers could help alleviate this situation in much of the area.

M. S. Hines, M. E. Janson, and P. R. Speer report that streams in the Quaternary alluvium along the Malden-Kennett Prairie and the west side of Sikestone Ridge, most of which are in dredged channels, have higher base flows than streams in any other part of the alluvium in the Mississippi embayment section of Missouri and northern Arkansas. Streams in the alluvium between lat 35° N. and 36° N. have the lowest base flows in the area studied. Streams that originate in Paleozoic rocks outside the embayment have well-sustained base flows except where the streams have not eroded deeply enough to expose productive Paleozoic aquifers. Sands of the Wilcox Group or Formation and the McNairy Sand in Crowleys Ridge are important contributors to the base flow of streams draining the east side of the ridge north of Marmaduke, Ark.

Upper Mississippi River basin

Reconnaissance studies by P. G. Olcott, in Missouri and Illinois, indicate that unconsolidated glacial-drift and alluvial aquifers in buried bedrock valleys beneath the Mississippi, Kaskaskia, and Cache Rivers and the headwater diversion canal generally yield from 400 to 1,000 gallons per minute of water to wells and are the aquifers of greatest yield in the upper Mississippi River basin south of the Illinois River. Bedrock aquifers consisting largely of dolomites of Cambrian Devonian age generally yield up to 400 gpm of water to wells in the area west of the Mississippi River. An abundant supply of ground water, greater than the present demand, is commonly available from these aquifers.

Ohio River valley

The Ohio River valley between Pittsburgh, Pa., and Marietta, Ohio, contains thick deposits of permeable gravel and sand, and the river provides a perennial source of recharge for development of large ground-water supplies. P. R. Jordan, in a reconnaissance survey, also found conditions favorable for ground-water development in the sandstones of the Mississippian System in the northern parts of the Beaver and Mohoning River basins. Most of the ground water in this area has mineral concentrations that are within limits recommended for drinking water, although softening and iron removal are generally desirable. The effect of gross contamina-

tion of surface water in the basin on the quality of the ground water cannot be assessed without additional data.

MINNESOTA

Statewide study of ground-water quality

A study of the chemical quality of water in the major aquifers in Minnesota by M. L. Maderak indicates that water in the Jordan Sandstone tends to be more uniform in chemical type from area to area and from year to year than water in the other Paleozoic aquifers or in the glacial drift. Water in any one Paleozoic aquifer generally is of a distinct type, whereas water in the glacial-drift aquifers is of several different types. Except for extreme southwestern Minnesota, the water in all aquifers tends to have a higher concentration of dissolved constituents in the southern part than in the central and northern parts of the State.

WISCONSIN

Water in Wisconsin River valley

In the lower part of the Wisconsin River valley, additional large-capacity wells yielding 1,000 gallons per minute or more of good-quality water can be constructed in thick alluvial sand and gravel, or in the underlying extensive sandstone aquifer, according to studies by L. J. Hamilton and T. R. Dosch. The flow of the lower Wisconsin River, stabilized by an upstream network of 23 reservoirs, is an additional large potential source of good-quality water. Decreased streamflow and declines of ground-water levels in aquifers during 1962-64 are attributed to deficiencies in precipitation rather than to increased pumpage of ground water. Withdrawal of ground water can be greatly increased without seriously affecting either streamflow or ground-water levels.

Ground water in Racine and Kenosha Counties

Piezometric mapping of the deep sandstone aquifer in Racine and Kenosha Counties, in southeastern Wisconsin, by R. D. Hutchinson indicates that the ground-water divide between the Milwaukee and Chicago cones of depression extends from east to west through central Kenosha County. Water levels in wells tapping this important aquifer continue to drop 4 to 7 feet per year. Brackish water has been found in wells more than 2,000 feet deep in Racine. Mapping of water levels in the shallow Niagara aquifer indicates that the principal recharge area is near the Fox River valley, and that the water is confined in more than half the area by clay till that reaches a thickness of nearly 300 feet.

The quality of the water is generally acceptable for most uses.

MICHIGAN

Drought in southern Michigan

Streamflow records show that runoff for the 1964 water year in the southern half of the Lower Peninsula was generally the lowest since streamflow records began at the turn of the century. Mean monthly levels of Lakes Michigan and Huron were at a record low for each month of the 1964 calendar year, with the alltime record low occurring in March. Levels in 5 of 11 lakes that have been gaged for 15 years or more declined to new lows. Water levels in representative observation wells were at low or record-low stages in 1964. A protracted deficiency in precipitation is the principal cause of the drought. At Lansing the deficiency for the period January 1962 to June 1964 amounted to nearly 21 inches.

Water resources in Marquette iron range

Water-resources investigations in the Marquette iron range area, in northwestern Michigan, indicate total water supply ample for present and expected needs. Only about 1½ inches of the estimated annual average runoff (14 inches) is used at present (1964), and most of the water now used is available and fit for reuse. About 190 cubic feet per second is available in the principal streams of the 610-square-mile area during 90 percent of the time. With storage, these streams could supply a firm flow of about 450 cfs. The ground-water potential is best in the glacial-outwash deposits. In the extensive outwash plain south and east of Goose Lake, for example, it is estimated that at least 15 cfs could be developed from the northern part of the outwash plain alone. The quality of the surface and ground waters is generally good to excellent.

Suspended-sediment concentrations as high as 45,400 parts per million have been measured in the tailings discharge from the Humboldt mine, Marquette County, where the discharge enters the McKinnon Lake settling basin, according to Russell F. Flint. The suspended material had a specific gravity of 3.07; it was 33 percent sand, 60 percent silt, and 7 percent clay. Suspended-sediment concentrations in the outflow from the settling basin, measured a quarter of a mile downstream, were observed to be 92 ppm or less during 1961–64. Measurements of suspended sediment at the gaging station on the Black River near Republic, about 7 miles downstream from McKinnon Lake, indicated no significant increases in concentration or sediment yield during the period.

OHIO

Preglacial drainage in Miami River valley

A. M. Spieker has mapped the bedrock surface in the Great Miami River valley, between Middletown and Hamilton in southwestern Ohio, largely on the basis of the results of a seismic-refraction survey, and has found two buried valleys separated by a bedrock high. The older valley, 250 to 300 feet deep and extending through Trenton, Busenbark, and Overpeck, is believed to have been formed during an interglacial period prior to the onset of the Illinoian Glaciation. This valley is 1½ to 2 miles wide. The younger valley, which underlies the present course of the Great Miami River, is only about 100 feet deep and is believed to have been formed when the Wisconsin ice sheet blocked the older valley, situated about 2 miles to the northwest. The younger valley is but half a mile wide. Both valleys are filled with permeable sand and gravel and are considered to be excellent sources of ground water.

Cuyahoga River water quality in Cleveland area

Water quality in the navigable portion of the Cuyahoga River at Cleveland varies greatly within the 25-foot depth of the river as well as longitudinally and in cross section. According to recent work by M. E. Schroeder and C. R. Collier, stratification of the water is caused by industrial and municipal wastes but is complicated by seiching of Lake Erie. Stratification of the water is greatest in the lower 2-mile reach of the river. For example, in one set of measurements at the Center Street bridge, 4,500 feet upstream from the mouth of the stream, the conductance ranged from 1,560 micromhos at the surface to 530 micromhos at a depth of 25 feet; the corresponding range in temperature was 76°F to 62°F. In the upper reach of the navigable channel (2 to 6 miles from the mouth), however, conductance at that time ranged from 1,510 to 1,240 micromhos and temperature from 84°F to 76°F.

Ground water in Berea Sandstone

A regional study of the Berea Sandstone by J. J. Rau indicates that it consistently yields more water to wells than do the other consolidated-rock units of Devonian and Mississippian age in northeastern Ohio. The Berea Sandstone is an important aquifer in parts of Trumbull, Portage, Ashtabula, Geauga, Lake, Cuyahoga, Summit, and Lorain Counties, where its thickness ranges from a few feet to 235 feet and averages about 50 feet. The average specific capacity in 344 wells is 1.5 gallons per minute per foot of drawdown. The highest transmissibility

values were recorded in southeastern Ashtabula County and central Trumbull County, to the northeast, and the lowest values in Summit and Lorain Counties to the southwest. The average depth to which wells penetrate the aquifer is 30 feet, and the average yield of these wells as reported by drillers is 14 gpm.

Ground water in Muskingum and Hocking River basins

A reconnaissance study by G. D. Dove shows the Muskingum River basin in southeastern Ohio to be rich in ground water. Wells drilled into outwash deposits along the Walhonding and Tuscarawas Rivers, the principal tributaries of the Muskingum, yield as much as 1,200 and 2,000 gallons per minute, respectively, and as much as 1,500 gpm is obtained from the deposits along the course of the Muskingum River. In the central and eastern parts of the Muskingum River basin, away from the stream valleys, the principal sources of ground water are the sandstones of Pennsylvanian age. Most ground water in the basin is not highly mineralized and may be used for most purposes with little treatment.

Large supplies of ground water are also available along the course of the Hocking River. In the glaciated northwestern part of the Hocking River basin, near Lancaster, individual wells yield more than 500 gpm, but south of Lancaster the outwash deposits are thinner, less permeable, and consequently yield less water.

INDIANA

Water resources of Delaware County

R. E. Hoggatt and J. D. Hunn find that water supplies now available in Delaware County, in east-central Indiana, will be adequate for most foreseeable agricultural, municipal, and industrial purposes if developed properly. Most streams in the south half of the county, regardless of size, can be depended on to yield more than 0.013 million gallons per day per square mile 90 percent of the time. On the other hand many streams in the north half of Delaware County having less than 10 square miles of drainage area, including the headwaters of larger streams, go dry frequently. The principal sources of ground water are sand and gravel deposits of Pleistocene age and the upper 100 feet of the dolomite bedrock of Silurian age. The areas of greatest potential are in the south-central and southeastern parts of the county. Each aquifer is capable of yielding as much as 500 gallons per minute to individual wells in these areas.

MISSOURI

Surface-water quality

The overall quality of Missouri surface water is good, according to C. T. Taylor (1-64) in a statewide study of the quality of water. In general, the surface water is hard and moderately mineralized, but it may be used with limited treatment for most purposes. For irrigation purposes, the water is classified as a low-sodium water with medium to low salinity suitable for the irrigation of most plants without special practices for salinity control.

The most highly mineralized water is found in the northern glaciated plains and the western plains. Here the average dissolved-solids content found during a 12-month study was 274 parts per million, and the average hardness was 194 ppm. A distinctive feature of surface-water quality in this area is a relatively high sulfate content (average 59 ppm). The sulfate is believed to have been derived from shale and coal deposits in the bedrock, which is predominantly of Pennsylvanian age.

The western Ozark region, underlain by relatively pure Mississippian limestone, comprises an area in which a high (average 8.2) calcium-magnesium ratio is typical of the surface water. The average dissolved-solids content was 200 ppm, and the average hardness was 165 ppm. On the other hand, in the central and eastern Ozark highlands, a region underlain by Cambrian and Ordovician dolomite, a low calcium-magnesium ratio (average 1.0) is typical of the surface water. The average dissolved-solids content here was 168 ppm, the lowest in the State; the average hardness was 157 ppm.

Drainage anomalies

The basins of North Fork River and Bryant Creek, which adjoin one another in Douglas and Ozark Counties in south-central Missouri, have similar drainage areas, basin shapes, topography, and climate. However, the base flow of the lower reach of North Fork River is about twice that of Bryant Creek. E. J. Harvey and John Skelton find that karst features are well developed in the basin of North Fork River but poorly developed in that of Bryant Creek, and attribute the higher base flow to a more effective underground drainage system. A system of springs increases the base flow of North Fork River about 150 percent in a 500-foot reach of the stream in Ozark County.

ARKANSAS

Low flow of Arkansas streams

A study by M. S. Hines reveals the duration of

daily flow and the frequency of low flows at 65 gaging stations and 97 partial-record sites in Arkansas. Major streams such as the White, Saline, Ouachita, and Black Rivers, and tributary streams in the northeastern part of the coastal plain, in the Springfield-Salem Plateau, in the Novaculite uplift, and in the Athens-Piedmont Plateau, are most likely to be dependable sources of water supply during dry periods. Tributary streams in the central and southern part of the coastal plain, in the Boston Mountains, in the Arkansas Valley, and in the Fourche Mountains do not have dependable flow during dry periods, and many of them often go dry.

Ground water in Jackson and Independence Counties

A water-resources study of Jackson and Independence Counties, in northeastern Arkansas, by D. R. Albin and M. S. Hines has shown that as much as 2,000 gallons per minute can be obtained from individual wells in alluvial deposits but that yields from other deposits seldom will exceed 50 gpm. Ground water generally is available throughout both counties, but in the area within the Ozark Plateaus users requiring more than about 50,000 gallons of water per day generally will be limited to supplies developed from streams.

KENTUCKY

Fresh-water-saline-water interface

Mapping of the fresh-water-saline-water interface by H. T. Hopkins, as part of the larger study of saline water in Kentucky, shows that the ground-water flow pattern in the limestones of the Mississippian Plateau and Blue Grass regions, in central Kentucky, is controlled by the relatively deeply incised streams and that the interface is lowest below these streams. In the western and eastern coal fields, in contrast, and in the Jackson Purchase region, in westernmost Kentucky, the interface reflects structural lows and highs.

Water supply for Louisville is adequate

The water supply for the Louisville area appears to be adequate for many years to come, according to E. A. Bell. The principal sources of supply are the Ohio River and ground water from deposits of sand and gravel adjacent to the river. The minimum daily flow on record for the Ohio River was 1.3 billion gallons in 1930, more than 6 times the average daily use of 211 million gallons in 1962. Of the fresh water used in 1962, 40 mgd was from the ground and 171 mgd from the Ohio River. Public-supply systems furnished 91.3 mgd, and industries and commercial establishments supplied 119.7 mgd for their own use

from private systems. The principal water problems are problems of management, and are associated with the distribution of supplies, the quality of water, drainage, and waste disposal.

Ground-water movement in Mammoth Cave area

Continued investigation of the ground-water resources of the Mammoth Cave area by R. V. Cushman has furnished greater detail on the pattern of circulation in the cavernous limestones. The available information suggests that the gradient of the water table is controlled in part by the structure of the underlying Paleozoic rocks, and particularly the Chattanooga Shale, which forms the base of the zone of unconfined water. A marked steepening of the gradient of the water table in the vicinity of the Dripping Springs escarpment coincides with a monoclinical fold in the underlying bedrock. The regional gradient and direction of water movement in the carbonate rocks are from the Pennyroyal Plain northwest to the Green River. The pattern of the subsurface drainage is dendritic, the major part of the flow being concentrated in several large conduits as it nears points of discharge into the Green River.

Eocene aquifers in Jackson Purchase region

Geologic and hydrologic studies in the Jackson Purchase region, in westernmost Kentucky, by R. W. Davis, T. W. Lambert, and A. J. Hansen, Jr., show that Eocene aquifers of probable Claiborne age are saturated with water of good quality above the altitudes of most of the local streams and maintain perennial flows of water in Mayfield Creek downstream from Mayfield, Obion Creek downstream from Pryorsburg, and Bayou du Chien downstream from the vicinity of Water Valley. Most wells which tap Eocene aquifers yield more than enough water for domestic use: properly constructed wells should yield more than 100 gallons per minute.

Water in alluvium of Ohio River valley

An investigation by J. T. Gallaher and W. E. Price, Jr., shows the existence of a large perennially available supply of ground water in the alluvium of the Ohio River valley in Kentucky. Yields as great as 1,000 gallons per minute are obtained by induced infiltration from wells penetrating the coarse basal alluvium near the river. Poorest ground-water conditions are found where the relatively coarse Ohio Valley alluvium has been replaced by or mixed with the clay, silt, and fine sand of tributary streams, although in some of those areas there is a coarse-

grained basal deposit capable of yielding more than 200 gpm to properly developed wells.

TENNESSEE

New low ground-water levels

New alltime-low water levels recorded in Madison and Shelby Counties, in southwestern Tennessee, reflect increasing pumpage at Jackson and Memphis, respectively, according to R. H. Bingham. An observation well at Crosstown, 25 miles northeast of the center of withdrawal at Memphis, recorded effects of Memphis pumpage for the first time in 1964. The water level in this well reached a new alltime low of 199.33 feet below land surface and 0.77 feet below the previous low.

Ground-water development in Memphis area

Analysis of hydrologic data from the new Lichterman well field, in the "500-foot" sand aquifer at Memphis, by D. J. Nyman shows that the water-level gradient developed after the well field goes into operation will cause part of the aquifer within and to the east of the field to change from an artesian condition to either a semiartesian or a water-table condition. Because of the great variation in thickness of the capping clay, each of these three hydraulic conditions will exist at some places within a radius of a few miles of the center of pumping.

The natural isolation of the "500-foot" sand in the Memphis area from overlying and underlying aquifers affords a high degree of protection for the water, which is of generally excellent quality, but the possibility of recharge through adjacent aquifers cannot be ignored. Studies by E. A. Bell indicate that slight changes in chemical quality of the water in the "500-foot" sand, particularly increases in hardness and dissolved solids, correspond to changes in hydrologic conditions that reflect man's activities. The large extent of the "500-foot" sand and the slow rate of migration of water through this sand in the Memphis area are favorable for identifying potential changes of chemical quality of the water before it reaches points of discharge.

ALABAMA

Water-quality anomaly in Coker Formation

A study of the quality of artesian ground water in the Coker Formation of Late Cretaceous age in Tuscaloosa County, in west-central Alabama, by J. G. Newton shows the anomalous occurrence of ground water containing as much as 2,000 parts per million of chloride and 900 ppm hardness. The mineralized water is found in a well-defined belt 2 to 3 miles

wide that trends southwestward, parallel to the regional dip of the Cretaceous formations and to the trend of major faults and folds in the Paleozoic rocks of eastern Tuscaloosa County.

Surface water in southwestern Alabama

L. B. Peirce and S. M. Rogers have outlined present and probable future surface-water conditions in an area of 15 counties in southwestern Alabama.

The average annual rainfall in the area studied ranges from 51 to 65 inches, and annual runoff ranges from 15 to 28 inches. These values reflect the considerable range in climatic and physiographic conditions in the area. On the average, evapotranspiration consumes about two-thirds of the annual rainfall. The average flow of the streams ranges from 1 to 2 cubic feet per second per square mile of surface drainage area. Low flows are considerably more variable, however, reflecting the diverse lithology and structure of coastal-plain aquifers. The low flow for 7 consecutive days that occurs every other year, on the average, ranges from no flow to more than 1 cfs per sq mi. Streams draining permeable sand and gravel in the coastal counties have the highest dry-weather flow; streams draining impermeable chalk of the Prairie Belt frequently cease flowing. Floods are likely in any month but are most common from February through April. Floods on the Alabama and Tombigbee Rivers rise and fall slowly and may last from 1 to 3 weeks or longer.

The stream waters are generally of good chemical quality; they are soft and have low dissolved-solid content. With few exceptions the waters of the area are of a quality suitable for most industrial or domestic uses. Stream temperatures range from low values of 38°–44°F in January and February to high values of 80°–92°F in July and August. The Alabama and Tombigbee Rivers furnish an abundant year-round supply of water at temperatures suitable for most industrial purposes.

MISSISSIPPI

Deepest fresh ground water in Mississippi

The deepest fresh ground water in Mississippi is indicated at 3,500 feet by electric logs in Smith County, in south-central Mississippi, according to Roy Newcome, Jr. Deep-lying fresh-water aquifers constitute potential sources of large water supplies in most parts of the Pascagoula River basin. The Wilcox Group in the northern part of the basin is virtually untapped. Miocene formations in the southern part of the basin contain thick aquifers at depths greater than those of present wells.

Ground water in delta area

B. E. Wasson has concluded, on the basis of several aquifer tests in the Mississippi River alluvium of the delta area in northwestern Mississippi, that very large quantities of water can be developed from shallow wells (100–150 feet deep) in most of the 22 counties of the area. The aquifer is readily recharged by precipitation (which totals about 52 inches per year) and by perennial streams that cross the delta. The combination of wide areal occurrence, ability to transmit water, and high recharge potential make the Mississippi River alluvium one of the major aquifers in the Nation.

Fresh water in Cretaceous aquifers

Investigations of Cretaceous aquifers in northeastern Mississippi by E. H. Boswell have shown that large supplies of fresh ground water are available in most of an area of about 3,500 square miles delineated on previous maps¹⁷ as being underlain by highly mineralized water. Predictions based on interpretation of electric logs and hydrologic data have been confirmed by recent drilling which has resulted in the development of five new public water supplies and construction of wells in recreational areas; several additional public water-supply wells are being drilled or planned. The aquifers being developed are in sand and gravel of the Tuscaloosa Group. Wells range in depth from about 1,800 feet to more than 2,400 feet and yield several hundred gallons per minute. The dissolved-solids content of the water ranges from less than 500 parts per million to nearly 1,000 ppm. The aquifers contain fresh water considerably farther downdip than was previously thought possible, and at depths of more than 3,000 feet below the surface. The part of Mississippi in which aquifers yield less than 50 gpm or contain water having a dissolved-solids content higher than 2,000 ppm is now restricted to an area of less than 200 square miles in extreme northeastern Lauderdale and southeastern Kemper Counties, in the east-central part of the State.

High-production wells in Hancock County

Studies by Roy Newcome, Jr., and others reveal that water wells capable of yielding 5,000 gallons per minute or more each can be developed in deeplying aquifers in Hancock County. Recently completed wells at the National Aeronautics and Space Administration rocket-test facility are 1,500–1,900 feet deep and have been tested at discharge rates of

1,100–5,000 gpm. Static water levels stand 80 to 104 feet above the land surface. The temperature of water in the deepest aquifer is 100°F. Pumping tests reveal aquifer transmissibilities as great as 200,000 gallons per day per foot. Dissolved-solids content increases with depth but is less than 350 parts per million in all deep aquifers sampled.

LOUISIANA

Aquifers in Pointe Coupee Parish

Fresh ground water is available in southern Pointe Coupee Parish in south-central Louisiana to depths of about 2,600 feet below sea level, and in as many as 10 to 12 separate aquifers at one location. However, preliminary data reported by M. D. Winner, Jr., indicate that heavy withdrawals of ground water in the nearby Baton Rouge area are beginning to affect water levels in certain aquifers in southern Pointe Coupee Parish that are hydraulically connected with aquifers in the Baton Rouge area. Water-level declines of about 2 to 3 feet per year in excess of local pumping effects are being noticed in the southeastern corner of the parish.

High-chloride water in alluvium

Preliminary study of the Mississippi River alluvial aquifer in Franklin Parish, in northeastern Louisiana, by J. R. Marie indicates the presence of a zone of chloride-contaminated water within the aquifer. The chloride zone trends north-south and is 35 miles long; it is 2 to 4 miles wide and has a core of highly mineralized water flanked by zones of progressively less mineralized water. The greatest recorded chloride concentration is 4,090 parts per million.

Methane in ground water in Louisiana

A. H. Harder, and W. R. Holden of Louisiana State University (1–65), have described methods of detecting methane gas and determining its concentration in well water; and Harder, H. M. Whitman, and S. M. Rogers (2–65) have mapped the distribution of methane in the Chicot and Evangeline aquifers in southwestern Louisiana and have studied the potential explosion hazard of the gas. In most of the area studied the ground water now pumped from wells contains less than 1 part per million of methane. However, the concentrations found range from 0.0 to 40 ppm. Organic matter in the aquifers is believed to be the principal source of the methane, but some of it may have migrated from oil and gas sands near the producing fields. Theoretically, pumping water that contains more than 1.1 ppm methane at 74°F and at atmospheric pressure into poorly ventilated air spaces constitutes an explosion hazard.

¹⁷ C. L. McGuinness, 1963, The role of ground water in the national water situation: U.S. Geol. Survey Water-Supply Paper 1800, p. 445–446 and pl. 1.

Aeration of the water before use and adequate ventilation of air spaces in which the water is stored and used should eliminate the possibility of explosions of the methane.

ROCKY MOUNTAIN AREA

Inadequate distribution and supply continued to be the most serious water problems in the Rocky Mountain area, which includes all the States encompassing the Rocky Mountains, except Idaho, and the plains States from North Dakota to Texas. The most critical areas are those of large withdrawals of ground water for use in irrigation and, locally, for use in rapidly expanding municipalities.

The quality of both surface water and ground water has deteriorated locally because of (1) leaching of salts from the soils by irrigation, (2) induced flow from areas of high mineralization to areas of lower mineralization as a result of increased water-table gradients caused by heavy pumping, and (3) contamination by oil-field brines and other pollutants.

Increased competition between users of surface water and ground water in river valleys where the shallow ground water and the river constitute a single water resource has led to detailed studies of the entire water system in some basins. The data collected are now being used in the construction of analog models that simulate the system and facilitate more intelligent management of the supply.

Some of the significant results of investigations by the U.S. Geological Survey of water supplies in the Rocky Mountain area are described in the following section.

INTERSTATE STUDIES

Aquifer defined in Colorado River valley

A newly recognized aquifer of possible economic significance has been defined broadly by geologic mapping and test drilling in the Parker-Blythe-Cibola area, along the Colorado River in southwestern Arizona and adjacent California. Although a few wells near Parker have obtained part of their supply from this aquifer for many years, not until a recent study by D. G. Metzger (p. C203-C205) was the aquifer differentiated from much younger deposits and demonstrated to extend throughout the area. The aquifer, a conglomerate of possible Miocene age, crops out in the vicinities of Parker and Cibola, but is buried throughout most of the area beneath estuarine deposits of possible Pliocene age. Further tests are needed to estimate potential well yields and to define the water quality.

Suspended sediment in Niobrara River

Analysis by M. L. Maderak and H. D. Stephens of streamflow and suspended-sediment data from the Niobrara River basin, in Nebraska, South Dakota, and Wyoming, indicates that 98 percent of the time the mean daily suspended-sediment concentration of the main stem ranges from 14 to 500 parts per million; 50 percent of the time the concentration is 79-1,500 ppm; and 0.2 percent of the time the concentration is 5,000-6,800 ppm. Data on the chemical quality of water indicate that, because of the large amounts of ground-water inflow, the relative proportion of the individual constituents tends to remain fairly uniform regardless of the flow.

MONTANA

Industrial pumping will affect artesian pressure in aquifer west of Cedar Creek anticline

An investigation by O. J. Taylor of an aquifer in the Fox Hills Sandstone and basal part of the Hell Creek Formation, on the western flank of the Cedar Creek anticline in eastern Montana, indicates that proposed industrial use of ground water from the aquifer will affect the artesian head in wells west of the anticline and the water table in the outcrop area. The response of the hydrologic system to the withdrawals was calculated using a proposed pumping schedule and known aquifer characteristics. Values of drawdown to be expected after 5, 10, and 21 years of pumping under this schedule were plotted on maps. The pressure in some domestic and stock wells will be affected by the drawdown during the first 10 years of pumping. Municipal wells in the area will be affected only slightly. Owing to reduction in the pumping rate with time, the artesian pressure will largely have recovered by the end of the proposed 21-year pumping period.

Aquifer tests reveal presence of fracture permeability

Fracturing has caused an irregular distribution of permeability in the Virgelle Sandstone in Glacier and Toole Counties, in northwestern Montana. Aquifer tests by R. G. McMurtrey and E. A. Zimmerman revealed coefficients of transmissibility in excess of 20,000 gallons per day per foot in some parts of the consolidated sandstone aquifer. The Virgelle, which provides stock and domestic water to many and widespread wells, has recently been tapped to provide water for secondary recovery of oil. This industrial use has now been largely suspended because of concern over possible depletion and unsatisfactory well performance.

Artesian water available in Blood Creek syncline area

Artesian water suitable for livestock use can be obtained from wells tapping the Judith River Formation in the Blood Creek syncline area of north-central Montana. W. R. Osterkamp and N. J. King have compiled contour maps of the top of the Judith River Formation and of its potentiometric surface from measurements taken at numerous wells in the area. The study indicates that flowing wells with yields up to 15 gallons per minute can be developed at depths of 300 to 1,000 feet.

NORTH DAKOTA**Water-level decline near Minot**

According to W. A. Pettyjohn, the ground-water level in the Minot aquifer, in the Souris River valley at Minot, in north-central North Dakota, has declined more than 70 feet since 1915 and, in some areas, more than 20 feet since 1961. During 1963, water was withdrawn from the aquifer for the Minot municipal supply at an average rate of 3.8 million gallons per day, and the peak demand was over 7 mgd. Test drilling has shown that the areal extent of the aquifer is substantially less than was previously indicated. The aquifer will probably produce water at a rate of only 3 mgd under existing conditions without additional lowering of the water level.

Recharge to Heimdal outwash channel

Test drilling in Foster and Eddy Counties, in east-central North Dakota, has shown that segments of the present James River channel overlie the buried Heimdal outwash channel, which contains aquifers capable of large yields. Thus, flow in the James River may furnish part of the recharge to the Heimdal aquifers. According to Henry Trapp, Jr., two test holes drilled in the deep part of the Heimdal channel penetrated no significant thickness of sand or gravel, showing that the aquifers are not necessarily coextensive with the deepest parts of the channel.

Thick glacial drift overlies Missouri River valley

C. A. Armstrong reports that test drilling in Divide County, in the northwest corner of the State, revealed that as much as 638 feet of glacial drift overlies the ancestral Missouri River valley. Coefficients of transmissibility of buried outwash deposits range from about 13,500 to more than 10,000 gallons per day per foot. The ground water from the glacial drift is generally very hard and, except in areas of glacial outwash, contains more than 1,000 parts per million of dissolved solids.

Water levels decline near Fargo

R. L. Klausing reports that water levels in the West Fargo artesian aquifer, near Fargo, at the eastern edge of North Dakota, continue to decline because of increased industrial and municipal pumpage. The water level has declined from a high of about 26 feet in 1937 to the present record low of 108 feet below land surface. Near the center of the cone of depression the water level has declined below the top of the aquifer.

Undeveloped aquifer in Williams County

E. A. Ackroyd reports that alluvial deposits of sand and gravel in 120 square miles of the Little Muddy Creek valley, in northwestern North Dakota, yield ground water under artesian conditions. The maximum thickness of the aquifer is 137 feet. Only a small part of the available water is currently being withdrawn from the aquifer—mainly for domestic and stock use.

Productive aquifers found in Burleigh County

Studies by P. G. Randich show that aquifers adjacent to the Missouri River and in melt-water channels underlie an area of 230 square miles in southern Burleigh County in south-central North Dakota. Aquifer tests indicate that coefficients of transmissibility range from 10,000 to 350,000 gallons per day per foot. Approximately 7 million acre-feet of water is in storage in these aquifers, and it is estimated that 8,000 acre-feet per year moves from the aquifer to the Missouri River valley near Bismarck. According to J. L. Hatchett, most of the water has a high salinity hazard and a residual sodium carbonate classification of marginal to unsuitable for irrigation. These facts will require careful evaluation of the effect of this water on land and crops where it is currently being used for irrigation.

WYOMING**Ground water in Laramie County**

In water-resources study of Laramie County, in southeastern Wyoming, M. E. Lowry and M. A. Crist determined that the water table has declined significantly only in the vicinity of the Cheyenne municipal well fields. Withdrawal of ground water for irrigation in the Pine Bluffs area has increased from about 1,500 acre-feet in 1936 to an estimated 17,000 acre-feet in 1964, with no significant water-level decline. Deep wells drilled recently in the central part of the county show that yields as great as 300 gallons per minute can be developed in many areas where only small-yield stock wells had been drilled previously.

Extensive aquifer in Great Divide basin

A reconnaissance of a 10,500-square-mile area, chiefly in Sweetwater County, in southwestern Wyoming, by G. E. Welder and L. J. McGreevy, indicates that about a third of the area is underlain by water of fair to good quality. The most promising aquifer with respect to both quantity and quality of water is in the Battle Spring Formation, in the eastern Great Divide basin. Deep wells tapping the Battle Spring at favorable locations would probably yield more than 500 gallons per minute of water having less than 300 parts per million of dissolved solids.

SOUTH DAKOTA

Undeveloped artesian water supplies

Rocks of pre-Cretaceous age in the Black Hills and southern Williston basin have been described by M. J. Ellis and C. F. Dyer as constituting the greatest potential source for development of large, permanent water supplies in western South Dakota. Nine rock units are known to be water bearing, and three of them, the Whitewood Dolomite (and equivalent Red River Formation), the Pahasapa Limestone (and equivalent Madison Group), and the Minnelusa Formation, have good to excellent potential for development. The strata are, in general, deeply buried and have been reached by very few water wells except near the Black Hills.

Glacial-outwash aquifers in Campbell County

Preliminary studies by N. C. Koch and L. S. Hedges indicate that a fifth of Campbell County, in north-central South Dakota, is covered by glacial-outwash aquifers. A buried valley extends from the Oahe Reservoir near Pollock east-southeastward to Herreid and then southward to the vicinity of Mound City. The valley is eroded more than 100 feet below the general bedrock surface and is filled with glacial sediments.

Large ground-water storage in Clay County

J. C. Stephens reports that more than 5 million acre-feet of ground water is contained in transient storage in glacial and alluvial material that underlies three-fourths of Clay County, in southeastern South Dakota. Preliminary analysis of aquifer-test data indicates that the average coefficient of transmissibility of the aquifer may be as high as 500,000 gallons per day per foot.

Ground water in Beadle County

Aquifers more than 25 feet thick in the glacial drift in Beadle County, in east-central South

Dakota, contain more than 4,500,000 acre-feet of water in transient storage, according to Lewis Howells and J. C. Stephens; an additional 700,000 acre-feet of water is stored in aquifers 10 to 25 feet thick. Aquifer tests of the major aquifers in the drift disclosed that the coefficient of transmissibility ranges from 13,000 to 500,000 gallons per day per foot. The rate of withdrawal of water is estimated to have been more than 10 million gallons per day in 1962; of this total, nearly 3 mgd was from aquifers in the drift, and about 7 mgd was from bedrock aquifers.

NEBRASKA

Water level declines near Alliance

C. F. Keech reports that ground-water withdrawals for irrigation in an area north of Alliance, in western Nebraska, have lowered the water table as much as 40 feet since 1946. Recent measurements of water levels in 35 wells indicate that the water table has declined more than 20 feet over an area of 40 square miles and more than 40 feet in about 20 percent of the area.

UTAH

Large ground-water development in Utah

Ted Arnow reports that there are now 59 areas of known or potential ground-water development in Utah. The aquifers are consolidated rocks in 8 areas and unconsolidated rocks in 51 areas. Approximately 35,000 wells have been drilled in the State, and all except about 500 obtain water from the unconsolidated aquifers. The withdrawal of water from wells in the unconsolidated aquifers of the 19 most important areas totals nearly 600,000 acre-feet annually.

Large water discharge in Sevier Desert

R. W. Mower and R. D. Feltis estimate the minimum ground-water discharge in the Sevier Desert basin during 1964 to have been 165,000 acre-feet; of this total only 29,000 acre-feet was discharged by wells and 136,000 acre-feet was discharged principally by evaporation and by native phreatophytes. Decline of water levels in observation wells since 1950 was partly because of increased draft by wells and partly because of below-normal precipitation and attendant smaller quantities of recharge. Although discharge by wells increased from 1,500 acre-feet in 1950 the ground-water reservoir can be developed further, and with judicious selection of well sites it may be possible to increase withdrawal severalfold over that of 1964, without serious consequences.

Increased recharge in Pavant Valley

Water levels declined less than half as much in observation wells in the Kanosh district of Pavant Valley during 1964 as during any of the 4 previous years, according to R. W. Mower, although there were no significant changes in pumpage. The less-than-normal decline apparently was due to greater-than-normal recharge caused by runoff from a local thundershower on September 13, 1963, and by melt water from a thick snowpack (estimated to have attained 150 to 200 percent of the normal) during the last week of March 1964. The amount of water reaching a low area near the west-central part of the district, that contains several small sinkholes, was estimated to be 3,000–5,000 acre-feet. The ponded water disappeared within a short time after it had collected, apparently chiefly by draining through the sinkholes to recharge the shallow underlying basalt aquifer. These observations confirm an earlier conclusion that this low area is a major recharge area.

Water quality in Skull Valley

Data on the quality of ground water in Skull Valley, which extends southward from the southwest side of Great Salt Lake, indicate a wide range of water quality. According to Osamu Hattori, ground water in the northern part of Skull Valley is a sodium chloride water having about 1,000–7,000 parts per million of dissolved solids. The southeastern fourth of the valley contains calcium-bicarbonate or calcium-sodium-bicarbonate waters having generally less than 500 ppm of dissolved solids.

Ground water in solution channels in Scipio Valley

Ground-water levels in Scipio Valley, a topographically closed subbasin in western Utah, are unusual in that their depths change abruptly near the middle of the valley from about 50 to about 300 feet below land surface. L. J. Bjorklund and G. B. Robinson, Jr., report that the deeper water levels are in a zone of sinkholes formed by the solution of limestone along faults and joints in the North Horn Formation and in the Flagstaff Limestone, and by the subsequent collapse of overlying sediments. The faults traverse the valley in a northeasterly direction, and the sinkholes are aligned along the faults. The deeper ground water moves along the solution channels toward the points of discharge at Moulten and Blue Springs, about 8 miles northeast of the middle of Scipio Valley. The springs control the elevation of the deeper water levels.

COLORADO

Water resources of Arkansas River valley

Additional data and the reevaluation of past records have furnished a dependable basis for the construction of an electrical analog model of the Arkansas River valley between Canon City and the State line, according to J. E. Moore and C. T. Jenkins. Measurements of coefficients of transmissibility and storage, water levels, the application of water for irrigation, withdrawals of ground water, and river gains and losses have been programmed into the analog model to simulate the hydrologic system. The amount of surface water applied, changes in ground-water level, and river gain and loss correspond closely. Stream reaches having the greatest loss are in the areas of greatest withdrawal of ground water and of greatest decline of ground-water level. Conversely, gaining reaches are in areas of little pumpage and steady or rising water level.

Ground-water supply and pumpage in Black Squirrel Valley

A study by E. D. Jenkins and H. E. McGovern defines the alluvium of Black Squirrel Creek valley in a part of El Paso County near Ellicott, in east-central Colorado. The sandy channels of Black Squirrel Creek and tributaries are dry most of the time but may carry large quantities of water during periods of excess precipitation, much of which percolates through the permeable sandy channel to the water table. Calculations show an annual recharge of about 10,000 acre-feet, with about 400,000 acre-feet of ground water in transient storage. During 1964, 7,000 acre-feet of ground water was pumped for irrigation and 5,200 acre-feet was pumped to Colorado Springs for municipal use. The chemical quality of the water is suitable for most uses.

KANSAS

Water quality in Walnut River

Four surveys were made to determine the chemical quality of base runoff of the Walnut River and most of its tributaries, in south-central Kansas, over a range of water discharge representing antecedent conditions of above-normal precipitation and of drought. As a result of the surveys, R. B. Leonard found that about 60–70 percent of the dissolved-solids load in the lower reaches of the Walnut River was derived from soluble rock. Most of the remainder, consisting largely of sodium and chloride, was probably residual salt leached from soil and shallow aquifers contaminated by oil-field-brine disposal installations, most of which have been abandoned.

Streamflow at most data-collection sites on the main stem was generally unsatisfactory for drinking water and was suitable for irrigation only on well-drained soils from which accumulated salts are leached periodically.

Seepage inflow to South Fork Solomon River

M. W. Busby reports a change in seepage inflow to the lower reach of South Fork Solomon River, in north-central Kansas, after development of the Webster Irrigation Unit. The inflow below the diversion dam increased from 7.8 cubic feet per second in 1959, before the Unit was in operation, to 10.5 cfs in 1964 after 4 years of operation. This represents a 35-percent increase in inflow during a period when there was a decrease in rainfall. Ground-water levels in the irrigated area adjacent to the stream were as much as 8 feet higher in 1964 than in 1959.

Ground water in Neosho Valley

W. L. Jungmann reports that test drilling and pumping tests in the Neosho River valley in Labette and Cherokee Counties, in the southeast corner of the State, suggest the general availability of as much as 50 gallons per minute of good-quality water in alluvial deposits of Illinoian and Wisconsin ages. This is an area in which only relatively low-yielding domestic and stock wells have been developed previously. As much as 50 gpm of water containing less than 500 parts per million of chloride is available from deep (about 800 feet) Ordovician rocks in a local area in southeastern Labette County. Elsewhere, these rocks generally yield highly saline water.

Ground water in Finney County

Unconsolidated deposits of Tertiary and Quaternary age constitute the most productive ground-water reservoir in Finney County, according to W. R. Meyer, E. D. Gutentag, and D. H. Lobmeyer. Pumpage from wells increased from approximately 71,000 acre-feet in 1940 to approximately 295,000 acre-feet in 1963, and the total discharge from 73,000 to 297,000 acre-feet. Total storage in the reservoir is computed to be approximately 20 million acre feet. Maps showing net changes in water level indicate that only 16 percent of the water pumped in the period 1940-64 was derived from the change in storage. The average recharge to the ground-water reservoir was about $2\frac{1}{2}$ inches per year or approximately 121,000 acre-feet per year for this period. The annual discharge from wells is now exceeding recharge and is causing an average decline of approximately 1 foot per year over the county, whereas in

the early portion of the period, recharge exceeded discharge and water levels were rising.

ARIZONA

Ground water resources of Kingman area

J. B. Gillespie and C. B. Bentley report that about 40,000 acre-feet of ground water is pumped annually for domestic and industrial supplies in the Sacramento and Hualpai basins of northwestern Arizona. There are 200 wells in the area; yields are as large as 2,000 gallons per minute. Water levels are from 400 to 1,000 feet below land surface in the central parts of the basins and 50 to 100 feet in some marginal areas.

NEW MEXICO

Paving a small watershed

Runoff after thunderstorms ranged from 16 to 67 percent of the precipitation on a 9-acre area that had been paved with asphaltic materials, according to W. C. Ballance. Runoff from an adjoining 9-acre tract having a natural surface ranged from 0 to 3.4 percent, and it occurred only when runoff from the paved area exceeded 40 percent of the precipitation. Most or all of the precipitation on the area with natural surface returned to the atmosphere through evapotranspiration; runoff from the paved area flowed into an excavated earthen tank, from which most of the water percolated to the ground-water reservoir.

Ground water in Roswell area

Restudy of subsurface geology of the Roswell artesian basin, in southeastern New Mexico, by G. E. Maddox and J. D. Hudson indicates that the shallow aquifer is not as large as was formerly believed. Many irrigation wells that were previously thought to be developed in the shallow aquifer are developed in an aquitard that lies between the shallow and artesian aquifers.

The direction of flow between the shallow and artesian aquifers in the Roswell basin has been reversed and now is from the shallow to the artesian aquifer. Ground-water discharge to the Pecos River in the Roswell basin now occurs only from the shallow aquifer.

OKLAHOMA

Ground water in cavernous gypsum

A study by C. E. Steele and J. E. Barclay indicates that ground-water resources in the southwestern corner of Oklahoma may now be overdeveloped. Cavernous gypsum in the Dog Creek Shale and Blaine Formation in Harmon and adjacent parts of

Greer and Jackson Counties forms one of the most productive solution-channel aquifers in the State. Although the aquifer contains about 336,000 acre-feet of water in storage in the irrigation area, annual pumpage of about 50,000 acre-feet appears to be greater than the average rate of replenishment. Water levels in the aquifer continued to decline during the relatively wet period from 1958 to 1962.

Ground water in Cleveland and Oklahoma Counties

P. R. Wood and L. C. Burton report that the principal source of ground water for municipal and industrial use in Cleveland and Oklahoma Counties, in the central part of the State, is the Garber Sandstone and the Wellington Formation. In these two counties the Garber and Wellington are estimated to contain about 34 million acre-feet of fresh water in available storage—more than 80 times the 400,000 acre-feet of water that has been withdrawn from the aquifer since pumping began in the early 1900's. Recharge averages about 72,000 acre-feet per year.

TEXAS

Iron in ground water in east Texas

In a study of the ground water in Camp, Franklin, Morris, and Titus Counties, in northeastern Texas, M. E. Broom, W. H. Alexander, Jr., and B. N. Myers found a general stratification of the ground water with respect to pH, hardness, and dissolved iron. The ground-water reservoir includes a shallow zone of oxidation, A; a deep zone of reduction, C; and an intermediate and unstable zone, B, in which the waters from above and below mix. Ground water from zones A and C generally is relatively free of iron, whereas water from zone B generally contains objectionable amounts of iron. Wells constructed to draw only from zones A or C should yield water relatively free of iron.

Vertical movement of ground water

E. T. Baker, Jr., found that in Jackson County, a heavily irrigated area along the Gulf coast of Texas, the shallow parts of the Gulf coast aquifer which are lightly pumped supply large quantities of water to the deeper, heavily pumped zone. The transfer of water to the heavily pumped zone from shallow sands was observed in a multiscreened irrigation well by spinner surveys using a deep-well current meter. When water levels were lowest, near the end of the irrigation season, the survey showed that 42 gallons per minute was continually being transferred through the well from the shallow sand to various deeper sands within the zone of heavy pumpage. After the irrigation wells in the area had been shut

down for several months the transfer of water to the heavily pumped zone through the well had diminished to 14 gpm. The surveys indicate that the heavily pumped zone was effectively being repressured or replenished with water between the irrigation seasons.

Water of good quality in San Jacinto River basin

L. S. Hughes and Jack Rawson found that surface water in the San Jacinto River basin, in eastern Texas, is of excellent chemical quality and is suitable for most uses. The dissolved-solids concentration of most streams in the basin is generally less than 200 parts per million. Runoff from the outcrop areas of the Catahoula Sandstone, Lagarto Clay, and Oakville Sandstone in the upper part of the basin has dissolved-solids concentrations ranging from 100 to 200 parts per million; water draining from the outcrop areas of younger formations is less mineralized.

PACIFIC COAST AREA

The Pacific coast area includes the States of Alaska, California, Hawaii, Idaho, Nevada, Oregon, and Washington. It is an area in which precipitation ranges from zero in some desert areas of southern California and Nevada to nearly 500 inches annually on the slopes of Mount Waialeale on the Island of Kauai in Hawaii. It includes much of the Great Basin, from which no flow reaches the sea, and regions of the contrasting heavy flows of much of the Columbia River system and of the Yukon. The area has known searing droughts and devastating floods and includes problem areas of land subsidence and of sea-water intrusion, both related to ground-water withdrawals. The first commercial application of geothermal steam power in the United States was made in the region.

Land use in the Pacific coast area ranges from intensive agriculture and heavy industrial development to great National Parks and Primitive Areas set aside to be left perpetually in their wild state. The activities of man have caused, and are continuing to cause, major changes in the hydrologic regimen of the region. Inevitably there are large and growing problems resulting from concentration of population in areas of relative water shortage, and from the quantities of waste that impose loads upon some streams greater than the streams can carry without massive deterioration in quality.

The U.S. Geological Survey is making water-resources studies in each of the States in the Pacific coast area. These studies cover a broad spectrum of

problems related to quantitative evaluation and continuing inventories of surface-water and ground-water supplies, of their chemical quality, and of the sediment loads of streams. Basic-data networks have been operative throughout the area for many years and are being expanded as need arises. The hydrology of selected river basins is being studied to develop a fuller understanding of the characteristics of the system to provide an improved base for planning of water-development projects. Special studies in limnology, glaciology, land subsidence, and geochemistry, for example, support the investigations of surface-water discharge, flood studies, ground-water evaluation, chemical quality of water, and sediment transport.

Some of the many water-resources activities in the Pacific coast area are described below.

ALASKA

Greater Anchorage area

The predicted water-supply demand of 60 million gallons per day for the year 1980 in the Greater Anchorage area can be supplied adequately through use of both surface and ground waters. Hydraulically connected streams and aquifers offer conditions suitable for the coordinated use of the total water resources, with an opportunity for artificial and induced ground-water recharge. Loss of considerable portions of the extensive artesian-aquifer system by sea-water intrusion and contamination through leakage of septic-tank effluents from the shallow water-table aquifer in the recharge areas pose the greatest potential hazards, according to D. A. Sommers.

Potential water sources on Annette Island, southeastern Alaska

At least two levels of raised beach deposits are present on Annette Island in extreme southeastern Alaska. M. V. Marcher points out that locally the beach deposits are the source of seeps and springs and may be a significant source of water to properly constructed wells or infiltration galleries. Test drilling has verified the presence of ground water in the sand and gravel beds.

Water from crystalline rocks

Although crystalline rocks have not been considered favorable for drilling in southeastern Alaska, M. V. Marcher reports that test wells on Annette Island show that ground water is available from highly fractured igneous rocks near Tamgas Harbor and Annette Island airport, and from phyllite and

schist underlying the village of Hoonah. In interior Alaska test drilling at the hotel in Mount McKinley National Park showed that small amounts of ground water can be obtained from schist in that area. At the Wonder Lake Ranger Station, also in the park, a well yielded about 1 gallon per minute from fractured schist below 190 feet of permafrost.

PACIFIC NORTHWEST

The Pacific Northwest comprises the States of Idaho, Oregon, and Washington. The mountainous areas are extensive and well watered, with precipitation reaching a high of about 200 inches per year in a few spots along the Washington coast and a peak of 100 to 120 inches elsewhere along the coast. East of the coastal mountains, however, precipitation declines rapidly, and in each of the three States there are areas which receive only about 8 inches of precipitation per year. Thus, runoff is abundant in many areas but semiarid conditions prevail in parts of each State. In December 1964 and January 1965 serious floods occurred in western Oregon.

Because of international agreements relative to water of the Columbia River and interstate agreements relative to water of the Snake River, the basic-data networks related to the discharge and chemical quality of these streams have long been important.

Hydrology of Snake River Plain, southern Idaho

A study by E. H. Walker suggests that the Snake River Plain aquifer of southern Idaho received about 3,260,000 acre-feet per year of recharge in the mid-1950's by infiltration from lands irrigated with surface water. These new estimates were made in cooperation with the U.S. Bureau of Reclamation, for design of an electric-analog model of the aquifer. The contribution from irrigation was about 47 percent of the total recharge, 6,940,000 acre-feet per year, and 80 percent as much as the total recharge to the aquifer in the early 1900's before man had appreciably altered the natural regimen of the aquifer. Present conditions are not greatly different from those of the mid-1950's.

Electric-analog model studies in southeastern Idaho

J. G. Crosthwaite, M. E. Mundorff, and E. H. Walker used a steady-state analog model analyzer developed by R. W. Stallman to estimate seepage losses from a proposed reservoir site on the lower Teton River in southeastern Idaho. The study indicates that seepage losses would not exceed 4 percent of the average annual flow of the river.

Ground water resources of Island County, Wash.

An appraisal by H. W. Anderson of the ground-water resources of Island County, in Puget Sound, shows that throughout most of the county, wells can supply enough water to satisfy municipal and rural domestic and irrigation demand. Virtually all ground water now developed or capable of being developed occurs in permeable glacial or interglacial deposits of Pleistocene age. A major water-bearing zone lies near sea level, extending just above and below it throughout much of the county.

Ground water in Baker County, Oreg.

According to G. L. Ducret and D. B. Anderson, large quantities of water are available for irrigation and other uses from wells tapping alluvial deposits in Baker Valley, Oreg., and from basalt aquifers in nearby areas east of the valley. Wells yielding as much as 1,000 gallons per minute tap sand and gravel aquifers in valley alluvium and alluvial-fan deposits, and wells yielding as much as about 1,500 gpm tap lava rock and cinders.

Donald Price found that the three main agricultural segments of the Burnt River valley, in Baker County in eastern Oregon, are underlain by fine-grained fluviolacustrine deposits of Miocene and Pliocene age that yield only small quantities of water to wells and springs in the valley. However, permeable gravel layers and intercalated lava flows known to be present locally may yield moderate to large quantities of water to wells, provided that the flows and layers extend below the water table.

Ground water in Eola-Amity Hills area, Oregon

Because of the diversity of geologic conditions the availability and quality of ground water vary considerably within the 230-square-mile Eola-Amity Hills area of the northern Willamette Valley, Oreg. Donald Price reports that young alluvium of the Willamette River yields large quantities of good-quality water to wells. In contrast, marine sedimentary rocks, which underlie much of the western half of the area, yield only small quantities of water to wells, and the water generally contains a relatively large concentration of chloride.

NEVADA

Nevada probably receives less moisture per square mile from precipitation than any other State in the Union. A small available supply is its first water problem. Streamflow from the Sierra Nevada and rights to share in water of the Colorado River partly alleviate the problems of supply. In addition, some valleys in the State are presently at a stage

of modest development which leaves existing supplies of ground water adequate for existing needs. Important basic-data networks were continued and augmented.

Reconnaissance ground-water studies continued

Reconnaissance ground-water studies have been completed or were in progress in 11 valleys during the year. Preliminary estimates of perennial yield, based on estimates of average annual recharge to and discharge from the ground-water reservoir for several of the valleys, are as follows: 10,000 acre-feet for Smith Creek valley, Lander County (D. E. Everett and F. E. Rush, 1-64); 6,000 acre-feet for Ione Valley, Nye County (Everett and Rush, 1-64); 13,000 acre-feet for Grass Valley, Pershing County (Philip Cohen and D. E. Everett, 5-64); 21,000 acre-feet for Monitor Valley, Nye and Lander Counties (F. E. Rush and D. E. Everett, 2-64); 4,000 acre-feet for Antelope Valley, Eureka County (Rush and Everett, 2-64); 16,000 acre-feet for Kobeh Valley, Eureka County (Rush and Everett, 2-64); 37,000 acre-feet for Upper Reese River valley, Lander and Nye Counties; and 2,000 acre-feet for locally derived ground water in the Oreana subarea of Lovelock Valley, Pershing County; and about 70,000 acre-feet for Spring Valley, White Pine County.

Regional ground-water system identified

A regional ground-water system in the White River valley area in eastern and southeastern Nevada has been defined more fully than before (T. E. Eakin, 1-64; T. E. Eakin and I. J. Winograd, 3-64); I. J. Winograd and T. E. Eakin, 1-64). The 7,700-square-mile area which comprises the system includes 13 valleys along and adjacent to the Pleistocene channel of the White River. Although inter-basin movement has been described in eastern and southern Nevada in recent years, the extent of the system was not previously understood. Carbonate rocks of Paleozoic age which occupy much of a large miogeosynclinal area of eastern and southern Nevada and western Utah and which have been subjected to a very complex structural history appear to be the principal media for the regional transmission of ground water. Superimposed over the bed-rock aquifer are alluvial aquifers of the Basin and Range type. Water-budget studies of the alluvial aquifers, the most important type in the State, are complicated by the fact that the carbonate rocks may carry water from one basin to another through the bounding mountains. Most alluvial valleys in the Basin and Range province can be considered to be

discrete hydrologic units except for such streams and shallow alluvial channels as may connect them.

Humboldt River project completed

The interagency Humboldt River Research Project study near Winnemucca, Nev., by Philip Cohen (2-64, 3-64) and others, completed in 1964, indicates that the average annual inflow to the area studied was about 250,000 acre-feet in water years 1949-62. Of this amount, about 68 percent was Humboldt River streamflow, 23 percent was precipitation on the area, 6 percent was subsurface ground-water inflow, and about 3 percent was tributary streamflow. The average annual surface-water outflow was about 155,000 acre-feet, or about 17,000 acre-feet less than the surface-water inflow. The total average annual evapotranspiration loss was about 115,000 acre-feet.

Surface-water resources of Nevada

R. D. Lamke and D. O. Moore estimate that the average annual runoff within Nevada is 3,500,000 acre-feet. The average annual streamflow for the 1912-63 reference period was 1,500,000 acre-feet entering the State and 850,000 acre-feet leaving the State. The estimated average annual water supply available to Nevada is 4,100,000 acre-feet. The estimated average annual precipitation over the entire State is slightly more than 9 inches, or about 54,000,000 acre-feet. The statewide runoff is thus about 61½ percent of the precipitation.

An inventory of the better-known Nevada springs indicated that the total annual springflow may average more than 300,000 acre-feet. There was 170,000,000 acre-feet of water stored in lakes and reservoirs in and bordering Nevada as of October 1, 1963. About 30,000,000 acre-feet was contained in lakes and reservoirs wholly within Nevada.

Ground-water storage depleted in Quinn River area

A study of the Quinn River valley area in northern Nevada by C. J. Huxel shows that since 1950, pumping of ground water for irrigation has resulted in depletion of ground-water storage by about 120,000 acre-feet and an average water-level decline of about 22 feet. The moderately large water-level decline is caused by the concentration of pumping in the southern part of Quinn River valley. During the past 5 years, pumpage has increased from 23,000 to 40,000 acre-feet per year.

Water regimen changing in Kings River valley

Investigation by G. T. Malmberg of the manageable water resources of Kings River valley in Nevada and Oregon indicates that the average annual

surface-water inflow is about 7,300 acre-feet; the average annual recharge to the ground-water reservoir is nearly 20,000 acre-feet. Although most of the surface water was diverted for irrigation, only about one-third is consumptively used. Most of the remaining 5,000 acre-feet infiltrates to the ground-water reservoir.

In 1964 the pumpage, about 22,000 acre-feet, was used to irrigate about 6,000 acres. During the period 1956-63, the net depletion by pumping and natural discharge of ground water in storage totaled about 60,000 acre-feet. The water table has been declining progressively at about 3 feet per year since pumping began, but the rate probably will increase due to the effects of bedrock boundaries. If pumping is continued at the same rate as in 1964, the water table at the pumping centers can be expected to drop about 30 feet in 10 years, thereby reversing the hydraulic gradient and causing practically all ground water in the northern half of the valley to move toward the pumped area.

CALIFORNIA

Water problems in California varied in degree rather than in kind during the year. In southern California, runoff continued below average. However, during December 1964 and January 1965 severe floods occurred in the north, causing loss of life and heavy destruction of private property and of public facilities. At one key station on the North Fork American River, discharge in December 1964 was 2,190 percent of the median for the month. Tens of gaging stations, part of the surface-water basic-data network, were damaged or totally destroyed by floodwaters. These have been restored to operation as rapidly as water and weather conditions allowed. Special attention has been given during the year to collection of basic data on ground-water levels and chemical quality in the Mojave Desert region.

Work continued on studies of land subsidence, effects of urbanization on the hydrology of small watersheds, and the hydrologic regimen of individual ground-water basins. The status of water supplies for numerous military installations throughout the State was again assessed as part of a continuing inventory made in cooperation with the U.S. Department of Defense.

Relation between ground water and surface water in Santa Ana River valley

In a study of the reach of the Santa Ana River between Riverside and Prado Dam, in southern

California, J. J. French, E. R. Hedman, and L. C. Dutcher found that the net ground-water contribution to surface flow has decreased during the period 1933 to 1964. Both surface-water inflow to the reach and evapotranspiration losses in the bottom lands along it have remained relatively constant during this period, but the water table has declined.

Multiple piezometers installed in central California

J. F. Poland reports that a multiple-piezometer observation well constructed in a subsidence bowl near Pixley, in the southern San Joaquin Valley, is providing data on artesian head in six well-defined individual beds of sand and gravel. These strata are the principal aquifers in an alluvial aquifer system, 500 feet thick, which is heavily pumped through fully perforated gravel-packed irrigation wells and is compacting as a result of head decline. The piezometers have demonstrated that significant head differences exist among the several aquifers, both in response to pumping and during prolonged intervals of no pumping. During the irrigation season a normally downward vertical gradient through the aquifer system is reversed.

Areal ground-water studies, Santa Barbara County

An investigation by K. S. Muir indicates that there is about 280,000 acre-feet of potable ground water in storage in the aquifers underlying the city of Santa Barbara and the adjacent area of Montecito, in coastal southern California. This ground water is stored in fine-grained deposits in a zone between the altitudes of 250 feet above sea level and 200 feet below sea level.

Study of inflow-outflow relationships in the Lompoc subarea of the Santa Ynez River basin by R. E. Evenson, for the period 1957-62, indicates a possible perennial supply of 22,000 acre-feet. The amount actually produced in the future may be less than this estimate, however, because of two factors: (1) reduction in amounts of water pumped for irrigation, and (2) changes in water quality due to recycling of irrigation water and the inflow of poor-quality connate water from the consolidated rocks that might require reduced draft in some parts of the subarea.

A ground-water appraisal in the Santa Ynez Upland area by G. F. LaFreniere has led to the following results: (1) the extent of the upland area as defined by J. E. Upson in 1952 has been increased about 40 square miles bordering Happy Canyon, by recognition of the hydrologic continuity of water-bearing deposits in the two areas; the upland is important as a recharge area in the Santa Ynez

subarea of the Santa Ynez River Valley; (2) evidence has been found that only a minor amount of recharge can be expected to the basin from Cachuma Reservoir which presently inundates the Paso Robles and Careaga Formations in Cachuma and Santa Cruz Canyons; (3) an estimate has been made of 10 million acre-feet of water in storage in the entire ground-water basin, but a perennial yield probably not exceeding 8,000 acre-feet.

HAWAII AND GUAM

The high permeability of most of the volcanic-rock aquifers of the Hawaiian Islands promotes rapid recharge but inhibits development of sizable surface runoff in most areas. All ground-water bodies in the islands are probably in contact with sea water. As a result it is necessary to maintain a safe and sometimes delicate balance between recharge and the withdrawal of water so as to minimize mixing of fresh and salty water and to preserve the usefulness of pumped supplies.

The U.S. Geological Survey maintains continuing data-collection programs to monitor surface-water, ground-water and chemical-quality conditions on the islands. Areal hydrologic studies were a significant part of the investigations. During the year, personnel of the Hawaii district also participated in water-supply studies on the Island of Guam. Results of some of the studies are presented below.

Ground water studies on Oahu, Hawaii

Between 7 and 15 million gallons per day of irrigation water is recharging the shallow ground-water body in the coastal part of the Mokuleia area, in northwestern Oahu. Aquifer tests and allied investigations by J. C. Rosenau, E. R. Lubke, and R. H. Nakahara indicate that up to 5 mgd of this might be intercepted through the pumping of large-diameter shallow wells dug in the permeable limestone and dune-sand deposits of the shore area. Beneath this shallow aquifer the deeply buried lavas of the Waianae Volcanic Series discharge perhaps as much as 10 mgd to the sea.

In a study of ground water in the coastal part of the Kahuku area in northern Oahu, K. J. Takasaki and Santos Valenciano have found that the water in the upper part of the fresh-water lens has a higher content of chloride and nitrate than water deeper in the lens. The high chloride content results from infiltration of irrigation water pumped from wells affected by sea-water intrusion; the nitrate is contributed by fertilizers applied to the irrigated sugarcane fields.

Water resources of the Hilo-Puna area, Hawaii

Although rainfall is abundant up to an altitude of 5,000 feet in the Hilo-Puna area of Hawaii, there is but little runoff of surface water, according to George Yamanaga and D. A. Davis, because of the highly porous nature of the lava-flow terrane. Most of the discharge into the sea is at many springs along the shore. In the Hilo area, the visible flow of springs amounts to several hundred million gallons of water having low salinity. Locally in the Puna area the ground water is characterized by high temperature and high salinity which are thought to be due to volcanic activity along the southeast rift of Kilauea.

Hydrology of Guam

Studies in Guam by P. E. Ward, S. H. Hoffard, and D. A. Davis¹⁸ show that the average runoff to the sea in the southern half of the island is roughly 250 million gallons per day. This amounts to about half the daily rainfall on the drainage basins, which are underlain by volcanic rock having generally low permeability. The northern half of the island is a highly permeable limestone terrane in which virtually all the flow of water to the sea is underground through a fresh-water lens near sea level. Sea-level fluctuations resulting from tides and storm waves are transmitted far inland through the permeable rock and cause mixing of the fresh water with sea water. Consequently, ground water in the limestone is brackish in most of the coastal area.

MANAGEMENT OF NATURAL RESOURCES ON THE PUBLIC LAND

In order for the standard of living to keep pace with population expansion, wise management of the Nation's resources is essential. The lands owned by the Federal Government make up an important percentage of this Nation's resource base (fig. 2) and as a result require increasing attention to keep up with development needs. Management control is exercised by several agencies of the Government, depending upon the type of resource, the kind of ownership, and the various legal authorizations involved. The Conservation Division of the U.S. Geological Survey performs a number of important tasks in this connection. Through its Branches of Mineral Classification and of Waterpower Classification the division classifies the public lands for leasable mineral values and for waterpower purposes. Through

¹⁸ P. E. Ward, S. H. Hoffard, and D. A. Davis, 1965, *Hydrology of Guam*: U.S. Geol. Survey Prof. Paper 403-H. [In press]

FIGURE 2.—Percentage of total United States mineral production that comes from Federal land.

its Branches of Mining Operations and of Oil and Gas Operations it supervises prospecting, development, and recovery of minerals from wells and mines under leases, permits, or licenses on Federal and Indian lands.

Field offices of the Division are listed on page A211. Geologic and hydrologic work in progress by geologists and engineers of the Conservation Division is given in the list of investigations starting on page A217, under the categories of geologic mapping, glaciology, waterpower classification, and various commodities such as coal and petroleum and natural gas. Scientific and economic results of these investigations are published as books and maps in the regular series of Geological Survey publications. Important scientific findings of current interest by Division geologists are included in the section "Regional Geology."

CLASSIFICATION OF MINERAL LANDS

The principal leasable minerals are oil, gas, oil shale, coal, phosphate, sodium, and potash. Since 1906, public lands believed to contain leasable minerals have been withdrawn from entry by the Secretary of the Interior, on the advice of the U.S. Geological Survey, pending their classification as mineral or nonmineral lands. Much of the task has been completed, but at present about 41 million acres of land is so withdrawn.

In order to classify land for its mineral value, the geology must be mapped in detail, usually at a scale of 1:24,000. Existing geologic maps are used where of suitable quality and scale. Mapping for mineral-land classification differs in some details from gen-

eral geologic mapping in that more measurements of stratigraphic sections per quadrangle are required to show thicknesses of coal or other leasable minerals, and more frequent sampling is necessary to determine mineral quality. Drill-hole cuttings or cores and electric or radioactivity logs provide subsurface data for classification. Samples obtained from surface and subsurface operations are analyzed in the laboratory. When the areal extent, thickness, and grade of mineral deposits are established, the land containing them is classified as to its value for mineral resources. The geologic maps produced for classification purposes are published in the standard map series of the Geological Survey.

WATERPOWER CLASSIFICATION

Classification of public lands to conserve and utilize water resources was begun in 1888, to preserve reservoir sites for irrigation. It has been continued chiefly for hydroelectric development. Present activity includes stream-basin investigations, a review of land classifications and reserves, and the measurement of selected glaciers.

The program of stream-basin investigations is a systematic search for basins and subbasins where water projects can be developed in the future to serve the needs of a growing population. Sites that meet the criteria for classification are set aside by the Secretary of the Interior on the recommendation of the Geological Survey, through procedures established by Congress and the Department of the Interior. Land set aside by these procedures may be disposed of only according to procedures which provide that it may be reacquired by the Government for purposes of waterpower development without cost. Some 16 million acres are presently involved in this program.

A program to reassess all land classifications and reserves was begun in 1956. Many classifications and reserves had been made before 1920 under Congressional and departmental urging for immediate protection of sites for waterpower development. Because of this urgency many of the classifications were based on inadequate mapping and hydrologic data. Under the present program all reserves are being reevaluated and classified by rigorous standards. Lands that do not qualify as waterpower sites are removed from the reserves and are opened to disposition under the public-land laws. Progress through fiscal 1964 on this program indicates that as much as 50 percent of lands now reserved for waterpower may eventually be eliminated from the reserves.

River and lake basins are mapped mostly at a scale of 1:24,000. Contours of lake bottoms are compiled by precise sounding surveys. The results of investigations are published as river survey maps, a regular Geological Survey map series.

A special project to gather information on the rates of ablation and the recession or advance of selected glaciers was started in 1951 on Nisqually Glacier in Mount Rainier National Park, Wash., and in 1944 on Grinnell and Sperry Glaciers in Glacier National Park, Mont. Annual measurements are made at monumented cross sections, and the glaciers are mapped completely at 5-year intervals. In 1961, at the request of and in cooperation with the U.S. Bureau of Reclamation, measurements were begun on Barrier Glacier, Mount Spurr, Alaska, to determine what effect the growth or shrinkage of the glacier has on fluctuations in the level of Lake Chakachamna.

SUPERVISION OF PROSPECTING, DEVELOPMENT, AND RECOVERY OF MINERALS

Supervision of operations under mineral leases entails the investigation of lands and deposits under application for mineral leases, oil and gas leases, and prospecting permits; recommendation of lease terms and unit areas; enforcement of operating regulations and measures to assure the safety and welfare of workmen; maintenance of production records; and determination and collection of royalties and rentals. The mineral supervisory branches of the Conservation Division also act as advisors to the Secretary of the Interior, to other bureaus of the Department, and to other Government agencies concerned with the administration of the Mineral Leasing Laws.

Royalties from public lands are distributed 52½ percent to the reclamation fund, 37½ percent to the States in which the minerals or fuels are produced (except for Alaska, which receives 90 percent), and 10 percent to the Federal Treasury. Royalties from other land categories are distributed in many different ways as provided by law, but the largest share of these royalties is returned directly to the Federal Treasury.

Mining operations

The Branch of Mining Operations supervises operations concerned with discovery, development, and production of coal, oil shale, phosphate, potassium, solid and semisolid bitumen, and sodium from public land; and of sulfur from public land in Louisi-

Mineral production, value, and royalty

Lands	Oil (barrels)	Gas (thousand cubic feet)	Gas liquids (gallons)	Other ¹ (tons)	Value (dollars)	Royalty (dollars)
Public-----	181,726,000	641,276,000	484,478,000	26,470,000	765,910,000	80,754,000
Acquired-----	9,747,000	24,067,000	342,000	321,000	43,223,000	4,620,000
Indian-----	35,584,000	100,143,000	129,204,000	9,371,000	139,711,000	17,157,000
Military-----	2,857,000	63,617,000	74,327,000	-----	19,755,000	3,789,000
Outer Continental Shelf.	128,656,000	636,150,000	-----	1,050,000	527,083,000	92,051,000
Naval Petroleum Reserve No. 2.	4,000,000	5,800,000	10,500,000	-----	14,873,000	1,925,000
Total-----	362,570,000	1,471,053,000	698,851,000	37,212,000	1,510,555,000	200,296,000

¹ All minerals except petroleum products; includes coal, potassium, sodium, and so forth.

ana and New Mexico. The Branch also supervises (1) the production of silica sand on certain lands in Nevada; (2) mercury on certain Spanish land grants; (3) all minerals except oil and gas on restricted, allotted, and tribal Indian lands; and (4) all minerals recoverable in commercial quantities, except oil and gas, on acquired lands.

The column entitled "other" in the preceding table shows production of minerals from leased Federal lands under supervision of the Branch of Mining Operations for the fiscal year ending June 30, 1965.

Oil and gas operations

The Branch of Oil and Gas Operations supervises the discovery, development, and production of crude oil and natural gas and associated products from leased public, acquired, Indian, Outer Continental Shelf, and Naval Petroleum Reserve lands.

The table shows the production of crude oil and gas, the value of all mineral products, and the royalties received from supervised leases on the various categories of Federal and Indian lands during fiscal 1965.

GEOLOGY AND HYDROLOGY APPLIED TO ENGINEERING AND PUBLIC HEALTH

INVESTIGATIONS RELATED TO NUCLEAR ENERGY

Nuclear reactors and underground nuclear explosions release radioactive products to the geologic and hydrologic environments. In the interest of public safety, the distribution, movement, and concentration of these products must be determined and the potential hazard evaluated. Large nuclear explosions may produce excessive ground motion and the geology of the affected areas must be evaluated in the planning stage in the interest of public concern. In addition, the engineering and construction of facilities for nuclear reactors and underground nuclear explosions at proposed sites present difficult problems requiring a thorough knowledge of the geologic and hydrologic conditions in a variety of natural environments.

Since 1956 the U.S. Geological Survey has provided geologic and hydrologic information and evaluation of sites for underground nuclear explosions at the Nevada Test Site, and at sites for PLOWSHARE (peaceful uses of nuclear explosions). In connection with these programs the Geological Survey performs a broad range of studies ranging from detailed evaluation of specific sites and the geologic and hydrologic effects of explosions at specific sites, to areal geologic studies of areas contiguous to test sites.

SITE EVALUATION

Tunnel-site development, Nevada Test Site

Many of the military effects of nuclear explosions are studied by means of explosions in tunnels at the Nevada Test Site. Very detailed geologic exploration and appraisal is required to ensure that the tunnels are properly located. Several examples of this type of appraisal are given below:

At one test site a shear zone was discovered during exploration of a proposed working point. On the basis of detailed study, R. E. Davis concluded that the proposed working point was unsuitable. A reappraisal of the geology of the area by W. L. Emerick and T. L. Prather established a new working-point location free of faulting.

In a tunnel on Rainier Mesa two sites were relocated in an area of the tunnel that is relatively free from jointing and faulting to allow tunnel construction with a minimum of mining difficulties.

J. R. Ege made the geologic studies necessary for the location and construction of a test site in the Climax granitic stock. During the course of exploratory drilling at the site he devised a system of drill-core classification which serves as a basis for determining the suitability of the underlying rock for tunnel construction. As a result of careful study of the fractures in the tunnel and vicinity, Ege was also able to determine a major joint set which was used to determine the position of a cavity constructed at this site.

Velocity layering of rock surrounding tunnels in the Climax stock

Seismic studies were made in tunnels in the Climax granitic stock, Nevada Test Site, to determine the thickness of a layer of low-velocity rock that has been found to exist around tunnel openings. It is believed that these layers develop by different mechanisms in different rock types. In compressible, porous rock such as sandstone and tuff, it has been shown that velocity depends on porosity and pressure. In these rock types the low-velocity layer probably develops as decreased pressure permits the rock to expand and interstitial porosity to increase. Secondary fracturing along joints and bedding planes probably also affects the velocity in the zone of expansion. In rocks of low interstitial porosity such as limestone and granite, the low-velocity layer is probably caused by the enlargement of existing fractures and the formation of new ones along planes of weaknesses as the rock expands inward in response to stress relief.

J. H. Scott and R. D. Carroll made in-hole and refraction seismic measurements to determine variations in the thickness and velocity of the low-velocity layer surrounding tunnels constructed in quartz monzonite and granodiorite of the Climax stock. Measurements were made in different qualities of rock, with fracturing ranging from minimum to

severe. In general the thickness of the low-velocity layer ranged from 0 to 8 feet, and its velocity ranged from 5,000 to 14,000 feet per second. Velocity of normal rock behind the low-velocity layer ranged from 14,000 to 25,000 fps. Ranges of thickness and velocity were about the same for quartz monzonite and granodiorite. The lowest velocities were associated with rock that was most severely fractured, and the thickness of the low-velocity layer appeared to be affected by the direction as well as the density of fracturing. The present preliminary study provided only semiquantitative relationships between fracturing, velocity, and thickness of the low-velocity layer. Future detailed studies are planned from which quantitative correlations of the parameters of interest in various rock types may be made.

Evaluation of the LONG SHOT site, Amchitka Island, Alaska

The geology of Amchitka Island in the Aleutian Islands was studied by L. M. Gard to evaluate the suitability of the island for Project LONG SHOT, which is a proposed underground nuclear explosion for improving detection of underground nuclear explosions. These field investigations aided in deciphering the subsurface geology at the site. Exploration of the site to a depth of 2,600 feet was accomplished by core holes drilled by the U.S. Army Corps of Engineers.

The site is located in a thick sequence of gently dipping Tertiary volcanic detritus cut by linear fracture zones. The rocks are partly altered basaltic volcanic breccias of the Oligocene to Miocene Banjo Point Formation containing two andesite sills near the working point. Hydrologic testing showed that the fracture zones and sills are much more permeable than the bulk of the Banjo Point Formation.

Byproducts of the fieldwork resulted in revision of knowledge of the stratigraphic sequence on Amchitka Island. The Banjo Point Formation, previously described as older than the Chitka Point Formation, was found to conformably overlie and interfinger with the Chitka Point Formation. Rocks of the Amchitka Formation, thought to be the oldest exposed on the island, because they are intensely altered and recrystallized, were found to be merely rocks of the Chitka Point and Banjo Point Formations that have been locally metamorphosed near large intrusive bodies.

The Chitka Point and Banjo Point Formations contain many turbidites (submarine mudflows) not previously described from the island.

The island was revisited shortly after the 7.5-magnitude (Richter scale) earthquake of February 3, 1965. Only slight damage was found to have occurred in the form of slumping of road fills. Evidence was seen that a tsunami (seismic sea wave) had moved flotsam about 6 feet above high tide.

PLOWSHARE PROGRAM

The aim of the PLOWSHARE Program of the U.S. Atomic Energy Commission is to develop peaceful uses of nuclear explosions. The U.S. Geological Survey participates in and contributes to this effort in various ways, including selection of sites potentially useful for experiments, making detailed studies of the geology and hydrology of explosion sites, and making feasibility studies of beneficial applications of nuclear explosions in the field of natural resources.

Applications of nuclear explosions to water-resources development

Potential applications of nuclear explosives to water-resource development and management have been appraised by A. M. Piper and F. W. Stead (1-65) and F. W. Stead (1-64). Nuclear explosions fully contained underground may offer very substantial advantages and economies under certain limited hydrologic situations; however, much more must be learned about the phenomena of such detonations before their effects can be adapted to each particular situation, with assurance that the seeming advantage can be realized in full while undesirable side effects are constrained as circumstances require.

GEOLOGIC AND HYDROLOGIC EFFECTS OF NUCLEAR EXPLOSIONS

Explosion-produced fractures in Yucca Flat, Nevada Test Site

Underground testing of nuclear devices has been, and is currently being, conducted in Yucca Flat, at the U.S. Atomic Energy Commission's Nevada Test Site. Yucca Flat is an alluvium-filled valley underlain by tuff which is in turn underlain by generally more competent sedimentary rocks. Most of the devices tested to date were detonated either in alluvium or tuff. D. D. Dickey has studied maps of the fracturing, at the surface, produced by the explosions and concludes that the surface fracture pattern is influenced by the rock surrounding the explosion. Tests in the northern part of Yucca Flat show particularly well a symmetrical pattern of radial and concentric fractures in alluvium for explosions in alluvium, whereas explosions in the underlying tuff produce fractures in alluvium with

a strong asymmetrical northeast-trending pattern, with many northeast-trending fractures. The northeast direction is parallel to a strong set of northeast-trending pre-explosion joints in the bedrock at the northeast edge of Yucca Flat. In instances where the cavity from an explosion in tuff has collapsed and a crater or sink allowed to form in the alluvium at the surface, the resulting fracture pattern is that of the characteristic northeast-trending pattern for explosions in tuff, superimposed upon a pattern characteristic for explosions in alluvium.

Activation of Yucca fault, Nevada Test Site, by underground nuclear explosions

Underground nuclear explosions in Yucca Flat have caused displacement along the Yucca fault. The formula $S = \frac{D}{W^{1/3}}$ can be applied to determine if the fault will show additional displacement as a result of a particular underground nuclear explosion. In the above formula, S is scaled distance, D is distance in feet from the explosion point to the fault, and W is the yield of the device in kilotons equivalent TNT. Where S for a particular explosion is less than 1,000 the Yucca fault is expected to be activated as a result. Certain factors, such as rock type, abundance of fractures, or any condition that affects the distribution of energy between the explosion and the fault, would require some adjustment of the formula. Also where displacement along the fault has occurred as a result of a past explosion, there appears to be a loss of sensitivity; in other words more energy may be required to trigger the fault.

It is presently believed that the fault movement is triggered by explosions that relieve tectonic stress. First-order surveys across the Yucca fault before and after a recent explosion which activated it showed not only a downdropping on the east side, but an absolute rise in elevation of 24/1000 of a foot on the west side. Release of tectonic stress also is suggested from comparison of U.S. Coast and Geodetic Survey calculations of seismic energy measured for different explosions. Two explosions with which accompanying fault movement was recorded had a greater percentage of seismic energy than did 17 others with which no fault movement was associated.

Hydrologic characteristics of the BILBY chimney

The hydrologic characteristics of a rubble chimney created by the BILBY event—the first nuclear explosion within a water-saturated zone at the Nevada Test Site—and the surrounding area were studied by W. E. Hale, W. A. Beitem, and M. S. Garber. The device, having a yield of more than 200 kilotons,

was detonated in September 1963 in a thick porous tuffaceous aquiclude with a fracture transmissibility of about 10 to 50 gallons per day per foot. The collapse which followed the explosion created a "rubble chimney," with a radius of about 280 feet, partly within the zone of saturation. Measurements of the rise of water in, and pumping tests of, the rubble chimney were made in a hole drilled and cased into the center of the chimney. Analyses of the rates of water-level rise before, during, and after 2 pumping tests indicate that the effective porosity of the rubble in the chimney—at a level about 700 feet above the point of detonation—is slightly less than 2 percent; the preshot effective fracture porosity was a fraction of a percent. Subsequent measurements of water-level change should yield estimates of the rubble porosity at shallower depths. The transmissibility of the rubble chimney is about 30,000 gpd per ft, or 10^3 that of the undisturbed rock.

Effect of the SALMON event, Tatum salt dome, Mississippi, on nearby aquifers and streams

The SALMON event, a nuclear explosion of about 5 kilotons, was fired at a depth of about 2,700 feet in the Tatum salt dome on October 22, 1964. A part of the Geological Survey's hydrologic program involved the measurement of water levels in nearby wells and flow of streams in the area through a period before and after the SALMON event.

Four wells within $1\frac{1}{4}$ miles of the emplacement hole were instrumented with pressure transducers to record short- and long-term water-pressure response to the SALMON event. The movement of water within each well was constrained for a few days prior to and following the detonation by placing packers just below the static water level. Two of the equipped wells, E-9 and HT-3, were finished in the calcite caprock in the depth interval of 950–1,100 feet, overlying the salt dome and at distances of 1,700 and 1,780 feet from the device hole. Another equipped well, HT-5, 1,800 feet from the device hole is finished in a sand aquifer about 100 feet thick and 200 feet above the calcite caprock. The remaining equipped well, HT-2, is off the dome, 6,180 feet southwest of the device hole. This well taps a limestone and sandstone aquifer in the depth interval of 1,809–2,059 feet.

W. E. Hale, M. S. Garber, and F. C. Koopman report that the initial pressure-pulse response in the different wells began in the time interval from 0.27 and 0.6 seconds after the detonation. The early response was in part cyclic, but a gradual and per-

sistent net increase in water pressure began within a second in the three closer wells. In well E-9, the net increase in pressure, equivalent to 15 feet of water, was reached in about 3 minutes; after a day the pressure began to decline and within a period of approximately 25 days had returned to the predetonation head. A somewhat similar history of water-pressure fluctuations was recorded in well HT-3, but the maximum rise in pressure was only equivalent to 8.8 feet of water. The maximum pressure increase in well HT-5 was equivalent to about 5 feet of water. The inferred cause of the net pressure increase in these three close-in wells is a slight compaction of the aquifer skeleton and concurrent compression of the water in the aquifer, resulting from ground motion caused by the detonation. In time, this water-pressure increase is dissipated by movement of water away from the compressed volume. The compaction of the aquifer, required to give increase in pressures equivalent to 5 to 15 feet of water, ranges between 1 and 2 thousandths of a percent.

In well HT-2, more than a mile from the dome, the initial pressure response was a net decrease equivalent to about 1 foot of water. During the following day the water pressure increased, equivalent to an estimated elevation of 1.7 feet above the pre-explosion level, and thereafter decreased to the preshot level within about 6 days. No interpretation of the pressure response in this well has been made.

Half Moon Creek, which drains the immediate area of Tatum dome, was instrumented with a recording gage at a point near the mouth of the stream about 1.5 miles north of the emplacement hole for the SALMON event. The gage was operated through a period several weeks before the detonation and for several days after the event to record changes in streamflow related to the detonation. Half Moon Creek is tributary to the westward-flowing Lower Little Creek which has several other tributaries in the vicinity of Half Moon Creek. A recording gage about 3 miles northwest of Tatum dome has been in operation on this stream for a few years.

C. P. Humphreys, Jr., and R. E. Taylor report that a temporary but significant increase in discharge past the recording gages on Half Moon Creek and Lower Little Creek occurred following the SALMON event. Prior to the SALMON event, the streamflow at the two recording gages was remarkably stable. No rain had fallen in the drainage basin since the heavy rains accompanying hurricane Hilda on October 4 and 5, 1964. Prior to the detonation

the base flow on Half Moon Creek was about 14 cubic feet per second and that on Lower Little Creek at the gage about 32 cfs.

On October 22, 1964, 1 hour and 35 minutes after the SALMON event, the discharge on Half Moon Creek began to increase. During the following interval of 3 hours and 45 minutes, the discharge increased a total of 1.5 cfs. A nearly steady flow of 15.6 cfs was maintained for the following 13 hours after which the discharge began to decrease. By noon on October 24, the discharge was near the predetonation rate. In all, the increased discharge amounted to about 150,000 cubic feet.

On Lower Little Creek, an increase in the discharge rate began about the same time as that on Half Moon Creek. The discharge increased by about 6.3 cfs during the following 9 hours. Thereafter the discharge began to decrease but did not decline to the predetonation discharge rate before additional runoff occurred in the area from showers. The increased discharge on Lower Little Creek attributable to effects from the SALMON event is estimated to be 1 million cubic feet.

Compaction of saturated near-surface sediments by the seismic motion from the SALMON event probably was the major cause for the observed temporary increased discharge at the two recording stations on the Lower Little Creek drainage basin. The compaction of sediments that produced the nearly instantaneous discharge of water from storage is believed to have been in areas where the water table was very near the surface.

BASIC GEOLOGIC STUDIES

Pahute Mesa, Nevada Test Site

Identification and correlation of subsurface geologic units from additional drill-hole data on Pahute Mesa are revealing an increasingly complex picture of its volcanic stratigraphy and structure. Positive identification of ash-flow tuffs and rhyolitic lavas, from the Black Mountain volcanic center to the west and the Timber Mountain caldera to the south, in the subsurface of Pahute Mesa has outlined further the distribution pattern of the ash-flow tuffs and has provided information about the paleotopographic surface of pre-Timber Mountain time.

A third major volcanic center, probably a caldera buried by the later volcanic flows, recently has been outlined under Pahute Mesa. The east-central part of Pahute Mesa was, for several years, believed to be the center of extrusion for the Belted Range Tuff. Subsurface data from approximately 20 exploratory

holes, gravity data from D. L. Healey and C. H. Miller, and detailed petrographic studies by F. M. Byers and K. A. Sargent, now indicate the existence of a buried probable caldera which centers in the SE $\frac{1}{4}$ of the Silent Butte quadrangle. The caldera is roughly circular with a rim diameter of approximately 12 miles and has a cumulative structural relief of as much as 7,000 feet. Calculation of the volume of rock believed to be related to the main subsidence of the caldera indicates that 80 cubic miles of tuff, mainly Belted Range Tuff, and a nearly equivalent volume of alkaline rhyolitic lavas are genetically related to the buried volcanic center. The lavas appear to be emplaced along ring fracture zones to the east and northeast of the center of the structure. Further work shows that rhyolitic and quartz latitic lava flows and tuffs intermediate in age between the Belted Range Tuff and the Paintbrush Tuff are present in the western part of the caldera and may have originated beneath Pahute Mesa. Limited correlation of the oldest rocks penetrated under Pahute Mesa has been made with ash-flow tuffs and rhyolites to the north in the Kawich, Belted, and Cactus Ranges.

Regional stratigraphy and structure of the Amargosa Desert area

A geologic map of the Amargosa Desert area is being prepared to provide a geologic framework for ground-water studies of the Nevada Test Site and vicinity. The map is being compiled from existing sources and supplementary reconnaissance mapping. The mountains surrounding the Amargosa Desert Basin, as well as the desert floor, that are being mapped include Bare Mountain and the Beatty area in the northwest, parts of the Grapevine, Funeral, and Greenwater Ranges in the west; the Resting Spring Range, northern Nopah Range, and the hills north of Pahrump in the south; and the northern Spring Mountains, Specter Range, and western Spotted Range in the east and northeast.

The stratigraphic section includes, above the crystalline Precambrian basement rocks, a virtually conformable sedimentary sequence of Precambrian to Permian age and includes, at the top, middle and late Tertiary sedimentary and volcanic rocks and Quaternary basin fill. Although thickness changes and facies relations are important in all parts of the Precambrian and Paleozoic sedimentary sequence, the same principal stratigraphic units are recognizable throughout the sequence except for the Mississippian System. In the eastern and southern part of the area, the Mississippian consists largely

of fossiliferous limestone; in the northern part of the area, the sequence is mainly argillite, quartzite, and chert-pebble conglomerate.

Tertiary stratigraphic units, principally ash-flow tuffs, are traceable across the northern part of the area from the Nevada Test Site to the Daylight Pass area in California. These volcanic rocks are underlain by an unnamed sedimentary sequence consisting principally of tuffaceous, locally fish-bearing, shale and sandstone. In the Funeral and Grapevine Mountains, this sequence in turn overlies lower Oligocene conglomerate. Tuffaceous shale and sandstone similar to those beneath the ash-flow sequence occur in the Ash Meadows area. Another Tertiary sedimentary unit, not yet correlated with certainty, is present in the southeastern Funeral Mountains.

The northeastern edge of the area is crossed by the Las Vegas Valley shear zone; the southwestern part of the area includes part of the Furnace Creek fault zone. Both of these structures are northwest-trending right-lateral shear zones. Another parallel but apparently smaller right-lateral fault has been recognized near the northern end of Stewart Valley. The amount of displacement on that fault has not yet been accurately determined, but there appears to be more offsetting by drag along the zone than by the faulting. This right-lateral shear zone projects northwestward through the central part of the Amargosa Desert Basin. The Precambrian and Paleozoic sequence is broken by many thrust faults including several of major displacement. There is some evidence that the crystalline Precambrian basement may have been involved in at least some of the major thrusting. The Tertiary rocks define a basin-range deformational style, with antithetic normal fault blocks tilted generally eastward.

Relation of gravity to mineralized areas, south-central Nevada

Geologic reconnaissance mapping of the Las Vegas Bombing and Gunnery Range by E. B. Ekren, R. E. Anderson, and C. L. Rogers has delineated a widespread Tertiary ash-flow tuff sheet that rests on an erosional surface of considerable relief. Several topographically high areas of hydrothermally altered and mineralized rocks were not covered by this tuff sheet and have been prospected and mined in the past. It seems likely that other mineralized areas may be concealed by this post-ore tuff sheet and by alluvium.

Studies made by D. L. Healey and others show that gravity anomalies here, as elsewhere in the

Basin and Range province, are generally high over the ranges and low over the basins. Gravity-high anomalies also correlate closely with known mineralized areas (Goldfield, Gold Crater, Cactus Range, and south of Cactus Range); this close correlation leads to the conclusion that additional mineralized areas may be postulated from the gravity data. One such anomaly is centered in Stonewall Flat, which is a structural basin. The surface geology of the areas surrounding Stonewall Flat offers no clues either to the presence of a structural high or to the rock type causing the anomaly. Depth calculations based on an aeromagnetic survey of the Las Vegas Bombing and Gunnery Range indicate a depth of 300 to 500 feet to the top of the rocks causing the gravity anomaly (G. D. Bath, written communication, 1964). The gravity high is apparently an expression of a structural dome that may reflect a buried volcanic center similar to that at Goldfield or Tonopah.

Magnetic properties of volcanic rocks, Nevada Test Site

Measurements by G. D. Bath of the remanent magnetization of volcanic rocks from the Nevada Test Site tend to confirm the current hypothesis that during geologic time the geomagnetic field has undergone periodic reversals. A rock cooling in the earth's magnetic field acquires a magnetism direction that is parallel to the field, and therefore a record of the fossil direction of the field is preserved in the magnetic properties of the rock. Investigations of properties of welded tuff, rhyolite, and basalt units reveal uniform directions of magnetism, but these directions may be either along or opposite to the present geomagnetic field.

The presence of rocks with normal and reversed polarities provided a basis for stratigraphic correlation that was applied successfully by (1) measurement of magnetic properties in the field, and (2) interpretation of the aeromagnetic survey of the Nevada Test Site. After establishing polarities for many important units, rapid techniques were used to determine polarities in the field while geological studies were underway. Strongly magnetized and thick units with reversed polarities give distinctive magnetic anomalies, and in several instances the effects of a particular unit could be seen in the aeromagnetic data. For example, the negative anomalies that follow the Reveille Range come from the tuff of White Blotch Spring, and those along the Kawich Range come from the tuff of Trailer Pass. Many of these anomalies arise from source rocks that are covered by alluvium or younger strata. Here the

anomalies not only identify but also give the approximate depth of the magnetized rock.

K. A. Sargent (p. B142-B145) found that magnetic-susceptibility data combined with grain-density measurements provided a useful tool for distinguishing between lithologically similar olivine basalt flows.

Zeolite studies, Nevada Test Site

X-ray diffraction studies by D. L. Hoover and A. O. Shepard of samples from drill holes and outcrops indicate that zeolite zoning is persistent throughout much of the Nevada Test Site. Chabazite forms an upper, sporadic zone. Clinoptilolite, mordenite, and analcime form progressively deeper, but overlapping zones on Pahute Mesa and Rainier Mesa. The mordenite zone has not been found in Yucca Flat, Frenchman Flat, or Jackass Flat. Thinning of zones with thinning sections of rocks susceptible to zeolitization and the presence of the same zones at greatly differing depths indicate that the zoning is probably related to downward changes in ground water rather than to temperature and pressure.

Compilation of bulk densities of vitric rocks and their zeolitized equivalents has shown that bulk densities of ash-fall tuffs increase and that those of dense lava flows decrease during zeolitization. Chemical analyses of selected pairs of vitric and zeolitized rocks corrected for differing bulk densities show not only major differences in Si, K, Na, Ca, Mg, and H₂O, but also 10 to 20 percent differences in Al.

DISPOSAL OF RADIOACTIVE WASTES

Research related to the disposal of radioactive wastes included (1) theoretical and analytical studies of flow phenomena controlling the time and space distribution of radiocontaminants in surface streams and through geologic media; (2) studies of the fate of radioactivity discharged from waste-generating facilities; and (3) geohydrologic evaluations of existing and planned nuclear installations. Most of the work was done in cooperation with the U.S. Atomic Energy Commission.

TRANSPORT OF RADIONUCLIDES

Diffusion in surface streams

In an exploratory laboratory study of lateral turbulent diffusion at the surface of an alluvial channel, W. W. Sayre and A. R. Chamberlain (1-64), by studying the spread of small floating polyethylene particles, demonstrated a means of evaluating successfully the lateral diffusion coefficient and of quantitatively estimating fundamental turbulence prop-

erties such as the intensity and LaGrangian integral scale of turbulence in an alluvial channel. They found that the coefficient of lateral turbulent diffusion increases initially with increasing distance downstream from the source, but that it tends toward a constant limiting value in a distance that is very small compared to the length scale of most diffusion phenomena in natural streams of practical interest.

Transport through earth materials

The transport of radionuclides through earth materials is a complex physicochemical phenomenon. Akio Ogata (1-64) studied the mechanics of dispersion analytically and experimentally and defined the problem in terms of the following major factors controlling dilution: (1) mechanical or hydrodynamic dispersion, (2) adsorption, and (3) heterogeneity. Correlation between experimental and theoretical results for the spread of a tracer was fair.

Ogata (2-64) also contributed to the solution of waste-transport problems in a theoretical analysis of the effect of a reacting medium on the dispersion of fluids. He considered the effects of the dispersion process to be of the same order of magnitude as those of the adsorptive process and, using an assumed linear adsorption isotherm, developed solutions to the differential equations that describe the two processes.

STUDIES AT NUCLEAR-ENERGY FACILITIES

Radioactivity in water, Brookhaven National Laboratory, New York

Wallace de Laguna (1-64) has studied chemical and radiochemical analyses of more than 300 water samples collected at Brookhaven National Laboratory, New York, between 1948 and 1953. The samples were intended to provide norms for problems of water contamination that might arise in the future. No samples of ground water showed alpha activity; only 14 showed beta activity, the maximum being 5×10^{-8} microcuries per milliliter. Beta activity found in 55 surface-water samples (maximum of 3.7×10^{-7} microcuries per milliliter) is attributed to fallout from nuclear-weapons tests. Widespread contamination of ground water from the leaching of fertilizer in intensively farmed areas, and local contamination from cesspools, are indicated by high nitrate concentrations in ground water.

Permeability of weathered crystalline rocks in Georgia

At the Georgia Nuclear Laboratory J. W. Stewart (2-64) studied the infiltration characteristics and

permeability of weathered crystalline rocks in relation to the use of seepage pits for disposal of low-level liquid radioactive wastes. Water was found to move along the schistosity, mostly along the strike but also along (up and down) the dip. Computations of ground-water velocity based on Darcy's law indicated rates of flow of 0.1 foot to 3 feet per day. Rates measured with chloride and nitrate tracers ranged from 0.4 foot to as much as 25 feet per day.

Extensive unsaturated deposits at National Reactor Testing Station, Idaho

The subsurface geology of the National Reactor Testing Station, in Idaho, was described by E. H. Walker (3-64). The station is underlain, to depths still unknown, by basalt flows of the Snake River Group. The basalt is overlain by surficial unsaturated unconsolidated sedimentary rocks more than 50 feet thick under about 175 square miles of the station; a total of more than 4 cubic miles of such material lies beneath the station. Since most of these materials are fine grained, they can be used in the relatively dry Idaho climate as temporary storage reservoirs for low-level liquid radioactive wastes. However, this practice is followed in only a very small part of the station.

FATE OF RADIOACTIVE LIQUID EFFLUENT

Radionuclides in Columbia River and Columbia River estuary

In the Columbia River below Pasco, Wash., W. H. Haushild, H. H. Stevens, Jr., and G. R. Dempster, Jr., found that about 90 percent of the combined annual load of the 6 dominant radionuclides is transported in solution and 10 percent is transported sorbed on sediments. However, for certain radionuclides, the annual load transported by sediments exceeds the load transported in solution by as much as 570 percent. A gross budget of the radionuclide load in the Columbia River indicates that the total load at Vancouver, Wash., is about 75 percent of the total load at Pasco, Wash. (Vancouver is 224 river miles downstream from Pasco). That part of the radionuclide load at Pasco which is not transported to Vancouver or accounted for by loss by radioactive decay varies with season and is largest during the annual spring high water.

In their studies of radionuclide movement in the Columbia River estuary, D. W. Hubbell, E. A. Prych, and J. L. Glenn have developed (1) a vibratory sampler for obtaining cores in sandy bed material, (2) a system for sampling large volumes of water-suspended sediment mixture at stream velocity, and

(3) a sled-mounted radiation-detection system for continuously measuring radioactivity in the bed along traverse lines. Radiation measurements indicate that the most abundant radionuclides in the bed material of the estuary are chromium-51 and zinc-65. The level of activity varies widely from place to place, and the highest activities are associated with the finest bed materials.

Radionuclides in the Clinch River, Tenn.

In an inventory of principal radionuclides in bottom sediments of the Clinch River downstream from White Oak Creek, P. H. Carrigan, Jr., found that 77 percent of the identified radioactivity was cesium-137 (155 curies). Quantities of other major radionuclides were: cobalt-60, 17.5 curies; ruthenium-106, 15.5 curies; strontium-90, 2.9 curies; and about 10 curies of rare earths. Approximately 95 percent of the radioactivity is in the 15 downstream miles of the 21-mile reach studied.

R. J. Pickering found that the vertical distribution of cobalt-60 in cores of Clinch River bottom sediment is similar to that of cesium-137. The pattern of distribution of cesium-137 reflects the pattern of annual releases of waste, but that of cobalt-60 does not. Hence, it is inferred that, unlike cesium, most of the cobalt-60 released to the river is not associated with sizes of suspended solids that will settle.

E. A. Jenne and S. S. Wahlberg report that most of the cobalt-60 scavenged by the sediment of White Oak Creek, Tennessee (through which low-level waste from the Oak Ridge National Laboratory passes to the Clinch River) is found in a metal-oxide reducible fraction. Reducible manganese and iron oxides contain approximately equal amounts of the scavenged cobalt-60. A small portion of the cobalt-60 is contained in an organic fraction.

Disposal of liquids in Bandelier Tuff

According to W. D. Purtyman and F. C. Koopman, the physical and hydrologic characteristics of the Tshirege Member of the Bandelier Tuff in New Mexico are favorable to the injection and storage of liquid wastes such as treated low-level radioactive wastes. An injection rate of about 28,400 gallons per day into a moderately welded tuff would be possible by use of a properly perforated 12-inch-diameter well 100 feet deep.

FLOW OF AIR IN TUFF

At the Los Alamos Scientific Laboratory, in New Mexico, the movement of air into and out of the ground under natural barometric gradients is being

studied in an effort to establish the feasibility of using the Bandelier Tuff for storage of gaseous radioactive wastes. The duration of air exhaust or intake cycles indicates that the pressure within the tuff does not adjust readily, but that instead the rock is compressed or expanded with these changes. Analysis of the effects of barometric pressure changes has also shown that there is a deformation of the rock caused by tidal effects. This deformation causes a change in the void ratio of the formation and influences the flow of air to or from the ground.

WATER-CONTAMINATION STUDIES

Development of water resources is restricted or complicated in many areas by the spread of waste products into water supplies. The continuing growth of population and industry leads to the production of an increasing volume and variety of waste products, and to increasing problems of contamination.

Patterns of contaminated zones in ground

The patterns of zones of ground water contaminated by wastes placed in the ground have been studied by H. E. LeGrand (2-65). Many factors, including the degree to which each contaminant decreases in potency by (1) decay or some other die-away mechanism, (2) sorption on earth materials, and (3) dilution, determine the size and shape of contaminated zones in the water-table aquifer. Most contaminated zones do not extend to wells or streams, although contaminated waters from some waste sites do reach enough wells to cause problems of major concern. Where the decrease in potency is only by dilution, the contaminated zone is likely to be greatly elongated in the direction of groundwater flow, and the contaminated water may reach perennial streams. In spite of the fact that many hundreds of thousands of contaminated zones occur in the water-table aquifers of the United States, only a few have been mapped.

Amino acid content of sewage greatly reduced by primary treatment

Raw sewage entering the Denver sewage treatment plant, and effluent from sewage-treatment facilities at Denver, Colo., Chicago, Ill., Trenton, N.J., and Hamilton County, N.J., were analyzed for amino acid content by Lloyd Kahn and C. H. Wayman (1-64). Total removal of amino acids in sewage was accomplished by means of activated-sludge treatment; primary treatment alone removed 65 percent of the proteinaceous matter. It is expected that even primary sewage treatment will sufficiently lower the

amino acid and protein content in sewage plants to prevent contamination of surface waters.

Long-term changes in quality of Passaic River, N.J.

P. W. Anderson and S. D. Faust (p. D214–D218) showed, by analysis of streamflow and chemical-quality data for the Passaic River at Little Falls, N.J., a general increase in content of dissolved solids and a 20-percent decrease in dissolved-oxygen content over the period 1947–64. They attributed this deterioration of quality to the disposal of increasing volumes of municipal and industrial wastes in the river.

DETERGENTS AND PESTICIDES IN WATER SUPPLIES

LABORATORY STUDIES

Bacterial degradation of ABS

A study by C. H. Wayman, J. B. Robertson, and C. W. Hall (2–64) of bacterial degradation of detergent compounds showed that alkylbenzenesulfonate (ABS) with a branched alkyl chain was not degraded significantly under either aerobic or anaerobic conditions. ABS with a straight alkyl chain, the new biodegradable type, was degraded very slowly under anaerobic conditions but somewhat more rapidly than the branched-chain ABS. The straight-chain ABS was degraded rapidly under aerobic conditions and in some instances was totally destroyed in 2 days. Degradation in all instances occurred faster at 35°C than at 20°C or 10°C. The study indicated that the new straight-chain ABS would be broken down in aerobic sewage-treatment processes but that in anaerobic systems it would be little broken down and would be a potential water pollutant.

Adsorption of ABS on clay minerals

A report by C. H. Wayman and J. B. Robertson (3–64) summarized a study of the adsorption of the common detergent compound, alkylbenzenesulfonate (ABS), on the soil clay minerals kaolinite, illite, and montmorillonite. The study showed that these minerals adsorbed only minor amounts of ABS as compared with activated carbon or colloidal alumina. Adsorption increased with increasing ABS concentrations, according to Freundlich or Langmuir isotherms. ABS with a C₁₅ alkyl chain was adsorbed more readily than that with a shorter C₁₂ chain. Adsorption was stronger at pH 4 than at pH 7 or 10. The presence of chloride salts of Al³⁺, Ca²⁺, and Na⁺ in solution increased ABS adsorption, the AlCl₃ salt having the greatest effect. It was con-

cluded that the minerals studied would not adsorb significant amounts of ABS under natural conditions.

Hydrogen isotopic-exchange reaction in the system aldrin–water

M. C. Goldberg and R. L. Wershaw have studied the hydrogen isotopic-exchange reaction in the system aldrin–water in order to determine whether there is detectable ionization of aldrin. The data obtained indicate that if aldrin ionizes, its ionization constant is less than 10^{–30}. Cyprax, a water-soluble fungicide that has detergent properties, was shown to be utilized as the sole source of carbon by an *Achromobacter* species and by a *Flavobacterium* species of soil bacteria. Degradation of this compound by soil micro-organisms was substantiated by exposing C¹⁴-labelled cyprax to river muds collected in the Denver, Colo., area. The bioactive river muds metabolized solid-state cyprax to CO₂, whereas sterile muds did not.

FIELD STUDIES

Movement of detergents in ground water

An investigation by J. C. Chemerys of the movement of contaminants from septic tanks in housing developments near Raleigh, N.C., suggests that chloride and nitrate ions move farther in the ground-water flow system than do detergents. However, in the areas studied, none of these constituents were found in concentrations approaching maximum safe concentrations recognized by the U.S. Public Health Service for public water supplies.

ACID MINE WATER

Anthracite mine drainage in Pennsylvania

Monitoring of the gravity overflow of mine water from water-filled anthracite coal mines under Scranton, Pa., for 3 years indicates that the total load of dissolved solids in the overflow increases after the levels of the mine-water pools rise and the inundated area becomes larger. W. T. Stuart reports that the overflow rate varies seasonally and that the smaller flows are more concentrated than larger flows. The yields of dissolved material decrease with time—rapidly at first and slowly thereafter—apparently as the acid-forming minerals in the mine are depleted. At one site, where the overflow was stopped and impounded for 6 weeks in 1963, the concentration of acid in the water almost doubled, and the sulfate, iron, and manganese increased about 10 percent after the overflow was resumed. The coal-bearing area contributing water to the abandoned mines

is about 56 square miles, and the area of the mine-water pool within the coal-bearing area is 34 square miles. In 1963, discharge of water from the pool area averaged 35,500 gallons per minute, and yielded 15 tons of sulfate per day, 1.1 tons of iron, 0.2 tons of manganese, and an equivalent of 2.6 tons of sulfuric acid if neutralized to pH 8.0.

Acid mine drainage in Maryland

In a reconnaissance study of the chemical characteristics of Maryland streams J. D. Thomas found that acid mine drainage in the headwaters of the Potomac River basin affects the quality of water in the Potomac River. Ground-water samples collected in that area were of the calcium bicarbonate type, but the surface water, of calcium sulfate type, showed the influence of acid mine drainage. In the vicinity of Hancock, in western Maryland, the Potomac water begins to change to a calcium bicarbonate sulfate type because of inflow from the South Branch Potomac River and the Cacapon River, which contain water of the calcium bicarbonate type. Still farther downstream, below the mouths of the Shenandoah River and of Antietam and Conococheague Creeks, the Potomac water is of the calcium bicarbonate type.

Acid mine drainage in Kansas

W. J. Seevers reports that acid mine drainage from abandoned underground coal mines is the principal source of acid pollution in the Cherry Creek drainage basin in Cherokee County, Kans. The principal sources of acid mine drainage were found, by measuring the pH and the water temperature at points along the stream, to be two mine-drainage shafts. An anomalous distribution of alternating riffles and pools, and peculiar meander patterns, found near these shafts appears to be the result of increased acidity. It is believed that the reduction of pH and the increase in streamflow directly below points of effluence cause flocculation and deposition of a portion of the stream load, which results in a change in meander patterns of the stream. Surface runoff from the extensive spoil dumps of strip mines in the Cherry Creek basin contributes only minor amounts of acid water to the streams.

INDUSTRIAL EFFECTS

Thermal loading of a river in Pennsylvania

Dissolved oxygen in the West Branch of the Susquehanna River in Pennsylvania was investigated by K. V. Slack and F. E. Clarke (p. C193-C195) at stations above and below a steam-electric power-

plant. The river receives strip-mine drainage and moderate quantities of municipal wastes above the powerplant. When studied in October, a diurnal fluctuation in dissolved-oxygen content occurred only above the plant. Tests with river water in light and dark bottles indicated no production of oxygen above the powerplant and a net loss of oxygen below the plant. A significant quantity of oxygen is thought to be added to the river above the powerplant by streambed plants but not by phytoplankton. Below the heated outfall, the reduced abundance of riverbed algae is probably a result of the effects of chemical and fine-sediment wastes from the plant. The shallow depth and slow rate of organic decomposition in the acid water also contribute to high concentrations of dissolved oxygen in this river.

BRINE AND SALINE WATER

Natural brine in Upper Brazos River basin, Texas

An investigation of the occurrence, movement, and discharge of natural brine in the Upper Brazos River basin, Texas, by P. R. Stevens and W. F. Hardt has shown that natural brine occurs at relatively shallow depth in rocks of Permian age throughout the area. The brine in most places is overlain by relatively fresh, circulating ground water. A sharp interface separates the brine and fresh water into discrete hydraulic units. The freshwater head influences the configuration of the interface but does not control flow conditions in the brine system, which is relatively static compared to the overlying system of circulating fresh ground water. Regional topography, climate, and geology are the principal controls on the movement and discharge of the brine; circulation of infiltrated rainwater in local systems is a minor factor in the evolution of the brine. Discharge of brine occurs where the interface between brine and fresh water intersects the land surface. Methods to reduce or prevent discharge of natural brine into the Brazos River and its tributaries, and thereby to improve the utility of Brazos River water, are receiving great impetus. The investigation thus far has shown that remedial measures based upon impoundment of the natural brine in its area of discharge, and other control measures based upon the assumption that brines are generated by circulation of infiltrated rainwater in local systems, cannot be effective.

Chemical quality of water in Neches River basin, Texas

In a reconnaissance of the chemical quality of surface water in the Neches River basin in Texas, L. S. Hughes and D. K. Leifeste found that streams drain-

ing a part of the East Texas oil field contained water of high chloride content and great acidity, with pH as low as 3.2. They showed that the source of the acidity could be hydrogen ions adsorbed on clay minerals near the land surface and transferred to shallow ground water as the result of base exchange with oil-field brine. Adsorbed hydrogen ions can be displaced from clay when the latter is wetted by a solution containing a preponderance of sodium ions. When clay samples from the area were leached with oil-field brine, which was neutral (neither acid nor alkaline), they yielded acid solutions similar in chemical composition to the acid water in the streams.

Contamination by brine in Caldwell County, Tex.

On the basis of a study of the ground-water resources of Caldwell County, Tex., C. R. Follett reports an example of contamination of fresh ground water by oil-field brine, and the later improvement in the quality of the water by stopping the contamination at the source. As early as 1928, brine which had been produced with oil was discharged on the surface of the Leona Formation, a highly permeable water-bearing unit. Contamination over a period of several years increased the chloride content of the water in the Leona in a triangular-shaped area whose apex was at the source of contamination and which extended at least $2\frac{1}{2}$ miles downgradient. The contaminant evidently traveled the $2\frac{1}{2}$ miles in less than 10 years, for in 1938 the chloride content of water from a well in the Leona, $2\frac{1}{2}$ miles from the source of contamination, was 211 parts per million, or more than twice the content of the normal uncontaminated water. By 1943, the chloride content of water in that well had increased to 604 ppm. After the source of contamination was discovered the oil field was closed and facilities were installed for injection of the brine into deep salt-water sands. After the contamination was stopped at the source the quality of the ground water steadily improved, and by 1963 the chloride content was 101 ppm—a quality approaching that of the normal uncontaminated water.

Salt load of Pecos River reduced by pumping from aquifer

E. R. Cox and J. S. Havens report that the salt load has been reduced in the Pecos River downstream from Malaga Bend in Eddy County, southeastern New Mexico, by the Malaga Bend Experimental Salinity Alleviation Project. The 3-mile reach of the river at Malaga Bend is the natural discharge area for an artesian-brine aquifer, at a depth

of about 200 feet, near the base of the Rustler Formation of Permian age. Brine from a well that taps the brine aquifer is now pumped to a nearby depression at a rate of 450 gallons per minute. The head in the aquifer has been lowered, and the brine-discharge rate has been reduced. The average downstream gain in chloride load in the Malaga Bend reach of the river declined from 168 tons per day (July 16–17, 1963, before pumping began) to 59 tons per day (August 12–13, 1964).

Brine contamination in Morrow County, Ohio

Contamination by oil-field brine was observed in Morrow County, in north-central Ohio, during 1964. According to P. G. Drake, Whetstone Creek, a tributary of the Scioto River which drains the area of greatest oil production, carried chloride concentrations of more than 700 parts per million during low-flow periods. Prior to oil-drilling operations the concentrations were about 30 ppm. Ground-water contamination also was observed, particularly in shallow wells adjacent to brine drainage pits and ditches. Two public-supply wells of the City of Cardington, in which chloride concentrations of 3,750 ppm and 500 ppm were observed, were shut down because of the brine contamination.

Saline ground water migrates upward into Ocala Limestone at Brunswick, Ga.

According to R. L. Wait, salt water (with chloride concentrations as high as 1,000 parts per million) is present in the Ocala Limestone in a small area at Brunswick, in southeastern Georgia, between depths of 560 and at least 1,300 feet below the surface. Fresh ground water occurs in the overlying Miocene rocks and in part of the underlying middle Eocene limestone. Evidence from C-14 analyses presented by B. B. Hanshaw, William Back, Meyer Rubin, and R. L. Wait (1–65) confirms earlier conclusions that the saline water is derived by upward flow from the underlying Claiborne rocks rather than from the ocean. This upward movement of salt water is also confirmed by a water-table decline map prepared by D. O. Gregg. Northward movement of the 150-ppm isochlor suggests movement of about half a mile since late 1962. An increased hydraulic gradient resulting from an increase of 36.7 million gallons per day in pumpage in late 1962 has caused the salt water to move more rapidly than previously.

Change in water quality after damming of Econfina Creek estuary, Florida

Water in a saline estuary of Econfina Creek, in Bay County in the Florida panhandle, freshened to

250 parts per million chloride in about 2 months after construction of a dam at Deer Point to form Deer Point Lake. The rate of freshening with respect to the volume of the lake and inflow to the lake was calculated by L. G. Toler, R. H. Musgrove, and J. F. Foster (1-64), and an empirical formula was derived which may have application to other similar water bodies. Some contamination of ground water occurred in the sand aquifer adjacent to the lake, but the saline water was flushed from the aquifer as a result of a general rise in ground-water levels surrounding the lake.

Sources of chloride in ground water, Nassau County, N.Y.

F. A. DeLuca, J. F. Hoffman, and E. R. Lubke have compiled records of chloride concentrations and temperatures of the waters of Nassau County, Long Island, N.Y., from about 1895 to 1962 and have concluded that most of the natural fresh ground water contains 10 parts per million or less of chloride. Ranges in chloride concentration in the ground water sampled were 2-9,200 ppm in the upper Pleistocene deposits, 2-12,900 ppm in the Jameco Gravel, 1-16,700 ppm in the Magothy (?) Formation, and 1-2,800 ppm in the Lloyd Sand Member of the Raritan Formation. Concentrations in excess of 10 ppm generally are attributable to 1 or more of 5 principal sources of contamination: (1) sea water, (2) fertilizer, (3) domestic wastes, (4) industrial wastes, and (5) salt used for dust or de-icing control on roads.

DISTRIBUTION OF MINOR ELEMENTS AS RELATED TO PUBLIC HEALTH

An increasing interest in the distribution of elements in plants, soils, rocks, and water, and the relation between this distribution and problems in public health has been made particularly evident in 1964 by the inclusion of symposia on these subjects at the meetings of both the American Public Health Association, and the American Association for the Advancement of Science. Scientists of the U.S. Geological Survey have been taking leading parts in field and laboratory studies of these relationships, and in arranging interdisciplinary meetings at which the results of studies are presented. Some current work by Survey scientists is summarized below.

Element distribution in plants and soils of New Mexico and Maryland

H. L. Cannon has compared the inorganic compositions of samples of soils and of native and culti-

vated vegetation from San Juan County, N. Mex., (an area probably low in incidence of cancer and heart disease) with the compositions of similar samples from Washington County, Md., (an area of higher cancer and heart disease rate). Iron, titanium, manganese, chromium, copper, lead, potassium and NO_3^{-1} contents are higher in the samples from Maryland.

Radium in ground water in Gas Hills uranium district, Wyoming

R. C. Scott and F. C. Armstrong have collected and analyzed for radium 226 20 samples of water from wells and mines of the Gas Hills uranium district. Three were of water from wells in the Cretaceous Cloverly Formation, 2 from wells in the Pennsylvanian Tensleep Sandstone, and 11 from mines and 4 from wells in the middle member of the Eocene Wind River Formation, the host rock of the uranium deposits.

Analyses of the samples by the Water Resources Division of the U.S. Geological Survey show that in this area the Cloverly is a source of water suitable for continuous domestic use. Ra^{226} content of water from the Cloverly ranged from 0.7 ± 0.1 to 1.8 ± 0.4 picocuries per liter (picocuries per liter = micro-microcuries per liter). The maximum permissible concentration of Ra^{226} recommended by the U.S. Public Health Service in domestic water for continuous use is 3 pc/liter.

In contrast, analyses of water from the Tensleep and Wind River Formations show that in this area these waters, with one exception, are suitable for industrial use, such as process water in the uranium mills, but not for continuous domestic use. The Ra^{226} content of the two samples from the Tensleep was 69 ± 14 and 120 ± 20 pc/liter. The Ra^{226} content of well water from the Wind River Formation in areas believed to be devoid of uranium mineralization ranged from 5.1 ± 1.0 to 23 ± 5 pc/liter. The Ra^{226} content of 10 mine-water samples from the Wind River Formation ranged from 70 ± 14 to $2,400 \pm 500$ pc/liter. One sample of water from the Wind River Formation, which contained only 1.9 ± 0.4 pc/liter of Ra^{226} , suggests that in some places in the Gas Hills the Wind River Formation may be a source of water suitable for continuous domestic use. The other samples from the Wind River Formation indicate, however, that water from this formation should be analyzed before use as a domestic supply, and that thereafter the supply should be sampled and analyzed periodically.

PROBLEMS IN ENGINEERING GEOLOGY AND HYDROLOGY

Geology of tunnels along Interstate Highway 70 in Colorado

Analysis of geologic and engineering data from the pilot bore for the Straight Creek highway tunnel beneath the Continental Divide shows that the magnitude of the rock load to be supported is directly related both to geologic factors and to engineering practices in tunneling. According to C. S. Robinson and F. T. Lee, geologic factors include density of fractures, amount and kind of clay minerals produced by alteration, and the width of fractured and altered zones. Tunneling practices resulting in excessive overbreak, a decrease in the rate of advance of the heading, and reexcavation of parts of the bore because of inadequate support all tend to increase rock loads over those predictable from the geologic conditions. The quantity of ground water entering the pilot bore was least in zones of alteration and where fractures are closely spaced, and greatest in relatively competent rock adjacent to sheared and altered zones. During construction of the pilot bore, fault gouge containing clay with high swelling pressures was encountered. Subsequent tests of this gouge in the laboratory of the Engineering Geology Branch and in the sedimentary mineralogy laboratory revealed a direct correlation between estimated montmorillonite content and calculated swelling pressures. If such tests were made during tunneling, the need for additional tunnel support might be anticipated.

Geologic mapping by M. H. Bergendahl has revealed that a shear zone consisting of highly fractured granite will be crossed for a distance of about 1,100 feet by a proposed highway tunnel on a projected route of Interstate Highway 70 through the Gore Range. If the zone continues at depth, it may cause problems in excavation and support of this part of the tunnel.

Forecast of coal-mine bumps at Sunnyside, Utah

General areas and times of severe bumps in the coal mines of the Sunnyside district between November 17 and 23, 1964, were forecast 4 days in advance by F. W. Osterwald and C. R. Dunrud. The forecast was based on a very pronounced increase in number and amplitude of seismic tremors recorded November 10 and 11 on the Sunnyside seismograph network. During this period the foci of many of the large-amplitude tremors were apparently along two faults 5 to 7 miles from an area later affected by severe bumps. On November 12, dangerous bumps

were forecast when epicenters of large-amplitude tremors shifted to map locations above abutment pillars and mined-out areas within the Sunnyside coal mines. These large tremors continued through November 16 and coincided with a period of decreasingly abundant but increasingly strong tremors. Dangerous bumps then followed in active mining areas and along haulageways from November 17 to 23.

Seismic tremors were very abundant during the first 3 months of 1965, when 215 tremors ranging in magnitude from 1.3 to 4.0 originated within or near the Sunnyside mining district. An increasing proportion of the epicenters of tremors larger than magnitude 2.5 were along or near faults 1 to 10 miles from the closest mine workings. Many tremors with epicenters near faults in the mining area failed to cause noticeable damage in the mines, which suggests that their foci probably were below the mine level.

Gravel deposits in Beltsville quadrangle, Maryland

Geologic mapping by C. F. Withington (p. B39-B42) in the northeastern part of the Washington, D. C., metropolitan area has outlined the distribution of gravel deposits within the Patuxent Formation of Early Cretaceous age. Knowledge of the distribution of this important resource will help insure its utilization before areas underlain by it are preempted by highways and housing developments.

Rockfalls, debris flows, and eruptions at Mount Rainier, Wash.

Rockfalls in December 1963, possibly triggered by a small steam explosion on the northeast side of Mount Rainier volcano, caused at least seven avalanches of rock debris. Some of these moved as far as 4.3 miles at a velocity of perhaps 100 to 300 miles per hour, and ended as close as 2,000 feet from an unoccupied campground in Mount Rainier National Park. Long distance of movement on a low gradient at the downvalley end of the avalanches was attributed by D. R. Crandell and R. K. Fahnestock to cushions of compressed air being formed beneath the debris as it hurtled off the steep front of a glacier.

Studies by D. R. Crandell and D. R. Mullineaux of debris flows, lava flows, and deposits of pumice and pyroclastic rock debris at Mount Rainier indicate that debris flows present the greatest hazard to man. If the spasmodic postglacial activity of the volcano continues, an eruption of pumice, lava, or rock debris may be expected no more often than about once every thousand years. Debris flows are

more hazardous because they are more frequent and some may travel many miles down valley floors where human habitations, campgrounds, and highways are concentrated.

LAND SUBSIDENCE

Studies of land subsidence caused by decline in artesian head are continuing in California, Nevada, Arizona, and Texas. These studies are contributing to knowledge of the physical and hydraulic properties of leaky aquifer systems, the storage characteristics of semipervious interbeds and confining beds, the change in the coefficient of storage with time and change in effective stress, and the inability of most well casings to resist the compacting forces.

Tilting and subsidence near Bakersfield, Calif.

According to B. E. Lofgren, the analysis of leveling of bench marks in the mountains south of the San Joaquin Valley, resurveyed periodically by the U.S. Coast and Geodetic Survey, indicates tectonic movement in this area where one of California's major earthquakes occurred in 1952. Available leveling data suggest a continuing tilt that increases northward toward the valley from an axis of flexure 2 miles south of Grapevine. This gradual tilting of the mountain block, together with the possibility of differential subsidence due to ground-water pumping in the valley, is of prime concern in the design and construction of two large pumping plants and an aqueduct system of the State in the immediate area.

Subsidence and earth cracks in southern Arizona

In the Salt River valley and at several other localities in southern Arizona, according to William Kam (p. B122-B125), decline of the water level has caused compaction of fine-grained valley fill and land-surface subsidence. He suggests differential compaction of such deposits as a possible cause of associated earth cracks. The sudden occurrence of these fissures intersecting natural drainage courses establishes new, locally lowered base levels, and permits increased headward erosion of the upstream segments of the drainage. Headward cutting along the drainageways produces gullies.

D. E. Peterson (1-64) in a study of earth fissures in the Picacho area of Pinal County in south-central Arizona, reports that the fissures appear to be the result of tensile and shear forces, with shear forces predominating, which are related to differential subsidence of the valley alluvium. The mountains and alluvium-covered pediments have remained stable while the valley alluvium has subsided. The subsidence has been caused or accelerated by large

changes in hydraulic head across fine-grained materials, largely silt and clay, which are the predominant sediments in the basin below a depth of approximately 600 feet. The horizontal permeability of these sediments appears to be significantly greater than the vertical permeability. As a consequence, the hydraulic gradients caused by intensive pumping increase the effective load on the fine-grained materials, thereby creating the compacting forces. Releveling of a U.S. Coast and Geodetic Survey level line between Picacho Peak and the Casa Grande Mountains by the U.S. Bureau of Reclamation in October 1964 suggests maximum subsidence of 5.1 feet since 1962. The bench mark of maximum subsidence is C 279, about 1½ miles northwest of Eloy. This is an area of intensive pumping and declining water levels. Subsidence has been occurring in this area for at least 20 years, but the rate apparently has accelerated considerably.

Rate of subsidence increasing in Houston area, Texas

R. K. Gabrysch has analyzed releveling by the U.S. Coast and Geodetic Survey in 1964 and reports that subsidence has accelerated slightly in the period 1959-64. Subsidence of the land surface in the Houston Ship Channel area increased from a rate of 1.0 foot per 5-year interval (1954-59) to 1.2 feet per 5 years (1959-64). Introduction of surface-water supplies to this area in 1954 permitted a reduction in pumpage. Recovery of water levels of as much as 25 feet between 1954 and 1959 occurred a few miles west of the ship-channel area. From 1959 to 1964, however, levels declined as much as 30 feet. It is not likely that there will be appreciable decrease in the present rate of decline of water levels in the next few years, because additional surface water will not be available for several years.

EVAPORATION SUPPRESSION

Monomolecular film increases net seepage rates from stock pond

A field test using monomolecular films to control evaporation from a 1-acre stock pond near Laredo, Tex., has been in progress since April 1964. The monomolecular films are formed from an aqueous dispersion of eicosanol and docosanol alcohols, and are dispensed from a new type of underwater dispenser which was designed by G. E. Koberg. Evaluation of the effectiveness of the film, by a comparison of the water budgets of the treated pond and of an untreated pond nearby, indicates that in this experiment the net seepage rates from the treated pond are increased more than the amount by which

the film reduces evaporation. A comparison of the water-surface temperatures of the two ponds shows that the treated water is generally 2°C warmer than the untreated water, which indicates an approximate reduction in evaporation of 25 percent. The increase in the net seepage rate for the treated pond is attributed to the transport of some alcohol into the ground with the water, and to the consequent reduction in surface tension and in resistance to flow as the alcohol moves through the voids. This effect seems to be greater in incompletely saturated ground (such as that inundated by a sudden rise in pond level) than in saturated ground, and G. E. Koberg believes that introduction of surfactants may present a possible method of increasing infiltration rates for short-duration storms.

FLOODS

Three major categories in the study of floods by the U.S. Geological Survey are (1) measurement of stage and discharge, (2) definition of the relation between the magnitude of floods and their frequency of occurrence, and (3) delineation of the extent of inundation of flood plains by specific floods or by floods having specific recurrence intervals. The following section, accordingly, is subdivided into discussions of outstanding floods of 1964–65, flood frequency, and flood mapping.

OUTSTANDING FLOODS OF 1964–65

Floods of December 1964 in far western States

The floods of December 1964 in the far western States, the most damaging in the history of the area, were outstanding for record-breaking peak discharges and for the unusually large area involved—Oregon, northern California, western Nevada and Idaho, and southern Washington—according to S. E. Rantz and A. M. Moore (2–65). The floods resulted from a series of storms in late December, but primarily from warm torrential rainfall of December 21–23. They took a total of 47 lives and caused damage of almost half a billion dollars.

Precipitation in the form of rain, ranging from 20 to 30 inches, fell in the Sierra Nevada up to altitudes of 10,000 feet during the major storm period, December 19–23, 1964. In Idaho, Washington, and parts of Oregon, melting snow augmented rain that fell on frozen ground. In December 1955 most of the same streams had rampaged wildly to create unprecedented disaster. The 1964 floods did not extend as far south as did those of 1955, but they covered a much larger area in Oregon and Idaho, and peaks at some gaging stations in the flood area were

considerably greater than those for the 1955 flood. Peaks on some streams in 1964 closely approached or even exceeded those from the almost-legendary floods of 1861–62, which occurred before the region was heavily settled.

Severe floods in Oregon, January 1965

Heavy rain and snowmelt, a repetition of conditions that had caused the disastrous floods in December 1964, again produced severe floods in Oregon during the last week in January. Peaks at some sites exceeded the December peaks, but major flooding in January was not areally so extensive as that during December.

Floods of April 1964 in Mississippi

Intense rain that totalled as much as 12 inches in a 12-hour period on April 5–6 fell in a band 10 miles wide that extends 60 miles across east-central Mississippi into Alabama. The rain produced the highest floods known on many small streams which drain areas covering as much as 500 square miles. Damage to highways and railroad fills was extensive. Peak discharge exceeded the 50-year discharge at many gaging stations.

During a 29-hour period on April 26–27, 11.69 inches of rain fell at Gulfport, and similar amounts fell along the coast from New Orleans, La., to Mobile, Ala. The recurrence interval for the peak discharge at many measured sites ranged between 25 and 50 years.

Floods of May 25 and June 16, 1964, in Nebraska

Heavy rains occurred May 25 in a band across eastern Nebraska from Genoa to Plattsmouth. The storm center was 8 miles northwest of Genoa, where unofficial reports indicate that a total of 8 to 9 inches of rain occurred during a 2-hour period. Emil Beckman reports peak runoff rates ranging from 1,180 to 1,670 cubic feet per second per square mile from small drainage basins. On June 16, 1964, thunderstorms extended along a path 10 to 20 miles wide across northeastern Nebraska from Boyd County to Cass County, a distance of over 200 miles. Hail, wind, and flood damage were locally severe. The most intense precipitation occurred in the Omaha metropolitan area, where 7 lives were lost and estimated property damages exceeded \$6 million. The maximum precipitation totals and peak runoff rates for these storms indicate recurrence intervals much greater than 100 years.

Floods of June 1964 in northwestern Montana

Floods that greatly exceeded previous maxima from mountain areas in northwestern Montana oc-

curred on June 8–10, 1964. The recurrence interval for peak flows at many sites exceeded 100 years. Antecedent conditions which set the stage for extreme floods were a snow cover 25 to 50 percent thicker than the average, below-average runoff in April and May, and extremely heavy rainfall concurrent with a rapid rise in temperature. During a 36-hour period, rainfall (locally as much as 16 inches) was centered along 100 miles of the Continental Divide in the northern half of Montana. Dams on two irrigation reservoirs failed, adding their contents to streams already in flood. The death toll reached 34, and property damage was estimated to be greater than \$62 million.

Texas floods of September 1964

Torrential rains on September 19–27 caused flooding in parts of eastern and southwestern Texas. Upper White Rock Creek at Dallas had record-breaking floods. On September 19–20 as much as 17 inches of rain fell in the upper Nueces River basin, causing near-record flooding.

Floods of September 1964 in eastern Virginia

Rains from tropical storm Cleo caused widespread flooding in eastern Virginia. The Hampton Roads area was swamped on September 1 following extremely heavy rains of as much as 13.7 inches in 24 hours at nearby Virginia Beach. Two persons lost their lives, and property damage was extensive. Elsewhere, flooding occurred in the Dan, middle Roanoke, and Chowan River basins. Several bridges on secondary roads were destroyed.

Hurricane floods of October 1964 in North Carolina and Louisiana

Rains associated with two great storms—hurricane Hilda, and hurricane Isabel—caused record-high and near-record floods on many streams, mostly those rising in the mountains of western North Carolina. More than 17 inches of rainfall occurred at Rosman, N.C., in the French Broad River basin, on October 4–5, during hurricane Hilda. The recurrence interval of flood peaks ranged in general from 50 to greater than 100 years.

According to a report by A. S. Lowe (1–64), torrential rains accompanying hurricane Hilda caused a flood on October 3–4 near Baton Rouge, La., that inundated an area similar to that inundated by floods of April 27–29, 1962. The recurrence interval for the peak discharges is about 50 years. Peak stages for the 1964 flood were higher at some points and lower at others than stages for the 1962 flood.

FLOOD FREQUENCY

Nationwide flood-frequency reports

Thirteen reports have been completed in the nationwide series of 19 reports on the magnitude and frequency of floods, and the remaining 6 reports are scheduled for completion by December 31, 1965. Each report is for a part corresponding to a drainage-basin subdivision used by the U.S. Geological Survey. Six reports had previously been published, and one was published this year: Part 1-A, Water-Supply Paper 1671 (A. R. Green, 1–64), for basins in the region from Maine to Connecticut.

Flood-frequency synthesis for small streams in Alabama

Methods of synthesizing flood-frequency curves for small watersheds are being studied in Alabama. According to L. B. Peirce a hydrometeorological approach using available long-term rainfall records to provide the necessary time sampling offers promise for study of 5- to 15-square-mile drainage basins. A relation between storm-rainfall characteristics and peak discharge has been defined on the basis of 3 to 10 years of concurrent rainfall and streamflow records. Amount and duration of rainfall, the antecedent rainfall, and a seasonal factor are used as independent variables from which an array of annual peaks is computed for a site by graphical multiple regression.

Flood-flow characteristics of small drainage basins in Massachusetts

A report by C. E. Knox and C. G. Johnson, Jr. (*in* A. R. Green, 1–64) describes methods for estimating the magnitude of a flood of any frequency between 1.1 and 100 years for any site, gaged or ungaged, on most unregulated streams in Massachusetts. Formulas were derived through multiple-regression techniques that relate flood peaks to drainage area, slope, and an orographic factor. The formulas apply only to sites for which the drainage area is more than 10 square miles and the usable upstream storage is less than 4.5 million cubic feet per square mile.

FLOOD MAPPING

Flood maps of urban areas

Flood-inundation maps showing the limits of inundation by major floods, flood profiles, stage-frequency relations, and descriptive text have been published during the current year as Hydrologic Investigations Atlases for the following areas: Mount Vernon, Ohio (HA–40, revised); Newark, Ohio (HA–44, revised); Chillicothe, Ohio (HA–45); Zanesville,

Ohio (HA-46); Circleville, Ohio (HA-48); Hinsdale, Ill. (HA-86); Libertyville, Ill. (HA-88); Joliet, Ill. (HA-89); Harvey, Ill. (HA-90); Baton Rouge, La. (HA-126); Jackson, Miss. (HA-127); Toa Alta, Toa Baja, and Dorado, P. R. (HA-128); Geneva, Ill. (HA-142); Lombard, Ill. (HA-143); and Wadsworth, Ill. (HA-144).

Effect of a bridge on flood elevations

Crest-water surfaces were mapped by K. V. Wilson and B. L. Neely, Jr., at four bridges in central Mississippi during the flood of April 6, 1964. Preliminary analysis of the data indicates that water-surface elevations at various places in the vicinity of a bridge may vary by several feet. Elevations at the downstream side (the usual location of a gaging station) were found to be as much as a foot higher than elevations along the downstream side of the highway fills near the bridge abutments. Results of the study emphasize the importance of careful site location when gages are established on or near bridges.

Flood mapping in Delaware River basin

A flood-inundation study of a 12.6-mile reach of the Delaware River in the vicinity of Easton, Pa., and Phillipsburg, N.J., by G. M. Farlekas and J. A. Bettendorf discloses a relatively small increase in areal extent of flooding with increase in flood magnitude. Floods ranging in recurrence interval from 11 to 130 years inundated areas ranging from 0.88 to 1.29 square miles. The range in stage for this range in flood magnitude was 6 to 16 feet above flood stage. Results show that for the reach studied, which is constricted by steep valley walls, depth of flooding is more significant than areal inundation.

Flood-plain inundation study, Princeton, N.J.

Height-frequency relations for New Jersey floods, as determined by D. M. Thomas (2-64), were verified by J. A. Bettendorf for the lower 10-mile reach of Story Brook in the vicinity of Princeton, N.J. Heights for floods of 1½- and 35-year recurrence intervals were estimated, using procedures developed by Thomas; the estimated heights were compared with measured heights based on flood profiles defined by surveyed high-water marks. Estimates for 9 of the 10 sites, which are within ± 23 percent of measured values for a 1½-year flood and within ± 13 percent for a 35-year flood, are significantly better than the expected limits of accuracy defined by Thomas.

EFFECTS OF THE 1964 ALASKA EARTHQUAKE

INVESTIGATIONS OF GEOLOGIC EFFECTS

Geological Survey contributes to reconstruction effort

In the wake of the 1964 Alaska earthquake the Federal Reconstruction and Development Planning Commission appointed the Scientific and Engineering Task Force. U.S. Geological Survey geologists were assigned to the task force and its field team, which had primary responsibility for making recommendations to the Commission as to where, and where not, Federal funds should be spent for reconstruction. In the carrying out of the program, fieldwork by R. W. Lemke was instrumental in selecting a new more stable location for dock and harbor facilities at the head of Resurrection Bay at Seward; investigations by H. W. Coulter and R. R. Migliaccio figured prominently in the decision to relocate the city of Valdez from its pre-earthquake site to the Mineral Creek area; and the U.S. Geological Survey geologists were instrumental in the selection of new sites required in the rebuilding of parts of Anchorage.

Lake seiches investigated

In a study of earthquake-induced seiches, and slides from the edges of deltas, at Kenai Lake on the Kenai Peninsula, D. S. McCulloch determined that water-level oscillations occurred at the natural period of the lake basin. The direction, run-up height, and inshore limit of wave action were determined; it was firmly established that subaqueous slides as well as seiches caused destructive local waves.

Submarine slides generate destructive sea waves

Studies by George Plafker of wave distribution along the shores of Prince William Sound and the Kenai Peninsula indicate that violent waves of local origin occurred at numerous localities during the earthquake. The surface work, supplemented by submarine studies by the U.S. Geological Survey and U.S. Coast and Geodetic Survey at selected localities and by eyewitness accounts, indicates that most of the waves were generated by precipitous earthquake-triggered submarine landslides. Using a sonarprobe loaned by G. G. Shor of the Scripps Institution of Oceanography, G. A. Rusnak was able to identify deposits of large submarine slides triggered within Prince William Sound. Seiche waves and waves generated by subaerial landslides apparently did not contribute significantly to the shoreline damage.

Inferred buried fault zone on Kenai lowland

Regional mapping of earthquake effects in the

Cook Inlet area by T. N. V. Karlstrom and H. L. Foster has delineated a linear fractured zone in the surficial deposits of the Kenai lowland. It appears to reflect a major buried sheared zone, or fault, that was reactivated during the 1964 earthquake. The fractured zone trends northeast or parallel to major steeply dipping thrust faults with recent right-lateral strike-slip displacements.

Recent movement in bedrock at depth beneath the fractured zone is suggested by a concentration of micro-aftershock activity beneath the zone, and by the zone's alignment with epicenters of past earthquakes. A buried fault of probable post-Cretaceous, pre-middle Tertiary age is suggested by scattered subsurface data. At one point beneath the zone, seismic evidence indicates a major fault of unknown strike but with large vertical offset.

The inferred fault passes close to the producing oil fields on the Kenai lowland. All production lies northwest of the inferred fault; numerous dry holes have been drilled southeast of the structure.

Land-sea level changes in Cook Inlet region

Observations of coastal geology by T. N. V. Karlstrom and personnel of the U.S. Fish and Wildlife Service indicate the following changes in sea-level datum along Cook Inlet: plus 4–6 feet near Hope; plus 3–4 feet near Chickaloon Bay; no change or less than 1 foot upward between the Point Possession and Anchor Point on the Kenai Peninsula and along the west shore of Cook Inlet near Tyonek; and plus 1–2 feet along the slumped front of the Susitna River delta. The geologic estimates of sea-level change at the localities along the south shore of Turnagain Arm are of the same order of magnitude as the changes determined for bedrock bench marks on the north shore by U.S. Coast and Geodetic Survey relevelling from datum control points at Seward and Valdez. The relevelling and observational results differ by about 1 foot from postearthquake, short-term records from tide gages, reported from both Anchorage and Nikishka. (Similar discrepancies between levelling and tide-gage records in the Cook Inlet region, observed in the past, were thought to reflect the presence of a sloping sea-level datum slope in Cook Inlet resulting from the restricted estuary environment and the high tidal range.)

Line of zero change in Cook Inlet

The combined geologic and relevelling data in Cook Inlet analyzed by T. N. V. Karlstrom indicate that the eastern part of the region subsided, with the maximum subsidence closely coinciding with the arcuate axis of the Chugach-Kenai Mountain range.

In the upper Turnagain Arm area subsidence reached a maximum of about 6 feet, and became progressively less to the northwest. The line of zero change separating the area of subsidence from the area of stability or minor uplift on the continental side is still not precisely located. Present data suggest that the line of zero change lies somewhat north of the Matanuska valley within the Talkeetna Mountains and passes southwestward across the Susitna River lowland through a point near Tyonek and into Cook Inlet. Here it appears to extend roughly parallel to the Aleutian Range coastline into the Shelikof Straits area.

Tilted lake basins

Observations on pre-earthquake high-water marks relative to present water levels in the basins of Tustumena, Skilak, and Eklutna Lakes by T. N. V. Karlstrom indicate that differential regional tilting to the southeast was probably less than 2–3 feet over the length of each lake.

Ground breakage in Copper River Basin area

Field investigations by O. J. Ferrians, Jr., indicate that ground breakage was widespread in the Copper River Basin area within a radius of 100 miles of the main epicenter of the earthquake. Ground cracks formed in flood plains of rivers, in deltas, and along the toes of alluvial fans. They also occurred locally in low terraces adjacent to flood plains, in highway and other fill material, along the margins of lakes, along the face of steep slopes of river bluffs and hillsides, and in areas cleared of vegetation for several years. Their overall distribution indicates that local geologic factors primarily controlled their distribution. The ground cracks were restricted to areas underlain by unconsolidated deposits where one or more of the following conditions existed: (1) permafrost was absent or deep lying, (2) the water table was near the surface, (3) bedrock was relatively deep lying, and (4) slopes were steep. In addition, at the time of the earthquake (March) the thickness of seasonal frost differed somewhat from place to place throughout the area.

Pre- and post-earthquake elevations related to gravity anomalies

Comparison of pre- and post-earthquake gravity observations by D. F. Barnes indicate that negative gravity changes are related to positive elevation changes, and suggests that the changes are accompanied by rearrangement of masses at depths shallow enough to affect the gravity observations. The gravity data do not seem to indicate that chemical

reactions, phase changes, or elastic changes not accompanied by horizontal deformation are causes of the elevation changes.

Barnes also reports that recently completed reconnaissance gravity surveys combined with previous years' surveys now provide coverage in a north-south traverse belt across the entire State. Preliminary analyses of these data and of geomorphic observations suggest that counterisostatic elevation changes similar to those during the Alaska earthquake are also probably occurring at much slower rates throughout southern Alaska as far north as the Yukon and Tanana Rivers. Thus positive isostatic anomalies are associated with increasing elevations in the Alaska Range and Talkeetna Mountains.

Deformation of Kodiak Island

On the basis of 23 estimates of land-level change on Kodiak Island, G. W. Moore defined the deformation that occurred there at the time of the Alaska earthquake. The island was downwarped along a broad northeast-trending asymmetrical syncline with its axis passing about 10 kilometers southeast of the city of Kodiak. The city was depressed 1.6 meters below its former level, whereas Narrow Cape to the southeast was elevated 1.2 m. The gentle northwest flank of the syncline slopes 2 centimeters per kilometer toward the axis, and the southeast flank slopes 20 cm per km. The synclinal axis plunges 1 cm per km to the northeast, and a parallel anticline lies next to the syncline off the southeast shore of Kodiak Island.

Origin and extent of Bootlegger Cove Clay

Failure of the Bootlegger Cove Clay in the Anchorage area during the Alaska earthquake, and the landsliding and catastrophic damage related to this failure, as reported by Arthur Grantz, George Plafker, and Reuben Kachadoorian¹⁹ renewed the interest of geologists and engineers in this deposit. Recent study of samples from exposures at the type locality by R. A. M. Schmidt²⁰ showed the presence of marine fossils except in the uppermost part of the section. P. J. Smith has examined samples from three cores, as an aid in clarifying the problem of the environment of deposition of the formation. Marine faunas were present throughout the sections studied, which extend through most of the formation. The faunas are similar to those of the present-day

cold shallow (± 25 meters) water of Cook Inlet. The water salinity inferred for the base of the section is normal for marine water, becoming lower toward the top.

F. W. Trainer and R. M. Waller (p. D167-D174), in a study of the glacial drift at Anchorage based chiefly on subsurface data, found that the Bootlegger Cove Clay is the youngest of several clays deposited in the area during Pleistocene time. Locally it is more than 200 feet thick, and in much of the area it is more than 100 feet thick. Its areal extent and the configuration of its base and upper surface show that it was deposited in, and retained the form of, a basin which had persisted from earlier in the Pleistocene.

Southern Alaska Continental Shelf tectonics

D. F. Barnes accompanied the USCGS ship *Surveyor* on a marine geophysical reconnaissance of the southern Alaska Continental Shelf following the 1964 earthquake. On this cruise the Patton Bay fault found by the U.S. Geological Survey on Montague Island was traced approximately 30 miles offshore, and similar evidence of old or recent faulting was found approximately 100 miles farther southwest along the approximate strike of the fault. Gravity data obtained on the same cruise show that a large free-air and isostatic gravity high covers the portion of the Continental Shelf where new emergent shorelines were found after the earthquake. Inasmuch as the earthquake did not decrease the gravity anomaly, the anomaly is believed to be the result of counterisostatic processes.

Crust deformed by 1964 Alaska earthquake

Studies of the tectonic deformation associated with the Alaska earthquake by George Plafker and other U.S. Geological Survey geologists indicate that: (1) vertical movement occurred over an area of at least 50,000 square miles, with a maximum observed uplift of 33 feet and a maximum subsidence of $7\frac{1}{2}$ feet; (2) the deformation consists of a coastal zone of uplift that includes much, if not all, of the Continental Shelf and is bordered on the inland side by a zone of subsidence; and (3) vertical displacements result largely from crustal warping with superimposed local surface faulting on Montague Island and the adjacent Continental Shelf. The main fault along which the earthquake occurred is not exposed at the surface. It is inferred to be related to the Aleutian Arc, and to dip at a low angle beneath the continental margin from the Aleutian Trench.

¹⁹ Arthur Grantz, George Plafker, and Reuben Kachadoorian, 1964, Alaska's Good Friday earthquake, March 27, 1964: U.S. Geol. Survey Circ. 491, 35 p.

²⁰ R. A. M. Schmidt, 1963, Pleistocene marine microfauna in the Bootlegger Cove Clay, Anchorage, Alaska: Science, v. 141, p. 350-351.

Uplift of Continental Shelf initiates ocean waves

George Plafker suggests that the train of long-period seismic sea waves that followed the earthquake was impulsively generated by rapid uplift of the Continental Shelf. Distribution of wave damage, and calculated travel distance of the initial crests, based upon reported arrival times along the coast of south-central Alaska, indicate that the major zone of wave generation is along a line at least 180 kilometers long that extends southwestward from Montague Island. The northern end of this zone coincides with the area of maximum known uplift off the southwest tip of Montague Island.

INVESTIGATIONS OF HYDROLOGIC EFFECTS

Effects in Alaska

The earthquake produced unusual and significant hydrologic effects in Alaska. According to R. M. Waller, preliminary analysis of water-level changes in artesian-aquifer systems composed of glacio-fluvial deposits indicates an apparently permanent lowering of artesian-pressure level in parts of some basins. The lowered levels appear to represent either increases in the transmissibility of the aquifers or increases in the rate of discharge from them. A. J. Feulner, using cyclic tidal stage-ratio and time-lag methods, has tentatively determined an increase in transmissibility at two wells at Anchorage. L. L. Dearborn and M. V. Marcher collected data on 100 wells in the Anchorage area and found that about 70 percent were affected by the earthquake: in 25 percent the water level had declined noticeably, while in about 40 percent the water had been muddied for periods ranging from a few hours to a month. A few wells were extensively damaged by breaking of the casing or screen or by the surging of mud up through open-end casing. R. M. Waller believes that shallow water-table aquifers, capped by seasonal frost, played an important role in extensive fracturing of surficial deposits and extrusion of sand and mud. The lower slopes of alluvial fans, the outer edges of deltas, the channels of braided streams, and swampy areas were the principal sites of this fracturing and extrusion. The ice covers of lakes and streams were broken or cracked to distances of at least 400 miles from the earthquake epicenter.

In many streams, flow was greatly reduced temporarily, according to M. J. Slaughter, as a result of (1) the loss of water into fractures, damming by landslides or snowslides; and (2) the sloshing of water from lakes. In other streams, flow increased, probably because of the release of water from ice-

choked reaches. Preliminary appraisals suggest no changes in long-term trends of stream discharge, however, except perhaps in areas of maximum uplift or subsidence. Several stream gages on Kodiak Island and in southeastern Alaska recorded seismic sea waves that travelled far upstream; and those gages on Kodiak Island now regularly record ocean tides, as a result of the subsidence of Kodiak Island. L. S. Leveen reports that the sediment load in the Matanuska River, which began its spring runoff immediately after the earthquake, increased to about five times that recorded at comparable rates of water discharge in previous years. The increase lasted through April; sediment discharge then returned to normal as most of the slide debris along the banks and the sand and silt extruded upon channel bars were removed.

Effects outside Alaska

The earthquake also caused important hydrologic changes outside Alaska. R. C. Vorhis has tabulated seismic water-level fluctuations recorded in 661 wells in 44 of the 50 States, and in 4 wells in Puerto Rico and 1 in the Virgin Islands. (Water-level fluctuations caused by this earthquake were also recorded in wells in Canada, Great Britain, Belgium, and Libya.) The largest recorded fluctuation, 23 feet, occurred in a well at Belle Fourche, S.Dak. The earthquake was recorded in 70 wells in Florida; in some wells, especially near Tampa, the movement was so violent that float cables were thrown from the pulleys of the recorders. Perhaps the most unusual hydrologic effect of this earthquake has been an apparently permanent change in the water-level at some localities. In the south Georgia basin, for example, water levels in some wells rose during a 3-week period after the quake, and the maximum rise recorded was about 3.3 feet. On the other hand, a well that penetrates a recharge area in Clay County, Fla., showed a reverse effect—the water level began to decline at the time of the quake and dropped 4 feet over a 3-week period. Similar local changes in water level following the earthquake occurred in Illinois, Michigan, Nevada, Pennsylvania, and Utah. These changes in water levels, which appear to be permanent, may reflect strains developed in the earth's crust as a result of the earthquake. A. M. Piper²¹ and others have suggested that earthquakes might cause strains that produced hydrologic changes.

²¹ A. M. Piper, 1933, Fluctuations of water-surface in observation-wells and at stream-gaging stations in the Mokelumne area, California, during the earthquake of December 20, 1932: *Am. Geophys. Union Trans.*, 14th Ann Mtg, p. 471-475.

The Alaska earthquake generated seismic seiches in many parts of the conterminous United States. Surface-water records collected by the Geological Survey and tabulated by R. C. Vorhis show that in Texas 69 gages recorded seiches; the maximum fluctuation had a double amplitude of 0.68 feet. In Georgia the maximum fluctuation recorded by 24 gages was 0.22 feet. Other States in which seiches

were recorded include Alabama, Arkansas, Illinois, Indiana, Kentucky, Louisiana, Missouri, Ohio, Pennsylvania and Tennessee. Most of the seiches recorded were in the south-central and southeastern States. Many swimming pools in the Gulf Coast States sloshed water at the time of the quake, but no reports of seiches have been received from the Atlantic Coast States.

REGIONAL GEOLOGY

Much of the geologic and geophysical work of the U.S. Geological Survey consists of the mapping of specific areas, mostly for publication as quadrangle maps at scales of 1:62,500 and 1:24,000. Some of these studies are for the purpose of extending the detailed geologic knowledge in areas of known economic interest; some are to gain detailed knowledge at localities or areas for engineering planning or construction. Still other mapping studies are carried on with paleontology, sedimentary petrology, or some other specialized topic as the primary objective.

The systematic description and mapping of rock units to show local and regional relations likewise constitute a major scientific objective. Mapping the geology of the United States is a mandate of the Organic Act establishing the Geological Survey, and the completion of geologic maps for the country at scales that will fulfill foreseeable needs and uses is a long-range goal. A summary of recent results of this mapping, especially in the fields of stratigraphy, structural geology, and regional geophysics, is discussed here according to subdivisions of the conterminous United States shown on figure 3.

FIGURE 3.—Index map of the conterminous United States, showing boundaries of regions referred to in discussion of regional geology.

INTERMEDIATE-SCALE GEOLOGIC MAPS

Geologic mapping at a scale of 1:250,000 on 1° by 2° topographic base maps comprises a new formal element of the U.S. Geological Survey's program. Geologic mapping at this scale has formed a small but significant part of the Survey's geologic studies for many years—notably in long-range programs to complete geologic coverage in Alaska, Oregon, Nevada, and the Colorado Plateau. The value of intermediate-scale geologic maps is also recognized by State Geologic Surveys, many of which have 1:250,000-scale geologic-map programs underway or completed. This joint attack by the Federal and State Surveys on a nationwide problem promises to provide complete intermediate-scale geologic map coverage of the United States at a rapid pace.

Coordinated geologic-mapping programs at this scale are well underway in the States of Alaska, Oregon, Nevada, Colorado, and Nebraska, and single-sheet 1° by 2° geologic maps have been started for Montana, Idaho, Wyoming, Washington, and the southern Appalachian region.

Although not a substitute for the standard inch-per-mile geologic map program, the 1:250,000-scale studies will indicate areas where the need for larger scale maps is most critical, and will direct attention to larger segments of the earth's crust. The ready availability of uniform coverage at intermediate scale promises more realistic priority assignments for new starts in inch-to-the-mile geologic mapping. The 1:250,000-scale maps have already proved to be ideal for geologic analysis of broad tectonic and stratigraphic problems, for analysis of mineral provinces, and for relating broad geophysical anomalies to surface geology.

MAPS OF LARGE REGIONS

Cooperative projects

Maps of national or international scope are prepared as a regular and important part of the Geological Survey program. Such maps utilize studies by Geological Survey personnel, published information and unpublished information supplied by State Geological Surveys, private companies, and universities. Some of the maps are prepared and published by the Geological Survey in collaboration with national and international scientific organizations.

A Bouguer gravity-anomaly map of the United States, exclusive of Alaska and Hawaii, scale 1:2,500,000, was published in November 1964. This map was compiled by the Special Committee for

Geophysical and Geological Study of the Continents, American Geophysical Union, G. P. Woollard, University of Hawaii, chairman, and was published by the Geological Survey, H. R. Joesting, coordinator. It shows relative gravity values, which reflect variations in thickness and density of rocks in the earth's crust; the contour interval is 10 milligals. The map is available in two editions: one is on translucent parchment paper for overlaying on the "Tectonic Map of the United States" or on other maps using the standard 1:2,500,000 base, and the other is on standard map paper.

Other cooperative maps currently in preparation that are to be published by the Geological Survey include:

1. Geologic map of North America, scale 1:5,000,000, multicolor in 2 sheets, each measuring 40 x 58 inches. This map was compiled by a committee of the Geological Society of America, E. N. Goddard, University of Michigan, chairman. It is planned for publication by the Geological Survey in late 1965, on a two-color base (black for culture and geography and blue for water). The base prepared by the Survey also shows depth of water in offshore areas by contours drawn at 200 meters, 500 meters, and at 500-meter intervals thereafter. More than 120 different geologic units are shown by overprinting screens and patterns of 6 colored inks.
2. Tectonic map of North America, scale 1:5,000,000. This map is being compiled for the Subcommittee for the Tectonic Map of the World, International Geological Congress, under the guidance of P. B. King, U.S. Geological Survey. Compilation of the southern half of the map is completed; the northern half is still in compilation stage. The work ultimately will contribute data about North America for the "Tectonic Map of the World," at a scale of 1:15,000,000, a cooperative international undertaking sponsored by the International Geological Congress.
3. Basement map of North America between latitudes 20° and 60° N., scale 1:5,000,000. Compilation of this map is by the Basement Rock Project Committee of the American Association of Petroleum Geologists, P. T. Flawn, University of Texas, chairman, using part of the new base map of North America. The map will show the altitude of the upper surface of the basement as determined from wells, geophysical measurements, and geologic inference. The basic well data compiled by the committee in preparing the basement map of North America will be pub-

lished by the American Association of Petroleum Geologists.

4. Basement rock map of the United States, scale 1:2,500,000. This map is sponsored by the Advanced Research Project Agency, Department of Defense, under the direction of W. R. Muehlberger, University of Texas. Compilation of the basement rock map is virtually completed. Colors and patterns will show the age and lithologic types of both the exposed and buried basement rocks. Some internal structures of the rocks, where known, will also be shown by symbols.

Paleotectonic maps

A long-term program to compile and publish paleotectonic maps of the conterminous United States for each of the geologic systems is continuing. The magnitude and sequence of tectonic events and the history of sedimentation are reconstructed on these maps by subdividing the system under investigation into intervals that can be broadly correlated for the country as a whole. Objective maps at a scale of 1:5,000,000 show the present extent, thickness, and lithofacies of each stratigraphic interval. Interpretive maps, most at a scale of 1:10,000,000, reconstruct the former extent of the interval, show the principal tectonic elements influencing deposition, identify some of the environments of deposition, and show other features of importance to regional geology. Folios of the Jurassic and Triassic Systems have been published; the folio for the Permian System has been compiled and is being readied for publication. Maps for the Pennsylvanian and Mississippian Systems are in preparation.

A map showing the total thickness of Pennsylvanian rocks has been compiled for the part of the conterminous United States west of long 95° W. Pennsylvanian rocks cover most of this area except for Washington, Oregon, California, and eastern Nevada, where Pennsylvanian rocks are present locally but for the most part have either been destroyed by igneous activity or are deeply buried by younger rocks. The map shows that Pennsylvanian rocks are thickest in a series of disconnected en echelon troughs lying in a northwest-trending band from southeastern Oklahoma to southern Idaho.

Catalog of aerial photographs

A catalog of aerial photographs in stereographic pairs illustrating geologic features in the 50 States and Puerto Rico is being prepared under the direction of C. S. Denny. Currently, entries have been made for about 300 sets of photographs; most entries include reference to a geologic report and to

a topographic and geologic map of the area shown in the photographs. A list of the photographs, where they can be obtained, and other pertinent information is in preparation.

National Atlas

The Geological Survey is one of several government agencies and private groups that are providing maps and text for a National Atlas. The National Atlas is being prepared under the direction of A. C. Gerlach, staff geographer, on loan from the Library of Congress and will be published under the auspices of the Department of the Interior. Maps will be at a scale of 1:7,500,000, or smaller. Maps of a geologic nature that are the responsibility of the Geological Survey and the geologists in charge of their preparation are as follows: (1) seismology and gravimetry tectonic features of the conterminous United States, and tectonic features of Alaska, P. B. King; (2) geologic history, L. C. Craig; (3) geologic map, D. M. Kinney; (4) glacial geology, C. S. Denny; and (5) karst areas and caves, W. E. Davies.

For further information see the writeup on "National Atlas" in the section, "Topographic Surveys and Mapping."

COASTAL PLAINS

ATLANTIC COASTAL PLAIN

Form of pre-Cretaceous basement in New Jersey

H. E. Gill reports that deep test drilling in Ocean and Burlington Counties, in the coastal plain of central New Jersey, has more precisely defined the pre-Cretaceous basement. In detail this surface is more irregular than was heretofore thought. Test wells penetrated the basement several hundred feet lower at Island Beach State Park in Ocean County, and higher to the west at Butlers Place, Burlington County, than was predicted from previously published data. The new data indicate the presence of a structural high, trending northwest-southeast, which had a strong influence on the lithologic character and distribution of deposits of the adjacent transitional sequence of Cretaceous marine deposits.

Miocene folding and faulting in Georgia

Structure contours of the top of the Suwanee Limestone of Oligocene age, drawn by C. W. Sever, show that this aquifer is folded into a northeast-plunging syncline near Meigs, in southwestern Georgia, and into a southwest-plunging anticline near Brunswick, at the Atlantic coast. A northeast-

trending high-angle fault named the Ochlockonee extends between these folds. Displacement across the fault is as much as 190 feet, with the southeast side upthrown. Thickening and displacement of the formations across the Ochlockonee fault show that the folding and faulting occurred periodically during early and middle Miocene time.

Surface expression of coastal-plain structures in New Jersey

Detailed mapping by J. P. Minard and J. P. Owens has outlined two well-defined domal structures in the coastal-plain sediments east of Trenton, N. J. The sediments known to be involved in the folding range in age from Late Cretaceous (Companian) through middle Miocene. The structures are elliptical and each measures approximately 1.5 by 4 miles. The long axes of the ellipses are aligned northeast-southwest, parallel to the regional strike of the formations. The presence of these folds and of other smaller structures throughout this area suggests widespread minor folding of the coastal-plain formations.

Ordovician shale found in basement rock in Florida

Recent examination by Jean Berdan of cuttings from a well in Suwannee County, in northern Florida, confirms the existence of a belt of Middle Ordovician black shale trending southeast-northwest beneath the middle of the county at depths between 3,000 and 4,000 feet. The Middle Ordovician rock is overlain to the northeast by black shales of Silurian age, and it is underlain to the southwest by quartzitic sandstones of Early Ordovician age.

Stratigraphic studies in South Carolina

A water well drilled near Orangeburg, in south-central South Carolina, penetrated 930 feet of coastal-plain sediments. Projection of known stratigraphic relationships into this region shows that the lower 500 feet of sediments should be Late Cretaceous in age, and approximately 400 feet should be assignable to the Tuscaloosa Formation (Woodbine age). G. E. Siple and W. D. Paradeses report, however, that the sediments are lithologically similar to the younger Black Creek or Ellenton Formations (Austin to Navarro in age). A pronounced thinning of the Tuscaloosa section in this region is thus indicated. A possible explanation is that the axis of deposition for the Upper Cretaceous sediments in this particular basin is aligned more nearly in a northeast-southwest direction than northwest-southeast, as had heretofore been postulated, and that the

sediments near Orangeburg represent a change to a thinner facies than the Tuscaloosa.

As part of a study of the distribution of permeable sands in central South Carolina, Siple and Paradeses have mapped hitherto unreported bodies of indurated coarse quartz sandstone with tan-colored claystone inclusions up to 20 centimeters in diameter, underlain by beds of claystone (prodelta clay?), in the Gilbert (15') and Batesburg S.E. (7½') quadrangles. Although the sand bodies are probably of Eocene age, they form residual hills standing about 25–30 feet above the surrounding plain which is formed of lower to middle Tertiary rocks. The sandstones are coarsely crossbedded and weathered to dark gray. Smooth-walled ovate cavities occur along the more or less horizontal bedding planes along the topsets or bottomsets, and their major axes are parallel to this bedding; presumably they were originally filled by clay balls.

Cretaceous microfaunas and heavy minerals along Chattahoochee River, Georgia and Alabama

During paleontologic and petrologic studies of the Blufftown Formation, Cusseta Sand, Ripley Formation, and Providence Sand in the section along the now-inundated Chattahoochee River in Georgia and Alabama, Harlan Bergquist and J. E. Johnston found an excellent microfaunal assemblage of about 60 species of Foraminifera and 7 or 8 species of ostracodes. A few planktonic species of Foraminifera occur in all the formations; arenaceous species appear to be most common in the Blufftown and Ripley Formations. Preliminary determinations indicate that the faunas are largely assemblages of Austin to Taylor age but represent Late Cretaceous stages from Santonian to Maestrichtian. Of particular note is a relatively high percentage of phosphate in the fine size of the heavy-mineral fractions of the Cusseta Sand.

Subsurface stratigraphic studies of Georgia coastal plain

Recent subsurface stratigraphic studies by S. M. Herrick show that Pliocene(?) sediments definitely underlie Glynn, Charlton, and Camden Counties, in southeastern Georgia, as indicated by the presence of certain varieties of the genus *Ammonia*. Between Liberty and Glynn Counties the lithologic section of upper Cretaceous (mid-Taylor age) to lower Tertiary (base of the middle Eocene) rocks changes from unconsolidated clastic rocks to a limestone facies. The sequence of carbonate rocks strongly resembles the limestones of similar age in peninsular Florida. Northeast of Liberty County, lower Eocene rocks are absent from sections penetrated in Geo-

logical Survey test holes in Chatham County, Ga., and on Parris Island, S.C. This absence may be due to up-faulting of these rocks and their subsequent removal by subaerial erosion.

Pollen in Pleistocene terrace deposits at Washington, D.C.

Palynological study of the lower terrace deposits at Washington, D.C., by A. S. Knox has revealed that several changes in climate and sea level occurred in the area during Pleistocene time. The evidence suggests that two interglaciations, the Sangamon and the Yarmouth, and two glaciations, the Wisconsin and the Illinoian, are represented. The interglacial deposits, with abundant pollen of broad-leaved deciduous trees, indicate a climate as warm as or warmer than that of the present; the glacial-age deposits contain abundant spruce pollen and indicate temperatures much lower than those of the present.

Buried Pleistocene(?) valley on Long Island, N.Y.

Julian Soren reports that deep drilling about a mile west of Lake Ronkonkoma, in the mid-island region of Suffolk County, Long Island, N.Y., has disclosed the presence of a deep buried valley cut in Cretaceous deposits. The valley trends about N. 45° W. from Long Island Sound, and is more than 300 feet below present sea level.

Significant finds of vertebrate fossils in Virginia and Florida

Two mammalian fossils recently given to the Survey for identification have raised questions concerning the correspondence of age classifications of the coastal-plain sediments based on vertebrate and on invertebrate fossils, respectively. A single lower horse molar, collected from the Yorktown Formation (upper Miocene) at Cobham Wharf, Va., has been identified by F. C. Whitmore, Jr., of the U.S. Geological Survey and M. F. Skinner, of the Frick Laboratory, American Museum of Natural History, as *Hipparion* cf. *H. eurystyle* Cope, of Clarendonian or early Hemphillian (early or early middle Pliocene) age. The possibility that part of the Yorktown may be of Pliocene age is further supported by the finding in 1960, at Hampton, Va., of a baleen whale of a type that seems too advanced for the Miocene.

Part of a horse skull, found in the Caloosahatchee Formation in its type area near Labelle, Fla., has been identified by Whitmore as *Equus*, probably a mid-Pleistocene species. This find is of particular interest because the rich invertebrate fauna of the Caloosahatchee is generally regarded as being of Pliocene age.

GULF COASTAL PLAIN AND MISSISSIPPI EMBAYMENT

Paleontologic data suggest northward Paleocene transgression across Mississippi embayment

Studies of Foraminifera by S. M. Herrick have shown that the Clayton Formation in Hardeman County, Tenn., although containing several "key" species that are reportedly limited elsewhere in the region to the Clayton Formation or its stratigraphic equivalents, contains 32 species that are reported by Cushman²² from the Matthews Landing Marl Member of the Porters Creek Formation that overlies the Clayton Formation in Alabama. These paleontologic data suggest to Herrick that the sea in which the Clayton was deposited was transgressive northward and that the Clayton Formation is of late Paleocene age in Tennessee.

Recently incised meanders and older terraces along Alabama River

The Alabama River between Montgomery and Selma, Ala., is reported by L. C. Conant to be bordered on one side or the other by 10- to 20-foot banks cut into the Eutaw Formation and Selma Group of Late Cretaceous age. The cut banks indicate that most or all of the present meanders are recently incised and are not subject to lateral shifting. Older downcutting is shown by the presence of broad thick terrace deposits of gravel, sand, and clay which were noted by Conant a few miles north of the flood plain, along this segment of the river, as much as 200 and 300 feet above the flood plain. Terraces of similar composition were earlier recognized by Monroe (p. 43-44²³) at comparable levels on the south side of the river but at considerably greater distances from the flood plain.

Jurassic stratigraphy, tri-state area of Arkansas, Louisiana, and Texas

Isopach mapping by K. A. Dickinson has revealed that the Buckner Member of the Haynesville Formation in northeast Texas, southwest Arkansas, and northwest Louisiana is composed of two genetically different units. The areas of deposition of both units were limited on the south by barriers consisting largely of two series of salt-cored gravity-flow anticlines. The barrier for the lower unit extends from the Linden oil field, in south-central Cass County, Tex., through the McKamie oil field in cen-

²² J. A. Cushman, 1951, Paleocene Foraminifera of the Gulf Coastal region of the United States and adjacent areas: U.S. Geol. Survey Prof. Paper 232, 75 p. [1952]

²³ W. H. Monroe, 1941, Notes on deposits of Selma and Ripley age in Alabama: Geol. Survey of Alabama Bull. 48, 150 p.

tral Lafayette County, Ark., and thence eastward through Columbia and Union Counties, Ark. The upper unit, which represents a less saline depositional environment than that of the lower unit, overlapped the barrier of the lower unit and extended southward to a barrier that trends from the Rodessa oil and gas field in northern Marion County, Tex., and Caddo Parish, La., eastward along the Louisiana State Line to the Haynesville oil field in northern Claiborne Parish, La.

NEW ENGLAND AND EASTERN NEW YORK

Glacial chronology of New England

The chronology of the glaciation of New England has been summarized in a review of published and unpublished data by J. P. Schafer and J. H. Hartshorn. The last glaciation (Wisconsin) was preceded by (1) a widespread glaciation that appears to be of post-Sangamon and pre-classical Wisconsin age, (2) by the deposition of the Gardiners Clay of Sangamon age, and (3) by the formation of still older drifts of uncertain correlation. The last ice sheet reached its maximum advance along the coast of southern New England, probably in early Woodfordian (Tazewell) time. Readvances in southern New England, northern Vermont, and southwestern Maine are not well dated, but may be approximately of Cary, Port Huron, and perhaps Valdres age. The last ice masses in New England were a possible local icecap on the White Mountains and in western Maine, and cirque glaciers on Mount Katahdin.

EASTERN NEW YORK

Five major units in northwest Adirondacks form refolded anticlinorium

Stratigraphic reconstruction by A. E. J. and C. G. Engel shows five major units in the northwest Adirondack Mountains. Three of the units are carbonate formations, one is a graywacke sandstone, and the other is a thick basal quartzite. These formations apparently form a refolded anticlinorium metamorphosed at progressively higher grades in the direction of the central Adirondack massif.

Migration of elements during metamorphism, northwest Adirondacks

The progressive metamorphism of marble, graywacke, pelite, quartzite, and basalt in the northwest Adirondacks involves progressive dehydration, decarbonation, and loss of alkalis, silica, and cogenetic trace elements. At the pressure and temperature

conditions of the upper amphibolite facies, anatectic magma is formed by partial melting of clastic sediments. The mineralogy of the metamorphic rocks has been explored in detail by A. E. J. and C. G. Engel. The volume of Mn, Fe, Ti, Pb, and Zn mobilized from the sedimentary column is in excess of that concentrated in the known economic concentrations of these elements in the northwest Adirondacks. Most mineral deposits in these rocks appear to be derived in large part by the metamorphic mobilization of the elements from the enveloping and subjacent country rock.

RHODE ISLAND

Geology of Chepachet quadrangle studied

Work by A. W. Quinn in the Chepachet quadrangle, northwestern Rhode Island, indicates that a series of syntectonic intrusive rocks invaded the Blackstone Series, and probably other older rocks. The intrusives range from diorite and amphibolite in the central part of the quadrangle to quartz diorite gneiss, granodiorite gneiss, and granite gneiss in the western half of the quadrangle. The gneisses appear, at least in part, to be equivalent to the Northbridge Granite Gneiss of Massachusetts.

MASSACHUSETTS

Stratigraphy and igneous geology in northeastern Massachusetts

R. O. Castle (p. C81-C86) has divided gneissic rocks in the South Groveland quadrangle, northeastern Massachusetts, into the Boxford Formation and the Fish Brook Gneiss. The Boxford is thought to be correlative with the Rye Formation of southeastern New Hampshire, and perhaps with the Marlboro Formation of eastern Massachusetts. The Fish Brook is either an isolated intrusive rock predating the subalkaline intrusive series in this area, or an ancient "dome" gneiss similar to those of western New England. The subalkaline intrusive series of C. H. Clapp²⁴ is revised into four units (p. C74-C80): (1) Sharpners Pond Tonalite, (2) Newburyport Quartz Diorite, (3) a "younger subalkaline series" defined by Priestley Toulmin 3d²⁵, and (4) Andover Granite. The Andover Granite is included with the subalkaline series rather than the alkaline series (C. H. Clapp²⁴) because of the continuity between the Andover and the pre-"alkalic" Sharpners Pond Tonalite.

²⁴ C. H. Clapp, 1921, *Geology of the igneous rocks of Essex County, Massachusetts*: U.S. Geol. Survey Bull. 704, 132 p.

²⁵ Priestley Toulmin 3d, 1964, *Bedrock geology of the Salem quadrangle and vicinity, Massachusetts*: U.S. Geol. Survey Bull. 1163-A, p. A1-A79.

Parting in gabbro at Nahant induced by shear during folding

Medium-grained gabbro at Nahant, Mass., examined by C. A. Kaye (p. C12–C19) is broken by closely spaced parallel partings (pseudobedding-plane partings) and the rocks have been deformed into a series of anticlines and synclines. The spacing of the partings varies with the tightness of the folding, and in the crests of small folds there are as many as eight partings per inch. The partings are thought to have been produced by shear failures on the concave side of flexed layers of gabbro. Initially, these layers were formed by well-developed sheeting-type jointing. Shear-induced tension may also have produced the sheeting joints and probably was caused by movement along a still-uncovered sizable reverse fault.

Reverse polarization noted in dikes of Boston Basin

Many mafic dikes in the Boston Basin have been found by C. A. Kaye to have a very high remanent polarization that is reverse to the present magnetic poles. The effect of local "hot spots" produces a complete reversal of a Brunton compass needle, even when held as far as 8 feet from outcrops.

Shelburne Falls dome exhibits "anomalous" gravity high

Continued analyses of the geophysical data of the Shelburne Falls gneissic dome and other similar structures in the western part of Massachusetts by R. W. Bromery indicate that the amphibolites associated with the Shelburne Falls dome produce an anomalous geophysical expression that contrasts with the patterns about other domes. For example, the Plainfield dome is characterized by a small gravity low and a pronounced magnetic low, whereas the Shelburne Falls dome has an associated pronounced gravity low and a pronounced magnetic high.

Folded early Paleozoic depositional basin in northwestern Massachusetts

A depositional basin involving rocks between the Cambrian Hoosac Schist and the Ordovician Moretown Formation has been found in part of the North Adams and Rowe quadrangles by N. L. Hatch. The basin is roughly $2\frac{1}{2}$ miles wide and 3,000 feet deep, and the rocks here are from 3 to 5 times thicker than in the surrounding area. The basin appears to have been the site of relatively intense folding, and difficulties in tracing units across this area are probably due to a combination of two or more stages of deformation as well as to complex sedimentary facies relations.

Fossil indicates Early Cambrian age for Weymouth Formation

A. R. Palmer has identified *Scenella*, a form not previously reported from the locality in a collection of fossils obtained by K. G. Bell from the Weymouth Formation in Nahant, northeastern Massachusetts. The fossils indicate an Early Cambrian age for the Weymouth Formation.

Fault separates Triassic rocks from metamorphic units in western Massachusetts

A dome outlined by three superadjacent distinctive metamorphic units has been mapped in the western part of the Southwick quadrangle, western Massachusetts, by R. W. Schnabel. The Triassic rocks covering the eastern two-thirds of the quadrangle are bounded on the west by a high-angle fault of at least 200-feet displacement.

Silurian fossils date Newbury Formation in northeastern Massachusetts

Two new fossil localities have been found by N. P. Cuppels in the Newbury Formation in the Georgetown quadrangle, northeastern Massachusetts. The fossils, which are whole and undeformed, include *Salopina* sp., *Protochonetes* sp., *Nuculites* sp., *Hermanino* sp., and indeterminate rhynchonellid and pteriod genera. They are identified as Late Silurian fauna, and the Newbury is thus probably equivalent to the Pembroke Formation of eastern Maine.

Artifacts and ventifacts in Cape Cod dunes

Wind-cut Indian artifacts associated with exhumed soil horizons in the Provincetown sand dunes were found by J. H. Hartshorn during a study of the Provincetown quadrangle. In the same area, ventifacts have been formed in a few tens of years on beaches that have been shut off artificially from the sea.

Deglaciation in Berkshires resulted in ice-dammed lakes

Continued investigation of glacial-lake features in the Berkshire Hills of western Massachusetts by G. W. Holmes has confirmed parts of F. B. Taylor's unpublished hypotheses of a complex of temporary ice-dammed lakes during deglaciation. A partial and tentative chronology is now suggested: (1) a small lake in the East Canaan, Conn., area, possibly connecting at a later stage with (2) a shallow lake in the Sheffield Falls Village lowland in the Housatonic valley; (3) glacial Lake Housatonic of F. B. Taylor, a large lake with several stages, dammed at the narrows of the Housatonic River near Glendale, and extending into Stockbridge Bowl and to the vicinity

of Tyringham; (4) a small lake or arm of a lake in the Dalton area, possibly connected at one time with Lake Bascom in the North Adams and Williamstown areas. Small glacial lakes probably also existed in the South Egremont, Hancock, Shaker Village, and Lake Onata areas.

Origin of stratified drift in northern Cape Cod

Mapping of the North Truro and Wellfleet quadrangles by Carl Koteff, R. N. Oldale, and J. H. Harts-horn suggests that all the stratified drift in that area originated from the South Channel ice lobe to the east. The deposits are thought to have been laid down in a glacial lake dammed by the Sandwich moraine to the south and ice to the east and north. The Highland clay is part of this lacustrine late-glacial sequence, and is not correlative with the Gardiners Clay on Long Island as suggested by earlier workers. Marine shell fragments in the drift, with radiocarbon ages of 20,000 years or more, are presumed to have been picked up during the advance of the ice over marine deposits.

CONNECTICUT

Aeromagnetic data indicate major structural break in Connecticut

Interpretation by R. W. Bromery of aeromagnetic data in eastern Connecticut shows a pronounced linear anomaly belt striking east through the northern parts of the Uncasville, Old Mystic, and Ashaway quadrangles. This magnetic feature is interpreted as indicating a major fault or geologic discontinuity because the magnetic anomaly patterns north and south of the linear anomaly belt are markedly different. The magnetic data show that this feature continues west and east of the boundaries of the Uncasville and Ashaway quadrangles, respectively.

Radiometric ages provide time scale for eastern Connecticut

Synthesis of radiometric and field data by R. E. Zartman, G. L. Snyder, T. W. Stern, R. F. Marvin, and R. C. Buckman (p. D1-D10) for eastern Connecticut indicate that: (1) the Tatnic Hill Formation and underlying rocks are probably Middle(?) Ordovician or older; (2) the Scotland Schist and Hebron Formation are Lower Devonian or older; (3) the Canterbury Gneiss, a related quartz monzonite dike, and the gabbro of Lebanon were intruded during the Acadian orogeny in Early Devonian time; and (4) older pegmatites were intruded over a time span ranging from possibly Middle Ordovician to Middle Devonian. The area was regionally dynamo-thermally metamorphosed in the Acadian (Devon-

ian) and thermally remetamorphosed in the Alleghenian (Permian) orogenies. Younger pegmatites were intruded in Permian(?) time.

A new member near the middle of the Brimfield Schist, believed to be equivalent to the Fly Pond Member of the Tatnic Hill Formation, has been mapped by G. L. Snyder in the Marlborough and Columbia quadrangles.

MAINE

Fossils indicate margin onlap during Llandovery to Ludlow time

Basal Silurian calcareous units that unconformably overlie Cambrian and Ordovician metamorphic rocks mapped by E. L. Boudette and D. S. Harwood in northwestern Maine contain diagnostic shelly faunas that have been studied in detail by A. J. Boucot. The spatial relationship of an assemblage containing *Strictlandia lens ultima* found on the south side of East Kennebago Mountain in the Kennebago Lake quadrangle to a younger assemblage containing *Eccentricosta* found near Parmachenee Lake in the Cupsuptic quadrangle indicates a north-westward margin onlap over Cambrian and Ordovician rocks in the region during a period spanning late Llandovery to Ludlow time. A distinctive, very coarse polymictic conglomerate, lithologically comparable to the Rangeley Conglomerate of Smith (1923)²⁶, is associated with the upper Llandovery rocks at Blanchard Ponds, Kennebago Lake quadrangle. A minimum late Llandovery age (assuming onlap) may thus be established for the hitherto undated Rangeley Conglomerate, which underlies a narrow belt that strikes east-northeast about 8 miles southeast of the dated locality.

Large faults may predate consolidation of Silurian strata

Faults of large displacement that probably predate metamorphism and possibly complete lithification of the strata they cut are major structures found in the Rangeley and Phillips quadrangles, west-central Maine; by R. H. Moench. One fault separates Ordovician(?) strata on the northwest side from Silurian strata; 5,000 to 10,000 feet of Silurian strata is missing. In one outcrop, the fault zone is about 6 inches thick; it has the appearance of a recrystallized, smeared mixture of sandy and pelitic material. Adjacent metasedimentary rocks show no sign of rupture; a small metasandstone dike extends from the adjacent bed to the southeast into the fault zone. These features suggest that the sands

²⁶ E. S. C. Smith, 1923, The Rangeley conglomerate: Am. Jour. Sci., 5th ser., v. 5, p. 147-154.

were unconsolidated after faulting. Two other faults of probably similar character are recognized in younger parts of the stratigraphic sections. They are defined by the absence of probably several thousands of feet of strata on their southeast sides; outcrops have not been found. It is possible that the faults are surfaces of gravity sliding, which took place when regions to the north were uplifted in Silurian and possibly Early Devonian time.

Silicic intrusions found to border Moxie mafic pluton on the east

Continuing study in the vicinity of the Moxie mafic pluton in west-central Maine by G. H. Espenshade shows that an irregular curving valley and lowland that trends eastward from Moosehead Lake for about 22 miles is underlain by gabbroic rocks of the Moxie pluton in the western 10 miles, by quartz diorite in the center, and by quartz monzonite at the eastern end. The quartz monzonite seems to be an offshoot of the Katahdin granite pluton to the north, and is probably the youngest of the three groups of rocks on the basis of K-Ar radiometric ages by Henry Faul; all the intrusions are probably Early Devonian. Contact-metamorphosed slates, siltstones, and sandstones of probable Silurian to Devonian age border the intrusions and form irregular ridges and knobs adjacent to the valley.

Brachiopods put limits on extent of pre-Silurian strata

Fossil brachiopods found by G. H. Espenshade a few thousand feet west of Bear Pond, in the northeastern part of First Roach Pond quadrangle, central Maine, have been identified by A. J. Boucot as late Llandovery to Wenlock in age. Presumably this would restrict the area of pre-Silurian rocks to the country north of Farrar Mountain; these older rocks were formerly thought to extend about 12 miles farther southwest.

Tin content of stream sediments highest near biotite granite

Tin has been determined spectrographically in some 3,800 stream-sediment samples taken by E. V. Post and W. H. Dennen in west-central and southeastern Maine. It is detectable in only 4 percent of the samples from west-central Maine but in more than 13 percent of the samples from southeastern Maine. In west-central Maine, tin is common only in sediment of streams draining the Katahdin Quartz Monzonite. In southeastern Maine, tin is widespread, but highest values are found in the sediment of streams that drain plutons of gray to pink biotite granite. This relationship appears significant

because of the presence in southern New Brunswick of the Mount Pleasant tin deposit near the north margin of the St. George biotite granite pluton.

Evidence for pre-Taconic orogeny in northern Maine

An orogeny of Cambrian(?) to Early Ordovician age (pre-Taconic) is suggested by R. B. Neuman on the basis of data from northern Maine and adjacent Quebec, together with information from the Shin Pond quadrangle to the south; this orogeny was probably responsible for much of the structural contrast generally attributed to the Taconic orogeny (Late Ordovician–Early Silurian). Taconic structural contrasts in the Shin Pond-Stacyville area are relatively minor, but at this time the major anticline of the area began to rise, and subsequent Silurian sedimentation was controlled by its presence.

Heavy metals in stream sediments related to granitic plutons

Anomalous amounts of heavy metals in stream sediments, according to work by E. V. Post, are distributed peripherally to granitic plutons east of Lincoln, Maine, and between the Passadumkeag Mountains (Saponac quadrangle) and Tomah Mountain (Waite quadrangle). These anomalies in east-central Maine appear to be a northeasterly continuation of a belt of anomalies found to the east of the Penobscot River and also considered to be related to granitic plutons.

New evidence proves presence of Ordovician and Silurian rocks

Reconnaissance geologic mapping by Louis Pavlides in the Mattawamkeag Lake quadrangle in eastern Maine has shown that, contrary to previous beliefs, Ordovician rocks are present in this area. Also, discovery of a Silurian graptolite locality in the northern part of the quadrangle represents the first paleontologic evidence supporting the presumed Silurian age for a wide belt of micaceous quartzites and siltstones in this region.

APPALACHIAN REGION

STRUCTURAL AND STRATIGRAPHIC STUDIES

Upper Cretaceous alluvial clay in southern Pennsylvania

Nonmarine lignitic clay containing pollen and spores identified by R. H. Tschudy as Late Cretaceous in age (see "Paleontology, Cretaceous pollen in the Appalachian Mountains") was collected by K. L. Pierce from dumps of a 19th-century iron-mining area at Pond Bank, near Chambersburg, Franklin County, in southern Pennsylvania. According to

mine records and published descriptions, the lignitic clay is 40 feet below ground and overlies at least 170 feet (possibly 335 feet) of clayey residuum on the Tomstown Dolomite. Pierce suggests (p. C152–C156), on the basis of the proportion of insolubles in the dolomite and of recent estimates of total erosion of the Appalachian Highlands since Cretaceous time, that at least 140 feet of residuum has accumulated since the lignitic clay was deposited. The Pond Bank deposit may be a remnant of a more extensive nonmarine alluvial cover deposited at least 1,400 feet above the present ground surface, or several hundred feet above the highest mountains in the area. If this is true, the deposit is considerably older than any of the existing geomorphic features.

Mauch Chunk red beds increase southeastward in eastern Pennsylvania

Stratigraphic studies in Luzerne County, in eastern Pennsylvania, by M. J. Bergin and J. F. Robertson show that southeastward thickening of the Mauch Chunk Formation is accompanied by a regular increase in the amount of red beds within the formation. In Susquehanna Gap, north of Pittston, the formation is 255 feet thick and contains no red beds. The thickness increases to 550 feet north of Kingston, where red-bed sequences as much as 20 feet thick make up about 25 percent of the formation. Red-bed sequences as much as 100 feet thick make up 40 to 50 percent of the formation north of Nanticoke and on Penobscot Mountain south of Ashley, where the Mauch Chunk is respectively 670 feet and 1,000 feet thick. The underlying Pocono Formation also thickens southeastward at the somewhat lower rate of about 25 feet per mile.

Underthrusting in Pennsylvania anthracite region

Structural analyses by G. H. Wood, Jr., indicate that the low-angle thrust and underthrust faults of the southern part of the Anthracite region of eastern Pennsylvania are located principally at horizons which separate thick sequences of competent and incompetent rocks. The forces that developed these faults were transmitted differentially according to the relative competency of the sequences. Competent sequences overlying less competent sequences commonly have been thrust over older rocks for considerable distances, the fault planes paralleling the attitudes of the overlying competent sequences and truncating underlying crumpled sequences. Where competent sequences lie below less competent ones, however, they commonly have been underthrust from the troughs of synclines toward the crests of anticlines. The planes of the underthrust

faults generally parallel the attitudes of the underlying competent sequences and truncate overlying crumpled, less competent sequences; this is the reverse of the relation common on the thrust faults.

Revision of faults by new mapping in southwestern Virginia

Geologic mapping by R. L. Miller and J. B. Roen in the coal fields of Wise and Lee Counties, in southwestern Virginia, has demonstrated that two large faults formerly thought to have eliminated up to 1,500 feet of beds of the Norton and overlying formations at the surface are nonexistent. In Wise County a sharp flexure with only minor fracturing and slippage of beds is present where the Pigeon Creek fault was thought to be. In Lee County, a similar sharp flexure where the North Fork fault was supposed to be manifests no fracturing or displacement whatever. The lower mined coals in Lee County, supposedly in the Wise Formation, are in reality in the Norton Formation. In Wise County there are no economically important coals close to the belt where the Pigeon Creek fault was supposed to be.

Probable presence of Pine Mountain fault in Wise County, Va.

Mapping by R. L. Miller has demonstrated, at the head of Powell Valley in Wise County, southwestern Virginia, the presence at the surface, over a wide area, of intensively deformed black shale of the Chattanooga shale (Devonian), which suggests strongly that the Pine Mountain fault or a branch of it is close to the surface in this region.

Pocahontas Formation extended in Virginia

Stratigraphic studies by K. J. Englund and A. O. DeLaney indicate that the Pocahontas Formation and several economically important coal beds within it are more widespread in southwest Virginia than was previously known. Recognition of beds containing fossil plants of the *Neuropteris pocahontas* zone (zone 4 of Read and Mamay²⁷) has been useful in the local and regional correlation of the Pocahontas Formation.

Barite deposits related to paleokarst in Virginia

From field examination of mines and prospects in the Clinch River barite district in Tazewell and Russell Counties, southwestern Virginia, Helmuth Wedow, Jr., concludes that bedrock occurrences of barite, fluorite, and sulfides, chiefly pyrite, are in

²⁷ C. B. Read and S. H. Mamay, 1964, Upper Paleozoic floral zones and floral provinces of the United States: U.S. Geol. Survey Prof. Paper 454-K, p. K1-K35.

solution-collapse breccias of the Kingsport and Longview Formations in the upper part of the Knox Group. As elsewhere in the southern Appalachian Valley, these breccias are localized along the interface between the uppermost thick limestone of the group and overlying fine-grained dolomite. The close association of the breccias with the post-Knox unconformity is well illustrated by the occurrence of red dolomites of the basal Middle Ordovician Blackford Formation as a breccia matrix and solution-channel filling through a zone several hundred feet thick.

Late Precambrian volcanism in southwestern Virginia

Volcanic rocks of the Mount Rogers Volcanic Group in the Blue Ridge province of southwestern Virginia and northwestern North Carolina are of late Precambrian age, and range from basalt through latite to rhyolite, according to D. W. Rankin. A major center for the extrusion of rhyolitic rocks was in the area of the Mt. Rogers massif, Virginia. Rankin has mapped two rhyolite units there: a lower one characterized by 0 to 10 percent phenocrysts; and an upper, porphyritic rhyolite containing 30 percent phenocrysts of quartz and alkali feldspar. The lower unit is about 2,000 feet thick and occurs largely as highly spherulitic lava flows with some welded-ash flows. The upper, porphyritic unit is probably thicker and consists largely of welded-ash flow sheets. A suggested volcanic history of the rhyolites begins with the relatively quiet eruption of hot, relatively dry magma, resulting in the lower, phenocryst-poor spherulitic lavas. With subsequent crystallization in the magma chamber, an increase in the vapor pressure may have given rise to relatively violent eruptions of porphyritic ash-flow sheets that constitute the upper rhyolite unit. Volcanic breccias consisting of fragments of the older, phenocryst-poor rhyolite in a matrix of porphyritic rhyolite crop out on the Mt. Rogers massif within the outcrop area of the younger rhyolite. These breccias which are interpreted as breccia pipes, support the general location of the volcanic center and the concept of late violent eruptions.

Clinchport fault in eastern Tennessee and southwestern Virginia

Mapping and detailed facies studies of the Cambrian rocks along the Clinchport and Hunter Valley faults in eastern Tennessee and southwestern Virginia by L. D. Harris (p. B49-B53) has helped to determine the structural relations between these faults. By comparing stratigraphic thicknesses of the Maryville Limestone and Rogersville Shale be-

tween Clinchport, Va., and Tazewell, Tenn., he has been able to show that the Clinchport thrust plate, with a thick section of Maryville and a thinner one of Rogersville, has overridden parts of the Hunter Valley thrust plate, in which these relative thicknesses are the reverse. This relation and reconnaissance mapping in other key areas suggest that the Clinchport fault is one of the major through-going faults of the southern part of the Valley and Ridge province.

Algal stromatolites in Copper Ridge Dolomite, Tennessee

Unusual wave-washed exposures of the Upper Cambrian Copper Ridge Dolomite, accessible during the low-water stages of Norris Lake in Claiborne and Union Counties, in northeastern Tennessee, were studied by L. D. Harris. These exposures reveal that perhaps as much as 60 percent of the formation in that area consists of algal stromatolites which form extensive bioherms 5 to 140 feet thick. Details of algal growth and relations to the enclosing clastic material are remarkably well displayed. Vugs occur locally in the central portions of algal heads and along growth layers within the heads. A petroliferous odor is commonly detected when the algal material is freshly broken, and the algal beds are therefore of special interest because of their possible bearing on the porosity and petroleum content of the dolomite.

Gold Hill fault zone in west-central North Carolina

Geologic mapping in the northwestern part of the Carolina slate belt in Rowan and Davidson Counties, N.C., by H. W. Sundelius, A. A. Stromquist, and A. R. Taylor has shown that most of the mines and prospects of the Gold Hill and Silver Hill mining districts lie between two fault zones within the slate-belt rocks. The rocks adjacent to and between the faults are highly sheared slate, phyllite, and schist which have been hydrothermally altered in many places. They locally contain small bodies of sheared, coarse-grained diorite. The faults cut across major folds in less metamorphosed volcanic and pelitic sedimentary rocks on the east. Structural relations of the rocks of the Charlotte belt adjoining on the west are still problematical.

Ordovician age of Rockmart Slate in Georgia established

Charles Cressler has found graptolites in the Rockmart Slate of Hayes (1894)²⁸ in Polk County, in northwestern Georgia, which are of middle Ordovician age. Prior age assignments for the Rockmart

²⁸C. W. Hayes, 1894, *Geology of a portion of the Coosa Valley in Georgia and Alabama*: Geol. Soc. America Bull., v. 5, p. 465-480.

were Ordovician and, after 1937, Mississippian. After this discovery R. A. Laurence learned from field notes that M. R. Campbell discovered graptolites in the Rockmart in 1890; but the find never was published and was forgotten. Thus, the confusion over the age of the Rockmart resulted because fossils were not rediscovered until the present.

GEOPHYSICAL STUDIES

Magnetic instability of Pennsylvania diabase

Geophysical investigations in the Gettysburg-Newark basin, Pennsylvania, by M. E. Beck, Jr., have provided a partial explanation for instability of remanent magnetic moments in the diabase of that area. Results to date indicate that (1) there is a marked correlation between instability of remanent magnetization and weak total magnetic moment, including both susceptibility and remanence; (2) unstable, weakly magnetized samples have a higher-than-average concentration of late magmatic or hydrothermal material, especially vermicular granophyre. These observations apply to comparisons between separate samples from the same locality as well as to contrasts between localities. Grain-size differences, variations in titanium content of magnetite, or a combination of these factors could also account for the observed range of magnetic properties.

Magnetic and radioactivity data in mapping

Virginia Piedmont

A detailed aeromagnetic and aeroradioactivity survey was completed for 980 square miles of the Piedmont province in eastern Virginia to aid in geologic mapping of an area of complex lithology and scarce outcrops. The area studied consists of sixteen 7½-minute quadrangles in Spotsylvania, Culpeper, Orange, Louisa, Caroline, Hanover, and Stafford Counties. S. K. Neuschel reports that correlation between airborne geophysical data and relatively recent 1:62,500 geologic mapping in the Piedmont of Maryland is excellent. On the Virginia State geological map (scale 1:500,000) information is of necessity very generalized, and the geophysical data exhibit many features which the State map does not explain. The results of detailed fieldwork to date in the Spotsylvania area, however, show good correlation with the geophysical data. There, high radioactivity and low magnetic intensity over granite distinguish this rock from granitic gneiss. Also, mafic rocks not previously mapped have been located on the basis of low radioactivity and high magnetic intensity. Much of the Spotsylvania area

consists of rocks having alternating plagioclase-muscovite-chlorite-magnetite and quartz-plagioclase-muscovite assemblages. These two rock types can be delineated by using the geophysical data in conjunction with geologic mapping.

Gravity and magnetic fields in eastern Tennessee and southeastern Kentucky

J. S. Watkins has described (1-64) a geologic interpretation of gravity and magnetic data for 10,000 square miles of eastern Tennessee and southeastern Kentucky. Some gravity and magnetic anomalies associated with basement rocks in this area trend more nearly northward than do surface structural features, reflecting the pre-Appalachian structural grain of the area. A northeast-trending basement lineament marked by gravity and magnetic anomalies parallels the western margin of the Valley and Ridge province from west-central Alabama to northern Tennessee. This lineament seems to be a transition zone between a tectonically active Appalachian crustal block and a stable cratonic block. Magnetic data and well records indicate that the basement surface in the Valley and Ridge province is relatively flat in this area and lies generally less than 15,000 feet below sea level. The absence of magnetic or gravity anomalies along the major thrust faults suggests that they do not penetrate the basement.

EASTERN PLATEAUS

PENNSYLVANIA

Coal-basin deposition in Washington area

Statistical studies by B. H. Kent, J. P. Schweinfurth, and J. B. Roen of more than 350 core logs of Pennsylvanian strata in the Washington area in southwestern Pennsylvania have led to preliminary conclusions concerning coal-basin sedimentation. The stratigraphic interval between the bases of successive major coal beds (for example, the 310-foot interval from the base of the Pittsburgh coal bed to the base of the Waynesburg coal bed and the 130-foot interval from the Waynesburg coal bed to the Little Washington-Washington coal complex) has most promise for revealing relationships between sediment-distribution patterns and the distribution, thickness, and character of coal beds. The thicknesses of these intervals vary rapidly and locally, and vary reciprocally with the thicknesses of constituent coal beds and limestone units; if the coal beds and limestone units are thick, the interval

is thin. Where the Pittsburgh-to-Waynesburg interval is thin, the Waynesburg coal zone is thick and unusually complex; characteristically, the Waynesburg coal zone contains several partings and thin beds of coal below the main bench, and the main bench is thicker than usual for that local area.

Mauch Chunk Formation and Pottsville Group in north-central Pennsylvania

Recent mapping by G. W. Colton in parts of Lycoming, Clinton, Tioga, and Potter Counties, in north-central Pennsylvania, has demonstrated that (1) the Upper Mississippian Mauch Chunk Formation and the Lower and Middle Pennsylvanian Pottsville Group underlie appreciably larger areas than are shown on the 1960 geologic map of Pennsylvania; (2) some, at least, of the Mauch Chunk and Pottsville sediments were transported into this area from the north; and (3) the boundary between the two systems is a disconformity or a gentle angular unconformity, as was shown by earlier workers to the west. These conclusions are based on detailed mapping of distinctive soil and float material, regional stratigraphic relationships, measurement of pebble diameters, and measurement of foreset beds in the scant exposures present.

KENTUCKY

Glacial drift of Nebraskan age in northern Kentucky

Studies by L. L. Ray have shown that glacial drift of Nebraskan age, overlain by drift of Kansan age, is present on the uplands of Boone County in northern Kentucky a few miles southwest of Cincinnati, Ohio. The Nebraskan drift was deposited by an ice sheet whose effects have been hitherto unrecognized east of the Mississippi River valley. The character of the advanced profiles of weathering developed on the drift indicates weathering in place under poor drainage conditions on the upland surface prior to deposition of the overlying Kansan drift. A similar profile of weathering on the younger Kansan drift was modified later under better drainage conditions resulting from dissection of the uplands in post-Kansan time. Along the Ohio River valley to the west, where there was some dissection of the uplands in post-Nebraskan, pre-Kansan (Aftonian) time, drift of Nebraskan age is present in Kentucky only on remnants of the dissected uplands. Its profile of weathering differs from that of the undissected uplands of Boone County, reflecting the better drainage conditions following deposition. Here, drift of Kansan age does not overlie the Nebraskan drift but

is present at subsummit positions in stream valleys tributary to the Ohio and on bedrock benches along the river valley. Present studies suggest that the maximum extent of glaciation in northern Kentucky was in Nebraskan time.

Shoals and islands during deposition of Pennsylvanian strata in eastern Kentucky

Geologic mapping by W. F. Outerbridge in the Paintsville, Offutt, and Inez quadrangles, in Johnson and Martin Counties in eastern Kentucky, indicates that a northeastward-trending zone of islands and shoals existed during Pennsylvanian time²⁹ and influenced deposition of the Magoffin Beds of Morse (1931)³⁰, which are in the Breathitt Formation. The shoals and islands formed on pre-Magoffin sediments and seemingly separated brackish-water marshes and lagoons on the northwest from marine waters to the southeast. Beach sediments (disarticulated and unoriented shells mixed with plant debris in a thin calcareous sandstone bed) are present north of the southwest end of the line of shoals and islands. The Magoffin Beds are thin and locally ripple marked where deposited along the crests of the shoals, and are missing where there were islands. Linguloid brachiopods are abundant and fresh-water clams are present at the northeast end of the shoals. They are associated with a 1-inch to 6-inch layer of canneloid coal indicative of deposition in a marsh. Brachiopods, crinoids, pelecypods, and gastropods have been found on both sides of the shoals, but cephalopods have been found only on the southeast side. Siderite is abundant on the northwest side of the shoal line and scarce on the southeast, suggesting the presence of brackish water to the northwest.

Volcanic origin of flint clay in eastern Kentucky

The Fire Clay coal bed of Pennsylvanian age contains a thin parting of flint clay that occurs extensively within the Breathitt Formation in eastern Kentucky and in correlative strata in adjacent states. In recent thin-section studies of the clay from near Hazard, Ky., V. M. Seiders (p. D52-D54) has recognized grains of quartz and clastic sanidine in a groundmass of clay, part of which P. D. Blackmon identified by X-ray diffraction as kaolinite. A volcanic origin of the flint clay is indicated by the

²⁹ W. F. Outerbridge, 1965, Magoffin Beds of Morse in three eastern Kentucky quadrangles [abs.]: Geol. Soc. America Southeastern Section Mtg., April 8-10, 1965, Program, p. 31.

³⁰ W. C. Morse, 1931, The Pennsylvanian invertebrate fauna of Kentucky in *Paleontology of Kentucky*: Kentucky Geol. Survey, ser. 6, v. 36, p. 293-348.

clearly clastic nature of the grains of sanidine, by embayed quartz, and by the absence of any obviously nonvolcanic debris except argillaceous matter. Seiders concludes that the fire clay was formed by the in-place alteration of volcanic ash.

Cyclical deposits in Mississippian limestones, western Kentucky

Depositional cycles in the upper part of the Ste. Genevieve Limestone and in the lower part of the overlying Girkin Formation, both of Late Mississippian age, have been recognized in western Hart County, Ky., by C. A. Sandberg and C. G. Bowles (2-65). Crossbedded calcarenite (clastic limestone), which forms the base of each cycle, is derived partly from rounded fragments of crinoids, echinoids, and other fossils, and partly from subangular limestone and chert fragments reworked from underlying beds. The highest bed of calcarenite in the Ste. Genevieve Limestone comprises noncurving crossbeds that are suggestive of torrential or deltaic deposition and have initial dips of 16° to 22° . The middle of each cycle consists of light-colored limestone that is partly fossiliferous and partly oolitic. The top of each cycle is characterized by dark limestone and limestone intraformational breccia containing round, accretive, armored nodules of broken black chert.

Movement of Lower Early Mississippian sediments across Cincinnati arch

The Borden Formation of Early Mississippian age, along the western flank of the Cincinnati arch in central Kentucky, consists of sediments that seemingly were transported southwestward across the site of the arch. The Borden in this area is made up of two members, the lower consisting of siltstone and shale, and the upper—the Muldraugh Member—consisting of siltstone, dolomite, and limestone. Geologic mapping by R. C. Kepferle and W. L. Peterson in Nelson, Hardin, and Larue Counties has delineated a 1- to 2-mile-wide zone across which the lower member thins southwestward from 280 feet to 55 feet as the overlying Muldraugh Member thickens from 60 feet to 280 feet. This zone trends N. 55° W. to N. 60° W. across the Elizabethtown, Nelsonville, Hodgenville, New Haven and Howardstown $7\frac{1}{2}$ -minute quadrangles. The inclined contact between the two members and the southwestward inclination of many beds of limestone within the Muldraugh Member are considered to represent initial dips which are perpendicular to the trend of the zone.

The strike of the zone is parallel to that of limestone reefs described by Thaden and others³¹ in the Fort Payne Formation, which is about the stratigraphic equivalent of the Borden Formation, west of Columbia in south-central Kentucky. The reefs occur in an arcuate belt, south of a linear body of sandstone (C. H. Maxwell and W. B. Turner, 1-64) that seemingly was deposited shoreward of the reefs (U.S. Geol. Survey, 11-64, p. A85). Along the east side of the arch, generally south of Berea, Ky., beds of siltstone in the Borden Formation have been shown by G. W. Weir, J. L. Gualtieri, and S. O. Schlanger to have a pronounced southward initial dip. A generally south and southwestward movement of sediments across the arch during Early Mississippian time is thus indicated.

Facies in Ordovician strata, Cincinnati arch

Mapping and stratigraphic studies of Middle and Upper Ordovician strata on and along the flanks of the Cincinnati arch in Kentucky have revealed complex facies within the rocks and demonstrated some of the facies relations. Work by E. R. Cressman, D. F. B. Black, W. C. MacQuown, J. S. Pomeroy, and others has shown, for example, that in the Frankfort area, Kentucky, a southeastward-striking band of crossbedded bioclastic calcarenite near the top of Middle Ordovician strata grades abruptly southwestward into interbedded shale and calcilutite. It interfingers northeastward with very fossiliferous argillaceous limestone which interfingers to the northeast with interbedded shale and sparsely fossiliferous calcilutite. Seemingly, the crossbedded calcarenite was deposited on the windward side and the fossiliferous limestone on the lee side of a low bank. The distribution of these facies was not controlled by the Cincinnati arch.

G. W. Weir, W. L. Peterson, and J. H. Peck have traced Upper Ordovician rocks, on lithologic evidence, from central Kentucky along the east and west flanks of the Cincinnati arch into areas of Ohio and Indiana where the classical work on paleontology and stratigraphy of the Upper Ordovician (Cincinnatian) strata was done. In general, dolomite, dolomitic mudstone, and limestone deposited in shallow water grade northward into shale and limestone that were deposited in deeper waters. The rocks of these facies strike generally eastward, and their distribution seems not to have been influenced by the Cincinnati arch.

³¹ R. E. Thaden, R. Q. Lewis, J. M. Cattermole, and A. R. Taylor, 1961, Reefs in the Fort Payne Formation of Mississippian age, south-central Kentucky: Art. 39 in U.S. Geol. Survey Prof. Paper 424-B, p. B88-B90.

TENNESSEE

Movement along Jacksboro fault at southwest edge of Cumberland overthrust

Work by K. J. Englund in Campbell County, Tenn., indicates that there has been approximately 11 miles of strike-slip movement along the Jacksboro fault at the southwest edge of the Cumberland overthrust sheet. The amount of movement, which has generally been considered by previous workers to be of smaller magnitude, was determined from stratigraphic studies and thickness maps of formations on opposite sides of the fault.

SHIELD AREA AND UPPER MISSISSIPPI VALLEY

Subsurface structure of western Lake Superior region

W. S. White has analyzed the subsurface structure of the western Lake Superior region using a combination of surface geology and aeromagnetic, gravity, and paleomagnetic data. Surface attitudes and map patterns suggest that the upper Keweenaw sedimentary rocks have the general form of a lens thickening southeastward away from a feathered edge along the Minnesota shore of Lake Superior. Graphic subtraction of the assumed gravitational effect of this sedimentary lens from the Bouguer anomalies of the region leaves a residual anomaly due primarily to the mafic lavas and intrusive rocks.

The analysis leads to recognition of the following stages in the tectonic history of the region. (1) Accumulation, during middle Keweenaw time, of a thick series of lava flows and mafic intrusives in two basins or troughs, separated by a positive area trending more or less north-south across the Bayfield Peninsula, Wis., in which the lavas are thin or absent. The Duluth Gabbro Complex, in the more westerly trough, probably pinches out southeastward beneath the lavas somewhere near the Minnesota shore. (2) Evolution of the present Lake Superior basin, with axis trending northeast, during late Keweenaw time. (3) Development of the Ashland syncline and the major faults of the region (Douglas, Keweenaw, Lake Owen) still later in Keweenaw time.

MICHIGAN

Geology in eastern part of Gogebic district reinterpreted

Regional stratigraphy and structure at the east end of the Gogebic iron-bearing district, Michigan, have been reinterpreted by C. E. Fritts using results of recent detailed mapping in conjunction with earlier data recorded by the Michigan Geological

Survey. The Tyler Slate of Animikie age trends generally eastward to Lake Gogebic where it wraps around the nose of an east-plunging anticline, continues westward for about 5 miles, and is terminated by a fault of southwest strike. The Tyler on the south limb of this fold conformably underlies a thick sequence of metasedimentary and metavolcanic rocks that probably conformably underlies the Michigamme Slate with which the Tyler is not equivalent as now interpreted. The sequence that is apparently between the Tyler and Michigamme Slates contains graded bedding and pillow structures that are right side up, and the overall structure is a south-dipping monocline rather than a fold in part overturned, as has been previously proposed. Lean iron-formation at the Banner exploration is stratigraphically above the Tyler Slate and is in a south-plunging open synclinal flexure rather than in a tight fold. Similarly, the so-called Turtle range and an inferred eastern extension of the Marenisco range are stratigraphically above the Tyler and are not belts of tightly folded rocks.

Three kinds of rock of different ages have been recognized in what formerly was mapped as Presque Isle Granite. A banded gneiss and an equigranular granite are unconformably overlain by the Tyler Slate and are interpreted as pre-Animikie in age. A granitic to quartz monzonitic gneiss intrudes Animikie strata, and an increase of metamorphic grade southeastward toward a broad area underlain in part by this gneiss indicates that metamorphism probably accompanied and perhaps followed its emplacement.

WISCONSIN

Paleomagnetism sets limit to age of gabbro in Wisconsin

Kenneth Books has found that paleomagnetic field directions in the gabbro intrusion and associated lava flows near Mellen, northern Wisconsin, throw light on the time of intrusion of the gabbro. After removal of extraneous components of magnetization from the paleofield directions by demagnetization runs on 40 samples of gabbro and lava from 8 sites, it was observed that the mean remanence directions from Fischer statistical analyses when uncorrected for the regional geologic dip of 60°–70° NW, are all scattered in a direction similar to the direction of remanence, also uncorrected for regional dip, in the Duluth Gabbro at Duluth, Minn.³² The agreement between all site mean remanence directions when uncorrected for regional geologic dip suggest (1)

³² P. M. Dubois, 1962, Paleomagnetism and correlation of Keweenaw rocks: Canada Geol. Survey Bull. 71, p. 42–43.

that the lava flows adjacent to the gabbro intrusion at Mellen were heated through the Curie point by the intrusion, and (2) that the gabbro intrusion at Mellen occurred near the end of regional deformation in the area.

Diatomite discovered in Driftless Area, Wisconsin

G. W. Andrews has found a 6-inch bed of diatomite in the late Pleistocene alluvium of the Trempealeau Valley near Hixton, Wis., the first diatomite deposit to be reported from the Driftless Area of the upper Mississippi Valley. The diatom assemblage consists of 23 genera and 74 species of which 3 species are new. The assemblage is unusual in that it is dominated by a large number of individuals of a single minute form—*Fragilaria construens* var. *venter*. The diatomite is closely associated with an underlying deposit of spruce wood and matted organic debris, dated at about 11,000 years B.P., by Meyer Rubin using the radiocarbon method. The diatoms were probably deposited in a shallow pond during regional cooling of the climate brought about by the ice advance of the Valdres Stade.

MINNESOTA

Aeroradioactivity map reflects glacial geology

An aeroradioactivity map compiled by S. K. Neuschel, covering a 10,000-square-mile area that centers about Minneapolis and St. Paul, shows good correlation with glacial geology; little or no correlation is shown with bedrock, which is exposed only at a few scattered places, mostly along major streams. Numerous small elongate areas of low radiation: (1) are present over morainal deposits; (2) have a north-south to northeast-southwest trend, as do most of the existing lakes; (3) average about 2 miles in width; (4) range from 5 to 10 miles in length; and (5) are best developed in the area northeast of the Mississippi River. Low radiation is present also over swampy areas and silted-in former glacial lakes. The map indicates that the number of lakes was formerly three to four times that of today, and also indicates the former courses of some of the existing drainage.

NEBRASKA

Map of bedrock geology and thickness of overburden compiled for southeastern Nebraska

A map showing bedrock geology and thickness of Quaternary overburden for an area of 8,400 square miles in southeastern Nebraska has been prepared cooperatively by G. E. Prichard, of the U.S. Geo-

logical Survey, and E. C. Reed, V. H. Dreezen, and R. R. Burchett, of the Nebraska Geological Survey.

INTERIOR HIGHLANDS AND EASTERN PLAINS

Age and correlation of old gravels in Val Verde County, Tex.

J. A. Sharps and V. L. Freeman have mapped extensive deposits of gravel in the southern half of the Feely quadrangle, Val Verde County, Tex. The oldest deposits, not terrace deposits related to the present river system, can be divided into two units of approximately the same age. One unit contains gravel composed solely of limestone and chert deposited from a southward-flowing stream. The other unit contains gravel from a limestone terrane as well as characteristic pebbles of fine-grained igneous and metamorphic rocks deposited from an eastward-flowing stream. The area 5 to 10 miles southeast of Comstock in the Feely quadrangle is probably the junction area of the 2 streams.

No direct evidence of age of these older gravels has been found. Their topographic position and the position of the Uvalde Gravel to the southeast suggests an equivalence. The Uvalde is generally called Pliocene(?) in age. Since the deposition of the Uvalde Gravel and the possible equivalent gravels of the Feely quadrangle there has been major uplift of the Edwards Plateau, including its westward extension the Stockton Plateau, and of the adjacent region in Mexico. The southern part of the Edwards Plateau, north of the Balcones Fault zone, probably has been tilted northward. The region along the Rio Grande, west of a line between Cline and Laredo, probably has been tilted eastward. This uplift and tilting are here considered Pleistocene in age.

Sedimentary slump(?) structure related to folding in Arkansas

A large-scale sedimentary flow feature in the lower part of the Savanna Formation of Middle Pennsylvanian age has been mapped by B. R. Haley in the western half of sec. 9, T. 7 N., R. 24 W., in Logan County, western Arkansas. Here a rock unit about 10 feet thick and composed mostly of sand along the limbs of the east-trending Paris syncline thickens along the axis of the syncline to a unit about 60 ft thick composed of mixed sand, silt, and clay. This feature is best explained by postulating preconsolidation sedimentary slump or flow immediately after the deposition of the sand. The sedimentary flow could have resulted from a gradient

established by the folding of the Paris syncline during Savanna deposition. Sandstone units in the underlying McAlester Formation become finer grained northward, either thin or pinch out northward, and converge northward across the syncline, indicating that the start of the folding in this area predates Savanna time.

Diagenesis of Pennsylvanian oolitic limestones in Arkansas

L. G. Henbest has found microtextural evidence indicating that diagenetic changes in the oolitic limestones of the Pennsylvanian Morrow Series in Washington County, Ark., can be grouped in the following 3 stages. (1) Formation of microstylolites, before lithification, by solution interpenetration at points of contact between oolites. (2) Secondary enlargement through growth as euhedral crystals of quartz sand grains that were nuclei of oolites or that touched oolites accompanied by etching of those quartz grains exposed to the matrix. (3) Because insufficient carbonate was mobilized in stages 1 and 2 to account for all the reconstruction of the matrix, a third and later stage of cementation is suggested.

These diagenetic processes play a prominent role in the evolution of marine limestones and influence their behavior as reservoirs of oil or of water.

NORTHERN ROCKY MOUNTAINS AND PLAINS

WASHINGTON

Covada Group of northeastern Washington revised

High-grade metamorphic rocks in the Twin Lakes quadrangle were originally included in the Covada Group, of probable Devonian to Mesozoic age, by J. T. Pardee,³³ but recent mapping by G. E. Becraft suggests that they are much older, possibly as old as Precambrian. These old rocks (high-grade gneisses interlayered with schists and white quartzite) are separated from younger less metamorphosed rocks of the Covada Group (phyllitic graywacke, phyllite, slate, and impure bluish-gray quartzite) by an eastward-dipping fault. Rocks of the Covada Group are intensely folded and faulted, but layering and foliation in the older high-grade metamorphic rocks dip gently to the west. The old metamorphic rocks are similar to the metamorphic rocks of Tenas Mary Creek in the Curlew quadrangle to the northeast.

³³ J. T. Pardee, 1918, *Geology and mineral deposits of the Colville Indian Reservation, Washington*: U.S. Geol. Survey Bull. 677, 186 p.

"Syncline" in northeastern part found to be a tilted fault block

R. C. Pearson has interpreted an 8- to 12-mile-wide belt of Tertiary rocks in the Bodie Mountain quadrangle, northeastern Washington, to be a tilted fault block rather than a syncline as shown on the tectonic map of the United States. The west side is bounded by numerous north- to north-northeast-trending faults with small displacement, and the east side is marked by high-angle faults with major movement as well as inferred low-angle normal (?) faults that may extend the length of the quadrangle and beyond.

IDAHO

Drainage reversed on Lemhi River during Pleistocene

The direction of ancestral drainage in the intermontane trenches of east-central Idaho has long been a subject of controversy, but E. T. Ruppel believes that stream patterns and the distribution of surficial deposits provide conclusive evidence for reversal of the Lemhi River in Pleistocene time. The Lemhi presently flows northwestward, and empties into the Salmon River. Before arching in Pleistocene time reversed the flow in the northern segment of the river it must have flowed southeastward, to empty into the Snake River. This conclusion suggests that the parallel and very similar Pahsimeroi River was reversed at the same time, and that the Salmon River was formed concurrently as erosion, stimulated by arching and faulting, combined segments of many streams into the present Salmon River canyon.

Pull-apart theory not applicable to Gem Valley

Although Gem Valley, in southeastern Idaho, has been thought to be a pull-apart or tensional rift on the back edge of a thrust plate, S. S. Oriel suggests (p. C1-C4) that the marked contrasts in thicknesses and facies of Paleozoic stratigraphic units flanking the central and southern parts of the valley make it unlikely that original sites of deposition were ever any closer together than they are now.

Volcanic rocks once considered Precambrian may be Tertiary

The Bannock Volcanic Formation, as redefined by Ludlum (1942),³⁴ was once thought to be of Precambrian age and the oldest unit in the vicinity of Pocatello, Idaho, but D. E. Trimble believes that field relations of the rocks indicate that they are much younger, perhaps as young as early Tertiary. In at least one locality the Bannock Volcanic Formation

³⁴ J. C. Ludlum, 1942, *Precambrian formations at Pocatello, Idaho*: Jour. Geology, v. 50, no. 1, p. 85-95.

overlies Precambrian quartzite. The volcanic rocks appear to be both undeformed and unaltered, except for extensive propylitization. Their attitude is discordant with adjoining rocks and they appear to have been erupted onto a topography that was not greatly different from that of the present.

First evidence of Precambrian rocks in Idaho batholith

Geologic studies by B. F. Leonard and isotope analyses by T. W. Stern provide the first evidence for Precambrian deformation and intrusion in old rocks preserved in the central part of the area now occupied by the Idaho batholith. Isotope dilution analysis by Stern gives a discordant Pb^{207}/Pb^{206} age of 680 million years for zircon from uncontaminated syenite of the pre-Idaho batholith Ramey Ridge Complex described by Leonard.³⁵ Other measured ages for the same zircon are: Pb^{206}/U^{238} 574 m.y., Pb^{207}/U^{235} 596 m.y., Pb^{208}/Th^{232} 538 m.y. The measured age of 680 m.y. is a minimum age for this intrusive complex, and for the tight northwest-trending folds in metasedimentary and metavolcanic rocks that it cuts. Hornblende from pegmatitic syenite of the complex has a K-Ar age of 93 m.y., according to R. F. Marvin; this is close to the Pb/α age of 108 ± 12 m.y. previously established by Larsen and others³⁶ for the Idaho batholith, which intrudes the Ramey Ridge Complex. If radiogenic lead loss is assumed to have occurred at 93 m.y., the discordia age of the syenite zircon is 725 m.y. Some lead loss may have occurred still earlier. Unless occult contamination affected the dated zircon, both the syenite complex and one major episode of folding are of Precambrian age.

MONTANA

Deformation and crustal shortening in eastern Rocky Mountains

The total crustal shortening on the thrust faults and folds in a section $17\frac{1}{2}$ miles long in the Sun River Canyon area of the disturbed belt of Montana have been computed by M. R. Mudge to be about 29 miles.

In the Wolf Creek area, R. G. Schmidt has mapped three small klippen that show that the El Dorado thrust sheet once extended at least 3 miles east of its present trace. The klippen consist of Meagher Limestone of Middle Cambrian age, and lie on rocks of the St. Mary River Formation of Late Cretaceous

age; they are thought to have been dragged along beneath a thrust plate that moved Belt rocks of Precambrian age eastward over rocks of Late Cretaceous age.

Sequence of thrusts records decreasing deformation

Mapping by G. D. Robinson and W. B. Myers in the northern Big Belt Mountains shows that the younger members of an overlapping sequence of major Laramide thrust faults have been deformed progressively less than the older by contemporaneous uplift of the Big Belt dome. The oldest thrust recognized dips northeastward on the northeast flank of the uplift and in places is very strongly deformed. Overlapping this, two successively younger thrusts show progressively less deformation. The El Dorado thrust, the youngest in the area, generally maintains the regional southwestern dip but is itself deformed on a small scale. Highly plastic deformation, notably in the axial region of the dome and in the floor of the oldest thrust, indicate both continued eastward thrusting and the uplift of the Big Belt dome.

Down-dropped blocks in diatremes near Bearpaw Mountains

Mapping southeast of the Bearpaw Mountains by B. C. Hearn, Jr., and C. P. Sabine has shown that diatremes as small as 150 feet in diameter contain sedimentary rocks of early Tertiary age 3,000 to 4,500 feet below their normal stratigraphic position. Hearn and Sabine suggest that diatreme-type volcanic activity is a possible mechanism for emplacement of isolated down-dropped fault blocks.

Correlation of rock units in western Montana

Reconnaissance mapping of Precambrian Belt Series rocks west of Anaconda by M. R. Klepper indicates that the Newland Formation of the Phillipsburg area is almost certainly an equivalent of the Helena Dolomite, and not of the Newland of the type area. A calc-hornfels unit of the Precambrian mapped by Klepper and H. W. Smedes in the Highland Mountains south of Butte probably is also an equivalent of the Helena Dolomite.

In the Alberton area, west of Missoula, a nearly pure pink and white quartzite, 60 to 100 feet thick, has been correlated tentatively with the Cambrian Flathead Quartzite by J. D. Wells. The quartzite rests, in part, on diabase sills and, in part, overlies with apparent regional unconformity different horizons of the upper part of the Missoula Group (gray-green micaceous siltite and argillite, and pink and white micaceous, feldspathic quartzite with conspicuous crossbedding). In nearby areas similar pure quartzite that has been included in the Pilcher

³⁵ B. F. Leonard, 1963, Syenite complex older than the Idaho batholith: U.S. Geol. Prof. Paper 450-E, p. 93-97.

³⁶ E. S. Larsen, Jr., and others, 1958, Lead-alpha ages of the Mesozoic batholiths of western North America: U.S. Geol. Survey Bull. 1070-B, p. 35-62.

Quartzite of Precambrian age is probably also correlative with the Flathead Quartzite.

WYOMING

Marker horizons in Eocene rocks

Recent stratigraphic studies of the Wind River Formation of early Eocene age in the western Wind River Basin by J. F. Murphy have revealed a continuous mappable 20- to 40-foot tuffaceous sandstone sequence 500 to 700 feet above the base of the formation, which can be used effectively to divide the Wind River Formation into two mappable units.

A distinctive assemblage of vertebrate and invertebrate fossils has been discovered by W. L. Roher and C. L. Gazin (p. D133-D138) in a claystone bed in the lower Eocene Willwood Formation about 750 feet below its upper contact in the Big Horn Basin about 35 miles east of Cody, Wyo. This bed is mappable and provides stratigraphic control where control has been lacking. The bed contains fossil vertebrates common to both the Gray Bull and Lysite faunal zones; the Gray Bull fauna occurs below this bed and the Lysite fauna occurs above. The invertebrates in the bed are fresh-water mollusks.

Present-day canyons have Eocene ancestry

J. F. Murphy has shown that the boulder deposits of the Wind River Formation of early Eocene age that are found along the eastern flank of the Wind River Range are actually fan deposits, the apices of which point into, or are in, present-day canyons. The position of these deposits indicates that many of the present canyons were cut before or during early Eocene time.

Underground connections in Daisy geyser group, Yellowstone Park

J. J. Rowe, R. O. Fournier, and G. W. Morey (p. B184-B186) used sodium iodide as a tracer to prove the existence of underground connections linking several geysers and a pool of the Daisy group at Yellowstone Park. After adding sodium iodide to Splendid Geyser, water samples were taken at frequent intervals at Splendid, Daisy, and Comet Geysers and at Bonita Pool. Iodine determinations made in the field using a battery-operated photoelectric colorimeter showed that: (1) connections exist among all four features, (2) the connection among Splendid, Bonita, and Comet are relatively shallow (indicated by the rapidity of transfer of water containing iodide), (3) a reservoir of about 71,000 gallons is situated under Splendid, and (4) a small geyser east of Bonita is not connected by shallow channels to the Daisy group.

Devonian fish support paleogeographic connection between formations

The discovery by C. A. Sandberg and J. F. Murphy of fossiliferous Lower Devonian rocks at Bull Lake Canyon, on the east flank of the Wind River Range, supports a paleogeographic connection, postulated by C. A. Sandberg,³⁷ between continental and marginal marine beds of the Beartooth Butte Formation on the east and nearshore marine beds of the Water Canyon Formation on the west. The fossils are well-preserved heterostracan fish remains collected from a channel-fill deposit of the Beartooth Butte Formation. This locality is 125 miles due south of the closest outcrop and type section of the Beartooth Butte Formation and 115 miles east of the nearest outcrop of the correlative Water Canyon Formation of Utah and Idaho.

Regional correlations of the Darby Formation

Regional stratigraphic studies of Devonian rocks by A. L. Benson, J. F. Murphy, and C. A. Sandberg in west-central Wyoming have demonstrated that the type Darby Formation³⁸ on Sheep Mountain, on the west flank of the Wind River Range, includes limestone and shale correlative with the Madison Limestone of Mississippian age and with a dark shale unit of Late Devonian and Early Mississippian age. The remainder of the type Darby is correlative with, and continuous with, the lower member of the Jefferson Formation (Sandberg, 1-65). On the east flank of the Wind River Range the dark shale unit at the top and beds equivalent to the Maywood Formation of Late Devonian age at the base have been mapped with the Darby by Murphy and Richmond.³⁹ Lithologic and paleontologic data demonstrate that the dark shale unit, which locally overlies older rocks in angular unconformity, is a basal facies of the Madison, and is in part Devonian in age.

Regional unconformity within the Mesaverde Formation

A widespread unconformity at the base of the Teapot Sandstone Member of the Upper Cretaceous Formation has been defined by J. R. Gill in the Big-horn, Wind River, and Hanna Basins. The unconformity increases in magnitude from east to west and appears to reflect broad gentle uplift in western Wyoming modified by intense local uplift in some areas such as the Lost Soldier-Lamont area at the

³⁷ C. A. Sandberg, 1961, Widespread Beartooth Butte Formation of Early Devonian age in Montana and Wyoming and its paleogeographic significance: *Am. Assoc. Petroleum Geologists Bull.*, v. 45, no. 8, p. 1301-1309.

³⁸ Eliot Blackwelder, 1918, New geological formations in western Wyoming: *Washington Acad. Sci. Jour.*, v. 8, no. 13, p. 417-426.

³⁹ J. F. Murphy and G. M. Richmond, 1965, Geologic map of Bull Lake West quadrangle: U.S. Geol. Survey Quad. Map GQ-432.

southeast end of the Wind River Range (Zapp and Cobban ⁴⁰). In south-central Wyoming, M. W. Reynolds finds that this unconformity cuts downward across the Mesaverde Formation from south to north; 1,400 feet of strata has been eroded below the Teapot Member between Rawlins and Separation Rim, and the Camp Creek syncline. Reynolds also recognizes an unconformity in post-Teapot, pre-Lewis time in which the Lewis Shale overlies successively older beds of the Mesaverde northward and ultimately rests on the Cody Shale. Both unconformities increase in magnitude toward the Sweetwater arch, suggesting that the arch was tectonically active in Late Cretaceous time.

According to Gill the regression in western Wyoming represented by the basal Teapot unconformity appears to have taken place concurrently with transgression of the sea into western Montana marginal to the Elkhorn Mountains volcanic field. This activity culminated during the range span of *Baculites compressus* (late Campanian) in the Upper Cretaceous. The absolute time of this event can be postulated from a potassium-argon date of about 75 million years obtained for biotite from a bentonite occurring in the zone of *Baculites compressus* reported by Folinsbee and others.⁴¹

Structural relief of Precambrian surface inferred from gravity data

Geologic and gravity profiles drawn by J. E. Case and W. R. Keefer transect the area from the deep part of the Wind River Basin to the crest of the Owl Creek Range, west of Wind River Canyon, and show that structural relief of the Precambrian surface along the line of profile from the basin to the crest of the uplift is on the order of 21,000 feet. The steep gravity gradient across the southern front of the Owl Creek Range indicates that the bordering major fault zone dips at a high angle. The gravity data support earlier interpretations, based upon regional stratigraphic and structural studies by Keefer, that movements between basin and mountain provinces in central Wyoming during the Laramide were predominantly vertical.

NORTH AND SOUTH DAKOTA

Precambrian stratigraphy in central Black Hills, S. Dak.

A working stratigraphic section of the Precam-

⁴⁰ A. D. Zapp and W. A. Cobban, 1962, Some Late Cretaceous strandlines in Wyoming: Art. 134 in U.S. Geol. Survey Prof. Paper 450-D, p. D52-D55.

⁴¹ R. E. Folinsbee, Halfdan Baadsgaard, and J. Lipson, 1961, Potassium-argon dates of Upper Cretaceous ashfalls, Alberta, Canada, in J. L. Kulp, ed., Geochronology of rock systems: New York Acad. Sci., Annals, v. 91, art. 2, p. 352-363.

brian rocks in the Hill City and Medicine Mountain quadrangles of the central Black Hills has been established by J. C. Ratté. The section is 30,000 to 50,000 feet thick and consists of the following units, in ascending order: (1) biotite-quartz schist, 1,000 to 1,500 feet thick; (2) a heterogeneous unit of quartz-feldspar gneiss and quartzite that includes metaconglomerate and arkose, tremolite-carbonate schist, and other metamorphosed calcareous sediments, graphitic quartzose schist, sugary quartzite, and beds of carbonate iron formation, 5,000 to 6,000 feet thick; (3) thin-bedded highly sheared phyllite and schist containing small bodies of metagraywacke, 20,000 to 30,000 feet thick; unconformity at base marked by irregularities in thickness and overlap of quartzite beds of unit 2; (4) principally thick-bedded metagraywacke and quartzose schists, 5,000 to 10,000 feet thick, correlated with the Bugtown Formation of the Fourmile and Berne quadrangles to the southwest. Unconformities may exist at the base of units 2 and 4. Pegmatitic granite intrudes the entire metasedimentary section. The recognition of this stratigraphic section will permit reasonable stratigraphic and structural interpretations of the central Black Hills.

Glacial features are channel fill, not lake deposits

C. H. Baker, Jr., has studied a series of short ridges of sand and gravel that lie above the highest shoreline of glacial Lake Agassiz. These ridges are part of a long sinuous deposit of sand and gravel, generally occupying a topographic low, which extends southeastward from the southernmost bend in the Sheyenne River in North Dakota to Lake Traverse, S. Dak. The sand and gravel seem to have been deposited by streams in an ice-margin channel rather than as beach deposits in hypothetical Lake Milnor as postulated by earlier workers. Baker suggests that the name "Lake Milnor" be changed to "Milnor Channel" and the name of the sand and gravel ridges that border the channel be changed from "Milnor beaches" to "Milnor Channel deposits."

SOUTHERN ROCKY MOUNTAINS AND PLAINS

GEOLOGY OF PRECAMBRIAN ROCKS

Mafic and ultramafic rocks in two alkalic complexes, Wet Mountains, Colo.

Detailed mapping by R. L. Parker and D. R. Shawe have shown that the mafic and ultramafic rocks associated with the alkalic intrusions of Gem Park and Iron Mountain east of Hillside, in the northern Wet

Mountains, Colo.,⁴² form two discrete stratiform complexes. The complex at Iron Mountain underlies about 3½ square miles and that at Gem Park about 2 square miles. Both complexes are distinctly layered. The layers consist of gabbro, olivine gabbro, pyroxenite, pyroxene-olivine rocks, and dunite and are inclined toward the centers of the complexes. The gabbro layers show many features such as rhythmic banding, scour-and-fill structures, and crossbedding which are typical of the classic Skaergaard intrusive, Greenland, and of the Stillwater Complex, Montana.

Metasedimentary rocks in northern Front Range, Colo.

Studies by W. A. Braddock have shown that the metasedimentary rocks in the vicinity of the Thompson Canyon in the northern Front Range, Colo., were folded and metamorphosed twice during Precambrian time. The early deformation produced large isoclinal folds with axial-plane schistosity. Although bedding in the hinges of the folds was obliterated by the axial plane schistosity, the folds can be identified by the orientation of graded bedding in the limbs. Metamorphism produced quartz-albite-muscovite phyllites in at least part of the area, but it is not known whether a zonal pattern of higher grade mineral assemblages was also formed during this first period.

During the second period of folding, the rocks were deformed primarily by pervasive displacements along two conjugate sets of slip cleavage planes. The beds that previously had been tilted to steep angles were deformed into steeply plunging folds whose axial planes make large angles with the axial planes of older folds. After this folding the rocks were again metamorphosed, producing a zonal pattern of mineral assemblages that probably reflects a temperature gradient. Biotite, garnet, staurolite, andalusite, sillimanite-muscovite, and sillimanite-microcline zones have been recognized.

Quartz diorite similar to the Boulder Creek Granodiorite probably was emplaced during the older period of deformation. Tonalite (Mount Olympus Granite) and granodiorite (Sherman Granite) was intruded after the second period of deformation; at a still later time, quartz monzonites of the St. Vrain and Log Cabin batholiths were intruded.

Isoclinal folding in central Front Range, Colo.

A symmetrical pattern of Precambrian metasedimentary rock units in the Squaw Pass quadrangle in

the central Front Range, Colo., is interpreted by D. M. Sheridan to be the result of an early period of isoclinal folding. The axial area of a major fold is believed to lie in a west-northwest-trending major belt of hornblende gneiss, which is bordered successively to the north and south by parallel major belts of sillimanitic biotite gneiss, hornblende gneiss, and microcline gneiss. Marker layers of cordierite-sillimanite-biotite gneiss and a distinctive feldspathic calc-silicate gneiss, apparently unique to this area, form a symmetrical pattern within the central belt of hornblende gneiss. Lineations of the area plunge predominantly west-northwest at a low to moderate angle parallel to the inferred major axis of folding. The isoclinal folding is considered to be the oldest of three periods of Precambrian deformation recognized in this region.

STRATIGRAPHY AND STRUCTURAL GEOLOGY

Age of Sangre de Cristo Formation, New Mexico

Detailed mapping by E. H. Baltz in the Montezuma and El Porvenir quadrangles in the Sangre de Cristo Mountains near Las Vegas, N. Mex., has shown that the upper part of the arkosic limestone member of the Madera Limestone grades northward into shales and arkoses of the lower part of the Sangre de Cristo Formation. Fusulinids collected from the arkosic limestone member indicate a Missouri and Virgil (Late Pennsylvanian) age for the part of the member that grades into the Sangre de Cristo, thus confirming the Pennsylvanian and Permian age of the Sangre de Cristo Formation in the region to the north. South of Las Vegas, the Madera Limestone was deformed into several large folds in latest Pennsylvanian or Early Permian time, and the Permian part of the Sangre de Cristo Formation lies with angular unconformity on the arkosic limestone member of the Madera. Major Laramide folds and faults are oriented generally parallel to the late Paleozoic folds.

Red-bed deposition in New Mexico

Red-bed deposition in the late Paleozoic began at different times in different places in central New Mexico. In the north-central Oscura Mountains, G. O. Bachman has found minor red beds in strata of Missouri (Late Pennsylvanian) age; in the Manzano Mountains, D. A. Myers reports that although the first red beds were deposited in early Virgil (Late Pennsylvanian) time, large-scale red-bed deposition did not occur until late Virgil or Early Permian time. In the southern San Andres Mountains, Bachman and Myers have found that the low-

⁴² R. L. Parker and F. A. Hildebrand, 1963, Preliminary report on alkalic intrusive rocks in the northern Wet Mountains, Colorado: Art. 181 in U.S. Geol. Survey Prof. Paper 450-E, p. E8-E10.

est red beds are above the Hueco Limestone of Permian age.

Widespread regional unconformity of Jurassic age

An unconformity marked by a distinctive chert-pebble zone has been traced by G. N. Pipiringos from the Black Hills in South Dakota throughout central and southeastern Wyoming, northwestern Colorado, eastern Uinta Mountains, and into the Colorado Plateau, where it had been recognized earlier by other workers. In southeast Wyoming the chert-pebble zone marks the base of the Canyon Springs Sandstone Member (Upper Jurassic) of the Sundance Formation. In central Wyoming it lies at the top of the Gypsum Spring Formation (early Middle Jurassic) and locally underlies beds of probable late Middle Jurassic age. In the Uinta Mountains the chert-pebble zone lies at the top of the Glen Canyon Sandstone and is overlain by either the Carmel Formation or the Entrada Sandstone. The chert-pebble zone is thus a regional feature in the Jurassic sequence from the Black Hills westward through the Uinta Mountains, and southward through the Colorado Plateau at least to the Zuni Mountains, N. Mex.

Late Cretaceous uplift and erosion in San Juan region, Colorado

An unconformity marking a period of uplift and erosion has been mapped by R. G. Dickinson within a sequence of Upper Cretaceous sedimentary and volcanic rocks in the Cimarron Ridge area, along the northwestern flank of the San Juan Mountains, Colo. The unconformity bevels Upper Cretaceous marine and nonmarine rocks and is overlain by newly discovered volcanic sediments dated as Late Cretaceous by plant fossils. After the Mancos sea had withdrawn (in late Campanian time) and at least 800 feet of conformable nonmarine sediments had been deposited, the area was uplifted and deformed. Erosion removed several thousand feet of rock from the uplifted area and cut a surface of low relief before volcanism began near the end of Cretaceous time and deposited an undetermined thickness of volcanic debris.

Stratigraphic relations between Morrison Formation and Dakota Sandstone

An unconformity separates the Morrison Formation and Dakota Sandstone throughout the North Park area, Colorado, but probably is absent farther south, in western Middle Park, according to W. J. Hail. As observed in the Lake Agnes and Gore Pass quadrangles south of Rabbit Ears and Muddy Passes, the contact appears to be conformable; claystones

typical of the Morrison Formation intertongue with sandy strata typical of the Dakota Sandstone.

Ilse fault in Wet Mountains, Colo., extended

Geologic reconnaissance by Q. D. Singewald in the northern Wet Mountains of Colorado extended the Ilse fault 6 miles northward from its previously known northern limit near Grape Creek into a previously known fault between Precambrian and Mesozoic rocks near Parkdale. The Ilse fault, long inferred to be the leading dislocation in the central part of the Wet Mountains, has Precambrian rocks on both walls along the segment hitherto mapped; hence neither the direction nor amount of displacement could be determined. The new data reveal that the east wall of the Ilse fault moved up relative to the west wall; the amount of vertical displacement remains indeterminate, but was of the magnitude of hundreds, if not thousands of feet.

Elk Mountain thrust fault south of Aspen, Colo.

In the Conundrum Creek valley 9 miles south of Aspen, Colo., Bruce Bryant has mapped a structurally complex block of hornfelsed sedimentary rocks underlying a much less deformed section of the Maroon Formation of Pennsylvanian and Permian age. On the southwest the complex structures in the lower rocks are truncated by a fault occupied by a sill, whereas on the north they pass upward transitionally into simpler structures of the Maroon Formation. The complexly deformed rocks below may represent a window beneath the Elk Range thrust fault, which crops out 7 miles to the southwest, and extends northwest-southeast for at least 25 miles. After faulting, Tertiary granodiorite invaded the fault zone and metamorphosed the rocks above and below.

Monoclines in southeastern Colorado

Reconnaissance studies by G. R. Scott have shown that the generally flat-lying strata in the La Junta 1° by 2° quadrangle in the Plains in southeastern Colorado are disturbed by a number of monoclinical folds of diverse trends. These folds range in length from 2 to 40 miles, and many have a sinuous pattern. Commonly the folds are paired, forming structural terraces. The monoclines are believed to be surface manifestations of deep-seated faults, although a laccolithic origin cannot be ruled out.

GEOLOGY OF VOLCANIC AND HYPABYSSAL INTRUSIVE ROCKS

New volcanic center in Wet Mountains, Colo.

Fieldwork by Q. D. Singewald in the Wet Mountains in Custer County, Colo. has disclosed a

previously unreported intrusive body more than a square mile in area, mainly in sec. 19, T. 23 S., R. 69 W. The intrusive opens upward and eastward into andesite that resembles Tertiary extrusive rocks farther south and west in the Wet Mountains, and thus may be one of the feeders for the volcanic material. A belt of the igneous rocks about 2,000 feet wide and 5,000 feet long is hydrothermally altered, impregnated by pyrite, and transected by pyritic veins containing small quantities of gold.

Age of volcanic activity in central Colorado

A black vitrophyre at the base of a thick rhyolitic welded tuff mapped by R. E. Van Alstine in the Poncha Springs quadrangle, central Colorado, was dated by the K-Ar method (analysts H. H. Thomas, R. F. Marvin, Paul Elmore, and H. Smith). An age of 34 million years (± 3 million years) was obtained both from a biotite separate and from the glass matrix, and the volcanic rocks are considered to have been erupted in the Oligocene. The welded tuff is more widespread than was previously realized and may correlate with similar rocks in southwestern South Park, dated by other workers as early Oligocene on the basis of vertebrate and invertebrate fossils in intertonguing sediments.

Dike systems in southern Colorado

Reconnaissance mapping by R. B. Johnson in the Trinidad 1° by 2° quadrangle in southern Colorado has shown that many dikes cut the sedimentary rocks of Cretaceous age east and northeast of the Spanish Peaks. These dikes belong to a large system of subparallel dikes that cuts the dikes of the Spanish Peaks radial system and trends normal to the strike of the folded sedimentary rocks of the Raton Basin. In contrast to the relatively siliceous porphyry dikes of the Spanish Peaks system, the younger dikes are mainly syenitic lamprophyres (minettes and vogesites) and several are more calcic lamprophyres and diabases. Some dikes of diabase near the New Mexico–Colorado boundary may be related to the Pleistocene and Recent flows, plugs, and volcanoes that make up the Raton Mesas and associated volcanic terrane.

GEOPHYSICAL INVESTIGATIONS

Gravity anomalies in San Luis Valley, Colo.

A series of echelon gravity highs mapped by J. R. Gava and D. E. Karig in the central and western San Luis Valley, Colo., have been interpreted as expressions of horsts of Precambrian rock covered by basin fill. Subparallel gravity lows in the eastern

part of the valley are believed to reflect a graben filled with perhaps as much as 30,000 feet of sedimentary and volcanic rocks. A gravity low over volcanic rocks in the Bonanza mining district, Colorado, at the northeast corner of the San Juan volcanic field was interpreted by Karig as a collapse caldera (p. B9–B12). The caldera is elliptical in outline with axes of 8 and 10 miles, and is filled with about 8,000 feet of low-density material.

Gravity survey of North Park, Colo.

Preliminary results of a gravity survey of the North Park basin in north-central Colorado by J. C. Behrendt disclosed gravity minimums of about -270 milligals over the basin, compared with about -215 mgal over the adjacent mountains. This suggests thicknesses of relatively low-density sedimentary rocks of about 4 kilometers within the basins.

QUATERNARY GEOLOGY

Pleistocene and Recent surfaces near Badger Creek, Colo.

Three pediment surfaces and a terrace have been recognized by C. T. Wrucke in the Badger Creek drainage in the Cameron Mountain quadrangle, central Colorado. The oldest and best developed pediment lies about 280 feet above Badger Creek and truncates Precambrian granite and Tertiary conglomerate; the intermediate and lower pediments, respectively 40 to 60 and 10 to 30 feet above the streams, bevel Tertiary tuff and conglomerate. The terrace is 2 to 10 feet above the streams. A strongly developed soil containing a reddish-brown B horizon formed on gravel resting on the highest surface indicates a pre-Wisconsin age for that pediment. The geomorphic position of the lower pediments suggests that they are of Wisconsin age. The terrace is probably of Recent age.

Glacial spillover through McClure Pass, Colo.

Glacial deposits along Colorado Highway 133 south of Glenwood Springs contain boulders derived from the Treasure Mountain dome, in the Crystal River drainage, as indicated by studies of L. H. Godwin. The boulders, at McClure Pass and westward down Lee Creek, indicate that ice from glaciers in the Crystal River canyon spilled through the pass more than twice. The spillovers are assigned to glaciations in the middle or early part of the Pleistocene sequence, when the Crystal River canyon was much shallower than now. As the canyon deepened, later glaciers were confined with it, and at least three existed since the last ice flowed through McClure Pass.

Pleistocene sequence near Leadville, Colo.

Studies by Ogden Tweto of samples from test holes drilled by the Bureau of Reclamation in the upper Arkansas Valley west of Leadville, Colo., show that the large, sharply defined horseshoe-shaped end moraine at Lake Fork is built up of as many as four different tills derived from the Sawatch Range to the west. These tills are each less than 50 to 150 feet thick and are separated by weathered zones. These end-moraine tills, which extend to depths of as much as 175 feet below the present valley floor, rest on a sequence of deposits of much greater areal extent derived wholly or in large part from the Mosquito Range to the east. From the top downward, this lower sequence consists of (1) Malta Gravel, generally less than 50 feet thick; (2) ancient glacial drift, with or without a heavy red gumbo zone at its top, depending on accidents of erosion or preservation; (3) local deposits of valley alluvium, some of which contain thin layers of volcanic ash, and bouldery channel deposits; and (4) the Pliocene Dry Union Formation. The stratigraphic relations established by superposition in the end-moraine and subjacent sequence confirm a Pleistocene history previously deduced indirectly from geographic and topographic relations at the surface.⁴³

Quaternary stratigraphy in Durango, Colo. area revised

G. M. Richmond (p. C137-C143) has restudied the type deposits of Quaternary stratigraphic units in the Durango area, San Juan Mountains, Colo., and has correlated them with the standard sequence of glacial events for the Rocky Mountain area as established in the Wind River Mountains of Wyoming. Moraines formerly called Wisconsin are of Pinedale age; some moraines formerly assigned to the Durango Glaciation are of Bull Lake age, but the type Durango moraine represents the earlier Sacajawea Ridge Glaciation. The Florida Gravel is outwash of Sacajawea Ridge age, and the Oxford Gravel is reworked colluvium derived from fluvio-glacial gravels of pre-Sacajawea Ridge age. The Bridgetimber Gravel has the characteristics of fluvio-glacial deposits, and probably was deposited during two pre-Sacajawea Ridge glaciations. Some deposits called Bayfield Gravel may correlate with the Bridgetimber Gravel, but others may be residual deposits weathered from underlying conglomerate beds in the Animas Formation.

⁴³ Ogden Tweto, 1961, Late Cenozoic events of the Leadville district and upper Arkansas Valley, Colorado: Art. 56 in U.S. Geol. Survey Prof. Paper 424-B, p. B133-B135.

COLORADO PLATEAU**GEOLOGIC STUDIES****Multiple periods of Precambrian deformation in Uncompahgre uplift**

In the continuing investigation of the Precambrian crystalline rocks comprising the core of the northwestern Uncompahgre uplift, Utah and Colorado, J. E. Case has recognized two ages of folding in the ancient feldspathic and biotitic gneisses and amphibolites. In the earlier deformation, folds formed around axes that trend approximately north. In the later deformation, the early folds were warped around axes that trend from east to northeast. A set of folds that trends northwest has not yet been related to either period of deformation.

Surface expression of basement structure in Paradox basin

Field evidence and subsurface studies of drill data lead R. J. Hite to suggest that a series of northeast-trending structures extend through and beyond the margins of the Paradox salt basin in southeastern Utah and southwestern Colorado. These structures are expressed at the surface by small faults, fractures, bands of mineralized and altered rock adjacent to fissures, and by zones of drag folds. From their geometry, they are believed to overlie basement wrench faults, at least some of which had large left-lateral displacement.

Northwest transport of coarse components of Moenkopi Formation

The coarse fraction of the Triassic sediments now comprising the Moenkopi Formation moved generally northwestward across the area of the Colorado Plateau according to R. A. Cadigan. The sediments were derived from crystalline rocks in areas inferred to have been rising slowly in southwestern Colorado, southwestern New Mexico, and southern Arizona. Each of the areas included local sources of tuff. This resultant direction of sediment transport is suggested by trend-surface analyses (computed linear surfaces) of statistical measures of grain-size distribution of many widely scattered representative samples. Median grain size, skewness, and kurtosis decrease in a northwest direction across the Colorado Plateau, and sorting improves.

Nonmarine fossils in Straight Cliffs and Wahweap Sandstone

The Tropic Shale and the overlying marine sandstone in the lower part of the Straight Cliffs Sandstone appear to record the last incursion and

withdrawal of the Late Cretaceous sea in the Nipple Buttes area of the Kaiparowits Plateau, Utah. Fossils collected by Fred Peterson, H. A. Waldrop, G. W. Horton, and R. L. Sutton from the lower part of the Straight Cliffs contain marine fossils diagnostic of the *Collignonicerias hyatti* faunal zone, according to W. A. Cobban. Overlying rocks in the middle and upper part of the Straight Cliffs Sandstone and in the shaly lower part of the overlying Wahweap Sandstone contain small but varied faunal assemblages including gastropods, pelecypods, ostracodes, and fish and reptile tooth and bone fragments. These fossils identified by Cobban, I. G. Sohn, and G. E. Lewis suggest nonmarine environments of deposition from the top of the lower part of the Straight Cliffs upward.

Key fossil zones in Lewis Shale

Key fossil zones of Late Cretaceous (Montana) age have been recognized in the Lewis Shale on the eastern side of the San Juan basin in northern New Mexico. At least 6 ammonite zones that are known in eastern Colorado and the northern Great Plains were identified by W. A. Cobban in fossils collected by Cobban, C. H. Dane, and E. R. Landis.

Southeast margin of Eocene Lake Uinta

The southeastern limit of oil-shale deposition in western Colorado has been identified by J. R. Donnell near the east end of Grand Mesa, Delta County. Well exposed at Mount Darlene are two 1-foot beds of lean oil shale separated by hundreds of feet of sandstone and siltstone. These two thin beds are the wedging edge of the 2,000 feet of continuous oil shale that was deposited in the center of the Piceance Creek lobe of Eocene Lake Uinta about 60 miles to the northwest.

Volcanic stratigraphy in High Plateaus

In the Awapa and Fish Lake Plateaus and adjacent Grass Valley, Sevier and Wayne Counties, Utah, P. L. Williams and R. J. Hackman have established details of volcanic stratigraphy which permit correlation of units between several of the High Plateaus and the Marysvale area to the west. The units are the Bullion Canyon Volcanics of Miocene (?) age, basaltic andesite erupted from three source areas, the Dry Hollow Formation of Pliocene (?) age, and surficial olivine basalt erupted from many vents on the Awapa Plateau. Some of the basalt is faulted and of preglacial age; some is apparently younger than the Bull Lake Glaciation.

Type Cerro Till found to be landslide material

The type Cerro Till is on Cerro Summit about 15

miles east of Montrose, Colo. Study by R. G. Dickinson (p. C147–C151) of the hummocky and bouldery surficial debris composing the deposit indicates that the material was emplaced primarily by landsliding, instead of by a piedmont glacier. The names Cerro Till and Cerro Glaciation are therefore abandoned by the Geological Survey.

Popularization of geology at Capitol Reef

Following publication of the final report of investigations in the Capitol Reef area, in south-central Utah near the west edge of the Colorado Plateau, by J. F. Smith and others,⁴⁴ Smith's help was requested by the National Park Service in identifying and describing rocks in the Capitol Reef National Monument. Smith also has served as geologic consultant in preparation of a short movie to be shown at the Monument. The movie, prepared by a private contractor, shows development of the Water-pocket fold and the present scenery. Included were diagrams and sketches by J. D. Stacy, scientific illustrator.

1:250,000-scale geologic map of Moab quadrangle, Colorado-Utah

The geology, structure, and uranium deposits of the Moab quadrangle, Colorado and Utah (P. L. Williams, 1–64), are shown on a new map of an area comprising 2° of longitude and 1° of latitude. Among other features the area includes the uranium deposits of the Uravan mineral belt and of Lisbon Valley, and also the oil field of the Lisbon Valley anticline, the La Sal group of laccolithic mountains, and the salt anticlines of the Paradox basin. The map, in two sheets, was compiled from more than 70 sources and is the second of a series of 10 planned to cover much of the Colorado Plateau.

GEOPHYSICAL STUDIES

Basement structure and salt thickness, Moab-Needles area, Utah

H. R. Joesting, J. E. Case, and Donald Plouff have completed interpretation of gravity and aeromagnetic surveys of the Moab-Needles area, Utah, which covers a large part of the new Canyon Lands National Park. In the northern third of the area, which lies in the deeper part of the Paradox basin, magnetic response is uniform, suggesting uniform Precambrian basement rocks. Pronounced gravity lows are associated with Moab Valley and Salt

⁴⁴ J. F. Smith, Jr., L. C. Huff, E. N. Hinrichs, and R. G. Luedke, 1963, Geology of the Capitol Reef area, Wayne and Garfield Counties, Utah: U.S. Geol. Survey Prof. Paper 363, 102 p.

Valley salt anticlines in the northeast corner of the area. From analysis of the gravity anomaly, the amplitude of the salt core of Moab Valley anticline is about 8,000 feet. In the southern third of the area, at the north end of Monument upwarp, the magnetic field is diverse, indicating a corresponding diversity of the basement rocks. In the central part of the area, anomalies are arranged systematically. A zone of steep gravity gradient trends eastward across the area and marks a major lithologic and structural boundary in the basement rocks. Just south of and approximately parallel to this zone are three prominent magnetic highs, each associated with a gravity high. These highs evidently are caused by large intrusive masses in the basement.

BASIN AND RANGE REGION

STRATIGRAPHY AND STRUCTURAL GEOLOGY

NEVADA AND EASTERN CALIFORNIA

Changes in stratigraphy across major faults in southern Great Basin

Stratigraphic studies of upper Precambrian and Lower Cambrian formations in the southern Great Basin by J. H. Stewart indicate abrupt changes in thickness and facies on opposite sides of the Death Valley and Furnace Creek fault complex in Inyo County, Calif., and across the Las Vegas shear zone in Clark County, Nev. The thicknesses of the Stirling Quartzite, Wood Canyon Formation, and Zabriskie Quartzite are approximately doubled across the Death Valley and Furnace Creek fault zone from the Panamint Range on the west side of Death Valley to the Funeral and Grapevine Mountains on the east side. Similar changes in these formations also were observed across the Las Vegas shear zone, although the relations there are less clear. Right-lateral strike-slip movement is suggested by the thickness and facies trends. Offsets may be about 50 miles on the Death Valley and Furnace Creek fault complex and about 30 miles on the Las Vegas shear zone.

Thrust fault in east-central California

A thrust fault which may have considerable regional significance has been recognized by J. H. Stewart, C. A. Nelson, and D. C. Ross in the Last Chance Range, Saline Range, and northern Inyo Mountains of California. The fault emplaced rocks of Precambrian and Early Cambrian age on rocks largely of Mississippian and younger age.

Imbricate thrust faults in Egan Range, Nev.

Mapping by A. L. Brokaw in the Reipetown quadrangle, Nevada, reveals imbricate thrust faults in rocks of Late Devonian to Pennsylvanian age. The rapid thinning of the Joana Limestone in the Robinson (Ely) district, postulated by earlier workers to be a depositional feature, is now believed to be due to tectonic slicing. Moreover, massive limestone considered by earlier workers to be part of the Chainman Shale has been recognized as a thrust segment of the lower part of the Joana Limestone. The structures mapped substantiate earlier conclusions, based on inadequate exposures, that significant faulting occurred in equivalent rocks throughout this part of the Egan Range.

Faults in Ward mining district, eastern Nevada

In the Ward mining district, south of Ely, Nev., mapping by T. L. Heidrick, Jr. and A. L. Brokaw shows that the productive part of the district is related to small monzonitic dikes and mineralized faults and fissures that cut beds in the lower part of the Ely Limestone. The area is a complexly faulted structural block that has been displaced upwards about 1,000 feet, relative to adjacent blocks, along a system of transverse bounding faults. The stratigraphic and structural studies indicate that exploration for deeper potentially favorable host rocks within the displaced block is economically feasible. Exploration targets outside the uplifted block are probably too deep to be economically attractive.

Zonation of Silurian and Devonian rocks by rugose corals

A workable zonation of Silurian and Devonian carbonate rocks in the Great Basin by rugose corals appears practicable, as the collections from areas of detailed geologic mapping are sectioned and studied, according to C. W. Merriam. In central Nevada the Lone Mountain Dolomite of Silurian age contains extensive carbonaceous dolomite bodies in its upper part which bear a coral fauna with bushy *Entellophyllum*. The overlying Devonian Nevada Formation includes seven rugose coral zones which are lettered A to G in ascending order. Zones A, B, and C are Lower Devonian, being characterized by Laccophyllidae, Halliidae, and Streptelasmatidae. Colonial rugose corals do not appear in great abundance until the higher Middle Devonian, as in zone F, which contains such genera as *Hexagonaria*, *Sociophyllum*, *Utaratuia*, and *Aphroidophyllum*. Solitary Digonophylloids and Acanthophylloids also occur in these beds which immediately underlie the *Stringocephalus* zone.

Thrusting in northern Nye County, Nev.

Additional mapping of Paleozoic rocks in the Hot Creek Range, central Nevada, by F. J. Kleinhampl, indicates that transitional facies Devonian rocks were thrust over miogeosynclinal carbonate rocks during the Antler orogeny in Late Devonian or earliest Mississippian time. Although rocks of Early Mississippian age are also involved in thrusting, they probably were deposited as an orogenic sediment overlapping autochthonous Devonian rocks and the earlier thrust plate, and subsequently were thrust in late Paleozoic or Mesozoic time.

UTAH

Extension of Mount Raymond thrust near Salt Lake City

M. D. Crittenden, Jr., has completed a reconnaissance study of a possible eastward extension of the Mount Raymond thrust zone east of Salt Lake City, Utah. This fault is a folded thrust which crosses the Wasatch Range from the mouth of Mill Creek near Salt Lake City to a point north of Park City, Utah, where it disappears beneath a cover of alluvium and volcanic rocks. Mapping near Peoa on the Weber River shows that the Crandall Canyon and Dry Canyon faults, previously described as high-angle normal faults, are accompanied by large-scale overturning and by changes in sedimentary facies that are indicative of thrusting. Inasmuch as earlier mapping by students at the University of Utah (N. C. Williams and J. H. Madsen⁴⁵) showed that these faults were overlapped by the Wanship Formation of Williams and Madsen (1959), the movement on the Mount Raymond thrust can be dated as early Late Cretaceous in age.

ARIZONA

Trend of Patagonia Mountains related to Late Cretaceous or early Tertiary faults

In southeastern Arizona, F. S. Simons has found that the northwest trend of the Patagonia Mountains probably was developed in Late Cretaceous or early Tertiary time by large faults that since have been obliterated by igneous intrusion. No sizable faults of this trend are now recognized along the west front of the mountains. Moreover, rocks like the oldest found within the range crop out on the broad alluvial plain as far as 3 miles west of the mountain front. Thus the present topography apparently is not related to basin-range block faulting in late Cenozoic time.

⁴⁵ N. C. Williams, and J. H. Madsen, Jr., 1959, Late Cretaceous stratigraphy of the Coalville area, Utah, in Intermountain Assoc. Petroleum Geologists Guidebook 10th Ann. Field Conf.: p. 122-125.

Abrigo Formation studied in southeastern Arizona

Stratigraphic studies of the Abrigo Formation in southeastern Arizona by M. H. Krieger reveal that the three members of the Abrigo have widespread distribution and lithologic continuity. The lower member, of late Middle Cambrian age, consists of mudstone, shale, fine-grained sandstone, and variable amounts of limestone and dolomite. The middle member is chiefly sandstone in the Holy Joe Peak and Christmas quadrangles, but to the south it is chiefly limestone or dolomite. The upper member consists of sandy dolomite and dolomitic sandstone in the lower part and thin-bedded dolomite in the upper part. The brachiopod *Billingsella*, which defines a narrow time zone in the middle Late Cambrian, has a widespread occurrence in the thin-bedded dolomite of the upper member.

EMPLACEMENT AND AGE OF INTRUSIVE ROCKS

Age of plutons in Elko County, Nev., and vicinity

In northeastern Nevada and adjacent parts of Idaho and Utah, scattered, relatively small plutons of silicic rocks intrude rocks as young as late Paleozoic, and are overlain by rocks as old as middle Tertiary. R. R. Coats, R. F. Marvin, and T. W. Stern report (p. D11-D15) that K-Ar ages of these rocks fall into three groups: Middle Jurassic, Late Cretaceous, and late Eocene to early Oligocene. Biotite-zircon pairs in samples from nine plutons have been dated by K-Ar and Pb-alpha methods. Results by the two methods are in close agreement in all except two plutons. No petrographic criteria have been found that may be correlated with the age assignments.

Age of intrusive and extrusive rocks in southeastern Arizona

S. C. Creasey reports that about 50 K-Ar ages have been determined from intrusive and extrusive rocks in southeastern Arizona thought to be either of Late Cretaceous to early Tertiary age (Laramide) or Tertiary age. The age determinations show a bimodal distribution in time. The older group, which includes predominantly intrusives but some extrusives, ranges in age from 50 to 75 million years. Most of the stocks associated with the porphyry copper deposits are within this group. The younger group, which is made up mainly of extrusive rocks, ranges in age from 20 to 30 million years. No large ore deposits are known to be associated with rocks of this younger group. Virtually no igneous activity

occurred during the period from 30 to 50 million years. These data, although incomplete, tend to confirm the concept that most of the economic copper deposits of southeastern Arizona are associated with stocks that were intruded during a restricted period of time.

Granodiorite stocks in central Yavapai County, Ariz.

In the northern part of the Mount Union quadrangle, Arizona, C. A. Anderson has found three granodiorite stocks of Cretaceous to early Tertiary age. These stocks crop out in an east-trending belt, and are aligned with a similar stock in the Copper basin to the west, which is mineralized with copper and molybdenum. Altered facies of the three stocks may be good targets for exploration.

Mesozoic plutons in south-central Mojave Desert, Calif.

In the Ord Mountains and Rodman Mountains quadrangles, California, T. W. Dibblee, Jr. (2-64, 3-64) has found at least two major granitic plutons of Mesozoic age. Both are intrusive into Precambrian gneiss and Paleozoic metasedimentary rocks. The older pluton contains deposits of iron ore along pendant margins of Paleozoic metasedimentary rocks; it is intruded by andesitic porphyry that locally contains deposits of copper, gold, tungsten, and molybdenum. These rocks are invaded by the younger pluton of Cretaceous age, which is unaltered and nearly devoid of metallic mineral deposits.

CENOZOIC VOLCANISM

Source of some exotic blocks in Keetley Volcanics, northern Utah

Mapping by M. D. Crittenden, Jr., near the center of the Keetley volcanic field, between Park City and Kamas, Utah, reveals what appears to be the top of a shallow porphyritic intrusive bounded by rectilinear faults and cutting both basement rocks of Triassic and Jurassic age and bedded andesitic volcanic rocks. The relations suggest that the faulted roof of an igneous mass intruded close to the base of the volcanic rocks, was lifted like a trap door, and then rafted outward in accompanying large-scale volcanic eruptions. Such sites are almost the only possible source for numerous building-size blocks of Nugget Sandstone and quartzite of the Gartra Grit Member of the Ankareh Formation that occur in the Keetley Volcanics between Park City and Wanship. It also suggests that the source of the Keetley Volcanics is in part buried beneath the volcanic field, and not somewhere outside it.

Possible source of rhyolitic lava flows and ash-flow tuffs in west-central Utah

Detailed mapping by H. T. Morris in the northwestern part of the Cherry Creek quadrangle, Utah, has disclosed a vitrophyre dike of rhyolitic composition nearly half a mile wide and more than 2½ miles long that was not recognized prior to the current studies in the area. This dike may be the long-sought eruptive center of extensive flows of rhyolitic lavas and ash-flow tuffs in this part of west-central Utah.

Volcanic units in Esmeralda County, Nev.

Recent studies of Tertiary volcanic rocks in Esmeralda County, Nev., by J. P. Albers show that many of the ash-flow tuffs and flow rocks have the composition of quartz latite, and thus are similar in composition to the plutonic rocks, which are dominantly quartz monzonite. In the Goldfield mining district a number of the volcanic units, including the Vindicator Rhyolite, Kendall Tuff, Morena Rhyolite, dacite vitrophyre, and Meda Rhyolite are now believed to be ash-flow tuffs instead of lava flows. The Meda Rhyolite and the dacite vitrophyre are considered to be parts of the same unit. The Sandstorm Rhyolite, as mapped by Ransome prior to 1909, includes rhyolite flows, ash-flow tuff, and shale.

GEOCHEMICAL EXPLORATION

Geochemical anomalies in northeastern Nevada

R. L. Erickson reports that reconnaissance geochemical sampling in the Pequop Mountains and Wood Hills in northeastern Nevada resulted in discovery of anomalously high metal values in a region generally considered unfavorable for prospecting. Anomalous Pb-Zn-Ag was found along a fault zone in dolomite of Silurian (?) age, and anomalous W-As-Sb probably originated in marble and schist of lower to middle Paleozoic age. Small pods of barite also were discovered at several localities in the range. Mercury, associated with the Pb-Zn-Ag anomaly, appears to increase in concentration vertically. The Guilmette Formation of Devonian age contains more Hg than the underlying base-metal-bearing Silurian dolomite. Thus, strong Hg anomalies with minor amounts of Pb-Zn-Ag in the Guilmette Formation may indicate concealed Pb-Zn-Ag deposits in the underlying Silurian dolomite.

Sampling of bedrock and soil in Stockton district, Utah

R. J. Roberts, E. W. Tooker, and W. J. Moore have collected approximately 1,500 bedrock and soil samples in an area of 4 square miles in the Stockton

mining district, Utah. One objective of the study is to establish the distribution of minor elements associated with lead-silver mineralization. The ore bodies of lead and silver are localized at intersections between northwest-trending breccia zones and thin argillaceous limestone units. A compilation of 400 analyses of bedrock from the northern half of the area shows zones of Cu-Mo, Ag-As, and Pb-Zn extending outward from several major fissures. The heavy-metal concentrations of 75 soil samples in the same area are uniformly low, and there is practically no correspondence with the metal zones established from bedrock samples. Both the Stockton deposits and the Bingham copper deposits, 12 miles northeast, occur in the predominantly clastic Bingham sequence of the Oquirrh Formation, as defined by Tooker and Roberts.⁴⁶

GEOFYSICAL STUDIES

Gravity survey in Carson Sink area, Nevada

Gravity anomalies in the Carson Sink area near Fallon, Nev., indicate that the subsurface bedrock topography is more irregular than the surface topography would suggest, according to R. R. Wahl. Gravity highs in the mountains to the north, west, and southwest of Carson Sink suggest an arcuate structure that could form a transition from northeast-trending anomalies in the northeastern two-thirds of the area to northwest-trending anomalies in the southwestern third of the area. Three gravity lows south of the West Humboldt Range and one gravity low in the Humboldt Sink have closures greater than 15 milligals. The indicated thickness of the valley fill at these localities ranges from 5,000 to 10,000 feet.

Geophysical studies in Safford Valley, Ariz.

Aeromagnetic and gravity studies by G. P. Eaton and G. E. Andreasen in Safford Valley, Ariz. show that the valley fill has a maximum thickness of 3,000 feet, but that the central axis of the valley lies several miles southwest of the present course of the Gila River. The data indicate that the valley is underlain by rocks with properties like those of the Precambrian gneisses and the Upper Cretaceous to lower Tertiary intrusive rocks exposed in the mountains to the southwest. Measurements of bulk density and grain density made on cores from the upper third of the valley fill indicate an average total porosity of 30 percent.

⁴⁶ E. W. Tooker, and R. J. Roberts, 1963, Comparison of Oquirrh Formation sections in the northern and central Oquirrh Mountains, Utah: Art. 188 in U.S. Geol. Survey Prof. Paper 450-E, p. E32-E36.

COLUMBIA PLATEAU AND SNAKE RIVER PLAIN

OREGON

Basalt extruded along dike swarm

The Picture Gorge Basalt of the Columbia River Group in the Aldrich Mountain quadrangle, Oregon, was extruded along a northwest-trending dike swarm according to T. P. Thayer and C. E. Brown. They have found that the basalt was as much as 6,000 feet thick near the dike swarm, but thinned to 1,500 feet to the west and 2,500 feet to the east. During the extrusion an elongate shield-type domical ridge accumulated to a height of 3,500–4,500 feet above the level of the surrounding plain and was flanked by the accumulating fluvial volcanic debris of the Mascall Formation of late Miocene age. The upper flows of the Picture Gorge Basalt probably intertongued with the sediments of the Mascall Formation, which must have exceeded 4,500 feet in thickness in the Picture Gorge area, about 10 miles to the west.

IDAHO

Buried topographic features interpreted from geophysical data

Evidence for two filled sedimentary depressions and two buried ridges in the eastern Snake River Plain, in the vicinity of the National Reactor Testing Station, Idaho, has been obtained by D. R. Mabey using gravity and aeromagnetic data collected in 1964 in cooperation with the U.S. Atomic Energy Commission. One of these features reported previously (U.S. Geol. Survey, 11-64, p. 109) is a depression filled with a thick sequence of sedimentary and volcanic rocks that is inferred to extend around the east and south sides of the Arco Hills. Another depression extending southwest from Mud Lake is interpreted by G. H. Chase to be a graben. An uplift of pre-Tertiary rocks is inferred to extend east and southeast from Big Southern Butte. A buried extension of the Beaverhead Mountains is suggested in the area of Circular Butte. Although a large amount of relief on the surface of the pre-Tertiary rock beneath the plain is indicated, there is no implication that the basin-range structure in adjoining regions is continuous across the plain.

D. A. Morris reports that the contours of a persistent zone of high permeability that crosses lithologic types appears to reflect the general alignment of the Mud Lake structure, and suggests control by secondary structures such as joint and fracture systems.

PACIFIC COAST REGION

WASHINGTON

Stratigraphy of Anarchist Series of north-central Okanogan Highlands

Four mappable units have been defined by C. D. Rinehart and K. F. Fox in the metamorphic rocks of the Permian Anarchist Series of Daly (1912)⁴⁷ in the Loomis area. Although the rocks are metamorphosed in the greenschist facies and are much folded and faulted, they retain enough sedimentary structures to permit determination of order of superposition. From oldest to youngest the units are: (1) interbedded phyllite, slate, and minor limestone; (2) limestone interbedded with substantial amounts of graywacke, slate, and conglomerate, and containing poorly preserved fossils locally near the top; (3) slate and graywacke with prominent and distinctive chert conglomerate; and (4) interbedded greenstone and quartzite. Units 1 and 4 are in general equal to divisions proposed by Waters and Krauskopf.⁴⁸ All units are apparently conformable although varied in thickness; aggregate thickness is more than 15,000 feet.

Permian brachiopods in Anarchist Series of northern Washington

A small suite of brachiopods from the Anarchist Series of Daly (1912),⁴⁷ Okanogan County, Wash., collected by C. D. Rinehart, includes several Permian forms. Genera identified by J. T. Dutro, Jr., are *Megousia*, *Yakovlevia* (*Muirwoodia* of earlier usage), *Spiriferina*, and *Rhynchopora*. Poorly preserved pelecypods, gastropods (including *Euphemites*?) and bryozoans complete the fossil assemblage. This collection confirms an earlier tentative Permian assignment by G. H. Girty, based on collections of A. C. Waters.⁴⁸ Clearly, Daly's Anarchist Series is at least partly equivalent to the Cache Creek Group of British Columbia.

Upside-down syncline in metamorphic rocks of North Cascades

The discovery of relict graded bedding and flame structure in southwest-dipping schists and gneisses near Holden, by F. W. Cater and C. A. Hopson, has made it possible to show that a large tight antiform on strike to the southeast (Lucerne quadrangle) is really an upside-down syncline.

⁴⁷ R. A. Daly, 1912, Geology of the North American Cordillera at the forty-ninth parallel: Canada Geol. Survey Dept. Mines Mem. 38, [1915?], Maps 10-12.

⁴⁸ A. C. Waters and K. B. Krauskopf, 1941, Protoclastic border of the Colville batholith: Geol. Soc. America Bull., v. 52, p. 1355-1418.

Age of Vashon Stade and origin of Puget Sound trough in Washington

Two recent papers, one by D. R. Crandell, D. R. Mullineaux, and H. H. Waldron (p. B132-B136) and the other by Mullineaux, Waldron, and Meyer Rubin (4-65) describe results of Pleistocene stratigraphic studies in the Puget Sound lowland, interpreted in the light of radiocarbon age determinations on a variety of materials. A sequence of late interglacial and early Vashon glacial deposits in Seattle indicates that the Puget glacier lobe of Vashon age did not occupy the Seattle area until after 15,000 years ago, several thousand years later than previously believed; the ice retreated from the area by about 13,500 years ago. Distribution of the deposits on both sides of Puget Sound indicates that a deep Puget Sound trough did not exist before the Puget lobe arrived; thus, the trough was eroded wholly by the glacier rather than modified by it from a preexisting valley.

Major tectonic break in northeastern Olympic Mountains

In the Mount Angeles quadrangle, R. W. Tabor and W. C. Cady have found the first evidence of a tectonic break separating lower (?) to middle Eocene mafic volcanic and basaltic clastic sedimentary rocks of the Crescent Formation (which forms the outer volcanic ring of the Olympic Mountains) from the generally slaty core rocks. These core rocks have been referred to, in part, as the Paleocene(?) or lower Eocene Soleduck Formation of Reagan (1909).⁴⁹ The Crescent Formation strikes west-northwest, dips steeply north, generally has its top to the north but is locally overturned. The core rocks, characterized by thick-bedded micaceous sandstone, strike more northwesterly and are overturned, chiefly to the northeast, except where involved in a large-scale dextral fold. In the core and converging northwestward with the tectonic break is a marker zone characterized by thin, discontinuous, mafic volcanic rocks and bearing an assemblage of mollusks dated as late Eocene(?) by W. O. Addicott. The tectonic break is marked by sheared-out isoclinal folds and appears to cut out about 12,000 feet of core section; further understanding of relative movement along the break depends on more conclusive dating of the core rocks.

Stratigraphy of Tertiary rocks in southwestern Washington

In the Grays River area, E. W. Wolfe and E. H.

⁴⁹ A. B. Reagan, 1909, Some notes on the Olympic Peninsula, Wash.: Kansas Acad. Sci. Trans., v. 22, p. 131-238.

U.S. Geological Survey, 1963, Geological Survey Research 1963, Summary of investigations: U.S. Geol. Survey Prof. Paper 475-A, p. A96.

McKee have mapped a nearly continuous section of marine sedimentary and volcanic rocks ranging in age from middle Eocene to early or middle Miocene and containing possible potential petroleum reservoirs in the upper Eocene and Miocene shallow-water sands. In this section, major volcanic episodes are recorded in the middle Eocene, upper Eocene, and middle(?) Miocene rocks. Heavy-mineral assemblages derived from pre-Tertiary crystalline rocks differ in successive lithologic units and may be useful as stratigraphic markers.

OREGON

Extrusive camptonite in central part of Oregon Coast Range

P. D. Snavely, Jr., has determined that flows and breccias interbedded with upper Eocene marine rocks of the Euchre Mountain quadrangle are of camptonitic composition (barkevikite-augite basalt) and were probably fed by camptonite dikes of the area. Thick sequences of alkalic basalt flows occur in the same stratigraphic position as the camptonite rocks in this part of the Coast Range and suggest that the latter were derived from the same parent magma as the alkalic basalt.

Geology of western Cascade Range

A paper by D. L. Peck and others (2-64) describes Eocene, Oligocene, and Miocene volcanic rocks. The interbedded flows and pyroclastic rocks crop out in an area of about 7,500 square miles, average about 12,000 feet in thickness, and interfinger with marine sedimentary rocks on the west. The volcanic rocks are warped into a series of northeast-trending en echelon folds. Although predominantly basaltic andesite, the rocks range in composition from basalt to rhyodacite, and the origin of sequence can most logically be explained by crystal fractionation of 3 or 4 successive magmas or groups of similar magmas.

CALIFORNIA

KLAMATH MOUNTAINS AND NORTHERN COAST RANGES

Discovery of graptolites in Klamath Mountains

In a recent article, Michael Churkin, Jr. (p. C72-C73), describes graptolites found in shale and siltstone of the Gazelle Formation. These fossils extend the age of the formation downward to include latest Early and Middle Silurian and promise to become a useful stratigraphic tool in this structurally complex area.

Major left-lateral faults in Coast Range in northern California

In the Colyear Springs quadrangle, E. H. Bailey and D. L. Jones have mapped several major faults along the eastern edge of the Klamath Mountain and Coast Range provinces. The northwesterly trending Elder Creek fault marks the northern extent of the Jurassic Knoxville Formation of the Great Valley sequence and has a stratigraphic throw of at least 20,000 feet, probably due to left-lateral movement of even greater magnitude. A parallel fault, 5 miles to the northeast, has similar movement sense and offset of more than a mile. Secondary faults trend northeasterly and have right-lateral separations of comparable magnitude. Renewal of movement on the secondary faults has offset the Pliocene Tehama Formation.

Pennsylvanian metamorphic event in southern Klamath Mountains

As reported in a recent article by M. A. Lanphere and W. P. Irwin (p. D27-D33), potassium-argon ages of hornblende, ranging from 270 to 286, million years, indicate that the Salmon Hornblende Schist was metamorphosed during a hitherto unrecognized Pennsylvanian orogeny rather than during either a pre-Silurian or a Late Jurassic (Nevadan) orogeny. The new age of metamorphism helps explain the paucity of Pennsylvanian sedimentary rocks in the area and indicates that Devonian and Mississippian sedimentary and volcanic rocks to the east, which are chemically similar to the schist, might be the protoliths.

Geologic significance of Franciscan Formation

The Franciscan Formation of the Coast Ranges, long under the scrutiny of California geologists, is described in an exhaustive monograph by E. H. Bailey, W. P. Irwin, and D. L. Jones (3-64). The Franciscan, a eugeosynclinal assemblage of rocks with a known outcrop area of more than 15,000 square miles and an inferred thickness greater than 50,000 feet, is predominantly graywacke with lesser amounts of shale, altered mafic volcanic rocks, chert, minor limestone, and metamorphic rocks; ultramafic rocks are important associates. The age of the Franciscan ranges from Late Jurassic to Late Cretaceous.

The metamorphic rocks are of the zeolite, blueschist, and eclogite facies and crop out both as elongate belts and as isolated tectonic blocks. Pressures for the formation of the blueschists must have exceeded 5 kilobars (greater than 8 kb for the

eclogitic rocks), although temperatures were less than 300°C. Local rapid downwarp and accumulations with equally rapid uplift may account for metamorphism in these facies.

The Franciscan rocks are pervasively folded and faulted; structural trends are generally northwest. In the northern Coast Ranges, major faults and serpentine masses separate the Franciscan from its miogeosynclinal equivalent, the Great Valley sequence, to the east; the latter may be thrust westward over the Franciscan rocks. K-feldspar content of the graywackes of the Great Valley sequence increases with a decrease in age, reflecting progressive unroofing of Late Jurassic to mid-Cretaceous Klamath Mountain and Sierra Nevada batholiths. In the southern Coast Ranges, the Franciscan is exposed both to the east of the San Andreas fault, where it occurs chiefly as piercements along the Diablo antiform, and to the west of the Nacimiento fault.

Two types of regional structural terranes are recognized: one has a crystalline basement like the metamorphic and granitoid rocks of the Sierra Nevada; the other, which includes all the Franciscan rocks, has an unknown basement. However, the abundance of ultramafic rocks in the Franciscan plus the lack of sialic crystalline inclusions in the ultramafic rocks indicates that the Franciscan may have been deposited directly on a basaltic crust or on peridotite.

SOUTHERN COAST RANGES

Stratigraphy of Pancho Rico Formation in Salinas Valley

D. L. Durham and W. O. Addicott described the Pliocene Pancho Rico Formation in a recent report.⁵⁰ The formation comprises sandy marine strata and interbedded siltstone and shale that generally overlie marine Miocene Monterey Shale and underlie the nonmarine Pliocene and Pleistocene(?) Paso Robles Formation in southern Monterey County. The molluscan fauna, consisting principally of species not previously reported from the area, is correlated with the fauna of the Jacalitos Formation, the traditional lower Pliocene standard of the Pacific coast. The nearshore mollusks of the Pancho Rico include many warm-water types, suggesting a marine climate comparable to that off western Baja California. Faunal distribution suggests that a Pliocene seaway connected the Salinas Valley area with the Santa Maria basin to the southeast.

⁵⁰ D. L. Durham, and W. O. Addicott, 1965, Pancho Rico Formation, Salinas Valley, California: U.S. Geol. Survey Prof. Paper 524-A. [In press]

Large fault in southern Salinas Valley

According to D. L. Durham in a recent article (p. D106-D111), the juxtaposition of contrasting units of the upper Miocene and Pliocene Santa Margarita and Pancho Rico Formations brought about by a large fault in the Bradley and Adelaida quadrangles suggests late or post-Pliocene right-lateral displacement of at least 11 miles.

Paleoenvironment of phosphates in Monterey Formation in San Luis Obispo County

Foraminifera of the Monterey Formation at Indian Creek show a marine regression in the middle Miocene (lower and upper Luisian stage of Kleinpell) according to P. J. Smith. The lower 1,000 feet of the formation was deposited at bathyal depths. The water became shallower during deposition of the upper 300 feet at which time an inner continental shelf environment was established, and potentially commercial phosphate was deposited.

New evidence of right-lateral separation along San Andreas fault

According to H. E. Clifton, the distribution, lithologic changes, and structures of basalt flows, and the directional structures of sedimentary rocks, as well as marine-nonmarine facies changes, indicate that there was a west-southwestward regional slope in the vicinity of the southeastern Caliente Range during the middle and late Miocene. The size of granite clasts in a nonmarine pebbly sandstone of probable late Miocene age increases east-northeastward toward the San Andreas fault. Extrapolation of the size increase according to Sternberg's law suggests a source just across the fault in an area now underlain by a thick sequence of middle Miocene deep-water marine sedimentary rocks. The closest granite mass that was exposed during the late Miocene lies 15 miles to the southeast along the fault, indicating at least this amount of right-lateral separation since deposition of the pebbly sandstone.

Décollement structure in Tertiary rocks south of Fox Mountain, Santa Barbara County

Fossil mollusks and corals from graded deep-water marine sedimentary rocks in the Santa Barbara Canyon area indicate that the rocks are middle to late Eocene in age, rather than Cretaceous as hitherto assigned. J. G. Vedder reports that the sequence, which may be more than 15,000 feet thick, forms the north limb of a broad syncline with an axis lying along the north side of Strawberry Peak. South of Fox Mountain and above the homoclinal

Eocene rocks, sedimentary rocks of middle Miocene age chiefly facies within the Monterey Shale are tightly folded. Rocks above and below the contact are jointed, fractured, and contorted, suggesting that the break is a décollement, perhaps related to a low-angle reverse fault dipping northward beneath Fox Mountain.

SIERRA NEVADA

Correlation of aeromagnetic data with major geologic units in central Sierra Nevada

Data from an aeromagnetic survey of a strip across the Sierra Nevada between lat 37° and $37^{\circ} 15'$ N. correlates well with major geologic units according to a preliminary study by H. W. Oliver. Roof pendants of Triassic and Jurassic metavolcanic rocks stand out as magnetic highs of 200–400 gammas, whereas Paleozoic metasedimentary rocks are lows relative to the surrounding granitic rocks. There are a number of examples of association of magnetic anomalies with individual plutons within the batholith, as for example a nearly universal correlation of magnetic lows with various separate plutons of the Cathedral Peak type.

Deformation of Dinkey Creek roof pendant of central Sierra Nevada

Metamorphosed sedimentary rocks in the Dinkey Creek roof pendant exhibit three systems of folds that can be distinguished from one another by their different styles and orientations according to a recent paper by P. C. Bateman and R. W. Kistler. They have shown that minor folds of each system are superimposed on the folds of older sets, indicating that the systems were formed successively and not concurrently. First folds are nearly isoclinal, and before the second deformation their axial surfaces had a strike of about N. 5° E. and a dip of 45° W.; second folds are open and their axial surfaces are vertical and strike N. 15° W.; only locally developed, the third folds also are open, and their axial surfaces are vertical and strike N. 60° W. The folds are continuous across the pendant and are truncated by bounding granitic plutons, indicating that the fold systems were formed before and independently of the emplacement of the plutons. Corresponding fold systems in adjacent pendants and in the south end of the western metamorphic belt suggest that the folds were formed during three periods of regional deformation.

Potassium-argon dating and paleomagnetism of Bishop Tuff

The Bishop Tuff has been redated and studied

paleomagnetically, according to a recent paper by G. B. Dalrymple, Allan Cox, and R. R. Doell (6–65). Duplicate potassium-argon analyses on three samples of sanidine, concentrated from primary pumice fragments from widely separated localities, give ages of 0.68, 0.71, and 0.74 million years, respectively. The latter two determinations are on samples from the basal ash fall, and the age of 0.71 m. y. was on a sample from the basal ash where it overlies Sherwin Till. Paleomagnetic studies on 103 oriented samples from 5 widely separated localities suggest that all the sampled outcrops cooled within several centuries or less. All the samples are normally magnetized, and the directions of magnetization from the more consolidated parts of the unit have less scatter than would be produced by geomagnetic secular variation over periods of time ranging from hundreds to thousands of years. Thus the Bishop Tuff probably erupted during a single igneous event about 0.7 m.y. ago, at a time when the earth's magnetic field was normal.

Complex history of Bishop Tuff

M. F. Sheridan, D. F. Crowder, and R. L. Smith have recognized that the Bishop Tuff consists of one ash-flow cooling unit in the west half of the White Mountain Peak quadrangle; they have mapped several lithologies within the tuff. These lithologies represent separate flows, detritus formed during brief erosional intervals, rocks altered by fumaroles, and zones of welding and vapor-phase alteration.

SAN JOAQUIN VALLEY

Stratigraphy of alluvial deposits in southern San Joaquin Valley

The fresh-water-bearing alluvial deposits of Tertiary and Quaternary age in the southern part of the San Joaquin Valley have been divided into three major units by M. G. Croft. Subsurface geologic studies have revealed that the two upper units are comparatively coarser grained than the lowermost unit. The uppermost unit is a veneer, rarely exceeding 60 feet in thickness. Each unit is further subdivided into: (1) oxidized alluvial-fan deposits and (2) reduced flood-plain and deltaic deposits, both of the Sierra Nevada and Coast Range provenances; and (3) flood-basin, lacustrine, and marsh deposits. The two lower units interfinger with eight well-defined lacustrine clay beds. Radiocarbon ages from the middle unit as determined by Meyer Rubin, one from wood collected 3 feet beneath the youngest lacustrine clay bed and one from wood in the upper part of the same clay bed, are $26,780 \pm 600$ years

and $9,040 \pm 300$ years, respectively. The clay beds probably were deposited in large lakes coeval with glaciation in the Sierra Nevada.

Geophysical studies in San Joaquin Valley

H. W. Oliver suggests that a serpentine body within the granitic basement lies under the Quaternary alluvium near Le Grand; there a magnetic high of about 300 gammas is centered over a gravity low of about 17 milligals. About 10 miles west of Chowchilla, the southern extension of the major magnetic ridge of the Sacramento Valley is expressed with a total relief of about 1,300 gammas and capped with elongate 100–600-gamma anomalies. The extension is coincident with the high gravity axis of the San Joaquin Valley.

Miocene macrofaunal chronology in southeastern San Joaquin Valley

In a recent report, W. O. Addicott (p. C101–C109) described macrofossils representative of the three principal time-stratigraphic divisions of the California Miocene chronology as defined by C. E. Weaver and others.⁵¹ Northeast of Bakersfield, marine mollusks referable to the Vaqueros stage are abundant in the Jewett Silt Member of Diepenbrock (1933)⁵² of the Temblor Formation; the Olcese Sand and the Round Mountain Silt, both of Diepenbrock (1933),⁵² yield the well-preserved Barker's Ranch fauna, the standard for the Temblor stage. In the northern Tejon Hills near Arvin, the Santa Margarita Formation bears mollusks of the Briones, Cierbo, and Neroly stages.

ALASKA

Figure 4 is an index map of Alaska showing the boundaries of the regions referred to in the following summary of scientific and economic findings of recent geologic and geophysical studies.

NORTHERN ALASKA

Triassic marine ostracodes in Alaska

Triassic marine ostracodes from the Shublik Formation or northeastern Alaska have been identified by I. G. Sohn (1–64). These ostracodes are among the first Triassic marine ostracodes to be recognized on the North American continent. They were collected by E. G. Sable in 1948 and C. L.

⁵¹ C. E. Weaver and others, 1944, Correlation of the marine Cenozoic formations of western North America: Geol. Soc. America Bull., v. 55, p. 569–598.

⁵² Alex Diepenbrock, 1933, Mount Poso oil field: California Oil Fields, Div. Oil and Gas, v. 19, no. 2, p. 4–35.

Whittington in 1952 from a locality on Dodo Creek 2 to 2.3 miles above its junction with the Sadlerochit River. The collections contain the following:

Hungarella sp. or spp.

Paracypris? sp. or spp.

Darwinula? sp.

Steinkerns unident.

Geologic study of Romanzof Mountains

E. G. Sable⁵³ has made a comprehensive study of the extreme northeastern part of Alaska, between the Jago and Hulahula Rivers. Rocks of sedimentary and of igneous origins are well represented. Sedimentary and metasedimentary bedrock in the area ranges in age from Middle Devonian(?) to Cretaceous. The sedimentary rocks include carbonate rocks, shale, siltstone, sandstone, and conglomerate; and graywacke, phyllite, argillite, and slate make up the metasedimentary rocks. Unconsolidated deposits include glacial drift representing five advances of the ice, and recent alluvial and colluvial deposits.

Intrusive rocks, mostly quartz monzonite to granite in composition, are exposed in a batholith and a stock in the Romanzof Mountains. Contacts with Neruokpuk Formation (Upper(?) Devonian or older) are mostly abrupt, concordant to cross-cutting, and locally adjoin tactite and hornfels. Contacts with Kekiktuk Conglomerate (Upper(?) Devonian or Mississippian) are gradational through a schistose zone. The lead-alpha age of zircon appears to be Late Devonian; the potassium-argon age of biotite, which is Cretaceous, may reflect updating by later reheating. Field age relationships are inconclusive but suggest pre-Kayak(?) (Late Devonian) granite emplacement. A late Paleozoic age is suggested for mafic volcanic(?) rocks and greenstone dikes in the granitic rocks and the Neruokpuk Formation.

The structural grain in the area strikes east-northeast, and south-dipping elements are common. Structures include relatively broad folds, small tight folds, and at least one large-scale overthrust fault. Although Mesozoic and Tertiary deformational features are dominant in northern Alaska, the Romanzof area may have been part of a Late Devonian orogenic belt continuous with one in northern Canada.

The mineral potential of the area is as yet largely unknown. Minor amounts of metallic sulfides and oxides are present in granite and Neruokpuk Formation rocks. Analyses of stream-silt samples suggest

⁵³ E. G. Sable, 1965, Geology of the Romanzof Mountains, Brooks Range, northeastern Alaska: U.S. Geol. Survey open-file report, 218 p., 7 pls., 23 figs., Aug. 10, 1965.

FIGURE 4.—Index map of Alaska, showing boundaries of regions referred to in discussion of Alaska geology.

the possibility of tin and beryllium potential. The Shublik Formation (Lower to Upper Triassic) contains rock phosphate.

New analysis of geologic structure of northern Alaska

E. H. Lathram has compiled a new geologic map of northern Alaska; his interpretation and analysis of the regional geology of this area suggest the following conclusions:

(1) Mid-Early Cretaceous intrabasin thrusting and gravity sliding, and Laramide thrusting, previously proposed in more restricted mapping, seem to have been widespread. The disposition of extensive thrust sheets transported along major thrust zones controls the present distribution of formations and facies throughout the Brooks Range and southern foothills of the range.

(2) Laramide detachment thrusts marked at depth by seismic discontinuities probably underlie the folds of the northern foothills of the Brooks Range. The thrusts appear not to crop out, except

perhaps along the faulted axes of northerly asymmetrical anticlines. These folds are the focus of past and current drilling.

(3) The thrusting has displaced the entire Brooks Range northward toward the Barrow high and the Romanzof Mountains.

These factors should be considered in exploration and in evaluation of the petroleum potential of northern Alaska.

Age of Ordovician and Devonian mafic rocks in northern Alaska

A single potassium-argon age of 475 million years, determined by M. A. Lanphere on hornblende from volcanic rocks in the Doonerak Mountain area of the south-central part of the Brooks Range, suggests that these volcanic rocks and associated slate and shale may be Ordovician rather than Devonian as had been previously thought.⁵⁴ Another single po-

⁵⁴ W. P. Brosgé and H. N. Reiser, 1960, Progress map of the geology of the Wiseman quadrangle, Alaska: U.S. Geol. Survey open-file report, 2 sheets.

tassium-argon age determination by Lanphere, a 373-m.y. age for hornblende from a mafic intrusive in the same area, substantiates the Devonian age of some intrusives in the Brooks Range.

Lisburne Hills largely fault sheets

Geologic mapping by I. L. Tailleux in the Point Hope quadrangle has disclosed that the central and northern Lisburne Hills consist of a number of broad, gently to moderately deformed allochthonous sheets of upper Paleozoic and Mesozoic predominantly marine sedimentary rocks. This structure is similar to that of the southern Lisburne Hills.⁵⁵ The apparent lower sheet is tightly crumpled and imbricately faulted along the arcuate east front of the hills, a structure analogous to that in the toe of a mud slide. Little stratigraphic order could be made of the graywacke and mudstone that underlie the area east of the Lisburne Hills; rocks probably equivalent to the Fortress Mountain Formation (Lower Cretaceous) predominate. Near the Lisburne Hills the graywacke and mudstone overlie fine-grained rocks of earliest Cretaceous age. To the east, along the upper Ipewik River, similar rocks overlie clay shales of Jurassic and Early Cretaceous age.

New interpretation of Tiglupuk Formation

Analysis of new paleontologic and field-stratigraphic data by D. L. Jones and Arthur Grantz (2-64) indicates that the Tiglupuk Formation of northern Alaska is a tectonic assemblage rather than a stratigraphic formation, and that it consists of thrust slices of a Jurassic and Early Cretaceous geosyncline. This interpretation of the Tiglupuk has major implications for any appraisal of the tectonics or petroleum possibilities of northern Alaska. Reconstruction of the paleogeography of the Tiglupuk assemblage suggests that: (1) the Jurassic and Early Cretaceous geosyncline of northern Alaska was compound, with a flysch and chert-ophiolite basin to the south, a medial intrabasin high, and a starved, pyritic-shale basin to the north; (2) this compound geosyncline was disrupted by pre-Albian thrusting (gravity sliding?) which produced the Tiglupuk assemblage; (3) the petroleum possibilities of the newly recognized mid-basin high should be tested, for such highs are a favorable habitat for petroleum. The area underlain by this high is potentially very large.

⁵⁵ R. H. Campbell, 1961, Thrust faults in the southern Lisburne Hills, northwest Alaska: Art. 354 in U.S. Geol. Survey Prof. Paper 424-D, p. D194-196.

WEST-CENTRAL ALASKA

Newly described marine sediments

Previously undescribed marine sediments in the White Mountain, McGrath area of Alaska have been reported by C. L. Sainsbury (p. C91-C95). Carbonate and clastic rocks about 5,000 feet thick have been dated as Middle (?) Ordovician and Devonian (?). These rocks are in fault contact with 8,100 feet of quartz conglomerate of Cretaceous (?) age. The rocks are unmetamorphosed and may be important in petroleum exploration.

Former warmer climate in Kotzebue Sound area

On the basis of (1) a buried discontinuous wood-rich zone in a now treeless area; (2) fossil beaver habitation beyond their modern range; (3) evidence for a period when ice wedges melted; and (4) buried soils well below the top of permafrost in the Kotzebue Sound-Seward Peninsula area, D. S. McCulloch and D. M. Hopkins postulate a postglacial warm interval of climate in northwestern Alaska. Nine radiocarbon dates suggest that this warm interval started about 10,000 years ago and ended 7,000 years ago.

Two large plutons mapped in Hughes quadrangle

In the Hughes quadrangle in west-central Alaska W. W. Patton, Jr., and T. P. Miller completed 1:250,000-scale mapping of two quartz monzonite-granodiorite plutons whose extent and lithologic character were previously poorly known. One pluton, about 100 square miles in area, is located in the Indian Mountains in the southeastern part of the quadrangle; the other, covering about 150 square miles, forms the core of the Zane Hills along the western margin of the quadrangle. Both plutons are of economic significance as they appear to be genetically related to important gold-placer deposits. Placer deposits within and along the eastern margin of the Indian Mountains pluton have been mined extensively in recent years, although present activity is limited to a small sluicing operation at the head of the Indian River. In the Zane Hills the Fairbanks Exploration Co. currently operates a small dredge on Bear Creek, along the eastern edge of the pluton.

Radiometric dating of deposits of Yarmouth age

The first radiometric ages of deposits of Yarmouth age on the North American continent have been obtained by D. S. McCulloch, D. W. Taylor, and Meyer Rubin (1-65). A fossil marine mollusk fauna containing more than 60 species has been collected from Yarmouth-age sediments of a marine delta which crops out on the Baldwin Peninsula and

north shore of Hotham Inlet in the Kotzebue Sound area. These delta sediments are stratigraphically well dated because they lie directly under Illinoian-age till, and have been deformed, presumably by the Illinoian glacier. Thorium-uranium age determinations on these shells indicate ages of $170,000 \pm 17,000$ years and $175,000 \pm 16,000$ years.

EAST-CENTRAL ALASKA

Area of Cambrian rocks expanded

Abundant specimens of *Oldhamia*, a fan-shaped trace fossil of probable Cambrian age, were found by Michael Churkin, Jr., and E. E. Brabb (p. D120–D124) in the Charley River quadrangle. The distinctive unit in which *Oldhamia* occurs is correlated with a nearby archaeocyathid-bearing unit of Early Cambrian age, thereby doubling the areal extent of known Cambrian rocks in east-central Alaska. This find, and two other occurrences of *Oldhamia*—in the Mt. Schwatka area⁵⁶ and in the Crazy Mountains—about 175 and 100 miles, respectively, west of the Charley River occurrence, suggests that rocks of Cambrian age extend to the central interior of Alaska.

Stratigraphy along Porcupine River

An almost complete section of the stratigraphic sequence from the upper part of the Ordovician through the Lower(?) Permian, on the Porcupine River, is provided by correlation of measured sections and by tentative age assignments of large fossil collections. W. P. Brosgé, H. N. Reiser, Michael Churkin, Jr., and J. T. Dutro, Jr., found that the pre-Permian part of this sequence differs greatly from that in the Brooks Range. It is only about 2,500 to 3,000 feet thick and consists mostly of shallow-water and near-shore shelf deposits. Ordovician, Silurian, and Devonian rocks are mainly fine-grained limestone and dolomite and coral-reef talus; Upper Silurian graptolitic shale a few hundred feet thick is a notable exception. Mississippian rocks are mainly shale (locally plant bearing) and shaly to sandy limestone and sandstone, and are only about 500 feet thick. Unconformities occur at the base of the Devonian and of the Mississippian. The Mississippian rocks in turn are overlain unconformably by 200 to 400 feet of Permian limestone and local basal sandstone, succeeded by a shale unit faunally similar to the Siksikpuk Formation and lithically similar to the southern calcareous Lower(?) Permian facies of the eastern Brooks Range.

⁵⁶ J. B. Mertie, Jr., 1937, The Yukon-Tanana region Alaska: U.S. Geol. Survey Bull. 872, 276 p., 15 pls.

Paleozoic stratigraphy of Tatonduk–Nation Rivers area

A sequence of Ordovician, Silurian, and Devonian rocks has been mapped in the vicinity of the Alaska-Yukon Territory border by E. E. Brabb and Michael Churkin, Jr., (1–64, 1–65; Churkin and Brabb, 1–65). The lower half of the sequence, 400 to 900 feet of Ordovician and Silurian graptolitic shale and chert, is considered to be a southwestern extension of the Road River Formation from the Richardson Mountains of the Yukon Territory. The McCann Hill Chert of Middle and Late Devonian age is 200 to 800 feet thick, consists mostly of chert and siliceous shale, and overlies the Road River Formation. In the vicinity of Jones Ridge, only 7 miles north of exposures of Road River Formation, a pure limestone section contains shelly faunas contemporaneous with the Road River Formation and McCann Hill Chert. A rapid change in conditions of sedimentation is suggested. The Tatonduk-Nation Rivers area seems to lie in a transitional zone between an early Paleozoic eugeosynclinal belt south of the Yukon River and a contemporaneous miogeosynclinal belt along the Porcupine River.

SOUTHWESTERN ALASKA

Sedimentary rocks on Nunivak Island

Geologic mapping by J. M. Hoare and W. H. Condon on Nunivak Island, in western Alaska, confirmed the occurrence of sedimentary rocks reported by Dall.⁵⁷ Several hundred feet of gently dipping sandstone and siltstone are exposed in cliffs surrounding an unnamed bay 7 miles west of Mikoryuk and underlie an area of about 25 square miles south of the bay. The rocks are largely nonmarine and contain good plant fossils identified by Jack Wolfe as being of Albian and Cenomanian age. Much of the sandstone here and on the west side of Nelson Island is laumontitized and probably forms a southwestern extension of the belt of laumontitized sediments which crops out north of the Yukon River, 150 miles farther northeast (J. M. Hoare, W. H. Condon, and W. W. Patton, Jr., 1–64). A second area of largely fine-grained sedimentary rocks was found on Nunivak Island on the south shore of Nash Harbor.

Jurassic age of Cook Inlet plutonism

R. L. Detterman, B. L. Reed, and M. A. Lanphere (p. D16–D21) have reported on the age of the plutonism in the Cook Inlet region of Alaska. Potassium-argon dates of hornblende and biotite indicate

⁵⁷ W. H. Dall and G. D. Harris, 1892, Correlation paper—Neocene: U.S. Geol. Survey Bull. 84, p. 245.

an age of 170 million years for emplacement of the Aleutian Range batholith; geologic mapping suggests emplacement between Early Jurassic and early Middle Jurassic time. The age of this batholith agrees closely with the age of the Kosina batholith in the Talkeetna Mountains about 250 miles to the northeast.

SOUTHERN ALASKA

Jarosite in McCarthy quadrangle

Hydrothermal alteration products, chiefly jarosite not previously reported, were found by E. M. MacKevett, Jr., near the Kennecott copper mines. The principal occurrence of the alteration products lies between the Jumbo and Erie mines in a zone that is several hundred feet long and as much as 10 feet thick. This zone consists mainly of a pulverulent mass of jarosite and clay minerals. It occurs along the contact between Nikolai Greenstone and the Chitistone Limestone in an area that was underlain by some of the large chalcocite lodes and at the approximate stratigraphic horizon of the lodes. Jarosite was also identified from altered limestone collected on the Bonanza mine dump and from altered limestone that was collected about 20 miles southeast of the Kennecott mines. Alteration zones of similar appearance have been noted elsewhere in the southern Wrangell Mountains. This alteration is significant because it should shed some light on the genesis of the Kennecott-type copper lodes, as well as provide target areas for exploration.

Brooks Lake and Naptowne glaciations correlated

R. L. Detterman, B. L. Reed, and Meyer Rubin (p. D34–D36) have established a minimum date of $8,520 \pm 350$ years B. P., based on radiocarbon analysis, for the second major advance of the Brooks Lake (Wisconsin age) Glaciation in the Iliamna Lake area. Four, or possibly five major advances are recorded by moraines in this area. These probably correspond to the Naptowne (Wisconsin age) Glaciation described by Karlstrom.⁵⁸

Tertiary fossils from Prince William Sound region

Marine megafossils collected by George Plafker from the greenstone unit of the Orca Group (previously considered Late Cretaceous (?) in age) near Galena Bay shed new light on the stratigraphy of the Prince William Sound region. The fossils include two species each of diagnostic crabs and Pelecypoda which, according to F. S. MacNeil, are of middle to

late Eocene age. They indicate that at least part of the Orca Group is of early Tertiary age, and that the Yakataga geosyncline, of the Gulf of Alaska Tertiary province, extended into the Prince William Sound region during Eocene time.

Continental Shelf of Alaska

Preliminary synthesis of the Quaternary history of the Gulf of Alaska Continental Shelf by T. N. V. Karlstrom suggests that the topography of the shelf, a series of arcuate ridges and associated deep troughs, represents the effects of deposition and excavation by Pleistocene glaciers. These glaciers repeatedly coalesced to form continuous piedmont ice sheets along the coast. The outermost submerged ridges near the shelf margin adjoining the Aleutian Trench are in line with, or rim, the seaward ends of the troughs that deepen landward to maximum depths between 100 and 400 fathoms. By this interpretation, during both the Knik (pre-classical Wisconsin age) and the Naptowne (classical Wisconsin age) maxima the shelf was virtually covered by ice except for Middleton and Chirikof Islands and a few isolated emergent areas near Kodiak Island. During the earlier, more extensive glaciations—Mount Susitna (Nebraskan? age), Caribou Hills (Kansan? age), and Eklutna (Illinoian? age)—the shelf was probably completely covered by ice that fronted in the ocean throughout the region.

SOUTHEASTERN ALASKA

Igneous activity in Chatham Strait area dated

Interpretation and analysis of more than 30 isotopic ages and of the Tertiary sedimentary and volcanic record in the area comprising eastern Chichagof Island, Baranof Island, and Admiralty Island have been completed by R. A. Loney and D. A. Brew. The results show that at least three post-Paleozoic igneous events have occurred and that probably 10,000 feet of vertical movement has taken place on the Chatham Strait and nearby faults since an Eocene intrusive event. An intrusion of Oligocene age heals a subsidiary fault zone, indicating that some of the vertical and accompanying lateral displacements occurred during the Eocene-Oligocene interval.

Isostatic rebound at Glacier Bay

E. H. Lathram estimates that a load of at least 3,000 feet of ice has been removed from the Glacier Bay area since 1790. Calculations by E. H. Lathram and M. D. Crittenden, Jr., using the time constants derived by Crittenden for unloading of Lake Bonne-

⁵⁸ T. N. V. Karlstrom, 1957, Tentative correlation of Alaskan glacial sequence, 1956: *Science*, v. 125, no. 3237, p. 73–74.

ville, Utah, indicate that this negative load should result in uplift on the order of 5 centimeters per year. Tidal-gauge records of the U.S. Coast and Geodetic Survey are reported (S. D. Hicks, written communication, 1964) to substantiate this result.

PUERTO RICO

Hydrogeologic map of Puerto Rico

A hydrogeologic map of Puerto Rico and adjacent islands was prepared by R. P. Briggs and J. P. Akers for inclusion in a report by D. B. Bogart, Ted Arnow, and J. W. Crooks (1-64) summarizing data and preliminary conclusions resulting from the first years of water resources investigations carried on in cooperation with the Commonwealth of Puerto Rico. The hydrogeologic map, which is also available as a Geological Survey Hydrologic Investigations Atlas (R. P. Briggs and J. P. Akers, 3-65), is a multicolor edition of the previously published provisional geologic map of Puerto Rico and adjacent islands, with surface and ground-water data added.

Variations with time in composition of volcanic rocks

Results of field and chemical studies by A. E. Nelson (1-65) suggests that from Early Cretaceous (Albian) to latest Cretaceous (Maestrichtian) time there was a generally continuous variation from basic to acidic in the chemical composition of volcanic effusions in the north-central part of the volcanic-plutonic core of Puerto Rico. Rocks of basaltic composition were deposited first; these were followed by andesite lavas and finally by pyroclastic rocks ranging in composition from dacite to rhyolite.

Wrench-fault deformation has resulted in large displacements

In the course of geologic mapping of the Aguas Buenas and Naranjito quadrangles and adjacent areas in northeastern Puerto Rico, M. H. Pease, Jr., has found evidence for regional large-scale wrench-fault deformation resulting from northeast-southwest compression. Although the displaced and deformed strata in all but a few instances reflect left-lateral movements, individual faults within the zone record both left-lateral and right-lateral movements. Summation of individual fault displacements shows an overall left-lateral displacement of possibly 20 kilometers. Right-lateral displacement on at least one fault within the zone may be as much as 15 km, and on one left-lateral wrench fault displacement may approach 30 km.

Isla Mona caves and guano deposits

During the late 19th and early 20th centuries, phosphatic guano was mined from caves in Isla Mona, a limestone island located in the Mona Passage, halfway between Puerto Rico and Hispaniola. Conflicting reports on the quantity of guano remaining caused the Puerto Rico Economic Development Administration to request an examination of these deposits and their geological environment by R. P. Briggs and V. M. Seiders. The largest cave systems were found in the base of the gently southwesterly-dipping Lirio Limestone adjacent to the cliffs that form most of the periphery of the island. Although rooms more than 40 meters across are common near the cliffs, large cave development only rarely extends more than 100 meters in from the cliff face; beyond this, passages pinch out. Higher in the Lirio Limestone, smaller ramifying caves occur, but no large rooms were found. No extensive cave systems were found in the dolomitic Isla Mona Limestone that underlies the Lirio. It is apparent that the zone where the lower part of the Lirio Limestone intersects the cliffs was the most favorable cave-forming locale on Isla Mona. On the basis of traverses through all large and many small cave areas, it is estimated that the total of guano originally available probably did not greatly exceed 100,000 short tons, about half of which has been removed in mining. The remaining guano is chiefly in relatively thin deposits in the more remote cave areas, or in isolated thicker pods left unmined for reasons not known.

Thickness of south-coast middle Tertiary basin from gravity studies

A detailed gravity survey of the south-coast middle Tertiary sedimentary basin has been completed by a geophysical party led by Andrew Griscom in cooperation with the Puerto Rico Economic Development Administration. Interpretation of the data indicates that the maximum nearshore thickness of rocks of Oligocene and younger ages probably does not exceed 7,000 feet (about 2,100 meters). Also indicated by the gravity data are major folds and a major fault trending N. 80° W. from Salinas for about 20 kilometers.

Aeromagnetic map of Puerto Rico Trench analyzed

Andrew Griscom, and W. H. Geddes, U.S. Naval Oceanographic Office, have completed analysis of an aeromagnetic map of the Puerto Rico Trench. The data indicate that there is probably no major fault on the north side of the trench. On the south side of the trench a major fault (possibly a thrust fault) with a vertical separation of at least 2 kilometers is

buried beneath the middle Tertiary and younger sedimentary rocks which extend north of Puerto Rico.

ANTARCTICA

The U.S. Geological Survey in cooperation with the U.S. Antarctic Research Program (USARP) of the National Science Foundation is continuing reconnaissance geologic mapping and topical studies in several localities in the western half of Antarctica (fig. 5). Topographic and planimetric base maps are compiled by the Topographic Division of the Geological Survey. (See section "Topographic Surveys and Mapping, Mapping in Antarctica.") Logistical support for the Antarctic field projects is provided by the U.S. Navy Operation Deep Freeze; helicopters and their crews for the northeastern Victoria Land study were provided by the U.S. Army.

Reconnaissance of northeastern Victoria Land

A geologic reconnaissance of 20,000 square miles of largely unexplored mountains was carried out by Warren Hamilton, D. F. Crowder, and D. A. Coates. The work was done by helicopter from Hallett Station and extended inland 200 miles, almost to the Rennick Glacier. The mainland consists largely of low-grade but highly deformed metagraywacke and metasiltstone, intruded by discordant plutons of

granodiorite. The clastic metasedimentary rocks lie unconformably upon high-grade schist and gneiss, which are exposed in the southwest and are probably products of Cambrian or Early Ordovician metamorphism. The metasedimentary rocks are overlain by little-deformed and unmetamorphosed Beacon Sandstone (Carboniferous? or Permian) intruded by diabase sills (Jurassic?). Deposition, deformation, and metamorphism of the clastic rocks as well as intrusion of the granodiorite thus occurred within the Ordovician to Carboniferous interval. Long high peninsulas along the Ross Sea coast (Adare, Hallett, and Daniell Peninsulas and Coulman Island) consist of Tertiary submarine lavas and breccias capped by Quaternary subaerial lavas. The marine rocks have been raised as high as 6,000 feet above sea level on faults bounding the peninsulas on the east (seaward) side. The Paleozoic rocks of the mainland have also been uplifted by faulting along the coast.

Ages of orogeny in the Pensacola Mountains

The stratigraphy and structure of the Pensacola Mountains (D. L. Schmidt, P. L. Williams, W. H. Nelson, and J. R. Ege, p. D112-D119; and 1-64), indicate that they are part of the fold-mountain belt coincident with the modern Transantarctic Mountains. In turn, the tectonic history of the Transantarctic fold-mountain belt is made clearer by the vivid details of geology found in the Pensacola Mountains.

Three major orogenies are recorded in the Pensacola Mountains: (1) During a late Precambrian orogeny, eugeosynclinal slate and graywacke rocks were folded. (2) During a post-Middle Cambrian orogeny (age based on fossils identified by A. R. Palmer), platform sediments consisting of limestone, clastic rocks, and volcanic sedimentary rocks were folded, and granitic magma intruded the Cambrian and older rocks. Granitic plutonic rocks associated with this event are isotopically dated by Rb-Sr whole-rock analysis as 510 ± 30 million years by Z. E. Peterman. (3) During a late Paleozoic orogeny, continental and shallow marine clastic sediments of uncertain age were folded. These clastic sedimentary rocks are younger than the granitic pluton but are probably older than flat-lying sedimentary rocks of Permian age (J. M. Schopf).⁵⁹ Biotite from the granitic pluton is dated isotopically by the K-Ar method as 265 ± 13 million years, or latest Carboniferous, by R. F. Marvin. This biotite

FIGURE 5.—Index map of Antarctica.

⁵⁹ J. M. Schopf, 1964, Paleobotanical studies in Antarctica [abs.]: *Geol. Soc. America Spec. Paper* 76, p. 317.

age presumably dates some part of the late Paleozoic orogeny.

Comagmatic igneous suites of Marie Byrd Land

Current petrochemical studies suggest the existence of at least four comagmatic suites in the Eights Coast-Marie Byrd Land region of West Antarctica according to E. L. Boudette and A. A. Drake, Jr. The oldest suite (1) consists of hypersthene-bearing rocks, including leuco-augite norite on the Eights Coast and quartz gabbro in the Executive Committee Range, types formerly known only in East Antarctica. These rocks have not been dated isotopically but are almost certainly pre-Mesozoic. The next younger suite of rocks (2) forms the Eights Coast-Thurston Island composite batholith, the characteristic rock of which is quartz diorite. Rocks of suite 2 have been recognized along the Eights Coast, where they are intrusive into rocks of suite 1. The rocks are at least 150 million years old, as determined by lead-alpha dating of zircon, but contain biotite that is thought to have crystallized during the Late Cretaceous to early Tertiary Andean orogeny, according to A. A. Drake, Jr., T. W. Stern, and H. H. Thomas (p. C50-C53). Suite 3 consists of an extensive series of intrusive and effusive rocks characterized by alkalic granite and rhyolite; a shallow intrusive member of suite 3 intrudes the quartz diorite of suite 2 along the Eights Coast. Members of suite 3 in the Executive Committee Range consist of folded rhyolitic volcanic rocks. Suite 3 is probably contemporaneous with Andean plutonic rocks of the Antarctic Peninsula that have an unknown structural relationship. The youngest suite (4) is volcanic, probably of Quaternary age, and consists of basalt and a distinctive anorthoclase-bearing

trachyte. These rocks previously have been considered to form a single effusive suite with the felsic volcanic rocks of suite 3. They are, however, not deformed and are petrochemically quite different. Deformation of the volcanic rocks of suite 3 observed by E. L. Boudette probably antedates the accumulation of the undeformed Quaternary volcanic rocks of suite 4.

Rubidium-strontium age studies by R. F. Marvin, F. G. Walthall, and C. E. Hedge on alkaline biotite granite collected by Boudette in the Clark Mountains of the Edsel Ford Ranges suggests a Late Jurassic or Early Cretaceous age for the granite-rhyolite suite.

Anorthoclase trachyte of Ross Sea-Marie Byrd Land region

Anorthoclase has been identified in trachyte from the Crary Mountains in Marie Byrd Land and on Ross Island by E. L. Boudette and A. B. Ford. Two samples of anorthoclase, $\text{Or}_{15.7}\text{Ab}_{74.2}\text{An}_{10.1}$ and $\text{Or}_{16.8}\text{Ab}_{64.8}\text{An}_{18.4}$, respectively, are homogeneous, transitional in structural state, and triclinic optically negative. The trachyte containing the mineral is the most distinctive rock in the Quaternary alkaline basalt-trachyte suite of the Ross Sea-Marie Byrd Land region. Feldspars of comparable chemical composition and petrographic relations in the Executive Committee Range and on Ross Island that are reported in the literature as plagioclase are probably also anorthoclase. The structural parameters of the anorthoclase from the Crary Mountains and Ross Island were calculated by computer methods in conjunction with work in progress by D. B. Stewart and T. L. Wright. (See the section "Mineralogic Studies and Crystal Chemistry, Physical Properties of the Feldspars.")

GEOLOGIC AND HYDROLOGIC INVESTIGATIONS IN OTHER COUNTRIES

For more than two decades the U.S. Geological Survey has assisted newly developing nations in appraising their natural resources, establishing geologic and hydrologic agencies, and training earth scientists. During this time the Geological Survey has sent 513 members of its staff on investigations and training assignments to 68 countries, and has provided specialized training in the United States for 787 scientists from 70 nations. Under the auspices of the U.S. Agency for International Development and other national and international agencies, this assistance has helped to strengthen

earth sciences programs or establish vigorous national geologic and hydrologic organizations in 24 countries.

In fiscal year 1965, the U.S. Geological Survey assigned 128 specialists to 21 countries at the request of other governments, and provided or helped arrange academic or on-the-job training in the United States for 92 geologists and engineers from other countries. The accompanying table summarizes the type of assistance given to each country by the Geological Survey during the year.

Technical assistance to other countries provided by the U.S. Geological Survey during fiscal year 1965

Country	USGS specialists assigned to other countries			Scientists from other countries trained in the United States	
	Number	Type	Type of activity ¹	Number	Field of training
Latin America					
Argentina-----	None	-----	-----	1	Geology and mining (drilling).
Bolivia-----	1	Chemist	A, C	1	Geochemical research.
Brazil-----	3	Geologist		None	
	9	do	A, C, D	2	Topographic mapping.
	1	Cartographer		1	Photogeology.
	1	Topographic engineer		1	Laboratory analysis and techniques.
	3	Hydrogeologist		1	Ground-water hydrology.
	3	Hydraulic engineer (SW)		1	Stratigraphy and sedimentology.
				3	General geology.
				1	Vertebrate paleontology.
				1	Mineral exploration.
British Guiana--	None	-----	-----	1	Ground-water exploration techniques.
Chile-----	None	-----	-----	1	Cartography and map preparation.
Colombia-----	5	Geologist	A	1	Engineering geology.
Costa Rica-----	3	do	B	None	
	5	Geophysicist			
	1	Hydrogeologist			
	1	Chemist			
Panama-----	1	Hydrogeologist	C	1	Igneous geology.
Venezuela-----	None	-----	-----	2	Surface-water techniques.
Africa					
Dahomey-----	2	Geologist	B	None	
Egypt-----	4	Hydrogeologist	A	4	Ground-water hydrology.
	1	Hydrochemist			
Ghana-----	None	-----	-----	1	Water development (drilling).
Liberia-----	5	Geologist	A, C	2	Mineral exploration.
	1	Administrative officer		1	Surface-water techniques.
Libya-----	1	Hydrogeologist	B	1	Ground-water hydrology.
Morocco-----	None	-----	-----	1	Mining engineering.
Nigeria-----	4	Hydrogeologist	A	None	
	2	Hydraulic engineer (SW)			
Sudan-----	None	-----	-----	1	Ground-water development.
				1	Photogrammetry and cartography.
				1	Rural water development.
				2	Ground-water hydrology.
Tunisia-----	1	Hydrogeologist	B	None	

Technical assistance to other countries provided by the U.S. Geological Survey during fiscal year 1965—Continued

Country	USGS specialists assigned to other countries			Scientists from other countries trained in the United States	
	Number	Type	Type of activity ¹	Number	Field of training
Near East-South Asia					
Afghanistan----	2	Hydraulic engineer (SW)	A	1	Photogrammetry and cartography.
				1	Hydrologic engineering.
				1	Cartography.
				1	Hydrology.
				2	Topography.
				1	Aerial photography and mapping.
				1	Surface-water hydrology.
India-----	None			3	Exploration and development of uranium ores.
Iran-----	None			1	Seismic instrumentation.
Jordan-----	1	Hydrogeologist	C	1	Spectrography.
				1	Well drilling; engineering.
				1	Hydrometry.
				1	Hydrology.
				1	Surface-water hydrology.
Nepal-----	2	Hydraulic engineer (SW)	B	None	
Pakistan-----	15	Geologist	A	1	Geochemistry; mineralogy; petrology.
	1	Geophysicist		2	Mineral exploration and development.
	1	Geochemist		7	Hydrology.
	2	Photogrammetrist		2	Geologic administration.
	2	Cartographer		1	Economic geology.
	1	Publications specialist		1	Laboratory techniques and analysis.
	1	Driller		3	Ground-water hydrology.
	2	Program assistant			
Saudi Arabia---	5	Hydrogeologist	A	1	Mass spectrometry.
	9	Geologist			
	1	Geochemist			
	5	Geophysicist			
	1	Chemist			
	1	Geodesist			
	1	Photogrammetrist			
	1	Topographic engineer			
	1	Program assistant			
	2	Driller			
	1	Transportation officer			
Turkey-----	1	Cartographer	C	1	Mapping; photogrammetry.
	2	Geologist		1	Topography.
	2	Hydrogeologist		4	Water resources and ground-water engineering.
Far East					
Indonesia-----	1	Geologist	B	1	General geology; geophysics.
				1	Ground-water investigations (drilling).
				1	General geology.
Japan-----	None			1	Mining geology.
				1	Geothermal conditions in the United States.
				1	Ash-flow tuffs.
				1	Sedimentary petrology.
Korea-----	1	Hydrogeologist	C	1	Geophysics; geochemical exploration.
Philippines-----	2	Geologist	A	1	Structural geology.
				2	Ground-water hydrology.
				1	Engineering geology.
				1	Stratigraphy; sedimentation; petroleum geology.
Thailand-----	3	Geologist	A, B	1	Hydrometeorology.
				1	Ground-water hydrology.

¹ A, broad program of advisory help in institutional development and direct help in resources appraisal; B, limited program of advisory help, training, and investigation in selected fields; C, advisory help or consultation, with specific training or investigational activity; D, geological education and assistance to universities.

An important result of the U.S. Geological Survey assistance to the newly developing countries has been the publication of maps and reports needed for exploration, development, and use of resources.

National geologic maps or summary resources reports have been issued for Ecuador, Indonesia, Libya, Pakistan, Peru, the Philippine Islands, Saudi Arabia, and Thailand. Many hundreds of publica-

tions resulting from the work of U.S. Geological Survey specialists and their counterparts have been issued by host-country agencies. In addition, 685 documents, resulting from technical assistance activities, have been published or issued by the Geological Survey. A new bibliography and index of the documents issued by the Geological Survey has been compiled and is now being prepared for publication.

SUMMARY BY COUNTRIES

AFGHANISTAN

Surface-water studies of watersheds expanded

The program of surface-water investigations was markedly expanded to include the Ghazni, Tarnak, Khash, Farah, Adraskand, and Hari Rud watersheds, in addition to existing streamflow, meteorologic, and sediment studies currently in progress in the Helmand River basin. Also, plans were made for an expanded network of snow courses as well as extensive snow surveys in the Helmand basin to improve the accuracy of forecasting for spring runoff. The U.S. Geological Survey team is making a study of recorded data on flood duration, low flow, and flood frequency in the Helmand River basin to assist the Helmand Valley Authority and the Water and Soils Survey Authority in water-management problems.

BOLIVIA

Economic geology

A report titled "The Gold Placer Deposits at the Foot of the Eastern Cordillera of Bolivia," by H. G. Freydanck has been issued by the Bolivian National Department of Geology as a result of the geologic-assistance program sponsored by the U.S. Agency for International Development. The report describes the distribution and extent of gold-bearing gravels in eastern Bolivia, which constitute one of Bolivia's major resources. Geologic exploration necessary to the study was supervised by C. M. Tschanz and J. F. Seitz, of the U.S. Geological Survey. Analyses of samples from mine dumps in the Berenguela district, taken by C. M. Tschanz and analyzed by the Bolivian National Department of Geology, show 3 to 5 percent cadmium. Zinc concentrates produced as a by-product in Comibol's Telamayo mill contain between 0.8 to 1.4 percent cadmium.

BRAZIL

Iron

J. V. N. Dorr 2d (1-64, 2-65) discusses, in two companion papers, the nature, origin, and reserves

of the supergene and hypogene iron ores, respectively, in Minas Gerais, Brazil. Each type resulted from enrichment of the original sedimentary iron formation (itabirite). Weathering related to peneplains, the highest and oldest of which may date back to Early Cretaceous time, removed quartz and hydrated some of the iron oxides to form "intermediate" ores averaging about 63 percent iron. Known reserves are estimated at 600 million tons. High-temperature solutions removed quartz and redistributed iron from the itabirite in favorable structural settings, (chiefly in folds, in a few places in thrust-fault zones, and in some localities, where regional linear structures steepen abruptly) to form masses of hard, nearly pure hematite averaging more than 68 percent iron. The ore textures and distribution of the deposits show clearly that enrichment occurred at depth during a severe Precambrian orogeny. Some individual deposits contain more than 400 million tons; in all, 40 ore bodies and closely associated groups of ore bodies are known.

Copper

Geologic and geochemical studies of the Caraiba copper deposit and the surrounding Curaca River basin area by R. W. Lewis, Jr., in cooperation with S. Q. Mattoso, R. J. P. Brim, and A. M. Santos, of the University of Bahia and the Departamento Nacional da Produção Mineral, have indicated the possibility of additional copper deposits near Caraiba in north-central Bahia. Geologic mapping and soil sampling revealed a potential cupriferous belt in highly metamorphosed early Precambrian rocks between Santa Rosa and Carrio Vermelho, 70 kilometers to the north. Copper mineralization is found in ultramafic intrusions, which are localized in small folds of an anticlinal arch. The cupriferous ultramafic masses are marked by well-defined copper anomalies in both soil and vegetation. These anomalies are from 50 to 100 times the background value for copper and are also associated with anomalies indicative of accumulation of copper of possible economic value.

Hydrology

In northeastern Brazil, hydrogeologic work that has been completed indicates favorable ground-water conditions in the alluvial aquifer of the Capibaribe basin, Pernambuco; in the limestone aquifer of the Irecé area; and in alluvial sandstone and limestone aquifers of the Acu Valley, Rio Grande, do Norte.

L. C. Davis reports that measurements to evaluate the flow of the Amazon River during the period July 1963 to August 1964 showed an average rate of flow

to the ocean of 7 million cubic feet per second. A maximum rate of flow measured at Obidos, Brazil, was found to be close to 12 million cfs, a figure considerably higher than that previously estimated (8.5 million cfs reported in Professional Paper 501-A).

Geologic-education campaign completed

In December 1964 the Brazilian "Campanha para a Formação de Geólogos" (Campaign for the Education of Geologists), begun in 1957, was completed. This project, carried out in cooperation with the Brazilian Ministry of Education and Culture under the sponsorship of the U.S. Agency for International Development, succeeded in firmly establishing or improving modern programs of geologic education at Brazilian universities at São Paulo, Porto Alegre, and Rio de Janeiro. Substantial assistance was also given universities at Ouro Preto, Recife, and Salvador. The U.S. Geological Survey furnished more than 30 man-years of teaching assistance over the 7-year period.

This cooperative effort has culminated in the establishment of six dispersed centers of geologic research. Faculties practicing research, and students working on their senior theses, currently are producing more geologic mapping than Brazil has enjoyed in many years, and the various journals inaugurated by these schools are greatly increasing the production of geologic literature and maps. This information already is being put to use, particularly in the compilation of national maps. Coupling this school effort with the very large absolute increase in geologic manpower, Brazil now is in a good position to staff its research organizations and to begin a significant attack on the enormous problem of studying the geology and resources of the largest nation in Latin America.

D. J. Cederstrom, U.S. Geological Survey, working with the Government of Brazil, has prepared a textbook of ground-water hydrology in the Portuguese language. It is entitled "Aqua Subterranea, Uma Introdução." D. F. Campbell, of the U.S. Geological Survey, and John Milne Albuquerque Forman, of the Brazilian National School of Geology, collaborated on the preparation of a syllabus of economic geology which has been published in Portuguese in two volumes by the National School of Geology at Rio de Janeiro, Brazil.

CHILE

Potassium-argon ages of ash flows in Chile

According to a report by R. J. Dingman (p. C63-C67), of the U.S. Geological Survey, ash-flow sheets

that cover many thousands of square kilometers in Chile, Bolivia, Argentina, and southern Peru are also one of the most extensive ash-flow series in the world. On the basis of potassium-argon determinations on biotite from four rock samples, the ash-flow sheets in the San Pedro de Atacama region of northern Chile are of Pliocene age. The ages determined range from 4.7 to 7.6 million years.

Mineralogy of Chilean nitrates

Mineralogic studies of the Chilean nitrate deposits by George Ericksen and Mary Mrose have brought to light the fact that potassium, which is of economic importance, occurs mainly as niter (KNO_3) and as a probably new sulfate mineral, on which further work is being done. Though earlier workers have inferred from chemical data that the potassium is in sylvite, polyhalite, and apthitalite, these minerals have never been identified in the nitrate ores, and thus probably are rare or absent. Darapskite, $\text{Na}_3(\text{NO}_3)(\text{SO}_4) \cdot \text{H}_2\text{O}$, is a widespread and relatively abundant constituent of the nitrate ores, including those high in chloride as well as those high in sulfate. Three new minerals, in addition to the potassium-bearing sulfate mineral, have been tentatively identified; two are chromate-bearing salts, and the other is a sulfate.

COLOMBIA

Economic geology

Preliminary investigations on the western side of the large Cretaceous geosyncline show phosphate rock possibilities in grades of 15 to 20 percent phosphate. The favorable geologic environment gives some hope for the possible discovery of higher grade material.

Stratigraphic studies of evaporites in the Cretaceous section near Bogotá indicate a large potential for rock salt and possibly some gypsum-anhydrite, potash, and pyrite.

The Colombian National Geological Service is mapping on a 1:25,000 scale in 4 areas selected for their mineral and metal possibilities. A systematic geologic study of mineral and metal resources of Colombia is being carried out by 22-25 Colombian geologists of the National Geological Service assisted by E. M. Irving, L. V. Blade, Tomas Feininger, and Donald McLaughlin, all of the U.S. Geological Survey.

GHANA

Ground water

H. E. Gill, reporting on a ground-water reconnaissance in the Republic of Ghana, has defined four

major geohydrologic provinces. In three of these, known as the Precambrian, Voltaian, and Coastal Block Fault provinces, large or even adequate ground-water supplies are difficult to develop from the prevailing crystalline and consolidated sedimentary rocks. However, small supplies (from a few gallons to 50 gallons per minute) of good quality water can generally be obtained in most parts from carefully located wells. Water supplies ranging from 50 to 200 gpm or more can be obtained from wells tapping sand and soft limestone aquifers of the Coastal Plain province near the Gulf of Guinea.

INDONESIA

Engineering geology

At the close of the year, R. J. Anderson completed manuscripts, in cooperation with counterparts, for publication by the Indonesian Geological Survey on engineering-geology investigations within the Tji-manuk River Project, West Java. A report on recent faulting at the Eretan damsite, West Java, and ground-water studies near Makassar, South Sulawesi are in preparation.

A geologic map of Indonesia, on a scale of 1:2,000,000, is in process of editing, with expectation of publication late in 1965.

NIGERIA

Ground water

William Ogilbee, working in the Sokoto artesian basin in northwestern Nigeria, put down 20 test wells ranging from 300 to 2,000 feet in depth and found high-pressure artesian aquifers in an area of about 2,700 square miles in the western part of the basin. Flowing wells have already markedly contributed to the economy of the region.

In the Chad basin, northeast Nigeria, 200 small-diameter artesian wells drilled in an area covering 13,500 square miles found sufficient water for the requirements of the seminomadic people and their cattle.

PAKISTAN

Geologic map of Pakistan printed

A new geologic map of Pakistan at a scale of 1:2,000,000, compiled by R. O. Jackson, U.S. Geological Survey, and M. Abu Bakr, Geological Survey of Pakistan, was published in December 1964 by the Geological Survey of Pakistan at the Survey of Pakistan press, Rawalpindi. This multicolor, 36- by 45-inch map was prepared as part of the Mineral Exploration and Development Program sponsored

by the Government of Pakistan and the U.S. Agency for International Development. Cartographic work was done by the Geological Survey of Pakistan, Quetta, under the overall supervision of J. T. Heare. M. E. Hanes assisted with the final preparation of the map and the supervision of the press run. This map is based on the compilation of all available published and unpublished geologic data as of late 1964, as well as upon results of several special projects undertaken jointly by members of the Geological Survey of Pakistan and the U.S. Geological Survey.

Lithofacies maps of Paleocene rocks in West Pakistan

C. R. Meissner, with Habib-ur-Rahman of the Geological Survey of Pakistan, prepared an isopach map and three lithofacies maps of the Paleocene rocks of West Pakistan. This study reveals the distribution, thickness, and lithology of the Paleocene rocks, economically significant as oil-bearing rocks of the Potwar Plateau in northern West Pakistan. The Paleocene reservoir rocks in the oil fields of the Potwar Plateau have a thickness of 300 to 500 feet or more, a clastic ratio of 1 or less, and a sand-shale ratio of less than 1:4. On the basis of these criteria for finding oil-reservoir rock in Paleocene sediments, the isopach and lithofacies maps show 3 areas where exploration might be concentrated: 2 in the Sulaiman Range and 1 in the northwestern Kirthar Range.

Interpretation of Himalayan structures

The Tertiary mountain system of West Pakistan has a series of four parallel reentrants and intervening festoonlike arcs in the mountain front. Analyses of metamorphic structures in the easternmost reentrant (Hazara-Kashmir), which were mapped by T. W. Offield, J. A. Calkins, and geologists of the Geological Survey of Pakistan, suggest that the structures were formed by large-scale southward movement of rock masses on either side of a zone in which such movement was impeded, probably by a subsurface buttress. Geophysical surveys and deep drilling reveal the presence of basement highs aligned with the axial zones of the reentrant. Regional southward thrusts have not yet been identified in Pakistan, but the festoonlike configuration of the ranges and the coincidence of basement highs with at least two of the reentrants lead to the speculation that the whole system formed as a result of regional supracrustal movement southward around deeply buried basement promontories.

Chromite studies in Hindubagh district

Mapping by Roger van Vloten of the ultramafic intrusive at a scale of 1:12,000 is nearing completion in the eastern part of the district. A special study

of serpentine diapirs indicated by circular lineaments within the mass was undertaken over the entire extension of the complex. Intensive sampling of the rocks at some diapir structures has shown areas of low-density rocks entirely surrounded by high-density rocks, supporting the idea that the circular lineaments are caused by upward thrust of lighter masses. A contour map showing variation in iron content of chromite samples over the entire complex indicates low-iron anomalies at several diapir sites. Evidence is accumulating that the complex is stratified, but disturbed by faulting and diapirism, especially in the eastern part.

To search the areas of circular lineaments for chromite ore bodies, gravity surveys were completed in 11 areas by Abul Farah, Arif Raza Jafree, and Mohammad Ali Mirza, of the Geological Survey of Pakistan, assisted by W. J. Dempsey. Several gravity anomalies were delineated, and two of them were drilled to a depth of 300 feet. No chromite ore bodies were found. Regional gravity surveys conducted across and around the mass seem to indicate that there is a subsurface extension of the ultramafic rocks to the south and to the east tens of miles beyond the outcrop limit, and that the body or bodies thin to the northwest in conformity with the observed regional stratification.

Mineral investigations in Chagai district

R. H. Nagell, with counterpart geologists of the Geological Survey of Pakistan, completed a study of mineralization in the Chagai district including occurrences of copper, iron, and travertine ornamental-stone deposits in the Chagai Hills. The only deposits of economic importance in this area are those of travertine. Deposits of sulfur and travertine and showings of copper, iron, lead, and zinc in other parts of the Chagai district were also investigated. Of these, the Koh-i-Sultan Volcano sulfur deposit, which produced about 500 tons of sulfur per month, and the Zard Kan, Mashki Chah, and Zeh travertine deposits are the most important economically.

Mineralogic studies of Ziarat laterite

J. J. Matzko, S. A. Stanin, and counterpart geologists of the Geological Survey of Pakistan completed a study on ferruginous "laterite" at the base of the Tertiary sequence near Ziarat, West Pakistan. Detailed mineralogic work was done on samples from 4 stratigraphic sequences of which 3 sections contain intervals 2, 6, and 19 feet thick that are sufficiently high in alumina to justify further investigation for possible bauxite ore. Aluminum in the samples is found in the oxide minerals—boehm-

ite, diaspore, and gibbsite. The laterite has a high iron content which is derived from admixed hematite.

Corrosion and encrustation mechanisms in water supplies of Indus Plain

Preliminary field and laboratory studies of ground water in reclamation projects of the Indus Plain, West Pakistan, by F. E. Clarke and I. K. Barnes, showed that both corrosion and mineral encrustation are contributing to early failure of dewatering wells. Thermodynamic studies showed the waters to be supersaturated in respect to calcite, siderite, and iron hydroxides, so that some encrustation probably is inevitable, even in the absence of pumping. Bacterial sulfate reduction and action of long-line differential-aeration cells appear to be the major corrosion processes, and there is evidence that corrosion is intensifying mineral-encrustation processes. Tentative recommendations for corrective action already are being implemented. New construction materials have been selected to minimize corrosion and bonding of encrustants; design changes are being made to eliminate long-line currents, and sterilization treatments are being applied.

PANAMA

Hydrology

As part of a hydrogeologic training program for the Panamanian Instituto de Recursos Hidraulicos y Electrificación, R. J. Dingman made a brief ground-water reconnaissance of the Azuero Peninsula of south-central Panama. The work points up favorable possibilities for the development of moderate- to large-capacity wells tapping conglomerate or sandstone aquifers near La Flora and Las Flores. In areas favorable for ground-water development, additional detailed geologic mapping and also some seismic surveys are needed because of the complexity of the geologic framework.

PHILIPPINE ISLANDS

Geologic map of Philippine Islands

A new geologic map of the Philippine Islands on a scale of 1:1,000,000 was published as a joint project of the Philippine Government, private petroleum companies, the U.S. Agency for International Development, and the U.S. Geological Survey. Data were assembled and compiled by Oscar Crispin and Froilan Geruasio of the Philippine Bureau of Mines. Twenty-three rock units that represent geologic periods from the pre-Jurassic basement complex to Recent deposits are shown. Among the major features illustrated on the map are the large northwest-

trending faults that cut almost every large island except Palawan. Major volcanic activity may be related to the faulting. Flows, tuffs, and ash cover large areas; further structural studies are needed for purposes of correlation. The geologic settings of deposits of copper, chromium, iron, and nickel are shown.

Chromite in Zambales Mountain Range

Detailed mapping by Darwin Rossman and geologists of the Philippine Bureau of Mines in the Zambales Mountain Range covered much of the gabbro-peridotite complex. Five 1:50,000-scale quadrangle sheets show that at least 90 percent of the chromite occurs in 2 dunite zones in peridotite near gabbro contacts in the vicinity of the Acoje and Masinloc mines. Layering in the rocks cannot be used as a reliable guide to prospecting. A report on this work is in preparation.

Santa Inez iron-ore deposit replaces limestone

Geological mapping and exploration by Laurence Andrews and geologists of the Philippine Bureau of Mines show that the Santa Inez iron-ore deposit is a replacement of a limestone member of a roof pendant of Eocene to Miocene sedimentary rocks that are surrounded by intrusive diorite and skarn. To date, 30 to 50 million tons of magnetite, with associated pyrite, pyrrhotite, and chalcopyrite, are indicated. A progress report by Andrews and Leonardo Antonio, Philippine Bureau of Mines, is in preparation.

Mineral-resource maps

Seven mineral-resource maps have been published to show the location of major mineral deposits. No geology is shown.

SAUDI ARABIA

Geochemical prospecting of wadi sediments

Geochemical prospecting for copper, zinc, molybdenum, and tungsten in igneous and metamorphic rocks along the mountains of the Red Sea scarp and the interior plateau of the Precambrian shield in Saudi Arabia by P. K. Theobald, Jr., and C. E. Thompson has disclosed that anomalous concentrations of these metals in wadi sediments are greatly diluted by wind-transported debris. The mean grain size of the diluting debris is about 100 to 150 mesh. Debris coarser than about 80 mesh does not seem to have travelled far by air. The geochemical effects of dilution of wadi sand by major air transport of fine-grained particles from one drainage area to another can be largely avoided by using coarse fractions of sediment for analysis. Tests for copper in

103 pairs of samples of sand disclosed copper in the range of 20 to 6,000 parts per million in the -30 to +80 mesh used in this study and 20 to 1,500 ppm in the -80 mesh ordinarily used in such investigations. At one locality represented by 91 pairs of samples, the amount of copper in the coarse fraction ranged from <10 to 600 ppm, with 75 ppm being the anomalous threshold; and the zinc ranged from 25 to 800 ppm, with 100 ppm and above considered anomalous. In the fine fraction the copper content ranged from 10 to 225 ppm, with the median at 75 ppm; and zinc ranged from <25 ppm to 300 ppm, with the median at 50 ppm.

Two other metal-bearing media seem useful for reconnaissance sampling: heavy minerals, and magnetite. Heavy minerals from wadi sand show distinctive variations in molybdenum and tungsten content, and magnetite displays strong regional variations in copper, zinc, and molybdenum. Both of these media are present as alluvial trains in the wadis, in contrast to the normal bulk wadi sediment, in which anomalous concentrations of metals are abruptly masked by local sediment.

Mineral discoveries in Arabian Shield

Scheelite and powellite were found for the first time in Saudi Arabia by C. S. Hummel, J. W. Mytton, and J. W. Whitlow, working with Hashim Hakim and A. O. Ankary, of the Directorate General of Mineral Resources, Royal Saudi Arabian Government. Scores of small occurrences, none of which are minable, were discovered in an area 20 to 30 miles wide extending southward about 125 miles from Bishah to Khamis Mushayt. Both minerals occur as disseminated grains in the youngest of the Precambrian granites. They are also disseminated in long, narrow roof pendants of greenstone, chlorite schist, and hornblende schist in the granite, but have not been observed in pendants of marble. Quartz veins and pegmatite dikes of simple mineral composition apparently genetically related to the granite also contain scheelite and powellite.

A possible occurrence of wolframite related to the youngest Precambrian granite has been discovered by Richard Goldsmith, of the U.S. Geological Survey, and Jameel Kouther, counterpart geologist with the Directorate General of Mineral Resources. Heavy-mineral concentrates from the area give no reaction for scheelite by ultraviolet light, but have been found by C. E. Thompson to contain as much as 1,500 parts per million of tungsten. This tungsten is tentatively attributed to wolframite in the concentrates.

Beryl has been discovered in Saudi Arabia by J. W. Whitlow at one locality about 125 miles northeast of Bishah. The beryl is in pegmatite in massive red granite of Precambrian age. The beryl-bearing pegmatite is roughly circular in plan and about 300 feet across. Field estimates indicate less than 0.1 percent beryl in the pegmatite.

Jasperoid in barite veins near Rabagh

An unusual dark-golden-brown jasperoid was found by D. A. Brobst in pods in the thicker parts of barite veins in fractures of the Red Sea rift system about 60 miles north of Jidda, Saudi Arabia. The barite deposits are about 20 miles inland from Rabagh on the Red Sea. The suite of trace elements in the jasperoid includes Be, Pb, Y, B, Mn, Fe, and Cu in anomalous amounts and some Mg and Mo. Field observations and the suite of trace elements detected suggest that the jasperoid is primary. Further investigations of the area containing barite veins may determine if these veins are part of a large zoned ore deposit.

Experimental electromagnetic investigations

Fractures and joint surfaces in zones of alteration and country rock around many of the ancient mines in Saudi Arabia are coated with copper carbonate. The prevalence of this coating in certain places suggests that massive sulfides may be associated with the gold and silver deposits that were formerly worked in these mines. Experimental electromagnetic investigations conducted over five ancient mines by W. E. Davis and R. V. Allen with N. N. Akhrass and R. H. Zeldan, of the Directorate General of Mineral Resources, Royal Saudi Arabian Government, revealed only small anomalies associated with the zones of alteration. This lack of a pronounced electromagnetic response over the zones indicates that the copper minerals are too disseminated to be detected readily by electromagnetic methods, and that the induced-polarization technique, which is generally more successful in searching for disseminated sulfide deposits, should be used in conjunction with other geophysical prospecting methods.

Topographic field operations

Gene Harbert, James Crabtree, and Thomas Taylor began the organization of a Topographic Mapping Section within the Ministry of Petroleum and Mineral Resources for the immediate purpose of furnishing topographic services to the joint minerals-exploration project. The topographic facility thus established is to continue its work after the

close of its present duties as a part of the national mapping effort, with strong emphasis on the specific mapping needs of the Ministry.

A photogrammetric and photographic laboratory has been constructed and equipped in Jidda. Tidal observations of the Red Sea have been observed and recorded at Jidda for most of 1 year. Fieldwork to establish a basic geodetic net of vertical and horizontal control for western Arabia has been started.

THAILAND

Economic geology

In the Loei-Chiengkarn area, northeast Thailand, preliminary examinations of 47 iron, base-metal, manganese, gold, barite, and gypsum prospects were completed. Detailed geologic, geochemical, and geophysical studies were conducted at 3 iron and 2 base-metal prospects. Diamond drilling is in progress at two iron deposits. At one of these, approximately 7 million tons of iron ore is indicated.

At Chaiyaphum, northeast Thailand, geologic investigation and diamond drilling of a salt deposit resulted in the development of 600 million tons of halite in a bedded deposit. Additional reserves are inferred.

Three economic-geology reports entitled "The Mae Jong Manganese Deposit of Northwestern Thailand," by L. S. Gardner and R. C. Smith; "Fluorspar Deposits of Thailand," by L. S. Gardner and R. C. Smith; and "Phichit Gypsum Deposits, Central Thailand," by L. S. Gardner were submitted to the Royal Thai Department of Mineral Resources for publication in Thailand.

INTERNATIONAL COOPERATION WITHIN THE REGION OF THE CENTRAL TREATY ORGANIZATION

CENTO symposia on geologic studies

In continuation of activities concerned with coordinating geologic information and geologic investigations within the framework of the CENTO countries, representatives of the U.S. Geological Survey took part in symposia and field trips and were actively engaged in several joint CENTO working groups on various geologic studies and problems common to Pakistan, Iran, and Turkey. A noteworthy contribution of the Second Field Symposium of the CENTO Stratigraphic Working Group resulted in a report entitled "Correlation of the Cretaceous System in Turkey, Iran, and Pakistan." This report describes the five major tectonic zones that extend through these CENTO

countries and the Cretaceous System as developed distinctively in each of these zones. Summaries of the historical geology of the Cretaceous System and of the main mineral and mineral-fuel deposits within this system are given for each country. The important mineral and mineral-fuel deposits are copper, chromite, iron, phosphate, and petroleum in Turkey; chromite, lead-zinc, iron, phosphate, and petroleum in Iran; and chromite, laterite, and iron in Pakistan. Attempts are made to relate known mineral deposits to their habitat so that mineral exploration in one country may be aided by the knowledge gained in others.

COOPERATIVE SHORT-TERM TOPICAL INVESTIGATIONS ABROAD

AUSTRALIA

Geomorphology

S. A. Schumm made preliminary studies of modern and ancient river channels located on the Riverine Plains of New South Wales and found that the morphology and sediment characteristics of the Murrumbidgee River differ markedly from those of the ancient aggraded channels still exposed on the surface of the alluvial plain. The modern river has adjusted its gradient for a major decrease in sediment load and changed discharge by a modification of the shape and sinuosity of the channel rather than by incision into the alluvial plain.

GREENLAND

Radiocarbon age determination of turf in Ivar Baardsons Glacier

Turf collected by J. H. Hartshorn and J. P. Schafer that is interbedded with ablation moraine deposits of the large moraine at the end of the Ivar Baardsons Glacier, Schuchert Dal, East Greenland, has a radiocarbon age of $1,490 \pm 250$ years (W-1378) according to Meyer Rubin. This moraine and similar moraines of nearby glaciers were formed by an ice advance that began much earlier than the advance of the 17th to 19th centuries, to which recent moraines around the North Atlantic have generally been attributed. The turf is from a locality nearly free of vegetation and underlain by dead ice; thus, there is a question as to the conditions under which the formerly continuous cover of vegetation was established. A colder climate presumably would be less favorable for the spread of vegetation; a warmer climate probably would produce accelerated melting of the buried dead ice and severe instability of the ground surface.

Meyer Rubin found the C^{14} age of pelecypod shells from marine deposits 43–50 meters above the present sea level at a nearby locality to be $7,900 \pm 350$ years (W-1381). This approximately fits Washburn and Stuivers' ⁶⁰ curve of uplift in Kong Oscars Fjord 80 kilometers to the north.

ICELAND

Ten Pleistocene glaciations recognized in Iceland

Stratigraphic, paleomagnetic, and paleontologic studies in Iceland suggest that the "standard" glacial sequences of the Alps, northwestern Europe, and north-central United States may record half or less of the glacial and interglacial episodes that took place during the Pleistocene Epoch. D. M. Hopkins, of the U.S. Geological Survey, and Thorleifur Kinorason, of the University Research Institute of Iceland, demonstrated a record of at least 10 and perhaps 12 distinct glaciations, separated from one another by interglacial episodes, in the course of their study during the summer of 1964 of Tjornes, northern Iceland, and of several other Icelandic localities. The tillites and intervening lavas upon which this record is based all lie above and are younger than marine beds containing an early Pleistocene marine fauna, according to determinations by P. F. MacNeil, paleontologist, U.S. Geological Survey. Paleomagnetic studies by R. R. Doell, of the U.S. Geological Survey, suggest, though, that the oldest glaciation took place at least 1.8 million years ago and more likely on the order of 2.5 to 3.0 million years ago.

JAPAN

Geophysics

Geophysical investigations of calderas in Japan are currently being made by H. R. Blank, Jr., U.S. Geological Survey, in cooperation with Japanese scientists. Airborne-magnetometer surveys of 18 calderas or calderalike structures were carried out during the first half of 1964 by a private company under contract to the U.S. Geological Survey. The data show that many of the calderas are located on discontinuities in the Neogene or pre-Neogene basement complex; in the case of Aso caldera, the discontinuity is interpreted as a west-northwest-trending fault which controls the alignment of post-caldera central cones. A number of volcanic bodies in the target areas produce negative aeromagnetic anomalies, owing to hitherto unsuspected inverse polarization. The polarization is attributed to

⁶⁰ A. L. Washburn and Minze Stuiver; 1962. Radiocarbon-dated post-glacial delevelling in northeast Greenland: *Arctic*, v. 15, p. 66–73.

thermal remanant magnetization acquired during the most recent geomagnetic field reversal, thus setting a minimum age for the bodies. Active solfataras also produce negative anomalies in some cases. The latter may result from destruction of magnetic minerals by chemical action, together with elevation of temperatures above the Curie point, in zones located within strongly positive regions.

LEEWARD ISLANDS

Paleontology

The first fossil evidence for an Eocene age of deposits on the island of Nevis, Leeward Islands, was obtained from a sample submitted by C. O. Hutton, of Stanford University. The sample, consisting of limestone fragments, contained two species of large Foraminifera identified as *Amphistegina* cf. *A. parvula* and *Helicostegina* cf. *H. dimorpha*. Both of these species are common in middle Eocene deposits elsewhere in the Caribbean and Gulf coast regions.

SINO-SOVIET GEOLOGIC STUDIES

Sino-Soviet geologic atlases

The U.S. Geological Survey, on the behalf of Advanced Research Projects Agency, is preparing interpretative geologic atlases of the Sino-Soviet bloc countries that will provide the basic earth-science information needed to discriminate between manmade and natural seismic events in these countries. One atlas, at a scale of 1:5,000,000, will depict the general geologic environment, major tectonic features, thickness of the crust, and regional seismic characteristics of these countries. Final completion is expected by September 1965. Another atlas, at a scale of 1:250,000, is being prepared of the Ulan-Ude quadrangle, a highly seismic area in southeastern Russia that is potentially suitable for underground clandestine nuclear tests. This atlas is the first of a series in which the basic information on the geology, hydrology, soils, vegetation, and cultural features of such areas is being compiled and its potential for nuclear test sites evaluated. The Ulan-Ude quadrangle will be completed by June 1965.

Coal production in China and USSR analyzed

According to a study by Paul Averitt,⁶¹ coal production in Communist China increased very rapidly in the 1957–60 period as a result of technical help

from the USSR. As a result, production exceeded that of the United States in 1960. Following 1960, the production dropped markedly, probably as a result of withdrawal of the technical assistance. Coal production in the USSR increased rapidly in the period 1952–58, and was leveled off since then, probably as a result of the increased production and use of petroleum.

STUDY OF IRAZU VOLCANO, COSTA RICA

Major damage to the economy of Costa Rica was caused by the 1963–65 eruption of Irazú Volcano (figs. 6 and 7). The erupted ash ruined pastures in the heart of the nation's dairy country, destroyed a substantial part of the coffee crop, and contaminated the water supply of major cities. Accelerated runoff of rainwater from the ash-covered slopes of the volcano caused destructive floods, mudflows, and landslides.

In March 1964, a 2-year project for the scientific study of Irazú and other active Costa Rican volcanoes (fig. 8) was started jointly by the Government of Costa Rica and by the Agency for International Development, U.S. Department of State. Two Costa Rican and two U.S. Geological Survey geologists were assigned to the project full time; more than a dozen others provided part-time assistance on different phases of investigation. The work was begun as an emergency project under the direction of K. J. Murata. H. R. Blank, Jr., established gravity stations. H. H. Waldron was

FIGURE 6.—Aerial view from the eastern (windward) side of the Irazú Volcano, which is explosively erupting steam and ash. The highest point (11,260 ft) on the volcano is the somma ridge on the left side of the photo. The ridge is 390 feet above the ephemeral lake of Playa Hermosa.

⁶¹ Paul Averitt, 1964, Recent trends in coal production in the USSR and Communist China: *Econ. Geology*, v. 59, no. 2, p. 323–324.

FIGURE 7.—View of erupting Irazú Volcano from the south-west flank, about 3 miles from the summit. A vertically rising column of explosively ejected steam and ash appears on the right skyline. A column erupted a few minutes earlier is being blown westward (left) by the prevailing wind and is dropping out much coarse ash.

responsible for engineering studies, W. D. Cardwell undertook water-resources investigations, and S. J. Gawarecki and R. M. Moxham directed remote-sensing infrared surveys.

Precise geodetic survey

Measurement of upheaval or tilting of ground in the summit region has been used as a means of monitoring the waxing and waning of magmatic pressure in active Hawaiian and Japanese volcanoes. The technique was applied to erupting Irazú through precise determinations of levels on bench marks set along the highway to the summit; the first survey in May 1964 showed that the upper half of the volcano upheaved 11 centimeters above levels determined originally in 1949. A repetition of the survey in September 1964 disclosed a subsidence of the volcano to about the 1949 configuration, indicating a drastic drop in magmatic pressure. Activity declined gradually thereafter, and ash eruption ceased completely in February 1965.

Gravity survey

Gravity observations at 51 monumented stations along the precise level line extending from Cartago to the summit of Irazú Volcano were made in June 1964, September 1964, and February 1965 in an attempt to measure any gravity changes that may have been associated with eruptive activity during these intervals. It had earlier been established in Mihara Volcano, Oshima Island, Japan, that gravity changes of as much as 0.5 milligal during a single

FIGURE 8.—Sketch map of Costa Rica and its volcanoes. 1, Turrialba; 2, Irazú; 3, Barba; 4, Poas; 5, Arenal; 6, Tenorio; 7, Miravalles; 8, Ricon de la Vieja; and 9, Orosi.

eruptive episode could be correlated with inflation and deflation of the volcanic edifice, as reported by I. Hayakawa in 1952. The gravity changes at Irazú were too large to be attributed to the elevation changes alone and were presumably caused by shallow movement of magma. However, no time-dependent changes exceeding the maximum uncertainty in the reduced gravity-station values, or about 0.1 mgal, were detected in surveying the Irazú level line. The absolute accuracy of the gravity measurement must be considerably improved, and the surveys must embrace an entire cycle, before the gravimetric method of monitoring volcanic activity can be fully evaluated for Irazú.

Aerial infrared survey

The method of aerial infrared survey which had been used successfully to locate hot spots at Kilauea Volcano, Hawaii, was applied to Irazú and other volcanoes of the Central Cordillera during May 1964. Infrared imagery obtained using a specially equipped airplane of the University of Michigan clearly delineated thermal anomalies (active and dormant craters and fumerolic areas) on Irazú, Poas, and Turrialba Volcanoes.

Several craters exist at the summit of Irazú; of these, only the younger and currently active crater is marked by thermal activity. Infrared imagery

shows the area of this thermal anomaly to be hour-glass shaped and elongate to the east. Fumaroles are present approximately $2\frac{1}{2}$ miles east-northeast and 0.8 mile northwest of the summit; those to the east-northeast form three small clusters, those to the northwest are in a northeasterly trending linear array. A large group of hot springs, located approximately 1.3 miles northwest of the summit, form the source of the Rio Sucio. Additional warm springs are present locally along the course of the Rio Sucio for a distance of several miles from its headwaters.

The two southernmost and youngest of the craters on the summit of Turrialba show fumarolic activity, but no thermal anomalies were detected in the other craters. Analysis of the infrared signals reveals that the fumaroles in the southernmost crater have the highest temperature. No other thermal anomalies were detected in the vicinity of Turrialba.

Hot springs south of Cartago, near the villages of Tobosi and Hervidero, were found to be more abundant than previously supposed. A minimum of 27 were located near Tobosi and 7 in the vicinity of Hervidero.

Geochemical studies

The composition of Irazú magma remained remarkably constant at about 55 percent SiO_2 throughout the eruption. Materials of similar composition have been erupted by other volcanoes of the circum-Pacific region, such as Parícutin (Mexico), Crater Lake (Oregon), and certain volcanoes in Hokkaido (Japan). Petrographically, fresh Irazú bombs were porphyritic labradorite pyroxene-olivine-basaltic andesite. Soluble sulfates and chlorides (up to 0.7 percent SO_3 and 0.2 percent Cl) were invariably present in the erupted ash. These substances were

formed through solfataric alteration of old rocks of the volcano prior to, or during, the eruption and reflect the abundance of fragments of such altered rocks in the erupted ash.

Engineering geology

The December 1963 flood on the Reventado River, on the southwest slope of Irazú, destroyed several hundred homes, a major highway, and a railroad in the city of Cartago. All through the following rainy season (April-November 1964) flash floods of muddy water and boulders continued to menace the city. Also, large landslides began along the banks of the Reventado and its tributaries.

Engineering-geology studies indicated that main remedial measures should be aimed at controlling the runoff of rainwater from the ash-covered slopes of the volcano. A force of 500 men was put to work on the volcano, ditching to lead water away from the slide areas, installing brush dams in gullies, and planting ash-resistant grasses. A bedrock map of the Reventado Valley was prepared for locating a series of check dams along the river.

Ground-water hydrology

The eruption of Irazú Volcano resulted in considerable damage to the water supplies of San Jose and nearby communities. A reconnaissance program by the U.S. Agency for International Development and U.S. Geological Survey showed possibilities for obtaining ground water from eight types of aquifers or water-bearing zones in the Meseta Central Occidental. These aquifers include three kinds of alluvial deposits and interflow zones between lava flows, as well as scoria, cinders, and ash. Further geohydrologic studies and extensive test drilling will be undertaken before large-scale ground-water development is attempted.

INVESTIGATIONS OF PRINCIPLES AND PROCESSES

A substantial part of the Geological Survey research program is primarily topical and involves the application of principles and analytical techniques largely developed in the laboratory to the elucidation of the evolution, composition, and structure of: (1) the earth as a whole, (2) its rocks and minerals, (3) its constituent elements, (4) its waters, and (5) its past and present living forms. The emphasis is upon quantitative measurements as a means of obtaining basic data having genetic significance. For the past several years the scope of the topical studies has been broadened to include investigations of the Moon and of materials of extra terrestrial origin under sponsorship of the National Aeronautics and Space Agency.

The program of topical studies is, by its nature, long term, but it has produced important current benefits. For example, the program has played an integral part in setting up and operating a nationwide nuclear-blast and earthquake-detection system, together with the volcano eruption warning system for the island of Hawaii. Studies of the stability relations and isotopic compositions of minerals have given insight into the ore-forming processes and have provided new guides for finding ore. Many new analytical techniques and methods of wide application, within the Geological Survey and without, have been developed in the fields of wet chemistry, emission spectroscopy, mineralogy, X-ray spectrometry, and the electron microprobe. Analytical services in these fields and in the fields of paleontology and geochronology are provided for the Geological Survey as a whole.

PALEONTOLOGY

Each paleontologist of the U. S. Geological Survey maintains specialized research interests which are applied to biostratigraphic problems in a number of ways. Systematic and paleoecological studies of particular groups of plants or animals have long-term biostratigraphic implications that often are realized years after the primary work is published. On-the-spot consulting services are accomplished by

visits to field mapping projects. These serve to bring the lithostratigraphic and biostratigraphic approaches into harmony by resolving discordant interpretations. Fossils collected by the project geologist, in turn, may lead the paleontologist into peripheral studies that can amplify his own stratigraphic interpretations. Recent paleontologic developments of particular significance, many as yet unpublished, are summarized under the following major taxonomic groups.

FORAMINIFERA

Carboniferous endothyrids from Idaho

Calcareous Foraminifera of Meramec, Chester, and Pennsylvanian ages were identified by B. A. Skipp from samples of the White Knob Limestone collected by W. J. Mapel in the Doublespring quadrangle of central Idaho. One of the forms, *Howchinia* Cushman, 1927, is recognized for the first time in North America. *Howchinia* is associated with the endothyrid foraminifers *Endothyra scitula* Toomey, *Endothyra plectogyra* (E. Zeller), and *Tournayella* spp. which are indicative of Mississippian (Meramec) age. The genus *Howchinia* has previously been reported from the Lower Carboniferous (Viséan) of England and Russia.

Triassic arenaceous forms from Wyoming and Idaho

A small fauna of Lower Triassic arenaceous Foraminifera representing 7 genera and 9 species has been obtained from insoluble residues of the Thaynes Limestone in eastern Idaho and western Wyoming by M. L. Schroeder. Five species of silicified Lower Triassic Foraminifera were previously found in northeastern Nevada by W. W. Schell and D. L. Clark in 1955 and 1956. The microfossils from the Thaynes Limestone were obtained by dissolving more than 2,000 samples of rock from 4 different localities. Several species were found near the top of the Thaynes in the Salt River Range, Lincoln County, Wyo., and relatively abundant Foraminifera were found in the upper part of a 6-foot dark-gray fine-grained carbonaceous limestone bed in the Thaynes in the Caribou Range, Bonneville County, eastern Idaho.

Cretaceous species from Pierre Shale, eastern Wyoming

J. F. Mello's study of samples taken at 50-foot intervals through an approximately 3,300-foot-thick complete section of the Pierre Shale at Red Bird, Wyo., indicates that the foraminifera from almost all the samples are agglutinated forms and are few in number of specimens and species. Because most species occur in many consecutive samples, the fauna has a conservative aspect. The long ranges of many species, and the general similarity of foraminiferal assemblages throughout the section, indicate similarity of environment for the entire Pierre Shale. The predominantly arenaceous nature of the faunas suggests deposition at inner neritic depths.

LOWER INVERTEBRATES**Cambrian archaeocyathid from Alaska**

A. R. Palmer reports an Early Cambrian archaeocyathid unlike any previously known form from a collection made by E. E. Brabb along the Tatonduk River, east-central Alaska. The organism has a linear series of barrel-shaped segments ornamented with longitudinal rows of pustules. Near each joint is a collar with a ring of long, slender spines curved parallel to the long axis. In cross section, both the "barrels" and the spines have the double walls and radiating partitions characteristic of archaeocyathids.

Devonian rugose corals, eastern North America

Studies by W. A. Oliver, Jr., of colonial rugose corals of Schoharie and Onondaga age from eastern North America, have shown that most of the genera that are conventionally considered to belong to the Disphyllidae actually comprise ptenophyllid as well as disphyllid species. Species previously assigned to genera on the basis of colony form and gross structure are being revised on the basis of detailed analyses of internal morphology.

Mississippian chaetetid from Kentucky

Material collected by R. D. Trace from the so-called *Chaetetes*—algal horizon near the base of the Kinkaid Limestone in Crittenden, Dixon, and Caldwell Counties, Ky., has been studied by H. M. Duncan who reports that the chaetetid hydrozoan in this faunal zone is a lamellar form named *Chaetetella*. This lamellar chaetetid is particularly characteristic of upper Viséan and lower Namurian rocks in Russia and Great Britain. According to Trace, it is a useful guide fossil inasmuch as it almost invariably occurs within 5 to 10 feet of the base of the Kinkaid.

Echinoderm ecology

P. M. Kier (U. S. National Museum) and R. E. Grant have completed a paper on the distribution and living habits of 17 species of echinoids in the Key Largo Coral Reef Preserve of Florida (a Federal and State undersea wildlife preserve). They report that echinoids live on all kinds of substrates, and that some are highly specialized for the particular kind of environment inhabited. Four species live on dead coral rocks (but two of these also live on sand). Two species live under slabs of rocks; 5 species burrow in sand; 3 species live in grass on sand; and 1 species moves in groups on the surface of grassless sand at relatively great depth. Because many details of distribution and behavior of echinoids are reported for the first time, these results should be useful in interpreting the habits of fossil species.

Bryozoans of Kentucky Upper Ordovician

Study of bryozoans from the Upper Ordovician of Kentucky by O. L. Karklins has resulted in the revisions of the generic assignment of some species, in changes of stratigraphic ranges, and in finding of some new species and, possibly, genera. About 30 genera have been identified from a continuous 240-foot rock sequence in Mason County, Orangeburg and Maysville East quadrangles. Recognizable changes in assemblages suggest that the bryozoans range from late Eden to early Richmond age. Another section, about 210 feet thick, along the Mason-Lewis County line in the same quadrangles, has a bryozoan assemblage of 19 genera, suggesting a Richmond age. An assemblage of 15 genera was found in a 125-foot section in Madison County, Richmond North quadrangle. This bryozoan assemblage resembles bryozoans occurring in the lower half of the section in Mason County. These three sections are invaluable for local and regional correlations of the Upper Ordovician.

Middle Ordovician of Nevada

Physical and paleontologic stratigraphy of the lower Middle Ordovician strata of southern Nevada and adjoining California indicates certain modifications in zonation and correlation, according to R. J. Ross, Jr. (1-64). Only the *Orthidiella* and *Anomalorthis* Zones of the Whiterock Stage appear to have significant time value. The *Palliseria* Zone may be partially controlled by lithic facies and the *Rhyostrophia* Zone is thought to be post Whiterock.

Ordovician brachiopods in Kentucky

Kentucky Ordovician brachiopods are becoming better known through R. B. Neuman's study of silici-

fied specimens, providing firmer bases for correlation than is possible with less well preserved material. For example, *Onniella fertilis* (Bassler) seems to be restricted to the lower part of the Lexington Limestone, where the species is abundant, and the large form *Heterorthis bassleri* (Foerste) is characteristic of the middle part of the Lexington. Higher in the succession, the traditional Richmond guide fossils, *Leptaena richmondensis* (Foerste) and *Platystrophia cypha* (James) are as abundant in some beds traditionally dated as Maysville in age as they are in those dated as Richmond. Studies of the brachiopod *Platystrophia* indicate a clear difference at the species level between collections from Maysville and Richmond beds. Specimens collected by G. C. Simmons from beds of Richmond age in the Palmer quadrangle are specifically distinct from *Platystrophia ponderosa*, which characterizes the Maysville strata in the Richmond North quadrangle.

Mississippian brachiopod zonation in Oklahoma and Missouri

Study by Mackenzie Gordon, Jr., of the Mississippian brachiopods of the Tri-State lead and zinc mining district, in connection with the investigations of E. T. McKnight and R. P. Fisher in the Wyandotte quadrangle, Oklahoma and Missouri, has led to the recognition of six major brachiopod zones in the pre-Chester Mississippian rocks of that region. More than 140 species of brachiopods occur in these rocks, including approximately 35 undescribed species. The major faunal zones can be recognized along the length of the southern flank of the Ozark uplift and permit correlation with the type section in the upper Mississippi River valley.

Evolution of brachiopod superfamily Stenosismatacea

A study of this conservative group of Paleozoic brachiopods, the Stenosismatacea, by R. E. Grant (1-65) shows that internal structures of the earliest Middle Devonian species differ but slightly from structures of species that lived near the end of the phyletic line in the Late Permian. On the basis of moderate, but significant, changes in external morphology, the superfamily has been divided into 2 families, 4 subfamilies, 11 genera, and more than 200 species. The group is retained within the Rhynchonelloidea, and stratigraphic implications of phylogenetic trends are indicated.

MOLLUSCA

Stratigraphic distribution of *Matthevia*

Specimens of *Matthevia*, a genus that has defied precise classification, are widespread in the Western

United States, according to E. L. Yochelson, J. F. McAllister, and Anthony Reso (p. B73-B78). Characteristically, specimens are silicified and may be identified on the outcrop by the presence of two narrow deep cavities in a solid pyramidal mass. The genus seems to be confined to rocks of Late Cambrian Trempealeau age. In the Funeral Mountains, Death Valley region, California, and in Pahrnagat Range, southern Nevada, extensive fieldwork has demonstrated the stratigraphic utility of *Matthevia* in correlating sections of complexly faulted rocks.

First belemnites from Permian Phosphoria Formation

Examination by Mackenzie Gordon, Jr., of slender rodlike longitudinally striated fossils from the Phosphoria Formation, previously thought to have been spines of fish, shows them to be belemnoid guards of the aulococerid type. They are the hard parts contained within the bodies of a specialized group of squidlike animals, the first Permian ones known in the United States. In fact, only one Permian species has been described previously, from the Upper Permian of Greenland. The Greenland species appears to be represented in the Upper Permian Retort Member of the Phosphoria Formation in western Montana. Another species, in the Lower Permian Meade Peak Member of the Phosphoria in southeastern Idaho, belongs in an undescribed genus and species. It is of considerable interest that in the Upper Permian these free-swimming organisms were common to Greenland and Montana waters.

Triassic bactritid cephalopods found in California

A slender transversely ribbed bactritid cephalopod is under study by Mackenzie Gordon, Jr. It was found by N. J. Silberling in a collection from the Upper Triassic (Karnian) Hosselkus Limestone of California. As the bactritids are supposed to have become extinct at the end of the Paleozoic, this Late Triassic form is of unusual interest from an evolutionary viewpoint because it extends the range of another fossil group into the Mesozoic.

New zonation of Triassic Shublik Formation, Alaska

Study of extensive fossil collections from the Shublik Formation in the Mt. Michelson quadrangle and adjacent areas, northern Alaska, by N. J. Silberling permits recognition of possibly eight different ages based mainly on pelecypods and cephalopods. These faunal subdivisions range from Ladinian through latest Triassic and, apparently, include a full development of Karnian and Norian assemblages.

New Jurassic pelecypod from Alaska

Ralph Imlay has identified the pelecypod *Otapiria* from collections obtained from the topmost part of the Shublik Formation in northern Alaska. The species present is similar to species from the basal Jurassic of New Zealand and of northeastern Siberia. The occurrence of *Otapiria*, plus the presence of *Inoceramus*, which is not known below Jurassic, indicates at least an Early Jurassic age for the top of the Shublik.

Late Tertiary gastropod evolution in Florida

Recent fieldwork by Druid Wilson has established the sequence of evolution in a group of fossils belonging to the gastropod subgenus *Fasciolaria* (*Cinctura*). The group commonly occurs in the upper Tertiary and Pleistocene of Florida. The progressive loss of sculpture of the early whorls from Caloosahatchee time through Fort Thompson time can be used for age determination in the field. A late Miocene species is related, but is not within the evolutionary sequence of the later occurring group.

Fresh-water mollusks from Idaho

In the course of studies of the molluscan fauna of the Pliocene and Pleistocene Glens Ferry Formation, southern Idaho, D. W. Taylor observed color pattern preserved in some of the shells. Suspecting that color pattern in the calcareous layers of the gastropod shell is a primitive character, Taylor surveyed several families to find out what color patterns exist and how variable they are. Virtual uniformity of pattern exists within groups that can be defined on grounds other than color pattern; and the groups are of useful size, generally at the family or subfamily level. Taylor has identified a Glens Ferry genus, previously considered extinct, as probably surviving in Yunnan Province, China. The number of related groups of fresh-water mollusks (both Tertiary and Recent) in northeastern Asia and northwestern America is mounting steadily. Faunally, the lake most similar to the Pliocene and Pleistocene lake in which the Glens Ferry was deposited, is Lake Biwa, Japan. Some of the poorly known lakes in Yunnan also have related molluscan assemblages.

Pacific coast Pleistocene paleoecology

W. O. Addicott has completed a study of late Pleistocene marine mollusks of the central-California-Oregon area and their paleozoogeographic and paleoecologic significance. Assemblages from the lowest marine terrace of the southwestern Santa

Cruz Mountains in California (lat 37° N.) show many similarities to mollusks whose modern ranges overlap in southern British Columbia (lat 48° N.) The late Pleistocene assemblages of northern California and Oregon collectively constitute a faunal province that is analogous to the modern Aleutian molluscan province (southern boundary lat 48° N.), and that had its southern boundary between lat 35° N. and 37° N. This suggests a substantially cooler marine hydroclimate from central California to Oregon in the late Pleistocene than at present.

ARTHROPODA

Alaskan Cambrian trilobites

Study of stratigraphically controlled collections from Cambrian rocks of the Alaska-Yukon boundary area shows the presence of at least nine distinct trilobite faunas representing all major parts of the Cambrian Period. More than 100 trilobite species have been identified and described by A. R. Palmer. The oldest fauna is a rich assemblage of trilobites almost entirely of Siberian aspect. A second Early Cambrian fauna has largely North American elements. The Middle Cambrian faunas have affinities with faunas from the southern Cordilleran region, Kazakhstan, eastern Australia, and boulder conglomerates in eastern Canada. The Late Cambrian faunas have affinities to faunas from the same regions and, in addition, some genera previously have been known only from China. Included are the first North American representatives of the widespread Asiatic genus *Proceratopyge*, the Chinese genus *Yupingia*, and other Asiatic forms. These trilobite assemblages are vital links in intercontinental correlations between Asia, Australia, and North America for the middle part of the Upper Cambrian.

Restudy of Silurian ostracode genus

The type specimens of ostracode species assigned to the genus *Beyrichia* by Ulrich and Bassler (1923) have been restudied by Jean Berdan, who finds that they belong to at least 3 genera (2 of which are new) and 2 subfamilies. These genera seem to have fairly restricted ranges and may prove useful in Silurian biostratigraphic studies.

Classification of ostracode superfamily Healdiacea

The shape of carapace, muscle-scar pattern, and inferred nature of soft parts of the fossil superfamily Healdiacea suggest that it should be referred to the order Podocopida, according to I. G. Sohn (p. B69-B72). On the basis of muscle-scar pattern, the genus *Pseudophanasymmetria* is assigned to the

Healdiacea and the generic diagnosis is emended to include dimorphism. *Pseudophanasymmetria? afoveata* is considered to be a dimorphic form of *P. foveata*.

Revision of ostracode families

J. E. Hazel has discovered, by dissection of living representatives, that the ostracode subfamily Campylocytherinae and the *Jugosocythereis-Hermanites* complex of genera, which are normally placed in the Trachyleberididae, are actually representatives of the Hemicytheridae. This has allowed the construction of a more valid phylogenetic classification for the important Hemicytheridae-Trachyleberididae subfamilies.

Oligocene crabs from Alaska

Two fossil crabs, previously known only from the Oligocene or late Eocene, were identified by F. S. MacNeil from beds of the Orca Group, Prince William Sound area, Alaska. These strata are in an area thought previously to consist entirely of upper Mesozoic rocks.

VERTEBRATES

Miocene horses from Panama

Additional collection in the Canal Zone has yielded two horse molars, representing the first Miocene horses to be found in Central America. F. C. Whitmore has identified two genera of forest-living, browsing horses from material collected in Panama in January 1965. *Anchitherium* and *Archaeohippus* are forms related to other Miocene species throughout the world. The species of *Anchitherium* is closest to a species from the lower Miocene of the High Plains. The *Archaeohippus* is most like species found in the late lower Miocene of Texas and late lower to early middle Miocene of Florida. These horses were three-toed, had low-crowned teeth, and probably subsisted for the most part on soft vegetation. Such food was probably also eaten by the other known members of the fauna whose remains are found in the Cucaracha Formation on the banks of the Panama Canal: the rhinoceros *Diceratherium*, an oreodont resembling the North American genus *Brachycrus*, and an unnamed artiodactyl most closely resembling members of the primitive family Protoceratidae. Presence of horses of this type in the Cucaracha fauna reinforces the earlier conclusion that, in the middle and probably in the early Miocene, Panama was connected to North America by a broad land mass and was separated from South America. Apparently, a single

continuous fauna lived along the northern and western shores of the Gulf of Mexico during the Miocene.

Locomotion of *Paleoparadoxia* analyzed

C. A. Repenning has made considerable progress in the preparation of the *Paleoparadoxia* specimen from the Stanford Accelerator Site, at Palo Alto, Calif. An analysis of its means of locomotion suggests that, because the heavy hind legs are short and the hind feet are obliquely articulated to the tibia, the animal could not stand erect. *Paleoparadoxia* must have moved on land with flexed legs in a ponderous hopping gait, much as do modern sea lions. In the water, the hind legs were probably used in a froglike kicking motion. Although relatively less shortened, the stout front legs also were modified toward paddling and were ill-adapted to walking.

Big Bone Lick site, Kentucky, dated

Matrix associated with the type skull of the extinct muskox *Bootherium bombifrons* from Big Bone Lick, Ky. was submitted by F. C. Whitmore to Meyer Rubin for age determination by C-14 dating and E. B. Leopold for pollen studies. Rubin reports a date of 17,200 years B. P. (climax of Tazewell ice advance), and Leopold found a preponderance of spruce pollen. The C-14 date is the oldest yet found at Big Bone Lick and indicates the probable existence in the area of a fauna older than those so far collected.

Giant Pleistocene cat found in Alaska

F. C. Whitmore has identified a jaw bone, probably washed out of peat or silt overlying high-level gravels on Lost Chicken Creek near Cathedral Bluffs, central Alaska, as *Felix atrox*, the lion-sized Pleistocene cat especially well-known from the Rancho la Brea tar pits of California. The fossil, collected by H. L. Foster, substantiates reports of large cat bones previously found in Alaska, although there are no published reports of the occurrence of the species this far north. The previous known range included California, Nevada, Idaho, Arizona, Texas, Nebraska, and Maryland.

PALEOBOTANY

Carboniferous iron bacteria identified

The structural remains of fossil Coal Measures bacteria may be resolved clearly by means of the electron microscope, using replicas prepared by delicate oxidation etching of polished pyrite. Two species of iron bacteria closely allied with distinctive modern forms have been identified by J. M. Schopf.

Alliance of the fossil iron bacteria with modern types strongly suggests a similar ecology. Specific microenvironments of Pennsylvanian age must have been similar to those which are widespread at the present time. Additional information is provided on specific deposition of authigenic pyrite and the origin of these common impurities in coal and other sediments. The specific nature of the pyrite also affects the process of Recent weathering that results in an acidic condition in ground water derived from coal-mine refuse areas. The micrographic replica method evidently may be extended with considerable benefit to the microbiology of other types of ancient deposits.

Callipterid reproductive methods studied

S. H. Mamay has completed research describing a new Permian plant genus *Tinsleya*, including the first demonstration of female reproductive parts among the callipterids. These findings have direct bearing on the question of pteridospermous reproduction and may be significant in problems of cycad and angiosperm evolution.

Early Tertiary wood from Atlantic Continental Shelf

A specimen of dicotyledonous wood dredged up by a fishing boat in the Georges Bank area of the Continental Shelf off New England has been identified by R. A. Scott. The specimen, submitted by J. C. Hathaway, represents the extant genus *Euptelea*, now growing in Japan, China, and parts of India. The only previous fossil record of the genus in North America is from the early Tertiary of Oregon. The Atlantic occurrence is evidence for widespread distribution of *Euptelea* on this continent during early Tertiary, or possibly Late Cretaceous time.

Cenomanian floras in southwest Alaska

Large collections of fossil leaves, made by J. M. Hoare during the summer of 1964, indicate the presence of Albian rocks on Nunivak Island and probable Cenomanian rocks on Nelson Island. According to Jack Wolfe's analysis, the collections from Nelson Island represent the first probable Cenomanian floras south of the Brooks Range; they are dominated by flowering plants and have some similarities to Cenomanian floras of the Atlantic Coastal Plain.

Pierre Shale palynomorphs from Wyoming studied

The pollen and spore flora of the Pierre Shale at Red Bird, Wyo., has been the subject of a study by E. B. Leopold and Bernadine Tschudy. The sequence has been dated by W. A. Cobban and J. R. Gill as late Campanian to early Maestrichtian in age, according

to baculite evidence. Though these marine sediments are poor in plant microfossils, a total of 191 forms were found. These include: 68 spore, 15 gymnosperm, 5 probable monocot and 45 dicot types, and 14 dinoflagellate algal, 18 hystrich, and 26 miscellaneous planktonic species. The seed plants include several groups whose modern relatives live in Australasia in the warm temperate and subtropical zones. Among the dicotyledons, 14 forms are assigned to the genus *Aquilapollenites* Rouse which, as far as is known, represents an extinct group. The fern types include several with living counterparts in the New World tropics. The flora seems to show little qualitative change through the late Campanian and early Maestrichtian; the primary changes are in pollen abundance in the sediments. The land-based flora of the marine Red Bird section corresponds to the pollen and spore flora described by K. R. Newman⁶² from the continental deposits of the Mesa-verde Group near Craig, Colo.

Cretaceous pollen in Appalachian Mountains

Samples of a black sandy clay from the Appalachian Mountains near Chambersburg, Pa., have yielded a suite of Late Cretaceous pollen and spores, according to R. H. Tschudy (p. B64-B68). Evidence from known stratigraphic ranges of some forms, and from the number and percentage of triplicate pollen, indicates an extreme possible age range from early Turonian through middle Campanian. A probable age range for the deposit is late Turonian through early Campanian. This locality is about 80 miles west of the nearest known Upper Cretaceous outcrops.

Megaspores from Tuscaloosa Group

R. H. Tschudy reports the identification of the megaspore genus *Ariadnaesporites* in a sample from the Tuscaloosa Group from Alabama. Structurally similar, but much smaller, spores were found tangled in the long hairlike processes of the megaspores. Tschudy believes them to be microspores of *Ariadnaesporites* and the tangling to be an adaptation to promote fertilization of the megagamete. This genus was reported previously only from an Upper Cretaceous coal in Greenland.

Mummified microfossils from Green River Formation

Continued study of the fresh-water microorganisms found in the rich oil shales of the Eocene Green River Formation of Wyoming by W. H. Bradley has revealed the presence of several genera of micro-

⁶² K. R. Newman, 1964, Palynologic correlations of Late Cretaceous and Paleocene formations, northwestern Colorado: Soc. Econ. Paleontologists and Mineralogists Spec. Publ. 11, p. 169-180.

scopic aquatic fungi. Several of these are specifically indistinguishable from those living today in the lakes of the Lake District of England. Others are specifically unlike any known living aquatic or terrestrial fungi, yet the genus is fixed beyond doubt. More remarkable as microfossils are the well-preserved remains of one rotifer, *Cephalodella* sp., and two midge larvae that must have just emerged from their eggs at the time they died. All these microfossils are unmineralized and, in reality, are mummified. This sort of preservation for the rotifer and the two midge larvae is remarkable because such delicate animal tissue ordinarily decays within hours of death.

Reclassification of diatoms

In connection with the preparation of the section of Diatomacea for the *Treatise on Invertebrate Paleontology*, K. E. Lohman has devised a new classification of the diatoms, consisting of 1 class, 4 orders, 11 suborders, 23 families, 54 subfamilies, and 320 genera. This reclassification includes all of the diatoms, living and fossil, marine and nonmarine.

Miocene diatoms from Washington

An extensive study by K. E. Lohman of diatoms from the Latah Formation near Spokane, Wash., has pointed up correlations with other middle and late Miocene deposits in the West. The material, collected by J. W. Hosterman, contains a number of species in common with the Latah Formation near Mica, Wash.; the Virgin Valley Beds of Humboldt County, Nev.; and the Mascall Formation of Central Oregon. The diatoms suggest that the Latah near Spokane was deposited in a temperate, fresh-water lake having a pH ranging from 6 to 8.

MARINE GEOLOGY AND HYDROLOGY

The marine program and related activities of the U.S. Geological Survey emphasize: (1) mapping and evaluation of the geologic environment and resources of the sea floor adjacent to the United States, (2) topical studies of geologic and hydrologic processes and environmental factors both near the coasts and in deep-sea areas, and (3) detailed investigations of oceanic islands under the jurisdiction of the United States. The major effort has continued to be concentrated on regional studies of the Atlantic Continental Shelf and Slope. Following the Alaska Earthquake of 1964, investigations of earthquake-triggered submarine slides and their effects were undertaken as part of the Geological Survey's contri-

bution to the disaster program. (See the section "Alaska Earthquake 1964.")

ATLANTIC CONTINENTAL SHELF AND SLOPE

During 1964, the 5-year reconnaissance studies of the east-coast continental margin by the Geological Survey entered their third year. These investigations, directed by K. O. Emery, of the Woods Hole Oceanographic Institution, are carried on in collaboration with the institution and with the U.S. Bureau of Commercial Fisheries. Nearly continuous spring, summer, and fall cruises were undertaken to collect systematically surficial sediments and rocks from all parts of the shelf and slope and to obtain sub-bottom structural profiles of the Gulf of Maine and adjacent portions of the shelf and slope. Complementary laboratory and office studies have continued to emphasize the compilation and evaluation of existing geologic and hydrologic data for the entire shelf and the analysis and interpretation of new samples and data from the Gulf of Maine and adjacent shelf.

Potential for new oil discoveries increases seaward

Chiefly on the basis of records and samples obtained from State geological surveys and from oil companies, J. C. Maher has prepared a longitudinal cross section and a series of transverse cross sections of the Atlantic Coastal Plain from Key West, Fla., to Long Island, N.Y., showing correlations of the subsurface stratigraphy. The sections clearly illustrate the seaward thickening of the formations and the progressive lithologic changes from nearshore to offshore marine facies. They provide a basis for correlation of rocks beneath the continental shelf with those beneath adjacent land and support the suggestion that the potential for discovery of new petroleum reserves improves seaward along the Atlantic coast.

Contour of floor of Gulf of Maine reflects basement rocks

On the basis of preliminary results of continuous sub-bottom profiling (sparker) surveys in the Gulf of Maine, Richard Tagg and Elazar Uchupi have concluded that unconsolidated sediments are generally less than 100 meters thick and that major topographic irregularities of the gulf are related to corrugations within the basement rocks. Triassic rocks, which form the basement in the northeastern part of the gulf, extend 120 kilometers westward from the Bay of Fundy and have been eroded, partly by glacial scour, to depths of 300 m and more below present sea level. Where the Triassic rocks are

deeply eroded, glacial and Recent sediments exceed 100 m in thickness.

Reworking of glacial sediments greatest north of Cape Cod

J. C. Hathaway, Jobst Hulsemann, J. S. Schlee, and J. V. A. Trumbull report that the amount of apparent reworking of unconsolidated glacial sediments differs in areas of similar topography in various parts of the Gulf of Maine and is greatest in and near the Wilkinson and Murray Basins, north of Cape Cod, Mass. R. M. Pratt has found that nearshore gravels consist chiefly of rocks from adjacent shores, whereas gravels of the central and southern gulf area have a more diversified lithology, probably caused by ice rafting from several source areas.

Oyster shells from shelf are 8,000–11,000 years old, according to C-14 dating

In the course of sediment sampling on the shelf and of dredging by the U.S. Bureau of Commercial Fisheries, shells of shallow-water (less than 40 meters) oysters have been recovered from depths of 145–165 m at 71 localities between North Carolina and Cape Cod. Radiocarbon measurements on 4 shells indicate ages of 8,000–11,000 years. A. S. Merrill, K. O. Emery, and Meyer Rubin have concluded that the oysters lived in lagoons or estuaries that were submerged when sea level rose at the end of the latest glacial epoch.

Gravels off New Jersey are potential construction material

J. S. Schlee (1-64), on the basis of textural analyses of sediment samples off the coast of New Jersey, has outlined an extensive area of Pleistocene sandy gravels and pebbly sands at depths of less than 40 meters. The sands and gravels provide a potential source of construction material for adjacent coastal areas. Iron staining of the grains suggests that the deposits are relict.

Salinity and temperature variations in Maine estuaries

R. H. Meade has collected hydrologic data in several areas where the existing data were scarce. Salinity and temperature profiles of the Penobscot and Kennebec River estuaries were made in July. One 12-hour anchor station on the Penobscot River at Winterport recorded a tidal range in surface salinity of 4.3 to 15.1‰. Two 12-hour stations in September near Georges Bank, one in Great South Channel and the other in Northeast Channel, also recorded tidal variations in salinity and temperature. In Northeast Channel, salinity ranged from

31.8 to 34.7‰ during one tidal cycle; the temperature of the bottom water during this time remained nearly constant at 7.1°C.

MARINE GEOLOGIC ENVIRONMENT AND PROCESSES

Cores from experimental Mohole relate fossils and vulcanism

A study of the composition and mineralogy of late Tertiary sediments, recovered from the experimental deep-sea Mohole drill hole (Guadalupe site), allowed quantitative interpretation of the relations between biogenic opal, biogenic calcite, and lithogenic matter. Large amounts of trachytic ash, rhyolitic ash, or decomposition products of basalt in a few samples record eruptions of nearby oceanic and continental volcanoes. Volcanic activity seems to influence organic productivity as suggested by the tendency of sediments rich in remains of siliceous and calcareous organisms to have above-average proportions of basaltic materials (K. J. Murata and R. C. Erd, 1-65).

Eocene Radiolaria from Puerto Rico Trench

K. N. Sachs, Jr., has identified Eocene Radiolaria from a dredge sample recovered from the north slope of the Puerto Rico Trench at a depth of 3,650 to 3,150 fathoms. The radiolarian fauna contained in one of four samples submitted by R. L. Chase, of Woods Hole Oceanographic Institution, has many elements in common with that from the Eocene of Saipan, described by W. R. Riedel.⁶³

Organic content of deep-sea sediments in North Pacific

J. G. Palacas and V. E. Swanson have analysed the organic matter from segments of three cores from the North Pacific Ocean, provided by G. W. Moore and collected during the 1963 cruise of the USCGS ship *Pioneer*. The clayey sediment contains 250–3,000 parts per million humic acids, 55–200 ppm amino acids, and 30–40 ppm bitumens. These 3 fractions account for more than 50 percent of the organic matter in gray clay from a depth of 6,800 meters in the Aleutian Trench, but for less than 25 percent in tan siliceous clay from a depth of 5,400 m in the central North Pacific.

FRESH-WATER–SALT-WATER MIXING

Sporadic low-oxygen concentrations, Delaware River estuary

In summarizing 14½ years of monthly measurements of a 44-mile tidal reach of the Delaware River

⁶³ W. R. Riedel, 1957, Eocene Radiolaria in *Geology of Saipan, Marianna Islands*, pt. 3, Paleontology: U.S. Geol. Survey Prof. Paper 280-G, p. 257–263, pls. 62, 63.

estuary W. B. Keighton has noted occasional low dissolved-oxygen concentrations, especially in that reach of the river adjacent to Philadelphia, Pa. At the Benjamin Franklin Bridge, Philadelphia, a third of the samples were less than 30 percent saturated with oxygen.

Biota zones due to offshore ground-water discharge, Florida

In studies of fresh ground-water discharge to the sea, F. A. Kohout and M. C. Kolipinske demonstrated that biological zonation in Biscayne Bay is related to ground-water discharge. Fresh ground water flowing seaward through the upper part of the Biscayne aquifer discharges through the bottom sediments of the bay in a littoral zone that extends about 400 feet seaward from the shoreline. Brackish-water species of both plants and animals predominate near shore and are progressively replaced seaward by typical marine species. The zonation of different sea grasses is recognizable by color differentiation on aerial photographs.

WATER MOVEMENT IN ESTUARIES

Theoretical analysis of flow in estuaries

Using the nonlinear hyperbolic partial-differential equations representing one-dimensional homogeneously dense unsteady flow in open channels, R. A. Baltzer and Chintu Lai have found it possible to model analytically the transient streamflow observed in rivers and some estuaries. Numerical solution techniques premised on power-series methods, characteristic methods, and implicit methods provide both accurate and economical means for determining transient-flow information when evaluation is carried out using a high-speed digital computer.

BIOHYDROLOGIC STUDIES OF ESTUARIES

Phytoplankton and tides, Duwamish River estuary, Washington

The diurnal variability of phytoplankton population (as indicated by diatom and chlorophyll concentrations) in the Duwamish River estuary adjacent to Puget Sound is inversely related to tidal stage, according to E. B. Welch and J. F. Santos. The relation is less clear at high than at low river discharges. High plankton concentrations at low tide (2 to 3 times the high-tide concentrations) appear to be a result of high velocities near the river bottom that carry benthic diatoms into suspension. This theory is supported by the improvement of the relation between tidal stage and plankton population as sampling depth increases.

Variation in attachment organisms, Patuxent River estuary, Maryland

In a 2-year study of attached marine organisms in the Patuxent River, R. L. Cory has observed that generally the attachment sequence is the same but that numbers of individuals and total biomass vary from station to station and year to year. At six of the sites, ash-free dry weight (organic carbon), a measure of the productivity of attached organisms, was significantly higher in 1964 than in 1963. At one site it was almost the same as in 1963, and two sites, both in the vicinity of a new powerplant, showed nearly a 50-percent reduction in biomass.

Cory believes that the higher biomass production over the entire estuary, with the exception of a 1-mile stretch in the vicinity of the powerplant, is probably due to the decrease in rainfall in 1964 which resulted in slightly higher salinities and a decrease in nutrient flushing time. The decrease in production at the plant site was probably due to higher turbidities observed during most of the summer while the intake and effluent channels were being dredged.

OCEANIC ISLANDS

Marine geologic studies, Palau to Guam

At the invitation of the U.S. Coast and Geodetic Survey, Gilbert Corwin and J. I. Tracey, Jr., participated in marine geological studies near the Palau Islands and Guam aboard the USCGS ship *Pioneer* on her return from the Indian Ocean. The studies included bottom sampling and geophysical measurements while underway across and along the arcuate ridges from which the islands rise and across and along the associated trenches. Preliminary analysis of sparker-probe measurements adjacent to the Palau Islands appears to confirm the importance of northwest-southeast faulting in development of the islands, and suggests the presence of at least three major longitudinal faults on the west side of the Palau trench. The identification of westward-dipping sediments on the east side of the trench and of their termination within the trench presents problems that merit further investigation.

Moderately indurated calcareous silt dredged from the west slope of the Palau Islands and poorly indurated limestone with volcanic pebbles from the east slope appear to be deposits of Pleistocene age that lack counterparts on the islands. If Pleistocene, these samples represent a potential source of information on intervals of otherwise unrecorded

local geologic history when sea level was lower than at present.

Samples of andesitic pumice, which included one piece with a 20-mm glomerocryst of calcic plagioclase and olivine, was dredged from the crest of a submarine volcano 20 miles northwest of Guam. The lack of appreciable alteration and the nature of the pumice suggest that it is the product of a Recent eruption and that the source magma was of the northern Mariana (high-alumina) type rather than southern Mariana (tholeiitic) type. Late Quaternary ash eruptions of the volcano may explain the presence of volcanic detritus in limestone of the northern plateau of Guam and the origin of the overlying thin lateritic soils.

On Guam, Corwin and Tracey discovered a deeply weathered volcanic plug near the center of the southern volcanic area of the island. The plug appears to be one of several probable sources of the patchy uppermost lava flows of Guam (Dandan flows of Miocene age).

ASTROGEOLOGY

The U.S. Geological Survey investigations in support of the space exploration program of the National Aeronautics and Space Administration include: (1) earth-based lunar investigations, (2) studies of terrestrial and experimental impact and cratering phenomena, (3) cosmochemistry and petrology, (4) unmanned lunar exploration studies, and (5) manned lunar exploration studies.

EARTH-BASED LUNAR INVESTIGATIONS

Lunar geologic mapping

Geologic mapping of the Moon at a scale of 1:1,000,000 forms the base for the lunar investigations of the U.S. Geological Survey. Figure 9 shows the status of this mapping program. Maps completed to date cover about 3 million square miles.

Major structure of Mare Vaporum region interpreted

A preliminary geologic map of the Mare Vaporum region, which lies within the peripheral belt of

FIGURE 9.—Index map of the Moon, showing status of geologic mapping at a scale of 1:1,000,000. ACIC number and name are given in each region.

structures related to the Mare Imbrium basin, has been completed by D. E. Wilhelms (U.S. Geol. Survey, 1-65, pt. A). Major structural elements within the region, mare depressions and terra high, are interpreted as synclines and anticlines, which are roughly concentric with the Mare Imbrium basin and in places are modified by faults. Most of the terra in the region is deformed by troughs and ridges of the "Imbrian sculpture" radial to the basin and is covered by the Fra Mauro Formation, which is interpreted as ejecta from the Mare Imbrium basin. Some of the structure may have been produced by interaction of deforming forces at the time the Mare Imbrium basin was formed on older structures related to the basin of Mare Serenitatis. Still other structures not obviously related to the Mare Imbrium basin, such as the Triesnecker rille system, may have been formed by isostatic readjustment of the lunar crust subsequent to the Imbrian event. Part of the Hyginus rille is aligned with Imbrian sculpture.

Stratigraphy of western Theophilus area unravelled

A complex stratigraphic sequence has been mapped in the terra of the western Theophilus region by D. J. Milton (U.S. Geol. Survey, 1-65, pt. A). Crater rims and other terrain cut by prominent scarps (Imbrian sculpture) trending northwest-southeast are believed to be pre-Imbrian. Terrain with a much finer texture of small rounded hills and valleys elongated in the same direction is correlated with the Fra Mauro Formation of Apenninian age. This is overlain by a smooth-surfaced plains-forming unit that partially fills basins. Adjacent higher areas with subdued relief are probably underlain by a thinner deposit of the same material. This material is tentatively considered to be of Apenninian age; it may, however, be Archimedian. A plateau-forming unit has a broadly rolling smooth surface with some steep-walled irregular depressions. This unit, in contrast to the plains-forming unit, does not tend to occupy basins but forms broad plateaus with the underlying relief completely concealed. The age of the plateau-forming unit is uncertain; it may be an age equivalent of the plains-forming unit.

Stratigraphic relations in Grimaldi region

Preliminary geologic mapping of the Grimaldi region by J. F. McCauley (U.S. Geol. Survey, 1-65, pt. A) establishes the stratigraphic relations among the regional materials surrounding the Mare Orientale and Mare Humorum basins. North of the crater Fontana, smooth-textured high-albedo material of the Cordillera Group (associated with the

Orientale basin) overlies darker, rougher textured material of the Humorum Group, which clearly is younger than the Humorum basin. Very dark mare material, extensive within the area, is characterized by a low spatial density of craters and a general lack of superimposed rays, suggesting that it may be younger than the mare material typical of the Procellarum Group.

Complex sequence of events recorded in Pitatus region

S. R. Titley (U.S. Geol. Survey, 1-65, pt. A) has mapped the stratigraphic and structural relations in the Pitatus region and adjacent areas. A complex sequence of events in this region includes basin formation, regional deposition, and tectonic activity. Certain critical structural and stratigraphic relations permit tentative interpretations of relative ages of the Mare Humorum, Mare Nubium, and Mare Imbrium basins. Ejecta(?) from Mare Humorum basin make up most of the old upland exposures around the west edge of Mare Nubium and, in the northwest part of the quadrangle, may underlie the Fra Mauro Formation, which consists of ejecta from the Mare Imbrium basin. Structural relations on the west edge of the quadrangle suggest that the circumferential structures about Mare Humorum are older than those about Mare Nubium.

Possible volcanic complex near Harbinger Mountains

A possible volcanic complex near the Harbinger Mountains of the Moon has been investigated by H. J. Moore (U.S. Geol. Survey, 1-65, pt. A). Abundant hills, craters, and rilles are present just west of the Harbinger Mountains. The close association of the craters and rilles suggest that the area is a volcanic complex. These lunar craters resemble terrestrial maars and rampart craters, and one is like a caldera. Lunar craters that are similar to maars and rampart craters are larger than their suggested terrestrial counterparts. Some of the sinuous rilles are crater chains; not all can be the result of erosion by nuées ardentes as suggested by other workers.

Hummocky deposit about Mare Humorum basin interpreted

S. R. Titley and R. E. Eggleton (U.S. Geol. Survey, 1-65, pt. A) have described an extensive hummocky deposit around the Mare Humorum basin. This material, which is uniform in its subdued topographic characteristics and normal albedo, is distributed about Mare Humorum. It occurs now as isolated patches and tongues but is believed to have surrounded Mare Humorum at one time as a nearly continuous blanket. In many respects it is

similar to hummocky Fra Mauro Formation. The extent and distribution of the hummocky material about Mare Humorum is comparable with those of the Fra Mauro Formation, when the relative sizes of the Humorum and Imbrium basins are considered. The material has all the characteristics of an extensive ejecta blanket and appears to be derived from the Mare Humorum basin.

THEORETICAL ANALYSES OF LUNAR PHENOMENA

Trajectories for Copernicus ray material calculated

Trajectories of objects producing Copernicus ray material on the crater Eratosthenes have been studied by M. H. Carr (U.S. Geol. Survey, 1-65, pt. A). In the region of Eratosthenes, areas lacking ray material are principally on slopes facing away from Copernicus. On the assumption that the absence of ray material in these areas is due mainly to topographic shielding, trajectories were computed for objects ejected from Copernicus to produce secondary craters and rays in Eratosthenes. Mean values of 7° for the ejection angle 1.1 kilometers per second for the ejection velocity were found.

Isostatic rebound of lunar craters interpreted

Harold Masursky (U.S. Geol. Survey, 1-65, pt. A) has reported on the isostatic rebound of lunar craters and a reinterpretation of Ptolemaeus. The rebound of lunar craters similar in size to Ptolemaeus indicates that the cover on the floor of the crater may be very thin; that is, only thick enough to have buried the central peaks. The present shallow depth of the crater is attributed chiefly to respringing of the crater floor.

Impact temperatures around Timocharis calculated

The history of the temperature regime around the crater Timocharis was computed by M. H. Carr (U.S. Geol. Survey, 1-65, pt. A), assuming an impact origin for the crater and shock equations of state similar to basalt for the target and projectile. For impact velocities greater than 8 kilometers per second, temperatures sufficiently high for melting extend a distance of 6 km below the crater. At lower impact velocities, melting temperatures are not reached. The transient high temperature source at the surface, immediately after impact, has an insulating effect, so that lunar internal heat cannot escape to the surface. This raises temperature at depths far below the impact-induced temperature anomaly. Lowering of the thermal conductivity near the surface by impact brecciation also has an insulating effect. The temperature rise at depth may be

great enough to produce melting and volcanism long after the impact event.

INSTRUMENTATION AND TECHNIQUES FOR LAND-BASED LUNAR STUDY

Excess infrared emission at lunar limb studied

Infrared emission from the illuminated Moon is being investigated by Kenneth Watson (U.S. Geol. Survey, 1-65, pt. A). Edison Pettit and S. B. Nicholson⁶⁴ observed that the infrared emission of the Moon near zero phase angle at the limb along the lunar equator exceeds the emission to be expected from a Lambert emitter. Pettit and Nicholson suggested that this excess emission at the limb is due entirely to the lack of shadows at full Moon; only illuminated surfaces contribute to the emission. This cannot be valid at the scale of lunar mountains and valleys, however. The average slopes at this scale are much too low to produce appreciable departure from Lambert emission in the infrared. At the other extreme or scale, reradiation between inclined surfaces less than a few centimeters apart also tends to produce uniform emission that follows the Lambert law. The observed excess emission of the limb may be caused by either (1) roughness on a scale between a centimeter and meter, and (2) non-Lambert emission for individual surface elements. In the first case, roughness may possibly be related to the high spatial density of secondary craters on the lunar surface.

Study of lunar polarimetry begun

D. E. Wilhelms (U.S. Geol. Survey, 1-65, pt. A) has initiated a program of lunar polarimetry with a Lyot visual polarimeter mounted on the 12-inch refractor of Lick Observatory. Preliminary results indicate that polarization of areas on the lunar surface 8 kilometers in diameter can be measured with these instruments under conditions of good viewing.

New high-speed camera reduces shutter errors

Existing high-speed photographic systems employed in lunar photographic photometry have, for the most part, contained large inherent errors. These errors can be traced to (1) shutters with variable speeds and (2) processing systems that employ non-uniform agitation of developing solutions. H. A. Pohn (U.S. Geol. Survey, 1-65, pt. A) has designed a high-speed photometric camera that largely overcomes shutter errors by adjustable slits that are pulled along mated Teflon ways. This system is coupled with a plate-processing system that holds the

⁶⁴ Edison Pettit and S. B. Nicholson, 1930, Lunar radiation and temperatures: *Astrophys. Jour.*, v. 71, p. 102-135.

internal variations in plate response to about 3 percent of mean plate density. A camera with the new design features will be used on the Geological Survey's 30-inch telescope to make photographs for a photoelectrically calibrated photographic atlas of the Moon.

TERRESTRIAL AND EXPERIMENTAL IMPACT AND CRATERING PHENOMENA

The U.S. Geological Survey program of study of natural and manmade terrestrial craters throws light on the origin of their structure, stratigraphy, and mineral composition and, what is perhaps more important, also provides clues to the makeup and origin of lunar craters.

FIELD STUDIES OF IMPACT STRUCTURES

Ray crater in Australia mapped and interpreted

D. J. Milton and F. C. Michel (p. C5-C11) have mapped the only terrestrial meteorite impact crater known to have rays of ejecta similar to those around lunar craters. Crater 3 of the Henbury crater field, Northern Territory, Australia, is a roughly circular crater about 200 feet in diameter with a relief of 15 feet from floor to rim crest and 5 feet from the surrounding surface to the crest. It is formed in alternating beds of well-cemented sandstone and weak shale dipping about 35° and overlain by a varying thickness of pediment gravel. Beds striking into the walls of the crater are deformed, predominantly by folding about almost vertical axes.

Specific types of ejecta when traced to specific outcrops of source beds in the crater permitted the directions and distances of throw-out to be measured. From these data the distribution of impact energy was calculated. The results showed that the effective energy decreased as a function of the radius of the shock-wave front at a greater rate than would be predicted from a simple shock-propagation model probably because the finite strength of the bedrock is not considered in such a model.

Rim of Flynn Creek cryptovolcanic structure studied

D. J. Roddy (U.S. Geol. Survey, 1-65, pt. B) found that the southeastern rim of the Flynn Creek structure, Tennessee, a feature of suspected impact origin, consists of a structurally complex sequence of Upper Ordovician rocks and contains what appears to be a part of a land surface that existed immediately before the deformation. A large mass of breccia covers the ancient land surface and is preserved in a downfaulted block. Field and laboratory studies

of the rocks underlying this surface strongly suggest that they are Late Ordovician in age and can probably be assigned to the Richmond Group. Calculations of the stratigraphic thicknesses in the southeastern rim suggest that not more than 60 meters of Late Ordovician strata was removed by erosion after the structure was formed and before later sediments of Late Devonian age were deposited. The maximum age of the Flynn Creek structure is therefore Late Ordovician. The minimum age is early Late Devonian.

A preliminary search did not reveal the high-pressure silica polymorphs coesite and stishovite that have been found in some known meteorite impact structures. Geophysical studies by Shawn Biehler and D. J. Roddy indicate that there is no large magnetic anomaly, as might be expected if an igneous plug were present below the structure.

LABORATORY STUDIES OF IMPACT-RELATED PHENOMENA

Stishovite synthesized by experimental shock metamorphism

P. S. DeCarli (U.S. Geol. Survey, 1-65, pt. B), Stanford Research Institute, and D. J. Milton separated small amounts of stishovite from explosively shocked sandstones, novaculite, and single-crystal quartz. Estimated peak pressures for the syntheses ranged from 150 to 280 kilobars and shock temperatures from 150° to 300° C. Coesite was not detected in any sample. This suggests that quartz can invert during shock to a short-range-order phase with sixfold coordination, a small part of which may develop the long-range order of stishovite. Part of the short-range-order phase may invert to coesite during the more protracted decay of the pressure pulse through the stability field of coesite accompanying meteorite-crater formation.

High-index plagioclase glass from meteorite described

M. B. Duke (U.S. Geol. Survey, 1-65, pt. B) found the average composition of maskelynite (natural plagioclase glass) from the Shergotty meteorite to be about An_{49} . Its density varies within a range near 2.57 grams per cubic centimeter. The refractive index ranges from 1.539 to 1.554 and is distinctly higher than the indices of synthetic plagioclase glass in the same compositional range. Experiments show that the refractive index can be markedly reduced by heating for a few minutes at temperatures as low as 450°C. The glass is believed to have been formed by solid-state transformation under shock pressure.

Argon loss from shocked granite, Ries crater

P. Signer (U.S. Geol. Survey, 1-65, pt. B) compared the argon content of unshocked and strongly shocked granite from the Ries crater, Germany, and found that the shocked granite contains far less radiogenic argon than the unshocked. New amorphous phases formed by shock absorb much greater quantities of atmospheric argon than their parent crystalline phases.

Two types of "shock" minerals from natural craters noted

E. C. T. Chao (U.S. Geol. Survey, 1-65, pt. B) described selective shock transformation of minerals in rocks from known and probable meteorite craters and showed that their special characteristics may be used as evidence of meteorite impact. The transformations are of two types: (1) from a crystalline phase of complex chemical composition to glass of the same composition and (2) from a crystalline phase of simple chemical composition to its high-pressure and high-density polymorph. Transformations of the first type, formed at high pressure and probably low temperatures, occurred in impact-metamorphosed sandstones from the Henbury craters of Australia and in biotite granite, gneiss, and schist in suevite from the Ries crater. Transformations from a low-density to a high-density polymorph, such as from quartz to coesite or stishovite, occurred (1) in impact-metamorphosed sandstones of Meteor crater, Ariz., and the Wabar crater, Saudi Arabia; (2) in biotite granite gneiss of the Ries crater; and in (3) metamorphic rocks of the Lake Bosumtwi crater, Ghana.

Experimental impact studies using high-energy missiles

A study by H. J. Moore, Reuben Kachadoorian, and H. G. Wilshire (U.S. Geol. Survey, 1-65, pt. B) of craters produced by the impact of missiles provides data on cratering by projectiles with large kinetic energies (2.3×10^{15} ergs). The amount of material displaced during crater formation by projectile impact is nearly the same as the amount of material displaced during crater formation by chemical explosives with shallow depths of burial when the kinetic energy of the projectile (corrected for the angle of impact) is equal to the equivalent energy of TNT of the chemical explosive.

Strain gages measure shock-wave speeds in sandstone

M. H. Carr (U.S. Geol. Survey, 1-65, pt. B) used strain gages to measure shock pressures and shock-wave speeds in sandstone. The Hugoniot curve of Coconino Sandstone for pressures from 0 to 150

kilobars shows shock-wave speeds from 3.0 to 4.0 kilometers per second. For similar shock energies in sandstone from the Moenkopi Formation, an elastic precursor with a speed of 3.1 km/sec and a shock wave with speeds from 2.5 to 2.9 km/sec are generated.

Crater shapes reflect porosity and cohesion of target

H. J. Moore, D. E. Gault, E. D. Heitowit, and R. V. Lugn (U.S. Geol. Survey, 1-65, pt. B) found that the shapes of craters produced by hypervelocity impacts differ according to the porosity and cohesion of the target. Depth-to-diameter ratios increase from 1 : 5 for nonporous targets (basalt) to about 1 : 1 for porous targets (pumice). Craters in weakly bonded sand targets are larger than those in sandstone and basalt, and craters in fine-grained weakly bonded sand targets are smaller than those in coarse-grained sand. Application of a defect theory of strength explains the differences between crater size and shape in sandstone and those in weakly bonded sand targets. The shapes of recent small lunar craters may be used to estimate cohesion of lunar surface materials.

Significant parts of the ejecta from craters in pumice are complex aggregates of crushed pumice and glass fibers. The aggregates resemble volcanic ash. It is postulated that, as the result of micro-meteoroid bombardment, the uppermost part of the lunar surface could be similar to volcanic ash.

COSMOCHEMISTRY AND PETROLOGY

Studies in cosmochemistry and petrology include (1) geological distribution and physical and chemical properties of tektites, (2) investigations of micro-analytical techniques for analyzing milligram and submilligram quantities of tektites, meteorites and, eventually, returned lunar samples, and (3) petrology and chemistry of meteorites.

Aerodynamically modified moldavite described

E. C. T. Chao⁶⁵ found indirect evidence of aerodynamic heating and ablation on a moldavite from Slavice, Czechoslovakia. No evidence for heating or ablation of moldavites had previously been recognized. The indirect evidence consists of the keel left by spallation of the heated and strained glass of the leading surface of a teardrop-shaped tektite. The keel is identical to those of australite indicators (partially spalled tektites). This is the first evidence of the probable entry velocity of moldavites into

⁶⁵ E. C. T. Chao, 1964, Spalled, aerodynamically modified moldavite from Slavice, Moravia, Czechoslovakia: *Science*, v. 146, No. 3645, p. 790-791.

Earth's atmosphere which must have been greater than about 5 kilometers/second to cause ablation.

Age and abundance of Australasian tektites

Field geologic studies of Australasian tektite occurrences by E. C. T. Chao (U.S. Geol. Survey, 1-65, pt. C) indicate that they formed since middle Pleistocene. The tektites are found primarily in laterite soils, where transportation probably has been minimal, and in gravel deposits, where reworking and transportation are probable. The maximum number of tektites in the Australasian strewn field is estimated to be about 2×10^{13} on the basis of their areal density in places where reworking is improbable. From estimates of the number of tektites concentrated by fluvial transportation in the Manila Bay area of the Philippine Islands, the minimum number of tektites is estimated to be about 4×10^9 for the entire strewn field. No terrestrial impact crater capable of producing the Australasian tektites has been discovered.

Analyses of small tektite samples by two spectrographic methods

C. S. Annell (U.S. Geol. Survey, 1-65, pt. C) analyzed 1-2-milligram samples of tektites for Al, Ca, Fe, Mg, and Ti by emission spectrography with both the rotating-disk-high-voltage-spark method and the gas-controlled-arc methods. The two solution methods give similar precision and accuracy, but the rotating-disk method is 3 to 5 times more sensitive and has wider application to the study of small quantities of silicate minerals. Working curves for the spectrographic determination of Na and K by the rotating-disk-high-voltage-spark method were also obtained. As little as 0.1 percent Na and 0.3 percent K now can be determined in silicate samples weighing 1 to 2 mg.

Mo, As, and Ir in tektite spherules imply meteoritic origin

Preliminary applications of the laser microprobe for emission spectrographic analysis and neutron-activation analysis of spherules from Philippine tektites by Frank Cuttitta, E. C. T. Chao, and C. S. Annell (U.S. Geol. Survey, 1-65, pt. C) indicate the presence of siderophile and chalcophile elements such as Mo, As, and Ir. These minor elements are generally much more abundant in iron meteorites than in most silica-rich rocks. The qualitative neutron-activation analysis gives further indication of the meteoritic origin of metallic spherules in tektites.

Older analyses for alkalis and TiO_2 of tektites in error

D. B. Tatlock (U.S. Geol. Survey, 1-65, pt. C) compared 55 older analyses of Australasian tektites

with 110 modern, precisely monitored analyses and found that more than half of the older alkali and titania determinations are decidedly inaccurate and misleading. Deviations of the older analyses from the restricted values of the modern analyses are comparable to the imprecisions shown by early analyses of G-1 granite and W-1 diabase. This suggests that a high percentage of older alkali and titania analyses, such as those of Washington's tables, are of questionable quality.

Luminescence of achondritic forsterite and synthetic diopside

A mineral that luminesces red under ultraviolet radiation in the Norton County, Kans., enstatite achondrite meteorite has been identified as forsterite by M. B. Duke and B. A. Tryba (U.S. Geol. Survey, 1-65, pt. C). Low forsterite content in the meteorite means that the luminescent efficiency of the total meteorite is very low. This appears to eliminate ultraviolet-excited enstatite achondrite as a possible lunar red fluorescent material. Electron-excited luminescence of synthetic diopside shows the major role played by minor-element content of the mineral. Pure diopside luminesces blue; Mn-bearing diopside, red; Ti-bearing, orange; and Ni-bearing, green.

Thermomagnetic studies of tektites

In continuation of studies of the magnetic properties of tektites, F. E. Senftle, A. N. Thorpe, and S. R. White (A. N. Thorpe and F. E. Senftle, 1-64) have made thermomagnetic experiments on tektites to establish the state of the iron in the glass. Metallic iron spherules can be formed from the dissolved iron in any tektite heated under strong reducing conditions to temperatures above $1,530^\circ\text{C}$ (melting point of iron). If the tektite is heated to lower temperatures under reducing conditions for several days, subsequent magnetic measurements show that the iron in solution, mainly as Fe^{+2} , is not reduced by this treatment. However, the magnetism of the surface regions of the tektite changes from paramagnetic to ferromagnetic after heating. The only reasonable explanation of this phenomenon is that a small amount of particulate iron present before heating, aggregates by a diffusion process to form ferromagnetic centers during heating. Further work is being done to see if the particulate matter is submicroscopic metallic spherules or an iron compound, as this is germane to the origin of tektites.

Electrical resistivity and viscosity of tektite glass

A. F. Hoyte, F. E. Senftle, and P. W. Wirtz (1-65) measured the electrical resistivity of 25 tektites from 7 different strewn fields as a function of temperature

over the interval 330–450°K, using a time-of-discharge technique. From these data they determined activation energy for each specimen, and constructed a viscosity curve for the temperature region between 1,370° and 1,770°K. The calculated curve agrees within the limits of error with the experimental curves for several examples. The method can be used to calculate the viscosity within half an order of magnitude of the correct value. Mean viscosity curves have been constructed from the experimental data for tektites from the seven known strewn fields (A. F. Hoyte, F. E. Senftle, and P. W. Wirtz, 1–65).

UNMANNED LUNAR EXPLORATION STUDIES

The unmanned lunar exploration studies are concerned primarily with the scientific exploitation of the imagery received from the unmanned spaceflight projects of the National Aeronautics and Space Administration. The work includes the development of flight-operations procedures for spacecraft television systems and techniques by which the data received can be reduced to provide detailed topographic information, terrain maps, and large-scale geologic maps of the lunar surface.

RANGER TELEVISION INVESTIGATIONS

Rangers VII, VIII, and IX have contributed much important new information on the fine structure of the lunar surface. The wealth of photographic data acquired (over 17,000 pictures) will require several years of systematic analysis. Craters are the dominant small relief features; small positive relief features on the lunar surface, other than crater rims, are relatively rare. Ranger-VII pictures contributed to a better understanding of the nature of the lunar rays. The ray areas have a high density of irregularly shaped, shallow secondary craters probably formed by the low-velocity impact of materials thrown out of larger primary craters. The bright walls of these telescopically unresolvable craters contribute to the high albedo of the rays and, in addition, the rim materials of the secondary craters have a higher albedo than the adjacent parts of the maria.

Craters less abundant than impact origin predicts

On the basis of the observed crater distribution on the lunar maria, the present rate of meteorite infall on the earth, and information on the distribution and age of ancient impact craters and eroded structures of possible impact origin on the earth, E. M. Shoemaker has constructed a detailed model for the fine structure of the maria. The present number of small primary craters on the Moon is considerably lower than the number expected to have been formed

by meteorite impact. The expected number of primary craters of a few meters in diameter and larger is more than two orders of magnitude greater than the observed number. A similar deficiency exists for small secondary craters.

It is impossible to observe the number of small craters of both primary and secondary impact origin expected to have been formed on the lunar maria. The total area of expected craters larger than 1 meter across is more than 10 times the total area of the mare surface. Small craters are lost by superposition of other craters of the same size or larger. In addition, the repeated formation of small craters obliterates larger ones. This interaction of craters produces a layer of shattered and pulverized rock that covers most of the mare. The layer is of varying thickness and rests with an irregular contact on the underlying mare material. Fragments in this layer have been derived for the most part by ejection of material from nearby craters, but some of the material has been derived from great distances. The grain size of the debris layer tends to decrease from bottom to top, because the fragments in the upper part have been shocked and broken a greater number of times during the formation of the more numerous smaller craters. The porosity of the debris layer may be as much as 90 percent at the surface, but it probably decreases rapidly at depth because of compaction by repeated impact events.

Ranger photograph permits detailed slope measurements

H. A. Pohn and L. C. Rowan made detailed slope measurements of the last Ranger-VII P3 frame using a modification of photoclinometric techniques employed in lunar terrain analysis. The calibrated transfer characteristics of the Ranger camera system were used, along with an adopted photometric function, to determine slopes from the density of the Ranger photographs. Topographic profiles were then constructed from the slope measurements, and the profiles were adjusted to a common datum plane to prepare a contour map. The final map shows an area approximately 30 by 25 meters at an approximate scale of 1:80 with a contour interval of 10 centimeters. It represents the most detailed information available on the distribution of lunar slopes and has been used to study problems of the landing dynamics of both manned and unmanned spacecraft.

SURVEYOR TELEVISION INVESTIGATIONS

The Surveyor television investigations by E. C. Morris, H. E. Holt, and R. M. Batson have been concerned with the development of test and calibration procedures for the Surveyor cameras and the

development of techniques for extracting photogrammetric, photometric, and geologic information from the television pictures to be received from the system on the lunar surface. This work is required to insure that the television system on the Surveyor spacecraft will function as an effective geologic tool and that the maximum amount of information about the surface of the Moon will be acquired from each successful flight.

Prototype of Surveyor television camera tested

A field test of a prototype Surveyor television camera was conducted during the summer of 1964 on the Bonito lava flow in Sunset Crater National Monument, 15 miles northeast of Flagstaff, Ariz. Experience gained in this test led to the development of an efficient command sequence for mission operations. The pictures obtained have been used to develop specialized cartographic techniques for handling, processing, and mosaicking the imagery. Photogrammetric analysis of the imagery obtained indicates that distance measurements can be made to an accuracy of ± 3 percent at a distance of 20 meters from the spacecraft; this figure may be exceeded by later flight-configured cameras.⁶⁶

Specialized instruments have been designed to aid in the geologic analysis and photogrammetric reduction of the Surveyor television imagery. These include a specialized stereoscope to view spherical mosaics prepared from large numbers of narrow-angle pictures taken by the Surveyor cameras and a special stereoranging instrument to be used as a focus analyzer or as a stereoplotter during mission operations.

LUNAR ORBITER INVESTIGATIONS

Kenneth Watson has solved the general equations for photoclinometric mapping of the Moon's surface from spacecraft imagery, and a computer program that will permit automatic digital production of topographic maps is being written. With these results and appropriate instrumentation, it will be possible to produce base maps very rapidly for terrain and geologic analysis from lunar Orbiter imagery.

LUNAR TERRAIN ANALYSIS

Slope measurements lead to classification of lunar surface

Lunar terrain analysis by J. F. McCauley and L. C. Rowan has led to a quantitative morphologic

classification of the lunar surface.⁶⁷ An automated photometric measuring technique developed by the Geological Survey was used to measure some 40,000 slope components in the east-west direction from the best available earth-based photographs. Slopes of 1 kilometer base length are measured. These measurements have been used as the basis for classifying the lunar surface within the equatorial belt ($\pm 10^\circ$ latitude, $\pm 60^\circ$ longitude). The primary terrain characteristics used for classification purposes have been the average or median slope and the 10th-percentile and the 90th-percentile slopes. The lunar equatorial belt was divided into 6 terrain units at a scale of 1:2,000,000. The smoothest of these units comprises relatively featureless parts of the maria that have a median slope of about 1.5° or less. The roughest terrain unit is in the lunar highlands and has a median slope of about 6° . This terrain is comparable to very rough terrestrial mountain areas such as the Sierra Nevada, in California. The terrain map has been used to help select the target areas for Rangers VII and VIII and for Surveyor test flights.

Extrapolation techniques have been used for the purpose of predicting the average slope of lunar terrain units over a broad range of slope lengths. The predicted median slope of the smooth mare terrain unit for slopes with a base length of 1 meter was within 1° of the median slopes measured with photoclinometric techniques from the last photograph obtained from Ranger VII.

STUDIES RELATED TO MANNED LUNAR EXPLORATION

Studies are being carried out to prepare for geological exploration of the lunar surface by manned space flight. The work includes geological training of the astronauts and investigation of methods of geological and geophysical exploration to be used by the astronauts on the lunar surface.

ASTRONAUT TRAINING

Program of instruction: first year

In July 1963, the U.S. Geological Survey, in cooperation with the Manned Spacecraft Center of the National Aeronautics and Space Administration, began a program to train astronauts in geology. The purpose of the program is to teach the astronauts to carry out geological studies on APOLLO missions. In the first year 13 classroom lectures were

⁶⁶ U.S. Geological Survey, 1964, Surveyor television investigations, Surveyor television experiment, field test no. 2, December 1964, Final report.

⁶⁷ U.S. Geological Survey, 1964, Terrain analysis of the lunar equatorial belt, July 1, 1964: Preliminary report.

given under the direction of E. D. Jackson on terrestrial and lunar aspects of the following subjects: geologic processes, geologic principles, stratified rocks, geologic structure and landforms, geologic mapping, and geophysics. Four field trips were held under the direction of A. H. Chidester.

At the Grand Canyon, in Arizona, training was concerned principally with field techniques and with the fundamentals of geology, such as the law of superposition, and laws of truncation and intersection. A field trip to west Texas provided training in observing and measuring stratigraphic sections and in mapping complexly folded and faulted rocks; the astronauts were also introduced to a variety of volcanic rocks. In the Sunset crater area, near Flagstaff, Ariz., the astronauts received training in hand-specimen petrology, morphology, stratigraphy, and structure of basaltic cinder cones and associated lava flows. A night at Kitt Peak Observatory, near Tucson, Ariz., was spent in lunar observations and mapping. At Philmont Ranch, New Mexico, the earlier training was applied in mapping exercises in three areas of varied geologic character, and an introduction was given in field geophysical techniques.

Program of instruction: second year

In the second year of the training program, classroom lectures were given by visiting experts in volcanic geology, impact structures, shock metamorphism, and lunar mapping. Field trips were oriented chiefly toward providing experience in a wide variety of volcanic geology and in crater geology, with emphasis on petrologic and stratigraphic relations, structural features, constructional forms, and genetic history. At Newberry crater, Oregon, under A. C. Waters, the astronauts studied (1) a basaltic shield volcano with a summit caldera in which the final stage of eruption was rhyolitic; and (2) the flanking fields of cinder cones on the shield, which are structurally controlled. At Valles caldera, New Mexico, under R. A. Bailey, they studied geologic and geophysical features of a caldera with a resurgent structural dome associated with rhyolite ash-flow tuffs. H. A. Powers conducted the astronaut group in a study of the shield volcanoes of Hawaii, and in the geological and geophysical research activities of the Hawaii Volcano Observatory. At the Nevada Test Site, W. J. Carr and R. L. Christiansen led a trip to Timber Mountain caldera, which had a complex history of ash-flow eruption after resurgence of the dome; and to Black Mountain

caldera, which has no resurgent dome, but has instead centrally located constructional volcanic forms. The astronauts studied nuclear craters that are structurally analogous to impact craters, and followed this with a study of Meteor crater, Arizona, all under the leadership of E. M. Shoemaker.

INVESTIGATIONS IN SUPPORT OF MANNED LUNAR EXPLORATION

A manned lunar-exploration investigations group has been organized under D. P. Elston to advise the National Aeronautics and Space Administration on the requirements for geologic exploration of the Moon, to help develop and test operations and instruments of potential use during lunar exploration, and to provide the scientific and technical capability to simulate specific types of lunar surface missions.

Instrument testing in the field

Field tests were directed by H. H. Schmitt and G. A. Swann on the Bonito lava flow and at Meteor crater, in Arizona, to evaluate several possible instrument systems. The following instrument systems were shown to have useful geologic applications in Project APOLLO and possible following stages of lunar exploration: (1) a hand-held surveying staff on which are mounted imaging, tracking, and orientation subsystems; (2) an exploration periscope capable of range, azimuth and vertical-angle determination from the lunar excursion module of the APOLLO spacecraft; (3) a surveillance television system with both automatic and manual tracking of an astronaut on the surface from the lunar excursion module; (4) a staff-mounted television camera capable of transmitting continuous high-resolution images of surface materials; and (5) an instrument and sample carrier for use on the lunar surface.

Field experiments by Yukio Yamamoto with a prototype ranging laser have confirmed the applicability of this instrument for surveying from a lunar roving vehicle.

Testing of survey procedures in the field

During the field tests carried out on the Bonito lava flow and at Meteor crater, data have been obtained that permit both subjective and statistical evaluations of the relation between time required for given geological and surveying operations and the quality and quantity of the data obtained. Motion picture films of these operations have been compiled by H. G. Stephens so that further time and motion studies can be made by interested agencies. Time and information studies conducted to date have led

to the following observations and recommendations:

(1) Pre-flight planning of sample sites, in the absence of an automatic tracking system, is not feasible in heterogeneous terrain because the large amount of time required to locate sample sites would eliminate other high-priority scientific tasks during early APOLLO landings. (2) If samples and photographs of lunar material are returned to earth, geologic description on the Moon should be limited to (a) features too subtle, large, or complex to be illustrated by a sample or photograph, (b) items necessary to place the sample, photograph, or experiment in the context of its environment, and (c) lunar surface features which identify a structural unit for future reference during the lunar mission. (3) Operational flexibility rather than operational programming should be emphasized in the geological training program for the astronauts, as the time and efficiency of description, sampling, and photography largely depend on the nature of the geologic features and will not be completely predictable.

The manned lunar exploration investigations group has assisted the field geology planning team for Project APOLLO in preparing the functional specifications for an APOLLO field-exploration system. This system will provide automatic imaging, tracking orientation, physical-properties sensing, and telemetry subsystems. Such a system will permit the astronauts to use their time on the lunar surface for description of the Moon's fine structure and for the selection of pertinent samples, photographs, and physical-properties measurements, rather than for many routine acts of mapping and data collection.

Development of geophysical instruments and techniques

Investigations under the direction of J. S. Watkins are concerned with devising geophysical instruments and techniques for use by man on the lunar surface and for developing guides for interpreting the data. Field investigations have been conducted at Mono craters, California; the Pisgah crater area near Barstow, Calif.; the S. P. crater flow and the Kana-a flow in the San Franciscan volcanic field near Flagstaff, Ariz.; Meteor crater, Arizona; and in the Hopi Buttes area of the Navajo Reservation, Ariz. Work to date has resulted in the development of a shallow seismic-refraction technique for use in Project APOLLO to determine the thickness and character of the uppermost layer of fragmental material on the lunar surface. Field investigations also have shown that seismic techniques can be used to detect shallow subsurface cavities.

GEOPHYSICAL INVESTIGATIONS

STUDIES OF THE CRUST AND UPPER MANTLE

Continuing interpretation and synthesis of seismic-refraction data from throughout the United States are expanding our knowledge of the variations of crustal and upper mantle structure within and among the various physiographic provinces, and are increasing our understanding of the nature of regional isostatic compensation. It is now possible to publish reasonably reliable maps of crustal and upper-mantle velocity structure in the continental United States (L. C. Pakiser and J. S. Steinhart, 3-64). Large-scale differences extending deep into the earth that are associated with major physiographic features at the earth's surface have been discovered. For example, studies of the rate of attenuation of P-waves that have penetrated the upper mantle support previous conclusions, derived from traveltime data alone, as to differences in the properties of the upper mantle beneath the Central and Western States.

The accumulated evidence shows that the difference between continents and oceanic basins, as well as the difference between active areas and stable areas on continents, clearly is related to the observed seismic characteristics of the crust and upper mantle. Recent analysis of surface-wave data, which has been one of the principal sources of information on mantle structure, indicates that the mantle low-velocity zone for shear waves varies significantly between continents and ocean basins. The evidence available from such studies suggests that the lateral variations in upper-mantle velocity reported by the U.S. Geological Survey in recent years extend to considerable depth, at least several hundred kilometers, and thus are indicative of the deep structure of the planet rather than simply indicating minor near-surface differences.

REGIONAL SEISMIC-REFRACTION STUDIES

Differences in mantle between Eastern and Western United States

A detailed seismic-refraction profile extending more than 1400 kilometers was made from Lake Superior to the vicinity of Denver, Colo., as a part of the international cooperative Lake Superior experiment. This profile contains some of the highest-quality seismic data ever recorded in this distance range. The data show an unusually low attenuation of P-waves that have traveled in the upper mantle, and give no indication of a shadow zone which is usually taken as evidence of a low-velocity

zone in the upper mantle. In fact, the low attenuation with distance implies an increase in velocity with depth sufficient to hold a large fraction of the seismic energy in shallow propagation paths.

By contrast, analysis of data from nuclear and chemical explosions in the Western United States (J. C. Roller, 2-65; Ronald Willden, p. C44-C50) shows a high attenuation of P-waves in the upper mantle and indicates that the mantle in the tectonically active Western United States is distinctly different from the mantle in the stable areas of the east. This difference had been predicted earlier from the traveltimes of P-waves, but the noted amplitude attenuation with distance is an important additional clue to the nature of the rocks composing the upper mantle.

Basalt-filled rift under Lake Superior suggested

Continuing analysis of data collected in the Lake Superior experiment shows that the crustal structure under Lake Superior is probably a continuation of the crustal structure associated with the midcontinent gravity high. This segment of the crust apparently was formed by the intrusion of great masses of basalt into a crustal rift which extended from Lake Superior to the southern limits of the midcontinent gravity high. In addition to the local geologic interest and associated economic implications of this structure, its existence illustrates that anomalous zones can persist in the crust for hundreds of millions of years and has an important bearing on the strength and stability of crustal rocks over long periods of time.

No marked crustal root under Southern Rockies

Two underground mining blasts detonated at Climax, Colo., in May 1964, have led to increased understanding of the nature of isostatic compensation in the Western United States. Interpretation of seismograms recorded at 17 locations in the Front Range uplift of the Southern Rocky Mountains suggests a crustal thickness north of Climax of approximately 54 kilometers, thinning abruptly by about 15 km in the area directly east of the Uinta Mountains (W. H. Jackson and L. C. Pakiser, p. D85-D92). Comparison of these results with those previously reported for the Basin and Range, Colorado Plateaus, Middle Rocky Mountains, and Great Plains provinces reveals that there is no pronounced crustal root under the Southern Rocky Mountains (as might be expected because of their high average elevations) and that isostatic compensation is related primarily to density variations in the upper

mantle and only secondarily to variations in crustal thickness.

Crust under Basin and Range province and Snake River Plain

D. P. Hill and L. C. Pakiser have completed a study of crustal structure along a line that extends northward from the Nevada Test Site across the Basin and Range province and the Snake River Plain to Boise, Idaho. Seismic waves generated by chemical explosions along profiles about 300 kilometers long and by nuclear explosions at the Nevada Test Site along the entire 1700-km-long profile were analyzed. The study indicates that the M-discontinuity is very irregular, with abrupt (possible vertical) changes in depth of several kilometers at several places along the profile. These abrupt changes in depth could be due to faults that extend downward into the upper mantle. Crustal thickness in the Basin and Range province ranges from 25 to 36 km. Crustal thickness in the Snake River Plain ranges from 42 to 49 km. An intermediate crustal layer, probably of basaltic composition, rises from a depth of about 20 km in the Basin and Range province to less than 10 km in the Snake River Plain. Amplitudes of seismic waves generated by the nuclear explosions fall off approximately as the inverse cube of distance, but are scattered through a factor of five. This extreme scatter in amplitudes may be caused by the irregular shape of the M-discontinuity and by complexities in crustal structure.

Crust thickens near Mogollon Rim

Analysis of a set of seismic recordings made at the Tonto Forest Seismological Observatory, Payson, Ariz., shows an abrupt thickening of the crust in the vicinity of the Mogollon Rim (J. C. Roller and D. H. Warren, written communication). The Mogollon Rim is generally considered to be the transition between the Basin and Range province and the Colorado Plateaus province and formerly was generally thought to be an erosional feature. The new seismic evidence suggests that the boundary represents a fundamental lateral change in the crust of the earth.

Break in crustal structure at Walker Lane indicated

An analysis of a seismic profile recorded between Lake Mead, Nev., and Mono Lake, Calif., clearly shows the thickening of the crust from the Basin and Range province to the Sierra Nevada, and suggests a break in crustal structure as the profile crosses the surface feature known as the Walker Lane, according to Lane Johnson.^{67a}

^{67a} Lane Johnson, 1965, Crustal structure between Lake Mead, Nevada, and Mono Lake, California: *Jour. Geophys. Research*. [In press]

High crustal velocities typical of rift zones on Hawaii

During August 1964 seismic-refraction profiles were recorded along the northeast, southeast, and west coasts of the triangular-shaped island of Hawaii. Shots were fired at 10-kilometer intervals from the U. S. Coast Guard cutter *Cape Small*, and were recorded along each coast by 5 refraction units spaced at approximately 25-km intervals. Interpretation of the seismograms by D. P. Hill indicates that the crust is 16 to 18 km thick under the west flanks of Mauna Loa and Hualalai, and 11 to 12 km thick under the northeast and southeast flanks of Kilauea. The crust has an intermediate thickness along the northeast flanks of Mauna Kea and the Kohala Mountains. The velocity of P waves in the upper crust increases with depth from 2.0 to as much as 6.0 km/sec; velocities in the upper crust are generally lower on the flanks of Kilauea than on the flanks of the other volcanoes. Clearly recorded arrivals indicate that a layer with velocities from 6.9 to 7.4 km/sec forms the lowermost 4–8 km of the crust under each of the coasts. The velocity of P_n under each of the coasts is about 8.2 km/sec. Anomalous high crustal velocities are associated with the major rift zones extending from the five volcanoes that form the island. This interpretation of the refraction data is consistent with gravity data on the island of Hawaii.

Seismic network at site of Hebgen Lake earthquake

The Yellowstone Park, Wyo.–Hebgen Lake, Mont. region—site of the major Hebgen Lake earthquake of August, 1959—provides a natural laboratory in which to study the relationships between volcanic activity, recent earthquakes, and surface geology. In the fall of 1963, J. P. Eaton of the U. S. Geological Survey, in cooperation with the National Park Service, initiated a detailed study of earthquakes in and near Yellowstone National Park. Since that time, equipment and operational procedures have been developed that permit continuous recording at remote stations throughout the year, and a network of 3 vertical-component seismometers and 2 three-component seismometers has been established.

Highlights of a preliminary study of earthquakes recorded from December 1964 to March 1965 are: (1) a swarm of about 70 earthquakes originating in the Upper Geyser Basin area near Old Faithful was recorded December 15, 1964; (2) a swarm of about 150 earthquakes originating along the eastern margin of West Yellowstone Basin was recorded February 19–20, 1965; (3) other earthquakes recorded during this period outline a zone of activity

trending roughly east-west from Hebgen Lake to Norris Junction.

Important earthquake zone northeast of Denver indicated

Seismic activity is continuing northeast of Denver, Colo. Geophysicists of the U. S. Geological Survey and the Colorado School of Mines are cooperating in a study of these earthquakes. To supplement the sparse network of seismic stations, the U. S. Geological Survey operated three additional portable seismic systems on a temporary basis during February and March, 1965. A study by Y. Wang of the Colorado School of Mines indicates that the most important earthquake zone is in the Commerce City area northeast of Denver. Hypocenters determined for approximately 150 earthquakes in this area range in depth from 5 to 40 kilometers.

COOPERATIVE EXPERIMENTS IN SEISMOLOGY**USGS takes part in Canadian study of crust in Alberta**

In July 1964, a 500-ton charge of TNT was detonated on the surface of the Watching Hill Range, Suffield Experimental Station, Alberta, Canada, for the Canadian Defense Research Board. The shot was recorded by seven U. S. Geological Survey low-frequency seismic-refraction systems along a line extending from High River, Alberta, in the Great Plains, westward into the Canadian Rocky Mountains at radial distances ranging from 220 to 300 kilometers from the shot. Other Geological Survey units that recorded the explosion were a conventional exploration reflection-refraction system located near Brooks, Alberta, about 100 km west of the shot point, and a prototype slow-speed tape system located about 350 km from the shot point in the Rocky Mountains north of Canal Flats, British Columbia. All seismic systems recorded air arrivals in addition to seismic arrivals. The work was conducted in cooperation with Dr. G. L. Cumming, associate professor of geophysics, University of Alberta, to assist in his study of the earth's crust in the Great Plains and Rocky Mountains of Alberta.

Crustal structure in Oklahoma under cooperative study

U.S. Geological Survey personnel also participated in a cooperative seismic study of the earth's crust and upper mantle in central Oklahoma as a joint Geological Survey–Advanced Research Projects Agency (ARPA) contribution to the transcontinental geophysical survey of the upper mantle project. The study was organized and directed by Dr. Eysteinn Tryggvason, Assistant Professor of Geophysics, University of Tulsa. Other participants were University of Wisconsin, Jersey Production

Research Co., and Seismograph Service Corp. A total of 10 shots, ranging in size from 1,000 to 10,000 pounds, were fired with shot-point communications and timing equipment operated by Geological Survey crews. Five Geological Survey seismic crews occupied presurveyed locations along the recording line, moving with each successive shot.

REGIONAL GRAVITY STUDIES

San Luis Valley believed to be a rift broken by horsts and grabens

A regional gravity survey in the San Luis Valley area of South Central Colorado has been completed by J. R. Gaca. Interpretation of the data indicates that the San Luis Valley is a large rift valley internally broken by horsts and grabens. Depths to Precambrian rock underlying a filled graben along the eastern edge of the valley range from a minimum of 16,000 feet near Villa Grove to more than 30,000 feet north of Mount Blanca. The Sangre de Cristo mountains are interpreted as being locally uncompensated, indicating that these mountains are an excess mass being supported by the strength of the earth's crust.

Similar crust under Mount Shasta and Lassen Peak noted

Gravity measurements in the Mount Shasta–Lassen Peak areas of northern California suggest certain similarities in the upper part of the earth's crust in these two regions (La Fehr, written communication; L. C. Pakiser, 1–64). Interpretation of the Bouguer anomaly field suggests that both regions are characterized by large volumes of low-density rocks at relatively shallow depths. Limiting depth estimates indicate that the anomalous masses cannot lie entirely below depths of 4 to 10 kilometers. Application of Gauss' theorem⁶⁸ to the residual gravity anomalies in these areas indicates that the amount of mass deficiency so obtained corresponds well with the topographic mass excess. This indicates that the topographic mass excess of the Mount Shasta–Lassen Peak area is in approximate isostatic equilibrium with the shallow, low-density rocks. Thus, the compensation of Mount Shasta may be entirely a local phenomenon. In the classic sense of Airy or Pratt there may be no compensation necessary for Mount Shasta or the rest of the high Cascades. A vertical column taken through the crust at Mount Shasta may have no net excess mass below the relatively shallow depth of 6 to 10 km (La Fehr, written communication).

⁶⁸ T. R. La Fehr, 1965, The estimation of the total amount of anomalous mass by Gauss' theorem: *Jour. Geophys. Research*, v. 70, no. 8, p. 1911–1919.

GEOHERMAL INVESTIGATIONS

Studies of terrestrial heat flow in the Appalachian Mountain system, in active orogenic regions such as Puerto Rico and Chile, in the Western United States, and in the Arctic Ocean basin are continuing.

Heat flow in Appalachian Mountains measured

W. H. Diment reports that measurements of terrestrial heat flow at six localities in the central and southern part of the Appalachian Mountain system range from 0.7 to 1.4 microcalories per square centimeter per second. If a regional pattern exists, these measurements would suggest that heat flow is low in the eastern part of the Valley and Ridge province, higher in the metamorphic terrain adjacent to the east, and possibly highest in the intruded core of the Appalachian Mountain system (W. H. Diment, R. Raspet, M. A. Mayhew, and R. W. Werre, 4–65).

Low heat flow found in two active orogenic regions

Terrestrial heat flow was found to be low at two localities in active orogenic regions. A value of 0.6 microcalorie per square centimeter per second was found near Mayaguez, Puerto Rico (W. H. Diment and J. D. Weaver, 2–64; W. H. Diment, 1–64), and a preliminary value of 0.7–1.0 microcalorie per square centimeter per second was found near Valenor, Chile. These low values are not entirely unexpected, and when compared with values in Japan and California, suggest that high lateral gradients of heat flow may be typical of active orogenic regions.

Average heat flow in the western conterminous United States and Alaska

A few generalized preliminary results are available for the thermal gradient and thermal conductivity measurements being carried out in the western conterminous United States and Alaska by A. H. Lachenbruch. If heat flow is defined as "average" if its value is between 1.2 and 1.6 microcalories per square centimeter per second, and "low" or "high" if it is respectively below or above this range, then the heat flows determined at the western edge and the eastern edge of the Basin and Range province are high. Heat flow at the southern margin of the Basin and Range province varies locally from low to high. One determination in the eastern Mojave region of California, suggests average heat flow, and average values are being obtained in other parts of the State, in the Los Angeles Basin and western San Joaquin Valley. A high value was obtained in the Coeur d'Alene district of Idaho, and evidently average values were obtained in arctic Alaska. The

significance of results such as these cannot be studied profitably until more data become available.

Heat flow in the Arctic Ocean

The Arctic Ocean lies in the center of what was formerly one of the largest geothermally uncharted regions on earth. Studies by A. H. Lachenbruch show that the heat flow throughout much of the Canada Basin is about 1.40 microcalories per square centimeter per second, and that at its northern limit (at the flank of the Alpha Rise) heat flow drops abruptly to less than 0.8 microcalorie per square centimeter per second. The anomaly has no simple explanation in terms of bottom topography, oceanic circulation, or inhomogeneity in source distributions or thermal properties of the crust or upper mantle.

Theoretical and experimental studies by A. H. Lachenbruch show that the formulas in general use for finding the thermal conductivity of oceanic sediments yield results on the order of 10 percent too low for material from the Arctic Ocean. This was determined from a series of 300 thermal conductivity tests of sediment from the Arctic Ocean. In addition, mathematical results have been obtained for the effect of a surface slope on surface heat flow. These results provide a means of making approximate topographic corrections to ocean-bottom heat-flow data.

THEORETICAL AND EXPERIMENTAL GEOPHYSICS

ROCK MAGNETISM

Time scale for geomagnetic-polarity epochs

The time scale, shown in the accompanying table, for reversals of the geomagnetic field has been developed by Allan Cox, R. R. Doell, and G. B. Dalrymple.⁶⁹ Ages, given in millions of years, were determined by the potassium-argon method.

Reversed geomagnetic polarity	Age (millions of years)	Normal geomagnetic polarity
	0.0	
	0.85 ± 0.15	Brunhes epoch
Matuyama epoch	2.4 ± 0.1	(Olduvai event, 1.8–1.9 m.y.)
(Mammoth event, 3.0–3.1 m.y.)	3.35 ± 0.1	Gauss epoch
Gilbert epoch		

Geomagnetic polarity epochs are time intervals during which the magnetic field of the earth was

⁶⁹ Allan Cox, R. R. Doell, and G. B. Dalrymple, 1964, Reversals of the earth's magnetic field: *Science*, v. 144, p. 1537–1543.

predominantly or entirely of one polarity. These are interrupted by brief fluctuations in polarity termed "polarity events," as indicated in the accompanying table. During the past year several gaps in the time scale have been filled with information from additional paleomagnetic-geochronometric studies of rocks from the Sierra Nevada, the Pribilof Islands of Alaska, the Galapagos Islands, and New Mexico.

One of the more important new results has been confirmation of the Olduvai normal event occurring 1.8–1.9 million years ago. Previously this event was based on a single determination from Africa. Allan Cox, D. M. Hopkins, and G. B. Dalrymple have found four additional normally magnetized flows with the same age in the Pribilof Islands. These flows are bracketed in radiometric age and in stratigraphic position by reversely magnetized flows, leaving little room for questioning that the Olduvai event represents a brief worldwide fluctuation in the polarity of the earth's field.

A close correlation of polarity epochs with geomorphology has been established in the Hawaiian Islands, in Alaska, and in the Galapagos Islands. In all these areas, primary volcanic features are still recognizable in volcanic terrain formed during the Brunhes normal-polarity epoch. In terrain formed during the Matuyama or earlier polarity epochs, there is much more erosion and faulting. One of the more important contributions of paleomagnetism to geology may, in fact, be to provide in the Matuyama-Brunhes boundary an easily recognizable time horizon from which variations in rates of erosion may be determined as a function of environment.

The polarity time scale has also been linked to other methods for geologically inferring age. The relation of polarity epochs to the Pliocene-Pleistocene boundary, as defined by the type section in Italy, is being reviewed. Most probably the Pliocene-Pleistocene boundary lies within the Gauss normal-polarity epoch, although this is subject to considerable uncertainty because of a paucity of paleomagnetic and geochronometric results from the type section. Despite the uncertainty of the link with the Pliocene-Pleistocene boundary, polarity epochs are already proving very useful in providing stratigraphic links between Alaska, Idaho, California, New Mexico, Hawaii, Iceland, France, and the U.S.S.R.

Geomagnetic secular variation linked to earth's mantle

At the present time magnetic north deviates from true north by an angle which, while highly variable,

is typically several tens of degrees. Only a small part of this secular variation is due to iron-bearing minerals in the earth's crust; the greater part of such variation is caused by a system of currents in the earth's liquid core. The irregular field produced at the earth's surface today by these currents varies from place to place. In the South Atlantic Ocean, Indian Ocean, and Siberia, it is pronounced; in the central Pacific it is almost absent. The question arises, have these geographical differences persisted over long periods of time? If the answer is yes, the highly significant geophysical conclusion may be drawn that some process or condition in the mantle exerts control on the system of currents in the core which produces the secular variation, giving us one of our first indications of lateral inhomogeneities in the earth's lower mantle. Allan Cox and R. R. Doell (2-64) have found from paleomagnetic studies in Alaska, the western conterminous United States, Hawaii, and the Galapagos Islands that many of the geographic controls apparent in the present pattern of geomagnetic secular variation have persisted for the past million years. The present dependence on latitude is even more pronounced. Moreover, the low value of secular variation in the central Pacific has persisted during this time. An explanation that was previously offered is that this represents inductive shielding by layers with high electrical conductivity in the mantle. Cox and Doell (2-64) have examined this quantitatively and found that it fails by a wide margin to explain all of the observations. They advance the alternative explanation that lateral temperature difference in the lower mantle may exert control on the pattern of fluid convection in the core.

Paleomagnetism of Mesozoic rocks

Consistent paleomagnetic directions have been obtained by Andrew Griscom for each of the following Mesozoic intrusions in Maine, New Hampshire, and Vermont: Cuttingsville, Ascutney, Merrymeeting, Pawtuckaway, Ossipee, Alfred, Tatnic, Cape Neddick, Sanford, and Agamenticus. The directions differ somewhat but are all consistent with the Mesozoic ages. Some are normally, and others reversely, magnetized.

Geophysical investigations in the Gettysburg-Newark basin, Pennsylvania, by M. E. Beck, Jr., have suggested a correlation between lithology and variations in the degree of instability of remanent magnetic moments in the diabase of that area. Results to date show a marked correlation of magnetic instability with samples having a higher than average concentration of late magmatic material, espe-

cially interstitial micropegmatite. These observations apply to comparisons between separate samples from the same locality as well as to contrasts between localities. Grain-size differences, variations in degree of oxidation, or the existence of a range of titanium substitution in magnetite, all may contribute to the observed differences in magnetic properties.

Computer technique for magnetic analysis as applied to seamounts

In the conventional method used at present the direction of internal magnetization is determined by a laboratory examination of hand specimens of rock. On the other hand, the computer method developed by B. F. Grossling permits a determination to be made even for large geologic bodies. As an example, he determined the internal magnetization for three seamounts off the coast of California, finding that the directions of magnetization are markedly different from that of the earth's present magnetic field.

The determination of the internal magnetization of seamounts can be a valuable tool in the study of the relative movements of segments of the earth's crust, and of the crust as a whole with respect to the earth's magnetic field. Because seamounts are numerous (their number has been estimated at about 10,000) and scattered throughout most of the oceans, they may be expected to yield, ultimately, a wealth of information.

Changes in position of a seamount, since the time of its cooling below the Curie point, with respect to the dipole field, will result in a discrepancy between the directions of the remanent and induced magnetizations. The fact that some seamounts are found to be magnetized uniformly throughout may indicate that the earth's dipole field has been relatively stable during the length of time required for the formation and sufficient cooling of the seamount. Seamounts might have formed at quite different times. Meanwhile, many crustal movements have taken place.

An area of particular interest for an investigation of crustal movements, using the magnetization of seamounts, is the area of the Pacific Ocean west of the United States. Vacquier and others believe that very large displacements, even larger than 1,000 kilometers, have taken place along certain linear fractures in the oceanic crust. Such large displacements near the continental block may require other secondary displacements, such as rotation of crustal blocks, or an entirely new genetic conception about the formation of the oceanic and continental crusts.

New method for gravity and magnetic interpretation

R. G. Henderson's work on depth calculation by downward continuation has led to development of the depth functional, a new and powerful concept in quantitative magnetic and gravity interpretation. With the aid of the functional, explicit formulas for depth and depth extent have been obtained for some configurations for the first time. The depth for more complex configurations can be found from a plot of the functional versus depth. An analytical method has been devised for locating contacts between contrasting magnetic rocks despite shifts in the anomaly, with respect to the body, due to horizontal components of magnetization. It is now possible to compute susceptibility without recourse to a model for size.

Some success has been achieved in computing bottom configurations for two- and three-dimensional gravity distributions by downward continuation and an improved version of Tsubois' condensation method.

PHYSICAL PROPERTIES OF MINERALS AND ROCKS

New method for calculation of elastic constants

Previous methods of calculating the average elastic constants of single crystal aggregates invoked unrealistic assumptions regarding the stress (or strain) distribution in the aggregate. These assumptions led to dissimilar values for the calculated averages, especially when the anisotropy of the single crystal was large.

Louis Peselnick and Robert Meister have devised a variational method for calculating the average elastic constants, based on an energy extremum, and not using the assumptions on stress or strain. The method was extended to aggregates having hexagonal single crystals and to aggregates of trigonal single crystals. The bounds of the effective moduli obtained in this way show a considerable improvement over the bounds found by using the conventional stress-strain assumptions.

Experimental compaction of carbonate sediments

The pressure required to compact carbonate sediments during diagenesis is estimated with difficulty from such evidence as absence of crushing of fossils. Even delicate aragonite needles, 1 micron long, do not begin to crush until the pressure exceeds 100 bars. Most fossil shells, moreover, retain integrity to 1 kilobar of pressure because the wet sediment flows into and around the shell and supports it. Studies by E. C. Robertson show that compaction, measured by the ratio of grain volume to bulk vol-

ume, varies linearly with log pressure from 0.7 to 1 kb, but compaction increases more rapidly at higher pressure, approaching 1.0 (zero porosity) at about 20 kb. In the linear range, other variables affecting compaction are initial water content, grain size, and rate of loading; in the high range, permeability and cohesive grain friction become more important.

Dielectric constant of moist rocks

An old controversy regarding the dielectric constant of moist rock has been settled, according to J. H. Scott, by the development of a new laboratory method which avoids errors inherent in previous methods. Past attempts to determine the dielectric constant of moist rock have been hampered by electrode-polarization errors which caused measured values to be too high. Some critics believed that values of dielectric constant measured at low frequencies (100 cycles per second), which have been reported in the order of 10^6 for porous, saturated rocks, were entirely erroneous, and that the true value was probably closer to 10 or 20, as it is for dry rock. The new laboratory method uses electrochemically reversible electrodes made of platinized platinum plates and blotter-paper discs that are saturated with a suspension of silver and silver chloride. Preliminary results indicate that measurements by previous investigators are in error by no more than an order of magnitude, and that the dielectric constant of moist rock at low frequencies is, indeed, quite high.

Seismic-wave attenuation in various rocks

The amplitudes of the P head wave have been measured by J. S. Watkins, J. C. DeBremaecker, and R. H. Godson on an aa lava flow, on cinders, and on limestone. The results can be treated by assuming a complex shear modulus μ , but a real second Lamé constant λ . The data are satisfied by an equation of the type $A = Nd^{-2} \exp(-\Pi d Q\alpha^{-1}\nu\alpha - 1)$, where A is the amplitude, N an arbitrary constant, d the distance, $Q\alpha$ the attenuation coefficient for P waves, ν the frequency, and α the P-wave velocity. The ratio of the attenuation coefficient for P waves and the attenuation coefficient for shear waves ($Q\alpha/Q\beta$) derived from the assumption of a complex shear modulus equals half the ratio of the squares of the P- and S-wave velocities ($0.5 \alpha^2/\beta^2$). This relationship is in reasonable agreement with published results of investigations of P- and S-wave attenuation.

Field versus laboratory measurements of wave velocities

C. R. Tuttle and E. J. Monk measured a mean longitudinal wave velocity along the strike of 16,290

feet per second for 6 selected NX core samples of quartz-albite-feldspar schist from Dartmouth, Mass. The velocity for the same rock type measured in the field under shallow overburden in a direction bearing an unknown relation to structure was 14,900 feet per second. The mean longitudinal wave velocity measured on 16 cores from 5 boreholes compared to the velocity measured in the field was in the ratio of 1:0.9.

AERIAL INFRARED SURVEYS

Temperature distribution at summit of Mount Rainier

Aerial infrared surveys of Mount Rainier, Wash., by R. M. Moxham, D. R. Crandell, and W. E. Marlatt (p. D93-D100) showed that the highest temperature fumaroles are associated with the youngest, east summit crater. Temperatures are lower around the periphery of the west crater; a reticulated pattern extends down the northwest flank of the present summit cone and is bounded by an arcuate pattern of isolated anomalies that perhaps coincides with a third, older crater that has not been previously recognized. The summit craters are offset to the southeast of the central axis of the volcano and inferred central plug. The maximum temperatures associated with each of the craters are found consistently on their northwest quadrant, coinciding with the nearest approach to the central plug.

Results of infrared survey of Irazú Volcano, Costa Rica, are summarized in the section "Investigations in Other Countries."

SOLID-STATE STUDIES

FIELD STUDIES OF THERMOLUMINESCENCE

Contact metamorphic effects near Marble, Colo.

C. R. Roach and G. R. Johnson have studied the thermoluminescence characteristics of two formations in an attempt to trace their thermal histories. Detailed studies have been made on the Yule marble (contact metamorphosed Leadville Limestone) and the Niobrara Limestone near the Treasure Mountain stock, in the vicinity of Marble, Colo. The marble samples near the granite-marble contact and the nonrecrystallized limestone farthest removed from the stock had the greatest thermoluminescence. Heat from the stock probably destroyed some of the natural thermoluminescence of the Niobrara Limestone near the intrusive. The cause of the higher thermoluminescence in the marble near the contact is not clear.

Characteristics of granite near a ring dike in Colorado

In a study of Precambrian granite near a ring-dike complex in the vicinity of Virginia Dale, Colo.,

C. R. Roach and G. R. Johnson found that samples of the granite adjacent to the outer ring of the complex do not show natural thermoluminescence initially. However, after irradiation with gamma rays, they exhibit the thermoluminescence characteristic of this granite elsewhere. These relations suggest that the natural thermoluminescence of the Precambrian granite near the outer ring was destroyed by a relatively recent thermal event.

LABORATORY STUDIES OF THERMOLUMINESCENCE

Emission of cesium iodide measured by fast-scan monochromometer

Using a fast-scan monochromometer developed and built in laboratories of the U.S. Geological Survey, Prudencio Martinez and F. E. Senftle have been able to make spectral-emission measurements of thermoluminescence peaks in the ultraviolet and visible regions in a matter of a few seconds. X-ray fluorescence measurements of cesium iodide show 3 emission bands—at 290 millimicrons, 325 $m\mu$, and a broad band centered at 400 $m\mu$. The decay times of the 290- $m\mu$ and 325- $m\mu$ bands are temperature dependent with maximum values at 100 nanoseconds and 500 nsec, respectively. The decay time of the 400- $m\mu$ emission is about 15 microseconds. The 290- $m\mu$ and 325- $m\mu$ emission is ascribed to the production of V_k centers by the X-rays. Due to an absorption band at 299 $m\mu$, the 290- $m\mu$ emission is strongly absorbed in thick crystals. The cause of the 400- $m\mu$ band has not yet been determined.

Background thermoluminescence reduced by water lapping

In a study of low-temperature thermoluminescence of halite, C. R. Roach and G. R. Johnson have found that low-temperature autoluminographs of halite wafers indicate that water-lapped wafers of halite have less background thermoluminescence than do emery-lapped wafers of halite. Low-temperature glow curves obtained on these wafers indicate that the intensity of background thermoluminescence of the water-lapped halite is, on the average, about 250 percent lower than that of emery-lapped halite. These studies indicate that the best way to prepare samples of halite for low-temperature thermoluminescence studies is to prepare the wafers first by emery-lapping techniques and then to finish the specimens by water lapping.

MAGNETIC-SUSCEPTIBILITY MEASUREMENTS

Magnetic properties of zircon due to iron oxide coatings

In magnetic susceptibility and magnetization measurements on more than 100 granitic zircon sam-

ples, A. N. Thorpe, F. E. Senftle, and Sherman White found that all but one sample showed weak ferromagnetic properties. Leaching, reduction, and oxidation experiments along with magnetization studies on some of these specimens indicate that the crystal grains are coated with one or more iron oxides and (or) iron oxide hydrates and that the ferromagnetism is due to either or both Fe_3O_4 and $\gamma\text{-Fe}_2\text{O}_3$. The Fe_3O_4 or $\gamma\text{-Fe}_2\text{O}_3$ are not contamination brought in from outside the zircon crystals, but are formed by oxidation of natural-impurity iron exposed to the atmosphere on the grain surface.

Magnetic susceptibility of palladium

A. N. Thorpe has measured the magnetic susceptibility of palladium over the temperature interval between 4°K and 300°K. Previous measurements did not cover the 4°K to 20°K region. His measurements confirm the previous work and show that the susceptibility is almost temperature independent in the 4°K to 20°K range. The palladium-hydrogen and palladium-deuterium measurements confirm Mott's hypothesis that the addition of hydrogen or deuterium to palladium fills the D band of the electronic energy configuration. However, the measurements also show that the systems palladium-hydrogen and palladium-deuterium are not completely two-phased at intermediate concentrations of hydrogen and deuterium.

GEOCHEMISTRY, MINERALOGY, AND PETROLOGY

FIELD STUDIES IN PETROLOGY AND GEOCHEMISTRY

SILICIC PLUTONIC ROCKS

Allanite and monazite from Mount Wheeler area, Nevada

D. E. Lee and Harry Bastron have studied the rare-earth elements in 20 samples of allanite and 13 samples of monazite separated from granitic and xenolithic rocks of the Mount Wheeler area, eastern Nevada. The degree of fractionation of the rare earths, as measured by the sum of the atomic percentages of La, Ce, and Pr, varies directly with the CaO content of the whole rock, which itself depends largely on proximity of limestone. Comparisons of CaO and rare-earth data between xenoliths and respective closely associated intrusive phases indicate that the degree of fractionation of rare earths responds to whole-rock composition over distances of a few yards or more. Assimilation of limestone

rather than magmatic differentiation must be the dominant process that produced the observed chemical gradients in these granitic rocks. Allanites from the most calcium-rich rocks show a degree of concentration of the most "basic" rare earths usually found only in minerals from alkalic rocks and carbonatites, and whole-rock concentrations of such rare constituents as total rare earths, Zr, Ti, P, Ba, and Sr increase sympathetically with whole-rock calcium. These observations suggest that the chemical processes that produced these intrusive rocks may be similar to processes that form alkalic rocks and their associated carbonatites.

Allanite in Boulder Creek batholith, Colorado

N. L. Hickling has made a chemical, optical, and X-ray study of the accessory allanite in the Boulder Creek batholith, Colorado, as part of a continuing investigation of the rocks and minerals in this complex Precambrian intrusion. Allanite was separated from 20 representative rocks ranging from mafic granodiorite to leuco-quartz monzonite.

Consistent with an inferred late (deuteric?) origin for the allanite based upon field and micro-evidence, specific rare earths and other constituents of the allanite when plotted against the same, or other, oxides in the containing rocks showed little or no systematic relationship. In certain of the variation diagrams, however, the field of allanite from rocks of relatively low apparent age (as dated by T. W. Stern using the lead-alpha method applied to zircon) was clearly separable from that of allanite from relatively high-age rocks, indicating a loss of constituents during some later event. The variation in indices of refraction of the allanite was found to be strongly dependent upon its thorium and uranium content, and only slightly related, if at all, to other major constituents. The main variation in indices, therefore, is attributed to metamictization. The resulting radiation damage is evident in X-ray powder patterns as a broadening of lines.

Distribution of tantalum in southern California batholith

The distribution of tantalum in comagmatic rocks and in coexisting minerals of the southern California batholith has been studied by David Gottfried and J. I. Dinnin (p. B96-B100). The study is based on low-level tantalum analyses by J. Thompson of the Imperial College of Science and Technology, University of London, using neutron-activation methods and by Dinnin, using chemical methods. The tantalum content of the batholith is estimated at 0.4 parts per million, ranging from 0.13-0.24 ppm in gabbro to 0.38-0.45 ppm in granodiorite and quartz monzo-

nite. About 80 percent of the tantalum in the rocks is contained in hornblende and biotite. In the more siliceous rocks the tantalum content of these minerals is greater, but the abundance of the minerals is reduced.

Microperthite in Boulder batholith, Montana

R. I. Tilling, during the course of a petrologic study of the Boulder batholith, Montana, has found that the potassic phase of microperthite within 2 to 3 miles of the margin of the Rader Creek granodiorite is monoclinic, whereas the potassic phase near the core of this granodiorite body exhibits simultaneous monoclinic and triclinic symmetry. These dual-symmetry feldspars appear to record an intermediate state in the discontinuous inversion of orthoclase to microcline.

Use of refractive index of glass beads to estimate silica content

Use of the index of refraction of glass beads fused from samples of volcanic rocks has been suggested for estimating the silica content of the rock. N. K. Huber and C. D. Rinehart have now demonstrated that maximum accuracy can only be obtained when separate curves are established for individual rock suites.

MAFIC AND ULTRAMAFIC ROCKS

Alkali content of alpine-type ultramafic rocks

A study of the sodium and potassium contents of alpine-type ultramafic rocks from 6 regions in the United States and 1 in Australia was made by Warren Hamilton and Wayne Mountjoy. A dilution-addition modification of the flame-spectrophotometric method was used to measure the very small quantities of alkalis present in these rocks. The median value found for Na_2O is only 0.004 percent, and that for K_2O is 0.003 percent. Derivation of basalt magma from upper-mantle material similar to such ultramafic rocks, as has been postulated, is precluded by the relative amounts of sodium and potassium, which are 200 to 600 times more abundant in basalt than in these ultramafic rocks. The ultramafic rocks might have originated either as magmatic crystal precipitates or as mantle residues left after melting and removal of basaltic magma.

Composition and origin of oceanic tholeiites

A. E. J. Engel and C. G. Engel have determined that the oceanic tholeiites that form many submerged volcanic features in the oceans are characterized by extremely low amounts of Ba, K, P, Pb, Rb, Sr, Th, U, and Zr, as well as $\text{Fe}_2\text{O}_3/\text{FeO} < 0.2$ and $\text{Na/K} > 10$ in unaltered samples. The oceanic

tholeiites also have rare-earth distribution patterns, and K/Rb (1,300) and $\text{Sr}^{87}/\text{Sr}^{86}$ (0.702) ratios that are similar to those of calcium-rich (basaltic) achondritic meteorites. In contrast, the alkali-rich basalts that cap submarine and island volcanoes are relatively enriched in Ba, K, La, Nb, P, Pb, Rb, Sr, Ti, Th, U, and Zr, and in ferric iron. The analytical data and field relations indicate that the alkali-rich basalts are derived from oceanic tholeiites via processes of magmatic differentiation, and that the oceanic tholeiites are the principal, or only primary magma generated in the upper mantle. The close compositional kinship between the oceanic tholeiites and calcium-rich achondrites is further evidence of the relatively primitive nature of the oceanic tholeiites. The chemical properties of oceanic tholeiites suggest that the upper mantle probably contains less than the following amounts (in parts per million): Ba, 10; K, 1,000; Pb, 0.4; Rb, 10; Th, 0.2; and U, 0.1. The $\text{Sr}^{87}/\text{Sr}^{86}$ ratio must be less than 0.7015, Th/U about 2, K/Rb about 1,500–2,000, and $\text{Fe}_2\text{O}_3/\text{FeO}$ less than 0.1.

Origin of aluminous basalt versus tholeiitic basalt

Warren B. Hamilton (4–64) proposed that basaltic magmas of the high-alumina association are generated by partial melting of the basaltic component of mantle rock at levels above transformation of mantle to eclogite or pyroxenite. The broad compositional spectrum within high-alumina basalts reflects varying degrees of partial melting over a broad temperature range. Tholeiite magmas, on the other hand, are thought to be generated beneath the level of transformation to eclogite or pyroxenite, and the narrow spectrum of their compositions reflect more complete melting of mantle material over a narrow temperature range.

Undevitrified Late Triassic glass from Alaska

Petrographic and X-ray study by D. A. Brew and L. J. P. Muffer (p. C38–C43) of glass-bearing aquagene tuff of Late Triassic (Norian) age from the Keku Strait area has shown that the basaltic glass is undevitrified and that it occurs together with other volcanic fragments as angular clasts in a matrix of sparry calcite. Glass fragments separated from the rock were analyzed by J. D. Obradovich and found to give a K-Ar age of 126 million years, indicating that some radiogenic argon has been lost from the glass. This glass is apparently one of the oldest undevitrified volcanic glasses ever reported.

X-ray curves give composition of Hawaiian olivine

K. J. Murata, Harry Bastron, and W. W. Brannock (p. C35–C37) have established an X-ray curve

for the determination of the composition of Hawaiian olivine. The curve deviates from that previously established by E. D. Jackson⁷⁰ for plutonic olivine. This deviation is related to the larger cell size of the Hawaiian olivine, which in turn is caused by the presence of calcium, a cation of large radius relative to other cations of olivine. The Hawaiian olivine contains 0.2–0.5 percent calcium, compared to a few hundredths of a percent in plutonic olivine.

SILICIC VOLCANIC ROCKS

Chemical characteristics of pantellerite glass, Nevada

D. C. Noble (p. B85–B90) has separated and described a pantellerite glass from the basal vitrophyre of a middle Pliocene ash-flow tuff in southern Nevada. Content of sodium, iron, zirconium, and halogens of the glass is extremely high for a silicic rock, and content of aluminum, magnesium, and calcium is abnormally low. Analyses of devitrified rocks of the same ash-flow tuff are low in sodium and halogens; these elements probably were leached from the pantellerite by ground water after devitrification. This pantellerite is tentatively believed to have been generated by the direct fractionation of a mafic or low-silica intermediate alkalic magma without the existence of trachyte as an intermediate magma.

Scapolite related to chlorine release in Bandelier Tuff, New Mexico

R. L. Smith and K. O. Dickson, during the course of a continuing study of the mineralogy of ash flows of the Bandelier Tuff, Jemez Mountains, N. Mex., have discovered that scapolite is a common and apparently characteristic mineral of the vapor-phase zone. The mineral is associated with cristobalite, tridymite, and alkali feldspar in pumice vesicles and is apparently precipitated from chlorine-rich gases during cooling of the tuff. X-ray studies show that the cell dimensions of the scapolite change systematically with degree of welding of the tuff, and may reflect differences in Si-Al ordering related to rate of cooling of the tuff or may reflect variable chemistry. The discovery provides clues to problems of chlorine release during crystallization of rhyolitic ash flows.

Nb, Th, and U decrease upward in Bandelier Tuff

David Gottfried has studied the variation in amount of Nb, Th, and U in several sections of the Bandelier Tuff in the Jemez Mountains, N. Mex. The

vitric zone at the base of a typical section of genetically related multiple-flow units contains the highest amounts of Nb (130 parts per million), Th (30 ppm), and U (10 ppm). There is a progressive decrease in the amounts of these elements in the flow units higher in each section. The uppermost units contain approximately 60 ppm Nb, 15 ppm Th, and 5 ppm U. These data support the concepts of R. L. Smith and R. A. Bailey that the Bandelier ash-flow sheets represent the inverted contents of zoned or layered magma chambers.

Three different niobium trends in volcanic rocks noted

In a general study of niobium in volcanic rocks, David Gottfried has isolated three fairly distinct trends of variation of niobium during magmatic differentiation: (1) an increase in niobium throughout differentiation (alkalic and alkalic-calcic volcanic suites from Augusta County, Va., and from the Big Bend area, Texas), (2) a decrease in niobium throughout differentiation (high calcic volcanic suites from Saipan, the Mount Garibaldi area of British Columbia, and the Strawberry Mountains, Oreg.), and (3) a slight increase in niobium in mafic rocks and a relatively constant amount in the intermediate and siliceous rocks (calc-alkalic suite from the Medicine Lake Highland, Calif.). The volcanic rocks of a low-lime alkalic suite show a 10- to 50-fold enrichment in niobium over the amounts found in corresponding rock types in a highly calcic suite.

METAMORPHIC ROCKS

New iron silicates reported from Cazadero area, California

R. G. Coleman, R. C. Erd, and D. E. Lee have identified two recently discovered⁷¹ iron silicates, deerite and howieite, in the Franciscan Formation of the Cazadero area, California, and suspect the presence of zussmanite also at this locality. These minerals appear to be restricted to iron-rich metasedimentary rocks of intermediate metamorphic grade within the glaucophane schist facies. The bulk composition of the metasedimentary rocks is similar to that of the Precambrian iron formations, except for high sodium values. Stilpnomelane, crossite-riebeckite, and spessartine-almandine garnet are the commonly associated minerals in the Cazadero area. Deerite is found in both quartz-rich and quartz-poor rocks. In contrast, howieite (and zussmanite?) is usually restricted to quartz-poor iron-

⁷⁰ E. D. Jackson, 1960. X-ray determinative curve for natural olivine of composition $\text{Fe}_{0.90-1.00}$; Art. 197 in U.S. Geol. Survey Prof. Paper 400-B, p. B432–B434.

⁷¹ S. O. Agrell, M. G. Gown, and D. McKie, 1965. Deerite, howieite, and zussmanite, three new minerals from the Franciscan of the Laytonville district, Mendocino Co., California [abs.]: *Am. Mineralogist*, v. 50, no. 1–2, p. 278.

stones. The apparent restriction of these new minerals to the glaucophane schist facies indicates that they may be the high-pressure analogs of greenalite, minnesotaite, and grunerite.

Composition of jadeite throws light on its genesis

R. C. Coleman (p. C25–C34) has found that the pyroxene from the metagraywackes of the Franciscan Formation in California is not a pure jadeite. The presence of significant amounts of pyroxene components other than jadeite in these jadeitic pyroxenes suggest formation at lower pressures and temperatures than those found experimentally for the conversion of albite to pure jadeite + quartz. Petrologic and chemical evidence indicates that the conversion of graywacke to the assemblage quartz-jadeitic pyroxene-lawsonite-muscovite-glaucophane takes place without the introduction of alkalies. The reaction $8 \text{ albite} + \text{chlorite} + \text{CaCO}_3 \rightarrow 5 \text{ jadeitic pyroxene} + 2 \text{ glaucophane} + \text{quartz} + 2\text{H}_2\text{O} + \text{CO}_2$ may be important in relating greenschist to glaucophane schist facies metamorphism.

Geothermal well data record present-day metamorphism

Petrologic and chemical data from the Salton Sea geothermal area have enabled D. E. White and L. J. P. Muffler (1–65) to conclude that upper Cenozoic sediments of the Colorado River delta are being metamorphosed in the present thermal environment to mineral assemblages of the greenschist facies. In the I.I.D. No. 1 geothermal well, the following generalized reaction occurs in the zone from 1,200 to 2,300 feet and in the general temperature range of 120°C to 200°C:

In the zone from 2,700 to 4,000 feet and at temperatures in the range from 200°C to more than 300°C, calcite breaks down and epidote is formed. This second relationship involves considerable cation exchange with the saline geothermal brine found in the geothermal field.

THERMAL WATER

Large magma chamber at Steamboat Springs, Nev., indicated

D. E. White, G. A. Thompson, and C. H. Sandberg, (2–64), in an exhaustive documentation of the detailed late Tertiary and Quaternary history of the Steamboat Springs thermal area, Washoe County, Nev., conclude that a magma chamber of at least 50 cubic kilometers at 6 to 9 km depth has evolved heat, water, and mineral matter for at least 100,000 years.

Structural control for most of the thermal activity of the area was provided by old north-trending east-dipping normal faults. Extensive deposits of siliceous sinter are localized around structurally favorable outlets.

Three patterns of alteration found in drill holes at Steamboat Springs, Nev.

Robert Schoen and D. E. White have recognized three patterns of alteration in two drill holes at Steamboat Springs, Nev.:

1. Near-surface acid leaching where H_2S oxidizes to form H_2SO_4 above the water table; opal, kaolinite, and alunite are characteristic products.
2. Near-surface alteration below the chloride-water table; hydrothermal K-feldspar is an especially abundant replacement of plagioclase of the volcanic rocks but also replaces plagioclase of granodiorite.
3. Zoned alteration adjacent to fractures at depths greater than 200 feet. In the outer zone, hornblende is replaced by illite-montmorillonite and chlorite, biotite by chlorite, and plagioclase by albite and calcite and (cr) illite-montmorillonite. In the most intensely altered rocks next to fractures, all earlier formed hydrothermal silicates and plutonic K-feldspar are replaced by illite (< 10 percent interlayered montmorillonite), quartz, and pyrite.

DIAGENETIC ROCKS

Lazurite, talc, and chlorite in Green River Formation, Wyoming

W. H. Bradley (2–64) has identified authigenic lazurite, talc, and chlorite in a salt bed of the Green River Formation, Wyoming. Talc and chlorite as authigenic minerals in lacustrine salt beds had not previously been reported. The presence of these silicates instead of the usual magnesium silicates of greater hydration, such as stevensite or saponite, suggests that the assemblage formed at a late stage of diagenesis, and may have replaced previously formed saponite or stevensite.

Alteration of rhyolitic glass related to topography

R. A. Sheppard and A. J. Gude 3rd (p. D44–D47) in a study of the diagenetic alteration of a gently dipping bedded tuff in the Pliocene Ricardo Formation of southeastern California, have documented a correlation between topographic elevation and stage of replacement of fresh rhyolitic glass by clinoptilolite, opal and (or) montmorillonite. The tuff above an elevation of 3,000 feet generally is fresh, but the

same tuff below 3,000 feet is altered. This suggests that alteration was accomplished by subsurface water after the formation had been tilted. Solution and hydrolysis of rhyolitic glass by downward-moving water would increase the pH and salinity of the water, thereby providing a chemical environment more favorable for the formation of zeolites.

MINERALOGIC STUDIES AND CRYSTAL CHEMISTRY

Crystal structure of scapolites

The crystal structure of mizzonite, a calcium and carbonate-rich scapolite, has been refined by J. J. Papike. The results of this refinement together with a previous analysis of the crystal structure of a sodium and chlorine-rich scapolite (J. J. Papike and T. Zoltai, 1-65) provide the following new crystal chemical information concerning the variation of the structure with composition:

The basic structure is the same throughout the solid-solution series, although there are some significant distortions in the tetrahedral framework.

Ordered marialite-rich scapolites have aluminum in only 1 of the 2 types of 4-membered rings of tetrahedra present in the structure, whereas meionite-rich scapolites have aluminum distributed in both types of rings. The aluminum distribution varies with the overall composition and can be associated with the variation of unit-cell dimensions in the isomorphous series.

The electron-density distribution indicates disorder of the CO_3^{2-} groups in the structure. The disorder does not appear to be rotational, but instead, the result of averaging among four distinct positions for each group.

Mineralogy of djurleite

D. E. Appleman and E. H. Roseboom have found that the new copper sulfide mineral djurleite, is orthorhombic with $a = 15.752$, $b = 26.895$, and $c = 13.554$ Å. The composition of synthetic djurleite lies between $\text{Cu}_{1.963}\text{S}$ and $\text{Cu}_{1.972}\text{S}$; the density is 5.747 ± 0.007 as measured on 2 large natural crystals. Previous workers had concluded incorrectly that djurleite was probably two phases. From Roseboom's work on the system Cu-S, it is clear that djurleite is a single phase. The most probable formula for djurleite is $128[\text{Cu}_{1.969}\text{S}]$, corresponding to a unit-cell content of $\text{Cu}_{252}\text{S}_{128}$. This represents a loss of four Cu atoms per unit cell over chalcocite, Cu_2S .

Aluminous enstatites

B. J. Skinner and F. R. Boyd (2-64) have calculated unit-cell volumes and densities for a series of

aluminous enstatites at regular intervals up to 15 weight percent Al_2O_3 . The extensive coupled substitution of (AlAl) for (SiMg) in orthorhombic enstatite (MgSiO_3) had been demonstrated earlier by F. R. Boyd and J. L. England⁷² in their high-pressure studies. They proposed that a pyroxene having a composition equivalent to pyrope garnet ($\text{Mg}_3\text{Al}_2\text{Si}_3\text{O}_{12}$) with 25 percent Al_2O_3 substitution might be prepared at high pressure. However, projecting the density-versus-composition relation determined for aluminous enstatites to the composition $\text{Mg}_3\text{Al}_2\text{Si}_3\text{O}_{12}$, a density of 3.28 ± 0.01 gram/cubic centimeter is predicted. When this is compared to the high measured density of 3.58 g/cm^3 for pyrope it suggests that increasing pressure will favor the assemblage pyrope plus enstatite over aluminous enstatites, a conclusion now verified by F. R. Boyd and J. L. England.⁷³

Relation of reedmergnerite to feldspar structures

Following the completion of least-squares refinement of 3,000 observed X-ray intensity measurements of reedmergnerite, NaBSi_3O_8 , D. E. Appleman and J. R. Clark compared the refined structure to the related low albite, maximum microcline, and anorthite crystal structures. Strikingly similar Si-O distances between like atoms in reedmergnerite and low albite confirm the assumption of completely ordered tetrahedral cations in both mineral structures. The sodium cations in reedmergnerite, unlike those in low albite, exhibit no apparent anisotropy. Also, the AlO_4 tetrahedra are found to be less regular than either the SiO_4 or BO_4 tetrahedra.

Polytypism in micas

Malcolm Ross, D. R. Wones, and Hiroshi Takeda have completed a systematic description of all possible 1-, 2-, 3-, and 4-layer mica polytypes, as well as all 5- and 6-layer forms which have 0° and $\pm 120^\circ$ stacking sequences. Unit-cell parameters, vector stacking symbols, space groups, and "structure presence criteria" have been derived for all of the above 62 possible crystal structures. With these data it is possible to predict the crystal structure of a large number of these polytypes merely by inspecting the single-crystal X-ray photographs. The accompanying table lists the various crystallographic parameters of all possible 1-, 2-, and 3-layer forms.

The $3\text{Tc}_2[02\bar{2}]$ and $4\text{M}_2[2220]$ forms, as well as many other more complex polytypes such as 8-, 11-, and 20-layer monoclinic and 8-, 14-, and 25-layer

⁷² F. R. Boyd and J. L. England, 1960, Aluminous enstatite, in *Minerals of the mantle*: Carnegie Inst. Washington Year Book, 1959-1960, p. 49-52.

⁷³ F. R. Boyd and J. L. England, 1964, The system enstatite-pyrope: Carnegie Inst. Washington Year Book 1963-1964, p. 157-161.

Information for systematic identification of all possible 1-, 2-, and 3-layer mica polytypes

Polytype vector symbol	a (Å)	b (Å)	d_{001} (Å)	α (degrees)	β (degrees)	γ (degrees)	Space group	Structure presence criteria
1M[0]-----	5.3	9.2	10.0	90	100.0	90	$C2/m$	--
2M ₁ [2 $\bar{2}$]-----	5.3	9.2	20.0	90	95.1	90	$C2/c$	$h3hl:l = 2n + h$
2M ₂ [1 $\bar{1}$]-----	9.2	5.3	20.0	90	98.7	90	$C2/c$	--
20[33]-----	5.3	9.2	20.0	90	90	90	$Ccmm$	--
3T[222]-----	5.3	--	30.0	90	90	120	$P3_112$	$hh2\bar{h}l:l = 3n$
3M ₁ [033]-----	5.3	9.2	30.0	90	93.4	90	$C2/m$	$Ok\bar{l}:l = 3n$
3M ₂ [1 $\bar{1}\bar{2}$]-----	5.3	9.2	30.0	90	93.4	90	$C2$	--
3Tc ₁ [01 $\bar{1}$]-----	5.3	9.2	30.0	98.7	91.7	90	$C\bar{1}$	$Ok\bar{l}:l = 3n$
3Tc ₂ [02 $\bar{2}$]-----	5.3	9.2	30.0	92.9	95.1	90	$C\bar{1}$	$hOl:l = 3n$
3Tc ₃ [123]-----	5.3	9.2	30.0	90	93.4	90	$C\bar{1}$	--

triclinic forms have been found by Malcolm Ross and D. R. Wones (2-65). Examination of more than 150 biotite crystals by Ross indicates that (1) most biotite samples consist of mixtures of many complex polytypes, and (2) all biotite structures consist of only 0° and $\pm 120^\circ$ stacking sequences. At the present time the following distribution of polytypes in micas is indicated: biotites, mixture of many polytypes and also 1M_r-120 forms (random $\pm 120^\circ$ layer rotations); phlogopites, 1M, rarely 2M₁, 3T; lithium micas, 1M, 2M₁, 2M₂, 3T, rarely more complex forms; muscovite and paragonite, 2M₁, rarely 1M, 3T; authigenic micas, 1M, 2M₁, plus possible disordered 1M_r-60, 1M_r-120, 1M_r-180, and two-dimensional forms.

Crystal structure of ulexite

The crystal structure of the Na-Ca borate mineral ulexite has been found by J. R. Clark and D. E. Appleman (3-64) to contain three structural groups: isolated borate polyanions, $[B_5O_6(OH)_6]^{-3}$; Ca-coordination polyhedra; and Na-coordination octahedra that share edges to form continuous chains parallel to the c direction. These three groups are joined by hydrogen bonds into a three-dimensional network. The structural formula for ulexite is $NaCaB_5O_6(OH)_6 \cdot 5H_2O$. The isolated polyanions, according to C. L. Christ's fourth rule⁷⁴, may polymerize into chains by linking and splitting out water. Such chains have been found by H. M. Kurbanov and others⁷⁵ in the crystal structure of the related mineral, probertite, $CaNaB_5O_7(OH)_4 \cdot 3H_2O$.

Crystal structure of zinc analog of hummerite

H. T. Evans, Jr. (1-65) has solved the crystal

structure of a complex potassium-zinc vanadate, $K_2Zn_2V_{10}O_{28} \cdot 16H_2O$, the zinc analog of hummerite. The structure was solved from a three-dimensional, sharpened Patterson map. The decavanadate group $(V_{10}O_{28})^{-6}$ consists of a cluster of 10 condensed VO_6 octahedra: 6 are arranged in a 2×3 rectangle by sharing horizontal edges, and 2 pairs of octahedra are inserted above and below the rectangle by sharing sloping edges. The $(V_{10}O_{28})^{-6}$ ion is known to exist in solutions although until the present X-ray study was completed the structure was not understood.

Coordination of uranyl ion in uranium (VI) oxide hydrates

C. L. Christ (1-65) has explained the spontaneous transformation of schoepite I ($UO_3 \cdot nH_2O$), with loss of water, to schoepite II or schoepite III, as due to a change from an unstable near-hexagonal coordination of the uranyl ion in schoepite I, to a pentagonal coordination of the uranyl ion in the less hydrous phases. A plan is proposed for determining the kind and extent of thermodynamic stability among these hydrates of UO_3 , and for relating the thermodynamic stability to the observed type of uranyl ion coordination (6-, 5- or 4-fold).

Crystallography of ganophyllite

Two polytypes of ganophyllite, a manganese aluminosilicate, have been recognized by M. L. Lindberg. Ganophyllite samples from Franklin Furnace, N. J. contain two forms with the following unit-cell data: (phase A) $a = 50.36$, $b = 27.08$, $c = 16.59$ Å, $\beta = 94^\circ 10'$, space group Cc or $C2/c$; and (phase B) $a = 24.70$, $b = 16.32$, $c = 22.20$ Å, $\beta = 94^\circ 20'$, space group Cc or $C2/c$. Phase A was also found in samples from the Harstig mine, Sweden. The following unit-cell parameters were measured from this specimen: $a = 50.34$, $b = 27.04$, $c = 16.60$ Å, $\beta = 94^\circ 10'$,

⁷⁴ C. L. Christ, 1960, Crystal chemistry and systematic classification of hydrated borate minerals: *Am. Mineralogist*, v. 45, p. 334-340.

⁷⁵ H. M. Kurbanov, I. M. Rumanova, and N. V. Belov, 1963, Crystal structure of probertite, $CaNa[B_5O_7(OH)_4] \cdot 3H_2O$: *Dokl. Akad. Nauk SSSR*, v. 152, p. 1100-1103. [in Russian]

space group *Cc* or *C2/c*. Both phases have subcells similar in projection on (010) to an "expanded" $2M_1$ mica. A relationship to the alleverdite structure is indicated.

Physical properties of feldspars

Refinement of X-ray powder diffraction data of the feldspar phases in perthites with sodic bulk compositions by D. B. Stewart, T. L. Wright, and Dora von Limbach yielded the unit-cell dimensions and structural states of the exsolved phases. Before exsolution takes place, the lattice becomes considerably distorted, particularly along the *a* axis. The distortion is relieved by a simultaneous change of composition and structural state as exsolution takes place. The low temperature assemblage ultimately consists of low albite of nearly pure $\text{NaAlSi}_3\text{O}_8$ composition, and maximum microcline of nearly pure KAlSi_3O_8 composition. Intermediate compositions and structural states do occur, but they are rare and characteristically show cell dimensions that are strongly distorted with reference to the standard curves for known series of intermediate structural states. The refined dimensions gave values for the reciprocal cell angles that were consistently different from those reported by W. S. MacKenzie and J. V. Smith⁷⁶ for the same samples. Their results were obtained by another X-ray method (J. V. Smith and W. S. MacKenzie⁷⁷) that apparently does not always yield correct values.

The unit-cell parameters of a number of natural ternary feldspars of high structural state (anorthoclases) were determined and compared with the determinative curves for the high albite-high sanidine series to determine how closely the cell parameters indicate the Or content and structural state. The very satisfactory agreement suggests that the determinative curves prepared for the pure alkali feldspars may be applicable to An-bearing ternary solid solutions as well.

Examination by D. B. Stewart, T. L. Wright, and Dora von Limbach of the relationship between the volume of calcic labradorite (An 67.2) and temperature indicates the possible existence of a reversible high-order transformation between 650° and 1050°C. The transformation may be the change from a body-centered lattice to a C-centered lattice. If verified by X-ray single-crystal measurements in this temperature range, the possibility of establishing an

equilibrium phase diagram for calcic plagioclase by high-temperature X-ray measurements is indicated.

D. B. Stewart resolved a mineralogical mystery by showing that the physical properties reported for "analbite" and "monalbite," supposedly polymorphs of $\text{NaAlSi}_3\text{O}_8$ feldspar, were actually those for the potassium-bearing sodic feldspars, anorthoclase and sanidine. This result considerably simplifies the phase diagram for alkali feldspars, and makes our understanding of feldspar order-disorder relations more complete.

EXPERIMENTAL GEOCHEMISTRY

BASIC STABILITY RELATIONSHIPS

Theory and applications of phase diagrams

E-an Zen has found in systems of $n + 3$ phases that a definite relationship exists between the permissible configurations of invariant points and univariant curves and the relative compositions of the phases, and that for some cases, a unique relationship exists. Zen also found that the possible "grids" may be derived from the projection of primitive polyhedra without body diagonals.

Heats of solution of alkali feldspars

Heats of solution of low albite, microcline, and 10 intermediate compositions on the $\text{NaAlSi}_3\text{O}_8$ - KAlSi_3O_8 binary, in 20 percent HF, have been measured by D. R. Waldbaum and R. A. Robie. The excess enthalpy of mixing is nearly symmetric and has a maximum value of $1,800 \pm 200$ calories. Refined unit-cell parameters provide information on the volume of mixing, and permit a thermodynamic analysis of the alkali feldspar solid-solution series.

STUDIES OF MAGMATIC PROCESSES

Crystal settling and convection in granitic magma

H. R. Shaw (1-65), in an examination of the conditions for natural and forced convection and crystal settling in magma of granitic composition, found that the behavior of such a magma cannot be treated as a static condition. Crystallization, differentiation, and heat flow must take account of mass and heat transfer by natural convection. Although more fundamental data are required concerning the mechanical behavior of the crust in which a magma is contained, dike propagation and eruption rates of rhyolite magma require the flow sequence: Incompressible homogeneous flow $\rightarrow$ compressible two-phase flow (liquid-gas) $\rightarrow$ particulate flow.

Absorption of H_2O by obsidian at high temperature and pressure

The rate of absorption of H_2O in obsidian at

⁷⁶ W. S. MacKenzie and J. V. Smith, 1962, Single crystal X-ray studies of crypto- and micro-perthites: *Norsk Geologisk Tidsskrift*, v. 42, p. 72-103.

⁷⁷ J. V. Smith and W. S. MacKenzie, 1955, The alkali feldspars: II. A simple X-ray technique for the study of alkali feldspars: *Am. Mineralogist*, v. 40, p. 773-747.

850°C and at fluid pressures between 100 and 2,000 bars has been determined by H. R. Shaw. The sorption curves at 850°C and 700 bars are complicated by sharp changes in slope, although the initial slopes of all the curves at a variety of conditions of pressure and temperature are similar. The data indicate that the average diffusion coefficients range from 1 to 7×10^{-8} cm²/sec⁻¹, that the Stokes-Einstein relationship is not applicable to the diffusion of H₂O in silicate melts, and that the gain or loss of H₂O by a diffusion mechanism in granitic magma is exceedingly slow. A driving potential of 2,000 bars H₂O pressure would move the hydration front in a dry granitic melt less than 50 meters in 1 million years.

Reactions of phlogopite and quartz

D. R. Wones and F. C. Dodge have demonstrated that the assemblage phlogopite + quartz reacts to form the two assemblages, sanidine + enstatite + gas, and melt + enstatite + gas. The univariant curves that represent the two reactions intersect at an univariant point at 840° ± 10°C and 400 ± 100 bars H₂O pressure. The fluorobiotite component may increase the thermal stability of the biotite phase, but the data indicate temperatures and H₂O pressures at which biotite and quartz may crystallize contemporaneously from a silicate melt. The data also prove that charnockitic metamorphic rocks result from low H₂O pressures during metamorphism, and probably represent either metamorphosed igneous rocks, or a metamorphic terrane dominated by decarbonation reactions.

The univariant reaction phlogopite + sanidine + quartz + H₂O ⇌ melt is less than 10°C below the sanidine + quartz + H₂O curve and indicates that high MgO contents of granitic magmas are due either to superheating during fusion or to contamination of a melt of "granite minimum" composition.

Pyrrhotite-biotite, a promising geothermometer

Compositions of coexisting pyrrhotite and biotite in the Mount Antero Granite, Chaffee County, Colo., are consistent with calculated values for magmatic temperatures, according to Priestley Toulmin III. Toulmin and D. R. Wones are undertaking a laboratory investigation to calibrate this promising geothermometer.

STUDIES OF METAMORPHIC PROCESSES

Low H₂O pressures calculated for metamorphism of marble

D. R. Wones calculated the H₂O pressures implied by biotite-potassium feldspar-magnetite assemblages

described by A. E. J. Engel and C. G. Engel⁷⁸ in gneiss and interbedded marble from the northwest Adirondacks of New York. The calculated pressures range between 0.1 and 10 bars, and agree with comparable calculations based on the amphibolite + clinopyroxene + orthopyroxene + quartz assemblages. The low H₂O pressures are consistent with a metamorphism in which the decarbonation of the marble was the dominant reaction. The CO₂ pressures, calculated from graphite equilibria, range from 500 to 5,000 bars and are consistent with the proposed model of the metamorphism.

Sedimentary-zeolitic-greenschist facies

A combination of pressure-temperature (P-T) diagrams and activity-activity diagrams may provide a sound basis for the interpretation of coexisting mineral parageneses. E-an Zen has shown that plots of the activity of H₂O, a_{H_2O} , and activity of metal oxides, a_{MO} , explain some relationships between the zeolite facies (high a_{H_2O} -high a_{MO}), sedimentary facies (high a_{H_2O} -low a_{MO}), and greenschist facies (low a_{H_2O} -high a_{MO}). As a consequence, the zeolite facies is not a necessary intermediate step between the sedimentary and greenschist facies.

SOLUBILITIES AND EXCHANGE REACTIONS

Flocculation of kaolinite suspensions

The flocculation of fine kaolinite suspensions in aqueous ionic solutions was found by J. B. Robertson and C. H. Wayman to occur over a range of pH values in weak ($10^{-1}N$) sodium, calcium, or aluminous chloride solutions. Maximum flocculation occurred at pH 4.0-2.0 for sodium and calcium ion-saturated kaolinite and at pH 5.0-2.5 for aluminum ion-saturated kaolinite. The flocculation was determined by measuring dispersion settling rates by a light-beam transmittance principle.

Solubility of silica glass at elevated temperature and pressure

R. O. Fournier and J. J. Rowe have determined the solubility of silica glass in H₂O at 200°C-380°C and at pressures of 15-1,000 bars. Logarithmic plots of H₂O fugacity and the concentration of dissolved silica suggest the presence of three different hydration states of the dissolved silica. The number of H₂O molecules associated with an SiO₂ group varies inversely with the density of the solution. Similar types of dissolved species coexist with quartz and silica glass at any given temperature and pressure.

⁷⁸ A. E. J. Engel and C. G. Engel, 1958, 1960, Progressive metamorphism and granitization of the major paragneiss, northwest Adirondack Mountains, New York; pt. I. Total rock; pt. 2. Mineralogy: Geol. Soc. America Bull., v. 69, p. 1369-1413, 1958; v. 71, p. 1-58, 1960.

Exchange constants for natural and synthetic glasses

Exchange constants for natural and synthetic glasses which involve the ions H^+ , Na^+ , K^+ , Ca^{+2} , and Mg^{+2} have been determined by A. H. Truesdell using an electrode method⁷⁹. The aluminosilicate glasses vary in content of Fe_2O_3 , FeO , MgO , CaO , Na_2O and K_2O , and these variations affect the exchange constants in a manner that may be explained by a model of ion-exchange selectivity.

Sulfate equilibria

G. W. Morey, J. J. Rowe, and C. C. Silber have shown, in their studies of the system K_2SO_4 – $CaSO_4$ – $MgSO_4$, that a high-temperature form of $MgSO_4$ exists and that a new compound, $CaSO_4 \cdot 3MgSO_4$ exists between 800° and 850°C. The latter compound extends to the field between langbeinite–calcium langbeinite.

Euhedral anhydrite, $CaSO_4$, has been synthesized in unseeded artificial sea water at 70°C by E-an Zen. The crystals exhibit skeletal growth and pinacoidal facies.

Activity-activity diagrams explain borate parageneses

C. L. Christ, A. H. Truesdell, and R. C. Erd have explained mineral parageneses in borate deposits of California and Nevada by representing the system Na_2O – MgO – CaO – B_2O_3 – H_2O in terms of activity-activity diagrams and the three ternary systems: (1) $4CaO \cdot 5B_2O_3 \cdot 7H_2O$ (priceite)– B_2O_3 – H_2O – H_2O ; (2) $Na_2B_4O_7 \cdot 4H_2O$ (kernite)– $2CaO \cdot 3B_2O_3 \cdot 5H_2O$ (colemanite)– H_2O ; and (3) $2CaO \cdot 3B_2O_3 \cdot 5H_2O$ – $2MgO \cdot 3B_2O_3 \cdot 7H_2O$ – H_2O .

Stability fields in systems K_2O - and Na_2O - Al_2O_3 - SiO_2 - H_2O

J. J. Hemley has found, in the system K_2O - and Na_2O - Al_2O_3 - SiO_2 - H_2O , that relations between kaolinite, muscovite, and K-feldspar are quite different in the aqueous chloride and aqueous sulfate environments. The muscovite field is both shifted to higher total acidities and reduced in size in the aqueous sulfate environment. The effect of silica activity on the weathering of kaolinite and muscovite to form bauxite; the analcite-albite relationships; and the coexistence of K-feldspar and kaolinite in alteration assemblages are all problems in which the silica activity plays an important part. Hemley has found that high silica activities cause kaolinite to react to form pyrophyllite or montmorillonite at temperatures above 200°C.

⁷⁹ A. H. Truesdell, 1962, Study of natural glasses through their behavior as membrane electrodes: *Nature*, v. 194, p. 77–79.

Clays as osmotic membranes may "float" overthrusts

Osmotic pressures of 500 bars can be obtained in clayey sediments by filtration of NaCl-saturated solutions through clays which act as osmotic membranes. E-an Zen and B. B. Hanshaw (2–65) has suggested that such situations (1) may exist in sections containing thick shale beds and evaporite deposits, and (2) could "float" sheets of unconsolidated sediments as much as 4 kilometers thick. Such mechanisms may be important in the mechanism of near-surface thrusting as suggested by M. K. Hubbert and W. W. Rubey.⁸⁰

STUDIES OF FLUID INCLUSIONS IN MINERALS

Salinity of fluid in inclusions found to increase with ore stage

Edwin Roedder and A. V. Heyl made a field and laboratory study of the conditions of ore deposition at the Mex-Tex mine, Hansonburg district, New Mexico. The ore is similar in some respects to the "coontail" ore of the southern Illinois lead-zinc-fluorite deposits. The salinity of the fluid in primary fluid inclusions progressively increased (hence freezing temperatures of the fluid decreased) throughout the first 6 of the 7 stages of ore deposition recognized. The salinity changed from about 9.5 percent NaCl equivalent brines in the inclusions in the earliest fluorite to about 17.5 percent in the 6th stage as represented by inclusions in the cores of late amethyst crystals. The 7th and last stage comprising inclusions in the outer rims of the late amethyst crystals contained only 12 percent NaCl equivalent brines. Numerous planes of secondary inclusions cutting these samples contained fluids equivalent to primary inclusions in later zones. Such a systematic increase in salinity with stage in ore deposition has never been reported before in ore deposits of any kind.

Evidence for separation of an immiscible water-rich fluid phase

A missing link in the chain of reasoning that connects hydrothermal fluids and granitic melts has long been an ignorance as to whether or not a sudden break takes place in the line of descent involving the separation of an immiscible water-rich fluid phase from a silicate-rich, fluid melt. Edwin Roedder, in cooperation with D. S. Coombs of the University of Otago, New Zealand, has found concrete evidence of such a break in the course of a study of the fluid

⁸⁰ M. K. Hubbert and W. W. Rubey, 1959, Role of fluid pressure in mechanics of overthrust faulting. I. Mechanics of fluid filled porous solids and its application to overthrust faulting: *Geol. Soc. America Bull.*, v. 90, p. 115–166.

inclusions in the minerals of ejected granitic blocks in volcanic breccias from Ascension Island, in the south Atlantic Ocean. Primary inclusions in these rocks have a phase assemblage indicating that immiscible globules of highly saline (about 50 percent NaCl) aqueous fluids with some CO₂ occurred in the rhyolitic silicate melt, and were trapped in the inclusions.

"Fetid" odor of calcite due to liquid H₂S in inclusions

Although mineral samples have been found that give off the "fetid" odor of H₂S on crushing, they are rare, and nothing unusual has ever been seen in their inclusions. It has been presumed that the odor was from exceedingly minute amounts of H₂S dissolved in the water-rich fluid inclusions. A sample of strongly fetid calcite from the U.S. Section (Lark mine, Lark, Utah) has been found by Edwin Roedder to contain inclusions with four mutually immiscible fluids, plus a crystal of NaCl. One of the fluids is believed to be liquid H₂S, on the basis of its index of refraction and phase relationships; it might easily be mistaken for water in a cursory examination. Although the evidence from fluid inclusions indicates H₂S to be a relatively minor constituent, and seldom found in hydrothermal fluids, these particular fluids had a very high concentration of H₂S.

GEOCHEMICAL DATA

Collection and synthesis of geochemical data are important parts of the program of the U.S. Geological Survey. Some types of collections, such as the "Data of Geochemistry," are of immediate value as reference works, not only to geologists but to workers in other scientific fields as well. Other collections, commonly those resulting from fieldwork, provide background for studies of geologic processes and environmental effects on animal and human health, or for programs in economic geology such as geochemical prospecting. (See also the sections "Distribution of Elements as Related to Public Health, Minor Elements," "Field Geochemistry and Petrology," "Experimental Geochemistry," "Geochemical Prospecting," and "Geochemistry of Water.")

Lithium in plants and soils of Nevada basins

Unusual amounts of lithium have been found in plants and soils from several basins in Nevada. H. L. Cannon reports that spectrographic analyses of samples of 20 species of plants from 27 sites in 9 basins has shown a content of lithium ranging from 300 to 1,500 parts per million in plant ash and as much as 3,000 ppm in soil. Concentrations of

lithium occur in springs, the salts of closed basin playas, and old lake beds. The most probable source is volcanic tuff, which is abundant in the study area. Further studies will be made of the source of the lithium and its secondary enrichment in plants and soils of both hydrologically "closed" and "open" basins.

Metal content of black shales

J. D. Vine reports that a comparison of the abundance and distribution of elements in shelf and eugeosynclinal black shales indicates that vanadium, zinc, chromium, nickel, copper, and silver are locally concentrated in both environments. Eugeosynclinal sediments were represented by analyses of samples of Ordovician and Silurian black siliceous shales from the western conterminous United States, and the shelf environment by analyses of several sets of transgressive marine black-shale samples associated with cyclothemic coal sequences. Shales of the shelf environment are especially characterized by local concentrations of molybdenum, and those of the eugeosynclinal environment by concentrations of barium. Copper and nickel show a consistent positive correlation with the organic carbon content of samples. Vanadium and chromium show a consistently strong positive correlation with each other, but the highest concentrations of these metals do not necessarily correlate with the highest concentration of organic carbon.

Extractable organic material in Cretaceous shales

H. A. Tourtelot and I. C. Frost (p. D73-D81) have extracted organic material from 5 samples of nonmarine carbonaceous shale, 10 samples of marine black shale, and 3 samples of marlstone, all from the Pierre Shale and equivalent stratigraphic units of Late Cretaceous age. The organic material was separated with activated silica and solvents into fractions interpreted as saturated hydrocarbons, aromatic hydrocarbons, and "nonhydrocarbons." Organic carbon content of the samples extracted ranges from 0.53 to 17.3 percent, but only 1 sample contained less than 2 percent. Extractable organic material content of the samples ranged from 0.008 to 0.354 percent and was roughly proportional to the organic carbon content. The ratio between saturated and aromatic hydrocarbon fractions is more than 1 except for 1 sample, and tends to be inversely proportional to the organic carbon content. The data on extractable material do not permit distinction between shales and marlstones of marine or nonmarine origins.

Comparison of the data on these samples with published data of samples with a wide range of ages and geologic histories suggests that post-depositional reactions increase the proportion of saturated and aromatic hydrocarbons compared to nonhydrocarbons.

Composition of magnetite as related to type of occurrence

Compilation by Michael Fleischer of analyses of magnetite from veins in basaltic trap rocks shows that such magnetite has a high content of MgO. Its bulk composition covers the greater part of the isomorphous series, magnetite-magnesioferrite (p. D82-D84). It is also characterized by having a much lower content of TiO₂ than magnetite from diabbases or basalts.

Magnetite from alkalic rocks and carbonatites usually has a high MgO, Al₂O₃ and MnO content; its composition is such that it would be expected to contain exsolved hercynite or spinel.

Tin content of silicate rocks

Claude Huffman, Jr., analyzed 111 plutonic, volcanic, and sedimentary rocks for tin, using the ion-exchange separation technique described in Professional Paper 501-A, p. A196, with results as follows:

<i>Rock</i>	<i>Number of samples</i>	<i>Range of tin (ppm)</i>	<i>Average tin (ppm)</i>
1. Pikes Peak Granite, Taryall lobe, Colorado-----	39	11-32	16
2. Beryliferous rhyolitic tuff, Utah-----	13	22-270	75
3. Pierre Shale, South Dakota and Montana-----	53	< 1-4	1
4. Porphyritic biotitic granite, Alaska-----	6	7-54	27

Sources of error in field geochemical data analyzed

A. T. Miesch has defined sampling and analytical error in field geochemical data, classified the types of error present, and described the effects of various error types in data interpretation by statistical and nonstatistical methods. The most serious error types are overall bias in the data, bias or precision that is variable from one sampling locality to another, and nonnormal error frequency distributions. The error effects are the same whether the error is due to sampling or analysis, or whether the method of interpretation is statistical or nonstatistical.

GEOCHEMISTRY OF WATER

Investigations of the geochemistry of water by the U.S. Geological Survey are directed mainly toward understanding the interrelations of the chemi-

cal character of water with the geologic and hydrologic environment. Some of the topics under study include the source of dissolved constituents in precipitation, the chemical content of stream water, the relation of the chemistry of ground water to the mineralogy of aquifer materials, the effects of local hydrology on the chemistry of lake sediments, and the use of isotopes in hydrologic investigations.

STUDIES OF ATMOSPHERIC PRECIPITATION

Dissolved solids in snow

R. J. Archer, A. M. LaSala, Jr., and W. E. Harding found concentrations of dissolved solids as high as 327 parts per million and of sulfate as high as 222 ppm in samples of a single snowfall occurring on February 16, 1964, in an area of 2,000 square miles in western New York, bordering Lake Erie and the Niagara River. The highest concentrations found were in samples collected downwind (east and northeast) of Buffalo.

Dissolved solids in rain

Studies by A. W. Gambell and D. W. Fisher have shown that over a 7,000-square-mile drainage area in North Carolina, for a 1-year period, rainfall contributed an amount of dissolved solids equal to one-third to two-thirds of the total dissolved solids carried by the streams. The amount of SO₄⁻² and NO₃⁻¹ brought down in rainfall exceeded that carried by the streams. The "excess" of SO₄⁻² and NO₃⁻¹ appears to be a normal occurrence for the area studied; it is thought that these constituents are taken up by plants, and that when the plants die and decay some sulfur and nitrogen are lost to the atmosphere in gaseous form.

Comparison of composition of dissolved solids in rain and in ground water

R. L. Laney (p. C187-C189) compared the chemical composition of dissolved solids in rain and in ground water in western North Carolina. The outstanding features he found are (1) an apparent lack of gradational members between typical analyses of rain water and of ground water, (2) a change from a calcium sulfate and calcium bicarbonate composition (rain) to a mixed-cation bicarbonate composition, and (3) a marked increase of silica in the ground water.

Tritium in precipitation and stream water

Concentrations of tritium in precipitation in the United States, observed by G. L. Stewart, T. A. Wyerman, and C. M. Hoffman, were in general

lower than those in 1963 but were still considerably higher than 1962 levels. Exceptions were found at Albuquerque, N. Mex., and at Lincoln, Nebr., where 1964 peaks were slightly higher than those in 1963. The highest concentrations were observed at Denver, Colo. (12,950 tritium units in 1963 and 9,700 T.U. in 1964); the lowest were at San Juan, P.R. (440 T.U. in 1963 and 290 T.U. in 1964). A spring rise and a peak in late spring and early summer were observed in 1964, as in previous years. The concentration of tritium in streams can be correlated with tritium rainout and with stream discharge and other stream characteristics. Concentrations in streams follow patterns in time similar to those in precipitation; those in 1964 were in general slightly lower than those in 1963.

STUDIES OF STREAMS AND LAKES

Saline lakes in Oregon

Studies by A. S. Van Denburgh have shown that Abert and Summer Lakes, two saline, closed-basin lakes in south-central Oregon, contain about 13 million and 0.6 million tons, respectively, of dissolved solids—mostly sodium, carbonate bicarbonate, and chloride. Salts stored within the top 5 feet of peripheral and lake-bottom sediments total about 60 million tons at Abert Lake (total area 75 square miles) and about 20 million tons at Summer Lake (85 sq mi).

Mineralogy of sediments exposed to brine

Extensive mineralogical examination by B. F. Jones of clay and fine-silt fractions from alluvial-fan materials, older lacustrine sediments, and modern playa deposits of the Deep Springs Valley, Calif., suggests that limited reaction takes place when the clay minerals are exposed for long periods to alkaline brine with a content of as much as 2½ percent K^{+1} . The apparent correlation of clay suites with sediment source materials suggests that original mica structures, after weathering, are rejuvenated by K^{+1} uptake, but that there is little effect of the brine on expandable clays originally derived from sources other than sheet-silicate minerals.

GROUND-WATER STUDIES

Exhaustion of cation-exchange capacity of glauconitic sand

Investigations by H. B. Wilder, in the Atlantic Coastal Plain in North Carolina, indicate that the cation-exchange capacity of glauconitic sands in some geologic formations has been exhausted by

hard ground water flowing through the formations. A test well in Craven County, N.C., which penetrated glauconitic Cretaceous sands of unusually high permeability, yielded water with a hardness of more than 200 parts per million (as $CaCO_3$). Normally, in this area, ground water in this sand would have less than 50 ppm hardness. Laboratory tests with samples of the glauconitic sands produced no softening of natural hard-water samples.

Estimation of dissolved-solids content of ground water

Using the many available electrical logs and chemical analyses, A. N. Turcan, Jr., developed an empirical field method for estimating the content of dissolved solids in potable water in most of the major aquifers in Louisiana. The method is based upon the averaged relation of the resistivity reading from the long normal curve on the electrical log to the laboratory resistivity reading for water from the aquifer. Results indicate that this relation, which is termed the field-formation resistivity factor, can be used for an aquifer over a large area. The factor varies with changes in the sand-clay ratio and in the basic type of water.

Uranium in ground water, Arkansas River valley, Colorado

Collection and analysis of 200 samples of ground water from the Arkansas River valley in Colorado by R. C. Scott and J. E. Moore showed that the ground water contains as much as 180 micrograms of uranium per liter, whereas water in the Arkansas River contains 2–3 micrograms per liter—the background amount of uranium for the area. Uranium concentrations in ground water from the valley-fill aquifer generally increase downstream. The uranium in the ground water is thought to have been introduced in phosphate fertilizers which have widespread use throughout the valley. Isolated anomalous highs are in areas where agricultural activity is or has been most intense and where well density is greatest. Anomalous lows are along the southern edge of the valley where ground water in the alluvium is diluted by recharge from adjacent dune sands. The downstream increase in uranium concentrations may be attributed to continuous reuse of irrigation water by successive farmers.

Velocity of ground-water flow determined from radiocarbon studies

Radiocarbon determinations on water from a series of wells in west-central Florida, made by R. N. Cherry, Meyer Rubin, B. B. Hanshaw, and William

Back, show lower concentrations (younger water) in recharge areas than in areas downgradient from the recharge areas. The calculated velocity of ground-water movement, 7 meters per year for 140 kilometers of travel, is in agreement with a velocity of 7 m per yr calculated independently from hydrologic equations.

CHEMICAL EQUILIBRIUM STUDIES

Behavior of manganese in water

J. D. Hem⁸¹ found a stability constant of $10^{4.24}$ for a 1:1 complex of divalent manganese with tannic acid. This type of metal-organic complex may influence the behavior of manganese and other metals in natural water. A similar but weaker complexing action occurs between gallic acid and manganese.

Chemical equilibria in Florida ground water

In a continuing study of chemical equilibrium between solid phases and water in the aquifer system of central Florida, William Back and B. B. Hanshaw have shown that ground water in areas of recharge in the limestone aquifer is undersaturated with respect to calcium, aragonite, dolomite, gypsum, and anhydrite; away from the recharge areas it may maintain a state of supersaturation with respect to calcite and dolomite.

Chemical equilibria in prairie potholes in North Dakota

During a study of two prairie potholes in North Dakota, H. T. Mitten (p. C176-C180) found that concentrations of dissolved oxygen and CO_3^{-2} and the ratio of the ion-activity product to the equilibrium constant for calcite varied diurnally in the water. Peak values for these three variables correlated with the peak (maximum or minimum) values of temperature. The water in both potholes was supersaturated with respect to calcite at all times.

THERMAL WATERS

Types of thermal waters in Yellowstone Park

C. L. Christ, R. M. Garrels (Harvard University), and K. L. Temple (Montana State College) analyzed samples of eight thermal waters from bubblers and pools in the area of the Artists Paint Pot, in Yellowstone National Park, Wyo. The samples were analyzed for sodium and potassium, by the electrode method in the field and by the flame-photometer method in the laboratory. Results obtained are shown in the accompanying table.

⁸¹ J. D. Hem, 1965, Reduction and complexing of manganese by gallic acids: U.S. Geol. Survey Water-Supply Paper 1667-D. [In press]

Sample	Method					
	Electrode			Flame		
	Na	K	Na/K	Na	K	Na/K
1-----	17.0	0.78	21.8	17.6	0.56	31.5
2-----	18.2	.77	23.6	18.4	.58	31.7
3-----	17.4	.72	24.2	17.5	.56	31.3
4-----	17.5	.74	23.6	18.0	.56	32.2
5-----	1.02	.79	1.29	¹ (2.42)	.58	4.17
6-----	1.50	1.42	1.06	1.32	.96	1.38
7-----	.77	.85	.90	.77	.62	1.24
8-----	² (0.91)	.83	1.09	.77	.56	1.38

Results are in milliequivalents. 1-4, alkaline waters, pH $\approx$ 9; 5-8, very acid waters, pH $\approx$ 2.7.

¹ Flame photometer value not accurate.

² Electrode value not accurate.

The agreement between results by these two methods is better for sodium than for potassium. The values obtained by the electrode method are in closer agreement with values found by E. T. Allen and A. L. Day⁸² than are those by the flame-photometer method.

Samples analyzed are representative of the waters throughout Yellowstone Park. There are two types of waters, an alkaline type and an acid type, each of which is remarkably consistent in terms of composition. This consistency suggests one common mechanism for the origin of the alkaline waters and one common mechanism for the origin of the acid waters. The alkaline waters are thought to result from the travel of solution through the rock and equilibration with it, probably at the three-phase point where montmorillonite, kaolinite, and muscovite exist at equilibrium; this would yield the Na:K:H ratio found. On the other hand, the acid waters can be explained by assuming an original solution of H_2S , which is not very corrosive toward rock, becoming oxidized to aqueous H_2SO_4 as the solution nears the surface, when it will then largely dissolve the rock except for quartz. At the surface, kaolinite will be precipitated.

INVESTIGATIONS AT THE HAWAIIAN VOLCANO OBSERVATORY

Eruption of Kilauea, March 1965

Kilauea erupted more than 30 million cubic yards of lava from fissures along nearly 8 miles of the upper east rift zone between March 5 and March 15, 1965. This adds another rift eruption to the series of 1962 and 1963 that broke from the rift without a preliminary summit eruption in the

⁸² E. T. Allen and A. L. Day, 1935, Hot springs of the Yellowstone National Park: Carnegie Inst. Washington Pub. 466, p. 453.

caldera. Each rift outbreak, however, has been associated with a short, abrupt collapse of the summit as evidenced by tilt and level observations, further substantiated this year by surface-distance measurement. The distance between two points across the summit caldera, somewhat more than 3 kilometers, was shortened during the summit collapse by 25–30 centimeters as determined by a series of tellurometer and geodimeter measurements.

This eruption filled Napau Crater to a depth of 40 feet in a few hours, but about half this drained back immediately into the vents. A lake more than 330 feet deep formed during 9 days of eruption in the west pit of Makaopuhi Crater; after draining back it still contains over 6 million cubic yards of lava in its 270-foot depth. Documentation of the cooling history of this new lake was started at once, with the benefit of previous experience on Kilauea Iki and Alae lakes.

Crystallization of Alae lava lake

The 50-foot-thick Alae lava lake, formed in the August 1963 eruption of Kilauea Volcano (D. L. Peck and others⁸³), solidified in August 1964, and its last interstitial melt crystallized in late September. Most aspects of the cooling history were watched closely by recurrent surveying, surface mapping, and core drilling in a total of 12 holes. Repeated leveling showed, unexpectedly, that the lake surface constantly rose over those parts of the lake still partly molten and subsided only over the solidified part during the year required for complete solidification. The rise was greatest (as much as 0.81 feet uplift) in a doughnut-shaped area that ringed the lake about halfway from center to shore. Uplift of the lake surface apparently was caused by continued exsolution of gas from lava undergoing crystallization; vesicles formed by the gas and trapped in the solidifying lava reduced the bulk density of the crust to less than that of the molten lava. Preliminary estimates based on the leveling data, density of drill core and a bulk thermal contraction rate of 0.001 percent per degree from leveling and temperature measurements near the margin of the lake, suggest that the melt in the lake in September 1963 had a density of about 2.76. The average density of the drill core used in the calculations, excluding the upper 7 feet, was 2.665 grams per cubic centimeter.

⁸³ D. L. Peck, J. G. Moore, and George Kojima, 1964, Temperatures in the crust and melt of Alae lava lake, Hawaii, after the August 1963 eruption of Kilauea Volcano—a preliminary report, in *Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D*, p. D1–D7.

Petrographic study of more than two dozen core and probe samples from the zone of crystallization at the base of the crust of the lake indicates a solidus temperature of $980 \pm 10^\circ\text{C}$. All core that was cooler than 980°C before drilling (disregarding that from the chilled surface of the lake) is completely crystalline except for 5 to 10 percent colorless interstitial glass with refractive index of 1.49. Core that was hotter than 980°C contains brown glass quenched from the melt, the volume percent and refractive index of the glass increasing proportionately with temperature; at 1030°C the lava contains 40 percent glass with index of 1.57, and at 1070°C , 65 percent glass with index of 1.61. The great change in refractive index may result from the late crystallization of opaque material, largely ilmenite.

All samples studied, whether pumice quenched at the point of eruption or wholly or partly crystallized core at all different depths, show inhomogeneous distribution of 1–5 percent olivine crystals. Chemical analyses of 16 samples spanning initially quenched pumice to core samples (partly crystallized and partly molten) from depths as great as 31 feet in the lava lake show no change in composition associated with position in the lake or with time of existence as a liquid after extrusion (see accompanying table). Apparently there was no

Chemical composition of material from Alae lava lake
[In weight percent]

Constituent	Maximum	Minimum	Average of 16 samples	Partly crystalline core drilled Sept. 1964, at 31-ft depth	Ooze from 19–20-ft depth, Dec. 30, 1963
SiO ₂ -----	50.51	50.32	50.44	50.47	53.30
Al ₂ O ₃ -----	13.82	13.41	13.64	13.69	12.37
TiO ₂ -----	2.85	2.63	2.73	2.71	3.36
FeO +0.9 Fe ₂ O ₃ -----	11.11	10.75	10.94	10.88	14.28
MgO -----	7.68	7.32	7.55	7.58	3.10
CaO -----	11.20	11.00	11.12	11.17	7.28
Na ₂ O -----	2.48	2.33	2.40	2.41	3.26
K ₂ O -----	.59	.53	.55	.54	1.57

detectable differential movement of plagioclase, pyroxene, or olivine crystals during the year of crystallizing. Five samples of the interstitial liquid that was filter pressed into the drill holes from the partly molten crystal mesh at the base of the crust show large changes in composition, particularly depletion of magnesia and lime and increase of silica, alkali, and titania.

Submarine pillow basalt from Mauna Kea rift zone

The U.S. Coast and Geodetic Survey hosted James G. Moore and Dallas I. Peck, U.S. Geological Survey,

in a dredging program from the research vessel *Surveyor*. Samples collected from the submarine part of the east rift zone of Mauna Kea volcano (to a depth of 3,200 meters) are fresh olivine-rich tholeiitic pillow basalt. In contrast is the cap of lava and cinder cones of alkalic basalt and hawaiite that covers, and may be restricted to, the upper exposed flanks and summit of Mauna Kea.

The vesicularity of the submarine samples from Mauna Kea is greater than the vesicularity of similar samples collected from the same depth on the submarine rift zones of the currently active Mauna Loa and Kilauea. This suggests that Mauna Kea has sunk about 1,000 meters relative to sea level since these vesicular submarine lavas erupted.

Seismic-refraction profiles yield crustal thicknesses at Hawaii

Three seismic-refraction profiles were investigated along the northeast, southeast, and west coasts of the triangular-shaped island of Hawaii. Two hundred twenty five seismograms were recorded by 5 refraction units spaced at about 25 kilometer intervals along each coast from 300-pound shots fired at 10-km intervals from the U.S. Coast Guard cutter *Cape Small*. Preliminary interpretation, by David P. Hill of the Hawaiian Volcano Observatory, indicates that the crust is about 16 km thick under the west flanks of Mauna Loa and Hualalai, is 11 km thick under the northeast and southeast flanks of Kilauea, and has an intermediate thickness along the northeast flanks of Mauna Kea and the Kohala Mountains. The velocity of P waves in the upper crust increases with depth from 2.0 to as much as 6.0 km/sec; velocities in the upper crust are generally lower on the flanks of Kilauea than on the flanks of the other volcanoes. Clearly recorded arrivals indicate that a layer with velocities from 6.9 to 7.3 km/sec forms the lowermost 4–8 km of the crust under each of the coasts. The velocity of P_n under each of the coasts is about 8.2 km/sec. Anomalously high crustal velocities are associated with the major rift zones extending from the five volcanoes that form the island. This interpretation of the refraction data is consistent with gravity data on the island of Hawaii.

ISOTOPE AND NUCLEAR STUDIES

GEOCHRONOLOGY

An interlaboratory standard muscovite for K-Ar analyses

Approximately 1,100 grams of an 81-million-year-old muscovite, P-207, has been prepared by M. A.

Lanphere and G. B. Dalrymple (1-65) as an interlaboratory standard for argon and potassium analyses. Twelve argon analyses and 10 potassium analyses made in the U.S. Geological Survey laboratory give mean values of 1.253×10^{-9} moles $\text{Ar}^{40}_{\text{rad}}/\text{gram}$ ($\sigma_x = 0.002$) and 10.19 percent K_2O ($\sigma_x = 0.03$).

This standard sample has been distributed to 21 age determination laboratories in Australia, Canada, England, Japan, Switzerland, and the United States. To date, 11 laboratories have reported the results of their analyses. The mean values of potassium and argon analyses in 11 laboratories and of rubidium and strontium analyses in 3 laboratories are:

$\text{Ar}^{40}_{\text{rad}} = 1.267 \times 10^{-9}$ moles/g ($\sigma_x = 0.009$); $\text{K}_2\text{O} = 10.34$ percent ($\sigma_x = 0.05$); $\text{Rb} = 9.37 \times 10^{-6}$ moles/g; $\text{Sr}^{87}_{\text{rad}} = 3.17 \times 10^{-9}$ moles/g; and common $\text{Sr} = 0.113 \times 10^{-6}$ moles/g.

Effect of weathering upon U-Th-Pb system in zircon

In an investigation of the possible effects of weathering on the U-Th-Pb system in zircon, T. W. Stern, S. S. Goldich, and M. F. Newell analyzed 3 samples of zircon from residual clay formed by weathering of the granite gneiss in the vicinity of Morton, Minn., and 5 samples of zircon from the fresh gneiss. The latter are somewhat discordant but define a chord intersecting the concordia curve at 3,600 million years and 1,900 m.y. The zircon samples from the residual clay all show lead loss of 50 percent or more, with the greatest loss for samples with the greatest content of uranium and thorium. A plot of these data on a concordia diagram clearly shows the superimposed effect of weathering on this zircon population. Geologic evidence shows that the loss of lead from zircon at Morton, Minn., came about only after prolonged weathering; nevertheless, the results emphasize that this factor should not be ignored, especially in studying zircon from metasedimentary rocks which may have gone through one or more sedimentary cycles.

Variations in $\text{U}^{238}/\text{U}^{234}$ and $\text{U}^{235}/\text{U}^{234}$ ratios of zircon

Zircon concentrates representing 13 distinct occurrences were analyzed for isotopic abundance of U^{238} , U^{235} , and U^{234} by Bruce Doe and M. F. Newell. Twenty determinations by mass spectrometric techniques and 5 cross checks on $\text{U}^{238}/\text{U}^{234}$ by alpha spectrometry show that the $\text{U}^{238}/\text{U}^{235}$ ratios do not vary more than 0.5 percent and that the $\text{U}^{235}/\text{U}^{234}$ ratio variation is 9 percent. Variations of $\text{U}^{235}/\text{U}^{234}$

cannot be correlated with age, grain size, percent discordance in isotopic ages, type of discordance (positive or negative), uranium content, or radiation damage. Sample inhomogeneity or chemical procedures may account for the nonreproducibility of U^{235}/U^{234} ratios in several of the samples. The range in U^{235}/U^{234} is not great enough to account for discordant U-Pb ages in zircons resulting from disequilibrium decay of U^{234} . Since most zircon concentrates are obtained from rock samples near the earth's surface, the weathering effect could be estimated by the U^{235}/U^{234} ratio; however, our observations suggest that near-surface incipient weathering has little or no effect on zircon.

Old age of Morton and Montevideo gneisses confirmed

An extensive dating study by C. E. Hedge of the granitic gneiss and associated plutonic rocks at Morton, Minn., has indicated a geologic history both long and complex. Whole-rock Rb-Sr ages of the Morton and Montevideo granite gneisses form a spectrum ranging from 2.5 billion years to 3.5 b.y. The older age limit is in agreement with earlier zircon age determinations by E. J. Catanzaro⁸⁴ and T. W. Stern,⁸⁵ and the 2.5-b.y. age represents a time of major metamorphism and plutonism as indicated by both K-Ar and Rb-Sr mineral ages. The Sacred Heart Granite yields concordant Rb-Sr, K-Ar, and U-Pb zircon ages of 2.5 b. y. Late granite is dated at 1.7 b.y. by all methods.

Rb-Sr dating of argillaceous metasedimentary rocks

Whole-rock Rb-Sr dating studies of Precambrian argillaceous metasedimentary rocks being conducted by Z. E. Peterman strongly suggest that such rock types do not remain closed systems after deposition and lithification. Those sedimentary rocks which have been metamorphosed in the upper greenschist facies or higher facies give ages which can reasonably be attributed to the time of metamorphism. Other sedimentary rocks which have not been appreciably metamorphosed yield ages which are significantly too young both for an age of deposition and an age of metamorphism.

Age of alkalic intrusive rocks from eastern and midcontinent States

R. E. Zartman and R. F. Marvin in cooperation with A. V. Heyl and Maurice Brock are continuing a geochronologic study of mica peridotites and re-

lated alkaline intrusive rocks from the eastern and midcontinent states. Biotites and phlogopites were dated by both the K-Ar and Rb-Sr methods with the following results:

(1) The syenitic bodies which intrude Ordovician rocks near Beemerville, N. J., yield a Late Ordovician age of 440 ± 20 million years.

(2) Diatremes of mica peridotite which cut Cambrian rocks in southeastern Missouri give an Early Devonian age of 385 ± 20 m.y.

(3) Dikes, sills, and explosion breccia in Mississippian carbonate rocks from the Rosiclare fluorite district of Illinois and Kentucky have an Early Permian age of 265 ± 15 m.y.

(4) Isolated mica peridotites, nepheline syenites, and alkalic lamprophyres occurring in various areas of Paleozoic rocks give Jurassic to Cretaceous ages. Included in this latter group are intrusions having an age of approximately 140 ± 10 m.y. along the western side of the Appalachian Mountains in Vermont, New York, Pennsylvania, and Virginia, and also distinctly younger alkalic rocks of Magnet Cove, Ark., and Silver City, Kans., where ages of 95 ± 5 m.y. are obtained.

Ground waters increase in age toward Great Salt Lake

Ground water along the line of flow from the Stansbury Mountains recharge area toward the southern end of Great Salt Lake, Tooele County, Utah, has been sampled by A. B. Tanner, and the radiocarbon content of the bicarbonate in seven of the water samples has been determined by Meyer Rubin. The results when plotted on a map show a progressive increase in apparent age down the alluvial fan toward Great Salt Lake. The rate of ground-water movement along the flow line averages 4.5 feet per year in the region of low salinity (<100 parts per million of dissolved solids) but decreases abruptly at the contact with highly saline waters here considered remnant from glacial Lake Bonneville. The apparent age of the water ranges from 2,000 years in the recharge area to more than 30,000 years in the highly saline area.

List of additional age determinations

A considerable part of the work of the geochronology laboratory of the U.S. Geological Survey is carried on in cooperation with Survey field geologists. Results of such service investigations throw light on local geologic problems and are reported under appropriate headings elsewhere in this chapter. A tabulation of these determinations for use in cross reference follows:

⁸⁴ E. J. Catanzaro, 1963, Zircon ages in southwestern Minnesota: *Jour. Geophys. Research*, v. 68, p. 2041-2048.

⁸⁵ T. W. Stern, 1964, Isotopic ages of zircon and allanite from the Minnesota River valley and La Sal Mountains, Utah [abs.]: *Am. Geophys. Union Trans.*, v. 45, no. 1, p. 116.

Samples dated	Method	State or area (from east to west by regions shown in table of contents)	Subdivision shown in table of contents
Oyster shells ----- Metamorphic and igneous rocks -----	C-14 Rb-Sr K-Ar U-Th-Pb Pb- α C-14 C-14	Atlantic shelf ----- Connecticut -----	Marine Geology and Hydrology. New England and Eastern New York.
Matrix, muskox skull ----- Spruce wood -----	C-14 C-14	Kentucky ----- Wisconsin -----	Paleontology. Shield Area and Upper Mississippi Valley.
Vitrophyre -----	K-Ar	Colorado -----	Southern Rocky Mountains and Plains.
Syenite -----	U-Th-Pb K-Ar	Idaho -----	Northern Rocky Mountains and Plains.
Silicic intrusives ----- Intrusives and extrusives ----- Variety of materials ----- Salmon and Abrams Schists ----- Bishop Tuff ----- Wood ----- Hornblende, igneous ----- Hornblende, biotite, igneous ----- Organic material ----- Mollusk shells -----	K-Ar K-Ar C-14 K-Ar K-Ar C-14 K-Ar C-14 Th-U disequilib- rium Rb-Sr K-Ar Pb- α K-Ar C-14 C-14	Nevada ----- Arizona ----- Washington ----- California ----- do ----- do ----- Alaska ----- do ----- do ----- do ----- Antarctica ----- do ----- do ----- Chile ----- Greenland ----- do -----	Basin and Range Region. Do. Pacific Coast Region. Do. Do. Do. Alaska (northern). Alaska (southwestern). Alaska (southern). Alaska (west-central). Antarctica. Do. Do. Geologic and Hydrologic Investiga- tions in Other Countries. Do. Do.
Whole rock, granitic ----- Biotite from granite ----- Zircon from quartz diorite ----- Ash flows ----- Turf in glacial moraine ----- Pelecypod shells -----	Rb-Sr K-Ar Pb- α K-Ar C-14 C-14	Antarctica ----- do ----- do ----- Chile ----- Greenland ----- do -----	Antarctica. Do. Do. Geologic and Hydrologic Investiga- tions in Other Countries. Do. Do.

RADIOACTIVE DISEQUILIBRIUM

Fractionation of uranium isotopes in sandstone

Isotopic ratios of U^{234}/U^{238} for 14 samples representing a sandstone-type uranium deposit that formed at, and just above, the present level of standing water in the Wentz mine, Natrona County, Wyo., show that the deposit has a relatively consistent U^{234} deficiency ranging from 7 to 29 percent (J. N. Rosholt, Jr., and C. P. Ferriera, p. C58-C62). Th^{230}/U^{234} and Pa^{231}/U^{235} ratios indicate that the mean time interval since uranium emplacement near the level of standing water occurred about 80,000 years ago and that the time interval generally is greater with increasing distance above standing water. The source uranium also appears to be deficient in U^{234} ; this deficiency was maintained by a relatively continuous preferential leaching of U^{234} .

Isotopic composition of uranium in surface water

Determinations of uranium isotopes in surface water that drained diverse terranes on opposite sides of the Shirley Basin, Wyo. indicated differences in the isotopic composition of uranium derived from different sources. R. C. Scott and K. W. Edwards found a considerable enrichment of U^{234} with respect to U^{238} in water from granitic terranes, two samples of which had an average U^{234} excess of 50 percent

above the radioactive equilibrium composition, whereas water that drained sedimentary terranes was characterized by only a slight excess of the U^{234} isotope.

Migration of uranium in soil profiles

Data collected on the isotopic composition of uranium and thorium in soil profiles provided information for interpreting the causes of uranium migration in soils and for constructing a tentative model to explain the isotopic evolution of uranium and thorium in this environment. The model proposed by J. N. Rosholt, Jr., and J. R. Dooley, Jr., indicates that (1) uranium was leached at depth in the profile, (2) preferential leaching of U^{234} continued with time in the subsoil, and (3) upward capillary migration of a fraction of the uranium with a higher than normal U^{234}/U^{238} ratio tended to make uranium of high U^{234} ratio available for isotopic exchange in upper soil horizons and for assimilation in organic complexes in surface soils which are rich in organic matter. Thus some of the organic-rich surface soils, which have had considerable time to develop, contain uranium with excess U^{234} compared to U^{238} . Interpretations from this model suggest that remixing of radioisotopes by geochemical processes can play a significant part, in addition to the usual physical processes of radio-

active growth and decay, in the production of an isotopic composition near radioactive equilibrium.

STABLE-ISOTOPE INVESTIGATIONS

LEAD

Isotopic composition of lead in basalt

The systematic study of the isotopic composition of lead in basalt by Mitsunobu Tatsumoto has included oceanic tholeiites from the Mid-Atlantic ridge and East Pacific rise, and basalt flows from the marine Tertiary sediments of the Newport Embayment, Oregon. The isotope composition of lead from these various samples is nonuniform, giving the following ranges in ratios:

Pb^{206}/Pb^{204} -----	17.8 -18.8
Pb^{206}/Pb^{207} -----	1.15- 1.20
Pb^{208}/Pb^{204} -----	36.6 -38.4
U^{238}/Pb^{204} -----	5 -13
Th^{232}/U^{238} -----	0.9 - 2.9

One of the tholeiite samples dredged from near Ascension Island, in the Atlantic Ocean, contains the least radiogenic lead among those basalts so far studied. The lead from tholeiites within the Miocene sediments of Oregon has an isotopic composition similar to basalt of the Columbia River Group, but the alkali basalt from this Oregon section shows large radiogenic lead aberrations similar to those of the Iwo-Jima volcanic rocks.⁸⁶

Lead isotopes in feldspar from Boulder batholith

Recent work by Bruce Doe, R. I. Tilling, and M. R. Klepper has been completed on the isotopic composition of lead in feldspar from the major intrusive units of the Boulder batholith, Montana. The ratios of lead isotopes (given in the following order: (a) Pb^{206}/Pb^{204} , (b) Pb^{206}/Pb^{207} , (c) Pb^{208}/Pb^{204}) from these units range widely [(a) 17.0 to 18.4, (b) 1.095 to 1.171, and (c) 38.0 to 39.0] and show unusual relationships between separate intrusive units. These units are internally uniform isotopically and contain B-type lead anomalies. Lead-isotope ratios characteristic of the oldest major intrusive granodiorite to the south [(a) 16.99, (b) 1.0948 and, (c) 38.04] are drastically different from those typical of the Unionville Granodiorite [(a) 18.08, (b) 1.150, and (c) 38.51] (Adolph Knopf⁸⁷), the oldest intrusive unit some 80 kilometers to the north. These two units, though similar in appearance and intrusive sequence, appear to be distinct in their genesis as testified by the lead-isotope ratios. Either these

geographically separate intrusives were generated by different magmas having distinct ratios, or the magmas came from the same source and subsequently acquired distinct ratios. The Butte Quartz Monzonite and its contained ores, younger associated granites, and the Clancy Granodiorite (Adolph Knopf⁸⁷) make up a uniform isotopic unit that includes 75 percent of the exposed batholith. This uniform unit gives the following average ratios: (a) 18.00, (b) 1.46, and (c) 38.54. Further interpretation of the lead-isotope data confirms the geologic observations that similar isotopic intrusive units were derived from a common source by differentiation.

Isotopic variation of ore lead in Colorado Rockies

Collaborative studies of A. P. Pierce, R. S. Cannon, Jr., and J. C. Antweiler on the isotopic nature of lead from the Colorado mineral belt have revealed an extremely complicated and yet unresolved geologic relationship that produces lead-isotope variations. Isotopic composition of these ore leads differs systematically in an unusual manner from those of ordinary ore-leads. The principal ores are low in uranium derived lead. Pb^{206}/Pb^{204} and Pb^{207}/Pb^{204} ratios are 18.37 to 17.57 and 15.71 to 15.58, respectively. Low content of uranium-derived lead gives the Laramide ores "model" age values that range continuously from 0 to as much as 750 million years. This distinctive isotopic variation has been named "Colorado mineral belt type" ore lead. Such unusual isotopic variations are believed to result from partial remobilization of lead from Precambrian rocks; however, other alternate hypotheses require testing.

One episode of Precambrian mineralization in the Rocky Mountains of Colorado is characterized by "Grenville"-type ore lead with a model age of about 1,650 m.y. Ore lead of this same approximate composition has been found abundantly in major ore deposits in other States in the Rocky Mountains region and elsewhere in the world. This statistical correlation may be considered encouraging to future search for a major deposit of this type even though other kinds of geologic evidence have not heretofore stimulated much exploration for such deposits in Colorado and Wyoming.

OXYGEN AND TRITIUM

Oxygen isotopes in carbonates from Pierre Shale

Shells of many specimens of cephalopods collected by H. A. Tourtelot (baculites, ammonites, nautiloids) from the Pierre Shale still consist of original arago-

⁸⁶ M. Tatsumoto, 1964, Isotopic composition of lead in volcanic rocks from Japan, Iwo-Jima, and Hawaii [abs.]: *Am. Geophys. Union Trans.*, v. 45, p. 109.

⁸⁷ Adolph Knopf, 1957, The Boulder batholith of Montana: *Am. Jour. Sci.*, v. 255, p. 81-103.

nite. The inner nacreous layer of many specimens of *Inoceramus* also consists of original aragonite, and the outer prismatic layer is calcite. Preservation of aragonite from Late Cretaceous time seems to depend on the specimens having been sealed in early diagenetic concretions or buried in claystone. Oxygen-isotope analyses by R. O. Rye of aragonite from baculites give temperatures of 15°–27°C, which bracket the temperatures determined on two specimens of Pierre belemnites (17°–22°C). Aragonite and calcite from individual *Inoceramus* valves have the same isotopic composition, but indicate temperatures of 29°–37°C. The temperatures indicated by nektonic baculites and other cephalopods are considerably lower than those indicated by benthonic pelecypods even where the cephalopods and pelecypods come from the same locality. This contrast between nekton (belemnites) and benthos has been reported before in material from widely scattered localities (mostly European). It has been attributed to a temperature control whereby the benthos grew shell only within the warmer part of the local temperature range while the nekton grew shell in other regions of the Cretaceous seas where the temperatures were markedly lower. Such an explanation is not satisfying for the Pierre material because it implies that present-day equatorial temperatures north of lat 40° N. existed in Cretaceous time, and there are no unknown parts of the Pierre sea basin in which temperatures as low as 15°C can be postulated.

Tritium fractionation

Fractionation of tritium occurs as a result of many physical and chemical processes. Such isotopic fractionation varies from only a few percent, as in liquid-vapor equilibria systems, up to a factor of 30 percent or more as in the electrolysis of tritiated water. G. L. Stewart has observed that isotopic fractionation occurs during the diffusion of water through porous materials. Laboratory investigations have shown that the degree of fractionation depends upon the tenacity with which water is held to mineral surfaces. Preliminary results show that tritiated water held at low tensions has a greater tritium concentration than water held at higher tensions. It has also been observed that fractionation occurs in different degrees depending upon the temperature at which water is removed from clay systems. Tritiated water extracted at various pressures and temperatures from sand and glass beads showed considerably less fractionation than tritiated water extracted from the clays kaolinite, illite, and montmorillonite.

HYDRAULIC AND HYDROLOGIC STUDIES

SURFACE WATER

Hydraulic studies of open-channel flow have helped define relationships among velocity, depth, slope, and roughness in straight and meandering channels, and have shed new light on the mechanics of transport and dispersal of solutes and solids in the flow.

Hydrologic studies have helped show how annual runoff, storm runoff, and peak flow are affected by natural and manmade features in the drainage basins.

Flow in open channels

Jacob Davidian found that an expression containing depth of flow (D), median grain size (d_{50}), and Froude number (F) gives some insight into scale effects in alluvial channels ranging in size from laboratory flumes to large rivers. The expression, which is $F^4(D/d_{50})^{3/2}$, has been used to predict regimes of flow (type of bed configuration) and to explain the inverse relation known to exist between Froude number and channel size in meandering streams.

Dispersion in open-channel flow

Although previous work by Nobuhiro Yotsukura showed that in flume experiments the longitudinal dispersion coefficient was a function of the Reynolds number, he has found the relation to be of little use in predicting the longitudinal-dispersion coefficient in natural channels, because the nonuniformity of the flow makes it nearly impossible to evaluate the convective effect. The lateral dispersion coefficient, on the other hand, is relatively free from the convective effect, and the formula $L = W^2/\beta D \sqrt{8f}$, which he developed from flume data, allows the prediction of the lateral dispersal length with reasonable accuracy. In this formula L represents the downstream distance necessary for complete lateral dispersal, and W , D , and f are the width, depth, and friction factors, respectively; β is a coefficient which ranges from 0.3 to 1.0 in natural streams.

Discharge coefficients for flow in open channels

Resistance to flow in alluvial channels is intimately related to the bed configurations that form there. Using laboratory and field data, D. B. Simons and E. V. Richardson (2–65) have defined the Chezy discharge coefficient, C , for plane bed conditions as $C/\sqrt{g} = 7.4 \log (D/d_{85})$; in this formula g is the force of gravity, D is the depth of flow, and d_{85} is the grain diameter which is coarser than 85 per-

cent of the grains on the bed. To determine the Chezy coefficient for ripple, dune, or antidune bed configuration, they applied a factor to the plane bed equation to compensate for the energy dissipation resulting from form roughness and from wave acceleration and deceleration. The correction factor was defined, on the basis of laboratory and field data, as a function of bed material size, slope, depth, and bed configuration.

Flow in curved channels

C. H. Hannum computed discharge for a concrete-lined flume by using the superelevation of the water surface in a bend as a measure of the velocity head. His results differed from discharge measured with a current meter by an average of only 6 percent. The flume was 40 feet wide, with vertical walls and an 8½-foot trapezoidal center section, and thus is comparable to a main channel with flood plains on each side. The bends were 36°, 47°, and 62°. Fifteen of the 18 determinations agreed with the measured discharge within 10 percent.

On the basis of a study of flow in seven bends of a trapezoidal concrete-lined canal, D. B. Simons and E. V. Richardson found that the usual method for designing sinuous canals based on the energy-loss concept is inadequate. A design method was developed which considers the effect of each bend on backwater and superelevation as well as on possible wave height.

Flow under ice cover

K. L. Carey found the configuration of the underside of the ice on the Saint Croix River near Danbury, Wis. to resemble the ripples and dunes found in streams with sandy bottoms. The dunes and ripples are oriented generally transverse to the flow and extend for half a foot to at least 4 feet before merging into similar forms that are out of phase with their neighbors. The ice dunes have an average wave length of 0.65 foot and an average amplitude of 0.07 foot. The configuration, which varies slowly with time, is assumed to be due to the turbulent, thermal, and hydraulic characteristics of the flow, because the suspended-sediment load is virtually nil.

Backwater in open channels

Although the pool elevation at a dam 29 miles downstream now submerges the low-water control at a bridge on the Neosho River in northeastern Oklahoma by 3 feet, L. L. Laine has found that the stage-discharge relation for high discharges is the same as before the dam was built; and that the

dam causes no decrease in the net operating head on the turbines at Pensacola Dam, half a mile upstream from the bridge, when the turbine discharge exceeds 7,000 cubic feet per second. From this he concludes that backwater in the tailrace of a power plant at low flow does not necessarily indicate a decrease in the net head on the turbines under operating conditions. The conveyance of the channel downstream from the bridge is obviously such that backwater from the downstream dam does not extend this far upstream at high discharges.

Rate of movement and dispersion of solutes in streams

Measurements of the time of travel of soluble contaminants were made in many stream reaches. A tracer, Rhodamine-B dye, is introduced into the stream and is detected with a fluorometer as it is carried successively farther downstream. The distribution of concentration in the dye cloud as it passes a monitored cross section is obtained from readings of the fluorometer repeated at short intervals or from a record of continuous monitoring. Time of travel in a reach, which is usually taken as the elapsed time between peak concentrations at the ends of the reach, varies with the discharge of the stream. For example, P. W. Anderson and G. M. Horwitz measured travel times of 5 days and 12 days at moderate and extremely low discharges, respectively, through a 31-mile reach of the Passaic River, N.J. Average velocities of peak concentrations of dye ranging from 0.3 feet per second to 0.5 feet per second were reported by J. D. Shell in 2 reaches of Big Black River, Miss.

Dye was introduced near one bank of the Missouri River at Jefferson City, Mo., when the discharge of the river was 37,000 cubic feet per second, which is the normal flow for the navigation season. Samples were taken at 5 lateral points at each of 4 sampling sites in a 76-mile reach. M. S. Petersen reported that lateral dispersion of the dye progressed slowly. The first sampling site to show dye throughout the cross section was 26 miles downstream from Jefferson City. Velocities in the sub-reaches ranged from 2.9 to 3.6 feet per second.

J. R. Williams injected dye into a tide-affected reach of the Duwamish River, Wash., and measured dye concentration continuously for 67 hours. Four peaks of concentration were recorded at one site, each peak concentration being appreciably less than the preceding one.

J. F. Wilson and W. E. Forrest measured the time of travel of the Potomac River in the 192-mile reach from Cumberland, Md., to Washington, D. C.

Discharge ranged from 400 cfs at Cumberland to 5,000 cfs at Washington. The time of travel was 15.6 days.

W. W. Evett compared measured time of travel through a 26-mile reach of the Catawba River, S. C., with that computed from velocities at 34 surveyed cross sections. Agreement between the two was good for the total reach, but variations of about 15 percent were found for subreaches.

Analysis of data obtained by W. W. Sayre and D. W. Hubbell in a Colorado canal demonstrated convincingly that the longitudinal dispersion of low concentrations (less than 500 parts per million) of colloiddally suspended bentonite particles did not differ significantly from the longitudinal dispersion of dye in the same flow.

In a series of flume experiments W. W. Sayre and F. M. Chang showed that the longitudinal dispersion coefficients for dye and suspended silt-size particles in open-channel flow were on the order of 10 times as large as the longitudinal dispersion coefficients for small polyethylene particles floating on the surface.

Effect of reservoirs on time of travel

Also using Rhodamine-B dye as a tracer, D. F. Farrell found that 10 low dams in a 50-mile reach of the Housatonic River in western Massachusetts so slowed the velocity of water that 12 days were required for the dye to travel through the 50-mile reach. River discharge ranged from 25 cubic feet per second at the injection point to 145 cfs at the sampling point farthest downstream.

T. J. Buchanan and J. E. McCall found that pools above the dams on the South Branch of the Raritan River downstream from Stanton, N.J., tend to speed the velocity of the flood wave and to slow the movement of water particles, so that the flood wave traveled 2.0 to 2.7 times as fast as the water as compared with 1.1 to 1.6 times as fast in the natural stream channel above Stanton. These figures confirm the expected effect of reservoirs, and provide factual data for use in studying the movement of contaminants in stream channels and for scheduling the release of water from upstream reservoirs for water supply 30 miles downstream.

Effect of reservoir release on stream temperature

P. W. Anderson and G. M. Horwitz found that following the release of water in July 1964 from Pepacton Reservoir, N.Y., on the East Branch of the Delaware River, the temperature of the river water 30 miles downstream decreased by about 20°F. A simultaneous drop in air temperature ac-

counts for only about 5°F of this decrease in stream temperature. At a point 58 miles downstream the water temperature decreased 10°F 3 days later.

Time-of-travel characteristics

Using field measurements of the time of travel of water in White Oak Creek in Tennessee, and relating the results to mean velocity, discharge, and the flow-duration curve, W. M. McMaster found in 4 out of 5 trials that the time required for water to travel through a 1.6-mile reach of the creek is between 2.5 and 5 hours. The median travel time in this creek, which drains from the Oak Ridge National Laboratory area, is 3.7 hours for the 1.6-mile reach.

Effect of urbanization on peak flow

In analysis of how change in land use affects peak discharge, the effect on the time distribution of storm runoff is usually more important than the effect on the percentage of the precipitation that becomes runoff. J. R. Crippen (p. D196-D198) studied discharge hydrographs of Sharon Creek, near Menlo Park, Calif., for storms in 1959, 1960, and 1963. Urbanization of the 0.38-square-mile basin between 1960 and 1963 brought the change from natural ground cover to suburban residences, light commercial facilities, and a golf course. Change in the distribution of runoff, as a result of this change in land use, increased the peak discharge by about 40 percent, and decreased by a similar amount the time required for nine-tenths of the storm runoff to leave the basin.

L. A. Martens found that a formula previously developed for streams near Washington, D.C., for relating mean annual flood to basin characteristics, is applicable to streams in the Piedmont region near Charlotte, N.C. The formula, by R. W. Carter,⁸⁸ is

$$Q_{MAF} = 223A^{0.85}T^{-0.45}K,$$

in which Q_{MAF} is the mean annual flood in cubic feet per second, A is the area of the drainage basin in square miles, T is the basin lag time in hours, and K represents relative imperviousness. Martens used a second formula, developed by D. G. Anderson, to determine T without the use of rainfall records. This formula is

$$T = 4.18(L/\sqrt{S})^{0.52},$$

where L is stream length and S is channel slope. The relation between mean annual floods and drainage area determined with Carter's formula agreed closely with a similar relation defined by gaging-station records. Because the difference was only 10 percent at 1 square mile and zero at 285 square miles,

⁸⁸ R. W. Carter, 1961, Magnitude and frequency of floods in suburban areas: Art. 5 in U.S. Geol. Survey Prof. Paper 424-B, p. B9-B11.

Martens concludes that the formula can be used to estimate the effect of urbanization on peak near Charlotte by substituting empirically determined values for T and K .

Effect of saltcedar on channel characteristics of Pecos River

Physical effects of increase in the area covered by the phreatophyte saltcedar (*Tamarix gallica*), in and along the Pecos River in Texas near the New Mexico boundary, described by R. U. Grozier (p. B175-B176), include a 35-percent reduction in channel conveyance, observed in the records of floods in August 1916 and June 1937, and a 15-percent reduction in channel cross section between December 1932 and March 1963.

Rainfall-runoff relations

E. E. Schroeder found that storm runoff from a 75-square-mile basin in northeastern Texas can be graphically related to storm rainfall, with good results, by using as factors (1) duration and amount of the rainfall, (2) season of the year, and (3) an index of soil-moisture conditions. The index of soil moisture was computed from antecedent precipitation adjusted by a factor of 0.9^t in which t is the number of days since the precipitation.

Effect of storage on runoff

F. W. Kennon and C. T. Welborn found that 11 floodwater-retarding structures in Texas reduce streamflow by an average of 34 percent during years of average precipitation (32 inches) and by about 10 percent when the annual rainfall is 40 inches. For years in which the annual precipitation is less than 20 inches all the runoff is usually retained in the structures. Evaporation losses account for about half of the reduction, and seepage and evapotranspiration losses for the other half.

Reservoir storage for flood reduction

Data on the volume and frequency of flood runoff at stream-gaging stations are used to determine the size of storage reservoirs required to control the amount of flow. By analyzing annual flood volumes at stream-gaging stations in Kansas, L. W. Furness, C. V. Burns, and M. W. Busby (1-64) found that for a 4-percent chance of inadequacy, a reservoir with a capacity of a 1-inch depth of runoff from the basin could hold the release rate to 0.05 cubic feet per second per square mile in western Kansas, as compared with 30 to 80 cfs per sq mi for small streams in eastern Kansas. By using the relations they developed, similar computation of reservoir capacity, flood outflow, and chance of inadequacy can be made

on any stream in Kansas that drains more than 100 square miles.

Relation of flow variability to geology

For drainage areas of 25 to 100 square miles in New Jersey, E. G. Miller found that an index of the slope of the duration curves of daily discharge could be used to classify the variability of the streamflow in different types of terrain. For the Coastal Plain he found the ratio of the discharge at the 20- and 80-percent flow-duration points to vary only from 2.2 to 3.6, as compared with much higher and more variable ratios in other parts of the State. Ratios of 15 to 20 for areas underlain by Triassic formations, in southern Hunterdon and northern Mercer Counties in northwestern New Jersey, were the highest in the State.

Estimation of mean runoff from ungaged mountain basins

H. C. Riggs and D. O. Moore (p. D199-D202) used a relation between surface runoff and basin area, for mountain basins in northern Nevada and southern Idaho, to estimate mean runoff from ungaged basins nearby. Because the magnitudes of both precipitation and evaporation in this mountainous region are related to altitude, runoff data used in the runoff-area relationship were for zones of altitude within each basin.

Time of low flow

Although some of the heaviest storm rainfall in Hawaii occurs during the cooler months of November to April, G. T. Hirashima found that for streams whose basins have little surface or ground-water storage the lowest flows of the year often occur during these months. He attributes this phenomenon to the fact that many periods of consecutive days with little or no rain occur in these months of decreased trade-wind activity.

Base-flow relations

In relating the base flow of small streams near Oak Ridge, Tenn., to that at nearby stream-gaging stations, W. M. McMaster found that satisfactory correlation could be made using the flow on concurrent days which were preceded by at least 10 days without rises. In 4 years of record he found more than 20 recessions for which this limitation was satisfied, and that the resulting correlation coefficient for 4 pairs of stations ranged from 0.91 to 0.97.

GROUND WATER

Research on ground water continued across a broad front, the range in subject matter being illus-

trated, on the one hand, by experiments on water movement in clay involving flows of as little as a few cubic centimeters per day, and on the other hand by studies of the influence on ground-water flow and chemistry of structures in lava rocks covering tens or hundreds of square miles. Temperature was recognized as a property having greater hydrologic significance than has been generally thought. There was renewed effort to correlate ground-water availability with topographic criteria as a method of prospecting. Seismic and other fluctuations of water levels in wells were scrutinized for their hydrologic significance. Substantial efforts continued in laboratory and field research on permeability and specific yield of porous media and on the usefulness of radioactive tracers. Carbonate rocks continued to receive attention as aquifers of both great economic importance and great hydrologic complexity. Both research on, and routine application of, electric-analog modeling of ground-water systems continued. Research on possible applications of "remote sensing" of hydrologic phenomena from the air was accelerated during the year. Techniques were developed for handling several types of problems involving recovery of fresh water in areas where salt-water encroachment is a threat.

Relation of permeability to particle size in a glacial-outwash aquifer

S. E. Norris and R. E. Fidler (p. D203-D206) investigated the relation between particle size and permeability in a glacial-outwash aquifer at Piketon, in southern Ohio. The aquifer contains two types of material, (1) medium to coarse sand, and (2) coarse sand to coarse gravel, which are characterized by different kinds of particle-size graphs. The first type of material, which is relatively less permeable, has an average uniformity coefficient of 6; the second type, more permeable, has an average uniformity coefficient of 13.9.

Darcy's law and water movement in kaolinite

H. W. Olsen (1-65) measured the flow of water through rigidly confined samples of kaolinite, as a followup of work that partially explained apparent deviations from Darcy's law in clays as due to atmospheric contamination of capillary tubes.^{88a} The new experiments, using a system in which the contamination problem is avoided, confirmed the validity of Darcy's law at hydraulic gradients of 0.2-40 in samples consolidated under loads of 0.5-350 kilograms per square centimeter.

^{88a} H. W. Olsen, 1965, Deviations from Darcy's law in saturated clays: Soil Sci. Soc. America. [In press]

Structural control of ground water in basalt

Studies by R. C. Newcomb of the effects on ground water of tectonic structures in basalt of the Columbia River Group in Washington, Oregon, and Idaho have confirmed preliminary conclusions that flow tends to be concentrated in synclines. Plateau areas lacking pronounced structures are characterized by widespread but meager supplies of high-level water in relatively unfractured basalt. Although the largest supplies are found in synclines, there are several disadvantages of using water from these areas: (1) water of poor quality may rise along faults or tensional joints along synclinal axes; and (2) the synclinal aquifers tend to be isolated from one another and limited in natural recharge potentiality. Artificial recharge is a promising means of increasing natural supplies.

Temperature as a geohydrologic tool

R. W. Stallman (1-65), R. A. McCullough, and E. E. Parshall, using temperature profiles as a measure of ground-water velocity, have found that (1) infiltration rates as low as 0.05 foot of water per day can be measured by analysis of the diurnal temperature wave originating at the land surface; (2) downdip flows as small as a few tenths of a foot per day in artesian aquifers can be calculated from temperature profiles in wells if the dip of the strata is known; and (3) steady vertical flow in an aquifer, or vertical leakage from one aquifer to another, as small as 10^{-3} foot per day can be detected by analysis of temperature profiles. Present equipment measures temperature reliably to 0.005°C , and improvement to 0.001°C is in prospect.

J. D. Bredehoeft and I. S. Papadopoulos (1-65) have developed a technique of estimating from the earth's thermal profile, vertical ground-water flow through semiconfining beds.

F. C. Koopman and W. D. Purtyman, in studies related to disposal of heat from radioactive materials stored underground, discovered that holes ending above the water table tend to gain heat energy from the atmosphere, whereas holes extending into the zone of saturation tend to lose heat to the atmosphere.

J. M. Weigle made the interesting temperature-related observation that, in southern New Hampshire, winter thaws following summer and fall droughts resulted in ground-water recharge through dry and therefore porous and permeable soil. Brief thaws following periods of normal precipitation found the soil pores filled with frozen water and resulted in direct runoff rather than ground-water recharge.

Topographic and other criteria for predicting ground-water conditions

Topographic criteria recognized in previous years as valuable in predicting well yields in crystalline rocks of the southern Piedmont (in general, wells in valleys yield more than wells on hills) are being extended to other areas and types of rocks. Other criteria, such as fracture patterns visible at the land surface, are being studied also.

C. W. Poth found that wells in crystalline rocks in upland areas of Chester and Delaware Counties, in southeastern Pennsylvania, were drilled deeper and reached water at greater depth, penetrated a thinner mantle of weathered rock, and had lower yields than wells in draws; wells on slopes were intermediate. L. D. Carswell found similar conditions in Triassic rocks in southernmost New York and adjacent New Jersey and made the additional observation that headwater valleys of high stream gradient tended to have a thicker and more permeable zone of fresh-water circulation than valleys farther downstream. Harold Meisler and A. E. Becher found wells in carbonate rocks in the Lancaster quadrangle, in southeastern Pennsylvania, to be both shallower and more productive in valleys than on ridges.

F. W. Trainer and R. L. Ellison, the latter of the University of Virginia, found that fracture traces seen on aerial photographs are more abundant in carbonate rocks in the Shenandoah Valley in western Virginia than in shale; in both types of rock the dip of the strata is an important control on both orientation and abundance of fractures, and hence probably on the occurrence of ground water.

J. J. Musser, in the Roaring Creek district of east-central West Virginia, has correlated depths of overburden above coal mines with ground-water conditions. Thick overburden is characterized by little fracturing and perched ground water; thin overburden tends to be fractured and drained of ground water.

Interpretation of water-level fluctuations in wells

Interest in seismically induced fluctuations of water levels in wells, stimulated by the Alaska earthquake of 1964, continued during the year. R. C. Vorhis (2-65) devised a technique of estimating the magnitude of earthquakes from the fluctuation of the water level and the location of the well with respect to the epicenter of the quake. H. H. Cooper, Jr., J. D. Bredehoeft, I. S. Papadopoulos, and R. R. Bennett developed an equation relating seismic fluctuations to the type and frequency of the seismic wave, the geometry of the well (diameter of well

and depth of water), and hydraulic characteristics of the aquifer (transmissibility and storage coefficients). (See also the section "Alaska Earthquake, 1964".)

K. D. Wahl observed "reverse" water-level fluctuations in a well in Pickens County, Ala. The water level in the well, in a confined aquifer 235 feet deep, rises when nearby wells tapping an aquifer 113 feet deep are pumped. The cause is not certain but may be the unloading effect on the deeper aquifer of removal of water from the shallower one. Explanation of the phenomenon might contribute to understanding of the hydraulic characteristics of the aquifers.

Permeability and specific yield

Following up research by Benjamin Reyes and A. I. Johnson on relation of permeability to particle size, Johnson, R. C. Prill, and D. A. Morris evaluated methods of determining unsaturated permeability. As an outgrowth of this study, A. H. Ludwig is developing a centrifuge method of measuring unsaturated permeability, and Ludwig and R. P. Moston are evaluating gas-permeability methods.

A report by Prill, and Morris Johnson⁸⁹ shows that time of drainage is an important factor in determining the specific yield of materials packed in long vertical columns, even rather permeable materials that might be expected to drain quickly. Centrifuge techniques for evaluating the effect of variation in time of drainage have been developed by Prill, Johnson, and Ludwig.

The concepts of material transport in ground water systems were evaluated by H. E. Skibitzke (1-64), who pointed out that the tensor characteristic of the permeability coefficient and its correlation with the dispersion coefficient provide a possible means of defining the flow regime with a given aquifer. Because of the large-scale changes in magnitude and direction of the tensor component of permeability, the concept of a mean line of dispersion may produce a more significant statement on the characteristics of flow than does the present concept of hydraulic potential and the streamline.

Use of radioactive substances as tracers in geohydrology

J. M. Cahill tested radiophosphorus (P-32) and an organic dye in a glycerin base against a weak sodium chloride solution for effectiveness as tracers of water movement. The tests were made in columns of silica sand. The tests show both P-32 and the dye to be well suited for laboratory use in studies of mass transport of liquids through porous media.

⁸⁹ R. C. Prill, A. I. Johnson, and D. A. Morris, 1965, Specific yield—laboratory experiments showing the effect of time on column drainage: U.S. Geol. Survey Water-Supply Paper 1662-B, 55 p.

C. W. Carlston reviewed the usability of tritium in geohydrologic research. Its short half life (12.26 years) limits its usefulness as a tracer of ground-water movement to small and perhaps unrepresentative sections of aquifers. Its usefulness for dating recharge is similarly limited because of mixing of waters of different ages. The presence of tritium is meaningless as an indication of ground-water availability because in a humid area it may be present in a shallow aquifer of small yield. Carbon-14 is more useful because its time span of dating, a few tens of thousands of years, appears to include the periods during which most of our ground-water resources accumulated.

Carbonate-rock hydrology

W. J. Powell, and D. B. Knowles, and P. E. La-Moreaux of the Geological Survey of Alabama, found a close correlation among water chemistry, thickness of residuum, and productivity of underlying limestone aquifers in northern Alabama. Where the residuum is thick and gravelly the water in it and in the underlying limestone tends to be softer and the limestone to be more permeable and productive than where the residuum is thin.

From the results of 25 pumping tests of wells in limestone in Tennessee, G. K. Moore developed a type curve that is useful in predicting well yield from "step" drawdown tests of a new well (pumping the well at several successively greater rates and measuring the water-level drawdown). (See also the section, "Water Resources, Pacific Coast Area, Nevada".)

Use of electrical analog models in water-table studies

N. D. White and W. F. Hardt analyzed an electrical-analog model of the San Simon Basin, Ariz., and predicted water-level declines that will occur by 1980—120 feet near Bowie and 160 feet near San Simon. The analysis was based on hypothetical amounts and areal distribution of pumping in the basin which are believed reasonable on the basis of present trends in the use of ground water in the basin.

Geochemical data used in outlining geohydrology of Sparta Sand

J. N. Payne, as a phase of his study of the hydrologic system formed by the Sparta Sand in Louisiana and adjacent States, found geochemical data useful in outlining subsurface geology and water movement. For example, channel sands oriented parallel to the direction of regional flow contain fresher water than interchannel sediments or channel sands oriented across the flow.

Remote sensing of hydrologic phenomena

H. E. Skibitzke and associates, expanding work already begun on a small scale, studied the design, construction, and testing of infrared sensors and the use of conventional film photography (including that on infrared film) as a means of gathering information on water. Color aerial photographs of the Everglades, in southern Florida, and of adjacent coastal waters show paths of surface flow and points of underwater discharge of ground water more effectively than other techniques tried to date. Film photography and infrared imagery of parts of Puerto Rico and the Virgin Islands are being analyzed for signs of offshore discharge of ground water.

Recovery of fresh water in saline-water areas

G. G. Wyrick and H. B. Wilder developed an interesting and effective method of pumping from a thin fresh-water lens at a campground in the Cape Hatteras National Seashore Recreational Area, N. C. The lens is only 3 feet thick, and fresh water is skimmed from the upper part of it by means of an infiltration gallery. The gallery is pumped at a sufficiently low rate to prevent the rise of the underlying salt water to the bottom of the gallery bottom for at least 12 hours. Twice a day, when the tide goes out, the adjacent salt water retreats shoreward and the fresh water, following behind it, sweeps away the cone of salt water that has been rising beneath the gallery.

R. A. Long tested a "seavenger-well" system in Ascension Parish in southern Louisiana. The well is screened in a fresh-water aquifer above and a saline aquifer below. A packer separates the two parts of the well; through it passes a pump extending into the lower part. Pumping the two parts of the well at the same time lowers the head in the salt-water aquifer. This lowering prevents the upward movement of saline water, through the confining bed separating the aquifers, that normally occurs here when the fresh-water level is lowered by pumping. It was concluded, however, that separate wells would be more effective than the dual-purpose well.

Rejuvenation of waste-disposal well

A deep waste-disposal well at the National Reactor Testing Station, Idaho, was rehabilitated by selective perforation of the casing, guided by study of geophysical logs, by releasing and following a radioactive tracer in the well, and by injection tests. According to D. A. Morris, the capacity of the well to absorb wastes was increased from less than 200

to more than 1,000 gallons per minute, at modest cost. Similar techniques could be used to increase the yield of production wells where the proper conditions exist.

Geochemistry of glacial and fluvial sediments

Waters from four glacial and fluvial rock units on the Kitsap Peninsula and adjacent islands in the Puget Sound, Wash., studied by A. S. Van Denburgh (p. D219–D221), differ sufficiently in average content of potassium, orthophosphate, and dissolved solids, and in ratios of these constituents to each other, to enable identifying sources of ground water and to help in working out patterns of movement.

RELATIONS BETWEEN GROUND WATER AND SURFACE WATER

Because the principles governing the movement and occurrence of surface water and ground water differ in many respects, the methods by which they are investigated likewise differ. For this reason they are commonly studied separately. In many areas, however, water in streams and lakes and water underground are closely related and interdependent. Included in U.S. Geological Survey hydrologic investigations, therefore, are studies of (1) the magnitude and source of ground-water contributions to streamflow, (2) methods of forecasting the low flow of streams by use of ground-water levels and aquifer characteristics, (3) the effects of changes in the stage of streams and surface impoundments on ground-water levels, and (4) other subjects concerning the relations of water in the two environments.

Forecasts of ground-water storage and inflow to streams of Columbia River basin

A method has been developed by M. I. Rorabaugh for forecasting the ground-water inflow to the Spokane River between Post Falls, Idaho, and Spokane, Wash. The forecasts include: (1) ground-water outflow from Rathdrum Prairie, indexed by an observation well; (2) reduction in outflow as the water level declines seasonally; (3) bank storage caused by rises in stage of the Spokane River; and (4) return flow associated with lowering stages of the river. Verification of monthly forecasts for the period August through March for 33 years shows that observed flows exceed the forecast 95 percent of the time. The portion of the outflow associated with the water level in the Prairie has had a range of 300 cubic feet per second during the period 1929–64.

A study of forecasts of assured flow for the October–March period in the Bitterroot Basin, Mont., by W. D. Simons, indicates that the record minimum runoff during the power season from October 1936 through March 1937, did not stem from a deficiency of water in the basin, but probably from unusually large ice storage. Analysis of data on the temperatures in western Montana indicates that in January 1937 both the minimum monthly temperature and the duration of days below freezing reached record or near-record values.

Separation of streamflow into components

Using a method devised by G. R. Kunkle⁹⁰, L. E. Bidwell and R. D. Cotter have separated the flow of the Pomme de Terre River in Minnesota during a 6-year period into base flow, basin-storage discharge, and overland runoff. A plot of these components along with total flow provides a meaningful summary of the flow characteristics.

L. G. Toler (p. C206–C208) has used the specific conductance of water to separate the base flow of Econfinia Creek, Fla., into components from the Floridan aquifer (cavernous limestone) and from the overlying sand aquifer. The components from the Floridan aquifer was correlated with the stages of lakes that recharge the aquifer.

G. R. Kunkle (p. D207–D210) used measurements of specific conductance and of discharge in a study of two streams in east-central Iowa. He separated the hydrograph of one stream into surface-runoff and ground-water components, and he computed the ground-water discharge from an aquifer to a particular reach of the other stream.

Ground-water inflow, bank and channel storage, and streamflow pickup

W. E. Clark (p. D211–D213) studied streamflow in an 8-mile reach of the Santa Fe River in north-central Florida. The pickup in flow in this reach is provided, entirely or nearly so, by ground-water runoff. After floods, part of the ground-water runoff from within the 8-mile reach is provided by water which drains back into the stream from bank storage.

Increase in base flow due to removal of vegetation

H. C. Riggs (p. C196–C198) has related the base flows of small streams in the Hazel River basin, Virginia, to the percentage of the stream valley that is cleared of trees and brush. His analysis, based on 8 sets of measurements on 9 streams, indicates that

⁹⁰ G. R. Kunkle, 1962, The baseflow-duration curve, a technique for the study of ground-water discharge from a drainage basin: *Jour. Geophys. Research*, v. 67, no. 4, p. 1543–1554.

clearing increases the yield considerably during droughts but only slightly when the base flow is high.

Runoff altered by coal mining

Coal mining in Randolph County, W.Va., has caused a part of the flow of Roaring Creek to be diverted into Grassy Run. According to J. J. Musser, some of the water from the Roaring Creek basin moves down the dip of a synclinal limb through abandoned mines into streams of the Grassy Run basin. The augmented base flow of Grassy Run, per square mile, is 20 to 30 times that of Roaring Creek.

New tunnel in Hawaii will decrease yield of Waihee tunnel

G. T. Hirashima reports that the Waihee tunnel reservoir in the Waihee Valley, Oahu, Hawaii, receives as much as half a million gallons of water a day from ground-water discharge whenever the stage in the tunnel is lowered sufficiently. A proposed new tunnel will capture this discharge of ground water, thereby decreasing the yield of the Waihee tunnel.

Estimating aquifer characteristics from changes in ground-water levels near clogged streambeds

A pumping test from a stream in the Little Plover River basin, in Portage County, in central Wisconsin, indicated that the stream was not freely connected to the contiguous aquifer. Treating the perimeter of the stream and the aquifer as a two-permeability system, A. G. Law has developed an analytical procedure to estimate the hydraulic characteristics of the aquifer and the effect of the clogged streambed on the response of ground-water levels to changes in stream stage. The procedure employs a type-curve technique.

Streamflow computed from ground-water recharge data

According to E. P. Weeks, D. W. Ericson, and C. L. R. Holt, Jr., the monthly mean streamflow of the Little Plover River in Portage County, Wis., can be computed from recharge data. The basin was divided into rectangular areas based on the drainage pattern, the location of the ground-water divides, and the geology of the basin. Average values for the coefficients of transmissibility and storage were then determined for each of the rectangular areas, and these values were used to compute the rates at which given volumes of recharge were discharged to the stream. The computed mean monthly streamflow for a 20-month period agreed with the measured streamflow within a few percent in all but 2 months, in which recharge and runoff were abnormally high.

SOIL MOISTURE AND EVAPOTRANSPIRATION

Permeability of soil to water determined from movement of water vapor

In a study near Cuba, in north-central New Mexico, R. W. Stallman, G. E. Ghering, and R. A. McCullough determined the vertical permeability of soils to water by measuring gas pressures at different depths below the surface in the unsaturated zone. The conductivity of the soil to gas is related to its conductivity to water, and moisture flow was estimated from the results of the measurements.

Storage of soil moisture in flood-plain deposits

A. O. Waananen measured an increase in soil-moisture content in Humboldt River sediments near Winnemucca, in north-central Nevada, after periods of high flow. This moisture was retained in storage during 2 years of near-normal rainfall and runoff but surely will become depleted during a very dry spell.

Magnitude of rise of soil moisture computed

R. W. Stallman and J. E. Reed described steady capillary conduction of moisture from the water table in a homogeneous medium, by equations which can be solved by different methods that use digital computers, analog computers, or graphical techniques. Applying the graphical methods for computing the steady flow of liquid between water table and land surface they found that the magnitude of rise through silts can attain as much as 100 feet, at a rate exceeding the demand on moisture made by the vegetation. Thus the assumption that a deep water table is isolated from evaporation may be subject to doubt.

Formula for soil-moisture change with time

M. R. Collings and R. M. Myrick, studying soil moisture changes in the Cibique Ridge basin in east-central Arizona, constructed an empirical equation describing the decrease in soil moisture with time, with which they can predict soil-moisture content at future dates with a satisfactory degree of confidence, for use in precipitation-runoff studies.

Infiltration in fine-textured soils favorable for forage grass

R. F. Miller, F. A. Branson, J. S. McQueen, and William Buller report that soil-moisture conditions favorable for the growth of good forage grasses on fine-textured soils on basin spreaders can be attained, provided that ponding is prevented. Coarse-textured soils have rates of infiltration and percolation too high for water to remain available for edible plants,

although at some places sediment deposits may, after a time, lower the infiltration and percolation rates.

Projected effect of dams on evapotranspiration rates

J. E. Reed, M. S. Bedinger, and H. H. Tanaka have found that construction of navigation dams along the Arkansas River will result in a change in evapotranspiration rates, with the effect that the water levels in the alluvium along the river will not rise as much as the stream stage. They used analog models to arrive at their conclusion.

Use of soil moisture by riparian vegetation

It is often assumed that destroying the riparian vegetation will result in an increase in ground-water supply because of the diminished use of ground water resulting from the destruction. R. C. Zimmerman reports that along the tributaries of the San Pedro River, in south-eastern Arizona, vegetation commonly depends not on ground water but on water retained in the alluvium along the stream following floods. Destruction of such vegetation would result in reduced river-channel stability and would deprive livestock of emergency browse.

J. S. McQueen and R. F. Miller, studying vegetation along the Gila River in Arizona, found indications that saltcedar may not use soil moisture at all but may live on ground water exclusively. However, cottonwood, mesquite, and willow seem to draw upon both sources.

Effect of Great Swamp, N. J., on streamflow

During periods of high evapotranspiration, streamflow from Great Swamp, N.J., is considerably less than inflow to the swamp. E. G. Miller (p. B177-B179) believes that evapotranspiration causes a significant reduction in outflow during the period from June to September.

Evapotranspiration studies

T. E. A. van Hylckama found, near Buckeye in southwestern Arizona, that saltcedar grown in tanks (evapotranspirometers) containing water of low salinity used more water than saltcedar grown in those with high-salinity water. This observation is surprising in view of the well-known tolerance of these plants for salt. The rate of water loss from tanks during hot days can be assigned to transpiration and evaporation factors by the use of harmonic analysis. Water-level records on bare and vegetated tanks both show daily fluctuations which are suggestive of transpiration curves but which apparently are due to daily barometric-pressure changes. Such curves cannot be used to estimate evapotranspiration.

T. W. Robinson and D. E. Donaldson, working with evapotranspirometers at Winnemucca, Nev., showed that a fluorescent dye, Pontacyl Brilliant Pink B, can be used to trace the path of water from the ground water to the leaves. Dye was detected (1) in the soil moisture, (2) in the leaves, and (3) in transpired water collected from the leaves in plastic bags.

T. W. Robinson also reports from Winnemucca that a clear-cut effect of air temperature and length of frost-free season was found on the use of water by four species of woody plants. Water use by greasewood, rabbitbrush, willows, and wildrose was 25 to 40 percent less in 1964 than in 1963. The growing season was both shorter and cooler in 1964 than in 1963: there were 102 frost-free days in 1964, as compared with 135 in 1963, and the average temperature for the period April 1 to November 1 was 1.7°F lower in 1964 than in 1963.

Computation of evapotranspiration

J. S. Meyers and A. M. Sturrock, Jr., compared use of the energy-budget and mass-transfer methods of computing evapotranspiration on two large reservoirs: Garrison Reservoir in North Dakota and Pomme de Terre Reservoir in Missouri. Using the energy-budget techniques developed for Lakes Hefner and Mead,⁹¹ they found good agreement for an entire season but not for short periods. The relationship of energy budget versus mass transfer, so well defined for Lakes Hefner and Mead, did not agree for the two reservoirs.

W. S. Eisenlohr, Jr., in a study of the hydrology of prairie potholes, found that the mass-transfer coefficient N , described by Harbeck,⁹² changes with change in vegetation. This effect provides a means of separating the water losses due to evaporation from those due to transpiration. The mass-transfer coefficient for evaporation is a constant; the one for transpiration has a curvilinear relation to the emerged height of the vegetation.

O. E. Leppanen reports that analysis of data describing the thermal-energy exchange over 12-foot-tall saltcedar thickets indicates that the energy-budget approach to evaluate evapotranspiration is workable. Computation of energy available for evapotranspiration at Buckeye, in southwestern Arizona, shows that evapotranspiration can be

⁹¹ U.S. Geological Survey, 1954, Water-loss investigations—Lake Hefner studies: U.S. Geol. Survey Prof. Paper 269, 158 p.

G. E. Harbeck, M. A. Kohler, G. E. Koberg, and others, 1958, Water-loss investigation—Lake Mead studies: U.S. Geol. Survey Prof. Paper 298, 100 p.

⁹² G. E. Harbeck, Jr., 1962, A practical field technique for measuring reservoir evaporation utilizing mass-transfer theory: U.S. Geol. Survey Prof. Paper 272-E, p. 101-105.

estimated within 20 percent of that measured by evapotranspirometers. Further refinement appears to be possible.

SEDIMENTATION

The scope of investigation termed "sedimentation" includes the sequence of events which begins with the separation of particles from parent rock and concludes with their consolidation into another rock. Sedimentation, therefore, involves consideration of sediment sources; of the weathering, erosion, transportation and deposition of sediments; and of environments of deposition and sedimentary deposits.

EROSION

Erosion near mouth of Rio Puerco

The general balance between erosional and depositional processes in a channel may be altered by the formation of a head cut and its movement upstream. Observations by C. F. Nordin, Jr. (4-64), indicated that aggradation and armoring of the bed of the Rio Puerco, in New Mexico, followed aggradation in the main stream, the Rio Grande. Local scour in the Rio Grande at the mouth of the Rio Puerco formed a head cut and a plunge pool, by erosion of the armored layer and an underlying layer of sand. In 1 year the head cut moved upstream in the Rio Puerco a distance of 1 mile. The depth of the plunge pool reached a maximum of 8 feet during this period.

Scour and fill in sand-bed streams

B. R. Colby (3-64) concluded from a review of general principals of scour and fill and a study of the behavior of a few types of natural streams that:

1. At any instant, the quantity of moving sand, even in deep and swift flows, is usually equivalent to an average depth of deposit of only a few hundredths of a foot on the streambed.
2. In a straight sand-bed channel that has uniform cross section, slope, roughness, and bed material, the streambed neither scours nor fills appreciably during the passage of a flood.
3. In a straight channel having uniform cross section but varying slope or roughness along the channel, fill during a rising stage is usually followed by fill on the falling stage, and scour during a rising stage is usually followed by scour on the falling stage.
4. If a change in roughness or in streamflow requires a change in the depth of flow, the depth generally adjusts through a change in elevation of the

water surface and not through a change in elevation of the streambed.

Rate of slope denudation in San Gabriel Mountains

V. C. LaMarche, Jr. (1-64) found, in the San Gabriel Mountains in southern California, that the exposed root system of a 1,400-year-old pine indicates a slope-denudation rate of about 4 acre-feet of sediment per square mile per year during the lifetime of the tree. Modern sediment yields computed from 40 years of reservoir-sedimentation records approach this rate.

Piping and collapse structures in near-surface deposits

In studies of basic processes of erosion, G. G. Parker⁹³ has shown that piping and collapse structures occur commonly in semiarid parts of the United States and under certain conditions in humid climates. Piping is the natural development of subsurface drainage tubes in relatively insoluble clastic materials. Collapse structures attack several kinds of low-density detrital materials, chiefly deposits of silt, loess, and volcanic ash having a very loose initial packing. Acting alone or together, piping and collapse structures often produce pseudokarst features.

TRANSPORTATION

Predicting sediment loads of rivers

In streams such as the Rio Puerco, in New Mexico, C. F. Nordin, Jr., (4-64) found that the concentrations of suspended sediment are so great that the properties of the fluid, per se, cannot be neglected and that both the flow and concentration change so rapidly with time that the usual assumptions of uniform flow and equilibrium transport are not applicable.

D. W. Hubbell and W. W. Sayre (2-64) concluded from a study of the transport and longitudinal dispersion of bed-material particles in the North Loup River in Nebraska that the discharge of bed-material particles can be computed by using data collected by the radioactive-tracer technique in conjunction with a continuity equation.

In a study of the fluvial characteristics of the Middle Rio Grande, in New Mexico, J. K. Culbertson and D. R. Dawdy (1-64) computed total sand transport using both the modified Einstein and the

⁹³ G. G. Parker, 1963, Piping, a geomorphic agent in landform development of the drylands: Internat. Assoc. Sci. Hydrology Pub. 65, Berkeley, Calif., p. 103-113, 3 figs.

G. G. Parker, L. M. Shown, and K. W. Ratzlaff, 1964, Officer's Cave, a pseudokarst feature in altered tuff and volcanic ash of the John Day Formation in eastern Oregon: Geol. Soc. America Bull., v. 75, no. 5, p. 393-402, 2 pls., 3 figs.

Bagnold methods. For the data available, the modified Einstein method gave consistent results, whereas the Bagnold method did not appear to be suitable for the field conditions met in the study.

A means of determining the bed-material discharge per foot of stream was presented by B. R. Colby (4-64) in terms of four dominant measures: total shear, shear velocity, mean velocity, and stream power. The graphical average relationship of bed-material discharge to mean velocity is the most convenient to apply. The effects of water temperature and of high concentration of fine sediment on the relationship were determined approximately.

VARIABILITY OF SEDIMENT LOADS IN RIVERS

Very high concentrations of suspended sediment

The highest observed suspended-sediment concentrations occur in streams that drain semiarid regions. J. P. Beverage and J. K. Culbertson (1-64) present data from the basins of the Rio Grande and the Colorado River for samples in which streamflow was more than 40 percent sediment by weight. They found no relation between stream discharge and the occurrence of these very high concentrations of sediment. The proportion of sand increases with increasing concentrations of suspended sediment, and the minimum concentration of the clay fraction, when there is more than 5 percent sand, appears to be about 15 percent. Increase in clay content decreases the fall velocity of the finer sand fractions⁹⁴ and thus facilitates the transport of sand.

Snowmelt and sediment loads

Studies by B. L. Jones of sediment loads in streams in small drainage basins in central and northern Pennsylvania indicate that a third to a half of the suspended-sediment discharge for the year occurs during periods of snowmelt runoff.

Flood runoff and sediment load

S. E. Rantz and A. M. Moore (2-65) report that floods in the far western States in December 1964 were outstanding for record-breaking peak discharges and accompanying maximum suspended-sediment concentrations. In tributary streams of the lower Snake and Columbia Rivers the maximum observed sediment concentrations ranged from 188,000 to 309,000 parts per million.

⁹⁴ C. F. Nordin, Jr., 1963, Sediment transport in alluvial channels, a preliminary study of sediment transport parameters, Rio Puerco near Bernardo, New Mexico: U.S. Geol. Survey Prof. Paper 462-C.

D. B. Simons, E. V. Richardson, and W. L. Haushild, 1963, Studies of flow in alluvial channels, some effects of fine sediment on flow phenomena: U.S. Geol. Survey Water-Supply Paper 1498-G.

Urbanization and sediment load

D. H. Carpenter reports that accelerating urban development has increased the sediment load in the Northwest Branch of the Anacostia River, Md., near Washington, D.C. The average annual sediment yield has increased from 470 tons per square mile in 1962 to 1,800 tons per square mile in 1964.

DEPOSITION

Effect of confining banks and channel encroachment by saltcedar

R. C. Culler and R. M. Myrick found spectacular sediment deposition and channel plugging in the Gila River in the upper part of the San Carlos Reservoir area, in Arizona. A dense cover of saltcedar on the riverbanks and flood plain had held the channel to the same location for a period of 20 years. Deposition on natural levees and in the channel caused the river bed to aggrade to a level above the adjacent flood plain. Beginning in 1962 the debris carried by summer floods jammed and became buried in sediment in the channel, and water flowed in rivulets over the natural levees. The channel plugging progressed upstream half a mile in 1963 and 2 miles in 1964.

Storage in reservoirs

Trap-efficiency investigations at K-79 Reservoir, Kiowa Creek basin, Colorado, indicate that the sediment discharge from the reservoir is limited to short periods of time. Robert Brennan reports that during the 8-year period, 1956-63, about 85 percent of the sediment discharge from the reservoir occurred during about 0.1 percent of the time. Sediment deposition in the reservoir has been reduced by deposition above upstream structures which were built in 1960.

George Porterfield and C. A. Dunnam (1-64) reported that the loss of original storage capacity in Lake Pillsbury, Calif., during the period 1921-59, due to accumulation of sediment, was 8.1 percent of the original total storage capacity. The elevation of the outlet sill of the dam is 1,740.7 feet above mean sea level. Fifty percent of the sediment, 3,810 acre feet, was deposited above the 1,794-foot contour.

Sedimentary structures formed by flow in stream channels

Results from laboratory and field studies of the mechanics of flow and sediment transport in alluvial channels have been used by D. B. Simons, E. V. Richardson, and C. F. Nordin, Jr., to establish flow regimes and channel-bed configurations; sizes and

shapes of ripples, dunes and antidunes; and sediment-transport rates associated with the various bed forms. Because sedimentary deposits contain ripple marks and crossbedding from the initiation and migration of configurations generated at a fluid-sediment interface by a current, the relations developed from flume and field studies have applications in interpreting the hydraulic environment of such deposits. Bed configuration is related to stream power and particle size, and the ratio of wavelength to amplitude of dunes is related to shear stress and particle size. If ripple marks or cross stratification in a sedimentary deposit permit identification of bed configuration, the range in velocity, depth, and unit water discharge associated with a particular sedimentary unit can be determined.

Sedimentary structures in streambed deposits

E. D. McKee determined through experiments on ripple lamination that such lamination is typically formed and preserved in river flood-plain deposits but is not preserved in tidal-flat deposits. Thus the abundance of ripple lamination constitutes a useful criterion for distinguishing between sediments of two environments that in general are similar.

Orientation of axes in sediments

A. E. Scheidegger (p. C164–C167) shows that the mean of a series of axes in space (normals to bedding planes, long axes of sand grains, and so forth) is not given by the vector mean of the individual axes, but should be calculated as the eigenvector of a certain symmetric matrix.

LIMNOLOGY

Studies of Pretty Lake, Ind.

J. F. Ficke and R. G. Lipscomb have studied stratification of water in Pretty Lake, LaGrange County, in northeastern Indiana. Chemical stratification of the water in winter, when the water density is near its maximum throughout the lake, is attributed to the action of algae. The algae remove nutrients from the upper part of the water; after death the algae settle to the bottom and decompose, enriching the bottom water.

During summer, Ficke and Lipscomb found diurnal changes in pH and dissolved-oxygen content. The greatest changes were in a shallow embayment where little mixing with water from the rest of the lake occurred. The changes are due to photosynthetic release of oxygen by aquatic plants during the day and to their consumption of oxygen at night.

Ficke described (p. C199–C202) the course of seasonal erasure of thermal stratification on the basis of changes in the temperature and dissolved-oxygen content of the water during late summer and autumn.

Hydrology of Great Salt Lake

In a summary of the physical and chemical hydrology of Great Salt Lake, Utah, D. C. Hahl, M. T. Wilson, and R. H. Langford (p. C181–C186) describe two recent significant discoveries: (1) the two-directional flow of brine through a causeway which separates the lake into two arms, and (2) the formation of a temporary fresh-water layer at the surface of part of the lake as a result of inflow from streams and of precipitation on the lake.

Effect of emergent vegetation on stratification of water in a prairie pothole

Study of a shallow 15-acre pothole in Dickey County, in southeastern North Dakota, showed the influence of emergent vegetation on water properties. During a windy day, H. T. Mitten found in areas of emergent plants a stratification of specific conductance, temperature, and dissolved oxygen between the surface and the bottom in water about 2 feet deep. Where the pothole was without emergent vegetation the water was mixed by wind action, and in this open area there was no significant difference in the properties of water between surface and bottom.

GEOMORPHOLOGY

Erosion rates and processes

Six years of observations of erosion pins, painted rocks, and erosion plots by L. B. Leopold, W. W. Emmett, and R. M. Myrick near Santa Fe, N. Mex., show that about 95 percent of the sediment produced from woodland hillslopes of low relief results from unconcentrated sheet erosion. Processes of mass movement and gully extension contribute a small percentage of the total sediment load under present conditions. Of the total sediment eroded, only about a third is caught in small reservoirs or contributed to aggrading channels.

In his investigations of the effect of exposure on slope erosion in three areas in Colorado and New Mexico, R. F. Hadley has found that the rate of removal of surficial material by sheet erosion is 50 to 75 percent greater on slopes that have southerly and southeasterly exposures than on slopes that have other exposures.

S. A. Schumm and R. J. Chorley (2-64), in a study of cliff recession and rock weathering, reported some interesting observations at Pueblo Bonito, Chaco Canyon, N. Mex. In 1941 a huge monolith of Cliff House Sandstone toppled and damaged a portion of the ruins. Measurements of the movement of the rock prior to its fall demonstrate that the movement of the sandstone block over the underlying shale is an exponential function of total precipitation and that the rate of movement was seasonal, being relatively faster during the winter. It is estimated that the rock began moving slowly away from the cliff about 550 B.C.

Circular patterns caused by exfoliation

In a study of aerial photographs of the Grandfather Mountain area on the Blue Ridge escarpment, in western North Carolina, J. T. Hack found numerous circular patterns from a quarter of a mile to more than a mile in diameter. They are marked by curving ravines, ridges, and streams. Generally circular basins and domical hills of similar diameter are also common. It is believed that the circular features are formed as a result of exfoliation. The tectonic joints in the area open when the stress on the rock, due to deep burial, is relieved by dilation and spalling along large and deep exfoliation fractures or sheets. The fracturing produces roughly circular sheets of rock of varying permeability. This process may help explain the irregular distribution of saprolite and the occurrence of shallow groundwater supplies in massive crystalline rocks.

Geometry of free meanders

C. W. Carlston has completed a study of the geometry of free stream meanders. The relation of free-meander wavelength to stream discharge is most closely approximated (S. E. = 11.8 percent) by the equation $L_m = 106.1\bar{Q}^{0.46}$, where L_m is meander wavelength, $\bar{Q}$ is mean annual discharge, and S.E. is the standard error. For bankfull discharge (Q_b), $L_m = 8.2Q_b^{0.62}$ and S. E. = 25 percent; for mean discharge of the month of maximum discharge ($\bar{Q}_{mm}$), $L_m = 80.0\bar{Q}_{mm}^{0.46}$ and S. E. = 15 percent. For meander width (W_m), $W_m = 65.8\bar{Q}^{0.47}$ and S. E. = 23 percent; for stream width (W) $W = 7\bar{Q}^{0.46}$ (approximately). Carlston concludes that the dominant discharge which controls meander wavelength is in a range of flows, possibly on falling stages, between $\bar{Q}_{mm}$ and $\bar{Q}$; and that meander migration takes place during these stages. Some field evidence indicates that slope also influences meander wavelength. Whatever the hydrodynamic cause of the wave form of meanders, the effect of

the tortuosity of meandering streams is to produce a low-gradient channel for average flows, while the higher-gradient flood plain is an efficient transit channel for large-scale floods.

Paleohydrologic studies along upper San Joaquin River, Calif.

Studies by R. J. Janda indicate that during periods of aggradation, in Pleistocene time, the San Joaquin River near Friant, Calif., carried approximately $2\frac{1}{2}$ times more sediment than the modern river. Channel alluvium deposited during periods of aggradation contains a greater percentage of coarse sand than the modern alluvium.

Paleohydrologic studies at Thorne Cave, Utah

The relationship of alluvial deposits to artifacts of early man has been studied by H. E. Malde at Thorne Cave, near Jensen, Utah. Valley alluvium which accumulated to a depth of 48 feet along Cliff Creek continued to build up another 13 feet while man occupied the cave, and then reached 30 feet higher, sealing in the signs of man. The cave was subsequently exposed by erosion that carved a deep arroyo in the valley. Radiocarbon dates indicate that the cave debris is about 4,000 years old, and they suggest that the valley alluvium was deposited during the period from 5,000 to 3,000 years B.P. The link of the alluvium with early man is archeologically important because the deposit implies a relatively wet environment.

PLANT ECOLOGY

Plants and their hydrologic and physical environment are intimately interrelated. The kinds of plants, their establishment and survival, and their growth rates are strongly affected by, and in turn affect, surficial deposits, geomorphic processes, and land use. Study of these complex interrelations is necessary to provide a better understanding of man's environment.

Historic changes in vegetation in southwestern States

The saguaro (*Carnegiea gigantea*) has failed to reproduce in some habitats where it formerly maintained dense stands. Previous studies have shown that most saguaros in the Saguaro National Monument, in southeastern Arizona, may be gone by the year 2000 because of the sparseness of seedlings in the population and the inevitable death of each remaining old cactus. According to recent studies by R. M. Turner, S. M. Alcorn, George Olin, and J. A. Booth, the pattern of mortality among the seedlings

indicates that most deaths may be attributed to (1) lack of shade, (2) small herbivores, and (3) insects. Lack of shade appears to be the most important of these three factors. Presumably, the amount of shade is affected by the volume of associated plants and may be altered by fluctuations in climate or by changes in man's use of the land.

Growth of trees related to local climatic variables

Growth studies of eastern deciduous trees by R. L. Phipps have shown that annual ring growth probably occurs first in the top of a tree and then, at an increasingly later time, down the trunk. Growth at any level is consequently a function of previous growth at higher levels and of environmental conditions at the time growth occurs at that level. Thus, an understanding of annual ring growth of trees in relation to environment necessitates growth determinations from several heights in the trunk. Better correlations with environmental variables have been obtained using ring data from trees that grew under relatively optimum conditions than from suppressed trees containing discontinuous or missing rings. A high degree of correlation has been obtained between ring-area increment and evapotranspiration deficit, a single parameter of environmental water and energy. Growth of an individual tree was found to reflect strongly the particular microclimate in which it grows.

Tree growth and precipitation

In studies of tree rings from Arizona, New Mexico, and Colorado, Lee Horner, Deric O'Bryan, and N. C. Matalas have found, in correlation of precipitation and ring width, that a match to within 60 percent of perfect correlation can be attained. The best results were obtained by use of three measures: (1) precipitation for the 12-month period November through October, (2) total annual precipitation, and (3) winter (November-April) precipitation. The use of ring area (especially where based on 4 radii instead of 1 radius) improved correlation of tree growth with November-October annual precipitation by an additional 10 percent over use of ring width. The yield of corn grown by Indians in nonirrigated fields also showed high correlation with the November-October annual precipitation.

Effects of coal strip mining on upland forests

Acid water from coal strip mines in Kentucky seems to cause trees irrigated by it to grow rapidly in radius. Studies by R. S. Sigafos (*in* C. R. Collier and others, 2-64, p. B73-B74) show that irrigated

trees grew more rapidly during the 5 years following cessation of mining than they had done previously. Preliminary analysis of data collected in 1964 from the same and additional trees after a second 5-year period shows continued rapid growth. Naturally seeded shortleaf, pitch, and Virginia pine saplings on mine tailings have grown rapidly, and abundant fruiting bodies of fungi among the pine trees suggest that mycorrhizae have developed in the tree roots. Eight-year-old pine saplings are producing viable seed, and second-generation seedlings are becoming established on the tailings.

Relation of flood-plain forests to land use and flooding

Past use of flood-plain land by man along the Potomac River in Maryland, near Washington, D.C., was found by R. S. Sigafos to have had little effect upon vegetation that has grown since abandonment of the activity. The present forest, at a site where a mill was operated until about 1900 and where the March 1936 flood later destroyed all summer cottages, is similar in species content and in form and appearance of the trees to forests on sites that were not used by man but are otherwise similar. Forests on sites of Chesapeake and Ohio Canal lock-keepers' gardens are like those on relatively undisturbed sites, even though most gardens were not abandoned until the 1920's. Floods that occur about once every 2 years or at greater intervals disturb the surface and tend to maintain an unstable environment in which erosion, deposition, and the development of vegetation are in dynamic equilibrium (R. S. Sigafos, 1-64, p. A31-A32). The most recent events—floods—seem to be the most effective agent in the life and death of the flood-plain forests.

Effects of soil moisture and land treatment upon establishment of grasslands

L. M. Shown, R. S. Aro, R. F. Miller, and F. A. Branson have studied the conversion of sagebrush land to grassland, a change which results in a more beneficial use of the water resource of some of the arid and semiarid regions of the intermountain west. They found that the establishment of grass is mostly a function of precipitation-soil relationships. The most productive seedlings, which signify efficient use of soil moisture, are found on areas receiving 10 to 15 inches of average annual precipitation, and on soils which have medium moisture-holding capacities per soil unit. The soil texture is usually sandy loam, silt loam, or loam. Seedlings either do not become established or soon die on coarser or finer soils, on thin rock, or in rocky soils, all of which do not hold enough water or are poorly

permeable. Seedling establishment is poor where growth of big sagebrush (*Artemisia tridentata*) is poor and where greasewood (*Sarcobatus vermiculatus*) and shadscale (*Atriplex confertifolia*) are predominant.

In a study of conversion of pinyon-juniper woodlands to grassland in semiarid areas of the West, R. S. Aro has found that the percentage of trees killed is directly proportional to the intensity of woodland treatment. Uprooting trees by chain and tractor was least effective, whereas chaining and bulldozing trees into windrows and burning the woodlands was the most effective treatment. Preliminary analyses suggest that benefits derived from increased grass in the more intensively treated areas exceed the higher cost.

Relationship between chemistry of bedrock and chemistry of plants

Vegetation and soil analyses of 17 sedimentary formations in Wyoming and Montana by F. A. Branson and R. F. Miller show that variation in chemical characteristics between formations is smaller than between different plant species. Potassium percentages in formations including the Eagle Sandstone, Wasatch Formation, Niobrara Formation, and Bearpaw Shale varied about 1 percent, but shrubs growing on these formations contained an average of more than 6 percent more potassium than did grasses. Causative factors, more important than chemical characteristics of geologic materials in determining kinds of plants present, are (1) soil textures, which affect infiltration and runoff rates; and (2) the quantities of water available for plant use. The great differences in uptake of certain elements by different species growing on the same soil indicates that upland vegetation may have significant effects on quality of surface water.

Effects of geomorphic process on desert vegetation

Investigation of winterfat (*Eurotia lanata*) and big sagebrush (*Artemisia tridentata*) along their contact zone in eastern Nevada by R. S. Aro has shown that soil, water, and forage are being lost by death of winterfat, which is valuable for livestock grazing. Erosion of the surface at the downslope boundary of winterfat is changing the slope from a concave to a convex one and resulting in the replacement of winterfat by sagebrush. Understanding of the nature and rate of this process should suggest appropriate corrective action to provide optimum utilization of the scarce water resource in this area and to help stabilize the soil.

GLACIOLOGY

Glaciology is an interdisciplinary field bearing on several aspects of earth science. Glaciers are important as winter resources in some Western States, such as Washington, where about 40 million acre-feet of water is currently stored in the form of glacier ice. The marked growth and decline of glaciers in response to slight changes in climate is, therefore, an important factor in water-resources planning. Conversely, the study of changes in the glaciers produces insight into the nature of climatic changes; traditionally this has been one of the most important reasons for the study of glaciers. The net mass budget, the crucial link between glaciers and climate, has received much recent scientific attention.

Net mass budget of South Cascade Glacier, Wash.

Ice-storage changes for South Cascade Glacier were determined by W. V. Tangborn for the 1957–64 period using basin runoff and precipitation measurements. Evaporation and condensation are shown to have negligible effect. Runoff, measured by a stream discharge station, averaged 4.05 meters per year; precipitation, determined by snow-accumulation measurements on the glacier and in storage gages, averaged 3.82 m/yr, resulting in a basin net loss of about 0.23 m/yr. During the same period, South Cascade Glacier net mass budgets were determined each year by conventional glaciological studies. The glaciological and hydrometeorological results are consistent both as to long-term averages and as continuous functions of time.

Data on relations between ice velocity, net mass budget, and changes in surface elevations, collected from 1957 to 1964, have been analyzed by M. F. Meier and W. V. Tangborn. Ice velocities range up to about 20 m/yr, net mass budgets averaged over the glacier surface range between -3.3 m of water equivalent (1957–58) and $+1.2$ m (1963–64), and the glacier became thinner (during the period 1958–61) at a rate averaging 0.93 m/yr. The algebraic sum of emergence velocity, net budget, and change of surface elevation was examined at four points on the glacier surface and as functions of distance along the length of the glacier. Ice discharge and thickness were calculated from these data. A steady-state distribution of net budget was used to calculate a steady-state discharge, which was found to be 2.2 times larger than the present discharge.

Studies by M. F. Meier indicate that an advance of South Cascade Glacier 4,700 years ago produced a massive moraine 880 m beyond the 1963 location of

the terminus. One subsequent advance in the 16th century and two in the 19th century produced small moraines 1,360, 1,250, and 1,130 m beyond the 1963 terminus. Since 1900 the glacier has receded. Approximate longitudinal profiles corresponding to these moraines were constructed and used with the present-day vertical variation of net mass budget to estimate steady-state net budgets at the times of moraine construction. A long-continued net budget only 90 centimeters more positive than the 1957–64 average would be sufficient to reproduce the greatest advance of this glacier since the Pleistocene.

Influence of 1964 Alaska earthquake on glaciers in State

The severe 1964 Alaska earthquake provided an ideal opportunity to test the theory that earthquake-induced avalanching of snow and ice can cause catastrophic glacier advances. An extensive aerial photographic study was made by A. S. Post in August 1964, when snow cover was at a minimum, and the results were compared with similar studies made in earlier years. Observed changes in the several thousand glaciers examined were largely those expected in a normal year. The only exceptions were 6 widely scattered glaciers that appeared to have begun very rapid movements; however, 4 other glaciers showed similar characteristics in 1963. One tributary of the Martin River Glacier may be responding to earthquake-induced avalanching, but the evidence is inconclusive. Thus there appears to be very little evidence of dynamic glacier response 5 months after an earthquake of unusual severity.

Movement and thickness of Teton Glacier, Wyo.

Resurvey of markers near the snowline on Teton Glacier, Grand Teton National Park, Wyo., by J. C. Reed, Jr. (p. B137–B141), showed that there the ice advanced as much as 37 feet between August 1963 and September 1964 while the terminus of the glacier remained in the same position. The ice is thought to have moved mostly as a rigid mass. It may be as thin as 50 feet near the snowline.

PERMAFROST

Permafrost map of Alaska

O. J. Ferrians, Jr.,⁹⁵ has prepared a map, at a scale of 1:2,500,000, showing the distribution and character of permafrost in Alaska. Because 85 percent of Alaska is within the permafrost region, the potential deleterious effects of building on perma-

frost pose serious engineering problems. This is the first such Statewide map produced in sufficient detail to be useful in the selection of sites, the determination of road alignments, and the design of engineering structures.

Inventory of pingos in part of central Alaska

Pingos, mounds or hills formed by updoming by hydrostatic pressure in areas underlain by perennially frozen ground, have received considerable attention in Alaska. More than 250 known pingos in the Yukon-Tanana Upland, central Alaska, were inventoried and described by G. W. Holmes, D. M. Hopkins, and H. L. Foster. These pingos, which occur in a zone of discontinuous permafrost, are evidence that pingos are not confined to areas of continuous permafrost, as was previously believed. All the pingos of the Yukon-Tanana Upland are of the open-system type (that is, the hydrostatic pressure results from hydraulic head related to an adjacent slope). Many show several stages of development, suggesting that sequences of growth, decay, and rejuvenation are common. The ages of the pingos are deduced to range from a few decades to several thousand years. Because the maximum age, based on radiocarbon dating of contained organic sediments, is about 7,000 years old, growth of the pingos is believed to be a product of recent climate.

Birch Creek pingo, Alaska

Studies by D. B. Krinsley (p. C133–C136) of sediments contained in a pingo along Birch Creek west of Circle, Alaska, indicate that permafrost was present during the period $6,950 \pm 400$ years B.P. to $5,720 \pm 65$ years B.P., which coincides with the postglacial thermal maximum (altithermal) episode. In the upper sediments of the pingo, lacustrine silt that was deposited in an oxbow of Birch Creek after $5,720 \pm 65$ years B.P. is now as much as 5 feet higher than the present level of Birch Creek. On the basis of this evidence the local base level is believed to have been reduced at least 5 feet in the last $5,720 \pm 65$ years, and the now-tilted sediments of the pingo are believed to have been updomed during this time.

ANALYTICAL METHODS

ANALYTICAL CHEMISTRY

Atomic absorption flame spectrophotometry

An extensive research program has been underway in the U.S. Geological Survey on testing the

⁹⁵ O. J. Ferrians, Jr., 1965, Permafrost map of Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-445. [In press]

applicability of atomic-absorption flame spectrophotometry to the analysis of geologic samples. This method is of considerable interest because of the reported high sensitivities for many elements. It is also less subject to interferences than ordinary emission flame photometry so that direct determinations with no separations are more often possible. Those elements for which high sensitivities have been reported and which are of geochemical interest are being studied systematically.

Edward Fennelly and Claude Huffman, Jr., were able to determine Pb ranging from 30 to 1,200 parts per million, and Zn from 50 to 500 ppm by atomic-absorption flame spectrophotometry in highly mineralized soil samples from Alaska. No preliminary separations were found necessary, the elements being directly determinable after dissolution of the samples in mineral acids.

The applicability of the method to the determination of Na, Ca, K, Mg, and Zn was demonstrated by Claude Hoffman, Jr., and Joseph Haffty. A few tenths of a part per million of Mg and Zn were found, concentrations too low to be determinable by emission flame spectrophotometry.

Claude Huffman, Jr., and J. D. Mensik have shown that silver can be determined quickly and reliably on a 1-gram sample to a threshold of about 1 ppm by the technique of atomic-absorption flame spectrophotometry. No difficulties were experienced on mineralized samples previously fire assayed and known to contain silver, or on selected ores and minerals known to contain many elements that interfere when other methods are used.

Ammonium ion in silicates

Robert Meyrowitz has developed a spectrophotometric procedure employing Nessler's reagent for the determination of ammoniacal nitrogen in small samples of silicate minerals. About 100 parts per million of ammonium oxide are determinable on a 50-milligram sample. The method as applied to illite, potassium feldspar, and celadonite separated from hydrothermally altered drill core from Steamboat Springs, Nev., yielded 950, 170, and 320 ppm of $(\text{NH}_4)_2\text{O}$, respectively. Muscovite from San Diego County, Calif., was found to contain 200 ppm of $(\text{NH}_4)_2\text{O}$.

Spectrophotometric determination of magnesium in phosphates

Spectrophotometric procedures based on Clayton Yellow which require no separations were developed by Robert Meyrowitz for rapid determination of

magnesium in phosphate rocks, and for the microdetermination of magnesium in phosphate minerals such as apatite, crandallite, whitlockite, morinite, and lewistonite. Interferences are masked by ethylenebis (2-oxyethylenenitrilo) tetracetic acid (EGTA) allowing 1 part of MgO to be determinable in the presence of up to 500 parts each of CaO and P_2O_5 .

Determination of silica after decomposition with HF

Irving May and J. J. Rowe have developed a method for determining silica in rocks and minerals after decomposition with hydrofluoric acid. The method avoids the loss of SiO_2 as volatile SiF_4 which customarily results when hydrofluoric decompositions are used. The samples are decomposed in a platinum-lined bomb (p. B165-B166), and the silica content of the sample solution is determined by a molybdenum-blue procedure using aluminum ion to complex interfering fluoride. Because no fluxes are introduced, both silica and alkalis can be determined on the same sample solution.

Detection of bismuth by fire assay

The routine fire assay for silver and gold has long been known to furnish indications of bismuth if this element is present in more than "trace" quantities. A series of cupellations with known amounts of bismuth made by O. M. Parker and L. B. Riley suggests that bismuth in concentrations as low as 0.1 percent may be detected during routine assaying. Furthermore, approximate bismuth percentages may be estimated by comparison with "standard" cupels having bismuth contents spaced geometrically in steps similar to the "6-step" spectrographic series, up to at least 7 percent bismuth (93 percent lead) in the lead assay button. Recognition of bismuth in samples during assaying is important as its presence may affect the silver values unless special precautions are taken.

Determination of fixed water in minerals by infrared absorption

A method was developed by I. A. Breger for the quantitative determination of fixed water (H_2O^{+1}) in rocks and minerals by infrared absorption analysis. Infrared absorption was also found useful in distinguishing between hydroxyl groups and water in minerals. Only 2 milligrams of sample is necessary for either of the infrared techniques, thus making it possible to reduce greatly the length of time spent in tedious handpicking of samples for analysis.

Removal of polonium from standard radium solutions

Polonium impurities in standard radium solutions can be easily removed by spontaneous electrochemical deposition on gold alloys. K. W. Edwards (p. D184–D188) found that when a gold-alloy disk is added to radium solutions which are 0.11–0.21 molar in hydrogen ion, 80–90 percent of the polonium is deposited within 48 hours. The deposition of polonium occurs with very slight contamination of the solution by gold, silver, and copper. No radium is deposited, and the solutions are therefore improved radium standards.

Bromine in brines and rocks

F. O. Simon and Irving May combined and refined several published procedures into a method for determining bromide in brines. Iodide is first oxidized to iodine with nitrite, and removed by boiling. Bromide is oxidized to bromate with hypochlorite, and the excess hypochlorite is destroyed by formate. Bromine is then generated by adding potassium bromide, which reacts with bromate. The bromine is determined spectrophotometrically by its bleaching effect on methyl orange. As little as 1 microgram of bromine can be determined reliably. The applicability of the method to the determination of bromine in rocks is being studied.

Confirmation of formula of langite

B. L. Ingram's determination of water in a small sample of langite confirms crystallographic evidence by Mary Mrose that the formula of langite is $\text{Cu}_4(\text{SO}_4)(\text{OH})_6 \cdot 2\text{H}_2\text{O}$. There has been uncertainty in the literature regarding the water content. The seventh edition of Dana lists langite as a monohydrate, although the original analysis by Maskelyne in 1865 and a subsequent one by Pierrot and Sainfield in 1958 lead to a dihydrate formula. Ingram's analysis also provides further evidence that posnjakite, $\text{Cu}_4(\text{SO}_4)_4(\text{OH})_6 \cdot 1\text{H}_2\text{O}$ is not the monoclinic dimorph of langite but a new mineral.

Trivalent vanadium in a chromian tourmaline

John Marinenko analyzed a chromian tourmaline from Mariposa County, Calif. Seventeen constituents were determined including ferrous iron, and vanadium in two valence states. Of interest is the finding that nearly all of the vanadium (about 5 percent vanadium in the original sample) is trivalent. The determination of valence states was a particularly difficult problem because the refractory nature of tourmaline required that the decomposi-

tion be made by fusing with powerful fluxes, without altering the valence states of the constituents.

OPTICAL SPECTROSCOPY

New device used to measure widths of intense lines

P. R. Barnett has designed and put into operation a device for use with certain types of densitometers that permits the spectroscopist to measure the widths of spectral lines recorded on plates or films. This measurement makes possible the use of lines for analytical purposes that could not otherwise be used because of their high density and broadened image. The inability to use these lines in the past has sometimes necessitated re-arc-ing the sample at a reduced exposure or after dilution. The measurements are precise and the analytical results are as accurate as those obtained by conventional densitometric methods.

Atmosphere control improves detectability of zinc

It is of special interest to the astrogeology program to be able to determine zinc in the concentration range of 1–100 parts per million. About 200 ppm is the limit normally attained in quantitative spectrochemical procedures. Using (1) a jet developed in the U.S. Geological Survey laboratory together with argo-arc atmosphere control,⁹⁶ (2) an undercut carbon sample-bearing electrode, (3) an auxiliary lens ahead of the slit, and (4) a higher current than normally used for the d-c arc, C. S. Annell and A. W. Helz have increased the detectability by several orders of magnitude so that zinc is determinable in concentrations as low as 1 ppm. The change from air to argon atmosphere control accounts for an improvement of about 10-fold in the detection limit; the use of carbon in place of graphite—2-fold; the introduction of the auxiliary lens to increase the light-gathering power of the instrument—4-fold; and the increase of the current from 15 to 25 amperes—1½-fold. Argon maintains a relatively cool steady arc; the volatilization of matrix material is minimized (lowering background), and the rate of volatilization of zinc is decreased (increasing efficiency of excitation).

X-RAY FLUORESCENCE ANALYSIS

Rock analysis by X-ray quantometer automated

Conversion to full automation of the X-ray quantometer for simultaneous determination of common rock-forming elements was completed by Leonard

⁹⁶ A. W. Helz, 1964, A gas jet for d-c arc spectroscopy: Art. 159 in U.S. Geol. Survey Prof. Paper 475-D, p. D176–D178.

Shapiro and Camillo Massoni (p. D178-D183). Steps automated include (1) positioning of a sample under the X-ray tube, (2) X-ray irradiation and X-ray fluorescence integration for a preset period of time, (3) termination of X-ray excitation and sequential read-out of integrated voltages, (4) printing of results, (5) positioning of a new sample, (6) repetition of the cycle, and (7) cutoff of instrument after a preset time. The unattended instrument can analyze 32 samples during the day and another 32 samples at night. In addition to virtually doubling productivity, the automated system yields more precise results as a result of a steady-state operation in which vacuum is maintained at a uniformly high level unattainable in manual operation.

X-ray fluorescence analysis of small samples

A combined X-ray fluorescence and chemical procedure has proved to be very useful for the total analysis of very small samples of minerals. In developing the method H. J. Rose and Frank Cuttitta sought to minimize the sample size required and to stress analyses that are otherwise unusually difficult and time consuming by chemical methods. In the procedure one milliliter of the sample solution is absorbed on powdered filter paper which is then pressed into a pellet for X-ray excitation. The method requires no more than 10 milligrams of sample and can detect microgram amounts of many elements. Applications include the determination of the zirconium-hafnium ratio in zircon and the determination of major elements in tektites, minerals, and alloys.

Rapid method for barium analyses of rocks and soils

Using an X-ray fluorescence method D. A. Brobst was able to run 200 barium analyses per day in the concentration range of 250 parts per million to 10 percent on rock and soil samples. The dry powdered samples were held for X-ray excitation in a disk 1 inch in diameter and having a suitably shaped plastic holder and cover. The X-ray fluorescence results obtained on 600 samples of Arkansas rocks when compared with barium results on the same samples obtained by spectrographic and (or) turbidimetric methods gave excellent agreement once allowance is made for the varying limits of sensitivity of the different methods.

Determination of chlorine and sulfur in fluid inclusions

X-ray fluorescence was used by P. J. Dunton for determining chlorine and sulfur in fluid inclusions. In 8 fluid inclusions in samples from Zacatacas, Mexico, chlorine concentrations ranged from 1 to 100 parts

per million and SO_4 from <1 to 10 ppm. Half a milliliter of the sample solutions was evaporated on aluminum planchettes for X-ray excitation.

Combined X-ray and chemical procedure used to measure magnesium

By first precipitating magnesium as the arsenate, then using X-ray fluorescence analysis to measure arsenic in the precipitate, W. W. Brannock and A. C. Bettiga were able to determine magnesium quantitatively. This new combined chemical and X-ray procedure promises to extend the range of X-ray fluorescence analysis to other elements that have a low atomic number and are, as a result, not now determinable.

ANALYSIS OF WATER

Atomic-absorption spectrophotometry

S. C. Downs is investigating atomic-absorption spectrophotometric techniques for the routine determination of several heavy metals. He has found that as little as 0.005 parts per million of zinc, 0.01 ppm of manganese, and 0.01 ppm of copper can be detected without enrichment techniques. Measurements of zinc are accurate and reproducible to within ± 0.01 ppm for concentrations from zero to 1 ppm and within ± 0.05 ppm for concentrations from 1 to 7.5 ppm. Measurements of manganese and copper are accurate and reproducible within ± 0.03 and ± 0.02 ppm, respectively, for concentrations from zero to 1 ppm, and within ± 0.15 and ± 0.10 ppm for concentrations from 1 to about 35 ppm. Analyses by atomic-absorption techniques require about a fourth as much time as analyses by standard chemical methods.

M. J. Fishman and K. A. Gardner have prepared tentative procedures for the determination of a number of major and minor cations in both fresh and brackish waters by atomic-absorption spectrophotometry. In general, atomic-absorption procedures are as accurate as conventional flame-photometric or titrimetric methods; in many cases they offer greater ease and simplicity of operation. As little as 5 parts per billion of magnesium can be determined directly; samples containing 10 ppm, or less, of calcium and magnesium can be analyzed much more quickly and accurately by atomic absorption than by titrimetric or colorimetric methods. The determination of strontium in both fresh water and in brines is greatly simplified, requiring neither separation nor preconcentration in order to achieve a sensitivity and accuracy of the order of 0.01 ppm.

Measurement of pesticides and herbicides by gas chromatography

W. L. Lamar, D. F. Goerlitz, and L. M. Law (1-64) have reported a convenient and extremely sensitive method (in the parts-per-trillion range) for the detection and measurement of chlorinated organic pesticides by electron-capture gas chromatography. A 4-liter sample of water is extracted with three 25-milliliter portions of purified hexane. The extracts are combined, and most of the hexane is removed by heating on a fluidized sand bath at 100°C. The extract is then transferred to a 10.00-ml volumetric flask, brought to volume with hexane, and about 50 milligrams of anhydrous sodium sulfate added. Five-microliter aliquots are then injected into an electron-capture gas chromatograph, using both Dow-11 and QF-1 columns to facilitate identification through the different partitioning effects of the two liquid substrates. Quantitative and qualitative pesticide standards are injected on the same day as the samples to aid identification and to provide corrections as necessary. The pesticides are usually identified by comparing retention times relative to aldrin on two columns. High purity of reagents and distilled water is essential; contamination of glassware is avoided by heating at 300°C overnight.

Identification and analysis of insecticides in water

A method for identifying and measuring chlorinated hydrocarbon insecticides in water was developed by Garland Stratton. The insecticides, with other organic compounds, are extracted from a 1-2-gallon sample with chloroform. The residue from evaporation of the chloroform extract is placed on a silica-gel column and eluted with hexane. The eluate is also dried. The resulting residue, dissolved in two drops of acetone, is spotted on silica-gel-loaded chromatograph paper and developed in hexane containing 1 percent acetone. The resulting chromatogram is cut into 4 sections, each of which is added to 1 milliliter of benzene to make sample concentrates for gas chromatographic analysis. This separates the insecticides into groups to facilitate analytical determination and enables detection and measurement of chlorinated hydrocarbon insecticides at very low concentrations.

D. F. Goerlitz and W. L. Lamar (1-65) also reported details of methods and techniques used to determine 5 to 10 nanograms (about 10 parts per trillion, using a 4-liter sample) of selected herbicides, including 2,4-D, 2,4,5-T, and Silvex. In tests made on 4-liter samples of distilled water containing 1.00 to 1.90 parts per billion of these herbicides, the

recoveries generally ranged from 73 to 95 percent. The analysis of samples from several canals in a National Wildlife Refuge gave values for Silvex which ranged from 0.13 to 51.5 ppb; for 2,4-D, from 0.10 to 0.25 ppb; and for 2,4,5-T, from 0.01 to 0.09 ppb.

Sodium-sensitive glass electrodes

B. F. Jones, A. H. Truesdell, and A. S. Van Denburgh (1-65) described a method using a glass electrode for the determination of sodium in closed basin waters of highly variable composition. In analyzing samples which ranged from 10 to 130,000 parts per million of sodium and from 102 to 332,000 ppm of total dissolved solids, they found in all cases that the electrode values compared favorably with conventional sodium determinations for both natural waters and stock solutions. They found no relation between differences in values obtained by glass-electrode and conventional methods and the contents of dissolved solids or of sodium. The electrode method offers good accuracy and the advantage of simplicity.

X-ray-fluorescence methods of analysis

Two X-ray-fluorescence methods have been developed and applied by P. J. Dunton to the quantitative and semiquantitative analysis of water samples for elements of atomic number 92 (U) through atomic number 14 (Si). The more sensitive methods consist of evaporating 0.5 milliliter of the sample onto aluminum foil or a filter-paper disk, which is then placed directly in the spectrograph. The sensitivity is 0.5 parts per million through atomic number 16 (S), and from 1 to 5 ppm for the lighter elements.

An alternate method must be used when the content of dissolved solids in the sample exceeds 1,000 ppm, because of the tendency for the dried residue to flake off the foil or paper disk. In this method a 4-ml sample aliquot is transferred to a sample cell fitted with a 1/4-mil Mylar film, and then is placed directly in the X-ray spectrograph. The sensitivity is 1 ppm for elements through atomic number 19 (K), and 5-10 ppm for elements lighter than potassium.

Determination of lead-210

A method for the determination of trace amounts of lead-210 has been developed by J. O. Johnson and K. W. Edwards. The lead is concentrated by coprecipitation with barium sulfate, purified by ion exchange, and finally reprecipitated as the chromate. The final precipitate is aged to allow ingrowth of 5-day bismuth-210; the beta activity of this daughter

nuclide is determined on a low-background counter. The detection limit (based on 95-percent confidence level) is 2 picocuries per liter for a 500-ml sample.

Method for determination of phosphate in water

M. J. Fishman and M. W. Skougstad (p. B167–B169) simplified an existing method to make it suitable for the rapid field and laboratory determination of phosphate in natural water. Replicate determinations are reproducible to ± 0.01 parts per million, and results agree to ± 0.01 ppm with those determined by the standard method in use in the U.S. Geological Survey.

Changes in composition of stored water samples

K. V. Slack and D. W. Fisher reported (p. C190–C192) changes in composition observed when water samples were stored in sealed bottles for periods as long as 5 months. Three different types of polyethylene containers were used. Following exposure for 3 months to alternating 12-hour light and dark periods, the pH of replicate samples ranged from 9.27 to 10.17, HCO_3^- was not detected in most samples, CO_3 concentrations ranged from 7 to 13 parts per million, and free CO_2 and NO_3^- were below detection. After an additional 2 months of dark storage, pH decreased 2 units relative to the illuminated condition, free CO_2 and HCO_3^- concentrations increased greatly, and CO_3 was below detection. The changes were attributed to photosynthesis by algae during illumination and to re-solution of substances following decay of the algae in the dark.

Carbonate brines

S. L. Rettig and B. F. Jones (1–64) analyzed samples of carbonate brines having a CO_2 content in the range of 0.5–9 percent by potentiometric titration and by manometric measurement of the total CO_2 evolved upon acidification. The potentiometric data show that the actual end points may differ significantly from the pH values of 8.2 and 4.5 commonly used as end points in such titrations. Some of the errors in the potentiometric method can be evaluated by comparison with the manometric method.

HYDROLOGIC MEASUREMENTS AND INSTRUMENTATION

Multiple-channel digital recording systems

R. N. Cherry (p. D222–D224) and G. F. Smoot have developed instruments which automatically record several hydrologic variables on punched tape for automatic data processing. These systems are battery powered for unattended operation over periods of several months at remote locations. Each

employs a digital recorder. Cherry's system records in sequence 2 values of water temperature and 2 values of the specific conductance of the water. Smoot has developed sensors to measure temperature, rainfall, water stage, specific conductance, and soil moisture, and is developing sensors for wind direction and velocity, and for water turbidity. A wide range of sensors and of time intervals for measurements can be used in a single system.

Instruments for measuring velocity of flow of water

A miniature current meter has been developed by G. F. Smoot for the measurement of longitudinal velocities and velocity fluctuations. This meter, utilizing a 30-mm-diameter Ott Minor propeller, has an increased sensitivity response provided by an improved bearing design and a drag-free electronic revolution-sensing system. The output from the sensing system is a series of pulses. Within the response range of the meter, minute rotational variations which are recorded as instantaneous frequency fluctuations in the sequence of pulses are proportional to the encountered turbulence. Initial field tests of the meter produced reasonable results.

Instruments which measure flow velocities by means of strain gages have been developed for use in surface streams by H. P. Guy, D. B. Simons, and J. B. Bole, and for use in deep water wells by J. E. Eddy. In these instruments the strain gages are components of wheatstone bridge circuits; the strain, expressed as electrical output, is proportional to the deforming force of the current and hence to the velocity. The instrument for surface streams, which has a drag-wire velocity probe, has been tested in flume studies to within 0.005 inch of the flume floor. The well instrument, in which the strain gage is mounted on a vane normal to the direction of flow, has been tested in wells at depths as great as 1,000 feet.

G. F. Smoot reports development of a continuous-tone sonar device utilizing the Doppler effect. A beam of ultrasonic energy projected into an inhomogeneous liquid is scattered by the suspended particulate matter in the liquid. Part of the energy, reflected to a receiver, shifts in frequency if there is a net movement of the inhomogeneities with respect to the transmitter or receiver. If the scatterers are stationary with respect to the liquid, the observed Doppler shift is proportional to the velocity of the liquid.

H. O. Wires and W. Smith have tested an acoustic-velocity meter which sends a cycled pulse diagonally back and forth across a stream. The difference in

transit time of the upstream and downstream pulses is related to water velocity. In the tests, on a canal and on the Snake River in Washington, measured discharges were within 5 percent and 2 percent, respectively, of known values.

Automatic data processing

Discharge in channels having flat and variable water-surface gradients can be computed from the water-surface stage and an index of the velocity provided by the deflection of a vane in the current. S. D. Leach reports that use of digital-punch equipment for recording stage and deflection at Florida stream-gaging stations has permitted rapid computation, by digital computer, of discharge at stations using deflection meters.

Measurement of stream flow by dilution method

A concentrated solution of radioactive gold was injected at a constant and known rate into an operating turbine penstock 15 feet in diameter, and the rate of water flow in the penstock was computed from observations of the time of passage of the tracer to a section 220 feet downstream from the injection point. B. J. Frederick reported that computed flow rates for 7 tests in the 1,100–1,650 cubic-foot-per-second range were within an average of about $2\frac{1}{2}$ percent of the discharge obtained from the powerplant integrator, which had been calibrated by the Gibson method.

J. F. Bailey and E. D. Cobb made 12 dye-dilution discharge measurements on 4 small streams in Maryland and Virginia. The average error for the 12 discharges, relative to discharge measured by current meter, was 5 percent. The dye was injected at a constant rate into discharges which ranged from 10 to 50 cfs. The reaches of channel used were from 200 to 1,800 feet long.

F. A. Kilpatrick reports favorable results from the dye-dilution method of measuring flows in open channels for discharges up to 3,000 cfs. The excessive channel length required in some instances for complete mixing, when using a point source of dye injection, suggests the desirability of using a line source.

Fluorometer correction curves

Temperature of sample and temperature of fluorometer compartment influence readings on the Turner-111 fluorometer. Bernard Dunn and D. E. Vaupel (p. D225–D227) have prepared correction curves for use when sample and compartment temperatures cannot be controlled.

Portable deep-well sampler

R. N. Cherry (p. C214–C216) has developed a portable deep-well sampler which isolates a section of well between two inflatable boots. The water sample is pumped from the isolated section by a submersible pump mounted between the boots. Auxiliary instruments can be used to measure temperature, specific conductance, or other characteristics of the water in the isolated interval during pumping.

Data storage, retrieval, and compilation

A system for the storage and retrieval of groundwater data has been devised by a committee of hydrologists under the chairmanship of S. M. Lang. The basic elements of the system are: (1) a well-schedule form for use in the field to record the data obtained during an inventory of wells, and (2) punch cards that are used for the actual storage and retrieval of the data. The schedule form is keyed to the punch cards in such a manner that a punch operator can transfer data conveniently from the schedule form to the cards. The system is being used on a trial basis in the States of Florida, Iowa, and Wyoming.

M. D. Edwards and T. H. Woodward have developed programs for compiling and tabulating water-quality data, and for computing such parameters as ionic strength of solution, activity coefficient, and activity of each component of the dissolved solids, from the analytical data.

J. M. McNellis and C. O. Morgan have developed digital-computer programs which compile tables of well records and well logs from data cards, and prepare maps of the thickness, saturated thickness, and other characteristics of geologic or geohydrologic units.

Concentration of water samples

One of the problems encountered in the study of trace elements and organic materials in natural waters is that of rapidly concentrating a sample to a residue without contamination or appreciable alteration of the dissolved solids. V. C. Kennedy has developed a successful technique for rapid concentration, using a combination of a commercial evaporator and a commercial freeze-dryer.

Measurement of dissolved oxygen

A low-cost small kit for measuring dissolved oxygen has been designed by K. V. Slack for use in water-quality studies. Oxygen determinations can be made more rapidly and safely, and with less chance of contamination, than is possible with the usual method. All necessary equipment and supplies

for performing the Alsterberg modification of the Winkler method can be packed into an index-card file box measuring $3 \times 5 \times 2\frac{3}{4}$ inches. The cost of the kit, exclusive of reagents, is about \$15.00. An oxygen determination for a 10-milliliter sample takes less than 10 minutes; precision is better than 0.5 ppm.

Measurement of cation-exchange rates

A method of measuring rates of cation exchange, developed by V. C. Kennedy and T. E. Brown, involves the use of specific-ion electrodes to monitor the rate at which monovalent cations are released into solution when a monovalent, monionic, solid exchanger is added to a solution of a divalent cation. As the monovalent cation is released from the solid by exchange with the dissolved divalent cation, the electromotive force of the specific-ion electrode reflects changes in the concentration of monovalent

cation in solution. The emf of the specific-ion electrode forms the input to a pH meter whose output is plotted on a potentiometric strip recorder. When calcium is used to exchange for sodium, about 90-percent exchange occurs in 1–2 seconds for kaolinite, montmorillonite, or illite in particles smaller than 2 microns. Exchange on sand-size stream sediment containing fine-grained rock fragments may take 10 minutes or more before 90-percent completion is attained.

R. L. Malcolm and V. C. Kennedy have used the exchange of barium for potassium in an attempt to distinguish various clay minerals on the basis of differences in rates of exchange. Results indicate that the rate-of-exchange curve for vermiculite differs significantly from that of kaolinite, montmorillonite, and illite. Thus, rate-of-exchange studies appear to be another tool which can be used in the characterization of clay minerals.

TOPOGRAPHIC SURVEYS AND MAPPING

MAPPING ACCOMPLISHMENTS

Objectives of National Topographic program

The major function of the Topographic Division of the U.S. Geological Survey is to prepare and maintain maps of the National Topographic Map Series covering the United States and other areas under the sovereignty of the United States of America. The individual series, at various scales, constitute a fundamental part of the background information needed to inventory, develop, and manage the natural resources of the country. Other Division functions include the production of special maps and research and development in techniques and instrumentation.

In addition to the maps described below, the Topographic Division prepares shaded-relief maps, United States base maps, special maps, and also a few planimetric maps.

Procedures for obtaining copies of the maps and map products of the Survey are given in the section "How To Order Geological Survey Publications."

Series and scales

All topographic surveys, except those in Alaska, conform to standards of accuracy and content required for publication at the scale of 1:24,000. Initial publication scale may be either 1:24,000 or 1:62,500, depending on the need. If 1:62,500-scale maps are published initially, the 1:24,000-scale surveys, in the form of photogrammetric compilation sheets, are available as advance prints and for future publication at the larger scale. For Alaskan maps, the publication scale is 1:63,360 or "inch-to-the-mile."

Coverage of the Nation

Approximately 63 percent of the total area of the 50 States, Puerto Rico, and the Virgin Islands (fig. 10) is covered by published standard quadrangle maps at scales of 1:20,000 (Puerto Rico only), 1:24,000, 1:62,500, and 1:63,360 (Alaska only). An additional 8 percent of the total area is covered by topographic surveys which are available as advance prints at these scales.

During fiscal year 1965, 1,078 maps were published covering previously unmapped areas equivalent to 3 percent of the total area referred to above. In addition, 392 new maps at the scale of 1:24,000, equivalent to approximately 1 percent of the total area, were published to replace 15-minute quadrangle maps (scale 1:62,500) not meeting present needs. For the extent and location of map coverage, see figure 11.

Map revision and maintenance

During fiscal 1965, about 42,000 square miles of 7½-minute mapping was added to the growing backlog of maps needing revision. Concurrently, the backlog was diminished by revising about 11,000 square miles of mapping, leaving about 403,000 square miles of 7½-minute mapping needing revision at the end of the year (fig. 12).

1:250,000-scale series

The 48 conterminous States and Hawaii are 99 percent covered by 1:250,000-scale maps originally prepared as military editions by the U.S. Army Map Service. As these maps are completed, certain changes and additions are incorporated to make them more suitable for civil use. This series of maps is being revised and maintained by the Topographic Division. Maps of Alaska at this scale are prepared and published by the U.S. Geological Survey. Coverage of the 50 States, Puerto Rico, and the Virgin Islands by 1:250,000-scale maps and the work in progress are shown on figure 13.

State maps

State maps are published at scales of 1:500,000 and 1:1,000,000, except for Alaska, which is covered by base maps published at scales of 1:1,584,000 and 1:2,500,000, and Hawaii, which is not yet covered by any of these types of maps.

Thirty-one maps covering 35 States and the District of Columbia, compiled to modern standards, have been published in a new series comprising as many as four editions: base; base and highways; base, highways, and contours; and shaded relief on a modified base. As shown on figure 14, other conterminous States are covered by an earlier series.

FIGURE 10.—Progress of 7 1/2- and 15-minute quadrangle topographic mapping in the 50 States, Puerto Rico, and the Virgin Islands.

FIGURE 11.—Status of 7½- and 15-minute quadrangle topographic mapping.

FIGURE 12.—Status of revision of large-scale mapping.

FIGURE 13.—Status of 1:250,000-scale mapping.

NOTE: The letter "P" on the pattern for "mapping in progress, new series (contour only)" in Oregon, Wyoming, and Iowa indicates that planimetric maps of these States have already been published.

FIGURE 14.—Status of State maps.

Metropolitan areas

Metropolitan-area maps are prepared by combining on one or more sheets the 7½-minute quadrangles that cover a metropolitan area. Maps of 59 metropolitan areas have been published, including 1 new map completed during fiscal year 1965. Work in progress includes the revision of 8 others. Maps in the metropolitan-area series include:

PUBLISHED

Albuquerque, N. Mex.	Louisville, Ky.
Anchorage, Alaska	Madison, Wis.
Atlanta, Ga.	Wilwaukee, Wis.
Austin, Tex.	Minneapolis-St. Paul, Minn.
Baton Rouge, La.	New Haven, Conn.
Boston, Mass.	New Orleans, La.
Bridgeport, Conn.	New York, N.Y. (8 sheets)
Buffalo, N.Y.	Norfolk-Portsmouth-Newport News, Va.
Champaign-Urbana, Ill.	Oakland, Calif.
Chattanooga, Tenn.	Peoria, Ill.
Chicago, Ill. (3 sheets)	Philadelphia, Pa. (2 sheets)
Cincinnati, Ohio	Pittsburgh, Pa.
Cleveland, Ohio	Portland-Vancouver, Oreg.-Wash.
Columbus, Ohio	Rochester, N.Y.
Davenport-Rock Island-Moline, Iowa-Ill.	Salt Lake City, Utah
Dayton, Ohio	San Diego, Calif.
Denver, Colo.	San Francisco, Calif.
Detroit, Mich. (2 sheets)	San Juan, P.R.
Duluth-Superior, Minn.-Wis.	Seattle, Wash.
Fort Worth, Tex.	Shreveport, La.
Gary, Ind.	Spokane, Wash.
Hartford-New Britain, Conn.	Tacoma, Wash.
Honolulu, Hawaii	Toledo, Ohio
Houston, Tex.	Washington, D.C.
Indianapolis, Ind.	Wichita, Kans.
Juneau, Alaska	Wilkes-Barre-Pittston, Pa.
Knoxville, Tenn.	Wilmington, Del.
Little Rock, Ark.	Worcester, Mass.
Long Beach, Calif.	Youngstown, Ohio
Los Angeles, Calif. (2 sheets)	

IN REVISION

Baton Rouge, La.	Salt Lake City, Utah
Cincinnati, Ohio	San Juan, P.R.
Cleveland, Ohio	Spokane, Wash.
Little Rock, Ark.	Washington, D.C.

National park maps

Maps of 40 of the 201 national parks, monuments, historic sites, and other areas administered by the National Park Service have been published and are available for distribution. These usually are made by combining all existing quadrangle maps of the area into one map sheet, but occasionally surveys are made covering only the park area. Most of the other parks, monuments, and historic sites are shown on maps of the standard quadrangle series. Published maps in the National Park series include:

Acadia National Park, Maine	Isle Royale National Park, Mich.
Bandelier National Monument, N. Mex.	Lassen Volcanic National Park, Calif.
Black Canyon of the Gunnison National Monument, Colo.	Mammoth Cave National Park, Ky.
Bryce Canyon National Park, Utah	Mesa Verde National Park, Colo.
Canyon de Chelly National Monument, Ariz.	Mount McKinley National Park, Alaska
Carlsbad Caverns National Park, N. Mex.	Mount Rainier National Park, Wash.
Cedar Breaks National Monument, Utah	Olympic National Park, Wash.
Colonial National Historical Park (Yorktown), Va.	Petrified Forest National Monument, Ariz.
Colorado National Monument, Colo.	Rocky Mountain National Park, Colo.
Crater Lake National Park, Oreg.	Scotts Bluff National Monument, Nebr.
Craters of the Moon National Monument, Idaho	Sequoia and Kings Canyon National Parks, Calif.
Custer Battlefield, Mont.	Shenandoah National Park, Va. (2 sheets)
Devils Tower National Monument, Wyo.	Vanderbilt Mansion National Historic Site, N.Y.
Dinosaur National Monument, Colo.-Utah	Vicksburg National Military Park, Miss.
Franklin D. Roosevelt National Historic Site, N.Y.	Wind Cave National Park, S. Dak.
Glacier National Park, Mont.	Yellowstone National Park, Wyo.-Mont.-Idaho
Grand Canyon National Monument, Ariz.	Yosemite National Park, Calif.
Grand Canyon National Park, Ariz. (2 sheets)	Yosemite Valley, Calif.
Grand Teton National Park, Wyo.	Zion National Park (Kolob section), Utah
Great Sand Dunes National Monument, Colo.	Zion National Park (Zion Canyon section), Utah
Great Smoky Mountains National Park, N.C.-Tenn. (2 sheets)	

Million-scale maps

The worldwide million-scale series of topographic quadrangle maps was originally sponsored by the International Geographical Union and designated the International Map of the World on the Millionth Scale (IMW). The conterminous United States will be covered by 53 maps, 17 of which were produced before 1955. At that time the U.S. Army Map Service began a military series at 1:1,000,000 scale. Eventually this military series will be modified slightly and published in the IMW series (fig. 15).

Three of the maps, Hudson River, Mississippi Delta, and San Francisco Bay, are no longer available as IMW maps, but the areas are covered by maps in the military series. Both the IMW and military series are available for Boston, Chesapeake Bay, Hatteras, Mount Shasta, and Point Conception. In addition, the American Geographical Society has

FIGURE 15.—Status of 1:1,000,000-scale topographic mapping.

published the Sonora, Chihuahua, and Monterrey maps; and Canada, the Regina and Montreal maps. Puerto Rico is covered by two maps, compiled by the American Geographical Society and published by both the Society and the Army Map Service.

Some maps of the military series have been modified for broader civil use by changing them to conform to the IMW sheet lines and sheet numbering system, but they do not meet IMW specifications in all respects. These maps are recognized by the United Nations Cartographic Office as provisional editions in the IMW series.

Work in progress includes four new maps: Blue Ridge, Lake Superior, Mt. Whitney, and Pikes Peak.

Official United States wall map

A revised edition of the official United States wall map, prepared at the direction of the 88th Congress, was published in January 1965. This 42- by 65-inch, 12-color map, at the scale of 1:3,168,000 (1 inch equals 50 miles), shows national parks and monuments, national forests, Indian reservations, wildlife refuges, public lands, historical boundaries, and 7,500 selected cities, towns, rivers, and lakes. Insets at smaller scales show Alaska, Hawaii, the Canal Zone, and Puerto Rico and the Virgin Islands. (Copies of this map are available only from the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402, at \$2 each.)

MAPPING IN ANTARCTICA

The topographic mapping of Antarctica, conducted as a part of the U.S. Antarctic Research Program (USARP) of the National Science Foundation, was continued during fiscal year 1965. Four topographic engineers went to Antarctica during the austral summer of 1964-65 to obtain geodetic control for the topographic mapping program and to execute surveys in support of other scientific disciplines. Also, a specialist in aerial photography was again assigned to Christchurch, New Zealand, for photographic liaison duty with the U.S. Navy.

Topographic field operations

E. R. Soza, Alfred Zavis, C. E. Morrison, and D. E. Reed made two helicopter-supported electronic-distance traverses in the David Glacier and Northern Victoria Land areas (fig. 17) to establish control for topographic mapping and to establish geodetic ties between previously established control nets. This party also completed the precise Byrd Station glaciological strain net initiated in the 1963-64 austral season, so that this net now extends eastward from

New Byrd Station to the divide between the Filchner Ice Shelf and the Ross Ice Shelf. Snow accumulation markers were set along the new traverse, and accumulation measurements were made on the markers and poles established the previous year. The total traverse is a 106-mile-long chain of 52 quadrilaterals. To obtain the required survey accuracy of 1 part in 60,000, the engineers used theodolites for measuring angles and electronic equipment for measuring distances. This party also established glaciological strain nets at mileposts 60 and 120 of Army-Navy Drive (the sledge trail between Little America and Byrd Station) and determined new geographic positions for these two mileposts. In addition, positions were determined for four ice holes in McMurdo Sound for University of Arizona biologists using scuba diving techniques to investigate polar adaptation of the Weddell seal.

Supported by a U.S. Army helicopter detachment, E. R. Soza and C. E. Morrison established control by electronic-distance traverse from McMurdo Station around the perimeter of Ross Island. They also completed an electronic-distance traverse near Byrd Station to determine the position of the University of Washington's VLF (very low frequency) long-wire antenna facility, established to study the properties of the earth's magnetic field.

Alfred Zavis and D. E. Reed obtained 5 stellar observations and 8 sets of solar observations at the previously established New Byrd Astronomic Station. Also at Byrd Station, the geographic position of Stanford University's readout center of the NASA POGO (polar orbiting geophysical observatory) satellite was determined. In addition, these engineers extended a second-order triangulation net at McMurdo for detailed mapping in anticipation of new construction at the main United States base.

Aerial photography

U.S. Navy Air Development Squadron 6 (VX-6) obtained aerial photographs for mapping in accordance with U.S. Geological Survey specifications. W. R. MacDonald was assigned to the U.S. Navy Photographic Laboratory at Christchurch, New Zealand, to advise on the quality of developed photographs, to assist with further planning, and to recommend reflights when necessary. MacDonald also served as visual navigator on flights over troublesome and high-priority areas.

This season's aerial photography program was the most successful in the history of Antarctic photographic operations (fig. 16). This is attributed to the fact that, in addition to the C-121J Super Con-

FIGURE 16.—Plan for optical aerial mapping photography of Antarctica, prepared by the National Science Foundation and the Geological Survey, May 1965.

FIGURE 17.—Index map of Antarctica, showing status of topographic mapping by the U.S. Geological Survey as of June 30, 1965.

stellation aircraft previously used in Antarctica photographic missions, an LC-130F Hercules aircraft, specially designed for aerial photographic work, was also used. The LC-130F is ski equipped, permitting the aircraft to make emergency landings on unprepared snow and ice surfaces. Also, it accommodates a removable 3,600-gallon internal gas tank which increase its flight time by approximately 5 hours. As a result, a final analysis of the season's photography indicates that areas totaling about 182,500 square miles were photographed acceptably for use in the Geological Survey mapping program (as compared to a net total of 244,500 square miles obtained between 1959 and 1964).

Cartographic activities

Six 1:250,000-scale topographic maps of the McMurdo area were published in shaded-relief editions, making a total of 19 sheets at this scale now available (fig. 17). Mapping at the same scale is in progress for 8 sheets in the Queen Maud Range, 6 in the Britannia Range, 9 in the Queen Alexandra Range, and 18 in Victoria Land.

A 1:500,000-scale shaded-relief planimetric sketch map was published of Northern Victoria Land, showing an area of 50,000 square miles north of lat 73° S., and between long 158° and 171° E. A 1:40,000-scale planimetric manuscript was compiled of the Neptune Range and used in revising the 1:500,000-scale planimetric sketch map of the Pensacola Mountains.

RESEARCH AND DEVELOPMENT

To support the National Topographic Program, continuing research and development are needed to improve techniques, instruments, and materials. The chief objectives are to improve the quality and usability of the maps and to reduce the cost of producing them.

ESTABLISHMENT OF TEST SITE FOR MAPPING RESEARCH

Many research projects in topographic surveying and mapping require a group effort by engineers and cartographers with different backgrounds and talents to achieve useful and meaningful results. A project of this type, completed in 1965 by the combined effort of field and photogrammetric engineers, was the establishment of a test site near Phoenix, Ariz., with a high density of control points. Within the test area, a 16-mile square, 31 points had been established and marked with permanent targets by

the U.S. Army Map Service. Many more points, however, were needed to provide adequate source data for testing and evaluating photogrammetric techniques and instrumentation.

Additional elevations were established at quarter-mile intervals within a 2- by 3-mile area, at half-mile intervals within a 3- by 5-mile area, and at mile intervals within an 8- by 8-mile area. Within the 8- by 8-mile area, additional horizontal points were established at mile intervals. Half of the total 16- by 16-mile area contains control points at intervals of 2 miles or less, and the rest contains control points at 4-mile intervals.

Each of the 223 new points was marked with a permanent monument for future recovery and also was temporarily marked with polyethylene sheeting to provide square target images on aerial photographs. Three sizes of targets were used: 6 by 6 feet, 9 by 9 feet, and 12 by 12 feet. Including the 31 Army Map Service points, the test area contains a total of 254 control points.

Horizontal positions were extended by second-order triangulation and electronic-distance traverse. Elevations were extended by fourth-order fly leveling, with closure errors held to 0.1 foot or less.

After the points had been surveyed and marked with targets, aerial photographs were taken at several flight heights, ranging from 3,000 to 24,000 feet. The photographs were taken with wide-angle (90°) and super-wide-angle (120°) cameras, on film and glass plates, with panchromatic and infrared emulsions. The photographs and survey data are now being used in an evaluation of super-wide-angle photogrammetric instrumentation and in a test of analytical aerotriangulation techniques.

FIELD SURVEYS

Bench-mark stability tests

A research project was started in July 1964 to investigate the stability and the comparative cost of various methods of setting bench marks in different types of soil. A test course about 3,000 feet long was established, containing 41 points marked in 5 different ways: by tablets embedded (1) in boulders and (2) in concrete posts, by tablets crimped (3) on special copper-coated rods and (4) on ordinary reinforcing steel, and (5) by copper nails in tree roots. The stability of the marks was determined by repeated leveling of super-first-order accuracy at 1- to 2-month intervals.

Although the tests are not completed, the preliminary results indicate that the copper-coated steel rods provide the most stable setting for the tablets.

Compared to conventional concrete posts, they cost only 50 percent as much and are much easier to transport and to set. They are obtained in 5-foot sections which can be joined by crimping with special tools, and can be driven with a gasoline-powered hammer. The tablets are fastened to the rods by crimping. Arrangements have been made to use this type of bench-mark setting in all topographic fieldwork.

A yearly cycle of variation in elevations has been observed. All marks showed upheaval in spring and summer and settling in fall and winter, in some cases as much as 5 millimeters. The tests are being continued to learn more about the long-range effects of this cyclic variation.

Military evaluation of ABC system

The U.S. Geological Survey supplied equipment components and engineer advisers for a military evaluation of the Survey-developed AirBorne Control System near Fort Sill, Okla., in September and October 1964. The U.S. Army Artillery Board was responsible for the evaluation, assisted by the Armor Board, the 30th Engineers, and Geodesy, Intelligence and Mapping Research and Development Agency (GIMRADA).

Summary conclusions of the evaluation are that the ABC System is suitable for establishing control for topographic mapping but that it is not an alternative to either of the military systems with which it was compared, the Visual Airborne Target Locator System and the Long Range Survey System. GIMRADA is now studying the adaptability of ABC System components to military design and the possible use of available military components.

Level-rod graduation strips

The evaluation of photographically prepared invar graduation strips for precise level rods, begun last year, is continuing. Several prototype strips have been calibrated. Sections of some of them have exceeded the specified accuracy. The reasons for failure of the prototype strips to meet specifications fully are being investigated, and improved testing and installation methods have been developed.

Commercially printed self-reading facing strips and check scales have been laboratory tested. These strips have withstood temperatures ranging from -65°F to $+155^{\circ}\text{F}$, and relative humidities from 16 percent to 100 percent. Strips of this material will be installed on rods to be used in the field as soon as possible.

An investigation of the possibility of using similar strips for stadia rods is underway, with a good

chance that a more scale-stable material can be found. Prospective savings over the cost of rebuilding and maintaining present silk-screen printing equipment will probably save several hundred dollars annually and will produce superior graduations.

PHOTOGRAMMETRY

Analytical aerotriangulation

The availability of high-speed electronic computers has led to the development of analytical systems of aerotriangulation for establishing the control needed to orient and position stereomodels for map compilation. The U.S. Geological Survey direct geodetic constraint method has been developed to solve the entire photogrammetric aerotriangulation problem mathematically. Input data for the solution consist of x and y photocoordinates measured on glass plates by means of a Mann monoscopic comparator.

To facilitate the measuring and recording of the great number of precise coordinates required for aerotriangulation, the Mann comparator has been recalibrated and modified by the addition of electric motor drives for the x and y motions, an analog-to-digital converter, an automatic electric typewriter, and an automatic card punch. These refinements have increased the speed and precision of operation and have virtually eliminated the possibility of gross errors in recording photocoordinates.

In an early test of the direct geodetic constraint method, a strip of 7 photographs taken at a flight height of 10,000 feet with a 6-inch camera was aerotriangulated with control limited to 2 horizontal points and 6 vertical points. The root-mean-square error (RMSE) for 13 vertical test points was 0.5 foot, and for 17 horizontal test points the RMSE was 1.2 feet.

Operational tests of the method are now in progress for the Middleburg and Arcola, Va., $7\frac{1}{2}$ -minute quadrangles. All the constraining ground-control points and all the test points for the project were marked with specially designed targets before the area was photographed on film and glass-plate negatives. Preliminary results indicate a high order of horizontal and vertical accuracy.

Super-wide-angle photogrammetric systems

The introduction of super-wide-angle cameras and stereoplotters has provided a potential for increased economy in photogrammetric mapping. Photographs taken with the new cameras (angular field, 120° ; focal length, 3.5 inches) cover 3 times the area of

those taken at the same flight height with wide-angle cameras (angular field, 90° ; focal length, 6 inches).

Comparative studies and production tests of available super-wide-angle stereoplotters continued during 1965, to determine the type most suitable for and adaptable to Geological Survey mapping operations. The accuracy and economy of super-wide-angle systems as compared with wide-angle systems for analytical aerotriangulation are also being investigated. Stereoplotters under test include the Wild Aviograph B8, the Kelsh K8, the Kern PG-2, the Bausch and Lomb Super Balplex (ER-32), and the Wild WH-6.

Automatic mapping instrumentation

As a means of investigating the possibilities and applications of automation in mapping, the U.S. Geological Survey in 1965 acquired a Stereomat III on indefinite loan from the U.S. Army Map Service. This instrument, designed for automatic stereocompilation, contains electronic image-correlating circuits that sense x and y parallax in the stereomodel. The error signals due to y parallax control servomotors attached to the optical projectors and automatically bring the projectors into proper orientation relative to each other. The x -parallax signals control servomotors which move the scanning head in the x , y , and z directions according to the selected mode of operation—contouring, profiling (in either x or y direction), or drainage compilation. It is expected that the profiling output of the Stereomat, in digital form, can be used to control the Orthophotoscope for automatic production of orthophotographs.

Multiband aerial photography

Simultaneous multiband photographs taken with a special six-unit camera were evaluated for potential utility in photogrammetric mapping. Four types of 70-mm film were used: standard color, false color (camouflage detection), infrared, and standard panchromatic (exposed through red, green, and blue filters). Coverage was obtained over typical areas of urban development, dense timber cover, and intricate hydrographic detail at flight heights of 6,000, 18,000, and 30,000 feet. The photographs were compared visually to assess their relative quality for photointerpretation of ground detail at the three flight heights.

The false color film was considered to have the greatest photointerpretive value, and the standard color film was rated a close second. The infrared film gave the best differentiation of ground detail in dense timber. Little falloff in photointerpretive quality was noted in the photographs taken at 30,000 feet as compared with those taken at 18,000 feet.

Orthophotomap research

The availability of instruments and practical techniques for preparing orthophotographs (that is, photographs corrected for image displacements due to tilt and relief) has led to the experimental development of a new map product, the orthophotomap. This new type of map presentation consists of a precise assembly of orthophotographs in standard quadrangle format, to which limited cartographic treatment in the form of symbols, names, boundaries, and marginal information is applied. The orthophotomap may be printed in multicolor form and is considered to have wide potential application in engineering and other scientific disciplines.

An orthophotomap of the Cave Creek 2 SE, Ariz., $7\frac{1}{2}$ -minute quadrangle has been prepared to demonstrate the additional map information provided by photographic imagery of natural terrain features in areas of sparse cultural detail. The color tones used in printing were adjusted to match very closely the natural colors of the terrain. (The original orthophotographs were prepared from normal black-and-white aerial photographs.) The orthophotomosaic was overprinted with contours, elevations, and selected cartographic symbols to augment the portrayal. The resultant presentation, which meets the National Map Accuracy Standards, contains much more detailed information about the terrain than can be expected of a conventional line-drawn map.

Color photomaps of large swamps

Two projects are underway to provide color photomaps of large swamp areas, the Everglades in Florida and the Okefenokee Swamp in southeastern Georgia. These projects differ from the orthophotomap project described above in that the range of topographic relief is too small to require the use of orthophotographs, and color aerial photographs are being used as part of the source data.

The shortcomings of line-drawn map representation of large swamps have long been recognized. Therefore, the new approach consists in preparing a precisely controlled photomosaic, adding cartographic delineation and accentuation where appropriate, and color-coding features of major interest. Some of the terrain features to be differentiated are pine keys, bay heads, mangrove, and sawgrass. Other features to be shown include conspicuous airboat trails, saline-water boundaries, and survival water holes, as well as standard planimetric features, such as roads, buildings, canals, and levees. Ground elevations will also be shown.

Three types of aerial photographs will be used in preparing the photomaps—color, infrared, and standard panchromatic. It is expected that the color photographs will be particularly useful in interpreting and delineating the various types of terrain.

NATIONAL ATLAS

Work on the National Atlas during 1965 consisted in compiling general reference maps and in preparing proof prints of the physical and thematic maps of the climatic and water sections. Parts of the historical section were started, and proofs of some of these pages are expected early in fiscal year 1966.

In addition, experimental printings were made in cooperation with the Publications Division of the U.S. Geological Survey to determine suitable proce-

dures, colors, and type of paper for the Atlas. Other investigations will lead to a final selection of type of binding, cover fabric, and type of plastic overlay sheets for the bound volumes.

During the year, the Bureau of the Budget issued a revision of Exhibit B to Circular A-16 which requires all interested Federal agencies to appoint liaison officers for the National Atlas Project. On February 16, 1965, a meeting of all the liaison officers was held. As a result of this initial meeting, activity on the atlas was increased significantly, and much planning of content and format was completed during the second half of the year.

For geologic subjects covered see "National Atlas," a subheading under "MAPS OF LARGE REGIONS" in the section "REGIONAL GEOLOGY."

COOPERATING AGENCIES FOR FISCAL YEAR 1965

FEDERAL AGENCIES

Agency for International Development
Air Force:
 Cambridge Research Center
 Technical Application Center
Army:
 Corps of Engineers, Waterways Experiment Station
 U.S. Army—Europe
Atomic Energy Commission:
 Division of Raw Materials
 Division of Reactor Development
 Idaho Operations Office
 Nevada Operations Office
 Research Division
 San Francisco Operations Office
 Savannah River Operations Office
Department of Agriculture:
 Forest Service
 Soil Conservation Service
Department of Commerce:
 Weather Bureau
Department of Defense:
 Advanced Research Projects Agency
 Defense Atomic Support Agency
 Defense Intelligence Agency
 Office of Scientific Research
Department of Health, Education and Welfare:
 Public Health Service
Department of the Interior:
 Bonneville Power Administration
 Bureau of Commercial Fisheries
 Bureau of Indian Affairs
 Bureau of Land Management
 Bureau of Mines
 Bureau of Reclamation
 Bureau of Sport Fisheries and Wildlife
 National Park Service
 Office of Minerals Exploration
 Office of Saline Water
 The Alaska Railroad
Department of State
District of Columbia
Executive Office of the President—Office of Emergency Planning
Federal Aviation Agency
Federal Power Commission
General Services Administration
National Aeronautics and Space Administration
National Science Foundation
Navy:
 Office of Naval Research

Navy—Continued
 Petroleum and Oil Shale Reserve
 Radiological Defense Laboratory
Tennessee Valley Authority

STATE, COUNTY, AND MUNICIPAL AGENCIES

Alabama:
 Alabama Highway Department
 City of Huntsville
 City of Mobile
 Department of Conservation
 Geological Survey of Alabama
 Water Improvement Commission
Alaska:
 City of Anchorage
 State Highway Department
Arizona:
 Arizona Highway Department
 Buckeye Irrigation Company
 City of Flagstaff
 City of Tucson
 City of Williams
 County of Apache, Superior Court
 Flood Control District of Maricopa County
 Gila Valley Irrigation District
 Maricopa County Municipal Water Conservation District
 No. 1
 Navajo Tribal Council
 Salt River Valley Water Users Association
 San Carlos Irrigation and Drainage District
 State Land Department
Arkansas:
 Arkansas Geological Commission
 Arkansas State Highway Commission
 University of Arkansas—Agricultural Experiment Station
California:
 Alameda County Flood Control and Water Conservation District
 Alameda County Water District
 Calaveras County Water District
 City of San Diego
 City of Santa Barbara
 Contra Costa County Flood Control and Water Conservation District
 County of Los Angeles Department of County Engineers
 Department of Conservation, Division of Mines and Geology
 East Bay Municipal Utility District
 East Kern Water Agency—Antelope Valley
 Georgetown Divide Public Utility District

California—Continued

Imperial Irrigation District
 Lake County Flood Control and Water Conservation District
 Metropolitan Water District of Southern California
 Montecito County Water District
 Monterey County Flood Control and Water Conservation District
 Orange County Flood Control District
 Orange County Water District
 Palo Verde Irrigation District
 Placer County Water Agency
 Sacramento Municipal Utility District
 San Bernardino County Flood Control District
 San Bernardino Valley Water Conservation District
 San Francisco City and County Public Utilities Commission
 San Francisco Water Department
 San Luis Obispo County Flood Control and Water Conservation District
 San Mateo County
 Santa Barbara County Water Agency
 Santa Clara County Flood Control and Water Conservation District
 Santa Cruz County Flood Control and Water Conservation District
 Santa Maria Valley Water Conservation District
 State Department of Fish and Game
 State Department of Water Resources
 State Water Pollution Control Board
 Ventura River Municipal Water District

Colorado:

Arkansas River Compact Administration
 City of Westminster
 Colorado River Water Conservation Board
 Colorado State Metal Mining Fund Board
 Colorado State University Agricultural Experiment Station
 Colorado Water Conservation Board
 Costilla Creek Compact Commission
 Denver Board of Water Commissioners
 Inter-county Regional Planning Commission
 Office of State Engineer, Division of Water Resources
 Rio Grande Compact Commission
 Southeastern Colorado Water Conservancy District

Connecticut:

City of Torrington—Engineering Department
 Connecticut Geologic and Natural History Survey
 Greater Hartford Flood Commission
 Hartford Department of Public Works
 Highway Department
 New Britain Board of Water Commissioners
 State Water Resources Commission

Delaware:

Delaware Geological Survey
 State Highway Department

District of Columbia:

District of Columbia Department of Sanitary Engineering

Florida:

Broward County—Board of County Commissioners
 Central and Southern Florida Flood Control District
 City of Boca Raton

Florida—Continued

City of Deerfield Beach
 City of Fort Lauderdale
 City of Jacksonville:
 City Commission
 Office of the City Engineer
 City of Miami—Department of Water and Sewers
 City of Miami Beach
 City of Naples
 City of Perry
 City of Pompano Beach
 City of Tallahassee
 Collier County—Board of County Commissioners
 Dade County—Board of County Commissioners
 Florida Geological Survey
 Hillsborough County—Board of County Commissioners
 Orange County—Board of County Commissioners
 Pinellas County—Board of County Commissioners
 Polk County—Board of County Commissioners
 State Board of Parks and Historic Memorials
 State Road Department
 Trustees of Internal Improvement Fund

Georgia:

Department of Mines, Mining and Geology, Division of Conservation
 State Highway Department

Hawaii:

City and County of Honolulu
 State Department of Land and Natural Resources

Idaho:

Idaho Department of Fish and Game
 Idaho Department of Highways
 Idaho Department of Reclamation

Illinois:

Division of Highways
 Division of Waterways
 Fountain Head Drainage District
 Metropolitan Sanitary District of Greater Chicago
 Northeastern Illinois Metropolitan Area Planning Commission
 Sanitary District of Bloom Township (Cook County)
 State Department of Public Works and Buildings
 State Department of Registration and Education

Indiana:

Department of Conservation
 Department of Natural Resources
 Flood Control and Water Resources Commission
 State Board of Health
 State Highway Commission

Iowa:

Agricultural Experiment Station, Iowa State University of Science and Technology
 City of Cedar Rapids
 City of Fort Dodge—Department of Utilities
 City of Iowa City
 City of Muscatine
 Iowa Geological Survey
 Iowa Institute of Hydraulic Research
 Iowa State Conservation Commission
 Iowa State Highway Commission
 Iowa State University
 Linn County—Board of Supervisors

Kansas:

Board of Agriculture, Division of Water Resources

Kansas—Continued

City of Wichita—Department of Public works
 Kansas State Geological Survey
 State Department of Health—Environment Health Services
 State Highway Commission
 State Water Resources Board

Kentucky:

Kentucky Geological Survey
 University of Kentucky

Louisiana:

Sabine River Compact Commission
 State Department of Conservation
 State Department of Highways
 State Department of Public Works

Maine:

Maine Public Utilities Commission
 State Highway Commission

Maryland:

Baltimore County—Department of Public Works
 Charles County
 City of Baltimore
 City of Cambridge
 City of Easton
 City of Salisbury
 Dorchester County—Board of County Commissioners
 Maryland Geological Survey
 State Board of Health
 State Planning Department
 State Roads Commission
 Talbot County—Board of County Commissioners
 Washington Suburban Sanitary Commission

Massachusetts:

Boston Metropolitan District Commission
 Massachusetts Department of Public Works:
 Division of Highways
 Division of Waterways
 Massachusetts Water Resources Commission

Michigan:

State Department of Conservation:
 Engineering and Architecture Section
 Fish Section
 Game Section
 Geological Survey Section
 Parks and Recreation Section
 State Highway Department
 State Water Resources Commission

Minnesota:

Board of County Commissioners of Hennepin County
 Department of Administration
 Department of Iron Range Resources and Rehabilitation
 State Department of Conservation, Division of Waters
 State Department of Highways

Mississippi:

City of Jackson
 Harrison County—Board of Supervisors and Development Commission
 Jackson County—Port Authority
 Mississippi Board of Water Commissioners
 Mississippi Research and Development Center
 Mississippi State Highway Department
 Pearl River Valley Water Supply District

Missouri:

Curators of the University of Missouri

Missouri—Continued

Division of Geological Survey and Water Resources
 State Department of Conservation:
 Conservation Commission
 Fisheries Division
 State Highway Commission
 Water Pollution Board

Montana:

Montana Bureau of Mines and Geology
 Montana State College—Endowment and Research Foundation
 State Engineer
 State Fish and Game Commission
 State Highway Commission
 State Water Conservation Board

Nebraska:

Department of Roads
 Department of Water Resources
 Nebraska Mid-State Reclamation District
 University of Nebraska—Conservation and Survey Division

Nevada:

Department of Conservation and Natural Resources
 Department of Highways
 Nevada Bureau of Mines

New Hampshire:

New Hampshire Water Resources Board

New Jersey:

Camden County Planning Board
 County of Bergen
 Delaware River Basin Commission
 Department of Agriculture
 Department of Conservation and Economic Development:
 Division of Fish and Game
 Division of Water Policy and Supply
 Department of Health
 North Jersey District Water Supply Commission
 Passaic Valley Water Commission
 Rutgers University, The State University of New Jersey

New Mexico:

Carlsbad Irrigation District
 Costilla Creek Compact Commission
 Interstate Stream Commission
 New Mexico Institute of Mining and Technology
 Pecos River Commission
 Pecos Valley Artesian Conservancy District
 Rio Grande Compact Commission
 State Engineer and School of Mines
 State Highway Department
 State Game and Fish Commission
 University of New Mexico

New York:

Board of Hudson River—Black River Regulating District
 Brighton Sewer District No. 2
 City of Albany—Department of Water and Water Supply
 City of Auburn—Water Department
 Dutchess County Board Supervisors
 Nassau County Department of Public Works
 New York City Board of Water Supply
 Office of Atomic and Space Development
 Onondaga County Department of Public Works
 Onondaga County Water Authority
 Oswegatchie River—Cranberry Reservoir Commission

New York—Continued

State Conservation Department:
 Division of Lands and Forests
 Division of Water Resources
 State Department of Commerce
 State Department of Health, Water Pollution Control Board
 State Department of Public Works
 Suffolk County Board of Supervisors
 Suffolk County Department of Public Works
 Suffolk County Water Authority
 Village of Nyack—Board of Water Commissioners
 Westchester County Department of Public Works

North Carolina:

City of Asheville
 City of Burlington
 City of Charlotte
 City of Durham
 City of Greensboro
 City of Winston-Salem
 North Carolina Department of Conservation and Development, Division of Mineral Resources
 Pitt County Board of Commissioners
 State Department of Water Resources
 State Highway Commission
 Town of Waynesville

North Dakota:

North Dakota Geological Survey
 Oliver County
 State Highway Department
 State Water Conservation Commission

Ohio:

City of Columbus—Department of Public Service
 Miami Conservancy District
 Ohio Department of Health
 Ohio Department of Highways
 Ohio Department of Natural Resources—Division of Water
 Ohio River Valley Water Sanitation Commission
 Scioto Conservancy District

Oklahoma:

City of Oklahoma City Water Department
 Oklahoma Department of Highways
 Oklahoma Geological Survey
 Oklahoma State Department of Health
 Oklahoma Water Resource Board

Oregon:

Board of Higher Education
 Burnt River Irrigation District
 City of Dallas
 City of Dalles City
 City of Eugene—Water and Electric Board
 City of McMinnville—Water and Light Department
 City of Portland—Bureau of Water Works
 City of Toledo
 Coos Bay—North Bed Water Board
 County Court of Douglas County
 County Court of Lane County
 County Court of Morrow County
 County of Coos—Board of Commissioners
 Mosier Irrigation District
 State Engineer, Water Resources Department
 State Game Commission
 State Highway Commission

Oregon—Continued

Vale Irrigation District
 Talent Irrigation District

Pennsylvania:

Chester County Soil and Water Conservation District
 City of Bethlehem
 City of Harrisburg
 City of Philadelphia
 Conestoga Valley Association, Inc.
 Department of Internal Affairs, Bureau of Topographic and Geologic Survey
 Pennsylvania Department of Agriculture
 Pennsylvania Department of Forests and Waters
 Pennsylvania Department of Health

Rhode Island:

Department of Public Works
 Rhode Island Water Resources Coordinating Board

South Carolina:

City of Spartanburg—Commissioners of Public Works
 State Development Board
 State Highway Department
 State Public Service Authority
 State Water Pollution Control Authority

South Dakota:

South Dakota Department of Highways
 South Dakota Geological Survey
 South Dakota Water Resources Commission

Tennessee:

City of Chattanooga
 City of Murfreesboro—Water and Sewer Department
 City of Oak Ridge
 Memphis Board of Light, Gas, and Water Commissioners—Water Division
 Metropolitan Government of Nashville and Davidson County—Department of Public Works
 Tennessee Department of Conservation:
 Division of Geology
 Division of Water Resources
 Tennessee Department of Highways
 Tennessee Department of Public Health
 Tennessee Game and Fish Commission

Texas:

City of Dallas
 City of Houston
 Highway Department
 Pecos River Commission
 Rio Grande Compact Commission
 Sabina River Compact Administration
 Texas Water Commission

Utah:

Bear River Compact Commission
 Salt Lake County
 State Road Commission of Utah
 Utah Geological and Mineralogical Survey
 Utah State Engineer
 Utah Water and Power Board

Vermont:

Department of Highways
 State Water Resources Board
 Vermont Geological Survey

Virginia:

City of Alexandria
 City of Charlottesville
 City of Newport News—Department of Public Utilities

Virginia—Continued

- City of Norfolk—Division of Water Supply
- City of Roanoke
- City of Staunton
- County of Chesterfield
- County of Fairfax
- Department of Conservation and Development—Division of Mineral Resources
- Department of Highways

Washington:

- City of Seattle
- City of Tacoma:
 - Department of Public Utilities
 - Department of Public Works
- King County Board of Commissioners
- Municipality of Metropolitan Seattle
- State Department of Conservation, Division of Mines and Geology
- State Department of Fisheries
- State Department of Game
- State Department of Highways
- State Pollution Control Commission
- Stevens County

West Virginia:

- Clarksburg Water Board
- Morgantown Water Commission
- State Department of Natural Resources
- State Geological and Economic Survey

West Virginia—Continued

- State Road Commission

Wisconsin:

- Madison Metropolitan Sewerage District
- Public Service Commission of Wisconsin
- Southeastern Wisconsin Regional Planning Commission
- State Committee on Water Pollution
- State Highway Commission
- University of Wisconsin—Geological and Natural History Survey

Wyoming:

- City of Cheyenne—Board of Public Utilities
- Geological Survey of Wyoming
- State Engineer
- Wyoming Highway Department
- Wyoming Natural Resource Board

Commonwealth of Puerto Rico:

- Aqueduct and Sewer Authority
- Department of Agriculture
- Department of Public Works
- Economic Development Administration
- Water Resources Authority

American Samoa:

- Government of American Samoa

Guam:

- Government of Guam

Virgin Islands:

- Government of the Virgin Islands

U.S. GEOLOGICAL SURVEY OFFICES

MAIN CENTERS

Main Office: General Services Building, 18th and F Streets NW., Washington, D.C. 20242; 343-1100

Rocky Mountain Center: Federal Center, Denver, Colo. 80225; BElmont 3-3611

Pacific Coast Center: 345 Middlefield Road, Menlo Park, Calif. 94025; DAvenport 5-6761

PUBLIC INQUIRIES OFFICES

<i>Location</i>	<i>Official in charge and telephone number</i>	<i>Address</i>
Alaska, Anchorage, 99501	Margaret I. Erwin (277-0577)	108 Skyline Bldg., 508 2d Ave.
California, Los Angeles, 90012	Lucy E. Birdsall (688-2850)	7638 Federal Bldg., 300 N. Los Angeles St.
San Francisco, 94111	Jean V. Molleskog (556-5627)	504 Custom House, 555 Battery St.
Colorado, Denver, 80202	Lorene C. Young (297-4169)	15426 Federal Bldg., 1961 Stout St.
Texas, Dallas, 75202	Mary E. Reid (Riverside 9-3230, ext. 3230).	602 Thomas Bldg., 1314 Wood St.
Utah, Salt Lake City, 84111	Maurine Clifford (524-5652)	8102 Federal Bldg., 125 South State St.
Washington, Spokane, 99204	Eva M. Raymond (TEmples 8-3361, ext. 121).	South 157 Howard St.

SELECTED FIELD OFFICES IN THE UNITED STATES AND PUERTO RICO

[Temporary offices not included; list current as of September 1, 1965. Correspondence to the following offices should be addressed to the Post Office Box, if one is given]

CONSERVATION DIVISION

<i>Location</i>	<i>Official in charge* and telephone number</i>	<i>Address</i>
Alaska, Anchorage, 99501	Leo H. Saarela (m) (277-0578), Alexander A. Wanek (c) (277-0570), W. J. Linton (o) (277-0570).	P.O. Box 259; Skyline Bldg., 218 E. St.
California, Los Angeles, 90012	Russell G. Wayland (c) (688-2846), D. W. Solanas (o) (688-2846).	7744 Federal Bldg., 300 N. Los Angeles St.
Sacramento, 95814	Richard N. Doolittle (w) (449-2203)	8030 Federal Bldg., 650 Capitol Ave.
Bakersfield, 93301	Harry Lee Wolf (o) (327-7274), E. E. Richardson (c) (327-7274).	309 Federal Bldg., 800 Truxtun Ave.
Colorado, Denver, 80202	G. G. Frazier (o) (297-3211), H. B. Lindeman (m) (297-4038).	15428 and 15444 Federal Bldg.
Denver, 80225	George H. Horn (c) (233-3611, ext. 8168)	Building 25, Federal Center.
Denver, 80202	W. C. Senkpiel (w) (297-3316)	15407 Federal Bldg., 1961 Stout St.
Durango, 81302	Jerry W. Long (o) (247-5144)	P.O. Box 1809; Jarvis Bldg., 125 W. 10th St.
Louisiana, New Orleans, 70113	Admiral D. Acuff (o) (527-6543)	T-6009 Federal Bldg., 701 Loyola Ave.
Lafayette, 70504	George Kinsel (o) (232-0239)	301 Federal Bldg.
Montana, Billings, 59103	Ray M. Bottomley (m) (245-6711, ext. 6368), Hillary A. Oden (o) (245-6711).	P.O. Boxes 1435 and 2550; 223 and 327 Federal Bldg.
Great Falls, 59401	Andrew F. Bateman (c) (454-3314), John A. Fraher (o) (454-3336).	P.O. Boxes 1215 and 2265; 510 First Ave.
New Mexico, Artesia, 88210	James A. Knauf (o) (746-4841)	Drawer U; 210 Carper Bldg., 105 S. 4th St.
Carlsbad, 88220	Robert S. Fulton (m) and Bruno R. Alto (c) (885-6454).	P.O. Box 1716; Federal Bldg., Fox and Haleguena St.
Farmington, 87401	Phillip T. McGrath (o) and J. E. Fassett (c) (325-4572).	P.O. Box 959; 409 Petroleum Club Plaza, 3535 East 30th St.

* The small letter in parentheses following each official's name denotes branch affiliation in the Conservation Division as follows: b—Branch of Connally Act Compliance, c—Branch of Mineral Classification, m—Branch of Mining Operations, o—Branch of Oil and Gas Operations, w—Branch of Waterpower Classification.

<i>Location</i>	<i>Official in charge* and telephone number</i>	<i>Address</i>
New Mexico—Continued		
Hobbs, 88240	Arthur R. Brown (o) (393-3612)	Box 1157; 205 N. Linam St.
Roswell, 88201	J. A. Anderson (o) and T. F. Stipp (c) (622-1332).	Drawer 1857; Farnsworth Bldg., 120 W. 2nd St.
Oklahoma, Holdenville, 74848	Gerhardt H. W. Schuster (o) (379-3840)	P.O. Box 789; 5 Federal Bldg.
McAlester, 74502	A. M. Dinsmore (m) (423-5030)	509 South 3rd St.
Miami, 74354	Andrew V. Bailey (m) (542-9481)	P.O. Box 509; 205 Federal Bldg.
Oklahoma City, 73102	Charley W. Nease (o) (236-2311)	4321 Federal Court House and Office Bldg., 220 N.W. 4th St.
Tulsa, 74103	Edward L. Johnson (c) and N. Orvis Frederick (o) (584-7161, ext. 638 and 632).	521 Wright Bldg., 115 W. 3rd St.
Oregon, Portland, 97208	Loyd L. Young (w) (226-3252)	P.O. Box 3087; 319 Post Office Bldg.
Texas, Kilgore, 75662	Warren W. Mankin (b) (5564)	P.O. Box 1230, Rader Bldg., 901-903 Broadway Blvd.
Midland, 79701	Everett H. Patterson (b) (684-6741)	P.O. Box 1830; 132 Western American Bldg., 305 N. Big Spring St.
Utah, Salt Lake City, 84111	Ernest Blessing (m) (524-5646), Harry McAndrews (c) (524-5643), Rodney A. Smith (o) (524-5650).	8402, 8422, and 8416 Federal Bldg.; 125 S. State St.
Washington, Tacoma, 98401	Gordon C. Giles (w) (383-5380)	P.O. Box 1152; 244 Federal Bldg.
Wyoming, Casper, 82602	J. R. Schwabrow (o) and Donald M. Van Sickle (c) (265-4310).	P.O. Box 400; 305 Federal Bldg.
Newcastle, 82701	Glenn E. Worden (o) (746-4554)	P.O. Box 219; 611 S. Summit St.
Rock Springs, 82901	John Duletsky (o) (362-6422), Arne A. Mattila (m) (362-7350).	P.O. Box 1170; 201 and 204 First Security Bank Bldg., 502 S. Front St.
Thermopolis, 82443	Charles P. Clifford (o) (864-3477)	P.O. Box 590; 202 Federal Bldg.

* See footnote, p. A211.

GEOLOGIC DIVISION

<i>Location</i>	<i>Geologist in charge and telephone number</i>	<i>Address</i>
Alaska, College, 99735	Robert M. Chapman (479-6725)	P.O. Box 580; Brooks Memorial Bldg.
Arizona, Flagstaff, 86002	Eugene M. Shoemaker (774-5081)	601 East Cedar Ave.
California, Los Angeles, 90424	Robert O. Castle (GRanite 3-0971, ext. 9881).	Geology Bldg., Univ. of California.
Hawaii, Hawaii National Park, 96718	Howard A. Powers (678-485)	Hawaiian Volcano Observatory.
Kansas, Lawrence, 66044	Windsor L. Adkison (Viking 3-2700)	c/o State Geological Survey, Lindley Hall, Univ. of Kansas.
Kentucky, Lexington, 40503	Paul W. Richards (4-2473)	496 Southland Drive.
Maryland, Beltsville, 20705	James C. Wright (GRanite 4-4800, ext. 470).	U.S. Geological Survey Bldg., Dept. of Agriculture Research Center.
Massachusetts, Boston, 02116	Lincoln R. Page (KENmore 6-1444)	Room 1, 270 Dartmouth St.
Michigan, Marquette, 49855	Jacob E. Gair (226-2110)	Industrial Lane.
New Mexico, Albuquerque, 87100	Charles B. Read (CHapel 7-0311, ext. 483).	P.O. Box 4083, Station A; Geology Bldg., Univ. of New Mexico.
Ohio, Columbus, 43200	James M. Schopf (AXminster 4-1810)	Orton Hall, Ohio State Univ., 155 Oval Drive.
Puerto Rico, Roosevelt, 00927	Watson H. Monroe (San Juan 6-5340)	P.O. Box 803.
Tennessee, Knoxville, 37906	Robert A. Laurence (2-7787)	13 Post Office Bldg.
Texas, Austin, 78701	D. Hoye Eargle (HObart 5-6501)	801 Federal Bldg.
Utah, Salt Lake City, 84111	Lowell S. Hilpert (524-5640)	8426 Federal Bldg.
Washington, Spokane, 99204	Albert E. Weissenborn (TEmples 8-2084)	South 157 Howard St.
Wisconsin, Madison, 53706	Carl E. Dutton (262-1854)	222 Science Hall, Univ. of Wisconsin.
Wyoming, Laramie, 82070	J. David Love (FRanklin 5-4495)	Geology Hall, Univ. of Wyoming.

TOPOGRAPHIC DIVISION

<i>Location</i>	<i>Engineer in charge and telephone number</i>	<i>Address</i>
California, Menlo Park, 94025	Robert O. Davis (415 325-6761, ext. 411)	345 Middlefield Rd.
Colorado, Denver, 80225	Roland H. Moore (303 233-3611, ext. 8551)	Federal Center, Bldg. 25.
Missouri, Rolla, 65401	Daniel Kennedy (314 364-3680)	P.O. Box 133; 9th and Elm Sts.
Virginia, Arlington, 22201	Charles F. Fuechsel (703 JAcson 5-7550)	1109 N. Highland St.

WATER RESOURCES DIVISION**AREA OFFICES**

<i>Location</i>	<i>Official in charge* and telephone number</i>	<i>Address</i>
Atlantic Coast Area:		
Arlington, Virginia, 20242 -----	George E. Ferguson, Area Hydrologist (202 343-4840).	George Washington Bldg., Arlington Towers, 1011 Arlington Blvd.
Mid-Continent Area:		
St. Louis, Missouri, 63103 -----	Harry D. Wilson, Jr., Area Hydrologist (314 622-4361).	1252 Federal Bldg., 1520 Market St.
Rocky Mountain Area:		
Denver, Colorado, 80225 -----	Sherman K. Jackson, Area Hydrologist (303 233-3611).	Federal Center, Bldg. 25.
Pacific Coast Area:		
Menlo Park, California, 94025 -----	Warren W. Hastings, Area Hydrologist (415 325-6761).	345 Middlefield Road.

DISTRICT OFFICES

Alabama, University, 35486 -----	William L. Broadhurst (w) (205 752-8105).	P.O. Box V; Oil and Gas Board Bldg., Univ. of Alabama.
Alaska, Anchorage, 99501 -----	Harry Hulsing (w) (907 277-5526, 5527)	P.O. Box 2480; 218 "E" St., Skyline Bldg.
Arizona, Tucson, 85717 -----	Horace M. Babcock (w) (602 623-7731) (ext. 5791).	P.O. Box 4070; Geology Bldg., Univ. of Arizona Campus.
Arkansas, Little Rock, 72201 -----	Richard T. Sniegocki (w) (501 372-5246)	2301 Federal Office Bldg., 700 West Capitol Ave.
California, Menlo Park, 94025 -----	Walter Hofmann (w) (415 325-6761)	345 Middlefield Road.
Colorado, Denver, 80215 -----	John W. Odell (s) (303 233-3611, ext. 644).	Rm. 22, 1455 Ammons St.
Denver, 80225 -----	Leonard A. Wood (g) (303 233-3611, ext. 8546).	Federal Center, Bldg. 25.
Connecticut, Hartford, 06101 -----	John Horton (s) (203 244-2528)	P.O. Box 715; 203 Federal Bldg.
Middletown, 06458 -----	John A. Baker (g) (203 346-5542)	204 Post Office Bldg.
Delaware, Dover, 19901 -----	Philip Pfannebecker (s) (302 734-2506)	604 Fairview Ave.
Florida, Tallahassee, 32304 -----	Clyde S. Conover (w) (305 377-4231)	P.O. Box 2315; Gunter Bldg. (Tennessee and Woodward Sts.)
Georgia, Atlanta, 30323 -----	Albert N. Cameron (w) (404 876-3311)	Rm. 164, Peachtree Seventh Bldg.
Hawaii, Honolulu, 96814 -----	Dan A. Davis (g) and Mearle M. Miller (s) (588-111, ext. 692, 693, 694).	332 and 330 First Insurance Bldg., 1100 Ward Ave.
Idaho, Boise, 83702 -----	Wayne I. Travis (s) (208 342-2711, ext. 531), Herbert A. Waite (g) (208 342- 2711, ext. 539).	Rms. 215 and 205, 914 Jefferson St.
Illinois, Champaign, 61820 -----	William D. Mitchell (s) (217 356-5221)	605 South Neil St.
Indiana, Indianapolis, 46204 -----	Malcom D. Hale (w) (317 633-7398)	Rm. 516, 611 North Park Ave.
Iowa, Iowa City, 52241 -----	Sulo W. Wiitala (s) (319 338-0581, ext. 475).	508 Hydraulic Laboratory
Iowa City, 52240 -----	Walter L. Steinhilber (g) (319 338-1173)	Geological Survey Bldg.
Kansas, Lawrence, 66045 -----	Robert J. Dingman (g) (913 864-3001)	c/o Univ. of Kansas.
Topeka, 66601 -----	Edward J. Kennedy (s) (913 234-8661, ext. 201).	P.O. Box 856; 403 Federal Bldg.
Kentucky, Louisville, 40202 -----	Robert V. Cushman (g) and Floyd F. Schrader (s) (502 582-5241, 5242, 5243).	310 Center Bldg. 522 West Jefferson St.
Louisiana, Baton Rouge, 70806 -----	Mack R. Stewart (s) and Russell M. McAvoy (q) (504 348-7181, ext. 281).	215 and 201 Prudential Bldg., 6554 Flor- ida Blvd.
Baton Rouge, 70803 -----	Rex R. Meyer (g) (504 348-7181, ext. 224).	P.O. Box GS, University Station; Rm. 223, Field House, Louisiana State Univ.
Maine, Augusta, 04330 -----	Gordon S. Hayes (s) and Glenn C. Pres- cott (g) (207 623-4511, ext. 708).	Vickery Hill Bldg., Court St.
Maryland, Towson, 21204 -----	John W. Wark (w) (301 828-7460)	724 York Road.

* The small letter in parentheses following each official's name signifies his affiliation in the Water Resources Division, as follows: g—Ground Water Branch; q—Quality of Water Branch; s—Surface Water Branch; w—Water Resources Division.

<i>Location</i>	<i>DISTRICT OFFICES—Continued</i> <i>Official in charge* and telephone number</i>	<i>Address</i>
Massachusetts, Boston, 02110	Charles E. Knox (w) (617 223-2824)	Rm. 205, 211 Congress St.
Michigan, Lansing, 48933	Arlington D. Ash (w) (517 372-1910, ext. 561 and 564).	700 Capitol Savings and Loan Bldg.
Minnesota, St. Paul, 55101	David B. Anderson (s) and Richmond F. Brown (g) (612 228-7842).	1610 and 1002 New Post Office Bldg.
Mississippi, Jackson, 39205	William H. Robinson (w) (601 354-2326, 2327).	P.O. Box 2052, 302 U.S. Post Office Bldg.
Missouri, Rolla, 65401	Anthony Homyk, Jr. (s) (314 364-1599) Edward J. Harvey (g) (314 364-1599).	P.O. Box 340; 103 W. 10th St.
Montana, Billings, 59601	Charles W. Lane (g) (406 259-2412)	P.O. Box 1818; Bell Bldg., 2 South 7th St., West.
Helena, 59601	Frank Stermitz (s) (406 442-9040, ext. 3263).	P.O. Box 1696; 409 Federal Bldg.
Nebraska, Lincoln, 68501	Kenneth A. MacKichan (w) (402 475-3326).	P.O. Box 591; 125 Nebraska Hall, 901 N. 17th St.
Nevada, Carson City, 89701	George F. Worts, Jr. (w) (702 882-1388)	222 E. Washington St.
New Jersey, Trenton, 08605	John E. McCall (s) (609 599-3511, ext. 214).	P.O. Box 967; 433 Federal Bldg.
Trenton, 08607	Allen Sinnott (g) (609 599-3511, ext. 213).	P.O. Box 1238; 432 Federal Bldg.
New Mexico, Albuquerque, 87106	Samuel W. West (g) and Jay M. Stow (q) (505 247-0311, ext. 2248 and 2249).	P.O. Box 4217; Geology Bldg., Univ. of New Mexico.
Santa Fe, 87501	Wilbur L. Heckler (s) (505 982-3801, ext. 3307).	P.O. Box 1750; Federal Bldg., Cathedral Place.
New York, Albany, 12201	Ralph C. Heath (w) (518 472-3107)	P.O. Box 948; 342 Federal Bldg.
North Carolina, Raleigh, 27602	Edward B. Rice (w) (919 828-9031, ext. 156).	P.O. Box 2857; 4th Floor, Federal Bldg.
North Dakota, Bismarck, 58502	Harlan M. Erskine (s) (701 255-4011, ext. 227), Delbert W. Brown (g) (701 255-4011, ext. 228).	P.O. Box 778; 348 New Federal Bldg., 3rd St. and Rosser Ave.
Ohio, Columbus, 43212	John J. Molloy (w) (614 221-6411, ext. 113).	85 Marconi Blvd.
Oklahoma, Oklahoma City, 73102	Alvin R. Leonard (g) (405 236-2412), Alexander A. Fischback, Jr. (s) (405 236-2257, 2258).	4011 and 4301 Federal Bldg., 200 North-west 4th St.
Oklahoma City, 73109	Richard P. Orth (q) (405 677-5022)	P.O. Box 95205; 2800 S. Eastern.
Oregon, Portland, 97208	Roy B. Sanderson (s) and Eugene R. Hampton (g) (503 226-3361, ext. 1248 and 1246).	P.O. Box 3418; 419 and 415 Federal Bldg., 701 NW. Glisan St.
	Lawrence Bodhaine (q) (503 234-3361, ext. 241).	P.O. Box 3202; 830 NE. Holladay.
Pennsylvania, Harrisburg, 17104	Joseph E. Barclay (g) (717 787-3420)	100 North Cameron St.
	Robert E. Steacy (s) (717 787-3917)	1224 Mulberry St.
Philadelphia, 19106	Norman H. Beamer (q) (215 597-4420)	1302 U.S. Custom House, 2d and Chestnut Streets.
Puerto Rico, Hato Rey, 00918	Dean B. Bogart (w) (809 766-3310)	12 Arroyo St.
Rhode Island, Providence, 02903	Joseph C. Rosenshein (g) (401 331-9312)	401-2 Federal Bldg. and U.S. Post Office.
South Carolina, Columbia, 29201	Albert E. Johnson (s) (803 253-8371, ext. 401).	121 Veterans Administration Regional Office Bldg., 1801 Assembly St.
Columbia, 29205	George E. Siple (g) (803 253-3405)	P.O. Box 5314; 627 Bull St.
South Dakota, Huron, 57350	John E. Powell (g) (605 352-8651, ext. 293).	P.O. Box 1412; 231 Federal Bldg.
Pierre, 57501	John E. Wagar (s) (605 224-7856)	P.O. Box 216; 207 Federal Bldg.
Tennessee, Nashville, 37203	Joseph S. Cragwall, Jr. (w) (615 242-8321, ext. 5924).	144 Federal Office Bldg.
Texas, Austin, 78701	Trigg Twichell (w) (512 476-6411)	Federal Bldg., 300 E. 8th Ave.
Utah, Salt Lake City, 84111	Milton T. Wilson (w) (801 524-5663)	8002 Federal Bldg., 125 S. State St.
Virginia, Charlottesville, 22903	James W. Gambrell (s) (703 296-5191 ext. 321).	P.O. Box 3327, University Station; Natural Resources Bldg., McCormick Road.
Washington, Tacoma, 98402	Leslie B. Laird (w) (206 383-2861, ext. 384).	Rm. 300, 1305 Tacoma Ave., So.

* See footnote, p. A213.

DISTRICT OFFICES—Continued

<i>Location</i>	<i>Official in charge* and telephone number</i>	<i>Address</i>
West Virginia, Charleston, 25301	William C. Griffin (s) (304 343-6181, ext. 311).	3303 New Federal Office Bldg. and U.S. Court House; 500 Quarrier St. East.
Morgantown, 26506	Porter E. Ward (g) (304 296-3338)	405 Mineral Industries Bldg., Univ. of West Virginia.
Wisconsin, Madison, 53706	Charles L. R. Holt, Jr. (g) (608 262-2488).	175 Science Hall, Univ. of Wisconsin.
Madison, 53705	Kenneth B. Young (s) (608 233-0195)	5001 University Ave.
Wyoming, Cheyenne, 82002	Ellis D. Gordon (g) (307 634-5920, ext. 2331), Leon A. Wiard (s) (307 634-5920, ext. 2317).	P.O. Box 2087; 2129 E. 8th Ave.
Worland, 82401	Thomas F. Hanley, (q) (307 347-2181)	1214 Big Horn Ave.

* See footnote, p. A213.

OFFICES IN OTHER COUNTRIES**GEOLOGIC DIVISION**

<i>Location</i>	<i>Official in charge</i>	<i>Address</i>
Bolivia, La Paz	James F. Seitz	U.S. Geological Survey, U.S. AID/Bolivia, c/o American Embassy, La Paz, Bolivia.
Brazil, Recife	John J. Matzko	U.S. AID/Recife, U.S. Department of State, Washington, D.C. 20521. STOP 27.
Rio de Janeiro	Alfred J. Bodenlos	U.S. Geological Survey, U.S. AID/RIO, c/o American Embassy, APO New York 09676.
Colombia, Barranquilla	Charles M. Tschanz	U.S. Geological Survey, c/o American Consul, American Consulate, Barranquilla, Colombia.
Bogotá	Earl M. Irving	U.S. Geological Survey, U.S. AID/UID, c/o American Embassy, Bogotá, Colombia.
Medellin	Lawrence V. Blade	U.S. Geological Survey, c/o American Consulate, Medellin, Colombia.
Costa Rica, San José	Frank D. Spencer	U.S. Geological Survey, c/o American Embassy, San José, Costa Rica.
Dahomey, Cotonou	Jules A. MacKallor	U.S. Geological Survey, U.S. AID/Cotonou, U.S. Department of State, Washington, D.C. 20521, STOP 27.
Liberia, Monrovia	Darwin L. Rossman	U.S. Geological Survey, U.S. AID/Monrovia, U.S. Department of State, Washington, D.C. 20521, STOP 27.
Pakistan, Quetta	Max G. White	U.S. Geological Survey, U.S. AID, c/o American Embassy, APO New York 09271.
Philippines, Manila	Laurence E. Andrews, Jr.	U.S. Geological Survey, U.S. AID, c/o American Embassy, APO San Francisco 96528.
Saudi Arabia, Jidda	Glen F. Brown	U.S. Geological Survey, c/o American Embassy, APO New York 09697.
Thailand, Bangkok	Charles T. Pierson	U.S. Geological Survey, USOM, c/o American Embassy, APO San Francisco 96346.

WATER RESOURCES DIVISION

<i>Location</i>	<i>Official in charge</i>	<i>Address</i>
Afghanistan, Bost	Vito J. Latkovich	U.S. Geological Survey, U.S. AID/Bost, c/o American Embassy, APO New York 09668.
Kabul	Arthur O. Westfall	U.S. Geological Survey, U.S. AID/Kabul, c/o American Embassy, APO New York 09668.

U.S. GEOLOGICAL SURVEY OFFICES

<i>Location</i>	<i>Official in charge</i>	<i>Address</i>
Brazil, Recife -----	Stuart L. Schoff -----	U.S. AID/Recife, U.S. Department of State, Washington, D.C. 20521, STOP 27.
Egypt, Cairo -----	Robert L. Cushman -----	U.S. Geological Survey, U.S. AID/Cairo, U.S. Department of State, Washington, D.C. 20521, STOP 27.
Nepal, Katmandu -----	Woodrow W. Evett -----	U.S. Geological Survey, U.S. AID, c/o American Embassy, APO New York 09674.
Nigeria, Kaduna -----	David A. Phoenix -----	U.S. Geological Survey, c/o Geological Survey of Nigeria, Kaduna South, Northern Nigeria.
Kano -----	Billy E. Colson -----	U.S. Geological Survey, U.S. AID Mission to Nigeria, c/o American Embassy, Lagos, Nigeria.
Maiduguri -----	G. C. Tibbitts -----	U.S. Geological Survey, c/o Geological Survey of Nigeria, Maiduguri, Northern Region, Nigeria.
Sokoto -----	Henry R. Anderson -----	U.S. Geological Survey, U.S. AID/GSN, P.O. Box 93, Sokoto, Northern Nigeria.
Pakistan, Lahore -----	Maurice J. Mundorff -----	U.S. Geological Survey, U.S. AID/L, c/o American Embassy, APO New York 09271.

INVESTIGATIONS IN PROGRESS IN THE GEOLOGIC, WATER RESOURCES, AND CONSERVATION DIVISIONS

Investigations in progress at the end of fiscal year 1965 are listed below, together with the names and headquarters of the individuals in charge of each. Headquarters at main centers are indicated by (W) for Washington, D.C., (D) for Denver, Colo., and (M) for Menlo Park, Calif.; headquarters in other cities are indicated by name (see list of offices, p. A211 for addresses). Inquiries regarding projects for which no address is given in the list of offices should be directed to the appropriate Division of the Geological Survey, Washington, D.C. 20242. The lowercase letter following the name of the project leader shows the Division technical responsibility: c, Conservation Division; w, Water Resources Division (g, Ground Water Branch; s, Surface Water Branch; q, Quality of Water Branch; h, General Hydrology Branch); no letter, Geologic Division.

The projects are classified by principal topic. Most geologic-mapping projects involve special studies of stratigraphy, petrology, geologic structure, or mineral deposits, but are listed only under "Geologic Mapping" unless a special topic of commodity is the primary justification for the project. A reader interested in investigations of volcanology, for example, should look under the heading "Geologic Mapping" for projects in areas of volcanic rocks, as well as under the heading "Volcanology." Likewise, most water-resources investigations involve special studies of several aspects of hydrology and geology, but are listed only under "Water Resources" unless a special topic—such as floods or sedimentation—is the primary justification for the project.

Areal geologic mapping is subdivided into mapping at scales smaller than 1 inch to 1 mile (for example, 1:250,000), and mapping at scales of 1 inch to 1 mile, or large (for example, 1:62,500; 1:24,000).

Abstracts. See Bibliographies and abstracts.

Analytical chemistry:

- Analytical methods—water chemistry (M. W. Skougstad, q, D)
- Analytical services and research (I. May, W; L. F. Rader, Jr., D; R. E. Stevens, M)
- Electron probe (R. H. Heidel, M)
- Organic geochemistry and infrared analysis (I. A. Bregger, W)
- Organic substances—pesticides—in water (W. L. Lamar, w, M)
- Physical chemistry of radioelements (K. W. Edwards, q, D)
- Radioactivation and radiochemistry (F. E. Senftle, W; H. T. Millard, D)
- Rock and mineral chemical analysis (J. J. Fahey, W)
- Rock chemical analysis:
 - General (L. C. Peck, D)
 - Rapid (L. Shapiro, W)
- Trace analysis methods:
 - Development (H. W. Lakin, D)
 - Research (F. N. Ward, D)
- Trace analysis service (F. N. Ward, D)

See also Spectroscopy.

Artificial recharge:

- Basalt aquifers, Oregon, Salem Heights and The Dalles (B. L. Foxworthy, g, Portland)
- Blended industrial waste, Arizona, Tucson area (W. G. Weist, w, Tucson)
- Treated sewage through an injection well, New York, Bay Park, Long Island (P. Cohen, w, Albany)
- Arkansas, Grand Prairie region (R. T. Sniegocki, w, Little Rock)

Asbestos:

- Southeastern United States, ultramafic rocks (D. M. Larrabee, W)
- Arizona, Blue House and McFadden Peak quadrangles (A. F. Shride, D)
- Vermont, north-central (W. M. Cady, D)

Barite:

- Arkansas (D. A. Brobst, D)

Base metals:

- Colorado:
 - Kokomo and Tenmile Range mining district (M. H. Bergendahl, D)
 - Wet Mountains (M. R. Brock, D)
- Montana, Philipsburg area (W. C. Prinz, W)
- Nevada, Antler Peak quadrangle (R. J. Roberts, M)
- Utah, San Francisco Mountains (D. M. Lemmon, M)

See also base-metal names.

Bauxite:

- Southeastern United States (E. F. Overstreet, W)
- Hawaii, Kauai (S. H. Patterson, W)

Beryllium:

- Western United States, volcanic and associated rocks (D. R. Shawe, D)
- Alaska, Lost River mining district (C. L. Sainsbury, D)
- Colorado:
 - Lake George district (C. C. Hawley, D)
 - Mt. Antero (W. N. Sharp, D)
- Nevada, Mt. Wheeler mine area (D. E. Lee, D)

Bibliographies and abstracts:

- Alaskan geology, index of literature (E. H. Cobb, M)
- Geochemical exploration abstracts (C. B. Davidson, D)
- Hydrology, bibliography (J. R. Randolph, w, W)
- North American geology, bibliography (M. Cooper, W)

Bibliographies and abstracts--Continued

Vanadium, geology and resources, bibliography (J. P. Ohl, D)

Borates:

Borate marshes, California, Nevada, and Oregon (W. C. Smith, M)

California:

Furnace Creek area (J. F. McAllister, M)

Searles Lake area (G. I. Smith, M)

Chromite. See Ferro-alloy metals.

Clay-water relations:

Clays, liquid movement in (H. W. Olsen, w, W)

Kaolinite, solubility (W. L. Polzer, w, M)

Clays:

Appalachia, northern part (J. W. Hosterman, W)

Florida and Georgia, Attapulgius-Thomasville fuller's earth deposits (S. H. Patterson, W)

Idaho, Greenacres quadrangle (P. L. Weis, W)

Maryland, statewide studies (M. M. Knechtel, W)

Washington:

Eastern part (J. W. Hosterman, W)

Greenacres quadrangle (P. L. Weis, W)

Coal:

Alabama, Warrior quadrangle (W. C. Culbertson, D)

Alaska:

Resources of State (F. F. Barnes, M)

Beluga-Yentna area (F. F. Barnes, M)

Bering River coal field (A. A. Wanek, c, Anchorage)

Nenana (C. Wahrhaftig, M)

Arizona, Navajo Reservation, fuels potential (R. B. O'Sullivan, D)

Arkansas:

Arkansas Basin (B. R. Haley, D)

Ft. Smith district (T. A. Hendricks, D)

California, Priest Valley SE quadrangle (E. E. Richardson, c, Bakersfield)

Colorado:

Animas River area (H. Barnes, D)

Carbondale coal field (J. R. Donnell, D)

Cerro Summit quadrangle (R. G. Dickinson, c, D)

Corral Bluffs quadrangle (P. E. Soister, c, D)

Elk Springs quadrangle (J. R. Dyni, c, D)

Hanover NW quadrangle (P. E. Soister, c, D)

Hot Sulphur Springs quadrangle (G. A. Izett, c, D)

Kremmling quadrangle (G. A. Izett, c, D)

Marble quadrangle (D. L. Gaskill, c, D)

Marcellina Mountain quadrangle (D. L. Gaskill, c, D)

Montrose 1 SE, 1 SW, and 4 NE quadrangles (R. G. Dickinson, c, D)

Oh-Be-Joyful quadrangle (D. L. Gaskill, c, D)

Placita SE quadrangle (L. H. Godwin, c, D)

Rangely 3 quadrangle (H. L. Cullins, c, D)

Trinidad coal field (R. B. Johnson, D)

Idaho, Driggs SE quadrangle (M. L. Schroeder, c, D)

Iowa, resources of State (E. R. Landis, D)

Kentucky:

Eastern part (K. J. Englund, W)

Jellico West and Ketchen quadrangles (K. J. Englund, W)

Montana:

Anaconda NW quadrangle (A. A. Wanek, c, Anchorage, Alaska)

Black Butte quadrangle (A. F. Bateman, Jr., c, Great Falls)

Gardiner SW quadrangle (G. D. Fraser, c, D)

Coal--Continued**Montana--Continued**

Girard coal field (G. E. Prichard, D)

Hardy quadrangle (K. S. Soward, c, Great Falls)

Hedstrom area (A. F. Bateman, Jr., c, Great Falls)

Jordan (30-minute) quadrangle (G. D. Mowat, c, Great Falls)

Montaqua quadrangle (E. D. Patterson, c, W)

Powder River coal fields (N. W. Bass, D)

Rocky Reef quadrangle (K. S. Soward, c, Great Falls)

Nevada, Coaldale area (R. G. Wayland, c, Los Angeles, Calif.)

New Mexico:

Animas River area (H. Barnes, D)

Fruitland Formation (J. E. Fassett, c, Farmington)

Johnson Trading Post quadrangle (J. S. Hinds, c, Farmington)

Mesa Portales quadrangle (J. E. Fassett, c, Farmington)

Raton coal basin:

Eastern part (G. H. Dixon, D)

Western part (C. L. Pillmore, D)

San Juan Basin, east side (C. H. Dane, W)

North Dakota:

Clark Butte and Clark Butte NE quadrangles (G. D. Mowat, c, Great Falls, Mont.)

Clark Butte NW and SW quadrangles (E. H. Gilmour, c, Great Falls, Mont.)

Dengate quadrangle (C. S. V. Barclay, c, D)

Glen Ullin quadrangle (C. S. V. Barclay, c, D)

Heart Butte and Heart Butte NW quadrangles (E. V. Stephens, c, D)

New Salem quadrangle (H. L. Smith, c, D)

North Almont quadrangle (H. L. Smith, c, D)

Oklahoma, Ft. Smith district (T. A. Hendricks, D)

Oregon:

Bandon SE quadrangle (R. G. Wayland, c, Los Angeles, Calif.)

Coquille SW quadrangle (R. G. Wayland, c, Los Angeles, Calif.)

Pennsylvania:

Anthraccite-mine drainage projects, geology in vicinity of (J. F. Robertson, D)

Anthraccite region, flood control (M. J. Bergin, W)

Bituminous coal resources of State (E. D. Patterson, W)

Southern anthracite field (G. H. Wood, Jr., W)

Washington County (B. H. Kent, D)

Western Middle anthracite field (H. Arndt, W)

South Dakota, Harding County and adjacent areas (G. N. Pippingos, D)

Tennessee:

Ivydell and Pioneer quadrangles (K. J. Englund, W)

Jellico West and Ketchen quadrangles (K. J. Englund, W)

Utah:

Gilbert Peak 1 NE quadrangle (J. R. Dyni, c, D)

Gunsight Butte quadrangle (F. Peterson, c, D)

Hurricane fault (southwestern Utah) (P. Averitt, D)

Jessen Butte quadrangle (J. R. Dyni, c, D)

Kaiparowits Peak 2 SE quadrangle (W. E. Bowers, c, D)

Kaiparowits Peak 4 quadrangle (H. D. Zeller, c, D)

Kolob Terrace coal field, southern (W. B. Cashion, D)

Navajo Reservation, fuels potential (R. B. O'Sullivan, D)

Coal--Continued

Utah--Continued

- Nipple Butte quadrangle (H. A. Waldrop, c, D)
- Phil Pico Mountain quadrangle (J. R. Dyni, c, D)

Virginia:

- Big Stone Gap district (R. L. Miller, W)
- Pocahontas coal beds (K. J. Englund, W)

Washington, Maple Valley, Cumberland and Hobart quadrangles (J. D. Vine, M)

Wyoming:

- Arlington quadrangle (H. J. Hyden, c, D)
- Bengough Hill quadrangle (H. J. Hyden, c, D)
- Bosler quadrangle (H. McAndrews, c, Salt Lake City, Utah)
- Ferris quadrangle (R. L. Rioux, c, W)
- Fish Lake quadrangle (W. L. Rohrer, c, D)
- Jackson (30-minute) quadrangle (D. A. Jobin, c, D)
- Kisinger Lakes quadrangle (W. L. Rohrer, c, D)
- Lake Ione quadrangle (H. McAndrews, c, Salt Lake City, Utah)
- McFadden quadrangle (H. J. Hyden, c, D)
- Oil Mountain quadrangle (W. H. Laraway, c, Casper)
- Pierce Reservoir quadrangle (H. J. Hyden, c, D)
- Pilot Knob quadrangle (W. L. Rohrer, c, D)
- Poison Spider quadrangle (W. H. Laraway, c, Casper)
- Reid Canyon (W. H. Laraway, c, Casper)
- Rock River quadrangle (H. J. Hyden, c, D)
- Square Top Butte quadrangle (W. H. Laraway, c, Casper)
- T-L Ranch quadrangle (H. J. Hyden, c, D)
- Taylor Mountain quadrangle (M. L. Schroeder, c, D)
- White Rock Canyon quadrangle (H. J. Hyden, c, D)

Construction and terrain problems:

- Deformation research (S. P. Kanizay, D)
- Lunar terrain studies (C. R. Warren, W)
- Miscellaneous site studies (D. J. Varnes, D)
- Mudflow studies (D. R. Crandell, D)
- Sino-Soviet terrain atlas (M. M. Elias, W)
- Water-resources development, potential applications of nuclear explosives (A. M. Piper, M; F. W. Stead, D)

Alaska:

- Northeastern Alaska coastal plain and foothills (C. R. Lewis, W)
- Origin and stratigraphy of ground ice in central Alaska (T. L. Péwé, College)
- Project CHARIOT (harbor construction) (G. D. Eberlein, M)
- Mt. Hayes D-3 and D-4 quadrangles (T. L. Péwé, College)
- Surficial and engineering geology:
 - Construction-materials sources (T. L. Péwé, College)
 - Anchorage-Matanuska Glacier area (T. N. V. Karlstrom, W)
 - Bristol Bay area (E. H. Muller, Ithaca, N.Y.)
 - Chitina Valley, lower part (L. A. Yehle, W)
 - Copper River basin:
 - Northeastern part (O. J. Ferrians, Jr., W)
 - Southeastern part (D. R. Nichols, W)
 - Southwestern part (J. R. Williams, W)
 - Denali Highway, eastern part (D. R. Nichols, W)
 - Johnson River district (H. L. Foster, W)
 - Kenai lowland (T. N. V. Karlstrom, W)
 - Kobuk River valley (A. T. Fernald, W)

Construction and terrain problems--Continued

Alaska--Continued

Surficial and engineering geology--Continued

- Mt. Chamberlain area (C. R. Lewis, W)
- Seward-Portage Railroad (T. N. V. Karlstrom, W)
- Slana-Tok area (H. R. Schmoll, W)
- Steese Highway area (W. E. Davies, W)
- Tanana River, upper part (A. T. Fernald, W)
- Taylor Highway area (H. L. Foster, W)
- Valdez-Tiekel belt (H. W. Coulter, W)
- Yukon-Koyukuk lowland (F. R. Weber, College)

California:

- Bodega Head reactor site (J. Schlocker, M)
- Faulting, recent (M. G. Bonilla, M)

Colorado:

- Air Force Academy (D. J. Varnes, D)
- Black Canyon of the Gunnison River (W. R. Hansen, D)
- Cheyenne Mountain, electrical properties (J. H. Scott, D)
- Gore Range (M. H. Bergendahl, D)
- Green River valley, upper part (W. R. Hansen, D)
- Roberts Tunnel (C. S. Robinson, D)
- Straight Creek tunnel (C. S. Robinson, D)
- Greenland, terrain studies (W. E. Davies, W)

Massachusetts:

- Application of geology and seismology to public-works planning (C. R. Tuttle, R. N. Oldale, Boston)
- Sea-cliff erosion studies (C. A. Kaye, Boston)

Montana, Wolf Point area (R. B. Colton, W)

Nebraska, Valley County (R. D. Miller, D)

Nevada:

Nevada Test Site:

- Pahute Mesa (P. P. Orkild, D)
- Site studies (R. E. Davis, D)

New Mexico, Nash Draw quadrangle (L. M. Gard, D)

South Dakota, Fort Randall Reservoir area (D. J. Varnes, D)

Utah:

- Coal-mine bumps (F. W. Osterwald, D)
- Green River valley, upper part (W. R. Hansen, D)
- Oak City area (D. J. Varnes, D)
- Virginia, Herndon quadrangle (R. E. Eggleton, Flagstaff, Ariz.)

See also Urban geology.

Contamination, water:

- Cadmium-chromium and detergent contamination in ground water, Nassau County, N.Y. (N. M. Perlmutter, w, Albany)
- Detergents, contamination at three public-supply well fields, Suffolk County, N.Y. (N. M. Perlmutter, w, Albany)
- Ground-water contamination (H. E. LeGrand w, W)
- Pesticides and insecticides, determination in water (G. Stratton, w, Columbus, Ohio)
- Pesticides and other pollutants, behavior in the hydrologic environment (M. C. Goldberg, L. Kahn, R. L. Wershaw, h, D)
- Sewage lagoon study (W. J. Powell, w, Tuscaloosa, Ala.)
- See also Radioactive-waste disposal.

Copper:

- Massive sulfide deposits (A. R. Kinkel, Jr., W)
- Sandstone copper deposits, Southwest United States (C. B. Read, Albuquerque, N. Mex.)
- Alaska, southern Brooks Range (W. P. Brosgé, M)

Copper--Continued**Arizona:**

Benson quadrangle (S. C. Creasey, M)
 Globe-Miami area (D. W. Peterson, M)
 Lochiel quadrangle (F. S. Simons, D)
 Mammoth quadrangle (S. C. Creasey, M)
 Nogales quadrangle (F. S. Simons, D)
 Ray district (S. C. Creasey, M)
 Twin Buttes area (J. R. Cooper, D)
 Colorado, Lisbon Valley area (G. W. Weir, Berea, Ky.)
 Michigan, Michigan copper district (W. S. White, W)
 Nevada, Ely district (A. L. Brokaw, D)
 New Mexico, Silver City region (W. R. Jones, D)
 Tennessee, Ducktown district and adjacent areas (R. M. Hernon, D)

Utah:

Bingham Canyon district (R. J. Roberts, M)
 Lisbon Valley area (G. W. Weir, Berea, Ky.)

Crustal studies. See Geophysics, regional.

Detergents. See Contamination, water.

Engineering geologic studies. See Construction and terrain problems; Urban geology.

Evaporation:

Evaporation and thermal-loading analysis, Hycro River basin, North Carolina (G. C. Goddard, W. L. Yonts, w, Raleigh)

Evaporation from Lake Helene, Fla. (R. B. Stone, w, Ocala)

Pond-evaporation study (F. N. Lee, s, Baton Rouge, La.)
 Reservoir evaporation, San Diego County, Calif. (W. Hofmann, w, M)

Evaporation suppression and evaporation:

Mechanics of (G. E. Koberg, h, D)

Evapotranspiration:

Comparison of methods of calculating evapotranspiration, using climatic data (R. W. Cruff, w, M)

Effect of removing riparian vegetation, Cottonwood Wash, Ariz. (J. E. Bowie, w, Tucson)

Evapotranspiration measurements, Deep Creek, Tex. (F. W. Kennon, w, Austin)

Hydrologic effects of vegetation modification (R. M. Myrick, w, Tucson, Ariz.)

Phreatophyte project, Gila River, Ariz. (R. C. Culler, w, Tucson)

Phreatophyte research (T. W. Robinson, w, M)

Extraterrestrial studies:

Astronauts, geologic-training program (R. E. Eggleton, Flagstaff, Ariz.)

Cratering, impact, and thermal investigations:

Experimental hypervelocity impact studies (H. J. Moore, M)

Impact metamorphism (E. C. T. Chao, W)

Shock-phase studies (D. J. Milton, M)

Tension fractures and thermal investigations (A. H. Lachenbruch, M)

Terrestrial impact structures (E. M. Shoemaker, Flagstaff, Ariz.)

Thermoluminescence and mass physical properties (C. H. Roach, D)

Lunar experiments:

Lunar physical properties, measuring techniques (E. M. Shoemaker, Flagstaff, Ariz.)

X-ray fluorescence equipment for lunar studies (I. Adler, W)

Lunar mapping:

Lunar photometry (E. C. Morris, Flagstaff, Ariz.)

Extraterrestrial studies--Continued**Lunar mapping--Continued**

Lunar stratigraphy and structure (R. J. Hackman, W; H. Masursky, M)

Lunar-terrain studies (C. R. Warren, W)

Tektite and meteorite investigations:

Chemistry of tektites (F. Cuttita, W)

Magnetic properties of tektites (A. N. Thorpe, W)

Mineralogy and petrology of meteorites and tektites (E. C. T. Chao, W)

Ferro-alloy metals:

Chromium resource studies (T. P. Thayer, W)

Manganese, geology and geochemistry (D. F. Hewett, M)

Manganese resource studies (J. V. N. Dorr II, W)

Molybdenum-rhenium resource studies (R. U. King, D)

Ultramafic rocks of the Southeastern United States (D. M. Larrabee, W)

California:

Chromite deposits, northern California (F. G. Wells, W)

Tungsten, Bishop district (P. C. Bateman, M)

Idaho, Blackbird Mountain area (J. S. Vhay, Spokane, Wash.)

Montana:

Chromite resources and petrology, Stillwater complex (E. D. Jackson, M)

Manganese deposits, Philipsburg area (W. C. Prinz, W)

Oregon, John Day area (T. P. Thayer, W)

Utah, San Francisco Mountains (D. M. Lemmon, M)

Flood characteristics of streams at selected sites:

Flood studies and bridge-site investigations (C. O. Ming, w, Tuscaloosa, Ala.)

Florida (W. C. Bridges, w, Ocala)

Illinois (W. D. Mitchell, s, Champaign)

Kentucky (C. H. Hannum, s, Louisville)

Nebraska (E. W. Beckman, w, Lincoln)

Puerto Rico (I. J. Hickenlooper, w, San Juan)

Tennessee (W. J. Randolph, w, Chattanooga)

Flood discharge from small drainage areas:

Arizona (B. N. Aldridge, w, Tucson)

Delaware (E. H. Mohler, Jr., w, Towson, Md.)

Idaho (C. A. Thomas, s, Boise)

Illinois (W. D. Mitchell, s, Champaign)

Iowa (H. H. Schwob, s, Iowa City)

Kansas (L. W. Furness, s, Topeka)

Maine (R. A. Morrill, s, Augusta)

Maryland (E. H. Mohler, Jr., w, Towson)

Massachusetts (C. G. Johnson, Jr., w, Boston)

Missouri (E. H. Sandhaus, s, Rolla)

Montana (F. C. Boner, s, Helena)

Nebraska (E. W. Beckman, w, Lincoln)

Nevada (R. D. Lamke, w, Carson City)

New Jersey (J. A. Bettendorf, s, Trenton)

North Dakota (O. A. Crosby, s, Bismarck)

Rhode Island (C. G. Johnson, Jr., w, Boston, Mass.)

South Dakota (R. E. West, s, Pierre)

Tennessee (W. J. Randolph, w, Chattanooga)

Nashville-Davidson County metropolitan area (L. G. Conn, w, Chattanooga)

Vermont (C. G. Johnson, Jr., w, Boston, Mass.)

Virginia (E. M. Miller, s, Charlottesville)

Flood frequency:

Comparison of flood-frequency studies for coastal basins in California (R. W. Cruff, S. E. Rantz, w, M)

Flood frequency, nationwide (A. R. Green, w, W)

Flood frequency--Continued

- Flood volume, duration, frequency (G. A. Kirkpatrick, w, W)
- Synthesis of flood frequency on small drainage areas from rainfall data (S. E. Rantz, w, M)
- Missouri River basin (H. F. Matthai, s, D)
- North Atlantic Slope basins (R. H. Tice, s, St. Louis, Mo.)
- Alabama (L. B. Peirce, w, Tuscaloosa)
- California (L. E. Young, w, M)
- Iowa (H. H. Schwob, s, Iowa City)
- Kansas (L. W. Furness, s, Topeka)
- Mississippi (K. V. Wilson, w, Jackson)
- North Carolina (H. G. Hinson, w, Raleigh)
- Ohio (W. P. Cross, w, Columbus)
- South Carolina (J. S. Stallings, s, Columbia)
- Tennessee (W. J. Randolph, w, Chattanooga)
- Wisconsin (D. W. Ericson, s, Madison)

Flood-inundation mapping:

- Flood-inundation maps (A. R. Green, J. O. Rostvedt, w, W)
- Hawaii, Oahu, Kahaluu area (M. M. Miller, s, Honolulu)
- Illinois, northeastern (W. D. Mitchell, s, Champaign)
- New Jersey (J. A. Bettendorf, s, Trenton)
- New York (K. I. Darmer, w, Albany)
- North Carolina (G. C. Goddard, L. A. Martens, w, Raleigh)
- Pennsylvania:
 - Red Clay Creek basin (M. A. Stoner, s, Harrisburg)
 - Schuylkill River from Conshohocken to Philadelphia (A. T. Alter, s, Harrisburg)
- Puerto Rico (w, San Juan):
 - Arecibo area (M. A. López)
 - Caguas area (M. A. López)
 - Manati area (M. A. López)
 - Mayaguez area (M. A. López)
 - Ponce area (M. A. López)
- Tennessee, Nashville-Davidson County metropolitan area (L. G. Conn, w, Chattanooga)
- Texas, Dallas, Bachman Branch, Joes Creek, and White Rock Creek (F. H. Ruggles, w, Austin)

Flood investigations, areal:

- Flood reports (J. O. Rostvedt, w, W)
- Far Western States, floods of December 1964 (S. E. Rantz, A. M. Moore, w, M)
- Arizona, Maricopa County, Flood investigations (B. N. Aldridge, w, Tucson)
- Arkansas, flood investigations (R. C. Christensen, w, Little Rock)
- Hawaii, Oahu, flood gaging (S. H. Hoffard, s, Honolulu)
- Kansas (L. W. Furness, s, Topeka)
- Kentucky, Ohio River, Floods of March 1964 (H. C. Beaber, s, Louisville)
- Nevada, floods of December 1964 (E. E. Harris, w, Carson City)
- Louisiana:
 - Sabine River near Logansport, flood profile (E. M. Miller, s, Baton Rouge)
 - Southwestern part—rainfall-runoff relations of floods (A. J. Calandro, s, Baton Rouge)
- New Jersey, flood warning (J. E. McCall, s, Trenton)
- New York, peak discharge of ungaged streams (S. H. Hladio, w, Albany)
- North Carolina, flood gaging (H. G. Hinson, w, Raleigh)
- South Carolina, Santee River basin (A. E. Johnson, s, Columbia)

Flood investigations, areal--Continued

- Tennessee:
 - Chattanooga Creek, flood profiles (A. M. F. Johnson, w, Chattanooga)
 - Nashville-Davidson County metropolitan area (L. G. Conn, w, Chattanooga)
- Texas, hydrologic effects of flood-retarding structures (F. W. Kennon, w, Austin)
- Utah, flood gaging (E. Butler, w, Salt Lake City)
- Virginia:
 - Fairfax County and Alexandria city, flood hydrology (D. G. Anderson, s, Charlottesville)
 - Flood investigations (E. M. Miller, s, Charlottesville)
- Wyoming, selected drainage areas under 10 square miles (G. S. Craig, Jr., s)

Fluorspar:

- Colorado, Bonanza and Poncha Springs quadrangles (R. E. Van Alstine, W)

Foreign nations, geologic investigations:

- Bolivia, mineral resources and geologic mapping, advising and training (C. M. Tschanz, La Paz)

Brazil:

- Base-metal resources (A. J. Bodenlos, Rio de Janeiro)
- Geologic education (A. J. Bodenlos, Rio de Janeiro)

- Chile, mineral resources and national geologic mapping (W. Danilchik, D)

- Colombia, minerals exploration and appraisal (E. Irving, Bogota)

- Costa Rica, volcanic studies (K. J. Murata, San Jose)
- Dahomey, minerals reconnaissance (J. A. MacKallor, Cotonou)

- Greenland, eastern, surficial geology, construction-site planning (W. E. Davies, W)

- Indonesia (R. F. Johnson, Bandung)

- Japan, calderas, aeromagnetic-gravity studies (H. R. Blank, Jr., M)

- Liberia (D. L. Rossman, Monrovia)

- Libya, industrial minerals and national geologic map (G. H. Goudarzi, W)

- Pakistan, mineral-resources development, advising and training (M. G. White, Quetta)

- Peru, mineral resources (W. W. Olive, Lima)

- Philippine Islands, iron, chromite, and nonmetallic mineral resources (J. F. Harrington, Manila)

- Saudi Arabia, crystalline shield, geologic and minerals reconnaissance (G. F. Brown, Jidda)

- Thailand, economic geology and mineral industry expansion, advising (L. S. Gardner, Bangkok)

Foreign nations, hydrologic investigations. See Water resources, other countries.**Fuels, organic. See Coal, Oil shale, Petroleum and natural gas.****Gas, natural. See Petroleum and natural gas.****Geochemical distribution of the elements:**

- Botanical exploration and research (H. L. Cannon, D)
- Coding and retrieval of geologic data (T. G. Lovering, D)

- Data of geochemistry (M. Fleischer, W)

- Data of rock analyses (M. Hooker, W)

- Geochemical sampling and statistical analysis of data (A. T. Miesch, D)

- Geochemistry of minor elements (G. Phair, W)

- Mineral fractionation and trace-element content of fine-grained sedimentary rocks (T. D. Botinelly, D)

- Minor-element distribution in black shale (J. D. Vine, M)

Geochemical distribution of the elements--Continued

- Minor elements in volcanic rocks (R. R. Coats, M)
- Organometallic complexes, geochemistry (I. A. Breger, W)
- Sedimentary rocks, chemical composition (H. A. Tourtelot, D)
- Synthesis of ore-mineral data (D. F. Davidson, D)
- California, Sierra Nevada batholith, geochemical study (F. Dodge, M)
- Colorado, Mount Princeton area (P. Toulmin III, W)
- Georgia, biogeochemical reconnaissance (H. T. Shacklette, D)
- Montana, Boulder batholith, petrochemistry (R. I. Tilling, W)
- Nevada, Mt. Wheeler mine area, beryllium distribution (D. E. Lee, M)
- Wisconsin, Driftless area, geochemical survey (H. T. Shacklette, D)

Geochemical prospecting methods:

- Botanical exploration and research (H. L. Cannon, D)
- Dispersion pattern of minor elements related to igneous intrusions (W. R. Griffiths, D)
- Geochemical exploration abstracts (C. Davidson, D)
- Instrument-development laboratory (W. W. Vaughn, D)
- Mineral exploration methods (G. B. Gott, D)
- Mobile spectrographic laboratory (F. N. Ward, D)
- Plant-analysis laboratory (F. N. Ward, D)
- Sulfides, accessory in igneous rocks (G. J. Neuerberg, D)
- Alaska, geochemical prospecting techniques (R. M. Chapman, College)
- Arizona, geochemical halos of mineral deposits (L. C. Huff, Manila, P.I.)
- Maine:
 - Geochemical mapping (E. V. Post, D)
 - The Forks quadrangle (F. C. Canney, E. V. Post, D)
- Michigan, Marquette County (K. Segerstrom, D)
- Nevada, geochemical halos of mineral deposits (R. L. Erickson, D)
- New Mexico, geochemical halos of mineral deposits (L. C. Huff, P.I.)
- Utah, geochemical halos of mineral deposits (R. L. Erickson, D)

Geochemistry, experimental:

- Alkali and alkaline-earth salt systems (E-an Zen, W)
- Environment of ore deposition (P. Toulmin III, W)
- Fluid inclusions in minerals (E. W. Roedder, W)
- Geologic thermometry (E. H. Roseboom, Jr., W)
- Hydrothermal silicate systems (P. Toulmin III, W)
- Hydrothermal solubility (G. W. Morey, W)
- Kinetics of igneous processes (I. Shaw, W)
- Late-stage magmatic processes (G. T. Faust, W)
- Metallic sulfides and sulfosalt systems (P. Toulmin III, W)
- Mineral equilibria, low-temperature (E-an Zen, W)
- Mineral fractionation and trace-element content of fine-grained sedimentary rocks (T. D. Botinelly, D)
- Organic geochemistry (J. G. Palacas, D)
- Organic geochemistry and infrared analysis (I. A. Breger, W)
- Organometallic complexes, geochemistry (I. A. Breger, W)
- Oxygen geothermometry (H. L. James, St. Paul, Minn.)
- Rock weathering and alteration (J. J. Hemley, M)
- Solubility of minerals in aqueous fluids (R. O. Fournier, P. Toulmin III, W)

Geochemistry, experimental--Continued

- Solution-mineral equilibria (C. L. Christ, W)
- Thermodynamic properties of minerals (E. H. Roseboom, Jr., W)

Geochemistry, water:

- Chemistry of atmospheric precipitation (A. W. Gambell, Jr., w, W)
- Geochemical controls of water quality (I. Barnes, w, M)
- Hydrology and geochemistry of the Atlantic Coast Continental Shelf and Slope (R. H. Meade, Jr., F. T. Manheim, w, Woods Hole, Mass.)
- Hydrosolic metals in natural water (J. D. Hem, w, M)
- Mineralogic controls of the chemistry of ground water (B. B. Hanshaw, w, W)
- Minor elements in fresh and saline waters of California, occurrence and distribution (W. D. Silvey, w, Sacramento)
- Radiochemical surveillance (V. J. Janzer, q, D)
- Radioelements, occurrence and distribution in water (R. C. Scott, q, D)
- Solute composition and minor-element distribution in lacustrine closed basins (B. F. Jones, w, W)
- Solute-solid relations in lacustrine closed basins of the alkali-carbonate type (B. F. Jones, w, W)
- Spatial distribution of chemical constituents in ground water (W. Back, w, W)

Geochemistry and petrology, field studies:

- Geochemical sampling and statistical analysis of data (A. T. Miesch, D)
- Geochemistry of minor elements (G. Phair, W)
- Gold, geochemistry and occurrence (J. C. Antweiler, D)
- Green River Formation, mineralogy and geochemistry (C. Milton, W)
- Humates, geology and geochemistry (V. E. Swanson, D)
- Igneous rocks of Southeastern United States (C. Milton, W)
- Inclusions in basaltic rocks (E. D. Jackson, M)
- Jasperoids (T. G. Lovering, D)
- Manganese, geology and geochemistry (D. F. Hewett, M)
- Mercury, geochemistry and occurrence (A. P. Pierce, D)
- Metamorphic rocks and ore deposits (R. C. Erd, M)
- Oceanic volcanics (A. E. J. Engel, La Jolla, Calif.)
- Ore lead, geochemistry and origins (R. S. Cannon, D)
- Pacific Coast basalts, geochemistry (K. J. Murata, M)
- Pierre Shale, chemical and physical properties, Montana, North Dakota, Nebraska, South Dakota, and Wyoming (H. A. Tourtelot, D)
- Rare-earth elements, resources and geochemistry (J. W. Adams, D)
- Sedimentary petrology laboratory (H. A. Tourtelot, D)
- Selenium, resources and geochemistry (D. F. Davidson, D)
- Thermal waters, origin and characteristics (D. E. White, M)
- Titanium, geochemistry and occurrence (N. Herz, W)
- Volcanoes, infrared studies (R. W. Moxham, W)
- Alaska, Katmai National Monument, petrology and volcanism (G. H. Curtis, M)
- California:
 - Burney area (G. A. MacDonald, Honolulu, Hawaii)
 - Coast Range ultramafic rocks (R. A. Loney, M)
 - Franciscan Formation, glaucophane schist (R. G. Coleman, M)
 - Kings Canyon National Park (J. G. Moore, M)
 - Ritter Range, metavolcanic rocks (R. S. Fiske, W)
 - Sierra Nevada batholith, geochemical study (F. Dodge, M)

Geochemistry and petrology, field studies--Continued**Colorado:****Front Range:**

Boulder Creek batholith (G. Phair, W)

Laramide intrusives (G. Phair, W)

Minturn quadrangle (T. S. Lovering, D)

Mt. Princeton area, distribution of elements (P. Toulmin III, W)

Wet Mountains, wallrock alteration (G. Phair, W)

Florida, Pamlico Sound area, organic geochemistry (H. L. Berryhill, Jr., D)

Hawaii, Hawaiian volcanology (H. A. Powers, Hawaii National Park, Hawaii)

Idaho, central Snake River plain, volcanic petrology (H. A. Powers, Hawaii National Park, Hawaii)

Montana:

Bearpaw Mountains, petrology (W. T. Pecora, W)

Boulder batholith, petrochemistry (R. I. Tilling, W)

Stillwater complex, petrology and chromite resources (E. D. Jackson, M)

Wolf Creek area, petrology (R. G. Schmidt, W)

New Mexico:

Grants area, mineralogy of uranium-bearing rocks (A. D. Weeks, W)

Valles Mountains (R. L. Smith, W)

New York, Gouverneur area, metamorphism and origin of mineral deposits (A. E. J. Engel, La Jolla, Calif.)

South Carolina, igneous and metamorphic rocks of the piedmont (W. C. Overstreet, Jidda, Saudi Arabia)

Texas, Duval and Karnes Counties, mineralogy of uranium-bearing rocks (A. D. Weeks, W)

Wisconsin, geochemical survey of the Driftless area (H. T. Schacklette, D)

Wyoming:

Green River Formation, geology and paleolimnology (W. H. Bradley, W)

Yellowstone Park, thermal waters and deposits (G. W. Morey, R. O. Fournier, W)

Geochronology:

Carbon-14 method (M. Rubin, W)

Definition of a tree-growth trend function (N. C. Matalas, L. Horner, w, W)

Dendrochronology (D. O'Bryan, w, W)

General statistical properties of sequences of tree-ring indices (N. C. Matalas, L. Horner, w, W)

Geologic time scale (R. E. Zartman, D)

Igneous rocks and deformational periods (R. W. Kistler, M)

Lead-alpha method (T. W. Stern, W)

Lead-uranium method (T. W. Stern, W)

Potassium-argon and rubidium-strontium methods (H. H. Thomas, C. E. Hedge, D; R. Kistler, M)

Radioactive-disequilibrium studies (J. N. Rosholt, D)

Alaska, southeastern part (G. D. Eberlein, M. A. Lanphere, M)

Colorado, Mesa Verde National Park, correlations between tree growth and climatic factors (D. O'Bryan, N. C. Matalas, w, W)

Maryland, Oakland, correlations between tree growth and climatic factors (D. O'Bryan, L. Horner, w, W)

See also Isotope and nuclear studies.**Geologic mapping:**

Map scale smaller than 1 inch to 1 mile:

Colorado Plateau:

Geologic maps (2-minute sheets) (D. G. Wyant, D)

Photogeologic mapping (A. B. Olson, W)

Geologic mapping--Continued

Map scale smaller than 1 inch to 1 mile--Continued

Sino-Soviet terrain atlas (M. M. Elias, W)

Alaska:

Compilation of geologic maps, 1:250,000 quadrangles (G. Gryc, M)

Geologic map of State (G. O. Gates, M)

Metallogenic map (C. L. Sainsbury, D)

Central and northern part, Cenozoic (D. M. Hopkins, M)

Northern part, petroleum investigations (G. Gryc, M)

Bristol Bay area, surficial geology (E. H. Muller, Ithaca, N.Y.)

Brooks Range, southern part (W. P. Brosge, M)

Buckland River area (W. W. Patton, Jr., M)

Charley River quadrangle (E. E. Brabb, M)

Delong Mountains quadrangle (I. L. Tailleir, M)

Fairbanks quadrangle (F. R. Weber, College)

Hughes-Shungnak area (W. W. Patton, Jr., M)

Huslia River area (W. W. Patton, Jr., M)

Iliamna quadrangle (R. L. Determan, M)

Kenai lowland, surficial geology (T. N. V. Karlstrom, W)

Klukwan iron district (E. C. Robertson, W)

Kobuk River valley (A. T. Fernald, W)

Livengood quadrangle (B. Taber, M)

Lower Yukon-Koyukuk area (W. W. Patton, Jr., M)

Lower Yukon-Norton Sound region (J. M. Hoare, M)

Nelchina area (A. Grantz, M)

Point Hope quadrangle (I. L. Tailleir, M)

Yukon-Koyukuk lowland, engineering geology (F. R. Weber, College)

Antarctica:

Western part, reconnaissance geology (E. L. Boudette, W)

Eights and Walgreen Coasts, reconnaissance geology (A. A. Drake, Jr., W)

Victoria Land, northeastern part (W. B. Hamilton, D)

California, San Raphael Primitive Area (H. D. Gower, M)

Colorado:

Oil-shale investigations (D. C. Duncan, W)

Durango 2-degree quadrangle (T. A. Steven, D)

Grand Junction 2-degree quadrangle (W. B. Cashion, D)

La Junta 2-degree quadrangle (G. R. Scott, D)

Pueblo 2-degree quadrangle (G. R. Scott, D)

Trinidad 2-degree quadrangle (R. B. Johnson, D)

Idaho:

Preston 2-degree quadrangle (S. S. Oriel, D)

Snake River plain, central part, volcanic petrology (H. E. Malde, D)

Spokane-Wallace region (A. B. Griggs, M)

Montana:

Butte 2-degree quadrangle (M. R. Klepper, W)

Spokane-Wallace region (A. B. Griggs, M)

Nevada:

Clark County (C. R. Longwell, M)

Douglas County (J. G. Moore, M)

Esmeralda County (J. P. Albers, M)

Eureka County (R. J. Roberts, M)

Lincoln County (C. M. Tschanz, La Paz, Bolivia)

Lyon County (J. G. Moore, M)

Geologic mapping--Continued**Map scale smaller than 1 inch to 1 mile--Continued****Nevada--Continued**

Nevada Test Site, reconnaissance (F. N. Houser, D)

Nye County:

Northern part (F. J. Kleinhampl, M)

Southern part (H. R. Cornwall, M)

Ormsby County (J. G. Moore, M)

Pershing County (D. B. Tatlock, M)

Ruby Mountains (C. R. Willden, D)

White Pine County (R. K. Hose, M)

New Mexico, geologic map (C. H. Dane, W)

North Carolina:

Knoxville 2-degree quadrangle (J. B. Hadley, W)

Winston-Salem 2-degree quadrangle (D. W. Rankin, G. H. Espenshade, W)

Oregon, geologic map (G. W. Walker, M)

South Carolina, Knoxville 2-degree quadrangle (J. B. Hadley, W)

Tennessee, Knoxville 2-degree quadrangle (J. B. Hadley, W)

Utah, Grand Junction 2-degree quadrangle (W. B. Cashion, D)

Virginia, Winston-Salem 2-degree quadrangle (D. W. Rankin, G. H. Espenshade, W)

Washington:

Grays Harbor basin, regional compilation (H. M. Beikman, M)

Spokane-Wallace region (A. B. Griggs, M)

Wyoming, Preston 2-degree quadrangle (S. S. Oriel, D)

Map scale 1 inch to 1 mile, and larger:

Alabama, Warrior quadrangle (W. C. Culbertson, D)

Alaska:

Project CHARIOT (harbor construction) (G. D. Eberlein, M)

Gulf of Alaska, Tertiary province (G. Plafker, M)

Northeastern part, coastal plain and foothills (C. R. Lewis, W)

Aleutian Islands:

Eastern part (R. E. Wilcox, D)

Western part (R. E. Wilcox, D)

Aleutian Trench - Trinity Island (G. W. Moore, M)

Anchorage - Matanuska Glacier area, surficial geology (T. N. V. Karlstrom, W)

Annette Island (H. C. Berg, M)

Beluga-Yentna area (F. F. Barnes, M)

Bering River coal field (A. A. Wanek, c, Anchorage)

Chitina Valley, lower part, surficial geology (L. A. Yehle, W)

Copper River basin:

Northeastern part, surficial geology (O. J. Ferrans, Jr., W)

Southeastern part, surficial geology (D. R. Nichols, W)

Southwestern part, surficial geology (J. R. Williams, W)

Denali Highway, eastern part, surficial geology (D. R. Nichols, W)

Heceta-Tuxekan area (G. D. Eberlein, M)

Iniskin-Tuxedni region (R. L. Detterman, M)

Johnson River district, surficial geology (H. L. Foster, W)

Geologic mapping--Continued**Map scale 1 inch to 1 mile, and larger--Continued****Alaska--Continued**

Katmai National Monument, petrology and volcanism (G. H. Curtis, M)

Lost River mining district (C. L. Sainsbury, D)

Mt. Chamberlain area, surficial geology (C. R. Lewis, W)

Mt. Hayes D-3 and D-4 quadrangles (T. L. Péwé, College)

Mt. Michelson area (E. G. Sable, Elizabethtown, Ky.)

Nenana coal investigations (C. Wahrhaftig, M)

Nome C-1 and D-1 quadrangles (C. L. Hummel, Bangkok, Thailand)

Seward-Portage Railroad, surficial geology (T. N. V. Karlstrom, W)

Slana-Tok area, surficial geology (H. R. Schmoll, W)

Steese Highway area, surficial geology (W. E. Davies, W)

Tanana River, upper part, surficial geology (A. T. Fernald, W)

Taylor Highway area, surficial geology (H. L. Foster, W)

Tofty placer district (D. M. Hopkins, M)

Valdez-Tiekel belt, surficial geology (H. W. Coulter, W)

Windy-Curry area (R. Kachadoorian, M)

Antarctica:

Horlick Mountains (A. B. Ford, W)

Pensacola Mountains (D. L. Schmidt, W)

Arizona:

Benson quadrangle (S. C. Creasey, M)

Blue Horse Mountain quadrangle (A. F. Shride, D)

Bradshaw Mountains (C. A. Anderson, M)

Carrizo Mountains area (J. D. Strobell, D)

Cibecue-Grasshopper area (T. L. Finnell, D)

Cochise County, southern part (P. T. Hayes, D)

Elgin quadrangle (R. B. Raup, D)

Empire Mountains (T. L. Finnell, D)

Gila River basin, upper part (R. B. Morrison, D)

Globe-Miami area (D. W. Peterson, M)

Heber quadrangle (E. J. McKay, D)

Holy Joe Peak quadrangle (M. H. Krieger, M)

Lochiel quadrangle (F. S. Simons, D)

McFadden Peak quadrangle (A. F. Shride, D)

Mammoth quadrangle (S. C. Creasey, M)

Mt. Wrightson quadrangle (H. Drewes, D)

Mustang Mountains (R. B. Raup, D)

Navajo Reservation, fuels potential (R. B. O'Sullivan, D)

Nogales quadrangle (F. S. Simons, D)

Prescott-Paulden area (M. H. Krieger, M)

Ray district, porphyry copper (S. C. Creasey, M)

Show Low quadrangle (E. J. McKay, D)

Twin Buttes area (J. R. Cooper, D)

Winkelman quadrangle (M. H. Krieger, M)

Arkansas:

Northern part, oil and gas investigations (E. E. Glick, D)

Arkansas Basin, coal investigations (B. R. Haley, D)

Ft. Smith district (T. A. Hendricks, D)

California:

Big Maria Mountains (W. B. Hamilton, D)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued
California--Continued

Bishop tungsten district (P. C. Bateman, M)
Blanco Mountain quadrangle (C. A. Nelson, Los Angeles)
Bucks Lake quadrangle (A. Hietanen-Makela, M)
Burney area (G. A. Macdonald, Honolulu, Hawaii)
Coast Range, ultramafic rocks (E. H. Bailey, M)
Condrey Mountain quadrangle (P. E. Hotz, M)
Cuyama Valley area (J. G. Vedder, M)
Death Valley (C. B. Hunt, Baltimore, Md.)
Furnace Creek area (J. F. McAllister, M)
Independence quadrangle (D. C. Ross, M)
Klamath Mountains, southern part (W. P. Irwin, M)
Little Maria Mountains (W. B. Hamilton, D)
Los Angeles area (J. T. McGill, D)
Los Angeles basin, eastern part (J. E. Schoellhamer, M)
Malibu Beach quadrangle (R. F. Yerkes, M)
Merced Peak quadrangle (D. L. Peck, Honolulu, Hawaii)
Mojave Desert:
 South-central part (T. W. Dibblee, Jr., M)
 Western part (T. W. Dibblee, Jr., M)
New York Butte quadrangle (W. C. Smith, M)
Oakland East quadrangle (D. H. Radbruch, M)
Palo Alto quadrangle (E. H. Pampeyan, M)
Panamint Butte quadrangle (W. E. Hall, W)
Point Dume quadrangle (R. H. Campbell, M)
Priest Valley SE quadrangle (E. E. Richardson, c, Bakersfield)
Riverside Mountains (W. B. Hamilton, D)
Sacramento Valley, northwest part (R. D. Brown, Jr., W)
Salinas Valley (D. L. Durham, M)
San Andreas fault (L. F. Noble, Valyermo)
San Francisco North quadrangle (J. Schlocker, M)
San Francisco South quadrangle (M. G. Bonilla, M)
San Mateo quadrangle (G. O. Gates, M)
Searles Lake area (G. I. Smith, M)
Shuteye Peak area (N. K. Huber, M)
Sierra foothills mineral belt (L. D. Clark, M)
Sierra Nevada batholith (P. C. Bateman, M)
Sierra tungsten belt, eastern (N. K. Huber, M)
White Mountain Peak quadrangle (D. F. Crowder, M)

Colorado:

Air Force Academy (D. J. Varnes, D)
Animas River area (H. Barnes, D)
Aspen quadrangle (B. Bryant, D)
Baggs area (G. E. Prichard, D)
Berthoud Pass quadrangle (P. K. Theobald, Jidda, Saudi Arabia)
Black Canyon of the Gunnison River (W. R. Hansen, D)
Black Hawk quadrangle (R. B. Taylor, W)
Bonanza quadrangle (R. E. Van Alstine, W)
Bottle Pass quadrangle (R. B. Taylor, W)
Boulder quadrangle (C. T. Wrukke, D)
Bull Canyon district (C. H. Roach, D)
Cameron Mountain quadrangle (C. T. Wrukke, D)
Carbondale coal field (J. R. Donnell, D)
Cerro Summit quadrangle (R. G. Dickinson, c, D)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued
Colorado--Continued

Cheyenne Mountain, electrical properties (J. H. Scott, D)
Corral Bluffs quadrangle (P. E. Soister, c, D)
Creede district (T. A. Steven, D)
Denver metropolitan area (R. M. Lindvall, D)
East Portal quadrangle (P. K. Theobald, Jidda, Saudi Arabia)
Eldorado Springs quadrangle (J. D. Wells, D)
Elk Springs quadrangle (J. R. Dyni, c, D)
Empire quadrangle (W. A. Braddock, Boulder)
Evergreen quadrangle (D. M. Sheridan, D)
Fraser quadrangle (P. K. Theobald, Jidda, Saudi Arabia)
Front Range:
 East-central part, mountain front area (D. M. Sheridan, D)
 Northeastern part, Fort Collins area (W. A. Braddock, Boulder)
Golden quadrangle (R. Van Horn, D)
Grand-Battlement Mesa (J. R. Donnell, D)
Green River valley, upper part (W. R. Hansen, D)
Hanover NW quadrangle (P. E. Soister, c, D)
Holy Cross quadrangle (O. Tweto, D)
Hot Sulphur Springs quadrangle (G. A. Izett, c, D)
Indian Hills quadrangle (D. M. Sheridan, D)
Kokomo mining district (M. H. Bergendahl, D)
Kremmling quadrangle (G. A. Izett, c, D)
La Sal area (W. D. Carter, W)
Lafayette quadrangle (K. B. Ketner, D)
Lake George district (C. C. Hawley, D)
Lisbon Valley area (G. W. Weir, Berea, Ky.)
Marble quadrangle (D. L. Gaskill, c, D)
Marcellina Mountain quadrangle (D. L. Gaskill, c, D)
Maybell-Lay area (M. J. Bergin, W)
Montrose 1 SE, 1 SW, and 4 NE quadrangles (R. G. Dickinson, c, D)
Morrison quadrangle (D. J. Gable, D)
Mt. Antero (W. N. Sharp, D)
Mt. Harvard quadrangle (M. R. Brock, D)
Nederland quadrangle (D. J. Gable, D)
Niwt quadrangle (M. McLachlan, D)
North Park:
 Eastern part (D. M. Kinney, W)
 Western part (W. J. Hail, D)
Oh-Be-Joyful quadrangle (D. L. Gaskill, c, D)
Park Range, northern part (G. L. Snyder, D)
Placita SE quadrangle (L. H. Godwin, c, D)
Poncha Springs quadrangle (R. E. Van Alstine, W)
Powderhorn area (J. C. Olson, D)
Pueblo and vicinity (G. R. Scott, D)
Ralston Buttes (D. M. Sheridan, D)
Rangely 3 quadrangle (H. L. Cullins, c, D)
Rico district (E. T. McKnight, W)
Rico-Animas area (W. P. Pratt, D)
Ruedie quadrangle (V. L. Freeman, D)
San Juan mining area (R. G. Luedke, W)
San Juan Mountains, western (A. L. Bush, D)
Slick Rock district (D. R. Shawe, D)
Squaw Pass quadrangle (D. M. Sheridan, D)
Straight Creek tunnel (C. S. Robinson, D)
Tenmile Range (M. H. Bergendahl, D)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Colorado--Continued

- Thornburg area (J. R. Dyni, c, D)
- Trinidad coal field (R. B. Johnson, D)
- Tungsten quadrangle (D. J. Gable, D)
- Wet Mountains (M. R. Brock, D)
- Woody Creek quadrangle (V. L. Freeman, D)

Connecticut:

Ashaway quadrangle:

- Bedrock geology (T. G. Feininger, Boston, Mass.)
- Surficial geology (J. P. Schafer, Boston, Mass.)
- Ashley Falls quadrangle, construction materials (G. W. Holmes, M)
- Bristol quadrangle, bedrock geology (H. E. Simpson, D)
- Broad Brook quadrangle (R. B. Colton, W)
- Columbia quadrangle, bedrock geology (G. L. Snyder, D)
- Danielson quadrangle, bedrock geology (H. R. Dixon, Boston, Mass.)
- Durham quadrangle (H. E. Simpson, D)
- Fitchville quadrangle, bedrock geology (G. L. Snyder, D)
- Hampton quadrangle, bedrock geology (H. R. Dixon, Boston, Mass.)
- Manchester quadrangle (R. B. Colton, W)
- Marlborough quadrangle, bedrock geology (G. L. Snyder, D)
- Meriden quadrangle, bedrock geology (P. M. Hanshaw, Boston, Mass.)
- Montville quadrangle, bedrock geology (R. Goldsmith, Boston, Mass.)
- Mystic quadrangle, bedrock geology (R. Goldsmith, Boston, Mass.)
- New Britain quadrangle, bedrock geology (H. E. Simpson, D)
- New Hartford quadrangle (R. W. Schnabel, D)
- New London quadrangle (R. Goldsmith, D)
- Niantic quadrangle (R. Goldsmith, D)
- Old Mystic quadrangle (R. Goldsmith, D)
- Plainfield quadrangle, bedrock geology (H. R. Dixon, Boston, Mass.)
- Scotland quadrangle, bedrock geology (H. R. Dixon, Boston, Mass.)
- Southwick quadrangle (R. W. Schnabel, D)
- Springfield South quadrangle (J. H. Hartshorn, C. Koteff, Boston, Mass.)
- Tariffville quadrangle:
 - Bedrock geology (R. W. Schnabel, D)
 - Surficial geology (A. D. Randall, g, Middletown)
- Thompson quadrangle (P. M. Hanshaw, H. R. Dixon, Boston, Mass.)
- Tolland Center quadrangle, construction materials (G. W. Holmes, M)
- Uncasville quadrangle, bedrock geology (R. Goldsmith, Boston, Mass.)
- Voluntown quadrangle, bedrock geology (T. G. Feininger, Boston, Mass.)
- Watch Hill quadrangle:
 - Bedrock geology (G. E. Moore, Jr., Columbus, Ohio)
 - Surficial geology (J. P. Schafer, Boston, Mass.)
- West Springfield quadrangle (R. B. Colton, J. H. Hartshorn, Boston, Mass.)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Connecticut--Continued

- Windsor Locks quadrangle, bedrock geology (R. W. Schnabel, D)
- District of Columbia, Washington metropolitan area (H. W. Coulter, C. F. Withington, W)

Florida:

- Land-pebble phosphate deposits (J. B. Cathcart, D)
- Attapulugus-Thomasville area, fuller's earth deposits (S. H. Patterson, W)

Georgia:

- Attapulugus-Thomasville area, fuller's earth deposits (S. H. Patterson, W)
- Brevard Belt, crystalline rocks (M. Higgins, Atlanta)
- Greenland, Schuchert Dal, East Greenland, glacial geology (J. S. Hartshorn, Boston, Mass.)

Idaho:

- Central part, radioactive placer deposits (D. L. Schmidt, W)
- American Falls region (D. E. Trimble, D)
- Aspen Range-Dry Ridge area (V. E. McKelvey, W)
- Bancroft quadrangle (S. S. Oriel, D)
- Bayhorse area (S. W. Hobbs, D)
- Big Creek quadrangle (B. F. Leonard, D)
- Clarks Fork quadrangle (J. E. Harrison, W)
- Coeur d'Alene mining district (S. W. Hobbs, D)
- Doublesprings quadrangle (W. J. Mapel, D)
- Driggs NE, SE, and SW quadrangles (M. L. Schroeder, c, D)
- Elmira quadrangle (J. E. Harrison, W)
- Garns Mountain NW and SW quadrangles (M. H. Staatz, c, D)
- Greenacres quadrangle (P. L. Weis, Spokane, Wash.)
- Hawley Mountain quadrangle (W. J. Mapel, D)
- Leadore quadrangle (E. T. Ruppel, D)
- Mt. Spokane quadrangle (A. E. Weissenborn, Spokane, Wash.)
- Mountain City quadrangle (R. R. Coats, M)
- Orofino area (A. Hietanen-Makela, M)
- Owyhee quadrangle (R. R. Coats, M)
- Packsaddle Mountain quadrangle (J. E. Harrison, W)
- Palisades Peak quadrangle (D. A. Jobin, c, D)
- Palisades Reservoir quadrangle (H. F. Albee, c, D)
- Patterson quadrangle (E. T. Ruppel, D)
- Pocatello quadrangle (D. E. Trimble, D)
- Poker Peak quadrangle (H. F. Albee, c, D)
- Riggins quadrangle (W. B. Hamilton, D)
- Soda Springs quadrangle (F. C. Armstrong, D)
- Upper Valley quadrangle (R. L. Rioux, c, W)
- Yellow Pine quadrangle (B. F. Leonard, D)
- Indiana, Owensboro quadrangle, Quaternary geology (L. L. Ray, W)

Kansas:

- Shawnee County (W. D. Johnson, Jr., Owensboro, Ky.)
- Wilson County (H. C. Wagner, M)

Kentucky:

Note: The entire State of Kentucky is being mapped geologically by 7½-minute quadrangles under a cooperative program with the Kentucky Geological Survey. 141 quadrangles have been published and 208 more are currently in progress. Project is

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Kentucky--Continued

under the supervision of P. W. Richards, Lexington, Ky. The following investigations are separate from the cooperative mapping program:

Appalachian folded belt, southern part (L. D. Harris, Knoxville, Tenn.)

Eastern part, coal investigations (K. J. Englund, W)

Jellico West quadrangle (K. J. Englund, W)

Ketchen quadrangle (K. J. Englund, W)

Owensboro quadrangle, Quaternary geology (L. L. Ray, W)

Maine:

Paleozoic stratigraphy, regional (R. B. Neuman, W)

Aroostook County, southern (L. Pavlides, W)

Attean quadrangle (A. L. Albee, Pasadena, Calif.)

Big Lake area (D. M. Larrabee, W)

Greenville quadrangle (G. H. Espenshade, W)

Kennebago Lake quadrangle (E. L. Boudette, W)

Moosehead gabbro (G. H. Espenshade, W)

Rangeley quadrangle (R. H. Moench, D)

Rumford quadrangle (R. H. Moench, D)

Stratton quadrangle, geophysical and geologic mapping (A. Griscom, W)

The Forks quadrangle (F. C. Canney, E. V. Post, D)

Maryland:

Chesapeake Bay area, upper part (J. P. Minard, W)

Washington, D. C., metropolitan area (H. W. Coulter, C. F. Withington, W)

Harford County (D. Southwick, W)

Massachusetts:

Assawompsett Pond quadrangle (C. Koteff, Boston)

Athol quadrangle (D. F. Eschman, Ann Arbor, Mich.)

Billerica quadrangle (R. H. Jahns, University Park, Pa.)

Blue Hills quadrangle (N. E. Chute, Syracuse, N. Y.)

Boston and vicinity (C. A. Kaye, Boston)

Clinton quadrangle, bedrock geology (R. F. Novotny, Boston)

Concord quadrangle (N. P. Cuppels, C. Koteff, Boston)

Duxbury quadrangle (N. E. Chute, Syracuse, N.Y.)

Georgetown quadrangle (N. P. Cuppels, Boston)

Heath quadrangle (A. H. Chidester, D; J. H. Hartshorn, Boston)

Lawrence quadrangle, bedrock geology (R. O. Castle, Los Angeles, Calif.)

Lowell quadrangle (R. H. Jahns, University Park, Pa.)

Norwood quadrangle (N. E. Chute, Syracuse, N.Y.)

Plainfield quadrangle, bedrock geology (P. H. Osberg, Orono, Maine)

Reading quadrangle:

Bedrock geology (R. O. Castle, Los Angeles, Calif.)

Surficial geology (R. N. Oldale, Boston)

Rowe quadrangle (A. H. Chidester, D; J. H. Hartshorn, Boston)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Massachusetts--Continued**Salem quadrangle:**

Bedrock geology (P. Toulmin III, W)

Surficial geology (R. N. Oldale, Boston)

Scituate quadrangle (N. E. Chute, Syracuse, N.Y.)

Shrewsbury quadrangle, bedrock geology (R. F. Novotny, Boston)

South Groveland quadrangle, bedrock geology (R. O. Castle, Los Angeles, Calif.)

Southwick quadrangle (R. W. Schnabel, D)

Springfield South quadrangle (J. H. Hartshorn, C. Koteff, Boston)

Taunton quadrangle (J. H. Hartshorn, Boston)

Tyngsboro quadrangle (R. H. Jahns, University Park, Pa.)

West Springfield quadrangle (R. B. Colton, D; J. H. Hartshorn, Boston)

Westford quadrangle (R. H. Jahns, University Park, Pa.)

Wilmington quadrangle, bedrock geology (R. O. Castle, Los Angeles, Calif.)

Michigan:

Dickinson County, southern (R. W. Bayley, M)

Gogebic Range, eastern (W. C. Prinz, W)

Iron County, eastern (K. L. Wier, D)

Iron River-Crystal Falls district (H. L. James, Minneapolis, Minn.)

Lake Algonquin drainage (J. T. Hack, W)

Marenisco - Watersmeet area, iron deposits (C. E. Fritts, D)

Marquette district, eastern (J. E. Gair, D)

Michigan copper district (W. S. White, W)

Negaunee quadrangle (J. E. Gair, D)

Palmer quadrangle (J. E. Gair, D)

Mississippi:

Homochitto National Forest (E. L. Johnson, c, Tulsa, Okla.)

Tatum salt dome (W. S. Twenhofel, D)

Missouri, Lesterville quadrangle (T. H. Kiilsgaard, W)

Montana:

Southwestern part, ore deposits (K. L. Wier, D)

Alberton quadrangle (J. D. Wells, W)

Anaconda NW quadrangle (A. A. Wanek, c, Anchorage, Alaska)

Bearpaw Mountains, petrology (W. T. Pecora, W)

Black Butte quadrangle (A. F. Bateman, Jr., c, Great Falls)

Boulder batholith area (M. R. Klepper, W)

Browning area, Quaternary geology (G. M. Richmond, D)

Cameron quadrangle (J. B. Hadley, W)

Clarks Fork quadrangle (J. E. Harrison, W)

Crazy Mountains Basin (B. A. Skipp, D)

Gardiner SW quadrangle (G. D. Fraser, c, D)

Girard coal field (G. E. Prichard, D)

Great Falls area (R. W. Lemke, D)

Hardy quadrangle (K. S. Soward, c, Great Falls)

Hedstrom area (A. F. Bateman, Jr., c, Great Falls)

Holter Lake quadrangle (G. D. Robinson, D)

Hughesville quadrangle (I. J. Witkind, D)

Jordan (30-minute) quadrangle (G. D. Mowat, c, Great Falls)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Montana--Continued

- Livingston--Trail Creek area (A. E. Roberts, D)
- Maudlow quadrangle (B. A. Skipp, D)
- Montaqua quadrangle (E. D. Patterson, c, W)
- Neihart 1 quadrangle (W. R. Keefer, D)
- Packsaddle Mountain quadrangle (J. E. Harrison, W)
- Philipsburg area, manganese deposits (W. C. Prinz, W)
- Powder River coal fields (N. W. Bass, D)
- Rocky Reef quadrangle (K. S. Soward, c, Great Falls)
- Sun River Canyon area (M. R. Mudge, D)
- Tepee Creek quadrangle (I. J. Witkind, D)
- Toston quadrangle (G. D. Robinson, D)
- Varney quadrangle (J. B. Hadley, W)
- Wise River quadrangle (G. D. Fraser, c, D)
- Wolf Creek area, petrology (R. G. Schmidt, W)
- Wolf Point area (R. B. Colton, D)

Nebraska:

- Omaha--Council Bluffs and vicinity (R. D. Miller, D)

Valley County (R. D. Miller, D)

Nevada:

- Antler Peak quadrangle (R. J. Roberts, M)
- Ash Meadows quadrangle (C. S. Denny, W)
- Beatty area (H. R. Cornwall, M)
- Bellevue Peak quadrangle (T. B. Nolan, W)
- Buffalo Mountain quadrangle (R. E. Wallace, M)
- Coaldale area (R. G. Wayland, c, Los Angeles, Calif.)
- Cortez quadrangle (J. Gilluly, D)
- Crescent Valley quadrangle (J. Gilluly, D)
- Ely district (A. L. Brokaw, D)
- Eureka quadrangle (T. B. Nolan, W)
- Garrison quadrangle (D. H. Whitebread, M)
- Horse Creek Valley quadrangle (H. Masursky, M)
- Humboldt Range, Unionville and Buffalo Mountain quadrangles (R. E. Wallace, M)
- Jiggs quadrangle (C. R. Willden, D)
- Kobeh Valley (T. B. Nolan, W; C. W. Merriam, M)
- Las Vegas--Lake Mead area (C. R. Longwell, M)
- Montello area (R. G. Wayland, c, Los Angeles, Calif.)

Mt. Lewis quadrangle (J. Gilluly, D)

Mountain City quadrangle (R. R. Coats, M)

Nevada Test Site:

Geologic studies (F. A. McKeown, D)

Pahute Mesa (F. N. Houser, D)

Site studies (R. E. Davis, D)

Owyhee quadrangle (R. R. Coats, M)

Pinto Summit quadrangle (T. B. Nolan, W)

Pioche district (C. M. Tschanz, La Paz, Bolivia)

Railroad district (J. F. Smith, Jr., D)

Schell Creek Range (H. D. Drewes, D)

Snake Range quadrangle (D. H. Whitebread, M)

Sonoma Range, northern, orogenic processes (J. Gilluly, D)

Spruce Mountain 4 quadrangle (G. D. Fraser, c, D)

Unionville quadrangle (R. E. Wallace, M)

Wheeler Peak quadrangle (D. H. Whitebread, M)

New Jersey:

Delaware River basin:

Lower part (J. P. Owens, W)

Middle part (A. A. Drake, Jr., W)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

New Mexico:

- Animas River area (H. Barnes, D)
- Carrizo Mountains area (J. D. Strobell, D)
- Franklin Mountains (R. L. Harbour, D)
- Gila River basin, upper part (R. B. Morrison, D)
- Grants area (R. E. Thaden, Columbia, Ky.)
- Johnson Trading Post quadrangle (J. S. Hinds, c, Farmington)
- Laguna district (R. H. Moench, D)
- Las Vegas quadrangle, western half (E. H. Baltz, g, Albuquerque)
- Manzano Mountains (D. A. Myers, D)
- Mesa Portales quadrangle (J. E. Fassett, c, Farmington)
- Nash Draw quadrangle (L. M. Gard, D)
- Oscura Mountains, southern part (G. O. Brachman, D)
- Raton coal basin:
 - Eastern part (G. H. Dixon, D)
 - Western part (C. L. Pillmore, D)
- San Andres Mountains, northern part (G. O. Bachman, D)
- San Juan Basin, east side (C. H. Dane, W)
- Silver City area (W. R. Jones, D)
- Valles Mountains, petrology (R. L. Smith, W)

New York:

- Dannemora quadrangle, surficial geology (C. S. Denny, W)
- Gouverneur area, metamorphism and origin of mineral deposits (A. E. J. Engel, La Jolla, Calif.)
- Mooers quadrangle (A. W. Postel, W)
- Ohio quadrangle (A. W. Postel, W)
- Plattsburgh quadrangle, surficial geology (C. S. Denny, W)
- Richville quadrangle (H. M. Bannerman, W)

North Carolina:

- Central Piedmont (H. Sundelius, W)
- Franklin quadrangle (F. G. Lesure, W)
- Grandfather Mountain (B. H. Bryant, D)
- Great Smoky Mountains (J. B. Hadley, W)
- Morganton area, geomorphic studies (J. T. Hack, W)
- Mount Rogers area (D. W. Rankin, W)
- Volcanic Slate series (A. A. Stromquist, D)

North Dakota:

- Clark Butte and Clark Butte NE quadrangles (G. D. Mowat, c, Great Falls, Mont.)
- Clark Butte NW and SW quadrangles (E. H. Gilmore, c, Great Falls, Mont.)
- Dengate quadrangle (C. S. V. Barclay, c, D)
- Glen Ullin quadrangle (C. S. V. Barclay, c, D)
- Heart Butte and Heart Butte NW quadrangles (E. V. Stephens, c, D)
- New Salem quadrangle (H. L. Smith, c, D)
- North Almont quadrangle (H. L. Smith, c, D)

Oklahoma, Ft. Smith district (T. A. Hendricks, D)

Oregon:

- Bandon SE quadrangle (R. G. Wayland, c, Los Angeles, Calif.)
- Coquille SW quadrangle (R. G. Wayland, c, Los Angeles, Calif.)
- John Day area (T. P. Thayer, W)
- Monument quadrangle (R. E. Wilcox, D)
- Newport Embayment (P. D. Snively, Jr., M)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Oregon--Continued

Ochoco Reservation, Lookout Mountain, Eagle Rock, and Post quadrangles (A. C. Waters, Baltimore, Md.)

Pacific Islands:

Bikini and nearby atolls (H. S. Ladd, W)
Western Pacific islands (G. Corwin, W)

Pennsylvania:

Anthracite mine-drainage projects, geology in the vicinity of (J. F. Robertson, D)

Anthracite region, flood control (M. J. Bergin, W)

Bituminous coal resources (E. D. Pattersen, W)

Devonian stratigraphy of State (G. W. Colton, W)

Allentown Northeast quadrangle (J. M. Aaron, W)

Delaware River basin:

Lower part (J. P. Owens, W)

Middle part (A. A. Drake, Jr., W)

Philadelphia district, Lower Cambrian (J. H. Wallace, W)

Shenango quadrangle, Mercer County (G. R. Schiner, g, Harrisburg)

Southern anthracite field (G. H. Wood, Jr., W)

Stoneboro quadrangle, Mercer County (G. R. Schiner, g, Harrisburg)

Western Middle anthracite field (H. Arndt, W)

Washington County (H. Berryhill, Jr., D)

Wind Gap and adjacent quadrangles (J. B. Epstein, W)

Puerto Rico (W. H. Monroe, San Juan)

Rhode Island:

Ashaway quadrangle:

Bedrock geology (T. G. Feininger, Boston, Mass.)

Surficial geology (J. P. Schafer, Boston, Mass.)

Carolina quadrangle, surficial geology (J. P. Schafer, Boston, Mass.)

Chepachet quadrangle, bedrock geology (A. W. Quinn, Providence)

Clayville quadrangle, bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Coventry Center quadrangle, bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Crompton quadrangle, bedrock geology (A. W. Quinn, Providence)

Kingston quadrangle, bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Newport quadrangle, bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Prudence Island quadrangle, bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Quonochontaug quadrangle, surficial geology (J. P. Schafer, Boston, Mass.)

Thompson quadrangle (P. M. Hanshaw, H. R. Dixon, Boston, Mass.)

Tiverton quadrangle, bedrock geology (A. W. Quinn, Providence)

Voluntown quadrangle (T. G. Feininger, Boston, Mass.)

Watch Hill quadrangle:

Bedrock geology (G. E. Moore, Jr., Columbus, Ohio)

Surficial geology (J. P. Schafer, Boston, Mass.)

Wickford quadrangle, bedrock geology (R. B. Williams, Lawrence, Kans.)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

South Dakota:

Black Hills, southern (G. B. Gott, D)

Fort Randall Reservoir area (D. J. Varnes, D)

Four Corners quadrangle (J. A. Van Lieu, Laramie, Wyo.)

Harding County and adjacent areas (G. N. Pipirinos, D)

Hill City pegmatite area (J. C. Ratté, D)

Keystone pegmatite area (J. J. Norton, W)

Rapid City area (E. Dobrovolsky, D)

Tennessee:

Eastern part, zinc studies (A. L. Brokaw, D)

Appalachian folded belt, southern part (L. D. Harris, W)

Ducktown district and adjacent areas (R. M. Herndon, D)

Great Smoky Mountains (J. B. Hadley, W)

Ivydell quadrangle (K. J. Englund, W)

Jellico West quadrangle (K. J. Englund, W)

Ketchen quadrangle (K. J. Englund, W)

Knoxville and vicinity (J. M. Cattermole, Columbia, Ky.)

Midway belt, western Tennessee (D. R. Rima, w, Chattanooga)

Mount Rogers area (D. W. Rankin, W)

Pioneer quadrangle (K. J. Englund, W)

Texas:

Coastal plain, geophysical and geological studies (D. H. Eargle, Austin)

North-central part, Pennsylvanian Fusulinidae (D. A. Myers, D)

Del Rio area (V. L. Freeman, D)

Franklin Mountains (R. L. Harbour, D)

San Antonio and vicinity (R. D. Miller, D)

Sierra Blanca area (J. F. Smith, Jr., D)

Sierra Diablo region (P. B. King, M)

Utah:

Coal-mine bumps (F. W. Osterwald, D)

Alta quadrangle (M. D. Crittenden, Jr., M)

Bingham Canyon district (R. J. Roberts, M)

Circle Cliffs area (E. S. Davidson, Tucson, Ariz.)

Confusion Range (R. K. Hose, M)

Crawford Mountains (W. C. Gere, c, D)

Garrison quadrangle (D. H. Whitebread, M)

Gilbert Peak 1 NE quadrangle (J. R. Dyni, c, D)

Green River valley, upper part (W. R. Hansen, D)

Gunsight Butte quadrangle (F. Peterson, c, D)

Hurricane fault, southwestern Utah (P. Averitt, D)

Jessen Butte quadrangle (J. R. Dyni, c, D)

Kaiparowits Peak 2 SE quadrangle (W. E. Bowers, c, D)

Kaiparowits Peak 4 quadrangle (H. D. Zeller, c, D)
Kolob Terrace coal field, southern part (W. B. Cashion, D)

La Sal area (W. D. Carter, W)

Lehi quadrangle (M. D. Crittenden, Jr., M)

Lisbon Valley area (G. W. Weir, Berea, Ky.)

Little Cottonwood area (G. M. Richmond, D)

Moab-Interriver area (E. N. Hinrichs, D)

Morgan quadrangle (T. E. Mullens, c, D)

Navajo Reservation (R. B. O'Sullivan, D)

Nipple Butte quadrangle (H. A. Waldrop, c, D)

Oak City area (D. J. Varnes, D)

Ogden 1 SE quadrangle (T. E. Mullens, c, D)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued
Utah--Continued

Ogden 4 NE quadrangle (T. E. Mullens, c, D)
 Ogden 4 NW quadrangle (R. J. Hite, c, D)
 Orange Cliffs area (F. A. McKeown, D)
 Park City area (M. D. Crittenden, Jr., M)
 Park City district (C. S. Bromfield, D)
 Phil Pico Mountain quadrangle (J. R. Dyni, c, D)
 Promontory Point (R. B. Morrison, D)
 Sage Plain area (L. C. Huff, Manila, P.I.)
 Salt Lake City and vicinity (R. Van Horn, D)
 San Francisco Mountains (D. M. Lemmon, M)
 San Rafael Swell (C. C. Hawley, D)
 Sheeprock Mountains, West Tintic district (H. T. Morris, M)
 Snake Range, Wheeler Peak and Garrison quadrangles (D. H. Whitebread, M)
 Strawberry Valley (A. A. Baker, W)
 Tintic lead-zinc district, eastern (H. T. Morris, M)
 Uinta Basin, oil shale (W. B. Cashion, D)
 Wasatch Mountains (A. A. Baker, W)
 Wheeler Peak quadrangle (D. H. Whitebread, M)

Vermont:

North-central part (W. M. Cady, D)
 Rowe and Heath quadrangles (A. H. Chidester, D;
 J. H. Hartshorn, Boston, Mass.)

Virginia:

Appalachian folded belt, southern part (L. D. Harris, W)
 Washington, D. C., metropolitan area (H. W. Coulter, C. F. Withington, W)
 Big Stone Gap district (R. L. Miller, W)
 Herndon quadrangle (R. E. Eggleton, Flagstaff, Ariz.)
 Mount Rogers area (D. W. Rankin, W)

Washington:

Bodie Mountain quadrangle (R. C. Pearson, D)
 Chewelah 1 quadrangle (L. D. Clark, M)
 Cumberland quadrangle (J. D. Vine, M)
 Glacier Peak quadrangle (D. F. Crowder, M)
 Grays Harbor basin, western part (H. C. Wagner, M)
 Grays River quadrangle (E. W. Wolfe, M)
 Hobart quadrangle (J. D. Vine, M)
 Holden quadrangle (F. W. Cater, D)
 Hunters quadrangle (A. B. Campbell, D)
 Inchelium quadrangle (A. B. Campbell, D)
 Loomis quadrangle (C. D. Rinehart, M)
 Lucerne quadrangle (F. W. Cater, D)
 Maple Valley, Hobart and Cumberland quadrangles (J. D. Vine, M)
 Mt. Spokane quadrangle (A. E. Weissenborn, Spokane)
 Olympic Peninsula:
 Eastern part (W. M. Cady, D)
 Northern part (R. D. Brown, Jr., W)
 Puget Sound Basin (D. R. Crandell, D)
 Republic-Curlew area (R. L. Parker, D)
 Seattle and vicinity (D. R. Mullineaux, D)
 Stevens County (R. G. Yates, M)
 Togo Mountain quadrangle (R. C. Pearson, D)
 Twin Lakes quadrangle (G. E. Becraft, D)
 Wilnot Creek quadrangle (G. E. Becraft, D)

Geologic mapping--Continued

Map scale 1 inch to 1 mile, and larger--Continued

Wisconsin, Florence County (C. E. Dutton, Madison)
Wyoming:

Arlington quadrangle (H. J. Hyden, c, D)
 Atlantic City district (R. W. Bayley, M)
 Baggs area (G. E. Prichard, D)
 Beartooth Butte quadrangle (W. G. Pierce, M)
 Bengough Hill quadrangle (H. J. Hyden, c, D)
 Bosler quadrangle (H. McAndrews, c, Salt Lake City, Utah)
 Bradley Peak quadrangle (R. W. Bayley, M)
 Clark quadrangle (W. G. Pierce, M)
 Cokeville quadrangle (W. W. Rubey, Los Angeles, Calif.)
 Crowheart Butte area (J. F. Murphy, D)
 Deep Lake quadrangle (W. G. Pierce, M)
 Devil Slide quadrangle (E. K. Maughan, D)
 Devils Tooth quadrangle (W. G. Pierce, M)
 Ferris quadrangle (R. L. Rioux, c, W)
 Fish Lake quadrangle (W. L. Rohrer, c, D)
 Fossil basin (J. I. Tracey, Jr., W)
 Four Corners quadrangle (J. A. Van Lieu, Laramie)
 Gas Hills district (H. D. Zeller, D)
 Grand Teton National Park (J. D. Love, Laramie)
 Jackson (30-minute) quadrangle (D. A. Jobin, c, D)
 Kisinger Lakes quadrangle (W. L. Rohrer, c, D)
 LaBarge 1 SW and 2 SE quadrangles (R. L. Rioux, c, W)
 Lake Ione quadrangle (H. McAndrews, c, Salt Lake City, Utah)
 Lamont-Baroil area (M. W. Reynolds, D)
 McFadden quadrangle (H. J. Hyden, c, D)
 Oil Mountain quadrangle (W. H. Laraway, c, Casper)
 Pat O'Hara quadrangle (W. G. Pierce, M)
 Pierce Reservoir quadrangle (H. J. Hyden, c, D)
 Pilot Knob quadrangle (W. L. Rohrer, c, D)
 Poison Spider quadrangle (W. H. Laraway, c, Casper)
 Reid Canyon quadrangle (W. H. Laraway, c, Casper)
 Rock River quadrangle (H. J. Hyden, c, D)
 Shirley Basin area (E. N. Harshman, D)
 Spence-Kane area (R. L. Rioux, c, W)
 Square Top Butte quadrangle (W. H. Laraway, c, Casper)
 Sweetwater County, Green River Formation (W. C. Culbertson, D)
 T-L Ranch quadrangle (H. J. Hyden, c, D)
 Taylor Mountain quadrangle (M. L. Schroeder, c, D)
 Teepee Creek quadrangle (I. J. Witkind, D)
 Wapiti quadrangle (W. G. Pierce, M)
 Wedding of Waters quadrangle (E. K. Maughan, D)
 Whalen-Wheatland area (L. W. McGrew, Laramie)
 White Rock Canyon quadrangle (H. J. Hyden, c, D)
 Wind River Basin, regional stratigraphy (W. R. Keefer, Laramie)
 Wind River Mountains, Quaternary geology (G. M. Richmond, D)

Subsurface:

Alabama, hydrogeologic study (W. J. Powell, w, Tuscaloosa)

Geomorphology:

- Clays, erosion characteristics (A. V. Jopling, w, Boston, Mass.)
- Erosion and resultant landform changes, basic processes (G. G. Parker, h, D)
- Fluvial morphology, effect of sediment characteristics (S. A. Schumm, h, D)
- Geomorphology and hydrology, basic research (C. W. Carlston, w, W)
- Hillslope erosion, study of (S. A. Schumm, h, D)
- Mathematical geomorphology (A. E. Scheidegger, h, Urbana, Ill.)
- Mudflow studies (D. R. Crandell, D)
- Relation of drainage networks and basin development to rock type and climate (R. F. Hadley, h, D)
- Slope morphology, effect of exposure (R. F. Hadley, h, D)
- Soil creep, mechanisms (R. L. Schiffman, w, Troy, N.Y.)
- Stream morphology and processes (R. K. Fahnestock, h, D)
- Colorado River, geologic history (C. B. Hunt, w, Baltimore, Md.)
- Ohio River valley, geologic development (L. L. Ray, W)
- Alabama, Russell Cave (J. T. Hack, W)
- Arizona, Tusayan Washes, study of channel and flood-plain aggradation (R. F. Hadley, h, D)
- California:
- Death Valley, morphologic changes on alluvial fans (L. K. Lustig, w, Sacramento)
 - Sierra Nevada, geomorphic studies (R. J. Janda, w, M)
 - White Mountains, denudation rates (V. C. LaMarche, Jr., w, M)
- Indiana, Owensboro quadrangle, Quaternary geology (L. L. Ray, W)
- Iowa, channel-geometry studies (H. H. Schwob, s, Iowa City)
- Kentucky, Owensboro quadrangle, Quaternary geology (L. L. Ray, W)
- Massachusetts, sea-cliff erosion studies (C. A. Kaye, Boston)
- Michigan, Lake Algonquin drainage (J. T. Hack, W)
- Montana, Browning area, Quaternary geology (G. M. Richmond, D)
- New Mexico, Santa Fe, particle movement and channel scour and fill of an ephemeral arroyo (L. B. Leopold, w, W)
- New York, northeast Adirondacks (C. S. Denny, W)
- North Carolina:
- Stream-channel characteristics and time-of-travel studies (N. O. Thomas, H. G. Hinson, w, Raleigh)
 - Morganton area (J. T. Hack, W)
- North Dakota, hydrology of prairie potholes (W. S. Eisenlohr, Jr., h, D)
- Wyoming, Wind River Mountains, Quaternary geology (G. M. Richmond, D)

See also Sedimentation.

Geophysics, regional:

- Aeroradioactivity surveys:
- Northeastern United States (P. Popenoe, W)
- California:
- San Andreas fault (J. H. Healy, D)
 - San Francisco (J. A. Pitkin, W)
- Colorado, Rocky Flats (J. A. MacKallor, Cotonou, Dahomey)
- Idaho, National Reactor Testing Station (R. G. Bates, W)

Geophysics, regional--Continued

- Aeroradioactivity surveys--Continued
- Illinois, Chicago (G. M. Flint, Jr., W)
 - Maryland, Belvoir area (S. K. Neuschel, W)
 - Minnesota, Elk River (J. A. Pitkin, W)
 - Ohio, Columbus (R. G. Bates, W)
 - Pennsylvania, Pittsburgh (R. W. Johnson, Knoxville, Tenn.)
 - Puerto Rico (J. A. Pitkin, W)
 - Texas, Fort Worth (J. A. Pitkin, W)
 - Virginia, Belvoir area (S. K. Neuschel, W)
- Cross-country aeromagnetic profiles (E. R. King, W)
- Crust and upper mantle:
- Analysis of traveltime data (J. C. Roller, D)
 - Geophysical studies (J. H. Healy, D)
 - Gravity surveying (D. J. Stuart, D)
 - Rocky Mountain seismic network (J. P. Eaton, D)
 - Seismic-refraction profiling (W. H. Jackson, D)
- Ultramafic intrusions, geophysical studies (G. A. Thompson, M)
- Antartica, Pensacola Mountains, geophysical studies (J. C. Behrendt, W)
- Arctic, geophysical studies (I. Zietz, W)
- Central United States, aeromagnetic surveys (J. W. Henderson, W)
- Colorado Plateau, regional geophysical studies (H. R. Joesting, W)
- Colorado Plateau and southern Rocky Mountains, aeromagnetic surveys (H. R. Joesting, W)
- Costa Rica, volcanic studies (K. J. Murata, San Jose, Costa Rica)
- Eastern Central United States, tectonic patterns (I. Zietz, W)
- Eastern United States, aeromagnetic surveys (R. W. Bromery, W)
- Gravity map of the United States (H. R. Joesting, W)
- Japan, calderas, aeromagnetic-gravity studies (H. R. Blank, Jr., M)
- Lake Superior region, geophysical studies (G. D. Bath, M)
- New England, geophysical studies (R. W. Bromery, W)
- Northeastern United States, gravity study (G. Simmons, Dallas, Tex.)
- Pacific Northwest:
- Aeromagnetic surveys (W. E. Davis, M)
 - Geophysical studies (W. E. Davis, M)
- Pacific Ocean, geophysical studies (D. F. Barnes, M)
- Pacific Southwest:
- Aeromagnetic surveys (D. R. Mabey, M)
 - Geophysical studies (D. R. Mabey, M)
- Pacific States, geophysical studies (A. Griscom, M)
- Tri-State eruptive-tectonic complex, Wyoming-Montana-Idaho, geophysical study (H. R. Blank, M)
- Alaska:
- Aeromagnetic surveys (G. E. Andreasen, W)
 - Regional gravity surveys (D. F. Barnes, M)
- Arizona:
- Central part, geophysical study (D. R. Mabey, M)
 - Safford Valley, geophysical studies (G. E. Andreasen, W)
 - Tombstone region, geophysical studies (G. E. Andreasen, W)
- California:
- Coast Range and Sacramento Valley, geophysical studies (G. D. Bath, M)

Geophysics, regional--Continued**California--Continued**

Los Angeles basin, gravity study (T. H. McCulloh, Riverside)

San Francisco Bay area, geophysical studies (G. D. Bath, M)

Sierra Nevada, geophysical studies (H. W. Oliver, M)

Colorado:

Arkansas Valley, geophysical study (J. E. Case, D)

Cheyenne Mountain, electrical properties (J. H. Scott, D)

Middle Park - North Park basins, geophysical studies (J. C. Behrendt, D)

Uncompahgre uplift, northwest portion, geophysical studies and geologic mapping (J. E. Case, D)

District of Columbia, Eastern Piedmont, geophysical studies (S. K. Neuschel, W)

Idaho, Snake River Plain, geophysical studies (D. R. Mabey, D)

Iowa, central, aeromagnetic survey (J. R. Henderson, W)

Maine:

Island Falls quadrangle, electromagnetic mapping (F. C. Frischknecht, W)

Stratton quadrangle, geophysical and geologic mapping (A. Griscom, W)

Maryland, Montgomery County, geophysical studies (A. Griscom, W)

Massachusetts:

Application of geology and seismology to public-works planning (C. R. Tuttle, R. N. Oldale, Boston)

Geophysical studies (R. W. Bromery, W)

Michigan:

Gogebic district, aeromagnetic study (J. E. Case, D)

Marquette district, aeromagnetic study (J. E. Case, D)

Minnesota, southern, aeromagnetic survey (I. Zietz, W)

Mississippi, Tatum salt dome (W. S. Twenhofel, D)

Missouri, southeast, aeromagnetic study (J. W. Allingham, W)

Montana:

Bearpaw Mountains, aeromagnetic study (K. G. Books, W)

Boulder batholith, aeromagnetic and gravity studies (W. E. Davis, M)

Nevada:

Central part, geophysical studies (D. R. Mabey, D)

Armagosa Desert, gravity surveys (R. G. Bates, D)

Clark County, gravity investigations (M. F. Kane, W)

Nevada Test Site, aeromagnetic surveys (J. W. Allingham, W)

New Jersey:

Gettysburg-Newark Basin, geophysical investigations (M. E. Beck, W)

New York-New Jersey Highlands, aeromagnetic studies (A. Jespersen, W)

New Mexico, Valles caldera, geophysical study (H. R. Joesting, W)

New York:

Adirondacks area, aeromagnetic studies (J. R. Balsley, Middletown, Conn.)

New York-New Jersey Highlands, aeromagnetic studies (A. Jespersen, W)

North Carolina, Concord quadrangle, geophysical studies (R. G. Bates, W)

Ohio, seismic survey for buried valleys (J. S. Watkins, Flagstaff, Ariz.)

Geophysics, regional--Continued

Oregon Cascades, geophysical study (H. R. Blank, M)

Pennsylvania:

Gettysburg-Newark Basin, geophysical investigations (M. E. Beck, W)

Gravity survey (R. W. Bromery, W)

Triassic area, aeromagnetic study (R. W. Bromery, W)

Puerto Rico, geophysical studies (A. Griscom, W)

South Dakota, Hills area, regional gravity studies (R. M. Hazelwood, D)

Texas, coastal plain, geophysical and geological studies (D. H. Eargle, Austin)

Utah:

Iron Springs, aeromagnetic survey (H. R. Blank, M)

Sheeprock Mountains, West Tintic district (D. R. Mabey, D)

Uncompahgre uplift, northwest portion, geophysical studies and geologic mapping (J. E. Case, D)

Washington:

Northeastern part, geophysical studies (W. T. Kinoshita, M)

Western part, gravity survey (D. J. Stuart, D)

Wisconsin, Florence County, aeromagnetic study (E. R. King, W)

Wyoming:

Anchor Reservoir, gravity survey (G. P. Eaton, D)

Black Hills area, regional gravity studies (R. A. Black, W)

Bradley Peak area (D. R. Mabey, D)

Jackson Hole region, geophysical studies (D. R. Mabey, D)

Mowry Shale and Frontier Formation, geophysical studies (G. P. Eaton, D)

Geophysics, theoretical and experimental:

Borehole geophysics as applied to geohydrology (W. S. Keys, h, D)

Earthquake - explosion effects (V. R. Wilmarth, W)

Earthquakes, local seismic studies (J. P. Eaton, D)

Elastic and inelastic properties of earth materials (L. Peselnick, W)

Electric and magnetic properties of minerals (A. N. Thorpe, W)

Electrical methods, development (C. J. Zablocki, D)

Electrical properties of rocks (G. V. Keller, D)

Electromagnetic radiation studies (W. A. Fischer, W)

Geophysical data, interpretation using electronic computers (R. G. Henderson, W)

Geothermal studies (A. H. Lachenbruch, M)

Gravity and magnetic anomalies, analysis (W. H. Diment, W)

Heat flow in the Appalachian Mountains (W. H. Diment, W)

Heat transfer in salt (E. C. Robertson, W)

Infrared and ultraviolet radiation studies (R. M. Moxham, W)

Magnetic and luminescent properties (F. E. Senftle, W)

Magnetic model studies (G. E. Andreasen, W)

Magnetic properties of crystals (A. N. Thorpe, W)

Magnetic properties of rocks (A. Griscom, W)

Radon, geologic behavior (A. B. Tanner, W)

Remanent magnetization of rocks (R. R. Doell, M)

Rock behavior at high temperature and pressure (E. C. Robertson, W)

Tension fractures and thermal investigation (A. H. Lachenbruch, M)

Geophysics, theoretical and experimental--Continued

- Thermodynamic properties of rocks (R. A. Robie, W)
- Ultramafic intrusions, geophysical studies (G. A. Thompson, M)

Glacial geology:

- Alaska, glacial map (H. W. Coulter, W)
- Antarctica, Pensacola Mountains (D. L. Schmidt, W)
- California, west-central Sierra Nevada (F. M. Fryxell, Rock Island, Ill.)
- Greenland, Schuchert Dal (J. S. Hartshorn, Boston, Mass.)

Glaciology:

- Glaciological research (M. F. Meier, w, Tacoma, Wash.)
- Alaska, Barrier Glacier (Mount Spurr) (G. C. Giles, c, Tacoma, Wash.)
- Montana, Glacier National Park:
 - Grinnell and Sperry Glaciers (A. Johnson, c, W)
 - Grinnell Glacier, hydrology (F. Stermitz, s, Helena)
- Washington, Mount Rainier National Park, Emmons and Nisqually Glaciers (G. C. Giles, c, Tacoma)

Gold:

- Geochemistry and occurrence (J. C. Antweiler, D)
- Gold deposits, United States (M. H. Bergendahl, D)
- Alaska:
 - Nome C-1 and D-1 quadrangles (C. L. Hummel, Bangkok, Thailand)
 - Tofty placer district (D. M. Hopkins, M)
- Colorado, Tenmile Range and Kokomo mining district (M. H. Bergendahl, D)
- Wyoming, Atlantic City district (R. W. Bayley, M)

Ground water—surface water relations:

- Bank-seepage studies (E. C. Pogge, s, Iowa City, Iowa)
- Flow losses in ephemeral stream channels (R. F. Hadley, h, D)
- Ground water — surface water interrelations (M. W. Busby, s, Topeka, Kans.)
- Streamflow in relation to aquifer characteristics (G. R. Kunkle, w, W)
- Florida, Lake Okeechobee, levee underseepage (F. W. Meyer, w, Tallahassee)
- Montana, Hungry Horse Reservoir, bank storage (A. F. Bateman, Jr., c, Great Falls)
- New Jersey, Ramapo River basin (J. Vecchioli, g, Trenton)
- New Mexico, White Sands Missile Range, research on paving a small watershed (W. C. Ballance, g, Albuquerque)
- Washington:
 - Cedar River loss study, surface and ground water (F. T. Hidaka, w, Tacoma)
 - Columbia River basin, relation of ground-water storage and streamflow (M. I. Rorabaugh, w, Tacoma)
- Tennessee, Upper Buffalo River (W. J. Perry, w, Chattanooga)
- Wisconsin, central Sand Plains, hydrology (E. P. Weeks, g, D. W. Ericson, s, Madison)

Hydraulics, ground water:

- Aquifer characteristics, determination by analysis of water-level data in an aquifer and an adjoining clogged stream (A. G. Law, h, Clemson, S. C.)
- Aquifer-test reevaluation, California (E. J. McClelland, w, Sacramento)
- Crystalline rocks, occurrence of ground water, Piedmont area of Maryland (E. G. Otton, w, Towson)
- Dielectric behavior of water-bearing sediments (W. O. Smith, w, W)

Hydraulics, ground water--Continued

- Fluid dynamics of the Bandelier Tuff, New Mexico (F. C. Koopman, g, Albuquerque)
- Geohydrologic environmental studies (J. N. Payne, h, Baton Rouge, La.)
- Ground-water mechanics, treatise (J. G. Ferris, w, Tucson, Ariz.)
- Limestone terranes, predicting well yields (G. K. Moore, w, Chattanooga, Tenn.)
- Mechanics of aquifers — principles of compaction and deformation (J. F. Poland, w, Sacramento, Calif.)
- Mechanics of fluid flow in porous media (A. Ogata, h, Honolulu, Hawaii)
- Mechanics of ground-water flow (H. H. Cooper, Jr., w, W)
- Permeability distribution study—Atlantic Coastal Plain (P. M. Brown, w, Raleigh, N.C.)
- Permeability of fractured rocks (F. W. Trainer, w, W)
- Permeability research, California (A. I. Johnson, g, D)
- Prototype electrical analog model of the ground-water system of western Long Island, N.Y. (N. M. Perlmutter, w, Albany)
- Regional hydrologic system analysis—hydrodynamics (R. R. Bennett, w, W)
- Regional hydrologic system analysis—permeability distribution (J. D. Bredehoeft, w, W)
- Research on laboratory and field methods (A. I. Johnson, g, D)
- Theory of multiphase flow — applications (R. W. Stallman, h, D)
- Transient flow in sediments (W. O. Smith, w, W)
- Unsaturated flow of water in sediments (W. O. Smith, w, W)
- Unsaturated-flow theory related to drainage and infiltration (J. Rubin, w, M)

Hydraulics, surface flow:

- Channel characteristics:
 - Large-scale roughness (J. Davidian, w, W)
 - Manning coefficient, determination from measured bed roughness in natural channels (L. E. Young, w, M)
 - Sand-channel streams, controls (F. A. Kilpatrick, w, W)
 - Stage-discharge relations from backwater profiles (H. A. Ray, w, M)
 - Vegetation and alluvial processes in a semiarid environment (R. G. Zimmermann, w, Baltimore, Md.)
- Channel constrictions:
 - Hydraulic factors, field measurement:
 - Performance of channel changes (P. O. Jefferson, w, Tuscaloosa, Ala.)
 - Performance of culverts (P. O. Jefferson, w, Tuscaloosa, Ala.)
 - Overall efficiency of bridges (K. V. Wilson, w, Jackson, Miss.)
 - Verification of hydraulic computation methods for bridge sites (C. O. Ming, w, Tuscaloosa, Ala.)
 - Verification of hydraulic techniques (W. J. Randolph, w, Chattanooga, Tenn.)
- Flow characteristics:
 - Dispersion by turbulent flow in open channels (N. Yotsukura, w, W)
 - Gaging streamflow through turbines (B. J. Frederick, w, Chattanooga, Tenn.)

Hydraulics, surface flow--Continued**Flow characteristics--Continued**

Mechanics of flow structure and fluid resistance--movable boundary (E. V. Richardson, h, Fort Collins, Colo.)

Unsteady flow in natural channels (R. A. Baltzer, w, W)

Vertical-velocity characteristics, Columbia River gaging stations, Washington and Oregon (J. Savini, G. L. Bodhaine, w, Tacoma, Wash.)

Laboratory studies:

Grain-size distribution and bedload transport (L. B. Leopold, G. Williams, w, W)

Laboratory studies of open-channel flow (H. J. Tracy, w, W)

Tests of crest-stage gage intake systems (J. Friday, s, Portland, Oreg.)

Time-of-travel studies:

Solutes (J. F. Wilson, Jr., w, W)

New Jersey (T. J. Buchanan, s, Trenton)

New York (B. Dunn, w, Albany)

Hydrologic-data collection and processing:

Automation systems and equipment for water (W. L. Isherwood, w, W)

Data-collection program, new criteria (M. A. Benson, w, W)

Data-processing methods, evaluation (A. I. Johnson, g, D)

Digital-computer method for computing sediment discharge (M. D. Edwards, w, Raleigh, N.C.)

Discharge determinations from a moving boat (T. J. Buchanan, s, Trenton, N.J.)

Drainage-area determinations:

Arkansas (R. C. Christensen, w, Little Rock)

Kentucky (H. C. Beaber, s, Louisville)

New Jersey, for gazetteer of streams (A. A. Vickers, s, Trenton)

South Carolina (B. H. Whetstone, s, Columbia)

Electrical analog model studies of areal hydrologic problems (J. D. Winslow, g, Lawrence, Kans.)

Extension of streamflow records (L. E. Carroon, s, D)

Hydrologic-data storage retrieval, and application by digital-computer techniques (C. O. Morgan, g, Lawrence, Kans.)

River-systems gaging (H. C. Riggs, w, W)

Sediment loads in streams--methods used in measurement and analysis (B. C. Colby, q, Minneapolis, Minn.)

Sediment manual (R. B. Vice, w, W)

Statistical inferences (N. C. Matalas, w, W)

Vigil Network Survey--observations of channel and slope processes (W. W. Emmett, L. B. Leopold, w, W)

Hydrologic instrumentation:

Acoustic velocity-measuring equipment--water (W. Hofmann, w, M)

Alluvial streams, controls and instrumentation for gaging (F. A. Kilpatrick, h, Fort Collins, Colo.)

Electronic-equipment development--water (J. E. Eddy, w, W)

Energy-budget evaporation studies, instruments (C. R. Daum, h, D)

Instrumentation research--water (H. O. Wires, w, Columbus, Ohio)

Laboratory research, instruments--water (G. F. Smoot, w, W)

Hydrologic instrumentation--Continued

Unstable flow in steep channels, instrumentation for study (W. Smith, w, M)

Hydrology, ground-water:

Aquifer systems, hydrogeology, Southeastern United States (V. T. Stringfield, w, W)

Aquifer test, Piketon, Ohio (S. E. Norris, R. E. Fidler, w, Columbus)

Geohydrologic environmental study (J. N. Payne, h, Baton Rouge, La.)

Gravity flow of water in soils and aquifers, western Kansas (R. C. Prill, g, Lawrence)

Problems in quantitative hydrology (M. I. Rorabaugh, w, Tacoma, Wash.)

Specific-yield studies, California (A. I. Johnson, g, D)

Hydrology, surface-water:

Changes in regimen, Santa Ana River, Calif. (M. B. Scott, w, Los Angeles)

Characteristics of flood hydrographs, Kansas (L. W. Furness, s, Topeka)

Flood-frequency synthesis for small streams, Alabama (L. B. Peirce, w, Tuscaloosa)

Flow probability, New Jersey streams (E. G. Miller, s, Trenton)

Hydrologic and hydraulic characteristics, Wragg Swamp canal (Montlemer Creek), Alabama (J. F. McCain, w, Tuscaloosa)

Hydrologic effects of small reservoirs, Sandstone Creek, Okla. (F. W. Kennon, w, Austin, Tex.)

Hydrology of small streams, New Hampshire (C. E. Hale, w, Boston, Mass.)

Lake mapping and stabilization, Indiana (D. C. Perkins, w, Indianapolis)

Long-term chronologies of hydrologic events (W. D. Simons, w, Tacoma, Wash.)

Peak inflow and outflow through ponds (J. E. McCall, s, Trenton, N.J.)

Rates of runoff from small rural watersheds, Alabama (L. B. Peirce, w, Tuscaloosa)

Surface-water hydrology of coastal basins, north of San Francisco to Eel River, Calif. (S. E. Rantz, w, M)

Unit-graph characteristics, Kansas (I. C. James, s, Topeka)

Variations in streamflow, Utah (G. L. Whitaker, w, Salt Lake City)

Variations in streamflow due to earthquakes, Utah (W. N. Jibson, w, Salt Lake City)

Water-quality and streamflow characteristics, Passaic River basin, New Jersey (P. W. Anderson, q, Trenton)

Industrial minerals:

Ultramafic rocks of the Southeast (D. M. Larrabee, W)
See also specific minerals.

Iron:

Clinton iron ores of the southern Appalachians (R. P. Sheldon, D)

Alaska, Klukwan iron district (E. C. Robertson, W)

Michigan:

Dickinson County, southern (R. W. Bayley, M)

Eust Marquette district (J. E. Gair, D)

Gogebic Range, eastern (W. C. Prinz, W)

Iron County, eastern (K. L. Wier, D)

Iron River-Crystal Falls district (H. L. James, Minneapolis, Minn.)

Iron--Continued**Michigan--Continued**

- Marenisco-Watersmeet area (C. E. Fritts, D)
- Negaunee and Palmer quadrangles (J. E. Gair, D)
- Montana, southwestern (K. L. Wier, D)
- Tennessee, Ducktown district and adjacent areas (R. M. Hernon, D)

Wisconsin, Florence County (C. E. Dutton, Madison)**Wyoming:**

- Atlantic City district (R. W. Bayley, M)
- Bradley Peak quadrangle (R. W. Bayley, M)

Isotope and nuclear studies:

- Instrument development (J. S. Stacey, D)
- Isotope geology of lead (A. P. Pierce, D)
- Isotope ratios in rocks and minerals (I. Friedman, W)
- Isotopic hydrology (G. L. Stewart, w, W)
- Isotopic studies of crustal processes (B. Doe, W)
- Isotopic studies of upper mantle (M. Tatsumoto, D)
- Light stable isotopes (I. Friedman, W)
- Nuclear irradiation (C. M. Bunker, D)
- Ore lead, geochemistry and origin (R. S. Cannon, D)
- Oxygen-isotope geothermometry (H. L. James, Minneapolis, Minn.)
- Tritium concentrations in precipitation, surface water, and ground water, coastal plain of New Jersey (E. C. Rhodehamel, g, Trenton)

See also Geochronology.

Lake levels:

- Elevations of Great Salt Lake, Utah (H. W. Chase, w, Salt Lake City)

Land subsidence:

- California, San Joaquin Valley (J. F. Poland, w, Sacramento)

Lead and zinc:

- Ore lead, geochemistry and origins (R. S. Cannon, D)
- Zinc resources of the world (T. H. Killsgaard, W)
- Arizona, Lochiel and Nogales quadrangles (F. S. Simons, D)
- California, Panamint Butte quadrangle (W. E. Hall, W)
- Colorado, Rico district (E. T. McKnight, W)
- Idaho, Coeur d'Alene mining district (S. W. Hobbs, D)
- Kansas, Picher lead-zinc district (E. T. McKnight, W)
- Missouri:
 - Picher lead-zinc district (E. T. McKnight, W)
 - Southeastern Missouri lead district (T. H. Killsgaard, W)

Nevada:

- Ely district (A. L. Brokaw, D)
- Pioche district (C. M. Tschanz, La Paz, Bolivia)
- New Mexico, Silver City area (W. R. Jones, D)
- Oklahoma, Picher lead-zinc district (E. T. McKnight, W)

Tennessee:

- Eastern part (A. L. Brokaw, D)
- Origin and depositional control of selected zinc deposits (H. Wedow, Jr., Knoxville, Tenn.)

Utah:

- East Tintic lead-zinc district (H. T. Morris, M)
- Park City district (C. S. Bromfield, D)
- West Tintic district, Sheeprock Mountains (H. T. Morris, M)
- Virginia, origin and depositional control of selected zinc deposits (H. Wedow, Jr., Knoxville, Tenn.)

Limestone-terrane hydrology (F. A. Swenson, h, D)**Limnology:**

- Chemical hydrology, Great Salt Lake, Utah (D. C. Hahl, w, Salt Lake City)

Limnology--Continued

- Hydrology and geochemistry of topographically closed lakes, south-central Oregon (K. N. Phillips, s, Portland)

- Organisms, effect on water quality of streams (K. V. Slack, w, W)

- Thermal and biological characteristics of lakes, Indiana (J. F. Ficke, w, Fort Wayne)

Low flow and flow duration:

- Alabama, low-flow analyses of streams (L. B. Peirce, w, Tuscaloosa)

- Arkansas, low-flow frequency studies (M. S. Hines, w, Little Rock)

Illinois:

- Low-flow frequency analyses (W. D. Mitchell, s, Champaign)

- Low-flow partial-record investigation (W. D. Mitchell, s, Champaign)

Iowa:

- Low-flow frequency studies (H. H. Schwob, s, Iowa City)

- Origin of base flow for small drainage basins (G. R. Kunkle, E. C. Pogge, s, Iowa City)

Kansas:

- Low-flow data collection (T. J. Irza, s, Topeka)

- Seepage flow of Kansas streams (M. W. Busby, s, Topeka)

- Massachusetts, low-flow characteristics (G. K. Wood, w, Boston)

- Mississippi, low-flow characteristics of streams (C. P. Humphreys, Jr., w, Jackson)

- Missouri, low-flow characteristics (J. S. Skelton, s, Rolla)

New York:

- Low-flow analysis for stream classification (O. P. Hunt, w, Albany)

- Low-flow frequency (O. P. Hunt, w, Albany)

- New Jersey, low-flow characteristics (E. G. Miller, s, Trenton)

Ohio:

- Flow duration (W. P. Cross, h, Columbus)

- Low-flow and storage requirements (W. P. Cross, w, Columbus)

- Pennsylvania, low-flow frequency analysis (W. F. Busch, s, Harrisburg)

- South Carolina, low-flow gaging (J. S. Stallings, W. M. Bloxham, s, Columbia)

- Tennessee, low-flow and flow-duration studies (J. S. Cragwall, Jr., w, Chattanooga)

- Texas, Little Cypress Creek, base flow, quantity and quality (J. H. Montgomery, w, Austin)

- Wisconsin, low-flow analyses (D. W. Ericson, s, Madison)

Lunar geology. See Extraterrestrial studies.

Manganese. See Ferro-alloy metals.

Marine geology:

- Atlantic Coastal Plain, regional synthesis (J. C. Maher, M)

- East coast continental shelf and margin (K. O. Emery, Woods Hole, Mass.)

- Gulf Coastal Plain, regional synthesis (J. C. Maher, M)

Marine hydrology:

- Effect of heated water, Patuxent River estuary, Maryland (R. L. Cory, w, W)

- Influence of industrial and municipal wastes on estuarine and offshore water quality, Washington (J. F. Santos, w, Tacoma)

Marine hydrology--Continued

- Minimum tides of the Delaware estuary (A. C. Lendo, s, Trenton, N.J.)
- Recording of maximum tides (J. A. Bettendorf, s, Trenton, N. J.)
- Tidal stage, discharge and velocity studies (A. C. Lendo, s, Trenton, N.J.)
- Zoogeography and ecology of offshore populations of amphioxus, Atlantic Continental Shelf of the United States (R. L. Cory, w, W)
- Washington, Willapa Bay project (A. O. Waananen, w, M)
- See also Sea-water intrusion.

Meteorites. See Extraterrestrial studies.**Mineral and fuel resources--compilations and topical studies:**

- Carbonate-rock resources (G. E. Ericksen, W)
- Drilling data, statistical techniques in the analysis of (H. Wedow, Knoxville, Tenn.)
- Energy resources of the United States (T. A. Hendricks, D)
- Massive sulfide deposits (A. R. Kinkel, Jr., W)
- Metallogenic maps, United States (P. W. Guild, W)
- Mineral exploration, Northwestern United States (D. R. MacLaren, Spokane, Wash.)
- Mineral fuel resources, United States (T. H. Kiilsgaard, W)
- Mineral-resource information and research (H. Kirkemo, W)
- Mineral-resource map, Utah (L. S. Hilpert, Salt Lake City, Utah)
- Mineral-resources appraisal, northern Wisconsin (C. E. Dutton, Madison)
- Oxygen isotope geothermometry (H. L. James, Minneapolis, Minn.)
- Resource data storage and retrieval (R. A. Weeks, W)
- Resource study techniques (R. A. Weeks, W)
- Uranium-bearing veins (G. W. Walker, D)
- Uranium deposits, formation and redistribution (K. G. Bell, D)
- Zinc deposits, origin and depositional control, Tennessee and Virginia (H. Wedow, Jr., Knoxville, Tenn.)
- Zoning of mineral deposits (D. A. Gallagher, M)
- See also specific minerals or fuels.

Mineralogy and crystallography, experimental:

- Crystal chemistry (H. T. Evans, Jr., W)
 - Borate minerals (J. R. Clark, C. L. Christ, W)
 - Phosphate minerals (M. E. Mrose, W)
 - Rock-forming silicate minerals (D. E. Appleman, W)
 - Uranium minerals (H. T. Evans, W)
- Mineralogic services and research (M. L. Lindberg, W)
- New minerals (D. E. Appleman, W)
- New minerals--micas and chlorites (M. D. Foster, W)
- Petrological services and research (C. Milton, W)
- Sedimentary mineralogy (P. D. Blackmon, D)
- See also Geochemistry, experimental.

Mining hydrology:

- Mining hydrology (W. T. Stuart, w, W)
- Study of the hydrologic and related effects of strip mining, Beaver Creek watershed, Kentucky (J. J. Musser, w, Columbus, Ohio)

Minor elements:

- Black shale (J. D. Vine, M)
- Dispersion pattern of minor elements related to igneous intrusions (W. R. Griffiths, D)
- Geochemistry (G. Phair, W)

Minor elements--Continued**Niobium:**

- Colorado, Wet Mountains (R. L. Parker, W)
- Phosphoria Formation, stratigraphy and resources (R. A. Gulbrandsen, M)
- Rare-earth elements, resources and geochemistry (J. W. Adams, D)
- Sedimentary rocks, mineral fractionation and fine-grained trace element content (T. D. Botinelly, D)
- Selenium resources and geochemistry (D. F. Davidson, D)
- Tantalum-niobium resources of the United States (R. L. Parker, W)
- Trace-analysis methods:
 - Development (H. W. Lakin, D)
 - Research (F. N. Ward, D)
- Volcanic rocks (R. R. Coats, M)

Model studies, hydrologic:

- Analog analysis of the hydrology of the Blue River basin, Nebraska (P. A. Emery, w, Lincoln)
- Analytical model of the land phase of the hydrological cycle (D. R. Dawdy, w, W)

Molybdenum. See Ferro-alloy metals.**Monazite:**

- Geology of monazite (W. C. Overstreet, Jidda, Saudi Arabia)
- Southeastern United States (W. C. Overstreet, Jidda, Saudi Arabia)

Moon studies. See Extraterrestrial studies.**Nickel.** See Ferro-alloy metals.**Nuclear explosions, hydrology:**

- Potential applications of nuclear explosives in development and management of water resources (A. M. Piper, F. W. Stead, w, M)
- Project CHARIOT, hydrology (A. M. Piper, w, M)
- Mississippi, Tatum salt dome area, water-resources evaluation (R. E. Taylor, w, Jackson)
- Nevada Test Site, hydrologic studies (I. J. Winograd, w, Carson City)

Oil shale:

- Alaska, north slope of Brooks Range (I. L. Tailleux, M)
- Colorado:
 - State resources (D. C. Duncan, W)
 - Grand-Battlement Mesa (J. R. Donnell, D)
 - Rangely 3 quadrangle (H. L. Cullins, c, D)
- Utah, Uinta Basin (W. B. Cashion, D)
- Wyoming, Green River Formation, Sweetwater County (W. C. Culbertson, D)

Paleobotany, systematic:

- Diatom studies (K. E. Lohman, W)
- Fossil wood and general paleobotany (R. A. Scott, D)
- Floras:
 - Cenozoic, Western United States (J. A. Wolfe, M)
 - Devonian (J. M. Schopf, Columbus, Ohio)
 - Pennsylvanian, Illinois and adjacent States (C. B. Read, Albuquerque, N. Mex.)
 - Permian (S. H. Mamay, W)
- Plant microfossils:
 - Cenozoic (E. B. Leopold, D)
 - Mesozoic (R. H. Tschudy, D)
 - Paleozoic (R. M. Kosanke, D)

Paleoecology:

- Green River Formation, Wyoming, geology and paleolimnology (W. H. Bradley, W)
- Coal-ball studies, Pennsylvanian (S. H. Mamay, W)

Paleoecology--Continued

Diatoms (K. E. Lohman, W)
Faunas, Late Pleistocene, Pacific Northwest (W. O. Addicott, M)

Foraminifera:

Ecology (M. R. Todd, W)
Cenozoic, larger forms (K. N. Sachs, Jr., W)
Recent, eastern Pacific (P. J. Smith, M)

Mollusks:

Tertiary nonmarine, biogeography, Snake River Plain and adjacent areas (D. W. Taylor, M)
Pacific Islands, biogeography (H. S. Ladd, W)
Ostracodes, Recent, North Atlantic (J. E. Hazel, W)
Paleoenvironment studies, Miocene, Atlantic Coastal Plain (T. G. Gibson, W)
Pollen, Recent, distribution studies (E. B. Leopold, D)
Tempskya, Southwestern United States (C. B. Read, Albuquerque, N. Mex.)

Vertebrate faunas, Ryukyu Islands, biogeography (F. C. Whitmore, Jr., W)

Paleontology, invertebrate, systematic:**Brachiopods:**

Carboniferous (M. Gordon, Jr., W)
Ordovician (R. B. Neuman, W; R. J. Ross, Jr., D)
Permian (R. E. Grant, W)
Upper Paleozoic (J. T. Dutro, Jr., W)

Bryozoans:

Ordovician (O. L. Karklins, W)
Upper Paleozoic (H. M. Duncan, W)

Cephalopods:

Jurassic (R. W. Imlay, W)
Triassic (N. J. Silberling, M)
Upper Cretaceous (W. A. Cobban, D)
Upper Paleozoic (M. Gordon, Jr., W)

Chitinozoans, Lower Paleozoic (J. M. Schopf, Columbus, Ohio)

Conodonts, Paleozoic (J. W. Huddle, W)

Corals, rugose:

Mississippian (W. J. Sando, W)
Silurian-Devonian (W. A. Oliver, Jr., W)

Foraminifera:

Fusuline and orbitoline (R. C. Douglass, W)
Cenozoic (R. Todd, W)
Cenozoic, California and Alaska (P. J. Smith, M)
Cretaceous (R. Todd, W)
Mississippian (B. A. L. Skipp, D)
Pennsylvanian-Permian, fusuline (L. G. Henbest, W)
Tertiary, larger (K. N. Sachs, Jr., W)

Gastropods:

Mesozoic (N. F. Sohl, W)
Miocene-Pliocene, Atlantic Coast (T. G. Gibson, W)
Oligocene, Mississippi (F. S. MacNeil, M)
Paleozoic (E. L. Yochelson, W)

Graptolites, Ordovician-Silurian (R. J. Ross, Jr., D)

Mollusks:

Cenozoic (F. S. MacNeil, M)
Late Cenozoic, nonmarine (D. W. Taylor, M)

Ostracodes:

Cenozoic (J. E. Hazel, W)
Lower Paleozoic (J. M. Berdan, W)
Upper Paleozoic (I. G. Sohn, W)

Pelecypods:

Inoceramid (D. L. Jones, M)
Jurassic (R. W. Imlay, W)
Oligocene, Mississippi (F. S. MacNeil, M)

Paleontology, invertebrate, systematic--Continued**Pelecypods--Continued**

Paleozoic (J. Pojeta, Jr., W)
Triassic (N. J. Silberling, M)
Radiolaria (K. N. Sachs, Jr., W)

Trilobites:

Cambrian (A. R. Palmer, W)
Ordovician (R. J. Ross, Jr., D)

Paleontology, stratigraphic:**Cenozoic:**

Coastal Plains (D. Wilson, W)
Pacific Coast, Miocene (W. O. Addicott, M)
Diatoms:
California and Nevada (K. E. Lohman, W)
Great Plains, nonmarine (G. W. Andrews, W)
Foraminifera, smaller, Pacific Ocean and islands (M. R. Todd, W)

Mollusks:

Alaska (F. S. MacNeil, M)
Oregon (E. J. Moore, M)
Western Pacific islands (H. S. Ladd, W)
Pollen and spores, Kentucky (R. H. Tschudy, D)

Vertebrates:

Pleistocene (G. E. Lewis, D)
Atlantic coast (F. C. Whitmore, Jr., W)
Pacific coast (C. E. Repenning, M)
Panama Canal Zone (F. C. Whitmore, Jr., W)

Mesozoic:

Pacific coast (D. L. Jones, M)
Pierre Shale, Front Range area (W. A. Cobban, G. R. Scott, D)

Cretaceous:

Foraminifera, Nelchina area, Alaska (H. R. Bergquist, W)
Gulf coast and Caribbean (N. F. Sohl, W)
Western interior United States (W. A. Cobban, D)
Jurassic, North America (R. W. Imlay, W)
Triassic, marine faunas and stratigraphy (N. J. Silberling, M)

Paleozoic:

Fusuline Foraminifera, Nevada (R. C. Douglass, W)
Paleobotany and coal studies, Antarctica (J. M. Schopf, Columbus, Ohio)
Subsurface rocks, Florida (J. M. Berdan, W)
Type Morrow Series, Washington County, Ark. (L. G. Henbest, W)
Cambrian (A. R. Palmer, W)

Mississippian:

Corals, northern Alaska (H. M. Duncan, W)
Stratigraphy and brachiopods, northern Rocky Mountains and Alaska (J. T. Dutro, Jr., W)
Stratigraphy and corals, northern Rocky Mountains (W. J. Sando, W)

Ordovician:

Stratigraphy and brachiopods, Eastern United States (R. B. Neuman, W)
Western United States (R. J. Ross, Jr., D)

Pennsylvanian:

Fusulinidae, north-central Texas (D. A. Myers, D)
Spores and pollen, Kentucky (R. M. Kosanke, D)

Permian:

Floras, Southwest United States (S. H. Mamay, W)
Stratigraphy and brachiopods, Southwest United States (R. E. Grant, W)

Paleontology, stratigraphic--Continued**Paleozoic--Continued****Silurian-Devonian:**

Corals, Northeastern United States (W. A. Oliver, Jr., W)

Great Basin and Pacific coast (C. W. Merriam, M)

Upper Silurian-Lower Devonian, Eastern United States (J. M. Berdan, W)

Upper Paleozoic, Great Basin (M. Gordon, Jr., W)

Paleontology, vertebrate, systematic:

Pleistocene fauna, Big Bone Lick, Ky. (F. C. Whitmore, Jr., W)

Artiodactyls, primitive (F. C. Whitmore, Jr., W)

Tritylodonts, American (G. E. Lewis, D)

Paleotectonic maps. See Regional studies and compilations.**Pegmatites:****North Carolina:**

Blue Ridge Mountains, southern part, mica deposits (F. G. Lesure, W)

Franklin quadrangle (F. G. Lesure, W)

South Dakota:

Hill City pegmatite area (J. C. Ratté, D)

Keystone pegmatite area (J. J. Norton, W)

Permafrost studies:

Distribution and general characteristics (W. E. Davies, W)

Alaska:

Ground ice in central Alaska (T. L. Péwé, College)

Ground water and permafrost (J. R. Williams, w, Boston, Mass.)

Petroleum and natural gas:

Organic geochemistry (J. G. Palacas, D)

Mesozoic rocks, Florida and the eastern Gulf coast (E. R. Applin, Jackson, Miss.)

Tuffs, Green River Formation (R. L. Griggs, D)

Upper Jurassic rocks, northeast Texas, southwest Arkansas, northwest Louisiana (K. A. Dickinson, D)

Williston Basin, Wyoming, Montana, North Dakota, South Dakota (C. A. Sandberg, D)

Alaska:

Northern part, petroleum (G. Gryc, M)

Gulf of Alaska Tertiary province (G. Plafker, M)

Iniskin-Tuxedni region (R. L. Detterman, M)

Lower Yukon-Koyukuk area (W. W. Patton, Jr., M)

Nelchina area (A. Grantz, M)

Arizona:

Navajo Reservation, fuels potential (R. B. O'Sullivan, D)

Arkansas:

Northern part (E. E. Glick, D)

Ft. Smith district (T. A. Hendricks, D)

California:

Eastern Los Angeles basin (J. E. Schoellhamer, M)

Salinas Valley (D. L. Durham, M)

Colorado:

Northwestern part, Upper Cretaceous stratigraphy (J. R. Gill, D)

Animas River area (H. Barnes, D)

Grand Junction 2-degree quadrangle (W. B. Cashion, D)

Rangely 3 quadrangle (H. L. Cullins, c, D)

Thornburg area (J. R. Dyni, c, D)

Kansas:

Sedgwick Basin (W. L. Adkison, Lawrence)

Shawnee County (W. D. Johnson, Jr., Owensboro, Ky.)

Wilson County (H. C. Wagner, M)

Petroleum and natural gas--Continued

Michigan, Michigan basin (G. V. Cohee, W)

Mississippi, Homochitto National Forest (E. L. Johnson, c, Tulsa, Okla.)

Nebraska, central Nebraska basin (G. E. Prichard, D)

New Mexico:

Animas River area (H. Barnes, D)

San Juan Basin, east side (C. H. Dane, W)

Oklahoma:

Ft. Smith district (T. A. Hendricks, D)

McAlester Basin (S. E. Frezon, D)

Utah:

Northeastern part, Upper Cretaceous stratigraphy (T. A. Hendricks, D)

Grand Junction 2-degree quadrangle (W. B. Cashion, D)

Navajo Reservation, fuels potential (R. B. O'Sullivan, D)

Virginia, Big Stone Gap district (R. L. Miller, W)

Washington:

Grays Harbor basin:

Regional compilation (H. M. Beikman, M)

Western part (H. C. Wagner, M)

Wyoming:

Upper Cretaceous regional stratigraphy (J. R. Gill, D)

Crowheart Butte area (J. F. Murphy, D)

LaBarge 1 SW and 2 SE quadrangles (R. L. Rioux, c, W)

Lamont-Baroil area (M. W. Reynolds, D)

Oil Mountain quadrangle (W. H. Laraway, c, Casper)

Poison Spider quadrangle (W. H. Laraway, c, Casper)

Reid Canyon quadrangle (W. H. Laraway, c, Casper)

Spence-Kane area (R. L. Rioux, c, W)

Square Top Butte quadrangle (W. H. Laraway, c, Casper)

Petrology. See Geochemistry and petrology.**Phosphate:**

Oriskany Formation (W. D. Carter, W)

Phosphoria Formation, stratigraphy and resources (R. A. Gulbrandsen, M)

Southeastern United States, phosphate resources (J. B. Cathcart, D)

California, Monterey Formation (H. D. Gower, M)

Florida, land-pebble phosphate deposits (J. B. Cathcart, D)

Idaho:

Aspen Range-Dry Ridge area (V. E. McKelvey, W)

Driggs NE, SE, and SW quadrangles (M. L. Schroeder, c, D)

Garns Mountain NW and SW quadrangles (M. H. Staatz, c, D)

Palisades Peak quadrangle (D. A. Jobin, c, D)

Palisades Reservoir quadrangle (H. F. Albee, c, D)

Poker Peak quadrangle (H. F. Albee, c, D)

Soda Springs quadrangle (F. C. Armstrong, D)

Upper Valley quadrangle (R. L. Rioux, c, W)

Montana:

South-central part (R. W. Swanson, Spokane, Wash.)

Wise River quadrangle (G. D. Fraser, c, D)

Nevada:

Montello area (R. G. Wayland, c, Los Angeles, Calif.)

Spruce Mountain 4 quadrangle (G. D. Fraser, c, D)

Utah:

Crawford Mountains (W. C. Gere, c, D)

Gilbert Peak 1 NE quadrangle (J. R. Dyni, c, D)

Jessen Butte quadrangle (J. R. Dyni, c, D)

Phosphate--Continued**Utah--Continued**

- Morgan quadrangle (T. E. Mullens, c, D)
- Ogden 1 SE quadrangle (T. E. Mullens, c, D)
- Ogden 4 NE quadrangle (T. E. Mullens, c, D)
- Ogden 4 NW quadrangle (R. J. Hite, c, D)
- Phil Pico Mountain quadrangle (J. R. Dyni, c, D)
- Vernal phosphate area (E. M. Schell, c, D)

Wyoming:

- Jackson (30-minute) quadrangle (D. A. Jobin, c, D)
- Taylor Mountain quadrangle (M. L. Schroeder, c, D)

Plant ecology:

- Basic research in vegetation and hydrology (R. S. Sigafos, w, W)
- Ecologic criteria for conversion of juniper-pinyon woodlands to grasslands (R. S. Aro, w, D)
- Vegetation changes in southwestern North America (R. M. Turner, w, Tucson, Ariz.)

See also Vegetation.

Potash:

- Colorado and Utah, Paradox basin (O. B. Raup, D)
- New Mexico, Carlsbad, potash and other saline deposits (C. L. Jones, M)

Public and industrial water supplies:

- Corrosion and encrustation mechanisms in water supplies (F. E. Clarke, w, W)
- Kentucky, statewide water-utilization studies (R. J. Pickering, w, Columbus, Ohio)
- New Mexico, use of water by municipalities (W. A. Mourant, J. A. Basler, g, Albuquerque)
- North Carolina, chemical and physical quality characteristics of public water supplies (J. C. Chermers, w, Raleigh)

Quality of water:

- Saline ground water of the United States (J. H. Feth, w, M)
- Delaware River, chemical characteristics (D. McCartney, q, Philadelphia, Pa.)
- Lower Columbia River, Washington - Oregon (L. B. Laird, w, Tacoma, Wash.)
- Alabama, compilation of water-quality records (J. R. Avrett, w, Tuscaloosa)
- California, effect of diversion works on the Trinity River (G. Porterfield, w, Sacramento)
- Delaware, chemical quality, statewide (E. F. McCarren, q, Philadelphia, Pa.)
- Indiana, saline-water resources (R. J. Pickering, w, Columbus, Ohio)
- Kansas:
 - South Fork Ninescaw River basin (A. M. Diaz, w, Lincoln, Nebr.)
- Kentucky, saline-water investigations (H. T. Hopkins, g, Louisville)
- Maryland, chemical-quality reconnaissance of streams (J. D. Thomas, w, Towson)
- Walnut River basin (R. F. Leonard, w, Lincoln, Nebr.)
- Nebraska, Niobrara River basin (M. L. Maderak, w, Lincoln)
- New Jersey, basic water-quality network (P. W. Anderson, q, Trenton)
- New Mexico, maps showing quality of water by counties (F. E. Busch, g, Albuquerque)
- New York, Glowegee Creek at AEC reservation near West Milton (F. H. Pauszek, w, Albany)

Quality of water--Continued**Ohio:**

- Maumee River basin (M. Deutsch, J. C. Wallace, w, Gahanna)
- Miami River basin (C. R. Collier, w, Columbus)
- Quality of surface and ground waters, statewide inventory (C. R. Collier, w, Columbus)
- Oklahoma, Washita River basin (J. J. Murphy, q, Oklahoma City)
- Pennsylvania, statewide studies (D. McCartney, q, Philadelphia)

Texas:

- Hubbard Creek basin (C. H. Hembree, w, Austin)
- Quality of base flow of streams (C. H. Hembree, w, Austin)
- Reconnaissance of streams, statewide (L. S. Hughes, w, Austin)
- Surface waters, statewide study (L. S. Hughes, w, Austin)
- Surface waters of the Brazos River basin (J. Rawson, H. B. Mendieta, w, Austin)
- Utah, chemical quality of ground water (A. H. Handy, w, Salt Lake City)

Washington:

- Chemical quality of ground water (A. S. VanDenburgh, w, Tacoma)
- Influence of natural gas on ground-water quality (L. B. Laird, w, Tacoma)
- Quality of surface water (N. F. Leibbrand, w, Tacoma)
- Water quality of Grays Harbor (J. P. Beverage, w, Tacoma)

See also Sedimentation and Water resources.

Quicksilver:

- Geochemistry (A. P. Pierce, D)
- Mercury deposits and mercury resources (E. H. Bailey, M)
- Alaska, southeast (E. M. MacKevett, M)
- California, Coast Range ultramafic rocks (E. H. Bailey, M)
- Oregon, Ochoco Reservation, Lookout Mountain, Eagle Rock, and Post quadrangles (A. C. Waters, Baltimore, Md.)

Radioactive materials, transport in water:

- Disposition of radionuclides, Lower Columbia River (W. L. Haushild, q, Portland, Oreg.)
- Distribution of radioactivity, Clinch River, Tenn., sediments (R. J. Pickering, w, Chattanooga)
- Exchange phenomena and chemical reactions of radioactive substances (E. A. Jenne, q, D)
- Infiltration of radioiodine (R. M. Richardson, w, Chattanooga, Tenn.)
- Mineralogy and exchange capacity of fluvial sediments (V. C. Kennedy, q, D)
- Movement of radionuclides, Columbia River estuary (D. W. Hubbell, q, Portland, Oreg.)
- Movement of radionuclides in water through earth materials (W. A. Beetem, q, D)
- Step length and rest periods of sediment in alluvial channels (W. W. Sayre, h, Fort Collins, Colo.)
- Clinch River, Tenn., study (P. H. Carrigan, w, Chattanooga)
- Savannah River study (A. E. Johnson, s, Columbia, S.C.)

Radioactive-waste disposal:

- Disposal of treated radioactive-waste effluents, Bandelier Tuff, New Mexico (W. D. Purtyman, g, Albuquerque)

Radioactive-waste disposal--Continued**Hydrogeologic studies:**

National Reactor Testing Station, Idaho (D. A. Morris, g, Boise)

Oak Ridge Reservation, Tenn. (W. M. McMaster, w, Chattanooga)

Savannah River Plant, S. C. (I. W. Marine, G. E. Siple, g, Columbia)

Laboratory investigations (C. R. Naeser, W)

Nuclear-irradiation studies (C. M. Bunker, D)

Waste-contamination studies, Los Alamos, N. Mex. — ground water (W. D. Purtyman, g, Albuquerque)

Rare-earth metals. See Minor elements.**Regional studies and compilations, large areas of the United States:**

Basement-rock map of the United States (R. W. Bayley, M)

Geologic map of the United States (P. B. King, M)

Geologic map of the United States between lats 35° N and 39° N, scale 1:1,000,000 (C. R. Willden, D)

Gravity map of the United States (H. R. Joesting, W)

Military intelligence studies (M. M. Elias, W)

National Atlas, water-resources section (H. E. Thomas, g, M)

Paleotectonic-map folios:

Mississippian System (L. C. Craig, D)

Pennsylvanian System (E. D. McKee, D)

Permian System (E. D. McKee, D)

Sino-Soviet terrain atlas (M. M. Elias, W)

Rhenium. See Minor elements and Ferro-alloy metals.**Saline minerals:**

Colorado and Utah, Paradox basin (O. B. Raup, D)

New Mexico, Carlsbad potash and other saline deposits (C. L. Jones, M)

Wyoming, Sweetwater County, Green River Formation (W. C. Culbertson, D)

Sea-water intrusion:

Salt-water intrusion in coastal streams (T. H. Woodard, w, Raleigh, N. C.)

Connecticut and New York, recognition of late glacial substages in coastal areas (J. E. Upson, w, W)

Delaware River basin, lower part (D. McCartney, q, Philadelphia, Pa.)

Florida:

Dade County and city of Miami (H. Klein, w, Tallahassee)

Everglades National Park, estuaries (K. A. MacKichan, w, Ocala)

Miami River (S. D. Leach, w, Ocala)

Georgia:

Brunswick area (R. L. Wait, w, Atlanta)

Savannah area (H. B. Counts, w, Atlanta)

Sedimentation:

Effect of sedimentation on the propagation of trout in small streams, Montana (A. R. Gustafson, q, Worland, Wyo.)

Evaluation of dependent and independent variables with respect to sediment transport and resistance to flow in alluvial channels (H. P. Guy, h, Fort Collins, Colo.)

Fall velocity of fluvial sediment particles as affected by size, shape, density, concentration, and turbulence (H. P. Guy, h, Fort Collins, Colo.)

General studies of erosion and sedimentation and evaluation of erosion-control practices (N. J. King, w, D)

Sedimentation--Continued

Some sedimentation characteristics of a sand-bed stream (D. M. Culbertson, w, Lincoln, Nebr.)

Sources, movement, and distribution of sediment in a small watershed (M. G. Wolman, w, Baltimore, Md.)

Statistical analysis of ripples, dunes, and antidunes (C. F. Nordin, Jr., h, Fort Collins, Colo.)

Transport properties of natural clays (R. G. Wolff, w, W) California, Eel and Mad River basins, sediment transport (G. Porterfield, w, Sacramento)

Colorado, Kiowa Creek, fluvial sedimentation and runoff (R. Brennan, q, D)

Indiana, reconnaissance of sediment yields in streams (R. F. Flint, w, Columbus, Ohio)

Kansas, Little Blue River basin, fluvial sediment and quality of water (J. C. Mundorff, w, Lincoln, Nebr.)

Missouri, St. Louis, special sediment investigations at Mississippi River (C. H. Scott, w, Lincoln, Nebr.)

Nebraska:

Little Blue River basin, fluvial sediment and quality of water (J. C. Mundorff, w, Lincoln)

Medicine Creek basin, erosion and deposition (J. C. Brice, w, Lincoln)

New Jersey:

Sediment reconnaissance, coastal plain streams (D. W. Moody, w, Philadelphia, Pa.)

Stony Brook watershed, fluvial sedimentation (J. R. George, q, Harrisburg, Pa.)

North Carolina, upper Yadkin River basin, sediment yield (H. E. Reeder, q, Raleigh)

Oregon:

Alsea River basin, sedimentation in forested drainage areas (R. C. Williams, q, Portland)

Fluvial sediment transport (R. C. Williams, q, Portland)

Sediment-transport characteristics of certain streams (R. C. Williams, q, Portland)

Pennsylvania:

Statewide studies (J. R. George, q, Harrisburg)

Bixler Run watershed, hydrology and sedimentation (B. L. Jones, q, Harrisburg)

Corey Creek and Elk Run watershed (B. L. Jones, q, Harrisburg)

Susquehanna River basin, fluvial sediment reconnaissance (J. R. George, q, Harrisburg)

Texas:

Reconnaissance sediment investigations (C. T. Welborn, w, Austin)

Upper Trinity River basin, sedimentation (C. H. Hembree, w, Austin)

Washington:

Chehalis River basin, sedimentation (P. A. Glancy, w, Tacoma)

Palouse River basin, fluvial sediment transport (P. R. Boucher, w, Pasco)

Walla Walla River basin, fluvial sediment transport (B. E. Mapes, w, Pasco)

Wisconsin, reconnaissance sediment investigations (C. R. Collier, w, Columbus, Ohio)

Wyoming, Wind River Basin, sediment transport (D. C. Dial, q, Worland)

See also Geomorphology and Quality of water.

Sedimentation, reservoirs:

- California, Stony Gorge Reservoir (J. M. Knott, w, Sacramento)
- Colorado, Kiowa Creek basin, K-79 reservoir (R. Brennan, q, D)
- Georgia, North Fork Broad River, subwatershed 14 near Avalon (B. F. Joyner, w, Ocala, Fla.)
- Louisiana, Bayou Dupont watershed, reservoir (R. L. McAvoy, q, Baton Rouge)
- Nevada, Peavine Creek (D. O. Moore, w, Carson City)
- New Jersey, Baldwin Creek reservoir, trap efficiency (J. R. George, q, Harrisburg, Pa.)
- Texas, Escondido Creek (C. H. Hembree, w, Austin)
- Utah, Paria River basin, Sheep Creek near Tropic sediment barrier (G. C. Lusby, w, D)

Selenium. See Minor elements.

Silica:

- Oriskany Formation (W. D. Carter, W)
- Tintic Quartzite (K. B. Ketner, D)

Silver:

- Reconnaissance and exploration (H. R. Cornwall, W)

Soil moisture:

- Development of field criteria for evaluating sites for flood waterspreading (R. F. Miller, w, D)
- Differences in patterns and modes of soil-moisture movement under and adjacent to riparian vegetation (R. F. Miller, w, D)
- Effect of grazing exclusion, Badger Wash area, Colorado (G. C. Lusby, w, D)
- Effect of mechanical treatment on arid lands, Western United States (F. A. Branson, w, D)
- Ion distribution, water movement in soils and vegetation (R. F. Miller, w, D)
- Plant and soil-water response to thermal gradient, Ogalla Formation, Nebraska-New Mexico (F. A. Branson, w, D)
- Plants as indicators of soil-moisture availability (F. A. Branson, w, D)
- Soil-moisture energy relationships under and adjacent to riparian vegetation (I. S. McQueen, w, D)
- Water application and use on a range water spreader, northeast Montana (F. A. Branson, w, D)

Spectroscopy:

- Mobile spectrographic laboratory (A. P. Marrinzino, D)
- Spectrographic analytical services and research (A. W. Helz, W; A. T. Myers, D; H. Bastrom, M)
- X-ray spectroscopy (H. J. Rose, Jr., W; W. W. Brannock, M)

Springs:

- Springs of California (C. F. Berkstresser, Jr., w, Sacramento)

Stratigraphy and sedimentation:

- Basement-rock map of United States (R. W. Bayley, M)
- Cave deposits, stratigraphy and mineralogy (W. E. Davies, W)
- Sedimentary environments, classification (E. J. Crosby, D)
- Sedimentary mineralogy (P. D. Blackmon, D)
- Sedimentary-petrology laboratory (H. A. Tourtelot, D)
- Sedimentary structures, model studies (E. D. McKee, D)
- Subsurface-data center (L. C. Craig, D)
- Middle and Late Tertiary history, Northern Rocky Mountains and Great Plains (N. M. Denson, D)
- Pennsylvanian and Permian stratigraphy, Front Range (E. K. Maughn, D)

Stratigraphy and sedimentation--Continued

- Upper Jurassic stratigraphy, northeast Texas, southwest Arkansas, northwest Louisiana (K. A. Dickinson, D)

Green River Formation, tuffs (R. L. Griggs, D)

Phosphoria Formation, stratigraphy and resources (R. A. Gulbrandsen, M)

Pierre Shale:

Chemical and physical properties, Montana, North Dakota, South Dakota, Wyoming, and Nebraska (H. A. Tourtelot, D)

Paleontology and stratigraphy, Front Range area (W. A. Cobban, G. R. Scott, D)

Atlantic Coastal Plain:

Regional synthesis (J. C. Maher, M)

Southern part (J. E. Johnston, W)

Colorado Plateau:

Lithologic studies (R. A. Cadigan, D)

San Rafael Group, stratigraphy (J. C. Wright, W)

Stratigraphic studies (R. A. Cadigan, D)

Triassic stratigraphy and lithology (J. H. Stewart, M)

East-coast continental shelf and margin (K. O. Emery, Woods Hole, Mass.)

Williston Basin, Wyoming, Montana, North Dakota, and South Dakota (C. A. Sandberg, D)

Alaska, Mesozoic stratigraphy (W. W. Patton, A. Grantz, M)

Arizona:

Dripping Spring quartzite (H. C. Granger, D)

Hermit and Supai Formations (E. D. McKee, D)

Redwall limestone (E. D. McKee, D)

California, Lower Cambrian strata of southern Great Basin (J. H. Stewart, M)

Colorado:**Northwestern part:**

Jurassic stratigraphy (G. N. Pipiringos, D)

Pennsylvanian evaporite (W. W. Mallory, D)

Upper Cretaceous stratigraphy (J. R. Gill, D)

Kansas, Sedgwick Basin (W. L. Adkison, Lawrence)

Massachusetts, central Cape Cod, subsurface studies (R. N. Oldale, C. R. Tuttle, and C. Koteff, Boston)

Nebraska, central Nebraska basin (G. E. Prichard, D)

Nevada, Lower Cambrian strata of southern Great Basin (J. H. Stewart, M)

New Mexico, Guadalupe Mountains (P. T. Hayes, D)

New York, Dunkirk Formation and related beds (W. de Witt, Jr., W)

Oklahoma:

Southern part, Permian stratigraphy (D. H. Eargle, Austin, Tex.)

McAlester Basin (S. E. Frezon, D)

Pennsylvania, Devonian stratigraphy (G. W. Colton, W)

Texas, northern, Permian stratigraphy (D. H. Eargle, Austin)

Utah, northeastern, Upper Cretaceous stratigraphy (J. R. Gill, D)

Washington, Grays Harbor basin, regional compilation (H. M. Beikman, M)

Wyoming:

Green River Formation, geology and paleolimnology (W. H. Bradley, W)

South-central part, Jurassic stratigraphy (G. N. Pipiringos, D)

Upper Cretaceous, regional stratigraphy (J. R. Gill, D)

Stratigraphy and sedimentation--Continued**Wyoming--Continued**

Lamont-Baroil area (M. W. Reynolds, D)

Wedding of Waters--Devil Slide quadrangles (E. K. Maughan, D)

See also Paleontology, stratigraphic, and specific areas under Geologic mapping.**Structural geology and tectonics:**

Deformation research (S. P. Kanizay, D)

Isotopic studies of crustal processes (M. Tatsumoto, W)

Rock behavior at high temperature and pressure (E. C. Robertson, W)

Alaska, tectonic map (G. Gryc, M)

California:

San Andreas fault (L. F. Noble, Valyermo)

Sierra foothills mineral belt (L. D. Clark, M)

Montana, Hebgen Lake earthquake investigations (J. B. Hadley, W; I. J. Witkind, D)

Nevada, orogenic processes northern Sonoma Range (J. Gilluly, D)

See also specific areas under Geologic mapping.**Talc:**

Southeast United States, ultramafic rocks (D. M. Larabee, W)

Vermont, north-central (W. M. Cady, D)

Tantalum. See Minor elements.**Temperature studies, water:**

Temperature distribution in natural streams (E. J. Jones, w, M)

Thermal characteristics of aquifer systems (R. Schneider, w, W)

Thorium:

Western States, thorium investigations (M. H. Staatz, D)

Colorado:

Gunnison County, Powderhorn area (J. C. Olson, D)

Wet Mountains (M. R. Brock, D)

Idaho, central, radioactive placer deposits (D. L. Schmidt, W)

Tin:

Geochemistry and occurrence (N. Herz, W)

Alaska:

Lost River mining district (C. L. Sainsbury, D)

Seward Peninsula (P. L. Killeen, W)

Tofty placer district (D. M. Hopkins, M)

Tungsten. See Ferro-alloy metals.**Uranium:**

Formation and redistribution of uranium deposits (K. G. Bell, D)

Uranium-bearing pipes, Colorado Plateau and Black Hills (C. G. Bowles, D)

Uranium-bearing veins (G. W. Walker, D)

Uranium in black shales, mid-continent area (D. H. Eargle, Austin, Tex.)

Uranium-vanadium deposits in sandstone, Colorado Plateau (R. P. Fischer, D)

Arizona, Dripping Spring quartzite (H. C. Granger, D)

Colorado:

Baggs area (G. E. Prichard, D)

Bull Canyon district (C. H. Roach, D)

Gypsum Valley district (C. F. Withington, W)

La Sal area (W. D. Carter, W)

Lisbon Valley area (G. W. Weir, Berea, Ky.)

Maybell-Lay area (M. J. Bergin, W)

Slick Rock district (D. R. Shawe, D)

Uravan district (R. L. Boardman, W)

Uranium--Continued

Idaho, Mt. Spokane quadrangle (A. E. Weissenborn, Spokane, Wash.)

New Mexico:

Northwestern part (L. S. Hilpert, Salt Lake City, Utah)

Ambrosia Lake district (H. C. Granger, D)

Grants area (R. E. Thaden, Columbia, Ky.)

Laguna district (R. H. Moench, D)

South Dakota:

Harding County and adjacent areas (G. N. Pippingos, D)

Southern Black Hills (G. B. Gott, D)

Texas, coastal plain, geophysical and geological studies (D. H. Eargle, Austin)

Utah:

La Sal area (W. D. Carter, W)

Lisbon Valley area (G. W. Weir, Berea, Ky.)

Sage Plain area (L. C. Huff, Manila, P. I.)

San Rafael Swell (C. C. Hawley, D)

Washington, Mt. Spokane quadrangle (A. E. Weissenborn, Spokane)

Wyoming:

Central part, selected uranium deposits (F. C. Armstrong, D)

Baggs area (G. E. Prichard, D)

Gas Hills district (H. D. Zeller, D)

Shirley Basin area (E. N. Harshman, D)

Urban geology:**California:**

Los Angeles area (J. T. McGill, D)

Malibu Beach quadrangle (R. F. Yerkes, M)

Oakland East quadrangle (D. H. Radbruch, M)

Palo Alto quadrangle (E. H. Pampeyan, M)

Point Dume quadrangle (R. H. Campbell, M)

San Francisco North quadrangle (J. Schlocker, M)

San Francisco South quadrangle (M. G. Bonilla, M)

San Mateo quadrangle (G. O. Gates, M)

Colorado:

Denver metropolitan area (R. M. Lindvall, D)

Golden quadrangle (R. Van Horn, D)

Pueblo and vicinity (G. R. Scott, D)

District of Columbia, Washington metropolitan area (H. W. Coulter, C. F. Withington, W)

Maryland, Washington, D. C., metropolitan area (H. W. Coulter, C. F. Withington, W)

Massachusetts, Boston and vicinity (C. A. Kaye, Boston)

Montana, Great Falls area (R. W. Lemke, D)

South Dakota, Rapid City area (E. Dobrovolsky, D)

Texas, San Antonio and vicinity (R. D. Miller, D)

Utah, Salt Lake City and vicinity (R. Van Horn, D)

Virginia, Washington, D. C., metropolitan area (H. W. Coulter, C. F. Withington, W)

Washington:

Puget Sound Basin (D. R. Crandell, D)

Seattle and vicinity (D. R. Mullineaux, D)

Urbanization, hydrologic effects:**Effect on flood flow:**

Kansas, Wichita area (M. W. Busby, s, Topeka)

North Carolina, Charlotte and Winston-Salem metropolitan areas (L. A. Martens, w, Raleigh)

Tennessee, Nashville--Davidson County metropolitan area (L. G. Conn, w, Chattanooga)

Hydrologic effects of urbanization (J. R. Crippen, w, M)

Urbanization, hydrologic effects--Continued

Maryland, Northwest Branch, Anacostia River basin (D. H. Carpenter, w, College Park)

Texas:

Turtle Creek (F. H. Ruggles, w, Austin)
Waller Creek (W. H. Espey, Jr., w, Austin)

Vanadium:

Commodity studies (R. P. Fischer, D)
Geology and resources, bibliography (J. P. Ohl, D)
Colorado Plateau, uranium-vanadium deposits in sandstone (R. P. Fischer, D)

Colorado:

Bull Canyon district (C. H. Roach, D)
La Sal area (W. D. Carter, W)
Lisbon Valley area (G. W. Weir, Berea, Ky.)
Slick Rock district (D. R. Shawe, D)
Uravan district (R. L. Boardman, W)

Utah:

La Sal area (W. D. Carter, W)
Lisbon Valley area (G. W. Weir, Berea, Ky.)
Sage Plain area (L. C. Huff, Manila, P. I.)

Vegetation:

Plant-analysis laboratory (F. N. Ward, D)
Alaska, vegetation map (L. A. Spetzman, W)
Pacific islands, vegetation (F. R. Fosberg, W)
See also Plant ecology.

Volcanic-terrane hydrology:

Columbia River Basalt (R. C. Newcomb, g, Portland, Oreg.)

Volcanology:

Pacific coast basalts, geochemistry (K. J. Murata, Costa Rica)
Silicic ash beds, correlation (H. A. Powers, Hawaii National Park, Hawaii)
Alaska, Katmai National Monument, petrology and volcanism (G. H. Curtis, M)
Costa Rica, volcanic studies (K. J. Murata, San Jose)
Hawaii, Hawaiian Volcano Observatory (H. A. Powers, Hawaii National Park)
Idaho, central Snake River Plain, volcanic petrology (H. A. Powers, Hawaii National Park, Hawaii)
Montana:
Bearpaw Mountains, petrology (W. T. Pecora, W)
Wolf Creek area, petrology (R. G. Schmidt, W)
New Mexico, Valles Mountains, petrology (R. L. Smith, W)

Water management:

Water-land relationships, Patuxent River basin, Maryland (D. O'Bryan, w, W)

Waterpower classification:

Western United States, waterpower resources (A. Johnson, c, W)
Wilderness and wild areas, classification of existing waterpower (J. D. Simpson III, c, W)

Alaska:

Burroughs Bay region, unnamed lake near Spur Mountain (J. B. Dugwyler, c, Tacoma, Wash.)
Eagle Lake, near Bradfield Canal (J. B. Dugwyler, c, Tacoma, Wash.)
Kenai Peninsula, Nellie Juan River (G. C. Giles, c, Tacoma, Wash.)

Kodiak Island:

Spiridon Lake (G. C. Giles, c, Tacoma, Wash.)
Terror Lake (G. C. Giles, c, Tacoma, Wash.)

Waterpower classification--Continued**Alaska--Continued****Lake Clark region:**

Kijik River (J. D. Simpson III, c, W)
Twin Lakes (R. Bondy, c, Tacoma, Wash.)

Arizona, Colorado River, Lake Powell section (H. D. Tefft, c, D)

California, statewide waterpower resources (R. N. Doolittle, c, Sacramento)

Colorado:

Elk River (H. D. Tefft, c, D)
King Solomon Creek (H. D. Tefft, c, D)
North Platte River basin (W. C. Senkpiel, c, D)

Idaho:

Statewide waterpower resources (L. L. Young, c, Portland, Oreg.)
Payette River (J. L. Colbert, c, Portland, Oreg.)
Salmon River basin (L. L. Young, c, Portland, Oreg.)
Weiser River basin (J. L. Colbert, c, Portland, Oreg.)
Montana, Flathead River (K. S. Soward, c, Great Falls)
Nevada, Duck Creek (K. W. Sax, c, Sacramento, Calif.)
Oklahoma, inventory of waterpower resources (W. C. Senkpiel, c, D)

Oregon:

Alsea River (L. L. Young, c, Portland)
Nehalem River (L. L. Young, c, Portland)
Siuslaw River (J. L. Colbert, c, Portland)

Utah:

Colorado River, Lake Powell section (H. D. Tefft, c, D)
Great Salt Lake basin (K. W. Sax, c, Sacramento, Calif.)
Green River, upstream from Flaming Gorge Dam (H. D. Tefft, c, D)
Sevier Lake basin (K. W. Sax, c, Sacramento, Calif.)

Washington:

Olympic Peninsula, review of waterpower withdrawals, Queets River and adjacent basins south and east to the Duckabush River basin (J. B. Dugwyler, c, Tacoma)
Statewide waterpower resources (G. C. Giles, c, Tacoma)

Wyoming, Green River, Flaming Gorge Reservoir section (H. D. Tefft, c, D)

Water resources:

Connecticut River basin — Vermont, New Hampshire, Massachusetts, Connecticut (D. J. Cederstrom, w, Boston, Mass.)

Lower Colorado basin, hydrology (C. C. McDonald, w, Yuma, Ariz.)

Mississippi embayment, hydrology (E. M. Cushing, w, Memphis, Tenn.)

Ohio River basin (M. Deutsch, w, Gahanna, Ohio)

Public domain:

Pacific coast area, water-supply exploration (C. T. Snyder, w, M)

Rocky Mountain area, water-supply exploration (C. E. Sloan, w, D)

Western States, areal hydrology (G. C. Lusby, w, D)

Upper Brazos River basin project, Permian Basin program (P. R. Stevens, w, Austin, Tex.)

Upper Mississippi River basin (P. G. Olcott, g, Madison, Wis.)

Water resources--Continued

Alabama (w, Tuscaloosa):

- Collection and compilation of water records (C. F. Hains)
- Hydrologic atlas of the State (C. F. Hains)
- Project planning, reports, and dissemination of information (W. J. Powell)
- Relation of oil and gas industry to water resources (W. J. Powell)
- Study of conservation lakes (C. F. Hains)
- Butler County, geologic and hydrologic profile along U.S. Highway 31 (J. G. Newton)

Ground water:

- Barbour County (J. G. Newton)
- Cullman County (R. J. Faust)
- Greene County (K. D. Wahl)
- Hale County (T. H. Sanford)
- Marion County (L. V. Causey)
- Marshall County (T. H. Sanford)
- Pickens County (K. D. Wahl)
- Sumter County (T. H. Sanford)

Hydrology:

- Southwest part (L. B. Pierce)
- Choctawhatchee-Escambia River basins (J. C. Scott)
- Huntsville area (W. F. Harris)
- Surface water, Tennessee River basin, low-flow investigations (J. R. Harkins)

Water resources:

- Southwest part, plan for investigation (J. R. Avrett)
- Natchez Trace Parkway, investigation along (W. J. Powell)

Alaska (w, Anchorage):

- Bridge-site investigations, Glacier Creek and Allen, Chulitna, Nome, and Tok Rivers (V. K. Berwick)
- Collection of basic data--streamflow, water quality, and sediment (H. Hulsing)
- Knik and Susitna Rivers, scour research program (V. K. Berwick)
- Tanana River, Nenana, bank-erosion study (V. K. Berwick)
- Nushagak Bay--Nilnilchik, sedimentation study (R. G. Schupp)

Ground water:

- Annette Island (M. V. Marcher)
- National parks (M. V. Marcher)
- Statewide inventory (M. V. Marcher)

Hydrographic studies:

- Kootznahoo Inlet (C. W. Boning)
- Taki Inlet (V. K. Berwick)

Hydrology:

- Anchorage area (M. V. Marcher)
- Cook Inlet--Knik Arm (R. G. Schupp)
- Tanana basin (M. V. Marcher)

Water resources:

- Angoon and Hoonah (M. V. Marcher)
- U.S. Air Force (A. J. Feulner)

American Samoa (Honolulu, Hawaii):

- Ground water (K. J. Takasaki, g)

Arizona (w, Tucson):

- East Verde River, transmountain diversion studies (E. S. Davidson)

Ground water:

- Analysis of water-level declines (N. D. White)

Water resources--Continued

Arizona--Continued

Ground water--Continued

- Beardsley area (W. Kam)
- Big Sandy Valley (W. Kam)
- Coconino County, southern (E. H. McGavock)
- Dateland-Hyder area (W. G. Weist)
- Kingman area (J. B. Gillespie)
- Navajo Indian Reservation (M. E. Cooley)
- Papago Indian Reservation (L. A. Heindl)
- Pinal County, northwestern (W. F. Hardt)
- Safford area (E. S. Davidson)
- Tucson basin (E. F. Pashley, Jr.)
- Willcox basin (S. G. Brown)
- Williams, analysis of public water supply (B. W. Thomsen)

Hydrology, alluvial basins (M. E. Cooley)

Water resources, Sycamore Creek basin (B. W. Thomsen)

Arkansas (w, Little Rock):

Ground water:

- Arkansas River valley (M. S. Bedinger)
- Lower Red River valley (M. S. Bedinger)

Water resources:

- Grant and Hot Spring Counties (H. N. Halberg)
- Independence and Jackson Counties (D. R. Albin)
- Pulaski and Saline Counties (R. O. Plebuch)
- White River basin (G. M. Hogensen)

California (w, Sacramento):

Ground water:

Antelope Valley:

- East Kern Water Agency (L. C. Dutcher)
- Eastern part (L. C. Dutcher)
- Barstow, Marine Corps Supply Center (G. Miller)
- Camp Pendleton Marine Corps Base (F. W. Giessner)
- China Lake, Naval Ordnance Test Station (F. W. Giessner)
- Death Valley National Monument, Saratoga Spring, reconnaissance (F. Kunkel)
- Edwards Air Force Base (F. W. Giessner)
- Elwood-Gaviota area (C. P. Zones)
- Fresno area (R. W. Page)
- Hanford-Visalia area (M. G. Croft)
- Orange County, coastal-area studies (L. C. Dutcher)
- Santa Ana area, upper part, south coastal area (L. C. Dutcher)
- Santa Ynez Uplands (G. F. LaFreniere)
- Summerland area (K. S. Muir)
- Twentynine Palms Marine Corps Training Center (F. W. Giessner)
- San Joaquin Valley, southern (M. G. Croft)
- Vandenberg Air Force Base (K. S. Muir)

Hydrology:

- Lompoc Plain, hydrologic inventory (R. E. Evenson)
- Point Reyes National Seashore (E. J. McClelland)

Colorado (Denver):

Ground water:

- Summary of pumping tests in State (W. W. Wilson, g)
- Bent County (J. H. Irwin, g)
- Big Sandy Valley, below Limon (D. L. Coffin, g)

Water resources--Continued

Colorado (Denver)--Continued

Ground water--Continued

Black Squirrel Creek Valley (H. E. McGovern, g, W)

Colorado High Plains, trends in ground-water development (A. J. Boettcher, g)

Denver Basin (J. A. McConaghy, g)

Denver Basin, ground-water trends (A. J. Boettcher, g)

Fremont and Pueblo Counties (H. E. McGovern, g, W)

Piceance Basin (D. L. Coffin, g)

Hydrology, Arkansas River basin-Canon City to State line (J. E. Moore, g)

Connecticut (q, s, Hartford; g, Middletown):

Ground water, Hamden-Wallingford area (A. M. LaSala, Jr., g)

Water resources:

Connecticut River basin (D. J. Cederstrom, w, Boston, Mass.)

Water resources of Connecticut:

Part 2, Shetucket River basin (M. P. Thomas, s)

Part 3, Thames River basin (C. E. Thomas, Jr., q)

Part 4, Southwestern coastal basins (R. B. Ryder, g)

Part 5, Lower Housatonic River basin (W. E. Wilson)

Florida (w, Tallahassee):

Statewide, special studies (C. S. Conover, K. A. MacKichan, R. W. Pride)

Water atlas (W. E. Kenner)

Alafia and Peace River basins, fluoride (L. G. Toler)

Southeastern Florida, water management effects (S. D. Leach)

Geohydrology:

Marion County (F. N. Visser)

Panama City area (J. B. Foster)

Sarasota County, Venice well field (W. E. Clark)

Ground water:

Baker, Duval, and Nassau Counties (G. W. Leve)

Dade County, special studies (H. Klein)

Fort Lauderdale area, special studies (H. Klein)

Water resources:

Broward County (C. B. Sherwood)

Everglades National Park (J. H. Hartwell)

Lower Hillsboro Canal area (R. G. Grantham)

Mid-Gulf basins (R. N. Cherry)

Myakka River basin (B. F. Joyner)

Orange County (W. F. Lichtler)

Volusia County, central (B. J. Bermes)

Georgia (w, Atlanta):

Ground water:

Floyd and Polk Counties (C. W. Cressler)

Grady County, Cairo area (C. W. Sever)

Water resources:

Cook County (C. W. Sever)

Pulaski County (R. C. Vorhis)

Rockdale County (M. J. McCollum)

Thomas County (C. W. Sever)

Guam (Honolulu, Hawaii):

Ground water (D. A. Davis, g)

Surface water (S. H. Hoffard, s)

Water resources--Continued

Hawaii (Honolulu):

Hydrologic studies (G. T. Hirashima, s)

Water resources:

Hawaii:

Hilo-Puna area, reconnaissance (G. Yamanaga, s)

Kau area, reconnaissance (D. A. Davis, g)

Kona area, reconnaissance (D. A. Davis, g)

Kauai, Waialaeale, rainfall (M. M. Miller, s)

Oahu:

Kahuku area (K. J. Takasaki, g)

Mokuleia-Waialua area (J. C. Rosenau, g)

Waianae district (C. P. Zones, g)

Windward Oahu (K. J. Takasaki, g)

Idaho (Boise):

Ground water:

Aberdeen-Springfield area (H. G. Sisco, g)

Mud Lake Basin (P. R. Stevens, g)

Rock Creek-Goose Creek area (E. G. Crosthwaite, g)

Salmon Falls Creek area (E. G. Crosthwaite, g)

Teton Basin, lower part (E. G. Crosthwaite, g)

Surface water:

Bruneau River basin, systems gaging (H. C. Riggs, s, W)

Snake River inflow, Milner to King Hill (C. A. Thomas, s)

Water resources:

Leadore, Big Springs Creek area (E. G. Crosthwaite, g; R. S. George, s)

Little Lost River basin (H. A. Waite, g; S. O. Decker, s)

Raft River basin (E. H. Walker, g; S. O. Decker, s)

Indiana (w, Indianapolis):

Ground water:

Northwestern part (J. D. Hunn)

West-central part (L. W. Cable)

Delaware County (R. E. Hoggatt)

Harrison County (J. D. Hunn)

Posey County (T. M. Robison)

Vanderburgh County (L. W. Cable)

Iowa (Iowa City):

Ground water:

Cretaceous aquifer of Iowa (P. D. Robinson, P. J. Horick, W. L. Steinhilber, g)

Mississippian aquifer of Iowa (P. J. Horick, W. L. Steinhilber, g)

South-central part, water from glacial deposits (J. W. Cagle, g)

Linn County (R. E. Hansen, g)

Muscatine Island (R. E. Hansen, g)

Kansas (Lawrence):

Analysis of hydrologic data (J. M. McNellis, g)

Ground water:

Northwestern part (S. W. Fader, g)

Southwestern part (W. R. Meyer, g)

Allen County (D. E. Miller, g)

Brown County (C. K. Bayne, g)

Butler County (J. M. McNellis, g)

Cherokee County (W. J. Seever, g)

Decatur County (W. G. Hodson, g)

Ellsworth County (C. K. Bayne, g)

Water resources--Continued

Kansas (Lawrence)--Continued

Ground water--Continued

Finney County (W. R. Meyer, g)
 Hamilton County (W. R. Meyer, g)
 Jefferson County (J. D. Winslow, g)
 Johnson County (H. G. O'Connor, g)
 Kearny County (W. R. Meyer, g)
 Labette County (W. L. Jungmann, g)
 Linn County (W. J. Seevers, g)
 Montgomery County (H. G. O'Connor, g)
 Neosho County (W. L. Jungmann, g)
 Pratt County (D. W. Layton, g)
 Republican River valley (S. W. Fader, g)
 Rush County (J. McNellis, g)
 Walnut River basin (J. M. McNellis, g)

Kentucky (Louisville):

Ground water:

Jackson Purchase area (R. W. Davis, g)
 Louisville area (E. A. Bell, g)
 Ohio River, alluvial terraces (W. E. Price, J. T. Gallaher, g)

Quality of surface and ground water -- statewide inventory (R. J. Pickering, q, Columbus, Ohio)

Water resources:

Bell County, upper Yellow Creek basin (D. S. Mull, g)

Mammoth Cave area (R. V. Cushman, g)

Louisiana (Baton Rouge):

Floods, southeastern Louisiana -- rainfall-runoff relations (A. J. Calandro, s)

Ground water:

Avoyelles Parish (J. R. Marie, g)
 Gramercy area (D. G. Moore, Jr., g)
 Greater New Orleans area (J. R. Rollo, g)
 Lower Red River valley (M. S. Bedinger, w, Little Rock, Ark.)
 Norco area (P. B. Bieber, g)

Surface water:

Southeastern Louisiana, unit-hydrograph studies (V. B. Sauer, s)
 Southwestern Louisiana, unit-hydrograph studies (V. B. Sauer, s)

Water resources:

Southwestern part (A. H. Harder, g; S. M. Rogers, q)
 Lake Pontchartrain study (G. T. Cardwell, g)
 Ouachita Parish (J. E. Rogers, g)
 Plaquemine-White Castle area (C. D. Whiteman, Jr., g)
 Pointe Coupee Parish (M. D. Winner, Jr., g)

Maine (Augusta):

Ground water:

Coastal area of southwestern Maine (G. C. Prescott, g)
 Lower Androscoggin basin (G. C. Prescott, g)
 Lower Penobscot basin (G. C. Prescott, g)

Maryland (w, Baltimore):

Ground water:

Baltimore County, use of ground water (C. P. Laughlin)
 Chesapeake and Ohio Canal (E. G. Otton)
 Susquehanna River basin (P. R. Seaber, g, Harrisburg, Pa.)

Water resources--Continued

Maryland (w, Baltimore)--Continued

Water resources:

Dorchester and Talbot Counties (F. K. Mack)
 Georges Creek basin, a corner of Appalachia (D. O'Bryan, J. W. Crooks, W)
 Patuxent River basin (J. W. Crooks, D. O'Bryan, w, W)
 Salisbury area (D. H. Boggess)

Massachusetts (w, Boston):

Ground water:

Highway salt, ground-water contamination from (R. G. Petersen)
 Assabet River basin (S. J. Pollock)
 Boston, central area (W. N. Palmquist)
 Cape Cod National Seashore (R. G. Petersen)
 Lower Merrimack valley (J. E. Cotton)
 Parker and Rowley Rivers drainage basins (E. A. Sammel)
 Ten Mile-North Taunton Rivers basin (J. R. Williams)

Water resources:

Connecticut River basin (D. J. Cederstrom)
 Housatonic River basin (R. F. Norvitch)
 Millers River basin (M. R. Collings)

Michigan (w, Lansing):

Ground water:

Dickinson County (G. E. Hendrickson)
 Kalamazoo County (J. B. Miller)
 Tri-County area (K. E. Vanlier)
 Surface water, North Branch Clinton River basin (R. G. Knutilla)

Water resources:

Branch County (L. E. Stoimenoff)
 Grand River basin (K. E. Vanlier)
 Marquette Iron Range (S. W. Wiitala)

Minnesota (St. Paul):

Ground water:

Grand Rapids area (E. L. Oakes, g)
 Kittson, Marshall, and Roseau Counties (G. R. Schiner, g)

Hydrology, Twin Cities metropolitan area, Anoka, Carver, Dakota, Goodhue, Hennepin, Ramsey, Scott, and Washington Counties (H. O. Reeder, g)

Water-resources reconnaissance of watershed units:

Chippewa River Unit, Chippewa, Douglas, Grant, Kandiyohi, Otter Tail, Stearns, and Stevens Counties (R. D. Cotter, g)
 Lac qui Parle River unit, Lac qui Parle, Lincoln, and Yellow Medicine Counties (R. D. Cotter, g)
 Mississippi Headwaters unit, Aitkin, Becker, Beltrami, Carlton, Cass, Clearwater, Crow Wing, Hubbard, Itasca, and St. Louis Counties (E. L. Oakes, g)
 Pomme de Terre River unit, Big Stone, Chippewa, Douglas, Grant, Otter Tail, Stevens, and Swift Counties (R. D. Cotter, g)
 Roseau River unit, Beltrami, Kittson, Lake of the Woods, Marshall, and Roseau Counties (T. C. Winter, g)
 Two Rivers unit, Kittson, Marshall, and Roseau Counties (R. W. Maclay, g)

Water resources--Continued

Mississippi (w, Jackson):

Geology and ground water:

- Big Black River basin (B. E. Wasson)
- Pascagoula River basin (R. Newcome, Jr.)
- Pearl River basin (J. W. Lang)

Ground water:

- Hancock County, National Aeronautics and Space Administration Test Facility (R. Newcome, Jr.)
- Pearl River basin, middle part (R. E. Taylor)

Water resources:

- Clay, Lowndes, Monroe, and Oktibbeha Counties (B. E. Wasson)
- Forrest, Green, Jones, Perry, and Wayne Counties (B. E. Wasson)
- Harrison and Stone Counties (R. Newcome, Jr.)

Missouri (Rolla):

- Joplin area (E. J. Harvey, g)
- White River basin (G. M. Hogenson, w, Little Rock, Ark.)

Montana (Billings):

Ground water:

- Cedar Creek anticline, west flank (O. J. Taylor, g)
- Deer Lodge Valley (R. L. Konizeski, g)
- Glacier and Toole Counties, Cut Bank area (E. A. Zimmerman, g)
- Lower Bighorn River valley, Hardin Unit (L. J. Hamilton, g)
- Missouri River bottoms, northeast Montana (W. B. Hopkins, g)
- Ravalli County, Bitterroot Valley (R. G. McMurtrey, g)
- Hydrology, Hungry Horse Reservoir (M. I. Rorabaugh, W. D. Simons, w, Tacoma, Wash.)

Nebraska (w, Lincoln):

Geology and hydrology, Saline County (P. A. Emery)

Ground water:

- Adams County (C. F. Keech)
- Box Butte County, water-level fluctuations (C. F. Keech)
- Fillmore County (C. F. Keech)
- York County (C. F. Keech)

Nevada (w, Carson City):

- Humboldt River valley, reconnaissance (T. E. Eakin, R. D. Lemke)

Ground water:

- Antelope Valley (F. E. Rush)
- Carico Lake valley (D. E. Everett)
- Eagle Valley (G. F. Worts, Jr.)
- Eldorado Valley (F. E. Rush)
- Grass Valley (P. Cohen)
- Huntington Valley (F. E. Rush, D. O. Moore)
- Kings River valley (G. T. Malmberg)
- Kobeh Valley (F. E. Rush)
- Lovelock area (D. E. Everett, F. E. Rush)
- Monitor Valley (F. E. Rush)
- Pahrump Valley (G. T. Malmberg)
- Piute Valley (F. E. Rush)
- Quinn River valley (C. J. Huxel, Jr.)
- Reese River valley, upper part (F. E. Rush)
- Spring Valley (F. E. Rush)
- Snake Valley (J. W. Hood, F. E. Rush)
- Steptoe Valley (T. E. Eakin, D. O. Moore, D. E. Everett)

Water resources--Continued

Nevada (w, Carson City)--Continued

Ground water--Continued

- Washoe Valley (G. F. Worts, Jr.)
- White River valley system (T. E. Eakin)
- Surface-water resources, statewide (R. D. Lamke, D. O. Moore)

New Hampshire (w, Boston, Mass.):

Ground water:

- Ashuelot River basin (J. M. Weigle)
- Lower Merrimack River basin (J. M. Weigle)
- Water resources, Connecticut River basin (D. J. Cederstrom)

New Jersey (Trenton):

- Drought in the Delaware River basin (J. E. McCall, s)
- Geohydrology, Raritan-Magothy aquifer system (H. E. Gill, D. Langmuir, g)

Ground water:

- Water-level fluctuations, 1963-67 (C. R. Austin, g)
- Camden County (E. Donsky, g)
- Cumberland County (J. G. Rooney, g)
- Essex County (W. D. Nichols, g)
- Hackensack River basin (L. D. Carswell, g)
- Ocean County (H. R. Anderson, g)
- Pine Barrens (E. C. Rhodehamel, g)
- Wharton Tract (E. C. Rhodehamel, g)

New Mexico (Albuquerque):

- Miscellaneous activities under the New Mexico State Engineer program (L. V. Davis, g)

State Planning Office Report (g):

- Pt. 1 -- Arkansas River basin (W. C. Ballance)
- Pt. 2 -- Southern High Plains (W. C. Ballance)
- Pt. 3 -- Pecos River basin (W. A. Maurant)
- Pt. 4 -- Central closed basins (F. B. Titus, Jr.)
- Pt. 5 -- Rio Grande basin (G. A. Dinwiddie)
- Pt. 6 -- Western closed basins (J. B. Cooper)
- Pt. 7 -- San Juan River basin (J. B. Cooper, F. D. Trauger)
- Pt. 8 -- Lower Colorado River basin (F. D. Trauger)
- Pt. 9 -- Southwestern closed basins (G. C. Doty)

Ground water:

- Water levels and artesian pressure in observation wells in New Mexico (F. E. Busch, g)

Eddy County:

- Major Johnson Springs aquifer, evaluation of hydraulic characteristics (R. L. Cushman, g)
- Malaga Bend area, evaluation of pumping effects (E. R. Cox, g)

Grant County (F. D. Trauger, g)

- Guadalupe County (A. Clebsch, Jr., G. A. Dinwiddie, g)

Luna County, southern (G. C. Doty, g)

McMillan delta area (J. S. Havens, g)

Manzano Mountains area (F. B. Titus, g)

Quay County (W. A. Maurant, g)

Roswell basin:

- Chaves County, well drilling and testing (G. E. Maddox, g)

- Chaves and Eddy Counties, quantitative analysis of the ground-water system (G. E. Maddox, g)

Sandia Mountains area (F. B. Titus, g)

- White Sands Missile Range, reconnaissance, ground-water resources at selected sites (G. C. Doty, g)

Water resources--Continued**New Mexico (Albuquerque)--Continued****Ground water--Continued**

White Sands Missile Range Headquarters well field, reappraisal (W. C. Ballance, g)

Water resources, Los Alamos (E. C. John, g)

New York (w, Albany):**Ground water:**

Nassau County (J. Isbister)

Orange County (M. H. Frimpter)

Queens County (J. Soren)

Rensselaer County, Schodack terrace (H. Stewart)

Suffolk County, mid-island area (J. Soren)

Susquehanna River basin (P. R. Seaber, g, Harrisburg, Pa.)

Syracuse area (I. H. Kantrowitz)

Ulster County (M. H. Frimpter)

Surface water:

Gazetteer of streams (F. L. Robison)

Statewide resources (J. C. Kammerer)

Susquehanna River basin, fluvial sediment (A. D. Randall)

Water resources:

Genesee River basin (B. K. Gilbert)

Lake Erie-Niagara area (A. L. LaSala)

Western Oswego River basin (W. J. Shampine)

North Carolina (w, Raleigh):

Stream sanitation and water supply of State (G. C. Goddard)

Ground water:

Cape Hatteras National Park, quality of ground water (H. B. Wilder)

Chowan County (O. B. Lloyd, Jr.)

Craven County (E. O. Floyd)

New Hanover County (G. Bain)

Pitt County (C. T. Sumsion)

Surface water:

Interpretation of surface-water data (G. C. Goddard)

Neuse River headwaters, surface-water resources (G. C. Goddard, J. F. Turner, Jr.)

North Dakota (Grand Forks):

Flood characteristics of streams at selected sites (O. A. Crosby, s, Bismarck)

Ground water:

Barnes County (T. E. Kelly, g)

Burleigh County (P. G. Randich, g)

Cass County (R. L. Klausling, g)

Divide County (C. A. Armstrong, g)

Eddy and Foster Counties (H. Trapp, g)

Grand Forks County (T. E. Kelly, g)

Minot area, artificial recharge (D. L. Hills, g)

Renville County (W. A. Pettyjohn, g)

Richland County (C. H. Baker, Jr., g)

Stutsman County (C. J. Huxel, g)

Ward County (W. A. Pettyjohn, q)

Wells County (P. G. Randich, g)

Williams County (E. A. Ackroyd, g)

Ohio (w, Columbus):

Flood characteristics of streams at selected sites (E. E. Webber)

Ground water:

Northeastern part, principal aquifers (J. L. Rau)

Lower Miami River basin (A. M. Spieker)

Lancaster municipal well field (S. E. Norris)

Water resources--Continued**Oklahoma (Oklahoma City):**

Special investigations and reports (P. R. Wood, g)

Ground water:

Fresh ground-water zone, thickness (D. L. Hart, Jr., g)

Arkansas River valley, eastern part of State (M. S. Bedinger, w, Little Rock, Ark.)

Arkansas and Verdigris River valleys (H. H. Tanaka, g)

Cleveland and Oklahoma Counties, Garber Sandstone and Wellington Formation (P. R. Wood, g)

Lower Red River valley (M. S. Bedinger, w, Little Rock, Ark.)

Woodward County (P. R. Wood, g)

Quality of water, upper Arkansas River basin (R. P. Orth, q)

Oregon (Portland):

Abert Lake and other closed-basin lakes, hydrology and geochemistry (A. S. Van Denburgh, q)

Ground water:

Eola-Amity Hills area (D. Price, g)

French Prairie area (D. Price, g)

Molalla-Salem slope area (E. R. Hampton, g)

Pennsylvania (g, Harrisburg; q, s, Philadelphia):**Ground water:**

Chester County, metamorphic and igneous rocks (C. W. Poth, g)

Lancaster County, carbonate rocks (H. Meisler, g)

Loysville quadrangle (H. E. Johnston, g)

Luzerne County, Wyoming Valley (J. R. Hollowell, g)

Mifflintown quadrangle (H. E. Johnston, g)

Monogahela River basin (B. M. Wilmoth, g)

New Oxford Formation (H. E. Johnston, g)

Susquehanna River basin (P. R. Seaber, g)

Quality of water:

Brandywine Creek basin, water-quality reconnaissance (A. N. Ott, q, Harrisburg)

Lehigh River basin, chemical quality (S. D. Faust, q)

Neshaminy Creek basin, chemical quality of surface waters (B. A. Malo, q)

Surface water, Philadelphia area, hydrology of streams (E. L. Smith, s)

Water resources, Schuylkill River basin (J. E. Biesecker, q)

Puerto Rico (w, San Juan):

Salinity reconnaissance and monitoring system, south coast (J. R. Díaz)

Water resources, Ponce area (N. E. McClymonds)

Rhode Island (w, Providence):

Ground water, Potowomut-Wickford area (J. Rosen-shein)

South Carolina (Columbia):**Ground water:****Coastal plain:**

Northeastern part, geology and ground-water resources (G. E. Siple, g)

Subsurface geology and hydrology (G. E. Siple, g)

Greenville County, geology and ground-water resources (N. C. Koch, g)

Leesville area, potential sand aquifers (G. E. Siple, W. D. Paradeses, g)

Water resources--Continued

South Carolina (Columbia)--Continued

Ground water--Continued

- Piedmont, alluvial aquifers (G. E. Siple, g)
- Savannah River Plant, operational effect on geologic and hydrologic environment (G. Siple, N. C. Koch, g)

Surface water:

- Analyses of streamflow characteristics (A. E. Johnson, J. S. Stallings, s)
- Compilation of streamflow records (A. E. Johnson, s)

South Dakota (Huron):

Ground water:

- Eastern part, hydrology of glacial drift in selected drainage basins: Big Sioux Basin from Sioux Falls to Brookings County line (M. J. Ellis, g)
- Studies of artesian wells and selected shallow aquifers (D. G. Adolphson, g)
- Beadle County (L. W. Howells, g)
- Campbell County (N. C. Koch, g)
- Clay County (J. C. Stephens, g)
- Dakota Sandstone (E. F. LeRoux, g)
- Pine Ridge Indian Reservation (M. J. Ellis, g)
- Rosebud Indian Reservation (M. J. Ellis, g)
- Standing Rock Indian Reservation, Bullhead, Little Eagle, and Wapala Indian villages (M. J. Ellis, g)

Tennessee (w, Chattanooga):

Ground water:

- Dickson County (R. H. Bingham)
- Germantown-Collierville area (D. J. Nyman)
- Highland Rim Plateau (O. T. Marsh)
- Lawrence County (R. H. Bingham)
- Lewis County (R. H. Bingham)
- Wayne County (R. H. Bingham)

Water resources:

- Memphis area (E. A. Bell)
- Montgomery County (W. J. Perry)
- Trace Creek (J. H. Criner, Jr.)

Texas (w, Austin):

Ground water:

- Atascosa County (R. C. Baker)
- Austin County (C. L. Wilson)
- Bastrop County (C. R. Follett)
- Bee County (B. N. Myers)
- Brazos River alluvium, occurrence and availability of ground water (J. G. Cronin)
- Brooks County (B. N. Myers)
- Ellis County (G. L. Thompson)
- El Paso area, continuing quantitative studies (M. E. Davis)
- Frio County (R. C. Baker)
- Galveston County (R. K. Gabrysch)
- Guadalupe County (G. H. Shafer)
- Harrison County (M. E. Broom)
- Houston County, occurrence and availability of ground water (G. H. Tarver)
- Houston district, continuing quantitative studies (R. K. Gabrysch)
- Jasper County (J. B. Wesselman)
- Kendall County (R. D. Reeves)
- Lee County (G. L. Thompson)
- Liberty County (R. B. Anders)
- Lower Red River valley (M. S. Bedinger, w, Little Rock, Ark.)

Water resources--Continued

Texas (w, Austin)--Continued

Ground water--Continued

- Maverick County (E. T. Baker, Jr.)
- Newton County (J. B. Wesselman)
- Nueces County (G. H. Shafer)
- Sabine County (R. B. Anders)
- San Antonio area (S. Garza)
- San Augustine (R. B. Anders)
- San Patricio County (G. H. Shafer)
- Tyler County (G. R. Tarver)
- Waller County (C. L. Wilson)
- Webb County (E. T. Baker, Jr.)
- Wood County (M. E. Broom)

Hydrologic investigations:

- Brazos, Colorado, San Antonio, and Trinity Rivers basins (C. R. Gilbert)
- Brazos River, tidal flow (R. E. Smith)
- Escondido Creek (F. W. Kennon)
- Guadalupe River, base-flow studies (J. T. Smith, H. L. Kunze)
- Little Elm Creek (E. E. Schroeder)
- Pin Oak Creek, small-watershed hydrology (J. T. Smith)

Utah (w, Salt Lake City):

Chemical characteristics of water resources of western Utah (O. Hattori)

Flood magnitude and frequency, The Great Basin (E. Butler)

Sedimentation of reservoirs, Sevier River basin, Mill Creek near Glenwood (R. E. Cabell)

Ground water:

- Reconnaissance of basins in western Utah--Snake Valley (J. W. Hood)
- Springs of Utah (J. C. Mundorff)
- Southwestern part, selected basins (G. W. Sandberg)
- Statewide ground-water conditions (T. Arnow)
- Colorado Plateaus, water from bedrock (R. D. Feltis)
- Juab Valley, northern part (L. J. Bjorklund)
- Sanpete Valley (G. B. Robinson, Jr.)
- Sevier Desert (R. W. Mower, R. D. Feltis)
- Sevier River basin between Yuba Dam and Leamington Canyon (L. J. Bjorklund)
- Sevier Valley, upper part (C. H. Carpenter, L. J. Bjorklund, G. B. Robinson)
- Tooele Valley (J. S. Gates)
- Utah Valley, southern part (C. H. Carpenter)

Surface water:

- Extension of streamflow records (J. K. Reid)
- Sevier Lake basin, reconnaissance of chemical-quality and fluvial-sediment characteristics of surface waters (D. C. Hahl, R. E. Cabell)

Water resources, Salt Lake County (W. V. Iorns)

Vermont:

Water resources, Connecticut River basin (D. J. Cederstrom, w, Boston, Mass.)

Virgin Islands (w, San Juan, P. R.):

Water resources (D. G. Jordan, O. J. Cosner)

Water supplies for Virgin Islands National Park (O. J. Cosner, D. G. Jordan)

Washington (w, Tacoma):

Influence of industrial and municipal wastes on estuarine and off-shore water quality (J. F. Santos)

Water resources--Continued

Washington (w, Tacoma)--Continued

Logging, effect on runoff, upper Green River watershed (D. Richardson)

Chehalis River basin, fluvial sediment transport (P. A. Glancy)

Columbia River basin, effect of fluvial sediments in recreational sites (P. R. Boucher, Pasco)

Grays Harbor, water quality (J. P. Beverage)

Lower Fleet Creek basin, hydrology (F. M. Veatch)

Ground water:

Influence of natural gas on ground water (L. B. Laird)

Statewide chemical quality of ground water (A. S. Van Denburgh)

Adams, Franklin, and Grant Counties (J. W. Bingham)

Island County (H. W. Anderson, Jr.)

King County, southwest (J. E. Luzier)

Mason County (D. Molenaar)

Odessa area, optimum ground-water withdrawal rate (A. A. Garrett)

Spokane County, northern (D. R. Cline)

Whitman County (K. L. Walters)

Water resources:

Colville River basin (W. R. Scott)

King County (D. Richardson)

Puget Sound and adjacent waters (D. Richardson)

West Virginia (Morgantown):

Ground water, Mason and Putnam Counties (B. M. Wilmoth, g)

Water resources, Monongahela River basin (P. E. Ward, g)

Wisconsin (Madison)

Ground water:

Kenosha and Racine Counties (R. D. Hutchinson, g)

Walworth County (J. H. Green, q)

Water resources:

Lower Wisconsin River valley (L. J. Hamilton, g; T. R. Dosch, s)

Upper Wisconsin River valley (R. W. Devaul, g)

Wyoming (g, Cheyenne; q, Worland):

Ground water:

Cheyenne area (L. J. McGreevy, g)

Green River structural basin, reconnaissance (G. E. Welder, g)

Water resources--Continued

Wyoming (g, Cheyenne; q, Worland)--Continued

Ground water--Continued

Laramie County (M. E. Lowry, g; J. R. Tilstra, q)

Wind River basin, reconnaissance (H. A. Whitcomb, g)

Wind River Indian Reservation (L. J. McGreevy, g)

Other countries:

Afghanistan, surface water, western part (A. O. Westfall, w, Kabul)

Brazil:

Hydrogeology, northeastern part (S. L. Schoff, w, Recife)

Surface water, northeastern part (L. J. Snell, w, Recife)

Libya, ground water, nationwide investigation and pilot development (J. R. Jones, w, Tripoli)

Nepal, surface water, nationwide investigations (W. W. Evett, w, Katmandu)

Nigeria:

Ground water, national program (D. A. Phoenix, w, Kaduna)

Hydrology:

Artesian water in Chad Basin (R. E. Miller, w, Maiduguri)

Sokoto Basin (W. Ogilbee, w, Skoto)

Surface water, northern part (C. R. Seiber, w, Kaduna)

Okinawa:

Ground water, southern part (D. A. Davis, g, Honolulu)

Sedimentation investigation, northern part (J. P. Beverage, q, M)

Pakistan, hydrologic investigations related to water-logging and salinity control in the Punjab Region (M. J. Mundorff, w, Lahore)

Tunisia, ground-water investigations and hydrogeologic mapping (V. C. Fishel, w, Tunis)

Turkey, nationwide ground-water investigations—advisory (C. R. Murray, w, Ankara)

United Arab Republic (Egypt), ground-water investigation and pilot development in the New Valley project (R. L. Cushman, w, Cairo)

Zeolites, southeastern California (R. A. Shappard, D)

Zinc. See Lead and zinc.

HOW TO ORDER U.S. GEOLOGICAL SURVEY PUBLICATIONS

All books, maps (exclusive of topographic quadrangle maps), and related publications published by the Geological Survey are listed in "Publications of the Geological Survey, 1879-1961," and in yearly supplements that keep the catalog up to date. New publications are announced each month in "New Publications of the Geological Survey." All of these lists of publications are free upon request to the GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242. They may be consulted at many public and educational-institution libraries, and at the Geological Survey offices named below.

Books, maps, charts, folios, and atlases that are out of print can no longer be purchased from any official source. They may be consulted at many libraries, and some can be purchased from dealers in second-hand books.

Ordering book reports

Professional papers, bulletins, water-supply papers, and miscellaneous book publications can be purchased from the SUPERINTENDENT OF DOCUMENTS, GOVERNMENT PRINTING OFFICE, WASHINGTON, D.C. 20402. Prepayment is required and may be made by money order or check payable to that office, or in cash—exact amount—at sender's risk. Postage stamps are not accepted. Book publications also may be purchased on an *over-the-counter* basis from the Geological Survey Public Inquiries Office listed on p. A211.

Circulars which are not out of print may be obtained free on application to the GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242.

Ordering maps and charts

Maps, charts, folios, and hydrologic atlases are sold by the Geological Survey. *Mail* orders for those covering areas east of the Mississippi River should be addressed to the GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242, and for areas west of the Mississippi River to the GEOLOGICAL SURVEY, FEDERAL CENTER, DENVER, COLO. 80225. Remittances should be sent by check or money order made payable to the Geological Survey or in cash—exact amount—at the sender's

risk. Postage stamps are not accepted. Retail prices are quoted in lists of publications and, for topographic maps, in indexes to topographic mapping for individual States. On an order amounting to \$10 or more at the retail price, 20 percent discount is allowed; on orders of \$60 or more, 40 percent is allowed. Most geologic maps are sent folded in envelopes unless flat copies are requested. Topographic maps are sent flat, except for orders of six maps or less; however, flat copies will also be sent for small orders on request. These publications also may be obtained on an area basis, by *over-the-counter sale* (but not by mail) from the Geological Survey Public Inquiries Offices listed on p. A211. Residents of Alaska may order Alaska maps from the GEOLOGICAL SURVEY, 310 FIRST AVE., FAIRBANKS, ALASKA 99701.

Indexes to topographic-map coverage of the various States and Guam and American Samoa are released periodically and are free on application. The release of revised indexes is announced in the monthly list of new publications of the Geological Survey. Each State index shows the areas mapped, with listings of special and United States maps, and gives lists of Geological Survey offices from which maps may be purchased and also of local agents who sell the maps.

Advance material available from current topographic mapping is indicated on quarterly releases of State index maps. This material, including such items as aerial photography, geodetic-control data, and preliminary maps in various stages of preparation and editing, is available for purchase. Information concerning the ordering of these items is given on the quarterly issues. Requests for indexes or inquiries concerning availability of advance materials should be directed to the MAP INFORMATION OFFICE, GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242.

Surface-water records of [State]

Pending publication of surface-water records at 5-year intervals in the Water-Supply Paper series on the basis of drainage basins, streamflow records and related data are being released in annual re-

ports on the basis of State boundaries. Distribution of these basic-data reports, which are free on request, is limited and primarily for local needs. Those interested should write to the GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242, specifying the State or States for which records are needed.

State water-resources investigations folders

A series of 8- by 10½-inch folders entitled "Water Resources Investigations in [State]" is a project of the Water Resources Division to inform the public about its current program in the 50 States and Puerto Rico, and Guam and American Samoa. As the State programs change, the folders are revised. Folders for all 50 States and Puerto Rico are available free on request to the GEOLOGICAL SURVEY, WASHINGTON, D.C. 20242.

Open-file reports

Open-file reports include unpublished manuscript reports, maps, and other material made available for public consultation and use. Arrangements can generally be made to reproduce them at private expense. The date and places of availability for consultation by the public are given in press releases or other forms of public announcement. In general, open-file reports are placed in one or more of the three Geological Survey libraries: ROOM 1033, GENERAL SERVICES BLDG., WASHINGTON, D.C.; BLDG. 25, FEDERAL CENTER, DENVER, COLO.; and 345 MIDDLEFIELD ROAD, MENLO PARK, CALIF. Other depositories may include one or more of the Geological Survey offices listed on pages A211 to A215, or interested State agencies. Many open-file reports are replaced later by formally printed publications.

CONTENTS OF GEOLOGICAL SURVEY RESEARCH 1965, CHAPTERS B, C, AND D

Listed below are the contents of Professional Papers 525-B, -C, and -D, comprising 128 articles, many of which are cited in the preceding pages. References to chapters B, C, and D are given in the test in the following form: Pakiser (p. B1-B8); the chapter is identified by the letter preceding the page number.

CHAPTER B

GEOLOGIC STUDIES

Geophysics

The basalt-eclogite transformation and crustal structure in the Western United States, by L. C. Pakiser.....	B1
Geophysical evidence of a caldera at Bonanza, Colo., by D. E. Karig.....	9
Hawaiian seismic events during 1963, by R. Y. Koyanagi and E. T. Endo.....	13

Economic geology

Correlation of zinc abundance with stratigraphic thickness variations in the Kingsport Formation, West New Market area, Mascot-Jefferson City mining district, Tennessee, by Helmuth Wedow, Jr., and J. R. Marie.....	17
Mercury-bearing antimony deposit between Big Creek and Yellow Pine, central Idaho, by B. F. Leonard.....	23
Suggestions for prospecting for evaporite deposits in southwestern Virginia, by C. F. Withington.....	29
Low-volatile bituminous coal of Mississippian age on the Lisburne Peninsula, northwestern Alaska, by I. L. Tailleux.....	34
Distribution of gravel in the Patuxent Formation in the Beltsville quadrangle, Prince Georges and Montgomery Counties, Md., by C. F. Withington.....	39

Structural geology and stratigraphy

Structure of the Timber Mountain caldera, southern Nevada, and its relation to basin-range structure, by R. L. Christiansen, P. W. Lipman, P. P. Orkild, and F. M. Byers, Jr.....	43
The Clinchport thrust fault—a major structural element of the southern Appalachian Mountains, by L. D. Harris.....	49
Correlation of Cretaceous and lower Tertiary rocks near Livingston, Mont., with those in other areas of Montana and Wyoming, by A. E. Roberts.....	54

Paleontology

An Upper Cretaceous deposit in the Appalachian Mountains, by R. H. Tschudy.....	64
Classification of the superfamily Healdiacea and the genus <i>Pseudophanasymmetria</i> Sohn and Berdan, 1952 (Ostracoda), by I. G. Sohn.....	69
Stratigraphic distribution of the Late Cambrian mollusk <i>Matthevia</i> Walcott, 1885, by E. L. Yochelson, J. F. McAllister, and Anthony Reso.....	73

Petrology and petrography

A classification for quartz-rich igneous rocks based on feldspar ratios, by J. T. O'Connor.....	79
Gold Flat Member of the Thirsty Canyon Tuff—a pantellerite ash-flow sheet in southern Nevada, by D. C. Noble.....	85
Precipitation and recycling of phosphate in the Florida land-pebble phosphate deposits, by Z. S. Altschuler.....	91

Geochemistry

Distribution of tantalum in some igneous rocks and coexisting minerals of the Southern California batholith, by David Gottfried and J. I. Dinnin.....	96
Metallic copper in stony meteorites, by M. B. Duke and Robin Brett.....	101

Geochronology

K-Ar and Rb-Sr ages of biotite from the Middle Jurassic part of the Carmel Formation, Utah, by R. F. Marvin, J. C. Wright, and F. G. Walthall.....	104
Potassium-argon ages of some plutonic rocks, Tenakee area, Chichagof Island, southeastern Alaska, by M. A. Lanphere, R. A. Loney, and D. A. Brew.....	108

Geomorphology and Pleistocene geology

Page

A large transitional rock glacier in the Johnson River area, Alaska Range, by H. L. Foster and G. W. Holmes.....	B112
Dissected gravels of the Río Copiapó valley and adjacent coastal area, Chile, by Kenneth Segerstrom.....	117
Earth cracks—a cause of gullyng, by William Kam.....	122
Glacial deposits of Nebraskan and Kansan age in northern Kentucky, by M. M. Leighton and L. L. Ray.....	126
Age and origin of the Puget Sound trough in western Washington, by D. R. Crandell, D. R. Mullineaux, and H. H. Waldron.....	132

Glaciology

Rate of ice movement and estimated ice thickness in part of the Teton Glacier, Grand Teton National Park, Wyo., by J. C. Reed, Jr.....	137
--	-----

Physical properties of rocks

Use of magnetic susceptibility and grain density in identification of basalt flows at the Nevada Test Site, by K. A. Sargent.....	142
The best value of porosity of lapilli tuff from the Nevada Test Site, by G. E. Manger.....	146

Analytical techniques

Gamma activation device for low-level beryllium analysis, by W. W. Vaughn, W. G. Cramer, and W. N. Sharp.....	151
Use of X-ray fluorescence in determination of selected major constituents in silicates, by H. J. Rose, Jr., Frank Cuttitta, and R. R. Larson.....	155
The problem of automatic plate reading and computer interpretation for spectrochemical analysis, by A. W. Helz.....	160
A platinum-lined bomb for the high-temperature decomposition of refractory minerals, by Irving May, J. J. Rowe, and Raymond Letner.....	165

HYDROLOGIC STUDIES**Quality of water**

Rapid field and laboratory determination of phosphate in natural water, by M. J. Fishman and M. W. Skougstad.....	167
Leachable silica and alumina in streambed clays, by E. C. Mallory, Jr.....	170

Surface water

Growth of salt cedar (<i>Tamarix gallica</i>) in the Pecos River near the New Mexico-Texas boundary, by R. U. Grozier.....	175
Effect of Great Swamp, N.J., on streamflow during base-flow periods, by E. G. Miller.....	177

Engineering hydrology

Shortening and protrusion of a well casing due to compaction of sediments in a subsiding area in California, by J. F. Poland and R. L. Ireland.....	180
---	-----

Experimental and theoretical hydrology

Use of sodium iodide to trace underground water circulation in the hot springs and geysers of the Daisy geyser group, Yellowstone National Park, by J. J. Rowe, R. O. Fournier, and G. W. Morey.....	184
The algebra of stream-order numbers, by A. E. Scheidegger.....	187

PUBLICATION TECHNIQUES**Technical illustrations**

Photographic copying using reflection-transmission illumination, by E. P. Krier.....	190
--	-----

CHAPTER C

GEOLOGIC STUDIES

Structural geology

Stratigraphic data bearing on inferred pull-apart origin of Gem Valley, Idaho, by S. S. Oriel, D. R. Mabey, and F. C. Armstrong.....	Page C1
Structure of a ray crater at Henbury, Northern Territory, Australia, by D. J. Milton and F. C. Michel.....	5
Folding of the Nahant gabbro, Massachusetts, by C. A. Kaye.....	12
Relation of laccolithic intrusion to faulting in the northern part of the Barker quadrangle, Little Belt Mountains, Mont., by I. J. Witkind.....	20

Mineralogy and petrology

Composition of jadeitic pyroxene from the California metagraywackes, by R. G. Coleman.....	25
X-ray determinative curve for Hawaiian olivines of composition For_{76-88} , by K. J. Murata, Harry Bastron, and W. W. Brannock.....	35
Upper Triassic undevitrified volcanic glass from Hound Island, Keku Strait, southeastern Alaska, by D. A. Brew and L. J. P. Muffer.....	38

Geophysics

Seismic-refraction measurements of crustal structure between American Falls Reservoir, Idaho, and Flaming Gorge Reservoir, Utah, by Ronald Willden.....	44
Seismic fluctuations in an open artesian water well, by J. D. Bredehoeft, H. H. Cooper, Jr., I. S. Papadopoulos, and R.R. Bennett.....	51

Geochemistry

Fractionation of uranium isotopes and daughter products in uranium-bearing sandstone, Gas Hills, Wyo., by J. N. Rosholt, Jr., and C. P. Ferreira.....	58
---	----

Geochronology

Pliocene age of the ash-flow deposits of the San Pedro area, Chile, by R. J. Dingman.....	63
Jurassic age of a mafic igneous complex, Christian quadrangle, Alaska, by H. N. Reiser, M. A. Lanphere, and W. P. Brosgé.....	68

Paleontology and stratigraphy

First occurrence of graptolites in the Klamath Mountains, Calif., by Michael Churkin, Jr.....	72
A proposed revision of the subalkaline intrusive series of northeastern Massachusetts, by R. O. Castle.....	74
Gneissic rocks in the South Groveland quadrangle, Essex County, Mass., by R. O. Castle.....	81
Stratigraphy of the upper part of the Yakima Basalt in Whitman and eastern Franklin Counties, Wash., by J. W. Bingham and K. L. Walters.....	87
Previously undescribed Middle(?) Ordovician, Devonian(?), and Cretaceous(?) rocks, White Mountain area, near McGrath, Alaska, by C. L. Sainsbury.....	91
Presence of the ostracode <i>Drepanellina clarki</i> in the type Clinton (Middle Silurian) in New York State, by J. M. Berdan and D. H. Zenger.....	96
Miocene macrofossils of the southeastern San Joaquin Valley, Calif., by W. O. Addicott.....	101

Glacial and Pleistocene geology

New evidence on Lake Bonneville stratigraphy and history from southern Promontory Point, Utah, by R. B. Morrison.....	110
Glaciation in the Nabesna River area, upper Tanana River valley, Alaska, by A. T. Fernald.....	120
Recent history of the upper Tanana River lowland, Alaska, by A. T. Fernald.....	124
Maximum extent of late Pleistocene Cordilleran glaciation in northeastern Washington and northern Idaho, by P. L. Weis and G. M. Richmond.....	128
Birch Creek pingo, Alaska, by D. B. Krinsley.....	133
Quaternary stratigraphy of the Durango area, San Juan Mountains, Colo., by G. M. Richmond.....	137
Distribution of Pleistocene glaciers in the White Mountains of California and Nevada, by V. C. LaMarche, Jr.....	144
Landslide origin of the type Cerro Till, southwestern Colorado, by R. G. Dickinson.....	147

Geomorphology

Geomorphic significance of a Cretaceous deposit in the Great Valley of southern Pennsylvania, by K. L. Pierce.....	152
--	-----

Economic geology	Page
Some potential mineral resources of the Atlantic continental margin, by K. O. Emery.....	C157
Marine geology	
Composition of basalts dredged from seamounts off the west coast of Central America, by C. G. Engel and T. E. Chase..	161
Data analysis	
On the statistics of the orientation of bedding planes, grain axes, and similar sedimentological data, by A. E. Scheidegger..	164
Analytical methods	
A spectrophotometric method for the determination of traces of gold in geologic materials, by H. W. Lakin and H. M. Nakagawa.....	168
A field method for the determination of silver in soils and rocks, by H. M. Nakagawa and H. W. Lakin.....	172

HYDROLOGIC STUDIES

Quality of water	
Diurnal variations of the chemical quality of water in two prairie potholes in North Dakota, by H. T. Mitten.....	176
Physical and chemical hydrology of Great Salt Lake, Utah, by D. C. Hahl, M. T. Wilson, and R. H. Langford.....	181
A comparison of the chemical composition of rainwater and ground water in western North Carolina, by R. L. Laney....	187
Light-dependent quality changes in stored water samples, by K. V. Slack and D. W. Fisher.....	190
Patterns of dissolved oxygen in a thermally loaded reach of the Susquehanna River, Pa., by K. V. Slack and F. E. Clarke..	193
Limnology and surface water	
Effect of land use on the low flow of streams in Rappahannock County, Va., by H. C. Riggs.....	196
Seasonal erasure of thermal stratification in Pretty Lake, Ind., by J. F. Ficke.....	199
Ground water	
A Miocene(?) aquifer in the Parker-Blythe-Cibola area, Arizona and California, by D. G. Metzger.....	203
Relation between ground water and surface water	
Use of specific conductance to distinguish two base-flow components in Econfinia Creek, Fla., by L. G. Toler.....	206
Relation between chemical quality and water discharge in Spring Creek, southwestern Georgia, by L. G. Toler.....	209
Hydrologic instrumentation	
A portable sampler for collecting water samples from specific zones in uncased or screened wells, by R. N. Cherry.....	214

CHAPTER D

GEOLOGIC STUDIES

	Page
Geochronology	
Implications of new radiometric ages in eastern Connecticut and Massachusetts, by Robert Zartman, George Snyder, T. W. Stern, R. F. Marvin, and R. C. Buckman	D1
Reconnaissance of mineral ages of plutons in Elko County, Nev., and vicinity, by R. R. Coats, R. F. Marvin, and T. W. Stern	11
Jurassic plutonism in the Cook Inlet region, Alaska, by R. L. Detterman, B. L. Reed, and M. A. Lanphere	16
Age and distribution of sedimentary zircon as a guide to provenance, by R. S. Houston and J. F. Murphy	22
Carboniferous isotopic age of the metamorphism of the Salmon Hornblende Schist and Abrams Mica Schist, southern Klamath Mountains, Calif., by M. A. Lanphere and W. P. Irwin	27
Radiocarbon dates from Iliamna Lake, Alaska, by R. L. Detterman, B. L. Reed, and Meyer Rubin	34
Mineralogy and petrology	
Magnetic spherules, colored corundum, and other unusual constituents of a heavy beach sand, Martha's Vineyard, Mass., by C. A. Kaye and M. E. Mrose	37
Zeolitic authigenesis of tuffs in the Ricardo Formation, Kern County, southern California, by R. A. Sheppard and A. J. Gude 3d	44
Thorium-bearing microcline-rich rocks in the southern Caballo Mountains, Sierra County, N. Mex., by M. H. Staatz, J. W. Adams, and N. M. Conklin	48
Volcanic origin of flint clay in the Fire Clay coal bed, Breathitt Formation, eastern Kentucky, by V. M. Seiders	52
Prehnite and hydrogarnet(?) in Precambrian rocks near Boulder, Colo., by C. T. Wrucke	55
Geochemistry	
Geochemical prospecting in the Browns Canyon fluor spar district, Chaffee County, Colo., by R. E. Van Alstine	59
Heat and free energy of formation of herzenbergite, troilite, magnesite, and rhodochrosite calculated from equilibrium data, by R. A. Robie	65
Extractable organic material in nonmarine and marine shales of Cretaceous age, by H. A. Tourtelot and I. C. Frost	73
Composition of magnetite as related to type of occurrence, by Michael Fleischer	82
Geophysics	
Seismic study of crustal structure in the Southern Rocky Mountains, by W. H. Jackson and L. C. Pakiser	85
Thermal features at Mount Rainier, Wash., as revealed by infrared surveys, by R. M. Moxham, D. R. Crandell, and W. E. Marlatt	93
Seismic investigations in the Harwich and Dennis quadrangles, Cape Cod, Mass., by R. N. Oldale and C. R. Tuttle	101
Structural geology	
Evidence of large strike-slip displacement along a fault in the southern Salinas Valley, Calif., by D. L. Durham	106
Stratigraphy and paleontology	
Upper Precambrian and Paleozoic stratigraphy and structure of the Neptune Range, Antarctica, by D. L. Schmidt, P. L. Williams, W. H. Nelson, and J. R. Ege	112
Occurrence and stratigraphic significance of <i>Oldhamia</i> , a Cambrian trace fossil, in east-central Alaska, by Michael Churkin, Jr., and E. E. Brabb	120
Late Devonian and Early Mississippian age of the Woodford Shale in Oklahoma, as determined from conodonts, by W. H. Hass and J. W. Huddle	125
Gray Bull and Lysite faunal zones of the Willwood Formation in the Tatman Mountain area, Bighorn Basin, Wyo., by W. L. Rohrer and C. L. Gazin	133
Tongues of the Green River and Wasatch Formations in the southeastern part of the Green River Basin, Wyo., by W. C. Culbertson	139

CHAPTER D

	Page
Economic geology	
Possible buried mineralized areas in Nye and Esmeralda Counties, Nev., R. E. Anderson, E. B. Ekren, and D. L. Healey	D144
Outlook for resumption of diatomite mining in southern Maryland and eastern Virginia, by M. M. Knechtel and J. W. Hosterman	151
Evaluation of the Martinsburg Shale and two younger formations as sources of lightweight aggregate in the Delaware River area, Pennsylvania-New Jersey, by A. A. Drake, Jr., M. V. Denny, and H. P. Hamlin	156
Lithium-bearing bentonite deposit, Yavapai County, Ariz., by J. J. Norton	163
Pleistocene geology	
Subsurface stratigraphy of glacial drift at Anchorage, Alaska, by F. W. Trainer and R. M. Waller	167
Marine geology	
Basins of the Gulf of Maine, by Elazar Uchupi	175
Analytical methods	
Automatic sample changer and controller for an X-ray quantometer, by Leonard Shapiro and Camillo Massoni	178
Selective removal of Po^{210} from aged radium standards, by K. W. Edwards	184
Use of bathocuproine in the quantitative determination of copper in soils, sediments, and rocks, by G. A. Nowlan	189
Use of arsenazo III in determination of thorium in rocks and minerals, by Irving May and L. B. Jenkins	192
HYDROLOGIC STUDIES	
Surface water	
Changes in character of unit hydrographs, Sharon Creek, Calif., after suburban development, by J. R. Crippen	196
A method of estimating mean runoff from ungaged basins in mountainous regions, by H. C. Riggs and D. O. Moore ..	199
Ground water	
Relation of permeability to particle size in a glacial-outwash aquifer at Piketon, Ohio, by S. E. Norris and R. E. Fidler	203
Relation between ground water and surface water	
Computation of ground-water discharge to streams during floods, or to individual reaches during base flow, by use of specific conductance, by G. R. Kunkle	207
Relation of ground-water inflow and of bank and channel storage to streamflow pickup in the Santa Fe River, Fla., by W. E. Clark	211
Quality of water	
Changes in quality of water in the Passaic River at Little Falls, N. J., as shown by long-term data, by P. W. Anderson and S. D. Faust	214
Chemical distinction between ground water of four sedimentary units on the Kitsap Peninsula and adjacent islands, Washington, by A. S. Van Denburgh	219
Hydrologic instrumentation	
Multiple hydrologic-parameter recording on a digital recorder, by R. N. Cherry	222
Effects of sample and fluorometer-compartment temperatures on fluorometer readings, by Bernard Dunn and D. E. Vaupel	225

PUBLICATIONS IN FISCAL YEAR 1965

A complete list of abstracts, papers, reports, and maps (exclusive of topographic maps) by U.S. Geological Survey authors published or otherwise released to the public during fiscal year 1965 (July 1, 1964–June 30, 1965) is given below. Publications are listed alphabetically by senior author. Each citation is identified by a number: for example, 1–64, which indicates the first entry for that author for 1964. The number is followed by the names of coauthors and the citation itself. References to this list are identified in the preceding text by author and serial number: for example, Adams and Boggess (1–64).

The list also includes some publications that were not listed in either Professional Papers 475–A or 501–A.

ADAMS, J. K.

- 1-64. (and BOGGESS, D. H.) Water-table, surface-drainage, and engineering soils map of the Ellendale quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-101, 1964.
- 2-64. (and BOGGESS, D. H.) Water-table, surface-drainage, and engineering soils map of the Harbeson quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-108, 1964.
- 3-64. (and BOGGESS, D. H.) Water-table, surface-drainage, and engineering soils map of the Hickman area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-100, 1964.
- 4-64. (and BOGGESS, D. H.) Water-table, surface-drainage, and engineering soils map of the Sharptown area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-84, 1964.
- 5-64. (and BOGGESS, D. H.) Water-table, surface-drainage, and engineering soils map of the Trap Pond area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-120, 1964.
- 6-64. (BOGGESS, D. H., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Clayton area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-83, 1964.
- 7-64. (BOGGESS, D. H., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Frankford area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-119, 1964[1965].
- 8-64. (BOGGESS, D. H., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Seaford East quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-106, 1964.
- 9-64. (BOGGESS, D. H., and DAVIS, C. F.) Water-table, surface-drainage, and engineering soils map of the Dover quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-139, 1964[1965].
- 10-64. (BOGGESS, D. H., and DAVIS, C. F.) Water-table, surface-drainage, and engineering soils map of the Lewes area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-103, 1964.

ADAMS, J. W.

- 1-64. Metallic mineral resources: Thorium and the rare earths, *in* Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 77-80, 1964; *also in* Mineral and water resources of Utah: *ibid*, p. 115-119, 1964.

ADAMS, J. W. --Continued

- 2-64. Metallic mineral resources: Titanium, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 119-120, 1964.
- 3-64. Miscellaneous metals: Rare earths, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., p. 127-132, 1964.
- 4-64. Nonmetallic and industrial minerals: Pegmatite minerals, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 169-173, 1964.
- 5-64. Nonmetallic and industrial minerals and materials resources: Pegmatite minerals, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 194-195, 1964.
- 1-65. Rare earths, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 88th, 1st sess., Comm. Print, p. 234-237, 1965.
- 2-65. The visible region absorption spectra of rare-earth minerals: *Am. Mineralogist*, v. 50, no. 3-4, p. 356-366, 1965.

ADDICOTT, W. O.

- 1-64. A late Pleistocene invertebrate fauna from southwestern Oregon: *Jour. Paleontology*, v. 38, no. 4, p. 650-661, 1964.
- 2-64. Pleistocene invertebrates from the Dume terrace, western Santa Monica Mountains, California: *Southern California Acad. Sci. Bull.*, v. 63, pt. 3, p. 141-150, 1964.
- 1-65. The enigmatic late Cenozoic gastropod *Schizopyga californiana* [abs.]: *Am. Malacological Union Ann. Rept.* for 1964, p. 44, 1965.
- 2-65. Some western American Cenozoic gastropods of the genus *Nassarius*: U.S. Geol. Survey Prof. Paper 503-B, p. B1-B21, 1965.

ADKISON, W. L.

- 1-64. Detailed description of sample logs for two wells in Shawnee County, Kansas: U.S. Geol. Survey open-file report, 34 p., 1964.

ADLER, Isidore

- 1-63. Electron-probe analysis of minerals: *Am. Soc. for Testing Materials, Spec. Tech. Pub.* 349, p. 183-192, 1963.
- 1-64. Standards in electron probe analysis of minerals, *in* Advances in X-ray analysis, v. 7, Denver Univ., Denver Research Inst., Application X-ray analysis, 12th Ann. Conf., Denver, Colo., 1963, *Proc.* New York, Plenum Press, p. 426-432, 1964.

ADOLPHSON, D. G.

- 1-64. (and ELLIS, M. J.) Basic hydrogeologic data, Skunk Creek-Lake Madison drainage basin, South Dakota: South Dakota Geol. Survey Water Resources Rept. 3, 70 p., 1964.

AKERS, J. P.

- 1-64. Geology and ground water in the central part of Apache County, Arizona: U.S. Geol. Survey Water-Supply Paper 1771, 107 p., 1964[1965].

ALBEE, A. L.

- 1-65. Chemical and modal analyses of the pre-Upper Silurian quartz monzonite and the post-Lower Devonian granodiorite, Attean quadrangle, Somerset County, Maine: U.S. Geol. Survey open-file report, 3 p., 1965.

ALBEE, H. F.

- 1-65. (and CULLINS, H. L.) Preliminary geologic map of the Poker Peak and Palisades Reservoir quadrangles, Bonneville County, Idaho, and Lincoln County, Wyoming: U.S. Geol. Survey open-file report, 1965.

ALBERS, J. P.

- 1-34. Structural and stratigraphic environment of granitic plutons in Esmeralda County, Nevada, in *Advancing Frontiers in Geology and Geophysics*, a volume in honour of M. S. Krishnan: Hyderabad, India, Osmania Univ. Press, p. 351-360, 1964[1965].

- 2-64. Tertiary and Quaternary rocks, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 30-32, 1964.

ALBIN, D. R.

- 1-64. Resume of the ground-water resources of Bradley, Calhoun, and Ouachita Counties, Arkansas: Arkansas Geol. Conserv. Comm. Water Resources Summary, no. 1, 5 p., 1964.

ALEXANDER, W. H., Jr.

- 1-64. (MYERS, B. N., and DALE, O. C.) Reconnaissance of the ground-water resources of the Guadalupe, San Antonio, and Nueces River basins, Texas: Texas Water Comm. Bull. 6409, 106 p., 1964.

ALLEN, H. E.

- 1-64. (and MAY, V. J.) Floods in Harvey quadrangle, Illinois: U.S. Geol. Survey Hydrol. Inv. Atlas HA-90, 1964.

- 2-64. (and MAY, V. J.) Floods in Lombard quadrangle, Illinois: U.S. Geol. Survey Hydrol. Inv. Atlas HA-143, 1964.

- 3-64. (and MAY, V. J.) Floods in West Chicago quadrangle, Northeastern Illinois: U.S. Geol. Survey open-file report, 9 p., 1964.

- 4-64. (and WYERMAN, Theodore A.) Floods in Joliet quadrangle, Illinois: U.S. Geol. Survey Hydrol. Inv. Atlas HA-89, 1964.

- 1-65. Floods in Blue Island quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 8 p., 1965.

- 2-65. Floods in Tinley Park quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 7 p., 1965.

- 3-65. (and MAY, V. J.) Floods in Naperville quadrangle, Illinois: U.S. Geol. Survey Hydrol. Inv. Atlas HA-154, map, 1965.

ALLEN, W. B.

- 1-65. (HAHN, G. W., and BRACKLEY, R. A.) Availability of ground water, upper Pawcatuck River basin, Rhode Island: U.S. Geol. Survey open-file report, 133 p., 1965.

ALLINGHAM, J. W.

- 1-64. Low-amplitude aeromagnetic anomalies in southeastern Missouri: *Geophysics*, v. 29, no. 4, p. 537-552, 1964.

ALTO, B. R.

- 1-65. (FULTON, R. S., and HAIGLER, L. B.) Salines, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 299-306, 1965.

- 2-65. (FULTON, R. S., and HAIGLER, L. B.) The potash industry, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 306-309, 1965.

ALTSCHULER, Z. S.

- 1-64. (CATHCART, J. B., and YOUNG, E. J.) Geology and geochemistry of the Bone Valley Formation and its phosphate deposits, west central Florida—Guidebook, Field Trip no. 6, Geological Society of America Ann. Mtg., Miami Beach, Nov. 1964: [New York, Geological Society of America], 68 p., 1964.

- 2-64. (DWORNIK, E. J., and KRAMER, Henry) Genesis of kaolinite from montmorillonite by weathering—Structural and morphological evidence of transformation [abs.], in *Clays and clay minerals*, 12th Natl. Conf., Atlanta, Ga., 1963, Proc.: New York, Macmillan Co., p. 197, 1964.

- 1-65. Precipitation and recycling of phosphate in the Florida land-pebble phosphate deposits, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B91-B95, 1965.

- 2-65. (and BOUDREAU, C. E.) A mineralogical and chemical study of the leached zone of the Bone Valley Formation of Florida: A progress report: U.S. Geol. Survey open-file report, TEI-102, 89 p., 1965.

ANDERSON, C. A.

- 1-65. Precambrian geology of the Prescott-Jerome area [abs.]: Arizona Geol. Soc. Mtg., Tucson, May 4, 1965 [abstract distributed at meeting - no formal publication.]

ANDERSON, A. L.

- 1-64. (and VAN ALSTINE, R. E.) Mineral resources: Fluorspar, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 79-84, 1964.

ANDERSON, L. A.

- 1-64. (ZANDLE, G. L., and others) Aeromagnetic map of Norman and part of Mahnom County, Minnesota: U.S. Geol. Survey Geophys. Inv. Map GP-325, 1964. [Originally published 1962.]

- 1-65. Electrical resistivity of the deep crust in the Columbia River Basin area: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 71, 1965.

ANDERSON, P. W.

- 1-65. (and GEORGE, J. R.) Water-quality characteristics of New Jersey streams: U.S. Geol. Survey open-file report, 113 p., 1965.

ANDERSON, Warren

- 1-65. (LICHTLER, W. F., and JOYNER, B. F.) Control of lake levels in Orange County, Florida: *Florida Geol. Survey Inf. Circ.* 47, 15 p., 1965.

ANDREASEN, G. E.

- 1-64. (BROMERY, R. W., and GILBERT, F. P.) Aeromagnetic map of the Glencoe-Ripley area, Payne County, Oklahoma: U.S. Geol. Survey Geophys. Inv. Map GP-469, 1964.

- 2-64. (BROMERY, R. W., and GILBERT, F. P.) Aeromagnetic map of the Hominy area, Osage County, Oklahoma: U.S. Geol. Survey Geophys. Inv. Map GP-470, 1964.

ANDREASEN, G. E. --Continued

3-64. (GRANTZ, Arthur, ZIETZ, Isidore, and BARNES, D. F.) Geologic interpretation of magnetic and gravity data in the Copper River Basin, Alaska: U.S. Geol. Survey Prof. Paper 316-H, p. H135-H153, 1964.

4-64. (PITKIN, J. A., and PETRAFESO, F. A.) Aeromagnetic map of the Long Beach-Santa Ana area, Los Angeles and Orange Counties, California: U.S. Geol. Survey Geophys. Inv. Map GP-464, 1964.

5-64. (PITKIN, J. A., and PETRAFESO, F. A.) Aeromagnetic map of western Los Angeles and vicinity, California: U.S. Geol. Survey Geophys. Inv. Map GP-466, 1964.

1-65. (MITCHELL, C. M., and TYSON, N. S.) Aeromagnetic map of Tombstone and vicinity, Cochise and Santa Cruz Counties, Arizona: U.S. Geol. Survey open-file report, with explanation, 1965.

ANDREWS, G. W.

1-65. Late Quaternary geologic history of the lower Chipewa Valley, Wisconsin: Geol. Soc. America Bull., v. 76, no. 1, p. 113-124, 1965.

2-65. Review of "Diatom floras in the Quaternary of the Gota River Valley (western Sweden)" by Urve Miller: Jour. Paleontology, v. 39, no. 3, p. 512-513, 1965.

ANNELL, C. S.

1-63. A spectrographic method for the determination of Cs, Rb, and Li in tektites, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963, Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 175-186, 1963.

2-63. Quantitative micro-determination of the major constituents of tektites by emission spectroscopy of solutions, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963, Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 170-174, 1963.

1-64. (CUTTITTA, Frank, CHAO, E.C.T., and FLETCHER, J. D.) Minor elements in selected australasian tektites [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 81, 1964.

APPLIN, E. R.

1-64. A microfauna from the Coker Formation, Alabama: U.S. Geol. Survey Bull. 1160-D, p. D65-D70, 1964.

APPLIN, P. L.

1-65. (and APPLIN, E. R.) The Comanche Series and associated rocks in the subsurface in central and south Florida: U.S. Geol. Survey Prof. Paper 447, 84 p., 1965.

ARCHER, Robert

1-63. (and SHAUGHNESSY, J. A.) Water quality in the Delaware River Basin, New York: New York State Dept. Commerce Bull. 5, 68 p., 1963.

ARMSTRONG, F. C.

1-64. Mineral resources: Uranium, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 233-239, 1964.

1-65. Northwestward projection of the Paris thrust fault, southeastern Idaho [abs.]: Geol. Soc. America Spec. Paper 82, p. 317, 1965.

2-65. (and ORIEL, S. S.) Tectonic development of Idaho-Wyoming thrust belt [abs.]: Mtn. Geologist, v. 2, no. 1, p. 45-46, 1965.

ARMSTRONG, J. R.

1-65. (CRANDELL, D. R., EASTERBROOK, D. J., and NOBLE, J. B.) Late Pleistocene stratigraphy and chronology in southwestern British Columbia and northwestern Washington: Geol. Soc. America Bull., v. 76, no. 3, p. 321-330, 1965.

ARMSTRONG, J. R. --Continued

2-65. (CRANDELL, D. R., EASTERBROOK, D. J., and NOBLE, J. B.) Late Pleistocene stratigraphic chronology in southwestern British Columbia and northwestern Washington [abs.]: Geol. Soc. America Spec. Paper 82, p. 237, 1965.

ARNOW, Ted, and others

1-64. Developing a state water plan—Ground-water conditions in Utah, spring of 1964: Utah Water and Power Board Coop. Inv. Rept. 2, 104 p., 1964.

2-64. (LANGFORD, R. H., and WILSON, M. T.) Developing a state water plan—A basic water-resource data program for Utah: Utah Water and Power Board Coop. Inv. Rept. no. 1, 23 p., 1964.

1-65. Groundwater in the Jordan Valley, Salt Lake County, Utah: Utah State Engineer Water Circ. 1, 11 p., 1965.

AVERITT, Paul

1-64. Mineral fuels and associated resources: Coal, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 39-51, 1964.

1-65. The future of coal production in the Rocky Mountains region: Econ. Geology, v. 60, no. 2, p. 351-359, 1965.

AVRETT, J. R.

1-64. (and CARROON, L. E.) Temperature of Alabama streams: Alabama Geol. Survey Div. Water Resources Inf. Ser. 35, 165 p., 1964.

AYER, G. R.

1-64. Float studies and temperature observations, Cumberland Bay area of Lake Champlain: U.S. Geol. Survey open-file report, 75 p., 1964.

BACHMAN, G. O.

1-65. Introduction, in *Mineral and water resources in New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 11, 1965.

2-65. Mineral industry in New Mexico, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong., 89th, 1st sess., Comm. Print, p. 13-17, 1965.

BACK, William

1-65. (and BARNES, Ivan) Relation of electrochemical potentials and iron content to ground-water flow patterns: U.S. Geol. Survey Prof. Paper 498-C, p. C1-C16, 1965.

BAILEY, E. H.

1-64. Metallic mineral resources: Mercury, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 119-123, 1964.

2-64. Mineral resources: Mercury, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 119-123, 1964.

3-64. (IRWIN, W. P., and JONES, D. L.) Franciscan and related rocks, and their significance in the geology of western California: California Div. Mines and Geology Bull. 183, 177 p., 1964.

4-64. (and SMITH, R. M.) Mercury—its occurrence and economic trends: U.S. Geol. Survey Circ. 496, 11 p., 1964.

BAKER, A. A.

1-65. (CALKINS, F. C., CRITTENDEN, M. D., Jr., and BROMFIELD, C. S.) Geologic map of the Brighton quadrangle, Utah: U.S. Geol. Survey open-file report, 1 sheet, 1965.

BAKER, E. T., Jr.

1-64. Geology and ground-water resources of Hardin County, Texas: Texas Water Comm. Bull. 6406, 179 p., 1964.

1-65. Ground-water resources of Jackson County, Texas: U.S. Geol. Survey open-file report, 314 p., 1965.

BAKER, J. A.

- 1-64. Geology and ground-water resources of Geauga County, Ohio: U.S. Geol. Survey open-file report, 113 p., 1964.
- 2-64. (and HEALY, H. G., and HACKETT, O. M.) Geology and ground-water conditions in the Wilmington-Reading area, Massachusetts: U.S. Geol. Survey Water-Supply Paper 1694, 80 p., 1964.
- 1-65. (LANG, S. M., and THOMAS, M. P.) Geology and hydrology of the Hartford Research Center, CANEL site, Middletown, Connecticut: U.S. Geol. Survey Bull. 1133-G, p. G1-G42, 1965.

BAKER, R. C.

- 1-64. (and BROWN, J. B.) Annotated bibliography of ground-water publications and open-file reports of the Texas Water Commission and U.S. Geological Survey for Texas through August 1963: Texas Water Comm. Circ. 64-02, 87 p., 1964.
- 2-64. (and DALE, O. C.) Ground-water resources of the lower Rio Grande Valley area, Texas: U.S. Geol. Survey Water-Supply Paper 1653, 56 p., 1964.
- 3-64. (HUGHES, L. S., and YOST, I. D.) Natural sources of salinity in the Brazos River, Texas—with particular reference to the Croton and Salt Croton Creek basins: U.S. Geol. Survey Water-Supply Paper 1669-CC, p. CC1-CC81, 1964.

BALLANCE, W. C.

- 1-65. Ground-water levels in New Mexico, 1963 and water levels in artesian wells in the Roswell area for period of record: New Mexico State Engineer Basic Data Rept., 143 p., 1965.
- 2-65. New Mexico State Planning Office Report—Arkansas River basin: U.S. Geol. Survey open-file report, 32 p., 1965.
- 3-65. (and TITUS, F. B., Jr.) New Mexico State Planning Office Report—Southern High Plains: U.S. Geol. Survey open-file report, 20 p., 1965.

BALSLEY, J. R.

- 1-64. (MEUSCHKE, J. L., and KIRBY, J. R.) Aeromagnetic map of parts of Matchwood, Ontonagon, Rockwood, and Greenwood quadrangles, Ontonagon County, Michigan: U.S. Geol. Survey open-file report, 1964.
- 1-65. (and BROMERY, R. W.) Aeromagnetic map of the Big Moose quadrangle, Herkimer and Hamilton Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-504, 1965.
- 2-65. (and BROMERY, R. W.) Aeromagnetic map of parts of the Bolton Landing, Glens Falls, and Whithall quadrangles, Warren and Washington Counties, New York, and Rutland County, Vermont: U.S. Geol. Survey Geophys. Inv. Map GP-509, 1965.
- 3-65. (and BROMERY, R. W.) Aeromagnetic map of parts of the Elizabethtown and Port Henry quadrangles, Essex County, New York: U.S. Geol. Survey Geophys. Inv. Map GP-506, 1965.
- 4-65. (and BROMERY, R. W.) Aeromagnetic map of part of the Lowville quadrangle, Lewis County, New York: U.S. Geol. Survey Geophys. Inv. Map GP-508, 1965.
- 5-65. (and BROMERY, R. W.) Aeromagnetic map of the McKeever quadrangle and part of the Port Leyden quadrangle, north-central New York: U.S. Geol. Survey Geophys. Inv. Map GP-510, 1965.
- 6-65. (and BROMERY, R. W.) Aeromagnetic map of the Number Four quadrangle, Herkimer and Lewis Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-502, 1965.

BALSLEY, J. R.--Continued

- 7-65. (and BROMERY, R. W.) Aeromagnetic map of the Old Forge quadrangle and part of the West Canada Lakes quadrangle, Herkimer and Hamilton Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-501, 1965.
- 8-65. (and BROMERY, R. W.) Aeromagnetic map of parts of Paradox Lake and Ticonderoga quadrangles, Essex and Warren Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-507, 1965.
- 9-65. (and BROMERY, R. W.) Aeromagnetic map of the Raquette Lake quadrangle, Hamilton County, New York: U.S. Geol. Survey Geophys. Inv. Map GP-503, 1965.
- 10-65. (and PETRAFESO, F. A.) Aeromagnetic map of parts of western Iron, southwestern Baraga, southern Houghton, southeastern Ontonagon, and northeastern Gogebic Counties, Michigan: U.S. Geol. Survey open-file report, 1965.
- 11-65. (and SMITH, C. W.) Aeromagnetic map of parts of northeastern Ontonagon, southern Houghton, and northwestern Baraga Counties, Michigan: U.S. Geol. Survey open-file report, 1965.
- 12-65. (and VARGO, J. L.) Aeromagnetic map of part of eastern Baraga County, Michigan: U.S. Geol. Survey open-file report, 1965.

BARKER, Fred

- 1-64. Reaction between mafic magmas and pelitic schist, Cortlandt, New York: Am. Jour. Sci., v. 262, no. 5, p. 615-634, 1964.

BARNES, D. F.

- 1-64. Mineral fuel resources: Coal, in Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 77-94, 1964.
- 1-65. Gravity changes during the Alaskan earthquake [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 45-46, 1965.
- 2-65. The U.S. Geological Survey's program of Alaskan gravity observations: Am. Geophys. Union Trans., v. 46, no. 1, p. 231-233, 1965.
- 3-65. (and ALLEN, R. V.) Alaskan gravity surveys: Progress report [abs.]: Geol. Soc. America Spec. Paper 82, p. 353-354, 1965.
- 4-65. (and ALLEN, R. V.) Reconnaissance marine and shoreline geophysical data from the area south and west of Kodiak Island, Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 354, 1965.
- 5-65. (LUCAS, W. H., and ALLEN, R. V.) Gravity measurements over the southern Alaskan Continental Shelf [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 106, 1965.

BARNES, Harley

- 1-63. Summary of geology, Jangle Ridge quadrangle, Nye and Lincoln Counties, Nevada: U.S. Geol. Survey Tech. Letter NTS-53, 12 p., 1963.
- 2-63. (HINRICHS, E. N., McKEOWN, F. A., and ORKILD, P. P.) U.S. Geological Survey investigations of Yucca Flat, Nevada Test Site, Pt. A., Geology of the Yucca Flat area: U.S. Geol. Survey Tech. Letter NTS-45, 196 p., 1963.
- 1-65. (CHRISTIANSEN, R. L., and BYERS, F. M., Jr.) Geologic map of the Jangle Ridge quadrangle, Nye and Lincoln Counties, Nevada: U.S. Geol. Survey Geol. Quad. Map GQ-363, 1965.

BARNES, Ivan

- 1-64. (and BACK, William) Geochemistry of iron-rich ground water of southern Maryland: Jour. Geology, v. 72, no. 4, p. 435-447, 1964.

BARNES, Ivan--Continued

- 1-65. Geochemistry of Birch Creek, Inyo County, California, a travertine depositing creek in an arid climate: *Geochim. et Cosmochim. Acta*, v. 29, no. 2, p. 85-112, 1965.

BAROSH, P. J.

- 1-64. Lower Permian stratigraphy of east-central Nevada and adjacent Utah: U.S. Geol. Survey open-file report, 144 p., 1964.
- 1-65. Permian stratigraphy of east-central Nevada [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 317-318, 1965.

BARTON, P. B., Jr.

- 1-64. (and TOULMIN, Priestley, 3d) Experimental determination of the reaction chalcopyrite+sulfur=pyrite+bornite from 350 to 500°C: *Econ. Geology*, v. 59, no. 5, p. 747-752, 1964.

BASS, N. W.

- 1-64. Mineral fuels and associated resources: Oil and gas, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 45-67, 1964.

- 2-64. Relationship of crude oils to depositional environment of source rocks in the Uinta Basin, in *Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse*, 13th annual field conference, 1964: Salt Lake City, Utah, Intermountain Association of Petroleum Geologists, p. 201-206, 1964.

BASSETT, A. M.

- 1-64. (and KUPFER, D. H.) A geologic reconnaissance in the southeastern Mojave Desert, California: California Div. Mines and Geology Spec. Rept. 83, 1964.

BATEMAN, A. F., Jr.

- 1-64. Montana's coal resources: U.S. Geol. Survey open-file report, 20 p., 1964.

BATES, R. G.

- 1-64. Natural gamma aero-radioactivity of the National Reactor Testing Station area, Idaho: U.S. Geol. Survey Geophys. Inv. Map GP-446, 1964.
- 1-65. Aeroradioactivity survey and areal geology of the National Reactor Testing Station area, Idaho (ARMS-1): U.S. Atomic Energy Comm., Civil Effects Test Operations, CEX-59.4.10, 22 p., 1965.

BATH, G. D.

- 1-64. (SCHWARTZ, G. M., and GILBERT, F. P.) Aero-magnetic and geologic map of east-central Minnesota: U.S. Geol. Survey Geophys. Inv. Map GP-474, 1964.
- 2-64. (SCHWARTZ, G. M., and GILBERT, F. P.) Aero-magnetic and geologic map of northwestern Minnesota: U.S. Geol. Survey Geophys. Inv. Map GP-471, 1964.

BAYLEY, R. W.

- 1-65. Preliminary geologic map of Bradley Peak quadrangle, Carbon County, Wyoming: U.S. Geol. Survey open-file report, 1965.

BECK, M. E., Jr.

- 1-64. (and MATTICK, R. E.) Interpretation of an aero-magnetic survey in western Pennsylvania and parts of eastern Ohio, northern West Virginia, and western Maryland: *Pennsylvania Geol. Survey Inf. Circ.* 52, 1964.

BECRAFT, G. E.

- 1-65. (and PEARSON, R. C.) Upper Paleozoic rocks in Ferry County and southern Stevens County, northeastern Washington: Preliminary discussion: *Geol. Soc. America Spec. Paper* 82, p. 318, 1965.

BEETEM, W. A.

- 1-63. (and JANZER, V. J.) Distribution coefficient data and preliminary estimates of movement of radionuclides, Tatum salt dome, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-31, 37 p., 1963.

BEHRENDT, J. C.

- 1-65. Densification of snow on the ice sheet of Ellsworth Land and the southern Antarctic Peninsula: *Jour. Glaciology*, v. 5, no. 40, p. 451-460, 1965.

BELL, K. G.

- 1-64. Mineral fuels and associated resources: Rock asphalt and solid asphaltic bitumens, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 73-75, 1964.
- 2-64. Structure and stratigraphy of the Nantasket locality, Trip H, in *Guidebook to field trips in the Boston area and vicinity*, Chestnut Hill, Mass., New England Intercollegiate Geol. Cong., 56th Ann. Mtg., 1964: [New Haven, Conn., Yale Univ., Dept. Geology] 1964: p. 115-120, 1964.

BENNETT, G. D.

- 1-64. (and PATTEN, E. P., Jr.) Constant-head pumping test of a multiaquifer well to determine characteristics of individual aquifers: U.S. Geol. Survey Water-Supply Paper 1536-G, p. G181-G203, 1964. (Originally published 1962.)

BENSON, M. A.

- 1-64. Discussion of "Hydrology of spillway design--Large structures--Adequate data", by F. F. Snyder: *Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour.*, v. 90, no. HY5, p. 297-300, 1964.

BERDAN, J. M.

- 1-65. Stratigraphy and faunas of subsurface lower Paleozoic rocks, Florida and adjacent states [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 10, 1965.

BERG, H. C.

- 1-64. (EBERLEIN, G. D., and MacKEVETT, E. M., Jr.) Metallic and nonmetallic mineral resources: Metallic mineral resources, in *Mineral and water resources of Alaska*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 95-125, 1964.

BERGENBACK, R. E.

- 1-64. Petrology of pre-Selma strata from core holes in western Alabama: U.S. Geol. Survey Bull. 1160-B, p. 9-53, 1964.

BERGENDAHL, M. H.

- 1-64. Metallic mineral resources: Complex precious- and base-metal deposits, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 77-84, 1964.
- 2-64. Metallic mineral resources: Cripple Creek District, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 85, 1964.
- 3-64. Mineral resources: Gold, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 93-101, 1964.
- 4-64. Metallic mineral resources: Gold, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong., 88th, 2d sess., Senate Doc. No. 87, p. 87-100, 1964.
- 5-64. Metallic mineral resources: Gold, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 83-89, 1964.
- 6-64. Metallic mineral resources: Gold and silver, in *Mineral and water resources of South Dakota*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 41-49, 1964.
- 7-64. (and SMITH, A. E.) Geology of the Fordsville quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-295, 1964.
- 1-65. Geology of the Cloverport quadrangle, Kentucky-Indiana, and the Kentucky part of the Cannelton quadrangle: U.S. Geol. Survey Geol. Quad. Map GQ-273, 1965.

BERGENDAHL, M. H. --Continued

- 2-65. Gold, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 131-139, 1965.
 - 3-65. Preliminary report on the geology of the eastern part of the Red-Buffalo Route of Interstate Highway 70, Summit County, Colorado: U.S. Geol. Survey open-file report, 10 p., 1965.
- BERGQUIST, H. R.
- 1-65. (and OVERSTREET, E. F.) Bauxite deposits of the Margerum district: Alabama: U.S. Geol. Survey Bull. 1199-D, p. D1-D23, 1965.
- BERRYHILL, H. L., Jr.
- 1-64. (and SCHWEINFURTH, S. P.) Geology of the Ellsworth quadrangle, Pennsylvania: U.S. Geol. Survey Geol. Quad. Map GQ-333, 1964.
 - 1-65. Geology of the Ciales quadrangle, Puerto Rico: U.S. Geol. Survey Bull. 1184, 116 p., 1965.
- BERWICK, V. K.
- 1-64. (CHILDERS, J. M., and KUENTZEL, M. A.) Magnitude and frequency of floods in Alaska south of the Yukon River: U.S. Geol. Survey Circ. 493, 15 p., 1964[1965].
- BEVERAGE, J. P.
- 1-64. (and CULBERTSON, J. K.) Hyperconcentrations of suspended sediment: Am. Soc. Civil Engineers Proc. Paper 4136, Hydraulics Div. Jour., v. 90, no. HY6, p. 117-128, 1964.
- BINGHAM, R. H.
- 1-64. Well water for home supplies in Lewis County: Tennessee Dept. Conserv. Div. Water Resources Hydrologic Atlas, no. 3, 12 p., 1964.
 - 1-65. (and MANN, K. D.) Well water for home supplies in Dickson County, Tennessee: U.S. Geol. Survey open-file report, 18 p., 1965.
 - 2-65. (and MANN, K. D.) Well water for home supplies in Wayne County, Tennessee: U.S. Geol. Survey open-file report, 18 p., 1965.
- BISHOP, A. A.
- 1-65. (SIMONS, D. B., and RICHARDSON, E. V.) Total bed-material transport: Am. Soc. Civil Engineers Proc., Jour. Hydraulics Div., v. 91, no. HY2, pt. 1, p. 175-191, 1965.
- BLACK, D. F. B.
- 1-64. Geology of the Versailles quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-325, 1964.
 - 1-65. (and MacQUOWN, W. C., Jr.) Itinerary: 1. Lithostratigraphy of the Ordovician Lexington limestone and Clays Ferry Formation of the central Bluegrass area near Lexington, Kentucky; 2. Excursion to the crypto-explosive structure near Versailles, Kentucky, by D. F. B. Black, in Annual Spring Field Conference of the Geological Society of Kentucky: Kentucky Geol. Soc., 51 p., 1965.
- BLACK, R. A.
- 1-63. Magnetic properties of magnetite-bearing grout in drill holes WDH 11 and WDH 12, U18a site: U.S. Geol. Survey Tech. Letter, DANNY BOY-2, 8 p., 1963.
 - 2-63. (and CUNNINGHAM, D. R.) Seismic determination of the elastic constants of dacite in the pilot air-storage chamber, Area 401, Nevada Test Site: U.S. Geol. Survey Tech. Letter, PLUTO-23, 9 p., 1963.
- BLADE, L. V.
- 1-64. Geology of the Cuba quadrangle in Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-322, 1964.
- BLAKE, M. C., Jr.
- 1-64. (and BAILEY, E. H.) Translation of part of "Outline of the analysis of igneous rocks and the determination of their chemical types" by A. N. Zavaritsky, USSR, 1941: Internat. Geology Rev., v. 6, no. 1, p. 1-17, 1964.

BLAKE, Paul

- 1-65. Survey instruments and methods in the United States of America: Surveying and Mapping, v. 25, no. 2, p. 244-252, 1965.
 - 2-65. Surveying and photogrammetry in the search for Pakistan's mineral resources: Surveying and Mapping, v. 25, no. 1, p. 69-77, 1965.
- BLANK, H. R., Jr.
- 1-65. Southwest Oregon gravity data: U.S. Geol. Survey open-file report, map, 1965.
 - 2-65. Regional gravity survey in southwestern Oregon: Preliminary results [abs.]: Geol. Soc. America Spec. Paper 82, p. 241, 1965.
- BODHAINE, G. L.
- 1-64. Drainage basins and their effect on bridge and culvert design: Arizona Conf. Roads and Streets, 13th Ann., Univ. Arizona, April 1964, Proc., p. 42-51, 1964.
- BOETTCHER, A. J.
- 1-64. Geology and ground-water resources in eastern Cheyenne and Kiowa Counties, Colorado: U.S. Geol. Survey Water-Supply Paper 1770N, p. N1-N16, 1964.
- BOGART, D. B.
- 1-64. (ARNOW, Ted, and CROOKS, J. W.) Water resources of Puerto Rico—a progress report: Puerto Rico Water Resources Authority Water Resources Bull. 4, [120] p., 1964.
- BOGGESE, D. H.
- 1-64. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Bethany Beach area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-122, 1964[1965].
 - 2-64. (and ADAMS, J. K.) Water-table, surface-drainage and engineering soils map of the Greenwood quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-99, 1964.
 - 3-64. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Laurel area, Delaware, U.S. Geol. Survey Hydrol. Inv. Atlas HA-123, 1964[1965].
 - 4-64. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Seaford West area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-105, 1964.
 - 5-64. (DAVIS, C. F., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Milford quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-133, 1964[1965].
 - 6-64. (ADAMS, J. K., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Milton quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-102, 1964.
 - 7-64. (ADAMS, J. K., and DAVIS, C. F.) Water-table, surface-drainage, and engineering soils map of Georgetown quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-107, 1964.
 - 8-64. (ADAMS, J. K., and DAVIS, C. F.) Water-table, surface-drainage, and engineering soils map of the Rehoboth Beach area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-109, 1964.
 - 1-65. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Kenton area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-138, 1965.
 - 2-65. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Little Creek quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-134, 1965.
 - 3-65. (and ADAMS, J. K.) Water-table, surface-drainage, and engineering soils map of the Millsboro area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-121, 1965.

- BOGGESE, D. H. --Continued
 4-65. (DAVIS, C. F., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Wyoming quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-141, 1965.
- BONILLA, M. G.
 1-65. Geologic map of the San Francisco South quadrangle, California: U.S. Geol. Survey open-file report, 1965.
 2-65. (and McCULLOCH, D. S.) Engineering geologic effects of the March 27, 1964, earthquake on the Alaska Railroad [abs.]: Geol. Soc. America Spec. Paper 82, p. 15-16, 1965.
- BOOKS, K. G.
 1-65. Natural gamma aeroradioactivity map of parts of the San Francisco region, California: U.S. Geol. Survey Geophys. Inv. Map GP-483, 1965.
- BOSWELL, E. H.
 1-65. Significance of dissolved solids in ground-water supplies in Mississippi: Mississippi Water News, v. 11, no. 9, p. 1, 3, 1965.
 2-65. (MOORE, G. K., MacCARY, L. M., and others) Cretaceous aquifers in the Mississippi Embayment, with discussion of quality of the water by H. G. Jeffery: U.S. Geol. Survey Prof. Paper 448-C, p. C1-C37, 1965.
- BOUCOT, A. J.
 1-64. (GRISCOM, Andrew, and ALLINGHAM, J. W.) Geologic and aeromagnetic map of northern Maine: U.S. Geol. Survey Geophys. Inv. Map GP-312, 1964.
- BOWERS, W. E.
 1-63. Some effects of alteration of a vitric ash-fall tuff from the Twin Peaks area, Rainier Mesa quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-14, 10 p., 1963.
 2-63. (and EMERICK, W. L.) Preliminary report on the geology of the U12p tunnel, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-6, 13 p., 1963.
 3-63. (and EMERICK, W. L.) Supplementary geologic mapping in the 1500 re-entry tunnel, Area 15, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-15-4, 5 p., 1963.
 4-63. (and HOUSER, F. N.) Lithologic log of the U12k.01 vertical drill hole, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Area-12-11, 10 p., 1963.
 5-63. (MEYER, G. L., SNYDER, R. P., and HOOVER, D. L.) Abridged lithologic log of Pahute Mesa Exploratory Hole, 6,541 to 7,563 feet, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-2, supp. 2, 5 p., 1963.
- BOWIE, J. E.
 1-65. (and KAM, William) Use of water by riparian vegetation, Cottonwood Wash, Arizona--A summary: U.S. Geol. Survey open-file report, 4 p., 1965.
- BOWLES, C. G.
 1-64. Nonmetallic and industrial mineral resources: Salt and brine, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 137-140, 1964.
- BOWMAN, Isaiah
 1-65. Well-drilling methods: U.S. Geol. Survey Water-Supply Paper 257, 139 p., 1965. (Originally published 1911.)
- BOYNTON, G. R.
 1-64. (and GILBERT, F. P.) Aeromagnetic map of the Cupsuptic quadrangle, Oxford and Franklin Counties, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-477, 1964.
- BOYNTON, G. R. --Continued
 2-64. (and GILBERT, F. P.) Aeromagnetic map of the Oquossoc quadrangle, Oxford and Franklin Counties, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-478, 1964.
 3-64. (and GILBERT, F. P.) Aeromagnetic map of the Phillips quadrangle, Franklin County, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-479, 1964.
 4-64. (and GILBERT, F. P.) Aeromagnetic map of the Rangeley quadrangle and part of the Kennebec Lake quadrangle, Franklin and Oxford Counties, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-480, 1964.
 5-64. (and VARGO, J. L.) Aeromagnetic map of parts of Gogebic and Ontonagon Counties, Michigan: U.S. Geol. Survey open-file report, map, 1964.
 1-65. (and PITILLO, D. R.) Natural gamma aeroradioactivity map of the Belvidere quadrangle, Warren County, New Jersey, and Northampton County, Pennsylvania: U.S. Geol. Survey open-file report, map, 1965.
 2-65. (and ZANDLE, G. L.) Aeromagnetic map of the Belvidere quadrangle, Warren County, New Jersey, and Northampton County, Pennsylvania: U.S. Geol. Survey open-file report, map, 1965.
- BRABB, E. E.
 1-64. (and CHURKIN, Michael, Jr.) Preliminary geologic map of the Charley River quadrangle, east central Alaska: U.S. Geol. Survey open-file report, map and explanation, 1964.
 1-65. (and CHURKIN, Michael, Jr.) Preliminary geologic map of the Eagle D-1 quadrangle, east-central Alaska: U.S. Geol. Survey open-file report, map, 1965.
- BRADLEY, Edward
 1-64. Geohydrologic analogies between the Jordan Valleys of Utah and the Holy Land: Internat. Assoc. Sci. Hydrology Bull., v. 9, no. 3, p. 12-23, 1964.
 2-64. Geology and ground-water resources of southeastern New Hampshire: U.S. Geol. Survey Water-Supply Paper 1659, 80 p., 1964.
- BRADLEY, W. H.
 1-64. Geology of Green River Formation and associated Eocene rocks in southwestern Wyoming and adjacent parts of Colorado and Utah: U.S. Geol. Survey Prof. Paper 496-A, p. A1-A86, 1964[1965].
 2-64. Lazurite, talc and chlorite in the Green River Formation of Wyoming: Am. Mineralogist, v. 49, nos. 5-6, p. 778-781, 1964.
- BREDEHOEFT, J. D.
 1-64. Variation of permeability in the Tensleep Sandstone in the Bighorn Basin, Wyoming, as interpreted from core analyses and geophysical logs, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. 166-170, 1964.
 2-64. (and FARVOLDEN, R. N.) Disposition of aquifers in intermontane basins of northern Nevada: Internat. Assoc. Sci. Hydrology Pub. 64, p. 197-212, 1964.
 1-65. (and PAPADOPULOS, I. S.) Rates of vertical ground-water movement estimated from the Earth's thermal profile: Water Resources Research, v. 1, no. 2, p. 325-328, 1965.
- BREGER, I. A.
 1-65. The geochemistry of lipids [abs.]: Jour. Am. Oil Chemists Soc., v. 42, no. 3, p. 150A, 1965.
- BRIGGS, R. P.
 1-65. A decade of geologic mapping in Puerto Rico--progress to date and work remaining [abs.]: Caribbean Geol. Conf., 4th, Trinidad, 1965, Program (mimeographed), 1965.

- BRIGGS, R. P. --Continued
- 2-65. The geologic history of Puerto Rico [abs.]: Caribbean Geol. Conf., 4th, Trinidad, 1965, Program (mimeographed), 1965.
 - 3-65. (and AKERS, J. P.) Hydrogeologic map of Puerto Rico and adjacent islands: U.S. Geol. Survey Hydrol. Inv. Map HA-197, 1965.
- BROBST, D. A.
- 1-64. Mineral resources: Barite, *in* Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 46-48, 1964.
 - 2-64. Nonmetallic and industrial minerals: Barite, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 157-160, 1964.
 - 3-64. Nonmetallic and industrial minerals and materials resources: Barite, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 154-157, 1964.
 - 1-65. Barite in the United States exclusive of Alaska and Hawaii: U.S. Geol. Survey Mineral Inv. Resource Map MR-43, 1965.
- BROCK, M. R.
- 1-65. (and BARKER, Fred) Intrusive welded tuff in the Sawatch Range, Colorado [abs.]: Geol. Soc. America Spec. Paper 82, p. 320-321, 1965.
- BRODERICK, G. N.
- 1-64. Nonmetallic and industrial minerals: Sulfur, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 177-180, 1964.
 - 1-65. Sulfur, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 309-312, 1965.
- BROECKER, M. E.
- 1-64. (and WINSLOW, J. D.) Ground-water levels in observation wells in Kansas, 1963: Kansas Geol. Survey Bull. 173, 94 p., 1964.
- BROMERY, R. W.
- 1-64. (and GRISCOM, Andrew) A gravity survey in the Mayaguez area of southwest Puerto Rico, *in* A study of serpentinite, the AMSOC core hole near Mayaguez, Puerto Rico: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 61-74, 1964.
 - 2-64. (and NATOFF, N. W.) Aeromagnetic map of the Forks quadrangle, Piscataquis and Somerset Counties, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-485, 1964.
 - 3-64. (PETTY, A. J., and GILBERT, F. P.) Natural gamma aeroradioactivity map of Bel Air and vicinity, Harford, Baltimore, and Cecil Counties, Maryland: U.S. Geol. Survey Geophys. Inv. Map GP-484, 1964.
 - 4-64. (PETTY, A. J., and SMITH, C. W.) Aeromagnetic map of Bel Air and vicinity, Harford, Baltimore, and Cecil Counties, Maryland: U.S. Geol. Survey Geophys. Inv. Map GP-482, 1964.
 - 5-64. (and SNAVELY, P. D., Jr.) Geologic interpretation of reconnaissance gravity and aeromagnetic surveys in northwestern Oregon: U.S. Geol. Survey Bull. 1181-N, p. N1-N13, 1964.
 - 1-65. Aeromagnetic map of the Albany-Newport area, Oregon, and its geologic interpretation: U.S. Geol. Survey Geophys. Inv. Map GP-481, 1965.
 - 2-65. (and McCASLIN, W. E.) Aeromagnetic map of Findlay, Ohio, and vicinity: U.S. Geol. Survey Geophys. Inv. Map GP-500, 1965.
- BROOM, M. E.
- 1-64. (ALEXANDER, W. H., Jr., and MYERS, B. N.) Ground-water resources of Camp Franklin, Morris, and Titus Counties, Texas: U.S. Geol. Survey open-file report, 260 p., 1964.
- BROSSE, W. P.
- 1-65. (and REISER, H. N.) Preliminary geologic map of the Arctic quadrangle, Alaska: U.S. Geol. Survey open-file report, 1 sheet, 1965.
- BROWN, Andrew
- 1-64. (and BREGER, I. A.) Evaluation of weathering in the Chattanooga shale by Fischer assay, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501C, p. 92-95, 1964.
- BROWN, D. A.
- 1-64. Fossil Bryozoa from drill holes on Eniwetok Atoll, *in* Bikini and nearby atolls, Marshall Islands: U.S. Geol. Survey Prof. Paper 260-EE, p. 1113-1116, 1964.
- BROWN, D. L.
- 1-65. Prospecting in Colorado—Guide to Colorado mineral deposits: Colorado School Mines Mineral Industries Bull., v. 8, no. 1, 11 p., 1965.
- BROWN, R. D., Jr.
- 1-64. Geologic map of the Stonyford quadrangle, Glenn, Colusa, and Lake Counties, California: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-279, 1964.
- BROWN, S. G.
- 1-64. Sulphur Spring Valley, Willcox basin [and] Douglas basin, *in* Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 23-28, 1964.
 - 1-65. (DAVIDSON, E. S., KISTER, L. R., and THOMSEN, B. W.) Water resources of Fort Huachuca Military Reservation, southeastern Arizona: U.S. Geol. Survey open-file report, 146 p., 1965.
- BROWN, T. E.
- 1-64. Measurement of relative cationic diffusion and exchange rates of montmorillonite, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. 96-99, 1964.
- BROWN, W. M.
- 1-64. (and REEVES, R. G.) Ferrous metals: Iron, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 93-98, 1964.
- BRUCE, R. L.
- 1-64. (and WEEKS, R. A.) Nonmetallic and industrial mineral resources: Sand and gravel, *in* Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 93-96, 1964.
- BUCHANAN, T. J.
- 1-64. Time of travel of soluble contaminants in streams: Am. Soc. Civil Engineers Proc. Paper 3932, Sanitary Eng. Div. Jour., v. 90, no. SA3, p. 1-12, 1964.
 - 1-65. (MILLER, E. G., and LUDLOW, J. M.) Base-flow relations for partial-record stations in New Jersey: U.S. Geol. Survey open-file report, 16 p., 1965.
- BULL, W. B.
- 1-64. A consolidation test for undisturbed sands: Materials Research and Standards, v. 4, no. 7, p. 347-351, 1964.
 - 2-64. Alluvial fans and near-surface subsidence in western Fresno County, California: U.S. Geol. Survey Prof. Paper 437-A, p. A1-A71, 1964.

BULL, W. B.--Continued

- 3-64. History and causes of channel trenching in western Fresno County, California [abs.]: Jour. Geophys. Research, v. 66, no. 5, p. 1552, 1964.
 - 4-64. Particle-size analysis of sand containing friable fragments: Materials Research and Standards, v. 4, no. 8, p. 407-410, 1964.
- BUNKER, C. M.
- 1-62. (BRADLEY, W. A., and SHUTLER, M. D.) Interim report on gamma radioactivity and bulk density measurements in drill holes, Area 401, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, PLUTO-20, 18 p., 1962.
 - 1-63. Gamma-ray spectral analyses of rock samples from the vicinity of the Marshmallow nuclear test: U.S. Geol. Survey Tech. Letter, MARSHMALLOW-7, 7 p., 1963.
 - 2-63. Preliminary radioisotope measurements, U12E.05 reentry tunnel, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-13, 5 p., 1963.
 - 3-63. (and BRADLEY, W. A.) Measurements of subsurface natural gamma radioactivity, Project Dribble, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-37, 14 p., 1963.
 - 4-63. (and BUSH, C. A.) Gamma-ray spectral analyses of rock samples from the 3B and 3E crosscuts, U16a reentry drift, Marshmallow site, Nevada Test Site: U.S. Geol. Survey Tech. Letter, MARSHMALLOW-8, 4 p., 1963.
 - 5-63. (and BUSH, C. A.) Gamma-ray spectral analyses of rock samples from the 3A crosscut, U16a reentry tunnel, Marshmallow site, Nevada Test Site: U.S. Geol. Survey Tech. Letter, MARSHMALLOW-9, 5 p., 1963.
 - 6-63. (and BUSH, C. A.) Gamma-ray spectral analyses of rock samples from the number 3 drift, U16a reentry tunnel, Marshmallow site, Nevada Test Site: U.S. Geol. Survey Tech. Letter, MARSHMALLOW-10, 4 p., 1963.
 - 7-63. (and BUSH, C. A.) Gamma-ray spectral analyses of rock samples from the 2A crosscut, U16a reentry tunnel, Marshmallow site, Nevada Test Site: U.S. Geol. Survey Tech. Letter, MARSHMALLOW-11, 5 p., 1963.
 - 1-64. Radioactivity- and density-measuring devices for oceanographic studies, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D65-D69, 1964.
 - 1-65. Subsurface measurements of natural gamma radioactivity, Nevada Test Site and vicinity, Nye, Lincoln, and Clark Counties, Nevada: U.S. Geol. Survey open-file report, TEI-858, 74 p., 1965.
- BURBANK, W. S.
- 1-64. (and LUEDKE, R. G.) Geology of the Iron-ton quadrangle, Colorado: U.S. Geol. Survey Geol. Quad. Map GQ-291, 1964.
 - 1-65. (and LUEDKE, R. G.) Volcanism and cauldron development in the western San Juan Mountains, Colorado [abs.]: Geol. Soc. America Spec. Paper 82, p. 321-322, 1965.
- BURNHAM, W. W.
- 1-64. Reverse-circulation drilling: Am. Soc. Testing Materials Spec. Tech. Pub. 351, p. 133-136, 1964.
- BURTON, L. C.
- 1-65. Ground water in terrace deposits of central Beckham County, Oklahoma: Oklahoma Water Resources Board Bull. 25, 30 p., 1965.
- BUSBY, M. W.
- 1-64. Relation of annual runoff to meteorological factors, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C188-C189, 1964.

BUSH, A. L.

- 1-64. Construction materials: Lightweight aggregates, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 197-202, 1964.
- BUTLER, A. P., Jr.
- 1-64. Metallic mineral resources: Uranium, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 161-165, 1964.
 - 2-64. Miscellaneous metals: Uranium, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 136-144, 1964.
- BUTLER, R. G.
- 1-65. Report of water levels in selected observation wells in Utah—April 1965: U.S. Geol. Survey open-file report, 1 p., 1965.
- BYERLY, P. E.
- 1-65. Geothermal surveys in sedimentary rocks near Grants and Laguna, New Mexico: Geophysics, v. 30, no. 3, p. 395-402, 1965.
- BYERS, F. M., Jr.
- 1-64. (and CUMMINGS, David) Geologic map of Scrugham Peak quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report, 1 p., 1964.
 - 2-64. (and MONK, E. F.) Geology and physical properties of rocks in Scrugham Peak quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-75, 15 p., 1964.
 - 3-64. (ROGERS, C. L., and LUFT, S. J.) Geologic map of the Buckboard Mesa quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report, 1 p., 1964.
- CABLE, L. W.
- 1-64. (and WATKINS, F. A., Jr., and ROBISON, T. M.) Hydrogeology of the principal aquifers in Vigo and Clay Counties, Indiana: U.S. Geol. Survey open-file report, 57 p., 1964.
 - 1-65. (and ROBISON, T. M.) Ground-water resources of Montgomery County, Indiana: U.S. Geol. Survey open-file report, 39 p., 1965.
 - 2-65. (and ROBISON, T. M.) Hydrogeology of rocks of Mississippian, Pennsylvanian, and Quaternary ages, Sullivan and Greene Counties, Indiana: U.S. Geol. Survey open-file report, 47 p., 1965.
- CAGLE, J. W., Jr.
- 1-64. (and NEWTON, J. G.) Geology and ground-water resources of Escambia County, Alabama: Geol. Survey Alabama, Div. Water Resources Bull. 74, 205 p., 1964.
- CALLAHAN, J. A.
- 1-64. Ground-water resources in Yazoo County, Mississippi: Mississippi Board Water Comm. Bull. 63-5, 42 p., 1964.
- CALLAHAN, J. E.
- 1-65. (and WAYLAND, R. G.) Geologic reconnaissance of the West Creek dam site near Skagway, Alaska: U.S. Geol. Survey Bull. 1211-A, p. A1-A13, 1965.
- CAMERON, C. C.
- 1-65. Mapping continental glacial drifts: An environmental approach [abs.]: Geol. Soc. America Spec. Paper 82, p. 25, 1965.
- CAMP, J. D.
- 1-65. Flood of 1962 near Baton Rouge, Louisiana: U.S. Geol. Survey Hydrol. Inv. Atlas HA-126, map, 1965.
- CAMPBELL, A. B.
- 1-65. General geology of the Hunters quadrangle and vicinity, northeastern Washington [abs.]: Geol. Soc. America Spec. Paper 82, p. 322, 1965.

- CAMPBELL, R. H.
1-65. Geologic map and structure sections of the Ogotoruk Creek area and vicinity, Alaska: U.S. Geol. Survey open-file report, map, 1965.
- CAMPBELL, W. J.
1-64. The wind-driven circulation of ice and water in a polar ocean [abs.]: *Am. Geophys. Union Trans.*, v. 45, no. 4, p. 619, 1964.
- CANNEY, F. C.
1-64. (and POST, E. V.) Suggested exploration target in west-central Maine, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D61-D64, 1964.
1-65. Geochemical prospecting: *Mining Eng.*, v. 17, no. 2, p. 87-89, 1965.
- CANNON, H. L.
1-63. Contributed to the section on "Geobotanical prospecting", in *Aridity and Man*: *Am. Assoc. Adv. Sci. Pub.* 74, p. 191-193, 1963.
1-64. Geochemistry of rocks and related soils and vegetation in the Yellow Cat area, Grand County, Utah: U.S. Geol. Survey Bull. 1176, 127 p., 1964.
1-65. (and FIDLER, D. A.) Comparison of geochemical environment in areas of high rate of cancer and heart disease in New York and Maryland with that of an area of low disease rate in New Mexico: U.S. Geol. Survey open-file report, 11 p., 1965; Abstract in *Geol. Soc. America Spec. Paper* 82, p. 366, 1965.
- CARLSTON, C. W.
1-64. Drainage density and streamflow: U.S. Geol. Survey Prof. Paper 422-C, p. C1-C8, 1964. (Originally published 1963.)
2-64. Use of tritium in hydrologic research—Problems and limitations: *Internat. Assoc. Sci. Hydrology Bull.*, v. 9, no. 3, p. 38-42, 1964.
1-65. Reply to discussion by B. R. Payne, "Use of tritium in hydrologic research—Problems and limitations": *Internat. Assoc. Sci. Hydrology Bull.*, v. 10, no. 1, p. 67, 1965.
- CARPENTER, C. H.
1-64. (ROBINSON, G. B., Jr., and BJORKLUND, L. J.) Selected hydrologic data, Upper Sevier River basin, Utah: Utah State Engineer's Office Basin Data Rept. 8, 29 p., 1964.
- CARR, M. H.
1-63. Shock equation of state, in *Astrogeologic studies*, annual progress report, August 25, 1962—July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 39-52, 1963.
2-63. The geology of the Timocharis Quadrangle, in *Astrogeologic studies*, annual progress report, August 25, 1962—July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 9-22, 1963.
- CARROLL, Dorothy
1-64. Ion-exchange capacity of sediments from the Experimental Mohole, Guadalupe site: *Jour. Sed. Petrology*, v. 34, no. 3, p. 537-542, 1964.
- CARROLL, R. D.
1-63. Determination of porosity from density logs in alluvium, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-51, 4 p., 1963.
2-63. Interpretation of geophysical logs—Pahute Mesa Exploratory Hole, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-5, 9 p., 1963.
3-63. Interpretation of geophysical logs, test well F, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-54, 22 p., 1963.
4-63. Interpretation of geophysical logs—test well 8, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-56, 13 p., 1963.
- CARROLL, R. D. --Continued
5-63. Interpretation of preshot geophysical logs, Sedan site, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-47, 11 p., 1963.
6-63. (and DICKEY, D. D.) On the relationship between sonic log and core resonance velocities in tuff, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-42, 8 p., 1963.
7-63. (and SCOTT, J. H.) Density-porosity of tuff, northern Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-51, 5 p., 1963.
1-64. Electrical resistivity of earth sediments near Upshot-Knothole sites, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Proj. -11, 33 p., 1964.
- CARRON, M. K.
1-63. Determination of ferrous iron in tektites by a semi-micro-volumetric method, in *Astrogeologic studies*, annual progress report, August 25, 1962—July 1, 1963, Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 187-196, 1963.
1-65. (MROSE, M. E., and REISER, H. N.) New data on cymrite, a hydrated silicate of barium and aluminum [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 26, 1965.
- CARTER, M. D.
1-65. Gem materials, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 267-276, 1965.
- CARTER, R. F.
1-64. (and GANNON, W. B.) Surface-water resources of the Yellow River basin in Gwinnett County, Georgia: *Georgia Geol. Survey Inf. Circ.* 22, 34 p., 1964.
- CARTER, W. D.
1-64. Construction materials: Sand and gravel, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 205-211, 1964.
2-64. (and SAVAGE, C. N.) Mineral resources: Silica, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 174-181, 1964.
1-65. Sand and gravel, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 353-361, 1965.
- CASE, J. E.
1-65. Gravitational evidence for a batholithic mass of low density along a segment of the Colorado mineral belt [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 26, 1965.
2-65. Principal facts for gravity stations in the Wind River Basin area, Wyoming: U.S. Geol. Survey open-file report, 2 p., 1965.
3-65. The U.S. Geological Survey's gravity program in Arizona, Colorado, New Mexico, and Utah: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 227-231, 1965.
4-65. (and GAIR, J. E.) Aeromagnetic map of parts of Marquette, Dickinson, Baraga, Alger, and Schoolcraft Counties, Michigan, and its geologic interpretation: U.S. Geol. Survey Geophys. Inv. Map GP-467, with 10-page text, 1965.
- CASHION, W. B.
1-64. Mineral fuels and associated resources: Oil shale, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 61-63, 1964.
2-64. Mineral fuels and associated resources: Other bituminous substances, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 63-68, 1964.

CASHION, W. B.--Continued

- 3-64. The distribution and quality of oil shale in the Green River Formation of the Uinta Basin, Utah-Colorado, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D86-D89, 1964; also in Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse, 13th annual field conference, 1964: Salt Lake City, Utah, Intermountain Association of Petroleum Geologists, p. 209-212, 1964.

CASTLE, R. O.

- 1-64. Geology of the Andover Granite and surrounding rocks, Massachusetts: U.S. Geol. Survey open-file report, 550 p., 1964.

CATHCART, J. B.

- 1-64. Economic geology of the Lakeland quadrangle, Florida: U.S. Geol. Survey Bull. 1162-G, p. G1-G128, 1964.

CEDERSTROM, D. J.

- 1-64. (and ASSAD, José Carlos) Observações hidrologicas no nordeste do Brasil: Brasil Divisão de Geologic E Mineralogia Notes Preliminares E Studies 20, 42 p., 1964.
2-64. (TRAINER, F. W., and WALLER, R. M.) Geology and ground-water resources of the Anchorage area, Alaska: U.S. Geol. Survey Water-Supply Paper 1773, 108 p., 1964.
1-65. (and HODGES, A. L.) A preliminary reconnaissance of the ground-water resources in the Connecticut River basin of New Hampshire and Vermont: U.S. Geol. Survey open-file report, 50 p., 1965.

CHAO, E. C. T.

- 1-64. (DWORNIK, E. J., and LITTLER, Janet) New data on the nickel-iron spherules from Southeast Asia tektites and their implications, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963, Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 53-83, 1963; also in Geochim. et Cosmochim. Acta, v. 28, no. 6, p. 971-980, 1964.
2-64. Spalled, aerodynamically modified moldavite from Slavic, Moravia, Czechoslovakia: Science, v. 146, no. 3645, p. 790-791, 1964.

CHASE, G. H.

- 1-64. Geologic considerations in selecting a site for a proposed deep exploratory well at the National Reactor Testing Station, Idaho [abs.]: Am. Geophys. Union Trans., v. 45, no. 4, p. 613, 1964.

CHINN, S. S. W.

- 1-64. Water-supply potential from an asphalt-lined catchment at Holualoa, Kona, Hawaii: U.S. Geol. Survey open-file report, 53 p., 1964.

CHRIST, C. L.

- 1-65. Phase transformations and crystal chemistry of schoepite: Am. Mineralogist, v. 50, nos. 1-2, p. 235-239, 1965.

CHRISTENSEN, R. C.

- 1-65. (and GILSTRAP, R. C.) Floodflow characteristics of Arkansas River at Interstate Highway 540, at Van Buren, Arkansas: U.S. Geol. Survey open-file report, 25 p., 1965.

CHRISTIANSEN, R. L.

- 1-65. (and NOBLE, D. C.) Black Mountain volcanism of southern Nevada [abs.]: Geol. Soc. America Spec. Paper 82, p. 246, 1965.
2-65. (LIPMAN, P. W., ORKILD, P. P., and BYERS, F. M., Jr.) Structure of the Timber Mountain caldera, southern Nevada, and its relation to Basin-Range structure, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B43-B48, 1965.

CHURKIN, Michael, Jr.

- 1-65. (and BRABB, Earl) Ordovician, Silurian, and Devonian biostratigraphy of east-central Alaska: Am. Assoc. Petroleum Geologists Bull., v. 49, no. 2, p. 172-185, 1965.

CHUTE, N. E.

- 1-64. Geology of the Norfolk Basin Carboniferous sedimentary rocks and the various igneous rocks of the Norwood and Blue Hills quadrangles, Trip G, in Guidebook to fieldtrips in the Boston area and vicinity, New England Intercollegiate Geol. Conf., 56th Ann. Mtg., Chestnut Hill, Mass., 1964: [New Haven, Conn., Yale Univ., Dept. Geology], p. 91-114, 1964.

CLARK, J. R.

- 1-64. Comment about the formula of veatchite—Discussion of paper by H. Jager and H. A. Lehmann (1963): Zeitschr. anorg. u. allgem. Chemie, v. 331, nos. 5-6, p. 348-349, 1964.
2-64. Comment on three proposed hydrate structures: Acta Cryst., v. 17, pt. 4, p. 459-461, 1964.
3-64. (and APPLEMAN, D. E.) Pentaborate polyanion in the crystal structure of ulexite, $\text{NaCaB}_5\text{O}_6(\text{OH})6.5\text{H}_2\text{O}$: Science, v. 145, no. 3638, p. 1295-1296, 1964.
1-65. Crystallographic data for the iron borate mineral, hulsite: Am. Mineralogist, v. 50, nos. 1-2, p. 249-254, 1965.
2-65. (and APPLEMAN, D. E.) Crystal structure of the sodium calcium borate, ulexite [abs.]: Am. Mineralogist, v. 50, nos. 1-2, p. 283, 1965.

CLARK, L. D.

- 1-64. Stratigraphy and structure of part of the western Sierra Nevada metamorphic belt, California: U.S. Geol. Survey Prof. Paper 410, 70 p., 1964[1965].
1-65. Bauxite deposits of the Springvale district, Georgia: U.S. Geol. Survey Bull. 1199-F, p. F1-F24, 1965.
2-65. (and CRITTENDEN, M. D., Jr.) Geology of the Mattingly quadrangle, Kentucky-Indiana: U.S. Geol. Survey Geol. Quad. Map GQ-361, 1965.

CLARK, W. E.

- 1-64. Possibility of salt-water leakage from proposed intra-coastal waterway near Venice, Florida well field: Florida Geol. Survey Rept. Inv. 38, 33 p., 1964.
2-64. (MUSGROVE, R. H., MENKE, C. G., and CAGLE, J. W., Jr.) Water-resources data for Alachua, Bradford, Clay, and Union Counties, Florida: Florida Geol. Survey Inf. Circ. 43, 154 p., 1964.

CLARKE, F. E.

- 1-64. Selection of metal components for long-term development of Egypt's corrosive ground waters: U.S. Geol. Survey open-file report, 39 p., 1964.
2-64. (and BARNES, Ivan) Preliminary evaluation of corrosion and encrustation mechanisms in tube wells of the Indus Plains, West Pakistan: U.S. Geol. Survey open-file report, 81 p., 1964.

CLARKE, J. W.

- 1-64. (CONANT, G. D., COOPER, Margaret, GRIMES, W. L., HACKMAN, B. S., JUSSEN, V. M., KING, R. R., LOUD, E. S., MEAD, M. C., OFTEDAHL, F. V., RANKIN, D. W., REES, V. E., and WHITE, W. S.) Bibliography of North American geology, 1960: U.S. Geol. Survey Bull. 1196, 777 p., 1964.
2-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, July 1964: U.S. Geol. Survey Geophys. Abs., no. 210, p. 519-599, 1964.
3-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, August 1964: U.S. Geol. Survey Geophys. Abs., no. 211, p. 601-685, 1964.

CLARKE, J. W. --Continued

- 4-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, September 1964: U.S. Geol. Survey Geophys. Abs. no. 212, p. 687-767, 1964.
- 5-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, October 1964: U.S. Geol. Survey Geophys. Abs. no. 213, p. 769-849, 1964.
- 6-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, November 1964: U.S. Geol. Survey Geophys. Abs. no. 214, p. 855-939, 1964.
- 7-64. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, December 1964: U.S. Geol. Survey Geophys. Abs. no. 215, p. 941-1027, 1964.
- 1-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, January 1965: U.S. Geol. Survey Geophys. Abs. no. 216, p. 1-81, 1965.
- 2-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, February 1965: U.S. Geol. Survey Geophys. Abs. no. 217, p. 83-167, 1965.
- 3-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, March 1965: U.S. Geol. Survey Geophys. Abs. no. 218, p. 169-255, 1965.
- 4-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, April 1965: U.S. Geol. Survey Geophys. Abs. no. 219, p. 257-343, 1965.
- 5-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, May 1965: U.S. Geol. Survey Geophys. Abs. no. 220, p. 345-431, 1965.
- 6-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Geophysical Abstracts, June 1965: U.S. Geol. Survey Geophys. Abs. no. 221, p. 433-521, 1965.
- 7-65. (VITALIANO, D. B., NEUSCHEL, V. S., and others) Index to Geophysical Abstracts Numbers 204-215: U.S. Geol. Survey Geophys. Abs., p. 1029-1293, 1965.

CLAYTON, R. N.

- 1-65. (FRIEDMAN, I., GRAF, D. L., MAYEDA, T., MEENTS, W. F., and SHIMP, N. F.) Origin of brines in the Illinois and Michigan Basins [abs.]: Geol. Soc. America Spec. Paper 82, p. 29, 1965.

CLEBSCH, Alfred, Jr.

- 1-63. (and PECKHAM, A. E.) Research by the United States Geological Survey related to the movement of radio-nuclides in the earth environment [with French abstract], La retention et la migration des ions radioactifs dans les sols—Colloque international, Saclay, France, 1962: Paris, Presses Universitaires de France, p. 189-197, 1963.

CLEMENT, W. G.

- 1-65. Complete Bouguer gravity map of the northern part of the San Francisco Bay area and its geologic interpretation: U.S. Geol. Survey Geophys. Inv. Map GP-468, 1965.

CLINE, D. R.

- 1-65. Geology and ground-water resources of Dane County, Wisconsin: U.S. Geol. Survey Water-Supply Paper 1779-U, p. U1-U64, 1965.

COBB, E. H.

- 1-64. Industrial minerals and construction materials occurrences in Alaska: U.S. Geol. Survey Mineral Inv. Resources Map MR-41, 1964.
- 2-64. Iron occurrences in Alaska: U.S. Geol. Survey Mineral Inv. Resources Map MR-40, 1964.
- 3-64. Lode gold and silver occurrences in Alaska: U.S. Geol. Survey Mineral Inv. Resources Map MR-32, 1964. (Originally published 1962.)

COBBAN, W. A.

- 1-64. (and SCOTT, G. R.) Multinodose scaphitid cephalopods from the lower part of the Pierre Shale and equivalent rocks in the conterminous United States: U.S. Geol. Survey Prof. Paper 483-E, p. E1-E13, 1964.

COBBAN, W. A. --Continued

- 1-65. (and GILL, J. R.) Stratigraphy of the Pierre Shale, Valley City and Pembina Mountain areas, North Dakota: U.S. Geol. Survey Prof. Paper 392-A, 20 p., 1965.

COHEE, G. V.

- 1-65. (WEST, W. S., CAMPBELL, R. H., and NELSON, C. A.) Changes in stratigraphic nomenclature by the U.S. Geological Survey, 1963: U.S. Geol. Survey Bull. 1194-A, p. A1-A35, 1965.

COHEN, Philip

- 1-64. A brief appraisal of the ground-water resources of the Grass Valley area, Humboldt and Pershing Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Ground-Water Resources—Reconn. Ser. Rept. 29, 40 p., 1964.
- 2-64. Preliminary results of hydrogeologic investigations in the valley of the Humboldt River near Winnemucca, Nevada: U.S. Geol. Survey Water-Supply Paper 1754, 590 p., 1964.
- 3-64. Water in the Humboldt River valley near Winnemucca, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 27, 68 p., 1964.
- 4-64. (and LOELTZ, O. J.) Evaluation of hydrogeology and hydrogeochemistry of Truckee Meadows area, Washoe County, Nevada: U.S. Geol. Survey Water-Supply Paper 1779-S, p. S1-S63; also in Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 28, 1964.
- 5-64. (and EVERETT, D. E.) Ground-water appraisal of Grass Valley, Pershing County, Nevada: Nevada Dept. Conserv. and Nat. Resources, Ground-Water Resources—Reconn. Ser. Rept. 29, 40 p., 1964.

COLBY, B. R.

- 1-64. Discharge of sands and mean-velocity relationships in sand-bed streams: U.S. Geol. Survey Prof. Paper 462-A, p. A1-A47, 1964.
- 2-64. Discussion of "Accuracy of current meter measurements", by R. W. Carter and I. E. Anderson: Am. Soc. Civil Engineers Proc. Hydraulics Div. Jour., v. 90, no. HY1, pt. 1, p. 349-352, 1964.
- 3-64. Scour and fill in sand-bed streams: U.S. Geol. Survey Prof. Paper 462-D, p. D1-D32, 1964.
- 4-64. Practical computations of bed-material discharge: Am. Soc. Civil Engineers Proc., v. 90, Paper 3843, Jour. Hydraulics Div., no. HY-2, pt. 1, p. 217-246, 1964.
- 1-65. (and SCOTT, C. H.) Effects of water temperature on the discharge of bed material: U.S. Geol. Survey Prof. Paper 462-G, p. G1-G25, 1965.

COLEMAN, R. G.

- 1-65. Compositional variation of garnets and pyroxenes in eclogites [abs.]: Geol. Soc. America Spec. Paper 82, p. 29-30, 1965.
- 2-65. (and COHAN, W. T.) Jadeite and asbestos in southern San Benito and western Fresno Counties, California—Pt. 2, in Geol. Soc. America Cordilleran Section, Guidebook 61st Ann. Meeting, Apr. 1965, Mercury, jadeite, and asbestos regions near Panoche Pass., Calif.: Fresno, California, p. 1-10, 1965.
- 3-65. (LEE, D. E., BEATTY, L. B., and BRANNOCK, W. W.) Eclogites and eclogites—their differences and similarities: Geol. Soc. America Bull., v. 76, no. 5, p. 483-508, 1965.

COLLIER, C. R.

- 1-64. (KRIEGER, R. A., and WHETSTONE, G. W.) Hydro-chemistry and sedimentation in Mammoth Cave [abs.]: Natl. Speleol. Soc. Bull., v. 26, no. 2, p. 83-84, 1964.
- 2-64. (and others) Influences of strip mining on the hydrologic environment in parts of Beaver Creek basin, Kentucky, 1955-59: U.S. Geol. Survey Prof. Paper 427-B, p. B1-B85, 1964.

- COLTON, R. B.
 1-63. Fault pattern in alluvium in Groom Mine SW quadrangle, Nevada: U.S. Geol. Survey Tech. Letter NTS-41, 5 p., 1963.
 2-63. Fault traces in alluvium and in rhyolite hills in Areas 3, 7, 9, 10, and 15, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-41, 5 p., 1963.
 3-63. Preliminary map of the close-in distribution of base surge and throwout material, Sedan event, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-39, 3 p., 1963.
 1-65. Pleistocene stratigraphy of Tudor Hill, South Windsor, Connecticut [abs.]: Geol. Soc. America Spec. Paper 82, p. 30, 1965.
- CONANT, L. C.
 1-64. General remarks on the pre-Selma Cretaceous strata of western Alabama: U.S. Geol. Survey Bull. 1160-F, p. F97-F101, 1964.
 1-65. Bauxite and kaolin deposits of Mississippi, exclusive of the Tippah-Benton district: U.S. Geol. Survey Bull. 1199-B, p. B1-B70, 1965.
- COOK, K. L.
 1-64. (and BERG, J. W., Jr.) Principal facts for gravity stations along the central and southern Wasatch Front, Utah: U.S. Geol. Survey open-file report, 3 p., 1964.
- COOLEY, M. E.
 1-64. (AKERS, J. P., and STEVENS, P. R.) Geohydrologic data on the Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah—Pt. 3, Selected lithologic logs, drillers' logs, and stratigraphic section: Arizona State Land Dept. Water Resources Rept. 12-C, 157 p., 1964.
 2-64. (HARSHBARGER, J. W., AKERS, J. P., and HARDT, W. F.) Regional hydrogeology of the Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah, with a section on Vegetation by O. N. Hicks: U.S. Geol. Survey open-file report, 245 p., 1964.
 3-64. (and HEVLY, R. H.) Geology and depositional environment of Laguna Salada, Arizona, in Prehistory of eastern Arizona, II, Chap. VIII: Fieldiana, Anthropology, v. 55, p. 188-200, 1964.
- COOPER, G. A.
 1-64. Brachiopods from Eniwetok and Bikini drill holes: U.S. Geol. Survey Prof. Paper 260-FF, p. FF1117-FF1120, 1964.
 2-64. (and GRANT, R. E.) New Premian stratigraphic units in Glass Mountains, west Texas: Am. Assoc. Petroleum Geologists Bull., v. 48, no. 9, p. 1581-1588, 1964.
- COOPER, J. B.
 1-65. New Mexico State Planning Office Report—Western closed basins: U.S. Geol. Survey open-file report, 25 p., 1965.
 2-65. (and JOHN, E. C.) Ground-water occurrence and geology of southeastern McKinley County, New Mexico: U.S. Geol. Survey open-file report, 263 p., 1965.
 3-65. (and TRAUGER, F. D.) New Mexico State Planning Office Report—San Juan River basin: U.S. Geol. Survey open-file report, 38 p., 1965.
- COOPER, J. R.
 1-64. (and SILVER, L. T.) Geology and ore deposits of the Dragoon quadrangle, Cochise County, Arizona: U.S. Geol. Survey Prof. Paper 416, 196 p., 1964.
- CORDOVA, R. M.
 1-65. (and SUBITZKY, Seymour) Groundwater in northern Utah Valley—A progress report for the period 1948-63: U.S. Geol. Survey open-file report, 61 p., 1965.
- CORLISS, J. B.
 1-64. (and MEADE, R. H.) Clay minerals from an area of land subsidence in the Houston-Galveston Bay area, Texas, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C79-C81, 1964.
- CORNWALL, H. R.
 1-64. Introduction, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 1, 1964.
 2-64. The mineral industry in Nevada, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 3-6, 1964.
 3-64. (and KLEINHAMPL, F. J.) Geology of Bullfrog quadrangle and ore deposits related to Bullfrog Hills caldera, Nye County, Nevada, and Inyo County, California: U.S. Geol. Survey Prof. Paper 454-J, p. J1-J25, 1964.
- COTTER, R. D.
 1-64. (and ROGERS, J. E.) Glacial geology of the Mountain Iron-Virginia-Eveleth area, Mesabi Iron Range, Minnesota, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C144-C146, 1964.
 2-64. (YOUNG, H. L., and WINTER, T. C.) Preliminary surficial geologic map of the Mesabi-Vermilion Iron Range area, northeastern Minnesota: U.S. Geol. Survey Misc. Geol. Inv. Map I-403, 1964.
- COULTER, H. W.
 1-65. (and MIGLIACCIO, R. R.) Engineering geologic effects of the March 27, 1964 earthquake at Valdez, Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 33, 1965.
 2-65. (HOPKINS, D. M., KARLSTROM, T. N. V., PÉWÉ, T. L., WAHRHAFTIG, Clyde, and WILLIAMS, J. R.) Map showing extent of glaciations in Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-415, 1965.
- COX, Allan
 1-64. Angular dispersion due to random vectors: Royal Astron. Soc. Geophys. Jour., v. 8, no. 4, p. 343-355, 1964.
 2-64. (and DOELL, R. R.) Long period variations of the geomagnetic field: Seismol. Soc. America Bull., v. 54, no. 6, pt. B, p. 2243-2270, 1964.
 3-64. (DOELL, R. R., and THOMPSON, George) Magnetic properties of the AMSOC serpentine core from Mayaguez, Puerto Rico, in A study of serpentinite, the AMSOC core hole near Mayaguez, Puerto Rico: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 49-60, 1964.
 1-65. Review of "Paleomagnetism and its application to geological and geophysical problems," by E. Irving: Science, v. 147, no. 3657, p. 494, 1965.
- COX, E. J.
 1-64. (and BOWLES, C. G.) Nonmetallic and industrial mineral resources: Gypsum and anhydrite, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 133-137, 1964.
- COX, E. R.
 1-65. Geology and hydrology between Lake McMillan and Carlsbad Springs, Eddy County, New Mexico: U.S. Geol. Survey open-file report, 139 p., 1965.
- CRANDELL, D. R.
 1-64. (and MILLER, R. D.) Post-hypsithermal glacier advances at Mount Rainier, Washington, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D110-D114, 1964.
 1-65. Alpine glaciers at Mount Rainier, Washington, during late Pleistocene and Recent time [abs.]: Geol. Soc. America Spec. Paper 82, p. 34-35, 1965.
 2-65. (MULLINEAUX, D. R., and WALDRON, H. H.) Age and origin of the Puget Sound trough in western Washing-

- CRANDELL, D. R. --Continued
 ton, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B132-B136, 1965.
- CREASEY, S. C.
 1-64. Review of "The tectonics of the Canadian Shield" by J. S. Stevenson: Science, v. 143, no. 3609, p. 945, 1964.
 1-65. Geology of the San Manuel area, Pinal County, Arizona, with a section on Ore deposits, by J. D. Pelletier and S. C. Creasey: U.S. Geol. Survey Prof. Paper 471, 64 p., 1965.
- CRESSLER, C. W.
 1-64. Geology and ground-water resources of the Paleozoic Rock area, Chattooga County, Georgia: Georgia Div. Conserv. Inv. Circ. 27, 14 p., 1964.
- CRESSMAN, E. R.
 1-64. Geology of the Tyrone quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-303, 1964.
 2-64. (and SWANSON, R. W.) Stratigraphy and petrology of the Permian rocks of southwestern Montana: U.S. Geol. Survey Prof. Paper 313-C, p. C275-C569, 1964.
- CRINER, J. H.
 1-64. (SUN, P.-C. P., and NYMAN, D. J.) Hydrology of aquifer systems in the Memphis area, Tennessee: U.S. Geol. Survey Water-Supply Paper 1779-O, p. O1-O54, 1964.
 1-65. Seismic surveying with firecrackers [abs.]: Geol. Soc. America SE section Ann. Mtg., Nashville, Tennessee, April 1965, Program, p. 17, 1965.
- CRIPPEN, J. R.
 1-65. Discussion of "Areal variations of mean annual runoff", by A. G. Hely: Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour., v. 91, no. HY2, p. 395-396, 1965.
 2-65. Natural water loss and recoverable water in mountain basins of southern California: U.S. Geol. Survey Prof. Paper 417-E, p. E1-E24, 1965.
- CRITTENDEN, M. D., Jr.
 1-64. Ferrous metals: Manganese, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 98-102, 1964.
 2-64. General geology of Salt Lake County, in Geology of Salt Lake County: Utah Geol. and Mineralog. Survey Bull. 69, p. 11-48, 1964.
 3-64. Geology: Structure, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 25-28, 1964.
 4-64. Metallic mineral resources: Manganese, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 113-119, 1964.
 5-64. Metallic mineral resources: Manganese, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 103-108, 1964.
 1-65. Geology of the Draper quadrangle, Utah: U.S. Geol. Survey Geol. Quad. Map GQ-377, 1965.
 2-65. Geology of the Dromedary Peak quadrangle, Utah: U.S. Geol. Survey Geol. Quad. Map GQ-378, 1965.
 3-65. Geology of the Mount Aire quadrangle, Salt Lake County, Utah: U.S. Geol. Survey Geol. Quad. Map GQ-379, 1965.
 4-65. Geology of the Sugar House quadrangle, Salt Lake County, Utah: U.S. Geol. Survey Geol. Quad. Map GQ-380, 1965.
 5-65. (CALKINS, F. C., and SHARP, B. J.) Geologic map of the Park City West quadrangle, Utah: U.S. Geol. Survey open-file report, 1 sheet, 1965.
 6-65. (and HOSE, R. K.) Geology of the Rome quadrangle in Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-362, 1965.
- CROFT, M. G.
 1-64. Geology and ground-water resources of Dade County, Georgia: Georgia Acad. Sci. Bull. V. XVIII, nos. 1 and 2, p. 13-14, 1964.
- CROSS, W. P.
 1-64. Floods of March 1963 in Ohio, and the flash flood of June 1963 in the vicinity of Cambridge: Ohio Div. Water Bull. 38, 82 p., 1964.
 2-64. Floods of March 1964 in Ohio: Ohio Div. Water Bull. 39, 58 p., 1964.
- CRUFF, R. W.
 1-64. (and RANTZ, S. E.) A comparison of flood-frequency studies for coastal basins in California: U.S. Geol. Survey open-file report, 116 p., 1964.
- CULBERTSON, J. K.
 1-64. (and DAWDY, D. R.) A study of fluvial characteristics and hydraulic variables, middle Rio Grande, New Mexico: U.S. Geol. Survey Water-Supply Paper 1498-F, p. F1-F74, 1964.
- CULLINS, H. L.
 1-65. (and BOWERS, W. E.) East Cortez coal area, Montezuma County, Colorado: U.S. Geol. Survey open-file report, 21 p., 1965.
- CUMMINGS, David
 1-63. Determination of flow direction in a rhyolite: U.S. Geol. Survey Tech. Letter NTS-41, 6 p., 1963.
 2-63. (SNYDER, R. P., and HOOVER, D. L.) Results of exploratory drilling and geologic studies at the proposed Ferris Wheel site, area 4, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-62, 47 p., 1963.
 1-65. Kink-bands: shock deformation of biotite resulting from a nuclear explosion: Science, v. 148, no. 3672, p. 950-952, 1965.
- CUMMINGS, T. R.
 1-65. Chemical character of surface waters of Oklahoma, 1961-1962: U.S. Geol. Survey open-file report, 203 p., 1965.
- CUPPELS, N. P.
 1-64. The Marlboro Formation in the Concord quadrangle, Trip F, in Guidebook to field trips in the Boston area and vicinity, New England Intercollegiate Geol. Cong., 56th Ann. Mtg., Chestnut Hill, Mass., [New Haven, Conn., Yale Univ., Dept. Geology] 1964: p. 81-89, 1964.
- CURTIS, L. W.
 1-64. (DOYLE, T. J., and WHETSTONE, G. W.) Discussion of "Annual variation of stream water temperature", by J. C. Ward: Am. Soc. Civil Engineers Proc., Sanitary Eng. Div. Jour., v. 90, no. SA4, p. 92-100, 1964.
- CUSHMAN, R. L.
 1-65. Evaluation of the hydraulic characteristics of the Major Johnson Springs aquifer, Eddy County, New Mexico: U.S. Geol. Survey open-file report, 38 p., 1965.
- CUSHMAN, R. V.
 1-64. (BAKER, J. A., and MEIKLE, R. L.) Records and logs of selected wells and test borings and chemical analyses of water in north-central Connecticut: Connecticut State Water Resources Comm., Connecticut Water Resources Bull. no. 4, 27 p., 1964.
 2-64. (KRIEGER, R. A., and McCABE, J. A.) Present and future water supply for Mammoth Cave National Park, Kentucky: U.S. Geol. Survey open-file report, 72 p., 1964.
- CUTHBERT, Margaret
 1-64. (and WARD, F. N.) Determination of iodine in vegetation, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C154-C156, 1964.

CUTTITTA, Frank

- 1-63. (CHAO, E. C. T., CARRON, M. K., LITTLER, J., FLETCHER, J. D., and ANNEL, C.) Some physical properties and the chemical composition of Australasian tektites, in *Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963*, Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 1-50, 1963.

- 1-64. (CHAO, E. C. T., CARRON, M. K., and LITTLER, J.) Some physical properties and the major chemical composition of selected australasian tektites [abs.]: *Am. Geophys. Union Trans.*, v. 45, no. 1, p. 81, 1964.

DALE, R. H.

- 1-64. (FRENCH, J. J., and WILSON, H. D., Jr.) The story of ground water in the San Joaquin Valley, California: U.S. Geol. Survey Circ. 459, 11 p., 1964.

DALRYMPLE, G. B.

- 1-64. Potassium-argon dates of three Pleistocene interglacial basalt flows from the Sierra Nevada, California: *Geol. Soc. America Bull.*, v. 75, no. 8, p. 753-757, 1964.

- 1-65. (and HIROOKA, Kimio) Variation of potassium, argon, and calculated age in a late Tertiary basalt flow [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 178, 1965.

- 2-65. (COX, Allan, and DOELL, R. R.) Potassium-argon age and paleomagnetism of the Bishop Tuff, California: *Geol. Soc. America Bull.*, v. 76, no. 6, p. 665-673, 1965.

DANE, C. H.

- 1-65. (and BACHMAN, G. O.) Topography and geology, in *Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print*, p. 19-39, 1965.

DASCH, M. D.

- 1-64. Metallic mineral resources: Antimony and other minor metals, in *Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 81-87, 1964.

- 2-64. Metallic mineral resources: Antimony, and other minor metals, in *Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 135-144, 1964.

- 3-64. Mineral resources: Garnet, in *Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 84-90, 1964.

- 4-64. Miscellaneous metals: Smelter byproducts—antimony, arsenic, bismuth, cadmium, selenium, and tellurium, in *Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 144-153, 1964.

- 5-64. Nonmetallic and industrial minerals and materials resources: Fluorine, in *Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 162-168, 1964.

- 6-64. Nonmetallic and industrial minerals and materials resources: Gem materials, in *Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print*, p. 169-177, 1964.

- 1-65. Antimony, arsenic, bismuth, and cadmium, in *Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print*, p. 365-372, 1965.

DAVIDSON, D. F.

- 1-64. Metallic mineral resources: Selenium and tellurium, in *Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87*, p. 134-137, 1964.

DAVIDSON, D. F.--Continued

- 1-65. (and GRANGER, H. C.) Selenium and tellurium, in *Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print*, p. 228-230, 1965.

DAVIDSON, E. S.

- 1-64. Duncan basin [and] Safford basin, in *Annual report ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19*, p. 17-18, 1964.

DAVIES, W. E.

- 1-64. Attapulgit from Carlsbad Caverns, New Mexico: U.S. Geol. Survey Prof. Paper 501-C, p. C82-C83, 1964; *abstract in Natl. Speleol. Soc. Bull.*, v. 26, no. 2, p. 74, 1964.

- 2-64. Glacial geology of Northern Greenland: *Polarforschung*, v. V, no. 1/2, p. 94-103, 1961[1964].

- 3-64. The future of the Potomac—a conflict in values: *Atlantic Naturalist*, v. 19, no. 4, p. 209-220, 1964.

- 1-65. Memorial to Ralph W. Stone, 1876-1964: *Natl. Speleol. Soc. Bull.*, v. 27, no. 1, p. 33-34, 1965.

DAVIS, C. F.

- 1-64. (and BOGGESE, D. H.) Water-table, surface-drainage, and engineering soils map of the Harrington quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-136, 1964[1965].

- 2-64. (and BOGGESE, D. H.) Water-table, surface-drainage, and engineering soils map of the Marydel area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-132, 1964[1965].

- 3-64. (and BOGGESE, D. H.) Water-table, surface-drainage, and engineering soils map of the Mispillion River quadrangle, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-137, 1964.

- 1-65. (BOGGESE, D. H., and COSKERY, O. J.) Water-table, surface-drainage, and engineering soils map of the Frederica area, Delaware: U.S. Geol. Survey Hydrol. Inv. Atlas HA-140, 1965.

DAVIS, G. H.

- 1-64. Management of water in arid lands: *Nat. History*, v. 73, no. 7, p. 26-33, 1964.

DAVIS, M. E.

- 1-65. Development of groundwater in the El Paso district, Texas, 1960-63—Progress report no. 9: *Texas Water Comm. Bull.* 6514, 34 p., 1965.

DAVIS, R. E.

- 1-63. Some physical properties of rocks penetrated in Pahute Mesa Exploratory Hole, 1,500 to 3,800 feet, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-3, 3 p., 1963.

- 1-64. Some physical properties of rocks penetrated in drill hole UE19d, 900 to 4,510 feet, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-17, 2 p., 1964.

DAVIS, R. W.

- 1-64. Availability of ground water in the Dexter quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-93, 1964.

- 2-64. Availability of ground water in the Lynn Grove quadrangle, Kentucky-Tennessee: U.S. Geol. Survey Hydrol. Inv. Atlas HA-112, 1964.

- 1-65. Availability of ground water in the Symsonia quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-157, 1965.

DAWDY, D. R.

- 1-64. (and O'DONNELL, Terence) Discussion of "Non-linear instantaneous unit-hydrograph theory", by Krishan

- DAWDY, D. R. --Continued
 Piara Singh: Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour., v. 90, no. HY5, p. 287-290, 1964.
- 1-65. Depth-discharge relations of alluvial streams—discontinuous rating curves: U.S. Geol. Survey Water-Supply Paper 1498-C, p. C1-C16, 1965. (Originally published 1961.)
- DeBREMAECKER, J. C.
 1-65. (GODSON, R. H., and WATKINS, J. S.) Attenuation measurements in the field [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 146, 1965.
- De CARLI, P. S.
 1-65. (and MILTON, D. J.) Stishovite—a synthesis by shock wave: Science, v. 147, p. 144-145, 1965.
- DECKER, R. W.
 1-65. Vertical ground displacements over the east rift of Kilauea volcano, Hawaii [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 185, 1965.
- De LAGUNA, Wallace
 1-64. Chemical quality of water, Brookhaven National Laboratory and vicinity, Suffolk County, New York: U.S. Geol. Survey Bull. 1156-D, p. D1-D73, 1964.
- DEMPSEY, W. J.
 1-65. (and VARGO, J. L.) Aeromagnetic map of parts of the Mt. Rose and the Virginia City quadrangles, Ormsby, Storey, and Washoe Counties, Nevada: U.S. Geol. Survey open-file report, 1965.
- DENNIS, P. E.
 1-64. (and HINDSON, L. L.) Contributions to the hydrology of Africa and the Mediterranean region: U.S. Geol. Survey Water-Supply Paper 1757-D, p. D1-D15, 1964.
- 2-64. (and HINDSON, L. L.) Ground-water provinces of southern Rhodesia: U.S. Geol. Survey Water-Supply Paper 1757-D, p. D1-D15, 1964.
- DENNY, C. S.
 1-64. Geology of the Brushart quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-324, 1964.
- 1-65. Alluvial fans in the Death Valley region, California and Nevada: U.S. Geol. Survey Prof. Paper 466, 62 p., 1965.
- 2-65. (and DREWES, Harald) Geology of the Ash Meadows quadrangle, Nevada-California: U.S. Geol. Survey Bull. 1181-L, p. L1-L56, 1965.
- DENSON, N. M.
 1-64. Geology, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 17-22, 1964.
- 2-64. The mineral industry in South Dakota, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 13-15, 1964.
- 3-64. (and MAPEL, W. J.) Summary and introduction, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 9-11, 1964.
- 4-64. (and RATTE, J. C.) Geology: Economic geology, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 33-37, 1964.
- 1-65. (and GILL, J. R.) Uranium-bearing lignite and carbonaceous shale in the southwestern part of the Williston basin—a regional study, with a section on Heavy minerals in Cretaceous and Tertiary rocks associated with uranium occurrences, by W. A. Chisholm: U.S. Geol. Survey Prof. Paper 463, 75 p., 1965.
- DEFTERMAN, R. L.
 1-65. (and REED, B. L.) Geochemical reconnaissance of stream sediments in the Iliamna quadrangle, Alaska: U.S. Geol. Survey open-file report, map, 1965.
- deWITT, Wallace, Jr.
 1-64. (and COLTON, G. W.) Bedrock geology of the Evitts Creek and Pattersons Creek quadrangles, Maryland, Pennsylvania, and West Virginia: U.S. Geol. Survey Bull. 1173, 90 p., 1964[1965].
- DIBBLEE, T. W., Jr.
 1-64. Geologic map of the Lucerne Valley quadrangle, San Bernardino County, California: U.S. Geol. Survey Geol. Inv. Map I-426, 1964.
- 2-64. Geologic map of the Ord Mountains quadrangle, San Bernardino County, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-427, 1964.
- 3-64. Geologic map of the Rodman Mountains quadrangle, San Bernardino County, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-430, 1964.
- 4-64. Geologic map of the San Geronio Mountain quadrangle, San Bernardino and Riverside Counties, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-431, 1964[1965].
- 1-65. Geologic map of the Cajon 7 1/2' quadrangle, San Bernardino County, California: U.S. Geol. Survey open-file report, 1965.
- 2-65. Geologic map of the Hesperia 15-minute quadrangle, San Bernardino County, California: U.S. Geol. Survey open-file report, 1965.
- DIMENT, W. H.
 1-64. Thermal conductivity of serpentinites from Mayaguez, Puerto Rico, and other localities, in A study of serpentinite, the AMSOC core hole near Mayaguez, Puerto Rico: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 92-106, 1964.
- 2-64. (and WEAVER, J. D.) Preliminary analysis of subsurface temperatures and heat flow in a hole near Mayaguez, Puerto Rico, in A study of serpentinite, the AMSOC core hole near Mayaguez, Puerto Rico: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 75-91, 1964.
- 1-65. (and WERRE, R. W.) Terrestrial heat flow in the central and southern Appalachians: Progress and problems [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 175, 1965.
- 2-65. (FRANCISCO ORTIZ, O., LOUIS SILVA, R., and CARLOS RUIZ, F.) Terrestrial heat flow at two localities near Vallenar, Chile [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 175, 1965.
- 3-65. (WERRE, R. W., BALDWIN, A. L., and SAUNDERS, E. D.) Heat flux through the bottom of a meromictic lake [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 175-176, 1965.
- 4-65. (RASPET, R., MAYHEW, M. A., and WERRE, R. W.) Terrestrial heat flow near Alberta, Virginia: Jour. Geophys. Research, v. 70, no. 4, p. 923-929, 1965.
- DINGMAN, R. J.
 1-64. Groundwater section of column "In the field": Kansas Univ. Newsletter, v. 64, no. 1, p. 5, 1964.
- DINGS, McC. G.
 1-65. (and WHITEBREAD, D. H.) Geology and ore deposits of the Metaline zinc-lead district, Pend Oreille County, Washington: U.S. Geol. Survey Prof. Paper 489, 109 p., 1965.
- DINWIDDIE, G. A.
 1-64. Municipal water supplies and uses, northeastern New Mexico: New Mexico State Engineer Tech. Rept. 29B, 64 p., 1964.
- 2-64. (and MOTTS, W. S.) Availability of ground water in parts of the Acoma and Laguna Indian Reservations, New Mexico: U.S. Geol. Survey Water-Supply Paper 1576-E, p. E1-E65, 1964.

DINWIDDIE, G. A. --Continued

- 1-65. New Mexico State Planning Office Report—Rio Grande basin: U.S. Geol. Survey open-file report, 42 p., 1965.

DIXON, H. R.

- 1-64. The Putnam Group of eastern Connecticut: U.S. Geol. Survey Bull. 1194-C, p. C1-C12, 1964.
1-65. (and SHAW, C. E., Jr.) Geologic map of the Scotland quadrangle, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-392, with text, 1965.

DOBROVOLNY, Ernest

- 1-65. (and SELKREGG, Lidia) Effects of the Good Friday earthquake on Anchorage, Alaska, and urban reconstruction [abs.]: Geol. Soc. America Spec. Paper 82, p. 356, 1965.
2-65. (MILLER, R. D., and HANSEN, W. R.) Engineering geologic effects of the March 27, 1964, earthquake at Anchorage, Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 46-47, 1965.

DOE, B. R.

- 1-65. (TILLING, R., and KLEPPER, M. R.) Lead isotope studies of major units of the Boulder batholith, Montana [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 165, 1965.
2-65. (TILTON, G. R., and HOPSON, C. A.) Lead isotopes in feldspars from selected granitic rocks associated with regional metamorphism: Jour. Geophys. Research, v. 70, p. 1947-1968, 1965.

DOELL, R. R.

- 1-65. (and COX, Allan) Measurement of the remanent magnetization of igneous rocks: U.S. Geol. Survey Bull. 1203-A, p. A1-A32, 1965.

DONNELLY, J. R.

- 1-64. Geology and oil-shale resources of the Green River Formation: Colorado School of Mines Quart. (First Symposium on Oil Shale), v. 59, no. 3, p. 153-163, 1964.
2-64. Mineral fuels and associated resources: Oil shale, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 67-73, 1964.

DORR, J. V. N., 2d

- 1-64. Supergene iron ores of Minas Gerais, Brazil: Econ. Geology, v. 59, no. 7, p. 1203-1240, 1964.
1-65. Manganese, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 183-195, 1965.
2-65. Nature and origin of the high-grade hematite ores of Minas Gerais, Brazil: Econ. Geology, v. 60, no. 1, p. 1-46, 1965.

DOTY, G. C.

- 1-65. New Mexico State Planning Office Report—Southwestern closed basins: U.S. Geol. Survey open-file report, 30 p., 1965.

DOUGLASS, R. C.

- 1-65. Larger foraminifers, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 20-25, 1965.
2-65. Photomicrography of thin sections, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 446-453, 1965.

DOYEL, W. W.

- 1-64. (and DINGMAN, R. J.) Hydrogeologic reconnaissance of the Republic of Korea, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D149-D152, 1964.

DRAKE, A. A., Jr.

- 1-64. (STEARN, T. W., and THOMAS, H. H.) Radiometric ages of zircon and biotite in quartz diorite, Eight Coast, Antarctica, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D50-D53, 1964.

DRAKE, A. A., Jr. --Continued

- 1-65. Carbonate rocks of Cambrian and Ordovician age, Northampton and Bucks Counties, eastern Pennsylvania and Warren and Hunterdon Counties, western New Jersey: U.S. Geol. Survey Bull. 1194-L, p. L1-L7, 1965.

DREWES, Harald

- 1-65. Thrust faults and glide faults in the Schell Creek Range near Ely, Nevada [abs.]: Geol. Soc. America Spec. Paper 82, p. 249, 1965.

DUCRET, G. L., Jr.

- 1-65. (and ANDERSON, D. B.) Records of wells, water levels and chemical quality of water in Baker Valley, Baker County, Oregon: Oregon State Engineer Ground Water Rept. 6, 34 p., 1965.

DUKE, M. B.

- 1-63. Metallic iron in basaltic achondrites, in Astrogeologic studies, Annual Progress Report, August 25, 1962-July 1, 1963—Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 197-207, 1963.

- 2-63. (and BRETT, Robin) Metallic copper in meteorites, in Astrogeologic studies, Annual Progress Report, August 25, 1962-July 1, 1963—Pt. C, Cosmochemistry and Petrography: U.S. Geol. Survey open-file report, p. 209-214, 1963.

- 1-65. Metallic iron in basaltic achondrites: Jour. Geophys. Research, v. 70, p. 1523-1527, 1965.

- 2-65. (and BRETT, Robin) Metallic copper in stony meteorites, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B101-B103, 1965.

DUNN, Bernard

- 1-65. Hydrology of the Irondequoit Creek basin, Rochester, New York: U.S. Geol. Survey open-file report, 40 p., 1965.

- 2-65. Time of travel studies, Fall Creek basin, Topkins County, New York: U.S. Geol. Survey open-file report, 13 p., 1965.

DUNRUD, C. R.

- 1-64. (and OSTERWALD, F. W.) Seismic study of coal mine bumps, Carbon and Emery Counties, Utah: U.S. Geol. Survey open-file report, 8 p., 1964.

DURFOR, C. N.

- 1-64. (and BECKER, Edith) Public water supplies of the 100 largest cities in the United States, 1962: U.S. Geol. Survey Water-Supply Paper 1812, 364 p., 1964.

- 2-64. (and BECKER, Edith) Selected data on public supplies of the 100 largest cities in the United States, 1962: Am. Water Works Assoc. Jour., v. 56, no. 3, p. 237-246, 1964.

- 1-65. (and BECKER, Edith) Chemical quality of public water supplies of the United States and Puerto Rico: U.S. Geol. Survey Hydrol. Inv. Atlas HA-200, 1965. (Originally published 1964.)

DURHAM, D. L.

- 1-64. (and ADDICOTT, W. O.) Upper Miocene and Pliocene marine stratigraphy in southern Salinas Valley, California: U.S. Geol. Survey Bull. 1194-E, p. E1-E7, 1964 [1965].

- 1-65. Geology of the Jolon and Williams Hill quadrangles, Monterey County, California: U.S. Geol. Survey Bull. 1181-Q, p. Q1-Q27, 1965.

DURY, G. H.

- 1-64. Principles of underfit streams: U.S. Geol. Survey Prof. Paper 452-A, p. A1-A67, 1964.

- 2-64. Subsurface exploration and chronology of underfit streams: U.S. Geol. Survey Prof. Paper 452-B, p. B1-B56, 1964.

- 1-65. Theoretical implications of underfit streams: U.S. Geol. Survey Prof. Paper 452-C, p. C1-C43, 1965.

DUTCHER, L. C.

- 1-65. Progress report on water studies in the Bloomington-Colton area, Upper Santa Ana Valley, California, 1964: U.S. Geol. Survey open-file report, 30 p., 1965.
- 2-65. Progress report on water studies in the San Timoteo-Smiley Heights area, Upper Santa Ana Valley, California, 1964: U.S. Geol. Survey open-file report, 31 p., 1965.
- 3-65. (and FRENCH, J. J.) Progress report on water studies in the Chino-Corona area, Upper Santa Ana Valley, California, 1964, Pt. I: U.S. Geol. Survey open-file report, 36 p., 1965.

DUTRO, J. T., Jr.

- 1-65. Brachiopods, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 44-48, 1965.

DYNI, J. R.

- 1-65. (and CULLINS, H. L.) Meeker and Loyd Sandstone Members of the Mancos Shale, Moffat and Rio Blanco Counties, Colorado: U.S. Geol. Survey Bull. 1194-J, p. J1-J7, 1965.

EAKIN, T. E.

- 1-64. Regional ground-water system in southeastern Nevada [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 51, 1964.
- 2-64. (and MOORE, D. O.) Uniformity of discharge of Muddy River Springs, southeastern Nevada, and relation to interbasin movement of ground water, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D171-D176, 1964.
- 3-64. (and WINOGRAD, I. J.) Interbasin movement of ground water in south-central Nevada—Some implications [abs.]: Geol. Soc. America Ann. Mtg., Miami, Florida, Nov. 1964, Program, p. 52, 1964.

EARGLE, D. H.

- 1-63. Southwest-northeast cross section, Tatum dome, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-29, 2 p., 1963.

EATON, G. P.

- 1-64. Preliminary gravity survey of areas of subsidence on a reservoir floor [abs.]: Assoc. Eng. Geologists Ann. Mtg. 1964, Sacramento, Calif., Program, p. 12, 1964.
- 1-65. Estimating average total porosity of an alluvial aquifer from gravity measurements [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 78, 1965.

EATON, J. P.

- 1-63. Volcano geophysics, *in* United States National Report 1960-1963: Internat. Union Geodesy and Geophysics, General Assembly, 13th, Berkeley, Calif., Aug. 1963, p. 507-508, 1963.

EBERLEIN, G. D.

- 1-64. Summary and introduction, *in* Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 9-12, 1964.

EDELLEN, G. W., Jr.

- 1-64. (RUGGLES, F. H., Jr., and CROSS, W. P.) Floods at Chillicothe, Ohio: U.S. Geol. Survey Hydrol. Inv. Atlas HA-45, 1964.
- 2-64. (RUGGLES, F. H., Jr., and CROSS, W. P.) Floods at Circleville, Ohio: U.S. Geol. Survey Hydrol. Inv. Atlas HA-48, 1964.
- 3-64. (RUGGLES, F. H., Jr., and CROSS, W. P.) Floods at Mount Vernon, Ohio: U.S. Geol. Survey Hydrol. Inv. HA-40 (revised) 1964.
- 4-64. (RUGGLES, F. H., Jr., and CROSS, W. P.) Floods at Newark, Ohio: U.S. Geol. Survey Hydrol. Inv. Atlas HA-44 (Revised) 1964.
- 5-64. (RUGGLES, F. H., Jr., and CROSS, W. P.) Floods at Zanesville, Ohio: U.S. Geol. Survey Hydrol. Inv. Atlas HA-46, 1964.

EGE, J. R.

- 1-62. (and JOHNSON, R. B.) Additional physical property data of rock samples from Area 401, Nevada Test Site: U.S. Geol. Survey Tech. Letter, PLUTO-21, supp. 1, 4 p., 1962.

EGGLETON, R. E.

- 1-63. Preliminary geology of the Rhipaeus quadrangle of the Moon and definition of the Fra Mauro Formation, *in* Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 46-62, 1963.

EISENHUTH, H. P.

- 1-65. Index of surface-water records to December 31, 1963—Pt. 7, Lower Mississippi River Basin: U.S. Geol. Survey Circ. 507, 55 p., 1965.
- 2-65. Index of surface-water records to December 31, 1963—Pt. 9, Colorado River basin: U.S. Geol. Survey Circ. 509, 49 p., 1965.
- 3-65. Index of surface-water records to December 31, 1963—Pt. 13, Snake River basin: U.S. Geol. Survey Circ. 513, 29 p., 1965.
- 4-65. Index of surface-water records to December 31, 1963—Pt. 14, Pacific slope basins in Oregon and lower Columbia River basin: U.S. Geol. Survey Circ. 514, 28 p., 1965.

EISENLOHR, W. S.

- 1-64. Discharge ratings for streams at submerged section controls: U.S. Geol. Survey Water Supply Paper 1779-L, p. L1-L32, 1964.

EKREN, E. B.

- 1-63. Geology of Silent Canyon SW quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-66, 14 p., 1963.
- 1-64. (ANDERSON, R. E., ORKILD, P. P., and HINRICH, E. N.) Geologic map of the Silent Butte quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report, 1 p., 1964.

- 2-64. (and SARGENT, K. A.) Geologic map of the Skull Mountain quadrangle at the Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey open-file report, 1964.

- 1-65. (and HOUSER, F. N.) Geology and petrology of the Ute Mountains area, Colorado: U.S. Geol. Survey Prof. Paper 481, 74 p., 1965.

ELLIS, M. J.

- 1-64. (and ADOLPHSON, D. G.) Hydrogeology of the glacial drift in the Skunk Creek-Lake Madison drainage basin, southeastern South Dakota: U.S. Geol. Survey open-file report, 40 p., 1964.

- 1-65. (and ADOLPHSON, D. G.) Ground-water resources at three towns on the Standing Rock Indian Reservation in South Dakota: U.S. Geol. Survey open-file report, 45 p., 1965.

ELSTON, D. P.

- 1-63. Pre-Imbrian stratigraphy of the Colombo quadrangle, *in* Astrogeologic studies, Annual Progress Report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 99-109, 1963.

- 2-63. (and MILTON, D. J.) A low velocity impact crater at Hellhole Bend, Little Colorado River, Arizona, *in* Astrogeologic studies, Annual Progress Report, August 25, 1962-July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 77-90, 1963.

- 1-65. Stratigraphic relations in eastern part of the Moon [abs.]: Geol. Soc. America Spec. Paper 82, p. 56, 1965.

EMERICK, W. L.

- 1-63. Geologic appraisal of the U20a drill hole area, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-7, 8 p., 1963.

EMERICK, W. L. --Continued

- 2-63. Preliminary report on the geology of the U12g tunnel, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Area-12-15, 19 p., 1963.
- 3-63. (and HASLER, J. W.) Geologic section through U12n tunnel, Rainier Mesa, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-10, 3 p., 1963.
- 4-63. (HOOVER, D. L., SNYDER, R. P., and HASLER, J. W.) Physical properties of tuff in chamber and workings of U12q drill hole, Rainier Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Area-12-9, 4 p., 1963.
- 5-63. (SNYDER, R. P., and BOWERS, W. E.) Outline of the geology of the U12b.10 tunnel, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-5, 11 p., 1963.
- 6-63. (SNYDER, R. P., and HOOVER, D. L.) Average physical properties of tuff in the vicinity of ground zero, U12g.01 tunnel, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-4, 3 p., 1963.
- 1-64. Exploratory drill holes UE1a, UE1b, and UE1c, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-54, 11 p., 1964.
- 1-65. Lithologic log of the U12g.05-U1 core drill hole, Area 12, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Area-12-17, 5 p., 1965.

EMERY, K. O.

- 1-65. (MERRILL, A. S., and TRUMBULL, J. V. A.) Geology and biology of the sea floor as deduced from simultaneous photographs and samples: *Limnology and Oceanography*, v. 10, no. 1, p. 1-21, 1965.

EMERY, P. A.

- 1-64. Basic-data report, Saline County, Nebraska: Nebraska Univ. Conserv. and Survey Div., Nebraska Water Survey Paper 15, 29 p., 1964.
- 2-64. Geology and ground-water resources of Richardson County, Nebraska: U.S. Geol. Survey Water-Supply Paper 1779-W, p. W1-W29, 1964.
- 1-65. Effect of ground-water pumping on stream flow and ground-water levels, Blue River basin, Nebraska: U.S. Geol. Survey open-file report, 11 p., 1965.
- 2-65. (and MALHOIT, M. M.) Water levels in observation wells in Nebraska, 1964: Nebraska Univ. Conserv. and Survey Div. Nebraska Water Survey Paper 17, 163 p., 1965; also in U.S. Geol. Survey open-file report, 165 p, 1965.

EMMETT, W. W.

- 1-64. (and LEOPOLD, L. B.) Discussion of "Geometry of river channels", by W. B. Langbein: *Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour.*, v. 90, no. HY5, p. 277-285, 1964.
- 2-64. (and WALLACE, J. R.) Errors in piezometric measurement: *Am. Soc. Civil Engineers Proc. Paper 4126*, Hydraulics Div. Jour., v. 90, no. HY6, p. 45-61, 1964.

ENGEL, A. E. J.

- 1-63. (and ENGEL, C. G.) Metasomatic origin of large parts of the Adirondack phacoliths: *Geol. Soc. America Bull.*, v. 74, no. 3, p. 349-354, 1963.
- 1-64. (and ENGEL, C. G.) Composition of basalts from the Mid-Atlantic Ridge: *Science*, v. 144, no. 3624, p. 1330-1333, 1964.
- 2-64. (and ENGEL, C. G.) Continental accretion and the evolution of North America, in *Advancing frontiers in geology and geophysics, a volume in honour of M. S. Kirshnan*: Hyderabad, India, Osmania Univ. Press, p. 17-37e, 1964[1965].
- 3-64. (and ENGEL, C. G.) Igneous rocks of the East Pacific Rise: *Science*, v. 146, no. 3643, p. 477-485, 1964.

ENGEL, A. E. J. --Continued

- 1-65. (and ENGEL, C. G.) Igneous rocks of the oceans [abs.]: *Geol. Soc. America Spec. Paper 82*, p. 56-57, 1965.

ENGEL, C. G.

- 1-63. (and ENGEL, A. E. J.) Basalts dredged from the northeastern Pacific Ocean: *Science*, v. 140, no. 3573, p. 1321-1324, 1963.

ENGLUND, K. J.

- 1-64. Geology of the Middlesboro South quadrangle, Tennessee-Kentucky-Virginia: U.S. Geol. Survey Geol. Quad. Map GQ-301, 1964.

ERD, R. C.

- 1-64. (WHITE, D. E., FAHEY, J. J., and LEE, D. E.) Buddingtonite, an ammonium feldspar with zeolitic water: *Am. Mineralogist*, v. 49, nos. 7-8, p. 831-849, 1964.
- 1-65. (LYON, R. J. P., and MADSEN, B. M.) Infrared studies of saline sulfate minerals: discussion: *Geol. Soc. America Bull.*, v. 76, no. 2, p. 271-282, 1965.

ERICKSEN, G. E.

- 1-65. Geology of the Chilean nitrate deposits [abs.]: *Geol. Soc. America Spec. Paper 82*, p. 57-58, 1965.

ERICKSON, R. L.

- 1-64. (and MARRANZINO, A. P.) Copper determinations on samples of alluvium and caliche from the Rocky Range, Beaver County, Utah: U.S. Geol. Survey open-file report, 1 p., 1964.
- 2-64. (MARRANZINO, A. P., ODA, Uteana, and JANES, W. W.) Geochemical exploration near the Getchell Mine, Humboldt County, Nevada: U.S. Geol. Survey Bull. 1198-A, p. A1-A26, 1964.
- 3-64. (MASURSKY, Harold, MARRANZINO, A. P., ODA, Uteana, and JANES, W. W.) Semiquantitative spectrographic and chemical analyses of rocks from the lower plate of the Roberts Thrust, north-central part of the Cortez quadrangle, Nevada: U.S. Geol. Survey open-file report, 2 p., 1964.

ERSKINE, C. F.

- 1-65. Landslides in the vicinity of the Fort Randell Reservoir, South Dakota: U.S. Geol. Survey open-file report, 317 p., 1965.

ESPENSHADE, G. H.

- 1-64. (and BOUDETTE, E. L.) Geology of the Greenville quadrangle, Maine: U.S. Geol. Survey Geol. Quad. Map GQ-330, 1964.

EVANS, H. T., Jr.

- 1-63. Uranyl ion coordination: *Science*, v. 141, p. 154-158, 1963.
- 1-64. (SHALLOW, E. G., and BARNES, W. H.) The structures of the decavanadate ion: *Jour. Am. Chemical Soc.*, v. 86, p. 4209, 1964.
- 1-65. The crystal structure of the decavanadate, $K_2Zn_2V_{10}O_{28} \cdot 16H_2O$, the chemical analog of hummerite pascoite [abs.]: *Am. Mineralogist*, v. 50, nos. 1-2, p. 284-285, 1965.

EVENSON, R. E.

- 1-64. Suitability of irrigation water and changes in ground-water quality in the Lompoc subarea of the Santa Ynez River basin, Santa Barbara County, California: U.S. Geol. Survey open-file report, 57 p., 1964.

EVERETT, D. E.

- 1-64. (and RUSH, F. E.) Ground-water appraisal of Smith Creek and Ione valleys, Lander and Nye Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources, Ground-Water Resources-Reconn. Ser. Rept. 28, 19 p., 1964.

FADER, S. W.

- 1-64. (GUTENTAG, E. D., LOBMEYER, D. H., and MEYER, W. R.) Geohydrology of Grant and Stanton

FADER, S. W. --Continued

Counties, Kansas: Kansas Geol. Survey Bull. 168, 147 p., 1964.

FAHEY, J. J.

1-63. Separation of stishovite and coesite from sandstone shocked by meteorite impact, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963--Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 31-38, 1963.

FAUST, G. T.

1-64. (and FAHEY, J. J.) The serpentine-group minerals: U.S. Geol. Survey Prof. Paper 384-A, p. A1-A92, 1964. (Originally published 1962.)

FEININGER, Tomas

1-65. Petrology of the Ashaway and Voluntown quadrangles, Connecticut-Rhode Island [abs.]: Dissert. Abs., v. 25, no. 8, p. 4649-4650, 1965.

FERAY, D. E.

1-65. The role of tectonic and environmental factors in the origin and distribution of sediments [abs.]: South Texas Geol. Soc. Bull., v. 5, no. 1, p. 7-8, 1965.

FERNALD, A. T.

1-64. Surficial geology of the central Kobuk River Valley, northwestern Alaska: U.S. Geol. Survey Bull. 1181-K, p. K1-K31, 1964.

2-64. (and HOROWITZ, A. S.) Bedrock geology of the Nunatarssuaq area, northwest Greenland: Medd. om Grønland, v. 172, no. 6, 44 p., 1964.

1-65. Late Quaternary chronology, upper Tanana River Valley, eastern Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 60-61, 1965.

FERRIS, J. G.

1-65. (KNOWLES, D. B., BROWN, R. H., and STALLMAN, R. W.) Theory of aquifer tests: U.S. Geol. Survey Water-Supply Paper 1536-E, p. E69-E174, 1965. (Originally published 1962.)

FETH, J. H.

1-64. Hidden recharge: Ground Water, v. 2, no. 4, p. 14-17, 1964.

2-64. Review and annotated bibliography of ancient lake deposits (Precambrian to Pleistocene) in the Western States: U.S. Geol. Survey Bull. 1080, 119 p., 1964.

3-64. (ROGERS, S. M., and ROBERSON, C. E.) Chemical composition of snow in the northern Sierra Nevada and other areas: U.S. Geol. Survey Water-Supply Paper 1535-J, p. J1-J39, 1964.

1-65. Calcium, sodium, sulfate, and chloride in stream water of the western conterminous United States to 1957: U.S. Geol. Survey Hydrol. Inv. Atlas HA-189, 1965.

FEULNER, A. J.

1-64. Galleries and their use for development of shallow ground-water supplies, with special reference to Alaska: U.S. Geol. Survey Water-Supply Paper 1809-E, p. E1-E16, 1964.

2-64. (and SCHUPP, R. G.) Temperature and chemical quality of water from a well drilled through permafrost near Bethel, Alaska, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D144-D148, 1964.

FINCH, W. I.

1-64. Epigenetic uranium deposits in sandstone in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D76-D78, 1964.

2-64. Geology of the Symsonia quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-326, 1964.

3-64. (OLIVE, W. E., and WOLFE, E. W.) Ancient Lake in Western Kentucky and Southern Illinois, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C130-C133, 1964.

FINCH, W. I. --Continued

1-65. Geology of the Mayfield quadrangle, Graves County, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-372, 1965.

FINNELL, T. L.

1-64. Geology of the Manchester quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-318, 1964.

FISCHER, R. P.

1-64. Ferrous metals: Vanadium, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 115-120, 1964.

2-64. Metallic mineral resources: Vanadium, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 165-175, 1964.

3-64. Metallic mineral resources: Vanadium, in *Mineral and water resources of South Dakota*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 80-81, 1964.

4-64. Mineral resources: Vanadium, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 240-253, 1964.

5-64. (and KING, R. U.) Trends in the consumption and supply of molybdenum and vanadium: Am. Inst. Mining Metall. Engineers Trans., preprint 64-K-2, 19 p., 1964.

6-64. (and VINE, J. D.) Metallic mineral resources: Vanadium, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 133-135, 1964.

1-65. Vanadium, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 226-228, 1965.

FISCHER, W. A.

1-64. (MOXHAM, R. M., POLCYN, F., and LANDIS, G. H.) Infrared surveys of Hawaiian volcanoes: Science, v. 146, no. 3645, p. 733-742, 1964.

1-65. Some uses of infrared imagery in water resources investigations [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 78, 1965.

FISCHER, W. L.

1-65. The search for nonfuel minerals: South Texas Geol. Soc. Bull., v. 5, no. 2, p. 6-12, 1965.

FISHMAN, M. J.

1-64. (and SKOUGSTAD, M. W.) Catalytic determination of vanadium in water: Anal. Chemistry, v. 36, no. 8, p. 1643-1646, 1964.

2-64. (SKOUGSTAD, M. W., and SCARBRO, G. F., Jr.) Diazotization method for nitrate and nitrite: Am. Water Works Assoc. Jour., v. 56, no. 5, p. 633-638, 1964.

1-65. (and SKOUGSTAD, M. W.) Rapid field and laboratory determination of phosphate in natural water, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B167-B169, 1965.

FLANAGAN, F. J.

1-64. Judging the analytical ability of rock analysts by Chisquared, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C157-C158, 1964.

FLEISCHER, Michael

1-64. Manganese oxide minerals, VIII, Hollandite, in *Advancing frontiers in geology and geophysics*, a volume in honour of M. S. Kirshnan: Hyderabad, India, Osmania Univ. Press, p. 221-232, 1964[1965].

2-64. Review of "Mineralogy, Spravochnik, Tom I and Tom II" (Handbook on Mineralogy): Am. Mineralogist, v. 49, p. 1781-1782, 1964.

3-64. Translation of "The problem of the abundance of minor and rare-earth elements and minerals in effusive rocks", by V. I. Fin'ko: Geochemistry International, v. 2, p. 390-394, 1964[1965].

FLEISCHER, Michael--Continued

1-65. Memorial to Herman Yagoda: *Am. Mineralogist*, v. 50, nos. 3-4, p. 559-562, 1965.

2-65. Review of "Mineralogy of the rare earths", by E. I. Semenov: *Am. Mineralogist*, v. 50, p. 525, 1965.

FLOYD, E. O.

1-65. Geology and ground-water resources of the Monroe area, North Carolina: North Carolina Dept. Water Resources Ground-Water Bull. 5, 109 p., 1965.

FOLLETT, C. R.

1-65. Ground-water resources of Caldwell County, Texas: U.S. Geol. Survey open-file report, 214 p., 1965.

FORD, A. B.

1-64. Cordierite-bearing, hypersthene-quartz monzonite porphyry in the Thiel Mountains and its regional importance, in *Antarctic geology*; Proc. First International Symposium on Antarctic Geology, Cape Town, Sept. 1963: New York, John Wiley and Sons, p. 429-441, 1964[1965].

2-64. Review of Antarctic geology: *Internat. Geophysics Bull.*, no. 82, p. 1-19, 1964.

FOSBERG, F. R.

1-64. Résumé of the Cycadaceae: *Federation Proc.*, v. 23, no. 6, pt. 1, p. 1340-1341, 1964.

FOSTER, H. L.

1-65. (and HOLMES, G. W.) A large transitional rock glacier in the Johnson River area, Alaska Range [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 64, 1965.

2-65. (and HOLMES, G. W.) A large transitional rock glacier in the Johnson River area, Alaska Range, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B112-B116, 1965.

FOSTER, M. D.

1-65. Studies of the zeolites—Composition of zeolites of the natrolite group and compositional relations among thomsonites, gonnardites, and natrolites: U.S. Geol. Survey Prof. Paper 504-D, -E, p. D1-D7, E1-E10, 1965.

FOURNIER, R. O.

1-65. (and HEMLEY, J. J.) Effects of supersaturated silica solutions during the hydrothermal alteration of feldspars [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 64, 1965.

FOXWORTHY, B. L.

1-64. (HOGENSON, G. M., and HAMPTON, E. R.) Records of wells and springs, water levels and chemical quality of ground water in the East Portland area, Oregon: *Oregon State Engineer Ground Water Rept.* 3, 79 p., 1964.

FRANK, F. J.

1-64. South Gila Valley, Yuma Mesa, and Yuma Valley area, in *Annual report on groundwater in Arizona, Spring 1963 to Spring 1964*: Arizona Land Dept. Water Resources Rept. 19, p. 46, 1964.

FRANKLIN, G. J.

1-65. Geology of the Hanson quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-365, 1965.

FREDERICK, B. J.

1-64. Measurement of turbine discharge with radioisotopes: U.S. Geol. Survey open-file report, TEI 855, 32 p., 1964.

FREEMAN, V. L.

1-64. Geologic map of the Langtry quadrangle, Val Verde County, Texas: U.S. Geol. Survey Misc. Geol. Inv. Map I-422, 1964.

2-64. Geologic map of the Shumla quadrangle, Val Verde County, Texas: U.S. Geol. Survey Misc. Geol. Inv. Map I-424, 1964.

1-65. Geologic map of the Bakers Crossing quadrangle, Val Verde County, Texas: U.S. Geol. Survey Misc. Geol. Inv. Map I-434, 1965.

FRENCH, J. J.

1-65. (DUTCHER, L. C., and DANA, S. W.) Progress report on water studies in the Chino-Corona area, Upper Santa Ana Valley, California, 1964—Pt. II: U.S. Geol. Survey open-file report, 29 p., 1965.

2-65. (and PEARSON, E. G.) A brief water-resources reconnaissance of Pala and Rincon Indian Reservations, San Diego County, California: U.S. Geol. Survey open-file report, 41 p., 1965.

FRITTS, C. E.

1-65. Bedrock geologic map of the Ansonia quadrangle, Fairfield and New Haven Counties, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-426, 1965.

2-65. Bedrock geologic map of the Milford quadrangle, Fairfield and New Haven Counties, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-427, 1965.

3-65. Stratigraphy, structure, and granitic rocks in the Marenisco-Watersmeet area, Michigan [abs.], in *Eleventh Annual Institute on Lake Superior Geology*: St. Paul, Minn., Univ. Minnesota, May 1965, p. 15, 1965.

FRYKLUND, V. C., Jr.

1-64. Ore deposits of the Coeur d'Alene district, Shoshone County, Idaho, with a section on The bleached rock in the Coeur d'Alene district, by P. L. Weiss: U.S. Geol. Survey Prof. Paper 445, 103 p., 1964.

FURNESS, L. W.

1-64. (BURNES, C. V., and BUSBY, M. W.) Kansas streamflow characteristics—Pt. 5, Storage requirements to control high flow: *Kansas State Water Resources Board Tech. Rept.* 5, 169 p., 1964.

GABRYSCH, R. K.

1-65. Discussion of "Effects of drought on ground-water supplies", by J. R. Mount, in *Symposium on consideration of droughts in water planning*: Texas Water Comm. Bull. 6512, p. 169-170, 1965.

GALLAHER, J. T.

1-64. Geology and hydrology of alluvial deposits along the Ohio River in the Henderson area, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-91, 1964[1965].

2-64. Geology and hydrology of alluvial deposits along the Ohio River in the Stanley area, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-110, sheets 1 and 2, 1964.

3-64. Geology and hydrology of alluvial deposits along the Ohio River between the Uniontown area and Wickliffe, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-129, sheets 1 and 2, 1964.

GAMBELL, A. W.

1-64. (and FISHER, D. W.) Occurrence of sulfate and nitrate in rainfall: *Jour. Geophys. Research*, v. 69, no. 20, p. 4203-4210, 1964.

GARBER, M. S.

1-63. Large rise and following decline of water-level in well 7, Area 3, Nevada Test Site—An effect from the Aardvark underground nuclear explosion: U.S. Geol. Survey Tech. Letter YUCCA-48, 9 p., 1963.

GARD, L. M.

1-63. (and HAZLEWOOD, R. M.) Seismic refraction surveys at Saylor Creek site, Idaho: U.S. Geol. Survey Tech. Letter SCHOONER-1, 3 p., 1963.

2-63. (and HAZLEWOOD, R. M.) Seismic refraction survey at Winnemucca site, Nevada: U.S. Geol. Survey Tech. Letter SCHOONER-2, 4 p., 1963.

3-63. (and HASLER, J. W.) Geology of proposed Schooner site, Bruneau River area, Owyhee County, Idaho (A preliminary investigation): U.S. Geol. Survey Tech. Letter SCHOONER-3, 24 p., 1963.

GARZA, Sergio, compiler

1-64. Ground-water discharge from the Edwards and associated limestones, San Antonio area, Texas, 1963: *Edwards Underground Water Dist. Bull.* 5, 3 p., 1964.

- GARZA, Sergio--Continued
- 1-65. Chemical analyses of water from observation wells in the Edwards and associated limestones, San Antonio area, Texas, 1964: Edwards Underground Water Dist. Bull. 7, 12 p., 1965.
- GATES, G. O.
- 1-64. Geology:Geologic and tectonic setting, *in* Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 32-39, 1964.
- GATES, J. W.
- 1-65. Reevaluation of the ground-water resources of Tooele Valley, Utah: U.S. Geol. Survey open-file report, 151 p., 1965.
- GATEWOOD, J. S.
- 1-64. (WILSON, Alfonso, THOMAS, H. E., and KISTER, L.R.) General effects of drought on water resources of the southwest: U.S. Geol. Survey Prof. Paper 372-B, p. B1-B55, 1964.
- GAULT, D. E.
- 1-64. (HEITOWIT, E. D., and MOORE, H. J.) Ejecta with complex structures from craters formed by hypervelocity impact [abs.]: Am. Geophys. Union Trans., v. 1, p. 74, 1964.
 - 2-64. (HEITOWIT, E. D., and MOORE, H. J.) Some observations of hypervelocity impacts with porous media, *in* The lunar surface layer--materials and characteristics: New York, Academic Press, p. 151-178, 1964.
 - 1-65. (and MOORE, H. J.) Scaling relationships for micro-scale to megascale impact craters: Tampa, Florida, Hypervelocity Impact Symposium, 7th, 1964, Proc., v. 6, p. 341-351, 1965.
- GERE, W. C.
- 1-64. Mineral resources:Phosphate rock, *in* Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 152-161, 1964.
 - 2-64. Nonmetallic and industrial minerals and materials resources:Phosphate, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 195-205, 1964.
 - 3-64. (HORTON, G. W., HARSTEAD, J. N., and RUSSELL, D. F.) Mineral and fuels and associated resources: Oil and natural gas, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 51-60, 1964.
- GERLACH, A. C.
- 1-64. New approaches to the geography of the United States--introduction: Assoc. Am. Geographers Annals, v. 54, no. 1, p. 1, 1964.
- GIBSON, T. G.
- 1-65. Foraminifera:Summary of investigations conducted in 1964: Woods Hole Oceanographic Inst. Ref. No. 65-13, p. 69-70, 1965.
 - 2-65. Stratigraphy and paleoenvironmental analysis of the Lee's Creek North Carolina phosphate mine [abs.]: Geol. Soc. America Spec. Paper 82, p. 72, 1965.
- GIESSNER, F. W.
- 1-64. A reconnaissance of the geology and water resources of the Mission Creek Indian Reservation, Riverside County, California: U.S. Geol. Survey open-file report, 31 p., 1964.
 - 1-65. Ground-water conditions during 1964 at the Marine Corps Base, Twentynine Palms, California: U.S. Geol. Survey open-file report, 30 p., 1965.
- GILBERT, B. K.
- 1-65. (and KAMMERER, J. C.) Summary of water-resources records at principal measurement sites in the Genesee River basin through 1963: U.S. Geol. Survey open-file report, 61 p., 1965.
- GILBERT, C. R.
- 1-64. (COMMONS, G. G., KOBERT, G. E., and KENNON, F. W.) Hydrologic studies of small watersheds, Honey Creek basin, Collin and Grayson Counties, Texas, 1953-59: U.S. Geol. Survey Water-Supply Paper 1779-F, p. F1-F98, 1964.
 - 1-65. Data collecting programs during droughts, *in* Symposium on consideration of droughts in water planning: Texas Water Comm. Bull. 6512, p. 35-45, 1965.
- GILL, H. E.
- 1-64. (and VECCHIOLI, John) Availability of ground water in Morris County, New Jersey: U.S. Geol. Survey open-file report, 71 p., 1964.
- GILL, J. R.
- 1-65. (and COBBAN, W. A.) Stratigraphy of the Pierre Shale, Valley City and Pembina Mountain areas, North Dakota: U.S. Geol. Survey Prof. Paper 392-A, p. A1-A20, 1965.
- GILLULY, James
- 1-65. Volcanism, tectonism, and plutonism in the western United States: Geol. Soc. America Spec. Paper 80, 69 p., 1965.
 - 2-65. (and GATES, Olcott) Tectonic and igneous geology of the northern Shoshone Range, Nevada, with sections on Gravity in Crescent Valley, by Donald Plouff and Economic geology, by K. B. Ketner: U.S. Geol. Survey Prof. Paper 465-B, 153 p., 1965.
- GILSTRAP, R. C.
- 1-64. (and CHRISTENSEN, R. C.) Floods of July 16-17, 1963, in vicinity of Hot Springs, Arkansas: U.S. Geol. Survey open-file report, 6 p., 1964.
- GIROUX, P. R.
- 1-64. (HENDRICKSON, G. E., STOINMENOFF, L. E., and WHETSTONE, G. W.) Water resources of Van Buren County, Michigan: Michigan Geol. Survey Water Inv. 3, 144 p., 1964; also in U.S. Geol. Survey open-file report, 140 p., 1964.
 - 2-64. (and HUFFMAN, G. C.) Summary of ground-water conditions in Michigan in 1963: Michigan Geol. Survey Div. [Water Supply Rept. 8], 100 p., 1964.
- GIUSTI, E. V.
- 1-65. (and SCHNEIDER, W. J.) The distribution of branches in river networks: U.S. Geol. Survey Prof. Paper 422-G, p. G1-G10, 1965.
- GLICK, E. E.
- 1-64. (FREZON, S. E., and GORDON, Mackenzie, Jr.) Witts Springs Formation of Marrow age in the Snowball quadrangle, north-central Arkansas: U.S. Geol. Survey Bull. 1194-D, p. D1-D16, 1964[1965].
- GODWIN, L. H.
- 1-64. (and GASKILL, D. L.) Post-Paleocene West Elk laccolithic cluster, west-central Colorado, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C64-C65, 1964.
- GOERLITZ, D. F.
- 1-65. (and LAMAR, W. L.) Microcoulometric gas chromatographic analysis of selected herbicides in water: U.S. Geol. Survey open-file report, 14 p., 1965.
- GOLDSMITH, Richard
- 1-64. Surficial geology of the Niantic quadrangle, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-329, 1964.
 - 1-65. Preliminary bedrock geologic map of the New London quadrangle, New London County, Connecticut, and Suffolk County, New York: U.S. Geol. Survey open-file report, 1965.
 - 2-65. Preliminary bedrock geologic map of the Niantic quadrangle, New London County, Connecticut: U.S. Geol. Survey open-file report, 1965.

- GORDON, E. D.
1-64. Cooperative ground-water investigations in Wyoming during 1963 and 1964, *in* 37th Bienn. Rept. of the State Engineer to the Governor of Wyoming: Wyoming State Engineer, p. 90-99, 1964.
- GORDON, MacKenzie, Jr.
1-64. California Carboniferous cephalopods: U.S. Geol. Survey Prof. Paper 483-A, p. A1-A27, 1964.
2-64. Carboniferous cephalopods of Arkansas: U.S. Geol. Survey Prof. Paper 460, 322 p., 1964[1965].
- GOTT, G. B.
1-64. Pre-Fall River folding in the southern part of the Black Hills, South Dakota, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D28-D29, 1964.
2-64. (and McCARTHY, J. H., Jr.) Distribution of mercury, silver, tellurium, arsenic, and antimony in the Rowe Canyon area, White Pine County, Nevada: U.S. Geol. Survey open-file report, 1 p., 1964.
3-64. (and PIPERINGOS, G. N.) Metallic mineral resources: Uranium, *in* Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 50-56, 1964.
- GOTTFRIED, David
1-65. (and DINNIN, J. I.) Distribution of tantalum in some igneous rocks and coexisting minerals of the Southern California batholith, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B96-B100, 1965.
- GOWER, H. D.
1-64. (and MADSEN, B. M.) The occurrence of phosphate rock in California, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D79-D85, 1964.
1-65. (and PEASE, M. H., Jr.) Geology of the Montesano quadrangle, Washington: U.S. Geol. Survey Geol. Quad. Map GQ-374, 1965.
- GRANT, R. E.
1-65. The brachiopod superfamily Stenosismataceae: Smithsonian Misc. Colln., v. 148, no. 2, 192 p., 1965.
- GRANTHAM, P. E.
1-63. (and WASSON, B. E.) Inventory of wells near calibration shot points in south Mississippi: U.S. Geol. Survey Tech. Letter DRIBBLE-36, 7 p., 1963.
- GRANTZ, Arthur
1-62. Geography and geology of Simeonof Island, Shumagin Islands Group, southwestern Alaska: U.S. Geol. Survey Tech. Letter SIMEONOF-1, 15 p., 1962.
1-64. Mineral fuel resources: Southern Alaska, *in* Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 44-62, 1964.
2-64. (and PATTON, W. W., Jr.) Mineral fuel resources: Petroleum and natural gas, *in* Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 43-44, 1964.
1-65. Geologic map and cross sections of the Nelchina area, south-central Alaska: U.S. Geol. Survey open-file report, 4 sheets, 1965.
2-65. (JONES, D. L., and LANPHERE, M. A.) Upper Mesozoic rocks of the lower Chitina Valley, Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 356, 1965.
3-65. (PLAFKER, George, and CASE, J. E.) Tectonics of Alaska's Good Friday earthquake [abs.]: Geol. Soc. America Spec. Paper 82, p. 77, 1965.
- GREEN, A. R.
1-64. Magnitude and frequency of floods in the United States—Pt. 1-A, North Atlantic slope basins, Maine to GREEN, A. R. --Continued
Connecticut: U.S. Geol. Survey Water-Supply Paper 1671, 260 p., 1964.
- GRIFFITTHS, W. R.
1-64. Metallic mineral resources: Beryllium, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 70-75, 1964.
2-64. Metallic mineral resources: Beryllium, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 71-75, 1964.
3-64. Metallic mineral resources: Lithium, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 113, 1964.
4-64. Miscellaneous metals: Beryllium, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 120-124, 1964.
5-64. (and SAVAGE, C. N.) Mineral resources: Beryllium, *in* Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 49-51, 1964.
1-65. Beryllium, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 196-200, 1965.
- GRIGGS, A. B.
1-65. Stratigraphic relations between the Columbia River basalt and the Latah Formation in Spokane area, Washington [abs.]: Geol. Soc. America Spec. Paper 82, p. 328, 1965.
- GRIGGS, R. L.
1-64. Geology and ground-water resources of the Los Alamos area, New Mexico: U.S. Geol. Survey Water-Supply Paper 1753, 104 p., 1964.
- GRISCOM, Andrew
1-64. Magnetic properties and aeromagnetic interpretation of serpentinite from southwest Puerto Rico, *in* A study of serpentinite: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 39-48, 1964.
- GROZIER, R. U.
1-65. Growth of salt cedar (*Tamarix gallica*) in the Pecos River near the New Mexico-Texas boundary, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B175-B176, 1965.
2-65. (BLAKEY, J. F., and HEMBREE, C. H.) Water delivery study: Pecos River, Texas, quantity and quality, 1964: U.S. Geol. Survey open-file report, 27 p., 1965.
- GUILLOU, R. B.
1-64. The aerial radiological measuring surveys (ARMS) program, *in* The natural radiation environment: Chicago, Illinois, Univ. Chicago Press (for Rice University), p. 705-721, 1964.
- GUY, H. P.
1-63. (JACKSON, N. E., JARVIS, Kenneth, JOHNSON, C. J., MILLER, C. R., and STEINER, W. W.) A program for sediment control in the Washington Metropolitan region: Interstate Comm. Potomac River Basin Tech. Bull. 1963-I, 48 p., 1963.
1-64. A system for monitoring fluvial sediment [abs.]: Am. Geophys. Union Trans., v. 45, no. 4, p. 612, 1964.
1-65. (SIMONS, D. B., and RICHARDSON, E. V.) Summary of alluvial channel data from flume experiments, 1956-1961: U.S. Geol. Survey open-file report, 383 p., 1965.
- HACK, J. T.
1-65. Geomorphology of the Shenandoah Valley, Virginia and West Virginia, and origin of the residual ore deposits: U.S. Geol. Survey Prof. Paper 484, 84 p., 1965.

HACK, J. T.--Continued

- 2-65. Postglacial drainage evolution and stream geometry in the Ontonagon area, Michigan: U.S. Geol. Survey Prof. 504-B, p. B1-B40, 1965.

HACKETT, O. M.

- 1-64. The father of modern ground water hydrology: Ground Water, v. 2, no. 2, p. 2-5, 1964.
2-64. Ground-water levels in the United States, 1957-61, north-central states: U.S. Geol. Survey Water-Supply Paper 1781, p. 1-148, 1964.
1-65. Ground-water levels in the United States, 1958-62, northeastern states: U.S. Geol. Survey Water-Supply Paper 1782, 252 p., 1965.

HACKMAN, R. J.

- 1-63. A lunar isotonal map, in Astrogeologic studies, Annual Progress Report, August 25, 1962-July 1, 1963—Pt. D, Studies for spaceflight program: U.S. Geol. Survey open-file report, p. 13-28, 1963.
2-63. Stratigraphy and structure of the Montes Apenninus quadrangle of the Moon, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 1-8, 1963.

HAHL, D. C.

- 1-64. (and LANGFORD, R. H.) Dissolved-mineral inflow to Great Salt Lake and chemical characteristics of the Salt Lake brine—Pt. 2, Technical report: Utah Geol. and Mineralog. Survey Water Resources Bull. 3, pt. 2, 40 p., 1964.

HAIL, W. J., Jr.

- 1-64. Geology of the Summer Shade quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-308, 1964.
1-65. Geology of northwestern North Park, Colorado: U.S. Geol. Survey Bull. 1188, 133 p., 1965.

HALBERG, H. N.

- 1-64. (and REED, J. E.) Ground-water resources of eastern Arkansas in the vicinity of U.S. Highway 70: U.S. Geol. Survey Water-Supply Paper 1779-V, p. V1-V38, 1964.

HALE, W. E.

- 1-63. (WINOGRAD, I. J., and GARBER, M. S.) Preliminary appraisal of close-in aquifer response to the Bilby event, Yucca Flat, Nevada: U.S. Geol. Survey Tech. Letter NTS-63, 32 p., 1963.

HALL, W. E.

- 1-65. U.S. Geological Survey participation in the Upper Mantle project: Progress report [abs.]: Geol. Soc. America Spec. Paper 82, p. 82, 1965.

HALLIDAY, James

- 1-64. Photogrammetric contouring of areas covered by evergreen forests, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C190-C194, 1964.

HAMILTON, Warren

- 1-64. A tectonic map of Antarctica—a progress report, in Antarctic geology; Proc. First International Symposium on Antarctic Geology, Cape Town, Sept. 1963: New York, John Wiley and Sons, p. 676-680, 1964[1965].
2-64. Diabase sheets differentiated by liquid fractionation, Taylor Glacier region, South Victoria land, in Antarctic geology; Proc. First International Symposium on Antarctic Geology, Cape Town, Sept. 1963: New York, John Wiley and Sons, p. 442-454, 1964[1965].
3-64. Geologic map of the Big Maria Mountains NE quadrangle, Riverside County, California, and Yuma County, Arizona: U.S. Geol. Survey Geol. Quad. Map GQ-350, 1964.
4-64. Origin of high-alumina basalt, andesite, and dacite magmas: Science, v. 146, no. 3644, p. 635-637, 1964.
1-65. Diabase sheets of the Taylor Glacier region, Victoria Land, Antarctica, with sections on Petrography, by

HAMILTON, Warren--Continued

- Warren Hamilton, P. T. Hayes, and Ronald Calvert, and sections on Chemistry, by Warren Hamilton, V. C. Smith, P. S. D. Elmore, P. R. Barnett, and Nancy Conklin: U.S. Geol. Survey Prof. Paper 456-B, p. B1-B71, 1965.
2-65. The Permian record and continental drift [abs.]: Soc. Econ. Paleontologists and Mineralogists, Permian Basin Section, 10th Ann. Mtg., 1965, p. 17-19, 1965.
3-65. (and MOUNTJOY, Wayne) Alkali content of alpine ultramafic rocks: Geochim. Cosmochim. Acta, v. 29, no. 6, p. 661-671, 1965.
4-65. (and PAKISER, L. C.) Geologic and crustal cross section of the United States along the 37th parallel, a contribution to the upper mantle project: U.S. Geol. Survey Misc. Geol. Inv. Map I-448, 1965.

HAMPTON, E. R.

- 1-64. Geologic factors that control the occurrence and availability of groundwater in the Fort Rock Basin Lake County, Oregon: U.S. Geol. Survey Prof. Paper 383-B, B1-B29, 1964.

HANLY, T. F.

- 1-64. Quality of water investigations in Wyoming, 1963-64, in 37th Bienn. Rept. of the State Engineer to the Governor of Wyoming: Wyoming State Engineer, p. 100-106, 1964.

HANSEN, A. J., Jr.

- 1-64. (and SCHINER, G. R.) Ground-water resources of Block Island, Rhode Island: Rhode Island Water Resources Coordinating Board Geol. Bull. 14, 35 p., 1964.

HANSEN, W. R.

- 1-64. Nonmetallic and industrial minerals and materials resources: Stone, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 222-228, 1964.
1-65. Geology of the Flaming Gorge area, Utah-Colorado-Wyoming: U.S. Geol. Survey Prof. Paper 490, 196 p., 1965.
2-65. Geology of the Flaming Gorge quadrangle, Utah-Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-75, 1965. (Originally published 1955.)

HANSHAW, B. B.

- 1-64. Cation-exchange constants for clays from electrochemical measurements, in Clays and clay minerals, 12th Natl. Conf., Atlanta, Ga., 1963, Proc.: New York, Macmillan Co., p. 397-421, 1964.
1-65. (BACK, William, MEYER, Rubin, and WAIT, R. L.) Relation of carbon-14 concentration to saline water contamination of coastal aquifers: Water Resources Research, v. 1, no. 1, p. 109-114, 1965.

HARBECK, G. E., Jr.

- 1-64. Estimating forced evaporation from cooling ponds: Am. Soc. Civil Engineers Proc. Paper 4061, Power Div. Jour., v. 90, no. P03, p. 1-9, 1964.
2-64. (MEYERS, J. S., and HUGHES, G. H.) The effect of an increased heat load on the thermal structure and evaporation of Lake Colorado City, Texas: U.S. Geol. Survey open-file report, 41 p., 1964.

HARDER, A. H.

- 1-65. (and HOLDEN, W. R.) Measurement of gas in groundwater: Water Resources Research, v. 1, no. 1, p. 75-82, 1965.
2-65. (WHITMAN, H. M., and ROGERS, S. M.) Methane in the fresh-water aquifers of southwestern Louisiana and theoretical explosion hazards: Louisiana Geol. Survey Water Resources Pamph. 14, 22 p., 1965.

HARDT, W. F.

- 1-64. (CATTANY, R. E., and KISTER, L. R.) Basic data for western Pinal County, Arizona: Arizona Land Dept. Water Resources Rept. 18, 59 p., 1964.

- HARRIS, D. D.
1-64. (and RICHARDSON, E. V.) Stream gaging control structure for the Rio Grande conveyance channel near Bernardo, New Mexico: U.S. Geol. Survey Water-Supply Paper 1369-E, p. E123-E154, 1964.
- HARRIS, E. E.
1-64. (and RANTZ, S. E.) Effect of urban growth on streamflow regimen of Permanente Creek, Santa Clara County, California: U.S. Geol. Survey Water-Supply Paper 1591-B, p. B1-B18, 1964.
- HARRIS, L. D.
1-64. Geology of the Sulphur Lick quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-323, 1964.
1-65. The Clinchport thrust fault—a major structural element of the southern Appalachian Mountains, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B49-B53, 1965.
2-65. (and MILLER, R. L.) Geology of the Duffield quadrangle, Virginia: U.S. Geol. Survey Geol. Quad. Map GQ-111, 1965. (Originally published 1958.)
- HARRIS, W. H.
1-64. (WILDER, H. B.) Ground-water supply of Cape Hatteras National Seashore Recreational Area, North Carolina—Pt. 3: North Carolina Dept. Water Resources Rept. Inv. 4, 22 p., 1964.
- HARRISON, J. E.
1-65. (and JOBIN, D. A.) Geologic map of the Packsaddle Mountain quadrangle, Idaho: U.S. Geol. Survey Geol. Quad. Map GQ-375, with 4-page text, 1965.
- HART, D. H.
1-65. (and NEWCOMB, R. C.) Geology and ground water of the Tualatin Valley, Oregon: U.S. Geol. Survey Water-Supply Paper 1697, 172 p., 1965.
- HART, D. L., Jr.
1-65. Groundwater in the alluvial deposits of the Washita River between Clinton and Anadarko, Oklahoma: Oklahoma Water Resources Board Bull. 26, 25 p., 1965.
- HARTWELL, J. H.
1-64. (KLEIN, Howard, and JOYNER, B. F.) Preliminary evaluation of hydrologic situation in Everglades National Park, Florida: U.S. Geol. Survey open-file report, 25 p., 1964.
- HARVEY, E. J.
1-65. (GOLDEN, H. G., and JEFFERY, H. G.) Water resources of the Pascagoula area, Mississippi: U.S. Geol. Survey Water-Supply Paper 1763, 135 p., 1965.
- HASLER, J. W.
1-63. Potential sites for the Buggy event, Bruneau River area, Owyhee County, Idaho: U.S. Geol. Survey Tech. Letter, BUGGY-1, 9 p., 1963.
2-63. (SNYDER, R. P., and BOWERS, W. E.) Preliminary outline of the geology of the U12e.07a tunnel, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Area-12-7, 8 p., 1963.
1-64. Geologic appraisal of the U20c and U20d sites, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-12, 10 p., 1964.
- HAUSHILD, W. L.
1-64. (and DEMPSTER, G. R.) Discussion of "Annual variation of stream water temperature", by J. C. Ward: *Am. Soc. Civil Engineers Proc., Sanitary Eng. Div. Jour.*, v. 90, no. SA4, p. 90-92, 1964.
- HAVENS, J. S.
1-64. Recharge studies on the High Plains in northern Lea County, New Mexico: U.S. Geol. Survey open-file report, 82 p., 1964.
- HAYES, P. T.
1-64. Geology of the Guadalupe Mountains, New Mexico: U.S. Geol. Survey Prof. Paper 446, 69 p., 1964.
- HAYES, P. T. --Continued
2-64. (and LANDIS, E. R.) Geologic map of the southern part of the Mule Mountains, Cochise County, Arizona: U.S. Geol. Survey Misc. Geologic Inv. Map I-418, 1964.
1-65. Nitrates and guano, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 372-374, 1965.
2-65. (and LANDIS, E. R.) Paleozoic stratigraphy of the southern part of the Mule Mountains, Arizona: U.S. Geol. Survey Bull. 1201-F, p. F1-F43, 1965.
3-65. (SIMONS, F. S., and RAUP, R. B.) Lower Mesozoic extrusive rocks in southeastern Arizona—the Canelo Hills Volcanics: U.S. Geol. Survey Bull. 1194-M, p. M1-M9, 1965.
- HAYNES, D. D.
1-64. Geology of the Mammoth Cave quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-351, 1964.
2-64. Reconnaissance for uranium in the Central Tucano Basin, Bahia, Brazil: U.S. Geol. Survey Bull. 1185-B, p. B1-B16, 1964.
3-64. Geology of the Glasgow North quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-339, 1964.
1-65. Geology of the Hiseville quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-401, 1965.
- HAYS, W. H.
1-64. Geology of the Grand Rivers quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-328, 1964.
- HAZEL, J. E.
1-65. Ostracoda: Summary of investigations conducted in 1964: Woods Hole Oceanographic Inst. Ref. No. 65-13, p. 73, 1965.
- HAZELWOOD, R. M.
1-63. Preliminary estimate of electrical properties of alluvium in Area 2, Nevada Test Site: U.S. Geol. Survey Tech. Letter, YUCCA-40, 4 p., 1963.
2-63. (HEALEY, D. L., and MILLER, C. H.) U.S. Geological Survey investigations of Yucca Flat, Nevada Test Site, Pt. B, Geophysical investigations: U.S. Geol. Survey Tech. Letter NTS-45, 53 p., 1963.
- HEALEY, D. L.
1-63. Correlation of gravity data and faulting in Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-43, 11 p., 1963.
2-63. (CURREY, F. E., and SMITH, R. C.) Interpretation of gravity data in the vicinity of ground water test well 8: U.S. Geol. Survey Tech. Letter NTS-55, 14 p., 1963.
3-63. (and MILLER, C. H.) Gravity survey of the Gold Meadows stock, Nevada Test Site, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-40, 7 p., 1963.
1-65. Principal facts and Bouguer anomaly values, Crooks Gap gravity survey, Fremont County, Wyoming: U.S. Geol. Survey open-file report, 1 p. text, 10 p. data sheets, 1965.
- HEALY, H. G.
1-64. Two-variable linear correlation analyses of water-level fluctuations in artesian wells in Florida, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D199-D202, 1964.
- HEALY, J. H.
1-65. (and PLOGER, S. L.) P_n arrivals recorded from nuclear explosions [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 156, 1965.
- HEARN, B. C., Jr.
1-64. (PECORA, W. T., and SWADLEY, W. C.) Geology of the Rattlesnake quadrangle, Bearpaw Mountains, Blaine County, Montana: U.S. Geol. Survey Bull. 1181-B, p. B1-B66, 1964[1965].

HEATH, J. A.

- 1-65. Bibliography of reports resulting from U.S. Geological Survey participation in the United States technical assistance program, 1940-65: U.S. Geol. Survey Bull. 1193, 51 p., 1965.

HEDGE, C. E.

- 1-64. Significance of radiogenic strontium in volcanic lavas [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 114, 1964.

HEIDEL, S. G.

- 1-65. Dissolved oxygen and iron in shallow wells at Salisbury, Maryland: Am. Water Works Assoc. Jour., v. 57, no. 2, p. 239-244, 1965.

HEINDL, L. A.

- 1-64. Cenozoic geology of Arizona—A 1960 resume: Arizona Geol. Soc. Digest, v. 5, p. 73-88, 1964.
 2-64. Should the term "Gila Conglomerate" be abandoned?: Arizona Geol. Soc. Digest, v. V, p. 73-88, 1964.
 3-64. (and McCLYMONDS, N. E.) Younger Precambrian Formations and the Bolsa(?) Quartzite of Cambrian age, Papago Indian Reservation, Arizona, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C43-C49, 1964.
 1-65. Mesozoic formations in the Vekol Mountains, Papago Indian Reservation, Arizona: U.S. Geol. Survey Bull. 1194-G, p. G1-G9, 1965.
 2-65. (and WHITE, N. D.) Hydrologic and drill-hole data, San Xavier Indian Reservation and vicinity, Pima County, Arizona: Arizona State Land Dept. Water-Resources Rept. 20, 48 p., 1965; also in U.S. Geol. Survey open-file report, 38 p., 1965.

HELY, A. G.

- 1-64. Areal variations of mean annual runoff: Am. Soc. Civil Engineers Proc. Paper 4030, Hydraulics Div. Jour., v. 90, no. HY5, p. 61-68, 1964.
 2-64. (and PECK, E. L.) Precipitation, runoff and water loss in the lower Colorado River—Salton Sea area: U.S. Geol. Survey Prof. Paper 486-B, p. B1-B16, 1964.

HELZ, A. W.

- 1-65. The problem of automatic plate reading and computer interpretation for spectrochemical analysis, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B160-B164, 1965.

HEMBREE, C. H.

- 1-64. (and BLAKEY, J. F.) Chemical quality of surface waters in the Hubbard Creek watershed, Texas—Progress report, September 1963: Texas Water Comm. Bull. 6411, 45 p., 1964.
 2-64. (KRIEGER, R. A., and JORDAN, P. R.) Chemical quality of surface waters, and sedimentation in the Grand River drainage basin, North and South Dakota: U.S. Geol. Survey Water-Supply Paper 1769, 78 p., 1964.

HENBEST, L. G.

- 1-65. Diagenetic phenomena in oolitic limestone, Marrow Series, Pennsylvanian, northwest Arkansas and northeast Oklahoma [abs.]: Geol. Soc. America Spec. Paper 82, p. 91, 1965.

HENDERSON, J. R.

- 1-64. (and SMITH, C. W.) Aeromagnetic map of the Old Speck Mountain quadrangle, Franklin and Oxford Counties, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-486, 1964.
 1-65. (and VARGO, J. L.) Aeromagnetic map of central Iowa: U.S. Geol. Survey Geophys. Inv. Map GP-476, 1965.

HENDRICKS, E. L.

- 1-64. Compilation of records of surface waters of Hawaii, July 1950 to June 1960: U.S. Geol. Survey Water-Supply Paper 1739, 136 p., 1964.

HENDRICKS, E. L. --Continued

- 2-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 1-A, North Atlantic slope basins, Maine to Connecticut: U.S. Geol. Survey Water-Supply Paper 1721, 317 p., 1964.
 3-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 1-B, North Atlantic slope basins, New York to York River: U.S. Geol. Survey Water-Supply Paper 1722, 578 p., 1964.
 4-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 2-A, South Atlantic slope basins, James River to Savannah River: U.S. Geol. Survey Water-Supply Paper 1723, 303 p., 1964.
 5-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 4, St. Lawrence River basin: U.S. Geol. Survey Water-Supply Paper 1727, 379 p., 1964.
 6-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 6-A, Missouri River basin above Sioux City, Iowa: U.S. Geol. Survey Water-Supply Paper 1729, 507 p., 1964.
 7-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. II, Pacific slope basins in California: U.S. Geol. Survey Water-Supply Paper 1735, 715 p., 1964.
 8-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 12, Pacific slope basins in Washington and upper Columbia River basin: U.S. Geol. Survey Water-Supply Paper 1736, 415 p., 1964.
 9-64. Compilation of records of surface waters of Alaska, October 1950 to September 1960: U.S. Geol. Survey Water-Supply Paper 1740, 86 p., 1964.

HENDRICKS, T. A.

- 1-63. Petroleum resources of the United States [abs.]: Tulsa Geol. Soc. Digest, v. 31, p. 257, 1963.
 1-65. Section of beds overlying the Springer(?) Formation in Ti Valley, Pittsburg County, Oklahoma: Oklahoma Geology Notes, v. 25, no. 1, p. 6-10, 1965.

HENDRICKSON, G. E.

- 1-64. (and KRIEGER, R. A.) Geochemistry of natural waters of the Blue Grass region, Kentucky: U.S. Geol. Survey Water-Supply Paper 1700, 135 p., 1964.
 1-65. Water management and recreation on the upper Au Sable River Michigan: U.S. Geol. Survey open-file report, 103 p., 1965.

HERNON, R. M.

- 1-64. Geologic maps and sections of the Ducktown, Isabella, and Persimmon Creek quadrangles, Tennessee and North Carolina: U.S. Geol. Survey open-file report, 3 sheets, 1964.
 2-64. (JONES, W. R., and MOORE, S. L.) Geology of the Santa Rita quadrangle, New Mexico: U.S. Geol. Survey Quad. Map GQ-306, 1964.

HERRICK, E. H.

- 1-65. (and DAVIS, L. V.) Availability of ground water in Tularosa Basin and adjoining areas, New Mexico and Texas: U.S. Geol. Survey Hydrol. Inv. Map HA-191, 1965.

HERRICK, S. M.

- 1-64. Offshore extension of the upper Eocene to Recent stratigraphic sequence in southeastern Georgia, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C61-C63, 1964.
 2-64. Upper Eocene smaller foraminifera from Shell Bluff and Griffin Landings, Burke County, Georgia, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C64-C65, 1964.

HERRICK, S. M. --Continued

- 1-65. A subsurface study of Pleistocene deposits in coastal Georgia: Georgia Geol. Survey Inf. Circ. 31, 8 p., 1965.
 - 2-65. (and WAIT, R. L.) Subsurface stratigraphy of coastal Georgia and South Carolina [abs.]: Am. Assoc. Petroleum Geologists Ann. Mtg., 50th, New Orleans, April 1965, Program, p. 68, 1965.
- HERZ, Norman
- 1-64. (and DUTRA, C. V.) Cobalt, nickel, chromium, scandium and niobium in biotite and the scandium geological thermometer: Soc. Brasileira Geologia Bol., v. 13, nos. 1-2, p. 23-42, 1964[1965].
- HEWETT, D. F.
- 1-64. Veins of hypogene manganese oxide minerals in the southwestern United States: Econ. Geology, v. 59, no. 8, p. 1429-1472, 1964.
 - 1-65. (RADTKE, Arthur, and TAYLOR, Charles) Black calcite—a new source of silver: Mining Cong. Jour., v. 51, no. 6, p. 78, 1965.
- HEYL, A. V.
- 1-64. Oxidized zinc deposits of the United States—Pt. 3, Colorado: U.S. Geol. Survey Bull. 1135-C, p. C1-C98, 1964.
 - 2-64. (HOSTERMAN, J. W., and BROCK, M. R.) Clay-mineral alteration in the Upper Mississippi Valley zinc-lead district, in Clays and clay minerals, 12th Nat. Conf., Atlanta, Ga., 1963, Proc.: New York, Macmillan Co., p. 445-453, 1964.
- HICKLING, N. L.
- 1-65. Allanites from the Boulder Creek Batholith, Colorado: U.S. Geol. Survey open-file report, 48 p., 1965.
- HIETANEN, Anna
- 1-65. Two episodes of metamorphism in the Belt Series northwest of the Idaho batholith [abs.]: Geol. Soc. America Spec. Paper 82, p. 332, 1965.
- HILL, D. P.
- 1-65. The U.S. Geological Survey's gravity program in Washington, Idaho, Montana, and Wyoming: Am. Geophys. Union Trans., v. 46, no. 1, p. 214-217, 1965.
- HILPERT, L. S.
- 1-64. Introduction, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 7-8, 1964.
 - 2-64. Geology: Introduction, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 17, 1964.
 - 3-64. Metallic mineral resources: Introduction, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 71, 1964.
 - 4-64. Metallic mineral resources: Mercury, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 108-111, 1964.
 - 5-64. Mineral fuels and associated resources: Introduction, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 39, 1964.
 - 6-64. Nonmetallic and industrial minerals and minerals resources: Introduction, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 151, 1964.
 - 7-64. (and DASCH, M. D.) Metallic mineral resources: Uranium, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 124-132, 1964.
 - 8-64. (and ROBERTS, R. J.) Geology: Economic geology, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 28-34, 1964.
 - 1-65. Uranium, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 209-226, 1965.
- HINDS, J. S.
- 1-64. Btu values of Fruitland Formation coal deposits in Colorado and New Mexico, as determined from rotary-drill cuttings, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D90-D94, 1964.
- HINES, M. S.
- 1-65. Water-supply characteristics of selected Arkansas streams: U.S. Geol. Survey open-file report, 73 p., 1965.
- HINKLE, Margaret
- 1-65. (LEONG, Kam Wo, and WARD, F. N.) Field determination of nanogram quantities of mercury in soils and rocks: U.S. Geol. Survey open-file report, 14 p., 1965.
- HINRICHS, E. N.
- 1-63. Geology of the Ammonia Tanks quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-59, 12 p., 1963.
 - 1-64. (KRUSHENSKY, R. D., LUFT, S. J., BROWN, R. D., and SIMONS, F. S.) Geologic map and sections of the Ammonia Tanks quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report, 1964.
 - 1-65. (and McKAY, E. J.) Geologic map of the Plutonium Valley quadrangle, Nye and Lincoln Counties, Nevada: U.S. Geol. Survey Geol. Quad. Map GQ-384, 1965.
- HITE, R. J.
- 1-64. Nonmetallic and industrial minerals and materials resources: Salines, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 206-215, 1964.
- HOARE, J. M.
- 1-64. (CONDON, W. H., and PATTON, W. W., Jr.) Occurrence and origin of laumontite in Cretaceous sedimentary rocks in western Alaska, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C74-C78, 1964.
- HOBBS, H. H., Jr.
- 1-64. (and BEDINGER, M. S.) A new troglitic crayfish of the genus *Cambarus* (Decapoda, Astacidae) from Arkansas with a note on the range of *Cambarus cryptodytes* Hobbs: Biol. Soc. Washington Proc. 1964, v. 77, art. 3, p. 9-15, 1964.
- HOBBS, S. W.
- 1-64. Ferrous metals: Tungsten, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 108-115, 1964.
 - 2-64. Metallic mineral resources: Tungsten, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 71-74, 1964.
 - 3-64. Mineral resources: Tungsten, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 223-233, 1964.
 - 1-65. Tungsten, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 241-255, 1965.
- HODGES, E. B.
- 1-64. Surface-water runoff, storage, and diversions, in Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 10-12, 1964.
- HOFFARD, S. H.
- 1-64. (and VAUDREY, W. C.) An investigation of floods in Hawaii through June 30, 1964: Hawaii State Div. Water and Land Devel. Prog. Rept. 7, 165 p., 1964.
- HOFMANN, Walter
- 1-64. (and RANTZ, S. E.) Arid lands hydrology: Duquesne Sci. Counselor, v. 27, no. 2, p. 34-40, 1964.
- HOGENSON, G. M.
- 1-65. (and FOXWORTHY, B. L.) Ground water in the East Portland area, Oregon: U.S. Geol. Survey Water-Supply Paper 1793, 78 p., 1965.

HOLLAND, P. H.

- 1-65. Base-flow studies, Guadalupe River, Comal County, Texas—Quantity, March 1962: Texas Water Comm. Bull. 6503, 6 p., 1965.
- 2-65. (and MENDIETA, H. B.) Base-flow studies, Llano River, Texas—Quantity and quality: Texas Water Comm. Bull. 6505, 20 p., 1965.
- 3-65. (and WELBORN, C. T.) Base-flow studies, Cibolo Creek, Texas—Quantity and quality, March 5-7, 1963: Texas Water Comm. Bull. 6511, 13 p., 1965.

HOLLOWELL, J. R.

- 1-65. Ground water in the alluvium of Elk Creek Basin, Oklahoma: Oklahoma Water Resources Board Bull. 28, 12 p., 1965.

HOLMES, G. W.

- 1-64. A Quaternary chronology of Ullsfjord, northern Norway [abs.], in *Internat. Geog. Cong.*, 20th, United Kingdom, 1964, Abs. Papers: London, Thomas Nelson and Sons Ltd., p. 94, 1964.
- 1-65. Geologic reconnaissance along the Alaska Highway, Delta River to Tok Junction, Alaska: U.S. Geol. Survey Bull., 1181-H, p. H1-H19, 1965.
- 2-65. Preliminary materials map, Otis quadrangle, Massachusetts: U.S. Geol. Survey open-file report, map, 6 data sheets, 1965.
- 3-65. Preliminary materials map, Pittsfield West quadrangle, Massachusetts-New York: U.S. Geol. Survey open-file report, 1965.
- 4-65. Preliminary materials map, Massachusetts portion of the Canaan quadrangle, New York-Massachusetts: U.S. Geol. Survey open-file report, 1965.
- 5-65. Preliminary materials map, Massachusetts portion Berlin quadrangle, New York-Massachusetts-Vermont: U.S. Geol. Survey open-file report, 1965.
- 6-65. Preliminary materials map, Windsor quadrangle, Massachusetts: U.S. Geol. Survey open-file report, 1965.
- 7-65. (and PEWE, T. L.) Geologic map of the Mount Hayes D-3 quadrangle, Alaska: U.S. Geol. Survey Geol. Quad. Map GQ-366, 1965.

HOLT, C. L. R., Jr.

- 1-64. (YOUNG, K. B., and CARTWRIGHT, W. H.) The water resources of Wisconsin: Univ. Wisconsin, Geol. and Nat. History Survey, 1964 Blue Book, p. 178-198, 1964.

HOOD, J. W.

- 1-65. (and HERRICK, E. H.) Water resources of the Three Rivers area, Otero and Lincoln Counties, New Mexico: U.S. Geol. Survey Hydrol. Inv. Atlas HA-192, 1965.

HOOKER, Marjorie

- 1-65. Origin of the term nuée ardente [abs.]: *Am. Mineralogist*, v. 50, nos. 1-2, p. 286-287, 1965.

HOOVER, D. L.

- 1-63. Geology and lithologic log of Pahute Mesa Exploratory Hole No. 1: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-6, 61 p., 1963.
- 2-63. Geologic appraisal of the U19a site, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-9, 12 p., 1963.
- 3-63. Geologic appraisal of the U20b site, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-8, 12 p., 1963.
- 4-63. (and MEYER, G. L.) Abridged lithologic log of Pahute Mesa Exploratory Hole, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-2, 7 p., 1963.
- 5-63. (MEYER, G. L., and SNYDER, R. P.) Abridged lithologic log of Pahute Mesa Exploratory Hole, 3,800 to 6,541 feet, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-2, supp. 1, 6 p., 1963.

HOOVER, D. L. --Continued

- 1-64. Geologic appraisal of the U20e site, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-II, 10 p., 1964.

- 1-65. (and SHEPARD, A. O.) Zeolite zoning in tuffs, Nevada Test Site, Nye County, Nevada [abs.]: *Am. Mineralogist*, v. 50, nos. 1-2, p. 287, 1965.

HOPKINS, D. M.

- 1-65. (MacNEIL, F. S., MERKLIN, R. L., and PETROV, O. M.) Quaternary correlations across Bering Strait: *Science*, v. 147, no. 3662, p. 1107-1114, 1965.

HORR, C. A.

- 1-64. Chemical quality of the ground water, in *Geology and ground-water resources in eastern Cheyenne and Kiowa Counties, Colorado*: U.S. Geol. Survey Water-Supply Paper 1779-N, p. N17-N30, 1964.

HORTON, G. W.

- 1-64. Penetration chart of the oil and gas fields of the Uinta Basin and the Uncompahgre Uplift, Utah, in *Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse*, 13th annual field conference, 1964: Salt Lake City, Utah, Intermountain Association of Petroleum Geologists, p. 186, 1964.
- 2-64. (and GERE, W. C.) Relation of stratigraphy and structure to gas production in the San Arroyo and East Canyon fields, Utah, in *Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse*, 13th annual field conference, 1964: Salt Lake City, Utah, Intermountain Association of Petroleum Geologists, p. 191-195, 1964.

HOSMAN, R. L.

- 1-64. The Carrico Sand, a potential aquifer in south-central Arkansas, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D158-D160, 1964 [1965].

HOSTERMAN, J. W.

- 1-64. (HEYL, A. V., and JOLLY, J. L.) Qualitative X-ray emission analysis studies of enrichment of common elements in wallrock alteration in the Upper Mississippi Valley zinc-lead district, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D54-D60, 1964.

- 2-64. (and PRATER, L. S.) Mineral resources: Clays, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 51-57, 1964.

HOSTETLER, P. B.

- 1-65. (COLEMAN, R. G., MUMPTON, F. A., and EVANS, B. W.) Brucite in Alpine serpentinites [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 183-184, 1965.

HOTZ, P. E.

- 1-64. Nickeliferous laterites in southwestern Oregon and northwestern California: *Econ. Geology*, v. 59, no. 3, p. 355-396, 1964.

HOUSER, F. N.

- 1-63. (and WILLIAMS, W. P.) Structure of U3ams sink, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-45, 12 p., 1963.

- 1-64. (and EMERICK, W. L.) Preliminary structural analysis of explosion-produced fractures, HARDHAT event, Area 15, Nevada Test Site, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C100-C102, 1964.

HOYTE, A.

- 1-65. (SENFLE, F., and WIRTZ, P.) Electrical resistivity and viscosity of tektite glass: *Jour. Geophys. Research*, v. 70, no. 8, p. 1985-1994, 1965.

HUBBELL, D. W.

- 1-64. Apparatus and techniques for measuring bedload: U.S. Geol. Survey Water-Supply Paper 1748, 74 p., 1964.

- HUBBELL, D. W.--Continued
 2-64. (and SAYRE, W. W.) Sand transport studies with radioactive tracers: *Am. Soc. Civil Engineers Proc.*, v. 90, Jour. Hydraulics Div., no. HY3, pt. 1, p. 39-68, 1964.
- HUBBERT, M. K.
 1-65. *Geology*, Chap. 6, *in* Listen to leaders in science: Atlanta, Ga., Tupper and Love, p. 73-83, 1965.
- HUBER, N. K.
 1-65. (and RINEHART, C. D.) The Devils Postpile National Monument: California Div. Mines and Geology Mineral Inf. Service, v. 18, no. 6, p. 109-118, 1965.
- HUFF, L. C.
 1-63. Contributed to the section on 'Geochemical methods', *in* Aridity and man: *Am. Assoc. Adv. Sci. Pub.* 74, p. 194-197, 1963.
- HUFFMAN, Claude, Jr.
 1-64. (and BARTEL, A. J.) Ion-exchange separation of tin from silicate rocks, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D131-D133, 1964.
- HUGHES, L. S.
 1-65. (and BLAKEY, J. F.) Chemical composition of Texas surface waters, 1962: Texas Water Comm. Bull. 6501, 167 p., 1965.
- HUNN, J. D.
 1-64. (and ROSENSHEIN, J. S.) Ground-water hydrology, development, and potential of rock units of Quaternary age, St. Joseph County, Indiana: U.S. Geol. Survey open-file report, 36 p., 1964.
- HUNT, O. P.
 1-65. Low-flow summary at stream-gaging stations on Long Island, New York: U.S. Geol. Survey open-file report, 20 p., 1965.
 2-65. Surface water resources in the vicinity of North Lake public campsite near Haines Falls, New York: U.S. Geol. Survey open-file report, 6 p., 1965.
- HURLEY, P. M.
 1-65. (BATEMAN, P. C., FAIRBAIRN, H. W., and PINSON, W. H., Jr.) Investigation of initial $\text{Sr}^{87}/\text{Sr}^{86}$ ratios in the Sierra Nevada plutonic province: *Geol. Soc. America Bull.*, v. 76, no. 2, p. 165-174, 1965.
- HYDEN, H. J.
 1-65. (McANDREWS, Harry, and TSCHUDY, R. H.) The Foote Creek and Dutton Creek Formations, two new formations in the north part of the Laramie basin, Wyoming: U.S. Geol. Survey Bull. 1194-K, p. K1-K12, 1965.
- HYLAND, J. B.
 1-64. (and KEECH, C. F.) Ground water in Cedar Rapids Division of lower Platte River basin, Nebraska, *with a section on* Chemical quality of water, by P. G. Rosene: U.S. Geol. Survey Water-Supply Paper 1779-H, p. H1-H12, 1964.
- IMLAY, R. W.
 1-64. Marine Jurassic pelecypods from central and southern Utah: U.S. Geol. Survey Prof. Paper 483-C, p. C1-C42, 1964.
 2-64. Middle Bajocian ammonites from the Cook Inlet region, Alaska: U.S. Geol. Survey Prof. Paper 418-B, p. B1-B61, 1964.
 3-64. Upper Jurassic mollusks from eastern Oregon and western Idaho: U.S. Geol. Survey Prof. Paper 483-D, p. D1-D21, 1964.
 1-65. Review of "Studies zur Stammesgeschichte der Ammoniten," by O. H. Schindewolf: *Jour. Paleontology*, v. 39, no. 3, p. 513-515, 1965.
- INTERIOR MISSOURI BASIN FIELD COMMITTEE
 1-64. Missouri River Basin progress report, October-December: Billings, Montana, Interior Missouri Basin Field Comm., 78 p., 1964.
- INTERIOR MISSOURI BASIN FIELD COMM.--Continued
 1-65. Missouri River Basin progress report, January-March: Billings, Montana, Interior Missouri Basin Field Comm., 58 p., 1965.
- IORNS, W. V.
 1-64. (HEMBREE, C. H., PHOENIX, D. A., and OAKLAND, G. L.) Water resources of the Upper Colorado River Basin--Basic data: U.S. Geol. Survey Prof. Paper 442, 1036 p., 1964.
- IRELAN, Burdge
 1-64. (and MENDIETA, H. B.) Chemical quality of surface waters in the Brazos River in Texas: U.S. Geol. Survey Water-Supply Paper 1779-K, p. K1-K70, 1964.
- IRWIN, J. H.
 1-65. *Geology and availability of ground water on the Ute Mountain Indian Reservation, Colorado and New Mexico*: U.S. Geol. Survey open-file report, 185 p., 1965.
- IRWIN, W. P.
 1-64. Late Mesozoic orogenies in the ultramafic belts of northwestern California and southwestern Oregon, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C1-C9, 1964.
- IZETT, G. A.
 1-64. (and BARCLAY, C. S. V.) Preliminary geologic map of the Hot Sulphur Springs SW quadrangle, Grand County, Colorado: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-291, 1964.
 1-65. Preliminary geologic map of the northwest quarter of the Hot Sulphur Springs quadrangle, Grand County, Colorado: U.S. Geol. Survey open-file report, 1965.
- JACKSON, E. D.
 1-64. Primary features of stratiform chromite deposits, *in* Methods of prospecting for chromite: Organization for Econ. Cooperation and Development, Paris, p. 111-134, 1964.
- JEFFERSON, P. O.
 1-65. Performance of channel changes: Alabama Highway Research HPR Rept. No. 9 (Interim Report), 25 p., 1965.
- JENNE, E. A.
 1-63. Chemical characterization of selected core samples from the vicinity of Tatum salt dome, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-35, 13 p., 1963.
- JESPERSEN, Anna
 1-64. Aeromagnetic interpretation of the Globe-Miami Copper district, Gila and Pinal Counties, Arizona, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D70-D75, 1964.
- JOBIN, D. A.
 1-64. (and SCHROEDER, M. L.) *Geology of the Conant Valley quadrangle, Bonneville County, Idaho*: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-277, 1964.
 2-64. (and SCHROEDER, M. L.) *Geology of the Irwin quadrangle, Bonneville County, Idaho*: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-287, 1964.
 1-65. Preliminary geologic map of the Palisades Peak quadrangle, Bonneville County, Idaho, and Teton County, Wyoming: U.S. Geol. Survey open-file report, 1965.
- JOHN, E. C.
 1-64. Petrology and petrography of the intrusive igneous rocks of the Lanai area, Juab County, Utah: Brigham Young Univ. Geology Studies, v. 11, p. 67-96, 1964.
- JOHNSON, Arthur
 1-64. (and SENKPIEL, W. C.) Water resources: Water-power, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 268-273, 1964.
 1-65. Glacier observations, Glacier National Park, Montana, 1964: U.S. Geol. Survey open-file report, 17 p., 1965.

- JOHNSON, A. I.
1-63. (and GNAEDINGER, J. P.) Bibliography, in Symposium on Soil Exploration: Am. Soc. Testing and Materials Spec. Tech. Pub. 351, p. 137-155, 1963.
1-64. Conversion factors and miscellaneous tables: U.S. Geol. Survey open-file report, 38 p., 1964.
1-65. Computer processing of hydrologic and geologic data for water-resources investigations: U.S. Geol. Survey open-file report, 25 p., 1965.
- JOHNSON, F. A.
1-64. Discussion of paper "Weather modification" by R. A. Schleusener: Jour. Hydraulics Div., Am. Soc. Civil Engineers, v. 90, no. HY5, pt. 1, p. 187-196, 1964.
- JOHNSON, J. H.
1-64. Eocene algae from Ishigaki-shima, Ryūkyū-rettō: U.S. Geol. Survey Prof. Paper 399-C, 13 p., 1964.
- JOHNSON, R. B.
1-63. (EGE, J. R., and HOOVER, D. L.) Geology of a prospective tunnel site for the underground storage of air near Wahmonie Flat, Area 401, Nevada Test Site, Nevada: U.S. Geol. Survey Tech. Letter, PLUTO-24, 19 p., 1963.
- JOHNSTON, R. H.
1-64. Ground water in the Niagara Falls area, New York, with emphasis on the water-bearing characteristics of the bedrock: New York State Conserv. Dept. Water Resources Comm. Bull. GW-53, 93 p., 1964.
- JOHNSTON, W. D., Jr.
1-65. The first geological map of the world: GW—The George Washington Univ. Mag., v. 2, no. 1, p. 8-11, 1965.
- JOLLY, J. L.
1-65. (and HEYL, A. V.) Mercury in sphalerite from central Kentucky and central Tennessee [abs.]: Geol. Soc. America, S. E. Sec., Ann. Mtg., Nashville, Tenn., April 1965, Program, p. 23, 1965.
- JONES, B. F.
1-65. (TRUEDELL, A. H., and VAN DENBURGH, A. S.) Glass electrode determination of sodium in closed basin waters [abs.]: Geol. Soc. America Spec. Paper 82, p. 103-104, 1965.
- JONES, C. L.
1-65. Petrography of evaporites from the Wellington Formation near Hutchinson, Kansas: U.S. Geol. Survey Bull. 1201-A, p. A1-A70, 1965.
- JONES, D. L.
1-64. (and BERG, H. C.) Cretaceous stratigraphy of the McCarthy A-4 quadrangle, southern Alaska: U.S. Geol. Survey Bull. 1180-A, p. A1-A18, 1964.
2-64. (and GRANTZ, Arthur) Stratigraphic and structural significance of Cretaceous fossils from the Tiglukpuk Formation, northern Alaska: Am. Assoc. Petroleum Geologists Bull., v. 48, no. 9, p. 1462-1474, 1964.
- JONES, J. R.
1-64. Ground-water maps of the Kingdom of Libya: U.S. Geol. Survey open-file report, 11 p., 1964.
- JONES, P. H.
1-63. (and SKIBITZKE, H. E.) Dispersion of aqueous radioactive waste in a heterogeneous aquifer, in La rétention et la migration des ions radioactifs dans les sols—Colloque international, Saclay, France, 1962: Paris, Presses Universitaires de France, p. 213, 1963.
- JONES, W. R.
1-65. Copper, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 160-176, 1965.
- JOPLING, A. V.
1-64. Laboratory study of sorting processes related to flow separation: Jour. Geophys. Research, v. 69, no. 16, p. 3403-3418, 1964.
- JORDAN, P. R.
1-65. Upper Ohio drainage area, in Ground-water distribution and potential in the Ohio River basin: U.S. Geol. Survey open-file report, 22 p., 1965.
- JORDAN, P. R. --Continued
- KACHADOORIAN, Reuben
1-65. (PLAFKER, George, and GRANTZ, Arthur) Effects of Alaska's Good Friday earthquake on land [abs.]: Geol. Soc. America Spec. Paper 82, p. 105-106, 1965.
- KAHN, Lloyd
1-64. (and WAYMAN, C. H.) Amino acids in raw sewage and sewage effluents: Water Pollution Control Federation Jour., v. 36, no. 11, p. 1368-1371, 1964.
- KAM, William
1-65. Earth cracks—A cause of gully, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B122-B125, 1965.
- KANE, M. F.
1-64. (and CARLSON, J. E.) Bouguer gravity anomaly map of Clark County, Nevada, and Gravity observations and Bouguer anomaly values for Clark County, Nevada: U.S. Geol. Survey open-file report, 34 p., 1964.
2-64. (and SNYDER, G. L.) Shape and structure of a gabbro body near Lebanon, Connecticut, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C22-C27, 1964.
- KANTROWITZ, I. H.
1-64. Bedrock topography in the Oneida Lake area, New York, in New York State Geol. Assoc., 36th Ann. Mtg., 1964, Guidebook: New York, City College of City Univ., Dept. Geology, p. 39-41, 1964.
2-64. Ground-water resources of the Syracuse area, in New York State Geol. Assoc., 36th Ann. Mtg., 1964, Guidebook: New York, City College of City Univ., Dept. Geology, p. 36-38, 1964.
- KARIG, D. E.
1-65. Geophysical evidence of a caldera at Bonanza, Colorado, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B9-B12, 1965.
- KARLSTROM, T. N. V.
1-64. (and others) Surficial geology of Alaska: U.S. Geol. Survey Misc. Geologic Inv. Map I-357, 1964.
- KARREN, R. J.
1-64. Recent studies of leveling instrumentation and procedures: Surveying and Mapping, v. 24, no. 3, p. 383-397, 1964.
- KAYE, C. A.
1-64. Illinoian and early Wisconsin moraines of Martha's Vineyard, Massachusetts, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C140-C143, 1964.
2-64. Outline of Pleistocene geology of Martha's Vineyard, Massachusetts, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C134-C139, 1964.
- KEECH, C. F.
1-64. Basic data report, York County, Nebraska: Nebraska Univ. Conserv. and Survey Div., Nebraska Water Survey Paper 16, 38 p., 1964.
2-64. Ground-water resources of Mirage Flats, Nebraska: U.S. Geol. Survey Water-Supply Paper 1779-BB, p. BB1-BB36, 1964.
3-64. Ground-water conditions in the proposed waterfowl refuge area near Chapman, Nebraska: U.S. Geol. Survey Water-Supply Paper 1779-E, p. E1-E55, 1964.
4-64. (and DREESZEN, V. H.) Availability of ground water in Hall County, Nebraska: U.S. Geol. Survey Hydrol. Inv. Atlas HA-131, map with text, 1964.
- KEEFER, W. R.
1-64. Preliminary report on the structure of the southeast Gros Ventre Mountains, Wyoming, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D22-D27, 1964.
1-65. Geologic history of the Wind River basin, central Wyoming [abs.]: Mtn. Geologist, v. 2, no. 1, p. 43, 1965.

- KEHN, T. M.
 1-64. Geology of the Madisonville West quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-346, 1964.
 2-64. Geology of the Slaughters quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-360, 1964.
- KEIGHTON, W. B.
 1-64. Delaware River water quality, Bristol to Marcus Hook, Pennsylvania, August 1949 to December 1963: U.S. Geol. Survey open-file report, 107 p., 1964.
- KELLY, T. E.
 1-64. Geology and groundwater resources, Barnes County, North Dakota—Pt. 2, Ground-water basic data: North Dakota Geol. Survey Bull. 43, 156 p., 1964.
 2-64. Geohydrology of the Spiritwood aquifer, Stutsman and Barnes Counties, North Dakota, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D161-D165, 1964.
- KENNEDY, V. C.
 1-64. A freezing method for collecting samples of unconsolidated sediment [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 107, 1964.
 2-64. Base-exchange capacity and clay mineralogy of some modern stream sediments: Internat. Assoc. Sci. Hydrology Pub. 64, p. 95-105, 1964.
- KEPFERLE, R. C.
 1-64. (and PETERSON, W. L.) Lithologic and radioactivity log of Belle Scott Quarry drill hole, Meade County, Kentucky: U.S. Geol. Survey open-file report, 1 p., 1964.
- KEROHER, G. C.
 1-64. Glossary of stratigraphic terms, in Upper Paleozoic floral zones and floral provinces of the United States: U.S. Geol. Survey Prof. Paper 454-K, p. K19-K32, 1964.
- KETNER, K. B.
 1-64. Nonmetallic and industrial minerals and materials resources: Refractory minerals, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 205-206, 1964.
 2-64. Nonmetallic and industrial minerals and materials resources: Silica, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 218-222, 1964.
- KIER, P. M.
 1-64. Fossil echinoids from the Marshall Islands: U.S. Geol. Survey Prof. Paper 260-GG, p. GG1121-GG1126, 1964.
 1-65. (GRANT, R. E., and YOCHELSON, E. L.) Whiten-fing fossils, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 453-456, 1965.
- KILSGAARD, T. H.
 1-64. Silver: Mining Cong. Jour., v. 50 no. 9, p. 60-64, 1964.
 2-64. Mineral resources: Coal, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 58-66, 1964.
 3-64. Mineral resources: Silver, lead, and zinc, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 181-194, 1964.
 4-64. (and HEYL, A. V.) Metallic mineral resources: Lead, zinc, and silver, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 96-103, 1964.
- KILBURN, Chabot
 1-64. Ground water in the upper part of the Teton Valley, Teton Counties, Idaho and Wyoming: U.S. Geol. Survey Bull. 1789, 60 p., 1964[1965].
- KILPATRICK, F. A.
 1-64. Source of base flow of streams, in Symposium—Surface waters, Gen. Assembly of Berkeley of I. U. G. G., 1963: Internat. Assoc. Sci. Hydrology Pub. 63, p. 329-339, 1964.
- KIMREY, J. O.
 1-64. The Pungo River Formation, a new name for middle Miocene phosphorites in Beaufort County, North Carolina: Southeastern Geology, v. 5, no. 4, p. 195-205, 1964.
- KINDSVATER, C. E.
 1-65. (CARTER, R. W., and TRACY, H. J.) Computation of peak discharge at contractions: U.S. Geol. Survey Circ. 284, 36 p., 1965. (Originally published 1953.)
- KING, E. R.
 1-65. (ZIETZ, Isidore, and ALLDREDGE, L. R.) Reply to Dr. Ostenso's comments on "The genesis of the Arctic Ocean basin": Science, v. 147, no. 3661, p. 1055-1056, 1965.
- KING, P. B.
 1-64. Further thoughts on tectonic framework of southeastern United States, in Tectonics of the southern Appalachians: Virginia Polytech. Inst. Dept. Geol. Sci. Mem. 1, p. 5-31, 1964.
 2-64. The Cordillera of western North America: Abstract of the geology technical session (Tectonic Symposium), Ann. Western Mtg., CIM, and Pacific NW Minerals Conf., AIME, 1964.
 3-64. The tectonic map of North America [with text and legend] (also French edition): Washington, D. C., U.S. Geological Survey, 26 p., 1964.
 1-65. Permian: Encyclopedia Britannica, v. 17, p. 532-534 (1963 ed.), 1965.
- KING, R. H.
 1-64. Report on three dolomite deposits near Helena, Montana: U.S. Geol. Survey open-file report, 35 p., 1964.
- KING, R. R.
 1-65. (LOUD, E. S., MEAD, M. C., JUSSEN, V. M., and CONANT, G. D.) Bibliography of North American Geology, Pt. 1, Bibliography, Pt. 2, Index: U.S. Geol. Survey Bull. 1195, v. 1, p. 1-1012; v. 2, p. 1013-1790; v. 3, p. 1791-2796; v. 4, p. 2797-4025, 1965.
- KING, R. U.
 1-64. Ferrous metals: Molybdenum, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 102-108, 1964.
 2-64. Metallic mineral resources: Molybdenum, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 74-77, 1964.
 3-64. Metallic mineral resources: Molybdenum, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 111-115, 1964.
 4-64. Mineral resources: Molybdenum, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 133-138, 1964.
 1-65. Molybdenum, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 201-207, 1965.
 2-65. Rhenium, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 375-386, 1965.
- KIRKEMO, Harold
 1-64. Metallic mineral resources: Copper, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 81-87, 1964.
- KISTER, L. R.
 1-65. (BROWN, S. G., SCHUMANN, H. H., and JOHNSON, P. W.) Maps showing fluoride content and salinity of ground water in the Willcox basin, Graham and Cochise Counties, Arizona: U.S. Geol. Survey open-file report, 39 p., 1965.
 2-65. (and HARDT, W. F.) Salinity of the ground water in western Pinal County, Arizona: U.S. Geol. Survey open-file report, 60 p., 1965.
- KISTLER, R. W.
 1-65. (BATEMAN, P. C., and BRANNOCK, W. W.) Isotopic ages of minerals from granitic rocks of the central

- KISTLER, R. W. --Continued
Sierra Nevada and Inyo Mountains, California: Geol. Soc. America Bull., v. 76, no. 2, p. 155-164, 1965.
- KLASNER, J. S.
1-64. A study of buried bedrock valleys near South Haven, Michigan, by the gravity method: U.S. Geol. Survey open-file report, 39 p., 1964.
- KLEIN, Howard
1-64. (SCHROEDER, M. C., and LICHTLER, W. F.) Geology and ground-water resources of Glades and Hendry Counties, Florida: Florida Geol. Survey Rept. Inv. 37, 134 p., 1964.
- KLEINHAMPL, F. J.
1-64. Metallic mineral resources: Silver, lead, and zinc, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 138-150, 1964.
- KLEMIC, Harry
1-64. (and WEST, W. S.) Geology of the Belmont and Calamine quadrangles, Wisconsin: U.S. Geol. Survey Bull. 1123-G, p. G361-G435, 1964.
1-65. Geology of the Honey Grove quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-376, 1965.
- KNOX, C. E.
1-64. (and JOHNSON, C. G., Jr.) Flood frequency formulas for Massachusetts: U.S. Geol. Survey open-file report, map, 1964.
- KOBERG, G. E.
1-64. Elimination of thermal stratification by an air-bubbling technique in Lake Wohlford, California, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D190-D192, 1964.
1-65. The effect of droughts on evaporation rates, in Symposium on consideration of droughts in water planning: Texas Water Comm. Bull. 6512, p. 49-58, 1965.
2-65. (and FORD, M. E., Jr.) Elimination of thermal stratification in reservoirs and the resulting benefits, with special emphasis on study of Lake Wohlford, California: U.S. Geol. Survey Water-Supply Paper 1809-M, p. M1-M28, 1965.
- KOHOUT, F. A.
1-64. Problems of hydrogeology in southern Florida [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 111, 1964.
- KONSELMAN, A. S.
1-63. (and HASLER, J. W., compilers) Mineral deposits suitable for PLOWSHARE mining experiments: U.S. Geol. Survey Tech. Letter, Spec. Proj. -7, 45 p., 1963.
- KOOPMAN, F. C.
1-63. (JOHNSON, A. I., ARMSTRONG, C. A., and TAYLOR, R. E.) Results of aquifer tests at Hydrologic Test Sites 1 and 2, Tatum Dome, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-32, 31 p., 1963.
- KOSANKE, R. M.
1-64. Applied Paleozoic palynology, in Palynology in oil exploration--A symposium, San Francisco, Calif., 1962: Soc. Econ. Paleontologists and Mineralogists Spec. Pub. 11, p. 75-89, 1964.
1-65. Palynological investigations in the Pennsylvanian of Kentucky--I: U.S. Geol. Survey open-file report, 38 p., 1965.
2-65. Palynological studies of the Hazard No. 7 and Francis coals of Kentucky [abs.]: Geol. Soc. America Spec. Paper 82, p. 112-113, 1965.
3-65. Plant microfossils of the Hazard No. 7 coal, Perry County, Kentucky: U.S. Geol. Survey open-file report, 3 p., 1965.
- KOSCHMANN, A. H.
1-65. (LOUGHLIN, G. F., and others) Mine maps of the Cripple Creek district, Colorado: U.S. Geol. Survey open-file report, 523 maps, 1965.
- KOTEFF, Carl
1-64. Geology of the Assawompset Pond quadrangle, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-265, 1964.
2-64. Surficial geology of the Concord quadrangle, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-331, 1964.
- KOYANAGI, R. Y.
1-64. Seismicity of the lower east rift zone of Kilauea volcano, Hawaii, January 1962-March 1963, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C103-C105, 1964.
2-64. (OKAMURA, A. T., KINOSHITA, W. T., MOORE, J. G., and POWERS, H. A.) Hawaiian Volcano Observatory Summary, July, August, and September 1963: U.S. Geol. Survey Hawaiian Volcano Observatory Summ. 31, 40 p., 1964.
1-65. (and ENDO, E. T.) Hawaiian seismic events during 1963, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B13-B16, 1965.
2-65. (OKAMURA, A. T., and POWERS, H. A.) Hawaiian Volcano Observatory Summary, January, February, and March 1964: U.S. Geol. Survey Hawaiian Volcano Observatory Summ. 33, 44 p., 1965.
- KRINSLEY, D. B.
1-64. Multiple glaciation of south-west Yukon [abs.], in Internat. Geog. Cong., 20th, United Kingdom, 1964, Abs. Papers: London, Thomas Nelson and Sons Ltd., p. 94, 1964.
1-65. Pleistocene geology of the south-west Yukon Territory, Canada: Jour. Glaciology, v. 5, no. 40, p. 385-397, 1965; abstract in Geol. Soc. America Spec. Paper 82, p. 114-115, 1965.
- KRIER, E. P.
1-65. Photographic copying using reflection-transmission illumination, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B190-B191, 1965.
- KRINSLEY, David
1-65. (and HAMILTON, Warren) Permo-Carboniferous tillites from South Africa, Australia, and Antarctica [abs.]: Geol. Soc. America Spec. Paper 82, p. 115, 1965.
- KRIVOY, H. L.
1-64. (KOYANAGI, R. Y., OKAMURA, Arnold, and KOJIMA, George) Hawaiian Volcano Observatory Summary 28, October, November, and December, 1962: U.S. Geol. Survey Hawaiian Volcano Observatory Summary 28, p. 6-51, 1964.
1-65. (KINOSHITA, W. T., OKAMURA, A. T., and KOYANAGI, R. Y.) Hawaiian Volcano Observatory Summary, April, May, and June 1963: U.S. Geol. Survey Hawaiian Volcano Observatory Summ. 30, 32 p., 1965.
- LACHENBRUCH, A. H.
1-64. (and MARSHALL, B. V.) Heat flux from the Arctic Ocean basin, preliminary results [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 123, 1964.
1-65. The effect of a slope on surface heat flow [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 176, 1965.
- LaCORNU, E. J.
1-65. (HANSON, R. L., and CRUFF, R. W.) Comparison of discharge measurements made by the point-velocity and the velocity-integration methods: U.S. Geol. Survey open-file report, 17 p., 1965.
- LADD, H. S.
1-65. Collecting fossils on tropical Pacific Islands, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 163-168, 1965.
2-65. Tertiary fresh-water mollusks from Pacific Islands: Malacologia, v. 2, no. 2, p. 189-198, 1965.
- LaHEIST, B. A.
1-64. Metallic mineral resources: Arsenic, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 65-70, 1964.
2-64. Mineral resources: Antimony, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 41-46, 1964.

- LaHEIST, B. A. --Continued
 3-64. Mineral resources: Smelter byproducts, *in* Mineral water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 194-211, 1964.
- LAI, Chintu
 1-65. Discussion of "Mathematical model for transient river flow", by Eivind Kindingstad: *Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour.*, v. 91, no. HY1, p. 167-172, 1965.
- LAIRD, L. B.
 1-64. Chemical quality of the surface waters of the Snake River basin: U.S. Geol. Survey Prof. Paper 417-D, p. D1-D47, 1964.
- LAKIN, H. W.
 1-65. (and NAKAGAWA, H. M.) A spectrophotometric method for the determination of gold useful in mineral exploration: U.S. Geol. Survey open-file report, 13 p., 1965.
- LAMAR, W. L.
 1-64. (GOERLITZ, D. F., and LAW, L. M.) Detection and measurement of chlorinated organic pesticides in water by electron capture gas chromatography: U.S. Geol. Survey open-file report, 25 p., 1964.
- LaMARCHE, V. C., Jr.
 1-64. Determination of denudation rates from measurements of root exposure [abs.]: *Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program*, p. 117, 1964.
- LAMBERT, T. W.
 1-64. Availability of ground water in the New Concord quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-118, 1964.
 2-64. (and MacCARY, L. M.) Geology of the Briensburg quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-327, 1964.
 1-65. Availability of ground water in the Fairdealing quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-156, 1965.
- LANDEN, David
 1-64. Alaska earthquake, 27 March 1964: *Science*, v. 145, no. 3627, p. 74-76, 1964.
- LANDIS, E. R.
 1-64. Lignitic coal of North Dakota, *in* The mineral resources of North Dakota: Grand Forks, Univ. North Dakota, p. 1-13, 1964.
 2-64. Mineral fuels and associated resources: Coal, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 35-45, 1964.
 3-64. Mineral fuel resources: Coal, *in* Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 147-151, 1964.
 1-65. Coal resources of Iowa [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 118, 1965.
- LANG, J. W.
 1-64. (and NEWCOME, Roy, Jr.) Status of salt-water encroachment in aquifers along the Mississippi Gulf Coast, 1964: Mississippi Board of Water Commissioners Bull. 64-5, 16 p., 1964.
- LANG, S. M.
 1-65. (and LEONARD, A. R.) Instructions for using the punch-card system for the storage and retrieval of ground-water data: U.S. Geol. Survey open-file report, 52 p., 1965.
- LANGBEIN, W. B.
 1-64. Hydrologic tools in paleoecological reconstruction, *in* The reconstruction of past environments—Fort Burgwin Conf. on Paleoecology, 1962, Proc. Fort Burgwin Research Center Pub. 3, p. 37-39, 1964.
 2-64. (and DAWDY, D. R.) Occurrence of dissolved solids in surface waters in the United States, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D115-D117, 1964.
- LANGFORD, R. H.
 1-64. The chemical quality of the groundwater, *in* Ground water resources and geology of northern and western Crook County, Wyoming: U.S. Geol. Survey Water-Supply Paper 1698, p. 52-77, 1964.
- LANPHERE, M. A.
 1-64. (MacKEVETT, E. M., Jr., and STERN, T. W.) Potassium-argon and lead-alpha ages of some plutonic rocks, Bokan Mountain area, southeastern Alaska: *Science*, v. 145, no. 3633, p. 705-707, 1964.
 1-65. (and DALRYMPLE, G. B.) An interlaboratory standard muscovite for argon and potassium analyses [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 178, 1965.
 2-65. (LONEY, R. A., and BREW, D. A.) Potassium-argon ages of some plutonic rocks, Tenakee area, Chichagof Island, southeastern Alaska, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B108-B111, 1965.
- LaROCQUE, G. A., Jr.
 1-65. General availability of ground water and depth to water level in the Missouri River Basin: U.S. Geol. Survey open-file report, map, 1965.
- LARRABEE, D. M.
 1-64. Bedrock geologic map of the Big Lake quadrangle, Washington County, Maine: U.S. Geol. Survey Geol. Quad. Map GQ-358, 1964.
 2-64. Reconnaissance bedrock geology of the Wabassua Lake quadrangle, Maine: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-282, 1964.
- LARSEN, E. S.
 1-64. (and BERMAN, Harry) The microscopic determination of the nonopaque minerals, second edition: U.S. Geol. Survey Bull. 848, 266 p., 1964. (originally printed 1934.)
- LARSON, R. R.
 1-64. (DWORNIK, E. J., and ADLER, I.) Electronprobe analysis of "cosmic" particles, *in* Cosmic dust: New York Acad. Sci. Annals, v. 119, art. 1, p. 282-286, 1964.
- LaSALA, A. M., Jr.
 1-64. Geology and ground-water resources of the Bristol-Plainville-Southington area, Connecticut: U.S. Geol. Survey Water-Supply Paper 1578, 70 p., 1964[1965].
 2-64. (and MIEKLE, R. L.) Records and logs of selected wells and test borings and chemical analyses of water in the Bristol-Plainville-Southington area, Connecticut: Connecticut State Water Resources Comm. Bull. no. 5, 18 p., 1964.
 3-64. (HARDING, W. E., and ARCHER, R. J.) Water resources of the Lake Erie-Niagara area, New York, a preliminary appraisal: New York Div. of Water Resources Bull. GW-52, 1964.
- LATHRAM, E. H., compiler
 1-65. Preliminary geologic map of northern Alaska: U.S. Geol. Survey open-file report, 1965.
 2-65. (POMEROY, J. S., BERG, H. C., and LONEY, R. A.) Reconnaissance geology of Admiralty Island, Alaska: U.S. Geol. Survey Bull. 1181-R, p. R1-R48, 1965.
- LEE, D. R.
 1-64. Daylight star observations in Antarctica: *Am. Soc. Civil Engineers Proc.*, v. 90, Jour. Surveying and Mapping Div., no. SU 2, p. 75-82, 1964.
- LEE, F. T.
 1-65. (ROBINSON, C. S., and CARROLL, R. D.) Loveland Basin landslide, Colorado [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 336, 1965.
- LeGRAND, H. E.
 1-64. Hydrogeologic framework of the Gulf and Atlantic Coastal Plain: *Southeastern Geology*, v. 5, no. 4, p. 177-194, 1964.
 2-64. Management aspects of ground-water contamination: *Water Pollution Control Federation Jour.*, v. 36, no. 9, p. 1133-1145, 1964.

LeGRAND, H. E. --Continued

3-64. System for evaluation of contamination potential of some waste disposal sites: *Am. Water Works Assoc. Jour.*, v. 56, no. 8, p. 959-974, 1964.

1-65. Environmental framework of ground-water contamination: *Ground Water*, v. 3, no. 2, p. 11-15, 1965.

2-65. Patterns of contaminated zones of water in the ground: *Water Resources Research*, v. 1, no. 1, p. 83-95, 1965.

3-65. Spectra of ground-water conditions [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 75, 1965.

LEIFESTE, D. K.

1-65. (and SMITH, J. T.) Base-flow studies, San Gabriel River, Texas—Quantity and quality, March 16-18, 1964: *Texas Water Comm. Bull.* 6510, 17 p., 1965.

LEIGHTON, M. M.

1-65. (and RAY, L. L.) Glacial deposits of Nebraskan and Kansan age in northern Kentucky, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, B126-B131, 1965.

LEMKE, R. W.

1-65. Engineering geologic effects of the March 27, 1964, earthquake at Seward, Alaska [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 121-122, 1965.

LEMMON, D. M.

1-64. Metallic mineral resources: Tungsten, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 121-124, 1964.

1-65. Geology and exploration: *Mining Eng.*, v. 17, no. 2, p. 83-84, 1965.

LEO, G. W.

1-65. (ROSE, H. J., Jr., and WARR, J. J.) Chromian muscovite from the Serra de Jacobina, Bahia, Brazil: *Am. Mineralogist*, v. 50, nos. 3-4, p. 392-402, 1965.

LEONARD, B. F.

1-64. (and BUDDINGTON, A. F.) Ore deposits of the St. Lawrence County magnetite district, northwest Adirondacks, New York: U.S. Geol. Survey Prof. Paper 377, 259 p., 1964.

1-65. Mercury-bearing antimony deposit between Big Creek and Yellow Pine, central Idaho, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B23-B28, 1965.

2-65. Tertiary dike swarm on Little Pistol Creek, Yellow Pine quadrangle, central Idaho [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 336, 1965.

LEONARD, R. B.

1-64. Cooperative water-quality investigation of the Walnut River basin: *Kansas Water News*, v. 7, no. 5, p. 5-6, 1964.

2-64. Results of four chemical-quality surveys in the Walnut River basin, Kansas (December 1961 to October 1963): U.S. Geol. Survey open-file report, 90 p., 1964.

3-64. (and SHAH, Kanti) The relation of the chemical quality of the Whitewater River at Towanda to that of the Walnut River at Winfield, Kansas, during the 1963 water year: U.S. Geol. Survey open-file report, 19 p., 1964.

LEOPOLD, E. B.

1-64. Reconstruction of Quaternary environments using palynology, in *The reconstruction of past environments*, assembled by J. J. Hester and James Schoenwetter: Santa Fe, Fort Burgwin Conf. Paleocology Proc., 1962, no. 3, p. 43-52, 1964.

2-64. (and PAKISER, H. M.) A preliminary report on the pollen and spores of the pre-Selma Upper Cretaceous strata of western Alabama: U.S. Geol. Survey Bull. 1160-E, p. E71-E95, 1964.

1-65. (and TSCHUDY, Bernadine) Plant and miscellaneous microfossils of the Pierre Shale: U.S. Geol. Survey open-file report, 4 p., 1965.

LEOPOLD, L. B.

1-64. International hydrological decade: *Geotimes*, v. 9, no. 4, p. 16, 1964.

2-64. (WOLMAN, M. G., and MILLER, J. P.) Fluvial processes in geomorphology: San Francisco, Calif., W. H. Freeman and Company, 522 p., 1964.

1-65. Water supply and management concepts: *Indus. Water Eng.*, v. 2, no. 1, p. 10-13, 22, 1965.

2-65. Ecological systems and the water resource: U.S. Geol. Survey Circ. 414-D, p. 21-26, 1965. (Originally published 1960.)

3-65. The challenge of water management: U.S. Geol. Survey Circ. 414-B, p. B7-B13, 1965. (Originally published 1960.)

LEPP, Henry

1-64. (and GOLDICH, S. S.) Origin of Precambrian iron formations: *Econ. Geology*, v. 59, no. 6, p. 1025-1060, 1964.

LESURE, F. G.

1-64. Mineral resources: Mica and associated pegmatite minerals, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 123-133, 1964.

2-64. Nonmetallic mineral resources: Pegmatite minerals, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 223-230, 1964.

1-65. Pegmatite minerals, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 278-290, 1965.

LEVE, G. W.

1-64. Analysis of current-meter data from wells by flow-distribution curves: *Ground Water*, v. 2, no. 2, p. 12-17, 1964.

LEWANDOWSKI, R. A.

1-65. Blade mounting, in *Handbook of paleontological techniques*: San Francisco, Calif., W. H. Freeman and Company, p. 256-257, 1965.

LEWIS, R. Q., Sr.

1-64. (and THADEN, R. E.) Geology of the Montpelier quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-337, 1964.

1-65. (and CAMPBELL, R. H.) Geology and uranium deposits of Elk Ridge and vicinity, San Juan County, Utah: U.S. Geol. Survey Prof. Paper 474-B, p. B1-B69, 1965.

2-65. (and THADEN, R. E.) Geology of the Russell Springs quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-383, 1965.

LIEBER, Maxim

1-64. (PERLMUTTER, N. M., and FRAUENTHAL, H. L.) Cadmium and hexavalent chromium in Nassau County ground water: *Am. Water Works Assoc. Jour.*, v. 56, no. 6, p. 739-747, 1964.

LINDBERG, M. L.

1-64. Crystallography of faheyite, Sapucaia pegmatite mine, Minas Gerais, Brazil: *Am. Mineralogist*, v. 49, nos. 3-4, p. 395-398, 1964.

LINEDALL, R. M.

1-64. Construction materials: Stone, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 211-215, 1964.

1-65. Stone, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 361-365, 1965.

LIPMAN, P. W.

1-64. (and McKAY, E. J.) Geology of the Topopah Spring SW quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-72, 8 p., 1964.

1-65. Chemical comparison of glassy and crystalline volcanic rocks: U.S. Geol. Survey Bull. 1201-D, p. D1-D24, 1965.

LIPMAN, P. W. --Continued

- 2-65. (QUINLIVAN, W. D., CARR, W. J., and ANDERSON, R. E.) Geologic map and sections of the Thirsty Canyon SE quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report TEI-859, 1965.

LIVINGSTONE, D. A.

- 1-64. Chemical composition of rivers and lakes, in Data of geochemistry, 6th ed.: U.S. Geol. Survey Prof. Paper 440-G, p. G1-G64, 1964. (Originally published 1963.)

LLOYD, O. B., Jr.

- 1-65. (and WILDER, H. B.) Ground-water supply of Cape Hatteras National Seashore Recreational area, North Carolina, Pt. 4: U.S. Geol. Survey open-file report, 23 p., 1965.

LOEBLICH, A. R., Jr.

- 1-64. (and TAPPAN, Helen) The species and stratigraphic distribution of *Caucasina* and *Aeolomorphella*, new genus (Foraminifera): Tulane Studies Geology, v. 2, no. 3, p. 69-88, 1964.

LOHMAN, K. E.

- 1-64. Stratigraphic and paleoecologic significance of the Mesozoic and Cenozoic diatoms of California and Nevada, in Palynology in oil exploration—A symposium, San Francisco, Calif., 1962: Soc. Econ. Paleontologists and Mineralogists Spec. Pub. 11, p. 58-64, 1964.

LOHMAN, S. W.

- 1-65. Geology and artesian water supply of the Grand Junction area, Colorado: U.S. Geol. Survey Prof. Paper 451, 149 p., 1965.
- 2-65. The geologic story of Colorado National Monument: Colorado and Black Canyon Nat. Hist. Assoc., 56 p., 1965.

LOHR, E. W.

- 1-64. (and LOVE, S. K.) The industrial utility of public water supplies in the United States, 1952—Pt. 1, States east of the Mississippi River: U.S. Geol. Survey Water-Supply Paper 1299, 639 p., 1964. (Originally published 1954.)
- 2-64. (and LOVE, S. K.) The industrial utility of public water supplies in the United States, 1952—Pt. 2, States west of the Mississippi River: U.S. Geol. Survey Water-Supply Paper 1300, 462 p., 1964. (Originally published 1952.)

LONEY, R. A.

- 1-64. Stratigraphy and petrography of the Pybus-Gambier area, Admiralty Island, Alaska: U.S. Geol. Survey Bull. 1178, 103 p., 1964 [1965].
- 1-65. Structural analysis of the Pybus-Gambier area, Admiralty Island, Alaska: Calif. Univ. Pubs. Geol. Sci., v. 46, no. 2, p. 33-80, 1965.

LONG, R. A.

- 1-64. Scavenger well system—A solution to the problem of vertical salt-water encroachment [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 123-124, 1964.

LOPEZ, M. A.

- 1-64. Floods at Toa Alta, Toa Baja, and Dorado, Puerto Rico: U.S. Geol. Survey Hydrol. Inv. Atlas HA-128, 1964.

LOVE, J. D.

- 1-64. Uraniferous phosphatic lake beds of Eocene age in Intermontane basins of Wyoming and Utah: U.S. Geol. Survey Prof. Paper 474-E, p. E1-E66, 1964.

LOVE, S. K.

- 1-64. Colorado River Basin to Pacific Slope Basins in Oregon and Lower Columbia River Basin—Pts. 9-14: U.S. Geol. Survey Water-Supply Paper 1945, 691 p., 1964.
- 2-64. Hudson Bay and Upper Mississippi River Basins, and Missouri River Basin—Pts. 5 and 6: U.S. Geol. Survey Water-Supply Paper 1943, 413 p., 1964.

LOVE, S. K. --Continued

- 3-64. Lower Mississippi River Basin and Western Gulf of Mexico Basins—Pts. 7 and 8: U.S. Geol. Survey Water-Supply Paper 1944, 645 p., 1964.

- 4-64. Ohio River Basin and St. Lawrence River Basin—Pts. 3 and 4: U.S. Geol. Survey Water-Supply Paper 1942, 322 p., 1964.

- 5-64. Quality of surface waters of the United States, 1958: Geol. Survey Water-Supply Paper 1574, 487 p., 1964.

- 6-64. Quality of surface waters of the United States, 1962—Pts. 1 and 2, North Atlantic slope basins and south Atlantic slope and eastern Gulf of Mexico basins: U.S. Geol. Survey Water-Supply Paper 1941, 434 p., 1964.

- 1-65. Quality of surface waters of the United States 1959—Pts. 1 and 2, North Atlantic slope basins and South Atlantic slope and eastern Gulf of Mexico basins: U.S. Geol. Survey Water-Supply Paper 1641, 460 p., 1965.

- 2-65. Quality of surface waters of the United States 1959—Pts. 3 and 4, Ohio River basin and St. Lawrence River basin: U.S. Geol. Survey Water-Supply Paper 1642, 303 p., 1965.

- 3-65. Quality of surface waters of the United States 1959—Pts. 7 and 8, Lower Mississippi River basin and western Gulf of Mexico basins: U.S. Geol. Survey Water-Supply Paper 1644, 508 p., 1965.

LOVING, H. B.

- 1-64. The airborne control system [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 31, 1964.

LOW, Doris

- 1-64. Redescription of *Anomalina eaglefordensis* Moreman: Cushman Found. Foram. Research Contr., v. 15, pt. 3, no. 286, p. 122-123, 1964.

LOWE, A. S.

- 1-64. Floods of 1964 near Baton Rouge, La.: U.S. Geol. Survey open-file report, 1 p., 1964.

LOWELL, W. R.

- 1-65. Geologic map of the Bannack-Grayling area, Beaverhead County, Montana: U.S. Geol. Survey Misc. Geol. Inv. Map I-433, with 6-page text, 1965.

LOWRY, M. E.

- 1-64. (and CUMMINGS, T. R.) Ground-water resources of Sheridan County, Wyoming: U.S. Geol. Survey open-file report, 1964.

- 2-64. (and GORDON, E. D.) Ground-water investigations in Yellowstone National Park, October 1960 to October 1963: U.S. Geol. Survey open-file report, 39 p., 1964.

LUFT, S. J.

- 1-63. Some physical property data for basaltic rocks, basalt flows of Fortymile Canyon, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-44, supp. 1, 5 p., 1963.

- 2-63. (and BYERS, F. M., Jr.) Preliminary report on the basalt flows of Fortymile Canyon, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-44, 14 p., 1963.

- 1-64. Mafic lavas of Dome Mountain, Timber Mountain caldera, southern Nevada, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D14-D21, 1964.

LUGN, R. V.

- 1-63. Photogrammetric mapping of experimental craters, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. D, Studies for Space Flight Program: U.S. Geol. Survey open-file report, p. 64-70, 1963.

LUSTIG, L. K.

- 1-64. Geomorphic parameters and the sediment yield of drainage basins in the San Gabriel Mountains, California [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 125, 1964.

- 1-65. (and BUSCH, R. D.) Sediment transport in Cache Creek drainage basin in Coast Ranges west of Sacramento,

- LUSTIG, L. K. --Continued
California: U.S. Geol. Survey open-file report, 101 p., 1965.
- LUZIER, J. E.
1-64. Ground-water supply for Mount Rainier National Park headquarters site near Ashford, Washington: U.S. Geol. Survey open-file report, 35 p., 1964.
- LYONS, J. B.
1-64. Distribution of thorium and uranium in three early Paleozoic plutonic series of New Hampshire: U.S. Geol. Survey Bull. 1144-F, p. F1-F43, 1964.
- MABEY, D. R.
1-64. Aeromagnetic survey of the Antler Peak quadrangle, Humboldt and Lander Counties, Nevada: U.S. Geol. Survey Prof. Paper 459-A, p. A70-A71, 1964[1965].
1-65. Regional gravity and aeromagnetic surveys applied to groundwater studies in the southwestern United States [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 78, 1965.
- McANDREWS, Harry
1-64. Geologic map of the Big Judson quadrangle, Albany County, Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-390, 1964[1965].
2-64. Geologic map of the Cooper Lake South quadrangle, Albany County, Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-391, 1964.
1-65. Geologic map of the Cooper Lake North quadrangle, Albany County, Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-430, 1965.
- McCARTHY, J. H.
1-65. (LOVERING, T. S., and LAKIN, H. W.) Density comparison method for the measurement of isotopic variations in prepared waters: U.S. Geol. Survey open-file report, 38 p., 1965.
- McCARTHY, L. T., Jr.
1-64. (and KEIGHTON, W. B.) Quality of Delaware River water at Trenton, New Jersey: U.S. Geol. Survey Water-Supply Paper 1779-X, p. X1-X51, 1964.
- McCARTNEY, D.
1-65. (and BEAMER, N. H.) Instrumentation in stream quality studies: *Indus. Water Eng.*, v. 2, no. 2, p. 10-13, 1965.
- MacCARY, L. M.
1-64. Availability of groundwater in the Briensburg quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-114, 1964[1965].
2-64. Availability of ground water in the Hardin quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-115, 1964[1965].
3-64. Availability of ground water in the Hazel quadrangle, Kentucky-Tennessee: U.S. Geol. Survey Hydrol. Inv. Atlas HA-124, 1964[1965].
4-64. (and LAMBERT, T. W.) Geology of the Briensburg quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-327, 1964.
- McCAULEY, J. F.
1-63. A preliminary report on the geology of the Hevelius, in *Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963*—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 74-84, 1963.
2-63. The stratigraphy of the Mare Orientale Region of the Moon, in *Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963*—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 86-97, 1963.
1-65. Terrain analysis of the lunar equatorial belt: U.S. Geol. Survey open-file report, 44 p., 1965.
- MACLAY, R. W.
1-64. Reconnaissance of the geology and ground-water resources in the Aurora area, St. Louis County, Minnesota: U.S. Geol. Survey open-file report, 50 p., 1964.
- MACLAY, R. W. --Continued
1-65. (WINTER, T. C., and PIKE, G. M.) Water resources of the Middle River watershed, northwestern Minnesota: U.S. Geol. Survey Hydrol. Inv. Atlas HA-201, 1965.
- McCLELLAND, E. J.
1-65. (and BADER, J. S.) A well-numbering grid: U.S. Geol. Survey open-file report, 6 p., 1965.
- McCLYMONDS, N. E.
1-64. (and HEINDL, L. A.) Stratigraphic sections of younger Precambrian and Paleozoic Formations, Papago Indian Reservation, Arizona: U.S. Geol. Survey open-file report, 118 p., 1964.
- McCOLLUM, M. J.
1-64. Salt-water movement in the principal artesian aquifer of the Savannah area, Georgia and South Carolina: *Ground Water*, v. 2, no. 4, p. 4-8, 1964.
2-64. (and COUNTS, H. B.) Relation of salt-water encroachment to the major aquifer zones, Savannah area, Georgia and South Carolina: U.S. Geol. Survey Water-Supply Paper 1613-D, p. D1-D26, 1964.
3-64. (and HERRICK, S. M.) Offshore extension of the Tertiary stratigraphic section in southeastern Georgia [abs.]: *Georgia Acad. Sci. Bull.*, v. 22, no. 8, p. 15, 1964.
4-64. (and HERRICK, S. M.) Offshore extension of the upper Eocene to recent stratigraphic sequence in southeastern Georgia, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C61-C63, 1964.
- McCONAGHY, J. A.
1-64. (CHASE, G. H., BOETTCHER, A. J., and MAJOR, T. J.) Hydrogeologic data of the Denver Basin, Colorado: Colorado Water Conserv. Board, Ground Water Basic Data Release 15, 224 p., 1964.
2-64. (and COLBURN, G. W.) Records of wells in Colorado: Colorado Water Conserv. Board, Ground Water Basic Data Release 17, 384 p., 1964.
- McCULLOCH, D. S.
1-65. (TAYLOR, Dwight, and RUBIN, Myer) Stratigraphy, nonmarine mollusks and radiometric dates from Quaternary deposits in the Kotzebue Sound area, western Alaska: *Jour. Geology*, v. 73, no. 3, p. 442-453, 1965.
- McCULLOCH, T. H.
1-65. Geologic map of the Nebo and Yermo quadrangles, San Bernardino County, California: U.S. Geol. Survey open-file report, 1965.
- MacDONALD, G. A.
1-64. Geology of the Prospect Peak quadrangle, California: U.S. Geol. Survey Geol. Quad. Map GQ-345, 1964.
1-65. (and KATSURA, Takashi) Eruption of Lassen Peak, Cascade Range, California, in 1915: example of mixed magmas: *Geol. Soc. America Bull.*, v. 76, no. 5, p. 475-482, 1965.
- McGAVOCK, E. H.
1-64. Apache County, Navajo County, [and] Coconino County, in *Annual report on ground water in Arizona, Spring 1963 to Spring 1964*: Arizona Land Dept. Water Resources Rept. 19, p. 49-52, 1964.
- McGOVERN, H. E.
1-64. Geology and ground-water resources of Washington County, Colorado: U.S. Geol. Survey Water-Supply Paper 1777, 46 p., 1964.
2-64. (GREGG, D. O., and BRENNAN, Robert) Hydrogeologic data of the alluvial deposits in Pueblo and Fremont Counties, Colorado: Colorado Water Conserv. Board Ground Water Basic Data Release 18, 27 p., 1964.
- McGUINNESS, C. L.
1-64. Carbonate rocks as aquifers [abs.]: *Geol. Soc. America Ann. Mtg.*, Miami Beach, Florida, Nov. 1964, Program, p. 131-132, 1964.
2-64. Generalized map showing annual runoff and productive aquifers in the conterminous United States: U.S. Geol. Survey Hydrol. Inv. Atlas HA-194, 1964.

McGUINNESS, C. L. --Continued

- 3-64. Ground-water research of the U.S. Geological Survey: U.S. Geol. Survey Circ. 492, 7 p., 1964.
- 1-65. Ground-water—A key resource: GroundWater, v. 3, no. 1, p. 24-29, 1965.
- 2-65. (and MEYER, Gerald) West Virginia's water situation—The role of groundwater: West Virginia Geol. Survey Newsletter, 7th issue, 5 p., 1965.

MacKALLOR, J. A.

- 1-65. Aeroradioactivity survey and geology of the GNOME (Carlsbad) area, New Mexico and Texas (ARMS-L): U.S. Atomic Energy Comm., Civil Effects Test Operations, CEX-59, 4, 24, 1965.
- 2-65. Natural gamma aeroradioactivity of the GNOME (Carlsbad) area, New Mexico and Texas: U.S. Geol. Survey Geophys. Inv. Map GP-462, 1965.

McKAY, E. J.

- 1-63. Geology of Jackass Flats quadrangle: U.S. Geol. Survey Tech. Letter NTS-57, 6 p., 1963.
- 2-63. Hydrothermal alteration in the Calico Hills, Jackass Flats quadrangle, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-43, 6 p., 1963.
- 1-64. (and WILLIAMS, W. P.) Geology of the Jackass Flats quadrangle, Nye County, Nevada: U.S. Geol. Survey Geol. Quad. Map GQ-368, 1964.

McKEE, E. D.

- 1-64. Inorganic sedimentary structures, *in* Approaches to paleoecology: New York, John Wiley and Sons, Inc., p. 275-295, 1964.
- 2-64. Problems of the recognition of arid and of hot climates of the past, *in* Problems in palaeoclimatology: New York, John Wiley and Sons, p. 367-377, 1964.

McKELVEY, V. E.

- 1-64. Future energy demand and supply: World Power Conf., 6th, Melbourne, Australia, Oct. 1962, Proc., v. 11, p. 4275, 1964.
- 2-64. Progress in the science of ore-finding: Mining Cong. Jour., v. 50, no. 2, p. 45-49, 1964.
- 3-64. Science and technology—Keys to abundance or instruments of destruction, *in* Issues in conservation: a series of eight discussions on conservation issues facing Wisconsin as set within the overall national context: Madison, Wisconsin Univ., Extension Div., p. 171-211, 1964.
- 1-65. (and DUNCAN, D. C.) United States and world resources of energy, *in* Symposium on fuel and energy economics...: Am. Chem. Soc., Div. Fuel Chemistry, Preprints of papers presented at 149th National Meeting, Detroit, Michigan, Apr. 4-9, 1965, v. 9, no. 2, p. 1-17, 1965.

MacKEVETT, E. M., Jr.

- 1-64. Ore controls at the Kathleen-Margaret (Maclaren River) copper deposit, Alaska, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C117-C120, 1964.
- 2-64. (and BLAKE, M. C., Jr.) Geology of the Sumdum copper-zinc prospect, southeastern Alaska: U.S. Geol. Survey Bull. 1108-E, p. E1-E31, 1964.
- 1-65. Preliminary geologic map of the McCarthy C-6 quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-444, 1965.

McMASTER, W. M.

- 1-65. Geology of the Elkmont quadrangle, Alabama-Tennessee: U.S. Geol. Survey Misc. Geol. Inv. Maps I-419, with 7-page text, 1965.
- 2-65. Geology of the Salem quadrangle, Alabama-Tennessee: U.S. Geol. Survey Misc. Geol. Inv. Map I-420, with 10-page text, 1965.

MacNEIL, F. S.

- 1-64. Eocene megafossils from Ishigaki-shima Ryukuretto: U.S. Geol. Survey Prof. Paper 399-B, p. B1-B14, 1964.

MacNEIL, F. S. --Continued

- 1-65. Evolution and distribution of the genus *Mya*, and Tertiary migrations of Mollusca: U.S. Geol. Survey Prof. Paper 483-G, p. G1-G51, 1965.

McQUEEN, I. S.

- 1-64. (MILLER, R. F., and BULLER, William) Calibration and evaluation of a wide range method for measuring moisture stress in field soil samples [abs.]: Am. Soc. Agronomy and Soil Sci. Soc. America Ann. Mtg., Kansas City, Missouri, Nov. 1964, Program "Agronomy Abstracts", p. 2, 1964.

MacQUOWN, W. C.

- 1-65. Geologic map of part of the Clintonville quadrangle, Kentucky: U.S. Geol. Survey open-file report, 1965.

MADERAK, M. L.

- 1-64. Relation of chemical quality of water to recharge to the Jordan sandstone in the Minneapolis-St. Paul area, Minnesota, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C176-C179, 1964.

MALDE, H. E.

- 1-64. The ecologic significance of some unfamiliar geologic processes, *in* The reconstruction of past environments—Fort Burgwin Conf. on Paleoecology, 1962, Proc.: Fort Burgwin Research Center Pub. 3, p. 7-15, 1964.
- 2-64. Environment and man in arid America: Science, v. 145, no. 3628, p. 123-129, 1964.
- 3-64. (and SCHICK, A. P.) Thorne Cave, northeastern Utah: Geology: Am. Antiquity, v. 30, no. 1, p. 60-73, 1964.

MALLORY, E. C., Jr.

- 1-65. Leachable silica and alumina in streambed clays, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B170-B174, 1965.

MALTBY, C. S.

- 1-65. An improved heat shield for precise levels: Surveying and Mapping, v. 25, no. 1, p. 102, 1965.

MAMAY, S. H.

- 1-65. Collecting coal balls, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 192-193, 1965.

MANGER, G. E.

- 1-65. The best value of porosity of lapilli tuff from the Nevada Test Site, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B146-B150, 1965.

MAPEL, W. J.

- 1-64. (and PILLMORE, C. L.) Geology of the Upton quadrangle, Crook and Weston Counties, Wyoming: U.S. Geol. Survey Bull. 1181-J, p. J1-J54, 1964.

MARCHER, M. V.

- 1-64. (BINGHAM, R. H., and LOUNSBURY, R. E.) Ground-water geology of the Dickson, Lawrenceburg, and Waverly areas in the western Highland Rim, Tennessee: U.S. Geol. Survey Water-Supply Paper 1764, 50 p., 1964.
- 1-65. A summary of water-supply problems in Alaska: Alaskan Sci. Conf., 15th, Sept. 1964, Proc., p. 375-379, 1965.

MAREK, Ladislav

- 1-64. (and YOCHELSON, E. L.) Paleozoic mollusk—Hyo-lithes: Science, v. 146, no. 3652, p. 1674-1675, 1964.

MARIE, J. R.

- 1-65. Occurrence of water with high hardness and high chloride content in the alluvial deposits of Franklin Parish, Louisiana [abs.]: Geol. Soc. America, SE section, Ann. Mtg., Nashville, Tennessee, April 1965, Program, p. 27-28, 1965.

MARINE, I. W.

- 1-64. Hydrogeology of crystalline rock beneath the Atlantic Coastal Plain at the Savannah River Plant, South Carolina [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 128, 1964.

- MARINE, I. W. --Continued
 2-64. (and PRICE, Don) Geology and ground-water resources of the Jordan Valley, Utah: Utah Geol. and Mineralog. Survey Water-Resources Bull. 7, 68 p., 1964.
- MARTIN, R. O. R.
 1-64. Use of precipitation records in the correlation of streamflow records: Internat. Assoc. Sci. Hydrology Bull., v. 9, no. 4, p. 24-31, 1964.
- MARTINSSON, Anders
 1-65. (and BERDAN, J. M.) The Beyrichiacean ostracode genus *Drepanellina*: Geol. Foren, Stockholm Forh., v. 86, p. 395-403, 1965.
- MARVIN, R. F.
 1-65. (WRIGHT, J. C., and WALTHALL, F. G.) K-Ar and Rb-Sr ages of biotite from the Middle Jurassic part of the Carmel Formation, Utah, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B104-B107, 1965.
- MARZOLF, J. E.
 1-65. Stratigraphy of the Mobil Oil Company Paleozoic test well No. 22-19-G, Tip-top unit, Sublette County, Wyoming: U.S. Geol. Survey open-file report, 39 p., 1965.
- MASIMER, G. E.
 1-65. Inclusions in quartz crystals from De Luz, California: Gems and Minerals, no. 329, p. 32-39, 1965.
- MATALAS, N. C.
 1-64. (and JACOBS, Barbara) A correlation procedure for augmenting hydrologic data: U.S. Geol. Survey Prof. Paper 434-E, p. E1-E7, 1964.
- MATTHES, F. E.
 1-65. Glacial reconnaissance of Sequoia National Park, California: U.S. Geol. Survey Prof. Paper 504-A, p. A1-A58, 1965.
- MATTICK, R. E.
 1-65. Aeromagnetic and generalized geologic map of the Bingham quadrangle, Somerset County, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-499, 1965.
- MATTSON, P. H.
 1-64. Petrography and structure of serpentinite from Mayagüez, Puerto Rico, in A study of serpentinite: Natl. Acad. Sci. --Natl. Research Council Pub. 1188, p. 7-24, 1964.
 1-65. (and NELSON, A. E.) Chemical variations in a volcanic rock suite, Puerto Rico [abs.]: Caribbean Geol. Conf., 4th, Trinidad (mimeographed), 1965.
- MAUGHAN, E. K.
 1-65. Freezeout and Ervay Members of the Goose Egg Formation in eastern Wyoming [abs.]: Geol. Soc. America Spec. Paper 82, p. 338, 1965.
- MAXWELL, C. H.
 1-64. (and TURNER, W. B.) Geology of the Cane Valley quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-369, 1964.
 1-65. Geology of the Dunnville quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-367, 1965.
 2-65. Geology of the Phil quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-395, 1965.
- MAY, Irving
 1-64. (DINNIN, J. I., and ROSENBAUM, Fred) A simple oxygen sheath for flame photometry, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C152-C153, 1964.
 1-65. (and ROWE, J. J.) Solution of refractory minerals in acids at high temperatures and pressures [abs.]: Geol. Soc. America Spec. Paper 82, p. 129, 1965.
 2-65. (ROWE, J. J., and LETNER, Raymond) A platinum-lined bomb for the high-temperature decomposition of refractory minerals, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B165-B166, 1965.
- MAY, J. R.
 1-64. (YANCHOSEK, J. J., and JEFFERY, H. G.) Chemical analyses of water from selected wells in the Arkansas River valley from the mouth to Fort Smith, Arkansas: U.S. Geol. Survey open-file report, 55 p., 1964.
 1-65. (YANCHOSEK, J. J., BEDINGER, M. S., and EMMETT, L. F.) Depth to water measurements in wells in the alluvium of the Arkansas River valley between Little Rock, Arkansas, and the Mississippi River—V. I, Arkansas County; V. II, Desha County; V. III, Jefferson and Lincoln Counties; and V. IV, Lonoke, Pulaski, and Saline Counties: U.S. Geol. Survey open-file report, 197 p., 1965.
 2-65. (YANCHOSEK, J. J., BEDINGER, M. S., and EMMETT, L. F.) Logs of selected test holes and wells in the alluvium of the Arkansas River valley between Little Rock, Arkansas, and the Mississippi River—V. I, Arkansas County; V. II, Desha County; V. III, Jefferson County; V. IV, Lincoln County; and V. V, Lonoke, Pulaski, and Saline Counties: U.S. Geol. Survey open-file report, 365 p., 1965.
- MAY, V. J.
 1-64. (and ALLEN, H. E.) Floods in Elgin quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 13 p., 1964.
 2-64. (and ALLEN, H. E.) Floods in Streamwood quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 7 p., 1964.
 3-64. (and ALLEN, H. E.) Floods in Wheaton quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 9 p., 1964.
- MAYES, J. L.
 1-64. (and CULBERTSON, Don) Chemical quality of surface waters in Kansas—1962 water year: Kansas State Dept. Health, Environmental Health Services Bull. 1-4, 61 p., 1964.
- MEADE, R. H.
 1-64. Removal of water and rearrangement of particles during the compaction of clayey sediments—review: U.S. Geol. Survey Prof. Paper 497-B, p. B1-B23, 1964.
- MEIER, M. F.
 1-64. Ice and glaciers, in Handbook of applied hydrology: New York, McGraw-Hill Book Co., p. 16-1—16-32, 1964.
 2-64. The recent history of advance-retreat and net budget of South Cascade Glacier [abs.]: Am. Geophys. Union Trans., v. 45, no. 4, p. 608, 1964.
- MEKLE, R. L.
 1-65. (and BAKER, J. A.) Ground-water levels in Connecticut, 1960-64: U.S. Geol. Survey open-file report, 25 p., 1965.
- MEINZER, O. E.
 1-65. Outline of ground-water hydrology, with definitions: U.S. Geol. Survey Water-Supply Paper 494, 71 p., 1965. (Originally published 1923.)
 2-65. The occurrence of groundwater in the United States, with a discussion of principles: U.S. Geol. Survey Water-Supply Paper no. 489, 321 p., 1965. (Originally pub. 1923.)
- MENKE, C. G.
 1-64. (MERIDITH, E. W., and WETTERHALL, W. S.) Water-resources records of Hillsborough County, Florida: Florida Geol. Survey Inf. Circ. no. 44, 95 p., 1964.
- MERRIAM, C. W.
 1-64. Cambrian rocks of the Pioche mining district, Nevada: U.S. Geol. Survey Prof. Paper 469, 59 p., 1964.
- MESNIER, G. N.
 1-64. (and ISERI, K. T.) Selected techniques in water resources investigations: U.S. Geol. Survey Water-Supply Paper 1669-Z, p. Z1-Z64, 1964. (Originally published 1963.)
- MEUSCHKE, J. L.
 1-64. (and VARGO, J. L.) Aeromagnetic map of the Amity quadrangle, Aroostook County, Maine: U.S. Geol. Survey Geophys. Inv. Map GP-487, 1964.

- MEUSCHKE, J. L.
1-65. (and LONG, C. L.) Aeromagnetic map of part of the Lanes Creek quadrangle, Caribou County, Idaho: U.S. Geol. Survey Geophys. Inv. Map GP-490, 1965.
- MEYER, G. L.
1-63. (and THORDARSON, William) Hydrology of the Pahute Mesa Exploratory Hole, Nye County, Nevada: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-4, 11 p., 1963.
- MIESCH, A. T.
1-63. (and DAVIS, C. W.) Computer analysis of areal microphotometer data from lunar photographs, *in* Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 85-114, 1963.
- MILLER, C. H.
1-63. (and HEALEY, D. L.) Configuration of the buried surface on pre-Tertiary rocks for an area immediately south of the Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-39, 11 p., 1963.
- MILLER, E. G.
1-65. Effect of Great Swamp, New Jersey, on streamflow during base-flow periods, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B177-B179, 1965.
- MILLER, G. A.
1-64. (and EVENSON, R. E.) Utilization of ground water in the Santa Maria Valley area, California: U.S. Geol. Survey open-file report, 67 p., 1964.
1-65. Ground-water conditions, U.S. Naval Missile Facility, Point Arguello, California, July 1963-June 1964: U.S. Geol. Survey open-file report, 20 p., 1965.
- MILLER, R. D.
1-64. Geology of the Omaha-Council Bluffs area, Nebraska-Iowa: U.S. Geol. Survey Prof. Paper 472, 70 p., 1964[1965].
2-64. (VAN HORN, Richard, DOBROVOLNY, Ernest, and BUCK, L. P.) Geology of Franklin, Webster, and Nuckolls Counties, Nebraska: U.S. Geol. Survey Bull. 1165, 91 p., 1964[1965].
- MILLER, R. L.
1-65. Pennsylvanian System of southwest Virginia [abs.]: Geol. Soc. America Spec. Paper 82, p. 306, 1965.
2-65. The mineral position of Guatemala in 1961, with special reference to industrial minerals: U.S. Geol. Survey open-file report, 40 p., 1965.
- MILLER, W. T.
1-64. Use of digital recorders and application of computers to processing of water data: U.S. Geol. Survey open-file report, 11 p., 1964.
- MILLS, W. B.
1-64. (McGILL, H. N., and FLUGARTH, M. W.) Hydrologic studies of small watersheds, Deep Creek, Colorado River basin, Texas, 1951-61: U.S. Geol. Survey open-file report, 164 p., 1964.
1-65. (and RAWSON, Jack) Base-flow studies, Lampasas River, Texas—Quantity and quality, June 3-6, 1963: Texas Water Comm. Bull. 6506, 16 p., 1965.
- MILTON, Charles
1-65. (INGRAM, Blanche, CLARK, J. R., and DWORNIK, E. J.) McKelveyite, a new hydrous sodium barium rare-earth uranium carbonate mineral from the Green River Formation, Wyoming [abs.]: Geol. Soc. America Spec. Paper 82, p. 133-134, 1965.
- MILTON, D. J.
1-63. The Campo del Cielo Meteorite Crater Field, Argentina, *in* Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 91-97, 1963.
1-64. Fused rock from Köfels, Tyrol: *Tschermaks Mineralog. Petrog. Mitt.*, v. 9, no. 12, 3d ser., p. 86-94, 1964.
- MILTON, D. J. --Continued
2-65. Structure of Henbury meteorite craters, Australia [abs.]: Geol. Soc. America Spec. Paper 82, p. 266, 1965.
- MINARD, J. P.
1-64. Geology of the Roosevelt quadrangle, New Jersey: U.S. Geol. Survey Geol. Quad. Map GQ-340, 1964.
1-65. Geologic map of the Woodstown quadrangle, Gloucester and Salem Counties, New Jersey: U.S. Geol. Survey Geol. Quad. Map GQ-404, 1965.
- MISER, H. D.
1-64. (and MILTON, Charles) Quartz, rectorite and cook-eite from the Jeffrey Quarry, near North Little Rock, Pulaski County, Arkansas: Arkansas Geol. Commission Bull. 21, 29 p., 1964.
- MITCHELL, C. M.
1-65. (KNOWLES, F. F., and PETRAFESO, F. A.) Aeromagnetic map of the Pocatello-Soda Springs area, Bannock and Caribou Counties, Idaho: U.S. Geol. Survey Geophys. Inv. Map GP-521, 1965.
2-65. (and ZANDLE, G. L.) Aeromagnetic map of the Casa Grande area, Maricopa and Pinal Counties, Arizona: U.S. Geol. Survey open-file report, 1965.
- MITTIN, Ivan
1-65. Translation of "The study of some hydrothermal and sedimentary heulandites," by N. V. Longvineko, G. I. Kulesko, and S. I. Shumenko: *Geochemistry Internat.*, v. 1964, no. 1, p. 132-190, 1965.
- MIXON, R. B.
1-65. Geology of the Krypton quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-389, 1965.
- MOENCH, R. H.
1-64. Geology of the Dough Mountain quadrangle, New Mexico: U.S. Geol. Survey Geol. Quad. Map GQ-354, 1964.
2-64. Geology of Precambrian rocks, Idaho Springs district, Colorado: U.S. Geol. Survey Bull. 1182-A, p. A1-A70, 1964[1965].
3-64. Geology of the South Butte quadrangle, New Mexico: U.S. Geol. Survey Geol. Quad. Map GQ-355, 1964.
1-65. (SCHLEE, J. S., and BRYAN, W. B.) Geologic map of the La Gotera quadrangle, Sandoval and Valencia Counties, New Mexico: U.S. Geol. Survey Geol. Quad. Map GQ-371, 1965.
- MONROE, W. H.
1-64. General description of cores of pre-Selma Cretaceous strata in western Alabama: U.S. Geol. Survey Bull. 1160-A, p. A1-A8, 1964.
2-64. The origin and interior structure of mogotes [Puerto Rico], *in* Internat. Geog. Cong., 20th, United Kingdom, 1964, Abs. Papers: London, Thomas Nelson and Sons Ltd., p. 108, 1964.
3-64. Lithological control and tropical karst topography [abs.], *in* Internat. Geog. Cong., 20th, United Kingdom, 1964, Abs. Papers: London, Thomas Nelson and Sons Ltd., p. 107-108, 1964.
4-64. (BERGENBACK, R. E., SOHL, N. F., APPLIN, E. R., LEOPOLD, E. B., PAKISER, H. M., and CONANT, L. C.) Studies of pre-Selma Cretaceous core samples from the outcrop area in western Alabama: U.S. Geol. Survey Bull. 1160, 108 p., 1964.
- MOORE, A. M.
1-64. Compilation of water-temperature data for Oregon streams: U.S. Geol. Survey open-file report, 134 p., 1964.
2-64. Correlation and analysis of water-temperature data for Oregon streams, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D185-D189, 1964; *also in* Geol. Survey open-file report, 83 p., 1964.

- MOORE, E. J.
1-63. Review of "Marine molluscan genera of Western North America: an illustrated key," by A. M. Keen: *Am.-Midland Naturalist*, v. 70, no. 2, p. 512, 1963.
- MOORE, F. B.
1-64. Lithologic and radioactivity log of Summit area drill hole, Hardin County, Kentucky: U.S. Geol. Survey open-file report, 1 sheet, 1964.
- MOORE, G. E., Jr.
1-64. Bedrock geology of the Kingston quadrangle, Rhode Island: U.S. Geol. Survey Bull. 1158-E, p. E1-E21, 1964.
- MOORE, G. K.
1-65. (and BINGHAM, R. H.) Availability of ground water in the western Highland Rim of Tennessee: *Tennessee Acad. Sci. Jour.*, v. 40, no. 1, p. 22-26, 1965.
- MOORE, G. W.
1-64. Abrupt change in cave history when ventilation begins [abs.]: *Natl. Speleol. Soc. Bull.*, v. 26, no. 2, p. 76, 1964.
2-64. Magnetic events which preceded the 1964 Alaska earthquake: *Nature*, v. 302, no. 4944, p. 508-509, 1964.
1-65. Tensional origin of the Aleutian Arc [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 134, 1965.
- MOORE, H. J.
1-63. Density of small craters on the lunar surface, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 34-50, 1963.
2-63. The geology of the Aristarchus quadrangle of the Moon, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 33-44, 1963.
3-63. (GAULT, D. E., and HEITOWIT, E. D.) Change in effective target strength with increasing size of hypervelocity impact craters, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 52-61, 1963.
1-65. (GAULT, D. E., and HEITOWIT, E. D.) Change of effective target strength with increasing size of hypervelocity impact craters: Tampa, Florida, Hypervelocity Impact Symposium, 7th, 1964, *Proc.*, v. 4, p. 35-45, 1965.
- MOORE, J. G.
1-64. Giant submarine landslides on the Hawaiian Ridge, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D95-D98, 1964.
2-64. The 1962 flank eruption of Kilauea: U.S. Geol. Survey Hawaiian Volcano Observatory Summ. 28, p. 1-5, 1964.
1-65. Petrology of deep-sea basalt near Hawaii: *Am. Jour. Sci.*, v. 263, no. 1, p. 40-52, 1965.
2-65. Properties of Hawaiian submarine basalt [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 267, 1965.
3-65. (and AULT, W. U.) Historic littoral cones in Hawaii: *Pacific Science*, v. 19, no. 1, p. 3-11, 1965.
- MOORE, L. C.
1-65. Cartographic scribing materials, instruments, and techniques: *Am. Cong. Surveying and Mapping, Cartographic Div.*, Tech. Pub. no. 3, 1965.
- MOORE, R. H.
1-64. (and BRYAN, Wendell) The practical application of research on visual factors in stereoplotting: *Photogramm. Eng.*, v. 30, no. 6, p. 991-999, 1020, 1964.
- MOORE, S. L.
1-64. Geology of the Freedom quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-349, 1964.
1-65. (and MILLER, R. C.) Geology of the Glasgow South quadrangle, Barren County, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-416, 1965.
- MOORE, S. L. --Continued
2-65. (and MILLER, R. C.) Geology of the Temple Hill quadrangle, Barren County, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-402, 1965.
- MOREY, G. W.
1-64. Phase-equilibrium relations of the common rock-forming oxides except water: U.S. Geol. Survey Prof. Paper 440-L, p. L1-L159, 1964.
- MORGAN, J. H.
1-64. Availability of ground water in the Elva quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-117, 1964[1965].
2-64. Availability of ground water in the Farmington quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-92, 1964.
3-64. Availability of ground water in the Kirksey quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-113, 1964.
4-64. Availability of ground water in the Oak Level quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-116, 1964.
5-64. Pumping test of an Eocene aquifer near Mayfield, Kentucky: U.S. Geol. Survey open-file report, 17 p., 1964.
1-65. Availability of groundwater in the Hico quadrangle, Kentucky: U.S. Geol. Survey Hydrol. Inv. Map HA-158, 1965.
2-65. Availability of groundwater in the parts of the Little Cypress and Calvert City quadrangles in the Jackson Purchase region, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-155, 1965.
- MORRIS, E. C.
1-63. Stratigraphic relations in the vicinity of the crater Julius Caesar, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 31-32, 1963.
2-63. (RING, J., and STEPHENS, H. G.) Photographic and photoelectric investigations of the Earth-Moon libration regions L₄ and L₅ from Mt. Chacaltaya, Bolivia, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 71-74, 1963.
- MORRIS, H. T.
1-64. Nonmetallic and industrial minerals and materials resources: Limestone and dolomite, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 188-194, 1964.
2-64. (and SHEPARD, W. M.) Evidence for a concealed tear fault with large displacement in the central East Tintic Mountains, Utah, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C19-C21, 1964.
- MORRIS, R. H.
1-65. Geology of the Burtonville quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-396, 1965.
2-65. Geology of the Charters quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-293, 1965.
- MORSE, E. K.
1-64. (STULIK, R. S., and SMITH, C. R.) Use of ground water, in *Annual report on groundwater in Arizona, Spring 1963 to Spring 1964*: Arizona Land Dept. Water Resources Rept. 19, p. 57-60, 1964.
- MOSTON, R. P.
1-64. (and JOHNSON, A. I.) Ultrasonic dispersion of samples of sedimentary deposits, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C159-C160, 1964.
- MOTTS, W. S.
1-64. (and CUSHMAN, R. L.) An appraisal of the possibilities of artificial recharge to ground-water supplies

- MOTTS, W. S. --Continued
in part of the Roswell basin, New Mexico: U.S. Geol. Survey Water-Supply Paper 1785, 85 p., 1964.
- MOULDER, E. A.
1-64. (and JENKINS, C. T.) Methods for controlling the ground-water regime, exploitation and conservation: Internat. Assoc. Sci. Hydrology Pub. 64, p. 329-342, 1964.
- MOUNT, Priscilla
1-64. Metallic mineral resources: Bismuth, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 75-78, 1964.
2-64. Nonmetallic and industrial minerals and materials resources: Sulfur, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 228-232, 1964.
- MOURANT, W. A.
1-65. New Mexico State Planning Office Report—Pecos River basin: U.S. Geol. Survey open-file report, 32 p., 1965.
- MOWER, R. W.
1-64. (and FELTIS, R. D.) Ground-water data, Sevier Desert, Utah: Utah State Engineer Basic Data Rept. 9, 34 p., 1964.
- MOXHAM, R. M.
1-64. Radioelement dispersion in a sedimentary environment and its effect on uranium exploration: Econ. Geology, v. 59, no. 2, p. 309-321, 1964.
2-64. Some aerial observations on the terrestrial component of environmental α radiation, *in* The natural radiation environment: Chicago, Illinois, Univ. Chicago Press (for Rice Univ.), p. 737-746, 1964.
1-65. New developments in geophysics: Am. Mining Cong. Jour., v. 51, no. 1, p. 50-53, 1965.
2-65. (FOOTE, R. S., and BUNKER, C. M.) Gamma-ray spectrometer studies of hydrothermally altered rocks: Econ. Geology, v. 60, no. 4, p. 653-671, 1965.
- MOYLE, W. R., Jr.
1-64. Data on water wells in the western part of the Antelope Valley area, Los Angeles and Kern Counties, California: U.S. Geol. Survey open-file report, 29 p., 1964.
- MROSE, M. E.
1-65. New specific refractive energy values for CuO and Sc_2O_3 [abs.]: Am. Mineralogist, v. 50, nos. 1-2, p. 288, 1965.
- MUDGE, M. R.
1-65. Bedrock geologic map of the Sawtooth Ridge quadrangle, Teton and Lewis and Clark Counties, Montana: U.S. Geol. Survey Geol. Quad. Map GQ-381, 1965.
- MUFFLER, L. J. P.
1-64. Geology of the Frenchie Creek quadrangle, north-central Nevada: U.S. Geol. Survey Bull. 1179, 99 p., 1964[1965].
- MUIR, K. S.
1-64. Water levels in observation wells in Santa Barbara County, California, 1963: U.S. Geol. Survey open-file report, 30 p., 1964.
- MULL, D. S.
1-65. Ground-water resources of the Jenkins-Whitesburg area, Kentucky: U.S. Geol. Survey Water-Supply Paper 1809-A, p. A1-A36, 1965.
- MULLENS, T. E.
1-64. Geology of the Cuba City, New Diggings, and Shullsburg quadrangles, Wisconsin and Illinois: U.S. Geol. Survey Bull. 1123-H, p. H437-H531, 1964[1965].
2-64. (and IZETT, G. A.) Geology of the Paradise quadrangle, Cache County, Utah: U.S. Geol. Survey Bull. 1181-S, p. S1-S32, 1964.
- MULLINEAUX, D. R.
1-64. (NICHOLS, T. C., and SPEIRER, R. A.) A zone of montmorillonitic weathered clay in Pleistocene deposits at Seattle, Washington, *in* Geological Survey Research
- MULLINEAUX, D. R. --Continued
1964: U.S. Geol. Survey Prof. Paper 501-D, p. D99-D103, 1964.
1-65. Geologic map of the Auburn quadrangle, King and Pierce Counties, Washington: U.S. Geol. Survey Geol. Quad. Map GQ-406, 1965.
2-65. Geologic map of the Black Diamond quadrangle, King County, Washington: U.S. Geol. Survey Geol. Quad. Map GQ-407, 1965.
3-65. Geologic map of the Renton quadrangle, King County, Washington: U.S. Geol. Survey Geol. Quad. Map GQ-405, 1965.
4-65. (WALDRON, H. H., and RUBIN, Meyer) Stratigraphy and chronology of late interglacial and early Vashon glacial time in Seattle area, Washington: U.S. Geol. Survey Bull. 1194-O, p. O1-O10, 1965.
- MUNDORFF, J. C.
1-64. Fluvial sediment in Kiowa Creek basin, Colorado: U.S. Geol. Survey Water-Supply Paper 1798-A, p. A1-A70, 1964.
2-64. (CROSTHWAITE, E. G., and KILBURN, Chabot) Ground water for irrigation in the Snake River basin in Idaho: U.S. Geol. Survey Water-Supply Paper 1654, 224 p., 1964.
3-64. (and SCOTT, C. H.) Fluvial sediment in the lower Kansas River basin, 1957-60—A progress report: Kansas Water Resources Board Bull. 7, 67 p., 1964.
- MURATA, K. J.
1-65. (and ERD, R. C.) Composition of sediments from the experimental Mohole Project (Guadalupe Site): Geol. Soc. America Spec. Paper 82, p. 136, 1965.
- MUSGROVE, R. H.
1-65. (BARRACLOUGH, J. T., and GRANTHAM, R. G.) The Pensacola area's water: Florida Geol. Survey Leaflet 3, 14 p., 1965.
2-65. (BARRACLOUGH, J. T., and GRANTHAM, R. G.) Water resources of Escambia and Santa Rosa Counties, Florida: Florida Geol. Survey Rept. Inv. 40, 102 p., 1965.
- MYER, G. H.
1-65. X-ray determinative curve for epidote: Am. Jour. Sci., v. 263, no. 1, p. 78-86, 1965.
- MYERS, B. M.
1-65. (and DALE, O. C.) Ground-water resources of Bee County, Texas: U.S. Geol. Survey open-file report, 165 p., 1965.
- MYERS, D. A.
1-65. Geology of the Wayland quadrangle, Stephens and Eastland Counties, Texas: U.S. Geol. Survey Bull. 1201-C, C1-C63, 1965.
2-65. Rhythmic sedimentation in upper part of Madera Limestone, northern Manzano Mountains, New Mexico [abs.]: Am. Assoc. Petroleum Geologists Bull., v. 49, no. 3, p. 353-354, 1965.
- MYERS, W. B.
1-64. Geology of the Petroleum quadrangle, Kentucky-Tennessee: U.S. Geol. Survey Geol. Quad. Map GQ-352, 1964.
- NACE, R. L.
1-64. The International Hydrological Decade: Am. Geophys. Union Trans., v. 45, no. 3, p. 413-421, 1964.
2-64. Water—essential factor of economic development: Impact, v. 24, no. 1, p. 39-55, 1964.
1-65. Global thirst and the International Hydrological Decade: Sci. Teacher, v. 32, no. 1, p. 13-16, 1965.
2-65. Water—a common problem: Bull. Atomic Scientists, p. 32-34, 1965.
3-65. Water management, agriculture, and ground-water supplies: U.S. Geol. Survey Circ. 415, 12 p., 1965. (Originally published 1960.)

- NAKAGAWA, H. M.
1-65. (and LAKIN, H. W.) A field method for the determination of silver in soils and rocks: U.S. Geol. Survey open-file report, 13 p., 1965.
- NASSAR, E. G.
1-65. Effects of wind and atmospheric pressure on storage measurements at Roosevelt Lake, Washington: U.S. Geol. Survey open-file report, 33 p., 1965.
- NEELY, B. L., Jr.
1-64. Floods of 1962 in Mississippi: Mississippi Board Water Comm. Bull. 64-6, 97 p., 1964.
- NELSON, A. E.
1-65. Trends of Cretaceous volcanism in north-central Puerto Rico [abs.]: Caribbean Geol. Conf., 4th, Trinidad, 1965, (mimeographed), 1965.
- NELSON, W. H.
1-64. Geology of the Pleasant Green Hill quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-321, 1964.
- NEUMAN, R. B.
1-64. Fossils in Ordovician tuffs, northeastern Maine, with a section on Trilobites, by H. B. Whittington: U.S. Geol. Survey Bull. 1181-E, p. E1-E38, 1964.
1-65. Collecting in metamorphic rocks, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 159-163, 1965.
- NEWCOME, Roy, Jr.
1-64. (and CALLAHAN, J. A.) Water for industry in the Corinth area, Mississippi: Mississippi State Board Water Comm. Bull. 64-2, 24 p., 1964.
2-64. (and GOLDEN, H. G.) Status of water resources in Jackson County, Mississippi: Mississippi State Board of Water Comm. Bull. 64-3, 17 p., 1964.
- NICHOLAS, G.
1-64. (and MOORE, G. W.) Out of phase seasonal temperature fluctuations in Cathedral Cave, Kentucky [abs.]: Natl. Speleol. Soc. Bull., v. 26, no. 2, p. 84, 1964.
- NOBLE, D. C.
1-64. (and McKAY, E. J.) Geology of the Deadhorse Flat quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-78, 11 p., 1964.
1-65. A rapid conoscopic method for measurement of 2V on the spindle stage: Am. Mineralogist, v. 50, nos. 1-2, p. 180-186, 1965.
2-65. Determination of composition and structural stage of plagioclase with the five-axis universal stage: Am. Mineralogist, v. 50, nos. 3-4, p. 367-381, 1965.
3-65. Gold Flat Member of the Thirsty Canyon Tuff—A pantellerite ash-flow sheet in southern Nevada, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B85-B90, 1965.
4-65. Ground-water leaching of sodium from quickly cooled volcanic rocks [abs.]: Am. Mineralogist, v. 50, nos. 1-2, p. 289, 1965.
5-65. (KISTLER, R. W., CHRISTIANSEN, R. L., LIPMAN, P. W., and POOLE, F. G.) Close association in space and time of alkalic, calc-alkalic, and calcic volcanism in southern Nevada: Geol. Soc. America Spec. Paper 82, p. 143-144, 1965.
- NOEHRE, A. W.
1-64. Floods in Wadsworth quadrangle, Illinois-Wisconsin: U.S. Geol. Survey Hydrol. Inv. Atlas HA-144, 1964[1965].
2-64. (and WALTER, G. L.) Floods in Romeoville quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 13 p., 1964.
3-64. (and WALTER, G. L.) Floods in Sag Bridge quadrangle, Illinois: U.S. Geol. Survey open-file report, 10 p., 1964.
4-64. (MAY, V. J., and WALTER, G. L.) Floods in Fox Lake quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 5 p., 1964.
1-65. Floods in Mokena quadrangle, northeastern Illinois: U.S. Geol. Survey open-file report, 11 p., 1965.
- NOEHRE, A. W. --Continued
2-65. (and WALTER, G. L.) Floods in Geneva quadrangle, Illinois: U.S. Geol. Survey Hydrol. Inv. Atlas HA-142, 1965.
- NOLAN, T. B.
1-64. Annual report of the Director, U.S. Geological Survey, to the Secretary of the Interior: Reprinted from the Annual Report of the Secretary of the Interior, for the fiscal year ended June 30, 1963: Washington, D. C., U.S. Dept. of Interior, p. 355-400, 1964.
- NORDIN, C. F., Jr.
1-64. Aspects of flow resistance and sediment transport, Rio Grande near Bernalillo, New Mexico: U.S. Geol. Survey Water-Supply Paper 1498-H, 41 p., 1964.
2-64. Discussion of "Practical computations of bed-material discharge," by B. R. Colby: Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour., v. 90, no. HY6, p. 309-313, 1964.
3-64. Discussion of "Sediment investigations—Middle Rio Grande," by E. L. Pemberton: Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour., v. 90, no. HY5, p. 273-175, 1964.
4-64. Study of channel erosion and sediment transport: Am. Soc. Civil Engineers Proc., v. 90, paper 3984, Jour. Hydraulics Div., no. HY 4, pt. 1, p. 173-192, 1964.
1-65. (and BEVERAGE, J. P.) Sediment transport in the Rio Grande, New Mexico: U.S. Geol. Survey Prof. Paper 462-F, p. F1-F35, 1965.
- NORTON, J. J.
1-64. Nonmetallic and industrial mineral resources: Pegmatite minerals, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 112-133, 1964.
- NORVITCH, R. F.
1-63. (and SMITH, R. E.) Yield of and quality of water from test well drilled at the Aurora site, Clark County, Nevada: U.S. Geol. Survey Tech. Letter NTS-49, 11 p., 1963.
1-64. Geology and ground-water resources of Nobles County and parts of Jackson County, Minnesota: U.S. Geol. Survey Water-Supply Paper 1749, 70 p., 1964.
- NYMAN, D. J.
1-65. Predicted hydrologic effects of pumping from the new Lichterman well field in the Memphis area, Tennessee: U.S. Geol. Survey open-file report, 48 p., 1965.
- O'CONNOR, J. T.
1-65. (ANDERSON, R. E., and LIPMAN, P. W.) Geologic map of the Thirsty Canyon quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report TEI-861, 1965.
2-65. A classification for quartz-rich igneous rocks based on feldspar ratios, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B79-B84, 1965.
- ODELL, J. W.
1-64. (COFFIN, D. L., and LANGFORD, R. H.) Water resources, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 233-283, 1964.
- OGATA, Akio
1-64. The spread of a dye stream in an isotropic granular medium: U.S. Geol. Survey Prof. Paper 411-G, p. G1-G11, 1964.
2-64. Mathematics of dispersion with linear adsorption isotherm: U.S. Geol. Survey Prof. Paper 411-H, H1-H9, 1964.
- OKAMURA, A. T.
1-65. (KOYANAGI, R. Y., and KINOSHITA, W. T.) Hawaiian Volcano Observatory summary, April, May, and June 1964: U.S. Geol. Survey Hawaiian Volcano Observatory Summary 34, 47 p., 1965.

- OLCOTT, P. G.
1-64. Geology and water resources of Winnebago County, Wisconsin: U.S. Geol. Survey open-file report, 122 p., 1964.
- OLDALE, R. N.
1-64. Surficial geology of the Salem quadrangle, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-271, 1964.
1-65. Subsurface geology of Cape Cod, Massachusetts [abs.]: Geol. Soc. America Spec. Paper 82, p. 144, 1965.
- OLIVE, W. W.
1-65. Geology of the Hico quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-332, 1965.
- OLIVER, H. W.
1-65. The U.S. Geological Survey's gravity program in California, Hawaii, Nevada, and Oregon: Am. Geophys. Union Trans., v. 46, no. 1, p. 219-222, 1965.
- OLIVER, W. A., Jr.
1-65. Corals, as illustrated by Paleozoic forms, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Co., p. 36-40, 1965.
- OLSEN, H. W.
1-65. Darcy's law in kaolinite [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 74, 1965.
- OLSON, J. C.
1-64. (and OVERSTREET, W. C.) Geologic distribution and resources of thorium: U.S. Geol. Survey Bull. 1204, 61 p., 1964.
- OLTMAN, R. E.
1-65. What is a river?: Sci. Teacher, v. 32, no. 1, p. 22-25, 1965.
- O'REAR, D. M.
1-64. Ground-water levels in Alabama in 1959 and 1960: Alabama Geol. Survey Circ. 23, 123 p., 1964.
- ORIEL, S. S.
1-65. Bringham, Langston, and Ute Formations in Portneuf Range, southeastern Idaho [abs.]: Geol. Soc. America Spec. Paper 82, p. 341, 1965.
- ORKILD, P. P.
1-63. Geology of Mine Mountain quadrangle: U.S. Geol. Survey Tech. Letter NTS-58, 12 p., 1963.
1-64. Geologic map of the Mine Mountain quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report TEI-854, 1964.
- ORTH, D. J.
1-64. Domestic geographic names activity in Federal mapping: Am. Cong. Surveying and Mapping Quart. Jour., v. 24, no. 4, p. 603-606, 1964.
- OSTERWALD, F. W.
1-64. (and DUNRUD, C. R.) Geology applied to study of coal mine bumps and mining methods at Sunnysdale, Utah: U.S. Geol. Survey open-file report, 8 p., 1964.
1-65. Structural control of uranium-bearing vein deposits and districts in the conterminous United States: U.S. Geol. Survey Prof. Paper 455-G, p. G121-G146, 1965.
- O'SULLIVAN, R. B.
1-65. Geology of the Cedar Mesa-Boundary Butte area, San Juan County, Utah: U.S. Geol. Survey Bull. 1186, 128 p., 1965.
- OUTERBRIDGE, W. F.
1-64. Geology of the Offutt quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-348, 1964[1965].
- OVERSTREET, E. F.
1-64. Geology of the southeastern bauxite deposits: U.S. Geol. Survey Bull. 1199-A, p. A1-A19, 1964.
- OVERSTREET, W. C.
1-65. (and BELL, Henry, III) Geologic map of the crystalline rocks of South Carolina: U.S. Geol. Survey Misc. Geol. Inv. Map I-413, 1965.
2-65. (and BELL, Henry, III) The crystalline rocks of South Carolina: U.S. Geol. Survey Bull. 1183, 126 p., 1965.
- OWENS, J. P.
1-64. (and MINARD, J. P.) Pre-Quaternary geology of the Bristol quadrangle, New Jersey-Pennsylvania: U.S. Geol. Survey Geol. Quad. Map GQ-342, 1964.
2-64. (and MINARD, J. P.) Pre-Quaternary geology of the Trenton East quadrangle, New Jersey-Pennsylvania: U.S. Geol. Survey Geol. Quad. Map GQ-341, 1964.
- PAGE, L. V.
1-64. (and MAY, H. G.) Water resources of Bossier and Caddo Parishes, Louisiana: Louisiana Geol. Survey Water Resources Bull. 5, 105 p., 1964.
2-64. (and PREÉ, H. L., Jr.) Water resources of De Soto Parish, Louisiana: U.S. Geol. Survey Water-Supply Paper 1774, 152 p., 1964[1965].
- PAKISER, L. C.
1-64. Gravity, volcanism, and crustal structure in the southern Cascade Range, California: Geol. Soc. America Bull., v. 75, no. 7, p. 611-620, 1964.
2-64. (KANE, M. F., and JACKSON, W. H.) Structural geology and volcanism of Owens Valley region, California—A geophysical study: U.S. Geol. Survey Prof. Paper 438, 68 p., 1964.
3-64. (and STEINHART, J. S.) Explosion seismology in the western hemisphere, Chap. 5, *in* Research in geophysics—V. 2. Solid earth and interface phenomena: Cambridge, Mass., The MIT Press, p. 123-147, 1964.
1-65. The basalt-eclogite transformation and crustal structure in the Western United States, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B1-B8, 1965.
2-65. (and ZIETZ, Isidore) Transcontinental crustal and upper-mantle structure [abs.]: Geol. Soc. America Spec. Paper 82, p. 146-147, 1965.
- PALMER, A. R.
1-64. An unusual Lower Cambrian trilobite fauna from Nevada: U.S. Geol. Survey Prof. Paper 483-F, p. F1-F13, 1964.
2-64. Pioche Shale faunules, *in* Cambrian rocks of the Pioche mining district, Nevada: U.S. Geol. Survey Prof. Paper 469, p. 25-27, 1964.
1-65. Evolutionary patterns in North American Cambrian trilobites [abs.]: Geol. Soc. America Spec. Paper 82, p. 147, 1965.
2-65. Preparation of plates for paleontologic publications, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Co., p. 456-459, 1965.
3-65. Trilobites of the Late Cambrian Pteroccephaliid bioterm in the Great Basin, United States: U.S. Geol. Survey Prof. Paper 493, 105 p., 1965.
- PANGBORN, M. W., Jr.
1-64. Review of "Earthquakes and earthquake structure," by J. H. Hodgson: Prof. Geographer, v. 16, no. 5, p. 36, 1964.
- PAPIKE, J. J.
1-65. (and ZOLTAI, Tibor) The crystal structure of a marialite scapolite: Am. Mineralogist, v. 50, May-June, nos. 5-6, p. 641-655, 1965.
- PARKER, G. G.
1-64. Piping, a geomorphic agent in landform development of the drylands: Internat. Assoc. Sci. Hydrology Pub. 65, p. 103-113, 1964.
2-64. (HELY, A. G., KEIGHTON, W. B., OLMSTED, F. H., and others) Water resources of the Delaware River basin: U.S. Geol. Survey Prof. Paper 381, 200 p., 1964[1965].
- PARKER, R. L.
1-64. Mineral resources: Niobium and tantalum, *in* Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 138-143, 1964.
2-64. Miscellaneous metals: Niobium and tantalum, *in* Mineral and water resources of Colorado: Washington, D.C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 124-127, 1964.

PARKER, R. L. --Continued

3-64. Nonmetallic and industrial minerals and materials resources: Alunite, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 151-154, 1964.

4-64. (and CALKINS, J. A.) Geology of the Curlew quadrangle, Ferry County, Washington: U.S. Geol. Survey Bull. 1169, 95 p., 1964[1965].

1-65. Niobium and tantalum, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 290-294, 1965.

PATTEN, E. P., Jr.

1-64. Uses of electric analog models in ground-water hydrology [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 150, 1964.

PATTERSON, S. H.

1-64. Construction materials: Clays, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 183-190, 1964.

2-64. Halloysite under clay and amorphous inorganic matter in Hawaii, in Clays and clay minerals, 12th Natl. Conf., Atlanta, Georgia, 1963, Proc.: New York, Macmillan Co., p. 153-172, 1964.

3-64. Nonmetallic and industrial minerals and materials resources: Clays, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 157-162, 1964.

4-64. (and COX, E. J.) Nonmetallic and industrial mineral resources: Clays, bentonite, and lightweight aggregate, in Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 105-112, 1964.

1-65. Diatomite, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 322-324, 1965.

2-65. (and HOLMES, R. W.) Clays, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 312-322, 1965.

PATTON, W. W., Jr.

1-64. Mineral fuel resources: Northern and interior Alaska, in Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 62-75, 1964.

2-64. (and TAILLEUR, I. L.) Geology of the Killik-Itkillik region, Alaska: U.S. Geol. Survey Prof. Paper 303-G, p. 409-500, 1964.

PAULSON, Q. F.

1-64. Geologic factors affecting discharge of the Sheyenne River in southeastern North Dakota, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D177-D181, 1964.

2-64. North Dakota's ground-water resources, in The mineral resources of North Dakota: Grand Forks, Univ. North Dakota, General Ext. Div., p. 53-59, 1964.

3-64. (and AKIN, P. D.) Ground-water resources of the Devils Lake area, Benson, Ramsey, and Eddy Counties, North Dakota: North Dakota State Water Conserv. Comm., North Dakota Ground-Water Studies no. 56, 211 p., 1964.

1-65. (and ZIMMERMAN, T. V.) Geology and ground-water resources of the Two Medicine unit and adjacent areas, Blackfeet Indian Reservation, Montana; with a section on chemical quality of water by R. R. Langford: U.S. Geol. Survey open-file report, 126 p., 1965.

PAUSZEK, F. H.

1-64. (and RUGGLES, F. H.) Effects of waste water from AEC plant on the hydrology of Glowegee Creek at West Milton, New York, 1958-61: U.S. Geol. Survey open-file report, 83 p., 1964.

1-65. (and EDMONDS, R. J.) A preliminary appraisal of water quality in the Housatonic River Basin in Connecticut: Connecticut Water Resources Comm. Bull. 6, 70 p., 1965.

PAUSZEK, F. H. --Continued

2-65. (and RUGGLES, F. H.) Effects of waste water from A. E. C. plant on the hydrology of Glowegee Creek at West Milton, New York, 1958-61: U.S. Geol. Survey Water-Supply Paper 1809-N, p. N1-N29, 1965.

PAVLIDES, Louis

1-64. The Hovey Group of northeastern Maine: U.S. Geol. Survey Bull. 1194-B, p. B1-B6, 1964.

2-64. (MENCHER, Ely, NAYLOR, R. S., and BOUCOT, A. J.) Outline of the stratigraphic and tectonic features of northeastern Maine, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C28-C38, 1964.

PEACE, R. R., Jr.

1-64. Geology and ground-water resources of the Russellville area, Alabama: Alabama Geol. Survey Bull. 77, 83 p., 1964.

PECK, D. L.

1-64. Preliminary geologic map of the Merced Peak quadrangle, California: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-281, 1964.

2-64. (GRIGGS, A. B., SCHLICKER, H. G., WELLS, F. G., and DOLE, H. M.) Geology of the central and northern parts of the western Cascade Range in Oregon: U.S. Geol. Survey Prof. Paper 449, 56 p., 1964.

3-64. (MOORE, J. G., and KOJIMA, George) Temperatures in the crust and melt of Alae Lava Lake, Hawaii, after the August 1963 eruption of Kilauea Volcano—A preliminary report, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D1-D7, 1964.

PEIRCE, L. B.

1-64. Reservoir temperatures in north-central Alabama: Alabama Geol. Survey Bull. 82, 103 p., 1964.

PERLMUTTER, N. M.

1-64. (and LIEBER, Maxim, and FRAUENTHAL, H. L.) Contamination of ground water by detergents in a suburban environment—south Farmingdale area, Long Island, New York, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C170-C175, 1964.

1-65. (and TODD, Ruth) Correlation and Foraminifera of the Monmouth Group (Upper Cretaceous), Long Island, New York: U.S. Geol. Survey Prof. Paper 483-I, p. 11-124, 1965.

PERRY, W. J.

1-65. (and MOORE, G. K.) Availability of water for industry in Montgomery County: Tennessee Div. Water Resources Hydrol. Atlas 4, 16 p., 1965.

PESSL, Fred, Jr.

1-63. Emerged marine features in the Mesters Vig area, East Greenland [abs.]: Coastal Research Notes, no. 5, p. 8-9, 1963.

1-65. Surficial geologic map of the Fitchville quadrangle, New London County, Connecticut: U.S. Geol. Survey open-file report, 1965.

PETERSON, D. E.

1-65. Bouguer anomaly map and principal facts for gravity stations for the Central Arizona project: U.S. Geol. Survey open-file report, 2 p., 57 data sheets, 5 maps, 1965.

PETTIJOHN, F. J.

1-64. Chemical composition of sandstones—excluding carbonate and volcanic sands, in Data of geochemistry, 6th ed.: U.S. Geol. Survey Prof. Paper 440-S, p. S1-S21, 1964. (Originally published 1963.)

PETTY, A. J.

1-65. (PETRAFESO, F. A., and MOORE, F. C., Jr.) Aeromagnetic map of the Savannah River Plant area, South Carolina and Georgia: U.S. Geol. Survey Geophys. Inv. Map GP-489, 1965.

PETTYJOHN, W. A.

1-65. (and HILLS, D. L.) Geohydrology of the Souris River valley in the vicinity of Minot, North Dakota—Ground-

- PETTYJOHN, W. A. --Continued
water basic data: U.S. Geol. Survey open-file report, 108 p., 1965.
- PÉWÉ, T. L.
1-64. (and HOLMES, G. W.) Geology of the Mt. Hayes D-4 quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-394, 1964.
- PHAIR, George
1-64. (and GOTTFRIED, David) The Colorado Front Range, Colorado, U.S.A., as a uranium and thorium province, *in* The natural radiation environment: Chicago, Illinois, Univ. Chicago Press (for Rice Univ.), p. 7-38, 1964.
- PHIBBS, E. J., Jr.
1-64. Chemical and physical character of surface waters of North Carolina, 1961-62: North Carolina Dept. Water Resources, Div. Stream Sanitation and Hydrol. Bull. 1, v. 6, 218 p., 1964.
1-65. Chemical and physical character of municipal water supplies in North Carolina: North Carolina Div. Stream Sanitation and Hydrology Bull. 2, Supp. 3, 48 p., 1965.
- PHILBIN, P. W.
1-64. (and KIRBY, J. R.) Aeromagnetic map of the Nyack quadrangle and part of the White Plains quadrangle, Bergen County, New Jersey, and Rockland and Westchester Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-493, 1964.
2-64. (and KIRBY, J. R.) Aeromagnetic map of the Park Ridge quadrangle, Bergen County, New Jersey and Rockland County, New York: U.S. Geol. Survey Geophys. Inv. Map GP-494, 1964.
3-64. (and KIRBY, J. R.) Aeromagnetic map of parts of the Hackensack and Paterson quadrangles, Bergen and Passaic Counties, New Jersey: U.S. Geol. Survey Geophys. Inv. Map GP-492, 1964.
4-64. (and KIRBY, J. R.) Aeromagnetic map of parts of the Yonkers and Mount Vernon quadrangles, Bergen County, New Jersey, and Bronx, Rockland, and Westchester Counties, New York: U.S. Geol. Survey Geophys. Inv. Map GP-495, 1964.
5-64. (PETRAFESO, F. A., and LONG, C. L.) Aeromagnetic map of the Georgia Nuclear Laboratory area, Georgia: U.S. Geol. Survey Geophys. Inv. Map GP-488, 1964.
1-65. (LONG, C. L., and MOORE, F. C.) Aeromagnetic map of the Columbus-Dayton area, Ohio and Indiana: U.S. Geol. Survey Geophys. Inv. Map GP-491, 1965.
- PHILLIPS, K. N.
1-65. (NEWCOMB, R. C., SWENSON, H. A., and LAIRD, L. B.) Water for Oregon: U.S. Geol. Survey Water-Supply Paper 1649, 150 p., 1965.
- PIERCE, A. P.
1-65. Carbon dioxide, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 125-130, 1965.
2-65. Helium, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 120-124, 1965.
- PILLMORE, C. L.
1-65. Geologic map of the Catskill NE quadrangle, Colfax County, New Mexico: U.S. Geol. Survey open-file report, 2 sheets, 1965.
2-65. Geologic map of the Catskill SE quadrangle, Colfax County, New Mexico: U.S. Geol. Survey open-file report, 1965.
- PIPER, A. M.
1-65. (and STEAD, F. W.) Potential applications of nuclear explosives in development and management of water resources—principles: U.S. Geol. Survey Rept. TEI 857, 128 p., 1965.
- PIPIRINGOS, G. N.
1-65. (CHISHOLM, W. A., and KEPFERLE, R. C.) Geology and uranium deposits in the Cave Hills area, Harding County, South Dakota: U.S. Geol. Survey Prof. Paper 476-A, p. A1-A64, 1965.
- PISTRANG, M. A.
1-64. (and KUNKEL, Fred) A brief geologic and hydrologic reconnaissance of the Furnace Creek Wash area, Death Valley National Monument, California: U.S. Geol. Survey Water-Supply Paper 1779-Y, p. Y1-Y35, 1964.
- PITKIN, J. A.
1-64. (NEUSCHEL, S. K., and BATES, R. G.) Aeroradioactivity surveys and geologic mapping, *in* The natural radiation environment: Chicago, Illinois, Chicago Univ. Press (for Rice Univ.), p. 723-736, 1964.
- PLAFKER, George
1-64. (and BERG, H. C.) Metallic and nonmetallic mineral resources: Nonmetallic mineral resources, *in* Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 125-146, 1964.
1-65. Tectonic deformation associated with the 1964 Alaska earthquake: Science, v. 148, no. 3678, p. 1675-1687, 1965.
2-65. (MAYO, L. R., and RUSNAK, G. A.) Effects of Alaska's Good Friday earthquake on coastal areas [abs.]: Geol. Soc. America Spec. Paper 82, p. 152, 1965.
3-65. (GRANTZ, Arthur, KACHADOORIAN, Reuben, MAYO, L. R., and GRYC, George) Geologic effects of the Good Friday earthquake in southern Alaska [abs.]: Geol. Soc. America Spec. Paper 82, p. 361, 1965.
- PLEBUCH, R. O.
1-64. Ground-water temperatures in the Coastal Plain of Arkansas: Arkansas Geol. and Conserv. Comm. Water Resources Summ. 2, 2 p., 1964.
- PLOUFFE, Donald
1-64. Gravity observations and Bouguer anomaly values near Tucson, Arizona: U.S. Geol. Survey open-file report, 30 p., 1964.
- PLUHOWSKI, E. J.
1-64. (and KANTROWITZ, I. H.) Hydrology of the Babylon-Isip area, Suffolk County, Long Island, New York: U.S. Geol. Survey Water-Supply Paper 1768, 119 p., 1964.
- POLAND, J. F.
1-64. Shortening and protrusion of well casings caused by compaction of sediments in subsiding areas [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 152-153, 1964.
1-65. (and IRELAND, R. L.) Shortening and protrusion of a well casing due to compaction of sediments in a subsiding area in California, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B180-B183, 1965.
- POLLOCK, S. J.
1-64. Bedrock geology of the Tiverton quadrangle, Rhode Island-Massachusetts: U.S. Geol. Survey Bull. 1158-D, p. D1-D16, 1964.
2-64. (and FLECK, W. B.) Assabet River basin—Records of wells and test holes, materials tests, and chemical analyses of water in the Assabet River basin, Massachusetts: U.S. Geol. Survey Massachusetts Basic-Data Rept. 8, Ground-Water ser., 45 p., 1964.
- POMERENE, J. B.
1-64. Geology of the Barthell quadrangle and part of the Oneida North quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-314, 1964.
2-64. Geology of the Portersburg quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-359, 1964.
- POOLE, F. G.
1-63. (DAVIS, R. E., EGE, J. R., and HOOVER, D. L.) Geology of a strip between Cane Spring road and Mara

- POOLE, F. G. --Continued
Wash, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-52, 2 p., 1963.
- 1-64. Paleowinds in the western United States, *in* Problems in palaeoclimatology: New York, John Wiley and Sons, p. 394-405, 423, 424, 1964.
- 2-64. (ELSTON, D. P., and CARR, W. J.) Geology of the Cane Spring quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report TEI-851, 1964.
- 3-64. (and STEWART, J. H.) Chinle Formation and Glen Canyon sandstone in northeastern Utah and northwestern Colorado, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D30-D39, 1964; *also in* Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse, 13th annual field conference, 1964: Salt Lake City, Utah, Intermountain Assoc. Petroleum Geologists, p. 93-104, 1964.
- POPENOE, Peter
- 1-64. (BOYNTON, G. R., and ZANDLE, G. L.) Aero-magnetic map of the East Lee quadrangle, Berkshire County, Massachusetts: U.S. Geol. Survey Geophys. Inv. Map GP-452, 1964.
- 2-64. (BOYNTON, G. R., and ZANDLE, C. L.) Aero-magnetic map of the North Adams quadrangle, Berkshire and Franklin Counties, Massachusetts, and Bennington County, Vermont: U.S. Geol. Survey Geophys. Inv. Map GP-454, 1964.
- PORTERFIELD, George
- 1-64. (and DUNNAM, C. A.) Sedimentation of Lake Pillsbury, Lake County, California: U.S. Geol. Survey Water-Supply Paper 1619-EE, p. EE1-EE46, 1964.
- POST, A. S.
- 1-64. Influence of the 1964 Good Friday earthquake on Alaskan glaciers [abs.]: Am. Geophys. Union Trans., v. 45, no. 4, p. 610-611, 1964.
- POWELL, J. E.
- 1-64. (WAGAR, J. E., and PETRI, L. R.) Water resources, *in* Mineral and water resources of South Dakota: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 161-210, 1964.
- POWELL, W. J.
- 1-64. (and KNOWLES, D. B.) Relationship of the geologic structure to the occurrence of ground water in a limestone terrane in the Tennessee Valley, Alabama [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 153-154, 1964.
- PRATT, W. P.
- 1-65. Geologic map of the Hurley West quadrangle, Grant County, New Mexico: U.S. Geol. Survey open-file report, 1965.
- PRICE, Don
- 1-64. Ground-water reconnaissance in the Burnt River Valley, Baker County, Oregon: U.S. Geol. Survey open-file report, 48 p., 1964.
- 1-65. (HART, D. H., and FOXWORTHY, B. L.) Artificial recharge in Oregon and Washington, 1962: U.S. Geol. Survey Water-Supply Paper 1594-C, p. C1-C65, 1965.
- PRICE, W. E., Jr.
- 1-64. Geology and hydrology of alluvial deposits along the Ohio River between Prospect and southwestern Louisville, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-130, sheet 1, 1964.
- 2-64. Geology and hydrology of alluvial deposits along the Ohio River between southwestern Louisville and West Point, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-111, sheet 1, 1964.
- 3-64. Fence diagrams of alluvial deposits along the Ohio River between Prospect and southwestern Louisville, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-130, sheet 2, 1964.
- 4-64. Geology and hydrology of alluvial deposits along the Ohio River between Catlettsburg and South Portsmouth, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-75, sheet 1, 1964.
- PRICE, W. E., Jr. --Continued
- 5-64. Sections and fence diagrams of alluvial deposits along the Ohio River between Catlettsburg and south Portsmouth, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-75, sheet 2, 1964.
- 6-64. Sections and fence diagrams of alluvial deposits along the Ohio River between southwestern Louisville and West Point, Kentucky: U.S. Geol. Survey Hydrol. Inv. Atlas HA-111, sheet 2, 1964.
- PRICHARD, J. I.
- 1-65. (KELLER, G. V., and ANDERSON, L. A.) Magnetic and telluric-electric studies at the Hawaiian Volcano Observatory, Hawaii [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 70-71, 1965.
- PRINZ, W. C.
- 1-64. Diagram for determining mineral composition in the system $MnCO_3$ - $CaCO_3$ - $MgCO_3$, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C84-C85, 1964.
- 2-64. Mineral resources: Manganese, *in* Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 116-119, 1964.
- PROMMEL, H. C.
- 1-64. (and HOPKINS, P. M.) Metallic mineral resources: Placer gold, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 85-93, 1964.
- PUFFETT, W. P.
- 1-64. Geology of the Hazard South quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-343, 1964.
- PUMPELLY, J. W.
- 1-64. Cartographic treatments in the production of orthophotomaps: Am. Cong. Surveying and Mapping Quart. Jour., v. 24, no. 4, p. 567-571, 1964.
- PYATT, E. E.
- 1-64. On determining pollutant distribution in tidal estuaries: U.S. Geol. Survey Water-Supply Paper 1586-F, 56 p., 1964.
- QUINLIVAN, W. D.
- 1-63. A stratigraphic summary and some physical properties of the tuffs of Timber Mountain: U.S. Geol. Survey Tech. Letter NTS-61, 22 p., 1963.
- 1-65. (and LIPMAN, P. W.) Compositional variations in some Cenozoic ashflow tuffs, southern Nevada [abs.]: Geol. Soc. America Spec. Paper 82, p. 342, 1965.
- RADBRUCH, D. H.
- 1-64. Log for field trip through Caldecott Tunnel, Berkeley Hills, California: U.S. Geol. Survey open-file report, 3 p., 1964.
- RAINEY, H. C., III
- 1-65. Geology of the Auburn quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-415, 1965.
- RAINWATER, F. H.
- 1-65. (and THATCHER, L. L.) Methods for collection and analysis of water samples: U.S. Geol. Survey Water-Supply Paper 1454, 301 p., 1965. (Originally published 1960.)
- RAMSPOTT, L. D.
- 1-65. Earliest effects of weathering in Elberton granite: Georgia Acad. Sci. Bull., v. 23, no. 1, p. 34-35, 1965.
- RANDALL, A. D.
- 1-64. Records and logs of selected wells and test borings, records of springs, and chemical analyses of water in the Farmington-Granby area, Connecticut: Connecticut Water Resources Comm. Water Resources Bull. 3, 25 p., 1964.
- 1-65. Preliminary surficial geologic map of the Danielson quadrangle, Connecticut: U.S. Geol. Survey open-file report, 1965.
- 2-65. (THOMAS, M. P., and THOMAS, C. E., Jr.) Water resources inventory of Connecticut—Pt. I, Quinebaug River basin: U.S. Geol. Survey open-file report, 305 p., 1965.

- RANDICH, P. G.
1-64. Map of Burleigh County, North Dakota, showing the location of aquifers and potential yields: U.S. Geol. Survey open-file report, 1964.
1-65. Geology and ground water resources of Burleigh County, North Dakota—Pt. 2, Ground water basic data: North Dakota Geol. Survey Bull. 42, 273 p., 1965.
- RANDOLPH, W. J.
1-64. (and FRYE, P. M.) Floods on Mill Creek near Antioch, Tennessee: U.S. Geol. Survey open-file report, 20 p., 1964.
- RANKIN, D. W.
1-65. The Matagamon sandstone—A new Devonian formation in north-central Maine: U.S. Geol. Survey Bull. 1194-F, p. F1-F9, 1965.
- RANTZ, S. E.
1-64. Stream hydrology related to the optimum discharge for King Salmon spawning in the northern California coast ranges: U.S. Geol. Survey Water-Supply Paper 1779-AA, p. AA1-AA16, 1964.
1-65. Floods of December 1964 in Redwood areas of north coastal California: U.S. Geol. Survey open-file report, 39 p., 1965.
2-65. (and MOORE, A. M.) Floods of December 1964 in the far western States: U.S. Geol. Survey open-file report, 112 p., 1965.
- RASETTI, France
1-65. (and PALMER, A. R.) Trilobites, in *Handbook of paleontological techniques*: San Francisco, Calif., W. H. Freeman and Co., p. 61-64, 1965.
- RASMUSSEN, W. C.
1-64. Permeability and storage of heterogeneous aquifers in the United States: *Internat. Assoc. Sci. Hydrology Pub.* 64, p. 317-325, 1964.
- RATTE, J. C.
1-64. Nonmetallic and industrial minerals: Alunite, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 155-157, 1964.
2-64. (and NORTON, J. J.) Metallic mineral resources: Tin, in *Mineral and water resources of South Dakota*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 68-71, 1964.
- RAUP, O. B.
1-64. Nonmetallic and industrial minerals: Salts of sodium and potassium, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 173-177, 1964.
- RAY, H. A.
1-64. Definition of stage-discharge relation at a stream-flow gaging station by standard step backwater analysis [abs.]: *Am. Geophys. Union Trans.*, v. 45, no. 4, p. 612, 1964.
1-65. Definition of stage-discharge relations by step-backwater analysis: U.S. Geol. Survey open-file report, 20 p., 1965.
- REED, J. C., Jr.
1-64. Chemistry of Greenstone of the Catoctin formation in the Blue Ridge of central Virginia, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C69-C73, 1964.
2-64. Geology of the Linville Falls quadrangle, North Carolina: U.S. Geol. Survey Bull. 1161-B, p. B1-B53, 1964.
3-64. Recent retreat of the Teton Glacier, Grand Teton National Park, Wyoming, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C147-C151, 1964.
4-64. (and BRYANT, Bruce) Evidence for strike-slip faulting along the Brevard zone in North Carolina: *Geol. Soc. America Bull.*, v. 75, no. 12, p. 1177-1196, 1964.
- REED, J. C., Jr. --Continued
1-65. Rate of ice movement and estimated ice thickness in part of the Teton Glacier, Grand Teton National Park, Wyoming, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B137-B141, 1965.
- REED, J. E.
1-64. (DEUTSCH, Morris, and WIITALA, S. W.) Induced recharge of an artesian glacial drift aquifer at Kalama-zoo Michigan: U.S. Geol. Survey open-file report, 65 p., 1964.
- REEDER, H. O.
1-64. Tritium content as an indicator of age and movement of ground water in the Roswell Basin, New Mexico, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C161-163, 1964.
- REEVES, R. G.
1-64. Geology of the Hima quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-319, 1964.
2-64. Metallic mineral resources: Iron, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong., 88th, 2d sess., Senate Doc. no. 87, p. 101-112, 1964.
3-64. Metallic mineral resources: Iron, in *Mineral and water resources of Utah*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 89-96, 1964.
1-65. Geology and mineral resources of the Monlevade and Rio Piracicaba quadrangles, Minas Gerais, Brazil: U.S. Geol. Survey open-file report, 173 p., 1965.
- REEVES, W. E.
1-65. (ROHNE, P. B., BLAKEY, J. F., and GILBERT, C. R.) Base-flow studies, Nueces River, Texas—Quantity and quality, November 23-25, 1964: U.S. Geol. Survey open-file report, 13 p., 1965.
- REPENNING, C. A.
1-65. An extinct shrew from the early Pleistocene of South Africa: *Jour. Mammalogy*, v. 46, no. 2, p. 189-196, 1965.
2-65. (COOLEY, M. E., and AKERS, J. P.) Stratigraphy of the Chinle and Moenkopi Formations, Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah: U.S. Geol. Survey open-file report, 146 p., 1965.
- RETTIG, S. L.
1-64. (and JONES, B. F.) Determination of carbonate, bicarbonate, and total CO₂ in carbonate brines, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D134-D137, 1964.
- RETTMAN, P. L.
1-65. (and LEGGAT, E. R.) Ground-water resources of Gaines County, Texas: U.S. Geol. Survey open-file report, 302 p., 1965.
- RICE, C. L.
1-64. Geology of the Lancer quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-347, 1964.
- RICHARDS, P. W.
1-64. Geology of the Smiths Grove quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-357, 1964.
- RICHARDSON, Donald
1-65. Effect of logging on runoff in upper Green River Basin, Washington: U.S. Geol. Survey open-file report, 44 p., 1965.
- RICHMOND, G. M.
1-64. Glaciation of Little Cottonwood and Bells Canyons, Wasatch Mountains, Utah: U.S. Geol. Survey Prof. Paper 454-D, p. D1-D41, 1964.
2-64. Three pre-Bull Lake tills in the Wind River Mountains, Wyoming—A reinterpretation, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D104-D109, 1964.
- RICHTER, D. H.
1-64. (AULT, W. U., EATON, J. P., and MOORE, J. G.) The 1961 eruption of Kilauea Volcano, Hawaii: U.S. Geol. Survey Prof. Paper 474-D, p. D1-D34, 1964.

RIGGS, H. C.

- 1-64. The base-flow recession curve as an indicator of ground water, *in* Symposium—Surface waters, Gen. Assembly of Berkeley of I. U. G. G., 1963: Internat. Assoc. Sci. Hydrology Pub. 63, p. 352-363, 1964.
- 2-64. Stream discharge regressions using precipitation, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C185-C187, 1964.
- 1-65. Discussion of "Areal variations of mean annual runoff," by A. G. Hely: Am. Soc. Civil Engineers Proc., Hydraulics Div. Jour., v. 91, no. HY2, p. 396-399, 1965.

RIMA, D. R.

- 1-64. (COSKERY, O. J., and ANDERSON, P. W.) Ground-water resources of southern New Castle County, Delaware: U.S. Geol. Survey Water-Supply Paper 1756 (Delaware Geol. Survey Bull. 11), 54 p., 1964.

ROBERTS, A. E.

- 1-65. Correlation of Cretaceous and Lower Tertiary rocks near Livingston, Montana, with those in other areas of Montana and Wyoming, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B54-B63, 1965.

ROBERTS, H. B.

- 1-64. Fossil decapod crustaceans from the Marshall Islands: U.S. Geol. Survey Prof. Paper 260-HH, p. HH1127-HH1131, 1964.

ROBERTS, R. J.

- 1-64. Economic geology, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 39-48, 1964.
- 2-64. Metallic mineral resources: Copper, *in* Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 75-83, 1964.
- 3-64. Paleozoic rocks, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 22-26, 1964.
- 4-64. Stratigraphy and structure of the Antler Peak quadrangle, Humboldt and Lander Counties, Nevada: U.S. Geol. Survey Prof. Paper 459-A, p. A1-A93, 1964[1965].
- 1-65. Generalized map of Paleozoic and Mesozoic facies in north-central Nevada: U.S. Geol. Survey open-file report, 1965.
- 2-65. (and ARNOLD, D. C.) Ore deposits of the Antler Peak quadrangle, Humboldt and Lander Counties, Nevada: U.S. Geol. Survey Prof. Paper 459-B, p. B1-B94, 1965.
- 3-65. (and THOMASSON, M. R.) Comparison of late Paleozoic tectonic history of northern Nevada and central Idaho [abs.]: Geol. Soc. America Spec. Paper 82, p. 343, 1965.

ROBERTSON, E. C.

- 1-64. Experimental consolidation of carbonate mud [abs.]: Abstracts of AAPG-SEPM-GAC-MAC papers, Toronto, Canada, May 18-21, 1964, p. 544, 1964.
- 2-64. Strengths and elastic moduli of serpentinite from Mayagüez, Puerto Rico, *in* A study of serpentinite, the AMSOC core hole near Mayagüez, Puerto Rico: Natl. Acad. Sci.-Natl. Research Council Pub. 1188, p. 118-125, 1964.
- 3-64. Viscoelasticity of rocks, *in* State of stress in the Earth's crust—International Conference, Santa Monica, California, 1963, Proc.: New York, America Elsevier Pub. Co., p. 180-233, 1964.

ROBIE, R. A.

- 1-65. Vacuum jacketed hydrofluoric acid solution calorimeter: Rev. Sci. Instruments, v. 36, no. 4, p. 484-486, 1965.

ROBINOVE, C. J.

- 1-65. Ground-water studies and analog models: U.S. Geol. Survey Circ. 468, 12 p., 1965. (Originally published 1962.)

ROBINSON, C. S.

- 1-64. (and LEE, F. T.) Geologic research at the Straight Creek tunnel site, Colorado: Highway Research Record no. 57, p. 18-34, 1964.

ROBINSON, C. S. --Continued

- 2-64. (MAPEL, W. J., and BERGENDAHL, M. H.) Stratigraphy and structure of the northern and western flanks of the Black Hills uplift, Wyoming, Montana, and South Dakota: U.S. Geol. Survey Prof. Paper 404, 134 p., 1964.

- 1-65. (and LEE, F. T.) Preliminary report on the engineering geology of the Straight Creek Tunnel pilot bore, Clear Creek and Summit Counties, Colorado: U.S. Geol. Survey open-file report, 34 p., 1965.

- 2-65. (and LEE, F. T.) The validity of geologic projection, a successful example: The Straight Creek Tunnel pilot bore, Colorado, with an appendix of examples of types of data and methods of analysis used: U.S. Geol. Survey open-file report, 65 p., 1965.

ROBINSON, F. M.

- 1-64. Core tests Simpson area, Alaska—Exploration of naval petroleum reserve No. 4 and adjacent areas, northern Alaska, 1944-53—Pt. 5, Subsurface geology and engineering data: U.S. Geol. Survey Prof. Paper 305-L, p. L645-L730, 1964.

ROBINSON, G. M.

- 1-64. The evaluation of water-resources development by analog techniques: Internat. Assoc. Sci. Hydrology Pub. 64, p. 442-454, 1964.

ROBINSON, T. W.

- 1-64. Phreatophyte research in the western States, March 1959 to July 1964: U.S. Geol. Survey Circ. 495, 31 p., 1964.

- 1-65. Introduction, spread and areal extent of saltcedar (Tamarix) in the western States: U.S. Geol. Survey Prof. Paper 491-A, p. A1-A12, 1965.

- 2-65. (and JOHNSON, A. I.) Selected bibliography on evaporation and transpiration: U.S. Geol. Survey Water-Supply Paper 1539-R, p. R1-R25, 1965. (Originally published 1961.)

ROBINSON, F. L.

- 1-65. Floods in New York—1964: U.S. Geol. Survey open-file report, 22 p., 1965.

ROBISON, T. M.

- 1-64. Occurrence and availability of ground water in Ohio County, West Virginia: West Virginia Geol. and Econ. Survey Bull. 27, 57 p., 1964.

RODDY, D. J.

- 1-63. Geologic section across the Flynn Creek structure, *in* Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 53-74, 1963.

ROEDDER, Edwin

- 1-65. (and COOMBS, D. S.) Immiscibility in granitic melts, indicated by fluid inclusions in ejected granitic blocks from Ascension Island volcanic breccias [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 187, 1965.
- 2-65. (and SMITH, R. L.) Liquid water in pumice vesicles, a crude but useful dating method [abs.]: Geol. Soc. America Spec. Paper 82, p. 164, 1965.

ROGERS, J. E.

- 1-64. Ground-water resources in relation to Louisiana agriculture: Louisiana Assoc. of Agronomists Proc., v. 5, p. 37-50, 1964.

- 1-65. (and CALANDRO, A. J.) Water resources of Vernon Parish, Louisiana: Louisiana Geol. Survey Water Resources Bull. 6, 104 p., 1965.

ROHRER, W. L.

- 1-65. Geologic map of the Adam Weiss Peak quadrangle, Hot Springs and Park Counties, Wyoming: U.S. Geol. Survey Geol. Quad. Map GQ-382, 1965.

ROLFE, B. N.

- 1-64. (and HADLEY, R. F.) Weathering and transport of sediment in the Cheyenne River Basin, eastern Wyoming, *in* Clays and clay minerals, 12th Natl. Conf., Atlanta, Georgia, 1963, Proc.: New York, Macmillan Co., p. 649-670, 1964.

ROLLER, J. C.

- 1-65. (HEALY, J. H., and JACKSON, W. H.) A seismic-refraction profile extending from Lake Superior, Wisconsin, to Agate, Colorado [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 155, 1965.

- 2-65. Crustal structure in the eastern Colorado Plateaus province from seismic-refraction measurements: *Seismol. Soc. America Bull.*, v. 55, no. 1, p. 107-119, 1965.

RORABAUGH, M. I.

- 1-64. Estimating changes in bank storage and ground-water contribution to streamflow, in *Symposium—Surface waters*, Gen. Assembly of Berkeley of I. U. G. G., 1963; *Internat. Assoc. Sci. Hydrology Pub.* 63, p. 432-441, 1964.

ROSE, H. J., Jr.

- 1-63. Semi-micro X-ray fluorescence analysis of tektites using 50 milligram samples, in *Astrogeologic studies*, annual progress report, August 25, 1962-July 1, 1963—Pt. D, *Studies for Space Flight Program*: U.S. Geol. Survey open-file report, p. 75-82, 1963; also in U.S. Geol. Survey Prof. Paper 475-D, p. D171-D173, 1964.

- 1-65. (and CUTTITTA, Frank) Microanalysis by X-ray fluorescence Determination of selected major constituents in silicates [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 166, 1965.

- 2-65. (CUTTITTA, Frank, and LARSON, R. R.) Use of X-ray fluorescence in determination of selected major constituents in silicates, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B155-B159, 1965.

ROSEBOOM, E. H., Jr.

- 1-65. Reactions in some synthetic and natural copper sulfides between 25° and 200°C [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 166-167, 1965.

ROSENE, P. G.

- 1-64. Chemical quality of the water, in *Ground-water conditions in the proposed waterfowl refuge area near Chapman, Nebraska*: U.S. Geol. Survey Water-Supply Paper 1779-E, p. E28-E38, 1964.

ROSENSHEIN, J. S.

- 1-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Fulton County*: Indiana Dept. Water Resources Bull. 20, 83 p., 1964.

- 2-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Jasper County*: Indiana Div. Water Resources Bull. 25, 83 p., 1964.

- 3-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Marshall County*: Indiana Dept. Water Resources Bull. 19, 157 p., 1964.

- 4-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Newton County*: Indiana Div. Water Resources Bull. 26, 69 p., 1964.

- 5-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Pulaski County*: Indiana Div. Water Resources Bull. 24, 71 p., 1964.

- 6-64. (and HUNN, J. D.) Ground-water resources of northwestern Indiana—Preliminary report, *Starke County*: Indiana Div. Water Resources Bull. 22, 87 p., 1964.

ROSHOLT, J. N., Jr.

- 1-63. Isotopic fractionation of uranium in fluvial sediments [abs.]: *Am. Geophys. Union Trans.*, v. 44, no. 4, p. 893, 1963.

- 1-64. (HARSHMAN, E. N., SHIELDS, W. R., and GARNER, E. L.) Isotopic fractionation of uranium related to roll features in sandstone, *Shirley Basin, Wyoming*: *Econ. Geology*, v. 59, no. 4, p. 570-585, 1964.

- 1-65. (BUTLER, A. P., GARNER, E. L., and SHIELDS, W. R.) Isotopic fractionation of uranium in sandstone, *Powder River Basin, Wyoming, and Slick Rock district, Colorado*: *Econ. Geology*, v. 60, no. 2, p. 199-213, 1965.

- 2-65. (DOE, B. R., and TATSUMOTO, M.) Isotopic composition of uranium in soil profiles [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 167-168, 1965.

ROSHOLT, J. N., Jr. --Continued

- 3-65. (TATSUMOTO, M., and DOOLEY, J. R., Jr.) Radioactive disequilibrium studies in sandstone: *Powder River Basin, Wyoming, and Slick Rock district, Colorado*: *Econ. Geology*, v. 60, no. 3, p. 477-484, 1965.

ROSS, C. P.

- 1-64. (REID, R. R., and WEISSENBORN, A. E.) *Geology, in Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 23-40, 1964.

ROSS, D. C.

- 1-65. *Geology of the Independence quadrangle, Inyo County, California*: U.S. Geol. Survey Bull. 1181-O, p. O1-O64, 1965.

ROSS, Malcolm

- 1-65. (and EVANS, H. T., Jr.) Studies of the torbernite minerals (III): Role of the interlayer oxonium, potassium, and ammonium ions, and water molecules: *Am. Mineralogist*, v. 50, nos. 1-2, p. 1-12, 1965.

- 2-65. (and WONES, D. R.) Polytypism in biotites [abs.]: *Am. Mineralogist*, v. 50, nos. 1-2, p. 291, 1965.

ROSS, R. J., Jr.

- 1-64. Middle and Lower Ordovician formations in southernmost Nevada and adjacent California: U.S. Geol. Survey Bull. 1180-C, p. C1-C101, 1964.

- 2-64. Relations of Middle Ordovician time and rock units in Basin Ranges, western United States: *Am. Assoc. Petroleum Geologists Bull.*, v. 48, no. 9, p. 1526-1554, 1964.

- 3-64. (and LONGWELL, C. R.) Paleotectonic significance of Ordovician sections south of the Las Vegas shear zone, in *Middle and Lower Ordovician formations in southernmost Nevada and adjacent California*: U.S. Geol. Survey Bull. 1180-C, p. C88-C93, 1964.

ROSTVEDT, J. O.

- 1-65. Summary of floods in the United States during 1960: U.S. Geol. Survey Water-Supply Paper 1790-B, p. B1-B147, 1965.

ROWE, J. J.

- 1-65. (FOURNIER, R. O., and MOREY, G. W.) Use of sodium iodide to trace underground water circulation in the hot springs and geysers of the Daisy Geyser Group, *Yellowstone National Park*, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B184-B186, 1965.

- 2-65. (MOREY, G. W., and FOURNIER, R. O.) System $\text{H}_2\text{O}-\text{Na}_2\text{O}-\text{SiO}_2$ at 200°, 250°, and 300°C [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 169-170, 1965.

- 3-65. (MOREY, G. W., and HANSEN, Ingrid) The binary system $\text{K}_2\text{SO}_4-\text{CaSO}_4$: *Jour. Inorganic and Nuclear Chemistry*, v. 27, no. 1, p. 53-58, 1965.

RUBIN, Jacob

- 1-64. (and STEINHARDT, R.) Soil water relations during rain infiltration, III, Water uptake at incipient ponding: *Soil Sci. Soc. America Proc.*, v. 28, no. 5, p. 614-619, 1964.

RUGGLES, F. H., Jr.

- 1-64. Floods on Bachman Branch and Joes Creek at Dallas, Texas: U.S. Geol. Survey open-file report, 18 p., 1964.

RUPPEL, E. T.

- 1-64. Strike-slip faulting and broken basin-ranges in east-central Idaho and adjacent Montana, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C14-C18, 1964.

- 1-65. Strike-slip faulting and broken basin-ranges in east-central Idaho and adjacent Montana [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 344, 1965.

RUSH, F. E.

- 1-64. Ground-water appraisal of the Meadow Valley area, Lincoln and Clark Counties, Nevada: *Nevada State Dept. Conserv. and Nat. Resources, Ground-Water Resources—Reconn. Ser. Rept.* 27, 43 p., 1964.

RUSH, F. E.--Continued

- 2-64. (and EVERETT, D. E.) Ground-water appraisal of Monitor, Antelope, and Koebe Valleys, Nevada: Nevada Dept. Conserv. and Nat. Resources Ground-Water Resources--Reconn. Ser. Rept. 30, 45 p., 1964.

RYALL, Alan

- 1-62. (and BIRTILL, John) Digital processing of array seismic recordings: U.S. Geol. Survey Tech. Letter [Crustal Studies], no. 2, 20 p., 1962.

RYAN, J. D.

- 1-64. Geology of the Edgemont quadrangle, Fall River County, South Dakota: U.S. Geol. Survey Bull. 1063-J, p. 379-426, 1964.

SABLE, E. G.

- 1-64. Geology of the Constantine quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-302, 1964.

SACHS, K. N., Jr.

- 1-64. Multilocular embryonites in *Lepidocyclina* (*Eulepidina*) *undosa* Cushman from Puerto Rico: *Micropaleontology*, v. 10, no. 3, p. 323-329, 1964.

- 2-64. (CIFELLI, Richard, and BOWEN, V. T.) Ignition to concentrate shelled organisms in plankton samples: *Deep-Sea Research*, v. 11, no. 4, p. 621-622, 1964.

SAINSBURY, C. L.

- 1-64. Association of beryllium with tin deposits rich in fluorite: *Econ. Geology*, v. 59, no. 5, p. 920-926, 1964.

- 1-65. Planetable maps and drill logs of the Camp Creek and Bessie-Mapel beryllium-fluorspar deposits, Lost River area, Alaska: U.S. Geol. Survey open-file report, 32 p., 1965.

- 2-65. (and JAHNS, R. H.) Tin, in *Mineral and water resources of New Mexico*: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 237-240, 1965.

SAMMEL, E. A.

- 1-64. Evaluation of numerical-analysis methods for determining variations in transmissibility: *Internat. Assoc. Sci. Hydrology Pub.* 64, p. 239-251, 1964.

- 2-64. (BRACKLEY, R. A., and PALMQUIST, W. N., Jr.) Synopsis of water resources of the Ipswich River Basin, Massachusetts: U.S. Geol. Survey Hydrol. Inv. Atlas HA-196, 1964.

- 1-65. (BAKER, J. A., and BRACKLEY, R. A.) Water resources of the Ipswich River Basin, Massachusetts: U.S. Geol. Survey open-file report, 197 p., 1965.

SAMPLE, R. D.

- 1-65. Geology of the Princeton West quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-385, 1965.

SANDBERG, C. A.

- 1-64. Precambrian to Mississippian paleotectonics of the southern Williston Basin [abs.]: *Internat. Williston Basin Symposium*, 3d, Geol. Soc. North Dakota-Saskatchewan Geol. Soc., Regina, Saskatchewan Program, Sept., p. 37-38, 1964.

- 2-64. (and McMANNIS, W. J.) Occurrence and paleogeographic significance of the Maywood Formation of late Devonian age in the Gallatin Range, southwestern Montana, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-C, p. C50-C54, 1964.

- 3-64. (and PRICHARD, G. E.) Mineral fuel resources: Petroleum and natural gas, in *Mineral and water resources of South Dakota*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 151-160, 1964.

- 1-65. Nomenclature and correlation of lithologic subdivisions of the Jefferson and Three Forks Formations of southern Montana and northern Wyoming: U.S. Geol. Survey Bull. 1194-N, p. N1-N18, 1965.

- 2-65. (and BOWLES, C. G.) Geology of the Cub Run quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-386, 1965.

SANDO, W. J.

- 1-64. Stratigraphic importance of corals in the redwall limestone, northern Arizona, in *Geological Survey Re-*

SANDO, W. J.--Continued

- search 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C39-C42, 1964.

- 1-65. *Faviphyllum rugosum* Hall, 1852 (Anthozoa, Rugosa): Proposed suppression under the plenary powers of generic and specific names: *Zool. Nomenclature*, v. 22, pt. 1, p. 55-56, 1965.

- 2-65. Revision of certain Paleozoic coral species from western United States [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 172, 1965.

SANTOS, J. F.

- 1-65. Quality of surface waters in the lower Columbia River basin: U.S. Geol. Survey Water-Supply Paper 1784, 78 p., 1965.

SARGENT, K. A.

- 1-63. Summary geology of the Skull Mountain quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-60, 11 p., 1963.

- 1-64. Geology of Quartet Dome quadrangle, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-69, 11 p., 1964.

- 1-65. Use of magnetic susceptibility and grain density in identification of basalt flows at the Nevada Test Site, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B142-B145, 1965.

- 2-65. (LUFT, S. J., GIBBONS, A. B., and HOOVER, D. L.) Geologic map of the Quartet Dome quadrangle, Nye County, Nevada: U.S. Geol. Survey open-file report, TEI-860, 1965.

- 3-65. Heavy minerals for correlation and identification of Tertiary rhyolites at the Nevada Test Site [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 344, 1965.

SATO, Motoaki

- 1-65. (and CHRIST, C. L.) Electrode properties of semiconductive binary sulfides and their possible geologic application [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 173, 1965; *abstract also in Econ. Geology*, v. 59, no. 7, p. 1422, 1965.

SAUER, S. P.

- 1-64. (and RAWSON, Jack) Base flow studies, Sabine River, Texas and Louisiana, quantity and quality, Sept. 4-5, 1963: U.S. Geol. Survey open-file report, 55 p., 1964.

- 1-65. (and BLAKEY, J. F.) Water delivery study, Nueces River, Texas, quantity and quality, August 1963: *Texas Water Comm. Bull.* 6509, 22 p., 1965.

- 2-65. Hydrologic studies of small watersheds: Mukewater Creek, Colorado River basin, Texas 1952-60: U.S. Geol. Survey open-file report, 97 p., 1965.

SAUER, V. B.

- 1-64. Floods in Louisiana--Magnitude and frequency, 2d ed.: Baton Rouge, Louisiana, Louisiana Dept. of Highways, 402 p., 1964.

- 2-64. Magnitude and frequency of storm runoff in southeastern Louisiana and southwestern Mississippi, in *Geological Survey Research 1964*: U.S. Geol. Survey Prof. Paper 501-D, p. D182-D184, 1964.

SAVAGE, C. N.

- 1-64. (and La HEIST, B. A.) Mineral resources: Peat, in *Mineral and water resources of Idaho*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 143-152, 1964.

SAYRE, W. W.

- 1-64. (and CHAMBERLAIN, A. R.) Exploratory laboratory study of lateral turbulent diffusion at the surface of an alluvial channel: U.S. Geol. Survey Circ. 484, 18 p., 1964.

SCHALLER, Waldemar

- 1-65. (VLISIDIS, A. C., and MROSE, M. C.) Macallisterite, $2\text{MgO} \cdot 6\text{B}_2\text{O}_3 \cdot 15\text{H}_2\text{O}$, a new hydrous magnesium borate mineral from Death Valley region, Inyo County, California [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 173-174, 1965.

SCHEIDEGGER, A. E.

- 1-64. Dynamic similarity conditions in geomorphology [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 175, 1964.
- 1-65. A theory of varve formation: Internat. Assoc. Sci. Hydrology Bull., v. 10, no. 1, p. 68-73, 1965.
- 2-65. The algebra of stream-order numbers, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B187-B189, 1965.

SCHELL, E. M.

- 1-64. Coal in the Uinta Basin, Utah, in Guidebook to the geology and mineral resources of the Uinta Basin, Utah's hydrocarbon storehouse, 13th annual field conference 1964: Salt Lake City, Utah, Intermountain Assoc. Petroleum Geologists, p. 253-257, 1964.

SCHER, M. B.

- 1-64. Mapmaking applications of orthophotography, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D138-D140, 1964.
- 2-64. Research in orthophotography: Photogramm. Eng., v. 30, no. 5, p. 756-762, 1964.
- 3-64. (and HALLIDAY, James) Photogrammetric contouring of areas covered by evergreen trees: Nachrichten aus dem Karten- und Vermessungswesen, Reihe V: Sonderhefte—No. 9, 1964.

SCHINER, G. R.

- 1-64. (and SCHNEIDER, Robert) Geology and ground-water conditions of the Redwood Falls area, Redwood County, Minnesota: U.S. Geol. Survey Water-Supply Paper 1669-R, p. R1-R46, 1964.
- 1-65. (and GONTHIER, J. B.) Ground-water map of the Prudence Island and Newport quadrangles: Rhode Island Water Resources Coordinating Board, Ground-Water Map GWM-20, 1965.
- 2-65. (and GONTHIER, J. B.) Ground-water map of the Tiverton and Sakonnet Point quadrangles, Rhode Island, and the Rhode Island portion of the Westport quadrangle, Massachusetts: Rhode Island Water Resources Coordinating Board Ground-Water Map GWM-21, 1965.

SCHLANGER, S. O.

- 1-64. Petrology of the limestones of Guam, with a section on Petrography of the insoluble residues, by J. C. Hathaway and Dorothy Carroll: U.S. Geol. Survey Prof. Paper 403-D, p. D1-D52, 1964.
- 1-65. Geology of the Maretburg quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-338, 1965.

SCHLEE, John

- 1-64. New Jersey offshore gravel deposit: Pit and Quarry, v. 57, no. 6, p. 80-81, 1964.
- 2-64. (UCHUPI, Elazar, and TRUMBULL, J. V. A.) Statistical parameters of Cape Cod beach and eolian sands, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D118-D122, 1964.

SCHLOCKER, Julius

- 1-63. Petrology and mineralogy of Tatum dome, Lamar County, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-28, 120 p., 1963.
- 1-65. (PAMPEYAN, E. H., and BONILLA, M. G.) Approximate trace of the main surface rupture in the San Andreas fault zone between Pacifica and Saratoga, California, formed during the earthquake of April 18, 1906: U.S. Geol. Survey open-file report, map, 1965.

SCHMIDT, D. L.

- 1-64. (DOVER, J. H., FORD, A. B., BROWN, R. D.) Geology of the Patuxent Mountains, in Antarctic geology, Proc. First International Symposium on Antarctic Geology, Cape Town, Sept. 1963: New York, John Wiley and Sons, p. 276-283, 1964[1965].
- 1-65. (WILLIAMS, P. L., NELSON, W. H., and EGE, J. R.) Geology of the Neptune Range, Pensacola Mountains, Antarctica [abs.]: Geol. Soc. America Spec. Paper 82, p. 175-176, 1965.

SCHMIDT, R. G.

- 1-64. (PECORA, W. T., and HEARN, B. C., Jr.) Geology of the Cleveland quadrangle, Bearpaw Mountains, Blaine County, Montana: U.S. Geol. Survey Bull. 1141-P, P1-P26, 1964.

SCHNABEL, R. W.

- 1-64. Geologic appraisal of the U20g site, Pahute Mesa, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Studies-1-15, 10 p., 1964.
- 2-64. (and ERIC, J. H.) Bedrock geologic map of the Windsor Locks quadrangle, Hartford County, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-388, 1964.
- 3-64. (and MacKALLOR, J. A.) Geology of the Fenton quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-317, 1964.
- 1-65. (and ERIC, J. H.) Bedrock geologic map of the Tariffville quadrangle, Hartford County, Connecticut, and Hampden County, Massachusetts: U.S. Geol. Survey Geol. Quad. Map GQ-370, 1965.

SCHNEIDER, Robert

- 1-64. Cenomanian-Turonian aquifer of central Israel—Its development and possible use as a storage reservoir: U.S. Geol. Survey Water-Supply Paper 1608-F, p. F1-F20, 1964.

SCHNEIDER, W. J.

- 1-65. (and others) Water resources of the Appalachian Region, Pennsylvania to Alabama: U.S. Geol. Survey Hydrol. Inv. Map HA-198, 1965.

SCHOEN, Robert

- 1-64. Origin of ironstones in the Clinton Group [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 177, 1964.
- 2-64. Clay minerals of the Silurian Clinton iron-stones, New York State: Jour. Sed. Petrology, v. 34, no. 4, p. 855-863, 1964.

SCHOFF, S. L.

- 1-64. (SALSO, J. H., and GARCÍA, José) Source of heat in a deep artesian aquifer, Bahia Blanca, Argentina, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D153-D157, 1964.
- 2-64. (and SAYAN M., Juan Luis) Hydrologic data for the Lambayeque Valley, northern Peru (basic records): U.S. Geol. Survey open-file report, 52 p., 1964.

SCHOPF, J. M.

- 1-64. Organization and affinity of Vertebraria [abs.]: Internat. Botanical Cong., 10th, Edinburgh, August 1964, p. 33, 1964.
- 2-64. Practical problems and principles in study of plant microfossils, in Palynology in oil exploration—A symposium, San Francisco, California, 1962: Soc. Econ. Paleontologists and Mineralogists Spec. Pub. 11, p. 29, 57, 1964.
- 3-64. Report on dating of plant fossils from the Mapleton sandstone, App. 1, in Reconnaissance bedrock geology of the Presque Island quadrangle, Maine: Maine Geol. Survey Quad. Mapping Ser., no. 2, p. 100-103, 1964.
- 4-64. Russian palynology today—Literature and application to exploration, in Palynology in oil exploration—A symposium, San Francisco, California, 1962: Soc. Econ. Paleontologists and Mineralogists Spec. Pub. 11, p. 181-200, 1964.
- 5-64. Middle Devonian plant fossils from northern Maine, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D43-D49, 1964.
- 1-65. A method for obtaining small acid-resistant fossils from ordinary solution residues, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Co., p. 301-304, 1965.
- 2-65. Review of "Advances in palynology," ed. by P. K. K. Nair: Science, v. 148, no. 3671, p. 831, 1965.

SCHULTZ, L. G.

- 1-64. Quantitative interpretation of mineralogical composition from X-ray and chemical data for the Pierre Shale: U.S. Geol. Survey Prof. Paper 391-C, p. C1-C31, 1964.

SCHUMM, S. A.

- 1-64. Seasonal variations of erosion rates and processes on hillslopes in western Colorado, *in* Fortschritte der Internationalen Hangforschung 1964 [Advances in hillslope investigation 1964]: Zeitschrift für Geomorphologie, Suppl., v. 5, p. 215-238, 1964.
- 2-64. (and CHORLEY, R. J.) The fall of threatening rock: Am. Jour. Sci., v. 262, no. 9, p. 1041-1054, 1964.
- 1-65. (and LICHTY, R. W.) Time, space, and causality in geomorphology: Am. Jour. Sci., v. 263, no. 2, p. 110-119, 1965.

SCHWALEN, H. C.

- 1-64. Chino Valley, *in* Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 53-54, 1964.
- 2-64. Upper Santa Cruz basin [and] Avra-Marana area, *in* Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 30-35, 1964.

SCHWOB, H. H.

- 1-64. Water resources of the English River, Old Mans Creek, and Clear Creek basins in Iowa: U.S. Geol. Survey open-file report, 57 p., 1964.

SCOTT, G. R.

- 1-64. Nonmetallic and industrial minerals: Gem stones, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 165-168, 1964.
- 2-64. (and COBBAN, W. A.) Stratigraphy of the Niobrara Formation at Pueblo, Colorado: U.S. Geol. Survey Prof. Paper 454-L, p. L1-L30, 1964.

SCOTT, J. C.

- 1-64. Ground-water resources of Russell County, Alabama—A reconnaissance: Alabama Geol. Survey Bull. 75, 77 p., 1964.

SCOTT, J. H.

- 1-63. (CARROLL, R. D., and BLACK, R. A.) Postshot electrical resistivity measurements at the Sedan site, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-49, 10 p., 1963.
- 1-64. Electrical resistivity of earth sediments near dipole antenna site, southern Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter, Spec. Proj.-10, 17 p., 1964.
- 2-64. (CARROLL, R. D., and CUNNINGHAM, D. R.) Dielectric constant and electrical conductivity of moist rock from laboratory measurements: U.S. Geol. Survey Tech. Letter, Spec. Proj. -12, 41 p., 1964.

SEABER, P. R.

- 1-64. Ground-water resources in the Susquehanna River basin: Susquehanna River Basin Coordinating Comm., 3d Mtg., Baltimore, Md., Mar. 6, 1964, Minutes 13 p., 1964.
- 2-64. (and HOLLYDAY, E. F.) Hardness of ground water in the glacial deposits of the Susquehanna River basin, New York and Pennsylvania [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov., 1964, Program, p. 181, 1964.

- 1-65. Variation in chemical character of water in the Englishtown Formation, New Jersey: U.S. Geol. Survey Prof. Paper 498-B, p. B1-B35, 1965.

SEGERSTROM, Kenneth

- 1-64. (and HENRIQUEZ, Hugo) Cavities, or "Tafoni," in rock faces of the Atacama Desert, Chile, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C121-C125, 1964.
- 2-64. (and MORAGA BRITO, Aldo) Carta geológica de Chile—Cuadrángulo Chanarcillo, Provincia de Atacama: Chile Inst. Inv. Geol. Carta 13, 1964.

SEGERSTROM, Kenneth--Continued

- 1-65. Dissected gravels of the Rio Copiapo Valley and adjacent coastal area, Chile, *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B117-B121, 1965.

- 2-65. Preliminary results of geochemical prospecting north of the Marquette iron range, Michigan [abs.], *in* Eleventh Annual Institute on Lake Superior Geology: St. Paul, Minn., Univ. Minnesota, May, p. 30, 1965.

SEIDERS, V. M.

- 1-64. Geology of the Hazard North quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-344, 1964.

SENFTLE, F. E.

- 1-63. (THORPE, A. N., and LEWIS, R. R.) Magnetic properties of nickel-iron spherules in tektites from Isabel, Philippine Islands, *in* Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. C, Cosmochemistry and petrography: U.S. Geol. Survey open-file report, p. 85-97, 1963.

SENKPIEL, W. C.

- 1-65. Waterpower, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 432-437, 1965.

SEVER, C. W.

- 1-64. Fuller's earth clay found in Cook County, Georgia [abs.]: Georgia Acad. Sci. Bull., v. 22, no. 8, p. 17, 1964.

- 2-64. Geologic evidence of a buried Miocene fault in Thomas County, southwestern Georgia [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 181-182, 1964.

- 3-64. Geology and ground-water resources of crystalline rocks, Dawson County, Georgia: Georgia Geol. Survey Inf. Circ. 30, 32 p., 1964.

- 4-64. Ground-water conduits in the Ashland Mica Schist, northern Georgia, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D141-D143, 1964.

- 5-64. Ground-water resources and geology of Seminole, Decatur, and Grady Counties, Georgia: U.S. Geol. Survey open-file report, 73 p., 1964.

- 6-64. Prospecting for limestone in Thomas County, Georgia [abs.]: Georgia Acad. Sci. Bull., v. 22, no. 8, p. 17, 1964.

- 1-65. Ground-water resources of Bainbridge, Georgia: Georgia Geol. Survey Inf. Circ. 32, 10 p., 1965.

SHACKLETTE, H. T.

- 1-64. (and SAUER, H. I.) The association of cardiovascular mortality rates in Georgia with the abundance and distribution of certain elements in rocks, soils, and plants: U.S. Geol. Survey open-file report, 21 p., 1964.

- 1-65. A leafy liverwort hydrosere on Yakobi Island, Alaska: Ecology, v. 46, no. 3, p. 377-378, 1965.

- 2-65. Bryophytes associated with mineral deposits and solutions in Alaska: U.S. Geol. Survey Bull. 1198-C, p. C1-C18, 1965.

SHARP, W. N.

- 1-64. (McKAY, E. J. McKEOWN, F. A., and WHITE, A. M.) Geology and uranium deposits of the Pumpkin Buttes area of the Powder River Basin, Wyoming: U.S. Geol. Survey Bull. 1107-H, p. H541-H638, 1964.

SHAW, H. R.

- 1-64. (and WONES, D. R.) Fugacity coefficients for hydrogen gas between 0° and 1000°C, for pressures to 3,000 atm: Am. Jour. Sci., v. 262, no. 7, p. 918-929, 1964.

- 1-65. Comments on viscosity, crystal settling and convection in granitic magmas: Am. Jour. Sci., v. 263, no. 2, p. 120-152, 1965.

SHAW, D. R.

- 1-64. (MOUNTJOY, Wayne, and DUKE, Walter) Lithium associated with beryllium in rhyolitic tuff at Spor Mountain, western Juab County, Utah, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C86-C87, 1964.

- SHAW, D. R. --Continued
 1-65. (and GRANGER, H. C.) Uranium ore rolls—an analysis: *Econ. Geology*, v. 60, no. 2, p. 240-250, 1965.
- SHELDON, R. P.
 1-64. Exploration for phosphorite in Turkey—A case history: *Econ. Geology*, v. 59, no. 6, p. 1159-1175, 1964.
 2-64. Paleolatitudinal and paleogeographic distribution of phosphorite, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C106-C113, 1964.
 1-65. Barrier Islands and lagoonal iron sedimentation in the Silurian of Alabama [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 182, 1965.
- SHELL, J. D.
 1-64. Flood on Pearl River at Jackson, Mississippi in 1961: U.S. Geol. Survey Hydrol. Inv. Atlas HA-127, 1964.
- SHEN, John
 1-65. Use of analog models in the analysis of flood runoff: U.S. Geol. Survey Prof. Paper 506-A, p. A1-A24, 1965.
- SHEPPARD, R. A.
 1-64. Geology of the Portsmouth quadrangle, Kentucky-Ohio, and parts of the New Boston and Wheelersburg quadrangles, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-312, 1964.
 2-64. (and GUDE, A. J., 3d.) Reconnaissance of zeolite deposits in tuffaceous rocks of the western Mojave Desert and vicinity, California, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C114-C116, 1964.
 1-65. (GUDE, A. J., 3d, and MUNSON, E. L.) Chemical composition of diagenetic zeolites from tuffaceous rocks of the Mojave Desert and vicinity, California: *Am. Mineralogist*, v. 50, nos. 1-2, p. 244-249, 1965.
- SHERIDAN, M. F.
 1-64. (and CROWDER, D. F.) Geologic map of the southwest quarter of the White Mountain Peak quadrangle, Mono County, California: U.S. Geol. Survey open-file report, 1964.
 2-64. (and CROWDER, D. F.) Geologic map of the northwest quarter of the White Mountain Peak quadrangle, Mono County, California: U.S. Geol. Survey open-file report, 1964.
- SHERWOOD, C. B.
 1-64. (and KLEIN, Howard) Surface- and ground-water relation in a highly permeable environment, in *Symposium—Surface waters*, Gen. Assembly of Berkeley of I. U. G. G., 1963: *Internat. Assoc. Sci. Hydrology Pub.* 63, p. 454-468, 1964.
- SHOEMAKER, E. M.
 1-63. (and EGGLETON, R.) Re-examination of the stratigraphy and structure of Sierra Madera, Texas, in *Astrogeologic studies, annual progress report*, August 25, 1962-July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 98-106, 1963.
- SHOWN, L. M.
 1-64. (MILLER, R. F., McQUEEN, I. S., and BULLER, William) Comparability of soil saturation percentage with soil moisture equivalent measured in a cooled, humidified centrifuge [abs.]: *Am. Soc. Agronomy and Soil Sci. Soc. America Ann. Mtg.*, Kansas City, Missouri, Nov. 1964, Program "Agronomy Abstracts", p. 50, 1964.
- SHOWS, T. N.
 1-64. (and GOLDEN, H. G.) Water resources studies in southeast Mississippi: U.S. Geol. Survey open-file report, 9 p., 1964.
- SIEMS, P. L.
 1-65. The geology of the Tertiary rocks of the central and southern parts of the Rosita quadrangle, Colorado: U.S. Geol. Survey open-file report, 11 p., 1965.
- SIGAFOOS, R. S.
 1-64. Botanical evidence of floods and flood-plain deposition: U.S. Geol. Survey Prof. Paper 485-A, p. A1-A35, 1964.
- SILBERLING, N. J.
 1-64. Mesozoic rocks, in *Mineral and water resources of Nevada*: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 27-30, 1964.
- SIMONS, D. B.
 1-65. (RICHARDSON, E. V., and NORDIN, C. F., Jr.) Bedload equation for ripples and dunes: U.S. Geol. Survey Prof. Paper 462-H, p. H1-H9, 1965.
 2-65. (and RICHARDSON, E. V.) Resistance to flow in alluvial channels: U.S. Geol. Survey open-file report, 186 p., 1965.
- SIMONS, F. S.
 1-64. Geology of the Klondyke quadrangle, Graham and Pinal Counties, Arizona: U.S. Geol. Survey Prof. Paper 461, 173 p., 1964.
- SIMS, P. K.
 1-64. Geologic map of the Central City quadrangle, Colorado: U.S. Geol. Survey Geol. Quad. Map GQ-267, 1964.
 2-64. (and GABLE, D. J.) Geology of Precambrian rocks, Central City district, Colorado: U.S. Geol. Survey Prof. Paper 474-C, p. C1-C52, 1964[1965].
 3-64. (SHERIDAN, D. M., KING, R. U., MOORE, E. B., RICHTER, D. H., and SCHLOTTMAN, J. D.) Geology of the uranium deposits in the Front Range, Colorado: U.S. Geol. Survey Bull. 1159, 116 p., 1964.
- SINCLAIR, W. C.
 1-63. (and LOELTZ, O. J.) Ground-water conditions in the Fernley-Wadsworth area, Churchill, Lyon, Storey, and Washoe Counties, Nevada: U.S. Geol. Survey Water-Supply Paper 1619-AA, p. AA1-AA22, 1963; also in *Nevada Dept. Conserv. and Nat. Resources Water Resources Bull.* 17, 1963.
- SIPLE, G. E.
 1-64. Artesian aquifer systems of the central Savannah River area, South Carolina-Georgia [abs.]: *Geol. Soc. America Ann. Mtg.*, Miami Beach, Florida, Nov. 1964, Program, p. 187, 1964.
 2-64. Geohydrology of storage of radioactive waste in crystalline rocks at the AEC Savannah River plant, South Carolina, in *Geological Survey Research* 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C180-C184, 1964.
- SISCO, H. G.
 1-64. (and LUSCOMBE, R. W.) Records of wells and water-level fluctuations in the Aberdeen-Springfield area, Bingham and Power Counties, Idaho, in 1963: U.S. Geol. Survey open-file report, 28 p., 1964.
- SKELTON, John
 1-63. Surface water investigations in Lower Little Creek basin, Lamar and Marion Counties, Mississippi: U.S. Geol. Survey Tech. Letter, DRIBBLE-33, 41 p., 1963.
- SKIBITZKE, H. E.
 1-64. Extending Darcy's concept of ground-water motion: U.S. Geol. Survey Prof. Paper 411-F, p. F1-F6, 1964.
- SKINNER, B. J.
 1-63. (and APPLEMAN, D. E.) Melanophlogite, a cubic polymorph of silica: *Am. Mineralogist*, v. 48, p. 854-867, 1963.
 2-63. (and FAHEY, J. J.) Observations on the inversion of stishovite to silica glass, in *Astrogeologic studies, annual progress report*, August 25, 1962-July 1, 1963—Pt. B, Crater investigations: U.S. Geol. Survey open-file report, p. 1-28, 1963.
 1-64. (BOYD, F. R., and ENGLAND, J. L.) A high-pressure polymorph of chalcocite, Cu₂S [abs.]: *Am. Geophys. Union Trans.*, v. 45, no. 1, p. 121-122, 1964.
 2-64. (and BOYD, F. R.) Aluminous enstatite: *Carnegie Inst. Washington, Year Book* 1963-1964, p. 163, 1964.
 1-65. The system arsenic-antimony: *Econ. Geology*, v. 60, no. 2, p. 228-239, 1965.
 2-65. The system Cu-Ag-S [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 188, 1965.

- SLOSS, Raymond
1-65. (and BROUSSARD, W. L.) Surface-water resources, Jefferson Davis Parish, Louisiana, *in* Jefferson Davis Parish resources and facilities: Louisiana Dept. Public Works, p. 31-41, 1965.
- SMITH, C. R.
1-64. Current publications of the Arizona district, *in* Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 3-8, 1964.
2-64. Gila County, *in* Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 56, 1964.
- SMITH, G. I.
1-64. Geology and volcanic petrology of the Lava Mountains, San Bernardino County, California: U.S. Geol. Survey Prof. Paper 457, 97 p., 1964[1965].
2-64. (and HAINES, D. V.) Character and distribution of nonclastic minerals in the Searles Lake evaporite deposit, California: U.S. Geol. Survey Bull. 1181-P, p. P1-P58, 1964.
- SMITH, J. H.
1-64. Geology of the sedimentary rocks of the Morrison quadrangle, Colorado: U.S. Geol. Survey Misc. Geol. Inv. Map I-428, 1964.
- SMITH, M. B.
1-64. Map showing distribution and configuration of basement rocks in California: U.S. Geol. Survey Oil and Gas Inv. Map OM-215, 1964.
- SMITH, P. B.
1-64. Ecology of benthonic species, *in* Recent foraminifera off Central America: U.S. Geol. Survey Prof. Paper 429-B, p. B1-B55, 1964.
2-64. Foraminifera from the Bootlegger Cove clay, Anchorage, Alaska, *in* Report on Anchorage area soil studies, Alaska: Seattle, Washington, Shannon and Wilson, Inc., App., p. J1-J5, 1964.
- SMITH, W. C.
1-64. Nonmetallic mineral resources: Borates, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 180-184, 1964.
- SMITH, W. O.
1-64. Review of "Theory of ground-water movement," by Poludarinova-Kochina: Am. Geophys. Union Trans., v. 44, no. 3, p. 732-733, 1964.
- SNARELY, P. D., Jr.
1-64. (RAU, W. W., and WAGNER, H. C.) Miocene stratigraphy of the Yaquina Bay area, Newport, Oregon: Ore Bin, v. 26, no. 8, p. 133-151, 1964.
2-64. (and WAGNER, H. C.) Geologic sketch of northwestern Oregon, *with* Geologic interpretation of reconnaissance gravity and aeromagnetic surveys in northwestern Oregon, by R. W. Bromery and P. D. Snarely, Jr.: U.S. Geol. Survey Bull. 1181-M, p. M1-M17, 1964.
3-64. (RAU, W. W., and WAGNER, H. C.) Miocene stratigraphy of the Yaquina Bay area, Newport, Oregon: The Ore Bin, v. 26, no. 8, p. 133-151, 1964.
- SNIEGOCKI, R. T.
1-65. (BAYLEY, F. H., 3d, ENGLER, Kyle, and STEPHENS, J. W.) Testing procedures and results of studies of artificial recharge in the Grand Prairie region, Arkansas: U.S. Geol. Survey Water-Supply Paper 1615-G, p. G1-G56, 1965.
- SNYDER, C. T.
1-64. (HARDMAN, George, and ZDENEK, F. F.) Pleistocene lakes in the Great Basin: U.S. Geol. Survey Misc. Geol. Inv. Map I-416, 1964.
- SNYDER, G. L.
1-64. Bedrock geology of the Willimantic quadrangle, Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-335, 1964.
- SNYDER, G. L. --Continued
2-64. Petrochemistry and bedrock geology of the Fitchville quadrangle, Connecticut: U.S. Geol. Survey Bull. 1161-I, p. I1-I63, 1964.
- SNYDER, R. P.
1-63. Bibliography of U.S. Geological Survey unclassified reports of investigations at nuclear test sites in the United States, fiscal years 1958-1962: U.S. Geol. Survey Tech. Letter, Spec. Proj. -6, 58 p., 1963.
2-63. Bibliography of U.S. Geological Survey unclassified reports of investigations at nuclear test sites in the United States, fiscal year 1963: U.S. Geol. Survey Tech. Letter, Spec. Proj. -8, 42 p., 1963.
3-63. (HOOVER, D. L., HASLER, J. W., WILLIAMS, W. P.) Geology of the U3cw site and the area around Well E, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-52, 18 p., 1963.
- SOHL, N. F.
1-64. Gastropods from the Coffee Sand (Upper Cretaceous) of Mississippi: U.S. Geol. Survey Prof. Paper 331-C, p. C345-C394, 1964.
2-64. Neogastropoda, Opisthobranchia, and Basommatophora from the Ripley, Owl Creek, and Prairie Bluff Formations: U.S. Geol. Survey Prof. Paper 331-B, p. B153-B344, 1964.
3-64. Pre-Selma larger invertebrate fossils from well core samples in western Alabama: U.S. Geol. Survey Bull. 1160-C, p. C55-C64, 1964.
4-64. (and KAUFFMAN, E. G.) Giant Upper Cretaceous oysters from the Gulf coast and Caribbean: U.S. Geol. Survey Prof. Paper 483-H, p. H1-H22, 1964.
1-65. Collecting in unconsolidated sediments, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 155-159, 1965.
2-65. (KAUFFMAN, E. G., and YOCHELSON, E. L.) Gastropods and pelecypods, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Co., p. 49-53, 1965.
- SOHN, I. G.
1-64. Significance of Triassic ostracodes from Alaska and Nevada, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D40-D42, 1964.
2-64. The ostracode genus *Cytherelloidea*, a possible indicator of paleotemperature: Pub. Stazione Zoologica di Napoli, v. 33, supp., p. 529-534, 1964[1965].
1-65. Classification of the superfamily Healdiacea and the genus *Pseudophanasymmetria* Sohn and Berdan, 1952 (Ostracoda), *in* Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B69-B72, 1965.
2-65. Late Quaternary ostracodes from the southern part of the Dead Sea, Israel, *in* Ocean Sci. and Ocean Eng. Conf., Washington, 1965: Marine Technology Soc. and Am. Soc. Limnology Oceanography Trans., v. 1, p. 82-94, 1965.
3-65. Significance of new marine Triassic ostracodes from North America and Eurasia [abs.]: Geol. Soc. America Spec. Paper 82, p. 190, 1965.
4-65. (BERDAN, J. M., and PECK, R.) Ostracodes, *in* Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Co., p. 75-89, 1965.
- SOISTER, P. E.
1-65. Geologic map of the Fort Lupton quadrangle, Weld Adams Counties, Colorado: U.S. Geol. Survey Geol. Quad. Map GQ-397, 1965.
2-65. Geologic map of the Hudson quadrangle, Weld and Adams Counties, Colorado: U.S. Geol. Survey Geol. Quad. Map GQ-398, 1965.
3-65. Geologic map of the Platteville quadrangle, Weld County, Colorado: U.S. Geol. Survey Geol. Quad. Map GQ-399, 1965.

SOUTHWICK, D. L.

- 1-64. Petrography of the basement gneiss beneath the Coastal Plain sequence, Island Beach State Park, New Jersey, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C55-C60, 1964.

SPEER, P. R.

- 1-64. (GOLDEN, H. G., and PATTERSON, J. F.) Low-flow characteristics of streams in the Mississippi embayment in Mississippi and Alabama, *with a section on* Quality of the water, by W. J. Welborne: U.S. Geol. Survey Prof. Paper 448-I, p. I1-I47, 1964[1965].
- 2-64. (HINES, M. S., CALANDRO, A. J., and others) Low-flow characteristics of streams in the Mississippi Embayment in southern Arkansas, northern Louisiana, and northeastern Texas, *with a section on* Quality of the water by H. G. Jeffery: U.S. Geol. Survey open-file report, 132 p., 1964.
- 3-64. (HINES, M. S., JANSON, M. E., and others) Low-flow characteristics of streams in the Mississippi Embayment in northern Arkansas and in Missouri, *with a section on* Quality of the water, by H. G. Jeffery: U.S. Geol. Survey open-file report, 101 p., 1964.

SPENCER, F. D.

- 1-64. Geology of the Boltsfork quadrangle and part of the Burnaugh quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-316, 1964.
- 2-64. Geology of the Tell City quadrangle, Kentucky-Indiana: U.S. Geol. Survey Geol. Quad. Map GQ-356, 1964.
- 3-64. Geology of parts of the Rockport and Lewisport quadrangles, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-315, 1964.

SPICER, H. C.

- 1-64. A compilation of deep earth temperature data: U.S.A. 1910-1945: U.S. Geol. Survey open-file report, 3 p., 1964.
- 2-64. Geothermal gradients and heat flow in the Salt Valley anticline, Utah: Boll. Geofisica Teor. ed Appl., v. 6, no. 23, p. 263-282, 1964.

SPIEKER, A. M.

- 1-64. Analog model study of the effect of future pumping from Pleistocene deposits in the Great Miami River Valley southwest of Hamilton, Ohio [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 193, 1964.
- 2-64. Effect of increased pumping of ground water in the Fairfield-New Baltimore area, Ohio—a prediction by analog model study: U.S. Geol. Survey open-file report, 105 p., 1964.
- 1-65. Hydrogeologic aspects of an analog model study of the Fairfield-New Baltimore area, Ohio: U.S. Geol. Survey open-file report, 121 p., 1965.

STAATZ, M. H.

- 1-64. Geology of the Bald Knob quadrangle, Ferry and Okanogan Counties, Washington: U.S. Geol. Survey Bull. 1161-F, p. F1-F79, 1964.
- 2-64. Metallic mineral resources: Thorium and the rare earths, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 150-154, 1964.
- 3-64. Miscellaneous metals: Thorium, *in* Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 132-135, 1964.
- 4-64. (and CARR, W. J.) Geology and mineral deposits of the Thomas and Dugway Ranges, Juab and Tooele Counties, Utah: U.S. Geol. Survey Prof. Paper 415, 188 p., 1964.
- 1-65. Thorium, *in* Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 230-234, 1965.

STAGER, H. K.

- 1-64. Geology of the Blackwater quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-304, 1964.

STALLMAN, R. W.

- 1-64. Multiphase fluids in porous media—A review of theories pertinent to hydrologic studies: U.S. Geol. Survey Prof. Paper 411-E, p. E1-E51, 1964.

STALLMAN, R. W.--Continued

- 1-65. Steady one-dimensional fluidflow in a semi-infinite porous medium with semisoidal surface temperature: Jour. Geophys. Research, v. 70, no. 12, p. 2821-2827, 1965.

STEAD, F. W.

- 1-64. Distribution in groundwater of radio nuclides from underground nuclear explosions, *in* Engineering with nuclear explosives—Proceedings of the Third Plowshare Symposium, April 21-23, 1964: U.S. Atomic Energy Comm. Rept. TID-7695, p. 127-138, 1964.

STEECE, F. V.

- 1-64. (and HOWELLS, Lewis) Geology and ground-water supplies in Sanborn County, South Dakota: U.S. Geol. Survey open-file report, 334 p., 1964.

STEELE, C. E.

- 1-64. (and BARCLAY, J. E.) Ground-water resources of Harmon County and adjacent parts of Greer and Jackson Counties, Oklahoma: U.S. Geol. Survey open-file report, 425 p., 1964.

STEPHENS, J. G.

- 1-64. Geology and uranium deposits at Crooks Gap, Fremont County, Wyoming, *with a section on* Gravity and seismic studies in the Crooks Gap area by D. L. Healey: U.S. Geol. Survey Bull. 1147-F, p. F1-F82, 1964.

STERN, T. W.

- 1-65. (GOLDICH, S. S., and NEWELL, M. F.) Effects of weathering on the U-Pb and Th-Pb systems in zircon [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 164-165, 1965.
- 2-65. (NEWELL, M. F., KISTLER, R. W., and SHAW, D. R.) Zircon uranium-lead and thorium-lead ages and mineral potassium-argon ages of La Sal Mountains rocks, Utah: Jour. Geophys. Research, v. 70, no. 6, p. 1503-1507, 1965.

STEWART, D. B.

- 1-65. Review of "Aspects of theoretical mineralogy in the U.S.S.R.": Am. Scientist, v. 53, no. 2, p. 262A, 1965.

STEWART, F. H.

- 1-63. Marine evaporites, *in* Data of geochemistry, 6th ed.: U.S. Geol. Survey Prof. Paper 440-Y, p. Y1-Y53, 1963.

STEWART, J. H.

- 1-64. Precambrian and Lower Cambrian rocks, *in* Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 21, 1964.

STEWART, J. W.

- 1-64. (CALLAHAN, J. T., and CARTER, R. F.) Geologic and hydrologic investigation at the site of the Georgia Nuclear Laboratory, Dawson County, Georgia: U.S. Geol. Survey Bull. 1133-F, p. F1-F90, 1964.

- 2-64. Infiltration and permeability of weathered crystalline rocks, Georgia Nuclear Laboratory, Dawson County, Georgia: U.S. Geol. Survey Bull. 1133-D, 57 p., 1964.

STRINGFIELD, V. T.

- 1-64. Relation to surface-water hydrology to the principal artesian aquifer in Florida and southeastern Georgia, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C164-C169, 1964.

- 2-64. (and LeGRAND, H. E.) Hydrology of limestone terranes in the Coastal Plain of the southeastern states [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 200, 1964.

STROMQUIST, A. A.

- 1-64. (WHITE, A. M., and McHUGH, J. B.) A geochemical investigation of the High Rock quadrangle, North Carolina, *in* Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C88-C91, 1964.

STUART, D. J.

- 1-65. Gravity data and Bouguer-gravity map for western Washington: U.S. Geol. Survey open-file report, 7 p., 46 gravity sheets, 1965.

- STULIK, R. S.
 1-64. Effect of ground-water withdrawal, 1954-63, in the lower Harquahala Plains area, Maricopa County, Arizona: U.S. Geol. Survey open-file report, 10 p., 1964.
 2-64. Effects of ground-water withdrawal, 1954-63, in the lower Harquahala Plains, Maricopa County, Arizona: Arizona State Land Dept. Water Resources Rept., 17, 8 p., 1964.
 3-64. Salt River Valley, Waterman Wash, Gila Bend, Harquahala Plains, McMullen Valley, Palomas Plain, Ranegras Plain areas, Big Sandy Valley, [and] Kingman-Hackberry area, in Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona Land Dept. Water Resources Rept. 19, p. 35-48, 1964.
- SUNDELIUS, H. W.
 1-64. (and BELL, Henry, 3d) An unusual radioactive, rare-earth-bearing sulfide deposit in Cabarrus County, North Carolina: Southeastern Geology, v. 5, no. 4, p. 207-221, 1964.
- SWAIN, F. M.
 1-64. (and BROWN, P. M.) Cretaceous Ostracoda from wells in the southeastern United States: North Carolina Dept. Conserv. Div. Mineral Resources Bull., no. 78, 55 p., 1964.
- SWANSON, V. E.
 1-64. (and BERRYHILL, H. L., Jr.) Geology of the Washington East quadrangle, Pennsylvania: U.S. Geol. Survey Geol. Quad. Map GQ-334, 1964.
 2-64. (and LANDIS, E. R.) Geology of a uranium-bearing black shale of Late Devonian age in north-central Arkansas: Arkansas Geol. and Conserv. Comm. Inf. Circ. 22, 16 p., 1964.
 1-65. (and PALACAS, J. G.) Humate in coastal sands of northwest Florida: U.S. Geol. Survey Bull. 1214-B, p. B1-B29, 1965.
- SWENSON, H. A.
 1-64. Sediment in streams: Jour. Soil and Water Conserv., v. 19, no. 6, p. 223-226, 1964.
 1-65. Minerals in water: Sci. Teacher, v. 32, no. 1, p. 17-21, 1965.
 2-65. (and BALDWIN, H. L.) A primer on water quality: U.S. Geol. Survey Misc. Rept., 27 p., 1965.
- SWINDEL, G. W.
 1-64. (WILLIAMS, M. R., and GEURIN, J. W.) Water in Alabama; revised by H. L. Baldwin, 1963: U.S. Geol. Survey Water-Supply Paper 1765, 89 p., 1964.
- SWITZER, George
 1-65. (CLARKE, R. S., Jr., SINKANKAS, John, and WORTHING, H. W.) Fluorine in hambergite: Am. Mineralogist, v. 50, nos. 1-2, p. 85-95, 1965.
- TAILLEUR, I. L.
 1-65. Low-volatile bituminous coal of Mississippian age on the Lisburne Peninsula, northwestern Alaska, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B34-B38, 1965.
- TANAKA, H. H.
 1-64. (and HOLLOWELL, J. R.) Hydrology of the alluvium of the Arkansas River, Muskogee, Oklahoma, to Fort Smith, Arkansas: U.S. Geol. Survey open-file report, 92 p., 1964.
- TANGBORN, W. V.
 1-64. Glacier mass budget measurements by hydro-meteorological means [abs.]: Am. Geophys. Union Trans., v. 45, no. 4, p. 610, 1964.
- TANNER, A. B.
 1-64. Physical and chemical controls on distribution of radium-226 and radon-222 in ground water near Great Salt Lake, Utah, in The natural radiation environment: Chicago, Illinois, Univ. Chicago Press (for Rice Univ.), p. 253-278, 1964.
 2-64. Radon migration in the ground—A review, in The natural radiation environment: Chicago, Illinois, Univ. Chicago Press (for Rice Univ.), p. 161-190, 1964.
- TARVER, G. R.
 1-65. Ground-water resources of Houston County, Texas: U.S. Geol. Survey open-file report, 133 p., 1965.
- TATLOCK, D. B.
 1-64. Nonmetallic mineral resources: Kyanite group aluminous minerals, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 213-217, 1964.
- TATSUMOTO, M.
 1-65. Isotopic composition of lead in basalts [abs.]: Am. Geophys. Union Trans., v. 46, p. 165, 1965.
 2-65. Uranium, thorium, and lead abundances in volcanic rocks from Hawaii, Iwo Jima, and Japan [abs.]: Geol. Soc. America Spec. Paper 82, p. 205, 1965.
- TAYLOR, A. R.
 1-64. Geology of the Rewey and Mifflin quadrangles, Wisconsin: U.S. Geol. Survey Bull., 1123-F, p. F279-F360, 1964.
- TAYLOR, C. T.
 1-64. Chemical quality of Missouri surface water: Missouri Water Pollution Board, 28 p., 1964.
- TAYLOR, G. C., Jr.
 1-65. Hydrology—the science of water: Sci. Teacher, v. 32, no. 1, p. 26-29, 1965.
- TAYLOR, H. F., Jr.
 1-64. (DUKE, M. B., SILVER, L. T., and EPSTEIN, S.) Oxygen isotopic studies of minerals in stony meteorites [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 112, 1964.
- TAYLOR, O. J.
 1-65. Ground-water resources along Cedar Creek anticline in eastern Montana: Montana Bur. Mines and Geology Mem. 40, 99 p., 1965.
- TAYLOR, R. E.
 1-63. (and GRANTHAM, P. E.) Basic data for domestic supply wells in a five-mile radius of Tatum Salt Dome Lamar County, Mississippi—A supplement to Technical Letter DRIBBLE-30: U.S. Geol. Survey Tech. Letter, DRIBBLE-38, 3 p., 1963.
- TEICHERT, Curt
 1-64. Recent German work on the Cambrian and Saline series of the Salt Range, West Pakistan: Geol. Survey of Pakistan Records, v. 11, pt. 1, 8 p., 1964.
 2-64. Some biological and paleogeographic factors in the evaluation of ancient climates, in Problems in paleo-climatology: New York, John Wiley and Sons, p. 597-600, 1964.
- THADEN, R. E.
 1-64. (TRITES, A. F., Jr., and FINNELL, T. L.) Geology and ore deposits of the White Canyon area, San Juan and Garfield Counties, Utah: U.S. Geol. Survey Bull. 1125, 166 p., 1964[1965].
- THAYER, T. P.
 1-64. Geologic features of podiform chromite deposits, in Methods of prospecting for chromite: Organization for Econ. Cooperation and Development, Paris, p. 135-148, 1964.
 2-64. Principal features and origin of podiform chromite deposits, and some observations on the Guleman-Soridag district, Turkey: Econ. Geology, v. 59, no. 8, p. 1497-1524, 1964.
 3-64. (and BROWN, C. E.) Pre-Tertiary orogenic and plutonic intrusive activity in central and northeastern Oregon: Geol. Soc. America Bull., v. 75, no. 12, p. 1255-1261, 1964.
- 1-65. Review of "Les peridotites serpentinisées en France, Group 1, Peridotites intracrystallines, Pt. 4, Massif Central Median: Bassin du Haut-Allier" by F. H. Forestier: Science, v. 147, no. 3662, p. 1137, 1965.
- THEOBALD, P. K.
 1-65. Preliminary geologic map of the Berthoud Pass quadrangle, Clear Creek and Grand Counties, Colorado: U.S. Geol. Survey Misc. Geol. Inv. Map I-443, 1965.

- THOMAS, D. M.
 1-64. Floods in New Jersey, magnitude and frequency: New Jersey Dept. Conserv. and Econ. Devel., Div. of Water Policy and Supply, Water Resources Circ. 13, 145 p., 1964.
 2-64. Flood-depth frequency in New Jersey: New Jersey Dept. of Conserv. and Econ. Devel., Div. of Water Policy and Supply, Water Resources Circ. 14, 14 p., 1964.
- THOMAS, H. E.
 1-64. Cultural control of water development: CIBA Jour., no. 31, p. 25-32, 1964.
 2-64. Water resources, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 273-314, 1964.
 1-65. Career, anyone?: Ground Water, v. 3, no. 1, p. 2-4, 1965.
 2-65. Our cultural approach to water rights: Ground Water, v. 3, no. 2, p. 19-21, 1965.
- THOMPSON, G. A.
 1-64. (and WHITE, D. E.) Regional geology of the Steamboat Springs area, Washoe County, Nevada: U.S. Geol. Survey Prof. Paper 458-A, p. A1-A52, 1964.
- THOMPSON, M. M.
 1-64. (and LEWIS, J. G.) Practical improvements in stereoplotting instruments: Photogramm. Eng., v. 30, no. 5, p. 802-809, 1964.
 2-64. (and SPEERT, J. L.) Mapping the surface of the Earth: Nat. History, v. 73, no. 8, p. 30-37, 1964.
- THOMSON, B. W.
 1-64. (and SCHUMANN, H. H.) Water resources of the Sycamore Creek watershed, Maricopa County, Arizona, a progress report: U.S. Geol. Survey open-file report, 28 p., 1964.
- THORPE, A. N.
 1-63. (and SENFTLE, F. E.) Submicroscopic spherules and color of tektites, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. C, Cosmochemistry and petrography: U.S. Geol. Survey open-file report, p. 109-136, 1963.
 1-64. Magnetic properties of Pd, Pd-H, and Pd-D from 300°K to 4.2°K: U.S. Geol. Survey open-file report, 56 p., 1964.
- TILLING, R. I.
 1-64. Variation in modes and norms of an "homogeneous" pluton of the Boulder batholith, Montana, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D8-D13, 1964.
- TILTON, G. R.
 1-64. (GAST, P. W., and HEDGE, C. E.) Variation in the isotopic composition of strontium and lead from basalts [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 109, 1964.
 1-65. (and DOE, B. R.) Lead isotopes and the age of the Earth [abs.]: Geol. Soc. America Spec. Paper 82, p. 207, 1965.
- TITLEY, S. R.
 1-63. A summary of the geology of the Mare Humorum quadrangle of the Moon, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 64-72, 1963.
- TITUS, F. B., Jr.
 1-65. New Mexico State Planning Office Report—Central closed basins: U.S. Geol. Survey open-file report, 36 p., 1965.
- TOCHER, Don
 1-64. Earthquakes and rating scales: Geotimes, v. 8, no. 8, p. 15-18, 1964.
- TODD, D. K.
 1-65. Annotated bibliography on artificial recharge of ground water through 1954: U.S. Geol. Survey Water-Supply Paper 1477, 115 p., 1965. (Originally published 1959.)
- TODD, Ruth
 1-64. Planktonic Foraminifera from deep-sea cores off Eniwetok Atoll: U.S. Geol. Survey Prof. Paper 260-CC, p. CC1067-CC1100, 1964.
 2-64. Recent literature on the Foraminifera: Cushman Found. Foram. Research Contr., v. 15, pt. 1, p. 39-44, 1964; ibid., pt. 2, p. 73-77; ibid., pt. 3, p. 124-129;
 1-65. Ibid., v. 16, pt. 1, p. 46-52; ibid., pt. 3, p. 87-94, 1965.
 2-65. (LOW, Doris, and MELLO, J. F.) Smaller foraminifers, in Handbook of paleontological techniques: San Francisco, Calif., W. H. Freeman and Company, p. 14-20, 1965.
- TOLER, L. G.
 1-64. (MUSGROVE, R. H., and FOSTER, J. F.) Freshening of Deer Point Lake, Bay County, Florida: Am. Water Works Assoc. Jour., v. 56, no. 8, p. 984-990, 1964.
- TOOKER, E. W.
 1-64. (and STEWART, J. H.) Geology: Stratigraphy, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 19-25, 1964.
 2-64. (and STEWART, J. H.) Geology: Topography, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 17-18, 1964.
- TOULMIN, L. D.
 1-64. (LAMOREAUX, P. E., and NEWTON, J. G.) Profile showing geology along the Chattahoochee River, Alabama: Alabama Geol. Survey Map 28, 1 sheet, 1964.
- TOULMIN, Priestley, 3d
 1-64. Igneous rocks of the Salem area, Massachusetts, Trip E, in Guidebook to field trips in the Boston area and vicinity: New England Intercollegiate Geol. Conf., 56th Ann. Mtg., Yale Univ., Dept. Geology, p. 67-79, 1964.
- TOURTELOT, H. A.
 1-64. Bauxite deposits of the Tippah-Benton district, Mississippi: U.S. Geol. Survey Bull. 1199-C, p. C1-C31, 1964[1965].
 2-64. Minor-element composition and organic carbon content of marine and nonmarine shales of Late Cretaceous age in the western interior of the United States: Geochim. et Cosmochim. Acta, v. 28, no. 10, p. 1579-1604, 1964.
- TRACEY, J. I., Jr.
 1-64. (SCHLANGER, S. O., STARK, J. T., and others) General geology of Guam: U.S. Geol. Survey Prof. Paper 403-A, p. A1-A104, 1964.
- TRACY, H. J.
 1-65. Discharge characteristics of broad-crested weirs: U.S. Geol. Survey Circ. 397, 15 p., 1965. (Originally published 1957.)
- TRAINER, F. W.
 1-64. (and EDDY, J. E.) A periscope for the study of borehole walls, and its use in ground-water studies in Niagara County, New York, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D203-D207, 1964.
- TRAUGER, F. D.
 1-64. (and BUSHMAN, F. X.) Geology and ground water in the vicinity of Tucumcari, Quay County, New Mexico: New Mexico State Engineer Tech. Rept. 30, 178 p., 1964.
 1-65. New Mexico State Planning Office Report—Lower Colorado River basin: U.S. Geol. Survey open-file report, 41 p., 1965.
- TRAVIS, W. I.
 1-64. (WAITE, H. A., and SANTOS, J. F.) Water resources: Introduction, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 255-308, 1964.
- TRENT, V. A.
 1-65. Geology of the Matewan quadrangle in Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-373, 1965.

TREXLER, J. P.

- 1-64. (WOOD, G. H., Jr., and ARNDT, H. H.) Progress of U.S. Geological Survey investigations in the Anthracite region of Pennsylvania: *Pennsylvania Acad. Sci. Proc.* 1963, v. 37, p. 295-299, 1964.

TRIMBLE, D. E.

- 1-65. (and SCHAEFFER, F. E.) Stratigraphy of the Precambrian rocks and lowest Cambrian rocks in the Pocatello area, Idaho [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 349, 1965.

TRUEDELL, A. H.

- 1-65. (JONES, B. F., and Van DENBURGH, A. S.) Glass electrode determination of sodium in closed basin waters: *Geochim. Cosmochim. Acta*, v. 29, no. 9, p. 725-737, 1965.

TSCHUDY, R. H.

- 1-64. Palynology and time-stratigraphic determinations, in *Palynology in oil exploration—A symposium*, San Francisco, California 1962: *Soc. Econ. Paleontologists and Mineralogists Spec. Pub.* 11, p. 18-28, 1964.
 1-65. An Upper Cretaceous deposit in the Appalachian Mountains, in *Geological Survey Research 1965*: U.S. Geol. Survey Prof. Paper 525-B, p. B64-B68, 1965.
 2-65. Palynological investigations in the Upper Cretaceous and Tertiary of the Mississippi Embayment region—I: U.S. Geol. Survey open-file report, 63 p., 1965.
 3-65. Plant and miscellaneous microfossils from the Parachute Creek Member of the Green River Formation: U.S. Geol. Survey open-file report, 3 p., 1965.

TURCAN, A. N., Jr.

- 1-65. Calculation of water quality from electrical logs—Theory and practice [abs.]: *Am. Geophys. Union Trans.* v. 46, no. 1, p. 76-77, 1965.
 2-65. Ground water for rice irrigation in southwestern Louisiana: *The Rice Jour.*, v. 68, no. 3, p. 32-33, 1965.

TWENTER, F. R.

- 1-65. (and COBLE, R. W.) The water story in central Iowa: *Iowa Geol. Survey Water Atlas*, 89 p., 1965.

TWETO, Ogden

- 1-64. Geology, in *Mineral and water resources of Colorado*: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 11-27, 1964.

ULRICH, G. E.

- 1-65. New evidence for the origin of the Williams Range thrust fault, Front Range, Colorado [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 210-211, 1965.

UNITED STATES DEPARTMENT OF THE INTERIOR

- 1-64. Geological Survey in progress: *Missouri River Basin Field Comm. Program Issue*, p. 12-17, 1964.

UNITED STATES GEOLOGICAL SURVEY

- 1-64. Bibliography of North American geology, 1960: U.S. Geol. Survey Bull. 1196, 777 p., 1964.
 3-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 1-A, North Atlantic slope basins, Maine to Connecticut: U.S. Geol. Survey Water-Supply Paper 1721, 317 p., 1964.
 4-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 1-B, North Atlantic slope basins, New York to York River: U.S. Geol. Survey Water-Supply Paper 1722, 578 p., 1964.
 5-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 3-B, Cumberland and Tennessee River basins: U.S. Geol. Survey Water-Supply Paper 1726, 269 p., 1964.
 6-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 4, St. Lawrence River basin: U.S. Geol. Survey Water-Supply Paper 1727, 379 p., 1964.
 7-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 5, Hudson Bay and upper Mississippi River basins: U.S. Geol. Survey Water-Supply Paper 1728, 576 p., 1964.

UNITED STATES GEOLOGICAL SURVEY--Continued

- 8-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 6-A, Missouri River basin above Sioux City, Iowa: U.S. Geol. Survey Water-Supply Paper 1729, 507 p., 1964.
 9-64. Compilation of records of surface waters of the United States, October 1950 to September 1960—Pt. 7, Lower Mississippi River basin: U.S. Geol. Survey Water-Supply Paper 1731, 552 p., 1964.
 10-64. Geological Survey, in *Missouri River Basin progress annual report 1964*: Billings, Montana, Interior Missouri River Basin Field Comm., 191 p., 1964.
 11-64. Geological Survey research 1964—Chap. A: U.S. Geol. Survey Prof. Paper 501-A, p. A1-A367, 1964.
 12-64. Mineral and water resources of Colorado: U.S. Cong., 88th, 2d sess., Senate Comm. Interior and Insular Affairs, Comm. Print, 302 p., 1964.
 13-64. Summary of floods in the United States during 1959: U.S. Geol. Survey Water-Supply Paper 1750-B, p. B1-B101, 1964.
 15-64. Astrogeologic studies, annual progress report, August 25, 1962, to July 1, 1963: U.S. Geol. Survey open-file report, 578 p., 1964.
 16-64. Mineral and water resources of Idaho: Washington, D. C., U.S. Cong., 88th, 2d sess., Senate Comm. Interior and Insular Affairs, Comm. Print, 335 p., 1964.
 17-64. Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 85, 314 p., 1964.
 1-65. Astrogeologic Studies, Annual Progress Report, July 1, 1963 to July 1, 1964—Pts. A, B, C, and supplement: U.S. Geol. Survey open-file report, 450 p., 1965.
 2-65. Good Friday quake revisited: *California Div. Mines and Geology Mineral Inf. Service*, v. 18, no. 5, p. 100-101, 1965.
 3-65. Quality of surface waters of the United States, 1959—Pts. 1-2, North Atlantic slope basins and south Atlantic slope and eastern Gulf of Mexico basins: U.S. Geol. Survey Water-Supply Paper 1641, 460 p., 1965.
 4-65. Ground-water levels in the United States, 1958-62, Northeastern States: U.S. Geol. Survey Water-Supply Paper 1782, 252 p., 1965.
 5-65. Quality of surface waters of the United States 1959—Pts. 3-4, Ohio River basin and St. Lawrence River basin: U.S. Geol. Survey Water-Supply Paper 1642, 303 p., 1965.
 6-65. Mineral potential of eastern Montana: Washington, D. C., U.S. Cong. 89th, 1st sess., Senate Doc. no. 12, 77 p., 1965.
 7-65. Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Senate Comm. Interior and Insular Affairs, Comm. Print, 437 p., 1965.
- UNITED STATES GEOLOGICAL SURVEY, BRANCH OF COMPUTATION
- 1-64. Computer programs for the B220: Washington, D. C., U.S. Geol. Survey, 38 p., 1964.
- UNITED STATES GEOLOGICAL SURVEY, CONSERVATION DIVISION
- 1-64. Reported occurrences of selected minerals in Wyoming—Includes most metals and nonmetals; does not include mineral fuels, sodium and potassium salts, or phosphate: U.S. Geol. Survey Mineral Inv. Resource Map MR-42, 1964.
- UNITED STATES GEOLOGICAL SURVEY, GEOLOGIC DIVISION
- 1-64. Location of geologic field projects as of June 30, 1964: Washington, D. C., U.S. Geol. Survey open-file report, 1964.
 2-64. Location of geologic field projects in Alaska as of June 30, 1964: Washington, D. C., U.S. Geol. Survey open-file report, 1964.
 1-65. Field investigations in California, 1965: *California Div. Mines and Geology Mineral Inf. Service*, v. 18, no. 5, p. 81-85, 1965.

UNITED STATES GEOLOGICAL SURVEY, TOPOGRAPHIC DIVISION

- 1-64. Cartographic tables: U.S. Geol. Survey Topographic Instructions, Book 5, Chaps. 5B1 and 5B2, 142 p., 1964.
- UPSON, J. E.
1-64. (and SPENCER, C. W.) Bedrock valleys of the New England coast as related to fluctuations of sea level: U.S. Geol. Survey Prof. Paper 454-M, p. M1-M44, 1964.
- VAN ALSTINE, R. E.
1-64. Nonmetallic and industrial minerals: Fluorspar, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 160-165, 1964.
1-65. Fluorspar, in Mineral and water resources of New Mexico: Washington, D. C., U.S. Cong. 89th, 1st sess., Comm. Print, p. 260-267, 1965.
- VAN HORN, Richard
1-64. Nonmetallic and industrial minerals and materials resources: Lightweight aggregate, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 185-188, 1964.
2-64. Nonmetallic and industrial minerals and materials resources: Sand and gravel, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 215-218, 1964.
1-65. Section 2—Late Wisconsin and Recent stratigraphy in downtown Salt Lake City, Utah: Utah Geol. and Mineralog. Survey Spec. Studies, no. 11, p. 29-35, 1965.
- VAN HOUTEN, F. B.
1-64. Tertiary geology of the Beaver Rim area, Fremont and Natrona Counties, Wyoming: U.S. Geol. Survey Bull. 1164, 99 p., 1964[1965].
- VAN LIER, K. E.
1-63. Ground-water in Menominee County: Michigan Geol. Survey Div. Water Inv. 2, 42 p., 1963.
1-64. Ground water in the Tri-County Region, Michigan—Chap. 1, Tri-County regional transportation study-Natural resource problem study: Lansing, Michigan, Tri-County Regional Planning Comm., 19 p., 1964.
- VAN LIEU, J. A.
1-64. (and PATTERSON, E. D.) Coal resources of Lawrence County, Pennsylvania: U.S. Geol. Survey Bull. 1143-B, p. B1-B33, 1964.
- VAUGHN, W. W.
1-64. (and MCCARTHY, J. H., Jr.) An instrumental technique for the determination of submicrogram concentrations of mercury in soils and rocks: U.S. Geol. Survey open-file report, 10 p., 1964.
2-64. (and MCCARTHY, J. H., Jr.) An instrumental technique for the determination of submicrogram concentrations of mercury in solids, rocks, and gas, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D123-D127, 1964.
1-65. (CRAMER, W. G., and SHARP, W. N.) Gamma activation device for low-level beryllium analysis, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B151-B154, 1965.
- VEDDER, J. G.
1-65. Review of "The San Andreas fault zone from the Temblor Mountains to Antelope Valley, southern California" by J. C. Crowell: Am. Assoc. Petroleum Geologists Bull., v. 49, no. 3 (pt. 1), p. 320-321, 1965.
- VHAY, J. S.
1-64. Mineral resources: Cobalt and nickel, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 66-68, 1964.
2-64. Mineral resources: Copper, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 68-74, 1964.
- VICE, R. B.
1-64. Sedimentary program of the Geological Survey: U.S. Geol. Survey open-file report, 9 p., 1964.
- VINE, J. D.
1-65. Metals in Pennsylvanian black shale from Kentucky and Indiana [abs.]: Geol. Soc. America Spec. Paper 82, p. 212, 1965.
- VINE, J. D. --Continued
2-65. Spectrographic analyses of Paleozoic black shale samples: U.S. Geol. Survey open-file report, 5 p., 1965.
- VISHER, F. N.
1-64. Fresh water and salt water in southern Oahu, Hawaii [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, 1964, Program, p. 213, 1964.
1-65. (and WETTERHALL, W. S.) Desalination of aquifers in Florida [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 75, 1965.
- VLISSIDES, S. D.
1-64. Map of Texas showing oil and gas fields, pipelines, and areas of exposed basement rocks: U.S. Geol. Survey Oil and Gas Inv. Map OM-214, 1964.
2-64. (and QUIRIN, B. A.) Oil and gas fields of the United States, exclusive of Alaska and Hawaii: U.S. Geol. Survey Spec. Map. 2 sheets, 1964.
- VOEGELI, P. T., Sr.
1-65. Ground-water resources of North Park and Middle Park, Colorado—A reconnaissance: U.S. Geol. Survey Water-Supply Paper 1809-G, p. G1-G54, 1965.
2-65. (and HERSHEY, L. A.) Geology and ground-water resources of Prowers County, Colorado: U.S. Geol. Survey Water-Supply Paper 1772, 101 p., 1965.
- VORHIS, R. C.
1-64. Earthquake-induced water-level fluctuations from a well in Dawson County, Georgia: Seismol. Soc. America Bull., v. 54, no. 4, p. 1023-1034, 1964.
2-64. Earthquake-induced water-level fluctuation in a Dawson County well [abs.]: Georgia Acad. Sci. Bull., v. 22, no. 8, p. 17, 1964.
3-64. Ground-water data from the Prince William Sound earthquake [abs.]: Earthquake Notes, v. 35, nos. 3-4, p. 47, 1964.
1-65. Bibliography of publications relating to ground water prepared by the Geological Survey and cooperating agencies, 1946-55: U.S. Geol. Survey Water-Supply Paper 1492, 203 p., 1965. (Originally published 1957.)
2-65. Earthquake magnitudes from hydroseismic data: Ground Water, v. 2, no. 1, p. 12-20, 1965.
3-65. Notes on the stratigraphy of Pulaski County, Georgia [abs.]: Geol. Soc. America SE section, Ann. Mtg., Nashville, Tenn., April 1965, Program, p. 37, 1965.
- WAANANEN, A. O.
1-64. Water resources, with a section on Waterpower by G. C. Giles; and a section on Storage by A. O. Waananen, in Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 149-179, 1964.
- WAHLBERG, J. S.
1-65. (and DEWAR, R. S.) Comparison of distribution coefficients for strontium exchange from solutions containing one and two competing cations: U.S. Geol. Survey Bull. 1140-D, p. D1-D10, 1965.
- WAHLSTROM, E. E.
1-64. The validity of geologic projection—A case history: Econ. Geology, v. 59, no. 3, p. 465-474, 1964.
- WAHRHAFTIG, Clyde
1-64. (and GATES, G. O.) Geology: Physiographic setting, in Mineral and water resources of Alaska: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 27-32, 1964.
- WALDROP, H. A.
1-65. (and PETERSON, Fred) Preliminary geologic map of the southeast quarter of the Nipple Butte quadrangle, Kane County, Utah, and Coconino County, Arizona: U.S. Geol. Survey open-file report, 1965.
- WALKER, E. H.
1-64. Ground water in the Sandpoint region, Bonner County, Idaho: U.S. Geol. Survey Water-Supply Paper 1779-I, p. 11-129, 1964.
2-64. Mineral resources: Salt, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 165-169, 1964.

- WALKER, E. H. --Continued
 3-64. Subsurface geology of the National Reactor Testing Station, Idaho: U.S. Geol. Survey Bull. 1133-E, 22 p., 1964.
 1-65. Ground water in the upper Star Valley, Wyoming: U.S. Geol. Survey Water-Supply Paper 1809-C, p. C1-C17, 1965.
- WALL, J. R.
 1-65. (and DUTCHER, L. C.) Progress report on water studies in the Orange County coastal area, California: U.S. Geol. Survey open-file report, 37 p., 1965.
- WALLACE, R. E.
 1-64. Geology: Topography: Stratigraphy, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong. 88th, 2d sess., Senate Doc. no. 87, p. 11-20, 1964.
 2-64. Structural evolution, in Mineral and water resources of Nevada: Washington, D. C., U.S. Cong., 88th, 2d sess., Senate Doc. no. 87, p. 32-39, 1964.
 3-64. (and SILBERLING, N. J.) Westward tectonic over-riding during Mesozoic time in north-central Nevada, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C10-C13, 1964.
- WALLER, R. M.
 1-64. Ground-water conditions in the Anchorage, Alaska slide area [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 216, 1964.
 2-64. Hydrology and the effects of increased ground-water pumping in the Anchorage area, Alaska: U.S. Geol. Survey Water-Supply Paper 1779-D, p. D1-D36, 1964.
 1-65. (THOMAS, H. E., and VORHIS, R. C.) Effects of the Good Friday earthquake on water supplies: Am. Water Works Assoc. Jour., v. 57, no. 2, p. 123-131, 1965.
- WALTER, L. S.
 1-64. (and CARRON, M. K.) Vapor pressure and vapor fractionation of silicate melts of tektite composition, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963--Pt. C, Cosmochemistry and petrography: U.S. Geol. Survey open-file report, p. 140-168, 1963; also in Geochim. et Cosmochim. Acta, v. 28, no. 6, p. 937-951, 1964.
- WALTERS, K. L.
 1-65. (and KIMMEL, G. E.) Ground-water occurrence and stratigraphy of unconsolidated deposits, central Pierce County, Washington: U.S. Geol. Survey open-file report, 803 p., 1965.
- WARD, F. N.
 1-64. (and McHUGH, J. B.) Determination of mercury in vegetation with dithizone--A single extraction procedure, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D128-D130, 1964.
- WARD, P. E.
 1-64. (and TRUXES, L. S.) Water wells in Puerto Rico: Commonwealth of Puerto Rico Water Resources Bull. 3, 248 p., 1964.
- WARREN, D. H.
 1-65. (ROLLER, J. C., and JACKSON, W. H.) Seismic-refraction survey in the vicinity of the Tonto Forest Seismological Observatory, Arizona [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 155, 1965.
- WARREN, W. C.
 1-65. (BRIDGE, Josiah, and OVERSTREET, E. F.) Bauxite deposits of Virginia: U.S. Geol. Survey Bull. 1199-K, p. K1-K17, 1965.
 2-65. (and CLARK, L. D.) Bauxite deposits of the Eufaula district, Alabama: U.S. Geol. Survey Bull. 1199-E, p. E1-E31, 1965.
- WASSON, B. E.
 1-64. Source and development of public and industrial water supplies in northwestern Mississippi: U.S. Geol. Survey open-file report, 139 p., 1964.
 2-64. (GOLDEN, H. G., and GAYDOS, M. W.) Water for industrial development in Clay, Lowndes, Monroe and Oktibbeha Counties, Mississippi: U.S. Geol. Survey open-file report, 48 p., 1964.
- WATKINS, F. A., Jr.
 1-64. (and JORDAN, D. G.) Ground-water resources of west-central Indiana--Preliminary report, Parke County: Indiana Div. Water Resources Bull. 23, 125 p., 1964.
 2-64. (and JORDAN, D. G.) Ground-water resources of west-central Indiana--Preliminary report, Putnam County: Indiana Div. Water Resources Bull. 21, 83 p., 1964.
- WATKINS, J. S.
 1-64. Regional geologic implications of the gravity and magnetic fields of a part of eastern Tennessee and southern Kentucky: U.S. Geol. Survey Prof. Paper 516-A, p. A1-A17, 1964.
 2-64. Translation of "Evaluation of the depth of penetration of gravity exploration during the search for ore deposits," by A. Ya Yarosh: Internat. Geology Rev., v. 6, no. 3, p. 400-404, 1964.
 3-64. (and BAILEY, N. G.) Refraction seismic survey of parts of the Little Miami River and Scioto River Valleys, Ohio: U.S. Geol. Survey open-file report, 7 p., 1964.
 1-65. (and GODSON, R. H.) *In situ* seismic velocities observed in volcanic rocks in northern Arizona and eastern California [abs.]: Am. Geophys. Union Trans., v. 46, no. 1, p. 156-157, 1965.
 2-65. (and SPIEKER, A. M.) Seismic refraction surveys in ground-water investigations--Some experience in southwestern Ohio: Am. Geophys. Union Trans., v. 46, no. 1, p. 77-78, 1965.
- WATSON, Kenneth
 1-63. Lunar infrared emission study, in Astrogeologic studies, annual progress report, August 25, 1962-July 1, 1963--Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 29-33a, 1963.
- WAYLAND, R. G.
 1-65. The correlation of coal beds in Squaw Basin and part of Eden Ridge, T. 33 S., R. 11 W., W. M., southwestern Oregon: U.S. Geol. Survey open-file report, 34 p., 1965.
- WAYMAN, C. H.
 1-64. (and ROBERTSON, J. B.) Zero point of charge of kaolinite surfaces in acid solution [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 218, 1964.
 2-64. (ROBERTSON, J. B., and HALL, C. W.) Biodegradation of surfactants under aerobic and anaerobic conditions: Proc. of the 18th Indus. Waste Conf., 1963: Eng. Bull. Purdue Univ., Ext. Ser. 115, v. XLVIII, no. 31, p. 578-588, 1964.
 3-64. (and ROBERTSON, J. B.) Adsorption of ABS on soil minerals: Proc. of the 18th Indus. Waste Conf. 1963: Eng. Bull. Purdue Univ., Ext. Ser. 115, v. XLVIII, no. 31, p. 523-533, 1964.
 1-65. A hard look at soft detergents: Bull. Atomic Scientists, v. 21, no. 4, p. 22-26, 1965.
- WEDDOW, Helmuth, Jr.
 1-65. Possible statistical criteria for distinguishing between tectonic and collapse breccias in bedded rocks [abs.]: Geol. Soc. America Spec. Paper 82, p. 312, 1965.
 2-65. (and MARIE, J. R.) Correlation of zinc abundance with stratigraphic thickness variations in the Kingsport Formation, West New Market area, Mascot-Jefferson City mining district, Tennessee, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B17-B22, 1965.
- WEEKS, E. P.
 1-64. Field methods for determining vertical permeability and aquifer anisotropy, in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-D, p. D193-D198, 1964.
 2-64. Hydrologic conditions in the Wheatland Flats area, Platte County, Wyoming: U.S. Geol. Survey Water-Supply Paper 1783, 79 p., 1964.
 3-64. (ERICSON, D. W., and HOLT, C. L. R., Jr.) Hydrology of the little Plover River basin, Portage County, Wisconsin, and the effects of water resource development: U.S. Geol. Survey open-file report, 152 p., 1964.

- WEEKS, R. A.
1-64. (and HILPERT, L. S.) The mineral industry in Utah, in Mineral and water resources of Utah: Washington, D. C., U. S. Cong. 88th, 2d sess., Comm. Print, p. 9-15, 1964.
- WEIGLE, J. M.
1-64. Ground-water favorability map of the Salem-Plaistow area, New Hampshire: New Hampshire Water Resources Board, 1964.
- WEIR, J. E., Jr.
1-64. (CRIPPEN, J. R., and DUTCHER, L. C.) A progress report and proposed test-well drilling program for the water-resources investigation of the Antelope Valley-East Kern Water Agency area, California: U.S. Geol. Survey open-file report, 127 p., 1964.
1-65. Ground-water inventory for 1963, Edwards Air Force Base, California: U.S. Geol. Survey open-file report, 28 p., 1965.
- WEIS, P. L., Jr.
1-64. Bleached rocks in the Coeur d'Alene district, section, in Ore deposits of the Coeur d'Alene district, Shoshone County, Idaho: U.S. Geol. Survey Prof. Paper 445, p. 83-96, 1964.
2-64. (and THEOBALD, P. K.) Geology of the Mont quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-305, 1964.
- WEISSENBORN, A. E.
1-64. Introduction and summary, in Mineral and water resources of Idaho: Washington, D. C., U. S. Cong. 88th, 2d sess., Comm. Print, p. 9-11, 1964.
2-64. The mineral industry in Idaho, in Mineral and water resources of Idaho: Washington, D. C., U. S. Cong. 88th, 2d sess., Comm. Print, p. 13-22, 1964.
- WEIST, W. G., Jr.
1-65. Reconnaissance of the ground-water resources in parts of Larimer, Logan, Morgan, Sedgwick, and Weld Counties, Colorado, with a section on The chemical quality of the water, by Robert Brennan: U.S. Geol. Survey open-file report, 59 p., 1964; also in U.S. Geol. Survey Water-Supply Paper 1809-L, p. L1-L24, 1965.
- WELD, B. A.
1-64. (GRIFFIN, M. S., and BRETT, G. W.) Reports and maps of the Geological Survey released only in the open files, 1963: U.S. Geol. Survey Circ. 488, 13 p., 1964.
- WELDER, G. E.
1-64. Summary of test drilling at Firehole Canyon and Buckboard Crossing, Sweetwater County, Wyoming: U.S. Geol. Survey open-file report, 43 p., 1964.
1-65. (and MCGREEVY, L. J.) Ground-water reconnaissance of the Great Divide and Washakie basins and some adjacent areas, southwestern Wyoming: U.S. Geol. Survey open-file report, 48 p., 1965.
2-65. (and WEEKS, E. P.) Hydrologic conditions near Glendo, Platte County, Wyoming: U.S. Geol. Survey Water-Supply Paper 1791, 82 p., 1965.
- WELLS, J. D.
1-64. (SHERIDAN, D. M., and ALBEE, A. L.) Relationship of Precambrian quartzite-schist sequence along Coal Creek to Idaho Springs Formation, Front Range, Colorado: U.S. Geol. Survey Prof. Paper 454-O, p. O1-O25, 1964.
- WELLS, J. W.
1-64. Fossil corals from Eniwetok Atoll: U.S. Geol. Survey Prof. Paper 260-DD, p. DD1101-DD1111, 1964.
- WEST, S. W.
1-65. (and BALDWIN, H. L.) The water supply of El Morro National Monument: U.S. Geol. Survey Water-Supply Paper 1766, 32 p., 1965.
2-65. (CUSHMAN, R. L., STOW, J. M., and BECKLER, W. L.) Water resources, in Mineral and water resources of New Mexico: Washington, D. C., U. S. Cong. 89th, 1st sess., Comm. Print, p. 387-432, 1965.
- WETTERHALL, W. S.
1-65. Reconnaissance of springs and sinks in west-central Florida: Florida Geol. Survey Rept. Inv. 39, 42 p., 1965.
- WHITCOMB, H. A.
1-64. (CUMMINGS, T. R., and McCULLOUGH, R. A.) Ground-water resources and geology of northern and central Johnson County, Wyoming: U.S. Geol. Survey open-file report, 195 p., 1964.
2-64. (and MORRIS, D. A.) Ground-water resources and geology of northern and western Crook County, Wyoming: U.S. Geol. Survey Water-Supply Paper 1698, 92 p., 1964.
1-65. Ground-water resources and geology of Niobrara County, Wyoming, with a section on Chemical quality of the ground water by T. R. Cummings: U.S. Geol. Survey Water-Supply Paper 1788, 101 p., 1965.
- WHITE, D. E.
1-63. (and WARING, G. A.) Volcanic emanations, in Data of geochemistry, 6th ed.: U.S. Geol. Survey Prof. Paper 440-K, p. K1-K29, 1963.
1-64. (HEM, J. D., and WARING, G. A.) Chemical composition of subsurface waters, in Data of geochemistry, 6th ed.: U.S. Geol. Survey Prof. Paper 440-F, p. F1-F67, 1964. (Originally published 1963.)
2-64. (THOMPSON, G. A., and SANDBERG, C. H.) Rocks, structure, and geologic history of Steamboat Springs thermal area, Washoe County, Nevada: U.S. Geol. Survey Prof. Paper 458-B, p. B1-B63, 1964.
1-65. (and MUFFLER, L. J. P.) Metamorphism of upper Cenozoic sediments to greenschist mineral assemblages; Salton Sea geothermal area, California [abs.]: Geol. Soc. America Spec. Paper 82, p. 221-222, 1965.
- WHITE, M. G.
1-64. Uranium at Morro do Vento, Serra de Jacobina, Brazil: U.S. Geol. Survey Bull. 1185-A, p. A1-A18, 1964.
- WHITE, N. D.
1-64. (and HARDT, W. F.) Further analysis of hydrologic data for San Simon basin, Cochise and Graham Counties, Arizona, including analysis by electrical-analog model: U.S. Geol. Survey open-file report, 81 p., 1964.
2-64. (STULIK, R. S., MORSE, E. K., and others) Annual report on ground water in Arizona, Spring 1963 to Spring 1964: Arizona State Land Dept. Water Resources Rept. 19, 60 p., 1964.
3-64. (STULIK, R. S., and RAUH, C. L.) Effects of ground water withdrawal in part of central Arizona projected to 1969: Arizona State Land Dept. Water Resources Rept. 16, 23 p., 1964.
1-65. (and SMITH, C. R.) Basic hydrologic data for San Simon basin, Cochise and Graham Counties, Arizona, and Hidalgo County, New Mexico: Arizona State Land Dept. Water-Resources Rept. 21, 42 p., 1965.
- WHITE, W. S.
1-65. Tectonics of the Keweenaw basin, western Lake Superior region [abs.], in Eleventh Annual Institute on Lake Superior Geology, May 1965: St. Paul, Minn., Univ. Minnesota, p. 34, 1965.
- WHITEHEAD, H. C.
1-64. (and FETH, J. H.) Chemical composition of rain, dry fallout, and bulk precipitation at Menlo Park, California, 1957-59: Jour. Geophys. Research, v. 69, no. 16, p. 3319-3333, 1964.
- WHITMORE, F. C., Jr.
1-65. (and STEWART, R. H.) Miocene mammals and Central America seaways: Science, v. 148, no. 3667, p. 180-185, 1965.
- WHITTINGTON, C. L.
1-65. (and FERM, J. C.) Geology of the Oldtown quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-353, 1965.
- WIARD, L. A.
1-64. Cooperative surface-water investigations in Wyoming during 1963 and 1964: State Engineer of Wyoming, 37th Biennial Rept., p. 82-89, 1964.
- WIER, K. L.
1-65. Preliminary geologic map of the Black Butte iron deposit, Madison County, Montana: U.S. Geol. Survey open-file report, 1965.

- WIESNET, D. R.
 1-64. (and ASSELSTINE, E. S.) Hydrologic and hydrogeologic mapping in the United States [abs.]: Assoc. Am. Geographers Annals, v. 54, no. 3, p. 441, 1964.
 2-64. (and WARMAN, J. C.) Classification of hydrologic maps [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 224, 1964.
- WIITALA, S. W.
 1-64. Magnitude and frequency of floods in the United States—Pt. 4, St. Lawrence River Basin: U.S. Geol. Survey open-file report, 81 p., 1964.
- WILFORD, G. E.
 1-64. (and BARNES, V. E.) Brunei tektites [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 82, 1964.
- WILHELM, D. E.
 1-63. Progress in mapping the Taruntius quadrangle, in Astrogeologic studies, annual progress report, August 25, 1962–July 1, 1963—Pt. A, Lunar and planetary investigations: U.S. Geol. Survey open-file report, p. 24–29, 1963.
 2-63. A photometric technique for measurement of lunar slopes, in Astrogeologic studies, annual progress report, August 25, 1962–July 1, 1963—Pt. D, Studies for space flight program: U.S. Geol. Survey open-file report, p. 1–12, 1963.
- WILLDEN, Ronald
 1-64. Geology and mineral deposits of Humboldt County, Nevada: Nevada Bur. Mines Bull. 59, 1964[1965].
 2-64. Geology of the Christmas quadrangle, Gila and Pinal Counties, Arizona: U.S. Geol. Survey Bull. 1161-E, p. E1–E64, 1964.
- WILLIAMS, J. R.
 1-64. Drainage-area data for eastern Washington: U.S. Geol. Survey open-file report, 18 p., 1964.
- WILLIAMS, P. L.
 1-64. Geology, structure, and uranium deposits of the Moab quadrangle, Colorado and Utah: U.S. Geol. Survey Misc. Geol. Inv. Map I-360, 1964.
- WILLIAMS, R. C.
 1-64. Sedimentation in three small forested drainage basins in the Alsea River basin, Oregon: U.S. Geol. Survey Circ. 490, 16 p., 1964.
- WILLIAMS, W. P.
 1-63. (EMERICK, W. L., and DAVIS, R. E.) Geologic appraisal of the U-3cs site and vicinity, Area 3, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-53, 13 p., 1963.
 2-63. (and HOOVER, D. L.) The lithology of the U-3cn drill hole site, Area 3, Nevada Test Site: U.S. Geol. Survey Tech. Letter YUCCA-50, 12 p., 1963.
 1-64. Surface effects from underground tests at the U9ao and U9ay sites, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-80, 14 p., 1964.
 2-64. Surface effects of the Bilby event, U3cn site, September 1963, Area 3, Yucca Flat, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-68, 16 p., 1964.
- WILMOTH, B. M.
 1-64. Estimated water budget for Pocatalico River basin, West Virginia [abs.]: West Virginia Acad. Sci. Proc., v. 36, p. 117–118, 1964.
- WILSHIRE, H. G.
 1-64. Geology of the New Concord quadrangle and part of the Buchanan quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-313, 1964.
- WILSON, K. V.
 1-64. Floods of 1961 in Mississippi: Mississippi Board Water Comm. Bull. 64-4, 93 p., 1964.
- WILSON, M. T.
 1-64. (LANGFORD, R. H., and ARNOW, Ted) Water resources, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 239–268, 1964.
- WILSON, M. T. --Continued
 2-64. (and THOMAS, H. E.) Hydrology and hydrogeology of Navajo Lake, Kane County, Utah: U.S. Geol. Survey Prof. Paper 417-C, p. C1–C26, 1964.
- WINOGRAD, I. J.
 1-63. Availability of ground water for a proposed townsite in Rock Valley, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-65, 13 p., 1963.
 1-64. (and EAKIN, T. E.) Interbasin movement of ground water in south-central Nevada—The evidence [abs.]: Geol. Soc. America Ann. Mtg., Miami Beach, Florida, Nov. 1964, Program, p. 227, 1964.
- WITCOMB, H. A.
 1-65. Ground-water resources and geology of Niobrara County, Wyoming, with a section on Chemical quality of the groundwater, by T.R. Cummings: U.S. Geol. Survey Water-Supply Paper 1788, 101 p., 1965.
- WITHINGTON, C. F.
 1-64. Construction materials: Gypsum and anhydrite, in Mineral and water resources of Colorado: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 190–197, 1964.
 2-64. Mineral resources: Gypsum and anhydrite, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 101–104, 1964.
 3-64. Nonmetallic and industrial minerals and materials resources: Gypsum and anhydrite, in Mineral and water resources of Utah: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 177–185, 1964.
 1-65. Distribution of gravel in the Patuxent Formation in the Beltsville quadrangle, Prince Georges and Montgomery Counties, Maryland, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B39–B42, 1965.
 2-65. Suggestions for prospecting for evaporite deposits in southwestern Virginia, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B29–B33, 1965.
- WITKIND, I. J.
 1-64. Geology of the Abajo Mountains area, San Juan County, Utah: U.S. Geol. Survey Prof. Paper 453, 110 p., 1964.
 1-65. Fracture-controlled laccoliths, Abajo Mountains, Utah [abs.]: Geol. Soc. America Spec. Paper 82, p. 351–352, 1965.
- WOLFE, E. W.
 1-64. Geology of the Fairdealing quadrangle, Kentucky: U.S. Geol. Survey Geol. Quad. Map GQ-320, 1964.
- WOLFE, J. A.
 1-64. Miocene floras from Fingerrock Wash, southwestern Nevada: U.S. Geol. Survey Prof. Paper 454-N, p. N1–N36, 1964.
 1-65. (HOPKINS, D. M., and LEOPOLD, E. B.) Tertiary stratigraphy and paleobotany of the Cook Inlet region, Alaska: U.S. Geol. Survey open-file report, 77 p., 1965.
- WOLFF, R. G.
 1-65. Infrared absorption patterns (OH region) of several clay minerals: Am. Mineralogist, v. 50, no. 1, p. 240–244, 1965.
- WOLMAN, M. G.
 1-65. (and LEOPOLD, L. B.) River flood plains: Some observations on their formation: U.S. Geol. Survey Prof. Paper 282-C, p. 87–107, 1965. (Originally published 1957.)
- WONES, D. R.
 1-65. Biotite-K feldspar—magnetite assemblages as indicators of H₂O pressure during metamorphic and igneous processes [abs.]: Geol. Soc. America Spec. Paper 82, p. 228–229, 1965.
- WOOD, G. H., Jr.
 1-65. (TREXLER, J. P., KEHN, T. M., SOREN, Julian, and YELENOSKY, Andy) Geologic maps of the west-

WOOD, G. H., Jr. --Continued

central part of the Southern Anthracite field, Pennsylvania: U.S. Geol. Survey open-file report, 10 maps with explanation, 1965.

WOOD, L. A.

- 1-65. (and GABRYSCH, R. K.) Analog model study of ground water in the Houston district, Texas, with a section on Design, construction, and use of electric analog models, by E. P. Patten, Jr.: Texas Water Comm. Bull. 6508, 103 p., 1965.

WOOD, P. R.

- 1-64. (and DALE, R. H.) Geology and ground-water features of the Edison-Maricopa area, Kern County, California: U.S. Geol. Survey Water-Supply Paper 1656, 108 p., 1964.
- 2-64. (and JOHNSTON, H. E.) Hydrology of the New Oxford Formation in Adams and York Counties, Pennsylvania: Pennsylvania Geol. Survey Bull. W21, 4th ser., 66 p., 1964.
- 3-64. (and MOELLER, M. D.) Ground-water levels in observation wells in Oklahoma, 1961-62: Oklahoma Water Resources Board Bull., 119 p., 1964.
- 1-65. (and BURTON, L. C.) Ground water in Cleveland and Oklahoma Counties, Oklahoma, with special reference to the Garber Sandstone and Wellington Formation: U.S. Geol. Survey open-file report, 132 p., 1965.
- 2-65. (and STACY, B. L.) Geology and ground-water resources of Woodward County, Oklahoma: Oklahoma Water Resources Board Bull. 21, [115 p.], 1965.

WOOLLARD, G. P., Chairman

- 1-64. (and JOESTING, H. R., coordinator) Bouguer gravity anomaly map of the United States (exclusive of Alaska and Hawaii): U.S. Geol. Survey Spec. Map, 2 sheets, 1964.

WORTS, G. F., Jr.

- 1-64. Hydrogeologic reconnaissance of Poro Point and vicinity, Luzon Island, Philippines: U.S. Geol. Survey Water-Supply Paper 1608-E, p. E1-E14, 1964.

WRIGHT, R. R.

- 1-64. (and COLLINGS, M. R.) Application of fluorescent tracing techniques to hydrologic studies: Am. Water Works Assoc. Jour., v. 56, no. 6, p. 748-754, 1964.

WRIGHT, T. L.

- 1-64. The alkali feldspars of the Tatoosh pluton in Mount Rainier National Park: Am. Mineralogist, v. 49, nos. 5-6, p. 715-735, 1964.
- 2-64. X-ray determination of composition and structural state of alkali feldspar [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 127, 1964.
- 1-65. (and FLEISCHER, Michael) Geochemistry of the platinum metals: U.S. Geol. Survey Bull. 1214-A, p. A1-A24, 1965.

YATES, R. G.

- 1-64. Geologic map and sections of the Deep Creek area, Stevens and Pend Oreille Counties, Washington: U.S. Geol. Survey Misc. Geol. Inv. Map I-412, 1964[1965].

YEVDJEVICH, V. M.

- 1-64. Bibliography and discussion of flood-routing methods and unsteady flow in channels: U.S. Geol. Survey Water-Supply Paper 1690, 235 p., 1964.

YOCHELSON, E. L.

- 1-64. The Early Ordovician gastropod *Ceratopea* from East Greenland: Meddel. om Grønland, v. 164, no. 7, p. 1-11, 1964.
- 1-65. Paleontological Society: Geotimes, v. 9, no. 7, p. 10-12, 1965.
- 2-65. Review of "Frameworks for dating fossil man," by K. P. Oakley: Jour. Paleontology, v. 39, no. 3, p. 515-516, 1965.

YOCHELSON, E. L. --Continued

- 3-65. Review of "The tentaculites of Bohemia: their morphology, taxonomy, phylogeny, and biostratigraphy," by Bedrich Boucek: Jour. Paleontology, v. 39, no. 3, p. 509-510, 1965.

- 4-65. (McALLISTER, J. F., and RESO, Anthony) Stratigraphic distribution of the Late Cambrian mollusk *Matthevia* Walcott, 1885, in Geological Survey Research 1965: U.S. Geol. Survey Prof. Paper 525-B, p. B73-B78, 1965.

YOTSUKURA, Nobuhiro

- 1-64. (and FIERING, M. B.) Numerical solution to a dispersion equation: Am. Soc. Civil Engineers Proc., v. 90, Paper 4046, Jour. Hydraulics Div., no. HY5, pt. 1, p. 83-104, 1964.

YOUNG, E. J.

- 1-65. Effect of isomorphic substitutions on the Omega index of refraction of apatite [abs.]: Geol. Soc. America Spec. Paper 82, p. 231, 1965.

YOUNG, L. L.

- 1-64. Summary of developed and potential waterpower of the United States and other countries of the world, 1955-62: U.S. Geol. Survey Circ. 483, 38 p., 1964.
- 2-64. Water resources: Waterpower, in Mineral and water resources of Idaho: Washington, D. C., U.S. Cong. 88th, 2d sess., Comm. Print, p. 317-328, 1964.

YOUNG, R. A.

- 1-63. Recommended sites for two water-supply wells in the Jackass Flats well field, Nye County, Nevada: U.S. Geol. Survey Tech. Letter NTS-48, 6 p., 1963.
- 2-63. (and WINOGRAD, I. J.) Availability of ground water for a proposed Public Health Service Experimental Farm in the Mid Valley area, Nevada Test Site: U.S. Geol. Survey Tech. Letter NTS-70, 12 p., 1963.

ZABLOCKI, C. J.

- 1-64. Electrical properties of serpentinite from Mayagüez, Puerto Rico, in A study of serpentinite, the AMSOC core hole near Mayagüez, Puerto Rico: Natl. Acad. Sci. - Natl. Research Council Pub. 1188, p. 107-117, 1964.

ZARTMAN, R. E.

- 1-64. The isotopic composition of lead in microclines from the Llano uplift, Texas [abs.]: Am. Geophys. Union Trans., v. 45, no. 1, p. 109, 1964.

ZELLER, H. D.

- 1-64. (and STEPHENS, E. V.) Geologic map of the Continental Peak quadrangle, Fremont and Sweetwater Counties, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-292, 1964.
- 2-64. (and STEPHENS, E. V.) Geologic map of the Dickie Springs quadrangle, Fremont and Sweetwater Counties, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-293, 1964.
- 3-64. (and STEPHENS, E. V.) Geologic map of the Parting of the Ways quadrangle, Sublette and Sweetwater Counties, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-296, 1964.
- 4-64. (and STEPHENS, E. V.) Geologic map of the Tule Butte quadrangle, Sweetwater County, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-297, 1964.
- 5-64. (and STEPHENS, E. V.) Geology of the NE 1/4 of the Essex Mountain quadrangle, Sweetwater County, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-285, 1964.
- 6-64. (and STEPHENS, E. V.) Geology of the NE 1/4 of Freighter Gap quadrangle, Sweetwater County, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-288, 1964.
- 7-64. (and STEPHENS, E. V.) Geology of the NW 1/4 of Freighter Gap quadrangle, Sweetwater County, Wyoming:

ZELLER, H. D. --Continued

U.S. Geol. Survey Mineral Inv. Field Studies Map MF-289, 1964.

8-64. (and STEPHENS, E. V.) Geologic map of the Hay Meadow Reservoir quadrangle, Sublette, Fremont, and Sweetwater Counties, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-295, 1964.

9-64. (and STEPHENS, E. V.) Geology of the Pinnacles NW quadrangle, Sweetwater County, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-286, 1964.

10-64. (and STEPHENS, E. V.) Geologic map of the Pacific Springs quadrangle, Fremont and Sweetwater Counties, Wyoming: U.S. Geol. Survey Mineral Inv. Field Studies Map MF-294, 1964.

ZEN, E-an

1-64. (and ALBEE, A. L.) Coexistent moscovite and paragonite in pelitic schists: *Am. Mineralogist*, v. 49, nos. 7-8, p. 904-925, 1964.

1-65. Solubility measurements in the system $\text{CaSO}_4\text{-NaCl-H}_2\text{O}$ at 35°, 50°, and 70°C and one atmosphere pressure: *Jour. Petrology*, v. 6, no. 1, p. 124-164, 1965.

2-65. (and HANSHAW, B. B.) Osmotic equilibrium and mechanics of overthrust faulting [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 232-233, 1965.

ZIETZ, Isidore

1-65. (and GEDDES, Wilburt) Crustal study of a transcontinental stripe east of the Rocky Mountains [abs.]: *Am. Geophys. Union Trans.*, v. 46, no. 1, p. 160, 1965.

2-65. (GEDDES, Wilburt, and LIDIAK, E. G.) Geophysical data from Denver, Colorado, to Washington, D. C., and its relationship to basement lithology [abs.]: *Geol. Soc. America Spec. Paper* 82, p. 233-234, 1965.

ZIMMERMAN, E. A.

1-64. Geology and water resources of the Bluewater Springs area, Carbon County, Montana: U.S. Geol. Survey Water-Supply Paper 1779-J, p. J1-J24, 1964.

ZONES, C. P.

1-63. Groundwater in the alluvium of Kings River Valley, Humboldt County, Nevada: U.S. Geol. Survey Water-Supply Paper 1619-L, p. L1-L38, 1963; also in Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 16, 38 p., 1963.

ZUBOVIC, Peter

1-64. (STADNICHENKO, Taisia, and SHEFFEY, N. B.) Distribution of minor elements in coal beds of the eastern interior region: U.S. Geol. Survey Bull. 1117-B, p. B1-B41, 1964.

PUBLICATIONS INDEX

- Absolute age, dates
 Basalt
 Tertiary, variations in K, Ar, and calculated ages: Dalrymple, G. B., 1-65.
- Absolute age, methods
 Potassium-argon
 Muscovite, interlaboratory standard: Lanphere, M. A., 1-65.
- Africa
 Hydrogeology
 General: Dennis, P. E., 1-64.
- Alabama
 Areal geology
 Elkmont quadrangle: McMaster, W. M., 1-65.
 Escambia County: Cagle, J. W., Jr., 1-64.
 Russellville area: Peace, R. R., Jr., 1-64.
- Economic geology
 Bauxite, Eufaula district: Warren, W. C., 2-65.
 Margerum district: Bergquist, H. R., 1-65.
- Geophysical surveys
 North-central, reservoir temperatures: Pierce, L. B., 1-64.
- Hydrogeology
 Escambia County, ground water: Cable, J. W., Jr., 1-64.
 General: Swindel, G. W., 1-64.
 Ground water levels, 1959 and 1960: O'Rear, D. M., 1-64.
 Mississippi embayment, low-flow characteristics of streams: Speer, P. R., 1-64.
 Russell County, ground water: Scott, J. C., 1-64.
 Russellville area ground water: Peace, R. R., Jr., 1-64.
 Streams, temperature: Avrett, J. R., 1-64.
 Tennessee Valley, limestone terrane: Powell, W. J., 1-64.
- Maps
 Elkmont quadrangle, geologic: McMaster, W. M., 1-65.
 Salem quadrangle, geologic: McMaster, W. M., 2-65.
- Paleontology
 Invertebrata, pre-Selma: Sohl, N. F., 3-64.
 Microfauna, Coker Formation: Applin, E. R., 1-64.
 Palynomorphs, Cretaceous: Leopold, E. B., 2-64.
- Petrology
 Western, core holes, pre-Selma strata: Bergenback, R. E., 1-64.
- Sedimentary petrology
 Silurian iron ores: Sheldon, R. P., 1-65.
- Stratigraphy
 Cretaceous, cores: Monroe, W. H., 1-64, 4-64.
- Alabama--Continued
 Stratigraphy--Continued
 Cretaceous, pre-Selma: Conant, L. C., 1-64.
- Structural geology
 Chattahoochee River, geologic profile: Toulmin, L. D., 1-64.
- Alaska
 Absolute age
 Bokan Mountain area, plutonic rocks: Lanphere, M. A., 1-64.
 Tenakee area, Paleozoic: Lanphere, M. A., 2-65.
- Areal geology
 Admiralty Island, reconnaissance: Lathram, E. H., 2-65.
 Anchorage area: Cederstrom, D. J., 2-64.
 Central Kobuk River Valley: Fernald, A. T., 1-64.
 Cub Run quadrangle: Sandberg, C. A., 2-65.
 Delta River to Tok Junction, reconnaissance: Holmes, G. W., 1-65.
 Killik-Itkillik region: Patton, W. W., Jr., 2-64.
 Kotzebue Sound area: McCulloch, D. S., 1-65.
 Simeonof Island: Grantz, Arthur, 1-62.
- Structural and tectonic history:
 Gates, G. O., 1-64.
- Sumdum copper-zinc prospect:
 MacKevett, E. M., Jr., 2-64.
- West Creek damsite, geologic reconnaissance: Callahan, J. E., 1-65.
- Earthquakes
 Gravity changes: Barnes, D. F., 1-65.
 Magnetic events, 1964: Moore, G. W., 2-64.
 March 27, 1964: Bonilla, M. G., 2-65; Coulter, H. W., 1-65; Dobrovolsky, Ernest, 1-65; Landen, David, 1-64; United States Geological Survey, 2-65.
- Effects: Kachadoorian, Reuben, 1-65.
- Effects on coastal areas:
 Plafker, George, 2-65.
- Effects on glaciers: Post, A. S., 1-64.
- Engineering geology: Lemke, R. W., 1-65.
- Geologic effects: Plafker, George, 3-65.
- Tectonic deformation: Plafker, George, 1-65.
- Tectonics: Grantz, Arthur, 3-65.
- 1964, engineering-geologic effects: Dobrovolsky, Ernest, 2-65.
- Economic geology
 Bryophytes associated with ore deposits: Shacklette, H. T., 2-65.
 Coal: Barnes, D. F., 1-64.
- Alaska--Continued
 Economic geology--Continued
 Coal, Lisburne Peninsula, Mississippian: Tailleux, I. L., 1-65.
 Copper, Kathleen-Margaret deposits: MacKevett, E. M., Jr., 1-64.
 Fluorspar, Lost River area: Sainsbury, C. L., 1-65.
- Industrial minerals and construction material occurrences:
 Cobb, E. H., 1-64.
- Iron: Cobb, E. H., 2-64.
- Metallic mineral resources: Berg, H. C., 1-64.
- Mineral fuel resources: Grantz, Arthur, 1-64; Patton, W. W., Jr., 1-64.
- Mineral resources: Eberlein, G. D., 1-64.
- Nonmetallic mineral resources:
 Plafker, George, 1-64.
- Engineering geology
 Petroleum reserve no. 4: Robinson, F. M., 1-64.
- General
 Leafy liverwort hydrosere, Yakobi Island: Shacklette, H. T., 1-65.
- United States Geological Survey, location of field projects: U.S. Geological Survey, Geologic Division, 2-64.
- Geomorphology
 Johnson River area, rock glacier: Foster, H. L., 1-65, 2-65.
- Physiographic setting: Wahrhaftig, Clyde, 1-64.
- Geophysical surveys
 Copper River Basin, magnetic and gravity: Andreasen, G. E., 3-64.
- Gravity: Barnes, D. F., 2-65, 3-65.
- Continental shelf: Barnes, D. F., 5-65.
- Marine and shoreline, south and west of Kodiak Island: Barnes, D. F., 4-65.
- Hydrogeology
 Anchorage area, effects of increased ground-water pumping: Waller, R. M., 2-64.
- Ground water resources:
 Cederstrom, D. J., 2-64.
- Anchorage slide area: Waller, R. M., 1-64.
- Bethel area, wells drilled through permafrost: Feulner, A. J., 2-64.
- Compilation of records, surface waters, 1950-60: Hendricks, E. L., 9-64.
- Galleries, development of shallow ground-water supplies:
 Feulner, A. J., 1-64.
- Resources: Waananen, A. O., 1-64.
- South of Yukon River, magnitude and frequency of floods: Berwick, V. K., 1-64.

- Alaska--Continued
Hydrogeology--Continued
Water resources: Eberlein, G. D., 1-64.
Water supply problems: Marcher, M. V., 1-65.
- Maps
Arctic quadrangle, geologic: Brosgé, W. P., 1-65.
Charley River quadrangle, geologic: Brabb, E. E., 1-64.
Eagle D-1 quadrangle, geologic: Brabb, E. E., 1-65.
Glaciations, extent of: Coulter, H. W., 2-65.
Iliamna quadrangle, geochemical, stream sediments: Detterman, R. L., 1-65.
Industrial minerals and construction material occurrences: Cobb, E. H., 1-64.
Lode gold and silver occurrences: Cobb, E. H., 3-64.
McCarthy C-6 quadrangle, geologic: MacKevett, E. M., Jr., 1-65.
Mineral resources, iron, occurrences: Cobb, E. H., 2-64.
Mount Hayes D-3 quadrangle, geologic: Holmes, G. W., 7-65.
Mount Hayes D-4 quadrangle, geologic: Péwé, T. L., 1-64.
Nelchina area, geologic: Grantz, Arthur, 1-65.
Northern, geologic: Lathram, E. H., 1-65.
Ogotoruk Creek area, geologic: Campbell, R. H., 1-65.
Surficial geology: Karlstrom, T. N. V., 1-64.
- Mineralogy
Laumontite, origin and occurrence: Hoare, J. M., 1-64.
- Paleontology
Ammonoidea, Bajocian: Imlay, R. W., 2-64.
Foraminifera, Bootlegger Cove clay: Smith, P. B., 2-64.
Ostracoda, northern, Triassic: Sohn, I. G., 1-64.
- Petrography
Pybus-Gambier area: Loney, R. A., 1-64.
- Stratigraphy
Cook Inlet region: Wolfe, J. A., 1-65.
Cretaceous, McCarthy A-4 quadrangle: Jones, D. L., 1-64.
Tiklukpuk Formation: Jones, D. L., 2-64.
Mesozoic, Chitina Valley: Grantz, Arthur, 2-65.
Ordovician, Silurian, Devonian, biostratigraphy: Churkin, Michael, Jr., 1-65.
Pybus-Gambier area: Loney, R. A., 1-64.
Quaternary, Tanana River Valley: Fernald, A. T., 1-65.
- Structural geology
Aleutian Arc, tensional origin: Moore, G. W., 1-65.
- Alaska--Continued
Structural geology--Continued
Pybus-Gambier area: Loney, R. A., 1-65.
- Alluvial fans
California and Nevada
Death Valley region: Denny, C. S., 1-65.
- Alunite
Colorado
San Juan Mountains: Ratté, J. C., 1-64.
- Utah
General: Parker, R. L., 3-64.
- Ammonoidea
Evolution: Imlay, R. W., 1-65.
- Analytical techniques
Spectrochemical analysis
Suggested automation: Helz, A. W., 1-65.
- Anhydrite
Colorado
Resources: Withington, C. F., 1-64.
- Idaho
Resources: Withington, C. F., 2-64.
- Utah
General: Withington, C. F., 3-64.
- Antarctica
Absolute age
Eight Coast area, quartz diorite: Drake, A. A., Jr., 1-64.
- Areal geology
Neptune Range: Schmidt, D. L., 1-65.
Patuxent Mountains: Schmidt, D. L., 1-64.
Review: Ford, A. B., 2-64.
- General
Daylight star observations: Lee, D. R., 1-64.
- Glacial geology
Densification of snow: Behrendt, J. C., 1-65.
- Maps
Tectonic: Hamilton, Warren, 1-64.
- Petrology
Thiel Mountains, cordierite-hypersthene-quartz monzonite: Ford, A. B., 1-64.
Victoria Land, diabase: Hamilton, Warren, 2-64, 1-65.
- Anthozoa
Classification
Faviphyllum rugosum Hall: Sando, W. J., 1-65.
Paleozoic species, western United States: Sando, W. J., 2-65.
- Paleozoic
Handbook: Oliver, W. A., Jr., 1-65.
- Antimony
Idaho
Big Creek-Yellow Pine area: Leonard, B. F., 1-65.
General: LaHeist, B. A., 2-64.
- South Dakota
General: Dasch, M. D., 1-64.
- Antimony--Continued
Utah
General: Dasch, M. D., 2-64.
- Appalachians
Heat flow
Central and southern: Diment, W. H., 1-65.
- Hydrogeology
Pennsylvania to Alabama, water resources: Schneider, W. J., 1-65.
- Paleontology
Palynomorphs, Cretaceous: Tschudy, R. H., 1-65.
- Arctic Ocean
Genesis: King, E. R., 1-65.
- Geophysical surveys
Heat flux: Lachenbruch, A. H., 1-64.
- Argentina
Hydrogeology
Bahia Blanca area: Schoff, S. L., 1-64.
- Meteorites
Campo del Cielo field: Milton, D. J., 1-63.
- Arizona
Areal geology
Apache County, central: Akers, J. P., 1-64.
Cenozoic, 1960 resume: Heindl, L. A., 1-64.
Christmas quadrangle: Wilden, Ronald, 2-64.
Dragoon quadrangle: Cooper, J. R., 4-64.
Klondyke quadrangle: Simons, F. S., 1-64.
Prescott-Jerome area: Anderson, C. A., 1-65.
San Manuel area: Creasey, S. C., 1-65.
- Economic geology
Dragoon quadrangle, ore deposits: Cooper, J. R., 4-64.
San Manuel area, ore deposits: Creasey, S. C., 1-65.
- Engineering geology
Low velocity impact crater, Hell-hole Bend: Elston, D. P., 2-63.
- Geomorphology
Southern: Kam, William, 1-65.
- Geophysical surveys
Globe-Miami Copper district, aeromagnetic: Jespersen, Anna, 1-64.
Gravity, United States Geological Survey program: Case, J. E., 3-65.
Tonto Forest Seismological Observatory, seismic: Warren, D. H., 1-65.
Tucson area, gravity: Plouff, Donald, 1-64.
Volcanic rocks, velocity determined in place: Watkins, J. S., 1-65.
- Hydrogeology
Apache, Navajo, and Coconino Counties, ground water: McGavock, E. H., 1-64.

Arizona--Continued

Hydrogeology--Continued

- Apache County, central, ground-water: Akers, J. P., 1-64.
 Central, ground-water withdrawal projection: White, N. D., 3-64.
 Chino Valley: Schwalen, H. C., 1-64.
 Cottonwood Wash, water use, phreatophytes: Bowie, J. E., 1-65.
 Current publications: Smith, C. R., 1-64.
 Duncan and Safford basins: Davidson, E. S., 1-64.
 Fort Huachuca Military Reservation, water resources: Brown, S. G., 1-65.
 Gila County: Smith, C. R., 2-64.
 Ground water, annual report, 1963-64: White, N. D., 2-64.
 Use: Morse, E. K., 1-64.
 Ground water in various areas: Stulik, R. S., 3-64.
 Harquahala Plains area, ground-water: Stulik, R. S., 1-64.
 Maricopa County, ground-water: Stulik, R. S., 2-64.
 Sycamore Creek watershed: Thomson, B. W., 1-64.
 Navajo and Hopi Indian Reservations: Cooley, M. E., 1-64, 2-64.
 Pinal County, salinity of ground-water: Kister, L. R., 2-65.
 Western, basic data: Hardt, W. F., 1-64.
 San Simon basin, data analysis: White, N. D., 1-64, 1-65.
 San Xavier Indian Reservation: Heindl, L. A., 2-65.
 Santa Cruz basin, upper, and Avra-Marana area: Schwalen, H. C., 2-64.
 South Gila Valley, Yuma Mesa, and Yuma Valley: Frank, F. J., 1-64.
 Surface water, 1963-64: Hodges, E. B., 1-64.
 Willcox and Douglas basins: Brown, S. G., 1-64.

Maps

- Big Maria Mountains NE quadrangle, geologic: Hamilton, Warren, 3-64.
 Casa Grande area, aeromagnetic: Mitchell, C. M., 2-65.
 Central, gravity: Peterson, D. L., 1-65.
 Mule Mountains, geologic: Hayes, P. T., 2-64.
 Nipple Butte quadrangle, geologic: Waldrop, H. A., 1-65.
 Tombstone and vicinity, aeromagnetic: Andreasen, G. E., 1-65.
 Willcox basin, ground water chemistry: Kister, L. R., 1-65.
 Sedimentary petrology
 Laguna Salada: Cooley, M. E., 3-64.
 Stratigraphy
 Cambrian, Papago Indian Reservation: Heindl, L. A., 3-64.

Arizona--Continued

Stratigraphy--Continued

- Chinle and Moenkopi Formations, Navajo and Hopi Indian Reservations: Repenning, C. A., 2-65.
 Mesozoic, Lower, southeastern: Hayes, P. T., 3-65.
 Vekol Mountains: Heindl, L. A., 1-65.
 Mule Mountains, southern part: Hayes, P. T., 2-65.
 Northern, corals in Redwall Limestone: Sando, W. J., 1-64.
 Precambrian and Paleozoic, Papago Indian Reservation: McClymonds, N. E., 1-64.

Arkansas

Areal geology

- North-central, uranium-bearing black shale, Devonian: Swanson, V. E., 2-64.
 Hydrogeology
 Arkansas River valley: Tanaka, H. H., 1-64.
 Water levels: May, J. R., 1-65.
 Water quality: May, J. R., 1-64.
 Well logs: May, J. R., 2-65.
 Bradley, Calhoun, and Ouachita Counties, ground water: Albin, D. R., 1-64.
 Coastal Plain, ground water temperatures: Plebuch, R. O., 1-64.
 Eastern, vicinity of U.S. Highway 70, ground water resources: Halberg, H. N., 1-64.
 Grand Prairie region, artificial recharge: Sniegocki, R. T., 1-65.
 Hot Springs, floods, 1963: Gilstrap, R. C., 1-64.
 South-central, Carrico Sand, potential aquifer: Hosman, R. L., 1-64.
 Streams, water-supply characteristics: Hines, M. S., 1-65.

Mineralogy

- Quartz, rectorite, cookeite, near North Little Rock: Miser, H. D., 1-64.
 Paleontology
 Cephalopoda, Carboniferous: Gordon, Mackenzie, Jr., 2-64.
 Sedimentary petrology
 Morrow Series, diagenesis: Henbest, L. G., 1-65.

Stratigraphy

- Witts Springs Formation, Snowball quadrangle: Glick, E. E., 1-64.

Arsenic

Nevada

- General: La Heist, B. A., 1-64.

Arthropoda

Decapod crustaceans

- Marshall Islands: Roberts, H. B., 1-64.

Associations

Paleontological Society

- History: Yochelson, E. L., 1-65.

Astrogeology

- United States Geological Survey, annual progress report, August 1962-July 1963: United States Geological Survey, 15-64.
 United States Geological Survey, annual progress report, July 1963-July 1964: United States Geological Survey, 1-65.

Atlantic Coastal Plain

- Hydrogeology: LeGrand, H. E., 1-64.

Atlantic Ocean

Petrology

- Mid-Atlantic Ridge, basalt: Engel, A. E. J., 1-64.

Australia

Structural geology

- Henbury meteorite craters: Milton, D. J., 1-65.

Austria

Petrology

- Köfels, fused rocks: Milton, D. J., 1-64.

Automatic data processing

- Computer programs: U.S. Geological Survey, Branch of Computation, 1-64.

Spectrochemical analysis

- Potential use: Helz, A. W., 1-65.

Barite

- Colorado: Brobst, D. A., 2-64.
 Idaho: Brobst, D. A., 1-64.
 United States: Brobst, D. A., 1-65.
 Utah: Brobst, D. A., 3-64.

Basalts

Composition

- Mid-Atlantic Ridge: Engel, A. E. J., 1-64.

Bauxite

Alabama

- Eufaula district: Warren, W. C., 2-65.
 Margerum district, Gordo Formation, Late Cretaceous: Bergquist, H. R., 1-65.

Georgia

- Springvale district: Clark, L. D., 1-65.

Mississippi

- Tippah-Benton district: Tourtelot, H. A., 1-64.

United States

- Southeastern: Overstreet, E. F., 1-64.

Virginia

- Staunton area, Valley and Ridge province: Warren, W. C., 1-65.

Bering Strait

- Quaternary correlations: Hopkins, D. M., 1-65.

Beryllium

Analysis

- Gamma activation device: Vaughn, W. W., 1-65.

Colorado

- General: Griffiths, W. R., 4-64.

Geochemistry

- Tin deposits rich in fluorite: Sainsbury, C. L., 1-64.

Beryllium--Continued

- Idaho
General: Griffiths, W. R., 5-64.
- Nevada
General: Griffiths, W. R., 1-64.
- New Mexico
General: Griffiths, W. R., 6-65.
- Utah
General: Griffiths, W. R., 2-64.
- Bibliography
- Evaporation and transpiration:
Robinson, T. W., 2-65.
- Hydrogeology
Texas, ground-water publications
and open-file reports: Baker,
R. C., 1-64.
- Lake deposits, ancient
Western states: Feth, J. H., 2-64.
- North America
Geology, 1950-59: King, R. R., 1-65.
1960: U.S. Geological Survey,
1-64.
- Soils
Engineering properties: Johnson,
A. I., 1-63.
- United States Technical Assistance
Program, U.S. Geological Sur-
vey: Heath, J. A., 1-65.
- Biography
Stone, Ralph W.: Davies, W. E., 1-65.
Yagoda, Herman: Fleischer,
Michael, 1-65.
- Bismuth
Nevada
General: Mount, Priscilla, 1-64.
- Borates
Nevada
General: Smith, W. C., 1-64.
- Brachiopoda
General
Handbook: Dutro, J. T., Jr., 1-65.
Stenoscismatacea: Grant, R. E.,
1-65.
- Brazil
Areal geology
Monlevade and Rio Piracicaba
quadrangles: Reeves, R. G.,
1-65.
- Economic geology
Iron, Minas Gerais: Dorr, J. V. N.,
2d, 1-64, 2-65.
Mineral resources, Monlevade and
Rio Piracicaba quadrangles:
Reeves, R. G., 1-65.
Uranium, central Tucano Basin:
Haynes, D. C., 2-64.
Morro do Vento area, Serra de
Jacobina: White, M. G., 1-64.
- Hydrology
Northeast: Cederstrom, D. J., 1-64.
- Mineralogy
Muscovite, chromian, Serra de Ja-
cobina: Leo, G. W., 1-65.
- Breccia
Tectonic and collapse
Distinguishing criteria: Wedow, Hel-
muth, Jr., 1-65.
- British Columbia
Stratigraphy
Pleistocene: Armstrong, J. R., 1-65,
2-65.

Calderas

- Gravity studies
Colorado: Karig, D. E., 1-65.
- Structural geology
Nevada, Timber Mountain cal-
dera, Tertiary: Christiansen,
R. L., 2-65.
- California
Absolute age
Bishop Tuff: Dalrymple, G. B.,
2-65.
Potassium-argon, Pleistocene
interglacial basalt flows:
Dalrymple, G. B., 1-64.
Sierra Nevada-Inyo Mountains:
Kistler, R. W., 1-65.
- Areal geology
Ash Meadows quadrangle: Denny,
C. S., 2-65.
Berkeley Hills, Caldecott Tunnel:
Radbruch, D. H., 1-64.
Bullfrog quadrangle: Cornwall,
H. R., 3-64.
Devils Postpile National Monu-
ment: Huber, N. K., 1-65.
Furnace Creek Wash area, re-
connaissance: Pistrang, M. A.,
1-64.
Independence quadrangle: Ross,
D. C., 1-65.
Jolon and Williams Hill quad-
rangles: Durham, D. L., 1-65.
Lava Mountains: Smith, G. I.,
1-64.
Mojave Desert, southeastern, re-
connaissance: Bassett, A. M.,
1-64.
San Francisco Bay area, gravity:
Clement, W. G., 1-65.
- Economic geology
Bullfrog quadrangle, ore de-
posits: Cornwall, H. R., 3-64.
Nickeliferous laterites: Hotz,
P. E., 1-64.
Phosphate rock, occurrence:
Gower, H. D., 1-64.
- Engineering geology
Land subsidence, well casing dis-
tortion: Poland, J. F., 1-65.
- General
Field investigations 1965: United
States Geological Survey, Geo-
logic Division, 1-65.
- Geochemistry
Birch Creek, travertine deposi-
tion: Barnes, Ivan, 1-65.
Sr-87/Sr-86 ratios, Sierra Nevada
plutonic province: Hurley,
P. M., 1-65.
- Geomorphology
Alluvial fans: Denny, C. S., 1-65.
Fresno County: Bull, W. B., 2-64.
- Geophysical surveys
Owens Valley, gravity: Pakiser,
L. C., 2-64.
United States Geological Survey
gravity program: Oliver,
H. W., 1-65.
Volcanic rocks, velocity deter-
mined in place: Watkins, J. S.,
1-65.

California--Continued

- Glacial geology
Sequoia National Park: Matthes,
F. E., 1-65.
- Hydrogeology
Antelope Valley area, well data:
Moyle, W. R., Jr., 1-64.
Antelope Valley-East Kern Water
Agency area: Weir, J. E., Jr.,
1-64.
Bloomington-Colton area, water
studies, 1964: Dutcher, L. C.,
1-65.
Chino-Corona area, water studies,
1964: Dutcher, L. C., 3-65.
Coastal basins, flood-frequency
studies: Cruff, R. W., 1-64.
Edwards Air Force Base, 1963:
Weir, J. E., Jr., 1-65.
Fresno County, western, channel
trenching: Bull, W. B., 3-64.
Furnace Creek Wash area, recon-
naissance: Pistrang, M. A.,
1-64.
Kern County, Edison-Maricopa
area: Wood, P. R., 1-64.
Lakes, thermal stratification,
elimination: Koberg, G. E.,
1-64, 2-65.
Lower Colorado River-Salton Sea
area: Hely, A. G., 2-64.
Marine Corps Base, Twentynine
Palms, ground water conditions,
1964: Giessner, F. W., 1-65.
Northern coastal ranges: Rantz,
S. E., 1-64.
Orange County coastal area: Wall,
J. R., 1-65.
Pala and Rincon Indian Reserva-
tions: French, J. J., 2-65.
Permanente Creek: Harris, E. E.,
1-64.
Point Arguello, U.S. Naval Mis-
sile Facility, ground-water con-
ditions: Miller, G. A., 1-65.
Redwood areas, floods: Rantz, S. E.,
1-65.
Riverside County, Mission Creek
Indian Reservation: Giessner,
F. W., 1-64.
San Joaquin Valley, ground-water:
Dale, R. H., 1-64.
San Timoteo-Smiley Heights area,
water studies, 1964: Dutcher,
L. C., 2-65.
Santa Ana Valley, upper, water
studies: French, J. J., 1-65.
Santa Barbara County, water levels,
1963: Muir, K. S., 1-64.
Santa Maria Valley area, ground
water, utilization: Miller, G. A.,
1-64.
Santa Ynez River basin, ground-
water quality: Evenson, R. E.,
1-64.
Sierra Nevada, northern, snow,
chemistry: Feth, J. H., 3-64.
Southern, water loss and recover-
able water in mountain basins:
Crippen, J. R., 2-65.

California--Continued

Maps

- Basement rocks, distribution and configuration: Smith, M. B., 1-64.
- Big Maria Mountains NE quadrangle, geologic: Hamilton, Warren, 3-64.
- Cajon quadrangle, geologic: Dibblee, T. W., Jr., 1-65.
- Hesperia quadrangle, geologic: Dibblee, T. W., Jr., 2-65.
- Long Beach-Santa Ana, aeromagnetic: Andreasen, G. E., 4-64.
- Los Angeles, western and vicinity, aeromagnetic: Andreasen, G. E., 5-64.
- Lucerne Valley quadrangle, geologic: Dibblee, T. W., Jr., 1-64.
- Merced Peak quadrangle, geologic: Peck, D. L., 1-64.
- Nebo and Yermo quadrangles, geologic: McCulloh, T. H., 1-65.
- Ord Mountains quadrangle, geologic: Dibblee, T. W., Jr., 2-64.
- Prospect Peak quadrangle, geologic: MacDonald, G. A., 1-64.
- Radman Mountains quadrangle, geologic: Dibblee, T. W., Jr., 3-64.
- San Francisco Bay area, gravity: Clement, W. G., 1-65.
- San Francisco region, radioactivity: Books, K. G., 1-65.
- San Francisco South quadrangle, geologic: Bonilla, M. G., 1-65.
- San Geronio Mountain quadrangle, geologic: Dibblee, T. W., Jr., 4-64.
- Stonyford quadrangle, geologic: Brown, R. D., Jr., 1-64.
- White Mountain Peak quadrangle, geologic: Sheridan, M. F., 1-64, 2-64.

Mineralogy

- Jadeite and asbestos: Coleman, R. G., 2-65.
- Macallisterite: Schaller, Waldemar, 1-65.
- Quartz crystals, inclusions: Masiemer, G. E., 1-65.

Paleontology

- Cephalopoda, Carboniferous: Gordon, MacKenzie, Jr., 1-64.
- Diatoms, Mesozoic and Cenozoic: Lohman, K. E., 1-64.
- Invertebrata, Pleistocene, Dumeret race: Addicott, W. O., 2-64.
- Mollusca, Cambrian: Yochelson, E. L., 4-65.

Petrology

- Igneous rocks, tantalum distribution: Gottfried, David, 1-65.
- Mojave Desert and vicinity, zeolite deposits: Sheppard, R. A., 2-64, 1-65.
- Salton Sea area, metamorphism, Cenozoic sediments: White, D. E., 1-65.

Precipitation

- Menlo Park, 1957-59, composition: Whitehead, H. C., 1-64.

California--Continued

Sedimentary petrology

- Cache Creek drainage basin: Lustig, L. K., 1-65.
- Lake Pillsbury: Porterfield, George, 1-64.
- San Gabriel Mountains: Lustig, L. K., 1-64.
- Searles Lake: Smith, G. I., 2-64.
- Stratigraphy
- Franciscan Formation: Bailey, E. H., 3-64.
- Miocene and Pliocene, southern Salinas Valley: Durham, D. L., 1-64.
- Ordovician, paleotectonic significance: Ross, R. J., Jr., 3-64.
- Ordovician formations: Ross, R. J., Jr., 1-64.
- Sierra Nevada metamorphic belt, western: Clark, L. D., 1-64.
- Structural geology
- Cascade Range, gravity and volcanism: Pakiser, L. C., 1-64.
- Northwestern, ultramafic belts: Irwin, W. P., 1-64.
- San Andreas fault: Vedder, J. G., 1-65.
- Surface trace: Schlocker, Julius, 1-65.
- Sierra Nevada metamorphic belt, western: Clark, L. D., 1-64.
- Volcanology
- Lassen Peak, 1915 eruption: MacDonald, G. A., 1-65.

Calorimeter

- Vacuum jacketed hydrofluoric acid solution type: Robie, R. A., 1-65.

Canada

- Structural geology
- Shield, tectonics: Creasey, S. C., 1-64.

Caribbean region

- Paleontology
- Pelecypoda, Cretaceous, oysters: Sohl, N. F., 4-64.

Cartography

- Instruments
- Stereoplotting: Thompson, M. M., 1-64.
- Instruments and techniques: Moore, L. C., 1-65.
- Orthophotography: Scher, M. B., 1-64, 2-64.
- Orthophotomaps: Pumpelly, J. W., 1-64.
- Photogrammetric contouring
- Areas covered by evergreen trees: Scher, M. B., 3-64.
- Stereoplotting: Moore, R. H., 1-64.
- Surface of Earth
- Mapping: Thompson, M. M., 2-64.
- Tables: U. S. Geological Survey, Topographic Division, 1-64.

Caves

- Evolution
- Changes caused by ventilation: Moore, G. W., 1-64.

Caves--Continued

Kentucky

- Cathedral Cave, seasonal temperature fluctuations: Nicholas, G., 1-64.

Central America

Paleontology

- Foraminifera, Recent, offshore: Smith, P. B., 1-64.

Cephalopoda

Carboniferous

- Arkansas: Gordon, Mackenzie, Jr., 2-64.
- California: Gordon, Mackenzie, Jr., 1-64.

Chemical analyses

- Carbonate brines: Rettig, S. L., 1-64.
- Rocks
- Chi-squared method of checking: Flanagan, F. J., 1-64.

Chile

Economic geology

- Nitrate deposits: Ericksen, G. E., 1-65.

Geomorphology

- Atacama Desert, cavities in rock faces: Segerstrom, Kenneth, 1-64.
- Rio Copiapo Valley area: Segerstrom, Kenneth, 1-65.

Heat flow

- Vallenar: Diment, W. H., 2-65.

Maps

- Chanarcillo quadrangle, geologic: Segerstrom, Kenneth, 2-64.

Chromite

Occurrence

- Podiform deposits: Thayer, T. P., 1-64, 2-64.

Stratiform deposits

- Primary features: Jackson, E. D., 1-64.

Clay mineralogy

Areal studies

- New York, Silurian Clinton ironstones: Schoen, Robert, 2-64.

Experimental studies

- Cation-exchange constants: Hanshaw, B. B., 1-64.

- Compaction process: Meade, R. H., 1-64.

- Infrared absorption patterns: Wolff, R. G., 1-65.

- Kaolinite: Wayman, C. H., 1-64.

- Montmorillonite: Brown, T. E., 1-64.

Mineral descriptions

- Kaolinite and montmorillonite, genesis by weathering: Altschuler, Z. S., 2-64.

Clays

- Colorado: Patterson, S. H., 1-64.
- Idaho: Hosterman, J. W., 2-64.
- New Mexico: Patterson, S. H., 2-65.
- South Dakota: Patterson, S. H., 4-64.
- Texas: Corliss, J. B., 1-64.
- Utah: Patterson, S. H., 3-64.

Climate

- Weather modification: Johnson, F. A., 1-64.

Coal

- Alaska
Resources: Barnes, D. F., 1-64.
- Colorado
East Cortez area: Cullins, H. L., 1-65.
Fruitland Formation, Cretaceous: Hinds, J. S., 1-64.
Resources: Landis, E. R., 2-64.
- Idaho
General: Kiilsgaard, T. H., 2-64.
- Iowa
Resources: Landis, E. R., 1-65.
- Montana
Resources: Bateman, A. F., Jr., 1-64.
- New Mexico
Fruitland Formation, Cretaceous: Hinds, J. S., 1-64.
- Pennsylvania
Lawrence County: Van Lieu, J. A., 1-64.
- Rocky Mountains
Future of production: Averitt, Paul, 1-65.
- South Dakota
General: Landis, E. R., 3-64.
- Utah
Resources: Averitt, Paul, 1-64.
Uinta Basin: Schell, E. M., 1-64.
- Cobalt
Idaho: Vhay, J. S., 1-64.
- Colorado
Areal geology
Central City district, Precambrian rocks: Sims, P. K., 2-64.
Cheyenne and Kiowa Counties: Boettcher, A. J., 1-64.
Colorado National Monument: Lohman, S. W., 2-65.
Eastern part of Red-Buffalo Route: Bergendahl, M. H., 3-65.
Flaming Gorge area: Hansen, W. R., 1-65.
Front Range, uranium deposits: Sims, P. K., 3-64.
General: Tweto, Ogden, 1-64.
Grand Junction area: Lohman, S. W., 1-65.
Green River Formation: Donnell, J. R., 1-64.
Associated Eocene rocks: Bradley, W. H., 1-64.
Idaho Springs district, Precambrian rocks: Moench, R. H., 2-64.
Moab quadrangle: Williams, P. L., 1-64.
North Park, northwestern, Late Permian to Late Tertiary: Hail, W. J., Jr., 1-65.
Rosita quadrangle, central and southern Tertiary: Siems, P. L., 1-65.
Straight Creek tunnel site: Robinson, C. S., 1-64.
Ute Mountain Indian Reservation: Irwin, J. H., 1-65.
Ute Mountains area: Ekren, E. B., 1-65.
Washington County: McGovern, H. E., 1-64.

Colorado--Continued

- Economic geology
Alunite, San Juan Mountains: Ratté, J. C., 1-64.
Barite: Brobst, D. A., 2-64.
Beryllium: Griffiths, W. R., 4-64.
Bitumens: Bell, K. G., 1-64.
Clays: Patterson, S. H., 1-64.
Coal, East Cortez area: Cullins, H. L., 1-65.
Fruitland Formation, Cretaceous: Hinds, J. S., 1-64.
General: Landis, E. R., 2-64.
Construction materials: Lindvall, R. M., 1-64.
Cripple Creek District, resources: Bergendahl, M. H., 2-64.
Fluorspar: VanAlstine, R. E., 1-64.
Gem stones: Scott, G. R., 1-64.
Gold, placer: Prommell, H. C., 1-64.
Gypsum and anhydrite: Withington, C. F., 1-64.
Iron: Brown, W. M., 1-64.
Lightweight aggregates: Bush, A. L., 1-64.
Manganese: Crittenden, M. D., Jr., 1-64.
Mineral deposits, guide: Brown, D. L., 1-65.
Molybdenum: King, R. U., 1-64.
Niobium and tantalum: Parker, R. L., 2-64.
Oil shale: Donnell, J. R., 2-64.
Uinta Basin, Green River Formation: Cashion, W. B., 3-64.
Oil shale resources, Green River Formation: Donnell, J. R., 1-64.
Pegmatite minerals: Adams, J. W., 4-64.
Petroleum and natural gas: Bass, N. W., 1-64.
Precious- and base-metal deposits: Bergendahl, M. H., 1-64.
Rare earths: Adams, J. W., 3-64.
Resources, general: U.S. Geological Survey, 12-64.
Salts of sodium and potassium: Raup, O. B., 1-64.
Sand and gravel, construction materials: Carter, W. D., 1-64.
Smelter byproducts, general: Dasch, M. D., 4-64.
Sulfur, nonmetallic and industrial: Broderick, G. N., 1-64.
Thorium, general: Staatz, M. H., 3-64.
Tungsten: Hobbs, S. W., 1-64.
Uranium: Butler, A. P., Jr., 2-64.
Uranium and thorium, Front Range: Phair, George, 1-64.
Vanadium: Fischer, R. P., 1-64.
Zinc: Heyl, A. V., 1-64.
- Engineering geology
Straight Creek Tunnel pilot bore: Robinson, C. S., 1-65, 2-65.
- Geochemistry
Slick Rock district, radioactivity disequilibrium: Rosholt, J. N., 3-65.

Colorado--Continued

- Geochemistry--Continued
Slick Rock district, uranium, isotope fractionation: Rosholt, J. N., 1-65.
- Geomorphology
Loveland Basin landslide: Lee, F. T., 1-65.
Western, hillslope erosion rates: Schumm, S. A., 1-64.
- Geophysical surveys
Bonanza mining area, caldera, gravity: Karig, D. E., 1-65.
Colorado mineral belt, gravity: Case, J. E., 1-65.
Gravity, United States Geological Survey program: Case, J. E., 3-65.
- Hydrogeology
Cheyenne and Kiowa Counties, ground-water: Boettcher, A. J., 1-64; Horr, C. A., 1-64.
Denver Basin: McConaghy, J. A., 1-64.
Grand Junction area: Lohman, S. W., 1-65.
Kiowa Creek basin, fluvial sediment: Mundorff, J. C., 1-64.
North Park and Middle Park, ground water: Voegeli, P. T., Sr., 1-65.
Northeastern, ground water: Weist, W. G., Jr., 1-65.
Prowers County, ground water: Voegeli, P. T., Sr., 2-65.
Pueblo and Fremont Counties, alluvial deposits: McGovern, H. E., 2-64.
Resources, general: U.S. Geological Survey, 12-64.
Upper Colorado River Basin, water resources: Iorns, W. V., 1-64.
Ute Mountain Indian Reservation, ground water, availability: Irwin, J. H., 1-65.
Washington County, ground water: McGovern, H. E., 1-64.
Water resources, general: Odell, J. W., 1-64.
Well records: McConaghy, J. A., 2-64.
- Maps
Berthoud Pass quadrangle, geologic: Theobald, P. K., 1-65.
Central City quadrangle, geologic: Sims, P. K., 1-64.
Cripple Creek district, mine: Koschmann, A. H., 1-65.
Fort Lupton quadrangle, geologic: Soister, P. E., 1-65.
Hot Sulphur Springs quadrangle, geologic: Izett, G. A., 1-65.
Hot Sulphur Springs SW quadrangle, geologic: Izett, G. A., 1-64.
Hudson quadrangle, geologic: Soister, P. E., 2-65.
Iron-ton quadrangle, geologic: Burbank, W. S., 1-64.
Morrison quadrangle, geologic, sedimentary rocks: Smith, J. H., 1-64.

Colorado--Continued

Maps--Continued

- Platteville quadrangle, geologic:
Soister, P. E., 3-65.
Rosita quadrangle, central and
southern, geologic: Siems,
P. L., 1-65.

Mineralogy

- Allanite, Boulder Creek batho-
lith: Hickling, N. L., 1-65.

Petrology

- Sawatch Range, welded tuff:
Brock, M. R., 1-65.
Ute Mountains area: Ekren, E. B.,
1-65.
West Elk laccolith cluster: God-
win, L. H., 1-64.

Stratigraphy

- Cretaceous, Mancos Shale: Dyni,
J. R., 1-65.
Niobrara Formation, Pueblo
area: Scott, G. R., 2-64.
Northwestern: Poole, F. G., 3-64.

Structural geology

- Williams Range thrust fault: Ul-
rich, G. E., 1-65.

Volcanism

- San Juan Mountains: Burbank,
W. S., 1-65.

Colorado Plateau

Geophysical surveys

- Crustal structure, seismic:
Roller, J. C., 2-65.

Connecticut

Areal geology

- Bristol-Plainville-Southington
area, bedrock formation: La-
Sala, A. M., Jr., 1-64.

- Fitchville quadrangle: Snyder,
G. L., 2-64.

- Middletown area: Baker, J. A.,
1-65.

- Niantic quadrangle: Goldsmith,
Richard, 2-65.

- Scotland quadrangle, metamor-
phic rocks, pre-Pennsylvan-
ian age: Dixon, H. R., 1-65.

Hydrogeology

- Bristol-Plainville-Southington
area, ground water: LaSala,
A. M., Jr., 1-64, 2-64.

- Farmington-Granby area: Ran-
dall, A. D., 1-64.

- Ground-water levels, 1960-64:
Meikle, R. L., 1-65.

- Housatonic River basin, water
quality: Pauszek, F. H., 1-65.

- Middletown area: Baker, J. A.,
1-65.

- North-central, well data: Cush-
man, R. V., 1-64.

- Quinebaug River basin: Randall,
A. D., 2-65.

Maps

- Ansonia quadrangle, geologic,
bedrock: Fritts, C. E., 1-65.
Danielson quadrangle, surficial
geology: Randall, A. D., 1-65.
Fitchville quadrangle, surficial
geology: Pessl, Fred, Jr., 1-65.

Connecticut--Continued

Maps--Continued

- Milford quadrangle, geologic:
Fritts, C. E., 2-65.

- New London quadrangle, geologic:
Goldsmith, Richard, 1-65.

- Niantic quadrangle, surficial
geology: Goldsmith, Richard,
1-64.

- Scotland quadrangle, geologic:
Dixon, H. R., 1-65.

- Tariffville quadrangle, geologic:
Schnabel, R. W., 1-65.

- Willamantic quadrangle, geologic:
Snyder, G. L., 1-64.

- Windsor Locks quadrangle:
Schnabel, R. W., 2-64.

Petrology

- Ashaway and Voluntown quad-
rangles: Feininger, Tomas,
1-65.

- Lebanon, gabbro body: Kane,
M. F., 2-64.

Stratigraphy

- Precambrian, quartzite-schist
sequence: Wells, J. D., 1-64.

- Putnam Group: Dixon, H. R.,
1-64.

- Quaternary, South Windsor: Col-
ton, R. B., 1-65.

Continental drift

- Permian indicators: Hamilton
Warren, 2-65.

Copper

- Idaho: Vhay, J. S., 2-64.

- Nevada: Kirkemo, Harold, 1-64.

- New Mexico: Jones, W. R., 1-65.

- Utah: Roberts, R. J., 2-64.

Cosmic dust

- Electron probe analysis: Larson,
R. R., 1-64.

Cratering

- Ejecta with complex structures:
Gault, D. E., 1-64.

- Photogrammetric mapping: Lugin,
R. V., 1-63.

- Porous media: Gault, D. E., 2-64.

- Scaling relationship: Gault, D. E.,
1-65.

Shock

- Equation of state: Carr, M. H.,
1-63.

- Target strength: Moore, H. J.,
3-63, 1-65.

Crustaceans

- Troglobitic crayfish
Arkansas: Hobbs, H. H., Jr.,
1-64.

Crystal chemistry

Apatite

- Isomorphous substitution: Young,
E. J., 1-65.

Crystal structure

- Decavanadate: Evans, H. T., Jr.,
1-65.

- Faheyite: Lindberg, M. L., 1-64.

- Hulsite: Clark, J. R., 1-65.

- Hydrates: Clark, J. R., 2-64.

- Scapolite: Papike, J. J., 1-65.

- Ulexite: Clark, J. R., 2-65.

Czechoslovakia

Paleontology

- Tentaculites: Yochelson, E. L.,
3-65.

Deformation

Biotite

- Kink-bands, nuclear explosions:
Cummings, David, 1-65.

Delaware

Hydrogeology

- Delaware River basin, water re-
sources: Parker, G. G., 2-64.

- New Castle County, southern,
ground water: Rima, D. R.,
1-64.

Maps

- Bethany Beach area, ground wa-
ter: Boggess, D. H., 1-64.

- Clayton area, ground water:
Adams, J. K., 6-64.

- Dover quadrangle, ground water:
Adams, J. K., 9-64.

- Ellendale quadrangle, ground
water: Adams, J. K., 1-64.

- Frankford area, ground water:
Adams, J. K., 7-64.

- Frederica area, ground water:
Davis, C. F., 1-65.

- Georgetown quadrangle, ground
water: Boggess, D. H., 7-64.

- Greenwood quadrangle, ground
water: Boggess, D. H., 2-64.

- Harbeson quadrangle, water and
soils: Adams, J. K., 2-64.

- Harrington quadrangle, ground
water: Davis, C. F., 1-64.

- Hickman area, ground water:
Adams, J. K., 3-64.

- Kenton area, ground water: Bog-
gess, D. H., 1-65.

- Laurel area, ground water: Bog-
gess, D. H., 3-64.

- Lewes area, ground water:
Adams, J. K., 10-64.

- Little Creek quadrangle, ground
water: Boggess, D. H., 2-65.

- Marydel area, ground water:
Davis, C. F., 2-64.

- Milford quadrangle, ground wa-
ter: Boggess, D. H., 5-64.

- Millsboro area, ground water:
Boggess, D. H., 3-65.

- Milton quadrangle, ground wa-
ter: Boggess, D. H., 6-64.

- Mispillion River quadrangle,
ground water: Davis, C. F.,
3-64.

- Rehoboth Beach area, ground
water: Boggess, D. H., 8-64.

- Seaford East quadrangle, ground
water: Adams, J. K., 8-64.

- Seaford West area, ground water:
Boggess, D. H., 4-64.

- Sharptown area, ground water:
Adams, J. K., 4-64.

- Trap Pond area, ground water:
Adams, J. K., 5-64.

- Wyoming quadrangle, ground
water: Boggess, D. H.,
4-65.

- Diatoms
Quaternary
Sweden: Andrews, G. W., 2-65.
- District of Columbia
Hydrogeology
Sediment control: Guy, H. P., 1-63.
- Drainage patterns
Tributaries
Distribution of branches: Giusti, E. V., 1-65.
- Earth
Temperature
Data compilation: Spicer, H. C., 1-64.
- Earthquakes
Magnitude
Hydroseismic data: Vorhis, R. C., 2-65.
Rating scales: Tocher, Don, 1-64.
Review: Pangborn, M. W., Jr., 1-64.
- Ecology
Geologic processes
Past environments reconstructed: Malde, H. E., 1-64.
- Economic geology
Experimental
Plowshare mining: Konselman, A. S., 1-63.
Future energy demand and supply: McKelvey, V. E., 1-64.
- Egypt
Hydrogeology
Ground water, corrosive, metals selection: Clarke, F. E., 1-64.
- Elastic properties
Rocks
Viscoelastic properties: Robertson, E. C., 3-64.
- Elastic waves
 P_n arrivals
Nuclear explosions: Healy, J. H., 1-65.
- Electrical properties
Moist rock: Scott, J. H., 2-64.
Sulfides: Sato, Motoaki, 1-65.
- Engineering geology
Land subsidence
Analysis, consolidation of undisturbed sands: Bull, W. B., 1-64.
Reservoir floors: Eaton, G. P., 1-64.
- Tunnels
Geologic projection: Wahlstrom, E. E., 1-64.
- Waste disposal
Adsorption on soil minerals: Wayman, C. H., 3-64.
Aquifers: Jones, P. H., 1-63.
Biodegradation: Wayman, C. H., 2-64.
Detergents, soft: Wayman, C. H., 1-65.
- Eniwetok
Paleontology
Corals: Wells, J. W., 1-64.
Foraminifera, deep-sea cores: Todd, Ruth, 1-64.
- Erosion
Rate
Determination from root exposure: LaMarche, V. C., Jr., 1-64.
- Eurasia
Paleontology
Ostracoda, marine Triassic: Sohn, I. G., 3-65.
- Evaporation
Cooling ponds
Forced, estimation: Harbeck, G. E., Jr., 1-64.
- Rate
Effect of drought: Koberg, G. E., 1-65.
- Faults
Thrust
Mechanics, osmotic equilibrium: Zen, E-an, 2-65.
- Flame photometry
Oxygen sheath: May, Irving, 1-64.
- Flora
Cretaceous
Pierre Shale: Leopold, E. B., 1-65.
- Tertiary
Green River Formation: Tschudy, R. H., 3-65.
- Vertebraria: Schopf, J. M., 1-64.
- Florida
Areal geology
Glades and Hendry Counties: Klein, Howard, 1-64.
- Economic geology
Bone Valley Formation, geology and geochemistry: Altschuler, Z. S., 1-64.
Lakeland quadrangle: Cathcart, J. B., 1-64.
- Geomorphology
West-central, springs and sinks: Wetterhall, W. S., 1-65.
- Hydrogeology
Alachua, Bradford, Clay, and Union Counties, water resources data: Clark, W. E., 2-64.
Aquifer desalinization: Visser, F. N., 1-65.
Artesian aquifer: Stringfield, V. T., 1-64.
Artesian wells, analysis of water-level fluctuations: Healy, H. G., 1-64.
Deer Point Lake, freshening: Toler, L. G., 1-64.
Escambia and Santa Rosa Counties, water resources records: Musgrove, R. H., 2-65.
Everglades National Park: Hartwell, J. H., 1-64.
Glades and Hendry Counties, ground water resources: Klein, Howard, 1-64.
Hillsborough County, water-resources records: Menke, C. G., 1-64.
Intracoastal waterway, salt-water leakage: Clark, W. E., 1-64.
Orange County, lake levels: Anderson, Warren, 1-65.
Pensacola area, water supply: Musgrove, R. H., 1-65.
Southern: Kohout, F. A., 1-64.
- Mineralogy
Bone Valley Formation, leached zone: Altschuler, Z. S., 2-65.
- Petrology
Land-pebble phosphate deposits: Altschuler, Z. S., 1-65.
- Florida--Continued
Sedimentary petrology
Coastal sands, humate: Swanson, V. E., 1-65.
- Stratigraphy
Comanche Series: Applin, P. L., 1-65.
Paleozoic, lower: Berdan, J. M., 1-65.
- Fluid inclusions
Granitic rocks
Immiscibility in melts: Roedder, Edwin, 1-65.
- Pumice
Dating method: Roedder, Edwin, 2-65.
- Fluorine
Utah
General: Dasch, M. D., 5-64.
- Fluorspar
Colorado: Van Alstine, R. E., 1-64.
Idaho: Anderson, A. L., 1-64.
New Mexico: Van Alstine, R. E., 1-65.
- Foraminifera
Anomalina eaglefordensis: Low, Doris, 1-64.
Caucasina and Aeolomorphella: Loeblich, A. R., Jr., 1-64.
- General
Investigations, 1964: Gibson, T. G., 1-65.
Larger, handbook: Douglass, R. C., 1-65.
Recent literature: Todd, Ruth, 2-64.
- France
Petrology
Peridotites: Thayer, T. P., 1-65.
- Garnet
Idaho
General: Dasch, M. D., 3-64.
- Gas, natural
Alaska
Mineral fuel resources: Grantz, Arthur, 2-64.
- South Dakota
General: Sandberg, C. A., 3-64.
- Utah
General: Gere, W. C., 3-64.
- Gastropoda
Nassarius
Cenozoic: Addicott, W. O., 2-65.
Schizopyga californiana
Cenozoic: Addicott, W. O., 1-65.
- Gems
Utah
General: Dasch, M. D., 6-64.
- General
Angular dispersion
Random vectors: Cox, Allan, 1-64.
Leaders in science: Hubbert, M. K., 1-65.
Science and technology: McKelvey, V. E., 3-64.
United States Geological Survey, research 1964: U. S. Geological Survey, 11-64.
United States Geological Survey open-file reports and maps: Weld, B. A., 1-64.

- Geochemical prospecting
 General: Canney, F. C., 1-65.
 Geobotanical: Cannon, H. L., 1-63.
 Mercury
 Soils and rock: Hinkle, Margaret, 1-65.
 Silver
 Field determination: Nakagawa, H. M., 1-65.
 Geochemistry
 Clays
 Streambed, silica and alumina, leachable: Mallory, E. C., Jr., 1-65.
 Diseases
 Environments of high- and low-rates compared: Cannon, H. L., 1-65.
 Ground water: White, D. E., 1-64.
 Electrochemical potentials: Back, William, 1-65.
 Iron, southern Maryland: Barnes, Ivan, 1-64.
 Lipids
 General: Breger, I. A., 1-65.
 Methods
 Arid regions: Huff, L. C., 1-63.
 Processes
 Travertine deposition: Barnes, Ivan, 1-65.
 Radionuclides
 Research by the United States Geological Survey: Clebsch, Alfred, Jr., 1-63.
 Volcanic emanations: White, D. E., 1-63.
 Geography
 Federal mapping
 Names activity: Orth, D. J., 1-64.
 Geologic thermometry
 Scandium: Herz, Norman, 1-64.
 Geological exploration
 General: Lemmon, D. M., 1-65.
 Nonfuel minerals: Fischer, W. L., 1-65.
 Geological maps
 World: Johnston, W. D., Jr., 1-65.
 Geomorphology
 Deserts
 Piping: Parker, G. G., 1-64.
 Dynamic similarity conditions:
 Scheidegger, A. E., 1-64.
 Flood plains
 Formation: Wolman, M. G., 1-65.
 Fluvial processes: Leopold, L. B., 2-64.
 Processes
 Time, space, and causality:
 Schumm, S. A., 1-65.
 Underfit streams
 Subsurface exploration and chronology: Dury, G. H., 2-64.
 Theory: Dury, G. H., 1-64, 1-65.
 Geophysical exploration
 Airborne control system: Loving, H. B., 1-64.
 Geophysics
 General
 Geophysical Abstracts: Clarke, J. W., 2-64, 3-64, 4-64, 5-64, 6-64, 7-64, 1-65, 2-65, 3-65, 4-65, 5-65, 6-65, 7-65.
 Geophysics--Continued
 New developments: Moxham, R. M., 1-65.
 Georgia
 Areal geology
 Coastal plain, Pleistocene deposits: Herrick, S. M., 1-65.
 Dade County: Croft, M. G., 1-64.
 Dawson County, crystalline rocks: Sever, C. W., 3-64.
 Georgia Nuclear Laboratory: Stewart, J. W., 1-64.
 Paleozoic Rock area: Cressler, C. W., 1-64.
 Economic geology
 Bauxite, Springvale district: Clark, L. D., 1-65.
 Fuller's earth clay: Sever, C. W., 1-64.
 Limestone: Sever, C. W., 6-64.
 Geochemistry
 Brunswick, ground water, C-14 and D content, relation to salt water intrusion: Hanshaw, B. B., 1-65.
 Cardiovascular mortality rates: Shacklette, H. T., 1-64.
 Hydrogeology
 Artesian aquifer: Stringfield, V. T., 1-64.
 Ashland mica schist, Dawson County: Sever, C. W., 4-64.
 Bainbridge, ground water: Sever, C. W., 1-65.
 Brunswick, Ocala Limestone, salt-water intrusion from Claiborne Group: Hanshaw, B. B., 1-65.
 Central Savannah River area, artesian aquifers: Siple, G. E., 1-64.
 Dade County, ground water: Croft, M. G., 1-64.
 Dawson County, ground water: Sever, C. W., 3-64.
 Paleozoic Rock area: Cressler, C. W., 1-64.
 Savannah area, saline-water intrusion: McCollum, M. J., 1-64, 2-64.
 Seminole, Decatur, and Grady Counties: Sever, C. W., 5-64.
 Water-level fluctuations, earthquake effects: Vorhis, R. C., 1-64, 2-64.
 Yellow River basin, surface-water resources: Carter, R. F., 1-64.
 Maps
 Georgia Nuclear Laboratory area, aeromagnetic: Philbin, P. W., 5-64.
 Savannah River Plant area, aeromagnetic: Petty, A. J., 1-65.
 Paleontology
 Foraminifera, Upper Eocene, Burke County: Herrick, S. M., 2-64.
 Stratigraphy
 Eocene to Recent, southeastern: Herrick, S. M., 1-64.
 Georgia--Continued
 Stratigraphy--Continued
 Pulaski County: Vorhis, R. C., 3-65.
 Subsurface, coastal: Herrick, S. M., 2-65.
 Tertiary, offshore extension: McCollum, M. J., 3-64, 4-64.
 Structural geology
 Southwestern, Thomas County, buried Miocene fault: Sever, C. W., 2-64.
 Weathering
 Dawson County, Georgia Nuclear Laboratory, crystalline rocks: Stewart, J. W., 2-64.
 Elberton granite: Ramspott, L. D., 1-65.
 Glaciation
 Continental drift
 Mapping, environmental approach: Cameron, C. C., 1-65.
 Glaciers
 Ice
 General review: Meier, M. F., 1-64.
 Mass budget
 Measurement by hydrometeorology: Tangborn, W. V., 1-64.
 Washington
 South Cascade Glacier: Meier, M. F., 2-64.
 Gold
 Colorado
 Placer: Prommell, H. C., 1-64.
 Exploration
 Spectrophotometric determination: Lakin, H. W., 1-65.
 Idaho
 General: Bergendahl, M. H., 3-64.
 Nevada
 General: Bergendahl, M. H., 4-64.
 New Mexico
 General: Bergendahl, M. H., 2-65.
 South Dakota
 Resources: Bergendahl, M. H., 6-64.
 Utah
 General: Bergendahl, M. H., 5-64.
 Gravity exploration
 Depth of penetration: Watkins, J. S., 2-64.
 Great Lakes area
 Structural geology
 Keweenaw basin: White, W. S., 1-65.
 Greenland
 Areal geology
 Nunatarssuaq area: Fernald, A. T., 2-64.
 Geomorphology
 Mesters Vig area, emerged marine features: Pessl, Fred, Jr., 1-63.
 Glacial geology
 Northern: Davies, W. E., 2-64.
 Paleontology
 Gastropoda, Ordovician, eastern: Yochelson, E. L., 1-64.

- Guam
 Areal geology: Tracey, J. I., Jr., 1-64.
 Petrology
 Limestones: Schlanger, S. O., 1-64.
- Guatemala
 Economic geology
 Review, 1961: Miller, R. L., 2-65.
- Gulf Coastal Plain
 Hydrogeology: LeGrand, H. E., 1-64.
- Gullying
 Earth cracks as a cause
 Southern Arizona, Kam, William, 1-65.
- Gymnosperms
 Cycadaceae
 Review: Fosberg, F. R., 1-64.
- Gypsum
 Colorado
 Resources: Withington, C. F., 1-64.
- Idaho
 Resources: Withington, C. F., 2-64.
- Utah
 General: Withington, C. F., 3-64.
- Virginia
 Occurrence and origin: Withington, C. F., 2-65.
- Hawaii
 Geomorphology
 Oahu, submarine landslides: Moore, J. G., 1-64.
- Geophysical surveys
 United States Geological Survey gravity program: Oliver, H. W., 1-65.
- Volcanoes, magnetic and telluric-electric studies: Prichard, J. I., 1-65.
- Hydrogeology
 Floods, through June 30, 1964: Hoffard, S. H., 1-64.
- Oahu, saline-water intrusion: Visher, F. N., 1-64.
- Surface waters, records, 1950-60: Hendricks, E. L., 1-64.
- Hydrology
 Kona, Holualoa, water-supply potential: Chinn, S. S. W., 1-64.
- Petrology
 Deep-sea basalt: Moore, J. G., 1-65.
- Submarine basalt: Moore, J. G., 2-65.
- Sedimentary petrology
 Halloysite underclay: Patterson, S. H., 2-64.
- Littoral cones, historic: Moore, J. G., 3-65.
- Volcanology
 Infrared surveys: Fischer, W. A., 1-64.
- Kilauea, 1961 eruption: Richter, D. H., 1-64.
- 1962 eruption: Moore, J. G., 2-64.
- seismicity, January 1962-March 1963: Koyanagi, R. Y., 1-64.
- Hawaii--Continued
 Volcanology--Continued
 Kilauea, temperature studies: Peck, D. L., 3-64.
- Vertical ground displacements: Decker, R. W., 1-65.
- Observations, 1962: Krivoy, H. L., 1-64.
- 1963: Koyanagi, R. Y., 2-64; Krivoy, H. L., 1-65.
- 1964: Koyanagi, R. Y., 2-65; Okamura, A. T., 1-65.
- Seismic events 1963: Koyanagi, R. Y., 1-65.
- Heat flow
 Lakes, metamictic: Diment, W. H., 3-65.
- Surface
 Slope effect: Lachenbruch, A. H., 1-65.
- Hydrogen
 Fugacity
 Between 0° and 1000°C: Shaw, H. R., 1-64.
- Hydrogeology
 Aquifers
 Carbonate rocks: McGuinness, C. L., 1-64.
- Constant-head pumping test: Bennett, G. D., 1-64.
- Electrochemical potentials: Back, William, 1-65.
- Porosity determined from gravity measurements: Eaton, G. P., 1-65.
- Tests, theory: Ferris, J. G., 1-65.
- Arid regions: Hofmann, Walter, 1-64.
- Automatic data processing
 Applications: Miller, W. T., 1-64.
- Correlation procedure: Matalas, N. C., 1-64.
- Base flow
 Source: Kilpatrick, F. A., 1-64.
- Bed-material discharge
 Computations: Colby, B. R., 4-64.
- Bibliography
 Flood-routing methods: Yevjevich, V. M., 1-64.
- Ground water, artificial recharge, through 1954: Todd, D. K., 1-65.
- Ground water publications, 1946-55: Vorhis, R. C., 1-65.
- Channels
 Changes, performance: Jefferson, P. O., 1-65.
- Chemistry
 Material soluble in water: Swenson, H. A., 1-65.
- Current meter data
 Analysis, flow-distribution curves: Leve, G. W., 1-64.
- Darcy's law
 Kaolinite: Olsen, H. W., 1-65.
- Depth-discharge relations
 Alluvial streams: Dawdy, D. R., 1-65.
- Discharge characteristics
 Broad-crested weirs: Tracy, H. J., 1-65.
- Hydrogeology--Continued
 Discharge measurement
 Use of radioisotopes: Frederick, B. J., 1-64.
- Dispersion equation
 Numerical solution: Yotsukura, Nobuhiro, 1-64.
- Drainage basins
 Bridge and culvert design: Bodhaine, G. L., 1-64.
- Drainage density
 Streamflow: Carlston, C. W., 1-64.
- Drought
 Data collecting programs: Gilbert, C. R., 1-65.
- Ground water supplies:
 Gabrysch, R. K., 1-65.
- United States, southwest: Gatewood, J. S., 1-64.
- Earthquake effects
 Water supply, Alaskan earthquake of March 27, 1964: Waller, R. M., 1-65.
- Engineering
 Spillway design: Benson, M. A., 1-64.
- Estuaries
 Pollutant distribution: Pyatt, E. E., 1-64.
- Experimental studies
 Analog models: Patten, E. P., Jr., 1-64; Robinove, C. J., 1-65; Wood, L. A., 1-65.
- Phreatophyte research: Robinson, T. W., 1-64.
- Phreatophytes, saltcedar: Robinson, T. W., 1-65.
- United States Geological Survey: McGuinness, C. L., 3-64.
- Exploration
 Infrared imagery: Fischer, W. A., 1-65.
- Floods
 Analog models: Shen, John, 1-65.
- Botanical evidence: Sigafos, R. S., 1-64.
- General
 Career opportunities: Thomas, H. E., 1-65.
- Conversion factors: Johnson, A. I., 1-64.
- International decade: Leopold, L. B., 1-64; Nace, R. L., 1-64, 1-65.
- Jordan Valley, Utah and Holy Land: Bradley, Edward, 1-64.
- Modern: Hackett, O. M., 1-64.
- Primer on water quality: Swenson, H. A., 2-65.
- Review: Taylor, G. C., Jr., 1-65.
- Hydrographs
 Accuracy, current meters: Colby, B. R., 2-64.
- Nonlinear instantaneous unit-theory: Dawdy, D. R., 1-64.
- Ground water
 A key resource: McGuinness, C. L., 1-65.

Hydrogeology--Continued

Ground water--Continued

- Contaminated zones, patterns:
LeGrand, H. E., 2-65.
- Contamination, environmental
framework: LeGrand, H. E.,
1-65.
- Management: LeGrand, H. E.,
2-64.
- General review: Meinzer, O. E.,
1-65.
- Management: Nace, R. L., 3-65.
- Exploitation and conservation:
Moulder, E. A., 1-64.
- Motion, Darcy's concept: Ski-
bitzke, H. E., 1-64.
- Movement, theory: Smith, W. O.,
1-64.
- Piezometric measurement, er-
rors: Emmett, W. W., 2-64.
- Radio nuclides from underground
nuclear explosions: Stead,
F. W., 1-64.
- Spectra of conditions: LeGrand,
H. E., 3-65.
- Vertical movements, Earth's
thermal profile: Bredehoeft,
J. D., 1-65.

Maps

- Classification: Wiesnet, D. R.,
2-64.

Nuclear explosions

- Water development and manage-
ment: Piper, A. M., 1-65.

Paleoecology

- Reconstructions: Langbein, W. B.,
1-64.

Peak discharge: Kindsvater, C. E.,
1-65.

Pollution

- Amino acids in sewage: Kahn,
Lloyd, 1-64.

Porous media

- Dispersion, mathematics: Ogata,
Akio, 2-64.

Multiphase fluids: Stallman, R. W.,
1-64.One-dimensional fluid flow: Stal-
lman, R. W., 1-65.Spread of dye stream: Ogata,
Okio, 1-64.

Rainfall

- Sulfate and nitrate: Gambell,
A. W., 1-64.

Recharge

- Hidden: Feth, J. H., 1-64.

Rivers, soluble contaminants

- Time of travel: Buchanan, T. J.,
1-64.

Rivers and lakes

- Chemical composition: Living-
stone, D. A., 1-64.

Runoff

- Areal variations: Crippen, J. R.,
1-65; Hely, A. G., 1-64.

Meteorological factors: Busby,
M. W., 1-64.

Saline-water intrusion

- Scavenger well system: Long,
R. A., 1-64.

Sedimentation

- Effect of water temperature:
Colby, B. R., 1-65.

Hydrogeology--Continued

Sedimentation--Continued

- Sand-bed streams: Colby, B. R.,
1-64.

Soil-moisture

- Infiltration of rain water: Rubin,
Jacob, 1-64.

Stream flow

- Alluvial channels: Simons, D. B.,
2-65.

- Measurement, submerged sec-
tion controls: Eisenlohr, W. S.,
1-64.

- Records correlated with pre-
cipitation: Martin, R. O. R.,
1-64.

- Supply and management concepts:
Leopold, L. B., 1-65.

Surface water

- Bank storage: Rorabaugh, M. I.,
1-64.

- Discharge measurements: La-
Cornu, E. J., 1-65.

- Highly permeable environment:
Sherwood, C. B., 1-64.

- Runoff, areal variations: Riggs,
H. C., 1-65.

- Sediment in streams: Swenson,
H. A., 1-64.

Technique

- Aquifer permeability: Weeks,
E. P., 1-64.

- Collection and analysis of water
samples: Rainwater, F. H.,
1-65.

- Data processing, water resources:
Johnson, A. I., 1-65.

- Fluorescent tracing: Wright, R. R.,
1-64.

- Ground-water, base-flow reces-
sion curve: Riggs, H. C., 1-64.

- Numerical-analysis methods,
transmissibility variations:
Sammel, E. A., 1-64.

- Ground-water data, storage and
retrieval: Lang, S. M., 1-65.

- Measuring bedload: Hubbell,
D. W., 1-64.

- Resource studies: Mesnier, G. N.,
1-64.

- Stream quality instrumentation:
McCartney, D., 1-65.

- Surface-water, stream-discharge
regressions using precipita-
tion: Riggs, H. C., 2-64.

- Stage-discharge relation, de-
finition: Ray, H. A., 1-64, 1-65.

- Water-resources development,
analog evaluation: Robinson,
G. M., 1-64.

Temperature

- Streams, annual variation: Curtis,
L. W., 1-64; Haushild, W. L.,
1-64.

Tritium, uses

- Problems and limitations: Carl-
ston, C. W., 2-64, 1-65.

Turbulent flow

- Laboratory study: Sayre, W. W.,
1-64.

Waste disposal

- Contamination potential evalua-
tion: LeGrand, H. E., 3-64.

Hydrogeology--Continued

Waste disposal

- Contamination potential evalua-
tion: LeGrand, H. E., 3-64.

- Radioactive, crystalline rocks:
Siple, G. E., 2-64.

Water analysis

- Herbicides: Goerlitz, D. F.,
1-65.

- Nitrate and nitrite, diazotization
method: Fishman, M. J., 2-64.

- Pesticides, chlorinated organic:
Lamar, W. L., 1-64.

- Vanadium, catalytic determina-
tion: Fishman, M. J., 1-64.

Water development

- Cultural control: Thomas, H. E.,
1-64.

- Water management: Leopold,
L. B., 3-65.

- Arid lands: Davis, G. H., 1-64.

Water resources

- Ecological systems: Leopold,
L. B., 2-65.

Water rights

- Cultural approach: Thomas, H. E.,
2-65.

Water supply

- Economic development: Nace,
R. L., 2-64.

- Problems: Nace, R. L., 2-65.

Hydrothermal alteration

Feldspar

- Supersaturated silica solutions:
Fournier, R. O., 1-65.

Idaho

Areal geology

- General: Ross, C. P., 1-64.

- National Reactor Testing Station:
Walker, E. H., 3-64.

- National Reactor Testing Station
area: Bates, R. G., 1-65.

- National Reactor Testing Station,
deep well site selection, geo-
logy: Chase, G. H., 1-64.

- Owyhee County, Schooner site:
Gard, L. M., 3-63.

Economic geology

- Antimony, Big Creek-Yellow
Pine area: Leonard, B. F.,
1-65.

- General: La Heist, B. A., 2-64.

- Barite: Brobst, D. A., 1-64.

- Beryllium: Griffiths, W. R., 5-64.

- Clays: Hosterman, J. W., 2-64.

- Coal: Kiilsgaard, T. H., 2-64.

- Cobalt and nickel: Vhay, J. S.,
1-64.

- Coeur d'Alene district: Fryklund,
V. C., Jr., 1-64.

- Bleached rocks: Weis, P. L., Jr.,
1-64.

- Copper: Vhay, J. S., 2-64.

- Fluorspar: Anderson, A. L., 1-64.

- Garnet: Dasch, M. D., 3-64.

- General: Weissenborn, A. E., 1-64.

- Gold: Bergendahl, M. H., 3-64.

- Gypsum and anhydrite: Withing-
ton, C. F., 2-64.

- Manganese: Prinz, W. C., 2-64.

- Mercury: Bailey, E. H., 2-64.

- Mica and pegmatite minerals:
Lesure, F. G., 1-64.

Idaho--Continued

- Economic geology--Continued
 Mineral industry: Weissenborn, A. E., 2-64.
 Molybdenum: King, R. U., 4-64.
 Niobium and tantalum: Parker, R. L., 1-64.
 Peat: Savage, C. N., 1-64.
 Phosphate rock: Gere, W. C., 1-64.
 Resources, general: U.S. Geological Survey, 16-64.
 Salt, resources: Walker, E. H., 2-64.
 Silica: Carter, W. D., 2-64.
 Silver, lead, and zinc: Kiilsgaard, T. H., 3-64.
 Smelter byproducts: LaHeist, B. A., 3-64.
 Tungsten: Hobbs, S. W., 3-64.
 Uranium: Armstrong, F. C., 1-64.
 Vanadium: Fischer, R. P., 4-64.
 Engineering geology
 Buggy event, potential sites: Hasler, J. W., 1-63.
 Geophysical surveys
 National Reactor Testing Station area, radioactivity: Bates, R. G., 1-65.
 Saylor Creek site, seismic: Gard, L. M., 1-63.
 Hydrogeology
 Aberdeen-Springfield area, water-level data, 1963: Sisco, H. G., 1-64.
 Resources, general: U.S. Geological Survey, 16-64.
 Sandpoint region, ground water: Walker, E. H., 1-64.
 Snake River basin, ground water: Mundorff, M. J., 2-64.
 Teton Valley, ground water: Kilburn, Chabot, 1-64.
 Water resources: Travis, W. I., 1-64.
 Waterpower: Young, L. L., 2-64.
 Maps
 Conant Valley quadrangle, geologic: Jobin, D. A., 1-64.
 Irwin quadrangle, geologic: Jobin, D. A., 2-64.
 Lanes Creek quadrangle, aeromagnetic: Meuschke, J. L., 1-65.
 National Reactor Testing Station area, radioactivity: Bates, R. G., 1-64.
 Packsaddle Mountain quadrangle, geologic: Harrison, J. E., 1-65.
 Palisades Peak quadrangle, geologic: Jobin, D. A., 1-65.
 Pocatello-Soda Springs area, aeromagnetic: Mitchell, C. M., 1-65.
 Poker Peak-Palisades Reservoir quadrangles, geologic: Albee, H. F., 1-65.
 Paleontology
 Mollusca, Jurassic: Imlay, R. W., 3-64.
 Petrology
 Belt Series, metamorphism: Hietanen, Anna, 1-65.

Idaho--Continued

- Petrology--Continued
 Yellow Pine quadrangle, Tertiary dike swarm: Leonard, B. F., 2-65.
 Stratigraphy
 Brigham, Langston, and Ute Formations: Oriel, S. S., 1-65.
 Precambrian and Cambrian, Pocatello area: Trimble, D. E., 1-65.
 Structural geology
 Late Paleozoic tectonic history: Roberts, R. J., 3-65.
 Paris thrust fault: Armstrong, F. C., 1-65.
 Strike-slip faulting: Ruppel, E. T., 1-64, 1-65.
 Thrust belt, tectonic development: Armstrong, F. C., 2-65.
 Igneous rocks
 Classification
 Feldspar ratios: O'Connor, J. T., 1-65.
 Translation from Russian of Zavaritsky's system: Blake, M. C., Jr., 1-64.
 Effusive
 Minor and rare-earth elements: Fleischer, Michael, 3-64.
 Ocean areas: Engel, A. E. J., 1-65.
 Rhyolite
 Flow direction: Cummings, David, 1-63.
 Ultramafic
 Alkali content: Hamilton, Warren, 3-65.
 Volcanics
 Chemical composition: Lipman, P. W., 1-65.
 Ground-water leaching: Noble, D. C., 4-65.
 Illinois
 Areal geology
 Cuba City, New Diggings, and Shullsburg quadrangles, geologic: Mullens, T. E., 1-64.
 Geochemistry
 Brines, Illinois basin, origin: Clayton, R. N., 1-65.
 Geomorphology
 Lakes, ancient, Finch, W. I., 3-64.
 Hydrogeology
 Blue Island quadrangle, floods: Allen, H. E., 1-65.
 Elgin quadrangle, floods: May, V. J., 1-64.
 Fox Lake quadrangle, floods: Noehre, A. W., 4-64.
 Mokena quadrangle, floods: Noehre, A. W., 1-65.
 Romeoville quadrangle, floods: Noehre, A. W., 2-64.
 San Bridge quadrangle, floods: Noehre, A. W., 3-64.
 Streamwood quadrangle, floods: May, V. J., 2-64.
 Tinley Park quadrangle, floods: Allen, H. E., 2-65.
 West Chicago quadrangle, floods: Allen, H. E., 3-64.

Illinois--Continued

- Hydrogeology--Continued
 Wheaton quadrangle, floods: May, V. J., 3-64.
 Maps
 Geneva quadrangle, floods: Noehre, A. W., 2-65.
 Harvey area, floods: Allen, H. E., 1-64.
 Joliet quadrangle, floods: Allen, H. E., 4-64.
 Lombard area, floods: Allen, H. E., 2-64.
 Naperville quadrangle, floods: Allen, H. E., 3-65.
 Wadsworth quadrangle, floods: Noehre, A. W., 1-64.
 Indiana
 Geochemistry
 Metals in black shale: Vine, J. D., 1-65.
 Hydrogeology
 Fulton County, ground water: Rosenshein, J. S., 1-64.
 Jasper County, ground water: Rosenshein, J. S., 2-64.
 Marshall County, ground water: Rosenshein, J. S., 3-64.
 Montgomery County, ground water resources: Cable, L. W., 1-65.
 Newton County, ground water: Rosenshein, J. S., 4-64.
 Parke County, ground water: Watkins, F. A., Jr., 1-64.
 Pulaski County, ground water: Rosenshein, J. S., 5-64.
 Putnam County, ground water: Watkins, F. A., Jr., 2-64.
 Saint Joseph County, Quaternary beds: Hunn, J. D., 1-64.
 Starke County, ground water: Rosenshein, J. S., 6-64.
 Sullivan and Greene Counties, Mississippian, Pennsylvanian, and Quaternary: Cable, L. W., 2-65.
 Vigo and Clay Counties, aquifers: Cable, L. W., 1-64.
 Maps
 Cloverport quadrangle, geologic: Bergendahl, M. H., 1-65.
 Columbus-Dayton area, aeromagnetic: Philbin, P. W., 1-65.
 Mattingly quadrangle, geologic: Clark, L. D., 2-65.
 Tell City quadrangle, geologic: Spencer, F. D., 2-64.
 Infrared analysis
 Saline sulfate minerals: Erd, R. C., 1-65.
 Iodine
 Geochemistry
 Vegetation: Cuthbert, Margaret, 1-64.
 Iowa
 Areal geology
 Omaha-Council Bluffs area: Miller, R. D., 1-64.
 Economic geology
 Coal, resources: Landis, E. R., 1-65.

Iowa--Continued

Hydrogeology

- Central: Twenter, F. R., 1-65.
English River, Old Mans Creek,
and Clear Creek basins: Sch-
wob, H. H., 1-64.

Maps

- Central, aeromagnetic:
Henderson, J. R., 1-65.

Iron

Alaska

- General: Cobb, E. H., 2-64.

Brazil

- Minas Gerais, high-grade hema-
tite: Dorr, J. V. N., 2d, 2-65.

Colorado

- Resources: Brown, W. M., 1-64.

Nevada

- General: Reeves, R. G., 2-64.

Utah

- General: Reeves, R. G., 3-64.

Isotopes

Water

- Variations, measurement: Mc-
Carthy, J. H., 1-65.

Israel

Hydrogeology

- Cenomanian-Turonian aquifer:
Schneider, Robert, 1-64.

Paleontology

- Ostracoda, Quaternary: Sohn,
I. G., 2-65.

Japan

Paleontology

- Algae, Eocene: Johnson, J. H.,
1-64.
Tertiary megafossils: MacNeil,
F. S., 1-64.

Kansas

Hydrogeology

- Floods, control, storage require-
ments: Furness, L. W., 1-64.
Grant and Stanton Counties: Fader,
S. W., 1-64.
Ground water: Dingman, R. J.,
1-64.
Ground-water levels, 1963:
Broeker, M. E., 1-64.
Kansas River basin, fluvial sedi-
ment, 1957-60: Mundorff,
J. C., 3-64.
Surface-water quality, 1962:
Mayes, J. L., 1-64.
Rivers: Leonard, R. B., 3-64.
Walnut River basin, water quality:
Leonard, R. B., 1-64, 2-64.

Petrology

- Hutchinson area, Wellington For-
mation, evaporites: Jones,
C. L., 1-65.

Stratigraphy

- Shawnee County, sample logs:
Adkison, W. L., 1-64.

Karst

- Lithological control: Monroe,
W. H., 3-64.

Kentucky

Areal geology

- Fordsville quadrangle: Bergen-
dahl, M. H., 7-64.
Middlesboro South quadrangle:
Englund, K. J., 1-64.

Kentucky--Continued

Geochemistry

- Metals in black shale: Vine,
J. D., 1-65.

Geomorphology

- Lakes, ancient: Finch, W. I.,
3-64.

Geophysical surveys

- Bell Scott Quarry drill hole, logs:
Kepferle, R. C., 1-64.
Hardin County, Summit area
drill hole, radioactivity log:
Moore, F. B., 1-64.
Regional implications: Watkins,
J. S., 1-64.

Glacial geology

- Northern part, Nebraskan and
Kansan: Leighton, M. M., 1-65.

Hydrogeology

- Beaver Creek basin, strip min-
ing effects: Collier, C. R., 2-64.
Blue Grass region, geochemistry:
Hendrickson, G. E., 1-64.
Jenkins-Whitesburg area, ground
water: Mull, D. S., 1-65.
Lynn Grove quadrangle, ground
water, availability: Davis,
R. W., 2-64.
Mammoth Cave National Park,
present and future water supply:
Cushman, R. V., 2-64.
Mayfield, aquifer test: Morgan,
J. H., 5-64.

Maps

- Auburn quadrangle, geologic:
Rainey, H. C., III, 1-65.
Barthell and Oneida North quad-
rangle, geologic: Pomerene,
J. B., 1-64.
Blackwater quadrangle, geologic:
Stager, H. K., 1-64.
Boltsfork and Burnaugh quad-
rangles, geologic: Spencer,
F. D., 1-64.
Briensburg quadrangle, geologic:
Lambert, T. W., 2-64, Mac-
Cary, L. M., 4-64.
Geologic and ground water:
MacCary, L. M., 1-64.
Brushart quadrangle, geologic:
Denny, C. S., 1-64.
Burtonville quadrangle, geologic:
Morris, R. H., 1-65.
Cane Valley quadrangle, geologic:
Maxwell, C. H., 1-64.
Charters quadrangle, geologic:
Morris, R. H., 2-65.
Clintonville quadrangle, geolog-
ic: MacQuown, W. C., 1-65.
Cloverpart and Cannelton quad-
rangles, geologic: Bergen-
dahl, M. H., 1-65.
Constantine quadrangle, geologic:
Sable, E. G., 1-64.
Cuba quadrangle, geologic: Blade,
L. V., 1-64.
Dexter quadrangle, ground water:
Davis, R. W., 1-64.
Dunnville quadrangle, geologic:
Maxwell, C. H., 1-65.
Elva quadrangle, ground water:
Morgan, J. H., 1-64.

Kentucky--Continued

Maps--Continued

- Fairdealing quadrangle, geologic:
Wolfe, E. W., 1-64.
Ground water: Lambert, T. W.,
1-65.
Farmington quadrangle, ground
water: Morgan, J. H., 2-64.
Fenton quadrangle, geologic:
Schnabel, R. W., 3-64.
Freedom quadrangle, geologic:
Moore, S. L., 1-64.
Glasgow North quadrangle,
geologic: Haynes, D. D., 1-64.
Glasgow South quadrangle,
geologic: Moore, S. L., 1-65.
Grand Rivers quadrangle,
geologic: Hays, W. H., 1-64.
Hanson quadrangle, geologic:
Franklin, G. J., 1-65.
Hardin quadrangle, geologic and
ground water: MacCary, L. M.,
2-64.
Hazard North quadrangle, geo-
logic: Seiders, V. M., 1-64.
Hazard South quadrangle, geo-
logic: Puffett, W. P., 1-64.
Hazel quadrangle, ground water:
MacCary, L. M., 3-64.
Henderson area, geologic and
hydrologic: Gallaher, J. T.,
1-64.
Hico quadrangle, geologic:
Olive, W. W., 1-65.
Ground water: Morgan, J. H., 1-65.
Hima quadrangle, geologic:
Reeves, R. G., 1-64.
Hiseville quadrangle, geologic:
Haynes, D. D., 1-65.
Honey Grove quadrangle, geo-
logic: Klemic, Harry, 1-65.
Kirksey quadrangle, ground wa-
ter: Morgan, J. H., 3-64.
Krypton quadrangle, geologic:
Mixon, R. B., 1-65.
Lancer quadrangle, geologic:
Rice, C. L., 1-64.
Little Cypress and Calvert City
quadrangles, ground water:
Morgan, J. H., 2-65.
Madisonville West quadrangle,
geologic: Kehn, T. M., 1-64.
Mammoth Cave quadrangle, geo-
logic: Haynes, D. C., 1-64.
Manchester quadrangle, geologic:
Finnell, T. L., 1-64.
Maretburg quadrangle, geologic:
Schlanger, S. O., 1-65.
Matewan quadrangle, geologic:
Trent, V. A., 1-65.
Mattingly quadrangle, geologic:
Clark, L. D., 2-65.
Mayfield quadrangle, geologic:
Finch, W. I., 1-65.
Mont quadrangle, geologic: Weis,
P. L., Jr., 2-64.
Montpelier quadrangle, geologic:
Lewis, R. Q., Sr., 1-64.
New Boston and Wheelersburg
quadrangles, geologic: Shep-
pard, R. A., 1-64.

Kentucky--Continued

Maps--Continued

New Concord and Buchanan quadrangles, geologic: Wilshire, H. G., 1-64.

New Concord quadrangle, ground water: Lambert, T. W., 1-64.

Oak Level quadrangle, ground water: Morgan, J. H., 4-64.

Offutt quadrangle, geologic: Outerbridge, W. F., 1-64.

Ohio River between Catlettsburg and South Portsmouth, geologic and hydrologic: Price, W. E., Jr., 4-64.

Ohio River between Prospect and southwestern Louisville, geologic and hydrologic: Price, W. E., Jr., 1-64.

Ohio River between southwestern Louisville and West Point, geologic and hydrologic: Price, W. E., Jr., 2-64.

Oldtown quadrangle, geologic: Whittington, C. L., 1-65.

Petroleum quadrangle, geologic: Myers, W. B., 1-64.

Phil quadrangle, geologic: Maxwell, C. H., 2-65.

Pleasant Green Hill quadrangle, geologic: Nelson, W. H., 1-64.

Portersburg quadrangle, geologic: Pomerene, J. B., 2-64.

Portsmouth quadrangle, geologic: Sheppard, R. A., 1-64.

Princeton West quadrangle, geologic: Sample, R. D., 1-65.

Rockport and Lewisport quadrangles, geologic: Spencer, F. D., 3-64.

Rome quadrangle, geologic: Crittenden, M. D., Jr., 6-65.

Russell Springs quadrangle, geologic: Lewis, R. Q., Sr., 2-65.

Slaughters quadrangle, geologic: Kehn, T. M., 2-64.

Smiths Grove quadrangle, geologic: Richards, P. W., 1-64.

Stanley area, geologic and hydrologic: Gallaher, J. T., 2-64.

Sulphur Lick quadrangle, geologic: Harris, L. D., 1-64.

Summer Shade quadrangle, geologic: Hail, W. J., Jr.

Symsonia quadrangle, geologic: Finch, W. I., 2-64.

Ground water: Davis, R. W., 1-65.

Tell City quadrangle, geologic: Spencer, F. D., 2-64.

Temple Hill quadrangle, geologic: Moore, S. L., 2-65.

Tyrone quadrangle, geologic: Cressman, E. R., 1-64.

Uniontown and Wickliffe areas, geologic and hydrologic: Gallaher, J. T., 3-64.

Versailles quadrangle, geologic: Black, D. F. B., 1-64.

Paleontology

Flora, Hazard No. 7 coal:

Kosanke, R. M., 3-65.

Palynomorphs, coals: Kosanke, R. M., 2-65.

Kentucky--Continued

Paleontology--Continued

Palynomorphs, Pennsylvanian: Kosanke, R. M., 1-65.

Sedimentation

Mammoth Cave: Collier, C. R., 1-64.

Stratigraphy

Ohio River between Catlettsburg and south Portsmouth alluvial deposits: Price, W. E., Jr., 5-64.

Ohio River between Prospect and southwestern Louisville alluvial deposits: Price, W. E., Jr., 3-64.

Ohio River between southwestern Louisville and West Point, alluvial deposits: Price, W. E., Jr., 6-64.

Ordovician, Lexington Limestone and Clays Ferry Formation: Black, D. F. B., 1-65.

Korea

Hydrogeology

Reconnaissance study: Doyel, W., 1-64.

Lakes

Ancient deposits

Western states, Precambrian to Pleistocene: Feth, J. H., 2-64.

Geochemistry

Great Salt Lake: Hahl, D. C., 1-64.

Temperature

Lake Champlain, Cumberland Bay area: Ayer, G. R., 1-64.

Landslides

South Dakota

Fort Randell Reservoir vicinity: Erskine, C. F., 1-65.

Lead

Abundance

Hawaii and Japan: Tatsumoto, M., 2-65.

Idaho

General: Kiilsgaard, T. H., 3-64.

Isotopes

Age of Earth: Tilton, G. R., 1-65.

Basalt: Tatsumoto, M., 1-65; Tilton, G. R., 1-64.

Feldspars, regional metamorphism: Doe, B. R., 2-65.

Montana, Boulder batholith: Doe, B. R., 1-65.

Nevada

General: Kleinhampl, F. J., 1-64.

Utah

General: Kiilsgaard, T. H., 4-64.

Leveling

Instruments

Heat shield: Maltby, C. S., 1-65.

Recent studies: Karren, R. J., 1-64.

Libya

Maps

Ground water: Jones, J. R., 1-64.

Lithium

Nevada

General: Griffiths, W. R., 3-64.

Louisiana

Hydrogeology

Aquifers, methane: Harder, A. H., 2-65.

Louisiana--Continued

Hydrogeology--Continued

Bossier and Caddo Parishes, water resources: Page, L. V., 1-64.

De Soto Parish, water resources: Page, L. V., 2-64.

Floods, 1964: Lowe, A. S., 1-64.

Magnitude and frequency: Sauer, V. B., 1-64.

Franklin Parish, alluvial deposits: Marie, J. R., 1-65.

Ground water: Roberts, J. E., 1-64.

Jefferson Davis Parish, surface water: Sloss, Raymond, 1-65.

Sabine River, base-flow studies, 1963: Sauer, S. P., 1-64.

Southeastern, storm runoff: Sauer, V. B., 2-64.

Southwestern: Turcan, A. N., Jr., 2-65.

Ground water, methane content: Harder, A. H., 1-65.

Vernon Parish: Rogers, J. E., 1-65.

Maps

Baton Rouge area, floods, 1962: Camp, J. D., 1-65.

Magmas

Granite

Immiscibility, fluid inclusions: Roedder, Edwin, 1-65.

Viscosity and crystal settling: Shaw, H. R., 1-65.

Water pressure

Indicators: Wones, D. R., 1-65.

Magnetic field, Earth

Variations

Long period: Cox, Allan, 2-64.

Magnetic properties

Igneous rocks

Remanent magnetization: Doell, R. R., 1-65.

Magnetite-bearing grout

Drill holes: Black, R. A., 1-63.

Palladium

300°K to 4.2°K: Thorpe, A. N., 1-64.

Maine

Areal geology

Northeastern: Pavlides, Louis, 2-64.

Geochemistry

West-central part, lead-zinc anomalies: Canney, F. C., 1-64.

Maps

Amity quadrangle, aeromagnetic: Meuschke, J. L., 1-64.

Big Lake quadrangle, geologic: Larrabee, D. M., 1-64.

Bingham quadrangle, aeromagnetic and geologic: Mattick, R. E., 1-65.

Cupsuptic quadrangle, aeromagnetic: Boynton, G. R., 1-64.

Forks quadrangle, aeromagnetic: Bromery, R. W., 2-64.

Greenville quadrangle, geologic: Espenshade, G. H., 1-64.

Northern, geologic and aeromagnetic: Boucot, A. J., 1-64.

Old Speck Mountain quadrangle, aeromagnetic: Henderson, J. R., 1-64.

- Maine--Continued
 Maps--Continued
 Oquossoc quadrangle, aeromagnetic: Boynton, G. R., 2-64.
 Phillips quadrangle, aeromagnetic: Boynton, G. R., 3-64.
 Rangeley and Kennebec Lake quadrangles, aeromagnetic: Boynton, G. R., 4-64.
 Wabassua Lake quadrangle, geologic: Larrabee, D. M., 2-64.
 Paleontology
 Flora, Mapleton Sandstone: Schopf, J. M., 3-64, 5-64.
 Ordovician tuffs, fossils: Neuman, R. B., 1-64.
 Petrology
 Granodiorite, Attean quadrangle: Albee, A. L., 1-65.
 Stratigraphy
 Devonian, Matagamon Sandstone: Rankin, D. W., 1-65.
 Hovey Formation: Pavlides, Louis, 1-64.
 Mammalia
 Tertiary
 Central American seaways: Whitmore, F. C., Jr., 1-65.
 Man, fossil
 Dating: Yochelson, E. L., 2-65.
 Manganese
 Colorado
 Resources: Crittenden, M. D., Jr., 1-64.
 Idaho
 General: Prinz, W. C., 2-64.
 Nevada
 General: Crittenden, M. D., Jr., 4-64.
 New Mexico
 General: Dorr, J. V. N., 2d., 1-65.
 Utah
 General: Crittenden, M. D., Jr., 5-64.
 Mantle
 Upper Mantle Project
 United States Geological Survey participation: Hall, W. E., 1-65.
 Marshall Islands
 Paleontology
 Echinoids: Kier, P. M., 1-64.
 Einwetok Atoll, fossil Bryozoa: Brown, D. A., 1-64.
 Maryland
 Areal geology
 Evitts Creek and Pattersons Creek quadrangles, stratigraphy and geologic history: deWitt, Wallace, Jr., 1-64.
 Economic geology
 Gravel deposits, Beltsville quadrangle: Withington, C. F., 1-65.
 Geophysical surveys
 Aeromagnetic: Beck, M. E., Jr., 1-64.
 Hydrogeology
 Salisbury, shallow wells: Heidel, S. G., 1-65.
 Southern, iron-rich ground water: Barnes, Ivan, 1-64.
 Maps
 Bel Air and vicinity, aeromagnetic: Bromery, R. W., 4-64.
 Maryland--Continued
 Maps--Continued
 Bel Air and vicinity, radioactivity: Bromery, R. W., 3-64.
 Massachusetts
 Areal geology
 Andover Granite: Castle, R. O., 1-64.
 Cape Cod, subsurface: Oldale, R. N., 1-65.
 Nantasket area: Bell, K. G., 2-64.
 Norwood and Blue Hills quadrangles: Chute, N. E., 1-64.
 Tiverton quadrangle: Pollock, S. J., 1-64.
 Glacial geology
 Martha's Vineyard: Kaye, C. A., 1-64.
 Hydrogeology
 Assabet River basin: Pollock, S. J., 2-64.
 Ipswich River basin: Sammel, E. A., 1-65.
 Wilmington-Reading area, ground water conditions: Baker, J. A., 2-64.
 Maps
 Assawompset Pond quadrangle, geologic: Koteff, Carl, 1-64.
 Berlin quadrangle: Holmes, G. W., 5-65.
 Canaan quadrangle: Holmes, G. W., 4-65.
 Concord quadrangle, geologic: Koteff, Carl, 2-64.
 East Lee quadrangle, aeromagnetic: Popenoe, Peter, 1-64.
 Ipswich River Basin, hydrogeology: Sammel, E. A., 2-64.
 North Adams quadrangle, aeromagnetic: Popenoe, Peter, 2-64.
 Otis quadrangle: Holmes, G. W., 2-65.
 Pittsfield West quadrangle: Holmes, G. W., 3-65.
 Salem quadrangle, surficial geology: Oldale, R. N., 1-64.
 Tarrifville quadrangle, geologic: Schnabel, R. W., 1-65.
 Windsor quadrangle: Holmes, G. W., 6-65.
 Petrology
 Salem area, igneous rocks: Toulmin, Priestley 3d, 1-64.
 Sedimentation
 Cape Cod, beach and eolian sands: Schlee, John, 2-64.
 Stratigraphy
 Marlboro Formation: Cuppels, N. P., 1-64.
 Quaternary, Martha's Vineyard: Kaye, C. A., 2-64.
 Mercury
 Analysis
 Vegetation: Ward, F. N., 1-64.
 Geochemical prospecting
 Instrumental technique: Vaughn, W. W., 1-64, 2-64.
 Soils and rock: Hinkle, Margaret, 1-65.
 Idaho
 General: Bailey, E. H., 2-64.
 Mercury--Continued
 Nevada
 General: Bailey, E. H., 1-64.
 Resources
 General: Bailey, E. H., 4-64.
 Utah
 General: Hilpert, L. S., 4-64.
 Metamorphic rocks
 Eclogite
 Garnet and pyroxene compositional variations: Coleman, R. G., 1-65.
 Varieties: Coleman, R. G., 3-65.
 Iron formations
 Precambrian, origin: Lepp, Henry, 1-64.
 Pelitic schists
 Coexisting muscovite and paragonite: Zen, E-an, 1-64.
 Metamorphism
 Regional
 Lead isotopes, feldspars: Doe, B. R., 2-65.
 Water pressure
 Indicators: Wones, D. R., 1-65.
 Meteorites
 Composition
 Achondrites, metallic iron: Duke, M. B., 1-63, 1-65.
 Metallic copper: Duke, M. B., 2-63, 2-65.
 Oxygen, isotopes: Taylor, H. F., Jr., 1-64.
 Michigan
 Areal geology
 Marenisco-Watersmeet area: Fritts, C. E., 3-65.
 Geochemistry
 Brines, Michigan basin, origin: Clayton, R. N., 1-65.
 Marquette iron range, prospecting: Segerstrom, Kenneth, 2-65.
 Geomorphology
 Ontonagon area, postglacial drainage: Hack, J. T., 2-65.
 Geophysical surveys
 Buried bedrock valleys, gravity: Klasner, J. S., 1-64.
 Lake Superior and Burt Lake, bottom sediments, radioactivity, measuring device: Bunker, C. M., 1-64.
 Hydrogeology
 Au Sable River, water management: Hendrickson, G. E., 1-65.
 Ground-water summary, 1963: Giroux, P. R., 2-64.
 Menominee County, ground water: Vanlier, K. E., 1-63.
 Recharge, glacial drift aquifer: Reed, J. E., 1-64.
 Tri-County region, ground water: Vanlier, K. E., 1-64.
 Van Buren County, water resources: Giroux, P. R., 1-64.
 Maps
 Baraga County, eastern, aeromagnetic: Balsley, J. R., 12-65.
 Gogebic and Ontonagon Counties, aeromagnetic: Boynton, G. R., 5-64.

Michigan--Continued

Maps--Continued

Iron, Baraga, Houghton, Ontonagon, and Gogebic Counties, aeromagnetic: Balsley, J. R., 10-65.

Matchwood, Ontonagan, Rockwood, and Greenwood quadrangles, aeromagnetic: Balsley, J. R., 1-64.

Ontonagon, Houghton, and Baraga Counties, aeromagnetic: Balsley, J. R., 11-65.

Upper Peninsula, aeromagnetic: Case, J. E., 4-65.

Microscope technique

Nonopaque minerals: Larsen, E. S., 1-64.

2V measurement: Noble, D. C., 1-65.

Universal stage

Plagioclase determination: Noble, D. C., 2-65.

Mineral descriptions

Biotite

Polytypism: Ross, Malcolm, 2-65.

Trace elements: Herz, Norman, 1-64.

Brucite

Alpine serpentinites: Hostetler, P. B., 1-65.

Buddingtonite: Erd, R. C., 1-64.

Calcite

Black, source of silver: Hewett, D. F., 1-65.

Chlorite

Wyoming, Green River Formation: Bradley, W. H., 2-64.

Coesite

Separation from sandstone shocked by meteorite impact: Fahey, J. J., 1-63.

Cymrite

New data: Carron, M. K., 1-65.

Enstatite

Aluminous: Skinner, B. J., 2-64.

Epidote

X-ray analysis: Myer, G. H., 1-65.

Feldspars

Composition and structure: Wright, T. L., 2-64.

Hambergite

Fluorine content: Switzer, George, 1-65.

Heulandite: Mittin, Ivan, 1-65.

Hollandite: Fleischer, Michael, 1-64.

Hulsite: Clark, J. R., 1-65.

Laumontite

Alaska: Hoare, J. M., 1-64.

Lazurite

Wyoming, Green River Formation: Bradley, W. H., 2-64.

Macallisterite: Schaller, Waldemar, 1-65.

McKelveyite: Milton, Charles, 1-65.

Schoepite

Phase transitions: Christ, C. L., 1-65.

Serpentine: Faust, G. T., 1-64.

Mineral descriptions--Continued

Stishovite

Separation from sandstone shocked by meteorite impact: Fahey, J. J., 1-63.

Synthesis by shock waves: DeCarli, P. S., 1-65.

Talc

Wyoming, Green River Formation: Bradley, W. H., 2-64.

Torbernite: Ross, Malcolm, 1-65.

Ulexite: Clark, J. R., 3-64, 2-65.

Veatchite

Formula: Clark, J. R., 1-64.

Zeolites: Foster, M. D., 1-65.

Mineral exploration

Recent progress: McKelvey, V. E., 2-64.

Mineralogy

General

Book review: Stewart, D. B., 1-65.

Identification

MnCO₃-CaCO₃-MgCO₃, determination of composition: Prinz, W. C., 1-64.

Refractive energy

CuO and Sc₂O₃: Mrose, M. E., 1-65.

Review of "Mineralogy of the rare earths," Fleischer, Michael, 2-65.

Review of "Mineralogy, Spravochnik, Tom I and Tom II": Fleischer, Michael, 2-64.

Minnesota

Areal geology

Aurora area: Maclay, R. W., 1-64.

Nobles and Jackson Counties, geologic: Norvitch, R. F., 1-64.

Redwood Falls area: Schiner, G. R., 1-64.

Glacial geology

Mountain Iron-Virginia-Eveleth area: Cotter, R. D., 1-64.

Hydrogeology

Aurora area: Maclay, R. W., 1-64.

Minneapolis-St. Paul area: Maderak, M. L., 1-64.

Nobles and Jackson Counties, ground water: Norvitch, R. F., 1-64.

Maps

East-central, aeromagnetic and geologic: Bath, G. D., 1-64.

Mesabi-Vermilion Iron Range area, geologic: Cotter, R. D., 2-64.

Middle River watershed, water resources: Maclay, R. W., 1-65.

Normal and part of Mahnomen Counties, aeromagnetic: Anderson, L. A., 1-64.

Northwestern, aeromagnetic and geologic: Bath, G. D., 2-64.

Mississippi

Economic geology

Bauxite, Tippah-Benton district: Tourtelot, H. A., 1-64.

Bauxite and kaolin deposits: Conant, L. C., 1-65.

Mississippi--Continued

Geochemistry

Cores, vicinity of Tatum salt dome: Jenne, E. A., 1-63.

Geophysical surveys

Lamar County, Project Dribble, radioactivity: Bunker, C. M., 3-63.

South, wells near shot points:

Grantham, P. E., 1-63.

Hydrogeology

Corinth area: Newcome, Roy, Jr., 1-64.

Cretaceous aquifers: Boswell, E. H., 2-65.

Floods, 1961: Wilson, K. V., 1-64.

1962: Neely, B. L., Jr., 1-64.

Ground water, dissolved solids:

Boswell, E. H., 1-65.

Jackson County, water resources, status: Newcome, Roy, Jr., 2-64.

Lower Little Creek basin:

Skelton, John, 1-63.

Mississippi embayment, low-

flow characteristics of streams:

Speer, P. R., 1-64.

Northwestern, water supplies:

Wasson, B. E., 1-64.

Pascagoula area, water re-

sources: Harvey, E. J., 1-65.

Pearl River at Jackson, floods, 1961: Shell, J. D., 1-64.

Salt water intrusion, 1964: Lang, J. W., 1-64.

Southeastern, water resources:

Shows, T. N., 1-64.

Southwestern, storm runoff:

Sauer, V. B., 2-64.

Tatum Dome, aquifer tests: Koop-

man, F. C., 1-63.

Tatum Salt Dome, movement of radionuclides: Beetem, W. A., 1-63.

Tatum Salt Dome area: Taylor, R. E., 1-63.

Water supply, industrial develop-

ment: Wasson, B. E., 2-64.

Yazoo County, ground water re-

sources: Callahan, J. A., 1-64.

Paleontology

Gastropoda, Coffee Sand, Cre-

taceous: Sohl, N. F., 1-64.

Late Cretaceous: Sohl, N. F., 2-64.

Petrology

Tatum Dome: Schlocker, Julius, 1-63.

Structural geology

Tatum Dome, Lamar County:

Eargle, D. H., 1-63.

Mississippi embayment

Hydrogeology

Surface water, low flow charac-

teristics: Speer, P. R., 2-64,

3-64.

Mississippi Valley

Economic geology

Lead-zinc district, clay-mineral alteration: Heyl, A. V., 2-64.

Missouri

Geophysical surveys

Low-amplitude aeromagnetic anomalies: Allingham, J. W., 1-64.

Hydrogeology

Surface water, chemical quality: Taylor, C. T., 1-64.

Mohole

Sediments

Guadalupe site, ion-exchange capacity: Carroll, Dorothy, 1-64.

Mohorovičić discontinuity

Basalt-eclogite transformation zone, relation to: Pakiser, L. C., 1-65.

Mollusca

Hyolithes

Paleozoic: Marek, Ladislav, 1-64.

Mya

Evolution and distribution, Tertiary: MacNeil, F. S., 1-65.

Molybdenum

Colorado

General: King, R. U., 1-64.

General

Supply and consumption: Fischer, R. P., 5-64.

Idaho

General: King, R. U., 4-64.

New Mexico

General: King, R. U., 1-65.

South Dakota

General: King, R. U., 2-64.

Utah

General: King, R. U., 3-64.

Montana

Areal geology

Black Hills uplift, northern and western flanks: Robinson, C. S., 2-64.

Blackfeet Indian Reservation: Paulson, Q. F., 1-65.

Bluewater Springs area: Zimmerman, E. A., 1-64.

Cleveland quadrangle: Schmidt, R. G., 1-64.

Rattlesnake quadrangle, geologic: Hearn, B. C., Jr., 1-64.

Economic geology

Black Butte iron deposit: Wier, K. L., 1-65.

Coal, resources: Bateman, A. F., Jr., 1-64.

Dolomite, Henena area: King, R. H., 1-64.

Eastern, potential: U.S. Geological Survey, 6-65.

Geochemistry

Boulder batholith, lead isotopes: Doe, B. R., 1-65.

Glacial geology

Glacier National Park, observations, 1964: Johnson, Arthur, 1-65.

Hydrogeology

Blackfeet Indian Reservation, ground water: Paulson, Q. F., 1-65.

Bluewater Springs area: Zimmerman, E. A., 1-64.

Cedar Creek anticline, ground water: Taylor, O. J., 1-65.

Montana--Continued

Maps

Bannack-Grayling area, geologic: Lowell, W. R., 1-65.

Black Butte iron deposit: Wier, K. L., 1-65.

Sawtooth Ridge quadrangle, geologic: Mudge, M. R., 1-65.

Petrology

Boulder batholith, variation in modes and norms: Tilling, R. I., 1-64.

Southwestern, Permian rocks: Cressman, E. R., 2-64.

Stratigraphy

Cretaceous and Tertiary, Livingston area: Roberts, A. E., 1-65.

Devonian, Maywood Formation: Sandberg, C. A., 2-64.

Jefferson and Three Forks Formations: Sandberg, C. A., 1-65.

Permian rocks, southwestern: Cressman, E. R., 2-64.

Structural geology

Strike-slip faulting: Ruppel, E. T., 1-64, 1-65.

Moon

Areal geology

Aristarchus quadrangle: Moore, H. J., 2-63.

Colombo quadrangle, pre-Imbrian stratigraphy: Elston, D. P., 1-63.

Eastern, stratigraphic relations: Elston, D. P., 1-65.

Hevelius area: McCauley, J. F., 1-63.

Mare Humorum quadrangle: Titley, S. R., 1-63.

Mare Orientale region: McCauley, J. F., 2-63.

Riphaeus quadrangle: Eggleton, R. E., 1-63.

Taruntius quadrangle: Wilhelm, D. E., 1-63.

Timocharis quadrangle: Carr, M. H., 2-63.

Craters

Density of small craters: Moore, H. J., 1-63.

Infrared emission: Watson, Kenneth, 1-63.

Maps

Isotonal: Hackman, R. J., 1-63.

Microphotometer measurements:

Miesch, A. T., 1-63.

Photographic and photoelectric studies: Morris, E. C., 2-63.

Stratigraphy

Julius Caesar area: Morris, E. C., 1-63.

Montes Apenninus quadrangle: Hackman, R. J., 2-63.

Surface features

Slopes, photometric measurement: Wilhelm, D. E., 2-63.

Terrain analysis

Equatorial belt: McCauley, J. F., 1-65.

Nebraska

Areal geology

Franklin, Webster, and Nuckolls Counties: Miller, R. D., 2-64.

Omaha-Council Bluffs area: Miller, R. D., 1-64.

Richardson County: Emery, P. A., 2-64.

Hydrogeology

Blue River basin, ground water-surface water relations: Emery, P. A., 1-65.

Chapman area, chemical quality of water: Rosene, P. G., 1-64.

Ground-water conditions: Keech, C. F., 3-64.

Hall County, availability of ground water: Keech, C. F., 4-64.

Lower Platte River basin, ground water: Hyland, J. B., 1-64.

Mirage Flats, ground-water resources: Keech, C. F., 2-64.

Observation wells, water levels, 1964: Emery, P. A., 2-65.

Richardson County, ground water: Emery, P. A., 2-64.

Saline County, basic-data report: Emery, P. A., 1-64.

York County, basic-data report: Keech, C. F., 1-64.

Nevada

Areal geology

Ammonia Tanks quadrangle: Hinrichs, E. N., 1-63.

Ash Meadows quadrangle: Denny, C. S., 2-65.

Black Mountain volcanic: Christiansen, R. L., 1-65.

Bullfrog quadrangle: Cornwall, H. R., 3-64.

Deadhorse Flat quadrangle: Noble, D. C., 1-64.

Frenchie Creek quadrangle: Muffler, L. J. P., 1-64.

General: Wallace, R. E., 1-64.

Humboldt County: Willden, Ronald, 1-64.

Jackass Flats quadrangle: McKay, E. J., 1-63.

Jangle Ridge quadrangle: Barnes, Harley, 1-63.

Mine Mountain quadrangle: Orkild, P. P., 1-63.

Nevada Test Site: Barnes, Harley, 2-63; Bowers, W. E., 2-63, 3-63, 4-63, 5-63.

Cane Spring road to Mara Wash: Poole, F. G., 1-63.

Ferris Wheel site: Cummings, David, 2-63.

Pahute Mesa Exploratory Hole: Hoover, D. L., 1-63, 4-63, 5-63.

U12b.10 tunnel: Emerick, W. L., 5-63.

U12e.07a tunnel: Hasler, J. W., 2-63.

U12g tunnel: Emerick, W. L., 2-63.

U12g.05-U1 drill hole: Emerick, W. L., 1-65.

U12n tunnel: Emerick, W. L., 3-63.

U19a site: Hoover, D. L., 2-63.

U20a drill hole area: Emerick, W. L., 1-63.

Nevada--Continued

Areal geology--Continued

Nevada Test Site--Continued

U20b site: Hoover, D. L., 3-63.

U20c and U20d sites: Hasler, J. W., 1-64.

U20e site: Hoover, D. L., 1-64.

U20g site: Schnabel, R. W., 1-64.

U3ams sink: Houser, F. N., 1-63.

U3cn site: Williams, W. P., 2-63.

U3cs site: Williams, W. P., 1-63.

U3cw site: Snyder, R. P., 3-63.

Wahmonie Flat area: Johnson, R. B., 1-63.

Quartet Dome quadrangle: Sargent, K. A., 1-64.

Scrugham Peak quadrangle: Byers, F. M., Jr., 2-64.

Shoshone Range, tectonics and igneous rocks: Gilluly, James, 2-65.

Silent Canyon SW quadrangle: Ekren, E. B., 1-63.

Skull Mountain quadrangle: Sargent, K. A., 1-63.

Steamboat Springs area: Thompson, G. A., 1-64; White, D. E., 2-64.

Topopah Spring SW quadrangle: Lipman, P. W., 1-64.

Economic geology

Antler Peak quadrangle, ore deposits: Roberts, R. J., 2-65.

Arsenic: La Heist, B. A., 1-64.

Beryllium: Griffiths, W. R., 1-64.

Bismuth: Mount, Priscilla, 1-64.

Borates: Smith, W. C., 1-64.

Bullfrog quadrangle, related ore deposits: Cornwall, H. R., 3-64.

Carbon dioxide: Pierce, A. P., 1-65.

Copper: Kirkemo, Harold, 1-64.

Gold: Bergendahl, M. H., 4-64.

Helium: Pierce, A. P., 2-65.

Humboldt County, mineral deposits: Wildden, Ronald, 1-64.

Iron: Reeves, R. G., 2-64.

Kyanite group: Tatlock, D. B., 1-64.

Lithium: Griffiths, W. R., 3-64.

Manganese: Crittenden, M. D., Jr., 4-64.

Mercury: Bailey, E. H., 1-64.

Mineral resources: Roberts, R. J., 1-64.

General: Cornwall, H. R., 1-64, 2-64.

Pegmatites: Lesure, F. G., 2-64.

Resources, general: U.S. Geological Survey, 17-64.

Rowe Canyon area, mineral resources: Gott, G. B., 2-64.

Selenium and tellurium: Davidson, D. F., 1-64.

Shoshone Range: Gilluly, James, 2-65.

Silver, lead, and zinc: Kleinhampl, F. J., 1-64.

Thorium and rare earths: Staatz, M. H., 2-64.

Uranium: Butler, A. P., Jr., 1-64.

Vanadium: Fischer, R. P., 2-64.

Nevada--Continued

Engineering geology

HARDHAT event, explosion-produced fractures: Houser, F. N., 1-64.

Nevada Test Site, nuclear explosion base surge and throw-out material: Colton, R. B., 3-63.

Surface effects of Bilby event: Williams, W. P., 2-64.

Surface effects of underground test: Williams, W. P., 1-64.

Geochemistry

Getchell Mine, exploration: Erickson, R. L., 2-64.

Geomorphology

Alluvial fans: Denny, C. S., 1-65.

Geophysical surveys

Alluvium, electrical properties: Hazlewood, R. M., 1-63.

Antler Peak quadrangle, aeromagnetic: Mabey, D. R., 1-64.

Clark County, gravity: Kane, M. F., 1-64.

Crecent Valley, gravity: Gilluly, James, 2-65.

Dacite, elastic constants, seismic determination: Black, R. A., 2-63.

Nevada Test Site, boreholes, physical properties of rocks: Davis, R. E., 1-63, 1-64.

Density log, porosity: Carroll, R. D., 1-63.

Density-porosity of tuff: Carroll, R. D., 7-63.

Electrical: Carroll, R. D., 1-64.

Electrical: Scott, J. H., 1-63, 1-64.

Geophysical logs: Carroll, R. D., 2-63, 3-63, 4-63, 5-63, 6-63.

Gravity: Healey, D. L., 2-63.

Gravity and faulting: Healey, D. L., 1-63.

Gold Meadows stock: Healey, D. L., 3-63.

Physical properties of rocks: Ege, J. R., 1-62.

Radioactivity logging: Bunker, C. M., 1-62.

Reentry tunnel, radioactivity: Bunker, C. M., 2-63.

Rock samples, radioactivity: Bunker, C. M., 1-63, 4-63, 5-63, 6-63, 7-63.

Subsurface measurements, radioactivity: Bunker, C. M., 1-65.

U12q drill hole, physical properties of tuff: Emerick, W. L., 4-63.

U12g, 01 tunnel, physical properties of tuff: Emerick, W. L., 6-63.

UE1a, UE1b, and UE1c drill holes: Emerick, W. L., 1-64.

United States Geological Survey investigations: Hazlewood, R. M., 2-63.

United States Geological Survey gravity program: Oliver, H. W., 1-65.

Nevada--Continued

Geophysical surveys--Continued

Winnemucca site, seismic: Gard, L. M., 2-63.

Hydrogeology

Aquifer response, Bilby event: Hale, W. E., 1-63.

Aquifers, intermontane basins: Bredehoeft, J. D., 2-64.

Aurora Site, test well: Norvitch, R. F., 1-63.

Fernley-Wadsworth area: Sinclair, W. C., 1-63.

Grass Valley area, ground-water: Cohen, Philip, 1-64, 5-64.

Humboldt River valley: Cohen, Philip, 2-64, 3-64.

Jackass Flats, well sites: Young, R. A., 1-63.

Kings River Valley, groundwater: Zones, C. P., 1-63.

Meadow Valley area, ground water: Rush, F. E., 1-64.

Mid Valley area, ground water: Young, R. A., 2-63.

Monitor, Antelope, and Kobeh Valleys, ground-water: Rush, F. E., 2-64.

Muddy River Springs, surface water: Eakin, T. E., 2-64.

Nevada Test Site, ground-water fluctuations, Aardvark explosion: Garber, M. S., 1-63.

Pahute Mesa Exploratory Hole: Meyer, G. L., 1-63.

Resources, general: U.S. Geological Survey, 17-64.

Rock Valley, ground water: Winograd, I. J., 1-63.

Smith Creek and Ione Valleys, ground-water appraisal: Everett, D. E., 1-64.

South-central, interbasin: Eakin, T. E., 3-64.

Interbasin ground-water movement: Winograd, I. J., 1-64.

Southeastern, ground water system: Eakin, T. E., 1-64.

Truckee Meadows area, evaluation: Cohen, Philip, 4-64.

Water resources, general: Cornwall, H. R., 1-64; Thomas, H. E., 2-64.

Maps

Ammonia Tanks quadrangle, geologic: Hinrichs, E. N., 1-64.

Buckboard Mesa quadrangle, geologic: Byers, F. M., Jr., 3-64.

Cane Spring quadrangle, geologic: Poole, F. G., 2-64.

Jackass Flats quadrangle, geologic: McKay, E. J., 1-64.

Jangle Ridge quadrangle, geologic: Barnes, Harley, 1-65.

Mine Mountain quadrangle, geologic: Orkild, P. P., 1-64.

Mount Rose and Virginia City quadrangles, aeromagnetic: Dempsey, W. J., 1-65.

North-central, Paleozoic and Mesozoic facies: Roberts, R. J., 1-65.

Nevada--Continued

Maps--Continued

- Plutonium Valley quadrangle, geologic: Hinrichs, E. N., 1-65.
 Quartet Dome quadrangle, geologic: Sargent, K. A., 2-65.
 Scrugham Peak quadrangle, geologic: Byers, F. M., Jr., 1-64.
 Silent Butte quadrangle, geologic: Ekren, E. B., 1-64.
 Skull Mountain quadrangle, geologic: Ekren, E. B., 2-64.
 Thirsty Canyon quadrangle, geologic: O'Connor, J. R., 1-65.
 Thirsty Canyon SE quadrangle, geologic: Lipman, P. W., 2-65.

Paleontology

- Diatoms, Mesozoic and Cenozoic: Lohman, K. E., 1-64.
 Flora, Miocene, Fingerrock Wash: Wolfe, J. A., 1-64.
 Mollusca, Cambrian: Yochelson, E. L., 4-65.
 Ostracoda, southern, Triassic: Sohn, I. G., 1-64.
 Pioche Shale faunules: Palmer, A. R., 2-64.
 Trilobites, Cambrian: Palmer, A. R., 1-64.

Petrology

- Granitic plutons, Esmeralda County: Albers, J. P., 1-64.
 Jackass Flats quadrangle, hydrothermal alteration: McKay, E. J., 2-63.
 Nevada Test Site, basalt: Luft, S. J., 1-63.
 Basalt, heavy minerals: Sargent, K. A., 3-65.
 Basalt, identification by magnetic properties: Sargent, K. A., 1-65.
 Lapilli tuff, porosity: Manger, G. E., 1-65.
 Zeolite zoning: Hoover, D. L., 1-65.
 Nye County, Thirsty Canyon Tuff: Noble, D. C., 3-65.
 Roberts Thrust rocks, chemical analysis: Erickson, R. L., 3-64.
 Southern, Cenozoic tuffs: Quinlivan, W. D., 1-65.
 Tertiary and Quaternary rocks: Albers, J. P., 2-64.
 Timber Mountain caldera, mafic lavas: Luft, S. J., 1-64.
 Tuffs: Quinlivan, W. D., 1-63.
 Vitric tuff, alteration, Rainier Mesa quadrangle: Bowers, W. E., 1-63.
 Volcanic rocks, space and time associations, southern: Noble, D. C., 5-65.

Stratigraphy

- Antler Peak quadrangle: Roberts, R. J., 4-64.
 Cambrian, Pioche mining district: Merriam, C. W., 1-64.
 General: Wallace, R. E., 1-64.
 Mesozoic: Silberling, N. J., 1-64.
 Ordovician, paleotectonic significance: Ross, R. J., Jr., 3-64.
 Ordovician formations: Ross, R. J., Jr., 1-64.
 Paleozoic: Roberts, R. J., 3-64.

Nevada--Continued

Stratigraphy--Continued

- Permian: Barosh, P. J., 1-65.
 Lower, east-central: Barosh, P. J., 1-64.
 Precambrian and Lower Cambrian: Stewart, J. H., 1-64.
 Structural geology
 Antler Peak quadrangle: Roberts, R. J., 4-64.
 General: Wallace, R. E., 2-64.
 Groom Mine SW quadrangle, faults in alluvium: Colton, R. B., 1-63.
 Late Paleozoic tectonic history: Roberts, R. J., 3-65.
 Nevada Test Site, buried pre-Tertiary surface: Miller, C. H., 1-63.
 Faults in alluvium: Colton, R. B., 2-63.
 North-central, overthrusting: Wallace, R. E., 3-64.
 Schell Creek Range, thrust and glide faults: Drewes, Harald, 1-65.
 Timber Mountain caldera: Christiansen, R. L., 2-65.

New England

Geomorphology

- Bedrock valleys, coast: Upson, J. E., 1-64.

New Hampshire

Areal geology

- Southeastern: Bradley, Edward, 2-64.

Geochemistry

- Th and U, early Paleozoic plutonic series: Lyons, J. B., 1-64.

Hydrogeology

- Connecticut River basin, ground water: Cederstrom, D. J., 1-65.
 Southeastern, ground water resources: Bradley, Edward, 2-64.

Maps

- Salem-Plaistow area, ground water: Weigle, J. M., 1-64.

New Jersey

Economic geology

- Gravel: Schlee, John, 1-64.

Hydrogeology

- Delaware River, quality of water: McCarthy, L. T., Jr., 1-64.
 Englishtown Formation, water chemistry: Seaber, P. R., 1-65.
 Floods, depth frequency: Thomas, D. M., 2-64.
 Magnitude and frequency: Thomas, D. M., 1-64.
 Great Swamp, surface water: Miller, E. G., 1-65.
 Morris County, ground water availability: Gill, H. E., 1-64.
 Stream-flow records, base-flow relations: Buchanan, T. J., 1-65.
 Surface water, quality of streams: Anderson, P. W., 1-65.

New Jersey--Continued

Maps

- Belvidere quadrangle, aeromagnetic: Boynton, G. R., 2-65.
 Radioactivity: Boynton, G. R., 1-65.
 Bristol quadrangle, geologic: Owens, J. P., 1-64.
 Hackensack and Paterson quadrangles, aeromagnetic: Philbin, P. W., 3-64.
 Nyack and part of White Plains quadrangles, aeromagnetic: Philbin, P. W., 2-64.
 Park Ridge quadrangle, aeromagnetic: Philbin, P. W., 2-64.
 Roosevelt quadrangle, geologic: Minard, J. P., 1-64.
 Trenton East quadrangle, geologic, pre-Quaternary: Owens, J. P., 2-64.
 Yonkers and Mount Vernon quadrangles, aeromagnetic: Philbin, P. W., 4-64.
 Woodstown quadrangle, geologic: Minard, J. P., 1-65.

Petrology

- Island Beach State Park, basement gneiss: Southwick, D. L., 1-64.

Stratigraphy

- Cambrian-Ordovician, Warren and Hunterdon Counties, carbonate rocks: Drake, A. A., Jr., 1-65.

New Mexico

Areal geology

- General: Dane, C. H., 1-65.
 Guadalupe Mountains: Hayes, P. T., 1-64.
 Lake McMillan and Carlsbad Springs: Cox, E. R., 1-65.
 Los Alamos area: Griggs, R. L., 1-64.
 McKinley County, southeastern: Cooper, J. B., 2-65.
 Tucumcari vicinity: Trauger, F. D., 1-64.
 Ute Mountain Indian Reservation: Irwin, J. H., 1-65.
 Economic geology
 Antimony, arsenic, bismuth, and cadmium: Dasch, M. D., 1-65.
 Beryllium: Griffiths, W. R., 6-65.
 Clays: Patterson, S. H., 2-65.
 Coal, Fruitland Formation, Cretaceous: Hinds, J. S., 1-64.
 Construction material: Lindvall, R. M., 1-65.
 Copper: Jones, W. R., 1-65.
 Diatomite: Patterson, S. H., 1-65.
 Fluorspar: Van Alstine, R. E., 1-65.
 Gem materials: Carter, M. D., 1-65.
 General: Bachman, G. O., 1-65.
 Gold: Bergendahl, M. H., 2-65.
 Manganese: Dorr, J. V. N., 2d, 1-65.
 Mineral industry: Bachman, G. O., 2-65.
 Molybdenum: King, R. U., 1-65.
 Niobium and tantalum: Parker, R. L., 1-65.

New Mexico--Continued

Economic geology--Continued

Nitrates and guano: Hayes, P. T., 1-65.

Pegmatite: Lesure, F. G., 1-65.

Potash: Alto, B. R., 2-65.

Rare earths: Adams, J. W., 1-65.

Resources: U. S. Geological Survey, 7-65.

Rhenium: King, R. U., 2-65.

Salts: Alto, B. R., 1-65.

Sand and gravel: Carter, W. D., 1-65.

Selenium and tellurium: Davidson, D. F., 1-65.

Sulfur: Broderick, G. N., 1-65.

Thorium: Staatz, M. H., 1-65.

Tin: Sainsbury, C. L., 2-65.

Tungsten: Hobbs, S. W., 1-65.

Uranium, general: Hilpert, L. S., 1-65.

Vanadium: Fischer, R. P., 1-65.

Waterpower: Senkpiel, W. C., 1-65.

Geophysical surveys

Carlsbad area, radioactivity: MacKallor, J. A., 1-65.

Grants and Laguna areas, geothermal: Byerly, P. E., 1-65.

Gravity, United States Geological Survey program: Case, J. E., 3-65.

Hydrogeology

Acoma and Laguna Indian Reservations, ground water: Dinwiddie, G. A., 2-64.

Arkansas River basin, planning: Ballance, W. C., 2-65.

Central closed basins, surface water: Titus, F. B., Jr., 1-65.

Colorado River basin, surface water, lower: Trauger, F. D., 1-65.

El Morro National Monument: West, S. W., 1-65.

Ground-water levels, 1963: Ballance, W. C., 1-65.

Lake McMillan and Carlsbad Springs: Cox, E. R., 1-65.

Lea County, recharge studies: Havens, J. S., 1-64.

Los Alamos area, ground water: Griggs, R. L., 1-64.

Major Johnson Springs aquifer: Cushman, R. L., 1-65.

McKinley County, ground water: Cooper, J. B., 2-65.

Municipal water supplies: Dinwiddie, G. A., 1-64.

Navajo and Hopi Indian Reservations: Cooley, M. E., 1-64, 2-64.

Pecos River basin, planning report: Mourant, W. A., 1-65.

Resources: U. S. Geological Survey, 7-65; West, S. W., 2-65.

Rio Grande, hydraulic variables: Culbertson, J. K., 1-64.

Rio Grande basin, planning report: Dinwiddie, G. A., 1-65.

Rio Grande conveyance channel: Harris, D. D., 1-64.

New Mexico--Continued

Hydrogeology--Continued

Roswell basin, artificial recharge: Motts, W. S., 1-64.

Tritium content: Reeder, H. O., 1-64.

San Juan River basin, planning report: Cooper, J. B., 3-65.

Southern High Plains, planning: Ballance, W. C., 3-65.

Southwestern closed basins, planning report: Doty, G. C., 1-65.

Tucumcari vicinity, ground water: Trauger, F. D., 1-64.

Ute Mountain Indian Reservation, ground water, availability: Irwin, J. H., 1-65.

Western closed basins, planning report: Cooper, J. B., 1-65.

Maps

Carlsbad area, radioactivity: MacKallor, J. A., 2-65.

Catskill NE quadrangle, geologic: Pillmore, C. L., 1-65.

Catskill SE quadrangle, geologic: Pillmore, C. L., 2-65.

Dough Mountain quadrangle, geologic: Moench, R. H., 1-64.

Hurley West quadrangle, geologic: Pratt, W. P., 1-65.

La Gotera quadrangle, geologic: Moench, R. H., 1-65.

South Butte quadrangle, geologic: Moench, R. H., 3-64.

Tularosa Basin, ground-water: Herrick, E. H., 1-65.

Santa Rita quadrangle, geologic: Hernon, R. M., 2-64.

Three Rivers area, water resources: Hood, J. W., 1-65.

Mineralogy

Attapulgit, Carlsbad Caverns: Davies, W. E., 1-64.

Sedimentary petrology

Madera Limestone, rhythmic sedimentation: Myers, D. A., 2-65.

Rio Grande area: Nordin, C. F., Jr., 1-64, 1-65.

Stratigraphy

Chinle and Moenkopi Formations, Navajo and Hopi Indian Reservations: Repenning, C. A., 2-65.

Weathering

Chaco Canyon National Monument, fall of threatening rock: Schumm, S. A., 2-64.

New York

Economic geology

Iron, St. Lawrence County: Leonard, B. F., 1-64.

Geomorphology

Oneida Lake area, bedrock topography: Kantrowitz, I. H., 1-64.

Hydrogeology

Babylon-Islip area: Pluhowski, E. J., 1-64.

Brookhaven National Laboratory and vicinity, chemical quality of water: DeLaguna, Wallace, 1-64.

New York--Continued

Hydrogeology--Continued

Delaware River basin, water quality: Archer, Robert, 1-63.

Fall Creek basin, time of travel studies: Dunn, Bernard, 2-65.

Floods, 1964: Robinson, F. L., 1-65.

Genesee River basin, water resource records: Gilbert, B. K., 1-65.

Glacial deposits, ground water hardness: Seaber, P. R., 2-64.

Glowegee Creek, AEC plant waste water, 1958-61: Pauszek, F. H., 1-64, 2-65.

Haines Falls area, surface water resources: Hunt, O. P., 2-65.

Irondequoit Creek basin: Dunn, Bernard, 1-65.

Lake Erie-Niagara area: LaSala, A. M., Jr., 3-64.

Long Island, low-flow summary: Hunt, O. P., 1-65.

Nassau County, ground water, chemistry: Lieber, Maxim, 1-64.

Niagara County, borehole periscope, fracture study: Trainer, F. W., 1-64.

Niagara Falls area, ground water: Johnston, R. H., 1-64.

South Farmingdale area: Perlmutter, N. M., 1-64.

Syracuse area, ground water: Kantrowitz, I. H., 2-64.

Maps

Berlin quadrangle: Holmes, G. W., 5-65.

Big Moose quadrangle, aeromagnetic: Balsley, J. R., 1-65.

Bolton Landing, Glens Falls, and Whithall quadrangle, aeromagnetic: Balsley, J. R., 2-65.

Canaan quadrangle: Holmes, G. W., 4-65.

Elizabethtown and Port Henry quadrangles, aeromagnetic: Balsley, J. R., 3-65.

Lowville quadrangle, aeromagnetic: Balsley, J. R., 4-65.

McKeever and Port Leyden quadrangles, aeromagnetic: Balsley, J. R., 5-65.

New London quadrangle, geologic: Goldsmith, Richard, 1-65.

Number Four quadrangle, aeromagnetic: Balsley, J. R., 6-65.

Nyack and part of White Plains quadrangles, aeromagnetic: Philbin, P. W., 1-64.

Old Forge and West Canada Lakes quadrangles, aeromagnetic: Balsley, J. R., 7-65.

Paradox Lake and Ticonderoga quadrangles, aeromagnetic: Balsley, J. R., 8-65.

Park Ridge quadrangle, aeromagnetic: Philbin, P. W., 2-64.

Pittsfield West quadrangle: Holmes, G. W., 3-65.

Raquette Lake quadrangle, aeromagnetic: Balsley, J. R., 9-65.

New York--Continued

Maps--Continued

Yonkers and Mount Vernon
quadrangles, aeromagnetic:
Philbin, P. W., 4-64.

Paleontology

Monmouth Group, Upper Cre-
taceous, correlation and
Foraminifera: Perlmutter,
N. M., 1-65.

Petrology

Adirondacks, phacolith, metaso-
matic origin: Engel, A. E. J.,
1-63.

Cortlandt complex, contact ef-
fects: Barker, Fred, 1-64.

Sedimentary petrology

Silurian, Clinton ironstones,
clay minerals: Schoen, Robert,
2-64.

Nickel

Idaho

General: Vhay, J. S., 1-64.

North America

Geophysical surveys

Explosion seismology: Pakiser,
L. C., 3-64.

Paleontology

Mollusca, marine genera:
Moore, E. J., 1-63.

Ostracoda, marine Triassic:
Sohn, I. G., 3-65.

Structural geology

Continental accretion: Engel,
A. E. J., 2-64.

Tectonic map: King, P. B., 3-64.

Western, Cordillera: King, P. B.,
2-64.

North Carolina

Areal geology

Ducktown, Isabella, and Per-
simmon Creek quadrangles:
Hernon, R. M., 1-64.

Linville Falls quadrangle: Reed,
J. C., Jr., 2-64.

Monroe area: Floyd, E. O.,
1-65.

Geochemistry

Cabarrus County, radioactive
sulfide deposit: Sundelius,
H. W., 1-64.

High Rock quadrangle: Stromquist,
A. A., 1-64.

Hydrogeology

Cape Hatteras area: Harris, W. H.,
1-64; Lloyd, O. B., Jr., 1-65.

Monroe area, ground-water:
Floyd, E. O., 1-65.

Municipal water supplies: Phibbs,
E. J., Jr., 1-65.

Surface waters, quality, 1961-62:
Phibbs, E. J., Jr., 1-64.

Stratigraphy

Lee's Creek, phosphate mine:
Gibson, T. G., 2-65.

Pungo River Formation: Kimrey,
J. O., 1-64.

Structural geology

Brevard zone, strike-slip move-
ment: Reed, J. C., Jr., 4-64.

North Dakota

Areal geology

Burleigh County: Randich, P. G.,
1-65.

North Dakota--Continued

Economic geology

Lignitic coal: Landis, E. R., 1-64.

Hydrogeology

Barnes County: Kelly, T. E.,
1-64.

Burleigh County, ground water:
Randich, P. G., 1-65.

Devils Lake area, ground water:
Paulson, Q. F., 3-64.

Grand River drainage basin:
Hembree, C. H., 2-64.

Ground water resources: Paulson,
Q. F., 2-64.

Sheyenne River, relation of dis-
charge to geology: Paulson,
Q. F., 1-64.

Souris River valley: Pettyjohn,
W. A., 1-65.

Spiritwood aquifer: Kelly, T. E.,
2-64.

Maps

Burleigh County, ground water:
Randich, P. G., 1-64.

Stratigraphy

Pierre Shale: Cobban, W. A.,
1-65.

Valley City, and Pembina
Mountain areas: Gill, J. R.,
1-65.

Quaternary, Ullsfjord, chrono-
logy: Holmes, G. W., 1-64.

Williston basin, paleotectonics:
Sandberg, C. A., 1-64.

Oceanography

Circulation, wind-driven

Polar ocean, ice and water:
Campbell, W. J., 1-64.

Ohio

Areal geology

Geauga County: Baker, J. A.,
1-64.

Geophysical surveys

Aeromagnetic: Beck, M. E., Jr.,
1-64.

Ground water, seismic: Watkins,
J. S., 2-65.

Little Miami River and Scioto
River valleys, seismic:
Watkins, J. S., 3-64.

Glacial geology

Northern part: Leighton, M. M.,
1-65.

Hydrogeology

Cambridge, floods, 1963: Cross,
W. P., 1-64.

Circleville, floods: Edelen,
G. W., Jr., 2-64.

Fairfield-New Baltimore area,
analog model study: Spieker,
A. M., 1-65.

Increased pumping effects:
Spieker, A. M., 2-64.

Floods, March 1964: Cross,
W. P., 2-64.

Geauga County, ground water:
Baker, J. A., 1-64.

Great Miami River Valley,
Hamilton area: Spieker,
A. M., 1-64.

Upper Ohio drainage area,
ground water distribution:
Jordan, P. R., 1-65.

Ohio--Continued

Maps

Chillicothe, floods: Edelen, G. W.,
Jr., 1-64.

Columbus-Dayton area, aeromag-
netic: Philbin, P. W., 1-65.

Findlay and vicinity, aeromag-
netic: Bromery, R. W., 2-65.

Mount Vernon, floods: Edelen,
G. W., Jr., 3-64.

Newark, floods: Edelen, G. W.,
Jr., 4-64.

Portsmouth quadrangle, geologic:
Sheppard, R. A., 1-64.

Zanesville, floods: Edelen, G. W.,
Jr., 5-64.

Oil shale

Colorado

Green River Formation, resources:
Donnell, J. R., 1-64.

Mineral fuels and associated re-
sources: Donnell, J. R., 2-64.

Uinta Basin, Green River For-
mation: Cashion, W. B., 3-64.

Utah

Uinta Basin: Bass, N. W., 2-64;
Cashion, W. B., 1-64, 3-64.

Oklahoma

Areal geology

Woodward County: Wood, P. R.,
2-65.

Hydrogeology

Arkansas River, alluvium: Tanaka,
H. H., 1-64.

Beckham County, ground water,
terrace deposits: Burton, L. C.,
1-65.

Cleveland and Oklahoma Counties,
ground water: Wood, P. R., 1-65.

Elk Creek Basin, ground water in
alluvium: Hollowell, J. R., 1-65.

Ground water levels in wells:
Wood, P. R., 3-64.

Harmon County: Steele, C. E., 1-64.

Surface water chemistry, 1961-62:
Cummings, T. R., 1-65.

Washita River area, ground water,
alluvial deposits: Hart, D. L.,
Jr., 1-65.

Woodward County, ground water:
Wood, P. R., 2-65.

Maps

Beckham County, geologic: Bur-
ton, L. C., 1-65.

Glencoe-Ripley area, aeromag-
netic: Andreasen, G. E., 1-64.

Hominy area, aeromagnetic:
Andreasen, G. E., 2-64.

Sedimentary petrology

Morrow Series, diagenesis:
Henbest, L. G., 1-65.

Stratigraphy

Beds overlying Springer(?) For-
mation: Hendricks, T. A.,
1-65.

Oregon

Areal geology

Central and northeastern, oro-
genic and plutonic activity:
Thayer, T. P., 3-64.

Fort Rock Basin, Lake County:
Hampton, E. R., 1-64.

Northwestern: Snively, P. D.,
Jr., 2-64.

Oregon--Continued

Areal geology--Continued

Tualatin Valley: Hart, D. H., 1-65.

Western Cascade Range, central and northern: Peck, D. L., 2-64.

Economic geology

Nickeliferous laterites: Hotz, P. E., 1-64.

Geophysical surveys

Northwestern, gravity and aeromagnetic: Bromery, R. W., 5-64.

Southwestern, gravity: Blank, H. R., Jr., 2-65.

United States Geological Survey gravity program: Oliver, H. W., 1-65.

Hydrogeology

Artificial recharge: Price, Don, 1-65.

Baker Valley, well data: Ducret, G. L., Jr., 1-65.

Burnt River valley, ground water reconnaissance: Price, Don, 1-64.

East Portland area: Foxworthy, B. L., 1-64.

Ground water: Hogenson, G. M., 1-65.

Fort Rock Basin, Lake County, occurrence and availability of ground water: Hampton, E. R., 1-64.

Pacific slope basins, surface-water records: Eisenhugh, H. P., 4-65.

Stream-temperature data compilation: Moore, A. M., 1-64.

Surface water, temperature studies: Moore, A. M., 2-64.

Tualatin Valley, ground water: Hart, D. H., 1-65.

Water supply: Phillips, K. N., 1-65.

Maps

Albany-Newport area, aeromagnetic and geologic: Bromery, R. W., 1-65.

Southwestern, gravity: Blank, H. R., Jr., 1-65.

Paleontology

Mollusca, Jurassic: Imlay, R. W., 3-64.

Pleistocene invertebrate fauna, southwestern: Addicott, W. O., 1-64.

Sedimentary petrology

Alsea River basin, sedimentation: Williams, R. C., 1-64.

Stratigraphy

Coal beds, correlation: Wayland, R. G., 1-65.

Tertiary, Yaquina Bay area: Snively, P. D., Jr., 1-64, 3-64.

Structural geology

Southwestern, ultramafic belts: Irwin, W. P., 1-64.

Ostracoda

Beyrichiacean: Martinsson, Anders, 1-65.

Ostracoda--Continued

Cytherelloidea

Paleotemperature, possible indicator: Sohn, I. G., 2-64.

Healdiace

Classification: Sohn, I. G., 1-65.

Investigations, 1964: Hazel, J. E., 1-65.

Pseudophanasymmetria

Classification: Sohn, I. G., 1-65.

Oxygen

Isotopes

Meteorites: Taylor, H. F., Jr., 1-64.

Pacific Islands

Paleontology

Brachiopoda, Eniwetok and Bikini drill holes: Cooper, G. A., 1-64.

Fossil collecting: Ladd, H. S., 1-65.

Mollusca, Tertiary: Ladd, H. S., 2-65.

Petrology

Basalt, northeastern: Engel, C. G., 1-63.

East Pacific Rise, igneous rocks: Engel, A. E. J., 3-64.

Sedimentary petrology

Mohole sediments, Guadalupe Site: Murata, K. J., 1-65.

Pakistan

Economic geology

Surveying and photogrammetry: Blake, Paul, 2-65.

Hydrogeology

Corrosion and encrustation mechanisms in tube wells: Clarke, F. E., 2-64.

Stratigraphy

Cambrian and Sabine series: Teichert, Curt, 1-64.

Paleoclimatology

Arid and hot climates of past: McKee, E. D., 2-64.

Biological and paleogeographic factors: Teichert, Curt, 2-64.

Paleomagnetism

General

Geological and geophysical applications: Cox, Allan, 1-65.

Paleontology

Technique

Acid-resistant fossils: Schopf, J. M., 1-65.

Blade mounting: Lewandowski, R. A., 1-65.

Collecting coal balls: Mamay, S. H., 1-65.

Collecting, metamorphic rocks: Neuman, R. B., 1-65.

Foraminifera: Todd, Ruth, 2-65.

Gastropoda and Pelecypoda: Sohl, N. F., 2-65.

Ignition to concentrate shelled organisms: Sachs, K. N., Jr., 2-64.

Ostracoda: Sohn, I. G., 4-65.

Paleozoic corals: Oliver, W. A., Jr., 1-65.

Photomicroscopy: Douglass, R. C., 2-65.

Plate preparation: Palmer, A. R., 2-65.

Paleontology--Continued

Technique--Continued

Trilobita, handbook: Rasetti, France, 1-65.

Whitening fossils: Kier, P. M., 1-65.

Palynology

Applied

Paleozoic: Kosanke, R. M., 1-64.

General

Review of book: Schopf, J. M., 2-65.

Petroleum exploration: Tschudy, R. H., 1-64.

Russian literature and application: Schopf, J. M., 4-64.

Pegmatites

Colorado

General: Adams, J. W., 4-64.

South Dakota

General: Norton, J. J., 1-64.

Utah

General: Adams, J. W., 5-64.

Pennsylvania

Areal geology

Evitts Creek and Pattersons

Creek quadrangles, stratigraphy and geologic history: deWitt, Wallace, Jr., 1-64.

Economic geology

Coal, Lawrence County: Van Lieu, J. A., 1-64.

Studies in Anthracite region: Trexler, J. P., 1-64.

Geophysical surveys

Aeromagnetic: Beck, M. E., Jr., 1-64.

Hydrogeology

Delaware River, water quality 1949-63: Keighton, W. B., 1-64.

Glacial deposits, ground water hardness: Seaber, P. R., 2-64.

New Oxford Formation: Wood, P. R., 2-64.

Maps

Belvidere quadrangle, aeromagnetic: Boynton, G. R., 2-65.

Radioactivity: Boynton, G. R., 1-65.

Bristol quadrangle, geologic: Owens, J. P., 1-64.

Ellsworth quadrangle, geologic: Berryhill, H. L., Jr., 1-64.

Southern Anthracite field, west-central, geologic: Wood, G. H., Jr., 1-65.

Trenton East quadrangle, geologic, pre-Quaternary: Owens, J. P., 2-64.

Washington East quadrangle, geologic: Swanson, V. E., 1-64.

Stratigraphy

Cambrian-Ordovician, Northampton and Bucks Counties, carbonate rocks: Drake, A. A., Jr., 1-65.

Permeability

Sandstone

Tensleep sandstone, Wyoming: Bredehoeft, J. D., 1-64.

Permian

General review: King, P. B., 1-65.

- Permo-Carboniferous
Glacial geology
South Africa, Australia, Antarctica, tillites: Krinsley, David, 1-65.
- Peru
Hydrogeology
Northern, Lambayeque Valley, data: Schoff, S. L., 2-64.
- Petroleum
Alaska
Mineral fuel resources: Grantz, Arthur, 2-64.
- Colorado
Resources: Bass, N. W., 1-64.
- Exploration
Microfossils: Schopf, J. M., 2-64.
- South Dakota
General: Sandberg, C. A., 3-64.
- United States
Resources: Hendricks, T. A., 1-63.
- Utah
General: Gere, W. C., 3-64.
- Phase equilibria
Arsenic-antimony: Skinner, B. J., 1-65.
- CaSO₄-NaCl-H₂O: Zen, E-an, 1-65.
- Chalcocite
High-pressure polymorph: Skinner, B. J., 1-64.
- Chaleopyrite+sulfur=pyrite+bornite, 350 to 500°C: Barton, P. B., Jr., 1-64.
- Copper sulfides
Reactions between 25° and 200°C: Roseboom, E. H., Jr., 1-65.
- Cu-Ag-S: Skinner, B. J., 2-65.
- H₂O-Na₂O-SiO₂: Rowe, J. J., 2-65.
- K₂SO₄-CaSO₄: Rowe, J. J., 3-65.
- Melanophlogite
Cubic polymorph of silica: Skinner, B. J., 1-63.
- Rock-forming oxides: Morey, G. W., 1-64.
- Stishovite
Inversion to silica glass: Skinner, B. J., 2-63.
- Philippines
Hydrogeology
Poro Point and vicinity, reconnaissance: Worts, G. F., Jr., 1-64.
- Phosphate
California
Occurrence: Gowen, H. D., 1-64.
- Field and laboratory determination
Natural water: Fishman, M. J., 1-65.
- Florida
Precipitation and recycling: Altschuler, Z. S., 1-65.
- Idaho
General: Gere, W. C., 1-64.
- Occurrence
Paleolatitudinal and paleogeographic distribution: Sheldon, R. P., 2-64.
- Turkey
Exploration: Sheldon, R. P., 1-64.
- Utah
General: Gere, W. C., 2-64.
- Photography
Methods
Contouring, evergreen forest areas: Halliday, James, 1-64.
- Photography--Continued
Methods--Continued
Reflection-transmission illumination: Krier, E. P., 1-65.
- Platinum
Geochemistry: Wright, T. L., 1-65.
- Puerto Rico
Areal geology
Ciales quadrangle: Berryhill, H. L., Jr., 1-65.
- Geologic history: Briggs, R. P., 2-65.
- Geologic mapping, progress to date: Briggs, R. P., 1-65.
- Geomorphology
Mogotes: Monroe, W. H., 2-64.
- Geophysical surveys
Mayagüez area, gravity: Bromery, R. W., 1-64.
- Mayagüez, magnetic properties, serpentine: Cox, Allan, 3-64.
- Southwest, magnetic and aeromagnetic: Griscom, Andrew, 1-64.
- Heat flow
Mayagüez area: Diment, W. H., 1-64, 2-64.
- Hydrogeology
Public water supply, chemical quality: Durfor, C. N., 1-65.
- Water resources, progress report: Bogart, D. B., 1-64.
- Water wells: Ward, P. E., 1-64.
- Maps
Ciales quadrangle, geologic: Berryhill, H. L., Jr., 1-65.
- Hydrogeologic: Briggs, R. P., 3-65.
- Toa Alta, Toa Baja, and Dorado, floods: Lopez, M. A., 1-64.
- Paleontology
Lepidocyclina (Eulepidina) undosa Cushman: Sachs, K. N., Jr., 1-64.
- Petrography
Mayagüez area, serpentinite: Mattson, P. H., 1-64.
- Serpentine, electrical properties: Zablocki, C. J., 1-64.
- Serpentinite, strength and elastic moduli: Robertson, E. C., 2-64.
- Volcanic rocks, chemical variations: Mattson, P. H., 1-65.
- Volcanology
Trends of Cretaceous: Nelson, A. E., 1-65.
- Quaternary
Environment
Reconstruction, palynology: Leopold, E. B., 1-64.
- Radioactivity
Instruments
Oceanographic studies: Bunker, C. M., 1-64.
- Radioactivity exploration
ARMS program: Guillou, R. B., 1-64.
- Geologic mapping: Pitkin, J. A., 1-64.
- Hydrothermally altered rock
Gamma-ray spectrometer study: Moxham, R. M., 2-65.
- Radioactivity exploration--Continued
Terrestrial gamma radiation
Aerial observations: Moxham, R. M., 2-64.
- Radon
Migration in ground: Tanner, A. B., 2-64.
- Rare earths
Colorado
General: Adams, J. W., 3-64.
- New Mexico
General: Adams, J. W., 1-65.
- Refractory minerals
Apparatus
Decomposition: May, Irving, 2-65.
- Solution
High temperatures and pressure: May, Irving, 1-65.
- Rhenium
New Mexico
General: King, R. U., 2-65.
- Rhode Island
Areal geology
Tiverton quadrangle: Pollock, S. J., 1-64.
- Hydrogeology
Block Island, ground-water: Hansen, A. J., Jr., 1-64.
- Upper Pawcatuck River basin, ground water: Allen, W. B., 1-65.
- Maps
Kingston quadrangle, geologic: Moore, G. E., Jr., 1-64.
- Prudence Island and Newport quadrangles, ground water: Schiner, G. R., 1-65.
- Tiverton and Sakonnet Point quadrangles, ground water: Schiner, G. R., 2-65.
- Petrology
Ashaway and Voluntown quadrangles: Feininger, Tomas, 1-65.
- Rhodesia
Hydrogeology
Southern, ground water provinces: Dennis, P. E., 2-64.
- Rivers
Channel geometry: Emmett, W. W., 1-64.
- Contaminants, soluble
Time of travel: Buchanan, T. J., 1-64.
- Drainage patterns
Distribution of branches: Giusti, E. V., 1-65.
- Education: Oltman, R. E., 1-65.
- Hydraulics
Flow resistance, alluvial channels: Simons, D. B., 2-65.
- Stream order
New system of numbering: Scheidegger, A. E., 2-65.
- Transient flow
Mathematical model: Lai, Chintu, 1-65.
- Rocky Mountains
Economic geology
Coal: Averitt, Paul, 1-65.

- Salt
Virginia
Occurrence and origin:
Withington, C. F., 2-65.
- Sand and gravel
Colorado
General: Carter, W. D., 1-64.
New Mexico
General: Carter, W. D., 1-65.
- Sandstone
Chemical composition: Petti-
john, F. J., 1-64.
- Saskatchewan
Structural geology
Williston basin, paleotectonics:
Sandberg, C. A., 1-64.
- Sedimentary petrology
Technique
Unconsolidated sediments,
collecting: Sohl, N. F., 1-65.
- Sedimentary rocks
Ironstones
Silurian, Clinton Group:
Schoen, Robert, 1-64.
Marine evaporites: Stewart,
F. H., 1-63.
- Shale
Pierre Shale, mineralogy:
Schultz, L. G., 1-64.
- Sedimentary structures
Inorganic: McKee, E. D., 1-64.
Ripples and dunes
Bedload equation: Simons,
D. B., 1-65.
- Varves
Theory of formation: Scheideg-
ger, A. E., 1-65.
- Sedimentation
Experimental studies
Channel erosion and transport:
Nordin, C. F., Jr., 4-64.
Flume, alluvial channel data:
Guy, H. P., 1-65.
Sorting processes and flow
separation: Jopling, A. V.,
1-64.
Texas, Rio Grande River:
Nordin, C. F., Jr., 3-64.
Flow resistance and transport,
Rio Grande area: Nordin,
C. F., Jr., 1-64.
- Land subsidence
Alluvial fans, California:
Bull, W. B., 2-64.
- Rates
Computation of bed-material
discharge: Nordin, C. F., Jr.,
2-64.
Transport monitoring system:
Guy, H. P., 1-64.
- Scour and fill
Sand-bed streams: Colby, B. R.,
3-64.
- Size analysis
Sand with friable fragments:
Bull, W. B., 4-64.
- Stream transport
Effect of water temperature:
Colby, B. R., 1-65.
Rio Grande, New Mexico: Nordin,
C. F., Jr., 1-65.
Sand-bed streams: Colby, B. R.,
1-64.
- Sedimentation--Continued
Stream transport--Continued
Suspended sediment: hypercon-
centrations: Beverage, J. P.,
1-64.
Total bed material: Bishop,
A. A., 1-65.
- Sediments
Base-exchange capacity
Modern stream sediments:
Kennedy, V. C., 2-64.
Carbonate mud
Experimental consolidation:
Robertson, E. C., 1-64.
- Clay
Compaction process: Meade,
R. H., 1-64.
- Fluvial
Monitoring system: Guy, H. P.,
1-64.
- Ion-exchange capacity
Mohole, Guadalupe site: Car-
roll, Dorothy, 1-64.
- Origin
Tectonic and environmental
factors: Feray, D. E., 1-65.
- Sand
Size analysis: Bull, W. B., 4-64.
Undisturbed, consolidation test:
Bull, W. B., 1-64.
- Technique
Sample dispersion, ultrasonic:
Moston, R. P., 1-64.
- Unconsolidated
Sample collection, freezing
method: Kennedy, V. C., 1-64.
- Seismic exploration
Attenuation measurements in field:
DeBreaecker, J. C., 1-65.
- Methods
Surveying with firecrackers:
Criner, J. H., 1-65.
- Technique
Digital processing: Ryall, Alan,
1-62.
- Selenium
Nevada
General: Davidson, D. F., 1-64.
New Mexico
General: Davidson, D. F., 1-65.
- Serpentine
Puerto Rico
Strength and elastic moduli:
Robertson, E. C., 2-64.
- Silica
Idaho
General: Carter, W. D., 2-64.
- Silver
Black calcite
New source: Hewett, D. F., 1-65.
- Idaho
General: Kiilsgaard, T. H., 3-64.
- Nevada
General: Kleinhampl, F. J., 1-64.
- Resources
Review: Kiilsgaard, T. H., 1-64.
- South Dakota
Resources: Bergendahl, M. H.,
6-64.
- Utah
General: Kiilsgaard, T. H., 4-64.
- Sodium
Analysis
Closed-basin waters: Jones,
B. F., 1-65; Truesdell,
A. H., 1-65.
- Soils
Engineering properties
Bibliography: Johnson, A. I.,
1-63.
Moisture stress measurement:
McQueen, I. S., 1-64.
Experimental studies
Saturation measurement: Shown,
L. M., 1-64.
- South Africa
Paleontology
Shrew, Pleistocene: Repenning,
C. A., 1-65.
- South Carolina
Areal geology
Piedmont: Overstreet, W. C.,
2-65.
- Hydrogeology
Central Savannah River area,
artesian aquifers: Siple,
G. E., 1-64.
Savannah area, saline-water
intrusion: McCollum, M. J.,
1-64, 2-64.
Savannah River Plant: Marine,
I. W., 1-64.
Radioactive waste: Siple,
G. E., 2-64.
- Maps
Piedmont, geologic: Overstreet,
W. C., 1-65.
Savannah River Plant area,
aeromagnetic: Petty, A. J.,
1-65.
- Stratigraphy
Subsurface, coastal: Herrick,
S. M., 2-65.
- South Dakota
Areal geology
Black Hills uplift, northern and
western flanks: Robinson,
C. S., 2-64.
Edgemont quadrangle: Ryan,
J. D., 1-64.
General: Denson, N. M., 1-64.
- Economic geology
Antimony: Dasch, M. D., 1-64.
Clays: Patterson, S. H., 4-64.
Coal: Landis, E. R., 3-64.
Gold and silver: Bergendahl,
M. H., 6-64.
Gypsum and anhydrite: Cox,
E. J., 1-64.
Industrial minerals: Bowles,
C. G., 1-64.
Mineral industry: Denson,
N. M., 2-64.
Mineral resources, general:
Denson, N. M., 3-64.
Molybdenum: King, R. U., 2-64.
Pegmatites: Norton, J. J., 1-64.
Petroleum and natural gas:
Sandberg, C. A., 3-64.
Resources, general: Denson,
N. M., 4-64.
Sand and gravel: Bruce, R. L.,
1-64.

- South Dakota--Continued
 Economic geology--Continued
 Thorium and the rare earths:
 Adams, J. W., 1-64.
 Tin: Ratté, J. C., 2-64.
 Tungsten, Black Hills area:
 Hobbs, S. W., 2-64.
 Uranium: Gott, G. B., 3-64.
 Cave Hills area: Pippingos,
 G. N., 1-65.
 Vanadium: Fischer, R. P.,
 3-64.
 Geomorphology
 Landslides, Fort Randall Re-
 servoir: Erskine, C. F.,
 1-65.
 Hydrogeology
 Grand River drainage basin:
 Hembree, C. H., 2-64.
 Sanborn County, ground water:
 Steece, F. V., 1-64.
 Skunk Creek-Lake Madison
 basin, glacial drift: Ellis,
 M. J., 1-64.
 Skunk Creek-Lake Madison
 drainage basin: Adolphson,
 D. G., 1-64.
 Standing Rock Indian Reserva-
 tion, ground water: Ellis,
 M. J., 1-65.
 Water resources: Powell, J. E.,
 1-64; Denson, N. M., 3-64.
 Structural geology
 Black Hills: Gott, G. B., 1-64.
 Spectroscopy
 Absorption
 Rare-earth minerals: Adams,
 J. W., 2-65.
 Electron probe
 Minerals: Adler, Isidore, 1-63,
 1-64.
 Stratigraphy
 Nomenclature
 Changes, United States Geol-
 ogical Survey: Cohee, G. V.,
 1-65.
 Gila Conglomerate, use of
 term: Heindl, L. A., 2-64.
 Streams
 Suspended sediment
 Hyperconcentrations: Beverage,
 J. P., 1-64.
 Stone
 Utah
 General: Hansen, W. R., 1-64.
 Strontium
 Analysis
 Distribution coefficients, one
 and two competing cation
 solutions: Wahlberg, J. S.,
 1-65.
 Isotopes
 Basalt: Tilton, G. R., 1-64.
 Radiogenic
 Volcanic lavas: Hedge, C. E.,
 1-64.
 Submarine geology
 Photography and sampling:
 Emery, K. O., 1-65.
 Subsidence
 Well casing protrusion: Poland,
 J. F., 1-64.
 Sulfur
 New Mexico
 General: Broderick, G. N., 1-65.
 Sulfur--Continued
 Utah
 General: Mount, Priscilla, 2-64.
 United States Geological Survey
 Annual report of Director:
 Nolan, T. B., 1-64.
 Location of field projects: U.S.
 Geological Survey, Geologic
 Division, 1-64.
 Tantalum
 Distribution
 Southern California batholith:
 Gottfried, David, 1-65.
 Tektites
 Brunei: Wilford, G. E., 1-64.
 Composition
 Cesium, rubidium, and lithium:
 Ansell, C. S., 1-63.
 Ferrous iron, determination:
 Carron, M. K., 1-63.
 Major constituents, emission
 spectroscopy: Ansell, C. S.,
 2-63.
 Minor elements: Ansell, C. S.,
 1-64.
 Nickel-iron spherules, south-
 east Asia: Chao, E. C. T., 1-64.
 Electrical resistivity and viscos-
 ity: Hoyte, A., 1-65.
 Experimental studies
 Silicate melts: Walter, L. S.,
 1-64.
 Magnetic properties
 Nickel-iron spherules: Senftle,
 F. E., 1-63.
 Moldavites
 Aerodynamically modified:
 Chao, E. C. T., 2-64.
 Physical properties and chemical
 composition, Australasian:
 Cuttitta, Frank, 1-63, 1-64.
 Submicroscopic spherules and
 color: Thorpe, A. N., 1-63.
 X-ray fluorescence analysis:
 Rose, H. J., Jr., 1-63.
 Tellurium
 Nevada
 General: Davidson, D. F., 1-64.
 New Mexico
 General: Davidson, D. F., 1-65.
 Tennessee
 Areal geology
 Ducktown, Isabella, and Per-
 simmon Creek quadrangles:
 Hernon, R. M., 1-64.
 Elkmont quadrangle: McMaster,
 W. M., 1-65.
 Middlesboro South quadrangle:
 Englund, K. J., 1-64.
 Economic geology
 Zinc, Mascot-Jefferson City
 mining district: Wedow,
 Helmuth, Jr., 2-65.
 Geophysical surveys
 Regional implications: Watkins,
 J. S., 1-64.
 Hydrogeology
 Dickson County, well water sup-
 ply: Bingham, R. H., 1-65.
 Highland Rim, western: Marcher,
 M. V., 1-64.
 Lewis County, well water supply:
 Bingham, R. H., 1-64.
 Lynn Grove quadrangle, ground wa-
 ter, availability: Davis, R. W., 2-64.
 Tennessee--Continued
 Hydrogeology--Continued
 Memphis area, aquifer sys-
 tems: Criner, J. H., 1-64.
 Lichterman well field: Ny-
 man, D. J., 1-65.
 Mill Creek area, floods:
 Randolph, W. J., 1-64.
 Montgomery County, availa-
 bility of water for industry:
 Perry, W. J., 1-65.
 Wayne County, well water sup-
 ply: Bingham, R. H., 2-65.
 Western Highland Rim, ground
 water, availability: Moore,
 G. K., 1-65.
 Maps
 Elkmont quadrangle, geologic:
 McMaster, W. M., 1-65.
 Hazel quadrangle, ground wa-
 ter: MacCary, L. M., 3-64.
 Petroleum quadrangle, geol-
 ogic: Myers, W. B., 1-64.
 Salem quadrangle, geologic:
 McMaster, W. M., 2-65.
 Paleontology
 Gastropoda, Late Cretaceous:
 Sohl, N. F., 2-64.
 Structural geology
 Flynn Creek structure: Roddy,
 D. J., 1-63.
 Texas
 Areal geology
 Hardin County: Baker, E. T.,
 Jr., 1-64.
 Sierra Madera: Shoemaker,
 E. M., 1-63.
 Wayland quadrangle: Myers,
 D. A., 1-65.
 Engineering geology
 Houston-Galveston Bay area,
 clay minerals: Corliss,
 J. B., 1-64.
 Geochemistry
 Llano uplift, lead isotopes in
 microcline: Zartman, R. E.,
 1-64.
 Geophysical surveys
 Carlsbad area, radioactivity:
 MacKallor, J. A., 1-65.
 Hydrogeology
 Bee County, ground water re-
 sources: Myers, B. M.,
 1-65.
 Bibliography: Baker, R. C., 1-
 64.
 Brazos River, surface waters,
 chemical quality: Irelan,
 Burdge, 1-64.
 Caldwell County, ground water:
 Follett, C. R., 1-65.
 Camp Franklin, Morris, and
 Titus Counties, ground wa-
 ter: Broom, M. E., 1-64.
 Cibolo Creek, base-flow
 studies: Holland, P. H., 3-65.
 Colorado River basin, Muke-
 water Creek: Sauer, S. P.,
 2-65.
 Comal County, base-flow
 studies: Holland, P. H., 1-65.
 Croton and Salt Croton Creek
 basins, natural sources of
 salinity: Baker, R. C., 3-64.

Texas--Continued

Hydrogeology--Continued

- Dallas area, floods: Ruggles, F. H., Jr., 1-64.
- Deep Creek, Colorado River basin, 1951-61: Mills, M.B., 1-64.
- Gaines County, ground water: Rettman, P. L., 1-65.
- Guadalupe, San Antonio and Nueces River basins, ground-water resources: Alexander, W. H., Jr., 1-64.
- Hardin County, ground water resources: Baker, E. T., Jr., 1-64.
- Honey Creek basin, small watersheds: Gilbert, C. R., 1-64.
- Houston County, ground water: Tarver, G. R., 1-65.
- Houston district, analog model study: Wood, L. A., 1-65.
- Hubbard Creek watershed, quality of surface waters: Hembree, C. H., 1-64.
- Jackson County, ground-water: Baker, E. T., Jr., 1-65.
- Lake Colorado City, thermal properties: Harbeck, G. E., Jr., 2-64.
- Lampasas River, base-flow studies, 1963: Mills, W. B., 1-65.
- Llano River, base-flow studies: Holland, P. H., 2-65.
- Lower Rio Grande Valley area, ground water: Baker, R. C., 2-64.
- Nueces River, base-flow studies, 1964: Reeves, W. E., 1-65.
- Water delivery: Sauer, S. P., 1-65.
- Pecos River, water delivery study: Grozier, R. U., 2-65.
- Sabine River, base-flow study, 1963: Sauer, S. P., 1-64.
- San Antonio area, ground-water discharge: Garza, Sergio, 1-64.
- Water in Edwards limestone: Garza, Sergio, 1-65.
- San Gabriel River area, base-flow studies, 1964: Leifeste, D. K., 1-65.
- Surface water, chemical composition, 1962: Hughes, L. S., 1-65.
- Western, surface water, growth of salt cedar: Grozier, R. U., 1-65.

Maps

- Bakers Crossing quadrangle, geologic: Freeman, V. L., 1-65.
- Carlsbad area, radioactivity: MacKallor, J. A., 2-65.
- Langtry quadrangle, geologic: Freeman, V. L., 1-64.
- Oil and gas fields, pipelines, and exposed basement rocks: Vlissides, S. D., 1-64.
- Shumla quadrangle, geologic: Freeman, V. L., 2-64.
- Tularosa Basin, ground-water: Herrick, E. H., 1-65.

Texas--Continued

Paleontology

- Brachiopoda, Permian, Glass Mountains: Cooper, G. A., 2-64.
- Sedimentary petrology
- Rio Grande, sedimentation: Nordin, C. F., Jr., 3-64.
- Thorium
- Abundance
- Hawaii and Japan: Tatsumoto, M., 2-65.
- Colorado
- General: Staatz, M. H., 3-64.
- Resources
- General review: Olson, J. C., 1-64.
- South Dakota
- Resources: Adams, J. W., 1-64.
- Utah
- Resources: Adams, J. W., 1-64.
- Tin
- Ion-exchange separation from silicate rocks: Huffman, Claude, Jr., 1-64.
- New Mexico
- General: Sainsbury, C. L., 2-65.
- South Dakota
- General: Ratté, J. C., 2-64.
- Titanium
- Utah
- Resources: Adams, J. W., 2-64.
- Trace-element analyses
- Black shale
- Spectrographic, Pennsylvanian: Vine, J. D., 2-65.
- Trilobita
- Cambrian
- Evolution, North America: Palmer, A. R., 1-65.
- General
- Handbook: Rasetti, France, 1-65.
- Tungsten
- Colorado
- General: Hobbs, S. W., 1-64.
- Idaho
- General: Hobbs, S. W., 3-64.
- New Mexico
- General: Hobbs, S. W., 1-65.
- South Dakota
- Black Hills area, Hobbs, S. W., 2-64.
- Utah
- General: Lemmon, D. M., 1-64.
- Turkey
- Economic geology
- Phosphate, exploration: Sheldon, R. P., 1-64.
- United States
- Economic geology
- Barite, general: Brobst, D. A., 1-65.
- Bauxite, southeastern: Overstreet, E. F., 1-64.
- Energy resources: McKelvey, V. E., 1-65.
- Manganese: Hewett, D. F., 1-64.
- Petroleum, resources: Hendricks, T. A., 1-63.
- Uranium, deposits in sandstone, general: Finch, W. I., 1-64.
- Structural control: Osterwald, F. W., 1-65.

United States--Continued

General

- Environment and man in arid regions: Malde, H. E., 2-64.
- Geography: Gerlach, A. C., 1-64.
- Nuclear test sites, bibliography: Snyder, R. P., 1-63, 2-63.
- Survey instruments and methods: Blake, Paul, 1-65.
- Geochemistry
- Eastern interior region, coal beds: Zubovic, Peter, 1-64.
- Western interior, shales, composition: Tourtelot, H. A., 2-64.
- Williston basin, uranium-bearing lignite and carbonaceous shale: Denson, N. M., 1-65.
- Geomorphology
- Great Basin, Pleistocene lakes: Snyder, C. T., 1-64.
- Geophysical surveys
- Columbia River Basin, electrical: Anderson, L. A., 1-65.
- Denver, Colorado to Washington, D. C.: Zietz, Isidore, 2-65.
- Southwestern, gravity and aeromagnetic: Mabey, D. R., 1-65.
- Transcontinental strip east of Rocky Mountains: Zietz, Isidore, 1-65.
- Washington, Idaho, Montana, and Wyoming, gravity: Hill, D. P., 1-65.
- Wisconsin to Colorado, seismic: Roller, J. C., 1-65.
- Hydrogeology
- Aquifers, permeability and storage, heterogeneous: Rasmussen, W. C., 1-64.
- Atlantic Coastal Plain, southern, limestone terrane: Stringfield, V. T., 2-64.
- Atlantic Slope and eastern Gulf of Mexico Basins, surface waters: Love, S. K., 6-64, 1-65.
- Colorado and Columbia River Basins, surface waters: Love, S. K., 1-64.
- Colorado River basin, surface-water records index: Eisenhuth, H. P., 2-65.
- Surface waters: Santos, J. F., 1-65.
- Cumberland and Tennessee River basins, surface water: United States Geological Survey, 5-64.
- Developed and potential water-power: Young, L. L., 1-64.
- East of Mississippi River, resources: Lohr, E. W., 1-64.
- Eastern, James River to Savannah River, surface waters: Hendricks, E. L., 4-64.
- New York to York River, surface waters: Hendricks, E. L., 3-64.
- Surface water: United States Geological Survey, 3-65.

United States--Continued

Hydrogeology--Continued

Floods, 1959: U.S. Geological Survey, 13-64.

1960: Rostveld, J. O., 1-65.

Ground-water occurrence:

Meinzer, O. E., 2-65.

Hudson Bay and upper Mississippi River basins, surface water: U.S. Geological Survey, 7-64.

Hudson Bay, Upper Mississippi River and Missouri River Basins, surface waters: Love, S. K., 2-64.

Lower Mississippi River and western Gulf of Mexico Basins, surface waters: Love, S. K., 3-64, 3-65.

Maine to Connecticut, surface waters: Hendricks, E. L., 2-64.

Mapping: Wiesnet, D. R., 1-64.

Mississippi River basin, lower, surface water: U.S. Geological Survey, 9-64.

Surface-water records: Eisenhuth, H. P., 1-65.

Missouri River basin: Interior Missouri Basin Field Committee, 1-64, 1-65; U.S. Department of the Interior, 1-64; U.S. Geological Survey, 10-64.

Missouri River basin above Sioux City, surface water: U.S. Geological Survey, 8-64.

Missouri River basin, ground water: LaRocque, G. A., Jr., 1-65.

Surface waters: Hendricks, E. L., 6-64.

North Atlantic slope basins, Maine to Connecticut, floods: Green, A. R., 1-64.

Surface water: U.S. Geological Survey, 3-64, 4-64.

North-central, ground-water levels, 1957-61: Hackett, O. M., 2-64.

Northeastern, ground-water levels, 1958-62: U.S. Geological Survey, 4-65.

Northeastern states, ground-water levels, 1958-62: Hackett, O. M., 1-65.

Occurrence of dissolved solids in surface waters: Langbein, W. B., 2-64.

Ohio and St. Lawrence Rivers basins, surface waters: Love, S. K., 4-64, 2-65.

Surface-water quality: U.S. Geological Survey, 5-65.

Pacific slope, California, surface waters: Hendricks, E. L., 7-64.

Pacific slope, Washington: Hendricks, E. L., 8-64.

Potomac River: Davies, W. E., 3-64.

Public water supplies, 1962: Durfor, C. N., 1-64, 2-64.

United States--Continued

Hydrogeology--Continued

Public water supply, chemical quality: Durfor, C. N., 1-65.

St. Lawrence River basin, floods: Wiitala, S. W., 1-64.

Surface waters: Hendricks, E. L., 5-64; U.S. Geological Survey, 6-64.

Snake River Basin, surface-water quality: Laird, L. B., 1-64.

Surface-water records index: Eisenhuth, H. P., 3-65.

Southwest, effect on water resources: Gatewood, J. S., 1-64.

Surface waters, quality: Love, S. K., 5-64.

Susquehanna River basin, ground water: Seaber, P. R., 1-64.

West of Mississippi River, water supply: Lohr, E. W., 2-64.

Western, calcium, sodium, sulfate, and chloride in stream water: Feth, J. H., 1-65.

Western States, 1964: Rantz, S. E., 2-65.

Phreatophyte research: Robinson, T. W., 1-64. Saltcedar: Robinson, T. W., 1-65.

Maps

Annual runoff and productive aquifers: McGuinness, C. L., 2-64.

Bouguer gravity anomaly: Woollard, G. P., 1-64.

Mineral resources, barite occurrences: Brobst, D. A., 1-65.

Oil and gas fields: Vlissides, S. D., 2-64.

37th parallel, geologic and crustal cross section: Hamilton, Warren, 4-65.

Paleoclimatology

Western, paleowinds: Poole, F. G., 1-64.

Paleontology

Cephalopoda, Pierre Shale: Cobban, W. A., 1-64.

Ostracoda, Cretaceous, southeastern: Swain, F. M., 1-64.

Palynomorphs, Upper Cretaceous and Tertiary, Mississippi embayment: Tschudy, R. H., 2-65.

Pelecypoda, Cretaceous, oysters: Sohl, N. F., 4-64.

Trilobites, Cambrian, Great Basin: Palmer, A. R., 3-65.

Stratigraphy

Ordovician, Basin Ranges: Ross, R. J., Jr., 2-64.

Paleozoic, terminology: Keroher, G. C., 1-64.

Structural geology

Appalachians, Clinchport thrust fault: Harris, L. D., 1-65.

United States--Continued

Structural geology--Continued

Crust, western: Pakiser, L. C., 1-65.

Transcontinental crust and upper-mantle structure:

Pakiser, L. C., 2-65.

Southeastern: King, P. B., 1-64.

Western, volcanism, tectonism, and plutonism: Gilluly, James, 1-65.

Uranium

Abundance

Hawaii and Japan: Tatsumoto, M., 2-65.

Colorado

General: Butler, A. P., Jr., 2-64.

Epigenetic deposits

Sandstones: Finch, W. I., 1-64.

Exploration

Radioelement dispersion:

Moxham, R. M., 1-64.

Geochemistry.

Isotopic fractionation, fluvial sediments: Rosholt, J. N., 1-63.

Uranyl ion coordination: Evans, H. T., Jr., 1-63.

Idaho

General: Armstrong, F. C., 1-64.

Isotopes

Soil profiles: Rosholt, J. N., 2-65.

Nevada

General: Butler, A. P., Jr., 1-64.

New Mexico

General: Hilpert, L. S., 1-65.

Occurrence

Ore rolls: Shawe, D. R., 1-65.

South Dakota

General: Gott, G. B., 3-64.

Utah

General: Hilpert, L. S., 7-64.

Utah

Absolute age

Carmel Formation, K-Ar and Rb-Sr ages, biotite: Marvin, R. F., 1-65.

La Sal Mountains rocks: Stern, T. W., 2-65.

Areal geology

Abajo Mountains area: Witkind, I. J., 1-64.

Cedar Mesa-Boundary Butte area, sedimentary rocks, Pennsylvanian to Recent: O'Sullivan, R. B., 1-65.

Elk Ridge area: Lewis, R. Q., Sr., 1-65.

Flaming Gorge area: Hansen, W. R., 1-65.

General: Hilpert, L. S., 2-64.

Green River Formation, associated Eocene rocks: Bradley, W. H., 1-64.

Jordan Valley: Marine, I. W., 2-64.

Moab quadrangle: Williams, P. L., 1-64.

Paradise quadrangle: Mullens, T. E., 2-64.

Utah--Continued

- Areal geology--Continued
Salt Lake County, general:
Crittenden, M. D., Jr., 2-64.
Thomas and Dugway Ranges:
Staatz, M. H., 4-64.
Thorne Cave, northeastern:
Malde, H. E., 3-64.
White Canyon area: Thaden,
R. E., 1-64.
Economic geology
Alunite: Parker, R. L., 3-64.
Antimony: Dasch, M. D., 2-64.
Barite: Brobst, D. A., 3-64.
Beryllium: Griffiths, W. R., 2-64.
Clays: Patterson, S. H., 3-64.
Coal, resources: Averitt, Paul,
1-64.
 Uinta Basin: Schell, E. M., 1-64.
Copper: Roberts, R. J., 2-64.
Fluorine: Dasch, M. D., 5-64.
Gem materials: Dasch, M. D.,
6-64.
Gold: Bergendahl, M. H., 5-64.
Gypsum and anhydrite: Withing-
ton, C. F., 3-64.
Iron: Reeves, R. G., 3-64.
Lead, zinc, and silver: Kiils-
gaard, T. H., 4-64.
Lightweight aggregate: Van
Horn, Richard, 1-64.
Limestone: Morris, H. T., 1-64.
Manganese: Crittenden, M. D.,
Jr., 5-64.
Mercury: Hilpert, L. S., 4-64.
Metallic mineral resources:
Hilpert, L. S., 3-64.
Mineral fuels and associated
resources: Cashion, W. B.,
2-64; Hilpert, L. S., 5-64.
Mineral resources, general:
Hilpert, L. S., 1-64, 6-64,
8-64; Weeks, R. A., 1-64.
Molybdenum: King, R. U., 3-64.
Natural gas, San Arroyo and
East Canyon fields: Horton,
G. W., 2-64.
Oil shale, Uinta Basin: Bass,
N. W., 2-64; Cashion, W. B.,
1-64, 3-64.
Ore deposits, White Canyon
area: Thaden, R. E., 1-64.
Pegmatite minerals: Adams,
J. W., 5-64.
Petroleum and natural gas:
Gere, W. C., 3-64.
 Uinta Basin and Uncompahgre
 Uplift: Horton, G. W., 1-64.
Phosphate: Gere, W. C., 2-64.
Refractory minerals: Ketner,
K. B., 1-64.
Salines: Hite, R. J., 1-64.
Sand and gravel: Van Horn,
Richard, 2-64.
Silica: Ketner, K. B., 2-64.
Stone: Hansen, W. R., 1-64.
Sulfur: Mount, Priscilla, 2-64.
Thorium and the rare earths:
Adams, J. W., 1-64.
Titanium: Adams, J. W., 2-64.
Tungsten: Lemmon, D. M., 1-64.
Uranium: Hilpert, L. S., 7-64.
 Elk Ridge area: Lewis, R. Q.,
 Sr., 1-65.

Utah--Continued

- Economic geology--Continued
Vanadium: Fischer, R. P., 6-64.
Engineering geology
Sunnydale, mine bumps:
Osterwald, F. W., 1-64.
Geochemistry
Copper, alluvium and caliche:
Erickson, R. L., 1-64.
Spor Mountain, lithium in
rhyolitic tuff: Shawe, D. R.,
1-64.
Uraniferous phosphatic lake
beds, Eocene: Love, J. D.,
1-64.
Yellow Cat area, Grand Coun-
ty, rocks, related soils,
and vegetation: Cannon, H.
L., 1-64.
Geomorphology
Topography: Tooker, E. W.,
2-64.
Geophysical surveys
Central and southern Wasatch
Front, gravity: Cook, K. L.,
1-64.
Gravity, United States Geol-
ogical Survey program:
Case, J. E., 3-65.
Salt Valley anticline, geo-
thermal gradient: Spicer,
H. C., 2-64.
Seismic, coal mine bumps:
Dunrud, C. R., 1-64.
Glacial geology
Little Cottonwood and Bells
Canyons, Wasatch Moun-
tains: Richmond, G. M., 1-64.
Hydrogeology
Great Salt Lake, brines: Hahl,
D. C., 1-64.
Ground-water conditions,
Spring 1964: Arnov, Ted,
1-64.
Jordan Valley: Bradley, Ed-
ward, 1-64; Marine, I. W.,
2-64.
Jordan Valley, ground water:
Arnov, Ted, 1-65.
Navajo and Hopi Indian Reser-
vations: Cooley, M. E., 1-
64, 2-64.
Navajo Lake: Wilson, M. T., 2-64.
Northern, Utah Valley, ground
water, 1948-63: Cordova,
R. M., 1-65.
Radon in ground water: Tanner,
A. B., 1-64.
Sevier Desert, ground water:
Mower, R. W., 1-64.
Tooele Valley, ground water:
Gates, J. W., 1-65.
Upper Sevier River basin, data:
Carpenter, C. H., 1-64.
Water levels, April 1965:
Butler, R. G., 1-65.
Water resources, general:
Hilpert, L. S., 1-64; Wilson,
M. T., 1-64.
Waterpower, general: Johnson,
Arthur, 1-64.
Water-resource data program:
Arnov, Ted, 2-64.

Utah--Continued

- Maps
Brighton quadrangle, geologic:
Baker, A. A., 1-65.
Draper quadrangle, geologic:
Crittenden, M. D., Jr.,
1-65.
Dromedary Peak quadrangle,
geologic: Crittenden, M. D.,
Jr., 2-65.
Flaming Gorge quadrangle,
geologic: Hansen, W. R.,
2-65.
Mount Aire quadrangle, ge-
ologic: Crittenden, M. D.,
Jr., 3-65.
Nipple Butte quadrangle, ge-
ologic: Waldrop, H. A.,
1-65.
Park City West quadrangle,
geologic: Crittenden, M. D.,
Jr., 5-65.
Sugar House quadrangle, ge-
ologic: Crittenden, M. D.,
Jr., 4-65.
Paleontology
Pelecypoda, Jurassic: Imlay,
R. W., 1-64.
Petrology
Lanan area, intrusive igneous
rocks: John, E. C., 1-64.
Stratigraphy
Chinle and Moenkopi Forma-
tions, Navajo and Hopi
Indian Reservations: Re-
penning, C. A., 2-65.
General: Tooker, E. W., 1-64.
Northeastern: Poole, F. G.,
3-64.
Permian, Lower: Barosh,
P. J., 1-64.
Quaternary, Salt Lake City:
Van Horn, Richard, 1-65.
Structural geology
Fracture-controlled lacco-
liths: Witkind, I. J., 1-65.
General: Crittenden, M. D.,
Jr., 3-64.
Tintic Mountains, tear fault:
Morris, H. T., 2-64.
Vanadium
Analysis
Determination in water: Fish-
man, M. J., 1-64.
Colorado
Resources, general: Fischer,
R. P., 1-64.
General
Supply and consumption:
Fischer, R. P., 5-64.
Geochemistry
Decavanadate ion, structure:
Evans, H. T., Jr., 1-64.
Idaho
Resources, general: Fischer,
R. P., 4-64.
Nevada
Resources, general: Fischer,
R. P., 2-64.
New Mexico
General: Fischer, R. P., 1-65.
South Dakota
Resources, general: Fischer,
R. P., 3-64.

Vanadium--Continued

Utah

Resources, general: Fischer, R. P., 6-64.

Vermont

Hydrogeology

Connecticut River basin, ground water resources: Cederstrom, D. J., 1-65.

Maps

Berlin quadrangle: Holmes, G. W., 5-65.

Bolton Landing, Glens Falls, and Whithall quadrangles, aeromagnetic: Balsley, J.R., 2-65.

North Adams quadrangle, aeromagnetic: Popenoe, Peter, 2-64.

Virginia

Areal geology

Middleboro South quadrangle: Englund, K. J., 1-64.

Economic geology

Bauxite, Staunton area, Valley and Ridge province: Warren, W. C., 1-65.

Gypsum and salt, occurrence and origin: Withington, C. F., 2-65.

Geochemistry

Catoctin Formation, greenstone: Reed, J. C., Jr., 1-64.

Geomorphology

Shenandoah Valley: Hack, J. T., 1-65.

Geophysical surveys

Alberta area, heat flow: Diment, W. H., 4-65.

Duffield quadrangle, geologic: Harris, L. D., 2-65.

Stratigraphy

Pennsylvanian, southwestern: Miller, R. L., 1-65.

Volcanoes

Hawaii

Kilauea, temperature studies: Peck, D. L., 3-64.

Vertical ground displacements: Decker, R. W., 1-65.

Volcanology

Geophysical study: Eaton, J. P., 1-63.

Nuée ardente

Origin of term: Hooker, Marjorie, 1-65.

Washington

Areal geology

Bald Knob quadrangle: Staatz, M. H., 1-64.

Curlew quadrangle: Parker, R. L., 4-64.

Hunters quadrangle: Campbell, A. B., 1-65.

Metaline zinc-lead district: Dings, M. G., 1-65.

Economic geology

Metaline zinc-lead district: Dings, M. G., 1-65.

Geophysical surveys

Western gravity: Stuart, D.J., 1-65.

Washington--Continued

Glacial geology

Mount Ranier, Alpine glaciers: Crandell, D. R., 1-65.

Post-hypsithermal: Crandell, D. R., 1-64.

Puget Sound area, Pleistocene: Crandell, D. R., 2-65.

Hydrogeology

Artificial recharge: Price, Don, 1-65.

Eastern, drainage-area data: Williams, J. R., 1-64.

Mount Rainier National Park, ground water: Luzier, J. E., 1-64.

Pierce County, ground water: Walters, K. L., 1-65.

Roosevelt Lake, storage measurements: Nassar, E. G., 1-65.

Upper Green River Basin, logging and runoff: Richardson, Donald, 1-65.

Maps

Auburn quadrangle, geologic: Mullineaux, D. R., 1-65.

Black Diamond quadrangle, geologic: Mullineaux, D. R., 2-65.

Deep Creek area, geologic: Yates, R. G., 1-64.

Montesano quadrangle, geologic: Gower, H. D., 1-65.

Renton quadrangle, geologic: Mullineaux, D. R., 3-65.

Western, Bouguer-gravity: Stuart, D. J., 1-65.

Petrology

Tatoosh pluton, alkali feldspars: Wright, T. L., 1-64.

Sedimentary petrology

Clay, Seattle, Capitol Hill: Mullineaux, D. R., 1-64.

Stratigraphy

Columbia River basalt and Latah Formation: Griggs, A. B., 1-65.

Paleozoic, upper, Ferry and Stevens Counties: Becraft, G. E., 1-65.

Pleistocene: Armstrong, J. R., 1-65, 2-65.

Seattle area: Mullineaux, D. R., 4-65.

Weathering

Chattanooga Shale

Fischer assay: Brown, Andrew, 1-64.

Zircon

U-Pb and Th-Pb systems: Stern, T. W., 1-65.

Well logging

Electrical

Water quality calculation: Turcan, A. N., Jr., 1-65.

Wells and drill holes

Drilling methods

General: Bowman, Isaiah, 1-65.

Reverse circulation: Burnham, W. W., 1-64.

Wells and drill holes

Numbering grid: McClelland, E. J., 1-65.

West Virginia

Areal geology

Evitts Creek and Pattersons Creek quadrangles, stratigraphy and geologic history: deWitt, Wallace, Jr., 1-64.

Geomorphology

Shenandoah Valley: Hack, J. T., 1-65.

Geophysical surveys

Aeromagnetic: Beck, M. E., Jr., 1-64.

Hydrogeology

Ground water: McGuinness, C. L., 2-65.

Ohio County, ground water: Robison, T. M., 1-64.

Pocatalico River basin, estimated water budget: Wilmoth, B. M., 1-64.

Wisconsin

Areal geology

Belmont and Calamine quadrangles: Klemic, Harry, 1-64.

Chippewa Valley, late Quaternary history: Andrews, G. W., 1-65.

Cuba City, New Diggings, and Shullsburg quadrangles, geologic: Mullens, T. E., 1-64.

Dane County: Cline, D. R., 1-65.

Rewey and Mifflin quadrangles: Taylor, A. R., 1-64.

Winnebago County: Olcott, P. G., 1-64.

Geochemistry

Upper Mississippi Valley, zinc-lead: Hosterman, J. W., 1-64.

Hydrogeology

Dane County, ground-water resources: Cline, D. R., 1-65.

Plover River basin: Weeks, E. P., 3-64.

Water resources: Holt, C. L. R., Jr., 1-64.

Winnebago County, water resources: Olcott, P. G., 1-64.

Maps

Wadsworth quadrangle, floods: Noehre, A. W., 1-64.

Wyoming

Areal geology

Beaver Rim area, Tertiary deposits: Van Houten, F. B., 1-64.

Black Hills uplift, northern and western flanks: Robinson, C. S., 2-64.

Crook County, northern and western: Whitcomb, H. A., 2-64.

Crooks Gap area: Stephens, J. G., 1-64.

Flaming Gorge area: Hansen, W. R., 1-65.

Wyoming--Continued

Areal geology--Continued

Green River Formation, associated Eocene rocks: Bradley, W. H., 1-64.

Johnson County, northern and central: Whitcomb, H. A., 1-64.

Niobrara County: Witcomb, H. A., 1-65.

Powder River Basin: Sharp, W. N., 1-64.

Upton quadrangle: Mapel, W. J., 1-64.

Wind River basin, geologic history: Keefer, W. R., 1-65.

Economic geology

Uranium, Powder River Basin: Sharp, W. N., 1-64.

Geochemistry

Powder River Basin, radioactivity disequilibrium: Rosholt, J. N., 3-65.

Uranium, isotope fractionation: Rosholt, J. N., 1-65.

Shirley Basin, fractionation of uranium related to roll features in sandstone: Rosholt, J. N., 1-64.

Uraniferous phosphatic lake beds, Eocene: Love, J. D., 1-64.

Geophysical surveys

Crooks Gap area, Fremont County, gravity: Healey, D. L., 1-65.

Wind River Basin area, gravity: Case, J. E., 2-65.

Glacial geology

Teton Glacier: Reed, J. C., Jr., 3-64, 1-65.

Wind River Mountains: Richmond, G. M., 2-64.

Hydrogeology

Bighorn Basin, variation of permeability from core analyses and geophysical logs: Bredehoeft, J. D., 1-64.

Cooperative ground water investigation, 1963, 1964: Gordon, E. D., 1-64.

Crook County, northern and western, ground water: Whitcomb, H. A., 2-64.

Glendo area: Welder, G. E., 2-65.

Ground water, chemical quality: Langford, R. H., 1-64.

Johnson County, ground-water resources: Whitcomb, H. A., 1-64.

Niobrara County, ground water: Whitcomb, H. A., 1-65.

Platte County, Wheatland Flats area: Weeks, E. P., 2-64.

Wyoming--Continued

Hydrogeology--Continued

Sheridan County, ground water: Lowry, M. E., 1-64.

Southwestern, Great Divide and Washakie basins: Welder, G. E., 1-65.

Star Valley, upper, ground water: Walker, E. H., 1-65.

Surface water, 1963-64: Wiard, L. A., 1-64.

Sweetwater County, test drilling summary: Welder, G. E., 1-64.

Teton Valley, ground water: Kilburn, Chabot, 1-64.

Water quality investigations, 1963-64: Hanly, T. F., 1-64.

Yellowstone National Park, ground water investigations, 1960-63: Lowry, M. E., 2-64.

Water circulation, tracing: Rowe, J. J., 1-65.

Maps

Adam Weiss Peak quadrangle, geologic: Rohrer, W. L., 1-65.

Big Judson quadrangle, geologic: McAndrews, Harry, 1-64.

Bradley Peak quadrangle, geologic: Bayley, R. W., 1-65.

Continental Peak quadrangle, geologic: Zeller, H. D., 1-64.

Cooper Lake North quadrangle, geologic: McAndrews, Harry, 1-65.

Cooper Lake South quadrangle, geologic: McAndrews, Harry, 2-64.

Dickie Springs quadrangle, geologic: Zeller, H. D., 2-64.

Essex Mountain quadrangle, geologic: Zeller, H. D., 5-64.

Flaming Gorge quadrangle, geologic: Hansen, W. R., 2-65.

Freighter Gap quadrangle, NE 1/4, geologic: Zeller, H. D., 6-64.

NW 1/4, geologic: Zeller, H. D., 7-64.

Hay Meadow Reservoir quadrangle, geologic: Zeller, H. D., 8-64.

Mineral: U.S. Geological Survey Conservation Division, 1-64.

Pacific Springs quadrangle, geologic: Zeller, H. D., 10-64.

Palisades Peak quadrangle, geologic: Jobin, D. A., 1-65.

Wyoming--Continued

Maps--Continued

Parting of the Ways quadrangle, geologic: Zeller, H. D., 3-64.

Pinnacles NW quadrangle, geologic: Zeller, H. D., 9-64.

Poker Peak-Palisades Reservoir quadrangles, geologic: Albee, H. F., 1-65.

Tule Butte quadrangle, geologic: Zeller, H. D., 4-64.

Mineralogy

Lazurite, talc, and chlorite, Green River Formation: Bradley, W. H., 2-64.

Sedimentation

Cheyenne River basin: Rolfe, B. N., 1-64.

Stratigraphy

Cretaceous and Paleocene, Foote Creek and Dutton Creek Formations: Hyden, H. J., 1-65.

Goose Egg Formation: Maughan, E. K., 1-65.

Jefferson and Three Forks Formations: Sandberg, C. A., 1-65.

Tip-top unit, Sublette County: Marzolf, J. E., 1-65.

Structural geology

Gros Ventre Mountains: Keefer, W. R., 1-64.

Thrust belt, tectonic development: Armstrong, F. C., 2-65.

X-ray fluorescence analysis

Silicates: Rose, H. J., Jr., 1-65, 2-65.

Yukon

Areal geology

Southwest, Pleistocene: Krinsley, D. B., 1-65.

Glacial geology

Southwest, multiple glaciation: Krinsley, D. B., 1-64.

Zinc

Colorado

Oxidized deposits: Heyl, A. V., 1-64.

Idaho

General: Kiilsgaard, T. H., 3-64.

Nevada

General: Kleinhampl, F. J., 1-64.

Tennessee

Correlation of abundance with rock thickness: Wedow, Helmuth, Jr., 2-65.

Utah

General: Kiilsgaard, T. H., 4-64.

SUBJECT INDEX

[Some discussions cover more than one page, but only the number of the first page is given. Page numbers in italic refer to the list of investigations in progress (see p. A217-A250). See also Index to List of Publications]

A		Alabama—Continued			
	Page		Page		Page
ABC system, for geodetic control, evaluation --	201	geologic mapping -----	224, 230	Alberta, Canada, seismic crustal studies -----	141
ABS, adsorption on clay minerals -----	53	geomorphology -----	231	Alcohol, use in evaporation suppression -----	58
bacterial degradation -----	53	ground water -- 30, 170, 171,	244	Aldrin-water system, hydrogen isotopic-exchange reaction -----	53
Absorption analysis, infrared, use in fixed-water determination --	182	hydrology -----	244	Algae, effect on lake water ----	177
Absorption spectra, use in identification of minor elements -----	14	low-flow studies -----	235	fossil, occurrence and age -----	77
Abstracts. <i>See</i> Bibliographies and abstracts.		paleobotany -----	126	Alkalic rocks, age determinations -----	162
Accessory minerals, as ore guides -----	16	paleontology and stratigraphy -----	70	magnetite in, composition -----	157
Acid mine drainage, effect on trees -----	179	quality of water -----	30, 239	Alkalies, in tektites, error in old analyses -----	135
occurrence -----	53, 54	surface water -----	30, 234, 244	in ultramafic rocks, amount -----	148
Acoustic device, for measuring water velocity --	186	terraces along Alabama River -----	71	Allanite, petrology and geochemistry -----	147
Activities, of elements on binary sulfides -----	7	water resources -----	244	Alteration, hydrothermal, in copper - lead - zinc deposits -----	17
Activity-activity diagrams, use in borate study --	155	Alaska, beryllium -----	11, 217	<i>See also</i> Diagenesis, Diagenetic rocks.	
use in metamorphic studies -----	154	bridge sites -----	244	Amchitka Island, Alaska, nuclear test site ----	46
Adirondack Mountains, New York, iron -----	3	coal -----	15, 218	American Samoa, cooperating agencies -----	209
New York, marble -----	154	construction and terrain problems -----	219	ground water -----	244
Aerial photographs, catalog --	69	continental shelf -----	104	Amino acids, in sewage, reduction -----	52
Aerial photography, in Antarctic studies -----	197, 200	cooperating agencies -----	205	Ammonia, association with mercury -----	6
in hydrologic-data collection -----	171	copper -----	219	Ammonites, occurrence and age -----	91
research -----	200, 201, 202	economic geology -----	100	Ammonium ion, determination in silicates -----	182
Aeromagnetic studies, district surveys -- 74, 78, 91, 95, 99, 105	231	geobotanical prospecting --	17	Analog models, use in water studies -----	35, 38, 155, 171, 174
investigations in progress. <i>See also</i> Magnetic studies.		geochemical prospecting --	222	Analyses. <i>See types of analyses under</i> Analytical chemistry.	
Aeroradioactivity studies, district surveys ----	78, 82	geochemistry -----	222	Analytical chemistry, investigations in progress -----	217
investigations in progress. <i>See also</i> types of analyses: Colorimetric, Gas chromatographic, Infrared absorption, Neutron activation, Optical spectroscopic, Spectrographic, Spectroscopic, Spectrophotometric, X-ray fluorescence.		geochronology -----	100, 101, 102, 103, 104, 223	results of investigations --	181
Aerotriangulation, analytical, for stereomodels -	201	geologic literature, index --	217	<i>See also types of analyses: Colorimetric, Gas chromatographic, Infrared absorption, Neutron activation, Optical spectroscopic, Spectrographic, Spectroscopic, Spectrophotometric, X-ray fluorescence.</i>	
Afghanistan, surface water. 111, 250		geologic mapping -- 68, 223, 225	231	Analytical services and research -----	217
Age determinations. <i>See</i> Geochronology and various methods.		geophysics -----	231	Anhydrite, euhedral, synthesis -----	155
Agency for International Development, results of USGS investigations for. 109, 118		glacial geology -----	104, 233	Antarctica, geochronology -- 106, 107	
Aggregate, lightweight, from slate -----	10	glaciology -----	181, 233	geologic mapping -----	223, 225
Air, flow in tuff -----	52	gold -----	233	geologic reconnaissance ----	106
Air temperature, effect on water use by plants -----	174	gravity studies -----	62, 63	geophysics -----	231
Alabama, coal -----	218	ground water -----	38, 244	glacial geology -----	233
cooperating agencies -----	205	heat flow -----	142		
floods -----	60, 221	hydrographic studies -----	244		
		hydrology -----	244		
		iron -----	234		
		mineralogy -----	8, 104		
		nuclear test site -----	46		
		oil shale -----	15, 236		
		paleobotany -----	126		
		paleoclimate -----	102		
		paleontology -- 63, 100, 102, 103, 104, 122, 123, 124, 125, 237	238		
		permafrost -----	181, 238		
		petroleum and natural gas -----	238		
		petrology -----	100, 102, 104, 148, 222		
		pingos -----	181		
		quicksilver -----	239		
		stratigraphy -----	46, 63, 100, 102, 103, 241		
		structural geology -----	61, 63, 64, 100, 101, 102, 104		
		surface water -----	38		
		tectonic map -----	241		
		tin -----	242		
		vegetation -----	243		
		volcanology -----	243		
		waterpower resources -----	243		
		Alaska earthquake, 1964, geologic and hydrologic effects --	61, 181		

See also Index to List of Publications

Antarctica—Continued	Page
paleontology	237
petrology	107
structural geology	106
topographic mapping	197
Antimony, as an ore guide for	
gold	16
associated with cinnabar ..	5
district studies	5, 6
Appalachia, clays	218
geologic mapping	68
hydrologic atlas	19
resource studies	2
<i>See also names of individual States.</i>	
Appalachian Mountains, heat	
flow	142
palynology	126
Aquifers, artesian, occurrence	32, 33, 34
basalt	169
contamination	23, 52, 53, 55, 56
correlation in Gulf coast	
area	25
delineation by seismic studies	21
determination of characteristics	173, 233
effects of earthquakes	64
effects of nuclear explosions ..	47
effects of particle size on permeability	169
relation to streamflow	233
sea-water intrusion, Hawaii ..	41
water-level fluctuations	170
Arabia. <i>See</i> Saudi Arabia.	
Aragonite, oxygen-isotope content	164
Archaeocyathids, occurrence and age	122
Archaeology, relation of artifacts to alluvium	178
Arctic, geophysics	231
Arctic Ocean, heat flow	143
Argon, loss from shocked granite	134
Arizona, artificial recharge	217
asbestos	217
beryllium minerals	10
coal	218
cooperating agencies	205
copper	4, 220
evapotranspiration studies	174, 220
floods	220, 221
geochemical prospecting	222
geochemistry	17
geochronology	93
geologic mapping	224
geomorphology	231
geophysics	95, 231
ground water	32, 36, 171, 244
hydrology	244
land subsidence	58
lead	235
petroleum and natural gas ..	15, 238
petrology	94
plant ecology	179
sedimentation	176
seismic crustal studies	140
soil-moisture studies	173, 174
stratigraphy	93, 241
structural geology	4, 93
topographic mapping, research	200
uranium	12, 242

Arizona—Continued	Page
water resources	243, 244
waterpower resources	243
zinc	235
Arkansas, artificial recharge	217
barite	217
coal	14, 218
cooperating agencies	205
evapotranspiration studies ..	174
floods	221
geologic mapping	224
ground water	26, 29, 244
low-flow studies	235
paleontology	237
petroleum and natural gas ..	238
stratigraphy	71, 241
structural geology	82
surface water	26, 28, 234
water resources	244
Arsenic, as an ore guide for	
gold	16
in tektites	135
Arthropoda, occurrence and age ..	124
Artificial recharge, investigation in progress	217
Asbestos, investigations in progress	217
Ascension Island, Atlantic Ocean, geochemistry	156, 164
Astrogeologic studies. <i>See</i> Extraterrestrial studies.	
Astronauts, geologic training program	137, 220
Atlantic coast region, water resources	18
<i>See also names of individual States.</i>	
Atlantic Coastal Plain, marine geology	235
paleoenvironment studies	237
paleontology	237
stratigraphy	127, 241
Atlantic Continental Shelf and Slope, marine geology	126, 127, 235
marine hydrology	236
stratigraphy	241
<i>See also</i> Sea-water intrusion.	
Atlantic Ocean. <i>See</i> Ascension Island, Mid-Atlantic ridge, North Atlantic.	
Atlases, geologic, Sino-Soviet National atlas	118, 69
Atmosphere control, in spectroscopy	183
Atomic energy. <i>See</i> Nuclear explosions, Radioactive minerals, and closely allied subjects following these topics; Salt domes.	
Attachment organisms, in estuaries	129
Australia, geomorphology	117
ray crater	133
Automation, of rock analyses ..	183
Axes, in sediments, calculation ..	177

B

Backwater, in open channels ..	166
Bacteria, iron, Carboniferous ..	125

	Page
Bank-seepage studies, investigations in progress	223
Barite, deposits, relation to paleokarst	76
investigations in progress ..	217
Barium, determination, X-ray fluorescence	184
in black shale	156
Basalt, aquifers, artificial recharge	217
aquifers, occurrence	169
composition and origin	148
in dike swarm, Columbia Plateau	95
lead in, isotopic composition ..	164
magnetite in, composition	157
submarine, near Hawaii	160
Basaltic glass, undevitrified	148
Base-flow studies, surface water ..	168, 172
Base metals, investigations in progress	217
<i>See also</i> Heavy metals, Light metals, and names of specific base metals.	
Basement-rock map, North America	68
United States	69
Basin and Range region, geological and geophysical studies	92, 140, 238
Batholiths, geologic studies	84, 147, 164
Bauxite, investigations in progress	217
Bed-load sediment discharge, computation	175, 176
Belemnites, occurrence and age ..	123
Bench marks, tests of materials for	200
Bentonite, rate of dispersion in water	167
Bertrandite, unusual mineral association	10
Beryl, in Saudi Arabia	116
Beryllium, district studies	10, 116
investigations in progress	217
Bibliographies and abstracts	217, 259
BILBY event, hydrologic studies ..	47
Biogeochemical studies, New Mexico and Maryland	56
Biota zones, near-shore, relation to ground-water discharge ..	129
Biotite, as a geothermometer ..	157
Bismuth, determination by fire assay	182
Black shale, metal content	156
minor-element distribution	222
uranium content	242
Bolivia, geologic mapping	221
metal deposits	111
mineral resources	221
Borates, crystal chemistry	236
district studies	8, 9
investigations in progress	218
mineral parageneses	155
Botanical exploration and research, investigations in progress	221

See also Index to List of Publications

	Page		Page		Page
Bouguer gravity-anomaly map, United States ---	68	California—Continued		Cesium-137, scavenged by river sediment -----	52
Brachiopods, evolution -----	123	paleontology -----	9,	Cesium iodide, thermolumines- cence -----	146
investigations in progress -----	237	97, 98, 100, 123, 124, 125		Chaetetids, occurrence and age -----	122
occurrence and age -----	75, 79, 93, 96, 122	petroleum and natural gas -----	238	Channel characteristics, inves- tigations in prog- ress -----	223
Brazil, base metals -----	221	petrology -----	97, 98, 99, 222	Channel encroachment, effect on sediment depo- sition -----	176
copper -----	111	phosphate -----	98, 238	Channel-fill deposits, glacial ---	86
geologic education -----	112, 221	quality of water -----	239	CHARIOT, Project, hydrology ---	236
hydrogeology -----	250	quicksilver -----	239	Chemical analyses. <i>See types of analyses:</i> Colori- metric, Gas chro- matographic, Infrared ab- sorption, Neu- tron activation, Optical spectro- scopic, Spectro- graphic, Spec- troscopic, Spec- trophotometric, X-ray fluores- cence.	
hydrology -----	111	reservoir sedimentation --	241	Chemical equilibria studies ---	159
iron -----	111	sedimentation -----	240	Chemistry. <i>See Analytical chemistry, Cosmo- chemistry, Crystal chem- istry, Water chemistry.</i>	
surface water -----	250	seismic crustal studies ---	140	Chile, geochronology -----	112
Bridge-site studies, investiga- tions in progress -----	220	springs -----	241	geologic mapping -----	221
Brine, as a water contaminant	54	stratigraphy -- 92, 97, 98, 99, 241		heat flow -----	142
bromine content, determi- nation -----	183	structural geology -- 92, 97, 98, 241		mineral resources -----	221
carbonate, chemical compo- sition -----	186	surface water ----- 40, 167, 234		mineralogy -----	112
effect on clay and silt -----	158	tectonics -----	241	China, geologic studies -----	118
relation to quicksilver and copper-lead-zinc- silver deposits --	6	tungsten -----	220	Chitinozoans, investigations in progress -----	237
<i>See also Saline water.</i>		urban geology -----	242	Chloride, in ground water, source -----	55
Bromine, as a salinity indicator in halite -----	8	waterpower resources -----	243	Chlorine, in fluid inclusions, de- termination -----	184
determination in brines and rocks -----	183	zeolites ----- 11, 250		in tuff, effect on scapolite --	149
Bryophytes, use in geobotanical prospecting -----	17	zinc -----	235	Chlorites, new minerals -----	236
Bryozoans, investigations in progress -----	237	Callipterids, reproductive meth- ods -----	126	occurrence in salt bed -----	150
occurrence and age -----	122	Cameras, high-speed, for lunar study -----	132	Chromatography. <i>See Gas chromatographic analysis.</i>	
Buddingtonite, new occurrence	6	super-wide-angle, for aerial photo- graphs -----	201	Chromite, investigations in progress -----	220, 221
Buried valleys, occurrence --	22, 27, 71	television, for lunar study -----	136, 137	Chromium, in black shale -----	156
C		Capitol Reef National Monu- ment, Utah, popu- lar geology -----	91	Cinnabar, associated with antimony ore -----	62
Cactus, disappearance in south- western States -----	178	Carbon-14, use as ground-water tracer -----	55, 158	Cities. <i>See Urban areas, Ur- banization.</i>	
Cadmium-chromium, contami- nation of ground water -----	219	Carbon-14 method, results of investigations -----	82, 104	Clay minerals, adsorption of ABS on -----	53
Calcite, black, association with silver -----	7	Carbonate rock, ground water in -----	171	cation-exchange rates, measurement -----	188
fluid inclusions in -----	156	resources -----	236	effect of brine on -----	158
oxygen-isotope content -----	164	Carbonate sediments, experi- mental compac- tion -----	145	Clay-water relations, investiga- tions in progress --	218
Calcium, determination, spec- trophotometric --	184	Carbonates, in cephalopod shells, oxygen-iso- tope content --	164	Clays, district studies -----	10
in Hawaiian olivine -----	148	Carbonatites, magnetite in, composition -----	157	effect on overthrusting ---	155
Calderas, geophysical studies, Japan -----	117	Caribbean region, paleontol- ogy -----	237	erosion characteristics ---	231
California, borates -----	8, 9, 218	<i>See also Isla Mona, Lee- ward Islands, Puerto Rico, Virgin Islands.</i>		investigations in progress --	218
chromite -----	220	Cat, Pleistocene, in Alaska ---	125	liquid movement in -----	218
coal -----	218	Cation disordered structure, system Cu-Ag-S-	6	<i>See also Flint clay.</i>	
construction and terrain problems -----	219	Cation-exchange capacity, glau- conitic sand --	158	Climate, effects. <i>See Precipita- tion, Rainfall, Snow.</i>	
cooperating agencies -----	205	Cation-exchange rates, meas- urement -----	188	Clinoptilolite, occurrence -----	11
evaporation studies -----	220	Caves, exploration for guano deposits -----	105		
floods -----	59, 220, 221	relation of artifacts to alu- vium -----	178		
geochemistry -----	155, 158, 222	Central Treaty Organization, geologic studies in member coun- tries -----	116		
geochronology -----	97, 99	Central United States, aero- magnetic surveys --	231		
geologic mapping -----	223, 225	<i>See also names of individ- ual States.</i>			
geomorphology -----	175, 178, 231	Cephalopods, investigations in progress -----	237		
geophysics -----	99, 100, 231	occurrence and age -----	123, 164		
geothermal well -----	150	shells, oxygen-isotope con- tent -----	164		
glacial geology -----	223				
gravity crustal studies ---	142				
ground water -----	32, 41, 223, 234, 244				
hydrology -----	244				
land subsidence -----	58, 235				
lead -----	235				
mineralogy -----	6,				
	147, 149, 150, 152				

See also Index to List of Publications

	Page		Page		Page
Coal, district studies	2, 14	Colorado River, geologic history	231	Covellite, natural coexistence with djurleite	8
investigations in progress	218	Colorimetric analysis, use in determination of gold and silver	17	Crabs, Oligocene, in Alaska	125
mine bumps	57	Columbia Plateau region, geologic studies	95	Cratering phenomena, terrestrial and experimental	133, 220
mining, effect on runoff	173	Columbia River Basalt, hydrology	243	Craters, lunar, distribution and abundance	136
effect on trees	179	Compaction, as cause of earth cracks	58	lunar, isostatic rebound	132
production, China and U.S.S.R.	118	experimental, of carbonate sediments	145	shape, affect of porosity and cohesion on	134
Coal balls, investigations in progress	236	Computers, use in studies, hydrologic	186, 187	Crinoids, occurrence and age	79
Cobalt-60, scavenged by river sediment	52	use in studies, lead-zinc	5	Crossbedding, relation to streamflow	176
Collapse structures, erosion caused by	175	magnetic	144	Crust, deformation by Alaska earthquake	63
Colombia, evaporites	112	<i>See also</i> Data processing.		Crust and upper mantle, geophysical studies	139, 142, 161
geologic mapping	112	Concentration, of dissolved solids in water, rapid method	187	Cryptovolcanic structures, Tennessee	133
minerals exploration	221	Connecticut, aeromagnetic studies	74	Crystal chemistry, results of investigations	151
phosphate	112	cooperating agencies	206	Crystal settling, in granitic magma	153
Colorado, base metals	217	geologic mapping	226	Crystalline rocks, as source of water	38
<i>See also</i> base-metal names.		ground water	245	permeability	51
beryllium	10, 11, 217	sea-water intrusion	240	Crystallography, experimental, investigations in progress	236
coal	218	stratigraphy and igneous geology	74	experimental. <i>See also</i> Geochemistry, experimental	
construction and terrain problems	57, 219	water resources	243, 245	Cu-Ag-S, system, high-temperature phases	8
cooperating agencies	206	Conodonts, investigations in progress	237	Current meters, for measuring water velocity	186
copper	220	Conservation, natural resources, activities of Conservation Division	42	Curved channels, flow in, theoretical studies	166
earthquake studies	141	Construction and terrain problems, district studies	57, 120	Cu ₂ S, high-pressure tetragonal, inversion	8
fluorspar	221	investigations in progress	219	Cyclic deposition, Kentucky	80
geochemistry	154, 164, 222, 223	<i>See also</i> Urban areas and under names of individual States.			
geochronology	223	Contamination, water, investigations in progress	217, 219	Dahomey, minerals reconnaissance	221
geologic mapping	68, 91, 223, 225	water, results of investigations	52	Dams, predicted effect on evapotranspiration	174
geomorphology	177	<i>See also</i> Radioactive wastes, Saline water.		Darcy's law, for water movement, deviations	169
geophysics	231, 232	Continental shelves, Alaska, tectonics	63, 64	Data collection and processing, geologic, coding and retrieval	221
glacial geology	89	Atlantic, marine geology	126, 127	hydrologic	186, 187, 234
gold	2, 15, 223	Convection, in granitic magma	153	<i>See also</i> Computers.	
gravity studies	89, 142	Cook Inlet, Alaska, earthquake effects	62	Debris flows, Mt. Rainier, Wash.	57
ground water	35, 244	Copper, determination, spectrophotometric	184	Deerite, new iron silicate	149
hydrology	245	district studies	4, 5	Deformation research, investigations in progress	219
intrusive rocks	89	in black shale	156	Delaware, cooperating agencies	206
lead	2, 235	investigations in progress	219	floods	220
mineralogy	147	Copper River Basin, Alaska, earthquake effects	62	quality of water	239
molybdenum	4	Corals, investigations in progress	237	Delaware River, chemical characteristics	239
niobium	236	occurrence and age	92, 122	Delaware River basin, drought in	247
oil shale	91, 236	Cores, Mohole project, geologic study	128	flood mapping	61
petroleum and natural gas	2, 15, 238	Cosmochemistry, results of investigations	134	sea-water intrusion	240
petrology	86, 87, 146, 223	Costa Rica, volcanic studies	118, 221, 231, 243	Deposition, sediments, results of investigations	176
plant ecology	179			Depth functional, use in magnetic and gravity interpretation	145
potash	239				
quality of water	158				
rare earths	14				
reservoir sedimentation	241				
resource compilation	2				
saline minerals	240				
sedimentation	176, 240				
seismic crustal studies	140				
silver	2				
soil-moisture studies	241				
stratigraphy	87, 88, 91, 241				
structural geology	87, 88, 90				
surface water	35				
thorium	13, 242				
uranium	12, 13, 242				
urban geology	242				
vanadium	243				
volcanic rocks	88				
water resources	244				
waterpower resources	243				
zeolites	11				
zinc	2, 235				
Colorado Plateau region, geological and geophysical studies	90, 231, 241				
vanadium	243				

See also Index to List of Publications

	Page		Page		Page
Detergents, contamination of ground water -----	219	Eastern United States, aero- magnetic surveys -----	231	Exploration, mineral, tech- niques for -----	15
laboratory and field stud- ies -----	53	paleontology -----	237, 238	Extraterrestrial studies, lunar, instrumentation and tech- niques -----	132
Diagenesis, oolitic limestone---	83	seismic mantle studies---	139	lunar, manned exploration---	158
zeolites -----	11	<i>See also</i> New England, Northeastern United States, Southeastern United States, and names of in- dividual States.		mapping and terrain studies -----	130, 136, 219, 220
Diagenetic rocks, mineralogy --	150	Echinoderms, ecology -----	122	theoretical analyses of various phe- nomena -----	132
Diatomite, occurrence -----	82	Education, geologic, in Brazil	112	unmanned exploration studies -----	136
Diatoms, effect on lake water---	177	geologic, popularization at National Monu- ment -----	91		
investigations in prog- ress -----	236, 237	Elastic constants, rocks and minerals, calcula- tion -----	145	F	
occurrence and age -----	82, 127	Electrical resistivity, tektite glass -----	135	Facies, sedimentary, coexisting mineral paragene- ses -----	134
reclassification -----	127	Electrodes, glass, for determi- nation of sodium in water -----	185	sedimentary, occurrence in Ordovician rocks in Kentucky --	80
Dielectric constant, of moist rock, determina- tion -----	145	Electron-probe studies, investi- gations in prog- ress -----	217	<i>See also</i> Maps, lithofacies.	
Differentiation, magmatic, vari- ation trends of niobium -----	149	Elevation changes, topographic, earthquake- caused -----	62, 64	Faults, activation by nuclear explosions -----	47
Digenite, coexistence with co- vellite and djur- leite -----	8	Emission, spectral, of cesium iodide -----	146	Feldspar, alkali, heat of solu- tion -----	153
Digital-computer systems, use in hydrologic studies -----	186, 187	Energy-budget method, com- puting evapo- transpiration --	174	lead in, isotopic composi- tion -----	164
Dilution method, of streamflow measurement -----	187	Energy resources, investiga- tions in progress.	236	physical properties -----	153
Diopside, synthetic, lumines- cence -----	135	Engineering geology. <i>See</i> Con- struction and ter- rain problems, Urban areas, geology.		<i>See also</i> K feldspar, Mi- croperthite, Pla- gioclase glass.	
Discharge, surface-water, co- efficients for open- channel flow --	172	Enstatite, crystal chemistry---	151	Ferro-alloy metals, investiga- tions in progress.	220
Dispersion, in open-channel flow -----	165	Equilibria studies, sulfides---	7, 8	Fire assay, use in bismuth de- termination -----	182
Dissolved oxygen, in lake water -----	177	Equipment, new or modified. <i>See</i> Instruments and equipment.	159	Fish, fossil, occurrence and age -----	85, 91
in river water -----	53, 54	Erosion, investigations in prog- ress -----	231	Fixed water, in minerals, de- terminations --	182
portable measuring kit---	187	results of investigations.	175, 177	Flint clay, volcanic origin-----	79
Dissolved solids, in ground water, estima- tion -----	158	Errors, in geochemical data, source -----	157	Flood-plain deposits, soil-mois- ture storage -----	173
in rain and snow -----	157	Estuaries, biohydrology -----	129	Floods, district studies-----	59
in river water -----	53	quality of water -----	55, 128	effects, on sediment load--	176
in water samples, concen- tration -----	187	water movement -----	129	on vegetation -----	179
District of Columbia, cooperat- ing agencies -----	206	Evaporation studies, investiga- tions in progress.	220	frequency -----	60
geologic mapping -----	226	Evaporation suppression, in- vestigations in progress -----	220	investigations in progress--	220
geophysics -----	232	results of investigations--	58	mapping -----	60
glacial geology -----	71	Evaporites, district studies---	8	reduction, effect of water storage -----	168
surface water -----	166	Evapotranspiration, investiga- tions in progress.	220	<i>See also under names of</i> <i>States.</i>	
urban geology -----	242	results of investigations.	173	Florida, clays -----	218
Djurleite, mineralogy -----	151	Evapotranspirometer, use in evaporation stud- ies -----	174	color photomaps of swamps -----	202
natural coexistence with covellite -----	8	Exchange constants, glass-----	155	cooperating agencies -----	206
Drift, glacial, Alaska-----	63	Exchange rates, cation, meas- urement -----	188	evaporation studies -----	220
Drought, effect on water levels, Michigan -----	27	Exchange reactions, results of investigations --	154	floods -----	220
Dye, use, in evapotranspiration studies -----	174	Exfoliation, as cause of circular patterns -----	178	geochemistry -----	223
use, in ground-water stud- ies -----	170			geohydrology -----	245
in surface-water studies -----	166, 187			geologic mapping -----	226
E				ground water -----	20, 23, 129, 245
Earth cracks, caused by com- paction -----	58			ground-water hydrology --	158
Earthquakes, Alaska, 1964-----	61			ground-water-surface- water relations.	172, 233
Colorado-Wyoming-Mon- tana -----	141			humate -----	15
effects, on glaciers-----	181			paleontology -----	71, 122, 124, 237
on water fluctuations in wells -----	170			petroleum and natural gas--	238
investigations in progress---	232			petrology -----	223
				phosphate -----	9, 238
				quality of water---	20, 24, 55, 159
				sea-water intrusion -----	240

See also Index to List of Publications

Florida—Continued	Page
stratigraphy	70
surface water	24
water resources	245
Flow, characteristics, investigations in progress	234
in estuaries, theoretical analysis	129
<i>See also</i> Streamflow.	
Fluid inclusions, chlorine and sulfur content, determination	184
in minerals	155
Flume studies	165, 166
Fluorescence, methods of analysis. <i>See</i> X-ray fluorescence.	
Fluorometer, correction curves	187
Fluorspar, investigations in progress	221
Foraminifera, investigations in progress	236, 237
occurrence and age	70, 121
Foreign nations, investigations in progress	221
results of investigations	109
Forsterite, luminescence in meteorite	135
Fractionation, of tritium	165
of uranium isotopes	163
Fractures, delineation on aerial photographs	170
effects, on aquifer permeability	32
effects, on rock erosion	178
produced by earthquakes	62
produced by nuclear explosions	46
Fuel resources, compilations and topical studies	236
Fuels, organic, results of investigations	14
<i>See also</i> Coal, Oil shale, Petroleum and natural gas.	

G

Ganophyllite, crystallography ..	152
Gas. <i>See</i> Petroleum and natural gas.	
Gas chromatographic analysis, use in herbicide and insecticide studies	182, 185
Gastropods, color patterns of shell	124
evolution	124
investigations in progress	237
occurrence and age	79, 91
Geobotanical prospecting. <i>See</i> Prospecting, geobotanical.	
Geochemical data, composition of rocks and minerals	156
use in geohydrologic study	171, 172
Geochemical exploration, abstracts	217
Geochemical prospecting. <i>See</i> Prospecting, geochemical.	
Geochemistry, distribution of the elements	221

Geochemistry—Continued	Page
distribution of the elements. <i>See also</i> under Maryland, New Mexico.	
experimental, investigations in progress	222
results of investigations	153
water, investigations in progress	222
results of investigations	157
<i>See also</i> Petrology and geochemistry and under names of States.	
Geochronology, investigations in progress	223
list of age determinations	163
results of investigations	93, 97, 99, 112, 117, 128, 161
<i>See also</i> Isotope and nuclear studies, Carbon-14 method, K/Ar method, Pb/a method, Rb/Sr method, and under names of States.	
Geodetic surveys, district	119
Geologic data, coding and retrieval. <i>See</i> under Data collection and processing.	
Geologic map of North America	68
Geologic mapping, intermediate-scale maps	68
investigations in progress	223
large regions	68
<i>See also</i> Maps, Mapping, and under names of States.	
Geomorphology, investigations in progress	231
results of investigations	177
<i>See also</i> under names of States.	
Geophysical instruments and techniques, for lunar exploration	139
Geophysics, investigations in progress	231, 232
results of investigations	139
<i>See also</i> Aeromagnetic studies, Magnetic studies, Aero-radioactivity studies, Gravity studies, and under names of States.	
Georgia, clays	218
color photomaps of swamps	202
cooperating agencies	206
geochemistry	222
geologic mapping	226
ground water	20, 23, 245
ground-water hydraulics	51
paleontology	70
phosphate	9

Georgia—Continued	Page
quality of water	20, 23, 55
radioactive-waste disposal	239
reservoir sedimentation	241
sea-water intrusion	240
stratigraphy	70, 77
structural geology	69
water resources	245
Geothermal studies, investigations in progress	232
<i>See also</i> Heat flow, Temperature studies.	
Geysers, interconnected underground	85
Ghana, ground water	112
Glacial deposits, hydrologic characteristics	169
submarine, reworking	128
<i>See also</i> Channel-fill deposits, Drift, Ice-contact deposits, Moraines.	
Glacial geology, investigations in progress	233
<i>See also</i> under names of States.	
Glaciers, study of growth and ablation	43
Glaciology, investigations in progress	233
results of investigations	180
Glass, exchange constants	155
optical properties, as measure of silica in volcanic rocks	148
<i>See also</i> Basaltic glass, Pantellerite glass, Plagioclase glass, Silica glass, Tektites.	
Glauconitic sand, cation-exchange capacity	158
Gold, association of tellurium with	15
association of trace metals with	16
determination, new colorimetric method	17
district studies	5, 6, 7
investigations in progress	233
radioactive, use in measuring streamflow	187
Granite, near ring dike, thermoluminescence	146
shocked, argon loss	134
Granitic magma, crystal settling and convection	153
Graptolites, investigations in progress	237
occurrence and age	75, 77, 97
Grasslands, relation to soil moisture	179
Gravel, district studies	57, 128
Gravity anomalies, map, United States	68
relation to earthquake-caused changes in elevation	62
relation to mineralized areas	49
Gravity studies, district surveys	8, 62, 63, 73, 86, 89, 91, 95, 105, 119, 142
investigations in progress	231
new interpretation method	145

See also Index to List of Publications

	Page
Great Basin. <i>See</i> Basin and Range region.	
Great Plains, paleontology	237
Great Salt Lake, Utah, physical and chemical hydrology	177
Green River Formation, investigations in progress	236, 238, 241
Greenland, construction and terrain problems	219
construction-site planning	221
geochronology	117
geologic mapping	226
glacial geology	233
glaciology	117
Greenschist facies, coexisting mineral paragenesis	154
Ground water, chemical equilibria	159
contamination, investigations in progress	219
occurrence	23, 53, 54, 55, 56
data storage and retrieval	187
dissolved-solids content	158
effects of earthquakes	64
flow, velocity	158
geochemistry	158
hydraulic and hydrologic studies	169
hydrology, investigations in progress	234
in crystalline rocks	233
in glacial deposits, effect of particle size	169
increase in age toward Great Salt Lake	162
offshore discharge	129
quality, compared with rainfall	157
<i>See also</i> Aquifers, contamination, and quality of water under names of States.	
relation to surface water	172, 233
sampling and recording devices	187
thermal characteristics	242
uranium content	158
vertical flow in wells	37, 41
Water Resources Division publications	18
Guam, cooperating agencies	209
marine geology	129
water resources	42, 245
Guano deposits, in caves on Isla Mona	105
Gulf coast, paleontology	237
petroleum and natural gas	238
Gulf Coastal Plain, marine geology	235
paleontology	237
Gulf of Maine, sea-floor topography	127

H

Halite, preparation for thermoluminescence study	146
Hardness, ground water, Susquehanna River basin	20

	Page
Hawaii, bauxite	217
cooperating agencies	206
floods	221
geochemistry	223
ground water	41, 173
hydrology	245
petrology	148, 160, 223
quality of water	41
seismic studies	141, 161
surface water	168
volcanology	159, 243
water resources	245
Hawaiian Volcano Observatory, results of investigations	159
Heat flow, results of investigations	142
Heat of solution, alkali feldspar	153
Heavy metals, district studies	3
Helium, occurrence in ground water	23
Herbicides, in water, determination	185
Highway tunnels, engineering geology	57
Horses, fossil, occurrence and age	71, 125
Howieite, new iron silicate	149
H ₂ S, in fluid inclusions	156
Humate, district studies	15
Hummerite, zinc analog of, crystal structure	152
Hydraulics, ground water, investigations in progress	233
surface water, investigations in progress	233
results of investigations	165
Hydrogeologic maps, Puerto Rico	105
Hydrologic data, collection and processing	186, 187, 234
Hydrologic instrumentation, investigations in progress	234
Hydrologic model studies, investigations in progress	236
Hydrology, bibliography	217
<i>See also</i> Ground water, Quality of water, Surface water.	

I

Ice, river, configuration on underside	166
<i>See also</i> Glaciology.	
Ice-contact deposits, as aquifers	21
Iceland, glacial geology	117
Idaho, antimony	6
clays	218
coal	218
cooperating agencies	206
ferro-alloy metals	220
floods	59, 220
geochemistry	223
geologic mapping	68, 223, 226
geomorphology	83
geophysics	95, 231, 232
ground water	38, 169, 171, 245
ground-water-surface-water relations	172

Idaho—Continued	Page
lead	2, 235
paleontology	121, 123, 124
petrology	84, 223
phosphate	238
quicksilver	5
radioactive-waste disposal	240
resource compilation	2
seismic crustal studies	140
silver	2
stratigraphy	51, 83
structural geology	6, 83, 95
surface water	38, 168, 245
thorium	242
tungsten	2
uranium	242
volcanology	243
water resources	245
waterpower resources	243
zinc	235
Ideality and nonideality, sulfide minerals	7
Illinois, cooperating agencies	206
floods	61, 220, 221
geophysics	231
ground-water	26
low-flow studies	235
paleobotany	236
phosphate	9
Impact phenomena, terrestrial and experimental	133
Impact structures, lunar, temperature regime	132
Indiana, cooperating agencies	206
geologic mapping	226
geomorphology	231
ground water	28, 245
limnology	177, 235
quality of water	177, 239
sedimentation	240
surface water	28, 234
water resources	245
Indonesia, construction and terrain problems	113
geologic investigations	221
geologic map	113
Industrial minerals, district studies	8
investigations in progress	234
Infrared analysis, investigations in progress	217
use in fixed-water determination	182
Infrared imagery, use in hydrologic-data collection	171
Infrared studies, district surveys	119, 146
lunar	132
Insecticides, in water, identification and analysis	182, 185, 219
Instruments and equipment, new or modified, automated X-ray quantummeter	183
new or modified, for hydrologic study	186, 234
for lunar study	132, 136, 137, 138, 139
Intrusive rocks, age determinations	162
Basin and Range region	93
Inversion, CuS to Cu ₂ S	8
Iowa, coal	218
cooperating agencies	206

See also Index to List of Publications

Iowa—Continued floods 220, 221 geomorphology 231 geophysics 232 ground water 245 ground-water-surface-water relations 172 low-flow studies 235 phosphate 9 water resources 245 Iran, minerals and mineral fuels 116 Irazú Volcano, Costa Rica, geologic studies 118 Iridium, in tektites 135 Iron, district studies 3 in ground water 37 investigations in progress 234 Iron-formation, Michigan 3 Iron oxide, effect on magnetization of zircon 146 Iron silicates, newly identified 149 Isla Mona, guano deposits in caves 105 Island regions. <i>See</i> Ascension Island, Guam, Isla Mona, Leeward Islands, Okinawa, Pacific Ocean, Palau Islands, Puerto Rico, Ryukyu Islands, Virgin Islands. Isostatic rebound, of lunar craters 132 Isotope and nuclear studies, investigations in progress 235 results of investigations 161 <i>See also</i> Geochronology. Isotopic analyses. <i>See</i> Neutron-activation analyses. J Jadeite, composition and origin 150 Japan, aeromagnetic-gravity studies 221, 231 geophysics 117 Jasperoid, association of tellurium and mercury with 16 in barite veins, Saudi Arabia 116 K K-Ar analyses, laboratory standard muscovite 161 K-Ar method, results of investigations 84, 89, 100, 101, 103, 104 K feldspar, relation to kaolinite and muscovite 155 Kansas, acid mine drainage 54 cooperating agencies 206 floods 220, 221 geologic mapping 226 ground water 36, 234, 246 hydrology 245 lead 235 low-flow studies 235	Kansas—Continued petroleum and natural gas 238 quality of water 35, 54, 239 sedimentation 240 stratigraphy 241 surface water 35, 168, 234 water resources 245 Kaolinite, movement of water through 169 relation to muscovite and K feldspar 155 solubility 218 suspensions, flocculation 154 Karst features. <i>See</i> Pseudokarst features. Kentucky, clays 79 coal 218 cooperating agencies 207 floods 220, 221 geologic mapping 226 geomorphology 231 geophysics 78 glacial geology 79 ground water 29, 246 mining hydrology 236 paleontology 122, 125, 237, 238 palynology 237 plant ecology 179 quality of water 239, 246 saline water 29 stratigraphy 79, 80 surface water 29, 234 water resources 239, 246 zinc 16 Key Largo Coral Reef Preserve, Florida, echinoderms 122 Kilauea Volcano, 1965 eruption 159 Kodiak Island, Alaska, structural geology 63 L Lake levels, investigations in progress 235 Lake Superior region, geophysics 231 seismic crustal studies 140 subsurface structure 81 <i>See also</i> Michigan, Minnesota, Wisconsin. Lakes, effects of earthquakes on 61, 62 evaporation studies 220 quality of water 159 saline 158 <i>See also</i> Limnology, Reservoirs. Land subsidence, district studies 58 investigations in progress 235 Land use, flood plains, effect on vegetation 179 Landslides, mistaken for glacial till 91 <i>See also</i> Debris flows, Rockfalls. Langite, confirmation of formula 183 Lapping, water, effect on thermoluminescence of halite 146 Laser microprobe, use in analysis of tektites 135 Lava lakes, crystallization 160	Lazurite, occurrence in salt bed 150 Lead, determination, in soil 182 determination, in water 185 district studies 5 investigations in progress 235 isotopic variation 165 Leeward Islands, paleontology 118 Leonhardite, occurrence in Colorado 11 Level rods, new types of graduation strips 201 Liberia, geologic investigations 221 Libya, geologic investigations 221 ground water 250 Light metals, district studies 8 Lightweight aggregate, district studies 10 Limestone, thermoluminescence 146 Limestone-terrane hydrology, investigations in progress 235 Limnology, investigations in progress 235 results of investigations 177 <i>See also</i> Lakes, Reservoirs. Lithium, in plants and soils 156 LONG SHOT site, Alaska, geology 46 Louisiana, cooperating agencies 207 floods 60, 61, 221, 246 ground water 25, 31, 171, 246 petroleum and natural gas 238 quality of water 31, 158 reservoir sedimentation 241 stratigraphy 25, 71, 241 surface water 246 water resources 246 Low-flow studies, district 21, 28, 168 <i>See also</i> under names of States. Luminescence, minerals in meteorites 135 <i>See also</i> Thermoluminescence. Lunar geology. <i>See</i> Extraterrestrial studies, lunar. M Mafic rocks, petrology and geochemistry 148 Magma. <i>See</i> Granitic magma. Magmatic processes, results of investigations 153 Magnesium, spectrophotometric determination 182, 184 X-ray-chemical determination 184 Magnetic studies, district surveys 3, 4, 8, 100, 116 results of investigations 143 <i>See also</i> Aeromagnetic studies, Magnetic susceptibility, Magnetization, Paleomagnetic studies. Magnetic susceptibility, measurement 146

See also Index to List of Publications

	Page
Magnetite, occurrence in New York	3
relation of composition to occurrence	157
Magnetization, remanent, occurrence	73, 78, 81
remanent, use in correlation	50
Maine, cooperating agencies	207
estuaries	128
floods	220
geochemical prospecting	17, 222
geologic mapping	227
geophysics	232
ground water	246
heavy metals in stream sediments	75
paleomagnetism	144
paleontology and stratigraphy	74, 75
quality of water	128
structural geology	74, 75
water resources	246
Management, natural resources on public lands	42
Manganese, behavior in water-determination, spectrophotometric	184
district studies	4
investigations in progress	220
See also Ferro-alloy metals.	
zoning in deposits	4
Manganese nodules, similarity to deep-sea type	4
Mantle, effect on geomagnetic variation	143
See also Crust and upper mantle.	
Mapping, floods	60
geologic	91, 111, 113
See also under names of States.	
lunar	136
topographic, in foreign countries	116
results of Topographic Division activities	116, 189
Maps, aeroradioactivity, Minnesota	82
basement rocks, North America	68
United States	69
Bouguer gravity anomaly, United States	68
geologic, Colorado-Utah	12
foreign countries	113, 114
North America	68
hydrogeologic, Puerto Rico	105
hydrologic, compiled by digital computers	187
isopach, Pakistan	113
lithofacies, Pakistan	113
mineral-resource, Philippines	115
orthophotomaps, research	202
paleotectonic, United States	69
permafrost, Alaska	181
tectonic, North America	68
See also Atlases.	
Marble, metamorphism	154
thermoluminescence	146

	Page
Marine geology and hydrology, investigations in progress	235
results of investigations	127
Maryland, acid mine drainage	54
clays	218
cooperating agencies	207
distribution of the elements in plants and soils	56
estuaries, attachment organisms	129
floods	220
geochronology	223
geologic mapping	227
geophysics	78, 231, 232
gravel deposits	57
ground water	20, 22, 246
hydrologic effects of urbanization	243
marine hydrology	235
plant ecology	179
quality of water	22, 54, 239
sedimentation	176
surface water	22, 166
urban geology	242
water resources	246
Maskelynite, occurrence in meteorite	133
Mass-transfer method, computing evapotranspiration	174
Massachusetts, artifacts and ventifacts	73
construction and terrain problems	219
cooperating agencies	207
floods	60, 220
geologic mapping	227
geomorphology	231
geophysics	232
glacial geology	73, 74
gravity studies	73
ground water	21, 246
low-flow studies	235
magnetic studies	73
paleontology	73
seismic studies	21
stratigraphy and igneous geology	72, 241
structural geology	73
surface water	167
urban geology	242
water resources	243, 246
Matthevia, stratigraphic distribution	123
Meanders, relation to stream discharge	178
Measurement techniques and instruments, hydrologic	186
Mercury, as an ore indicator	16
association with ammonia	6
domestic, economic evaluation	3
See Quicksilver for natural occurrences.	
Metallogenetic grid, for ore minerals	7
Metals, in black shale	156
in water, determination	184, 185
See also Base metals, Heavy metals, Light metals, Trace metals, and names of specific metals.	

	Page
Metamorphic rocks, petrology and geochemistry	149
Metamorphism, effect on thermoluminescence	146
marble	154
rocks in geothermal wells	150
zeolite-sedimentary-green-chist facies	154
Metasedimentary rocks, argillaceous, age determination	162
Meteorites, investigations in progress	220
luminescent minerals	135
plagioclase glass	133
Methane, in ground water	31
Metropolitan areas. See Urban areas, Urbanization.	
Micas, description of polytypes	151
new minerals	236
See also Biotite, Muscovite, Phlogopite.	
Michigan, cooperating agencies	207
copper	220
geochemical prospecting	222
geologic mapping	227
geomorphology	231
geophysics	232
ground water	27, 246
iron	3, 234
petroleum and natural gas	238
quality of water	27
surface water	27, 246
water resources	246
Microfossils, mummified	126
Microperthite, in Boulder batholith, Montana	148
Microprobe. See Electron-probe studies, laser microprobe.	
Mid-Atlantic ridge, geochemistry of lead	164
Midcontinent region, alkaline intrusives, age determinations	162
water resources	25
See also names of individual States.	
Mine bumps, in coal mines, prediction	57
Mineral analysis, investigations in progress	217
Mineral exploration, techniques	15
Mineral resources, compilations and topical studies	236
Mineralogy, experimental, investigations in progress	236
See also Geochemistry, experimental.	
results of investigations	151
See also under names of States.	
Minerals, on Federal lands, classification	42
on Federal lands, supervision	43
physical properties	145
Mining, acid mine drainage problem	53

See also Index to List of Publications

Mining—Continued	Page
coal, effect on trees.....	179
on Federal land, supervision	43
Mining hydrology, investigations in progress.....	236
Minnesota, cooperating agencies	207
geochronology	162
geophysics	82, 231, 232
ground water	26, 246
ground-water-surface-water relations ..	172
hydrology	246
quality of water	26
stratigraphy and structural geology	81
water resources	246
Minor elements, distribution, relation to public health	56
district studies	14
effects on system Fe-S.....	7
geochemical prospecting for	94
geochemistry	222
investigations in progress.....	236
See also Trace metals.	
Missiles, high-energy, use in impact studies ..	134
Mississippi, cooperating agencies	207
floods	59, 61, 221
geologic mapping	227
geophysics	232
ground water	30, 31, 47, 247
hydrologic effects of nuclear testing	47
low-flow studies	235
petroleum and natural gas	238
quality of water.....	31
surface water	48
water resources	236, 247
Mississippi embayment, hydrology	243
Paleocene transgressive sea	71
water resources	25
Mississippi River basin, water resources	243
Missouri, cooperating agencies.....	207
evapotranspiration studies	174
floods	220
geologic mapping	227
geophysics	232
ground water	26
lead	235
low-flow studies	235
paleontology	123
quality of water.....	28
sedimentation	240
surface water	28, 166
water resources	247
zinc	235
Missouri River, floods.....	221
Mizzonite, crystal structure.....	151
Mohole project, geologic study of cores	128
Moldavite, aerodynamically modified	134
Mollusks, investigations in progress	236
occurrence and age.....	85, 98, 100, 102, 123

Mollusks—Continued	Page
See also Belemnites, Cephalopods Gastropods, Pelecypods.	
Molybdenum, district studies.....	4
in black shale.....	156
in tektites	135
investigations in progress.....	220
Monazite, investigations in progress	236
rare-earth elements in	147
Monochrometer, use in thermoluminescence studies	146
Montana, base metals.....	217
See also base-metal names.	
chromite	220
coal	2, 218
construction and terrain problems	219
cooperating agencies	207
earthquake studies	141
floods	59, 220
geochemistry	164, 222, 223
geologic mapping	68, 223, 227
geomorphology	231
geophysics	231, 232
glaciology	233
ground water	2, 33, 247
ground-water-surface-water relations.....	172, 233
hot springs and geysers.....	85
hydrology	247
iron	235
manganese	4, 220
paleontology	123
petroleum and natural gas.....	3, 238
petrology	148, 223
phosphate	238
plant ecology	180
soil-moisture studies	241
stratigraphy	84, 241
structural geology	84, 242
tectonics	242
urban geology	242
volcanology	243
water resources	247
waterpower resources	243
Moon. See Extraterrestrial studies, lunar.	
Moraines, glacial	180
Mount Rainier, Wash., infrared surveys	146
Mudflow studies, investigations in progress	219
Mummified microfossils	126
Muscovite, laboratory standard for dating	161
relation to kaolinite and K feldspar	155
Muskox, Pleistocene, in Kentucky	125

N

National Atlas	69, 203
National parks and monuments, geologic studies ..	57, 91, 141, 159
topographic maps, list.....	195
National Topographic Program, objectives ..	189
Natural gas. See Petroleum and natural gas.	
Nebraska, construction and terrain problems ..	219

Nebraska—Continued	Page
cooperating agencies	207
floods	59, 220
geologic mapping	68, 228
ground water	34, 247
hydrology	247
petroleum and natural gas.....	238
quality of water.....	32, 239
sedimentation	175, 240
soil-moisture studies	241
stratigraphy	82, 240
water resources	247
Nepal, surface water.....	250
Net mass budget, glaciers.....	180
Neutron-activation analysis, for tantalum	147
of tektites	135
Nevada, antimony	5
base metals	217
See also base-metal names.	
beryllium	217
coal	218
construction and terrain problems	219
cooperating agencies	207
copper	5, 220
evapotranspiration studies	174
floods	59, 220, 221
geochemical prospecting ..	94, 222
geochemistry	155, 222
geochronology	93
geologic mapping	62, 223, 228
geophysics	49, 95, 232
gold	5, 6, 16
ground water	39, 40, 47, 247
lead	5, 235
lithium	156
mineralogy	50, 147
paleontology	92, 122, 237
petrology	48, 50, 94, 149
phosphate	238
plant ecology	180
prospecting by mineral association	16
reservoir sedimentation ..	241
resource compilation	2
seismic studies	45, 140
silver	5
soil-moisture studies	174
stratigraphy	5, 48, 92, 241
structural geology	5, 45, 46, 49, 92, 93, 242
surface water	40, 168, 247
tectonics	242
thermal water	150
water resources	247
waterpower resources	243
zinc	5, 235
Nevada Test Site, geologic and hydrologic studies ..	45, 46, 48, 236
New England, geophysics.....	231
glacial chronology	72
water resources	21
See also names of individual States.	
New Hampshire, cooperating agencies	207
ground water	21, 169, 247
paleomagnetism	144
surface water	234
water resources	243, 247
New Jersey, basement-rock surface	69
cooperating agencies	207
floods	61, 220, 221

New Jersey—Continued	Page
geohydrology	247
geologic mapping	228
geophysics	232
ground water	22, 170, 247
ground-water-surface- water relations	233
lightweight aggregate	10
low-flow studies	235
marine hydrology	236
mineralogy	152
offshore gravel	128
quality of water	53, 239
reservoir sedimentation	241
sedimentation	240
structural geology	70
surface water	22, 166, 167, 168, 174, 234
water resources	247
New Mexico, beryllium miner- als	10
coal	14, 218
construction and terrain problems	219
cooperating agencies	207
copper	220
distribution of the elements in plants and soils	56
erosion studies	175
geochemical prospecting	222
geochemistry	155, 223
geologic mapping	224, 228
geomorphology	177, 231
geophysics	231
ground water	36, 233, 247
ground-water-surface- water relations	233
lead	235
paleontology	91
permeability studies	173
petroleum and natural gas	2, 238
petrology	52, 149, 223
plant ecology	179
potash	239
quality of water	239
radioactive-waste disposal	239, 240
resource compilation	2
saline minerals	240
sedimentation	175
soil-moisture studies	241
stratigraphy	87, 241
structural geology	87
surface water	36, 248
thorium	13
uranium	242
volcanology	243
water resources	239
zinc	235
New York, buried valley on Long Island	71
cooperating agencies	207
floods	221
geochemistry	223
geologic mapping	228
geomorphology	231
geophysics	232
ground water	20, 21, 170, 233, 248
iron	3
low-flow studies	235
metamorphism	72, 154
petrology	223
precipitation studies	157
quality of water	20, 51, 56, 239
sea-water intrusion	240

See also Index to List of Publications

New York—Continued	Page
stratigraphy	72, 241
surface water	21, 167, 234, 248
water resources	248
Nickel, in black shale	156
Nigeria, ground water	113, 250
hydrology	250
surface water	250
Niobium, distribution in tuff	149
investigations in progress	236
variation during magmatic differentiation	149
North America, basement map	68
bibliography of geology	217
eastern, corals	122
geologic map	68
Jurassic paleontology	237
tectonic map	68
North Atlantic, ostracodes	237
North Atlantic Slope basins, flood studies	221
<i>See also names of individ- ual States.</i>	
North Carolina, cooperating agencies	208
evaporation studies	220
floods	60, 221
geologic mapping	224, 228
geomorphology	178, 231
geophysics	232
ground water	20, 153, 171, 248
hydrologic effects of urban- ization	242
pegmatites	238
phosphate	9
precipitation studies	157
quality of water	157, 158
sedimentation	240
structural geology	77
surface water	167, 248
water resources	239, 248
North Dakota, coal	218
cooperating agencies	208
evapotranspiration stud- ies	174
floods	220, 248
geomorphology	231
glacial geology	86
ground water	33, 248
limnology	159, 177
petroleum and natural gas	238
quality of water	159, 177
stratigraphy	33, 241
water resources	248
Northeastern United States, geophysics	231
gravity survey	231
paleontology	238
<i>See also New England and names of individ- ual States.</i>	
Northwestern United States, mineral explora- tion	236
<i>See also Pacific coast re- gion, Pacific Northwest, and names of indi- vidual States.</i>	
Nuclear explosions, hydrology	236
tunnel sites, selection	45
underground, detection	46
geologic and hydro- logic effects	46
use in water-resources de- velopment	46, 219

	Page
Nuclear studies. <i>See</i> Isotope and nuclear stud- ies.	
O	
Obsidian, absorption of water	153
Ohio, cooperating agencies	208
floods	60, 221, 248
geomorphology	27
ground water	26, 27, 28, 169, 234, 248
low-flow studies	235
quality of water	27, 55, 239
water resources	248
Ohio River basin, hydrology	243
water resources	20, 26
Ohio River valley, geologic his- tory	231
Oil. <i>See</i> Petroleum and natural gas.	
Oil-field brine, as water con- taminant	55
Oil shale, district studies	15, 91
investigations in progress	236
Okinawa, ground water	250
sedimentation	250
Oklahoma, coal	218
cooperating agencies	208
geologic mapping	228
ground water	36, 248
lead	235
paleontology	123
petroleum and natural gas	238
quality of water	239, 248
seismic crustal studies	141
stratigraphy	241
surface water	166, 234
water resources	248
waterpower resources	243
zinc	235
Olivine, Hawaiian, X-ray anal- ysis	148
Open channels, backwater	166
flow, theoretical studies	165
Optical spectroscopic analysis, results of investi- gations	183
Ore bodies, uranium roll-type, origin	13
Ore indicators	15
Ore minerals, sulfide, labora- tory studies	7
Oregon, artificial recharge	217
coal	14, 218
cooperating agencies	208
ferro-alloy metals	220
floods	59
geochemistry	164
geologic mapping	68, 224, 228
geophysics	232
ground water	39, 169, 248
ground-water-surface- water relations	172
limnology	235, 248
paleontology	124, 237
petrology	95, 97
quality of water	239
quicksilver	239
radioactive-waste dis- posal	239
saline lakes	158
sedimentation	240
stratigraphy	95, 97
surface water	234

- Oregon—Continued** Page
 water resources ----- 248
 waterpower resources ----- 248
Organic material, extractable,
 in shale ----- 156
Oriskany Formation, phos-
 phate ----- 238
 silica ----- 241
Orthophotomaps, research ----- 202
Osmotic pressure, in clayey
 sediments ----- 155
Ostracodes, classification --- 124, 125
 investigations in progress --- 237
 occurrence and age --- 70, 91, 100
Overthrusting, effect of clay
 on ----- 155
Oxygen, concentration in estu-
 aries ----- 128
 isotopes, in carbonates --- 164
See also Dissolved oxygen.
Oyster shell, radiocarbon age --- 128
- P**
- P waves, attenuation, deter-**
 mination ----- 145
Pacific coast region, floods --- 59
Pacific coast region, geochem-
 istry ----- 243
 geological and geophysical
 studies ----- 96
 geophysics ----- 231
 hydrology ----- 243
 paleoecology ----- 124, 238
 water resources ----- 37
See also names of individ-
ual States.
Pacific Northwest, geophysical
 studies ----- 231
 paleoecology ----- 236
 water resources ----- 38
See also names of individ-
ual States.
Pacific Ocean, eastern, Forami-
 nifera ----- 236
 geomagnetization studies --- 144
 geophysics ----- 231
 islands, biogeography --- 236
 geologic mapping --- 229
 vegetation ----- 243
See also Guam, Okina-
wa, Palau Islands,
Ryukyu Islands.
 organic content of sedi-
 ments ----- 128
Pacific Southwest, geophysics --- 231
See also names of individ-
ual States.
Pakistan, chromite ----- 113
 geologic maps ----- 113
 hydrology ----- 250
 mineral and mineral fuel
 resources --- 113, 114, 221
 mineralogy ----- 113
 quality of water ----- 114
 structural geology ----- 113
Palau Islands, marine geology --- 129
Paleobotany, investigations in
 progress ----- 236, 237
 occurrence and age of fos-
 sils ----- 103, 125
 systematic ----- 236
See also under names of
States.
Paleochannels. See Buried val-
 leys.
- Paleoecology, investigations in**
 progress ----- 236
 Pleistocene marine mol-
 lusks ----- 124
Paleohydrologic studies ----- 178
Paleomagnetic studies, dis-
 trict ----- 81, 99, 144
Paleontology, invertebrate, sys-
 tematic, investiga-
 tions in prog-
 ress ----- 237
 invertebrate, systematic,
 occurrence and
 age of fossils.
See classes of
invertebrate
fossils.
 stratigraphic, investiga-
 tions in progress --- 237
 vertebrate, systematic, in-
 vestigations in
 progress --- 237, 238
 systematic, occurrence
 and age of fossils --- 71,
 85, 91, 125
Paleoparadoxia, study of loco-
 motion ----- 125
Paleotectonic maps, United
 States ----- 69
Palladium, magnetic suscepti-
 bility ----- 147
Palynology. See Pollen, Spores.
Panama ground water ----- 114
 paleontology ----- 125
Panama Canal Zone, verte-
 brates ----- 235
Pantellerite glass, chemical
 characteristics --- 149
Particle size, effect on perme-
 ability ----- 169
Pb-a dating method, results of
 investigations --- 100, 107
Pb-Th dating method, results of
 investigations --- 84
Pb-U dating method, results of
 investigations --- 84
Pegmatites, investigations in
 progress ----- 238
Pelecypods, investigations in
 progress ----- 237
 occurrence and age --- 79, 91, 124
Pennsylvania, acid mine drain-
 age ----- 53
 coal ----- 14, 218
 coal-basin deposition --- 78
 cooperating agencies --- 208
 floods ----- 61, 221
 geologic mapping ----- 229
 geophysics ----- 78, 231, 232
 ground water --- 20, 26, 170, 248
 lightweight aggregate --- 10
 low-flow studies ----- 235
 paleobotany ----- 126
 paleomagnetism ----- 144
 paleontology ----- 75
 paleotectonic map ----- 69
 quality of water --- 20, 53, 239, 248
 sedimentation ----- 176, 240
 stratigraphy ----- 75, 79, 241
 structural geology ----- 76
 surface water ----- 248
 water resources ----- 248
Permafrost, investigations in
 progress ----- 238
 results of investigations --- 181
Permeability, crystalline rocks --- 51
 determination, in soil --- 173
 relation to fracturing, in
 aquifers ----- 32
- Permeability—Continued** Page
 relation to particle size --- 169
 unsaturated, determina-
 tion ----- 170
See also Darcy's law.
Perthite. See Micropertthite.
Peru, mineral resources --- 221
Pesticides, in water, determi-
 nation ----- 185
 in water, investigations in
 progress --- 217, 219
See also Herbicides, In-
secticides.
Petroleum and natural gas, dis-
 trict studies ----- 15
 investigations in progress --- 238
 on Federal land, supervi-
 sion ----- 44
 potential occurrence on At-
 lantic Continental
 Shelf ----- 127
Petrology, of extraterrestrial
 materials ----- 134
Petrology and geochemistry,
 field studies, re-
 sults of investi-
 gations ----- 147
Phase diagrams, theory and ap-
 plications ----- 153
Philippine Islands, chromite --- 115
 iron ----- 115
 maps, geologic ----- 114
 mineral-resource ----- 115
 mineral resources ----- 221
Phlogopite, reaction with
 quartz ----- 154
Phosphate, determination, in
 water ----- 185
 determination, of magne-
 sium in ----- 182
 district studies ----- 9
 investigations in progress --- 238
 origin ----- 9
 paleoenvironment, Califor-
 nia ----- 98
Phosphate minerals, crystal
 chemistry ----- 236
Phosphoria Formation, niobium
 stratigraphy ----- 241
Phosphorus. See Radiophos-
phorus.
Photogrammetry, instruments
 and techniques
 for lunar study --- 137
 results of investigations --- 201
Phreatophytes, investigations
 in progress ----- 220
Physical properties, rocks and
 minerals ----- 145
Phytoplankton, variation with
 tides ----- 129
Pierre Shale, paleontology --- 237
 stratigraphy ----- 241
Pingos, in central Alaska --- 181
Pioneer, USCGS ship, marine
 geologic studies --- 129
Pipes, uranium-bearing, Ari-
 zona ----- 12
Piping, erosion caused by --- 175
Plagioclase glass, from mete-
 orite ----- 133
Plant ecology, investigations in
 progress ----- 239
 results of investigations --- 178
See also Vegetation --- 239
Plant fossils. See Paleobotany.
Plants, changes in southwest-
 ern States ----- 178

See also Index to List of Publications

Plants—Continued	Page
chemistry, relation to soil chemistry	180
effect on lake water	177
lithium content	156
<i>See also</i> Trees, Vegetation.	
FLOWSHARE Program, geologic and hydrologic studies	46
Plutonic rocks, petrology and geochemistry	147
Polarimetry studies, lunar	132
Polarity, geomagnetic	143
Pollen, fossil, occurrence and age	71, 75, 126
investigations in progress	237
Polonium, removal from radium solutions	183
Porosity, of impact target, effect on crater shape	134
Potash, investigations in progress	239
Potassium, in alteration zone in ground water	17
Potassium feldspar. <i>See</i> K feldspar.	
Powellite, in Saudi Arabia	115
Prairie potholes, chemical equilibria	159
effect of vegetation on water	177
hydrology	174
Precipitation, chemical studies effect on tree growth	157
<i>See also</i> Rain, Rainfall, Snow.	
Preglacial drainage, Ohio	27
Primitive Areas, resource studies	2
Probe, drag-wire, for measuring water velocity	186
Prospecting, elements and minerals as ore guides	15
geobotanical	17
geochemical, investigations in progress	222
results of investigations	94, 115
Pseudokarst features, caused by piping and collapse structures	175
Public and industrial water supplies, investigations in progress	239
Public lands, classification	42
Publications, by USGS investigators in fiscal year 1965, list.	259
Water Resources Division	18
Puerto Rico, cooperating agencies	209
floods	61, 220, 221
geologic mapping	229
geophysics	105, 231, 232
heat flow	142
hydrogeologic map	105
petrology	105
structural geology	105
water resources	24, 248
Puerto Rico Trench, aeromagnetic study	105
radiolaria	128
Pyroxene. <i>See</i> Diopside, Enstatite.	
Pyrrhotite, use as geothermometer	154

Q	Page
Quality of water, analyses, results of investigations	184
data compilation and tabulation	187
recording and measuring devices	186, 187
relation to quicksilver and copper-lead-zinc-silver deposits	6
stored samples, changes in composition	186
Water Resources Division publications	18
<i>See also under names of States.</i>	
Quantometer. <i>See</i> X-ray quantometer.	
Quartz, reaction with phlogopite	154
Quaternary geology, investigations in progress	231
Quicksilver, district studies	5
investigations in progress	239
origin	6
R	
Radiation-detection system, for use on riverbed	52
Radioactivation, investigations in progress	217
Radioactive disequilibrium, results of investigations	163
Radioactive materials, transport in water, investigations in progress	239
Radioactive minerals, district studies	12
Radioactive wastes, disposal, investigations in progress	239
disposal, results of investigations	50
Radiocarbon. <i>See</i> Carbon-14.	
Radiochemistry, investigations in progress	217
Radioelements, physical chemistry	217
Radiolaria, investigations in progress	237
occurrence and age	128
Radionuclides, concentration in water	51
transport, investigations in progress	239
results of investigations	50, 51
Radiophosphorus, use in study of water movement	170
Radium, in ground water, Wyoming	56
standard solutions, removal of polonium	182
Rain, dissolved-solids content	157
tritium content	157
Rainfall, effect on runoff	168
Ranger spacecraft, use in lunar mapping	136
Rapid rock analyses, methods. <i>See</i> X-ray fluorescence analysis.	

	Page
Rare-earth elements, district studies	14
in allanite and monazite	147
resources and geochemistry	236
Ray crater, Australia	133
Ray material, lunar, calculation of trajectories	132
Rb-Sr method, results of investigations	106
Recharge data, ground-water, use in computing streamflow	173
Reedmergnerite, crystal chemistry	151
Refractive index, as measure of silica in volcanic rocks	148
Remanent magnetization, occurrence	73, 78, 81
use in correlation	50
Remote sensing. <i>See</i> Infrared imagery, Aerial photography.	
Reptiles, fossil, occurrence and age	91
Reservoirs, effects, on rate of movement of stream water	167
effects, on stream temperature	167
evaporation studies	220
sedimentation, investigations in progress	241
results of investigations	176
use for flood reduction	168
Resource compilation, Colorado, Idaho, Montana, Nevada, New Mexico	2
Rhenium, investigations in progress	220
<i>See also</i> Minor elements.	
Rhode Island, cooperating agencies	208
floods	220
geologic mapping	229
stratigraphy and igneous geology	72
water resources	248
Rift zones, crustal velocities	141
Ring dikes, granite associated with thermoluminescence	146
Ripple lamination, use in distinguishing sedimentary environments	177
Ripple marks, on underside of river ice	166
relation to streamflow	177
Rivers, sediment load, variability and prediction	175, 176
<i>See also</i> Estuaries, Streamflow, Surface water.	
Rock analyses, data of	221
investigations in progress	217
methods, X-ray	183
<i>See also</i> Analytical chemistry.	
Rockfalls, Mt. Rainier, Wash.	57
Rocky Mountain region, aeromagnetic surveys	231
coal	14

Rocky Mountain region— Continued	Page
seismic studies	140, 231
<i>See also names of individual States.</i>	
Roll-type uranium ore bodies, origin	13
Runoff, effect of coal mining on	173
effect of paved watershed on	36
relation to rainfall	168
Russia, geologic studies	118
Ryukyu Islands, biogeography ..	237

S

Sagebrush, relation to geomor- phic processes	180
Saline minerals, investigations in progress	240
Saline water, in aquifers .. 23, 55, 56	
interface with fresh water, Kentucky	29
investigations in progress ..	239
mixing with fresh water ..	128
occurrence, lakes	158
river estuaries	22, 149
streams	54, 55
<i>See also Brine.</i>	
SALMON event, hydrologic stud- ies	47
Salt, on highways, ground- water contamination by	246
Salt domes, use in nuclear test- ing	47, 236
Salt water. <i>See</i> Brine, Saline water.	
Saltcedar, effects, on sedimen- tation	176
effects on stream channels ..	167
water use	174
Samoa. <i>See</i> American Samoa.	
Sampler, for cores in sandy bed material	51
water, deep-well	187
Samples, water, changes in composition	186
water, concentration of dis- solved solids	187
Sand, as natural filter for water	20
glauconitic, cation-ex- change capacity	158
transport in rivers	175
Sandstone, copper deposits in, investigations in progress	219
fractionation of uranium isotopes in	163
uranium-bearing	12, 242
Saudi Arabia, geochemical prospecting	115
geologic investigations	221
magnetic surveys	116
mineral discoveries	115
topographic mapping	116
Scapolite, crystal structure ..	151
relation to chlorine release in tuff	149
Scheelite, in Saudi Arabia ..	115
Schoepite, transformation of types I, II, and III	152
Scour and fill, in sand-bed streams	175

See also Index to List of Publications

	Page
Sea-water intrusion, investiga- tions in progress	240
Seamounts, magnetization anal- ysis, computer method	144
Secular variation, geomagnetic, relation to mantle	143
Sediment, accumulation and discharge in res- ervoirs	176
deep-sea, Pacific Ocean ..	128
grain orientation in	175
load in rivers, variability and prediction .. 175, 176	
suspended, in Niobrara River	32
Sedimentary facies, coexisting mineral paragen- eses	154
Sedimentary mineralogy, in- vestigations in progress	241
Sedimentary petrology, investi- gations in prog- ress	241
Sedimentary rocks, trace-ele- ment content	222
Sedimentary structures, formed in streams	176
ripple marks	177
Sedimentation, investigations in progress	240
investigations in progress. <i>See also</i> Geomor- phology and Quality of water.	
results of investigations ..	175
Seiches, on lakes, earthquake- induced	61, 65
Seismic network, Rocky Moun- tains	231
Seismic studies, in ground- water explora- tion	21
in predicting mine bumps ..	57
investigations in progress ..	231
seismic-refraction investi- gations	139, 161
Seismic waves, attenuation, de- termination	145
Selenium, association with ura- nium	16
resources and geochem- istry	236
<i>See also</i> Minor elements.	
Sewage, lagoon studies	219
treatment for amino acids and proteins	52
Sewage and industrial waste, use in artificial recharge	217
Shale, extractable organic ma- terial	156
<i>See also</i> Black shale.	
"Shock" minerals, from natural craters	134
Shock waves, speed in sand- stone	134
Silica, determination in HF-de- composed rocks ..	182
investigations in progress ..	241
Silica glass, solubility	154
Silicate minerals, crystal chem- istry	236
Silicate rocks, tin content	157

	Page
Silicates, determination of am- monium ion in	182
Silicic rocks, petrology and geo- chemistry	147, 149
Silver, association with black calcite	7
determination, new color- imetric method ..	17
spectrophotometric	182
district studies	5
economic evaluation	3
in black shale	156
investigations in progress ..	241
Sino-Soviet studies, coal re- sources	118
terrain atlas	118, 219
Slides. <i>See</i> Debris flows, Land- slides, Rockfalls, Submarine slides.	
Slope denudation, rate	175
Slope measurements, lunar, from spacecraft photographs .. 136, 137	
Small samples, X-ray fluores- cence analysis	184
Snow, dissolved-solids content ..	157
Snowmelt, effect on sediment load	176
Sodium, determination, use of glass electrode	185
in ground water	159
Soil moisture, investigations in progress	241
results of investigations .. 173, 179	
Soils, barium content, X-ray de- termination	184
lithium content	156
migration of uranium	163
permeability, determina- tion	173
relation to chemistry of plants	180
Solid-state studies, results of investigations	146
Solubility studies, results of in- vestigations	154
Solutes, rate of movement in streams	166
Sonar device, for measuring water velocity	186
South Carolina, cooperating agencies	208
floods	221
geochemistry and petrol- ogy	223
geologic mapping	224
ground water	23, 248
helium in ground water ..	23
low-flow studies	235
radioactive-waste dis- posal	239, 240
stratigraphy	70
surface water	167, 234, 249
water resources	248
South Dakota, coal	218
construction and terrain problems	219
cooperating agencies	208
floods	220
geologic mapping	229
geophysics	232
glacial geology	86
ground water	34, 249
pegmatites	238
petroleum and natural gas ..	238
quality of water	32

See also Index to List of Publications

South Dakota—Continued	Page
stratigraphy	86, 88, 241
uranium	242
urban geology	242
water resources	249
Southeastern United States,	
asbestos	217
bauxite	217
ferro-alloy metals	220
ground water	234
industrial minerals	234
monzonite	236
phosphate	238
talc	242
<i>See also names of individual States.</i>	
Southwestern United States,	
changes in vegetation	178
paleoecology	237
paleontology	237
stratigraphy	237
<i>See also names of individual States.</i>	
Sparker-probe surveys, regional	127, 129
Specific conductance, use in studying base flow	172
Specific yield, effect of drainage time on	170
Spectral lines, measurement of width	183
Spectrographic analysis, tektites	135
Spectrometer, gamma-ray, use in uranium, thorium, and potassium analyses	17
Spectrophotometric analysis, atomic-absorption, heavy-metals determination in water	164
flame, determination of elements	181, 182
Spectroscopic, analysis, investigations in progress	241
<i>See also Optical spectroscopic analysis.</i>	
Sphalerite, mercury content	16
Spherules, in tektites	135
Spores, fossil, occurrence and age	75, 126
Springs, in Nevada	40
investigations in progress	241
Stability relationships, basic	153
State maps, topographic	189
Stereoplotters, super-wide-angle	201
Stishovite, synthesis in laboratory	133
Storage, of surface water, effect on runoff	168
of water samples, effect on composition	186
Stratigraphy, investigations in progress	241
<i>See also under names of States.</i>	
Streamflow, computed from ground-water data	173
effect of ground-water runoff on	172
effect of transpiration on	174

Streamflow—Continued	Page
effect of urbanization on	167
effect on ice cover	166
estimation in ungaged basins	168
low flow	21, 28, 168
rate of movement	166, 167
relation to aquifers	233
separation into components	172
theoretical, in open channels	165, 166
transport of solutes in	166
variability, relation to geology	168
velocity-measuring devices and methods	186, 187
Stromatolites, occurrence and age	77
Strip mining, hydrologic effects	236
Strontium, spectrophotometric determination	184
Structural geology, investigations in progress	241
<i>See also under names of States.</i>	
Submarine slides, earthquake-caused	61
Subsidence. <i>See</i> Land subsidence.	
Sulfate equilibria	155
Sulfide deposits, investigations in progress	236
Sulfide ore minerals, laboratory studies	7
Sulfides, accessory in igneous rocks	222
binary, activities of elements	7
massive deposits	219
Sulfur, determination in fluid inclusions	184
Surface water, bed load of, computation	175
contamination, occurrence	53, 54, 55
effect of earthquakes on	64, 65
effect of flood runoff on sediment load	176
effect of snowmelt on	176
hydraulic and hydrologic studies	165, 234
investigations in progress	233
occurrence of uranium in recording and measuring devices	186
relation to ground water	172
temperature distribution	242
transport of radionuclides in	50
tritium content	157
Water Resources Division publications	18
Surveyor spacecraft, development of cameras for	136
Swamps, color photomaps	202

T

Talc, investigations in progress	242
occurrence in salt bed	150
Tantalum, in Southern California batholith	147
investigations in progress	236
<i>See also Minor elements.</i>	

Tectonic maps, North America—World	68
Tectonics, investigations in progress	241
Tektites, investigations in progress	220
results of investigations	135
Television, use in mapping terrain on Moon	136
Tellurium, as an ore indicator	15, 16
Temperature studies, estuaries on Maine coast	128
ground water	169
lake water	177
lunar impact structures	132
oxygen isotopes in carbonates	164
rocks at Mt. Rainer, Wash.	146
stream water	54, 167
volcanoes in Costa Rica	119
<i>See also Heat flow, Thermal loading, and closely allied subjects following these topics.</i>	
Tennessee, coal	218
cooperating agencies	208
copper	220
cryptovolcanic structure	133
floods	220, 221
geologic mapping	224, 229
geophysics	78
ground water	30, 171, 249
ground-water-surface water relations	223
hydrologic effects of urbanization	242
iron	235
lead	5, 235
low-flow studies	235
paleontology	71, 77
quality of water	30, 52
radioactive-waste disposal	239
stratigraphy	71
structural geology	77, 81
surface water	167, 168
water resources	249
zinc	5, 235
Terrain analyses, lunar	136, 137
Texas, cooperating agencies	208
evaporation suppression	58
evapotranspiration	220
floods	60, 221
geochemistry	223
geologic mapping	229
geophysics	231, 232
ground water	25, 37, 249
hydrologic effects of urbanization	243
hydrology	249
land subsidence	58
low-flow studies	235
paleontology	237
petroleum and natural gas	238
petrology	223
quality of water	37, 54, 55, 239
reservoir sedimentation	241
sedimentation	240
stratigraphy	25, 71, 82, 241
surface water	167, 168
uranium	13, 242
urban geology	242
water resources	243, 249
zeolites	11
Th-U method, results of investigations	102

See also Index to List of Publications

	Page
Thailand, economic geology ---	116
geologic investigations ---	221
Thermal loading, Susquehanna River ---	54
Thermal stratification, lake water ---	177
Thermal water, composition, near ore deposits ---	6
effect on surrounding rocks ---	150
occurrence ---	159
origin ---	150
Thermoluminescence, investigations in progress ---	220
results of investigations ---	146
<i>See also</i> Luminescence.	
Thermomagnetic studies, tektites ---	135
Tholeiites, composition and origin ---	148
Thorium, distribution in tuff ---	149
district studies ---	13, 14
estimate of worldwide supply ---	3
in alteration zone ---	17
investigations in progress ---	242
Tides, effect on phytoplankton ---	129
Tilting, earthquake-caused ---	62, 63, 64
Tin, amount in silicate rocks ---	157
investigations in progress ---	242
Titanium dioxide, error in old analyses ---	135
Topographic Division, summary of activities ---	189
Topographic mapping. <i>See</i> Mapping, topographic.	
Topography, effect on groundwater occurrence ---	170
Tourmaline, chromian, trivalent vanadium content of ---	183
Trace analysis, methods ---	217
Trace elements, in sedimentary rocks ---	222
Trace fossils, occurrence and age ---	103
Trace metals, as ore guides for gold ---	16
Tracer studies, ground water ---	55, 158
sediment load of rivers ---	175
streamflow ---	166
water in plants ---	174
Transpiration, effect on streamflow ---	174
Transport, coarse sediments, in Colorado Plateau ---	90
radionuclides ---	50
sediments, results of investigations ---	175
solutes in streams ---	166
Tree rings, relation to precipitation ---	179
Tree roots, as indicator of slope denudation ---	175
Trees, effect of acid water on growth ---	179
relation of growth to climate ---	179
relation to land use and flooding on flood plains ---	179
<i>See also</i> Plants, Vegetation.	
Trilobites, investigations in progress ---	237
occurrence and age ---	124
Tritium, in precipitation and stream water ---	157

	Page
Tritium—Continued	
isotopic fractionation ---	165
limitations as water tracer ---	171
Tuff, air flow in ---	52
Tungsten, as an ore guide for gold ---	16
Tunisia, ground water ---	250
Tunnel sites, for nuclear testing, selection ---	45
Tunnels, effect on ground water highway, engineering geology ---	173
Turkey, ground water ---	57
minerals and mineral fuels ---	250
	116
U	
U-Th-Pb system, in zircon, effect of weathering ---	161
Ulexite, crystal structure ---	152
Ultramafic rocks, petrology and geochemistry ---	148
United Arab Republic, ground water ---	250
United States, energy resources ---	236
maps, basement-rock ---	68, 240
geologic ---	240
geologic. <i>See also</i> Mapping, geologic.	
gravity-anomaly ---	68, 240
metallogenic ---	236
paleotectonic ---	69, 240
topographic ---	189
mineral-fuel resources ---	236
tantalum-niobium resources ---	236
<i>See also</i> names of States and large regions.	
Uranium, district studies ---	12
distribution in tuff ---	149
in alteration zone ---	17
in ground water ---	158
investigations in progress ---	236, 242
isotope studies ---	161, 163
migration in soil ---	163
relation to selenium ---	16
types of occurrence ---	242
Uranium minerals, crystal chemistry ---	236
Uranium oxide hydrates, coordination of uranyl ion ---	152
Urban areas, flood mapping ---	60
geology, investigations in progress ---	242
topographic maps, list ---	195
Urbanization, effects, on runoff ---	36
effects, on sediment load ---	176
on streamflow ---	167
hydrologic effects, investigations in progress ---	242
U.S.S.R., geologic studies ---	118
Utah, base metals ---	217
base metals. <i>See also</i> base-metal names.	
beryllium minerals ---	10
coal ---	218
construction and terrain problems ---	57, 219
cooperating agencies ---	208
copper ---	220

	Page
Utah—Continued	
elevations of Great Salt Lake ---	235
evaporites ---	8
ferro-alloy metals ---	220
floods ---	221
geochemical prospecting ---	94, 222
geochemistry ---	156
geochronology ---	162
geologic mapping ---	91, 224, 229
geomorphology ---	178
geophysics ---	91, 232
ground water ---	34, 35, 162, 249
lead ---	235
limnology ---	177, 235
mineral-resource map ---	236
oil shale ---	236
paleontology ---	91
petroleum and natural gas ---	238
petrology ---	94
phosphate ---	9, 238
popularization of geology at National monument ---	91
potash ---	239
quality of water ---	35, 239
reservoir sedimentation ---	241
saline minerals ---	240
stratigraphy ---	91, 241
structural geology ---	90, 91, 93
surface water ---	234, 249
uranium ---	12, 242
urban geology ---	242
vanadium ---	243
water resources ---	249
waterpower resources ---	243, 249
zinc ---	235

V

Vanadium, bibliography of geology and resources ---	218
in black shale ---	156
investigations in progress ---	243
trivalent, in chromian tourmaline ---	183
Vegetation, effect on sedimentation ---	176
investigations in progress ---	243
removal, effect on base flow ---	172
riparian, use of soil moisture by ---	174
<i>See also</i> Plants, Trees.	
Velocity, of water, measuring devices and methods ---	186, 187
Velocity layering, in granitic rock ---	45
Vermont, asbestos ---	217
cooperating agencies ---	208
floods ---	220
geologic mapping ---	230
paleomagnetism ---	144
talc ---	242
water resources ---	243, 249
Vertebrates. <i>See</i> Paleontology, vertebrate.	
Virgin Islands, cooperating agencies ---	209
water resources ---	25, 249
Virginia, coal ---	219
construction and terrain problems ---	219
cooperating agencies ---	208
floods ---	60, 220, 221

See also Index to List of Publications

Virginia—Continued	Page
geologic mapping	224, 230
geophysics	78, 231
ground water	20, 170
lead	235
paleontology	71, 77
petroleum and natural gas	238
petrology	77
stratigraphy	76, 77
structural geology	76, 77, 170
surface water	172
urban geology	242
zinc	16, 235
Viscosity, tektite glass	135
Volcanic rocks, minor elements	222
occurrence	94
petrology and geochemistry	149
remanent magnetization	50
variation in composition with time	105
Volcanic-terrace hydrology, investigations in progress	243
Volcanoes, Costa Rica, geologic study	118
Hawaii, eruption	159
Washington, emission of steam	57
See also Lava lakes.	
Volcanology, investigations in progress	243

W

Walker Lane, Nevada, crustal structure	140
Washington, clays	10, 218
coal	219
construction and terrain problems	57
cooperating agencies	209
estuaries, biohydrology	129
floods	59
geochronology	96
geologic mapping	68, 224, 230
geophysics	232
glacial geology	96
glaciers	180
glaciology	233
ground water	39, 169, 250
ground-water-surface-water relations	172, 233
infrared surveys	146
marine hydrology	236
paleoecology	236
paleontology	96
petroleum and natural gas	238
quality of water	51, 172, 239, 249
radioactive-waste disposal	239
sedimentation	240
stratigraphy	83, 96, 241
structural geology	83, 96
surface water	166, 234
uranium	242
urban geology	242
water resources	249
waterpower resources	243
Water. See Ground water, Quality of water, Surface water.	

	Page
Water chemistry, analytical methods	217
analytical methods. See also Analytical chemistry, results of investigations.	
determination of insecticides and pesticides	182, 185, 217, 219
Water data, office for coordination	18
Water management, investigations in progress	243
Water resources, investigations in progress	243
nuclear explosives in development of	46, 219
Water supply, public and industrial, investigations in progress	239
Waterfalls, effect on stream-flow	21
Waterpower classification, investigations in progress	243
results of investigations	43
Wave velocity, in rock, measurement	145
Waves, sea, earthquake-caused	61, 64
Weathering, effect on U-Th-Pb system in zircon	161
Wells, deep, portable sampler-downward ground-water flow in	21
effect of earthquakes on water level	64, 170
geothermal, as source of data on metamorphism	150
"scavenger" type, for separation of fresh from saline water	171
waste-disposal, rejuvenation	171
water fluctuation in, interpretation	170
West Virginia, cooperating agencies	209
ground water	20, 23, 170, 250
ground-water-surface-water relations	173
surface water	22
water resources	250
Western United States, flood studies	221
heat flow	142
hydrology	243
paleobotany	236
paleontology	123, 237
seismic mantle studies	139
soil-moisture studies	241
thorium	242
waterpower resources	243
See also names of individual States.	
Wilderness Act, resource studies	2
Wilderness and wild areas, waterpower classification	243
Williston basin, stratigraphy	241
Winterfat, relation to geomorphic processes	180

	Page
Wisconsin, cooperating agencies	209
floods	221
geochemistry	222, 223
geologic mapping	230
geophysics	81, 232
ground water	26, 250
ground-water-surface-water relations	173, 233
iron	235
low-flow studies	235
mineral-resource appraisal	236
petrology	223
phosphate	9
sedimentation	240
stratigraphy and structural geology	82
surface water	166
water resources	250
Wolframite, in Saudi Arabia	115
Wood, fossil, from Atlantic Continental Shelf	126
Wyoming, coal	219
cooperating agencies	209
earthquake studies	141
floods	221
geochemistry	223
geologic mapping	68, 224, 230
geomorphology	85, 231
geophysics	231, 232
glaciers	181
gold	7, 233
gravity studies	86
ground water	33, 250
iron	235
mineralogy	150
oil shale	236
paleobotany	126, 127
paleoecology	236
paleontology	121, 122
petroleum and natural gas	238
petrology	223
phosphate	238
plant ecology	180
quality of water	32, 56, 159
saline minerals	240
sedimentation	240
stratigraphy	85, 88, 241
thermal water	159
uranium	12, 13, 16, 242
water resources	250
waterpower resources	243

X

X-ray analysis, Hawaiian olivine	148
X-ray fluorescence analysis, cesium iodide	146
rocks and soils	183
water	185
X-ray quantometer, automation	183

Y

Yellowstone National Park, Wyo., earthquake studies	141
thermal water	159

See also Index to List of Publications

Z	Page
Zeolite zoning, Nevada-----	50
Zeolites, district studies-----	11
investigations in progress--	250
Zeolitic facies, coexisting mineral parageneses--	154
Zinc, determination, optical spectroscopic ----	183
determination, spectrophotometric -----	181, 182
district studies -----	5

Zinc—Continued	Page
in black shale-----	156
investigations in progress -----	235, 236
mercury as an ore guide for -----	16
misleading concentration anomalies -----	17
Zircon, magnetic properties----	146
U-Th-Pb systems in, effect of weathering -----	161

Zircon—Continued	Page
variation in uranium-isotope ratios -----	161
Zoning, biological, relation to ground-water discharge -----	129
in beryllium-fluorite deposits -----	11
in manganese veins, significance -----	4
in uranium deposits-----	13
zeolite, Nevada -----	50

INVESTIGATOR INDEX

A

	Page
Ackroyd, E. A.	33
Adams, J. W.	13, 14
Addicott, W. O.	96, 98, 100, 124
Akers, J. P.	105
Akhrass, N. N.	116
Albers, J. P.	94
Albin, D. R.	29
Alcorn, S. M.	178
Alexander, W. H., Jr.	37
Allen, R. V.	116
Altschuler, Z. S.	9
Anderson, C. A.	94
Anderson, D. B.	39
Anderson, D. G.	68
Anderson, H. W.	39
Anderson, P. W.	53, 166, 167
Anderson, R. E.	49
Anderson, R. J.	113
Andreasen, G. E.	8, 95
Andrews, G. W.	82
Andrews, Laurence	115
Ankary, A. O.	115
Annell, C. S.	135, 183
Antonio, Leonardo	115
Antweiler, J. C.	7, 164
Appleman, D. E.	151, 152
Archer, R. J.	157
Armstrong, C. A.	33
Armstrong, F. C.	12, 13, 56
Arndt, H. H.	14
Arnold, D. C.	5
Arnow, Ted	34, 105
Aro, R. S.	180
Averitt, Paul	14, 118

B

Bachman, G. O.	87
Back, William	55, 158, 159
Bailey, E. H.	3, 97
Bailey, J. F.	187
Bailey, R. A.	138, 149
Baker, C. H., Jr.	86
Baker, E. T., Jr.	37
Baker, Roger	25
Bakr, M. A.	113
Ballance, W. C.	36
Baltz, E. H.	87
Baltzer, R. A.	129
Barclay, J. E.	36
Barker, Fred	4
Barnes, D. F.	62, 63
Barnes, I. K.	114
Barnett, P. R.	183
Barton, P. B., Jr.	7
Bastron, Harry	147, 148
Bateman, P. C.	99
Bath, G. D.	50
Batson, R. M.	136
Becher, A. E.	170
Beck, M. E., Jr.	78, 144
Beckman, Emil	59
Becraft, G. E.	83
Bedinger, M. S.	174
Beetem, W. A.	47

Page

Behrendt, J. C.	89
Bell, E. A.	29, 30
Bell, K. G.	73
Bennett, R. R.	170
Benson, A. L.	85
Bentley, C. B.	36
Berdan, Jean	70, 124
Bergendahl, M. H.	57
Bergin, M. J.	76
Bergquist, Harlan	70
Bettendorf, J. A.	61
Bettiga, A. C.	184
Beverage, J. P.	176
Bidwell, L. E.	172
Biehler, Shawn	133
Bjorklund, L. J.	35
Black, D. F. B.	80
Blackmon, P. D.	79
Blade, L. V.	112
Blank, H. R., Jr.	117, 118
Bogart, D. B.	105
Boggess, D. H.	22
Bole, J. B.	186
Books, Kenneth	81
Booth, J. A.	178
Boswell, E. H.	31
Boucot, A. J.	74, 75
Boudette, E. L.	74, 107
Bowles, C. G.	12, 80
Boyd, F. R.	151
Boynton, G. R.	4
Brabb, E. E.	103, 122
Braddock, W. A.	87
Bradley, W. H.	126, 150
Brannock, W. W.	148, 184
Branson, F. A.	173, 179, 180
Bredehoeft, J. D.	169, 170
Breger, I. A.	182
Bremaecker, J. C. de	145
Brennen, Robert	176
Brew, D. A.	104, 148
Briggs, R. P.	105
Brim, R. J. P.	111
Brobst, D. A.	116, 184
Brock, Maurice	4, 162
Brokaw, A. L.	92
Bromery, R. W.	73, 74
Broom, M. E.	37
Brosge, W. P.	103
Brown, C. E.	9, 95
Brown, T. E.	188
Bryant, Bruce	88
Buchanan, T. J.	167
Buckman, R. C.	74
Buddington, A. F.	3
Buller, William	173
Bunker, C. M.	17
Burns, C. V.	168
Burton, L. C.	37
Busby, M. W.	168
Byers, F. M.	49

C

Cadigan, R. A.	90
Cady, W. C.	96

Page

Cahill, J. M.	170
Calkins, J. A.	113
Campbell, D. F.	112
Campbell, R. H.	12
Canney, F. C.	17
Cannon, H. L.	56, 156
Cannon, R. S., Jr.	164
Cardwell, W. D.	119
Carey, K. L.	166
Carlston, C. W.	170, 178
Carpenter, D. H.	176
Carr, M. H.	132, 134
Carr, W. J.	138
Carrigan, P. H., Jr.	52
Carroll, R. D.	45
Carswell, L. D.	170
Case, J. E.	4, 86, 90, 91
Castle, R. O.	72
Catanzaro, E. J.	162
Cater, F. W.	96
Cathcart, J. B.	9
Cederstrom, D. J.	20, 112
Chamberlain, A. R.	50
Chang, F. M.	167
Chao, E. C. T.	134, 135
Chase, G. H.	95
Chase, R. L.	128
Chemerys, J. C.	53
Cherry, R. N.	24, 158, 186, 187
Chidester, A. H.	138
Chorley, R. J.	178
Christ, C. L.	7, 152, 155, 159
Christiansen, R. L.	138
Churkin, Michael	97, 103
Clark, D. L.	121
Clark, J. R.	151, 152
Clark, W. E.	172
Clarke, F. E.	54, 114
Clifton, H. E.	98
Coates, D. A.	106
Coats, R. R.	93
Cobb, E. D.	187
Cobban, W. A.	91, 126
Cohen, Philip	39, 40
Colby, B. R.	175, 176
Coleman, R. G.	149, 150
Collier, C. R.	27
Collings, M. R.	173
Colton, G. W.	79
Conant, L. C.	71
Condon, W. H.	103
Conklin, N. M.	13
Coombs, D. S.	155
Cooper, H. H., Jr.	170
Cooper, J. R.	4
Corwin, Gilbert	129
Cory, R. L.	129
Cosner, O. J.	25
Cotter, R. D.	172
Coulter, H. W.	61
Cox, Allan	99, 143, 144
Cox, E. R.	55
Crabtree, James	116
Craig, L. C.	69
Cramer, W. G.	1
Crandell, D. R.	57, 96, 146

See also List of Publications

	Page
Creasey, S. C.	5, 93
Cressler, Charles	23, 77
Cressman, E. R.	80
Crippen, J. R.	167
Crispin, Oscar	114
Crist, M. A.	33
Crittenden, M. D., Jr.	93, 94, 104
Croft, M. G.	99
Crooks, J. W.	23, 105
Crosthwaite, E. G.	38
Crowder, D. F.	99, 106
Culbertson, J. K.	175
Culler, R. C.	176
Cullins, H. L.	15
Cumming, G. L.	141
Cuppels, N. P.	73
Cushman, R. V.	29
Cuttitta, Frank	135, 184

D

Dalrymple, G. B.	99, 143, 161
Dane, C. H.	91
Davidian, Jacob	165
Davidson, D. F.	15
Davies, W. E.	69
Davis, D. A.	42
Davis, L. C.	111
Davis, R. E.	45
Davis, R. W.	29
Davis, W. E.	116
Dawdy, D. R.	175
Dearborn, L. L.	64
DeCarli, P. S.	133
DeLaney, A. O.	76
DeLuca, F. A.	56
Dempsey, W. J.	114
Dempster, G. R., Jr.	51
Dennen, W. H.	17, 75
Denny, C. A.	69
Denny, M. V.	10
Detterman, R. L.	103, 104
Deutsch, Morris	20
Dibblee, T. W., Jr.	94
Dickey, D. D.	46
Dickinson, K. A.	71
Dickinson, R. G.	88, 91
Dickson, K. O.	149
Diment, W. H.	142
Dingman, R. J.	112, 114
Dinnin, J. I.	147
Dodge, F. C.	154
Doe, Bruce	161, 164
Doell, R. R.	99, 117, 143, 144
Donaldson, D. E.	174
Donnell, J. R.	91
Dooley, J. R., Jr.	163
Dorr, J. V. N. 2d.	111
Dosch, T. R.	26
Dove, G. D.	20, 28
Downs, S. C.	184
Drake, A. A., Jr.	10, 107
Drake, P. G.	55
Ducet, G. L.	39
Duke, M. B.	133, 135
Duncan, H. M.	122
Dunn, Bernard	187
Dunnam, C. A.	176
Dunrud, C. R.	57
Dunton, P. J.	184, 185
Durham, D. L.	98
Dutcher, L. C.	41
Dutro, J. T., Jr.	96, 103
Dyer, C. F.	34

E

	Page
Eakin, T. E.	39
Eargle, D. H.	11, 13
Eaton, G. P.	95
Eaton, J. P.	141
Eddy, J. E.	186
Edwards, K. W.	163, 183, 185
Edwards, M. D.	187
Ege, J. R.	45, 106
Eggleton, R. E.	131
Eisenlohr, W. S., Jr.	174
Ekren, E. B.	49
Ellis, M. J.	34
Ellison, R. L.	170
Elmore, Paul	89
Elston, D. P.	138
Emerick, W. L.	45
Emery, K. O.	127
Emmett, W. W.	177
Engel, A. E. J.	72, 148, 154
Engel, C. G.	72, 148, 154
England, J. L.	151
Englund, K. J.	76, 81
Erd, R. C.	6, 128, 149, 155
Erickson, George	112
Erickson, R. L.	16, 94
Ericson, D. W.	173
Espenshade, G. H.	75
Evans, H. T., Jr.	152
Evenson, R. E.	41
Everett, D. E.	39
Evet, W. W.	167

F

Fahnestock, R. K.	57
Farah, Abul	114
Farlekas, G. M.	61
Farrell, D. F.	167
Faust, S. D.	53
Feininger, Tomas	112
Feltis, R. D.	34
Fennelly, Edward	182
Ferguson, H. G.	6
Ferrians, O. J., Jr.	62, 181
Ferriera, C. P.	163
Feulner, A. J.	64
Ficke, J. F.	177
Fidler, R. E.	169
Finnell, T. L.	12
Fisher, D. W.	157, 186
Fisher, R. P.	123
Fishman, M. J.	184, 185
Fleischer, Michael	157
Flint, R. F.	27
Follett, C. R.	55
Foot, R. S.	17
Ford, A. B.	107
Forman, J. M. A.	112
Forrest, W. E.	166
Foster, H. L.	62, 125, 181
Foster, J. B.	24
Foster, J. F.	56
Fournier, R. O.	85, 154
Fox, K. F.	96
Frederick, B. J.	187
Freeman, V. L.	82
French, J. J.	41
Freydanck, H. G.	111
Friel, E. A.	22
Fritts, C. E.	4, 81
Frost, I. C.	156
Furness, L. W.	168

G

	Page
Gabrysch, R. K.	58
Gaca, J. R.	89, 142
Gair, J. E.	3
Gallaher, J. T.	29
Gambell, A. W.	157
Garber, M. S.	47
Gard, L. M.	46
Gardner, K. A.	184
Gardner, L. S.	116
Gardner, R. A.	22
Garrels, R. M.	159
Gault, D. E.	134
Gawarecki, S. J.	119
Gazin, C. L.	85
Geddes, W. H.	105
Geruasio, Froilan	114
Ghering, G. E.	173
Gibbons, A. B.	12
Gilbert, B. K.	21
Gill, H. E.	69, 112
Gill, J. R.	85, 126
Gillespie, J. B.	36
Gilluly, James	6
Glenn, J. L.	51
Glover, R. E.	50
Godson, R. H.	145
Godwin, L. H.	89
Goerlitz, D. F.	185
Goldberg, M. C.	53
Goldich, S. S.	161
Goldsmith, Richard	115
Gordon, Mackenzie, Jr.	123
Gott, G. B.	15
Gottfried, David	147, 149
Granger, H. C.	13
Grant, R. E.	122, 123
Grantz, Arthur	63, 102
Green, A. R.	60
Gregg, D. O.	23, 55
Griffitts, W. R.	10
Griscom, Andrew	105
Grossling, B. F.	144
Grozier, R. U.	167
Gualtieri, J. L.	80
Gude, A. J., 3d.	11, 150
Gulbrandsen, R. A.	4
Gutentag, E. D.	36
Guy, H. P.	186

H

Habib-ur-Rahman	113
Hack, J. T.	178
Hackman, R. J.	91
Hadley, R. F.	177
Haffty, Joseph	182
Hahl, D. C.	177
Hail, W. J.	88
Hakim, Hashim	115
Hale, W. E.	47
Haley, B. R.	14, 82
Hall, C. W.	53
Hamilton, J. C.	16
Hamilton, L. J.	26
Hamilton, Warren	106, 148
Hamlin, H. P.	10
Hanes, M. E.	113
Hannum, C. H.	166
Hansen, A. J., Jr.	29
Hansen, H. J.	22
Hanshaw, B. B.	55, 155, 158, 159
Harbert, Gene	116
Harder, A. H.	31
Harding, W. E.	157
Hardt, W. F.	54, 171
Harris, L. D.	77

See also List of Publications

	Page
Harshman, E. N.	16
Hartshorn, J. H.	72, 73, 74
Hartwell, J. H.	24
Harvey, E. J.	28
Harwood, D. S.	74
Hatch, N. L.	73
Hatchett, J. L.	33
Hathaway, J. C.	126, 128
Hattori, Osamu	35
Haushild, W. H.	51
Havens, J. S.	55
Hawley, C. C.	10
Hayashi, Shoichiro	11
Hazel, J. E.	125
Healey, D. L.	49
Heare, J. T.	113
Hearn, B. C., Jr.	84
Hedge, C. E.	107, 162
Hedges, L. S.	34
Hedman, E. R.	41
Heidrick, T. L., Jr.	92
Heitowit, E. D.	134
Helz, A. W.	183
Hem, J. D.	159
Hemley, J. J.	153
Henbest, L. G.	83
Henderson, R. G.	145
Hendricks, T. A.	14
Herrick, S. M.	70, 71
Hewett, D. F.	2, 4, 7
Heyl, A. V.	16, 155, 162
Hickling, N. L.	147
Hicks, S. D.	104
Hill, D. P.	140, 141, 161
Hines, M. S.	26, 28
Hirashima, G. T.	68, 173
Hite, R. J.	9, 90
Hoare, J. M.	103, 126
Hodges, A. L., Jr.	20
Hoffard, S. H.	42
Hoffman, C. M.	157
Hoffman, J. F.	56
Hoggatt, R. E.	28
Holden, W. R.	31
Hollyday, E. F.	20
Holmes, G. W.	73, 181
Holt, C. L. R., Jr.	173
Holt, H. E.	136
Hoover, D. L.	50
Hopkins, D. M.	102, 117, 143, 181
Hopkins, H. T.	29
Hopson, C. A.	96
Horner, Lee	179
Horton, G. W.	91
Horwitz, G. M.	166, 167
Hosman, R. L.	25
Hosterman, J. W.	10, 127
Hotz, P. E.	6
Howells, Louis	34
Hoyte, A. F.	135
Hubbell, D. W.	51, 167, 175
Huber, N. K.	148
Hudson, J. D.	36
Huffman, Claude, Jr.	157, 182
Hughes, L. S.	37, 54
Hulsemann, Jobst	128
Hummel, C. S.	115
Humphreys, C. P., Jr.	48
Hunn, J. D.	28
Hutchinson, R. D.	26
Hutton, C. O.	118
Huxel, C. J.	40
Hylckama, T. E. A. van	174

I	Page
Imlay, Ralph	124
Ingram, B. L.	183
Irving, E. M.	112
Irwin, W. P.	97

J	Page
Jackson, E. D.	138, 149
Jackson, R. O.	113
Jackson, W. H.	140
Jafree, A. R.	114
Janda, R. J.	178
Janson, M. E.	26
Jenkins, C. T.	35
Jenkins, E. D.	35
Jenne, E. A.	52
Joesting, H. R.	68, 91
Johnson, A. I.	170
Johnson, C. G., Jr.	60
Johnson, G. R.	146
Johnson, J. O.	185
Johnson, Lane	140
Johnson, R. B.	89
Johnston, J. E.	70
Jolly, J. L.	16
Jones, B. F.	158, 185, 186
Jones, B. L.	176
Jones, D. L.	97, 102
Jordan, D. G.	25
Jordan, P. R.	20, 26
Joyner, B. F.	24
Jungmann, W. L.	36

K	Page
Kachadoorian, Reuben	63, 134
Kahn, Lloyd	52
Kam, William	58
Kammerer, J. C.	21
Kantrowitz, I. H.	21
Karig, D. E.	89
Karklins, O. L.	122
Karlstrom, T. N. V.	62, 104
Kaye, C. A.	73
Keech, C. F.	34
Keefer, W. R.	86
Keighton, W. B.	129
Keller, F. J.	21
Kennedy, V. C.	187, 188
Kennon, F. W.	168
Kent, B. H.	78
Kepferle, R. C.	80
Kier, P. M.	122
Kiilsgaard, T. H.	3
Kilburn, Chabot	25
Kilpatrick, F. A.	187
Kimrey, J. O.	9
King, N. J.	33
King, P. B.	68, 69
Kinney, D. M.	69
Kinorason, Thorleifur	117
Kistler, R. W.	99
Klausing, R. L.	33
Klein, Howard	24
Kleinhampl, F. J.	93
Klepper, M. R.	84, 164
Knowles, D. B.	171
Knox, A. S.	71
Knox, C. E.	60
Koberg, G. E.	58
Koch, N. C.	34
Kohout, F. A.	129
Kolipinske, M. C.	129
Koopman, F. C.	47, 52, 169
Koteff, Carl	74

L	Page
Kouther, Jameel	115
Krieger, M. H.	93
Krinslev, D. B.	181
Kunkle, G. R.	172

L	Page
Lachenbruch, A. H.	142, 143
La Fehr, T. R.	142
La Freniere, G. F.	41
Laguna, Wallace de	51
Lai, Chintu	129
Laine, L. L.	166
Lakin, H. W.	1, 7, 15, 16, 17
Lamar, W. L.	185
LaMarche, V. C., Jr.	175
Lambert, T. W.	25, 29
Lamke, R. D.	40
LaMoreaux, P. E.	171
Landis, E. R.	91
Laney, R. L.	157
Lang, S. M.	187
Langford, R. H.	177
Lanphere, M. A.	97, 101, 103, 157, 161
LaSala, A. M., Jr.	157
Lathram, E. H.	101, 104
Laurence, R. A.	78
Law, A. G.	173
Law, L. M.	185
Leach, S. D.	187
Lee, D. E.	147, 149
Lee, F. T.	57
LeGrand, H. E.	52
Leifeste, D. K.	54
Lemke, R. W.	61
Leonard, B. F.	3, 5, 6, 84
Leonard, R. B.	35
Leopold, E. B.	125, 126
Leopold, L. B.	177
Leppanen, O. E.	174
Leveen, L. S.	64
Lewis, G. E.	91
Lewis, R. Q.	12
Lewis, R. W., Jr.	111
Limbach, Dora von	153
Lindberg, M. L.	11, 152
Lipscomb, R. G.	177
Lobmeyer, D. H.	36
Lofgren, B. E.	58
Lohman, K. E.	127
Loney, R. A.	104
Long, A. T.	25
Long, R. A.	171
Lovering, T. G.	16
Lowe, A. S.	60
Lowry, M. E.	33
Lubke, E. R.	41, 56
Ludwig, A. H.	170
Lugn, R. V.	135

M	Page
Mabey, D. R.	95
McAllister, J. F.	8, 123
McCall, J. E.	167
McCarthy, J. H., Jr.	15, 16
McCauley, J. F.	131, 137
McClymonds, N. E.	24
McCollum, M. J.	23
McCoy, H. J.	24
McCulloch, D. S.	61, 102
McCullough, R. A.	169, 173
MacDonald, W. R.	197
McGovern, H. E.	35
McGreevy, L. J.	34
Mack, F. K.	22

See also List of Publications

	Page
McKee, E. D.	177
McKee, E. H.	97
MacKevett, E. M., Jr.	104
McKnight, E. T.	123
McLaughlin, Donald	112
McMaster, W. M.	167, 168
McMurtrey, R. G.	32
MacNeil, F. S.	104, 117, 125
McNellis, J. M.	187
McQueen, J. S.	173, 174
MacQuown, W. C.	80
Maddox, G. E.	36
Maderak, M. L.	26, 32
Maher, J. C.	127
Malcolm, R. L.	188
Malde, H. E.	178
Malmberg, G. T.	40
Mamay, S. H.	126
Mapel, W. J.	121
Marcher, M. V.	38, 64
Marie, J. R.	-5, 31
Marinenko, John	183
Marlatt, W. E.	146
Martens, L. A.	167
Martinez, Prudencio	146
Marvin, R. F.	74, 89, 93, 106, 107, 162
Massoni, Camillo	183
Masursky, Harold	132
Matalas, N. C.	179
Mattoso, S. Q.	111
Matzko, J. J.	114
Maxwell, C. H.	80
May, Irving	182, 183
Mayhew, M. A.	142
Meade, R. H.	128
Meier, M. F.	180
Meisler, Harold	170
Meissner, C. R.	113
Meister, Robert	145
Mello, J. F.	122
Mensik, J. D.	182
Merriam, C. W.	92
Merrill, A. S.	128
Metzger, D. G.	32
Meyer, F. W.	24
Meyer, W. R.	36
Meyers, B. N.	37
Meyers, J. S.	174
Meyrowitz, Robert	182
Michel, F. C.	133
Miesch, A. T.	157
Migliaccio, R. R.	61
Miller, C. H.	49
Miller, E. G.	168, 174
Miller, R. F.	173, 174, 180
Miller, R. L.	76
Miller, T. P.	102
Milton, D. J.	131, 133
Minard, J. P.	70
Mirza, M. A.	114
Mitten, H. T.	159, 177
Moench, R. H.	74
Monk, E. J.	145
Moore, A. M.	59, 176
Moore, D. O.	39, 40, 168
Moore, G. K.	171
Moore, G. W.	63, 128
Moore, H. J.	131, 134
Moore, J. E.	35, 158
Moore, W. J.	94
Morey, G. W.	85, 155
Morgan, C. O.	187
Morris, D. A.	95, 170, 171
Morris, E. C.	136
Morris, H. T.	94
Morrison, C. E.	197
Moston, R. P.	170

	Page
Mountjoy, Wayne	148
Mower, R. W.	34
Moxham, R. M.	17, 119, 146
Mrose, Mary	112, 183
Mudge, M. R.	84
Muffler, L. J. P.	148, 150
Muir, K. S.	41
Mullineaux, D. R.	57, 96
Mundorff, M. J.	38
Murata, K. J.	118, 128, 148
Murphy, J. F.	85
Musgrove, R. H.	56
Musser, J. J.	170, 173
Myers, D. A.	87
Myers, W. B.	84
Myrick, R. M.	173, 176, 177
Mytton, J. W.	115

N

Nagell, R. H.	114
Nakagawa, H. W.	1, 7, 17
Nakahara, R. H.	41
Neely, B. L., Jr.	61
Nelson, A. E.	105
Nelson, C. A.	92
Nelson, W. H.	106
Neuerburgh, G. J.	16
Neuman, R. B.	75, 122
Neuschel, S. K.	78, 82
Newcomb, R. C.	169
Newcome, Roy, Jr.	30, 31
Newell, M. F.	161
Newton, J. G.	30
Nicholson, S. B.	132
Noble, D. C.	149
Nordin, C. F., Jr.	175, 176
Norris, S. E.	169
Norvitch, R. F.	21
Nyman, D. J.	30

O

Obradovich, J. D.	148
O'Bryan, Deric	22, 179
Offield, T. W.	113
Ogata, Okio	51
Ogilbee, William	113
Olcott, P. G.	26
Oldale, R. N.	74
Olin, George	178
Oliver, H. W.	99, 100
Oliver, W. A., Jr.	122
Olsen, H. W.	169
Olson, J. C.	3
Oriel, S. S.	83
Osterkamp, W. R.	33
Osterwald, F. W.	57
O'Sullivan, R. B.	15
Outerbridge, W. F.	79
Overstreet, W. C.	3
Owens, J. P.	70

P

Pakiser, L. C.	139, 140, 142
Palacas, J. G.	15, 128
Palmer, A. R.	73, 106, 122, 124
Pampeyan, E. H.	6
Papadopoulos, I. S.	169, 170
Papike, J. J.	151
Paradeses, W. D.	70
Pardee, J. T.	83
Parker, G. G.	175
Parker, O. M.	182
Parker, R. L.	86

	Page
Parshall, E. E.	169
Patton, W. W., Jr.	102
Pavrides, Louis	75
Payne, J. N.	171
Pearson, R. C.	83
Pease, M. H., Jr.	105
Peck, D. L.	97
Peck, J. H.	80
Peirce, L. B.	30, 60
Pelletier, J. D.	5
Peselnick, Louis	145
Peterman, Z. E.	106, 172
Petersen, M. S.	166
Peterson, D. E.	58
Peterson, Fred	91
Peterson, W. L.	80
Petrafeso, F. A.	8
Pettit, Edison	132
Pettyjohn, W. A.	33
Phipps, R. L.	179
Pickering, R. J.	52
Pierce, A. P.	164
Pierce, K. L.	75
Pillmore, P. L.	14
Piper, A. M.	46, 65
Pipiringos, G. N.	88
Plaffer, George	61, 63, 64, 104
Plouff, Donald	91
Pohn, H. A.	132, 136
Poland, J. F.	41
Pollock, S. J.	21
Pomeroy, J. S.	80
Porterfield, George	176
Post, A. S.	181
Post, E. V.	17, 75
Poth, C. W.	170
Powell, W. J.	171
Powers, H. A.	138
Prather, T. L.	45
Pratt, R. M.	128
Price, Donald	39
Price, W. E., Jr.	29
Prichard, G. E.	82
Prill, R. C.	170
Prinz, W. C.	4
Prych, E. A.	51
Purtyman, W. D.	52, 169

Q

Quinn, A. W.	72
--------------	----

R

Radtke, Arthur	7
Randich, P. G.	33
Rankin, D. W.	77
Rantz, S. E.	59, 176
Raspet, R.	142
Ratté, J. C.	86
Rau, J. L.	27
Raup, O. B.	8
Rawson, Jack	37
Ray, L. L.	79
Reed, B. L.	103, 104
Reed, D. E.	197
Reed, J. C., Jr.	181
Reed, J. E.	173, 174
Reiser, H. N.	103
Repenning, C. A.	125
Reso, Anthony	123
Rettig, S. L.	186
Reyes, Benjamin	170
Reynolds, M. W.	86
Richardson, E. V.	165, 176
Richmond, G. M.	85, 87, 89
Riggs, H. C.	168, 172
Riley, L. B.	182

See also List of Publications

	Page
Rinehart, C. D.	96, 148
Roach, C. R.	146
Roberts, R. J.	1, 5, 6, 94
Robertson, E. C.	145
Robertson, J. B.	53, 154
Robertson, J. F.	76
Robie, R. A.	153
Robinson, C. S.	57
Robinson, G. B., Jr.	35
Robinson, G. D.	84
Robinson, T. W.	174
Roddy, D. J.	133
Roedder, Edwin	155, 156
Roen, J. B.	76
Rogers, C. L.	49
Rogers, S. M.	30, 31
Rohrer, W. L.	85
Roller, J. C.	140
Rorabaugh, M. I.	172
Rose, H. J.	184
Roseboom, E. H.	8, 151
Rosenau, J. C.	41
Rosholt, J. N., Jr.	163
Ross, D. C.	92
Ross, Malcolm	151, 152
Ross, R. J., Jr.	122
Rossmann, Darwin	115
Rowan, L. C.	136, 137
Rowe, J. J.	85, 154, 155, 182
Rubin, Meyer	55, 82, 96, 99, 102, 104, 117, 125, 128, 158, 162
Ruppel, E. T.	83
Rush, F. E.	39
Rusnak, G. A.	61
Rye, R. O.	164

S

Sabine, C. P.	84
Sable, E. G.	100
Sachs, K. N., Jr.	128
Sainsbury, C. L.	11, 102
Sandberg, C. A.	80, 85, 150
Santos, A. M.	111
Santos, J. F.	129
Sargent, K. A.	49, 50
Sato, Motoaki	7
Sayre, W. W.	50, 167, 175
Schafer, J. P.	72, 117
Scheidegger, A. E.	177
Schell, E. M.	9
Schell, W. W.	121
Schlanger, S. O.	80
Schlee, J. S.	128
Schmidt, D. L.	106
Schmidt, R. A. M.	63
Schmidt, R. G.	84
Schmidt, H. H.	138
Schnabel, R. W.	73
Schneider, W. J.	19
Schoen, Robert	150
Schopf, J. M.	125
Schroeder, E. E.	168
Schroeder, M. E.	27
Schroeder, M. L.	121
Schumm, S. A.	117, 178
Schweinfurth, J. P.	78
Scott, G. R.	88
Scott, J. H.	45, 145
Scott, R. A.	126
Scott, R. C.	56, 158, 163
Seaber, P. R.	20
Seevers, W. J.	54
Seiders, V. M.	79, 105
Seitz, J. F.	111
Senftle, F. E.	135, 146, 147
Sever, C. W.	23, 69
Shacklette, H. T.	17

Shapiro, Leonard	183
Sharp, W. N.	1, 11, 12
Sharps, J. A.	82
Shaw, H. R.	153
Shawe, D. R.	10, 11, 13, 86
Shell, J. D.	166
Shepard, A. O.	50
Sheppard, R. A.	11, 150
Sheridan, D. M.	13, 87
Sheridan, M. F.	99
Shoemaker, E. M.	136, 138
Shown, L. M.	179
Sigafoos, R. S.	179
Signer, P.	134
Silber, C. C.	155
Silberling, N. J.	123
Simmons, G. C.	123
Simon, F. O.	183
Simons, D. B.	165, 176, 186
Simons, F. S.	93
Simons, W. D.	172
Sims, P. K.	13
Singewald, Q. D.	13, 88
Siple, G. E.	23, 70
Skelton, John	28
Skibitzke, H. E.	170, 171
Skinner, B. J.	8
Skipp, B. A.	121
Skougstad, M. W.	185
Slack, K. V.	54, 186, 187
Slaughter, M. J.	64
Smedes, H. W.	84
Smith, H.	89
Smith, J. F.	91
Smith, P. J.	63, 98
Smith, R. C.	116
Smith, R. L.	99, 149
Smith, R. M.	3
Smith, W.	186
Smith, W. C.	9
Smoot, G. F.	186
Snavely, P. D., Jr.	97
Snyder, G. L.	74
Sohn, I. G.	91, 100, 124
Sommers, D. A.	38
Soren, Julian	71
Soza, E. R.	197
Speer, P. R.	26
Spieker, A. M.	27
Staat, M. H.	13, 14
Stacy, J. D.	91
Stallman, R. W.	38, 169, 173
Stanin, S. A.	114
Stead, F. W.	46
Steele, C. E.	36
Steinhart, J. S.	139
Stephens, H. D.	32
Stephens, H. G.	138
Stephens, J. C.	34
Stern, T. W.	74, 84, 93, 107, 147, 161, 162
Stevens, H. H., Jr.	51
Stevens, P. R.	54
Stewart, D. B.	107, 153
Stewart, G. L.	157, 165
Stewart, J. H.	92
Stewart, J. W.	51
Stratton, Garland	185
Stringfield, V. T.	20
Stromquist, A. A.	77
Stuart, W. T.	53
Sturrock, A. M., Jr.	174
Sundelius, H. W.	77
Sutcliffe, Horace, Jr.	24
Sutton, R. L.	91
Swann, G. A.	138
Swanson, V. E.	15, 128

T

	Page
Tabor, R. W.	96
Tagg, Richard	127
Tailleux, I. L.	15, 102
Takasaki, K. J.	41
Takeda, Hiroshi	151
Tanaka, H. H.	174
Tangborn, W. V.	180
Tanner, A. B.	162
Tatlock, D. B.	135
Tatsumoto, M.	164
Taylor, A. R.	77
Taylor, C. T.	28
Taylor, Charles	7
Taylor, D. W.	9, 102, 124
Taylor, O. J.	32
Taylor, R. E.	48
Taylor, Thomas	116
Tedford, R. H.	9
Temple, K. L.	159
Thaden, R. E.	12
Thayer, T. P.	95
Theobald, P. K., Jr.	115
Thomas, D. M.	61
Thomas, H. H.	89, 107
Thomas, J. D.	54
Thompson, C. E.	115
Thompson, G. A.	150
Thompson, J.	147
Thorpe, A. N.	135, 147
Tilling, R. I.	148, 164
Titely, S. R.	131
Toler, L. G.	56, 172
Tooker, E. W.	94
Toulmin, Priestley 3d	7, 154
Tourtelot, H. A.	15, 156, 164
Trace, R. D.	122
Tracey, J. I., Jr.	129, 130
Trainer, F. W.	21, 63, 170
Trapp, Henry, Jr.	33
Trimble, D. E.	83
Trites, A. F., Jr.	12
Truesdell, A. H.	155, 185
Trumbull, J. V. A.	128
Tryba, B. A.	135
Tryggvason, Eysteinn	141
Tschanz, C. M.	111
Tschudy, Bernadine	126
Tschudy, R. H.	75, 126
Turcan, A. N., Jr.	25, 158
Turner, R. M.	178
Turner, W. B.	80
Tuttle, C. R.	145
Tweto, Ogden	87, 90

U

Uchupi, Elazar	127
----------------	-----

V

Valenciano, Santos	41
Van Alstine, R. E.	89
Van Denburgh, A. S.	158, 172, 185
Vargo, J. L.	4
Vaughn, W. W.	1
Vaupel, D. E.	187
Vecchioli, John	22
Vedder, J. G.	98
Vine, J. D.	156
Visher, F. N.	23
Vloten, Roger van	113
Vorhis, R. C.	64, 65, 170

See also List of Publications

W	Page
Waananen, A. O.	173
Wahl, K. D.	170
Wahl, R. R.	95
Wahlberg, S. S.	52
Wait, R. L.	55
Waldbaum, D. R.	153
Waldron, H. H.	96, 118
Waldrop, H. A.	91
Walker, E. H.	38, 51
Wallace, J. C.	20
Waller, R. M.	63, 64
Walthall, F. G.	107
Wang, Y.	141
Ward, P. E.	42
Warren, D. H.	140
Warren, J. D.	24
Wasson, B. E.	31
Waters, A. C.	138
Watkins, J. S.	78, 138, 145
Watson, Kenneth	132, 137
Wayland, R. G.	14
Wayman, C. H.	53, 154
Weaver, J. D.	142
Webb, W. E.	22
Wedow, Helmuth, Jr.	5, 76
Weeks, A. D.	12, 13
Weeks, E. P.	173
Weigle, J. M.	21, 169
Weir, G. W.	80

	Page
Welborn, C. T.	168
Welch, E. B.	129
Welder, G. E.	34
Wells, J. D.	84
Werre, R. W.	142
Wershaw, R. L.	53
White, D. E.	6, 150
White, N. D.	171
White, S. R.	135
White, Sherman	147
White, W. S.	81
Whitlow, J. W.	115, 116
Whitman, H. M.	31
Whitmore, F. C., Jr.	71, 125
Whittington, C. L.	100
Wilder, H. B.	158, 171
Wilhelms, D. E.	131, 132
Willden, Ronald	140
Williams, J. R.	21, 166
Williams, P. L.	12, 19, 106
Wilmoth, B. M.	22
Wilshire, H. G.	134
Wilson, Druid	124
Wilson, J. F.	166
Wilson, K. V.	61
Wilson, M. T.	177
Winner, M. D., Jr.	31
Winograd, I. J.	39
Wires, H. O.	186
Wirtz, P. W.	135
Withington, C. F.	57

	Page
Witkind, I. J.	12
Wolfe, E. W.	96
Wolfe, Jack	103, 126
Wones, D. R.	151, 154
Wood, G. H., Jr.	76
Wood, P. R.	37
Woodward, T. H.	187
Wright, T. L.	107, 153
Wrucke, C. T.	89
Wyerman, T. A.	157
Wyrick, G. G.	171

Y

Yamamoto, Yukio	138
Yamanaga, George	42
Yochelson, E. L.	144
Yotsukura, Nobuhiro	165
Young, E. J.	9

Z

Zartman, R. E.	74, 162
Zavis, Alfred	197
Zeldan, R. H.	116
Zimmerman, E. A.	32
Zimmerman, R. C.	174
Zoltai, T.	151
Zen, E-an	153, 154, 155