CDF/PUB/ELECTROWEAK/PUBLIC/7725

Version 1.1

June 30, 2005

Proceedings of DIS05

Madison, Wisconsin, April 27-May 1, 2005

W Asymmetry and Z Rapidity Measurements

Y. S. Chung

Department of Physics and Astronomy, University of Rochester, Rochester, New York 14627-0171

Abstract. The measurements of W charge asymmetry and Z rapidity distributions are inputs to constrain the parton distribution functions at high Q^2 . The CDF and DØ experiments at the Tevatron have analyzed up to 340 pb^{-1} of Run 2 data to measure the W charge asymmetry and Z rapidity. The measurements are in generally good agreement with predictions.

Keywords: asymmetry rapidity PDF

PACS: 12.38.Qk

INTRODUCTION

The W and Z bosons are primarily produced by a quark and anti-quark annihilation in $p\bar{p}$ collisions at the Tevatron. Many precision measurements at the Tevatron are limited by the uncertainties on the parton distribution functions (PDFs). These include the W mass, cross section, transverse momentum distributions of W and Z bosons, and so on. For the W mass measurement in the CDF experiment, the current systematic uncertainty from PDF uncertainties is about $15 \text{ MeV}/c^2$ [1] and will become more significant as data accumulated.

The measurement of the W charge asymmetry provides input to the ratio of u and d quark components of PDFs, especially at the medium/high x . In the $p\bar{p} \rightarrow Z$ process, the momentum fraction of a parton is related to the Z rapidity (y)¹. Analysis of the high y region probes the PDFs at both high and low momentum fraction.

W ASYMMETRY MEASUREMENT

The $W^+(W^-)$ bosons at the Tevatron are produced primarily by the annihilation of $u(d)$ quarks in the proton and $\bar{d}(\bar{u})$ quarks in the anti-proton. Because u quarks carry, on average, more momentum than d quarks, the W^+ tend to follow the direction of the incoming proton and the W^- that of the anti-proton.

The $W^\pm$ charge asymmetry

$$A(y_W) = \frac{d\sigma_+/dy_W - d\sigma_-/dy_W}{d\sigma_+/dy_W + d\sigma_-/dy_W}, \quad (1)$$

where the subscript (+,-) denotes the charge of the W and y_W is the rapidity of the W bosons is a sensitive probe of the momentum fraction difference between u and d quarks

¹ $y = \frac{1}{2} \ln \frac{(E+P_z)}{(E-P_z)}$

in the $Q^2 \approx M_W^2$ region. A precise measurement of the W asymmetry serves as a valuable constraint on the u and d quark momentum distributions. For $p\bar{p}$ collisions in leading-order parton model, $A(y_W)$ is given approximately by

$$A(y_W) \approx \frac{u(x_1)d(x_2) - d(x_1)u(x_2)}{u(x_1)d(x_2) + d(x_1)u(x_2)}, \quad (2)$$

where $x_{1,2} = x_0 e^{\pm y_W}$, $x_0 = M_W/\sqrt{s}$, and s is the center of mass energy. The asymmetry, $A(y_W)$, is related to the slope of d/u at $Q^2 = M_W^2$ and in the moderate x region of W production.

The rapidity of the W boson is not measured because the longitudinal momentum of the neutrino can not be experimentally determined. Therefore, we measure the lepton asymmetry, which is a convolution of the W production charge asymmetry and the V-A couplings in the W decay. The lepton asymmetry is still sensitive to PDFs and is defined as:

$$A(y_l) = \frac{d\sigma_+/dy_l - d\sigma_-/dy_l}{d\sigma_+/dy_l + d\sigma_-/dy_l}. \quad (3)$$

The CDF experiment measured the forward-backward charge asymmetry of electrons from W boson decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using a data sample of 170 pb^{-1} of Run 2 data [2]. Candidate $W \rightarrow e\nu$ events were required to have exactly one electron candidate of $E_T > 25$ GeV, with $\cancel{E}_T > 25$ GeV and transverse mass in the range $50 \text{ GeV}/c^2 < M_T < 100 \text{ GeV}/c^2$.

The CDF experiment improved its sensitivity to the PDFs by reducing the decay asymmetry. The measurement separates the asymmetry measurement into two bins: $25 \text{ GeV} < E_T < 35 \text{ GeV}$ and $35 \text{ GeV} < E_T < 45 \text{ GeV}$ (Figure 1 and 2). This is done for the first time and the separation into different bins is a novel addition to the previous measurements. Predictions from CTEQ [3] and MRST [4] PDFs using a NLO RESBOS calculation [5] are shown for comparison.

The CDF Collaboration is additionally developing a new method to measure the W asymmetry directly by partially reconstructing W rapidity using a W mass.

Z RAPIDITY DISTRIBUTION

Measurement of the rapidity distribution of Drell-Yan pairs in the Z boson mass region provides a stringent test of PDFs. At leading order (LO), the momentum fraction, $x_+(x_-)$, carried by the parton from the proton (anti-proton) is related to the rapidity of the Z boson via the equation:

$$x_{\pm} = \frac{M_Z}{\sqrt{s}} e^{\pm y} \quad (4)$$

where M_Z is the mass of Z boson. Therefore the analysis of the high y region probes the high x as well as low x region.

The DØ experiment measures the rapidity distribution, $d\sigma/dy$, of the Drell-Yan process in the dielectron's mass range between 71 and 111 GeV/c^2 [6]. The data were collected with the DØ detector in Tevatron $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using a data

FIGURE 1. The measured lepton charge asymmetry for $25 \text{ GeV} < E_T < 35 \text{ GeV}$ with predictions from CTEQ6.1M (solid) and MRST02 (dashed) PDFs using a NLO RESBOS calculation. Both statistical and total (statistical+systematic) uncertainties are shown.

FIGURE 2. The measured lepton charge asymmetry for $35 \text{ GeV} < E_T < 45 \text{ GeV}$ with predictions from CTEQ6.1M (solid) and MRST02 (dashed) PDFs using a NLO RESBOS calculation. Both statistical and total (statistical+systematic) uncertainties are shown.

sample of 337 pb^{-1} of Run 2 data. The final result, with statistical and systematic errors, is shown in Figure 3. The measurement of $d\sigma/dy$ is compared with a prediction using an NNLO calculation based on the MRST 2001 PDFs [7]. The DØ forward calorimeters provide data over almost the entire rapidity range accessible at Tevatron. Measurements are in generally good agreement with the prediction.

FIGURE 3. The DØ measurement of the Z rapidity distribution. The outer error bars show the total error (statistical+systematic), while the inner error bars indicate the statistical error alone. The solid line shows the NNLO prediction based on the MRST 2001 PDFs.

CONCLUSIONS

The CDF experiment have measured the W charge asymmetry. Since the previous measurements upon which the current predictions are based are least constraining for $|\eta| > 1$ and do not separate the E_T dependence, inclusion of the new W asymmetry results will further constrain future fits and improve the predictions. The DØ measurement of the Z rapidity distribution is in generally good agreement with theory prediction. We expect that Tevatron Z and W measurements will provide important input to constrain the PDFs, especially in the high x region.

ACKNOWLEDGMENTS

I would like to thank the workshop organizers and working group leaders for their invitation and a wonderful workshop. Also many thanks to the electroweak conveners of both the CDF and DØ Collaborations for providing material and valuable comments.

REFERENCES

1. CDF 2 Collaboration, *CDF Run 2 W mass*, hep-ex/0505064 (2005), hep-ex/0506016 (2005).
2. CDF 2 Collaboration, D. Acosta *et al.*, *Phys. Rev. D* **71**, 051104(R) (2005).
3. J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. Nadolsky, and W. K. Tung, *JHEP* **0207** 012 (2002).
4. A. Martin, R. Roberts, W. Stirling, and R. Thorne, *Eur. Phys. J. C* **4**, 463 (1998).
5. C. Balazs and C.P. Yuan, *Phys. Rev. D*, **56**, 5558 (1997).
6. The DØ Collaboration, <http://www-d0.fnal.gov/Run2Physics/WWW/results/ew.htm>.
7. C. Anastasiou, L. Dixon, K. Melnikov and F. Petriello, *Phys. Rev. D* **69**, 094008 (2004).